

New Streaming Algorithms for High Dimensional EMD and MST

Xi Chen* Rajesh Jayaram† Amit Levi‡ Erik Waingarten§

November 8, 2021

Abstract

We study streaming algorithms for two fundamental geometric problems: computing the cost of a Minimum Spanning Tree (MST) of an n -point set $X \subset \{1, 2, \dots, \Delta\}^d$, and computing the Earth Mover Distance (EMD) between two multi-sets $A, B \subset \{1, 2, \dots, \Delta\}^d$ of size n . We consider the turnstile model, where points can be added and removed. We give a one-pass streaming algorithm for MST and a two-pass streaming algorithm for EMD, both achieving an approximation factor of $\tilde{O}(\log n)$ and using $\text{polylog}(n, d, \Delta)$ -space only. Furthermore, our algorithm for EMD can be compressed to a single pass with a small additive error. Previously, the best known sublinear-space streaming algorithms for either problem achieved an approximation of $O(\min\{\log n, \log(\Delta d)\} \log n)$ [AIK08, BDI⁺20]. For MST, we also prove that any constant space streaming algorithm can only achieve an approximation of $\Omega(\log n)$, analogous to the $\Omega(\log n)$ lower bound for EMD of [AIK08].

Our algorithms are based on an improved analysis of a recursive space partitioning method known generically as the Quadtree. Specifically, we show that the Quadtree achieves an $\tilde{O}(\log n)$ approximation for both EMD and MST, improving on the $O(\min\{\log n, \log(\Delta d)\} \log n)$ approximation of [AIK08, BDI⁺20].

*Columbia University. xichen@cs.columbia.edu. Supported by NSF IIS-1838154 and NSF CCF-1703925.

†Google Research. rkjayaram@google.com.

‡Part of this work was carried out while the author was a PhD student at the University of Waterloo. amit.levi@uwaterloo.ca.

§Stanford University. eaw@cs.columbia.edu. This material is based upon work supported by the National Science Foundation under Award No. 2002201.

Contents

1	Introduction	1
1.1	Our Results	2
1.2	Technical Overview	4
1.2.1	The Main Idea: Tree Embeddings with Data-dependent Edge Weights	4
1.2.2	Implementing Step 1: Quadtree Matching with Data-dependent Edge Weights	6
1.2.3	Implementing Step 2: From Quadtree to Sketching Algorithms	9
2	Preliminaries	13
3	Analysis of Quadtrees for EMD and MST	14
3.1	Approximation of EMD using Quadtrees	14
3.2	Approximation of MST using Quadtrees	18
4	Proof of Lemma 3.6	21
5	Streaming Preliminaries and Sketching Tools	26
5.1	Exponential Order Statistics	26
5.2	Geometric Streaming and Linear Sketching	29
5.3	Count-Sketch, ℓ_1 -sketch, and ℓ_1 -sampling	30
6	Linear Sketches for EMD	31
6.1	Preparation	32
6.2	Universe Reduction	34
6.3	Two-Round Linear Sketch of Lemma 6.6	36
6.4	Overview of the Linear Sketch of Lemma 6.5	36
6.5	Three Linear Sketches and their Performance Gurantees	39
6.6	Linear Sketch of Lemma 6.10	40
6.7	Linear Sketch of Lemma 6.11	42
6.8	Linear Sketch of Lemma 6.12	42
7	Linear Sketches for MST	44
7.1	Proof of Lemma 7.1	47
7.2	Proof of Lemma 7.3	49
7.3	Proof of Lemma 7.4	50
7.4	Proofs of Lemma 7.5 and Lemma 7.6	52
8	Lower Bound for Streaming MST	55

9	Sketching ℓ_p norms for p near 0	58
A	Embedding ℓ_p^d into $\{0, 1\}^d$	60
B	Bit Complexity and Randomness	63

1 Introduction

We study two fundamental geometric problems in high-dimensional spaces: the Earth Mover’s distance and minimum spanning tree. Let $(\mathcal{X}, d_{\mathcal{X}})$ be a metric space. Given two (multi-)sets $A, B \subset \mathcal{X}$ of size $|A| = |B| = n$, the Earth Mover’s distance (EMD) between A and B is

$$\text{EMD}_{\mathcal{X}}(A, B) = \min_{\substack{\text{matching} \\ M \subset A \times B}} \sum_{(a,b) \in M} d_{\mathcal{X}}(a, b).$$

Given a single multi-set $X \subset \mathcal{X}$ of size n , the minimum spanning tree (MST) of X is

$$\text{MST}_{\mathcal{X}}(X) = \min_{\substack{\text{tree } T \\ \text{spanning } X}} \sum_{(a,b) \in T} d_{\mathcal{X}}(a, b).$$

Computational aspects of EMD and MST consistently arise in multiple areas of computer science [RTG00a, HTF01, PC19], such as in computer vision [BVDPPH11, SDGP⁺15], image retrieval [RTG00b], biology [NW70], document similarity [KSKW15], machine learning [ACB17, MJ15, FCCR18], among other areas. Their centrality in both theory and practice has motivated the theoretical study of approximate and sublinear algorithms [Cha02, IT03, Ind04, FIS05, AIK08, ABIW09, HIM12, SA12, MS13, AS14, BI14, ANOY14, YO14, AKR15, BBD⁺17, She17, YV18, KNP19, BDI⁺20] in both low- and high-dimensional settings.

As an illustrative example, an important application for high-dimensional EMD comes from natural language processing, particularly document retrieval and classification. A document can be represented as a collection of vectors in Euclidean space by applying *word embeddings* [MSC⁺13, PSM14] to each of its words; these embeddings have the property that semantically similar words map to geometrically close vectors. In this context, computing the EMD between the embeddings of two documents yields a natural measure of similarity, aptly termed the *Word Mover’s Distance* [KSKW15].

In this paper, we study streaming and sketching algorithms for computing EMD and MST. Specifically, we consider the turnstile *geometric streaming model*, introduced by [Ind04], where the algorithm receives the input set $X \subset \mathcal{X}$ via an arbitrarily ordered sequence of insertions and deletions of points $p \in \mathcal{X}$. The goal is for the algorithm to approximate a fixed function of the implicit set of points X in small space, without storing X ; ideally, one would hope for space *polylogarithmic* in the number of points in $|X|$. We focus on the high-dimensional Euclidean space, where $\mathcal{X} = \{1, 2, \dots, \Delta\}^d$, and the distance between points is given by an ℓ_p norm for $p \in [1, 2]$. One can always reduce from the case of \mathbb{R}^d to $\{1, \dots, \Delta\}^d$ via standard embeddings (see Appendix A).

Prior Work on Sketching and Streaming EMD and MST. We briefly survey what is known for streaming and sketching EMD and MST. We emphasize that many aspects of the sketchability and streamability of EMD and MST remain open, and obtaining tight bounds for these tasks, as well as related geometric graph problems, still remains elusive.¹

Indyk [Ind04], building on work of [Cha02], was the first to formulate dynamic geometric streams and give algorithms for EMD and MST which achieved an $O(d \log \Delta)$ -approximation. The result for

¹See Open Problems 7 and 49 for sketching EMD in <https://sublinear.info/>

MST was improved to a $(1 + \epsilon)$ -approximation in [FIS05], however, the resulting space complexity is exponential in the dimension, making the algorithm suitable only in low-dimensional spaces. For EMD on the plane, [ABIW09] gave a $O(1/\epsilon)$ approximation at the cost of a Δ^ϵ dependence in the space complexity. The best lower bound on sketching EMD on the plane is due to [AKR15], where they show that one cannot have both a constant bit and constant approximation sketch. If the sketch proceeds by an embedding into ℓ_1 , [NS07] show the approximation must be $\Omega(\sqrt{\log \Delta})$. Parametrizing the approximation in terms of n , [BI14] gave embeddings of EMD on the plane into ℓ_1 with distortion $O(\log n)$.

For the high-dimensional regime, Andoni, Indyk, and Krauthgamer [AIK08] gave an algorithm for EMD (in fact, an embedding into ℓ_1) with approximation $O(\log n \log(d\Delta))$. Furthermore, building on an ℓ_1 -embedding lower bound of [KN06], they show that any s -bit sketch with approximation $\alpha > 1$ must have $s\alpha = \Omega(\log n)$. For sketching, the approximation of [AIK08] may be improved to $O(\log n \min\{\log n, \log(d\Delta)\})$ by the techniques in [BI14, BDI⁺20]. MST has not been formally considered in the high-dimensional regime, although we note that an $O(\log n \min\{\log n, \log(d\Delta)\})$ -approximate streaming algorithm readily applies here as well. For lower bounds on streaming high-dimensional MST, nothing was known, and (prior to this work) a constant-bit stream achieving a constant approximation was possible.

1.1 Our Results

In this work, we develop new algorithms and lower bounds for approximating EMD and MST in a stream. Specifically, we show that the approximation factor for these problems can be improved from $O(\log n \cdot \min\{\log n, \log(\Delta d)\})$ to $\tilde{O}(\log n)$. We now state the main results of this paper. In the theorem statements which follow, we consider a fixed setting of n, d and Δ . The metric space consists of points in $[\Delta]^d = \{1, \dots, \Delta\}^d$ with ℓ_p distance for any fixed $p \in [1, 2]$. We state the theorems in the *random-oracle model*, i.e., any random bits stored by the algorithm do not factor into the space complexity — we show that storing the random bits would incur at most an additive $d \cdot \text{polylog}(n, \Delta)$ bits of space (see Section 5.2, where we discuss removing the random oracle assumption).²

Theorem 1 (MST Streaming Algorithm). *There exists a turnstile streaming algorithm using at most $\text{polylog}(n, d, \Delta)$ bits of space which, given a set $X \subset [\Delta]^d$ of size n , outputs $\hat{\eta} \in \mathbb{R}$ satisfying*

$$\text{MST}_{\ell_p}(X) \leq \hat{\eta} \leq \tilde{O}(\log n) \cdot \text{MST}_{\ell_p}(X)$$

with high probability.

For EMD, our algorithm achieving an $\tilde{O}(\log n)$ -approximation requires two passes over the data. This arises from a technical issue in the approach for EMD which is not present in MST. We state the theorem in terms of two-pass streaming algorithms, and then show how to compress the two passes into one, at the cost of an additive error in the approximation.

Theorem 2 (EMD Two-Pass Streaming Algorithm). *Given two multi-sets $A, B \subset [\Delta]^d$ of size n there exists a two-pass turnstile streaming algorithm using $\text{polylog}(n, d, \Delta)$ bits of space which outputs $\hat{\eta} \in \mathbb{R}$ satisfying*

$$\text{EMD}_{\ell_p}(A, B) \leq \hat{\eta} \leq \tilde{O}(\log n) \cdot \text{EMD}_{\ell_p}(A, B)$$

²Also note that to even store a single update $p \in [\Delta]^d$, one requires $\Omega(d \log \Delta)$ bits of space.

with high probability.

Theorem 3 (EMD One-Pass Streaming Algorithm). *Given two multi-sets $A, B \subset [\Delta]^d$ of size n and any $\epsilon > 0$, there exists a turnstile streaming algorithm using $O(1/\epsilon) \cdot \text{polylog}(n, d, \Delta)$ bits of space which outputs $\hat{\eta} \in \mathbb{R}$ satisfying*

$$\text{EMD}_{\ell_p}(A, B) \leq \hat{\eta} \leq \tilde{O}(\log n) \cdot \text{EMD}_{\ell_p}(A, B) + \epsilon d \Delta n.$$

with high probability.

We encourage the reader to think of instances where A and B are size- n subsets of the hypercube $\{0, 1\}^d$ with ℓ_1 distance (i.e., $\Delta = 2$ and $p = 1$). This setting captures all the complexity encountered in this work. For $\Delta > 2$ and $p \in (1, 2]$, the algorithm first applies an embedding into $\{0, 1\}^d$ with ℓ_1 (see Appendix A).

Regarding the additive error in Theorem 3, while an appropriate setting of ϵ may absorb the additive error into relative error, we leave as an open problem whether this additive error may be removed completely in one-pass algorithms. For instance, if the points do not overlap almost always, i.e., when $|A \cap B|/|A \cup B| \leq 1 - \epsilon_0$, then $\text{EMD}_{\ell_p}(A, B) \geq \epsilon_0 n$, and ϵ may be set to $\epsilon_0/d\Delta$ in order to absorb the additive error into the relative error by increasing the space by a factor of $d\Delta$, and keeping a poly-logarithmic dependence on n . From a practical perspective, the fact that points do not overlap may be a reasonable assumption to make.

All of our streaming algorithms are *linear sketches*, meaning that they store only the matrix-vector product $\mathbf{S}f$ for some randomized $\mathbf{S} \in \mathbb{R}^{k \times n}$, where $f = f_X \in \mathbb{R}^{\Delta^d}$ is the indicator vector (with multiplicity) of X for the case of MST, and $f = f_{A,B} \in \mathbb{R}^{2 \cdot \Delta^d}$ is the indicator vector (with multiplicity) of A, B for EMD. Linear sketches are an important class of turnstile streaming algorithms, and have many well-known and studied advantages. For instance, such sketches directly resulted in algorithms for distributed computation such as the MPC model, as well as algorithms for multi-party communication. Our results, therefore, can be applied in a natural way to these models as well.

Improved Analysis of the Quadtree. The prior sketching algorithms are based on a hierarchical partitioning method known as the *Quadtree*.³ Here, we refer to quadtrees as a generic class of methods that embed points from \mathcal{X} into a randomized tree by recursively partitioning the space. At a high level, the Quadtree algorithm recursively and randomly partitions the space \mathcal{X} , which results in a rooted (randomized) tree. Each point in the set X for the case of MST, or $A \cup B$ for EMD, is sent down to a leaf of the tree. From there, a spanning tree or a matching, is constructed in a bottom-up fashion. Each point “walks up the tree” and is greedily connected (in the case of MST), or matched (in the case of EMD) as it encounters other points. This results in a very efficient *offline* (non-sketching) algorithm. The recent work of [BDI⁺20] study the quadtree algorithm explicitly, where they call it “Flowtree,” and showed it has favorable practical properties. From a theoretical point-of-view, the approximation incurred by these methods were the bottleneck in prior works for

³The name *Quadtree* is an artifact of the study of the algorithm originally in the planar (two-dimensional) case, in which the algorithm recursively partitions the plane into quadrants. Our Quadtrees, being in high dimensions, will partition space into more than 4 parts at a time. However, since they are the natural generalization of the planar case, it is common to refer to the generic method as Quadtree regardless of dimension.

sketching and streaming EMD and MST, here, we improve this analysis of [AIK08, BDI⁺20] from $O(\log n \min\{\log n, \log(d\Delta)\})$ to $\tilde{O}(\log n)$.

Theorem 4 (Quadtree Methods (Informal)). *Given two multi-sets $A, B \subset [\Delta]^d$ of size n , the “Flowtree” algorithm of [BDI⁺20] outputs an $\tilde{O}(\log n)$ -approximation to $\text{EMD}_{\ell_1}(A, B)$ with probability at least 0.9. Similarly, given a multi-set $X \subset [\Delta]^d$ of size n , the greedy, bottom-up spanning tree is an $\tilde{O}(\log n)$ -approximation to $\text{MST}_{\ell_1}(X)$ with probability at least 0.9.*

Lower bounds for MST. For lower bounds, [AIK08] shows that any randomized ℓ -bit streaming algorithm distinguishing $\text{EMD}_{\ell_1}(A, B) \geq r$ and $\text{EMD}_{\ell_1}(A, B) \leq r/\alpha$ with probability at least $2/3$ must satisfy $\alpha\ell = \Omega(d)$, where the instances used have $d = \log n$. For a qualitative comparison, estimating ℓ_1 norm does admit such $O(1)$ -approximation, $O(1)$ -bit space streaming algorithms (with public randomness), implying that EMD is a harder problem. We show an analogous lower bound for MST in the streaming model.

Theorem 5. *Any randomized ℓ -bit streaming algorithm which can distinguish whether a size- n set $X \subset \{0, 1\}^d$ has $\text{MST}_{\ell_1}(X) \geq nd/3$ or $\text{MST}_{\ell_1}(X) \leq nd/\alpha$ with probability at least $2/3$ must satisfy $\ell + \log \alpha = \Omega(\log n/\alpha)$. Moreover, this holds even in the insertion-only model, where points are only added to X in the stream.*

We emphasize that, prior to Theorem 5, there were no lower bounds known for streaming MST — not even an $\Omega(1)$ lower bound was known on the approximation of a constant-bit algorithm. We note that [AIK08] actually considers the (stronger) two-party communication setting for EMD, where each player receives one of the sets. The two-party communication game for MST where each player receives half of the set X is insufficient, as there is simple $O(1)$ -approximation, constant-bit protocol.⁴ Therefore, our theorem will crucially involve the streaming nature of the algorithm.

1.2 Technical Overview

1.2.1 The Main Idea: Tree Embeddings with Data-dependent Edge Weights

In [Ind04], Indyk described an approach for streaming a variety of graph problems (including MST and EMD) in discrete geometric spaces, leading to $O(d \log \Delta)$ -approximations for these problems in the metric space $[\Delta]^d$ with ℓ_1 distance. This approach, later refined in [AIK08], forms the basis of our work, so we give a very high level overview in order to highlight the new ideas. For simplicity, we describe it for EMD, as the high-level picture for MST is similar.

A streaming algorithm for EMD with sets $A, B \subset [\Delta]^d$ of size n may proceed in the following way:

1. Sample a recursive random partition of the space, broadly referred to as a quadtree, which specifies an embedding of the original space $[\Delta]^d$ into a rooted tree. For example, when $d = 2$, one may sample $\log_2 \Delta$ randomly shifted, nested square grids of side length $\Delta/2, \Delta/4, \dots, 1$ and arrange them into a rooted tree of depth $\log_2 \Delta + 1$. Each node corresponds to a region

⁴Intuitively, the players may compute the cost of their MST locally, and compute the distance between two arbitrary points. The sum of these quantities is a 3-approximation to the MST of the entire set.

of the space, where the root contains the entire space, and the children of a node have regions which partition the region of the parent. The points in A and B are assigned to leaves of this tree, according to the regions where points fall, and the quadtree implicitly defines a matching M between A and B given by the natural bottom-up greedy procedure. Having implicitly specified a matching M , the goal of the streaming algorithm will be to approximate the cost of M .

2. In order to do so, [Ind04, AIK08] maintains a high-dimensional vector which implicitly encodes the matching M . Specifically, the vector has a coordinate for each edge of the quadtree, and the entry in each coordinate is the number of points from A falling within the region of the child minus the number of points in B falling within the region of the child. Furthermore, the ℓ_1 -norm of the vector, where each coordinate of an edge is scaled by some edge weight (for example, by the size of the parent region) gives an approximation of the cost of M . Thus, this gives an ℓ_1 -embedding for EMD over $[\Delta]^d$, and known algorithms for streaming the ℓ_1 -norm can be applied.

With the above approach in mind, there are two steps involved in showing the approximation guarantee: (i) showing the matching M in Step 1 has approximately optimal cost, and (ii) showing that the appropriate scalings of coordinates reduce approximating the cost of the matching M to an ℓ_1 -computation. We note that even though the above presentation is a two-step procedure, [Ind04, AIK08] do not present it this way. In fact, de-coupling the matching M from the method to approximate the cost of M is an important conceptual contribution which is made explicit in [BDI⁺20], which led us to revisit the EMD problem.

Prior to our work, (i) proceeded by the method of tree embeddings. One assigns the edge weights to the quadtree and interprets it as a tree embedding of the metric $([\Delta]^d, \ell_1)$. By studying the distortion of this embedding, one bounds the cost of M . The edge weights chosen in [Ind04] (building on work of [Cha02]) embed $([\Delta]^d, \ell_1)$ with distortion $O(d \log \Delta)$, which will become the approximation. Refining the approach, [AIK08] show that another choice of edge weights (better suited for high-dimensional spaces) embeds subsets of $([\Delta]^d, \ell_1)$ with bounded average distortion which suffices for an $O(\log n \log(d\Delta))$ bound on the cost of M . Given the bound on M with respect to a fixed tree metric, (ii) is straight-forward: since the fixed tree metric specifies the scalings of the vector, and approximating the cost of M amounts to an ℓ_1 -norm computation.

Our main contribution is two-fold. First, we show how to go beyond the distortion argument in (i) to show that the cost of M is a $\tilde{\Theta}(\log n)$ -approximation to EMD with probability 0.9. To do so, we study a *data-dependent* notion: instead of fixing the edge weights as in [Ind04, AIK08], we allow the edge weights to depend on the input $A \cup B$. The use of data-dependent edge weights implies M is actually a better quality matching than what the method of tree embeddings specified. The data-dependent edge weights are (relatively) simple: the weight of an edge (u, v) , where u is the parent of v , is the average distance between a randomly sampled point of $A \cup B$ within the region of u and a randomly sampled point of $A \cup B$ within the region of v . However, the fact these data-dependent edge weights yield an improved upper bound on the cost of M constitutes the bulk of the work in Sections 3 and 4.

Unfortunately, the introduction of data-dependent edge weights breaks Step 2. Now, approximating the cost of M with the data-dependent weights is no longer as simple as an ℓ_1 -computation. The coordinates of the vector remain the same, however, the scaling of each coordinate depends on

additional structure of the points. Importantly, data-dependent edge weights do not result in an ℓ_1 -embedding, and we cannot use known ℓ_1 -sketching algorithms. This takes us to our algorithmic contribution, where we design the sketching algorithms for Step 2 with data-dependent edge weights. More generally, we introduce a two-step template for transforming data-dependent costs in the Quadtree into streaming algorithms. Conceptually, the approach generalizes the well-known ℓ_p sampling problem [MW10, AKO10, JST11, JW21] to ℓ_p -sampling with meta-data. For EMD, the high-level idea is the following: first, sample a coordinate of the vector proportional to the ℓ_1 -distribution (i.e., the ℓ_1 -sampling problem), and second, estimate the data-dependent edge weight for the coordinate sampled (the meta-data), so that we can scale the contribution of that coordinate appropriately.

1.2.2 Implementing Step 1: Quadtree Matching with Data-dependent Edge Weights

We begin by describing our improved analysis of the randomized space partitioning algorithm, *Quadtree*. For the sake of simplicity, we focus on its analysis in the context of approximating EMD; the same ideas work similarly for MST. We begin by more formally introducing the Quadtree in high-dimensional spaces. In what follows, we focus on the case when the metric space is the hypercube with the Hamming distance, i.e. $A, B \subset \{0, 1\}^d$ and $d(p, q) = \|p - q\|_1$ for $p, q \in \{0, 1\}^d$. For the approximation, this is without loss of generality. One may embed (\mathbb{R}^d, ℓ_p) into $\{0, 1\}^d$ by increasing the dimension (see Appendix A), which is irrelevant since the approximation we will show is dimension-independent.

Quadtree. The Quadtree algorithm creates a randomized tree \mathbf{T} with depth $h := \log_2 2d$ by recursively sub-dividing the hypercube $\{0, 1\}^d$. Therefore, each node u in \mathbf{T} will be associated with a subcube $S_u \subseteq \{0, 1\}^d$, where the root r has $S_r = \{0, 1\}^d$. To create these subcubes, each internal node u of \mathbf{T} at depth $j < h$ is labeled with an ordered tuple of 2^j coordinates $(i_1, \dots, i_{2^j}) \in [d]$ (which are not necessarily distinct), and has 2^{2^j} children. Each of the 2^{2^j} children of u will *uniquely* correspond to one of the 2^{2^j} fixings of the coordinates $(i_1, \dots, i_{2^j}) \in \{0, 1\}^{2^j}$. Specifically, each child v of u is assigned a unique bit-string $(b_1, \dots, b_{2^j}) \in \{0, 1\}^{2^j}$. The child v then corresponds to the subcube $S_v \subseteq S_u$ obtained by fixing the i_t -th coordinate to b_t , for each $t = 1, \dots, 2^j$. We now describe the procedure for generating a random Quadtree \mathbf{T} :

1. Uniformly sampling a tuple $(i_1, \dots, i_{2^j}) \in [d]^{2^j}$ of 2^j coordinates independently for each node u at depth $j \in \{0, 1, \dots, h - 2\}$ to use as its label.
2. Setting $(1, \dots, d)$ as the label of every node at depth $h - 1$.

A Quadtree \mathbf{T} defines a map φ from $\{0, 1\}^d$ to leaves of \mathbf{T} : $\varphi(p) = v$ if $p \in S_v$. Given A and B , we write A_v and B_v to denote $A \cap S_v$ and $B \cap S_v$ for each node v in \mathbf{T} .

Depth-greedy Matching from Quadtree. Given a random Quadtree \mathbf{T} , one obtains a natural depth-greedy matching as follows: We first map all points in $C = A \cup B$ to leaves of \mathbf{T} using φ . Then, we greedily match points between A and B in a bottom up fashion, by walking each point up the tree level-by-level, and at each node one arbitrarily matches as many of the unmatched points from A and B as possible. Let \mathbf{M} be any depth-greedy matching obtained from \mathbf{T} in this fashion.

The goal of our improved analysis of the Quadtree for EMD is to show that

$$\text{EMD}(A, B) \leq \mathbf{Cost}(\mathbf{M}) \stackrel{\text{def}}{=} \sum_{(a,b) \in \mathbf{M}} \|a - b\|_1 \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B)$$

with high probability (over the randomness of \mathbf{T}). Note that the first inequality is trivial.

Analysis of Quadtree via Tree Embeddings. Before presenting an overview of our new techniques, it will be helpful to begin with a recap of the analysis of [AIK08] which can be used to show that $\mathbf{Cost}(\mathbf{M}) \leq O(\log n \log d) \cdot \text{EMD}(A, B)$. The analysis of [AIK08] starts by assigning a weight of $d/2^i$ to each edge from a node at depth i to a node at depth $i + 1$ in \mathbf{T} . This defines a metric embedding $\varphi : A \cup B \rightarrow \mathbf{T}$ by mapping each point to a leaf of \mathbf{T} . The choice of edge weights is motivated by the observation that two points $x, y \in \{0, 1\}^d$ with $\|x - y\|_1 = d/2^i$ are expected to have their paths diverge for the first time at depth i . If this is indeed the case then $d_{\mathbf{T}}(\varphi(x), \varphi(y))$ would capture $\|x - y\|_1$ up to a constant.

To upperbound $\mathbf{Cost}(\mathbf{M})$, one studies the distortion of this embedding. Firstly, for any $\lambda > 1$ and $x, y \in \{0, 1\}^d$, it is easy to verify that distances in the tree metric do not contract much:

$$\Pr_{\mathbf{T}} \left[d_{\mathbf{T}}(\varphi(x), \varphi(y)) < \frac{1}{\lambda} \cdot \|x - y\|_1 \right] \leq \left(1 - \frac{\|x - y\|_1}{d} \right)^{1+2+\dots+2^{\lceil \log_2 \left(\frac{\lambda d}{\|x - y\|_1} \right) \rceil}} \leq 2^{-\Omega(\lambda)}.$$

Thus by a union bound, for all $x, y \in A \cup B$ we have

$$\|x - y\|_1 \leq O(\log n) \cdot d_{\mathbf{T}}(\varphi(x), \varphi(y)) \tag{1}$$

with probability at least $1 - 1/\text{poly}(n)$, which essentially means that we can assume (1) in the worst case. As a result, we have

$$\mathbf{Cost}(\mathbf{M}) = \sum_{(x,y) \in \mathbf{M}} \|x - y\|_1 \leq O(\log n) \sum_{(x,y) \in \mathbf{M}} d_{\mathbf{T}}(\varphi(x), \varphi(y)) \leq O(\log n) \sum_{(x,y) \in M^*} d_{\mathbf{T}}(\varphi(x), \varphi(y)),$$

where the last inequality holds for any matching M^* between A and B given that the depth-greedy matching is optimal under the tree metric. Setting M^* to be the optimal matching between A and B under the original ℓ_1 metric, we finish the proof by upperbounding $d_{\mathbf{T}}(\varphi(x), \varphi(y))$ using $O(\log d) \|x - y\|_1$. To see this, when $\|x - y\|_1 = \Theta(d/2^j)$, the probability that paths of x, y diverge at level $j - k$ is $\Theta(2^{-k})$ for each k , and when it does, $d_{\mathbf{T}}(\varphi(x), \varphi(y)) = \|x - y\|_1 \cdot \Theta(2^k)$. Since $j \leq h = O(\log d)$,

$$\mathbf{E}[d_{\mathbf{T}}(\varphi(x), \varphi(y))] \leq \|x - y\|_1 + \sum_{k=0}^j \Theta(2^{-k}) \cdot \|x - y\|_1 \cdot \Theta(2^k) = O(\log d) \cdot \|x - y\|_1. \tag{2}$$

Together they yield the aforementioned $O(\log n \log d) \cdot \text{EMD}(A, B)$ upper bound for $\mathbf{Cost}(\mathbf{M})$.⁵

Tree Embeddings with Data-dependent Edge Weights. We show how to go beyond the distortion arguments of [AIK08] by studying a tree embeddings with data-dependent edge weights. In what follows, for any vertex $u \in \mathbf{T}$, let $C_u = A_u \cup B_u$ be the set of all points which map through

⁵The reason that this analysis can achieve approximation $O(\min\{\log n, \log d\} \log n)$, as opposed to $O(\log n \log d)$ is that with probability $1 - 1/n$, every $x, y \in A \cup B$ with $\|x - y\|_1 = \Theta(d/2^j)$ diverges at depth after $j - O(\log n)$.

u (recall $C = A \cup B$). The weight we assign to each edge (u, v) ⁶ of \mathbf{T} will no longer be a fixed number $d/2^i$ but

$$\text{avg}_{u,v} \stackrel{\text{def}}{=} \mathbf{E}_{\substack{\mathbf{c} \sim C_u \\ \mathbf{c}' \sim C_v}} [\|\mathbf{c} - \mathbf{c}'\|_1],$$

i.e., the average distance between a point drawn randomly from C_u and a point drawn randomly from C_v ; when $C_v = \emptyset$ we define $\text{avg}_{u,v} = 0$ by default. Let $d_{\mathbf{T}}^*$ denote the tree metric under this new set of weights. Again, the depth-greedy matching \mathbf{M} we are interested in is optimal and the cost of \mathbf{M} under the new tree embedding can be expressed as

$$\mathbf{Value}_{\mathbf{T}}(A, B) \stackrel{\text{def}}{=} \sum_{(u,v) \in E_{\mathbf{T}}} \left| |A_u| - |B_v| \right| \cdot \text{avg}_{u,v}.$$

where $E_{\mathbf{T}}$ is the set of edges of T .⁷ On the one hand, $\mathbf{Value}_{\mathbf{T}}(A, B)$ is at least $\mathbf{Cost}(\mathbf{M})$ given that for any $x, y \in C$, we always have $\|x - y\|_1 \leq d_{\mathbf{T}}^*(\varphi(x), \varphi(y))$ by triangle inequality. On the other hand, $\mathbf{Value}_{\mathbf{T}}(A, B)$ is at most $\sum_{(a,b) \in M^*} d_{\mathbf{T}}^*(\varphi(a), \varphi(b))$ for any matching M^* and in particular, the optimal matching M^* under the ℓ_1 metric. As a result, it suffices to upperbound the cost of M^* under the data-dependent tree embedding by $\tilde{O}(\log n) \cdot \mathbf{Cost}(M^*)$ given that $\mathbf{Cost}(M^*) = \text{EMD}(A, B)$. To this end it suffices to show that the expectation of $d_{\mathbf{T}}^*(\varphi(a), \varphi(b))$ for any $a, b \in C$ can be bounded from above by $\tilde{O}(\log n) \cdot \|a - b\|_1$.

Inspector Payment. Fix $a, b \in C$. We introduce the following quantity as the *inspector payment* of (a, b) with respect to the Quadtree \mathbf{T} . (We imagine the process as first drawing the Quadtree and then an “inspector” who examines the tree to track down a and b , making payments accordingly.) Formally we let $(v_0(x), v_1(x), \dots, v_h(x))$ denote the root-to-leaf path of x in a Quadtree \mathbf{T} . Then

$$\mathbf{Pay}_{\mathbf{T}}(a, b) \stackrel{\text{def}}{=} \sum_{i \in [h]} \mathbf{1}\{v_i(a) \neq v_i(b)\} \cdot \left(\mathbf{E}_{\mathbf{c} \sim C_{v_{i-1}(a)}} [\|a - \mathbf{c}\|_1] + \mathbf{E}_{\mathbf{c} \sim C_{v_{i-1}(b)}} [\|b - \mathbf{c}\|_1] \right). \quad (3)$$

Intuitively, this payment scheme corresponds to an inspector who tracks down a and b from the root of T , and whenever a and b first diverge in the tree at node u , pays for a the average distance between a and a random point drawn from C_v for every node along the u -to-leaf path (including u); the inspector pays for b similarly. It again follows from triangle inequality that $2 \cdot \mathbf{Pay}_{\mathbf{T}}(a, b)$ is at least $d_{\mathbf{T}}^*(\varphi(a), \varphi(b))$. So it suffices to bound the expectation of $\mathbf{Pay}_{\mathbf{T}}(a, b)$ by $\tilde{O}(\log n) \cdot \|a - b\|_1$.

Before giving a sketch of this proof, which is the most challenging part of our Quadtree analysis, we note that the inspector payment (3) depends on the data A and B , as well as the Quadtree \mathbf{T} in two ways. The first is the depth when a and b first diverge, captured by the indicator $\mathbf{1}\{v_i(a) \neq v_i(b)\}$. The second is the average distance between a and C_v , which not only depends on a , but also on global properties of $C = A \cup B$. At a high level, incorporating this second aspect is the main novelty, since the average distance between a and C_v is an *average* notion of radii at v . Therefore, if the inspector pays a large amount, then an average point in C_v is far from a (as opposed to the farthest point implied by worst-case radii).

⁶We always use u in (u, v) to denote the parent and v to denote the child.

⁷We remark that one can define an analogous quantity for the case of MST, where given a single set $X \subset [\Delta]^d$, we set $\mathbf{Value}_{\mathbf{T}}(X) = \sum_{(u,v) \in E_{\mathbf{T}}} \mathbf{1}(|X_v|) \cdot \text{avg}_{u,v}$, where $\mathbf{1} : \mathbb{R} \rightarrow \{0, 1\}$ is the indicator function (i.e., $\mathbf{1}(x) = 0$ if and only if $x = 0$). It is this quantity that we will analyze in our results for MST.

Bounding Inspector Payments. Consider fixed $a, b \in C$ at distance $\|a - b\|_1 = \Theta(d/2^j)$, and we give some intuition behind our upper bound on the expectation of the a -part of the payment:

$$\sum_{i \in [h]} \mathbf{1}\{v_i(a) \neq v_i(b)\} \cdot \text{avg}_{a, i-1}, \quad \text{where} \quad \text{avg}_{a, i-1} \stackrel{\text{def}}{=} \mathbf{E}_{\mathbf{c} \sim C_{v_{i-1}(a)}} [\|a - \mathbf{c}\|_1].$$

We will ignore the indicator random variable $\mathbf{1}\{v_i(a) \neq v_i(b)\}$ and use linearity of expectation to focus on $\mathbf{E}_{\mathbf{T}}[\text{avg}_{a, i}]$. (With the indicator random variable, we need to consider the expectation of $\text{avg}_{a, i}$ conditioning on the event that a, b have diverged. The conditioning will not heavily influence the geometric intuition, so we will ignore this for the rest of this overview).

Let $v_i = v_i(a)$. Similar to worst-case bounds on radii, $\mathbf{E}_{\mathbf{T}}[\text{avg}_{a, i}]$ can still be $d/2^i \cdot \Omega(\log n)$. As an example, let i_1 be a relatively large depth and for some small $\epsilon \approx 10^{-6}$, consider a set P_1 of n^ϵ many points at distance $\epsilon \log n \cdot d/2^{i_1}$ around a . Then, at depth i_1 of a random Quadtree \mathbf{T} , a point in P_1 traverses down to node v_{i_1} with non-negligible probability, roughly $1/n^{-\epsilon}$. If no other points lie closer to a than those in P_1 , then $\mathbf{E}_{\mathbf{T}}[\text{avg}_{a, i_1}] = d/2^{i_1} \cdot \Omega(\epsilon \log n)$, since it is likely that some points of P_1 make it to v_{i_1} and significantly increase the average distance between a and $C_{v_{i_1}}$. If this happened on a for every depth i , the inspector would be in trouble, as there are $O(\log d)$ levels and a similar argument to that of worst-case radii would mean a payment of $O(\log d \log n) \cdot \|a - b\|_1$.

However, we claim if the arrangement of P_1 resulted in $\mathbf{E}_{\mathbf{T}}[\text{avg}_{a, i_1}] = d/2^{i_1} \cdot \Omega(\epsilon \log n)$, the same situation will be a lot more difficult to orchestrate for depth $i_2 \leq i_1 - O(\log \log n)$. In particular, at depth i_2 , in order to have $\mathbf{E}_{\mathbf{T}}[\text{avg}_{a, i_2}] = d/2^{i_2} \cdot \Omega(\epsilon \log n)$, there must be a set of points P_2 at distance $d/2^{i_2} \cdot \Omega(\epsilon \log n)$ which cause avg_{a, i_2} to be large. However, it is no longer enough to have $|P_2| = n^\epsilon$. The reason is that points of P_1 in v_{i_2} will help bring down the average distance. Since points in P_1 are at distance $\epsilon \log n \cdot d/2^{i_1} \ll d/2^{i_2}$ from a , there will oftentimes be $\Omega(n^\epsilon)$ points from P_1 in v_{i_2} . In order to significantly increase the average distance, v_{i_2} must oftentimes have at least $n^\epsilon / \text{polylog}(n)$ points from P_2 ; otherwise, avg_{a, i_2} will be mostly the average distance between a and points in P_1 . Since any given point from P_2 traverses down to v_{i_2} with probability roughly $1/n^\epsilon$, we must have $|P_2| \geq n^{2\epsilon} / \text{polylog}(n)$. This argument can only proceed for at most $O(1/\epsilon)$ depths before $|P_{O(1/\epsilon)}| > 2n$, in which case we obtain a contradiction, since all points are in $A \cup B$.

Generally, in order to increase the average distance between a and C_{v_i} *multiple times* as the depth i goes down, the number of points around a at increasing distances must grow very rapidly. More specifically, we show that if a depth i is “bad,” meaning that $\mathbf{E}_{\mathbf{T}}[\text{avg}_{a, i}] \geq \alpha \cdot d/2^i$ for some $\alpha = \omega(\log \log n)$, then the number of points within a ball of radius $d/(2^i \log n)$ around a and within a larger ball of radius $O(\log n \cdot d/2^i)$ around a must have increased by a factor of $\exp(\Omega(\alpha))$; this means the number of such depths i is at most $((\log n)/\alpha) \cdot \text{poly}(\log \log n)$. Combining this analysis and the fact that a and b must diverge in order to incur payment from the inspector, we obtain our upper bound that the expectation of $\text{Pay}_{\mathbf{T}}(a, b)$ is at most $\tilde{O}(\log n) \cdot \|a - b\|_1$.

1.2.3 Implementing Step 2: From Quadtree to Sketching Algorithms

By the prior discussion, after sampling a Quadtree \mathbf{T} , we know that the quantity $\text{Value}_{\mathbf{T}}(A, B)$ is a $\tilde{O}(\log n)$ approximation of the true cost $\text{EMD}(A, B)$. Specifically, we have:

$$\text{EMD}(A, B) \leq \text{Value}_{\mathbf{T}}(A, B) \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B) \tag{4}$$

Thus, the approach of our sketching algorithm is simply to approximate $\mathbf{Value}_{\mathbf{T}}(A, B)$. We will decompose $\mathbf{Value}_{\mathbf{T}}(A, B)$ based on its level: $\mathbf{Value}_{\mathbf{T}}(A, B) = \sum_{i=1}^h \mathbf{Value}_{\mathbf{T},i}(A, B)$, where

$$\mathbf{Value}_{\mathbf{T},i}(A, B) \stackrel{\text{def}}{=} \sum_{\substack{(u,v) \in E_{\mathbf{T}} \\ \mathbf{depth}(u,v)=i}} \left| |A_v| - |B_v| \right| \cdot \text{avg}_{u,v}$$

where $\mathbf{depth}(e)$ for an edge $e \in \mathbf{T}$ is the depth of the child vertex in e . We will attempt to estimate each $\mathbf{Value}_{\mathbf{T},i}(A, B)$ independently for each i , so in what follows we now fix any level $i \in [h]$.

We start with some notation. For any (non-root) vertex $v \in \mathbf{T}$, let $\pi(v)$ be the parent of v in \mathbf{T} . We then define the *discrepancy* vector for level i , denoted Δ^i , by $\Delta_v^i = |A_v| - |B_v|$ for every vertex v at depth i of the tree (i.e., Δ^i has a coordinate Δ_v^i for each vertex v at depth i). Next, for any vector $x \in \mathbb{R}^N$ and any $p \geq 0$, we define the ℓ_p distribution $\mathcal{D}_p(x)$ over the coordinates of x via $\mathcal{D}_p(x) = \left(\frac{|x_1|^p}{\|x\|_p^p}, \frac{|x_2|^p}{\|x\|_p^p}, \dots, \frac{|x_N|^p}{\|x\|_p^p} \right)$ for $p > 0$, and for $p = 0$ we define $\mathcal{D}_0(x)$ to be the uniform distribution over the support of x . Now observe:⁸

$$\mathbf{Value}_{\mathbf{T},i}(A, B) = \|\Delta^i\|_1 \cdot \mathbf{E}_{\mathbf{v} \sim \mathcal{D}_1(\Delta^i)} \left[\text{avg}_{\pi(\mathbf{v}), \mathbf{v}} \right]$$

Thus, we can write $\mathbf{Value}_{\mathbf{T},i}(A, B)$ as the ℓ_1 norm of Δ^i , multiplied by the expected value of $\text{avg}_{\pi(\mathbf{v}), \mathbf{v}}$ taken over drawing a vertex \mathbf{v} in level i with probability proportional to $|\Delta_{\mathbf{v}}| = \left| |A_{\mathbf{v}}| - |B_{\mathbf{v}}| \right|$. Note that the norm $\|\Delta^i\|_1$ can be easily estimated using the ℓ_1 sketches of Indyk [Ind06a]. Thus, this simple manipulation motivates the following approach: **(1)** sample a vertex \mathbf{v} from level i from the distribution $\mathcal{D}_1(\Delta^i)$, **(2)** recover the value $\text{avg}_{\pi(\mathbf{v}), \mathbf{v}}$, **(3)** repeat enough times so that the empirical mean of the variables $\text{avg}_{\pi(\mathbf{v}), \mathbf{v}}$ is a good approximation of the expectation $\mathbf{E}_{\mathbf{v} \sim \mathcal{D}_1(\Delta^i)} \left[\text{avg}_{\pi(\mathbf{v}), \mathbf{v}} \right]$.

For the last step, we note that it will be straightforward to bound the standard deviation of the variable $\text{avg}_{\pi(\mathbf{v}), \mathbf{v}}$ by $O(d \log n / 2^i)$, which is within a $O(\log n)$ factor of the error to which we will need to estimate the expectation. Thus, if we can carry out steps **(1)** and **(2)** which sample $\text{avg}_{\pi(\mathbf{v}), \mathbf{v}}$ from the correct distribution, we need only repeat them $\text{polylog}(n)$ times to estimate $\mathbf{Value}_{\mathbf{T},i}(A, B)$ to sufficiently small error.

Two-Pass Streaming Algorithms. We first describe how the above two steps can be carried out in *two-passes* over the data-stream. Perhaps unsurprisingly, our approach will be to carry out **(1)** on the first pass, obtaining a set of vertices v sampled from the correct distribution $\mathcal{D}_1(\Delta^i)$, and carry out **(2)** on the second pass, where we recover the actual value of $\text{avg}_{\pi(v), v}$ for the vertices v that were sampled.

More formally, our two-pass streaming algorithm proceeds as follows. First we draw a Quadtree \mathbf{T} (for which we may assume (4) holds) and then for each $i \in [h]$, we estimate $\mathbf{Value}_{\mathbf{T},i}(A, B)$ as follows. In the first pass we can estimate $\|\Delta^i\|_1$ to error $(1 \pm 1/2)$ with an ℓ_1 -sketch [Ind06b], and we also can sample $\mathbf{v} \sim \mathcal{D}_1(\Delta^i)$ via known algorithms for ℓ_1 -sampling [AKO10, JST11, JW21]. Furthermore,

⁸We remark that for the case of MST, the relevant quantity $\mathbf{Value}_{\mathbf{T},i}(X)$ below can be written as $\|\Delta^i\|_0 \cdot \mathbf{E}_{v \sim \mathcal{D}_0(\Delta^i)} \left[\text{avg}_{\pi(v), v} \right]$. Namely, we simply replace the ℓ_1 norm in both the scaling and the distribution by the ℓ_0 norm. Thus, the high-level approach to sketching MST will be similar. However, due to using the ℓ_0 instead of the ℓ_1 norm, an entirely different set of techniques will be required to implement each of the steps.

once a vertex v is fixed, we may estimate $\text{avg}_{\pi(v), \mathbf{v}}$ in the second round by a point in $C_{\pi(v)}$ and in C_v (via standard sub-sampling techniques) and approximating their distance using ℓ_1 sketches. By concurrently repeating this process $\text{polylog}(n)$ times, we obtain our desired approximation

The remaining challenge, however, is to produce $\mathbf{v} \sim \mathcal{D}_1(\Delta^i)$ and an estimate of $\text{avg}_{\pi(v), \mathbf{v}}$ *simultaneously* in a single pass over the data. This task is a special case of a problem we call *sampling with meta-data*, since the quantity $\text{avg}_{\pi(v), \mathbf{v}}$ will be the *meta-data* of the sample $\mathbf{v} \sim \mathcal{D}_1(\Delta^i)$ needed to estimate $\text{Value}_{\mathbf{T}, i}(A, B)$.

Sampling with Meta-Data and One-Pass Streaming The key task of sampling with meta-data is the following: for $n, k \in \mathbb{N}$, we are given a vector $x \in \mathbb{R}^n$ and collection of *meta-data* vectors $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{R}^k$, and the goal is to sample $i \in [n]$ with probability $|x_i|/\|x\|_1$ (or more generally, $|x_i|^p/\|x\|_p^p$), and output both i and an approximation $\hat{\lambda}_i \in \mathbb{R}^k$ of the vector λ_i . The challenge is to solve this problem with a small-space linear sketches of x and the meta-data vectors $\lambda_1, \dots, \lambda_n$. It is not hard to see that sampling with meta-data is exactly the problem we seek to solve for linear sketching of EMD.⁹

Our algorithm builds on a powerful sketching technique known as *precision sampling* [AKO10, JST11, JW21] for sampling an index $i \in [n]$ proportional to $|x_i|/\|x\|_1$ for a vector $x \in \mathbb{R}^n$ (or more generally, for $|x_i|^p/\|x\|_p^p$, but we focus on $p = 1$). The idea is to produce, for each $i \in [n]$ an independent exponential random variable $\mathbf{t}_i \sim \text{Exp}(1)$, and construct a “scaled vector” $\mathbf{z} \in \mathbb{R}^n$ with coordinates $\mathbf{z}_i = x_i/\mathbf{t}_i$. One then attempts to return the index $i_{\max} = \text{argmax}_{i \in [n]} \mathbf{z}_i$, since

$$\Pr_{\mathbf{t}_1, \dots, \mathbf{t}_n \sim \text{Exp}(1)} \left[\text{argmax}_{i' \in [n]} \frac{|x_{i'}|}{\mathbf{t}_{i'}} = i \right] = \frac{|x_i|}{\|x\|_1}.$$

To find the the index i_{\max} with a linear sketch, we can use a “heavy-hitters” algorithm, such as the Count-Sketch of [CCFC02].¹⁰ Specifically, Count-Sketch with error $\epsilon \in (0, 1)$ allows us to recover an estimate $\tilde{\mathbf{z}}$ to \mathbf{z} satisfying (roughly) $\|\tilde{\mathbf{z}} - \mathbf{z}\|_\infty \leq \epsilon \|\mathbf{z}\|_2$. Then one can show that $\text{argmax}_{i' \in [n]} |\tilde{\mathbf{z}}_{i'}|$ is close to being distributed as $|x_i|/\|x\|_1$.

In order to sample with meta-data, our sketch similarly samples independent exponential $\mathbf{t}_1, \dots, \mathbf{t}_n \sim \text{Exp}(1)$ and applies a Count-Sketch data structure on $\mathbf{z} \in \mathbb{R}^n$, where $\mathbf{z}_i = x_i/\mathbf{t}_i$, and obtains an estimate $\tilde{\mathbf{z}}$ of \mathbf{z} . In addition, we apply a Count-Sketch data structure with error ϵ for the vector \mathbf{w} with coordinates given by the values λ_i/\mathbf{t}_i , namely $\mathbf{w}_i = \lambda_i/\mathbf{t}_i$ (recall that we are assuming that the meta-data λ_i are scalars for this discussion). From this we obtain an estimate $\tilde{\mathbf{w}}$ of \mathbf{w} . The insight is the following: suppose the sample produced is $i^* \in [n]$, which means it satisfies $\tilde{\mathbf{z}}_{i^*} \approx \max_{i \in [n]} |x_i|/\mathbf{t}_i$. Then the value \mathbf{t}_{i^*} should be relatively small: in particular, one can show that we expect \mathbf{t}_{i^*} to be $\Theta(|x_{i^*}|/\|x\|_1)$, so that $\mathbf{z}_{i^*} \approx \Theta(\|x\|_1) = \Theta(\|\Delta^i\|_1)$. When this occurs, for each $\ell \in [k]$, the guarantees of Count-Sketch imply that the estimate $\mathbf{t}_{i^*} \cdot \tilde{\mathbf{w}}_{i^*}^\ell$ satisfies

$$|\mathbf{t}_{i^*} \cdot \tilde{\mathbf{w}}_{i^*} - \lambda_{i^*}| = \mathbf{t}_{i^*} |\tilde{\mathbf{w}}_{i^*} - \mathbf{w}_{i^*}| \leq \epsilon \mathbf{t}_{i^*} \|\mathbf{w}\|_2 \left(= O \left(\epsilon |x_{i^*}| \cdot \frac{\|\lambda\|_1}{\|x\|_1} \right) \text{ in expectation} \right)$$

⁹Namely, x is the vector Δ^i , and the meta-data vectors λ_v are $k = \text{polylog}(n)$ -dimensional ℓ_1 sketches of the values of $\text{avg}_{\pi(v), v}$. In the following discussion, for simplicity we omit the details on the ℓ_1 sketches for $\text{avg}_{\pi(v), v}$, since they proceed via somewhat standard techniques, and instead assume that the meta-data is exactly given by the scalars $\lambda_v \approx \text{avg}_{\pi(v), v}$.

¹⁰We do not explicitly use count-sketch in our one-pass algorithms, and instead apply a sketching procedure closely inspired by Count-Sketch.

where $\lambda \in \mathbb{R}^n$ is the vector with coordinates given by the meta-data $\lambda_1, \dots, \lambda_n$. In other words, if the size of λ_{i^*} is comparable to $|x_{i^*}|$, and if the ratio $\|\lambda\|_1/\|x\|_1$ of the meta-data norms to the norm of x is bounded, then $\mathbf{t}_{i^*}\tilde{\mathbf{w}}_{i^*}^\ell$ is a relatively good approximation to λ_{i^*} .

Unfortunately, in our application, the above will not always be the case. In particular, the norm of the meta-data $\|\lambda\|_1$ may be much, even $\text{poly}(n)$, larger than $\|x\|_1 = \|\Delta^i\|_1$. Intuitively, the issue is that each coordinate λ_v is a sketch of $\text{avg}_{\pi(v),v}$, which is a function both of the points in $C_{\pi(v)}$ and C_v . Thus, the size of the sketch of $\text{avg}_{\pi(v),v}$ depends on all the points in $C_{\pi(v)}$. Moreover, for *every* other sibling v' of v (meaning that $\pi(v') = \pi(v)$), the sketch of $\text{avg}_{\pi(v'),v'}$ will also have to take into account the same information from $C_{\pi(v)}$. Thus, this information is *duplicated* in the sketches of the meta-data, by a number of times equal to the number of children of $\pi(v)$. This duplication, or repetition of the same information in the sketch, results in a blow-up of the norm of λ so that $\|\lambda\|_1 = \Omega(\kappa \cdot \|\Delta^i\|_1)$, where κ is the maximum number of non-empty children of any parent in level $i - 1$. Since κ can be $\text{poly}(n)$, this is a non-trivial challenge.

Our solution to this, at a high level, is to develop a *two-step* precision sampling with meta-data algorithm to avoid duplication of meta-data. Instead of sampling the vertex \mathbf{v}^* directly, we first sample a parent \mathbf{u}^* from level $i - 1$ with probability proportional to the ℓ_1 -norm of Δ^i restricted to coordinates corresponding to the children of \mathbf{u}^* ; namely, we sample \mathbf{u}^* with probability proportional to $\sum_{v:\pi(v)=\mathbf{u}^*} |\Delta_v^i|$. Then, we use the precision sampling sketch which recovered \mathbf{u}^* to recover a sketch of the a randomly selected point in $C_{\mathbf{u}^*}$. Next, once we have \mathbf{u}^* , we apply precision sampling with meta-data once more, to sample a child \mathbf{v}^* of \mathbf{u}^* proportional to $|\Delta_{\mathbf{v}^*}^i|$, and then recover a sketch of a randomly selected point in $C_{\mathbf{v}^*}$. One can then put the two sketches from $C_{\mathbf{u}^*}, C_{\mathbf{v}^*}$ together to estimate $\text{avg}_{\mathbf{u}^*, \mathbf{v}^*}$.

To accomplish this two-part precision sampling scheme, we must generate a second set of exponentials $\{\mathbf{t}_v\}_v$, one for each child node v at depth i . In order to ensure that the sample produced by the second sketch actually returns a child \mathbf{v}^* of \mathbf{u}^* , and not a child of some other node, we crucially must scale the vector Δ^i by *both* the child exponentials $\{\mathbf{t}_v\}_v$ as well as the parent exponentials $\{\mathbf{t}_u\}_u$ from the first sketch. Thus, in the second sketch we analyze the twice-scale vector \mathbf{z} with coordinates $\mathbf{z}_v = \Delta_v^i/(\mathbf{t}_{\pi(v)}\mathbf{t}_v)$, and attempt to find the largest coordinate of \mathbf{z} . Importantly, notice that this makes the scaling factors in \mathbf{z}_v no longer independent: two children of the same parent share one of their scaling factors. Thus, executing this plan requires a careful analysis of the behavior of norms of vectors scaled by several non-independent variables with heavy-tailed distributions.

The advantage of this two-part scheme is that now there is no duplication of meta-data, since in the first step there is only one λ_u for each parent u , and in the second step, by conditioning on the parent exponential $\mathbf{t}_{\mathbf{u}^*}$ being sufficiently small, we ensure that the only meta-data that contributes non-trivially to the error of the sketch are the λ_v for children v of \mathbf{u}^* . This allows us, ultimately, to obtain our guarantees for one-pass streaming algorithms for EMD. The case of MST is similar at a high-level, however implementing the two-part precision sampling scheme requires an entirely different set of sketching tools, resulting from the fact that we now need to sample a vertex v from the ℓ_0 distribution $\mathcal{D}_0(\Delta^i)$.

2 Preliminaries

Given $n \geq 1$ we write $[n]$ to denote $\{1, \dots, n\}$. Given a vector $x \in \mathbb{R}^n$ and a real number $t \geq 0$, we define $x_{\leq t} \in \mathbb{R}^n$ to be the vector obtained by setting the largest $\lfloor t \rfloor$ coordinates of x in magnitude equal to 0 (breaking ties by using coordinates with smaller indices). For $a, b \in \mathbb{R}$ and $\epsilon \in (0, 1)$, we use the notation $a = (1 \pm \epsilon)b$ to denote the containment of $a \in [(1 - \epsilon)b, (1 + \epsilon)b]$.

For convenience, we will assume without loss of generality that d is always a power of 2 and write $h := \log_2 2d = \log_2 d + 1$. Given a node v in a rooted tree T , when v is not the root we use $\pi(v)$ to denote the parent node of v in T .

Next we give a formal definition of Quadrees used in this paper:

Definition 2.1 (Quadrees). *Fix $d \in \mathbb{N}$. A quadtree is a rooted tree T of depth $h := \log_2 2d$. We say a node v of T is at depth j if there are $j + 1$ nodes on the root-to- v path in T (so the root is at depth 0 and its leaves are at depth h). Each internal node v of T at depth $j < h$ is labelled with an ordered tuple of 2^j coordinates $i_1, \dots, i_{2^j} \in [d]$ (which are not necessarily distinct), and has 2^{2^j} children, each of which we refer to as the (b_1, \dots, b_{2^j}) -child of v with $b_1, \dots, b_{2^j} \in \{0, 1\}$. Every node at depth $h - 1$ is labelled with $(1, \dots, d)$. We write E_T to denote the edge set of T . Whenever we refer to an edge $(u, v) \in E_T$, u is always the parent and v is the child. A random quadtree \mathbf{T} is drawn by (1) sampling a tuple of 2^j coordinates uniformly and independently from $[d]$ for each node at depth $j < h - 1$ as its label; and (2) use $(1, 2, \dots, d)$ as the label of every node at depth $h - 1$. We use \mathcal{T} to denote this distribution of random quadrees.*

Given a quadtree T , each point $x \in \{0, 1\}^d$ induces a root-to-leaf path by starting at the root and repeatedly going down the tree as follows: If the current node v is at depth $j < h$ and is labelled with (i_1, \dots, i_{2^j}) , then we go down to the $(x_{i_1}, \dots, x_{i_{2^j}})$ -child of v . We write

$$\mathbf{v}_{0,T}(x), \mathbf{v}_{1,T}(x), \dots, \mathbf{v}_{h,T}(x)$$

to denote this root-to-leaf path, where each $\mathbf{v}_{j,T}$ is a map from $\{0, 1\}^d$ to nodes of T at depth j . We usually drop T from the subscript when it is clear from the context.

Alternatively we define a subcube $S_{v,T} \subseteq \{0, 1\}^d$ for each v : The set of the root is $\{0, 1\}^d$; If (u, v) is an edge, u is at depth j and is labelled with i_1, \dots, i_{2^j} , and v is the (b_1, \dots, b_{2^j}) -child of u , then

$$S_{v,T} = \{x \in S_{u,T} : (x_{i_1}, \dots, x_{i_{2^j}}) = (b_1, \dots, b_{2^j})\}.$$

Note that $S_{v,T}$'s of nodes v at the same depth form a partition of $\{0, 1\}^n$. The root-to-leaf path for $x \in \{0, 1\}^d$ can be equivalently defined as the sequence of nodes v that have $x \in S_{v,T}$.

Remark 6. *Both works of [AIK08, BDI⁺20] use a tree structure that is very similar to the Quadtree used in this paper. In particular, They consider a slightly different algorithm which at depth i , samples 2^i coordinates from $[d]$ and divides into 2^{2^i} branches according to settings of $\{0, 1\}$ to these 2^i coordinates (instead of each vertex independently sampling 2^i coordinates). For the sake of the analysis in Section 3, there will be no difference between independently sampling coordinates for each vertex in a level, and using the same sampled coordinates for each level. Thus, our analysis apply to trees of [AIK08, BDI⁺20] as well as the Quadrees defined here.*

3 Analysis of Quadrees for EMD and MST

Our goal in this section is to obtain expressions based on quadrees that are good approximations of EMD and MST. They will serve as the starting point of our sketches for EMD and MST later.

3.1 Approximation of EMD using Quadrees

Fix $n, d \in \mathbb{N}$ and let T be a quadtree of depth $h = \log_2 2d$. Let A and B be two multisets of points from $\{0, 1\}^d$ of size n each. For each node v in T , we define

$$A_{v,T} \stackrel{\text{def}}{=} \{a \in A : v_{i,T}(a) = v\} \quad \text{and} \quad B_{v,T} \stackrel{\text{def}}{=} \{b \in B : v_{i,T}(b) = v\}.$$

Equivalently we have $A_{v,T} = A \cap S_{v,T}$ and $B_{v,T} = B \cap S_{v,T}$. Let $C_{v,T} = A_{v,T} \cup B_{v,T}$. We give the definition of depth-greedy matchings.

Definition 3.1. *Let T be a quadtree. For any $a \in A$ and $b \in B$, let*

$$\text{depth}_T(a, b) \stackrel{\text{def}}{=} \text{depth of the least-common ancestor of leaves of } a, b \text{ in } T.$$

The class of depth-greedy matchings, denoted by $\mathcal{M}_T(A, B)$, is the set of all matchings $M \subseteq A \times B$ which maximize the sum of $\text{depth}_T(a, b)$ over all pairs $(a, b) \in M$. We write

$$\mathbf{Cost}(M) = \sum_{(a,b) \in M} \|a - b\|_1$$

to denote the cost of a matching M between A and B . Recall that $\text{EMD}(A, B)$ is defined as the minimum of $\mathbf{Cost}(M)$ over all matchings between A and B .

For each edge $(u, v) \in E_T$, we use $\text{avg}_{u,v,T}$ to denote the average distance between points of $C_{u,T}$ and $C_{v,T}$:

$$\text{avg}_{u,v,T} \stackrel{\text{def}}{=} \mathbf{E}_{\substack{\mathbf{c} \sim C_{u,T} \\ \mathbf{c}' \sim C_{v,T}}} [\|\mathbf{c} - \mathbf{c}'\|_1],$$

where both \mathbf{c} and \mathbf{c}' are drawn uniformly at random; we set $\text{avg}_{u,v,T}$ to be 0 by default when $C_{v,T}$ is empty. For notational simplicity, we will suppress T from the subscript when it is clear from the context. We are now ready to define the *value* of (A, B) in a quadtree T :

Definition 3.2. *Let T be a quadtree. The value of (A, B) in T is defined as*

$$\mathbf{Value}_T(A, B) \stackrel{\text{def}}{=} \sum_{(u,v) \in E_T} \left| |A_v| - |B_v| \right| \cdot \text{avg}_{u,v}. \tag{5}$$

We note that the right-hand side of (5) is data-dependent in two respects: the discrepancy between $|A_v|$ and $|B_v|$ and the average distance $\text{avg}_{u,v}$ between points in C_u and C_v .

Our main lemma for EMD shows that the value of (A, B) in a randomly chosen quadtree $\mathbf{T} \sim \mathcal{T}$ and the cost of any depth-greedy matching are all $\tilde{O}(\log n)$ -approximations to $\text{EMD}(A, B)$.

Lemma 3.3 (Quadtree lemma for EMD). *Let (A, B) be a pair of multisets of points from $\{0, 1\}^d$ of size n each. Let $\mathbf{T} \sim \mathcal{T}$. Then with probability at least 0.99, every $M \in \mathcal{M}_{\mathbf{T}}(A, B)$ satisfies we have*

$$\text{EMD}(A, B) \leq \mathbf{Cost}(M) \leq \mathbf{Value}_{\mathbf{T}}(A, B) \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B). \quad (6)$$

We start with the first inequality in (6). Indeed we will show that $\text{EMD}(A, B) \leq \mathbf{Value}_T(A, B)$ for any quadtree T (Lemma 3.4). To this end we prove that $\mathbf{Cost}(M) \leq \mathbf{Value}_T(A, B)$ for any depth-greedy matching between A and B obtained from T ; the latter by definition is at least $\text{EMD}(A, B)$.

Lemma 3.4. *Let T be any quadtree. Then $\mathbf{Cost}(M) \leq \mathbf{Value}_T(A, B)$ for any $M \in \mathcal{M}_T(A, B)$.*

Proof: Given an $M \in \mathcal{M}_T(A, B)$ and a pair $(a, b) \in M$, we write v and w to denote the leaves of a and b and use $v = u_1, u_2, \dots, u_k = w$ to denote the path from v to w in T . By triangle inequality,

$$\begin{aligned} \|a - b\|_1 &\leq \mathbf{E}_{\mathbf{c}_i \sim C_{u_i}} \left[\|a - \mathbf{c}_1\|_1 + \|\mathbf{c}_1 - \mathbf{c}_2\|_1 + \dots + \|\mathbf{c}_{k-1} - \mathbf{c}_k\|_1 + \|\mathbf{c}_k - b\|_1 \right] \\ &= \text{avg}_{u_1, u_2} + \dots + \text{avg}_{u_{k-1}, u_k}, \end{aligned}$$

where the equation follows from the fact the label of every node at depth $h-1$ is $(1, 2, \dots, d)$ and thus, all points at a leaf must be identical. Summing up these inequalities over all $(a, b) \in M$ gives exactly $\mathbf{Value}_T(A, B)$ on the right hand side. For this, observe that every M in $\mathcal{M}_T(A, B)$ has the property that, for any edge (u, v) in T , the number of $(a, b) \in M$ such that the path between their leaves contains (u, v) is exactly $\|A_v\| - \|B_v\|$. \blacksquare

Now it suffices to upperbound $\mathbf{Value}_{\mathbf{T}}(A, B)$ by $\tilde{O}(\log n) \cdot \text{EMD}(A, B)$ with probability at least 0.9 for a random quadtree $\mathbf{T} \sim \mathcal{T}$. For this purpose we let $C = A \cup B$ and define an *inspector payment* for any pair of points $a, b \in C$ ¹¹ based on a quadtree. Given $a, b \in C$, we let

$$\mathbf{Pay}_T(a, b) \stackrel{\text{def}}{=} \sum_{i \in [h]} \mathbf{1}\{v_i(a) \neq v_i(b)\} \cdot \left(\text{avg}_{a, i-1} + \text{avg}_{b, i-1} \right) \quad (7)$$

where

$$\text{avg}_{a, i-1} \stackrel{\text{def}}{=} \mathbf{E}_{\mathbf{c} \sim C_{v_{i-1}(a)}} [\|a - \mathbf{c}\|_1] \quad \text{and} \quad \text{avg}_{b, i-1} \stackrel{\text{def}}{=} \mathbf{E}_{\mathbf{c} \sim C_{v_{i-1}(b)}} [\|b - \mathbf{c}\|_1].$$

Intuitively $\mathbf{Pay}_T(a, b)$ pays for the average distance between a (or b) and points in $C_{v_i(a)}$ (or $C_{v_i(b)}$) along its root-to-leaf path but the payment only starts *at* the least-common ancestor of leaves of a and b . Note that $\mathbf{Pay}_T(a, b) = 0$ trivially if $a = b$.

We show that for any matching M between A and B , the total inspector payment from $(a, b) \in M$ is enough to cover $\mathbf{Value}_T(A, B)$:

Lemma 3.5. *Let T be any quadtree and M be any matching between A and B . Then we have*

$$\mathbf{Value}_T(A, B) \leq 2 \sum_{(a, b) \in M} \mathbf{Pay}_T(a, b). \quad (8)$$

¹¹While we will always have $a \in A$ and $b \in B$ in this subsection, this more general setting allows us to apply what we prove in this subsection to work on MST later.

Proof: Using the definition of $\mathbf{Value}_T(A, B)$, it suffices to show that

$$\sum_{(u,v) \in E_T} \left| |A_v| - |B_v| \right| \cdot \text{avg}_{u,v} \leq 2 \sum_{(a,b) \in M} \mathbf{Pay}_T(a, b).$$

By triangle inequality (and $\text{avg}_{a,h} = 0$ because every point in $C_{v_h(a)}$ is identical to a)

$$\begin{aligned} 2 \cdot \mathbf{Pay}_T(a, b) &\geq \sum_{i \in [h]} \mathbf{1}\{v_i(a) \neq v_i(b)\} \cdot \left(\text{avg}_{a,i-1} + \text{avg}_{a,i} + \text{avg}_{b,i-1} + \text{avg}_{b,i} \right) \\ &\geq \sum_{i \in [h]} \mathbf{1}\{v_i(a) \neq v_i(b)\} \cdot \left(\text{avg}_{v_{i-1}(a), v_i(a)} + \text{avg}_{v_{i-1}(b), v_i(b)} \right), \end{aligned}$$

i.e., $2 \cdot \mathbf{Pay}_T(a, b)$ is enough to cover $\text{avg}_{u,v}$ for every edge (u, v) along the path between the leaf of u and the leaf of v . The lemma then follows from the following claim: For every edge (u, v) in T , $\left| |A_v| - |B_v| \right|$ is at most the number of points $a \in A_v$ such that its matched point in M is not in B_v plus the number of points $b \in B_v$ such that its matched point in M is not in A_v . This follows from the simple fact that every $(a, b) \in M$ with $a \in A_v$ and $b \in B_v$ would get cancelled in $|A_v| - |B_v|$. This finishes the proof of the lemma. \blacksquare

By Lemma 3.5 the goal now is to upperbound the total inspector payment by $\tilde{O}(\log n) \cdot \text{EMD}(A, B)$ with probability at least 0.9 over a randomly picked quadtree \mathbf{T} . We consider a slight modification of the payment scheme given in (7) which we define next; the purpose is that the latter will be easier to bound in expectation, and most often exactly equal to (7).

Specifically, given any (a, b) with $a, b \in C$ and $i_0 \in [0 : h - 1]$, we let

$$\mathbf{Pay}_{i_0, T}^*(a, b) \stackrel{\text{def}}{=} \sum_{i > i_0}^h \mathbf{1}\{v_i(a) \neq v_i(b)\} \cdot \left(\text{avg}_{a,i-1}^* + \text{avg}_{b,i-1}^* \right), \quad (9)$$

where

$$\text{avg}_{a,i}^* \stackrel{\text{def}}{=} \mathbf{E}_{\mathbf{c} \sim C_{a,i}^*} [\|a - \mathbf{c}\|_1] \quad \text{and} \quad \text{avg}_{b,i}^* \stackrel{\text{def}}{=} \mathbf{E}_{\mathbf{c} \sim C_{b,i}^*} [\|b - \mathbf{c}\|_1]$$

and $C_{a,i}^*$ contains all points in $C_{v_i(a)}$ that is not too far away from a :

$$C_{a,i}^* \stackrel{\text{def}}{=} \left\{ c \in C_{v_i(a)} : \|a - c\|_1 \leq \frac{10d \log n}{2^i} \right\}.$$

The set $C_{b,i}^*$ is defined similarly. Roughly speaking, points in C that share the same node at depth i are expected to have distance around $d/2^i$ (given they have agreed on $2^i - 1$ random coordinates sampled so far); this is why we refer to points in $C_{a,i}^*$ as those that are not too far away from a .

The following is the crucial lemma for upperbounding the total expected payment according to an optimal matching M^* . We delay its proof to Section 4 and first use it to prove Lemma 3.3.

Lemma 3.6. *For any (a, b) with $a, b \in C$, $a \neq b$ and $i_0 \in [0 : h - 1]$ that satisfies*

$$i_0 \leq h_{a,b} \stackrel{\text{def}}{=} \left\lceil \log_2 \left(\frac{d}{\|a - b\|_1} \right) \right\rceil, \quad (10)$$

we have

$$\mathbf{E}_{\mathbf{T} \sim \mathcal{T}} \left[\mathbf{Pay}_{i_0, \mathbf{T}}^*(a, b) \right] \leq \left(\tilde{O}(\log n) + O(\log \log n) (h_{a,b} - i_0) \right) \cdot \|a - b\|_1.$$

Proof of Lemma 3.3 assuming Lemma 3.6: Let M^* be an optimal matching between A and B that achieves $\text{EMD}(A, B)$. Let $\mathbf{T} \sim \mathcal{T}$. Then we have from Lemma 3.5 that

$$\mathbf{Value}_{\mathbf{T}}(A, B) \leq 2 \sum_{\substack{(a,b) \in M^* \\ a \neq b}} \mathbf{Pay}_{\mathbf{T}}(a, b) \quad (11)$$

given that $\mathbf{Pay}_{\mathbf{T}}(a, b) = 0$ when $a = b$. Below we focus on the subset M' of M^* with $(a, b) \in M^*$ and $a \neq b$. For each $(a, b) \in M'$, let

$$0 \leq \ell_{a,b} \stackrel{\text{def}}{=} \max \{0, h_{a,b} - 2\lceil \log_2 n \rceil\} \leq h_{a,b}.$$

We show that with probability at least $1 - o(1)$ over the draw of \mathbf{T} , every $(a, b) \in M'$ satisfies

$$\mathbf{Pay}_{\mathbf{T}}(a, b) = \mathbf{Pay}_{\ell_{a,b}, \mathbf{T}}^*(a, b). \quad (12)$$

Combining (11) and (12), we have that with probability at least $1 - o(1)$ over the draw of \mathbf{T} ,

$$\mathbf{Value}_{\mathbf{T}}(A, B) \leq 2 \sum_{(a,b) \in M'} \mathbf{Pay}_{\ell_{a,b}, \mathbf{T}}^*(a, b). \quad (13)$$

By applying Lemma 3.6 to every $(a, b) \in M'$ with $i_0 = \ell_{a,b}$, as well as Markov's inequality, we have that with probability at least 0.99 over \mathbf{T} , the right hand side of (13) is at most

$$\tilde{O}(\log n) \sum_{(a,b) \in M'} \|a - b\|_1 = \tilde{O}(\log n) \cdot \text{EMD}(A, B).$$

By a union bound, $\mathbf{Value}_{\mathbf{T}}(A, B) \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B)$ with probability at least $.99 - o(1) \geq 0.9$.

It suffices to define an event that implies (12) and then bound its probability. The first part of the event requires that for every pair $(a, b) \in M'$, $\mathbf{v}_i(a) = \mathbf{v}_i(b)$ for every $i : 1 \leq i \leq \ell_{a,b}$. The second part requires that for any two distinct points $x, y \in A \cup B$ (not necessarily as a pair in M^* and not even necessarily in the same set), we have $\mathbf{v}_i(x) \neq \mathbf{v}_i(y)$ for all i with

$$2^i \geq \frac{10d \log n}{\|x - y\|_1}. \quad (14)$$

By the definition of $\mathbf{Pay}_{\ell_{a,b}, \mathbf{T}}^*(a, b)$ in (9), the first part of the event makes sure that we don't miss any term in the sum; the second part of the event makes sure that every $C_{a,i}^*$ is exactly the same as $C_{\mathbf{v}_i(a)}$ so that $\text{avg}_{a,i}^* = \text{avg}_{a,i}$ (and the same holds for b). It follows that this event implies (12).

Finally we show that the event occurs with probability at least $1 - o(1)$. First, for every $(a, b) \in M'$, if $\ell_{a,b} = 0$ then the first part of the event trivially holds. If $\ell_{a,b} > 0$ then $\ell_{a,b} = h_{a,b} - 2\lceil \log n \rceil$. The probability of $\mathbf{v}_i(a) \neq \mathbf{v}_i(b)$ for some $i : 1 \leq i \leq \ell_{a,b}$ is at most

$$1 - \left(1 - \frac{\|a - b\|_1}{d}\right)^{2^{\ell_{a,b}-1}} \leq 2^{\ell_{a,b}} \cdot \frac{\|a - b\|_1}{d} \leq \frac{1}{n^2}.$$

Hence, by a union bound over the at most n pairs $(a, b) \in M'$, the first part of the event holds with probability at least $1 - o(1)$. Furthermore, for any two distinct points $x, y \in A \cup B$, let

$$\ell^* = \left\lceil \log_2 \left(\frac{10d \log n}{\|x - y\|_1} \right) \right\rceil.$$

Then $v_i(x) = v_i(y)$ for some i that satisfies (14) would imply $v_{\ell^*}(x) = v_{\ell^*}(y)$ and $\ell^* \leq \log d$ (since $v_h(x) \neq v_h(y)$ given $x \neq y$). The event above happens with probability

$$\left(1 - \frac{\|x - y\|_1}{d}\right)^{2^{\ell^* - 1}} \leq \exp(-5 \log n) = \frac{1}{n^5}.$$

Via a union bound over at most $(2n)^2$ many pairs of x, y , we have that the second part of the event also happens with probability at least $1 - o(1)$. This finishes the proof of the lemma. \blacksquare

3.2 Approximation of MST using Quadrees

We will follow a similar strategy as we took in the previous subsection for EMD. Given a quadtree T of depth $h = \log_2 2d$, we define similarly $v_0(x), \dots, v_h(x)$ as the root-to-leaf path of $x \in \{0, 1\}^d$, and write S_v for each node v at depth i to denote the set of $x \in \{0, 1\}^d$ with $v_i(x) = v$.

Let $X \subseteq \{0, 1\}^d$ be a set of n points. We define X_v for each node v in T as $X \cap S_v$, and write L_i for each depth i to denote the set of nodes v at depth i such that $X_v \neq \emptyset$ and will refer to them as nonempty nodes.

We give the definition of depth-greedy spanning trees.

Definition 3.7. *Let T be a quadtree, and $X \subset \{0, 1\}^d$. For any DFS walk of the quadtree T starting at the root, let $\sigma: [n] \rightarrow X$ denote the order of points in X encountered during the walk, so that $v_h(\sigma(i))$ appears before $v_h(\sigma(i+1))$ for every $i \in [n-1]$. A depth-greedy spanning tree G obtained from a DFS walk is given by the edges $\{(\sigma(i), \sigma(i+1))\}_{i \in [n-1]}$. The class of depth-greedy spanning trees, denoted by $\mathcal{G}_T(X)$, is the set of all spanning trees G of X obtained from a DFS walk down the quadtree T . For any spanning tree G , we write*

$$\mathbf{Cost}(G) = \sum_{(a,b) \in E(G)} \|a - b\|_1$$

to denote the cost of a tree G (with $n-1$ edges) spanning points in X . Recall $\mathbf{MST}(X)$ is defined as the minimum of $\mathbf{Cost}(G)$ over all spanning trees G of X .

Similar to the previous subsection, for each edge $(u, v) \in E_T$, we write

$$\text{avg}_{u,v} \stackrel{\text{def}}{=} \mathbf{E}_{\substack{\mathbf{c} \sim X_u \\ \mathbf{c}' \sim X_v}} [\|\mathbf{c} - \mathbf{c}'\|_1].$$

when $X_v \neq \emptyset$, and $\text{avg}_{u,v} = 0$ when $X_v = \emptyset$. Recall $\pi(v)$ denotes the parent node of v in T . We are now ready to define the value of X in a quadtree T and then state the main lemma:

Definition 3.8. *Let T be a quadtree. The value of X in T is defined as*

$$\mathbf{Value}_T(X) \stackrel{\text{def}}{=} \sum_{i \in [h]} \mathbf{1}\{|L_i| > 1\} \cdot \sum_{v \in L_i} \text{avg}_{\pi(v), v}.$$

The main lemma for MST shows that the value of X for a random quadtree $\mathbf{T} \sim \mathcal{T}$ and the cost of any depth-greedy spanning tree $G \in \mathcal{G}_T(X)$ are $\tilde{O}(\log n)$ -approximations of $\mathbf{MST}(X)$.

Lemma 3.9 (Quadtrees lemma for MST). *Let $X \subseteq \{0, 1\}^d$ be a set of size n , and let $\mathbf{T} \sim \mathcal{T}$. Then with probability at least 0.99, for any $G \in \mathcal{G}_{\mathbf{T}}(X)$, we have that*

$$\frac{\text{MST}(X)}{2} \leq \frac{\text{Cost}(G)}{2} \leq \text{Value}_{\mathbf{T}}(X) \leq \tilde{O}(\log n) \cdot \text{MST}(X).$$

We start with the lower bound:

Lemma 3.10. *Let T be any quadtree and any depth-greedy spanning tree $G \in \mathcal{G}_T(X)$. Then $\text{Value}_T(X) \geq \text{Cost}(G)/2 \geq \text{MST}(X)/2$.*

Proof: Let w be the least common ancestor of leaves $\mathbf{v}_h(x)$, $x \in X$, and let T^* denote the subtree rooted at w that consists of paths from w to $\mathbf{v}_h(x)$, $x \in X$. Using T^* we can equivalently write

$$\text{Value}_T(X) = \sum_{(u,v) \in E_{T^*}} \text{avg}_{u,v}.$$

For each node $v \in T^*$ (note that $X_v \neq \emptyset$), we define ρ_v to be the center-of-mass of points in X_v :

$$\rho_v \stackrel{\text{def}}{=} \frac{1}{|X_v|} \sum_{x \in X_v} x.$$

By triangle inequality we have $\|\rho_u - \rho_v\|_1 \leq \text{avg}_{u,v}$ for every $(u, v) \in E_{T^*}$ and thus,

$$\sum_{(u,v) \in E_{T^*}} \|\rho_u - \rho_v\|_1 \leq \text{Value}_T(X).$$

We finish the proof by showing that any depth-greedy spanning tree G of X satisfies

$$\text{Cost}(G) \leq 2 \sum_{(u,v) \in E_{T^*}} \|\rho_u - \rho_v\|_1.$$

To this end we take a DFS walk of T^* from its root w and let $\sigma: [n] \rightarrow X$ be the order of points in X under which $\mathbf{v}_h(\sigma(1)), \dots, \mathbf{v}_h(\sigma(n))$ appear in the walk. Then we set G to be the spanning tree $\{(\sigma(i), \sigma(i+1))\}_{i \in [n-1]}$. For each $i \in [n-1]$, letting u_1, \dots, u_r be the part of DFS walk from $u_1 = \mathbf{v}_h(\sigma(i))$ to $u_r = \mathbf{v}_h(\sigma(i+1))$, we have from triangle inequality that

$$\|\sigma(i) - \sigma(i+1)\|_1 = \|\rho_{u_1} - \rho_{u_r}\|_1 \leq \|\rho_{u_1} - \rho_{u_2}\|_1 + \dots + \|\rho_{u_{r-1}} - \rho_{u_r}\|_1.$$

The lemma follows from the fact that a DFS walk visits each edge twice. ■

Now it suffices to upper bound $\text{Value}_{\mathbf{T}}(X)$ by $\tilde{O}(\log n) \cdot \text{MST}(X)$ with probability at least 0.9 for a random quadtree $\mathbf{T} \sim \mathcal{T}$. For this purpose, we use the same inspector payment defined in the last subsection (the only change is that the set C is now called X which is a set and has size n instead of $2n$). Recall that for any two points $x, y \in X$, we define

$$\text{Pay}_T(x, y) \stackrel{\text{def}}{=} \sum_{i \in [h]} \mathbf{1}\{\mathbf{v}_i(x) \neq \mathbf{v}_i(y)\} \cdot (\text{avg}_{x, i-1} + \text{avg}_{y, i-1}), \quad (15)$$

where

$$\text{avg}_{x, i-1} \stackrel{\text{def}}{=} \mathbf{E}_{\mathbf{c} \sim X_{\mathbf{v}_{i-1}(x)}} [\|x - \mathbf{c}\|_1] \quad \text{and} \quad \text{avg}_{y, i-1} \stackrel{\text{def}}{=} \mathbf{E}_{\mathbf{c} \sim X_{\mathbf{v}_{i-1}(y)}} [\|y - \mathbf{c}\|_1].$$

Next we show that the total payment from any spanning tree G is enough to cover $\text{Value}_T(X)$.

Lemma 3.11. *Let T be any quadtree and G be any spanning tree of X . Then we have*

$$\mathbf{Value}_T(X) \leq 2 \sum_{(x,y) \in E(G)} \mathbf{Pay}_T(x,y). \quad (16)$$

Proof: Let w be the least common ancestor of leaves $v_h(x)$, $x \in X$, and let T^* denote the subtree rooted at w that consists of paths from w to $v_h(x)$, $x \in X$. It suffices to show that

$$\sum_{(u,v) \in E_{T^*}} \text{avg}_{u,v} \leq 2 \sum_{(x,y) \in E(G)} \mathbf{Pay}_T(x,y).$$

By similar arguments in the proof of Lemma 3.5, $2 \cdot \mathbf{Pay}_T(x,y)$ is good enough to cover $\text{avg}_{u,v}$ for every edge along the path between the leaf of x and the leaf of y . The lemma follows by summing over all $(x,y) \in E(G)$ and noting that the $\text{avg}_{u,v}$ of each $(u,v) \in E_{T^*}$ is counted at least once. ■

To upperbound the total inspector payment from an optimal spanning tree by $\tilde{O}(\log n) \cdot \text{MST}(X)$, we similarly consider the modified payment scheme $\mathbf{Pay}_{i_0, T}^*(x,y)$ as in (9), replacing C by X . The same Lemma 3.6 applies and we use it to prove Lemma 3.9:

Proof of Lemma 3.9 assuming Lemma 3.6: The lower bound follows from Lemma 3.10. For the upper bound, let G^* be an optimal spanning tree of X and let $\mathbf{T} \sim \mathcal{T}$. By Lemma 3.11 we have

$$\mathbf{Value}_{\mathbf{T}}(X) \leq 2 \sum_{(x,y) \in E'(G^*)} \mathbf{Pay}_{\mathbf{T}}(x,y), \quad (17)$$

where $E'(G^*)$ denotes the set of edges (x,y) in G^* with $x \neq y$. For each $(x,y) \in E'(G^*)$, let

$$0 \leq \ell_{x,y} \stackrel{\text{def}}{=} \max \{0, h_{x,y} - 2\lceil \log_2 n \rceil\} \leq h_{x,y}.$$

By similar arguments as in the proof of Lemma 3.3, we have with probability at least $1 - o(1)$ over the draw of \mathbf{T} that every $(x,y) \in E'(G^*)$ satisfies

$$\mathbf{Pay}_{\mathbf{T}}(x,y) = \mathbf{Pay}_{\ell_{x,y}, \mathbf{T}}^*(x,y) \quad (18)$$

Combining (17) and (18), we have that with probability at least $1 - o(1)$ over the draw of \mathbf{T} ,

$$\mathbf{Value}_{\mathbf{T}}(X) \leq 2 \sum_{(x,y) \in E'(G^*)} \mathbf{Pay}_{\ell_{x,y}, \mathbf{T}}^*(x,y). \quad (19)$$

By applying Lemma 3.6 to every $(x,y) \in E'(G^*)$ with $i_0 = \ell_{x,y}$, as well as Markov's inequality, we have that with probability at least 0.99 over \mathbf{T} , the right hand side of (19) is at most

$$\tilde{O}(\log n) \sum_{(x,y) \in E'(G^*)} \|x - y\|_1 = \tilde{O}(\log n) \cdot \text{MST}(X).$$

By a union bound, $\mathbf{Value}_{\mathbf{T}}(X) \leq \tilde{O}(\log n) \cdot \text{MST}(X)$ with probability at least $.99 - o(1) \geq 0.9$. ■

4 Proof of Lemma 3.6

Recall $h = \log 2d$ is the depth of a quadtree. Let $(a, b) \in M^*$ with $a \neq b$ and $i_0 \in [0 : h - 1]$ with

$$i_0 \leq h_{a,b} \stackrel{\text{def}}{=} \left\lceil \log_2 \left(\frac{d}{\|a - b\|_1} \right) \right\rceil. \quad (20)$$

Our goal is to bound the expectation of the a -part of $\mathbf{Pay}_{i_0, \mathbf{T}}^*(a, b)$:

$$\sum_{i > i_0}^h \mathbf{1}\{\mathbf{v}_i(a) \neq \mathbf{v}_i(b)\} \cdot \text{avg}_{a,i-1}^* \quad (21)$$

over the draw of $\mathbf{T} \sim \mathcal{T}$; the bound for the b part is analogous. In what follows, all expectations are taken with respect to $\mathbf{T} \sim \mathcal{T}$ so we will skip \mathbf{T} in subscripts. In particular, we write $\mathbf{v}_i(a)$ to denote $\mathbf{v}_{i, \mathbf{T}}(a)$ just to emphasize that it is a random variable that depends on \mathbf{T} .

Given that we always have $\text{avg}_{a,i}^* \leq 10d \log n / 2^i$ for every i by definition, (21) is trivially $O(d \log n)$ and thus, the statement holds trivially when $\|a - b\|_1 = \Omega(d)$. So we assume in the rest of the proof that $\|a - b\|_1 \leq d/2$. To understand (21) for \mathbf{T} , we examine the sequence of sets $C_{\mathbf{v}_i(a)}$, where $\mathbf{v}_0(a), \dots, \mathbf{v}_h(a)$ is the root-to-leaf path of a in \mathbf{T} . Recall from the definition of \mathcal{T} that to draw $A_{\mathbf{v}_i(a)}$ and $B_{\mathbf{v}_i(a)}$, it suffices to consider independent draws of tuples

$$\mathbf{I}_0, \mathbf{I}_1, \dots, \mathbf{I}_{h-2}$$

with $\mathbf{I}_i \sim [d]^{2^i}$ uniformly, and then use them to define $C_{\mathbf{v}_i(a)}$ as follows:

$$C_{\mathbf{v}_i(a)} = \{x \in C : x_j = a_j \text{ for all } j \text{ that appears in } \mathbf{I}_0, \dots, \mathbf{I}_{i-1}\}$$

for each $i = 1, \dots, h - 1$; $C_{\mathbf{v}_0(a)} = C$ since $\mathbf{v}_0(a)$ is always the root; $C_{\mathbf{v}_h(a)}$ contains all copies of a in C . Next we let

$$\mathbf{D} = \{(i, \ell) : i \in \{0, \dots, h - 2\} \text{ and } \ell \in [2^i] \text{ such that } j = (\mathbf{I}_i)_\ell \text{ satisfies } a_j \neq b_j\}$$

be the set of index pairs of sampled coordinates where a and b disagree, and let

$$(\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}) \stackrel{\text{def}}{=} \begin{cases} \min \mathbf{D} & \mathbf{D} \neq \emptyset \\ (*, *) & \mathbf{D} = \emptyset \end{cases},$$

where the ordering in $\min \mathbf{D}$ is lexicographic.¹² Note, in particular, that the node $\mathbf{v}_{\mathbf{i}^{(s)}}(a) = \mathbf{v}_{\mathbf{i}^{(s)}}(b)$ is the least common ancestor of a and b in \mathbf{T} (hence, the “s” in $\mathbf{i}^{(s)}$ and $\boldsymbol{\ell}^{(s)}$ stands for “split”). The coordinate which is the first to satisfy $a_j \neq b_j$ is specified by the random variable

$$\mathbf{j}^{(s)} \stackrel{\text{def}}{=} \begin{cases} (\mathbf{I}_{\mathbf{i}^{(s)}})_{\boldsymbol{\ell}^{(s)}} & \text{if } (\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}) \neq (*, *) \\ * & \text{otherwise} \end{cases},$$

¹²All pairs (i, ℓ) with $i < \mathbf{i}^{(s)}$, or with $i = \mathbf{i}^{(s)}$ and $\ell < \boldsymbol{\ell}^{(s)}$, satisfy $a_j = b_j$ for $j = (\mathbf{I}_i)_\ell$.

where, notice that, $\mathbf{j}^{(s)} = *$ only if a and b satisfy $\mathbf{v}_k(a) = \mathbf{v}_k(b)$ for all $k \in \{0, \dots, h-1\}$. A trivial consequence of the above definitions is that for any $k \in \{0, \dots, h-1\}$,

$$\mathbf{1}\{\mathbf{v}_k(a) \neq \mathbf{v}_k(b)\} = \sum_{i=0}^{k-1} \sum_{\ell=1}^{2^i} \sum_{\substack{j \in [d] \\ a_j \neq b_j}} \mathbf{1}\{(\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}, \mathbf{j}^{(s)}) = (i, \ell, j)\}. \quad (22)$$

Recall the choice of $h_{a,b} \in [0 : h-1]$ in (20), we have that the expectation of (21) is at most

$$\sum_{k>i_0}^{h_{a,b}} \mathbf{E} \left[\mathbf{1}\{\mathbf{v}_k(a) \neq \mathbf{v}_k(b)\} \cdot \text{avg}_{a,k-1}^* \right] + O(\log n) \cdot \|a - b\|_1,$$

where we used the fact that $\text{avg}_{a,i}^* \leq 10d \log n / 2^k$ always holds. As a result $O(\log n) \cdot \|a - b\|_1$ as in the last term is enough to cover the sum over $k > h_{a,b}$ skipped in the above expression. Using (22), we may re-write the first summand above as

$$\begin{aligned} & \sum_{k>i_0}^{h_{a,b}} \mathbf{E} \left[\mathbf{1}\{\mathbf{v}_k(a) \neq \mathbf{v}_k(b)\} \cdot \text{avg}_{a,k-1}^* \right] \\ &= \sum_{k>i_0}^{h_{a,b}} \sum_{i=0}^{k-1} \sum_{\ell=1}^{2^i} \sum_{\substack{j \in [d] \\ a_j \neq b_j}} \Pr \left[(\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}, \mathbf{j}^{(s)}) = (i, \ell, j) \right] \cdot \mathbf{E} \left[\text{avg}_{a,k-1}^* \mid (\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}, \mathbf{j}^{(s)}) = (i, \ell, j) \right]. \end{aligned} \quad (23)$$

Notice that for each $i \in \{0, \dots, h-2\}$, $\ell \in [2^i]$, and $j \in [d]$ with $a_j \neq b_j$,

$$\Pr \left[(\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}, \mathbf{j}^{(s)}) = (i, \ell, j) \right] = \frac{1}{d} \left(1 - \frac{\|a - b\|_1}{d} \right)^{2^0 + \dots + 2^{i-1} + \ell - 1}, \quad (24)$$

which is $1/d$ when $i = 0, \ell = 1$. Consider for each $k \in \{i_0, \dots, h_{a,b} - 1\}$ and $j \in [d]$ with $a_j \neq b_j$,

$$Q_{k,j} \stackrel{\text{def}}{=} \max_{\substack{i \in \{0, \dots, k\} \\ \ell \in [2^i]}} \mathbf{E} \left[\text{avg}_{a,k}^* \mid (\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}, \mathbf{j}^{(s)}) = (i, \ell, j) \right],$$

so that invoking (24), we may upper bound (23) by

$$\sum_{\substack{j \in [d] \\ a_j \neq b_j}} \sum_{k=i_0}^{h_{a,b}-1} Q_{k,j} \cdot \sum_{i=0}^k \sum_{\ell=1}^{2^i} \frac{1}{d} \left(1 - \frac{\|a - b\|_1}{d} \right)^{2^0 + \dots + 2^{i-1} + \ell - 1} \leq \frac{2}{d} \sum_{\substack{j \in [d] \\ a_j \neq b_j}} \sum_{k=i_0}^{h_{a,b}-1} Q_{k,j} \cdot 2^k.$$

Therefore, it remains to show that for every $j \in [d]$ with $a_j \neq b_j$,

$$\sum_{k=i_0}^{h_{a,b}-1} Q_{k,j} \cdot 2^k = O(d) \cdot \left(\tilde{O}(\log n) + (h_{a,b} - i_0) \cdot \log \log n \right). \quad (25)$$

To prove this, we start with a lemma that shows that if $Q_{k,j}$ is large, then it must be the case that there are many points of distance between (roughly) $d/(2^k \log n)$ and $d \log n / 2^k$ from a in C .

Lemma 4.1. Fix $j \in [d]$ with $a_j \neq b_j$ and $k \in \{2, \dots, h_{a,b}\}$. Let $L \subseteq C_j$ be the multi-set

$$C_j \stackrel{\text{def}}{=} \{x \in C : x_j = a_j\} \quad \text{and} \quad L \stackrel{\text{def}}{=} \left\{ x \in C_j : \|a - x\|_1 \leq \frac{1}{\log n} \cdot \frac{d}{2^k} \right\}.$$

Suppose that for some $i = \{0, \dots, k\}$ and some $\ell \in [2^i]$, as well as some $\alpha \geq 800$, we have

$$\mathbf{E} \left[\text{avg}_{a,k}^* \mid (\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}, \mathbf{j}^{(s)}) = (i, \ell, j) \right] \geq \frac{d}{2^k} \cdot \alpha. \quad (26)$$

Then, the set

$$H \stackrel{\text{def}}{=} \left\{ x \in C_j : \|a - x\|_1 \leq 10 \log n \cdot \frac{d}{2^k} \right\} \quad \text{satisfies} \quad |H| \geq \frac{\exp(\alpha/8)}{\log n} \cdot |L|.$$

Proof: Let \mathcal{E} be the event of $(\mathbf{i}^{(s)}, \boldsymbol{\ell}^{(s)}, \mathbf{j}^{(s)}) = (i, \ell, j)$. For simplicity in notation, let $\mathbf{v} = \mathbf{v}_k(a)$ and

$$\mathbf{C} = \mathbf{C}_{a,k}^* \quad \text{where} \quad \mathbf{C}_{a,k}^* = \left\{ c \in \mathbf{C}_{\mathbf{v}_k(a)} : \|a - c\|_1 \leq \frac{10d \log n}{2^k} \right\}.$$

Every $x \in L$ is of distance at most $d/(2^k \log n)$ from a . Since $x \in \mathbf{C}$ whenever $\mathbf{v}_k(x) = \mathbf{v}$, we have

$$\mathbf{E} \left[|L \setminus \mathbf{C}| \mid \mathcal{E} \right] = \sum_{x \in L} \Pr \left[\mathbf{v}_k(x) \neq \mathbf{v} \mid \mathcal{E} \right] \leq |L| \cdot (2^k - 1) \cdot \frac{d/(2^k \log n)}{d - \|a - b\|_1} \leq \frac{2|L|}{\log n}, \quad (27)$$

because there are $2^k - 1$ coordinates sampled up to (but not including) depth k . The last inequality above also used the assumption that $\|a - b\|_1 \leq d/2$. Then, we have

$$\Pr \left[|\mathbf{C}| \geq \frac{|L|}{10} \mid \mathcal{E} \right] \geq \Pr \left[|L \cap \mathbf{C}| \geq \frac{|L|}{10} \mid \mathcal{E} \right] = 1 - \Pr \left[|L \setminus \mathbf{C}| \geq \frac{9|L|}{10} \mid \mathcal{E} \right] \geq 1 - \frac{20}{9 \log n}, \quad (28)$$

where the last inequality follows from Markov. Hence (noting $|\mathbf{C}| \geq 1$ since it always contains a)

$$\text{avg}_{a,k}^* = \frac{1}{|\mathbf{C}|} \sum_{x \in \mathbf{C}} \|a - x\|_1 \leq \frac{d\alpha}{2^{k+1}} + \frac{1}{|\mathbf{C}|} \sum_{x \in \mathbf{C}} \mathbf{1} \left\{ \|a - x\|_1 \geq \frac{d\alpha}{2^{k+1}} \right\} \cdot \|a - x\|_1.$$

Thus, we have (by splitting into two cases of $|\mathbf{C}| \geq |L|/10$ and $|\mathbf{C}| < |L|/10$ and applying (28))

$$\begin{aligned} \mathbf{E} \left[\text{avg}_{a,k}^* \mid \mathcal{E} \right] &\leq \frac{d\alpha}{2^{k+1}} + \mathbf{E} \left[\frac{1}{|\mathbf{C}|} \sum_{x \in \mathbf{C}} \mathbf{1} \left\{ \|a - x\|_1 \geq \frac{d\alpha}{2^{k+1}} \right\} \|a - x\|_1 \mid \mathcal{E} \right] \\ &\leq \frac{d\alpha}{2^{k+1}} + \frac{10}{|L|} \cdot \mathbf{E} \left[\sum_{x \in \mathbf{C}} \mathbf{1} \left\{ \|a - x\|_1 \geq \frac{d\alpha}{2^{k+1}} \right\} \|a - x\|_1 \mid \mathcal{E} \right] + \frac{20}{9 \log n} \cdot \frac{10d \log n}{2^k}, \end{aligned}$$

where the final term used the fact that $\|a - x\|_1$ is always at most $10d \log n/2^k$ for any $x \in \mathbf{C}$, by the definition of the latter. Combining (26) and the inequality above (and $\alpha \geq 800$), we have

$$\frac{\alpha}{40} \cdot \frac{d}{2^{k+1}} \cdot |L| \leq \mathbf{E} \left[\sum_{x \in \mathbf{C}} \mathbf{1} \left\{ \|a - x\|_1 \geq \frac{d\alpha}{2^{k+1}} \right\} \|a - x\|_1 \mid \mathcal{E} \right]. \quad (29)$$

We finish the proof of the lemma by upperbounding the right-hand side above in terms of the size of H . In particular, let H' be the set of points $x \in H$ with $\|a - x\|_1 \geq d\alpha/2^{k+1}$. Then

$$\mathbf{E} \left[\sum_{x \in \mathbf{C}} \mathbf{1} \left\{ \|a - x\|_1 \geq \frac{d\alpha}{2^{k+1}} \right\} \|a - x\|_1 \mid \mathcal{E} \right] \leq \sum_{x \in H'} \Pr [x \in \mathbf{C} \mid \mathcal{E}] \cdot \left(\frac{10d \log n}{2^k} \right). \quad (30)$$

We consider two cases: $i = k$ (see \mathcal{E}) and $i < k$. For the easier case of $i = k$, in order for $x \in \mathbf{C}$ to occur conditioned on \mathcal{E} , we have that all $2^k - 1$ coordinates sampled before depth k avoid separating x and a conditioning on not separating a and b . The probability of each sample is at most

$$\frac{d - \|a - x\|_1}{d - \|a - b\|_1} = 1 - \frac{\|a - x\|_1 - \|a - b\|_1}{d - \|a - b\|_1} \leq 1 - \frac{\|a - x\|_1 - \|a - b\|_1}{d}.$$

Therefore in this case we have

$$\Pr [x \in \mathbf{C} \mid \mathcal{E}] \leq \left(1 - \frac{\|a - x\|_1 - \|a - b\|_1}{d} \right)^{2^k - 1}.$$

Notice that by definition of $h_{a,b}$, and the fact that $k \leq h_{a,b}$, we have

$$\|x - a\|_1 - \|a - b\|_1 \geq \frac{d}{2^{k+1}}(\alpha - 2) \quad (31)$$

which implies that $\Pr[x \in \mathbf{C} \mid \mathcal{E}]$ is at most $\exp(-\alpha/8)$ using $\alpha \geq 800$.

Next we deal with the case when $i < k$. In order for $x \in \mathbf{C}$ to occur conditioned on \mathcal{E} , it needs to be the case that coordinates sampled before (i, ℓ) , of which there are $2^i + \ell - 2$ many, avoid separating x and a conditioning on not separating a and b ; the (i, ℓ) -th sample is j (which better not separate x and a ; otherwise the probability is trivially 0); and the remaining $2^k - 1 - 2^i - \ell + 1 = 2^k - 2^i - \ell$ coordinates do not separate x and a (but there will be conditioning on not separating a and b). So

$$\begin{aligned} \Pr [x \in \mathbf{C} \mid \mathcal{E}] &\leq \left(1 - \frac{\|a - x\|_1 - \|a - b\|_1}{d} \right)^{2^i + \ell - 2} \left(1 - \frac{\|a - x\|_1}{d} \right)^{2^k - 2^i - \ell} \\ &\leq \left(1 - \frac{\|a - x\|_1 - \|a - b\|_1}{d} \right)^{2^k - 2} \\ &\leq \left(1 - \frac{\alpha - 2}{2^{k+1}} \right)^{2^{k-1} - 1}, \end{aligned} \quad (32)$$

which is at most $\exp(-\alpha/8)$ using $k \geq 2$ and $\alpha \geq 800$. Hence, we can combine (29) and (30) to get

$$\frac{\alpha}{40} \cdot \frac{d}{2^{k+1}} \cdot |L| \leq |H| \cdot \exp\left(-\frac{\alpha}{8}\right) \cdot \frac{10d \log n}{2^k}.$$

Re-arranging the inequality, the lemma follows using $\alpha \geq 800$. ■

The next lemma helps upperbound the number of large $Q_{k,j}$'s.

Lemma 4.2. *Fix any $j \in [d]$ with $a_j \neq b_j$. When $\alpha \geq 20 \log \log n$, the set*

$$G_j(\alpha) = \left\{ k \in \{0, \dots, h_{a,b}\} : Q_{k,j} \geq \frac{d}{2^k} \cdot \alpha \right\} \quad \text{satisfies} \quad |G_j(\alpha)| \leq O \left(\left\lceil \frac{16 \log(2n)}{\alpha} \right\rceil \cdot \log \log n \right).$$

Proof: Assume for a contradiction that

$$|G_j(\alpha)| \geq \beta \cdot \lceil \log_2(10 \log^2 n) \rceil + 2, \quad \text{where } \beta \stackrel{\text{def}}{=} \left\lceil \frac{16 \log(2n)}{\alpha} \right\rceil.$$

Then there must be $k_1, \dots, k_\beta \in \{2, \dots, h_{a,b}\}$ with $k_1 > k_2 > \dots > k_\beta$ and every $t \in [\beta - 1]$ satisfies

$$k_t - k_{t+1} \geq \lceil \log_2(10 \log^2 n) \rceil.$$

This implies that every $t \in [\beta - 1]$ satisfies

$$\frac{10d \log n}{2^{k_t}} \leq \frac{d}{2^{k_{t+1}} \log n}.$$

If we consider for each $t \in [\beta]$, the following two multi-sets

$$L_t = \left\{ x \in C_j : \|a - x\|_1 \leq \frac{d}{2^{k_t} \log n} \right\} \quad \text{and} \quad H_t = \left\{ x \in C_j : \|a - x\|_1 \leq \frac{10d \log n}{2^{k_t}} \right\},$$

they satisfy

$$L_1 \subseteq H_1 \subseteq L_2 \subseteq H_2 \subseteq \dots \subseteq H_{\beta-1} \subseteq L_\beta \subseteq H_\beta, \tag{33}$$

but then invoking Lemma 4.1 (and using $20 \log \log n \geq 800$), we have that every $t \in [\beta]$ satisfy

$$|H_t| \geq \frac{|L_t| \exp(\alpha/8)}{\log n}.$$

Using $|L_1| \geq 1$ (since it contains a) and (33), we have

$$|H_\beta| \geq \left(\frac{\exp(\alpha/8)}{\log n} \right)^\beta > 2n,$$

using $\exp(\alpha\beta/8) \geq (2n)^2$, and $(\log n)^\beta < 2n$, by our choices of β and α . This is a contradiction, as we have $H_\beta \subseteq A \cup B$ and thus, $|H_\beta| \leq 2n$. ■

Finally we finish the proof of (25). Let $\alpha_0 = 20 \log \log n$. Use Lemma 4.2 we have

$$\begin{aligned} \sum_{k=i_0}^{h_{a,b}-1} Q_{k,j} \cdot 2^k &\leq (h_{a,b} - i_0) \cdot \alpha_0 d + \sum_{\kappa=0}^{\lceil \log_2(10 \log n) \rceil} |G_j(\alpha_0 2^\kappa)| \cdot \alpha_0 2^{\kappa+1} \cdot d \\ &\leq O(d) \cdot \left((h_{a,b} - i_0) \cdot \log \log n + \log n \cdot (\log \log n)^3 \right), \end{aligned}$$

where the upper limit of $\kappa \leq \lceil \log_2(10 \log n) \rceil$ comes from the fact that $G_j(10 \log n)$ is trivially empty since $\text{avg}_{a,k}^*$ is always at most $10d \log n / 2^k$ by definition. This finishes the proof of Lemma 3.6.

5 Streaming Preliminaries and Sketching Tools

5.1 Exponential Order Statistics

We review some properties of the order statistics of independent non-identically distributed exponential random variables. Let (t_1, \dots, t_n) be independent exponential random variables where t_i has mean $1/\lambda_i$ (equivalently, t_i has rate $\lambda_i > 0$), abbreviated as $t_i \sim \text{Exp}(\lambda_i)$. Recall that t_i is given by the cumulative distribution function (cdf) $\Pr[t_i \leq x] = 1 - e^{-\lambda_i x}$. Our algorithm will require an analysis of the distribution of values (t_1, \dots, t_n) . We begin by noting that constant factor scalings of an exponential variable result in another exponential variable.

Fact 5.1 (Scaling of exponentials). *Let $t \sim \text{Exp}(\lambda)$ and $\alpha > 0$. Then αt is distributed as $\text{Exp}(\lambda/\alpha)$.*

Proof: The cdf of αt is given by $\Pr[t < x/\alpha] = 1 - e^{-\lambda x/\alpha}$, which is the cdf of $\text{Exp}(\lambda/\alpha)$. ■

Definition 5.2. *Let $t = (t_1, \dots, t_n)$ be independent exponentials. For $k = 1, 2, \dots, n$, we define the k -th anti-rank $D_t(k) \in [n]$ of t to be the values $D_t(k)$ such that $t_{D_t(1)} \leq t_{D_t(2)} \leq \dots \leq t_{D_t(n)}$.*

Using the structure of the anti-rank vector, it has been observed [Nag06] that there is a simple form for describing the distribution of $t_{D_t(k)}$ as a function of $(\lambda_1, \dots, \lambda_n)$ and the anti-rank vector.

Fact 5.3 ([Nag06]). *Let $t = (t_1, \dots, t_n)$ be independently exponentials with $t_i \sim \text{Exp}(\lambda_i)$. Then $D_t(1) = i$ with probability $\lambda_i / \sum_{j \in [n]} \lambda_j$. Furthermore, the following two sampling procedures produce the same distribution over pairs in \mathbb{R}^2 :*

1. Sample $t = (t_1, \dots, t_n)$, where $t_i \sim \text{Exp}(\lambda_i)$, and output $(t_{D_t(1)}, t_{D_t(2)} - t_{D_t(1)})$.
2. Sample $i_1 \in [n]$ with $\Pr[i_1 = i] = \lambda_i / \sum_{j \in [n]} \lambda_j$, $E_1, E_2 \sim \text{Exp}(1)$ independently, and output

$$\left(\frac{E_1}{\sum_{j \in [n]} \lambda_j}, \frac{E_2}{\sum_{j \in [n] \setminus \{i_1\}} \lambda_j} \right).$$

Proof: This is a simple computation. We have that for any $r, r' \in \mathbb{R}_{\geq 0}$ and $i \in [n]$,

$$\begin{aligned} \Pr_t \left[\begin{array}{l} D_t(1) = i, \\ t_{D_t(1)} \geq r, \\ t_{D_t(2)} - t_{D_t(1)} \geq r' \end{array} \right] &= \int_{y:r}^{\infty} \lambda_i \exp(-\lambda_i y) \prod_{j \in [n] \setminus \{i\}} \Pr_{t_j \sim \text{Exp}(\lambda_j)} [t_j - y \geq r'] dy \\ &= \lambda_i \exp\left(-r' \sum_{j \in [n] \setminus \{i\}} \lambda_j\right) \int_{y:r}^{\infty} \exp\left(-y \sum_{j=1}^n \lambda_j\right) dy \\ &= \frac{\lambda_i}{\sum_{j=1}^n \lambda_j} \cdot \exp\left(-r' \sum_{j \in [n] \setminus \{i\}} \lambda_j\right) \cdot \exp\left(-r \sum_{j=1}^n \lambda_j\right) \\ &= \Pr \left[i_1 = i \wedge \frac{E_1}{\sum_{j \in [n]} \lambda_j} \geq r \wedge \frac{E_2}{\sum_{j \in [n] \setminus \{i_1\}} \lambda_j} \geq r' \right]. \end{aligned}$$

This finishes the proof. ■

Lemma 5.4. Fix $n \in \mathbb{N}$ and $x \in \mathbb{R}^n$ to any fixed vector, and consider independent draws $\mathbf{t}_1, \dots, \mathbf{t}_n \sim \text{Exp}(1)$. For any $\gamma \in (0, 1/2)$,

$$\Pr_{\mathbf{t}_1, \dots, \mathbf{t}_n} \left[\sum_{i \in [n]} \frac{|x_i|}{\mathbf{t}_i} \geq \frac{4 \log(n/\gamma)}{\gamma} \|x\|_1 \right] \leq 2\gamma.$$

Proof: We want to compute the expectation of $\sum |x_i|/\mathbf{t}_i$ and apply Markov's inequality; however, the above random variable does not have an expectation (since $1/\mathbf{t}_i$ may become too large when \mathbf{t}_i is small). To remedy that, we effectively truncate $1/\mathbf{t}_i$. Notice that for any $\alpha > 0$, the probability that any $\mathbf{t}_i \sim \text{Exp}(1)$ is less than α is at most α (simply by inspecting p.d.f of $\text{Exp}(1)$). We apply this in two ways: (i) for any $i \in [n]$, the probability that $1/\mathbf{t}_i > n/\gamma$ is at most γ/n , and (ii) for any $i \in [n]$ and $j \in \mathbb{N}$, the probability that $1/\mathbf{t}_i \in [2^j, 2^{j+1}]$ is at most $1/2^j$.

Letting \mathcal{Q}_i be the event that $1/\mathbf{t}_i > n/\gamma$, and by (i), we may union bound over all n to say \mathcal{Q}_i is never satisfied with probability at least $1 - \gamma$. So now we compute the expectation and apply Markov's inequality (conditioning on $\neg(\cup_i \mathcal{Q}_i)$ so the probability that $1/\mathbf{t}_i > n/\gamma$ is 0):

$$\begin{aligned} \mathbf{E}_{\mathbf{t}_1, \dots, \mathbf{t}_n} \left[\sum_{i=1}^n \frac{|x_i|}{\mathbf{t}_i} \mid \neg(\cup_i \mathcal{Q}_i) \right] &\leq \sum_{i=1}^n |x_i| \sum_{j=0}^{\infty} 2^{j+1} \cdot \Pr_{\mathbf{t}_i} \left[\frac{1}{\mathbf{t}_i} \in [2^j, 2^{j+1}] \mid \neg \mathcal{Q}_i \right] \\ &\leq \sum_{i=1}^n |x_i| \sum_{j=0}^{\log(n/\gamma)} 2^{j+1} \cdot \Pr_{\mathbf{t}_i} \left[\frac{1}{\mathbf{t}_i} \in [2^j, 2^{j+1}] \mid \neg \mathcal{Q}_i \right] \\ &\leq \sum_{i=1}^n |x_i| \sum_{j=0}^{\log(n/\gamma)} 2^{j+1} \cdot \frac{1}{2^j(1-\gamma)} \leq \|x\|_1 \cdot 2 \log(n/\gamma)/(1-\gamma). \end{aligned}$$

Then, by Markov's inequality,

$$\Pr_{\mathbf{t}_1, \dots, \mathbf{t}_n} \left[\sum_{i \in [n]} \frac{|x_i|}{\mathbf{t}_i} \geq \frac{4 \log(n/\gamma)}{\gamma} \|x\|_1 \mid \bigcup_i \neg \mathcal{Q}_i \right] \leq \gamma.$$

The proof follows from a union bound. ■

Lemma 5.5. For $n \in \mathbb{N}$, let $x \in \mathbb{R}^n$ be any fixed vector. Let $\mathbf{t}_1, \dots, \mathbf{t}_n \sim \text{Exp}(1)$ be i.i.d. exponentially distributed, and $\gamma > 0$ be smaller than some constant. Then, letting $i^* = \arg \max \frac{|x_i|}{\mathbf{t}_i}$ we have that

$$\frac{|x_{i^*}|}{\mathbf{t}_{i^*}} \geq \gamma \|x\|_1 \quad \text{and} \quad \frac{|x_{i^*}|}{\mathbf{t}_{i^*}} \geq (1 + \gamma) \max_{i \neq i^*} \left\{ \frac{|x_i|}{\mathbf{t}_i} \right\}, \quad (34)$$

holds with probability at least $1 - 4\gamma$.

Proof: Let \mathcal{E} is the first event in (34), and \mathcal{E}' is the second. By Fact 5.3, the quantity $|x_{i^*}|/\mathbf{t}_{i^*}$ is distributed as $\|x\|_1/\mathbf{E}_1$, where \mathbf{E}_1 is an exponential random variable. Using the cdf $1 - e^{-x}$ of an exponential, we have that $\Pr[\mathbf{E}_1 > 1/\gamma] < e^{-1/\gamma}$, thus $\Pr[\mathcal{E}] > 1 - e^{-1/\gamma} > 1 - \gamma$ for γ smaller than a constant.

We now bound $\Pr[\mathcal{E}']$. Again by Fact 5.3, we have that $\max_{i \neq i^*} \left\{ \frac{|x_i|}{t_i} \right\}$ is distributed as

$$\max_{i \neq i^*} \left\{ \frac{|x_i|}{t_i} \right\} \sim \frac{1}{\frac{\mathbf{E}_1}{\|x\|_1} + \frac{\mathbf{E}_2}{\|x\|_1 - |x_{i^*}|}} \leq \frac{\|x\|_1}{\mathbf{E}_1 + \mathbf{E}_2}.$$

Thus, \mathcal{E}' will hold so long as $\mathbf{E}_2 \geq 3\gamma\mathbf{E}_1$. The probability that this does not occur is given by

$$\Pr[\mathbf{E}_2 < 3\gamma\mathbf{E}_1] \leq \int_0^\infty e^{-x} \Pr \left[\mathbf{E}_1 > \frac{x}{3\gamma} \right] dx = \int_0^\infty e^{-x(1+\frac{1}{3\gamma})} dx = \frac{1}{(1+\frac{1}{3\gamma})} \leq 3\gamma.$$

Thus, we have $\Pr[\mathcal{E}'] > 1 - 3\gamma$, and so by a union bound the event in (34) holds with probability at least $1 - 4\gamma$, which completes the proof. \blacksquare

Given $x \in \mathbb{R}^n$ and an integer $\beta \geq 1$, we write $x_{-\beta} \in \mathbb{R}^n$ to denote the vector given by x where the β largest coordinates in magnitude are set to 0. When $\beta \geq 1$ is not an integer, $x_{-\beta}$ is interpreted as $x_{-\lfloor \beta \rfloor}$. We will need the following lemma which bounds the tail ℓ_2 and ℓ_1 norm of a vector after its entries are scaled independently by random variables

Lemma 5.6 (Generalization of Proposition 1 of [JW21]). *Fix $n \in \mathbb{N}$ and $c \geq 0$, and let $\mathcal{D}_1, \dots, \mathcal{D}_n$ be a sequence of distributions over \mathbb{R} satisfying*

$$\Pr_{t_i \sim \mathcal{D}_i} [t_i \geq y] \leq \frac{c}{y}, \quad \text{for all } i \in [n] \text{ and all } y > 0.$$

For any fixed vector $x \in \mathbb{R}^n$ and integer $\beta \geq 1$, consider the random vector $z \in \mathbb{R}^n$ given by letting

$$z_i \stackrel{\text{def}}{=} t_i \cdot x_i, \quad \text{where } t_i \sim \mathcal{D}_i \text{ independently for all } i \in [n].$$

Then we have

$$\|z_{-\beta}\|_2 \leq \frac{12c}{\sqrt{\beta}} \cdot \|x\|_1 \quad \text{and} \quad \|z_{-\beta}\|_1 \leq 9c \lceil \log_2 n \rceil \cdot \|x\|_1$$

with probability at least $1 - 3e^{-\beta/8}$ over the draws of $t_i \sim \mathcal{D}_i$.

Proof: Assume without loss of generality that $\beta \geq 8$; otherwise $1 - 3e^{-\beta/8} < 0$ and the statement is trivial. Define the following random sets \mathbf{I}_j for each $j = 0, 1, \dots, \lceil \log_2 n \rceil$:

$$\mathbf{I}_j = \left\{ i \in [n] : \frac{c\|x\|_1}{2^{j+1}} \leq |z_i| \leq \frac{c\|x\|_1}{2^j} \right\}$$

and notice that, for any $i \in [n]$, we have

$$\Pr_{t_i \sim \mathcal{D}_i} [i \in \mathbf{I}_j] \leq \Pr_{t_i \sim \mathcal{D}_i} \left[|t_i| \geq \frac{c\|x\|_1}{2^{j+1}|x_i|} \right] \leq \frac{2^{j+1}|x_i|}{\|x\|_1} \quad \text{implying} \quad \mathbf{E}_{t_1, \dots, t_n} [|\mathbf{I}_j|] \leq 2^{j+1}.$$

Let \mathcal{E}_1 denote the event that there exists $j \geq \lceil \log_2(\beta/8) \rceil$ such that $|\mathbf{I}_j| > 4 \cdot 2^{j+1}$. Then

$$\Pr_{t_1, \dots, t_n} [\mathcal{E}_1] \leq \sum_{j=\lceil \log_2(\beta/8) \rceil}^{\lceil \log_2 n \rceil} \Pr_{t_1, \dots, t_n} [|\mathbf{I}_j| > 4 \cdot 2^{j+1}] \leq \sum_{j=\lceil \log_2(\beta/8) \rceil}^{\lceil \log_2 n \rceil} \exp(-2^j) \leq 2 \exp(-\beta/8),$$

by a Chernoff bound. On the other hand, each $i \in [n]$ satisfies $|z_i| \geq 4c\|x\|_1/\beta$ with probability at most $\beta|x_i|/(4\|x\|_1)$ over the draw of $t_i \sim \mathcal{D}_i$. The event \mathcal{E}_2 that more than β indices $i \in [n]$ satisfy $|z_i| \geq 4c\|x\|_1/\beta$ happens with probability at most $\exp(-\beta/8)$. Whenever \mathcal{E}_1 and \mathcal{E}_2 do not occur, (which happens with probability at least $1 - 3e^{-\beta/8}$), we have

$$\|z_{-\beta}\|_2^2 \leq \sum_{j=\lceil \log_2(\beta/8) \rceil}^{\lceil \log_2 n \rceil} |\mathbf{I}_j| \cdot \frac{c^2\|x\|_1^2}{2^{2j}} + n \cdot \frac{c^2\|x\|_1^2}{n^2} \leq \frac{129}{\beta} \cdot c^2\|x\|_1 \quad \text{and}$$

$$\|z_{-\beta}\|_1 \leq \sum_{j=\lceil \log_2(\beta/8) \rceil}^{\lceil \log_2 n \rceil} |\mathbf{I}_j| \cdot \frac{c\|x\|_1}{2^j} + n \cdot \frac{c\|x\|_1}{n} \leq c\|x\|_1 (8\lceil \log_2 n \rceil + 1)$$

since once $\beta \geq n$, the bound trivially becomes 0. The lemma follows. \blacksquare

5.2 Geometric Streaming and Linear Sketching

All of the streaming algorithms in this paper will be *linear sketches*. We begin by formalizing what a linear sketch is in the context of geometric streaming algorithms. Recall that in the geometric streaming model, we receive a stream of updates $(p_1, \sigma_1), (p_2, \sigma_2), \dots, (p_m, \sigma_m)$, where $p_i \in \{0, 1\}^d$ and $\sigma_i \in \{1, -1\}$ indicates either an insertion or deletion of the point p_i from the active dataset $X \subset \{0, 1\}^d$. For the case of EMD, we also need to specify whether a given point p_i is being inserted or deleted from A or B , in which case the stream consists of updates of the form $(p_i, \sigma_i, \theta_i)$ where $\theta_i \in \{A, B\}$ indicates which of the two sets the update applies to.

We now observe that the above geometric streaming model is just a special case of the standard turnstile streaming model [BBD⁺02, M⁺05], which consists of a sequence of insertions and deletions to the coordinates of a high-dimensional vector $f \in \mathbb{R}^N$. Specifically, in the standard streaming model, the stream consists of updates $(i_1, \Delta_1), \dots, (i_m, \Delta_m)$, where $i_t \in [N]$ is a coordinate and $\Delta_t \in \{1, -1\}$. Here, the update (i_t, Δ_t) causes the change $f_{i_t} \leftarrow f_{i_t} + \Delta_t$. Thus, by simply setting $f = f_X \in \mathbb{R}^{2^d}$ to be the indicator vector (with multiplicity) of the multi-set X for the case of EMD, or setting $f = f_{A,B} \in \mathbb{R}^{2 \cdot 2^d}$ to be the indicator vector (with multiplicity) of the two sets A and B stacked together, then a stream of insertions and deletions of points p in the geometric model coincides to insertions and deletions to the coordinates of f . Given this connection, we can now define a linear sketch.

Definition 5.7. *Given a stream of updates to a vector $f \in \mathbb{R}^N$, a (one-pass) linear sketch generates a random matrix $\mathbf{S} \in \mathbb{R}^{k \times N}$, and stores only the matrix-vector product $\mathbf{S}f$ and the matrix \mathbf{S} . At the end of the stream, it answers a query based on $\mathbf{S}f, \mathbf{S}$. A two-pass linear sketch generates random matrix \mathbf{S}^1 , and on the first pass stores only $\mathbf{S}^1 f, \mathbf{S}^1$. After the first pass over the data, it generates a random matrix \mathbf{S}^2 (possibly depending on $\mathbf{S}^1 f, \mathbf{S}^1$), and after the second pass it answers a query based on $\mathbf{S}^1 f, \mathbf{S}^1, \mathbf{S}^2 f, \mathbf{S}^2$.*

The space used by a one-pass (resp. two-pass) linear sketch is the space required to store $\mathbf{S}f$ (resp. $\mathbf{S}^1 f, \mathbf{S}^2 f$.)

For simplicity, in the entirety of Sections 6 and 7, we will defer consideration of the *bit complexity* required to store entries of the sketches $\mathbf{S}f$, and instead focus on bounding the dimension of the

sketch $\mathbf{S}f$ — for instance, in both sections we will often work with real-valued random variables (with unbounded bit complexity). Then in Appendix B, we will handle the issue of bit-complexity, by demonstrating that we can generate all random variables to $\text{polylog}(n)$ -bits of precision. This will result in the space of the algorithm being within a factor of $\text{polylog}(n)$ of the dimension of the sketch.

Notice in the definition of a linear sketch, the space complexity did not depend on the space required to store the matrix \mathbf{S} , while the output the algorithm is allowed to depend on \mathbf{S} . This coincides with what is known as the *random oracle model* of streaming, or the *public coin* model of communication complexity. Formally:

Definition 5.8. *In the random oracle streaming model, the algorithm is given access to an arbitrarily long string of random bits which do not count against the space complexity. In particular, the space of a one-pass (resp. two-pass) linear sketching algorithm in the random oracle model is just the space required to store $\mathbf{S}f$ (resp. $\mathbf{S}^1f, \mathbf{S}^2f$.)*

Working in the random oracle model of streaming is common in the sketching literature, as nearly all lower bounds for streaming are derived from the public coin model of communication complexity (and therefore apply to the random oracle model). Moreover, in applications of sketching to distributed computation, the assumption of a random oracle is often founded.

On the other hand, we demonstrate that the random oracle model is not required for our algorithms. In particular, we show how the assumption of a random oracle can be removed in Appendix B, albeit at the cost of an additive $d \cdot \text{polylog}n$ in the space of the algorithm. Note that even to read a single update in the stream requires $\Omega(d)$ -bits of space, so any algorithm requires $\Omega(d)$ *working space* (see remark below). Thus, this additive $d \cdot \text{polylog}n$ in the space resulting from derandomization is comparable, up to log factors, to the space required to even read an update.

Since the issues of bit complexity and derandomization are handled in Appendix B, for the remainder of Sections 6 and 7 we assume the random oracle model, and analyze space in terms of the dimension of the sketch

Remark 7. *When discussing the space complexity of streaming algorithms, there are two separate notions: working space and intrinsic space. Oftentimes these two notations of space are the same (up to constant factors) for streaming algorithms, however for our purposes it will be useful to distinguish them, since a d -dimensional point itself requires $\Omega(d)$ -space to specify. The working space of a streaming algorithm \mathcal{A} is the space required to store an update (i_t, Δ_t) in the stream and process it. The intrinsic space is the space which the algorithm must store between updates. The intrinsic space coincides with the size of a message in the area of communication complexity, where two parties hold a fraction of the input, and must exchange messages to approximate a function of the entire input. The intrinsic space also coincides with the definition of space given in Definition 5.7. In this paper, we focus on intrinsic space, which we will hereafter just refer to as the space of the algorithm.*

5.3 Count-Sketch, ℓ_1 -sketch, and ℓ_1 -sampling

We now introduce several useful sketches from the streaming literature. We remark that all the sketches below are already derandomized (they do not require the random oracle model), and the

space complexity in bits is stated within the theorems.

Theorem 8 (Count-Sketch [CCFC02]). *Fix $n \in \mathbb{N}$ and $\epsilon \in (0, 1)$. There is a $O(\log n/\epsilon^2)$ -bits of space linear sketch that, given any input vector $x \in \mathbb{R}^n$, outputs $\hat{x} \in \mathbb{R}^n$ such that $\|\hat{x} - x\|_\infty \leq \epsilon \|x_{-1/\epsilon^2}\|_2$ with probability at least $1 - 1/\text{poly}(n)$.*

Theorem 9 (ℓ_1 -sketch [Ind06b, KNW10]). *Fix $n \in \mathbb{N}$ and $\epsilon, \delta \in (0, 1)$, and $s = O(\log(1/\delta)/\epsilon^2)$. Let Ω be an $s \times n$ matrix with independent Cauchy random variables $\Omega_{i,j} \sim \mathcal{C}$. For any $x \in \mathbb{R}^n$, let*

$$\eta = \frac{\text{median}\{(\Omega x)_j : j \in [s]\}}{\text{median}(|\mathcal{C}|)}.$$

Then we have $|\eta - \|x\|_1| \leq \epsilon \|x\|_1$ with probability at least $1 - \delta$ over the randomness of Ω . Moreover, the matrix Ω can be generated with limited independence so that it can be stored in $O(\log(1/\delta) \log(n)/\epsilon^2)$ -bits of space, and so that the above guarantees still hold.

Theorem 10 (Perfect ℓ_1 -sampling [JW21]). *For $m \in \mathbb{N}$, let $c > 1$ be an arbitrarily large constant and $t = O(\log^2 m)$. There exists a distribution $\mathcal{S}_a^1(m, t)$ supported on pairs $(\mathbf{S}, \text{Alg}_{\mathbf{S}}^1)$ where \mathbf{S} is an $t \times m$ matrix and $\text{Alg}_{\mathbf{S}}^1$ is an algorithm which receives as input a vector $y \in \mathbb{R}^t$ and outputs a failure symbol \perp with probability at most $1/3$, otherwise it returns an index $j \in [m]$. For any $x \in \mathbb{R}^m$ and $j \in [m]$,*

$$\left| \Pr_{(\mathbf{S}, \text{Alg}_{\mathbf{S}}^1) \sim \mathcal{S}_a^1(m, t)} [\text{Alg}_{\mathbf{S}}^1(\mathbf{S}x) = j \mid \text{Alg}_{\mathbf{S}}^1(\mathbf{S}x) \neq \perp] - \frac{|x_j|}{\|x\|_1} \right| \leq \frac{1}{m^c}.$$

Lastly, the matrix \mathbf{S} be generated with limited independence so that it can be stored in $O(\log^2 n (\log \log n)^2)$ bits of space, and so that the above guarantees still hold.

6 Linear Sketches for EMD

We give linear sketches for EMD in this section and prove the following theorems:

Theorem 11. *For $d, n \in \mathbb{N}$, there is a $O(1/\epsilon) \cdot \text{polylog}(n, d)$ -space linear sketching algorithm that, given multi-sets $A, B \subset \{0, 1\}^d$ with $|A| = |B| = n$, outputs a number η such that*

$$\text{EMD}(A, B) \leq \eta \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B) + \epsilon nd \tag{35}$$

with probability at least $2/3$.

Theorem 12. *For $d, n \in \mathbb{N}$, there is a 2-round $\text{polylog}(n, d)$ -space linear sketching algorithm that, given multi-sets $A, B \subset \{0, 1\}^d$ with $|A| = |B| = n$, outputs a number η such that*

$$\text{EMD}(A, B) \leq \eta \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B)$$

with probability at least $2/3$.

6.1 Preparation

Fix $d \in \mathbb{N}$, and fix a quadtree T of depth $h = \lceil \log_2 d \rceil$. For each $i \in [0 : h]$, we let V_i be the set of nodes of T at depth i ; for a node $v \in V_i$ for $i > 0$, we let $\pi(v)$ be the parent node of v in T (which is a node at depth $i - 1$). For each $i \in [0 : h]$, T induces a map $\mathbf{v}_i : \{0, 1\}^d \rightarrow V_i$: $\mathbf{v}_0(a), \mathbf{v}_1(a), \dots, \mathbf{v}_h(a)$ is the path of each point $a \in \{0, 1\}^d$ going down the quadtree T .

Fix $n \in \mathbb{N}$. Let $A, B \subseteq \{0, 1\}^d$ be the pair of input sets of size n each. We use A_v (or B_v) for each $v \in V_i$ to denote the set of $a \in A$ (or $b \in B$) with $\mathbf{v}_i(a) = v$ (or $\mathbf{v}_i(b) = v$), with $C_v = A_v \cup B_v$. Let

$$\Delta_i = \sum_{v \in V_i} \left| |A_v| - |B_v| \right|.$$

When $\Delta_i > 0$ (which is the main case we will work on), we write \mathcal{V}_i to denote the distribution over V_i where $v \in V_i$ is sampled with probability $|A_v| - |B_v| / \Delta_i$. For each $i \in [h]$, we let \mathcal{S}_i denote the distribution over subsets of $[d]$ which includes each coordinate independently with probability

$$\alpha_i \stackrel{\text{def}}{=} \frac{2^i}{d \log^2 n}.$$

Given $S \subset [d]$, the character $\chi_S : \{0, 1\}^d \rightarrow \{-1, 1\}$ is the function $\chi_S(x) = (-1)^{\sum_{k \in S} x_k}$.

We use the quadtree lemma for EMD (Lemma 3.3) in Section 3.1 to get the following lemma that is key to our linear sketches:

Lemma 6.1. *At least 90% of quadtrees T (as drawn from the distribution \mathcal{T}) satisfy*

$$\text{EMD}(A, B) \leq \sum_{i \in [h]} \mathcal{I}_i \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B) \quad \text{and} \quad \sum_{i \in [h]} \frac{\Delta_i d}{2^i} \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B) \quad (36)$$

where \mathcal{I}_i for each $i \in [h]$ is defined as $\mathcal{I}_i = 0$ when $\Delta_i = 0$ and

$$\mathcal{I}_i \stackrel{\text{def}}{=} \frac{2\Delta_i}{\alpha_i} \cdot \mathbf{E}_{\mathbf{v} \sim \mathcal{V}_i, \mathbf{S} \sim \mathcal{S}_i} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}}]$$

when $\Delta_i > 0$, where for each $v \in V_i$ and $u = \pi(v)$,

$$p_{u, v, S} \stackrel{\text{def}}{=} \Pr_{\mathbf{c}_u \sim C_u, \mathbf{c}_v \sim C_v} [\chi_S(\mathbf{c}_u) \neq \chi_S(\mathbf{c}_v)] \in [0, 1].$$

Proof: We show that 98% of quadtrees satisfy the first part of (36) and 98% of quadtrees satisfy the second part of (36) and the lemma follows. The second part follows from an analysis similar to (2) (although (2) only gives $O(\log d)$ on the right hand side instead of the $O(\log n)$ we need, one can improve it to $\min(\log n, \log d)$; see footnote 5). We focus on the first part of (36).

Recall from Lemma 3.3 that at least 99% of quadtrees T satisfy

$$\text{EMD}(A, B) \leq \mathbf{Value}_T(A, B) = \sum_{(u, v) \in E_T} \left| |A_u| - |B_u| \right| \cdot \text{avg}_{u, v} \leq \tilde{O}(\log n) \cdot \text{EMD}(A, B).$$

We can write $\mathbf{Value}_T(A, B)$ as the sum of \mathcal{I}_i^* over $i \in [h]$: $\mathcal{I}_i^* = 0$ when $\Delta_i = 0$ and

$$\mathcal{I}_i^* \stackrel{\text{def}}{=} \Delta_i \cdot \mathbf{E}_{\mathbf{v} \sim \mathcal{V}_i} \left[\text{avg}_{\pi(\mathbf{v}), \mathbf{v}} \right] = \Delta_i \cdot \mathbf{E}_{\mathbf{v} \sim \mathcal{V}_i} \left[\mathbf{E}_{\mathbf{c} \sim C_{\pi(\mathbf{v})}, \mathbf{c}' \sim C_{\mathbf{v}}} [\|\mathbf{c} - \mathbf{c}'\|_1] \right].$$

when $\Delta_i > 0$. Also recall from the proof of Lemma 3.3 that in at least $(1 - o(1))$ -fraction of T , every two points $c, c' \in C_{\pi(v)}$ for a node v at level i have $\|a - b\|_1 \leq O(d \log n / 2^i)$. When this happens, we have from the following claim that $\mathcal{I}^* \leq \mathcal{I} \leq 2\mathcal{I}^*$.

Claim 6.2. *Let $a, b \in \{0, 1\}^d$ with $\|a - b\|_1 = O(d \log n / 2^i)$. Then we have*

$$\frac{\alpha_i \|a - b\|_1}{2} \leq \Pr_{\mathbf{S} \sim \mathcal{S}_i} [\chi_{\mathbf{S}}(a) \neq \chi_{\mathbf{S}}(b)] \leq \alpha_i \|a - b\|_1.$$

Proof: Let D be the set of indices $i \in [d]$ with $a_i \neq b_i$. The second part follows from

$$\Pr_{\mathbf{S} \sim \mathcal{S}_i} [\chi_{\mathbf{S}}(a) \neq \chi_{\mathbf{S}}(b)] \leq \Pr_{\mathbf{S} \sim \mathcal{S}_i} [\mathbf{S} \cap D \neq \emptyset] \leq \alpha_i |D| = \alpha_i \|a - b\|_1.$$

The first part follows from

$$\Pr_{\mathbf{S} \sim \mathcal{S}_i} [\chi_{\mathbf{S}}(a) \neq \chi_{\mathbf{S}}(b)] \geq \Pr_{\mathbf{S} \sim \mathcal{S}_i} [|\mathbf{S} \cap D| = 1] = |D| \cdot \alpha_i \cdot (1 - \alpha_i)^{|D|} \geq \frac{\alpha_i |D|}{2}.$$

This finishes the proof of the claim. ■

It follows that the first part of (36) holds for 99% $- o(1)$ fraction of quadtrees T . ■

To prove Theorem 11 it suffices to prove the following lemma:

Lemma 6.3. *Fix a quadtree T , $i \in [h]$, and $\epsilon \in (0, 1)$. Then there is a $O(1/\epsilon) \cdot \text{polylog}(n, d)$ -space linear sketch that, given any (A, B) of size n each, outputs a number η_i such that*

$$\mathcal{I}_i - \frac{\text{end}}{2^i} \leq \eta_i \leq 2\mathcal{I}_i + \frac{\Delta_i d}{2^i} \tag{37}$$

with probability at least $2/3$.

Proof of Theorem 11 assuming Lemma 6.3: The linear sketch starts by sampling a quadtree T from \mathcal{T} and we assume that T satisfies (36), which happens with probability at least 0.9. For each $i \in [h]$, it repeats independently the linear sketch given in Lemma 6.3 for $O(\log d)$ times, and we use η_i to denote the median of $O(\log d)$ numbers returned by the reporting procedure. It follows from a Chernoff bound and a union bound that with probability $1 - o_d(1)$, η_i satisfies (37) for every $i \in [h]$. The reporting procedure returns $\boldsymbol{\eta} = \sum_{i \in [h]} \eta_i + \text{end}$. (35) follows directly from (36). ■

For Theorem 12 the following simpler lemma suffices:

Lemma 6.4. *Fix a quadtree T and any $i \in [h]$. There is a two-round $\text{polylog}(n, d)$ -space linear sketch that, given any (A, B) of size n each, outputs a number η_i such that*

$$\mathcal{I}_i \leq \eta_i \leq 2\mathcal{I}_i + \frac{\Delta_i d}{2^i} \tag{38}$$

with probability at least $2/3$.

Proof of of Theorem 12 using Lemma 6.4: The proof is exactly the same as that of Theorem 11, except that we don't need to add end in the estimate and no longer need the additive error. ■

6.2 Universe Reduction

Now we focus on the proof of Lemma 6.3. Fix a quadtree T and a depth $i \in [h]$. The first step is to perform a routine universe reduction.

Let $m = n^3$. Let H denote the following fixed bipartite graph between $U = [m]$ and $V = [m] \times [m]$: $(u, v) \in H$ if and only if $u = v_1$. Similarly we write $\pi(v)$ to denote the $u \in U$ with $(u, v) \in H$. A *universe reduction* between depths $i - 1$ and i of T consists of two functions $h_{i-1} : [V_{i-1}] \rightarrow [m]$ and $h_i : [V_i] \rightarrow [m]$, which can be used to induce a map \mathbf{v} from $\{0, 1\}^d$ to V : For each point $a \in \{0, 1\}^d$, $\mathbf{v}(a) = v = (v_1, v_2)$ if $h_{i-1}(\mathbf{v}_{i-1}(a)) = v_1$ and $h_i(\mathbf{v}_i(a)) = v_2$. We also use \mathbf{v} to map points to U using $\mathbf{u} : \{0, 1\}^d \rightarrow U$: For each $a \in \{0, 1\}^d$, $\mathbf{u}(a) = \mathbf{v}(a)_1$.

Given the input pair of points (A, B) , we similarly define A_v, B_v and C_v for each node $v \in V$ of H using \mathbf{v} and define A_u, B_u and C_u for each node $u \in U$ using \mathbf{u} (e.g., C_u is the set of $a \in A \cup B$ such that $\mathbf{u}(a) = u$). Let $\Delta = \sum_{v \in V} ||A_v| - |B_v||$. Similarly, whenever $\Delta > 0$, we define the distribution \mathcal{V} supported over V , where each $v \in V$ is sampled with probability $||A_v| - |B_v||/\Delta$. We also define

$$p_{u,v,S} := \Pr_{\mathbf{c}_u \sim C_u, \mathbf{c}_v \sim C_v} [\chi_S(\mathbf{c}_u) \neq \chi_S(\mathbf{c}_v)] \in [0, 1]$$

for each $v \in V$ and $u = \pi(v)$. Lemma 6.3 follows directly from the following lemma, which gives a linear sketch for approximating the expectation of $p(\pi(\mathbf{v}), \mathbf{v}, S)$ with $\mathbf{v} \sim \mathcal{V}$ for a given $S \subseteq [d]$.

Lemma 6.5. *Fix $\mathbf{v} : \{0, 1\}^d \rightarrow V$, $S \subseteq [d]$ and $\epsilon \in (0, 1)$. Then there is a $O(1/\epsilon) \cdot \text{poly}(\log n, \log d)$ space linear sketch that, on any input (A, B) of size n , outputs a number $\eta \in \mathbb{R}$ with the following property: Let $\tau = 1/\log^3 n$. Whenever $\Delta \geq \epsilon n / \log^3 n$, we have*

$$\eta = \mathbf{E}_{\mathbf{v} \sim \mathcal{V}} [p_{\pi(\mathbf{v}), \mathbf{v}, S}] \pm \tau$$

with probability at least $2/3$.

Proof of Lemma 6.3 assuming Lemma 6.5: Given a quadtree T , $i \in [h]$ and $\epsilon \in (0, 1)$, we start by sampling uniformly at random h_{i-1} and h_i and let \mathbf{v} denote the map they induce. We may assume that there is no collision with respect to (A, B) : Every two nodes $u, u' \in V_{i-1}$ with nonempty $C_u, C_{u'}$ in T satisfy $h_{i-1}(u) \neq h_{i-1}(u')$ and every $v, v' \in V_i$ with nonempty $C_v, C_{v'}$ satisfy $h_i(v) \neq h_i(v')$, and note the this event is violated with probability at most $2n^2/m$. When this happens, $\Delta = \Delta_i$ and \mathcal{I}_i can be defined equivalently using \mathbf{v} and (A, B) as follows: $\mathcal{I}_i = 0$ if $\Delta = 0$; otherwise ($\Delta > 0$), we have

$$\mathcal{I}_i = \frac{2\Delta}{\alpha_i} \cdot \mathbf{E}_{\mathbf{v} \sim \mathcal{V}, \mathbf{S} \sim \mathcal{S}_i} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}}].$$

To estimate \mathcal{I}_i , we use the linear sketch described in Figure 1, which uses $O(1/\epsilon) \cdot \text{poly}(\log n, \log d)$ space. For its correctness, we note that the following events happen with probability at least 0.9:

1. $\Delta = (1 \pm 0.01)\Delta$, which happens with probability at least 0.99;
2. If $\Delta \geq \epsilon n / \log^3 n$, then for every set \mathbf{S}_j , we have

$$\eta_j = \mathbf{E}_{\mathbf{v} \sim \mathcal{V}} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}_j}] \pm \frac{1}{\log^3 n},$$

which by a Chernoff bound and a union bound on j happens with probability $1 - o_n(1)$;

Linear Sketch:

1. Run the ℓ_1 -sketch from Theorem 9 on $(|A_v| - |B_v| : v \in V)$ (with $\epsilon = \delta = 0.01$).
2. Draw independently $O(\log^6 n)$ many \mathbf{S}_1, \dots from \mathcal{S}_i . For each \mathbf{S}_j , use \mathbf{v} to run the linear sketch in Lemma 6.5 (with $\tau = 1/\log^3 n$) independently $O(\log n)$ times.

Reporting Procedure:

1. Run the reporting procedure of the ℓ_1 -sketch of Theorem 9 to obtain Δ .
2. For each of the $O(\log^6 n)$ many j , run the reporting procedure to obtain a number for each of the $O(\log n)$ independent runs and let η_j denote their median.
3. If $\Delta < 2\epsilon n/\log^3 n$, output 0; otherwise output

$$\eta = \frac{3\Delta}{\alpha_i} \cdot \left(\text{avg}_j \{\eta_j\} + \frac{1}{8 \log^2 n} \right). \quad (39)$$

Figure 1: Linear sketch for Lemma 6.3

3. When $\Delta > 0$, we have that the

$$\text{avg}_j \left[\mathbf{E}_{\mathbf{v} \sim \mathcal{V}} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}_j}] \right] = \mathbf{E}_{\mathbf{v} \sim \mathcal{V}, \mathbf{S} \sim \mathcal{S}_i} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}}] \pm \frac{1}{\log^3 n},$$

which happens with probability at least 0.99. When the last two items hold, we have

$$\text{avg}_j \{\eta_j\} = \mathbf{E}_{\mathbf{v} \sim \mathcal{V}_i, \mathbf{S} \sim \mathcal{S}_i} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}}] \pm \frac{1}{8 \log^2 n}. \quad (40)$$

We finish the proof by showing that, whenever events above occur, η in (39) satisfies

$$\mathcal{I}_i - \frac{\epsilon n d}{2^i} \leq \eta \leq 2\mathcal{I}_i + \frac{\Delta d}{2^i}. \quad (41)$$

When $\Delta < 2\epsilon n/\log^3 n$ the inequality above is trivial because $\eta = 0$ and the LHS above is negative (using $\Delta \leq 2\Delta$). So we focus on the case when $\Delta \geq 2\epsilon n/\log^3 n$. In this case we have $\Delta \geq \epsilon n/\log^3 n$ and thus, (40) holds. The lower bound of (41) is trivial given (40) and $3\Delta > \Delta$. For the upper bound,

$$\eta \leq \frac{4\Delta}{\alpha_i} \cdot \left(\mathbf{E}_{\mathbf{v} \sim \mathcal{V}_i, \mathbf{S} \sim \mathcal{S}_i} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}}] + \frac{1}{4 \log^2 n} \right) \leq \frac{4\Delta}{\alpha_i} \cdot \mathbf{E}_{\mathbf{v} \sim \mathcal{V}_i, \mathbf{S} \sim \mathcal{S}_i} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}}] + \frac{\Delta d}{2^i} = 2\mathcal{I}_i + \frac{\Delta d}{2^i}.$$

This finishes the proof of the lemma. ■

For the two-round linear sketch, Lemma 6.4 follows from the following lemma:

Lemma 6.6. Fix $\mathbf{v} : \{0, 1\}^d \rightarrow V$ and $S \subseteq [d]$. Let $\tau = 1/\log^3 n$. Then there is a $\text{polylog}(n)$ -space two-round linear sketch that, on any input (A, B) of size n , outputs a number $\boldsymbol{\eta} \in \mathbb{R}$ such that

$$\boldsymbol{\eta} = \mathbf{E}_{\mathbf{v} \sim \mathcal{V}} [p_{\pi(\mathbf{v}), \mathbf{v}, S}] \pm \tau$$

with probability at least $2/3$.

Proof of Lemma 6.4 assuming Lemma 6.6: The first part of sampling h_{i-1} and h_i is the same as the proof of Lemma 6.3. Similar to the linear sketch of Lemma 6.3, we use the ℓ_1 -sketch of Theorem 9 to obtain an estimate Δ of Δ . We also sample $O(\log^6 n)$ many \mathbf{S}_1, \dots from \mathcal{S}_i and for each \mathbf{S}_j , repeat the two-round linear sketch of Lemma 6.6 $O(\log n)$ times to obtain the average as an estimate $\boldsymbol{\eta}_j$ of $\mathbf{E}_{\mathbf{v} \sim \mathcal{V}} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}_j}]$. The total space used is $\text{polylog}(n)$ and similar to the analysis of Lemma 6.3, we have that with probability at least 0.9 , $\Delta = (1 \pm 0.01)\Delta$ and

$$\text{avg}_j \{\boldsymbol{\eta}_j\} = \mathbf{E}_{\mathbf{v} \sim \mathcal{V}, \mathbf{S} \sim \mathcal{S}_i} [p_{\pi(\mathbf{v}), \mathbf{v}, \mathbf{S}}] \pm \frac{1}{8 \log^2 n}.$$

Finally, returning

$$\frac{3\Delta}{\alpha_i} \cdot \left(\text{avg}_j \{\boldsymbol{\eta}_j\} + \frac{1}{8 \log^2 n} \right)$$

satisfies the condition of Lemma 6.4. ■

6.3 Two-Round Linear Sketch of Lemma 6.6

We first give the easier two-round linear sketch of Lemma 6.6 as a warmup. In the first round, we apply the ℓ_1 -sampling linear sketch of Theorem 10 on the vector indexed by $v \in V$, where the entry indexed by v is $|A_v| - |B_v|$. The space used by this linear sketch is $O(\log^2 n)$. Let \mathbf{v} be the output of the linear sketch. Then either \mathbf{v} is the failure symbol \perp , which happens with probability at most $1/3$, or the distribution of $\mathbf{v} \in V$ has distance $1/n$ from \mathcal{V} in total variation.

Assuming $\mathbf{v} = v \in V$ from the first round and letting $u = \pi(v)$, the second round computes

$$p_{u, v, S} = \frac{|C_{u, S}|}{|C_u|} + \frac{|C_{v, S}|}{|C_v|} - \frac{|C_{u, S}|}{|C_u|} \cdot \frac{|C_{v, S}|}{|C_v|},$$

where we write $C_{u, S}$ to denote the set of points $a \in C_u$ with $\chi_S(a) = 1$; we define $C_{v, S}$ similarly. This can be done trivially by asking for these four numbers directly, with $O(\log n)$ space.

In summary, this two-round linear sketch uses $O(\log^2 n)$ -space and achieves the following: It either fails, with probability no more than $1/3$, or returns $p_{\pi(\mathbf{v}), \mathbf{v}, S}$ with \mathbf{v} drawn from a distribution that is $(1/n)$ -close to \mathcal{V} in total variation distance. Lemma 6.6 follows by repeating the two-round linear sketch independently and taking the average at the end.

6.4 Overview of the Linear Sketch of Lemma 6.5

We start with an overview of the linear sketch of Lemma 6.5. Fix $\mathbf{v} : \{0, 1\}^d \rightarrow V$, a set $S \subseteq [d]$, and parameters $\epsilon, \tau \in (0, 1)$. Let (A, B) be the input pair of n points each and we assume without

loss of generality that $\Delta \geq \epsilon n / \log^3 n$ (for the case when $\Delta < \epsilon n \log^3 n$ we only need to make sure that the reporting procedure returns a number and the same space upper bound applies, which will be trivial given the description of the linear sketch later).

We start with some notation. Let Q be the m^2 -dimensional nonnegative vector indexed by $v \in V$ with $Q_v = ||A_v| - |B_v||$ (so $\Delta = \sum_{v \in V} Q_v$). Let P be the m -dimensional nonnegative vector indexed by $u \in U$ with $P_u = \sum_{v: \pi(v)=u} Q_v$ (so $\Delta = \sum_{u \in U} P_u$ as well). We use \mathcal{U} to denote the distribution over U where each $u \in U$ is sampled with probability P_u/Δ ; recall that \mathcal{V} is the distribution where each $v \in V$ is sampled with probability Q_v/Δ . Ideally, $p_{\pi(\mathbf{v}), \mathbf{v}, S}$ with $\mathbf{v} \sim \mathcal{V}$ can be sampled using the following distribution \mathcal{D} supported on edges of H : $(\mathbf{u}, \mathbf{v}) \sim \mathcal{D}$ is sampled by first drawing $\mathbf{u} \sim \mathcal{U}$ and then drawing a $\mathbf{v} \sim \mathcal{V}$ conditioning on $\pi(\mathbf{v}) = \mathbf{u}$ (i.e. each node $v \in V$ with $\pi(v) = u$ is sampled with probability Q_v/P_u), and finally setting $\mathbf{p} = p_{\mathbf{u}, \mathbf{v}, S}$.

Our linear sketch for Lemma 6.5 starts by sampling independently two sequences of numbers $\mathbf{t}_u \sim \text{Exp}(1)$ and $\mathbf{t}_v \sim \text{Exp}(1)$ for each node $u \in U$ and $v \in V$. The presentation of our linear sketch and its analysis will proceed in the following three steps:

1. Given (A, B) and a tuple of positive numbers $(t_u, t_v : u \in U, v \in V)$, we define an edge (u^*, v^*) of H . We prove in Lemma 6.7 that, $(\mathbf{u}^*, \mathbf{v}^*)$ defined using $(\mathbf{t}_u, \mathbf{t}_v : u \in U, v \in V)$ when $\mathbf{t}_u, \mathbf{t}_v \sim \text{Exp}(1)$ independently has the same distribution as \mathcal{D} described above.
2. Next we define an event \mathcal{E} on a tuple of positive numbers $(t_u, t_v : u \in U, v \in V)$ with respect to (A, B) , and show that $(\mathbf{t}_u, \mathbf{t}_v : u \in U, v \in V)$ when $\mathbf{t}_u, \mathbf{t}_v \sim \text{Exp}(1)$ independently satisfies the event \mathcal{E} with respect to (A, B) with high probability (Lemma 6.8).
3. Finally, given \mathbf{v}, S , a tuple of positive numbers $(t_u, t_v : u \in U, v \in V)$ and $\epsilon, \tau \in (0, 1)$, we give a low-space linear sketch (Lemma 6.9) that, given any input pair (A, B) of size n each, returns a number $\boldsymbol{\eta} \in [0, 1]$ and satisfies the following property: Whenever $\Delta \geq \epsilon n / \log^3 n$ and $(t_u, t_v : u \in U, v \in V)$ satisfies \mathcal{E} with respect to (A, B) , we have $\boldsymbol{\eta} = p_{\mathbf{u}^*, \mathbf{v}^*, S} \pm \tau$ with probability at least $2/3$.

Lemma 6.5 follows (by repeating $O(1/\tau^2)$ many rounds: in each round we sample fresh $(\mathbf{t}_u, \mathbf{t}_v)$ and then run the linear sketch to obtain a number, and output the average at the end); see the proof of Lemma 6.5 at the end of this subsection.

For the first step, given a tuple of positive numbers $(t_u, t_v : u \in U, v \in V)$, we define $u^* \in U$ as the node $u \in U$ that maximizes P_u/t_u (breaking ties by taking the smallest such u). Given $\Delta > 0$, we must have $P_{u^*} > 0$. Next we define $v^* \in V$ as the $v \in V$ among those with $\pi(v) = u^*$ that maximizes Q_v/t_v (again breaking ties by taking the smallest such v). Similarly we have $Q_{v^*} > 0$.

The following lemma shows that $(\mathbf{u}^*, \mathbf{v}^*)$ obtained from $\mathbf{t}_u, \mathbf{t}_v \sim \text{Exp}(1)$ is distributed as \mathcal{D} .

Lemma 6.7. *Let (A, B) be a pair of n points. With $\mathbf{t}_u \sim \text{Exp}(1)$ and $\mathbf{t}_v \sim \text{Exp}(1)$ independently for each $u \in U$ and $v \in V$, we have that the distribution of $(\mathbf{u}^*, \mathbf{v}^*)$ is the same as \mathcal{D} .*

Proof: This follows from Fact 5.1 and Fact 5.3. ■

Next we describe the event \mathcal{E} in Figure 2, where we also introduce two new parameters γ and β .

Let γ and β be the following two parameters:

$$\gamma \stackrel{\text{def}}{=} \frac{\tau}{\log n} = \frac{1}{\log^4 n} \quad \text{and} \quad \beta \stackrel{\text{def}}{=} \left\lceil \frac{\log^5 n}{\epsilon \tau \gamma^3} \right\rceil = \frac{\text{poly}(\log n)}{\epsilon}. \quad (42)$$

We say a tuple of positive numbers $(t_u, t_v : u \in U, v \in V)$ satisfy event \mathcal{E} with respect to (A, B) if the following four events $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$ and \mathcal{E}_4 hold:

- Event \mathcal{E}_1 : The numbers $(t_u : u \in U)$ satisfy

$$\sum_{u \in U} \frac{P_u}{t_u} \leq \frac{4 \log(n/\gamma)}{\gamma} \cdot \Delta.$$

Moreover, u^* satisfies

$$\frac{P_{u^*}}{t_{u^*}} \geq \gamma \Delta \quad \text{and} \quad \frac{P_{u^*}}{t_{u^*}} \geq (1 + \gamma) \cdot \max_{u \neq u^*} \left\{ \frac{P_u}{t_u} \right\}.$$

- Event \mathcal{E}_2 : The numbers $(t_u, t_v : u \in U, v \in V)$ satisfy

$$\sum_{v \in V} \frac{Q_v}{t_{\pi(v)} t_v} \leq \frac{4 \log(n/\gamma)}{\gamma} \cdot \sum_{u \in U} \frac{P_u}{t_u}.$$

Moreover, v^* satisfies

$$\frac{Q_{v^*}}{t_{v^*}} \geq \gamma P_{u^*} \quad \text{and} \quad \frac{Q_{v^*}}{t_{v^*}} \geq (1 + \gamma) \cdot \max_{v: \pi(v)=u^*} \left\{ \frac{Q_v}{t_v} \right\}.$$

- Event \mathcal{E}_3 : The numbers $(t_u : u \in U)$ satisfy that

$$\sum_{u \in U} \frac{|C_u|}{t_u} \leq \frac{4 \log(n/\gamma)}{\gamma} \cdot n$$

and the ℓ_2 -norm of $(|C_u|/t_u : u \in U)_{-\beta}$ is at most $12n/\sqrt{\beta}$.

- Event \mathcal{E}_4 : The ℓ_2 -norm of $(|C_v|/(t_{\pi(v)} t_v) : v \in V)_{-\beta}$ is at most

$$\frac{12}{\sqrt{\beta}} \cdot \sum_{u \in U} \frac{|C_u|}{t_u}.$$

Figure 2: Event \mathcal{E} on $(t_u, t_v : u \in U, v \in V)$ with respect to (A, B)

Lemma 6.8. *Let (A, B) be a pair of n points. With $\mathbf{t}_u \sim \text{Exp}(1)$ and $\mathbf{t}_v \sim \text{Exp}(1)$ independently for each $u \in U$ and $v \in V$, \mathcal{E} holds with respect to (A, B) with probability at least $1 - O(\gamma) - e^{-\Omega(\beta)}$.*

Proof: The first part of event \mathcal{E}_1 follows from Lemma 5.4, and the second part follows from Lemma 5.5. Similarly, the first part of event \mathcal{E}_2 follows from Lemma 5.4 on the randomness of \mathbf{t}_v . The first part of \mathcal{E}_3 follows again by Lemma 5.4, and the second part of event \mathcal{E}_3 uses Lemma 5.6 (with $c = 1$). For event \mathcal{E}_4 , one can first use Lemma 5.6 (with $c = 1$) and randomness of \mathbf{t}_v to upperbound the ℓ_2 -norm of $(|C_v|/(\mathbf{t}_{\pi(v)}\mathbf{t}_v : v \in V))_{-\beta}$ by

$$\frac{12}{\sqrt{\beta}} \cdot \sum_{v \in V} \frac{|C_v|}{\mathbf{t}_{\pi(v)}} = \frac{12}{\sqrt{\beta}} \cdot \sum_{u \in U} \frac{|C_u|}{\mathbf{t}_u}.$$

Applying a union bound finishes the proof of the lemma. ■

Finally we state the performance guarantee of the main linear sketch:

Lemma 6.9. *Fix $\mathbf{v} : \{0, 1\}^d \rightarrow V$, $S \subseteq [d]$, a tuple of positive numbers $(t_u, t_v : u \in U, v \in V)$ and two parameters $\epsilon, \tau \in (0, 1)$. There is a $O(1/\epsilon) \cdot \text{poly}(\log n, \log d, 1/\tau)$ -space linear sketch that, on input (A, B) of size n each, outputs a number $\boldsymbol{\eta} \in [0, 1]$ with the following property: When $\Delta \geq \epsilon n / \log^3 n$ and (t_u, t_v) satisfies \mathcal{E} with respect to (A, B) , we have $\boldsymbol{\eta} = p_{u^*, v^*, S} \pm \tau$ with probability at least $2/3$.*

Proof of Lemma 6.5 assuming Lemma 6.9: The linear sketch repeats the following $O(1/\tau^2)$ rounds:

- Draw fresh $\mathbf{t}_u, \mathbf{t}_v \sim \text{Exp}(1)$ and repeat the linear sketch of Lemma 6.9 independently (setting τ in Lemma 6.9 to be a quarter of the τ in Lemma 6.5) for $O(\log(1/\tau))$ times.

For each round of the linear sketch, the reporting procedure uses the reporting procedure of Lemma 6.9 to get $O(\log(1/\tau))$ numbers in $[0, 1]$ and compute their median (in $[0, 1]$). It finally outputs the average of these $O(1/\tau^2)$ many medians.

To prove the correctness, it suffices to show that the expectation of $\boldsymbol{\eta}$ (the median) reported from each round is within $\mathbf{E}_{\mathbf{v} \sim \mathcal{V}}[p_{\pi(\mathbf{v}), \mathbf{v}, S}] \pm (\tau/2)$. To see this is the case, we compare the expectation of $\boldsymbol{\eta}$ with the ideal process of sampling first $(\mathbf{u}^*, \mathbf{v}^*) \sim \mathcal{D}$ and returning $p_{\mathbf{u}^*, \mathbf{v}^*, S}$, whose expectation is exactly $\mathbf{E}_{\mathbf{v} \sim \mathcal{V}}[p_{\pi(\mathbf{v}), \mathbf{v}, S}]$. First, the linear sketch may fail because $(\mathbf{t}_u, \mathbf{t}_v)$ sampled in this round does not satisfy \mathcal{E} , which happens with probability $o(\tau)$ and may shift the expectation by no more than $o(\tau)$. Second, the median may fail to fall inside the $\pm\tau/4$ interval centered at $p_{\mathbf{u}^*, \mathbf{v}^*, S}$, which happens with probability $\tau/8$ (because we repeat $O(\log(1/\tau))$ times in each round), which can shift the expectation by no more than $\tau/8$. Finally, even if the median is inside the interval $p_{\mathbf{u}^*, \mathbf{v}^*, S} \pm (\tau/4)$, the $\pm\tau/4$ may shift the expectation by $\tau/4$. Summing up these three cases, we conclude that the expectation can be shifted by no more than $\tau/2$. ■

6.5 Three Linear Sketches and their Performance Guarantees

Our linear sketch for Lemma 6.9 consists of three linear sketches whose goals are to recover node u^* , recover node v^* , and estimate $p_{u^*, v^*, S}$, respectively. For each $i \in [3]$, we write LS_i to denote the encoding algorithm of the i th linear sketch which uses its own randomness \mathbf{a}_i to generate the linear sketch $\text{LS}_i((A, B), \mathbf{a}_i)$. Each LS_i is paired with a decoding algorithm ALG_i (note that LS_i and ALG_i are deterministic) which has the following input and output:

1. ALG_1 takes \mathbf{a}_1 and $\text{LS}_1((A, B), \mathbf{a}_1)$ as input and outputs a vertex in U ;
2. ALG_2 takes a node in U , \mathbf{a}_2 and $\text{LS}_2((A, B), \mathbf{a}_2)$ as input and outputs a vertex in V ;
3. ALG_3 takes an edge in H , \mathbf{a}_3 and $\text{LS}_3((A, B), \mathbf{a}_3)$ as input and outputs a number in $[0, 1]$.

We now state their performance guarantees (where we fix $\mathbf{v}, S, (t_u, t_v), \epsilon$ and τ as in Lemma 6.9).

Lemma 6.10. *There is a $\text{poly}(\log n, 1/\gamma)$ -space linear sketch $(\text{LS}_1, \text{ALG}_1)$ which outputs a node in U and satisfies the following property: Whenever $\Delta > 0$ and (t_u, t_v) satisfies \mathcal{E}_1 with respect to (A, B) , ALG_1 returns u^* with probability at least $1 - 1/\text{poly}(n)$.*

Lemma 6.11. *There is a $\text{poly}(\log n, 1/\gamma)$ -space linear sketch $(\text{LS}_2, \text{ALG}_2)$ which outputs a node in V and satisfies the following properties: (1) If u is node given to ALG_2 , then the latter always returns a node \mathbf{v} with $\pi(\mathbf{v}) = u$; (2) Whenever $\Delta > 0$ and (t_u, t_v) satisfies \mathcal{E}_1 and \mathcal{E}_2 with respect to (A, B) , ALG_2 running on u^* , \mathbf{a}_2 and $\text{LS}_2((A, B), \mathbf{a}_2)$ returns v^* with probability at least $1 - \text{poly}(n)$.*

Lemma 6.12. *There is a $O(\beta \log n)$ -space linear sketch $(\text{LS}_3, \text{ALG}_3)$ which always outputs a number in $[0, 1]$ and satisfies the following property: Whenever $\Delta \geq \epsilon n / \log^3 n$ and (t_u, t_v) satisfies \mathcal{E}_3 and \mathcal{E}_4 with respect to (A, B) , ALG_3 running on (u^*, v^*) , \mathbf{a}_3 and $\text{LS}_3((A, B), \mathbf{a}_3)$ returns a number η that satisfies $\eta = p_{u^*, v^*, S} \pm \tau$ with probability at least $1 - 1/\text{poly}(n)$.*

Proof of Lemma 6.9 assuming Lemmas 6.10, 6.11 and 6.12: The main linear sketch of Lemma 6.9 flips coins $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$ and then encodes (A, B) as $(\text{LS}_i((A, B), \mathbf{a}_i) : i \in [3])$. The reporting procedure runs ALG_1 on \mathbf{a}_1 and $\text{LS}_1((A, B), \mathbf{a}_1)$ to obtain a vertex $\mathbf{u} \in U$, runs ALG_2 on \mathbf{u}, \mathbf{a}_2 and $\text{LS}_2((A, B), \mathbf{a}_2)$ to obtain a vertex $\mathbf{v} \in V$, and runs ALG_3 on $(\mathbf{u}, \mathbf{v}), \mathbf{a}_3$ and $\text{LS}_3((A, B), \mathbf{a}_3)$ to obtain η . By a union bound on errors of the three lemmas, we have $\eta = p_{u^*, v^*, S} \pm \tau$ with probability at least $2/3$. \blacksquare

6.6 Linear Sketch of Lemma 6.10

Let $\eta = O(\gamma^2 / \log n) = 1/\text{poly}(\log n)$. We describe $(\text{LS}_1, \text{ALG}_1)$ in Figure 3, which uses space

$$O\left(\frac{\log n}{\gamma^2}\right) \cdot O\left(\frac{\log n}{\eta^2}\right) = \text{poly}(\log n).$$

We start by showing that $\widehat{\mathbf{S}}^{(j)}$ is close to $\mathbf{S}^{(j)}$ in ℓ_∞ distance for each j . For this purpose, it suffices to upperbound $\|\mathbf{S}_{-1/\eta^2}^{(j)}\|_2$. Note that each entry of $\mathbf{S}^{(j)}$ is a sum with Cauchy random variables as coefficients. Using the 1-stability of the Cauchy distribution, each entry $\mathbf{S}_u^{(j)}$ can be equivalently generated by drawing a Cauchy variable α_u and setting

$$\mathbf{S}_u^{(j)} = \alpha_u \cdot \sum_{v:\pi(v)=u} \frac{Q_v}{t_u} = \alpha_u \cdot \frac{P_u}{t_u}.$$

Applying Lemma 5.6 and using the fact that $\Pr[\alpha \geq t] \leq 1/t$, we have from \mathcal{E}_1 that

$$\|\mathbf{S}_{-1/\eta^2}^{(j)}\|_2 \leq O(\eta) \cdot \sum_{u \in U} \frac{P_u}{t_u} \leq O\left(\frac{\eta \log n}{\gamma}\right) \cdot \Delta$$

Linear Sketch LS₁: We repeat the following procedure for $j = 1, \dots, O(\log n/\gamma^2)$ times:

1. For each $v \in V$, draw an independent Cauchy random variable α_v .
2. Define the following vector $\mathbf{S}^{(j)}$ indexed by $u \in U$: For each $u \in U$, let

$$\mathbf{S}_u^{(j)} := \sum_{v:\pi(v)=u} \alpha_v \cdot \frac{|A_v| - |B_v|}{t_u}.$$

3. Run the Count-Sketch of Theorem 8 on $\mathbf{S}^{(j)}$ (with $\epsilon = \eta$ and $n = m$).

Reporting Procedure ALG₁: For each j , use the reporting procedure of the Count-Sketch of Theorem 8 to compute a vector $\widehat{\mathbf{S}}^{(j)}$. Let $\widehat{\mathbf{S}}$ be the vector with

$$\widehat{\mathbf{S}}_u := \text{median} \left\{ \widehat{\mathbf{S}}_u^{(j)} : j = 1, \dots, O(\log n/\gamma^2) \right\}, \quad \text{for each } u \in U.$$

Output $\text{argmax}_{u \in U} \{\widehat{\mathbf{S}}_u\}$.

Figure 3: The linear sketch (LS₁, ALG₁) of Lemma 6.10

with probability $1 - 1/\text{poly}(n)$. Assuming this holds for every j , it follows from Theorem 8 that

$$\left\| \widehat{\mathbf{S}}^{(j)} - \mathbf{S}^{(j)} \right\| \leq O\left(\frac{\eta^2 \log n}{\gamma}\right) \cdot \Delta$$

with probability $1 - 1/\text{poly}(n)$. As a result (and given that we only apply a union bound on $O(\log n)$ many events), with probability at least $1 - 1/\text{poly}(n)$, we have for every $u \in U$:

$$\widehat{\mathbf{S}}_u = \text{median} \left\{ \mathbf{S}_u^{(j)} : j = 1, \dots, O(\log n/\gamma^2) \right\} \pm O\left(\frac{\eta^2 \log n}{\gamma}\right) \cdot \Delta.$$

On the other hand, for each $u \in U$, we have from Theorem 9 that

$$\text{median} \left\{ \mathbf{S}_u^{(j)} : j = 1, \dots, O(\log n/\gamma^2) \right\} = \text{median}(|\mathcal{C}|) \cdot \left(1 \pm \frac{\gamma}{4}\right) \frac{P_u}{t_u}$$

with probability at least $1 - \text{poly}(n)$. Assuming all these events occur and using \mathcal{E}_1 , every $u \neq u^*$ satisfies (letting c_0 be the constant $\text{median}(|\mathcal{C}|)$ below)

$$\widehat{\mathbf{S}}_u \leq c_0 \cdot \left(1 + \frac{\gamma}{4}\right) \cdot \frac{1}{1 + \gamma} \cdot \frac{P_{u^*}}{t_{u^*}} + O\left(\frac{\eta^2 \log n}{\gamma}\right) \cdot \Delta \leq c_0 \cdot \left(1 - \frac{\gamma}{2}\right) \cdot \frac{P_{u^*}}{t_{u^*}} + O\left(\frac{\eta^2 \log n}{\gamma}\right) \cdot \Delta.$$

and u^* satisfies

$$\widehat{\mathbf{S}}_{u^*} \geq c_0 \cdot \left(1 - \frac{\gamma}{4}\right) \frac{P_{u^*}}{t_{u^*}} - O\left(\frac{\eta^2 \log n}{\gamma}\right) \cdot \Delta.$$

We have

$$\widehat{\mathbf{S}}_{u^*} - \widehat{\mathbf{S}}_u \geq c_0 \cdot \frac{\gamma}{4} \cdot \frac{P_{u^*}}{t_{u^*}} - O\left(\frac{\eta^2 \log n}{\gamma}\right) \cdot \Delta > 0$$

using \mathcal{E}_1 and our choice of $\eta = O(\gamma^2/\log n)$.

Linear Sketch LS₂: Let S be the vector with

$$S_v := \frac{|A_v| - |B_v|}{t_{\pi(u)} t_v}.$$

Run the Count-Sketch of Theorem 8 on S with $\epsilon = \eta$ and $n = m^2$.

Reporting Procedure ALG₂ (given $u \in U$): Use the reporting procedure of the Count-Sketch of Theorem 8 to compute a vector $\widehat{\mathbf{S}}$, and output $\operatorname{argmax}_{v:\pi(v)=u} \{|\widehat{\mathbf{S}}_v|\}$.

Figure 4: The linear sketch (LS₂, ALG₂) of Lemma 6.11

6.7 Linear Sketch of Lemma 6.11

Let $\eta = O(\gamma^5 / \log^2 n)$. The pair (LS₂, ALG₂) is given in Figure 4. Assume both \mathcal{E}_1 and \mathcal{E}_2 hold, and assume that the node given to ALG₂ is u^* . Note that by \mathcal{E}_1 and \mathcal{E}_2 we have

$$\|S\|_1 = \sum_{v \in V} \frac{Q_v}{t_{\pi(v)} t_v} \leq O\left(\frac{\log n}{\gamma}\right) \cdot \sum_{u \in U} \frac{P_u}{t_u} \leq O\left(\frac{\log^2 n}{\gamma^2}\right) \cdot \Delta.$$

As a result, we have

$$\|\widehat{\mathbf{S}} - S\|_\infty \leq O\left(\frac{\eta \log^2 n}{\gamma^2}\right) \cdot \Delta.$$

Every v with $\pi(v) = u^*$ and $v \neq v^*$ satisfies

$$|\widehat{\mathbf{S}}_v| \leq \frac{Q_v}{t_{u^*} t_v} + O\left(\frac{\eta \log^2 n}{\gamma^2}\right) \cdot \Delta \quad \text{and} \quad |\widehat{\mathbf{S}}_{v^*}| \geq \frac{Q_{v^*}}{t_{u^*} t_{v^*}} - O\left(\frac{\eta \log^2 n}{\gamma^2}\right) \cdot \Delta.$$

As a result, using \mathcal{E}_1 and \mathcal{E}_2 we have

$$|\widehat{\mathbf{S}}_{v^*}| - |\widehat{\mathbf{S}}_v| \geq \frac{1}{t_{u^*}} \cdot \left(1 - \frac{1}{1 + \gamma}\right) \cdot \frac{Q_{v^*}}{t_{v^*}} - O\left(\frac{\eta \log^2 n}{\gamma^2}\right) \cdot \Delta \geq \Omega(\gamma^3) \cdot \Delta - O\left(\frac{\eta \log^2 n}{\gamma^2}\right) \cdot \Delta > 0$$

given our choice of $\eta = O(\gamma^5 / \log^2 n)$.

6.8 Linear Sketch of Lemma 6.12

Given S and $u \in U$, recall that $C_{u,S}$ denotes the set of points $a \in A \cup B$ with $\chi_S(a) = 1$, and $C_{v,S}$ is defined similarly. Then $p_{u,v,S}$ can be expressed as

$$\frac{|C_{u,S}|}{|C_u|} + \frac{|C_{v,S}|}{|C_v|} - \frac{|C_{u,S}|}{|C_u|} \cdot \frac{|C_{v,S}|}{|C_v|}.$$

We present the pair (LS₃, ALG₃) in Figure 5, which approximates the four numbers $|C_{u,S}|$, $|C_u|$ and $|C_{v,S}|$, $|C_v|$ and uses them to obtain an estimate of $p_{u,v,S}$. For correctness, it suffices to upperbound

$$\left| \frac{\widehat{\mathbf{S}}_{u^*}^{(2)}}{\widehat{\mathbf{S}}_{u^*}^{(1)}} - \frac{|C_{u^*,S}|}{|C_{u^*}|} \right| \quad \text{and} \quad \left| \frac{\widehat{\mathbf{S}}_{v^*}^{(4)}}{\widehat{\mathbf{S}}_{v^*}^{(3)}} - \frac{|C_{v^*,S}|}{|C_{v^*}|} \right|.$$

Linear Sketch LS₃: Let $S^{(1)}$ and $S^{(2)}$ the two vectors indexed by $u \in U$ with

$$S_u^{(1)} := \frac{|C_u|}{t_u} \quad \text{and} \quad S_u^{(2)} = \frac{|C_{u,S}|}{t_u}, \quad \text{for each } u \in U.$$

Let $S^{(3)}$ and $S^{(4)}$ be the two vectors indexed by $v \in V$ with

$$S_v^{(3)} := \frac{|C_v|}{t_{\pi(v)}t_v} \quad \text{and} \quad S_v^{(4)} = \frac{|C_{v,S}|}{t_{\pi(v)}t_v}, \quad \text{for each } v \in V.$$

Run the Count-Sketch of Theorem 8 on each $S^{(i)}$ (with $\epsilon = 1/\sqrt{\beta}$ and $n = m$ or m^2).

Reporting Procedure ALG₃ (given an edge (u, v) of H): Use the reporting procedure of the Count-Sketch of Theorem 8 to compute vectors $\widehat{\mathbf{S}}^{(i)}$, $i \in [4]$. Output

$$\frac{\widehat{\mathbf{S}}_u^{(2)}}{\widehat{\mathbf{S}}_u^{(1)}} + \frac{\widehat{\mathbf{S}}_v^{(4)}}{\widehat{\mathbf{S}}_v^{(3)}} - \frac{\widehat{\mathbf{S}}_u^{(2)}}{\widehat{\mathbf{S}}_u^{(1)}} \cdot \frac{\widehat{\mathbf{S}}_v^{(4)}}{\widehat{\mathbf{S}}_v^{(3)}} \quad (43)$$

after truncating it to $[0, 1]$.

Figure 5: The linear sketch (LS₃, ALG₃) of Lemma 6.12

We start with the first one. Note that by \mathcal{E}_3 we have

$$\|S_{-\beta}^{(2)}\|_2 \leq \|S_{-\beta}^{(1)}\|_2 = O\left(\frac{n}{\sqrt{\beta}}\right).$$

By Theorem 8, with probability at least $1 - \text{poly}(n)$, we have

$$\left\| \mathbf{S}^{(2)} - S^{(2)} \right\|_{\infty}, \left\| \mathbf{S}^{(1)} - S^{(1)} \right\|_{\infty} \leq O\left(\frac{n}{\beta}\right).$$

Focusing on $\widehat{\mathbf{S}}_{u^*}^{(1)}$ first, we claim that

$$\left| \widehat{\mathbf{S}}_{u^*}^{(1)} - S_{u^*}^{(1)} \right| \leq \frac{\tau}{16} \cdot S_{u^*}^{(1)}$$

This follows from our choice of $\beta = \log^5 n / (\tau \epsilon \gamma^3)$ and that

$$S_{u^*}^{(1)} \geq \frac{P_{u^*}}{t_{u^*}} \geq \gamma \Delta \geq \frac{\epsilon \gamma}{\log^3 n} \cdot n.$$

Therefore, we have

$$\frac{\mathbf{S}_{u^*}^{(2)}}{\mathbf{S}_{u^*}^{(1)}} = \frac{S_{u^*}^{(2)}}{S_{u^*}^{(1)}} \pm O\left(\frac{n}{\beta} \cdot \frac{1}{S_{u^*}^{(1)}}\right) = \left(1 \pm \frac{\tau}{8}\right) \cdot \frac{S_{u^*}^{(2)}}{S_{u^*}^{(1)}} \pm O\left(\frac{n}{\beta} \cdot \frac{\log^3 n}{\epsilon \gamma n}\right) = \frac{S_{u^*}^{(2)}}{S_{u^*}^{(1)}} \pm \frac{\tau}{6}.$$

The analysis for the other ratio is similar. First we have from \mathcal{E}_3 and \mathcal{E}_4 that

$$\|S_{-\beta}^{(4)}\|_2 \leq \|S_{-\beta}^{(3)}\|_2 \leq O\left(\frac{n \log n}{\sqrt{\beta\gamma}}\right).$$

By Theorem 8, with probability at least $1 - \text{poly}(n)$, we have

$$\|\mathbf{S}^{(3)} - S^{(3)}\|_\infty, \|\mathbf{S}^{(4)} - S^{(4)}\|_\infty \leq O\left(\frac{n \log n}{\beta\gamma}\right).$$

Focusing on $\widehat{\mathbf{S}}_{v^*}^{(3)}$ first, we have similarly that

$$\left|\widehat{\mathbf{S}}_{v^*}^{(3)} - S_{v^*}^{(3)}\right| \leq \frac{\tau}{16} \cdot S_{u^*}^{(3)}$$

using our choice of $\beta = \log^5 n / (\tau \epsilon \gamma^3)$ and that

$$S_{v^*}^{(3)} = \frac{Q_{v^*}}{t_{u^*} t_{v^*}} \geq \gamma^2 \Delta \geq \frac{\epsilon \gamma^2}{\log^3 n} \cdot n.$$

Therefore, we have

$$\frac{\mathbf{S}_{v^*}^{(4)}}{\mathbf{S}_{v^*}^{(3)}} = \frac{S_{v^*}^{(4)}}{S_{v^*}^{(3)}} \pm O\left(\frac{n \log n}{\beta\gamma} \cdot \frac{1}{S_{v^*}^{(3)}}\right) = \left(1 \pm \frac{\tau}{8}\right) \cdot \frac{S_{v^*}^{(4)}}{S_{v^*}^{(3)}} \pm O\left(\frac{n \log n}{\beta\gamma} \cdot \frac{\log^3 n}{\epsilon \gamma^2 n}\right) = \frac{S_{v^*}^{(4)}}{S_{v^*}^{(3)}} \pm \frac{\tau}{6}.$$

As a result we have (43) is

$$\left(\frac{S_{u^*}^{(2)}}{S_{u^*}^{(1)}} \pm \frac{\tau}{6}\right) + \left(\frac{S_{v^*}^{(4)}}{S_{v^*}^{(3)}} \pm \frac{\tau}{6}\right) - \left(\frac{S_{u^*}^{(2)}}{S_{u^*}^{(1)}} \pm \frac{\tau}{6}\right) \cdot \left(\frac{S_{v^*}^{(4)}}{S_{v^*}^{(3)}} \pm \frac{\tau}{6}\right) = \frac{S_{u^*}^{(2)}}{S_{u^*}^{(1)}} + \frac{S_{v^*}^{(4)}}{S_{v^*}^{(3)}} - \frac{S_{u^*}^{(2)}}{S_{u^*}^{(1)}} \cdot \frac{S_{v^*}^{(4)}}{S_{v^*}^{(3)}} \pm \tau.$$

7 Linear Sketches for MST

Fix $n, d \in \mathbb{N}$, and let $X = \{x_1, \dots, x_n\} \subset \{0, 1\}^d$. To settle on some notation, we let $h \stackrel{\text{def}}{=} \lceil \log_2 d \rceil$, and we consider a quadtree T of depth $h + 1$, whose levels are indexed by $i \in \{0, \dots, h\}$, starting at the root at depth $i = 0$, to the leaves at depth h . For $i \in \{0, \dots, h\}$, we let V_i be the set of nodes of T at depth i . For each $i \in \{0, \dots, h\}$, T induces a map $\mathbf{v}_i: \{0, 1\}^d \rightarrow V_i$, such that $\mathbf{v}_0(a), \mathbf{v}_1(a), \dots, \mathbf{v}_h(a)$ is the path of a point $a \in \{0, 1\}^d$ down the quadtree T . We say a maps to $v \in V_i$ if $\mathbf{v}_i(a) = v$.

For a node $v \in V_i$ for $i > 0$, we let $\pi(v)$ be the parent of v in T (which lies at depth $i - 1$). The subset $L_i \subset V_i$ is the set of nodes $v \in V_i$ which are *non-empty with respect to X* , i.e., some point $x \in X$ mapped to node v ; for $v \in L_i$, the set $X_v \subset X$ consists of the set of points of X which map to v . For a node $u \in V_{i-1}$, we let $C(u) \subset L_i$ be the set of child nodes of u . Our starting point is the fact that with probability at least 0.99 over the draw of $\mathbf{T} \sim \mathcal{T}$, two things occur: the first is that within any node $v \in L_i$, and any two $x, y \in X_v$, we have $\|x - y\|_1 \leq O(d \log n / 2^i)$. The second is that we can approximate the cost of the minimum spanning tree by the following sampling experiment.

1. For each $u \in L_{i-1}$, we sample a *parent representative* \mathbf{c}_u from X_u by sampling $\mathbf{v}' \sim C(u)$, and then sampling $\mathbf{c}_u \sim X_{\mathbf{v}'}$.
2. For each $v \in L_i$, we sample a *child representative* $\mathbf{r}_v \sim X_v$.

Then, we have that for every choice of \mathbf{c}_u and \mathbf{r}_v , we have

$$\text{MST}(X) \leq \sum_{i=1}^h \mathbf{1}\{|L_i| > 1\} \sum_{v \in L_i} \|\mathbf{r}_v - \mathbf{c}_{\pi(v)}\|_1 = \sum_{i=1}^h \mathbf{1}\{|L_i| > 1\} \cdot |L_i| \mathbf{E}_{\mathbf{v} \sim L_i} [\|\mathbf{r}_v - \mathbf{c}_{\pi(v)}\|_1] \quad (44)$$

In order to see why the first inequality in (44) holds, consider the following way to recursively build a spanning tree from a quadtree T : inductively assume that each node v contains a spanning tree of X_v , then if a node v is a child of u , place an edge between \mathbf{r}_v and \mathbf{c}_u , thereby connecting all spanning trees of children of u . On the other hand, the expectation of the above quantity (over the random choices of \mathbf{c}_u and \mathbf{r}_v) upper bounds $\text{MST}(X)$, up to a factor of $\tilde{O}(\log n)$. Specifically, by triangle inequality, as well as the distribution of \mathbf{c}_u , we have

$$\begin{aligned} & \sum_{i=1}^h \mathbf{1}\{|L_i| > 1\} \cdot |L_i| \mathbf{E}_{\mathbf{v} \sim L_i} \left[\mathbf{E}_{\mathbf{r}_v, \mathbf{c}_{\pi(v)}} [\|\mathbf{r}_v - \mathbf{c}_{\pi(v)}\|_1] \right] \\ & \leq \sum_{i=1}^h \mathbf{1}\{|L_i| > 1\} |L_i| \mathbf{E}_{\mathbf{v} \sim L_i} \left[\mathbf{E}_{\mathbf{r}_{\pi(v)} \sim X_{\pi(v)}} \left[\mathbf{E}_{\mathbf{r}_v, \mathbf{c}_{\pi(v)}} [\|\mathbf{r}_v - \mathbf{r}_{\pi(v)}\|_1 + \|\mathbf{r}_{\pi(v)} - \mathbf{c}_{\pi(v)}\|_1] \right] \right] \\ & = 2 \sum_{i=1}^h \mathbf{1}\{|L_i| > 1\} |L_i| \mathbf{E}_{\mathbf{v} \sim L_i} \left[\mathbf{E}_{\substack{\mathbf{r}_v \sim X_v \\ \mathbf{r}_{\pi(v)} \sim X_{\pi(v)}}} [\|\mathbf{r}_v - \mathbf{r}_{\pi(v)}\|_1] \right]. \end{aligned}$$

Furthermore, we know that this latter quantity satisfies

$$\sum_{i=1}^h \mathbf{1}\{|L_i| > 1\} \cdot |L_i| \mathbf{E}_{\mathbf{v} \sim L_i} \left[\mathbf{E}_{\substack{\mathbf{r}_v \sim X_v \\ \mathbf{r}_{\pi(v)} \sim X_{\pi(v)}}} [\|\mathbf{r}_v - \mathbf{r}_{\pi(v)}\|_1] + \frac{d}{2^i} \right] \leq \tilde{O}(\log n) \cdot \text{MST}(X).$$

Our goal is to design a linear sketch which uses $\text{poly}(\log d, \log n)$ space, and for every $i \in \{1, \dots, h\}$, outputs two values $\hat{\ell}_i, \hat{\mu}_i \in \mathbb{R}$, satisfying

$$|L_i| \leq \hat{\ell}_i \leq 1.5 \cdot |L_i| \quad \text{and} \quad \hat{\mu}_i \leq \mathbf{E}_{\substack{\mathbf{v} \sim L_i \\ \mathbf{r}_v, \mathbf{c}_{\pi(v)}}} [\|\mathbf{r}_v - \mathbf{c}_{\pi(v)}\|_1] \leq \hat{\mu}_i + \frac{d}{2 \cdot 2^i}. \quad (45)$$

If, for every $i \in \{1, \dots, h\}$, the estimates $\hat{\ell}_i$ and $\hat{\mu}_i$ all satisfy (45), then its not hard to see that

$$\sum_{i=1}^h \mathbf{1}\{\hat{\ell}_i > 1.5\} \cdot \hat{\ell}_i \cdot \left(\hat{\mu}_i + \frac{d}{2^i} \right)$$

is an $\tilde{O}(\log n)$ -approximation to $\text{MST}(X)$. We note that estimating $|L_i|$ is by now standard in the literature, via an ℓ_0 -sketch, and hence the main difficulty is producing estimates $\hat{\mu}_i$. Furthermore, if indeed every $x, y \in X_u$ at depth $i-1 \in \{0, \dots, h-1\}$ satisfies $\|x - y\|_1 \leq O(d \log n / 2^i)$, then

it suffices to design a linear sketch using $\text{poly}(\log n, \log d)$ space which can generate (approximate) samples from $\|\mathbf{r}_{\mathbf{v}} - \mathbf{c}_{\pi(\mathbf{v})}\|_1$ for $\mathbf{v} \sim L_i$ and $\mathbf{r}_{\mathbf{v}} \sim X_{\mathbf{v}}$ and $\mathbf{c}_{\pi(\mathbf{v})} \sim X_{\mathbf{v}'}$ for $\mathbf{v}' \sim C(\pi(\mathbf{v}))$ (we will formalize this notion soon in Lemma 7.1). This is true because each sample will be bounded by $O(d \log n / 2^i)$, so that the empirical mean of $\text{polylog}(n)$ such samples will be a desired additive $d/(2 \cdot 2^i)$ approximation.

We now specify how to generate samples for estimating $\|\mathbf{r}_{\mathbf{v}} - \mathbf{c}_{\pi(\mathbf{v})}\|_1$. One may think of sampling $\pi(\mathbf{v})$ and $\mathbf{c}_{\pi(\mathbf{v})}$ first, and then drawing \mathbf{v} and $\mathbf{r}_{\mathbf{v}}$ after. In order to do this, we let \mathcal{P}_i be the distribution supported on L_{i-1} given by $\mathbf{u} \sim \mathcal{P}_i$ being set to $u \in L_{i-1}$ with probability proportional to $|C(u)|$. Furthermore, we consider

$$\alpha_i \stackrel{\text{def}}{=} \frac{2^i}{d \log^3 n} \in [0, 1],$$

and we denote \mathcal{S}_i the distribution over subsets of $[d]$ which include each coordinate $k \in [d]$ i.i.d with probability α_i . For a subset $S \subset [d]$, the character $\chi_S: \{0, 1\}^d \rightarrow \{-1, 1\}$ is the function $\chi_S(x) = (-1)^{\sum_{k \in S} x_k}$. Then, we will estimate the distance $\|\mathbf{r}_{\mathbf{v}} - \mathbf{c}_{\mathbf{u}}\|_1$ by considering the values of $\chi_S(\mathbf{r}_{\mathbf{v}})$ and $\chi_S(\mathbf{r}_{\mathbf{v}'})$ for random choice of $\mathbf{S} \sim \mathcal{S}_i$. Formally, by setting of α_i and the fact $\|\mathbf{r}_{\mathbf{v}} - \mathbf{c}_{\mathbf{u}}\|_1 \leq O(d \log n / 2^i)$,

$$\begin{aligned} \mathbf{E}_{\substack{\mathbf{v} \sim L_i \\ \mathbf{r}_{\mathbf{v}}, \mathbf{c}_{\pi(\mathbf{v})}}} [\|\mathbf{r}_{\mathbf{v}} - \mathbf{c}_{\pi(\mathbf{v})}\|_1] &= \mathbf{E}_{\mathbf{u} \sim \mathcal{P}_i} \left[\mathbf{E}_{\substack{\mathbf{v}, \mathbf{v}' \sim C(\mathbf{u}) \\ \mathbf{r}_{\mathbf{v}} \sim X_{\mathbf{v}} \\ \mathbf{r}_{\mathbf{v}'} \sim X_{\mathbf{v}'}}} [\|\mathbf{r}_{\mathbf{v}} - \mathbf{r}_{\mathbf{v}'}\|_1] \right] \\ &= \mathbf{E}_{\mathbf{u} \sim \mathcal{P}_i} \left[\mathbf{E}_{\substack{\mathbf{v}, \mathbf{v}' \sim C(\mathbf{u}) \\ \mathbf{r}_{\mathbf{v}} \sim X_{\mathbf{v}} \\ \mathbf{r}_{\mathbf{v}'} \sim X_{\mathbf{v}'}}} \left[\mathbf{Pr}_{\mathbf{S} \sim \mathcal{S}_i} [\chi_S(\mathbf{r}_{\mathbf{v}}) \neq \chi_S(\mathbf{r}_{\mathbf{v}'})] \right] \right] \cdot \frac{d \log^3 n}{2^i} \pm o\left(\frac{d}{2^i}\right). \end{aligned}$$

Specifically, the sketch follows from approximately drawing samples according to the following lemma (Lemma 7.1). Even though the lemma statement is long, it simply states the fact that we can approximately sample from the distribution needed to estimate the quantity

$$\mathbf{E}_{\mathbf{u} \sim \mathcal{P}_i} \left[\mathbf{E}_{\substack{\mathbf{v}, \mathbf{v}' \sim C(\mathbf{u}) \\ \mathbf{r}_{\mathbf{v}} \sim X_{\mathbf{v}} \\ \mathbf{r}_{\mathbf{v}'} \sim X_{\mathbf{v}'}}} \left[\mathbf{Pr}_{\mathbf{S} \sim \mathcal{S}_i} [\chi_S(\mathbf{r}_{\mathbf{v}}) \neq \chi_S(\mathbf{r}_{\mathbf{v}'})] \right] \right] \quad (46)$$

up to additive $o(1/\log^3 n)$ error with $\text{polylog}(n)$ samples. One (minor) issue is that a $\text{poly}(\log n, \log d)$ -space sketch cannot specify the vertices \mathbf{v} or points $\mathbf{r}_{\mathbf{v}}$ because these would require $O(d)$ bits to specify. In order to specify points and vertices we consider a universe reduction step. This is simply a random function $c: V_{i-1} \cup V_i \cup \{0, 1\}^d \rightarrow \{0, 1\}^{\text{polylog}(n, d)}$ which is used to rename elements of L_{i-1}, L_i and X . In particular, since $|L_{i-1} \cup L_i \cup X|$ is small (at most $\text{poly}(n, \log d)$), the function will be injective on $L_{i-1} \cup L_i \cup X$ with high probability, and thus allows the sketch to refer to nodes implicitly. This is a very minor point, and we will abuse notation by assuming the universe reduction has been applied.

Lemma 7.1. Consider a fixed quadtree T , an index $i \in [h]$, as well as a small parameters $\epsilon, \delta > 0$. There exists a linear sketch which uses $\text{poly}(\log n, \log d, 1/\epsilon, \log(1/\delta))$ space and outputs a sample from a distribution \mathcal{D} supported on $L_{i-1} \times L_i^2 \times X^2 \times \{-1, 1\}^2$. The distribution \mathcal{D} is ϵ -close in total variation from one where a sample $(\xi_u, \xi_v, \xi_{v'}, \xi_{r_v}, \xi_{r_{v'}}, \xi_{\chi_v}, \xi_{\chi_{v'}}) \sim \mathcal{D}$ satisfies:

1. The value ξ_u represents a sample of \mathbf{u} , i.e., $\Pr[\xi_u = u] = |C(u)|/|L_i|$.
2. The value ξ_v represents a sample of \mathbf{v} , i.e., $\Pr[\xi_v = v \mid \xi_u = u] = 1/|C(u)|$.
3. The value ξ_{r_v} represents a sample of \mathbf{r}_v , i.e., $\Pr[\xi_{r_v} = r_v \mid \xi_v = v] = 1/|X_v|$.
4. The value $\xi_{v'}$ is independent of ξ_v , and represents a sample of \mathbf{v}' , i.e., $\Pr[\xi_{v'} = v' \mid \xi_u = u] = 1/|C(u)|$.
5. The value $\xi_{r_{v'}}$ represents a sample of $\mathbf{r}_{v'}$, i.e., $\Pr[\xi_{r_{v'}} = r_{v'} \mid \xi_{v'} = v'] = 1/|X_{v'}|$.
6. The value $\xi_{\chi_v}, \xi_{\chi_{v'}} \in \{-1, 1\}^2$ represent samples of $\chi_S(\mathbf{r}_v)$ and $\chi_S(\mathbf{r}_{v'})$, i.e., for any $(b_u, b_v) \in \{-1, 1\}^2$,

$$\Pr[(\xi_{\chi_v}, \xi_{\chi_{v'}}) = (b_u, b_v) \mid (\xi_{r_v}, \xi_{r_{v'}}) = (r_v, r_{v'})] = \Pr_{\mathbf{S} \sim \mathcal{S}_i}[(\chi_S(r_v), \chi_S(r_{v'})) = (b_u, b_v)].$$

Given Lemma 7.1, the sketch is straight-forward. We independently consider $t = \text{polylog}(n)$ draws from the distribution \mathcal{D} with $\epsilon = 1/\text{polylog}(n)$ and $\delta = 1/\text{poly}(n)$, and we count the number of times a sample $(\xi_u, \xi_v, \xi_{v'}, \xi_{r_v}, \xi_{r_{v'}}, \xi_{\chi_v}, \xi_{\chi_{v'}}) \sim \mathcal{D}$ satisfies $\xi_{\chi_v} \neq \xi_{\chi_{v'}}$. This count, divided by t , will be an appropriate estimate to (46), and completes the high-level plan of the sketch.

7.1 Proof of Lemma 7.1

We will prove Lemma 7.1 in steps, corresponding to the six itemized elements needed for the lemma. Hence, a sample $(\xi_u, \xi_v, \xi_{v'}, \xi_{r_v}, \xi_{r_{v'}}, \xi_{\chi_v}, \xi_{\chi_{v'}}) \sim \mathcal{D}$ will proceed in six steps (even though all are correlated). In order to handle the correlation, the first step of generating a sample is to perform the following procedure.

With the above sampling procedure and specified events, we will first show that the distribution of \mathbf{u}^* , \mathbf{v}^* , and \mathbf{v}^{**} generated above are appropriate, and that events \mathcal{E}_1 , \mathcal{E}_2 and \mathcal{E}_3 occur with high enough probability. Then, we show that there exists a linear sketch which can recover \mathbf{u}^* , \mathbf{v}^* , and \mathbf{v}^{**} ; and finally, we show how to sample the representative elements.

Lemma 7.2 (Parent and Child Distribution). *Let $u \in L_{i-1}$ and $v, v' \in C(u)$. Then, as per Figure 6, we have*

$$\Pr_{(t_u)_u}[\mathbf{u}^* = u] = \frac{|C(u)|}{|L_i|}, \quad \text{and} \quad \Pr_{\substack{(\mathbf{D}_{\kappa,j})_{\kappa,j} \\ (\mathbf{D}'_{\kappa,j})_{\kappa,j}}}[\mathbf{v}^* = v, \mathbf{v}^{**} = v' \mid \mathbf{u}^* = u] = \frac{1}{|C(u)|^2}.$$

Sample: Consider an independent sample $\mathbf{t}_u \sim \text{Exp}(1)$ for each $u \in V_{i-1}$. Furthermore, for each $\kappa \in \{0, \dots, \lceil \log_2 n \rceil\}$ and $j = O(\log(n))$, we sample two random subsets $\mathbf{D}_{\kappa,j}, \mathbf{D}'_{\kappa,j} \subset V_i$ by including each node $v \in V_i$ i.i.d with probability $1/2^\kappa$. We will consider the following three events, where we specify them with respect to a parameter $\gamma > 0$ which will be set to $1/\text{polylog}(n)$.

- Event \mathcal{E}_1 : we have

$$\sum_{u \in L_{i-1}} \frac{|C(u)|}{\mathbf{t}_u} \leq \frac{4 \log(n/\gamma)}{\gamma} \cdot |L_i|.$$

- Event \mathcal{E}_2 : let $\mathbf{u}^* \in L_{i-1}$ be the non-empty node of T which maximizes $|C(\mathbf{u}^*)|/\mathbf{t}_{C(\mathbf{u}^*)}$, and let \mathbf{u}^{**} be the non-empty node of T which has the second largest $|C(\mathbf{u}^{**})|/\mathbf{t}_{C(\mathbf{u}^{**})}$. Then, we have

$$\frac{|C(\mathbf{u}^*)|}{\mathbf{t}_{\mathbf{u}^*}} \geq \gamma \cdot |L_i| \quad \text{and} \quad \frac{|C(\mathbf{u}^*)|}{\mathbf{t}_{\mathbf{u}^*}} \geq (1 + \gamma) \cdot \frac{|C(\mathbf{u}^{**})|}{\mathbf{t}_{\mathbf{u}^{**}}}. \quad (47)$$

We note that we will aim to set $\xi_u = \mathbf{u}^*$.

- Event \mathcal{E}_3 : For every $\kappa \in \{0, \dots, \lceil \log_2 n \rceil\}$ and every j , we have for $\mathbf{R} \in \{\mathbf{D}_{\kappa,j}, \mathbf{D}'_{\kappa,j}\}$,

$$\sum_{u \in L_{i-1}} \frac{|\mathbf{R} \cap C(u)|}{\mathbf{t}_u} \leq \frac{\log^3 n}{2^\kappa \cdot \gamma} \sum_{u \in L_{i-1}} \frac{|C(u)|}{\mathbf{t}_u}.$$

Furthermore, there exists a maximum $\kappa^* \in \{0, \dots, \lceil \log_2 n \rceil\}$ and $j \geq j'$ (with respect to lexicographic ordering) such that

$$|\mathbf{D}_{\kappa^*,j} \cap \{v : v \in C(\mathbf{u}^*)\}| = |\mathbf{D}'_{\kappa^*,j'} \cap \{v' : v' \in C(\mathbf{u}^*)\}| = 1,$$

and this maximum satisfies $2^{\kappa^*} \geq |C(\mathbf{u}^*)|$. Let $\mathbf{v}^* \in C(\mathbf{u}^*)$ be that unique child of \mathbf{u}^* specified by $\mathbf{D}_{\kappa^*,j}$ and $\mathbf{v}^{**} \in C(\mathbf{u}^*)$ be the unique child of \mathbf{u}^* specified by $\mathbf{D}'_{\kappa^*,j'}$. We will aim to set $\xi_v = \mathbf{v}^*$ and $\xi_{v'} = \mathbf{v}^{**}$.

We note that the success of the entire sampling procedure is dependent on the collection $(\mathbf{t}_u)_{u \in V_{i-1}}$, $(\mathbf{D}_{\kappa,j})_{\kappa,j}$, and $(\mathbf{D}'_{\kappa,j})_{\kappa,j}$, and the events $\mathcal{E}_1, \mathcal{E}_2$, and \mathcal{E}_3 occurring. We will continually refer to $\mathbf{u}^*, \mathbf{v}^*$, and \mathbf{v}^{**} and κ^* as specified.

Figure 6: Sampling for the Parent and Children.

Proof: The first item follows from a simple calculation using the CDF of exponential random

variables

$$\begin{aligned}
\Pr_{(\mathbf{t}_m)_m} [\mathbf{u}^* = u] &= \Pr_{(\mathbf{t}_m)_m} \left[\frac{|C(u)|}{\mathbf{t}_u} \geq \frac{|C(u')|}{\mathbf{t}_{u'}} \quad \forall u' \in L_{i-1} \right] = \mathbf{E}_{\mathbf{t}_u} \left[\prod_{u' \in L_{i-1}} \Pr_{\mathbf{t}_{u'}} \left[\frac{|C(u)|}{\mathbf{t}_u} \geq \frac{|C(u')|}{\mathbf{t}_{u'}} \right] \right] \\
&= \mathbf{E}_{\mathbf{t}_u} \left[\exp \left(- \sum_{u' \neq u} \frac{|C(u')|}{|C(u)|} \cdot \mathbf{t}_u \right) \right] = \int_0^\infty \exp(-t) \exp \left(- \frac{t}{|C(u)|} \sum_{u' \neq u} |C(u')| \right) dt \\
&= \frac{|C(u)|}{|L_i|}.
\end{aligned}$$

The second item follows from the observation that conditioned on the event that some unique vertex $v \in C(\mathbf{u}^*)$ was been picked to $\mathbf{D}_{\kappa^*,j}$ (or equivalently, $\mathbf{D}'_{\kappa^*,j'}$), its distribution is uniform over $\{v : v \in C(\mathbf{u}^*)\}$, due to the fact that each $v \in V_i$ is picked independently with the same probability. \blacksquare

Lemma 7.3. *The events $\mathcal{E}_1, \mathcal{E}_2$, and \mathcal{E}_3 occur with probability at least $1 - 8\gamma - O(1/n)$ over the draw of $(\mathbf{t}_u)_{u \in V_{i-1}}$, $(\mathbf{D}_{\kappa,j})_{\kappa,j}$, and $(\mathbf{D}'_{\kappa,j})_{\kappa,j}$.*

We henceforth consider a fixed setting of $(\mathbf{t}_u)_{u \in V_{i-1}}$, as well as sets $(D_{\kappa,j})_{\kappa,j}$ and $(D'_{\kappa,j})_{\kappa,j}$ which satisfy events $\mathcal{E}_1, \mathcal{E}_2$ and \mathcal{E}_3 ; this also specifies the nodes u^*, v^* and v^{**} . We remove the boldfaced notation to indicate these are no longer random variables. The next lemmas show how to algorithmically recover the values of u^*, v^* , and v^{**} assuming that $\mathcal{E}_1, \mathcal{E}_2$ and \mathcal{E}_3 are satisfied, and how to sample their representative points.

Lemma 7.4 (Parent Recovery Lemma). *There exists a linear sketch which uses $\text{polylog}(n)$ space which, given access to the collection $(\mathbf{t}_u)_{u \in V_{i-1}}$ satisfying $\mathcal{E}_1 \wedge \mathcal{E}_2$, outputs an element $\xi_u \in V_{i-1}$ which is equal to u^* with high probability over internal randomness of the sketch.*

Lemma 7.5 (Child Recovery Lemma). *There exists a linear sketch which uses $\text{polylog}(n)$ space, which given access to the collection $(\mathbf{t}_u)_{u \in V_{i-1}}$ and $(D_{\kappa,j})_{\kappa,j}$ satisfying $\mathcal{E}_1 \wedge \mathcal{E}_2 \wedge \mathcal{E}_3$, outputs an element $\xi_v \in V_i$ which is equal to v^* with high probability over internal randomness of the sketch.*

Lemma 7.6 (Child Representative Sampling Lemma). *There exists a linear sketch which uses $\text{polylog}(n)$ space, which given access to the collection $(\mathbf{t}_u)_{u \in V_{i-1}}$ and $(D_{\kappa,j})_{\kappa,j}$, outputs an element ξ_{r_v} or “fail”. Whenever events $\mathcal{E}_1 \wedge \mathcal{E}_2 \wedge \mathcal{E}_3$ are satisfied, the sketch does not output “fail” with high probability, and the output $\xi_{r_v} \in X$ is uniformly distributed among X_{v^*} .*

Lemma 7.5 and Lemma 7.6 may be equivalently stated for v^{**} and $\xi_{r'_v}$. It will be clear from the proof of Lemma 7.5 and Lemma 7.6 that we may apply these same linear sketches to recover ξ_{χ_v} and $\xi_{\chi_{v'}}$ which are distributed as χ_S applied to ξ_{r_v} and $\xi_{r'_v}$ for any $S \subset [d]$.

7.2 Proof of Lemma 7.3

The proof follows from computing the probabilities of events $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$ and using a union bound. In order to compute the probability of event \mathcal{E}_1 we will use lemma 5.4, and use Lemma 5.5 to compute the probability of event \mathcal{E}_2 .

Lemma 7.7. For any $\gamma > 0$, event \mathcal{E}_3 holds with probability at least $1 - O(\gamma/\log n) - 1/\text{poly}(n)$.

Proof: Fix the set of exponential random variables $(t_u)_{u \in V_{i-1}}$ and let u^* be the node that maximizes $|C(u)|/t_u$ over all $u \in L_{i-1}$. For the first part of event \mathcal{E}_3 , consider any $\kappa \in \{0, \dots, \lceil \log_2 n \rceil\}$, and let $\mathbf{R} \subset V_i$ be a set which includes each $v \in V_i$ i.i.d with probability $1/2^\kappa$. Notice that

$$\mathbb{E}_{\mathbf{R}} \left[\sum_{u \in L_{i-1}} \frac{|C(u) \cap \mathbf{R}|}{t_u} \right] = \frac{1}{2^\kappa} \sum_{u \in L_{i-1}} \frac{|C(u)|}{t_u}.$$

Hence, the probability that this sum exceeds $\log^3 n/\gamma$ times the expectation is at most $\gamma/\log^3 n$, and we may therefore union bound over all $O(\log^2 n)$ possible draws of sets \mathbf{R} for all κ and j .

For the second part of event \mathcal{E}_3 , notice that for κ satisfying $2^\kappa \geq |C(u^*)| \geq 2^{\kappa-1}$, with constant probability over the draw of $\mathbf{R} \subset V_i$ generated by including each $v \in V_i$ i.i.d with probability $1/2^\kappa$, $|\mathbf{R} \cap C(u^*)| = 1$. Hence, by repeating for $j = O(\log n)$ times, \mathbf{R} intersects a unique element of $C(u^*)$ at least twice (thereby setting j and j'), and therefore the maximum κ^* for which this property holds furthermore satisfies $2^{\kappa^*} \geq |C(u^*)|$. ■

7.3 Proof of Lemma 7.4

We refer to $\Gamma \in \mathbb{Z}_{\geq 0}^{|L_i|}$ as the vector such that $\Gamma_v = |X_v|$ for each $v \in L_i$. We will refer to Figure 7. Note that Γ is a linear function of the input, and that the sketch is indeed a linear function of Γ which stores $k \cdot l = \text{polylog}(n)$ many ℓ_p -sketches of accuracy $(1 \pm O(1/\log n))$ for $p = \Omega(1/\log^2 n)$ which succeeds with high probability, and thus the entire sketch uses $\text{polylog}(n)$ space.

Proposition 7.8. Consider any fixed $j \in [l]$ and a fixed hash function $h_j: L_{i-1} \rightarrow [k]$, and any value of $q \in [k]$. Then, with probability at least $1 - 1/\text{poly}(n)$ over the randomness in the ℓ_p -sketches,

$$\mathbf{B}_{j,q} = (1 \pm \epsilon_0) \left((1 \pm \epsilon) \sum_{\substack{u \in L_{i-1} \\ h_j(u)=q}} \frac{|C(u)|}{t_u} \right)^{1/p}.$$

Proof: For $j \in [l]$ and $q \in [k]$, the vector whose ℓ_p norm we are estimating is $\tilde{\Gamma}_{j,q} \in \mathbb{R}^{|L_i|}$ where

$$(\tilde{\Gamma}_{j,q})_v \stackrel{\text{def}}{=} \begin{cases} \Gamma_v / t_{\pi(v)}^{1/p} & h_j(\pi(v)) = q \\ 0 & \text{o.w} \end{cases}$$

Therefore, by the correctness of the $(1 \pm \epsilon_0)$ -accuracy ℓ_p -sketches, we have

$$\mathbf{B}_{j,q} = (1 \pm \epsilon_0) \|\tilde{\Gamma}_{j,q}\|_p = (1 \pm \epsilon_0) \left(\sum_{\substack{u \in L_{i-1} \\ h_j(u)=q}} \frac{1}{t_u} \sum_{v \in C(u)} \Gamma_v^p \right)^{1/p}$$

with probability at least $1 - 1/\text{poly}(n)$. The claim then follows from the fact that $\Gamma_v \in \{0\} \cup [1, \dots, n]$, which means $\Gamma_v^p \in \{0\} \cup [1, 1 + \epsilon]$, since $p = \epsilon/(2 \log n)$. ■

Linear Sketch of Lemma 7.4: We receive as input a sequence of numbers $(t_u)_{u \in V_{i-1}}$. We set parameters $\epsilon, \epsilon_0 \in (0, 1)$ satisfying $\epsilon_0 < \epsilon < \gamma/10$, as well as $p = \epsilon/(2 \log n)$. (Recall the setting of $\gamma = 1/\text{polylog}(n)$).

- We will instantiate a Count-Sketch data structure with $l = O(\log n)$ independent hash functions into $k = O(\log(n/\gamma)/(\epsilon_0 \gamma^3))$ buckets. In other words, for every $j \in [l]$, we independently sample a hash function $\mathbf{h}_j: L_{i-1} \rightarrow [k]$, and we will maintain some memory corresponding to each hash function $j \in [l]$ and each bucket $q \in [k]$.
- For each $j \in [l]$ and $q \in [k]$, we will maintain a $(1 \pm \epsilon_0)$ -accuracy ℓ_p -sketch of the vector $\Gamma \in \mathbb{Z}_{\geq 0}^{|L_i|}$, where we rescale the entry Γ_v by $1/t_{\pi(v)}^{1/p}$ if $\mathbf{h}_j(\pi(v)) = q$, and 0 otherwise. Specifically, the ℓ_p -sketch proceeds by maintaining $t_0 = O(\log n/(\epsilon_0 p)^4)$ copies of the following linear function of Γ :
 - For each $t \in [t_0]$, we generate a sequence of independent p -stable random variables $(\alpha_v^{(t,j,q)})_{v \in L_i}$ and we maintain

$$\mathbf{A}_{j,q}^{(t)} = \sum_{u \in L_{i-1}} \sum_{\substack{v \in L_i \\ \mathbf{h}_j(v) = q}} \frac{\alpha_v^{(t,j,q)} \cdot \Gamma_v}{t_u^{1/p}}.$$

Reporting Procedure: Given the $k \cdot l$ many ℓ_p -sketches, corresponding to each hash function and bucket pair, we produce an output $u \in L_{i-1}$ (which we will show will be u^* with high probability when events \mathcal{E}_1 and \mathcal{E}_2 are satisfied).

- For each $j \in [l]$ and $q \in [k]$, we run the reporting procedure for the ℓ_p -sketch corresponding to the j -th hash function and q -th bucket. Let $(\mathbf{B}_{j,q})$ be the reported ℓ_p -sketches.
- For each $u \in L_{i-1}$, we consider the collection of l values $(\mathbf{B}_{j, \mathbf{h}_j(u)} : j \in [l])$ corresponding to reported ℓ_p -sketches of where u hashed into. We set $\tilde{\mathbf{B}}_u$ be the median of the values $(\mathbf{B}_{j, \mathbf{h}_j(u)})$, and we report $\arg\max_{u \in L_{i-1}} \tilde{\mathbf{B}}_u$.

Figure 7: Parent Recovery Linear Sketch for Lemma 7.4.

Lemma 7.9. Consider any setting of $(t_u)_{u \in V_{i-1}}$ where \mathcal{E}_1 and \mathcal{E}_2 hold, and let $K = |C(u^*)|/t_{u^*}$. For any $j \in [l]$, with probability at least $1 - 1/\text{poly}(n)$ over the randomness of the hash functions \mathbf{h}_j ,

$$\mathbf{B}_{j, \mathbf{h}_j(u^*)} \geq (1 - \epsilon_0)(1 - \epsilon)^{1/p} K^{1/p}.$$

Moreover, for any $u \neq u^*$ and $j \in [l]$, with probability at least $9/10 - 1/\text{poly}(n)$ over the draw of \mathbf{h}_j ,

$$\mathbf{B}_{j, \mathbf{h}_j(u)} < (1 - \epsilon_0)(1 - \epsilon)^{1/p} K^{1/p}.$$

Proof: By Proposition 7.8, for any $u \in L_{i-1}$, we have with probability $1 - 1/\text{poly}(n)$,

$$\mathbf{B}_{j, \mathbf{h}_j(u)} = (1 \pm \epsilon_0) \left((1 \pm \epsilon) \cdot \frac{|C(u)|}{t_u} + (1 \pm \epsilon) \sum_{\substack{u' \in L_{i-1} \setminus \{u\} \\ \mathbf{h}_j(u')=q}} \frac{|C(u')|}{t_{u'}} \right)^{1/p}. \quad (48)$$

When $u = u^*$, we obtain the first claim. For the second claim, notice that for any $u \neq u^*$, using \mathcal{E}_2 , we have that $|C(u)|/t_u < (1 - \gamma)K$, so plugging into (48) we have the upper bound

$$\mathbf{B}_{j, \mathbf{h}_j(u)} \leq (1 + \epsilon_0)(1 + \epsilon)^{1/p} \left((1 - \gamma)K + \sum_{\substack{u' \in L_{i-1} \setminus \{u\} \\ \mathbf{h}_j(u')=q}} \frac{|C(u')|}{t_{u'}} \right)^{1/p}. \quad (49)$$

Now note that by event \mathcal{E}_1 and \mathcal{E}_2 , and the fact that $\mathbf{h}_j(u)$ is a uniform hash function on to a universe of size k , we have

$$\mathbf{E}_{\mathbf{h}_j} \left[\sum_{\substack{u' \in L_{i-1} \setminus \{u\} \\ \mathbf{h}_j(u')=q}} \frac{|C(u')|}{t_{u'}} \right] \leq \frac{1}{k} \sum_{u' \in L_{i-1}} \frac{|C(u')|}{t_{u'}} \leq \frac{4 \log(n/\gamma)}{k\gamma^2} \cdot K < \frac{1}{10} \cdot \epsilon_0 \gamma K.$$

By applying Markov's inequality, with probability $9/10$, $\sum_{u' \in L_{i-1} \setminus \{u\}} |C(u')|/t_{u'} < \epsilon_0 \gamma K$. Hence, plugging back into (49),

$$\mathbf{B}_{j, \mathbf{h}_j(u)} < (1 + \epsilon_0)(1 + \epsilon)^{1/p} \left((1 - \gamma)K + \epsilon_0 \gamma K \right)^{1/p} < (1 + \epsilon_0)(1 + \epsilon)^{1/p} (1 - \gamma + \epsilon_0 \gamma)^{1/p} K^{1/p},$$

which implies our desired bound once $\gamma/10 > \epsilon > \epsilon_0$. \blacksquare

We are now ready to prove Lemma 7.4

Proof: [Proof of Lemma 7.4] By 7.12 and a Chernoff bound, for any $u \neq u^*$ we have $\tilde{\mathbf{B}}_u < (1 - \epsilon_0)(1 - \epsilon)^{1/p} K^{1/p}$ with probability at least $1 - \exp(-\Omega(\ell)) > 1 - 1/\text{poly}(n)$, and we can then union bound so that this holds for all $u \neq u^*$. Secondly, again by 7.12 and a Chernoff bound, we have that with probability at least $1 - 1/\text{poly}(n)$, $\tilde{\mathbf{B}}_{u^*} > (1 - \epsilon_0)(1 - \epsilon)^{1/p} K^{1/p}$. It follows that $u^* = \text{argmax}_u \tilde{\mathbf{B}}_u$ with high probability, which completes the proof. \blacksquare

7.4 Proofs of Lemma 7.5 and Lemma 7.6

We will prove Lemma 7.5 and Lemma 7.6 as corollaries of the following sketching lemma, which we prove next.

Lemma 7.10. *There exists a linear sketch using $\text{poly}(\log n)$ space which takes as input a sequence $(t_u)_{u \in V_{i-1}}$ satisfying events \mathcal{E}_1 and \mathcal{E}_2 , a set $D = D_{\kappa, j} \subset V_i$ for some κ, j satisfying \mathcal{E}_3 , and a set $P \subset \{0, 1\}^d$. Given u^* , the sketch recovers a vector $\mathbf{z} \in \mathbb{R}_{\geq 0}^{|P|}$ satisfying the following conditions with high probability:*

- If $v \in C(u^*) \cap D$, every $a \in X_v \cap P$ has $\mathbf{z}_a \geq 1/(3t_{u^*})^{1/p}$.
- If $v \notin C(u^*) \cap D$, then every $a \in X_v \cap P$ has $\mathbf{z}_a \leq 1/(3 \cdot 2^\kappa / |C(u^*)| \cdot t_{u^*})^{1/p}$.

From Lemma 7.10, we argue that Lemma 7.5 and Lemma 7.6 are straight-forward. To see why Lemma 7.5 follows, instantiate the above lemma with $P = \{0, 1\}^d$, and use the sketch to recover a vector $z^{(\kappa, j)} \in \mathbb{R}_{\geq 0}^{|X_{u^*}|}$ for every κ, j . Whenever $2^\kappa \geq |C(u^*)|$, there is a simple test to determine whether any $v \in C(u^*)$ happens to lie in $D_{\kappa, j}$: check whether $z_a^{(\kappa, j)} \geq 1/(3t_{u^*})^{1/p}$ for some $a \in \{0, 1\}^d$ whose i -th node in T is v . Even though the algorithm does not know $|C(u^*)|$ (and hence cannot check that $2^\kappa \geq |C(u^*)|$), it may start with the largest $\kappa = \lceil \log_2 n \rceil$ and work its way down, while recovering the sets $C(u^*) \cap D_{\kappa, j}$. By \mathcal{E}_3 , the maximum setting of κ and j, j' where $C(u^*) \cap D_{\kappa, j}$ and $C(u^*) \cap D_{\kappa, j'}$ contain a unique element also satisfies $2^\kappa \geq |C(u^*)|$. Hence, the maximum value of κ for which we find $C(u^*) \cap D_{\kappa, j}$ and $C(u^*) \cap D_{\kappa, j'}$ contain unique elements will recover v^* and v^{**} .

The proof of Lemma 7.6 follows similarly, except we change the setting of P . Namely, for every $\eta \in \{0, \dots, \lceil \log_2 n \rceil\}$, we instantiate the linear sketch of Lemma 7.10 with $\mathbf{P}_\eta \subset \{0, 1\}^d$ being a random set generated by including each point i.i.d with probability $1/2^\eta$. Then, we recover the vector $z^{(\eta)} \in \mathbb{R}_{\geq 0}^{|\mathbf{P}_\eta|}$ (obtained from Lemma 7.10 with $P = \mathbf{P}_\eta$), and we check the first time that $v \in C(u^*) \cap D$ contains a unique $a \in X_v \cap \mathbf{P}_\eta$ where $z_a^{(\eta)} \geq 1/(3t_{u^*})^{1/p}$. When this occurs, the point a is uniformly distributed among all X_v , and hence we use a as the child representative point r_v . Furthermore, it is simple to recover the evaluation of any Boolean function $f: \{0, 1\}^d \rightarrow \{-1, 1\}$ (in particular, χ_S) at r_v by utilizing Lemma 7.10: we instantiate the lemma again while letting $P = \mathbf{P}_\eta \cap f^{-1}(1)$, and we check whether we can recover the same point $a \in \mathbf{P}_\eta$. If so, then $f(a) = 1$, otherwise, $f(a) = -1$.

We now prove Lemma 7.10, where we specify the sketch in Figure 8.

Proposition 7.11. *Consider any fixed $j \in [l]$ and a fixed hash function $h_j: L_i \rightarrow [k]$, any value of $q \in [k]$. Then, with probability at least $1 - 1/\text{poly}(n)$ over the randomness in the ℓ_p -sketches,*

$$\mathbf{B}_{j,q} = (1 \pm \epsilon_0) \left((1 \pm \epsilon) \sum_{u \in L_{i-1}} \frac{|D \cap \{v \in C(u) : h_j(v) = q, X_v \cap P \neq \emptyset\}|}{t_u} \right)^{1/p}.$$

Proof: The proof mirrors that of Proposition 7.8. By the correctness of the $(1 \pm \epsilon_0)$ -accuracy ℓ_p -sketches, we have

$$\mathbf{B}_{j,q} = (1 \pm \epsilon_0) \|\tilde{\Gamma}_{j,q}\|_p = (1 \pm \epsilon_0) \left(\sum_{u \in L_{i-1}} \frac{1}{t_u} \sum_{\substack{v \in C(u) \\ h_j(v) = q}} \mathbf{1}\{v \in D\} \cdot |X_v \cap P|^p \right)^{1/p}$$

with probability at least $1 - 1/\text{poly}(n)$. The claim then follows from the fact that $|X_v \cap P| \in \{0\} \cup [1, \dots, n]$, which means $|X_v \cap P|^p \in \{0\} \cup [1, 1 + \epsilon]$, since $p = \epsilon/(2 \log n)$. ■

Lemma 7.12. *Consider any setting of $(t_u)_{u \in L_{i-1}}$ where $\mathcal{E}_1, \mathcal{E}_2$ are satisfied, and let D be some $D_{\kappa, j}$ where \mathcal{E}_3 is satisfied. For any $j \in [l]$ and any $v \in C(u^*) \cap D$, with probability at least $1 - 1/\text{poly}(n)$ over the randomness of the hash functions \mathbf{h}_j and ℓ_p -sketches,*

$$\mathbf{B}_{j, \mathbf{h}_j(v)} \geq 1/(3t_{u^*})^{1/p}.$$

Moreover, for any $v \in C(u^*) \setminus D$ and $j \in [l]$, with probability at least $1/2$ over the draw of \mathbf{h}_j and ℓ_p -sketches,

$$\mathbf{B}_{j, \mathbf{h}_j(v)} < 1/(3 \cdot 2^\kappa / |C(u^*)| \cdot t_{u^*})^{1/p}.$$

Linear Sketch for Lemma 7.10: We receive as input a sequence of numbers $(t_u)_{u \in V_{i-1}}$, a set $D \subset V_i$, and a set $P \subset \{0, 1\}^d$. We set parameters $\epsilon = \epsilon_0 = 1/2$ as well as $p = \epsilon/(2 \log n)$.

- We instantiate a Count-Sketch data structure with $l = O(\log n)$ independent hash functions into $k = O(\log^3 n \log(n/\gamma)/(\gamma^3))$ buckets.
- For each $j \in [l]$ and $q \in [k]$, we maintain a $(1 \pm \epsilon_0)$ -accuracy ℓ_p -sketch of the vector $\tilde{\Gamma}_{j,q} \in \mathbb{Z}_{\geq 0}^{|L_i|}$, where for every $v \in L_i$

$$(\tilde{\Gamma}_{j,q})_v = \begin{cases} \mathbf{1}\{v \in D\} \cdot \frac{|X_v \cap P|}{t_u^{1/p}} & \mathbf{h}_j(v) = q \\ 0 & \text{o.w.} \end{cases}.$$

Reporting Procedure: Given the $k \cdot l$ many ℓ_p -sketches, corresponding to each hash function and bucket pair, as well as the identity of u^* , we produce a vector $\mathbf{z} \in \mathbb{R}_{\geq 0}^{|P|}$.

- For each $j \in [l]$ and $q \in [k]$, we run the reporting procedure for the ℓ_p -sketch corresponding to the j -th hash function and q -th bucket. Let $(\mathbf{B}_{j,q})$ be the reported ℓ_p -sketches.
- For each $v \in C(u^*)$ and $a \in X_v \cap P$, we let \mathbf{z}_a be set to the minimum value of $(\mathbf{B}_{j,\mathbf{h}_j(v)} : j \in [l])$.

Figure 8: Child Recovery Linear Sketch for Lemma 7.5.

Proof: By Proposition 7.8, for any $v \in C(u^*) \cap D$, we have with probability $1 - 1/\text{poly}(n)$,

$$\begin{aligned} \mathbf{B}_{j,\mathbf{h}_j(v)} &\geq (1 \pm \epsilon_0) \left((1 \pm \epsilon) \cdot \frac{1}{t_{u^*}} + (1 \pm \epsilon) \sum_{u \in L_{i-1} \setminus \{u^*\}} \frac{|D \cap \{v' \in C(u) : \mathbf{h}_j(v') = q, X_{v'} \cap P \neq \emptyset\}|}{t_u} \right)^{1/p} \\ &\geq \frac{(1 - \epsilon_0)(1 - \epsilon)^{1/p}}{t_{u^*}^{1/p}} \geq \frac{1}{2} \cdot \frac{1}{(2t_{u^*})^{1/p}} \geq \frac{1}{(3t_{u^*})^{1/p}}. \end{aligned}$$

For the second claim, notice that for any $v \in C(u^*) \setminus D$, we have

$$\mathbf{B}_{j,\mathbf{h}_j(v)} \leq (1 + \epsilon_0)(1 + \epsilon)^{1/p} \left(\sum_{u \in L_{i-1}} \frac{|D \cap \{v' \in C(u) : \mathbf{h}_j(v') = \mathbf{h}_j(v), X_{v'} \cap P \neq \emptyset\}|}{t_u} \right)^{1/p}.$$

We now show that the right-hand side of the above inequality is small with probability at least $1/2$.

Using the fact that $\mathcal{E}_1, \mathcal{E}_2$, and \mathcal{E}_3 are satisfied,

$$\begin{aligned}
\mathbf{E}_{\mathbf{h}_j} \left[\sum_{u \in L_{i-1}} \frac{|D \cap \{v' \in C(u) : \mathbf{h}_j(v') = \mathbf{h}_j(v), X_{v'} \cap P \neq \emptyset\}|}{t_u} \right] \\
\leq \sum_{u \in L_{i-1}} \frac{|C(u) \cap D|}{k \cdot t_u} \leq \frac{\log^3 n}{k \cdot 2^{\kappa} \gamma} \sum_{u \in L_{i-1}} \frac{|C(u)|}{t_u} \\
\leq \frac{4 \log^3 n \log(n/\gamma)}{k \cdot 2^{\kappa} \cdot \gamma^2} \cdot |L_i| \leq \frac{4 \log^3 n \log(n/\gamma)}{k \cdot 2^{\kappa} \cdot \gamma^3} \cdot \frac{|C(u^*)|}{t_{u^*}} \\
\leq \frac{1}{30 \cdot 2^{\kappa} / |C(u^*)| \cdot t_{u^*}}.
\end{aligned}$$

Hence, with probability at least $2/3 - 1/\text{poly}(n) \geq 1/2$ over the choice of hash functions and ℓ_p -sketch, $\mathbf{B}_{j, \mathbf{h}_j(v)} \leq (3/2)(3/2)^{1/p} / (10 \cdot 2^{\kappa} / |C(u^*)| \cdot t_{u^*})^{1/p} \leq 1 / (3 \cdot 2^{\kappa} / |C(u^*)| \cdot t_{u^*})^{1/p}$. \blacksquare

8 Lower Bound for Streaming MST

Our main theorem is the following:

Theorem 13. *Let $n, d \in \mathbb{N}$ satisfy $n \leq 2^{3d/4}$ and $\ell \in \mathbb{N}$. A randomized ℓ -bit streaming algorithm which estimates the MST of n points in $\{0, 1\}^d$ up to multiplicative factor $\alpha > 1$ with probability at least $2/3$ must satisfy*

$$\ell \geq \frac{d}{\alpha} - \log \alpha.$$

In particular, if ℓ is a constant independent of n and d , then $\alpha = \tilde{\Omega}(\log n)$.

We first state a theorem of [AIK08] giving a lower bound for communication protocols for EMD over $\{0, 1\}^d$. Fix a $d \geq 1$ and $1 \leq \alpha \leq d$. Let $C \subset \mathbb{F}_2^d$ be a linear code of dimension at least $d/4$ and weight at least cd , for a constant $c \geq 0$. Furthermore, the set C^\perp is the orthogonal subspace (i.e., the set of vectors $z \in \{0, 1\}^d$ satisfying $\langle z, x \rangle = 0$ for all $x \in C$). Let \mathcal{T}_ϵ denote the distribution over $\{0, 1\}^d$ such that each bit of $z \sim \mathcal{T}_\epsilon$ is set to 1 with probability ϵ independently. We define a probability distribution \mathcal{D} supported on (Z, A, B) where $Z \in \{0, 1\}$ and A, B are subsets of size $n := |C^\perp| \leq 2^{3d/4}$.

Let $\epsilon := 1/(200\alpha)$. To draw $(\mathbf{Z}, \mathbf{A}, \mathbf{B}) \sim \mathcal{D}$,

1. We first draw $\mathbf{Z} \sim \{0, 1\}$ uniformly at random.
2. If $\mathbf{Z} = 1$, we sample $\mathbf{x} \sim \{0, 1\}^d$ and $\mathbf{z} \sim \mathcal{T}_\epsilon$, and set $\mathbf{y} = \mathbf{x} + \mathbf{z}$,

$$\mathbf{A} = \left\{ \mathbf{x} + c \in \{0, 1\}^d : c \in C^\perp \right\} \quad \text{and} \quad \mathbf{B} = \left\{ \mathbf{y} + c \in \{0, 1\}^d : c \in C^\perp \right\}.$$

3. If $\mathbf{Z} = 0$, we sample $\mathbf{x}, \mathbf{y} \sim \{0, 1\}^d$ independently and set

$$\mathbf{A} = \left\{ \mathbf{x} + c \in \{0, 1\}^d : c \in C^\perp \right\} \quad \text{and} \quad \mathbf{B} = \left\{ \mathbf{y} + c \in \{0, 1\}^d : c \in C^\perp \right\}.$$

The main lower bound of [AIK08] for EMD states the following.

Theorem 14 (Proposition 4.1 of [AIK08]). *Let Π be any deterministic 1-bit protocol between two parties which receive a subset of $\{0, 1\}^d$ as input. Then,*

$$\Pr_{(\mathbf{Z}, \mathbf{A}, \mathbf{B}) \sim \mathcal{D}} [\Pi(\mathbf{A}, \mathbf{B}) = \mathbf{Z}] \leq \frac{1}{2} + 2^{-\Omega(d/\alpha)}.$$

Given the above theorem, we now show that a too-good-to-be-true protocol for Euclidean MST on $\{0, 1\}^d$ would imply a protocol contradicting Theorem 14 above. Given a $k \geq 2$, we define the following distribution $\mathcal{D}^{(k)}$ supported on (Z, A_1, \dots, A_k) where $Z \in \{0, 1\}$ and A_1, \dots, A_k are subsets of $\{0, 1\}^d$ of size n :

1. We first draw $\mathbf{Z} \sim \{0, 1\}$ uniformly at random.
2. If $\mathbf{Z} = 1$, we draw $\mathbf{x} \sim \{0, 1\}^d$ and $\mathbf{z}_2, \dots, \mathbf{z}_k \sim \mathcal{T}_\epsilon$ independently. Set $\mathbf{x}_1 = \mathbf{x}$ and $\mathbf{x}_\ell = \mathbf{x}_{\ell-1} + \mathbf{z}_\ell$ for each $\ell = 2, \dots, k$, and

$$\mathbf{A}_\ell = \left\{ \mathbf{x}_\ell + c \in \{0, 1\}^d : c \in C^\perp \right\}.$$

3. If $\mathbf{Z} = 0$, we draw $\mathbf{x}_1, \dots, \mathbf{x}_k \sim \{0, 1\}^d$ independently and set

$$\mathbf{A}_\ell = \left\{ \mathbf{x}_\ell + c \in \{0, 1\}^d : c \in C^\perp \right\}.$$

Observation 15. When $k = 2$, $\mathcal{D}^{(k)}$ is exactly \mathcal{D} . For any $\ell < r \in [k]$, the marginal distribution of $(\mathbf{Z}, \mathbf{A}_\ell, \dots, \mathbf{A}_r)$ as $(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}$ is the same as $\mathcal{D}^{(r-\ell+1)}$.

Observation 16. For any $k \geq 2$, the following holds with probability at least $1 - o(1)$ over the draw of a sample $(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}$: the set $\mathbf{A} = \mathbf{A}_1 \cup \dots \cup \mathbf{A}_k$ has a minimum spanning tree of cost $O(\epsilon d n k + n d)$ if $\mathbf{Z} = 1$, and minimum spanning tree of cost at least $\Omega(d n k)$ if $\mathbf{Z} = 0$.

In our main technical lemma which follows, we consider k -party, one-way protocols. There are k players (labeled $1, \dots, k$), and player ℓ is allowed to send a single message to player $\ell + 1$. In particular, the protocol proceeds by the first player communicating to the second player, the second to third, and so on, until the k -th player, who produces an output.

Lemma 8.1 (Player Reduction Lemma). *Let ℓ be a positive integer and $k = 2^b + 1$ for some $b \geq 0$. Let Π be any deterministic ℓ -bit, k -party one-way protocol, in which each party receives a subset of $\{0, 1\}^d$. If*

$$\Pr_{(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}} [\Pi(\mathbf{A}_1, \dots, \mathbf{A}_k) = \mathbf{Z}] \geq \frac{1}{2} + \delta$$

for some $\delta > 0$, then there is a deterministic ℓ -bit, 2-party one-way protocol Π' such that

$$\Pr_{(\mathbf{Z}, \mathbf{A}_1, \mathbf{A}_2) \sim \mathcal{D}} [\Pi'(\mathbf{A}_1, \mathbf{A}_2) = \mathbf{Z}] \geq \frac{1}{2} + \frac{2\delta}{2^b} > \frac{1}{2} + \frac{2\delta}{k}.$$

We use this lemma to prove the main lower bound:

Proof of Theorem 13: Let $k = \Theta(\alpha)$ such that $k = 2^b + 1$ for some b and $(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}$. As per Observation 16, if Π^* is a randomized k -party, ℓ -bit one-way protocol for α -approximating MST, then one can derive from Π^* a deterministic k -party, ℓ -bit one-way protocol Π such that

$$\Pr_{(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}} [\Pi(\mathbf{A}_1, \dots, \mathbf{A}_k) = \mathbf{Z}] \geq \frac{2}{3} - o(1).$$

Applying Lemma 8.1, we obtain a deterministic two-party, ℓ -bit one-way protocol Π' such that

$$\Pr_{(\mathbf{Z}, \mathbf{A}_1, \mathbf{A}_2) \sim \mathcal{D}} [\Pi'(\mathbf{A}_1, \mathbf{A}_2) = \mathbf{Z}] \geq \frac{1}{2} + \Omega\left(\frac{1}{k}\right).$$

We now reduce the communication to 1 bit, at the cost of lowering the advantage by a factor of $O(2^\ell)$: the players guess a transcript of ℓ bits, and player 1 sends one bit to verify that their part of the guessed transcript agrees with Π' (also in Lemma 4.1 of [AIK08]). In particular, we get a deterministic two-party, one-bit one way protocol Π'' such that

$$\Pr_{(\mathbf{Z}, \mathbf{A}_1, \mathbf{A}_2) \sim \mathcal{D}} [\Pi''(\mathbf{A}_1, \mathbf{A}_2) = \mathbf{Z}] \geq \frac{1}{2} + \Omega\left(\frac{1}{k2^\ell}\right) \geq \frac{1}{2} + \Omega\left(\frac{1}{\alpha 2^\ell}\right).$$

The lower bound $\ell \geq (d/\alpha) - \log \alpha$ follows from Theorem 14 and Observation 15, since $\Omega(1/(\alpha 2^\ell)) \leq 2^{-\Omega(d/\alpha)}$. In particular, this implies that $\ell = \omega(1)$ when $\alpha = c \log n / \log \log n$ for some sufficiently small constant $c > 0$. \blacksquare

Lemma 8.1 is the corollary of the following lemma, which we prove next.

Lemma 8.2. *Let ℓ be a positive integer and $k = 2k' + 1$ for some positive integer k' . Let Π be any deterministic ℓ -bit, k -party one-way protocol, in which each party receives a subset of $\{0, 1\}^d$. If*

$$\Pr_{(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}} [\Pi(\mathbf{A}_1, \dots, \mathbf{A}_k) = \mathbf{Z}] \geq \frac{1}{2} + \delta$$

for some $\delta > 0$, then there is a deterministic ℓ -bit, $(k' + 1)$ -party one-way protocol Π' such that

$$\Pr_{(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_{k'+1}) \sim \mathcal{D}^{(k'+1)}} [\Pi'(\mathbf{A}_1, \dots, \mathbf{A}_{k'+1}) = \mathbf{Z}] \geq \frac{1}{2} + \frac{\delta}{2}.$$

Proof: Let \mathfrak{A} denote the collection of sets $\{C^\perp(x) : x \in \{0, 1\}^d\}$. It is easy to verify that $(\mathbf{Z}, \mathbf{A}_{k'+1})$ for $(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}$ is distributed equivalently as $\mathbf{Z} \sim \{0, 1\}$ and $\mathbf{A}_{k'+1} \sim \mathfrak{A}$ uniformly and independently. We introduce some notation. Given $A \in \mathfrak{A}$, $z \in \{0, 1\}$ and $M \in \{0, 1\}^\ell$, we write $p(A, z, M)$ to denote the probability of (below is the message of the k' th player)

$$\Pi_{k'}(\mathbf{A}_1, \dots, \mathbf{A}_{k'}) = M$$

when $(\mathbf{Z}, \mathbf{A}_1, \dots, \mathbf{A}_k) \sim \mathcal{D}^{(k)}$ conditioning on $\mathbf{Z} = z$ and $\mathbf{A}_{k'+1} = A$. Note that for any $A \in \mathfrak{A}$ and $z \in \{0, 1\}$, $\sum_M p(A, z, M) = 1$. We break up the analysis into two cases, corresponding to the value of the quantity

$$\xi := \frac{1}{4|\mathfrak{A}|} \sum_{A \in \mathfrak{A}} \sum_{M \in \{0, 1\}^\ell} |p(A, 0, M) - p(A, 1, M)|.$$

The first case occurs when $\xi \geq \delta/2$; the deterministic protocol Π' is obtained from Π as follows:

1. Given inputs $A_1, \dots, A_{k'}, A_{k'+1}$, the first k' players simulate Π ; let $M_{k'}$ denote the ℓ -bit message sent by player k' to player $k' + 1$.
2. After receiving $M_{k'}$, player $k' + 1$ returns 1 if $p(A_{k'+1}, 1, M_{k'}) \geq p(A_{k'+1}, 0, M_{k'})$ and returns 0 otherwise.

The probability that Π' succeeds can be expressed as

$$\begin{aligned}
& \frac{1}{2^{|\mathfrak{A}|}} \sum_{A \in \mathfrak{A}} \max \{p(A, 0, M), p(A, 1, M)\} \\
&= \frac{1}{2^{|\mathfrak{A}|}} \sum_{A \in \mathfrak{A}} \frac{p(A, 0, M) + p(A, 1, M) + |p(A, 0, M) - p(A, 1, M)|}{2} \\
&= \frac{1}{2} + \xi \geq \frac{1}{2} + \frac{\delta}{2}.
\end{aligned}$$

For the second case when $\xi < \delta/2$, we use Π to obtain the following randomized protocol Π^* :

1. Let $A_{k'+1}, \dots, A_k$ be the inputs of the $k' + 1$ players. The first player draws a message \mathbf{M} from the distribution $p(A_{k'+1}, 0, \cdot)$, i.e., it draws $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_{k'} \sim \mathfrak{A}$, and simulates the protocol to generate the message \mathbf{M} sent by the k' -th player.
2. The first player simulates Π as player $k' + 1$, using \mathbf{M} as the message from player k' ; the rest of the k' players simulate Π as players $k' + 2, \dots, k$.

Then one can bound the probability that Π^* succeeds by the probability that Π succeeds minus

$$\frac{1}{2^{|\mathfrak{A}|}} \sum_{A \in \mathfrak{A}} d_{\text{TV}}(p(A, 0, \cdot) - p(A, 1, \cdot)) = \xi \leq \frac{\delta}{2}.$$

As a result, there exists a deterministic protocol Π' that succeeds with probability at least $(1/2) + \delta - (\delta/2) \geq (1/2) + (\delta/2)$. ■

9 Sketching ℓ_p norms for p near 0

In this section, we compute the dependence on p for Indyk's ℓ_p -sketch using p -stable random variables. The reason for this section is that we will later utilize this sketch with $p \rightarrow 0$ as $n \rightarrow \infty$. We first define stable random variables.

Definition 9.1. *A distribution \mathcal{D} is said to be p -stable if whenever $\mathbf{x}_1, \dots, \mathbf{x}_n \sim \mathcal{D}$ are drawn independently and $\mathbf{a} \in \mathbb{R}^n$ is a fixed vector, we have*

$$\sum_{i=1}^n a_i \mathbf{x}_i \stackrel{d}{=} \|\mathbf{a}\|_p \mathbf{x}$$

where $\mathbf{x} \sim \mathcal{D}$ is again drawn from the same distribution, and the “ $\stackrel{d}{=}$ ” symbol above denotes distributional equality.

We refer to [Nol09] for a thorough discussion on p -stable random variables, and to [Ind06b] for their use in streaming/sketching computation. Our proof of correctness will utilize the following standard methods for generating p -stable random variables.

Proposition 9.2 ([CMS76]). *Fix any $p \in (0, 1)$. Then a draw from a p -stable distribution $\mathbf{x} \sim \mathcal{D}_p$ can be generated as follows: (i) generate $\boldsymbol{\theta} \sim [-\frac{\pi}{2}, \frac{\pi}{2}]$, (ii) generate $\mathbf{r} \sim [0, 1]$, and set*

$$\mathbf{x} = \frac{\sin(p\boldsymbol{\theta})}{\cos^{1/p}(\boldsymbol{\theta})} \cdot \left(\frac{\cos(\boldsymbol{\theta}(1-p))}{\ln(1/\mathbf{r})} \right)^{\frac{1-p}{p}}. \quad (50)$$

Remark 17. *By Proposition 9.2, in order to generate a draw of $|\mathbf{x}|$, for $\mathbf{x} \sim \mathcal{D}_p$, we may generate $\theta \sim [-\frac{\pi}{2}, \frac{\pi}{2}]$ and $r \sim [0, 1]$, and consider the magnitude of (50). Notice, furthermore, that $\cos(\theta(1-p)) = \cos(-\theta(1-p))$, so (50) is symmetric around the origin in θ . This means we may generate $|\mathbf{x}|$ by drawing $\mathbf{r} \sim [0, 1]$ and $\boldsymbol{\theta} \sim [0, \pi/2]$, and outputting $g(\mathbf{r}, \boldsymbol{\theta})$, for $g: (0, 1) \times (0, \pi/2) \rightarrow \mathbb{R}$ be*

$$g(r, \theta) \stackrel{\text{def}}{=} \frac{\sin(p\theta)}{\cos^{1/p}(\theta)} \left(\frac{\cos(\theta(1-p))}{\ln(1/r)} \right)^{\frac{1-p}{p}}. \quad (51)$$

One can check, via simple calculus, that the function $g: (0, 1) \times (0, \pi/2) \rightarrow \mathbb{R}$ is continuous and monotone increasing in both parameters.

Claim 9.3. *There is some setting $t^* \in [1/10, 9/10]$ satisfying*

$$g\left(t^*, \frac{\pi t^*}{2}\right) \stackrel{\text{def}}{=} \text{median}(|\mathcal{D}_p|) = \sup \left\{ z \in \mathbb{R} : \Pr_{\mathbf{x} \sim \mathcal{D}_p} [|\mathbf{x}| \leq z] \leq 1/2 \right\}.$$

Proof: Consider the CDF $F(z) = \Pr_{\mathbf{x} \sim \mathcal{D}_p} [|\mathbf{x}| \leq z]$ as well as the function $F(g(t, \pi t/2))$. Since $g(t, \pi t/2)$ is continuous and monotone increasing in both parameters and $F(z)$ is continuous and monotone increasing, the function $F(g(t, \pi t/2))$ is continuous and monotone increasing. Furthermore, since $g(\mathbf{r}, \boldsymbol{\theta})$ generates a draw from $|\mathbf{x}|$, where $\mathbf{x} \sim \mathcal{D}_p$ when $\mathbf{r} \sim [0, 1]$ and $\boldsymbol{\theta} \sim [0, \pi/2]$, for any $t \in [0, 1]$ we have that $t^2 \leq F(g(t, \pi t/2)) \leq 1 - (1-t)^2$. In other words, $F(g(1/10, \pi/20)) < 1/2 < F(g(9/10, 9\pi/20))$, so the result follows from the intermediate value theorem. ■

Lemma 9.4. *Fix any $p \in (0, 1)$ and $\epsilon \in (0, 1)$, and let $R = \text{median}(|\mathcal{D}_p|)$. Then there exists a small constant $c \in (0, 1)$ independent of p, ϵ such that*

$$\Pr_{X \sim \mathcal{D}_p} [R \leq |X| \leq (1+\epsilon)R] \geq cp^2\epsilon^2 \quad \text{and} \quad \Pr_{X \sim \mathcal{D}_p} [(1-\epsilon)R \leq |X| \leq R] \geq cp^2\epsilon^2$$

Proof: Let $t^* \in (1/10, 9/10)$ be the parameter satisfying $R = g(t^*, \pi t^*/2)$ as per Claim 9.3. Then, by the definition of $g(r, \theta)$ for generating a draw from $|\mathcal{D}_p|$, we have

$$\begin{aligned} \Pr_{\mathbf{x} \sim \mathcal{D}_p} [R \leq |\mathbf{x}| \leq (1+\epsilon)R] &\geq \Pr_{\substack{\boldsymbol{\theta} \sim [0, \pi/2] \\ \mathbf{r} \sim [0, 1]}} \left[\ln R \leq \ln g(\mathbf{r}, \boldsymbol{\theta}) \leq \ln R + \frac{\epsilon}{2} \right] \\ &\geq \Pr_{\mathbf{r} \sim [0, 1]} \left[t^* \leq \mathbf{r} \leq t^* + \frac{\epsilon}{2D} \right] \Pr_{\boldsymbol{\theta} \sim [0, \pi/2]} \left[\pi t/2 \leq \boldsymbol{\theta} \leq \pi t/2 + \frac{\pi\epsilon}{2D} \right] \gtrsim \frac{\epsilon^2}{D^2}. \end{aligned}$$

where $D = \max \left\{ \frac{d}{dt} [\ln g(t, \pi t/2)] (\ell) : \ell \in [1/100, 99/100] \right\}$, and we used the fact $g(r, \theta)$ is continuous and increasing in both variables. Hence, it remains to upper bound D , which we do next.

$$\frac{d}{dt} [\ln g(t, \pi t/2)] (\ell) = \frac{p\pi \cdot \cos(p\pi\ell/2)}{2 \cdot \sin(p\pi\ell/2)} + \frac{\pi \cdot \sin(\pi\ell/2)}{2p \cdot \cos(\pi\ell/2)} - \frac{(1-p)^2\pi \cdot \sin(\pi\ell/2(1-p))}{2p \cdot \cos(\pi\ell/2(1-p))} + \frac{1-p}{p\ell \ln(1/\ell)},$$

which is $O(1/p)$ for any $\ell \in [1/100, 99/100]$. The bound for the probability $|\mathbf{x}|$ lies between $(1-\epsilon)R$ and R follows similarly. \blacksquare

We are now ready to prove the correctness of the Indyk p -stable sketch.

Theorem 18 (Indyk's p -Stable Sketch). *Consider any $p, \epsilon, \delta \in (0, 1)$, and set $t = O(\log(1/\delta)/(p\epsilon)^4)$. Let $\mathbf{x}_1, \dots, \mathbf{x}_t \sim \mathcal{D}_p$ independently. Then, with probability at least $1 - \delta$,*

$$1 - \epsilon \leq \text{median}_{i \in [t]} \left\{ \frac{|\mathbf{x}_i|}{\text{median}(|\mathcal{D}_p|)} \right\} \leq 1 + \epsilon.$$

Proof: Set $R = \text{median}(|\mathcal{D}_p|)$. By Lemma 9.4, there exists a constant $c > 0$ such that for each trial $\mathbf{x}_i \sim \mathcal{D}_p$, the probability that $|\mathbf{x}_i| > (1 + \epsilon)R$ is at most $1/2 - cp^2\epsilon^2$, and the probability that $|\mathbf{x}_i| < (1 - \epsilon)R$ is at most $1/2 - cp^2\epsilon^2$. By Chernoff bounds, the number of trials $i \in [t]$ satisfying $|\mathbf{x}_i| > (1 + \epsilon)R$ is at most $t/2$ with probability at least $1 - \exp(-(\epsilon p)^4 t/3)$, and similarly, the number of trials $i \in [t]$ satisfying $|\mathbf{x}_i| < (1 - \epsilon)R$ is at most $t/2$ with probability at least $1 - \exp(-(\epsilon p)^4 t/3)$. When both of these events occur simultaneously, with probability at least $1 - 2e^{-\epsilon^4 p^4 t/3} > 1 - \delta$, the median falls within the range $(1 - \epsilon)R$ and $(1 + \epsilon)R$, and we obtain our desired bound. \blacksquare

Acknowledgments

We would like to thank David Woodruff, Ilya Razenshteyn, and Aleksandar Nikolov for illuminating discussions relating to this work.

A Embedding ℓ_p^d into $\{0, 1\}^d$

Lemma A.1. *Let $p \in (1, 2]$. There exists a distribution \mathcal{D}_p supported on \mathbb{R} which exhibits p -stability: for any $d \in \mathbb{N}$ and any vector $x \in \mathbb{R}^d$, the random variables \mathbf{Y}_1 and \mathbf{Y}_2 given by*

$$\begin{aligned} \mathbf{Y}_1 &= \sum_{i=1}^d x_i z_i && \text{for } z_1, \dots, z_d \sim \mathcal{D}_p \text{ independently,} \\ \mathbf{Y}_2 &= z \cdot \|x\|_p && \text{for } z \sim \mathcal{D}_p, \end{aligned}$$

are equal in distribution, and $\mathbf{E}_{z \sim \mathcal{D}_p}[\|z\|]$ is at most a constant C_p .

Consider fixed $R \in \mathbb{R}^{\geq 0}$ which is a power of 2, and for $p \in [1, 2]$, let $\mathbf{f}: \mathbb{R}^d \rightarrow \{0, 1\}$ be the randomized function given by sampling the following random variables

- $\mathbf{z} = (z_1, \dots, z_d)$ where $z_1, \dots, z_d \sim \mathcal{D}_p$. In the case $p = 1$, we let \mathcal{D}_1 be the distribution where $z_i \sim \mathcal{D}_1$ is set to 1 with probability $1/d$ and 0 otherwise.
- $\mathbf{h} \sim [0, R]$, and
- a uniformly random function $\mathbf{g}: \mathbb{Z} \rightarrow \{0, 1\}$.

and we evaluate

$$\mathbf{f}(x) = \mathbf{g} \left(\left\lfloor \frac{\sum_{i=1}^d x_i z_i - \mathbf{h}}{R} \right\rfloor \right) \in \{0, 1\}.$$

If $x, y \in \mathbb{R}^d$ with $\|x - y\|_p \leq R/t_p$ (for a parameter t_p which we specify later). When $p = 1$, we consider $\|x - y\|_\infty \leq R/t_1$. Then

$$\Pr_{\mathbf{z}, \mathbf{h}, \mathbf{g}} [\mathbf{f}(x) \neq \mathbf{f}(y)] = \frac{1}{2} \cdot \Pr_{\mathbf{z}, \mathbf{h}} \left[\left\lfloor \frac{\sum_{i=1}^d x_i z_i - \mathbf{h}}{R} \right\rfloor \neq \left\lfloor \frac{\sum_{i=1}^d y_i z_i - \mathbf{h}}{R} \right\rfloor \right].$$

Consider a fixed $\mathbf{z} = (z_1, \dots, z_d)$ and let

$$\mathbf{w} = \sum_{i=1}^d (x_i - y_i) z_i.$$

Then, we have

$$\Pr_{\mathbf{h} \sim [0, R]} \left[\left\lfloor \frac{\sum_{i=1}^d x_i z_i - \mathbf{h}}{R} \right\rfloor \neq \left\lfloor \frac{\sum_{i=1}^d y_i z_i - \mathbf{h}}{R} \right\rfloor \right] = \max \left\{ \frac{|\mathbf{w}|}{R}, 1 \right\},$$

so that

$$\Pr_{\mathbf{z}, \mathbf{h}, \mathbf{g}} [\mathbf{f}(x) \neq \mathbf{f}(y)] = \frac{1}{2} \mathbf{E}_{\mathbf{z}} \left[\max \left\{ \frac{|\mathbf{w}|}{R}, 1 \right\} \right].$$

When $p \in (1, 2]$, we notice that by Jensen's inequality and Lemma A.1,

$$\mathbf{E}_{\mathbf{z}} \left[\max \left\{ \frac{|\mathbf{w}|}{R}, 1 \right\} \right] \leq \max \left\{ \frac{C_p \|x - y\|_p}{R}, 1 \right\} = \frac{C_p \|x - y\|_p}{R},$$

and similarly, $\mathbf{E}_{\mathbf{z}} [\max\{|\mathbf{w}|/R, 1\}] \leq \|x - y\|_1 / (dR)$ when $p = 1$. Furthermore, for $p \in (1, 2]$, we lower bound the above quantity by

$$\mathbf{E}_{\mathbf{z}} \left[\max \left\{ \frac{|\mathbf{w}|}{R}, 1 \right\} \right] \geq \frac{\|x - y\|_p}{R} \cdot \mathbf{E}_{\mathbf{z} \sim \mathcal{D}_p} [\max\{|\mathbf{z}|, t_p\}] \geq \frac{\|x - y\|_p}{R} \cdot \frac{C_p}{2},$$

since $\mathbf{E}_{\mathbf{z} \sim \mathcal{D}_p} [\max\{|\mathbf{z}|, t_p\}] \rightarrow C_p$ as $t_p \rightarrow \infty$, which means that for some constant setting of high enough t_p , we obtain the above inequality. In particular, for $p \in (1, 2]$, we may choose t_p to be a large enough constant depending only on p . When $p = 1$,

$$\mathbf{E}_{\mathbf{z}} \left[\max \left\{ \frac{|\mathbf{w}|}{R}, 1 \right\} \right] \geq \sum_{i=1}^d \frac{1}{d} \left(1 - \frac{1}{d}\right)^{d-1} \max \left\{ \frac{|x_i - y_i|}{R}, 1 \right\} = \frac{1}{d} \left(1 - \frac{1}{d}\right)^{d-1} \frac{\|x - y\|_1}{R},$$

by setting $t_1 = 1$.

For $p \in (1, 2]$, we consider concatenating m independent executions of the above construction and consider that as an embedding of $\phi: \mathbb{R}^d \rightarrow \{0, 1\}^m$, we have that for any $x, y \in \mathbb{R}^d$ with $\|x - y\|_p \leq R/t_p$,

$$\begin{aligned} \mathbf{E}_{\phi} [\|\phi(x) - \phi(y)\|_1] &= m \cdot \mathbf{Pr}_{\mathbf{f}} [\mathbf{f}(x) \neq \mathbf{f}(y)] \\ &\asymp m \cdot \frac{\|x - y\|_p}{R}, \end{aligned}$$

where the notation \asymp suppresses constant factors. Suppose $A \cup B \subset \mathbb{R}^d$ is a subset of at most $2s$ points with aspect ratio

$$\Phi \stackrel{\text{def}}{=} \frac{\max_{x, y \in A \cup B} \|x - y\|_p}{\min_{\substack{x, y \in A \cup B \\ x \neq y}} \|x - y\|_p}.$$

Then, we let $R = t_p \cdot \max_{x, y \in A \cup B} \|x - y\|_p$, and

$$m = O(t_p \Phi \log s).$$

For any fixed setting of $x, y \in A \cup B$, the distance between $\phi(x)$ and $\phi(y)$ is the number of independent of executions of \mathbf{f} where $\mathbf{f}(x) \neq \mathbf{f}(y)$. Since executions are independent, we apply a Chernoff bound to say that with probability $1 - 1/s^{100}$ every $x, y \in A \cup B$ has $\|\phi(x) - \phi(y)\|_1$ concentrating up to a constant factor around its expectation, and therefore, we apply a union bound over $4s^2$ many pairs of points in $A \cup B$ to conclude ϕ is a constant distortion embedding. The case $p = 1$ follows similarly, except that

$$\mathbf{E}_{\phi} [\|\phi(x) - \phi(y)\|_1] \asymp m \cdot \frac{\|x - y\|_1}{dR},$$

and $R = 2 \max_{x, y \in A \cup B} \|x - y\|_{\infty}$. In summary, we have the following lemmas.

Lemma A.2. *Fix any $p \in (1, 2]$, $n, d \in \mathbb{N}$, and a parameter $\Phi \in \mathbb{R}^{\geq 0}$. There exists a distribution \mathcal{E}_p over embeddings $\phi: \mathbb{R}^d \rightarrow \{0, 1\}^{d'}$, where*

$$d' = O(\Phi \log n),$$

such that for any fixed set $X \subset \mathbb{R}^d$ of size at most n and aspect ratio in ℓ_p at most Φ , a draw $\phi \sim \mathcal{E}_p$ is a constant distortion embedding of X into the hypercube with Hamming distance with probability at least $1 - 1/n^{10}$.

Lemma A.3. *Fix any $n, d \in \mathbb{N}$, and a parameter $r, R \in \mathbb{R}^{\geq 0}$. There exists a distribution \mathcal{E}_1 over embeddings $\phi: \mathbb{R}^d \rightarrow \{0, 1\}^{d'}$, where*

$$d' = O\left(\frac{dR \log n}{r}\right),$$

such that for any fixed set $X \subset \mathbb{R}^d$ of size at most n and ℓ_{∞} distance at most R and ℓ_1 distance at least r , a draw $\phi \sim \mathcal{E}_1$ is a constant distortion embedding of X into the hypercube with Hamming distance with probability at least $1 - 1/n^{10}$.

B Bit Complexity and Randomness

In this section, we handle several details relating to the bit complexity and generation of randomness in the streaming model as raised in Section 5.2. In Sections 6 and 7, we analyzed one-and two pass *linear sketching* algorithms for EMD and MST. However, our algorithm assumed that we could generate real-valued random variables (e.g. Cauchy or exponential random variables). Of course, a small-space streaming algorithm can only generate such variables to finitely many bits of precision. Similarly, we assumed that all random variables generated at the start of the algorithm could be stored, without affecting the space of the algorithm — this is known as the random oracle model (Definition 5.8). In this section, we demonstrate how both of these details can be addressed by generating random variables to finite precision, and using pseudo-random generators to avoid the random oracle assumption.

Lemma B.1. *The one and two-pass linear sketching algorithms in Sections 6 and 7 can be implemented by generating each random variable to $\text{polylog}(n)$ bits of precision, so that the space required to store the linear sketches is within a $\text{polylog}(n)$ factor of the dimension of the sketch.*

Proof: We first handle the issue of bit-complexity. Specifically, we demonstrate that it suffices to generate each random variable used in the algorithms to $\delta = \text{polylog}(n)$ bits of precision. Since every entry of the linear sketches is a sum of at most n such variables, multiplied by integer coefficients of the vectors f_X or $f_{A,B}$ which are bounded in magnitude by n , the result will follow. Firstly, we argue that every random variable generated by the algorithms are bounded by $2^{\text{polylog}n}$ in magnitude with high probability. Note that the only unbounded distributions that we use in Sections 6 and 7 are the exponential distribution, and the p -stable distribution for $p \in \{1, \text{poly}(\frac{1}{\log n})\}$. Using the cdf of an exponential, $\text{poly}(n)$ exponentials generated in the sketches will be bounded by $\text{poly}(n)$ with probability $1 - \exp(-\text{poly}(n))$. Using tail bounds for p -stable variables [Nol09], the same is true for Cauchy (1-stable) random variables. Using the method for generating p -stable variables from Proposition 9.2, we have that a p -stable variable is given by:

$$\mathbf{x} = \sin(p\theta) \cdot \left(\frac{\cos(\theta(1-p))}{\cos^{\frac{1}{1-p}}(\theta) \ln(1/r)} \right)^{\frac{1-p}{p}}$$

where $\theta \sim [-\frac{\pi}{2}, \frac{\pi}{2}]$ and $r \sim [0, 1]$. Note that $|\theta| - \pi/2 > 1/\text{poly}(n)$ and $r > 1/\text{poly}(n)$ imply that $\frac{\cos(\theta(1-p))}{\cos^{\frac{1}{1-p}}(\theta) \ln(1/r)} < \text{poly}(n)$ for $p < 1/2$. It follows that $|\mathbf{x}| < 2^{\text{polylog}n}$ with probability $1 - 1/\text{poly}(n)$, and then one can union bound over all the p -stable variables generated.

By the above discussion, it follows that generating each random variable to δ -bits of precision results in an additive $1/2^{\text{polylog}n}$ error in each coordinate of the linear sketch $\mathbf{S}, \mathbf{S}f$. Moreover, generating the p -stables to δ bits of precision can be accomplished by generating r, θ to $\text{poly}(\delta)$ -bits (see argument in Appendix A6 of [KNW10]). It suffices to prove that this will not effect the output of the algorithm significantly. This argument is standard using Lipschitzness of the output of the linear sketching algorithm on the sketch (see, Appendix A6 of [KNW10]). The only step of the algorithm which is not Lipschitz in the sketch $\mathbf{S}f$ is the generation of a sampled edge in the Quadtree from the sketch in the one-pass algorithms (the two-pass algorithm from Section 6 uses the L_p sampler of [JW21] which already uses bounded bit complexity) — we argue that the probability that the

sample changes after truncating is small. To see this, note that our algorithms in Sections 6 and 7 are only required to recover whenever the events \mathcal{E}_1 and \mathcal{E}_2 hold. By the proofs of Lemma 6.11 (for EMD) and Lemma 7.9 (for MST) there will be a $1/\text{poly}(n)$ sized gap between the biggest and second largest estimated coordinate. As a result, whenever these two events hold, the additive $2^{-\text{polylog}n}$ error will not effect the sample obtained by the algorithm. Moreover, when \mathcal{E}_1 and \mathcal{E}_2 do not hold, recall that our algorithm is allowed to output any sampled edge (this error is absorbed into the variational distance of the algorithm). It follows that the remainder of the analysis of the sampling algorithm is unaffected by the additive $2^{-\text{polylog}n}$ to the coordinates of the sketch, which completes the proof. ■

Lemma B.2. *The one and two-pass linear sketching algorithms in Sections 6 and 7 can be derandomized with an additive $\tilde{O}(d \cdot \text{polylog}n)$ bits of space.*

Proof: We first handle the randomness required generation of the Quadtree. By Remark 6, in each depth $t \in [h]$ of the Quadtree we can sample the same set $(i_1, i_2, \dots, i_{2^t}) \sim [d]$ of coordinates for each vertex at that depth (instead of independently sampling coordinates in every vertex at the same depth). Since $h = \log 2d$, the total number of bits we must sample to define an entire quadtree is $2d \log d$. The algorithm can then generate this randomness and store it, resulting in an additive $O(d \log n)$ bits of space

Now that the Quadtree is generated, since the sketches used in Section 5.3 are already derandomized, it remains only to derandomize the exponential and p -stable random variables. Since these variables are independently drawn for separate edges of the tree, by the standard reordering argument of Indyk [Ind06a] for linear sketches, we can apply Nisan’s PRG [Nis92] to randomize these random variables with only a $\log(S)$ -blowup in the space complexity of the algorithm, where S is the space complexity of storing the sketch $\mathbf{S}f$. Since the latter is at most $\text{poly}(d, \log n)$, the blow-up is at most a $\log nd$ factor, which completes the proof. ■

References

- [ABIW09] Alexandr Andoni, Khanh Do Ba, Piotr Indyk, and David Woodruff. Efficient sketches for earth-mover distance, with applications. In *Proceedings of the 50th Annual IEEE Symposium on Foundations of Computer Science (FOCS '2009)*, 2009.
- [ACB17] Martin Arjovsky, Soumith Chintala, and Léon Bottou. Wasserstein generative adversarial networks. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pages 214–223, 2017.
- [AIK08] Alexandr Andoni, Piotr Indyk, and Robert Krauthgamer. Earth mover distance over high-dimensional spaces. In *Proceedings of the 19th ACM-SIAM Symposium on Discrete Algorithms (SODA '2008)*, pages 343–352, 2008.
- [AKO10] Alexandr Andoni, Robert Krauthgamer, and Krzysztof Onak. Streaming algorithms from precision sampling. *arXiv preprint arXiv:1011.1263*, 2010.

- [AKR15] Alexandr Andoni, Robert Krauthgamer, and Ilya Razenshteyn. Sketching and embedding are equivalent for norms. In *Proceedings of the 47th ACM Symposium on the Theory of Computing (STOC '2015)*, pages 479–488, 2015. Available as arXiv:1411.2577.
- [ANOY14] Alexandr Andoni, Aleksandar Nikolov, Krzysztof Onak, and Grigory Yaroslavtsev. Parallel algorithms for geometric graph problems. In *Proceedings of the 46th ACM Symposium on the Theory of Computing (STOC '2014)*, 2014.
- [AS14] Pankaj K. Agarwal and R. Sharathkumar. Approximation algorithms for bipartite matching with metric and geometric costs. In *Proceedings of the 46th ACM Symposium on the Theory of Computing (STOC '2014)*, pages 555–564, 2014.
- [BBD⁺02] Brian Babcock, Shivnath Babu, Mayur Datar, Rajeev Motwani, and Jennifer Widom. Models and issues in data stream systems. In *Proceedings of the twenty-first ACM SIGMOD-SIGACT-SIGART symposium on Principles of database systems*, pages 1–16. ACM, 2002.
- [BBD⁺17] MohammadHossein Bateni, Soheil Behnezhad, Mahsa Derakhshan, MohammadTaghi Hajiaghayi, Raimondas Kiveris, Solvio Lattanzi, and Vahab Mirrokni. Affinity clustering: Hierarchical clustering at scale. In *Proceedings of Advances in Neural Information Processing Systems 30 (NeurIPS '2017)*, 2017.
- [BDI⁺20] Arturs Backurs, Yihe Dong, Piotr Indyk, Ilya Razenshteyn, and Tal Wagner. Scalable nearest neighbor search for optimal transport. In *Proceedings of the 37th International Conference on Machine Learning (ICML '2020)*, 2020.
- [BI14] Arturs Bačkurs and Piotr Indyk. Better embeddings for planar earth-mover distance over sparse sets. In *Proceedings of the 41st International Colloquium on Automata, Languages and Programming (ICALP '2014)*, 2014.
- [BVDPPH11] Nicolas Bonneel, Michiel Van De Panne, Sylvain Paris, and Wolfgang Heidrich. Displacement interpolation using lagrangian mass transport. In *Proceedings of the 2011 SIGGRAPH Asia Conference*, pages 1–12, 2011.
- [CCFC02] Moses Charikar, Kevin Chen, and Martin Farach-Colton. Finding frequent items in data streams. *Automata, languages and programming*, pages 784–784, 2002.
- [Cha02] Moses Charikar. Similarity estimation techniques from rounding algorithms. In *Proceedings of the 34th ACM Symposium on the Theory of Computing (STOC '2002)*, pages 380–388, 2002.
- [CMS76] John M Chambers, Colin L Mallows, and BW Stuck. A method for simulating stable random variables. *Journal of the american statistical association*, 71(354):340–344, 1976.
- [FCCR18] Rémi Flamary, Marco Cuturi, Nicolas Courty, and Alain Rakotomamonjy. Wasserstein discriminant analysis. *Machine Learning*, 107(12):1923–1945, 2018.

- [FIS05] Gereon Frahling, Piotr Indyk, and Christian Sohler. Sampling in dynamic data streams and applications. In *Proceedings of the 21st ACM Symposium on Computational Geometry (SoCG '2005)*, 2005.
- [HIM12] Sariel Har-Peled, Piotr Indyk, and Rajeev Motwani. Approximate nearest neighbor: Towards removing the curse of dimensionality. *Theory of Computing*, 8(1):321–350, 2012.
- [HTF01] Trevor Hastie, Robert Tibshirani, and Jerome Friedman. *Elements of statistical learning: data mining, inference, and prediction*. Springer, 2001.
- [Ind04] Piotr Indyk. Algorithms for dynamic geometric problems over data streams. In *Proceedings of the 36th ACM Symposium on the Theory of Computing (STOC '2004)*, 2004.
- [Ind06a] Piotr Indyk. Stable distributions, pseudorandom generators, embeddings, and data stream computation. *Journal of the ACM (JACM)*, 53(3):307–323, 2006.
- [Ind06b] Piotr Indyk. Stable distributions, pseudorandom generators, embeddings, and data stream computation. *Journal of the ACM*, 53(3):307–323, 2006.
- [IT03] Piotr Indyk and Nitin Thaper. Fast color image retrieval via embeddings. In *Workshop on Statistical and Computational Theories of Vision (at ICCV)*, 2003.
- [JST11] Hossein Jowhari, Mert Sağlam, and Gábor Tardos. Tight bounds for lp samplers, finding duplicates in streams, and related problems. In *Proceedings of the Thirtieth ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems, PODS '11*, pages 49–58, New York, NY, USA, 2011. ACM.
- [JW21] Rajesh Jayaram and David Woodruff. Perfect l_p sampling in a data stream. *SIAM Journal on Computing*, 50(2):382–439, 2021.
- [KN06] Subhash Khot and Assaf Naor. Nonembeddability theorems via fourier analysis. *Mathematische Annalen*, 334(4):821–852, 2006.
- [KNP19] Andrey Boris Khesin, Aleksandar Nikolov, and Dmitry Paramonov. Preconditioning for the geometric transportation problem. In *Proceedings of the 35th International Symposium on Computational Geometry (SoCG '2019)*, 2019.
- [KNW10] Daniel M Kane, Jelani Nelson, and David P Woodruff. On the exact space complexity of sketching and streaming small norms. In *Proceedings of the twenty-first annual ACM-SIAM symposium on Discrete Algorithms*, pages 1161–1178. SIAM, 2010.
- [KSKW15] Matt Kusner, Yu Sun, Nicholas Kolkin, and Kilian Weinberger. From word embeddings to document distances. In *Proceedings of the 32nd International Conference on Machine Learning (ICML '2015)*, 2015.
- [M⁺05] Shanmugavelayutham Muthukrishnan et al. Data streams: Algorithms and applications. *Foundations and Trends[®] in Theoretical Computer Science*, 1(2):117–236, 2005.

- [MJ15] Jonas W Mueller and Tommi Jaakkola. Principal differences analysis: Interpretable characterization of differences between distributions. In *Advances in Neural Information Processing Systems*, pages 1702–1710, 2015.
- [MS13] Andrew McGregor and Daniel Stubbs. Sketching earth-mover distance on graph metrics. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, pages 274–286. Springer, 2013.
- [MSC⁺13] Tomas Mikolov, Ilya Sutskever, Kai Chen, Greg S Corrado, and Jeff Dean. Distributed representations of words and phrases and their compositionality. In *Proceedings of Advances in Neural Information Processing Systems (NIPS '2013)*, pages 3111–3119, 2013.
- [MW10] Morteza Monemizadeh and David P Woodruff. 1-pass relative-error lp-sampling with applications. In *Proceedings of the twenty-first annual ACM-SIAM symposium on Discrete Algorithms*, pages 1143–1160. SIAM, 2010.
- [Nag06] HN Nagaraja. Order statistics from independent exponential random variables and the sum of the top order statistics. *Advances in Distribution Theory, Order Statistics, and Inference*, pages 173–185, 2006.
- [Nis92] Noam Nisan. Pseudorandom generators for space-bounded computation. *Combinatorica*, 12(4):449–461, 1992.
- [Nol09] John P Nolan. Stable distributions: Models for heavy tailed data. <http://academic2.american.edu/~jpnolan/stable/chap1.pdf>, 2009.
- [NS07] Assaf Naor and Gideon Schechtman. Planar earthmover is not in L_1 . *SIAM Journal on Computing*, 37(3):804–826, 2007. An extended abstract appeared in FOCS’06.
- [NW70] Saul B Needleman and Christian D Wunsch. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *Journal of molecular biology*, 48(3):443–453, 1970.
- [PC19] Gabriel Peyré and Marco Cuturi. Computational optimal transport: With applications to data science. *Foundations and Trends® in Machine Learning*, 11(5–6):355–607, 2019.
- [PSM14] Jeffrey Pennington, Richard Socher, and Christopher Manning. Glove: Global vectors for word representation. In *Proceedings of the 2014 conference on empirical methods in natural language processing (EMNLP)*, pages 1532–1543, 2014.
- [RTG00a] Yossi Rubner, Carlo Tomasi, and Leonidas J. Guibas. The Earth Mover’s Distance as a metric for image retrieval. *International Journal of Computer Vision*, 40(2):99–121, 2000.
- [RTG00b] Yossi Rubner, Carlo Tomasi, and Leonidas J Guibas. The earth mover’s distance as a metric for image retrieval. *International journal of computer vision*, 40(2):99–121, 2000.

- [SA12] R. Sharathkumar and Pankaj K. Agarwal. A near-linear time ϵ -approximation algorithm for bipartite geometric matching. In *Proceedings of the 42nd ACM Symposium on the Theory of Computing (STOC '2012)*, 2012.
- [SDGP⁺15] Justin Solomon, Fernando De Goes, Gabriel Peyré, Marco Cuturi, Adrian Butscher, Andy Nguyen, Tao Du, and Leonidas Guibas. Convolutional wasserstein distances: Efficient optimal transportation on geometric domains. *ACM Transactions on Graphics (TOG)*, 34(4):1–11, 2015.
- [She17] Jonah Sherman. Generalized preconditioning and undirected minimum cost flow. In *Proceedings of the 28th ACM-SIAM Symposium on Discrete Algorithms (SODA '2017)*, 2017.
- [YO14] Arman Yousefi and Rafail Ostrovsky. Improved approximation algorithms for earth-mover distance in data streams. *arXiv preprint arXiv:1404.6287*, 2014.
- [YV18] Grigory Yaroslavtsev and Adithya Vadapalli. Massively parallel algorithms and hardness for single-linkage clustering under ℓ_p -distances. In *Proceedings of the 35th International Conference on Machine Learning (ICML '2018)*, 2018.