

Sharp ill-posedness for the generalized Camassa-Holm equation in Besov spaces

Jinlu Li¹, Yanghai Yu^{2,*} and Weipeng Zhu³

¹ School of Mathematics and Computer Sciences, Gannan Normal University, Ganzhou 341000, China

² School of Mathematics and Statistics, Anhui Normal University, Wuhu 241002, China

³ School of Mathematics and Big Data, Foshan University, Foshan, Guangdong 528000, China

January 24, 2022

Abstract: In this paper, we consider the Cauchy problem for the generalized Camassa-Holm equation that includes the Camassa-Holm as well as the Novikov equation on the line. We present a new and unified method to prove the sharp ill-posedness for the generalized Camassa-Holm equation in $B_{p,\infty}^s$ with $s > \max\{1 + 1/p, 3/2\}$ and $1 \leq p \leq \infty$ in the sense that the solution map to this equation starting from u_0 is discontinuous at $t = 0$ in the metric of $B_{p,\infty}^s$. Our results cover and improve the previous work given in [22], solving an open problem left in [22].

Keywords: Generalized Camassa-Holm equation; Ill-posedness; Besov space.

MSC (2010): 35Q53, 37K10.

1 Introduction

In this paper, we consider the Cauchy problem for the generalized Camassa-Holm-Novikov (gCHN) equation which was proposed by Anco, Silva and Freire [1] as follows

$$\begin{cases} m_t + u^k m_x + (k+1)u^{k-1}u_x m = 0, & (x, t, k) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{Z}^+, \\ m = u - u_{xx}, \\ u(0, x) = u_0(x), & x \in \mathbb{R}. \end{cases} \quad (1.1)$$

The gCHN is an evolution equation with $(k+1)$ -order nonlinearities, and can be regarded as a subclass of the generalized Camassa-Holm (g-kbCH) equation considered in [15, 17]

$$m_t + u^k m_x + bu^{k-1}u_x m = 0, \quad k \in \mathbb{Z}^+, b \in \mathbb{R}.$$

*E-mail: lijnlu@gnnu.edu.cn; yuyanghai214@sina.com (Corresponding author); mathzwp2010@163.com

When $k = 1$, (1.1) reduces to the classical Camassa-Holm (CH) equation [8–12]

$$u_t - u_{xxt} + 3uu_x = 2u_x u_{xx} + uu_{xxx}, \quad (1.2)$$

which was originally derived as a bi-Hamiltonian system by Fokas and Fuchssteiner [14] in the context of the KdV model and gained prominence after Camassa-Holm [3] independently re-derived it as an approximation to the Euler equations of hydrodynamics. (1.2) is completely integrable [3, 7] with a bi-Hamiltonian structure [6, 14] and infinitely many conservation laws [3, 14]. Also, it admits exact peaked soliton solutions (peakons) of the form $ce^{-|x-ct|}$ with $c > 0$, which are orbitally stable [13] and models wave breaking (i.e., the solution remains bounded, while its slope becomes unbounded in finite time [5, 9, 10]).

When $k = 2$, (1.1) becomes the famous Novikov equation [18, 19, 23–28, 30]

$$u_t - u_{xxt} + 4u^2 u_x = 3uu_x u_{xx} + u^2 u_{xxx}. \quad (1.3)$$

Home-Wang [20] proved that the Novikov equation with cubic nonlinearity shares similar properties with the CH equation, such as a Lax pair in matrix form, a bi-Hamiltonian structure, infinitely many conserved quantities and peakon solutions given by the formula $u(x, t) = \sqrt{c}e^{-|x-ct|}$.

Setting $\Lambda^{-2} = (1 - \partial_x^2)^{-1}$, then we transform (1.1) equivalently into the following nonlinear transport type equation

$$\begin{cases} u_t + u^k u_x = \mathbf{P}(u) + \mathbf{Q}(u), \\ u(0, x) = u_0(x), \end{cases} \quad (1.4)$$

where

$$\mathbf{P}(u) := -\partial_x \Lambda^{-2} \left(\frac{2k-1}{2} u^{k-1} u_x^2 + u^{k+1} \right) \quad \text{and} \quad \mathbf{Q}(u) := -\frac{k-1}{2} \Lambda^{-2} (u^{k-2} u_x^3). \quad (1.5)$$

As shown in [1, 15, 17], the gCHN equation (1.1) admits a local conservation law, possesses single peakons of the form $u(x, t) = c^{1/k} e^{-|x-ct|}$ as well as multi-peakon solutions and exhibits wave breaking phenomena (see [29] and the references therein).

In recent years, the issue of well-posedness in different spaces for the g-kbCH equation has been a fascinating object of research due to its abundant physical and mathematical properties and a series of achievements have been made in the study of the g-kbCH equation. For general k , using a Galerkin-type approximation scheme, Himonas-Holliman [17] established the local well-posedness of the g-kbCH equation in the Sobolev space $H^s(\mathbb{R} \text{ or } \mathbb{T})$. Zhao-Li-Yan [31] extended the above well-posedness result to the Besov space $B_{p,r}^s(\mathbb{R})$ with $s > \max\{1 + \frac{1}{p}, \frac{3}{2}\}$ and $1 \leq p, r \leq \infty$. However, for $r = \infty$, they established the continuity of the data-to-solution map in a weaker topology. Subsequently, Chen-Li-Yan [4] solved the critical case for $(s, p, r) = (\frac{3}{2}, 2, 1)$. Guo, Liu, Molinet and Yin [16] established the ill-posedness of the Camassa-Holm equation in the critical Sobolev space $H^{3/2}(\mathbb{R} \text{ or } \mathbb{T})$ and even in the Besov space $B_{p,r}^{1+1/p}(\mathbb{R} \text{ or } \mathbb{T})$ with $p \in [1, \infty]$, $r \in (1, \infty]$ by proving the norm inflation. In our recent paper [22], we proved the solution map to the Camassa-Holm equation starting from u_0 is discontinuous at $t = 0$ in the metric of $B_{p,\infty}^s(\mathbb{R})$, which implies the ill-posedness for this equation in $B_{p,\infty}^s(\mathbb{R})$. More precisely, we established

Theorem 1.1 (See [22]) *Let $s > 2 + \max \{3/2, 1 + 1/p\}$ with $1 \leq p \leq \infty$. There exists $u_0 \in B_{p,\infty}^s(\mathbb{R})$ and a positive constant ε_0 such that the data-to-solution map $u_0 \mapsto \mathbf{S}_t(u_0)$ of the Cauchy problem (1.4)-(1.5) with $k = 1$ satisfies*

$$\limsup_{t \rightarrow 0^+} \|\mathbf{S}_t(u_0) - u_0\|_{B_{p,\infty}^s} \geq \varepsilon_0.$$

In addition, for the Novikov equations, we proved that Theorem 1.1 holds for only $p = 2$ in [22], while for the case $p \neq 2$ the difficulty lies mainly in the construction of initial data u_0 due to the appearance of $u_0^2 \partial_x u_0$. Naturally, the method in [22] seems to be invalid when proving the ill-posedness for (1.4)-(1.5) with $k \geq 3$ in $B_{p,\infty}^s(\mathbb{R})$ ($p \neq 2$) since the construction of initial data make the computation of $u_0^k \partial_x u_0$ more difficult.

In this present paper, we shall develop a new and unified method to study the ill-posedness problem for (1.4)-(1.5) with general $k \in \mathbb{Z}^+$. Our main aim is to prove the solution map to the Cauchy problem (1.4)-(1.5) starting from u_0 is discontinuous at $t = 0$ in the metric of $B_{p,\infty}^s(\mathbb{R})$, which implies the ill-posedness for this equation in $B_{p,\infty}^s(\mathbb{R})$. Furthermore, we expect that the sharp index pair (s, p) satisfies that $s > \max \{3/2, 1 + 1/p\}$ with $1 \leq p \leq \infty$. Now let us state our main result of this paper.

Theorem 1.2 *Let $k \in \mathbb{Z}^+$ be fixed. Assume that*

$$s > \max \left\{ \frac{3}{2}, 1 + \frac{1}{p} \right\} \quad \text{with} \quad 1 \leq p \leq \infty. \quad (1.6)$$

There exists $u_0 \in B_{p,\infty}^s(\mathbb{R})$ and a positive constant ε_0 such that the data-to-solution map $u_0 \mapsto \mathbf{S}_t(u_0)$ of the Cauchy problem (1.4)-(1.5) satisfies

$$\limsup_{t \rightarrow 0^+} \|\mathbf{S}_t(u_0) - u_0\|_{B_{p,\infty}^s} \geq \varepsilon_0.$$

Remark 1.1 *As mentioned above, system (1.4)-(1.5) unifies the Camassa-Holm and Novikov equations. In [22], we only obtained the ill-posedness for the Novikov equation in $B_{2,\infty}^s$ with $s > \frac{7}{2}$ due to the technical difficulty. Thus Theorem 1.2 covers our recent results on both the Camassa-Holm and Novikov equations in [22]. Also, our method here is new and makes the proof simpler.*

The Cauchy problem for the Degasperis-Procesi equation reads as

$$\begin{cases} \partial_t u + uu_x = -\frac{3}{2} \partial_x (1 - \partial_x^2)^{-1} (u^2), & (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \\ u(0, x) = u_0(x), & x \in \mathbb{R}. \end{cases} \quad (1.7)$$

Remark 1.2 *We should mention that the H^1 norm of solutions to the g -kbCH is conserved if and only if $b = k + 1$, which naturally excludes the Degasperis-Procesi equation for the case $k = 1$ and $b = 3$. However, following the procedure in the proof of Theorem 1.2 with suitable modification, we can prove Theorem 1.2 holds for the Degasperis-Procesi equation.*

2 Preliminaries

Notation The notation $A \lesssim B$ (resp., $A \gtrsim B$) means that there exists a harmless positive constant c such that $A \leq cB$ (resp., $A \geq cB$). Given a Banach space X , we denote its norm by $\|\cdot\|_X$. For $I \subset \mathbb{R}$, we denote by $\mathcal{C}(I; X)$ the set of continuous functions on I with values in X . Sometimes we will denote $L^p(0, T; X)$ by $L_T^p X$.

Next, we will recall some facts about the Littlewood-Paley decomposition, the nonhomogeneous Besov spaces and some of their useful properties (For more details, see [2]). Let $\mathcal{B} := \{\xi \in \mathbb{R} : |\xi| \leq \frac{4}{3}\}$ and $\mathcal{C} := \{\xi \in \mathbb{R} : \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$. There exist two radial functions $\chi \in C_c^\infty(\mathcal{B})$ and $\varphi \in C_c^\infty(\mathcal{C})$ both taking values in $[0, 1]$ such that

$$\chi(\xi) + \sum_{j \geq 0} \varphi(2^{-j}\xi) = 1 \quad \forall \xi \in \mathbb{R}^d.$$

Definition 2.1 (See [2]) *For every $u \in \mathcal{S}'(\mathbb{R})$, the Littlewood-Paley dyadic blocks Δ_j are defined as follows*

$$\Delta_j u = \begin{cases} 0, & \text{if } j \leq -2; \\ \chi(D)u = \mathcal{F}^{-1}(\chi \mathcal{F}u), & \text{if } j = -1; \\ \varphi(2^{-j}D)u = \mathcal{F}^{-1}(\varphi(2^{-j}\cdot) \mathcal{F}u), & \text{if } j \geq 0. \end{cases}$$

Definition 2.2 (See [2]) *Let $s \in \mathbb{R}$ and $(p, r) \in [1, \infty]^2$. The nonhomogeneous Besov space $B_{p,r}^s(\mathbb{R})$ is defined by*

$$B_{p,r}^s(\mathbb{R}) := \left\{ f \in \mathcal{S}'(\mathbb{R}) : \|f\|_{B_{p,r}^s(\mathbb{R})} < \infty \right\},$$

where

$$\|f\|_{B_{p,r}^s(\mathbb{R})} = \begin{cases} \left(\sum_{j \geq -1} 2^{sjr} \|\Delta_j f\|_{L^p(\mathbb{R})}^r \right)^{\frac{1}{r}}, & \text{if } 1 \leq r < \infty, \\ \sup_{j \geq -1} 2^{sj} \|\Delta_j f\|_{L^p(\mathbb{R})}, & \text{if } r = \infty. \end{cases}$$

Remark 2.1 *It should be emphasized that the fact $B_{p,\infty}^s(\mathbb{R}) \hookrightarrow B_{p,\infty}^t(\mathbb{R})$ with $s > t$ will be often used implicitly.*

Finally, we give some important properties which will be also often used throughout the paper.

Lemma 2.1 (See [2]) *Let $(p, r) \in [1, \infty]^2$ and $s > \max\{1 + \frac{1}{p}, \frac{3}{2}\}$. Then we have*

$$\|uv\|_{B_{p,r}^{s-2}(\mathbb{R})} \leq C \|u\|_{B_{p,r}^{s-2}(\mathbb{R})} \|v\|_{B_{p,r}^{s-1}(\mathbb{R})}.$$

Lemma 2.2 (See [2]) *For $(p, r) \in [1, \infty]^2$, $B_{p,r}^{s-1}(\mathbb{R})$ with $s > 1 + \frac{1}{p}$ is an algebra. Moreover, for any $u, v \in B_{p,r}^{s-1}(\mathbb{R})$ with $s > 1 + \frac{1}{p}$, we have*

$$\|uv\|_{B_{p,r}^{s-1}(\mathbb{R})} \leq C \|u\|_{B_{p,r}^{s-1}(\mathbb{R})} \|v\|_{B_{p,r}^{s-1}(\mathbb{R})}.$$

Remark 2.2 Let $(p, r) \in [1, \infty]^2$ and $s > \max\{1 + \frac{1}{p}, \frac{3}{2}\}$, using Lemmas 2.1-2.2, we have

- for the terms $\mathbf{P}(u)$ and $\mathbf{P}(v)$, there holds

$$\|\mathbf{P}(u) - \mathbf{P}(v)\|_{B_{p,r}^{s-1}} \lesssim \|u - v\|_{B_{p,r}^{s-1}} (\|u\|_{B_{p,r}^{s-1}} + \|v\|_{B_{p,r}^{s-1}})^{k-1} (\|u\|_{B_{p,r}^s} + \|v\|_{B_{p,r}^s}). \quad (2.8)$$

- for the terms $\mathbf{Q}(u)$ and $\mathbf{Q}(v)$ (notice that $\mathbf{Q}(u) = 0$ for $k = 1$), there holds

$$\|\mathbf{Q}(u) - \mathbf{Q}(v)\|_{B_{p,r}^{s-1}} \lesssim \|u - v\|_{B_{p,r}^{s-1}} (\|u\|_{B_{p,r}^{s-1}} + \|v\|_{B_{p,r}^{s-1}})^{k-2} (\|u\|_{B_{p,r}^s} + \|v\|_{B_{p,r}^s})^2. \quad (2.9)$$

Lemma 2.3 (See [2]) For $1 \leq p \leq \infty$ and $s > 0$. There exists a constant C , depending continuously on p and s , we have

$$\left\| 2^{js} \|[\Delta_j, v] \partial_x f\|_{L^p} \right\|_{\ell^\infty} \leq C (\|\partial_x v\|_{L^\infty} \|f\|_{B_{p,\infty}^s} + \|\partial_x f\|_{L^\infty} \|\partial_x v\|_{B_{p,\infty}^{s-1}}),$$

where we denote the standard commutator $[\Delta_j, v] \partial_x f = \Delta_j(v \partial_x f) - v \Delta_j \partial_x f$.

3 Proof of Theorem 1.2

3.1 Construction of Initial Data

We need to introduce smooth, radial cut-off functions to localize the frequency region. Precisely, let $\widehat{\phi} \in \mathcal{C}_0^\infty(\mathbb{R})$ be an even, real-valued and non-negative function on \mathbb{R} and satisfy

$$\widehat{\phi}(\xi) = \begin{cases} 1, & \text{if } |\xi| \leq \frac{1}{4}, \\ 0, & \text{if } |\xi| \geq \frac{1}{2}. \end{cases}$$

Lemma 3.1 Define the function $f_n(x)$ by

$$f_n(x) = \phi(x) \cos\left(\frac{17}{12} 2^n x\right) \quad \text{with } n \gg 1.$$

Then we have

$$\Delta_j(f_n) = \begin{cases} f_n, & \text{if } j = n, \\ 0, & \text{if } j \neq n. \end{cases}$$

Proof. See [21].

Lemma 3.2 Define the initial data $u_0(x)$ as

$$u_0(x) := \sum_{n=0}^{\infty} 2^{-ns} \phi(x) \cos\left(\frac{17}{12} 2^n x\right).$$

Then for any $s > \max\{\frac{3}{2}, 1 + \frac{1}{p}\}$ and $k \in \mathbb{Z}^+$, we have for some n large enough

$$\begin{aligned} \|u_0\|_{B_{p,\infty}^s} &\leq C, \\ \|u_0^k \partial_x \Delta_n u_0\|_{L^p} &\geq c 2^{n(1-s)}, \end{aligned}$$

where C and c are some positive constants.

Proof. By the definition of Besov space and the support of $\varphi(2^{-j}\cdot)$, we have

$$\|u_0\|_{B_{p,\infty}^s} \leq C.$$

Using Lemma 3.1 yields

$$\Delta_n u_0(x) = 2^{-ns} \phi(x) \cos\left(\frac{17}{12} 2^n x\right),$$

equivalently,

$$\partial_x \Delta_n u_0 = 2^{-ns} \phi'(x) \cos\left(\frac{17}{12} 2^n x\right) - \frac{17}{12} 2^n 2^{-ns} \phi(x) \sin\left(\frac{17}{12} 2^n x\right).$$

Thus, we have

$$u_0^k \partial_x \Delta_n u_0 = 2^{-ns} u_0^k(x) \phi'(x) \cos\left(\frac{17}{12} 2^n x\right) - \frac{17}{12} 2^n 2^{-ns} u_0^k(x) \phi(x) \sin\left(\frac{17}{12} 2^n x\right).$$

Since $u_0^k(x)$ is a real-valued and continuous function on \mathbb{R} , then there exists some $\delta > 0$ such that

$$|u_0^k(x)| \geq \frac{1}{2} |u_0^k(0)| = \frac{1}{2} \left(\phi(0) \sum_{n=0}^{\infty} 2^{-ns} \right)^k = \frac{2^{sk} \phi^k(0)}{2(2^s - 1)^k} \quad \text{for any } x \in B_\delta(0). \quad (3.10)$$

Thus we have from (3.10)

$$\begin{aligned} \|u_0^k \partial_x \Delta_n u_0\|_{L^p} &\geq C 2^n 2^{-ns} \left\| \phi(x) \sin\left(\frac{17}{12} 2^n x\right) \right\|_{L^p(B_\delta(0))} - C 2^{-ns} \left\| \phi'(x) \phi^k(x) \cos\left(\frac{17}{12} 2^n x\right) \right\|_{L^p} \\ &\geq (c 2^n - C) 2^{-ns}. \end{aligned}$$

We choose n large enough such that $C < \frac{c}{2} 2^n$ and then finish the proof of Lemma 3.2.

3.2 Error Estimates

Proposition 3.1 *Assume that $\|u_0\|_{B_{p,\infty}^s} \lesssim 1$. Let $u \in L_T^\infty B_{p,\infty}^s$ be the solution of the Cauchy problem (1.4), then under the assumptions of Theorem 1.2, we have*

$$\|\mathbf{S}_t(u_0) - u_0\|_{B_{p,\infty}^{s-1}} \lesssim t. \quad (3.11)$$

Furthermore, there holds

$$\|\mathbf{w}\|_{B_{p,\infty}^s} \lesssim t^2, \quad (3.12)$$

here and in what follows we denote

$$\mathbf{w} := \mathbf{S}_t(u_0) - u_0 - t \mathbf{v}_0 \quad \text{with} \quad \mathbf{v}_0 := \mathbf{P}(u_0) + \mathbf{Q}(u_0) - u_0^k \partial_x u_0.$$

Proof. For simplicity, we denote $u(t) := \mathbf{S}_t(u_0)$ here and in what follows. Due to the fact $B_{p,\infty}^s \hookrightarrow \text{Lip}$, we know that there exists a positive time $T = T(\|u_0\|_{B_{p,\infty}^s})$ such that

$$\|u(t)\|_{L_T^\infty B_{p,\infty}^s} \leq C\|u_0\|_{B_{p,\infty}^s} \leq C.$$

Using the Mean Value Theorem and Remark 2.2 with $v = 0$, we obtain from (1.4) that

$$\begin{aligned} \|u(t) - u_0\|_{B_{p,\infty}^{s-1}} &\leq \int_0^t \|\partial_\tau u\|_{B_{p,\infty}^{s-1}} d\tau \\ &\leq \int_0^t (\|\mathbf{P}(u)\|_{B_{p,\infty}^{s-1}} + \int_0^t \|\mathbf{Q}(u)\|_{B_{p,\infty}^{s-1}}) d\tau + \int_0^t \|u^k \partial_x u\|_{B_{p,\infty}^{s-1}} d\tau \\ &\lesssim t \|u\|_{L_t^\infty B_{p,\infty}^s}^{k+1} \\ &\lesssim t \|u_0\|_{B_{p,\infty}^s}^{k+1} \\ &\lesssim t. \end{aligned} \tag{3.13}$$

By the Mean Value Theorem and (1.4), then we obtain that

$$\begin{aligned} \|\mathbf{w}\|_{B_{p,\infty}^{s-2}} &\leq \int_0^t \|\partial_\tau u - \mathbf{v}_0\|_{B_{p,\infty}^{s-2}} d\tau \\ &\lesssim \int_0^t (\|\mathbf{P}(u) - \mathbf{P}(u_0)\|_{B_{p,\infty}^{s-2}} + \|\mathbf{Q}(u) - \mathbf{Q}(u_0)\|_{B_{p,\infty}^{s-2}}) d\tau \\ &\quad + \int_0^t \|u^k \partial_x u - u_0^k \partial_x u_0\|_{B_{p,\infty}^{s-2}} d\tau. \end{aligned} \tag{3.14}$$

Using Remark 2.2 again yields

$$\int_0^t (\|\mathbf{P}(u) - \mathbf{P}(u_0)\|_{B_{p,\infty}^{s-2}} + \|\mathbf{Q}(u) - \mathbf{Q}(u_0)\|_{B_{p,\infty}^{s-2}}) d\tau \lesssim t^2, \tag{3.15}$$

Notice that the simple fact

$$a^{k+1} - b^{k+1} = (k+1)(a-b)\xi^k, \quad \xi \text{ is between } a \text{ and } b,$$

we obtain from Lemma 2.2 and (3.13) that

$$\begin{aligned} \|u^k \partial_x u(\tau) - u_0^k \partial_x u_0\|_{B_{p,\infty}^{s-2}} &\lesssim \|u^{k+1}(\tau) - u_0^{k+1}\|_{B_{p,\infty}^{s-1}} \\ &\lesssim \|u(\tau) - u_0\|_{B_{p,\infty}^{s-1}} \|u_0\|_{B_{p,\infty}^{s-1}}^k \\ &\lesssim \tau. \end{aligned} \tag{3.16}$$

Inserting (3.15) and (3.16) into (3.14) yields

$$\|\mathbf{w}\|_{B_{p,\infty}^{s-2}} \lesssim t^2.$$

Thus, we finish the proof of Proposition 3.1.

Now we present the proof of Theorem 1.2.

Proof of Theorem 1.2. Notice that

$$\mathbf{S}_t(u_0) - u_0 = t\mathbf{v}_0 + \mathbf{w}(t, u_0) \quad \text{and} \quad \mathbf{v}_0 = \mathbf{P}(u_0) + \mathbf{Q}(u_0) - u_0^k \partial_x u_0,$$

by the triangle inequality and Propositions 3.1, we deduce that

$$\begin{aligned} \|\mathbf{S}_t(u_0) - u_0\|_{B_{p,\infty}^s} &\geq 2^{ns} \|\Delta_n(\mathbf{S}_t(u_0) - u_0)\|_{L^p} \\ &= 2^{ns} \|\Delta_n(t\mathbf{v}_0 + \mathbf{w}(t, u_0))\|_{L^p} \\ &\geq t2^{ns} \|\Delta_n(\mathbf{v}_0)\|_{L^p} - 2^{2n} 2^{n(s-2)} \|\Delta_n(\mathbf{w}(t, u_0))\|_{L^p} \\ &\geq t2^{ns} \|\Delta_n(u_0^k \partial_x u_0)\|_{L^p} - t2^{ns} \|\Delta_n(\mathbf{P}(u_0) + \mathbf{Q}(u_0))\|_{L^p} \\ &\quad - C2^{2n} \|\mathbf{w}(t, u_0)\|_{B_{p,\infty}^{s-2}} \\ &\geq t2^{ns} \|u_0^k \partial_x \Delta_n u_0\|_{L^p} - t2^{ns} \|[\Delta_n, u_0^k] \partial_x u_0\|_{L^p} \\ &\quad - t \|\mathbf{P}(u_0) + \mathbf{Q}(u_0)\|_{B_{p,\infty}^s} - C2^{2n} t^2 \\ &\geq t2^{ns} \|u_0^k \partial_x \Delta_n u_0\|_{L^p} - Ct \|2^{ns} \|[\Delta_n, u_0^k] \partial_x u_0\|_{L^p}\|_{\ell^\infty} \\ &\quad - t \|\mathbf{P}(u_0) + \mathbf{Q}(u_0)\|_{B_{p,\infty}^s} - C2^{2n} t^2. \end{aligned} \tag{3.17}$$

By Lemmas 2.2-2.3, one has

$$\|\mathbf{P}(u_0) + \mathbf{Q}(u_0)\|_{B_{p,\infty}^s} \lesssim \|u_0^{k-1}(\partial_x u_0)^2 + u_0^{k+1}\|_{B_{p,\infty}^{s-1}} + \|u_0^{k-2}(\partial_x u_0)^3\|_{B_{p,\infty}^{s-1}} \lesssim 1$$

and

$$\|2^{ns} \|[\Delta_n, u_0^k] \partial_x u_0\|_{L^p}\|_{\ell^\infty} \lesssim \|\partial_x(u_0^k)\|_{L^\infty} \|u_0\|_{B_{p,\infty}^s} + \|\partial_x u_0\|_{L^\infty} \|\partial_x(u_0^k)\|_{B_{p,\infty}^{s-1}} \lesssim 1.$$

Gathering all the above estimates and Lemma 3.2 together with (3.17), we obtain

$$\|\mathbf{S}_t(u_0) - u_0\|_{B_{p,\infty}^s} \geq ct2^n - Ct - C2^{2n} t^2.$$

Taking large n such that $c2^n \geq 2C$, we have

$$\|\mathbf{S}_t(u_0) - u_0\|_{B_{p,\infty}^s} \geq \frac{c}{2} t2^n - C2^{2n} t^2.$$

Thus, picking $t2^n \approx \varepsilon$ with small ε , we have

$$\|\mathbf{S}_t(u_0) - u_0\|_{B_{p,\infty}^s} \geq \frac{c}{2} \varepsilon - C\varepsilon^2 \geq c_1 \varepsilon.$$

This completes the proof of Theorem 1.2.

Acknowledgements. J. Li is supported by the National Natural Science Foundation of China (11801090 and 12161004) and Postdoctoral Science Foundation of China (2020T130129 and 2020M672565). Y. Yu is supported by the National Natural Science Foundation of China (12101011) and Natural Science Foundation of Anhui Province (1908085QA05). W. Zhu is partially supported by the National Natural Science Foundation of China (11901092) and Natural Science Foundation of Guangdong Province (2017A030310634).

Conflict of interest The authors declare that they have no conflict of interest.

References

- [1] S.C. Anco, P.L. da Silva, I.L. Freire, A family of wave-breaking equations generalizing the Camassa-Holm and Novikov equations, *J. Math. Phys.*, 56 (2015) 091506.
- [2] H. Bahouri, J. Y. Chemin, R. Danchin, *Fourier Analysis and Nonlinear Partial Differential Equations*, Grundlehren der Mathematischen Wissenschaften, Springer, Heidelberg, 2011.
- [3] R. Camassa, D. Holm, An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.*, 71 (1993) 1661-1664.
- [4] D. Chen, Y. Li, W. Yan, On the Cauchy problem for a generalized Camassa-Holm equation, *Discr. Contin. Dyn. Syst.*, 35(3) (2015) 871-889.
- [5] A. Constantin, Existence of permanent and breaking waves for a shallow water equation: a geometric approach, *Ann. Inst. Fourier*, 50 (2000), 321-362.
- [6] A. Constantin, The Hamiltonian structure of the Camassa-Holm equation, *Exposition. Math.*, 15 (1997), 53-85.
- [7] A. Constantin, On the scattering problem for the Camassa-Holm equation, *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.*, 457 (2001), 953-970.
- [8] A. Constantin, J. Escher, Global existence and blow-up for a shallow water equation, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 26 (1998), 303-328.
- [9] A. Constantin, J. Escher, Well-posedness, global existence, and blowup phenomena for a periodic quasi-linear hyperbolic equation, *Comm. Pure Appl. Math.*, 51 (1998), 475-504.
- [10] A. Constantin, J. Escher, Wave breaking for nonlinear nonlocal shallow water equations, *Acta Math.*, 181 (1998), 229-243.
- [11] A. Constantin, J. Escher, Particle trajectories in solitary water waves, *Bull. Amer. Math. Soc.*, 44 (2007), 423-431.
- [12] A. Constantin, J. Escher, Analyticity of periodic traveling free surface water waves with vorticity, *Ann. Math.*, 173 (2011), 559-568.
- [13] A. Constantin, W. A. Strauss, Stability of peakons, *Comm. Pure Appl. Math.*, 53 (2000), 603-610.
- [14] A. Fokas, B. Fuchssteiner, Symplectic structures, their bäcklund transformation and hereditary symmetries, *Phys. D*, 4 (1981/1982) 47-66.
- [15] K. Grayshan, A. Himonas, Equations with peakon traveling wave solutions, *Adv. Dyn. Syst. Appl.* 8 (2013) 217-232.
- [16] Z. Guo, X. Liu, L. Molinet, Z. Yin, Ill-posedness of the Camassa-Holm and related equations in the critical space, *J. Differential Equations*, 266, 1698-1707 (2019).
- [17] A. Himonas, C. Holliman, The Cauchy problem for a generalized Camassa-Holm equation, *Adv. Differ. Equ.*, 19 (2014) 161-200.

- [18] A. Himonas, C. Holliman, The Cauchy problem for the Novikov equation, *Nonlinearity*, 25 (2012), 449-479.
- [19] A. Himonas, C. Holliman, C. Kenig, Construction of 2-peakon solutions and ill-posedness for the Novikov equation, *SIAM J. Math. Anal.*, 50(3) (2018), 2968-3006.
- [20] A. Home, J. Wang, Integrable peakon equations with cubic nonlinearity, *J Phys A.*, 41 (2008) 372002: 1-11.
- [21] J. Li, Y. Yu, W. Zhu, Non-uniform dependence on initial data for the Camassa-Holm equation in Besov spaces, *J. Differ. Equ.*, 269 (2020) 8686-8700.
- [22] J. Li, Y. Yu, W. Zhu, Ill-posedness for the Camassa-Holm and related equations in Besov spaces, *J. Differ. Equ.*, 306 (2022), 403–417.
- [23] J. Li, M. Li, W. Zhu, Non-uniform dependence on initial data for the Novikov equation in Besov spaces, *J. Math. Fluid Mech.* 22:50 (2020), 10pp.
- [24] L. Ni, Y. Zhou, Well-posedness and persistence properties for the Novikov equation, *J. Differ. Equ.*, 250 (2011) 3002-3021.
- [25] V. Novikov, Generalization of the Camassa-Holm equation, *J. Phys. A* 42 (2009) 342002.
- [26] X. Wu, Y. Yu, Y. Xiao, Non-uniform dependence on initial data for the generalized Camassa-Holm-Novikov equation in Besov space, *J. Math. Fluid Mech.*, (2021) 23:104.
- [27] X. Wu, Z. Yin, Well-posedness and global existence for the Novikov equation, *Ann. Sc. Norm. Super. Pisa Classe Sci. Ser. V* 11 (2012) 707-727.
- [28] X. Wu, Z. Yin, A note on the Cauchy problem of the Novikov equation, *Appl. Anal.*, 92 (2013) 1116-1137.
- [29] K. Yan, Wave breaking and global existence for a family of peakon equations with high order nonlinearity, *Nonlinear Anal. Real World Appl.*, 45 (2019) 721-735.
- [30] W. Yan, Y. Li, Y. Zhang, The Cauchy problem for the integrable Novikov equation, *J. Differ. Equ.*, 253 (2012) 298-318.
- [31] Y. Zhao, Y. Li, W. Yan, Local well-posedness and persistence property for the generalized Novikov equation, *Discr. Contin. Dyn. Syst.*, 34(2) (2014) 803-820.