

Two-type Estimates for the Boundedness of Generalized Riesz Potential Operator in the Generalized Weighted Local Morrey Spaces

Abdulhamit Kucukaslan^{a,b, 1}

^b*Institute of Mathematics of Czech Academy of Sciences,
115 67, Prague, Czech Republic*

^a*School of Applied Sciences, Pamukkale University, 20680, Denizli, Turkey*

Abstract

In this paper, we prove the Spanne-type boundedness of the generalized Riesz potential operator I_ρ from the one generalized weighted local Morrey spaces $M_{p,\varphi_1}^{\{x_0\}}(w^p, \mathbb{R}^n)$ to the another one $M_{q,\varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)$ with $w^q \in A_{1+\frac{q}{p}}$ for $1 < p < q < \infty$ and from the generalized weighted local Morrey spaces $M_{1,\varphi_1}^{\{x_0\}}(w, \mathbb{R}^n)$ to the weak generalized weighted local Morrey spaces $WM_{q,\varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)$ with $w \in A_{1,q}$ for $1 < q < \infty$. We also prove the Adams-type boundedness of the operator I_ρ from the weighted spaces $M_{p,\varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)$ to the another one $M_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)$ with $w \in A_{p,q}$ for $1 < p < q < \infty$ and from the weighted spaces $M_{1,\varphi}(w, \mathbb{R}^n)$ to the weak weighted spaces $WM_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)$ with $w \in A_{1,q}$ for $1 < q < \infty$.

AMS Mathematics Subject Classification: 42B20, 42B25, 42B35.

Key words: Generalized Riesz potential operator, generalized weighted local Morrey spaces, generalized weighted Morrey spaces, Muckenhoupt-Weeden classes.

¹Corresponding author.

The research of Abdulhamit Kucukaslan was supported by the grant of The Scientific and Technological Research Council of Turkey, Grant TUBITAK-1059B191600675.
E-mail address: kucukaslan@pau.edu.tr (A. Kucukaslan).

1 Introduction

Morrey spaces $M_{p,\lambda}(\mathbb{R}^n)$ were introduced by Morrey in [18] and defined as follows: For $0 \leq \lambda < n, 1 \leq p \leq \infty, f \in M_{p,\lambda}(\mathbb{R}^n)$ if $f \in L_p^{loc}(\mathbb{R}^n)$ and

$$\|f\|_{M_{p,\lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,r))} < \infty$$

holds. Morrey spaces found important applications to potential theory [1], elliptic equations with discontinuous coefficients [3] and Schrödinger equations [26].

On the other hand, on the weighted Lebesgue spaces $L_p(w, \mathbb{R}^n)$, the boundedness of some classical operators were obtained by Muckenhoupt [19], Muckenhoupt and Wheeden [20], and Coifman and Fefferman [4].

Recently, weighted Morrey spaces $M_{p,\kappa}(w, \mathbb{R}^n)$ were introduced by Komori and Shirai [13] as follows: For $1 \leq p \leq \infty, 0 < \kappa < 1$ and w be a weight, $f \in M_{p,\kappa}(w, \mathbb{R}^n)$ if $f \in L_p^{loc}(w, \mathbb{R}^n)$ and

$$\|f\|_{M_{p,\kappa}(w, \mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} w(B(x,r))^{-\frac{\kappa}{p}} \|f\|_{L_p(w, B(x,r))} < \infty.$$

They studied the boundedness of the aforementioned classical operators in these spaces. These results were extended to several other spaces in [11]. Weighted inequalities for fractional operators have applications to potential theory and quantum mechanics.

For a fixed $x_0 \in \mathbb{R}^n$ the generalized weighted local Morrey spaces $M_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)$ are obtained by replacing a function $\varphi(x_0, r)$ instead of r^λ in the definition of weighted local Morrey space, which is the space of all functions $f \in L_p^{loc}(w, \mathbb{R}^n)$ with finite norm

$$\|f\|_{M_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)} = \sup_{r > 0} \varphi(x_0, r)^{-1} w(B(x_0, r))^{-\frac{1}{p}} \|f \chi_{B(x_0, r)}\|_{L_p(w, \mathbb{R}^n)}.$$

During the last decades, the theory of boundedness of classical operators of the harmonic analysis in the generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$ have been well studied by now, we refer the readers to [9, 14, 15, 21] and [23].

For a measurable function $\rho : (0, \infty) \rightarrow (0, \infty)$ the generalized Riesz potential operator (or generalized fractional integral operator) I_ρ is defined by

$$I_\rho f(x) = \int_{\mathbb{R}^n} \frac{\rho(|x-y|)}{|x-y|^n} f(y) dy$$

for any suitable function f on \mathbb{R}^n . If $\rho(t) \equiv t^\alpha$, then we get the Riesz potential operator I_α . The generalized Riesz potential operator I_ρ was initially investigated in [22]. Nowadays many authors have been culminating important observations about I_ρ especially in connection with Morrey spaces. Nakai [22] proved the boundedness of I_ρ from the generalized Morrey spaces $M_{1,\varphi}(\mathbb{R}^n)$ to the spaces $M_{1,\psi}(\mathbb{R}^n)$ for suitable functions φ and ψ . The boundedness of I_ρ from the generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$ to the spaces $M_{q,\psi}(\mathbb{R}^n)$ is studied by Eridani [5], Guliyev et al [9], Kucukaslan et al [14, 15], Kucukaslan [16, 17], Nakai [23] and Nakamura [24].

Spanne-type and Adams-type boundednesses of generalized fractional maximal operator M_ρ in the generalized weighted local Morrey spaces $M_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)$ and generalized weighted Morrey spaces $M_{p,\varphi}(w, \mathbb{R}^n)$ were studied in [17]. But, Spanne-type and Adams-type boundedness of the generalized Riesz potential operator I_ρ in the spaces $M_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)$ and $M_{p,\varphi}(w, \mathbb{R}^n)$ have not been studied, yet.

Spanne [25] and Adams [1] studied boundedness of the Riesz potential in Morrey spaces. Their results, can be summarized as follows.

Theorem A. (*Spanne, but published by Peetre [25]*) Let $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$, $0 < \lambda < n - \alpha p$. Moreover, let $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$ and $\frac{\lambda}{p} = \frac{\mu}{q}$. Then for $p > 1$, the operator I_α is bounded from $M_{p,\lambda}(\mathbb{R}^n)$ to $M_{q,\mu}(\mathbb{R}^n)$ and for $p = 1$, I_α is bounded from $M_{1,\lambda}(\mathbb{R}^n)$ to $WM_{q,\mu}(\mathbb{R}^n)$.

Theorem B. (*Adams [1]*) Let $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$, $0 < \lambda < n - \alpha p$ and $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$. Then for $p > 1$, the operator I_α is bounded from $M_{p,\lambda}(\mathbb{R}^n)$ to $M_{q,\lambda}(\mathbb{R}^n)$ and for $p = 1$, I_α is bounded from $M_{1,\lambda}(\mathbb{R}^n)$ to $WM_{q,\lambda}(\mathbb{R}^n)$.

In the following theorems which were proved in [9], we give Spanne and Adams type results for the boundedness of operator I_ρ on the generalized local Morrey spaces $M_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ and on the generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$, respectively.

Theorem C. (*Spanne type result [9]*) Let $x_0 \in \mathbb{R}^n$, $1 \leq p < q < \infty$, the function ρ satisfy the conditions (3.1), (3.2) and [(3.4) for $w = 1$]. Let also (φ_1, φ_2) satisfy the conditions

$$\operatorname{ess\,inf}_{t < s < \infty} \varphi_1(x_0, s) s^{\frac{n}{p}} \leq C \varphi_2\left(x_0, \frac{t}{2}\right) t^{\frac{n}{q}},$$

$$\int_r^\infty \left(\operatorname{ess\,inf}_{t < s < \infty} \varphi_1(x_0, s) s^{\frac{n}{p}} \right) \frac{\rho(t)}{t^{\frac{n}{p}+1}} dt \leq C \varphi_2(x_0, r),$$

where C does not depend on x_0 and r . Then the operator I_ρ is bounded from

one generalized local Morrey spaces $M_{p,\varphi_1}^{\{x_0\}}(\mathbb{R}^n)$ to another one $M_{q,\varphi_2}^{\{x_0\}}(\mathbb{R}^n)$ for $p > 1$ and from the spaces $M_{1,\varphi_1}^{\{x_0\}}(\mathbb{R}^n)$ to the weak space $WM_{q,\varphi_2}^{\{x_0\}}(\mathbb{R}^n)$ for $p = 1$.

Theorem D. (Adams type result [9]) Let $1 \leq p < \infty$, $q > p$, $\rho(t)$ satisfy the conditions (3.2) and [(3.4) for $w = 1$]. Let also $\varphi(x, t)$ satisfy the conditions

$$\sup_{r < t < \infty} \varphi(x, t) \leq C \varphi(x, r),$$

$$\int_r^\infty \varphi(x, t)^{\frac{1}{p}} \frac{\rho(t)}{t} dt \leq C \rho(r)^{-\frac{p}{q-p}},$$

where C does not depend on $x \in \mathbb{R}^n$ and $r > 0$. Then the operator I_ρ is bounded from the one generalized Morrey space $M_{p,\varphi^{\frac{1}{p}}}(\mathbb{R}^n)$ to another one $M_{q,\varphi^{\frac{1}{q}}}(\mathbb{R}^n)$ for $p > 1$ and from the space $M_{1,\varphi}(\mathbb{R}^n)$ to the weak space $WM_{q,\varphi^{\frac{1}{q}}}(\mathbb{R}^n)$ for $p = 1$.

In this study, by using the method given in [10], we prove the Spanne and Adams type estimates for the boundedness of generalized Riesz potential operator I_ρ on the generalized weighted local Morrey spaces $M_{p,\varphi}^{\{x_0\}}(w^p, \mathbb{R}^n)$ with $1 \leq p < q < \infty$, $w^q \in A_{1+\frac{q}{p}}$ belonging to Muckenhoupt-Weeden class $A_{p,q}$. We find conditions on the triple $(\varphi_1, \varphi_2, \rho)$ which ensure the Spanne-type boundedness of the operator I_ρ from one generalized weighted local Morrey spaces $M_{p,\varphi_1}^{\{x_0\}}(w^p, \mathbb{R}^n)$ to another $M_{q,\varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)$ with $w^q \in A_{1+\frac{q}{p}}$ for $1 < p < q < \infty$ and from $M_{1,\varphi_1}^{\{x_0\}}(w, \mathbb{R}^n)$ to the weighted weak space $WM_{q,\varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)$ with $w \in A_{1,q}$ for $1 < q < \infty$ (see Theorem 3.3). We also find conditions on the pair (φ, ρ) which ensure the Adams-type boundedness of I_ρ from $M_{p,\varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)$ to $M_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)$ for $1 < p < q < \infty$, $w \in A_{1+\frac{q}{p}}$ and from $M_{1,\varphi}(w, \mathbb{R}^n)$ to $WM_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)$ for $1 < q < \infty$, $w \in A_{1,q}$ (see Theorem 4.1).

In all cases the conditions for the boundedness of I_ρ are given in terms of Zygmund-type integral inequalities on the all φ functions and r which do not assume any assumption on monotonicity of $\varphi_1(x, r)$, $\varphi_2(x, r)$ and $\varphi(x, r)$ in r .

By $A \lesssim B$ we mean that $A \leq CB$ with some positive constant C independent of appropriate quantities. If $A \lesssim B$ and $B \lesssim A$, we write $A \approx B$ and say that A and B are equivalent.

2 Preliminaries

Let $x \in \mathbb{R}^n$ and $r > 0$, then we denote by $B(x, r)$ the open ball centered at x of radius r , and by ${}^cB(x, r)$ denote its complement. Let $|B(x, r)|$ be the Lebesgue measure of the ball $B(x, r)$. A weight function is a locally integrable function on \mathbb{R}^n which takes values in $(0, \infty)$ almost everywhere. For a weight w and a measurable set E , we define $w(E) = \int_E w(x) dx$, the Lebesgue measure of E by $|E|$ and the characteristic function of E by χ_E . If w is a weight function, for all $f \in L_1^{loc}$ we denote by $L_p^{loc}(w, \mathbb{R}^n)$ the weighted Lebesgue space defined by the norm

$$\|f\chi_{B(x,r)}\|_{L_p(w, \mathbb{R}^n)} = \left(\int_{B(x,r)} |f(x)|^p w(x) dx \right)^{\frac{1}{p}} < \infty,$$

when $1 \leq p < \infty$ and by

$$\|f\chi_{B(x,r)}\|_{L_1(w, \mathbb{R}^n)} = \operatorname{ess\,sup}_{x \in B(x,r)} |f(x)w(x)| < \infty,$$

when $p = 1$.

We recall that a weight function w belongs to the Muckenhoupt-Wheeden class $A_{p,q}$ (see [19]) for $1 < p < q < \infty$, if

$$\sup_B \left(\frac{1}{|B|} \int_B w(x)^q dx \right)^{\frac{1}{q}} \left(\frac{1}{|B|} \int_B w(x)^{-p'} dx \right)^{\frac{1}{p'}} \leq C,$$

where the supremum is taken with respect to all balls B and $C > 0$. Note that, for all balls B by Hölder's inequality we get

$$|B|^{\frac{1}{p} - \frac{1}{q} - 1} \|w\|_{L_q(B)} \|w^{-1}\|_{L_{p'}(B)} \geq 1. \quad (2.1)$$

If $p = 1$, w is in the $A_{1,q}$ with $1 < q < \infty$ if

$$\sup_B \left(\frac{1}{|B|} \int_B w(x)^q dx \right)^{\frac{1}{q}} \left(\operatorname{ess\,sup}_{x \in B} \frac{1}{w(x)} \right) \leq C,$$

where the supremum is taken with respect to all balls B and $C > 0$.

The weight function w satisfies the reverse doubling condition if there exist constants $\alpha_1 > 1$ and $\alpha_2 < 1$ such that

$$w(B(x, r)) \leq \alpha_2 w(B(x, \alpha_1 r)) \quad (2.2)$$

for arbitrary $x \in \mathbb{R}^n$ and $r > 0$.

Lemma 2.1. [6, 8] If $w \in A_{p,q}$ with $1 < p < q < \infty$, then the following statements are true.

- (i) $w^q \in A_r$ with $r = 1 + \frac{q}{p}$.
- (ii) $w^{-p'} \in A_{r'}$ with $r' = 1 + \frac{p'}{q}$.
- (iii) $w^p \in A_s$ with $s = 1 + \frac{p}{q'}$.
- (iv) $w^{-q'} \in A_{s'}$ with $s' = 1 + \frac{q'}{p}$.

We find it convenient to define the generalized weighted local Morrey spaces in the form as follows.

Definition 2.2. Let $1 \leq p < \infty$ and $\varphi(x, r)$ be a positive measurable function on $\mathbb{R}^n \times (0, \infty)$. For any fixed $x_0 \in \mathbb{R}^n$ we denote by $M_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)$ the generalized weighted local Morrey space, the space of all functions $f \in L_p^{\text{loc}}(w, \mathbb{R}^n)$ with finite quasinorm

$$\|f\|_{M_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)} = \|f(x_0 + \cdot)\|_{M_{p,\varphi}(w, \mathbb{R}^n)}.$$

Also by $WM_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)$ we denote the weak generalized weighted local Morrey space of all functions $f \in WL_p^{\text{loc}}(w, \mathbb{R}^n)$ for which

$$\|f\|_{WM_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n)} = \|f(x_0 + \cdot)\|_{WM_{p,\varphi}(w, \mathbb{R}^n)} < \infty.$$

According to this definition, we recover the weighted local Morrey space $M_{p,\lambda}^{\{x_0\}}(w, \mathbb{R}^n)$ and weighted weak local Morrey space $WM_{p,\lambda}^{\{x_0\}}(w, \mathbb{R}^n)$ under the choice $\varphi(x_0, r) = r^{\frac{\lambda-n}{p}}$:

$$M_{p,\lambda}^{\{x_0\}}(w, \mathbb{R}^n) = M_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n) \Big|_{\varphi(x_0, r) = r^{\frac{\lambda-n}{p}}},$$

$$WM_{p,\lambda}^{\{x_0\}}(w, \mathbb{R}^n) = WM_{p,\varphi}^{\{x_0\}}(w, \mathbb{R}^n) \Big|_{\varphi(x_0, r) = r^{\frac{\lambda-n}{p}}}.$$

We denote by $L_\infty(w, (0, \infty))$ the space of all functions $g(t)$, $t > 0$ with finite norm

$$\|g\|_{L_\infty(w, (0, \infty))} = \sup_{t>0} w(t)g(t)$$

and $L_\infty(0, \infty) \equiv L_\infty(1, (0, \infty))$. Let $\mathfrak{S}(0, 1)$ be the set of all Lebesgue-measurable functions on $(0, 1)$ and $\mathfrak{S}^+(0, 1)$ its subset consisting of all non-negative functions on $(0, 1)$. We denote by $\mathfrak{S}^+(0, 1; \uparrow)$ the cone of all functions in $\mathfrak{S}^+(0, 1)$ which are non-decreasing on $(0, 1)$ and

$$\mathbb{A} = \left\{ \varphi \in \mathfrak{S}^+(0, 1; \uparrow) : \lim_{t \rightarrow 0+} \varphi(t) = 0 \right\}.$$

The following theorem was proved in [9] which we will use while proving our main results.

Theorem 2.3. *Let w_1, w_2 be non-negative measurable functions satisfying $0 < \|w_1\|_{L_1(t,1)} < 1$ for any $t > 0$. Then the identity operator I is bounded from $L_1(w_1, (0,1))$ to $L_1(w_2, (0,1))$ on the cone \mathbb{A} if and only if*

$$\left\| w_2 \left(\|w_1\|_{L_1(\cdot,1)}^{-1} \right) \right\|_{L_1(0,1)} < 1.$$

We will use the following statement on the boundedness of the weighted Hardy operator

$$H_w g(t) := \int_t^1 g(s) w(s) d\mu(s), \quad 0 < t < \infty,$$

where w is weight and $d\mu(s)$ is a non-negative Borel measure on $(0,1)$.

The following theorem was proved in [2].

Theorem 2.4. *Let w_1, w_2 and w be weights on $(0, \infty)$ and $w_1(t)$ be bounded outside a neighborhood of the origin. The inequality*

$$\operatorname{ess\,sup}_{t>0} w_2(t) H_w g(t) \leq C \operatorname{ess\,sup}_{t>0} w_1(t) g(t) \quad (2.3)$$

holds for some $C > 0$ for all non-negative and non-decreasing g on $(0,1)$ if and only if

$$B := \sup_{t>0} w_2(t) \int_t^\infty \frac{w(s) ds}{\operatorname{ess\,sup}_{s<\tau<\infty} w_1(\tau)} < \infty. \quad (2.4)$$

Moreover, the value $C = B$ is the best constant for (2.3).

Remark 2.5. In (2.3) and (2.4) it is assumed that $\frac{1}{1} = 0$ and $0 \cdot 1 = 0$.

3 Spanne-type result for the operator I_ρ in the spaces $M_{p,\varphi}^{\{x_0\}}(w^p, \mathbb{R}^n)$

We assume that

$$\int_1^1 \frac{\rho(t)}{t^n} \frac{dt}{t} < \infty, \quad (3.1)$$

so that the generalized Riesz potential $I_\rho f$ is well defined, at least for characteristic functions $1/|x|^{2n}$ of complementary balls:

$$f(x) = \frac{\chi_{\mathbb{R}^n \setminus B(0,1)}(x)}{|x|^{2n}}.$$

In addition, we shall also assume that ρ satisfies the growth condition: there exist constants $C > 0$ and $0 < 2k_1 < k_2 < 1$ such that

$$\sup_{r < s \leq 2r} \frac{\rho(s)}{s^n} \leq C \int_{k_1 r}^{k_2 r} \frac{\rho(t)}{t^n} \frac{dt}{t}, \quad r > 0. \quad (3.2)$$

This condition is weaker than the usual doubling condition for the function $\frac{\rho(t)}{t^n}$: there exists a constant $C > 0$ such that

$$\frac{1}{C} \frac{\rho(t)}{t^n} \leq \frac{\rho(r)}{r^n} \leq C \frac{\rho(t)}{t^n}, \quad (3.3)$$

whenever r and t satisfy $r, t > 0$ and $\frac{1}{2} \leq \frac{r}{t} \leq 2$.

In the sequel for the generalized Riesz potential operator I_ρ we always assume that ρ satisfies the conditions (3.2) and, then denote the set of all such functions by \tilde{G}_0 . We will write, when $\rho \in \tilde{G}_0$,

$$\tilde{\rho}(r) := Cr^n \int_r^\infty \frac{\rho(t)}{t^n} \frac{dt}{t}.$$

The following lemma is valid for the operator I_ρ .

Lemma 3.1. *[7] Let $w^q \in A_{1+\frac{q}{p}}$ satisfies (2.2), the function ρ satisfies the conditions (3.1)-(3.3), and $f \in L_1^{loc}(w, \mathbb{R}^n)$. Then there exist $C > 0$ for all $B(x, r) \subset \mathbb{R}^n$ such that the inequality*

$$\sup_{x \in \mathbb{R}^n, r > 0} \frac{\rho(r)}{r^n} \left(\int_{B(x,r)} w^q(x) dx \right)^{\frac{1}{q}} \left(\int_{B(x,r)} w(x)^{-p'} dx \right)^{\frac{1}{p'}} \leq C \quad (3.4)$$

is necessary and sufficient condition for the boundedness of generalized Riesz potential operator I_ρ from $L_p(w^p, \mathbb{R}^n)$ to $WL_q(w^q, \mathbb{R}^n)$ for $1 \leq p < q < 1$, and from $L_p(w^p, \mathbb{R}^n)$ to $L_q(w^q, \mathbb{R}^n)$ for $1 < p < q < 1$, $w^q \in A_{1+\frac{q}{p}}$, where the constant C does not depend on f .

The following is weighted local $L_p(\mathbb{R}^n)$ -estimate for the operator I_ρ .

Lemma 3.2. *Let fixed $x_0 \in \mathbb{R}^n$, and $1 \leq p < q < \infty$, $w \in A_{1+\frac{q}{p}}'$ and $\rho(t)$ satisfy the conditions (3.1) and (3.2).*

If the condition (3.4) is fulfilled, then the inequality

$$\begin{aligned} & \|I_\rho f \chi_{B(x_0, r)}\|_{L_q(w^q, \mathbb{R}^n)} \lesssim \|f \chi_{B(x_0, 2r)}\|_{L_p(w^p, \mathbb{R}^n)} \\ & + (w^q(B(x_0, r)))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} (w^q(B(x_0, r)))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t} \end{aligned} \quad (3.5)$$

holds for the ball $B(x_0, r)$ and for all $f \in L_p^{loc}(\mathbb{R}^n, w)$.

If the condition (3.4) is fulfilled, then for $p = 1$ the inequality

$$\begin{aligned} & \|I_\rho f \chi_{B(x_0, r)}\|_{WL_q(w^q)} \lesssim \|f \chi_{B(x_0, 2r)}\|_{L_1(w)} \\ & + (w^q(B(x_0, r)))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{B(x_0, t)}\|_{L_1(w)} (w^q(B(x_0, t)))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t} \end{aligned} \quad (3.6)$$

holds for the ball $B(x_0, r)$ and for all $f \in L_1^{loc}(w)$.

Proof. Let $1 \leq p < q < \infty$ and $w \in A_{1+\frac{q}{p}}'$. For fixed $x_0 \in \mathbb{R}^n$, set $B \equiv B(x_0, r)$ for the ball centered at x_0 and of radius r . Write $f = f_1 + f_2$ with $f_1 = f \chi_{2B}$ and $f_2 = f \chi_{\mathbb{C}_{(2B)}}$. Hence, by the Minkowski inequality we have

$$\|I_\rho f \chi_B\|_{L_q(w^q, \mathbb{R}^n)} \leq \|I_\rho f_1 \chi_B\|_{L_q(w^q, \mathbb{R}^n)} + \|I_\rho f_2 \chi_B\|_{L_q(w^q, \mathbb{R}^n)}.$$

Since $f_1 \in L_p(w^p, \mathbb{R}^n)$, $I_\rho f_1 \in L_q(w^q, \mathbb{R}^n)$ and from condition (3.4) we get the boundedness of I_ρ from $L_p(w^p, \mathbb{R}^n)$ to $L_q(w^q, \mathbb{R}^n)$ (see Lemma 3.1) and it follows that:

$$\|I_\rho f_1 \chi_B\|_{L_q(w^q, \mathbb{R}^n)} \leq \|I_\rho f_1\|_{L_q(w^q, \mathbb{R}^n)} \leq C \|f_1\|_{L_p(w^p, \mathbb{R}^n)} = C \|f \chi_{2B}\|_{L_p(w^p, \mathbb{R}^n)},$$

where constant $C > 0$ is independent of f .

It's clear that $z \in B$, $y \in \mathbb{C}_{(2B)}$ implies $\frac{1}{2}|x_0 - y| \leq |z - y| \leq \frac{3}{2}|x_0 - y|$. Then from conditions (3.1), (3.2) and by Fubini's theorem we have

$$|I_\rho f_2(z)| \lesssim \int_{\mathbb{C}_{(2B)}} \frac{\rho(|x - y|)}{|x - y|^n} |f(y)| dy \lesssim \int_{2r}^1 \int_{B(x_0, t)} |f(y)| dy \frac{\rho(t)}{t^n} \frac{dt}{t}.$$

Applying Hölder's inequality and from (2.1), we get

$$\begin{aligned} & \int_{\mathbb{C}_{(2B)}} \frac{\rho(|x - y|)}{|x - y|^n} |f(y)| dy \\ & \lesssim \int_{2r}^1 \|f \chi_{B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} \|w^{-1} \chi_{B(x_0, t)}\|_{L_{p'}(\mathbb{R}^n)} \frac{\rho(t)}{t^n} \frac{dt}{t} \\ & \lesssim \int_{2r}^1 \|f \chi_{B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} (w^q(B(x_0, t)))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t}. \end{aligned}$$

Moreover, for all $p \in [1, 1)$ the inequality

$$\begin{aligned} & \|I_\rho f_2 \chi_B\|_{L_q(w^p, \mathbb{R}^n)} \\ & \lesssim (w^q(B))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} (w^q(B(x_0, t)))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t} \end{aligned} \quad (3.7)$$

is valid. Thus

$$\begin{aligned} & \|I_\rho f \chi_B\|_{L_q(w^q, \mathbb{R}^n)} \lesssim \|f \chi_{2B}\|_{L_p(w^p, \mathbb{R}^n)} \\ & + (w^q(B))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} \|w^{-1} \chi_{B(x_0, t)}\|_{L_{p'}(\mathbb{R}^n)} \frac{\rho(t)}{t^n} \frac{dt}{t}. \end{aligned}$$

On the other hand,

$$\begin{aligned} & \|f \chi_{2B}\|_{L_p(w^p, \mathbb{R}^n)} \approx \frac{r^{\frac{n}{p}}}{\rho(r)} \|f \chi_{2B}\|_{L_p(w^p, \mathbb{R}^n)} \int_r^1 \frac{\rho(t)}{t^n} \frac{dt}{t} \\ & \leq \frac{r^{\frac{n}{p}}}{\rho(r)} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} \frac{\rho(t)}{t^n} \frac{dt}{t} \\ & \lesssim (w^q(B))^{\frac{1}{q}} \|w^{-1} \chi_B\|_{L_{p'}(\mathbb{R}^n)} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_p(w^p)} \frac{\rho(t)}{t^n} \frac{dt}{t} \\ & \lesssim (w^q(B))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} \|w^{-1} \chi_{B(x_0, t)}\|_{L_{p'}(\mathbb{R}^n)} \frac{\rho(t)}{t^n} \frac{dt}{t} \\ & \lesssim (w^q(B))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} (w^q(B(x_0, t)))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t}. \end{aligned} \quad (3.8)$$

Hence by

$$\|I_\rho f \chi_B\|_{L_q(w^p, \mathbb{R}^n)} \lesssim (w^q(B))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{B(x_0, t)}\|_{L_p(w^p, \mathbb{R}^n)} (w^q(B(x_0, t)))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t}$$

we get the inequality (3.5).

Now let $p = 1$ and $w \in A_{1,q}$. In this case by (3.4) we obtain

$$\begin{aligned}
\|I_\rho f_1 \chi_B\|_{WL_q(w^q, \mathbb{R}^n)} &\leq \|I_\rho f_1\|_{WL_q(w^q, \mathbb{R}^n)} \lesssim \|f_1\|_{L_1(w, \mathbb{R}^n)} = \|f \chi_B\|_{L_1(w, \mathbb{R}^n)} \\
&\approx \frac{r^n}{\rho(r)} \|f \chi_{2B}\|_{L_1(w, \mathbb{R}^n)} \int_r^1 \frac{\rho(t)}{t^n} \frac{dt}{t} \\
&\leq \frac{r^n}{\rho(r)} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_1(w, \mathbb{R}^n)} \frac{\rho(t)}{t^n} \frac{dt}{t} \\
&\lesssim (w^q(B))^{\frac{1}{q}} \|w^{-1} \chi_B\|_{L_1} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_1(w, \mathbb{R}^n)} \frac{\rho(t)}{t^n} \frac{dt}{t} \\
&\lesssim (w^q(B))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_1(w, \mathbb{R}^n)} \|w^{-1} \chi_{B(x_0, t)}\|_{L_1} \frac{\rho(t)}{t^n} \frac{dt}{t} \\
&\lesssim (w^q(B))^{\frac{1}{q}} \int_{2r}^1 \|f \chi_{2B(x_0, t)}\|_{L_1(w, \mathbb{R}^n)} (w^q(B(x_0, t)))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t}. \tag{3.9}
\end{aligned}$$

Then from (3.8) and (3.9) we get the inequality (3.6). \square

The following theorem one of the main result of our paper, in which we prove the Spanne-type estimate for the boundedness of generalized fractional integral operator I_ρ from generalized weighted local Morrey spaces $M_{p, \varphi_1}^{\{x_0\}}(w^p, \mathbb{R}^n)$ to $M_{q, \varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)$.

Theorem 3.3. *Let fixed $x_0 \in \mathbb{R}^n$, $1 \leq p < q < \infty$, $w \in A_{1+\frac{q}{p}}$, the function ρ satisfy the conditions (3.1), (3.2) and (3.4). Let also (φ_1, φ_2) satisfy the conditions*

$$\operatorname{ess\,inf}_{t < s < \infty} \varphi_1(x_0, s) s^{\frac{n}{p}} \leq C \varphi_2(x_0, \frac{t}{2}) t^{\frac{n}{q}}, \tag{3.10}$$

$$\int_r^\infty \frac{\operatorname{ess\,inf}_{t < s < \infty} \varphi_1(x_0, s) (w^p(B(x_0, s)))^{\frac{1}{p}} \rho(t)}{(w^q(B(x_0, t)))^{\frac{1}{q}} t^{\frac{n}{p}}} \frac{dt}{t} \leq C \varphi_2(x_0, r), \tag{3.11}$$

where C does not depend on x and r . Then the operator I_ρ is bounded from $M_{p, \varphi_1}^{\{x_0\}}(w^p, \mathbb{R}^n)$ to $M_{q, \varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)$ for $p > 1$ and from $M_{1, \varphi_1}^{\{x_0\}}(w, \mathbb{R}^n)$ to $WM_{q, \varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)$ for $p = 1$. Moreover, for $p > 1$

$$\|I_\rho f\|_{M_{q, \varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)} \lesssim \|f\|_{M_{p, \varphi_1}^{\{x_0\}}(w^p, \mathbb{R}^n)},$$

and for $p = 1$

$$\|I_\rho f\|_{WM_{q, \varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)} \lesssim \|f\|_{M_{1, \varphi_1}^{\{x_0\}}(w, \mathbb{R}^n)}.$$

Proof. Let $p > 1$. Then by Theorems 2.3, 2.4, Lemma 3.2 and conditions (3.10)-(3.11) we have that

$$\begin{aligned}
\|I_\rho f\|_{M_{q,\varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)} &= \sup_{r>0} \varphi_2(x_0, r)^{-1} (w^q(B(x_0, r)))^{-\frac{1}{q}} \|I_\rho f\|_{L_q(w^q, B(x_0, 2r))} \\
&= \sup_{r>0} \varphi_2(x_0, r)^{-1} (w^q(B(x_0, r)))^{-\frac{1}{q}} \|I_\rho f \chi_{B(x_0, 2r)}\|_{L_q(w^q, \mathbb{R}^n)} \\
&\lesssim \sup_{r>0} \varphi_2(x_0, r)^{-1} (w^q(B(x_0, r)))^{-\frac{1}{q}} \|f\|_{L_p(w^p, B(x_0, 2r))} \\
&\quad + \sup_{r>0} \varphi_2(x_0, r)^{-1} \int_r^1 \|f\|_{L_p(w^p, B(x_0, 2t))} \frac{\rho(t)}{t^{\frac{n}{p}+1}} dt \\
&\approx \sup_{r>0} \varphi_1(x_0, r)^{-1} (w^p(B(x_0, r)))^{-\frac{1}{p}} \|f\|_{L_p(w^p, B(x_0, r))} \\
&= \|f\|_{M_{p,\varphi_1}^{\{x_0\}}(w^p, \mathbb{R}^n)}.
\end{aligned}$$

Now let $p = 1$ then

$$\begin{aligned}
\|I_\rho f\|_{WM_{q,\varphi_2}^{\{x_0\}}(w^q, \mathbb{R}^n)} &= \sup_{r>0} \varphi_2(x_0, r)^{-1} (w^q(B(x_0, r)))^{-\frac{1}{q}} \|I_\rho f\|_{WL_q(w^q, B(x_0, 2r))} \\
&= \sup_{r>0} \varphi_2(x_0, r)^{-1} (w^q(B(x_0, r)))^{-\frac{1}{q}} \|I_\rho f \chi_{B(x_0, 2r)}\|_{WL_q(w^q, \mathbb{R}^n)} \\
&\lesssim \sup_{r>0} \varphi_2(x_0, r)^{-1} (w^q(B(x_0, r)))^{-\frac{1}{q}} \|f\|_{L_1(w, B(x_0, 2r))} \\
&\quad + \sup_{r>0} \varphi_2(x_0, r)^{-1} \int_r^1 \|f\|_{L_1(w, B(x_0, 2t))} \frac{\rho(t)}{t^{n+1}} dt \\
&\approx \sup_{r>0} \varphi_1(x_0, r)^{-1} (w(B(x_0, r)))^{-1} \|f\|_{L_1(w, B(x_0, r))} \\
&= \|f\|_{M_{1,\varphi_1}^{\{x_0\}}(w, \mathbb{R}^n)}.
\end{aligned}$$

Hence the proof is completed. \square

Corollary 3.4. *In the case $w \equiv 1$ from Theorem 3.3 we get Theorem C, in which we give Spanne-type result for generalized Riesz potential operator I_ρ on generalized local Morrey spaces $M_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ which was proved in [9] (Theorem 16, p.6).*

Corollary 3.5. *In the case $\rho(t) = t^\alpha, w \equiv 1, x \equiv x_0$ from Theorem 3.3 we get Spanne-type result for Riesz potential operator I_α on generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$ which was proved in [12] (Theorem 5.4, p.338).*

Corollary 3.6. *In the case $\rho(t) = t^\alpha$, $w \equiv 1$ and $\varphi(x_0, t) = t^{\frac{\lambda-n}{p}}$, $0 < \lambda < n$ from Theorem 3.3 we get Spanne result for Riesz potential operator I_α on local Morrey spaces $M_{p,\lambda}^{\{x_0\}}(\mathbb{R}^n)$ which is variant of Theorem A proved in [25].*

4 Adams-type result for the operator I_ρ in the spaces $M_{p,\varphi}(w)$

The following theorem is Adams-type estimate for generalized Riesz potential operator I_ρ on generalized weighted Morrey spaces $M_{p,\varphi}(w, \mathbb{R}^n)$.

Theorem 4.1. *Let $1 \leq p < \infty$, $q > p$, $w \in A_{1+\frac{q}{p}}$, $\rho(t)$ satisfy the conditions (3.1), (3.2) and (3.4). Let also $\varphi(x, t)$ satisfy the conditions*

$$c^{-1}\varphi(x, r) \leq \varphi(x, t) \leq c\varphi(x, r), \quad (4.1)$$

$$\int_r^\infty \frac{\operatorname{ess\,inf}_{t < s < \infty} (\varphi(x, s)w(B(x, s)))^{\frac{1}{p}}}{w(B(x, s))^{\frac{1}{q}}} \frac{\rho(t) dt}{t^n} \frac{1}{t} \leq C (\rho(r))^{-\frac{p}{q-p}}, \quad (4.2)$$

where C does not depend on $x \in \mathbb{R}^n$ and $r > 0$. Then the operator I_ρ is bounded from $M_{p,\varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)$ to $M_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)$ for $p > 1$ and from $M_{1,\varphi}(w, \mathbb{R}^n)$ to $WM_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)$ for $p = 1$. Moreover, for $p > 1$

$$\|I_\rho f\|_{M_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)} \lesssim \|f\|_{M_{p,\varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)},$$

and for $p = 1$

$$\|I_\rho f\|_{WM_{q,\varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)} \lesssim \|f\|_{M_{1,\varphi}(w, \mathbb{R}^n)}.$$

Proof. Let $1 < p < \infty$, $q > p$, $w \in A_{1+\frac{q}{p}}$ and $f \in M_{p,\varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)$. Write $f = f_1 + f_2$, where $B = B(x, r)$, $f_1 = f\chi_{2B}$ and $f_2 = f\chi_{\mathbb{R}^n \setminus 2B}$. Then we have

$$I_\rho f(x) = I_\rho f_1(x) + I_\rho f_2(x).$$

For $I_\rho f_1(y)$, $y \in B(x, r)$, following Hedberg's trick (see for instance [27], p. 354), we obtain

$$|I_\rho f_1(y)| \lesssim Mf(x)\rho(r),$$

(see [9], for more detail). Thus by taking $L_q(w, \mathbb{R}^n)$ -norm we get

$$\|I_\rho f_1\|_{L_q(B(x,r))(w, \mathbb{R}^n)} \leq w(B(x,r))^{\frac{1}{q}} \left(\int_{B(y,2r)} \left(\frac{\rho(|y-z|)}{|y-z|^n} |f(z)| \right)^q dz \right)^{1/q}.$$

For $I_\rho f_2(y)$, $y \in B(x, r)$ from (2.1) we have

$$\begin{aligned} |I_\rho f_2(y)| &\lesssim \int_{\mathbb{B}_{B(x,2r)}} \frac{\rho(|y-z|)}{|y-z|^n} |f(z)| dz \\ &\lesssim \int_{2r}^1 \|f \chi_{B(x,t)}\|_{L_p(w, \mathbb{R}^n)} w(B(x,r))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t}. \end{aligned} \quad (4.3)$$

Then from condition (4.2) and inequality (4.3) for all $y \in B(x, r)$ we get

$$\begin{aligned} |I_\rho f(y)| &\lesssim \rho(r) Mf(x) + \int_{2r}^1 \|f \chi_{B(x,t)}\|_{L_p(w, \mathbb{R}^n)} w(B(x,r))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t} \\ &\leq \rho(r) Mf(x) \\ &\quad + \|f\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)} \int_r^\infty \varphi(x, t)^{\frac{1}{p}} w(B(x, t))^{\frac{1}{p}} w(B(x, t))^{-\frac{1}{q}} \frac{\rho(t)}{t^n} \frac{dt}{t} \\ &\lesssim \rho(r) Mf(x) + \rho(r)^{-\frac{p}{q-p}} \|f\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)}. \end{aligned}$$

Hence choosing $\rho(r) = \left(\frac{\|f\|_{M_{p, \varphi^{1/p}}(w, \mathbb{R}^n)}}{Mf(x)} \right)^{\frac{q-p}{q}}$ for all $y \in B(x, r)$, we have

$$|I_\rho f(y)| \lesssim (Mf(x))^{\frac{p}{q}} \|f\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)}^{1-\frac{p}{q}}.$$

Consequently the statement of the theorem follows in view of the boundedness of the maximal operator M in $M_{p, \varphi^{\frac{1}{p}}}(\mathbb{R}^n)$ provided in [11] in virtue of condition (4.1), hence, for $1 < p < q < \infty$ we get

$$\begin{aligned} \|I_\rho f\|_{M_{q, \varphi^{\frac{1}{q}}}(w, \mathbb{R}^n)} &= \sup_{x \in \mathbb{R}^n, t > 0} \varphi(x, t)^{-\frac{1}{q}} w(B(x, t))^{-\frac{1}{q}} \|I_\rho f\|_{L_q(w, B(x, t))} \\ &\lesssim \|f\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)}^{1-\frac{p}{q}} \sup_{x \in \mathbb{R}^n, t > 0} \varphi(x, t)^{-\frac{1}{q}} w(B(x, t))^{-\frac{1}{q}} \|Mf\|_{L_p(w, B(x, t))}^{\frac{p}{q}} \\ &= \|f\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)}^{1-\frac{p}{q}} \left(\sup_{x \in \mathbb{R}^n, t > 0} \varphi(x, t)^{-\frac{1}{p}} (w(B(x, t)))^{-\frac{1}{p}} \|Mf\|_{L_p(w, B(x, t))} \right)^{\frac{p}{q}} \\ &= \|f\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)}^{1-\frac{p}{q}} \|Mf\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)}^{\frac{p}{q}} \\ &\lesssim \|f\|_{M_{p, \varphi^{\frac{1}{p}}}(w, \mathbb{R}^n)}, \end{aligned}$$

and for $1 < q < \infty$

$$\begin{aligned}
\|I_\rho f\|_{WM_{q,\varphi^{\frac{1}{q}}}(w,\mathbb{R}^n)} &= \sup_{x \in \mathbb{R}^n, t > 0} \varphi(x, t)^{-\frac{1}{q}} w(B(x, t))^{-\frac{1}{q}} \|I_\rho f\|_{WL_q(w, B(x, t))} \\
&\lesssim \|f\|_{M_{1,\varphi}(w,\mathbb{R}^n)}^{1-\frac{1}{q}} \sup_{x \in \mathbb{R}^n, t > 0} \varphi(x, t)^{-\frac{1}{q}} w(B(x, t))^{-\frac{1}{q}} \|Mf\|_{WL_1(w, B(x, t))}^{\frac{1}{q}} \\
&= \|f\|_{M_{1,\varphi}(w,\mathbb{R}^n)}^{1-\frac{1}{q}} \left(\sup_{x \in \mathbb{R}^n, t > 0} \varphi(x, t)^{-1} w(B(x, t))^{-1} \|Mf\|_{WL_1(w, B(x, t))} \right)^{\frac{1}{q}} \\
&= \|f\|_{M_{1,\varphi}(w,\mathbb{R}^n)}^{1-\frac{1}{q}} \|Mf\|_{M_{1,\varphi}(w,\mathbb{R}^n)}^{\frac{1}{q}} \\
&\lesssim \|f\|_{M_{1,\varphi}(w,\mathbb{R}^n)}.
\end{aligned}$$

Hence the proof is completed. \square

Corollary 4.2. *In the case $w \equiv 1$ from Theorem 4.1 we get Theorem D, in which we give Adams type result for generalized Riesz potential operator I_ρ on generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$ which was proved in [9] (Theorem 22, p.7).*

Corollary 4.3. *In the case $\rho(t) = t^\alpha$, $w \equiv 1$, $x \equiv x_0$ from Theorem 4.1 we get Adams type result for Riesz potential operator I_α on generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$ which was proved in [12] (Theorem 5.7, p.182).*

Corollary 4.4. *In the case $\rho(t) = t^\alpha$, $w \equiv 1$ and $\varphi(x_0, t) = t^{\frac{\lambda-n}{p}}$, $0 < \lambda < n$ from Theorem 4.1 we get Adams result for Riesz potential operator I_α on local Morrey spaces $M_{p,\lambda}^{\{x_0\}}(\mathbb{R}^n)$ which is variant of Theorem B proved in [25].*

Acknowledgement

The author would like to express his gratitude to the referees for their (his/her) very valuable comments.

Conflicts of Interest

The author declares that there are no conflicts of interest regarding the publication of this article.

References

- [1] D.R. Adams, *A note on Riesz potentials*, Duke Math., vol. **42**(4)(1975), 765-778.
- [2] M. Carro, L. Pick, J. Soria, V.D. Stepanov, *On embeddings between classical Lorentz spaces*, Math. Inequal. Appl. **4**(3)(2001), 397-428.
- [3] L. Cafarelli, *Elliptic second order equations*, Rend. Sem. Mat. Fis. Milano **58** (1998), (1990), 253-284. DOI 10.1007/BF02925245.
- [4] R. R. Coifman, C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, Tamkang J. Math., Studia Math. **51** (1974), 241-250.
- [5] A. Eridani, *On the boundedness of a generalized fractional integral on generalized Morrey spaces*, Tamkang J. Math., **33**(4)(2002), 335-340.
- [6] J. Garcia-Cuerva, J.L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Math. 16, Amsterdam, 1985.
- [7] I. Genebashvili, A. Gogatishvili, V. Kokilashvili, *Solution of two-weight problems for integral transforms with positive kernels*, Georgian Math. J., **3**(4)(1996), 31-342.
- [8] L. Grafakos, *Classical and Modern Fourier Analysis*, Pearson Education, Inc. Upper Saddle River, New Jersey, 2004.
- [9] V.S. Guliyev, A.F. Ismayilova, A. Kucukaslan, A. Serbetci, *Generalized fractional integral operators on generalized local Morrey spaces*, J. Funct. Spaces, Article ID 594323, (2015), 8 pages.
- [10] V.S. Guliyev, *Integral operators on function spaces on the homogeneous groups and on domains in \mathbb{R}^n* . [in Russian], Diss. Steklov Mat. Inst., (1994), Moscow.
- [11] V.S. Guliyev, *Generalized weighted Morrey spaces and higher order commutators of sublinear operators*, Eurasian Math. J., **3**(3)(2012), 33-61.
- [12] V.S. Guliyev, S.S. Aliyev, T. Karaman, and P. S. Shukurov, *Boundedness of sublinear operators and commutators on generalized Morrey Space*, Integral Equat. Oper. Theor., **71**(3)(2011), 327-355.

- [13] T.Y. Komori, S. Shirai, *Weighted Morrey spaces and a singular integral operator*, Math. Nachr. **282**(2)(2009), 219-231.
- [14] A. Kucukaslan, S.G. Hasanov, C. Aykol, *Generalized fractional integral operators on vanishing generalized local Morrey spaces*, Int. J. of Math. Anal., **11**(6)(2017) 277–291.
- [15] A. Kucukaslan, V.S. Guliyev, A. Serbetci, *Generalized fractional maximal operators on generalized local Morrey spaces*, Commun. Fac. Sci. Univ. Ank. Ser. A1. Math. Stat., **69**(1)(2020), pp. 73-87. DOI: 10.31801/cfsuasmas.
- [16] A. Kucukaslan, *Equivalence of norms of the generalized fractional fractional integral operator and the generalized fractional maximal operator on generalized weighted Morrey spaces*, Ann. Funct. Anal. **11**(2020) pp. 1007-1026. <https://doi.org/10.1007/s43034-020-00066-w>
- [17] A. Kucukaslan, *Two-type estimates for the boundedness of generalized fractional maximal operators on generalized weighted local Morrey spaces*, Turk. J. Math. Comput. Sci., **12**(1)(2020), pp. 57-66.
- [18] C.B. Morrey, *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc., **43** (1938), 126-166.
- [19] B. Muckenhoupt, R. Wheeden, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. , **165** (1972), 261-274.
- [20] B. Muckenhoupt, *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc. , **192** (1974), 207-226.
- [21] R. Mustafayev and A. Kucukaslan *An extension of Muckenhoupt-Wheeden theorem to generalized weighted Morrey spaces*, Georgian Math. Journal, DOI: <https://doi.org/10.1515/gmj-2020-2056>.
- [22] E. Nakai, *On generalized fractional integrals*, Taiwanese J. Math., **5** (2001), 587-602.
- [23] E. Nakai, *Generalized fractional integrals on generalized Morrey spaces*, Math. Nachr., **287**(2-3) (2014), 339–351.

- [24] S. Nakamura, *Generalized weighted Morrey spaces and classical operators*, Math. Nachr. **289**(1718)(2016). DOI:10.1002/mana.201500260.
- [25] J. Peetre, *On the theory of $M_{p,\lambda}$* , J. Funct. Anal., **4**(1969), 71-87.
- [26] A. Ruiz and L. Vega, *Unique continuation for Schrödinger operators with potentials in the Morrey class*, Publ. Math. **35**(2)(1991) 291-298, Conference of Mathematical Analysis (El Escorial, 1989).
- [27] E.M. Stein, *Harmonic Analysis: Real Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton Univ. Press, Princeton NJ (1993).