

WEIGHTED ESTIMATES FOR GENERALIZED RIESZ POTENTIALS

PABLO ROCHA

ABSTRACT. For $0 \leq \alpha < n$, $0 < p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, we obtain the boundedness from $H_w^p(\mathbb{R}^n)$ into $L_{w^{q/p}}^q(\mathbb{R}^n)$ of certain generalized Riesz potentials for certain weights w . This result is achieved via the infinite atomic decomposition developed in [16].

1. INTRODUCTION

Given $0 \leq \alpha < n$ and $m \in \mathbb{N} \cap (1 - \frac{\alpha}{n}, +\infty)$, we consider the operator $T_{\alpha,m}$ defined by

$$(1) \quad T_{\alpha,m}f(x) = \int_{\mathbb{R}^n} |x - A_1 y|^{-\alpha_1} \cdots |x - A_m y|^{-\alpha_m} f(y) dy, \quad (x \in \mathbb{R}^n),$$

where the α_j 's are positive constants such that $\alpha_1 + \cdots + \alpha_m = n - \alpha$ and the A_j 's are $n \times n$ invertible matrices. For the case $\alpha = 0$, we require that the matrices A_j satisfy the following additional condition: $A_i - A_j$ are invertible for $i \neq j$, $1 \leq i, j \leq m$.

The germ of this operator appears in [11], there F. Ricci and P. Sjögren obtained the boundedness on $L^p(\mathbb{H}_1)$, $1 < p \leq +\infty$, for a family of maximal operators on the three dimensional Heisenberg group \mathbb{H}_1 . To get this result, they studied the $L^2(\mathbb{R})$ boundedness of the operator

$$Tf(x) = \int_{\mathbb{R}} |x - y|^{\alpha-1} |(\beta - 1)x - \beta y|^{-\alpha} f(y) dy,$$

for $\beta \neq 0, 1$ and $0 < \alpha < 1$. A generalization of this operator on \mathbb{R}^n was studied by T. Godoy and M. Urciuolo in [3].

If $0 < \alpha < n$ and $m \geq 1$, then the operator $T_{\alpha,m}$ has the same behavior that the Riesz potential on $L^p(\mathbb{R}^n)$. Indeed

$$(2) \quad |T_{\alpha,m}f(x)| \leq C \sum_{j=1}^m \int_{\mathbb{R}^n} |A_j^{-1}x - y|^{\alpha-n} |f(y)| dy = C \sum_{j=1}^m I_{\alpha}(|f|)(A_j^{-1}x),$$

for all $x \in \mathbb{R}^n$, this pointwise inequality implies that $T_{\alpha,m}$ is a bounded operator from $L^p(\mathbb{R}^n)$ into $L^q(\mathbb{R}^n)$ for $1 < p < \frac{n}{\alpha}$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, and it is of type weak $(1, n/n - \alpha)$. We observe that if $A_j = id$ for all $j = 1, \dots, m$, then $T_{\alpha,m}$ is the Riesz potential I_{α} . Thus, for $0 < \alpha < n$ and $m \geq 1$, the operator $T_{\alpha,m}$ is a kind of generalization of the Riesz potential.

It is well known that the Riesz potential I_{α} is bounded from $H^p(\mathbb{R}^n)$ into $H^q(\mathbb{R}^n)$ for $0 < p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ (see [23], [6]). In [18], the author jointly with M. Urciuolo proved the $H^p(\mathbb{R}^n) - L^q(\mathbb{R}^n)$ boundedness of the operator $T_{\alpha,m}$ and we also showed that the $H^p(\mathbb{R}) - H^q(\mathbb{R})$ boundedness does not hold for $0 < p \leq \frac{1}{1+\alpha}$,

Key words and phrases: Weighted Hardy Spaces, Atomic Decomposition, Fractional Operators.

2.010 Math. Subject Classification: 42B25, 42B30.

$\frac{1}{q} = \frac{1}{p} - \alpha$ and $T_{\alpha,m}$ with $0 \leq \alpha < 1$, $m = 2$, $A_1 = 1$, and $A_2 = -1$. This is a significant difference with respect to the case $0 < \alpha < 1$, $n = m = 1$, and $A_1 = 1$.

For the case $\alpha = 0$ and $m \geq 2$, the operator $T_{0,m}$ was studied under the assumption that $A_i - A_j$ are invertible if $i \neq j$.

The behavior of this class of operators and generalizations of them on the spaces of functions $L^p(\mathbb{R}^n)$, $L_w^p(\mathbb{R}^n)$, $H^p(\mathbb{R}^n)$ was studied in [4], [12], [17], [18], and [14]. Their behavior on variable Hardy spaces $H^{p(\cdot)}(\mathbb{R}^n)$ and on variable Hardy-Morrey spaces $\mathcal{HM}_{p(\cdot),u}(\mathbb{R}^n)$ can be found in [19], [20], [15] and [24].

To obtain the boundedness of operators, like singular integrals or fractional type operators, in weighted Hardy spaces $H_w^p(\mathbb{R}^n)$, one can appeal to the atomic or molecular characterization of $H_w^p(\mathbb{R}^n)$, which means that a distribution in H_w^p can be represented as a sum of atoms or molecules; see [2], [22], [8], [7], and [16]. Then the boundedness of linear operators in H_w^p can be deduced, in principle, from their behavior on atoms. However, it must be mentioned that in the classical Hardy spaces, i.e.: $w \equiv 1$, M. Bownik in [1] has shown, by means of an example of Y. Meyer, that it not suffice to check that an operator from a Hardy spaces H^p , $0 < p \leq 1$, into some quasi Banach space X , maps atoms into bounded elements of X to establish that this operator extends to a bounded operator on H^p . This example is, in a certain sense, pathological. Now, if our operator T under study is bounded on L^s for some $1 < s < +\infty$, then the uniform boundedness of T on atoms in L^p norm (or H^p norm) implies that the operator extends to a bounded operator from H^p into L^p (or on H^p), this follows from the fact that one always can take an atomic decomposition which converges in the norm of L^s , see [25] and [13]. In [16] the author proved that if $w \in \mathcal{A}_\infty$, $0 < p \leq 1$ and $f \in \widehat{\mathcal{D}}_0$, then there exist a sequence of $w - (p, p_0, d)$ atoms $\{a_j\}$ and a sequence of scalars $\{\lambda_j\}$ with $\sum_j |\lambda_j|^p \leq c \|f\|_{H_w^p}^p$ such that $f = \sum_j \lambda_j a_j$, where the convergence is in $L^s(\mathbb{R}^n)$ for each $1 < s < \infty$ (see Theorem 13 below). This result allows to avoid any problem that could arise with respect to establish the boundedness of classical operators on $H_w^p(\mathbb{R}^n)$ (see [16], Theorems 2.10 and 3.4).

In this work, for certain weights w and by means of the atomic decomposition developed in [16], we will prove $H_w^p(\mathbb{R}^n) - L_{w^{q/p}}^q(\mathbb{R}^n)$ estimates for the operator $T_{\alpha,m}$ given by (1), where $0 < p \leq 1$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$.

To get our main results we will also consider the following condition for the weights: there exists $C > 0$ such that

$$(3) \quad w(A_j x) \leq C w(x), \quad \text{a.e. } x \in \mathbb{R}^n, \quad j = 1, \dots, m.$$

Next we give two nontrivial examples of weights that satisfy the condition (3). Observe that also power weights satisfy this condition.

Example 1. Let $w(x) = \begin{cases} \log(\frac{1}{|x|}) & |x| < \frac{1}{e} \\ 1 & |x| \geq \frac{1}{e} \end{cases}$. Then $w \in \mathcal{A}_1$ and satisfies (3).

Example 2. The nonnegative function $w(x) = |x|^a$, $x \in \mathbb{R}^n \setminus \{0\}$, is a weight in the Muckenhoupt class \mathcal{A}_p , $1 < p < \infty$, if and only if $-n < a < n(p-1)$ (see p. 506 in [5]). Such a weight satisfies (3).

In Section 2 we present the basics of weighted theory and briefly recall the atomic decomposition developed in [16]. In Section 3 appears our main results. We study the cases $\alpha = 0$ and $0 < \alpha < n$ separately. More precisely, we obtain:

Case $\alpha = 0$. If $w \in \mathcal{A}_\infty$ and satisfies (3), then $T_{0,m}$ can be extended to an $H_w^p(\mathbb{R}^n) - L_w^p(\mathbb{R}^n)$ bounded operator for each $0 < p \leq 1$ (see Theorem 17 below).

Case $0 < \alpha < n$. If $w^{\frac{n}{(n-\alpha)s}} \in \mathcal{A}_1$ with $0 < s < 1$, $\frac{r_w}{r_w-1} < \frac{n}{\alpha}$ and w satisfies (3), then $T_{\alpha,m}$ can be extended to an $H_{w^p}^p(\mathbb{R}^n)$ - $L_{w^q}^q(\mathbb{R}^n)$ bounded operator for each $s \leq p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ (see Theorem 18 below).

Notation: The symbol $A \lesssim B$ stands for the inequality $A \leq cB$ for some constant c . We denote by $B(x_0, r)$ the ball centered at $x_0 \in \mathbb{R}^n$ of radius r . For a measurable subset $E \subset \mathbb{R}^n$ we denote $|E|$ and χ_E the Lebesgue measure of E and the characteristic function of E respectively. Given a real number $s \geq 0$, we write $[s]$ for the integer part of s . As usual we denote with $\mathcal{S}(\mathbb{R}^n)$ the space of smooth and rapidly decreasing functions, with $\mathcal{S}'(\mathbb{R}^n)$ the dual space. If β is the multiindex $\beta = (\beta_1, \dots, \beta_n)$, then $|\beta| = \beta_1 + \dots + \beta_n$.

Throughout this paper, C will denote a positive constant, not necessarily the same at each occurrence.

2. PRELIMINARIES

A weight is a non-negative locally integrable function on \mathbb{R}^n that takes values in $(0, \infty)$ almost everywhere, i.e. : the weights are allowed to be zero or infinity only on a set of Lebesgue measure zero.

Given a weight w and $0 < p < \infty$, we denote by $L_w^p(\mathbb{R}^n)$ the spaces of all functions f satisfying $\|f\|_{L_w^p}^p := \int_{\mathbb{R}^n} |f(x)|^p w(x) dx < \infty$. When $p = \infty$, we have that $L_w^\infty(\mathbb{R}^n) = L^\infty(\mathbb{R}^n)$ with $\|f\|_{L_w^\infty} = \|f\|_{L^\infty}$. If E is a measurable set, we use the notation $w(E) = \int_E w(x) dx$.

Let f be a locally integrable function on \mathbb{R}^n . The function

$$M(f)(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy,$$

where the supremum is taken over all balls B containing x , is called the uncentered Hardy-Littlewood maximal function of f .

We say that a weight $w \in \mathcal{A}_1$ if there exists $C > 0$ such that

$$M(w)(x) \leq Cw(x), \quad a.e. x \in \mathbb{R}^n,$$

the best possible constant is denoted by $[w]_{\mathcal{A}_1}$. Equivalently, a weight $w \in \mathcal{A}_1$ if there exists $C > 0$ such that for every ball B

$$(4) \quad \frac{1}{|B|} \int_B w(x) dx \leq C \operatorname{ess\,inf}_{x \in B} w(x).$$

Remark 3. If $w \in \mathcal{A}_1$ and $0 < r < 1$, then by Hölder inequality we have that $w^r \in \mathcal{A}_1$.

For $1 < p < \infty$, we say that a weight $w \in \mathcal{A}_p$ if there exists $C > 0$ such that for every ball B

$$\left(\frac{1}{|B|} \int_B w(x) dx \right) \left(\frac{1}{|B|} \int_B [w(x)]^{-\frac{1}{p-1}} dx \right)^{p-1} \leq C.$$

It is well known that $\mathcal{A}_{p_1} \subset \mathcal{A}_{p_2}$ for all $1 \leq p_1 < p_2 < \infty$. Also, if $w \in \mathcal{A}_p$ with $1 < p < \infty$, then there exists $1 < q < p$ such that $w \in \mathcal{A}_q$. We denote by $\tilde{q}_w = \inf\{q > 1 : w \in \mathcal{A}_q\}$ the critical index of w .

Lemma 4. (Proposition 7.1.5 in [5]) If $w \in \mathcal{A}_p$ for some $1 \leq p < \infty$, then the measure $w(x)dx$ is doubling: precisely, for all $\lambda > 1$ and all ball B we have

$$w(\lambda B) \leq \lambda^{np} [w]_{\mathcal{A}_p} w(B),$$

where λB denotes the ball with the same center as B and radius λ times the radius of B .

Lemma 5. *If $w \in \mathcal{A}_\infty$ and A is a $n \times n$ invertible matrix such that $w(Ax) \lesssim w(x)$ a.e. $x \in \mathbb{R}^n$, then for each $M > 0$ there exists a constant $C > 0$ such that for every ball $B(x_0, r)$ holds*

$$w(B(Ax_0, 2Mr)) \leq Cw(B(x_0, r)).$$

Proof. Applying a change of variable and Lemma 4, the lemma follows. \square

Theorem 6. (Theorem 9 in [9]) *Let $1 < p < \infty$. Then*

$$\int_{\mathbb{R}^n} [Mf(x)]^p w(x) dx \leq C_{w,p,n} \int_{\mathbb{R}^n} |f(x)|^p w(x) dx,$$

for all $f \in L_w^p(\mathbb{R}^n)$ if and only if $w \in \mathcal{A}_p$.

Given $1 < p \leq q < \infty$, we say that a weight $w \in \mathcal{A}_{p,q}$ if there exists $C > 0$ such that for every ball B

$$\left(\frac{1}{|B|} \int_B [w(x)]^q dx \right)^{1/q} \left(\frac{1}{|B|} \int_B [w(x)]^{-p'} dx \right)^{1/p'} \leq C < \infty.$$

For $p = 1$, we say that a weight $w \in \mathcal{A}_{1,q}$ if there exists $C > 0$ such that for every ball B

$$\left(\frac{1}{|B|} \int_B [w(x)]^q dx \right)^{1/q} \leq C \operatorname{ess\,inf}_{x \in B} w(x).$$

When $p = q$, this definition is equivalent to $w^p \in \mathcal{A}_p$.

Remark 7. *From the inequality in (4) it follows that if a weight $w \in \mathcal{A}_1$, then $0 < \operatorname{ess\,inf}_{x \in B} w(x) < \infty$ for each ball B . Thus $w \in \mathcal{A}_1$ implies that $w^{\frac{1}{q}} \in \mathcal{A}_{p,q}$, for each $1 \leq p \leq q < \infty$.*

Given $0 < \alpha < n$, we define the fractional maximal operator M_α by

$$M_\alpha f(x) = \sup_{B \ni x} \frac{1}{|B|^{1-\frac{\alpha}{n}}} \int_B |f(y)| dy,$$

where f is a locally integrable function and the supremum is taken over all balls B containing x

The fractional maximal operators satisfies the following weighted inequality.

Theorem 8. (Theorem 3 in [10]) *If $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ and $w \in \mathcal{A}_{p,q}$, then*

$$\left(\int_{\mathbb{R}^n} [M_\alpha f(x)]^q w^q(x) dx \right)^{1/q} \leq C \left(\int_{\mathbb{R}^n} |f(x)|^p w^p(x) dx \right)^{1/p},$$

for all $f \in L_{w^p}^p(\mathbb{R}^n)$.

A weight w satisfies the reverse Hölder inequality with exponent $s > 1$, denoted by $w \in RH_s$, if there exists $C > 0$ such that for every ball B ,

$$\left(\frac{1}{|B|} \int_B [w(x)]^s dx \right)^{\frac{1}{s}} \leq C \frac{1}{|B|} \int_B w(x) dx;$$

the best possible constant is denoted by $[w]_{RH_s}$. We observe that if $w \in RH_s$, then by Hölder's inequality, $w \in RH_t$ for all $1 < t < s$, and $[w]_{RH_t} \leq [w]_{RH_s}$. Moreover, if $w \in RH_s$, $s > 1$, then $w \in RH_{s+\epsilon}$ for some $\epsilon > 0$. We denote by $r_w = \sup\{r > 1 : w \in RH_r\}$ the critical index of w for the reverse Hölder condition.

It is well known that a weight w satisfies the condition \mathcal{A}_∞ if and only if $w \in \mathcal{A}_p$ for some $p \geq 1$ (see corollary 7.3.4 in [5]). So $\mathcal{A}_\infty = \cup_{1 \leq p < \infty} \mathcal{A}_p$. Also, $w \in \mathcal{A}_\infty$ if and only if $w \in RH_s$ for some $s > 1$ (see Theorem 7.3.3 in [5]). Thus $1 < r_w \leq +\infty$ for all $w \in \mathcal{A}_\infty$.

Other remarkable result about the reverse Hölder classes was discovered by Stromberg and Wheeden, they proved in [21] that $w \in RH_s$, $1 < s < +\infty$, if and only if $w^s \in \mathcal{A}_\infty$.

Given a weight w , $0 < p < \infty$ and a measurable set E we set $w^p(E) = \int_E [w(x)]^p dx$. The following result is a immediate consequence of the reverse Hölder condition.

Lemma 9. *For $0 < \alpha < n$, let $0 < p < \frac{n}{\alpha}$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. If $w^p \in RH_{\frac{q}{p}}$, then*

$$[w^p(B)]^{-\frac{1}{p}} [w^q(B)]^{\frac{1}{q}} \leq [w^p]_{RH_{q/p}}^{1/p} |B|^{-\frac{\alpha}{n}},$$

for each ball B in \mathbb{R}^n .

Lemma 10. *(Lemma 4.6 in [16]) Let $0 < p < 1$. If $w^{1/p} \in \mathcal{A}_1$, then $p \cdot r_{w^p} \leq r_w \leq r_{w^p}$.*

Lemma 11. *(Lemma 4.7 in [16]) Let $0 < p < q$. If $w^q \in \mathcal{A}_1$, then $p \cdot r_{w^p} \leq q \cdot r_{w^q}$.*

We conclude these preliminaries with weighted Hardy theory. Topologize $\mathcal{S}(\mathbb{R}^n)$ by the collection of semi-norms $\|\cdot\|_{\alpha,\beta}$, with α and β multi-indices, given by

$$\|\varphi\|_{\alpha,\beta} = \sup_{x \in \mathbb{R}^n} |x^\alpha \partial^\beta \varphi(x)|.$$

For each $N \in \mathbb{N}$, we set $\mathcal{S}_N = \{\varphi \in \mathcal{S}(\mathbb{R}^n) : \|\varphi\|_{\alpha,\beta} \leq 1, |\alpha|, |\beta| \leq N\}$. Let $f \in \mathcal{S}'(\mathbb{R}^n)$, we denote by \mathcal{M}_N the grand maximal operator given by

$$\mathcal{M}_N f(x) = \sup_{t>0} \sup_{\varphi \in \mathcal{S}_N} |(t^{-n} \varphi(t^{-1} \cdot) * f)(x)|.$$

Given a weight $w \in \mathcal{A}_\infty$ and $p > 0$, the weighted Hardy space $H_w^p(\mathbb{R}^n)$ consists of all tempered distributions f such that

$$\|f\|_{H_w^p(\mathbb{R}^n)} = \|\mathcal{M}_N f\|_{L_w^p(\mathbb{R}^n)} = \left(\int_{\mathbb{R}^n} [\mathcal{M}_N f(x)]^p w(x) dx \right)^{1/p} < \infty.$$

Let $\phi \in \mathcal{S}(\mathbb{R}^n)$ be a function such that $\int \phi(x) dx \neq 0$. For $f \in \mathcal{S}'(\mathbb{R}^n)$, we define the maximal function $M_\phi f$ by

$$M_\phi f(x) = \sup_{t>0} |(t^{-n} \phi(t^{-1} \cdot) * f)(x)|.$$

For N sufficiently large, we have $\|M_\phi f\|_{L_w^p} \simeq \|\mathcal{M}_N f\|_{L_w^p}$, (see [22]). The following set

$$\widehat{\mathcal{D}}_0 = \{\phi \in \mathcal{S}(\mathbb{R}^n) : \widehat{\phi} \in C_c^\infty(\mathbb{R}^n), \text{ and } \text{supp}(\widehat{\phi}) \subset \mathbb{R}^n \setminus B(0, \delta) \text{ for some } \delta > 0\}$$

is dense in $H_w^p(\mathbb{R}^n)$, $0 < p < \infty$, if w is a doubling weight on \mathbb{R}^n (see [22], Theorem 1, pp. 103).

We adopt the weighted atomic decomposition developed in [16]. Let $w \in \mathcal{A}_\infty$ with critical index \tilde{q}_w and critical index r_w for reverse Hölder condition. Let $0 < p \leq 1$, $\max\{1, p(\frac{r_w}{r_w-1})\} < p_0 \leq +\infty$, and $d \in \mathbb{Z}$ such that $d \geq \lfloor n(\frac{\tilde{q}_w}{p} - 1) \rfloor$, a function $a(\cdot)$ is a $w - (p, p_0, d)$ atom centered in $x_0 \in \mathbb{R}^n$ if

$$(a1) \ a \in L^{p_0}(\mathbb{R}^n) \text{ with support in the ball } B = B(x_0, r).$$

$$(a2) \ \|a\|_{L^{p_0}(\mathbb{R}^n)} \leq |B|^{\frac{1}{p_0}} w(B)^{-\frac{1}{p}}.$$

$$(a3) \ \int x^\beta a(x) dx = 0 \text{ for all multi-index } \beta \text{ such that } |\beta| \leq d.$$

Remark 12. *The condition $\max\{1, p(\frac{r_w}{r_w-1})\} < p_0 < +\infty$ implies that $w \in RH_{(\frac{p_0}{p})}$. If $r_w = +\infty$, then $w \in RH_t$ for each $1 < t < +\infty$. So, if $r_w = +\infty$, it considers $\frac{r_w}{r_w-1} = 1$.*

The following theorem is crucial to get the main results.

Theorem 13. (Theorem 9 in [16]) *Let $f \in \hat{\mathcal{D}}_0$, and $0 < p \leq 1$. If $w \in \mathcal{A}_\infty$, then there exist a sequence of $w - (p, p_0, d)$ atoms $\{a_j\}$ and a sequence of scalars $\{\lambda_j\}$ with $\sum_j |\lambda_j|^p \leq c \|f\|_{H_w^p}^p$ such that $f = \sum_j \lambda_j a_j$, where the convergence is both in $L^s(\mathbb{R}^n)$ and pointwise for each $1 < s < \infty$.*

3. MAIN RESULTS

In the sequel, given a ball $B = B(x_0, r)$ and A_1, \dots, A_m $n \times n$ invertible matrices, we put $M = \max\{\|A_j\| : 1 \leq j \leq m\}$, $B_i^* = B(A_i x_0, 2Mr)$ for $i = 1, \dots, m$ and decompose $\mathbb{R}^n = (\bigcup_{i=1}^m B_i^*) \cup R$, where $R = \mathbb{R}^n \setminus (\bigcup_{i=1}^m B_i^*)$. Moreover, $R = \bigcup_{k=1}^m R_k$, where

$$R_k = \{x \in R : |x - A_k x_0| \leq |x - A_i x_0| \text{ for all } i \neq k\},$$

for $k = 1, \dots, m$.

Lemma 14. *For $0 \leq \alpha < n$ and $m \in \mathbb{N} \cap (1 - \frac{\alpha}{n}, +\infty)$, let $T_{\alpha, m}$ be the operator defined by (1), where the A_j 's are $n \times n$ invertible matrices. If $w \in \mathcal{A}_\infty$ and $a(\cdot)$ is a $w - (p, p_0, d)$ atom supported on the ball $B = B(x_0, r)$, then*

$$|T_{\alpha, m} a(x)| \lesssim [w(B)]^{-1/p} \sum_{k=1}^m \chi_{R_k}(x) \left(M_{\frac{\alpha}{n+d+1}}(\chi_B)(A_k^{-1}x) \right)^{\frac{n+d+1}{n}}, \quad \text{if } x \in R.$$

Proof. We denote $k(x, y) = |x - A_1 y|^{-\alpha_1} \cdots |x - A_m y|^{-\alpha_m}$. In view of the moment condition of $a(\cdot)$ we have

$$T_{\alpha, m} a(x) = \int_{B(x_0, r)} k(x, y) a(y) dy = \int_{B(x_0, r)} (k(x, y) - q_d(x, y)) a(y) dy, \quad (x \in R),$$

where q_d is the degree d Taylor polynomial of the function $y \rightarrow k(x, y)$ expanded around x_0 . By the standard estimate of the remainder term of the Taylor expansion, there exists ξ between y and x_0 such that

$$\begin{aligned} |k(x, y) - q_d(x, y)| &\lesssim |y - x_0|^{d+1} \sum_{k_1 + \dots + k_n = d+1} \left| \frac{\partial^{d+1}}{\partial y_1^{k_1} \dots \partial y_n^{k_n}} k(x, \xi) \right| \\ &\leq C |y - x_0|^{d+1} \left(\prod_{i=1}^m |x - A_i \xi|^{-\alpha_i} \right) \left(\sum_{l=1}^m |x - A_l \xi|^{-1} \right)^{d+1}. \end{aligned}$$

Now, we decompose $R = \bigcup_{k=1}^m R_k$ where

$$R_k = \{x \in R : |x - A_k x_0| \leq |x - A_i x_0| \text{ for all } i \neq k\}.$$

If $x \in R$ then $|x - A_i x_0| \geq 2Mr$, since $\xi \in B$ it follows that $|A_i x_0 - A_i \xi| \leq Mr \leq \frac{1}{2}|x - A_i x_0|$ so

$$|x - A_i \xi| = |x - A_i x_0 + A_i x_0 - A_i \xi| \geq |x - A_i x_0| - |A_i x_0 - A_i \xi| \geq \frac{1}{2}|x - A_i x_0|.$$

If $x \in R$, then $x \in R_k$ for some k and since $\alpha_1 + \dots + \alpha_m = n - \alpha$ we obtain

$$\begin{aligned} |k(x, y) - q_d(x, y)| &\leq C |y - x_0|^{d+1} \left(\prod_{i=1}^m |x - A_i x_0|^{-\alpha_i} \right) \left(\sum_{l=1}^m |x - A_l x_0|^{-1} \right)^{d+1} \\ &\leq C r^{d+1} |x - A_k x_0|^{-n+\alpha-d-1}, \end{aligned}$$

this inequality allows us to conclude that

$$\begin{aligned} |T_{\alpha,m}a(x)| &\leq C \|a\|_1 r^{d+1} |x - A_k x_0|^{-n+\alpha-d-1} \\ &\leq C |B|^{1-\frac{1}{p_0}} \|a\|_{p_0} r^{d+1} |x - A_k x_0|^{-n+\alpha-d-1}, \end{aligned}$$

since $\|a\|_{p_0} \leq |B|^{1/p_0} [w(B)]^{-1/p}$, we have

$$\begin{aligned} (5) \quad |T_{\alpha,m}a(x)| &\lesssim \frac{r^{n+d+1} |x - A_k x_0|^{-n+\alpha-d-1}}{[w(B)]^{1/p}} \\ &\lesssim \frac{\left(M_{\frac{\alpha n}{n+d+1}}(\chi_B)(A_k^{-1}x)\right)^{\frac{n+d+1}{n}}}{[w(B)]^{1/p}}, \quad \text{if } x \in R_k. \end{aligned}$$

This completes the proof. \square

Remark 15. In Lemma 14, we observe that the matrices A_j can be equal to each other.

We recall the definition of the critical indices for a weight w .

Definition 16. Given a weight w , we denote by $\tilde{q}_w = \inf\{q > 1 : w \in \mathcal{A}_q\}$ the critical index of w , and we denote by $r_w = \sup\{r > 1 : w \in RH_r\}$ the critical index of w for the reverse Hölder condition.

Theorem 17. For $\alpha = 0$ and $m \geq 2$, let $T_{0,m}$ be the operator defined by (1) where the A_j 's are $n \times n$ invertible matrices such that $A_i - A_j$ are invertible for $i \neq j$, $1 \leq i, j \leq m$. If $w \in \mathcal{A}_\infty$ and $w(A_j x) \lesssim w(x)$ a.e. $x \in \mathbb{R}^n$, $1 \leq j \leq m$, then $T_{0,m}$ can be extended to an $H_w^p(\mathbb{R}^n)$ - $L_w^p(\mathbb{R}^n)$ bounded operator for each $0 < p \leq 1$.

Proof. Theorem 6 in [20] asserts that the operator $T_{0,m}$ is bounded on $L^{p_0}(\mathbb{R}^n)$ for each $1 < p_0 < +\infty$. Given $0 < p \leq 1$, we take $d = \lfloor n(\frac{\tilde{q}_w}{p} - 1) \rfloor$ and $p_0 \in (1, +\infty)$ such that $\frac{r_w}{r_w-1} < p_0$. Thus, by Theorem 2.10 in [16], it suffices to show that the operator $T_{0,m}$ is uniformly bounded in L_w^p norm on all w -(p, p_0, d) atoms $a(\cdot)$.

Given a w -(p, p_0, d) atom $a(\cdot)$ supported on the ball $B = B(x_0, r)$, let $B_i^* = B(A_i x_0, 2Mr)$. We write

$$\begin{aligned} \int_{\mathbb{R}^n} |T_{0,m}a(x)|^p w(x) dx &= \int_{\bigcup_{i=1}^m B_i^*} |T_{0,m}a(x)|^p w(x) dx \\ &+ \int_{\mathbb{R}^n \setminus (\bigcup_{i=1}^m B_i^*)} |T_{0,m}a(x)|^p w(x) dx = I_1 + I_2 \end{aligned}$$

To estimate the term I_1 , we observe that $p \frac{r_w}{r_w-1} < p_0$ thus $w \in RH_{(\frac{p_0}{p})'}$ (see Remark 12), then Hölder's inequality applied with $\frac{p_0}{p}$, the condition (a2) of the atom $a(\cdot)$, and Lemma 5 give

$$\begin{aligned} I_1 &\lesssim \|T_{0,m}a\|_{p_0}^p \sum_{i=1}^m \left(\int_{B_i^*} [w(x)]^{(p_0/p)'} \right)^{1/(p_0/p)'} \\ &\lesssim \|a\|_{p_0}^p \sum_{i=1}^m |B_i^*|^{-p/p_0} w(B_i^*) \leq C. \end{aligned}$$

Taking account that $w(A_j x) \lesssim w(x)$ a.e. $x \in \mathbb{R}^n$, $1 \leq j \leq m$, and $w \in \mathcal{A}_{p \frac{n+d+1}{n}}$ ($p \frac{n+d+1}{n} > \tilde{q}_w$), from Lemma 14 with $\alpha = 0$ and Theorem 6, we get

$$I_2 \lesssim w(B)^{-1} \sum_{k=1}^m \int_{\mathbb{R}^n} [M(\chi_B)(x)]^{p \frac{n+d+1}{n}} w(A_k x) dx \leq C.$$

Thus, $T_{0,m}$ is uniformly bounded in L_w^p norm on every w -(p, p_0, d) atom $a(\cdot)$. \square

Theorem 18. For $0 < \alpha < n$ and $m \geq 1$, let $T_{\alpha,m}$ be the operator defined by (1) where the A_j 's are $n \times n$ invertible matrices. If $w^{\frac{n}{(n-\alpha)s}} \in \mathcal{A}_1$ with $0 < s < 1$, $\frac{r_w}{r_w-1} < \frac{n}{\alpha}$ and $w(A_j x) \lesssim w(x)$ a.e. $x \in \mathbb{R}^n$, $1 \leq j \leq m$, then $T_{\alpha,m}$ can be extended to an $H_{w^p}^p(\mathbb{R}^n)$ - $L_{w^q}^q(\mathbb{R}^n)$ bounded operator for each $s \leq p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$.

Proof. For $0 < p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, we have that $p < q \leq \frac{n}{n-\alpha}$. Now, the condition $w^{n/(n-\alpha)s} \in \mathcal{A}_1$, $0 < s < 1 < \frac{n}{n-\alpha}$, implies that w , $w^{1/p}$, w^p and w^q belong to \mathcal{A}_1 , for all $s \leq p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$.

We also have that $\tilde{q}_{w^p} = 1$, since $w^p \in \mathcal{A}_1$. We choose $d = \lfloor n(\frac{1}{p} - 1) \rfloor$ and take p_0 such that $\frac{r_w}{r_w-1} < p_0 < \frac{n}{\alpha}$, from Lemma 10, we have that $p \frac{r_{w^p}}{r_{w^p}-1} \leq \frac{r_w}{r_w-1} < p_0$ so $\max\{1, p(\frac{r_{w^p}}{r_{w^p}-1})\} < p_0$ (this last inequality is required in the definition of w^p -atom). By Theorem 13, given $f \in \widehat{\mathcal{D}}_0$ we can write $f = \sum \lambda_j a_j$ where the a_j 's are $w^p - (p, p_0, d)$ atoms, the scalars λ_j satisfies $\sum_j |\lambda_j|^p \lesssim \|f\|_{H_{w^p}^p}^p$ and the series converges in $L^{p_0}(\mathbb{R}^n)$. For $\frac{1}{q_0} = \frac{1}{p_0} - \frac{\alpha}{n}$, $T_{\alpha,m}$ is a bounded operator from $L^{p_0}(\mathbb{R}^n)$ into $L^{q_0}(\mathbb{R}^n)$, this follows from the pointwise estimate that appears in (2). Since $f = \sum_j \lambda_j a_j$ in $L^{p_0}(\mathbb{R}^n)$, we have that

$$(6) \quad |T_{\alpha,m} f(x)| \leq \sum_j |\lambda_j| |T_{\alpha,m} a_j(x)|, \quad \text{a.e. } x \in \mathbb{R}^n.$$

If $\|T_{\alpha,m} a_j\|_{L_{w^q}^q} \leq C$, with C independent of the $w^p - (p, p_0, d)$ atom $a_j(\cdot)$, then (6) allows us to obtain

$$\|T_{\alpha,m} f\|_{L_{w^q}^q} \leq C \left(\sum_j |\lambda_j|^{\min\{1, q\}} \right)^{\frac{1}{\min\{1, q\}}} \leq C \left(\sum_j |\lambda_j|^p \right)^{1/p} \lesssim \|f\|_{H_{w^p}^p},$$

for all $f \in \widehat{\mathcal{D}}_0$, so the theorem follows from the density of $\widehat{\mathcal{D}}_0$ in $H_{w^p}^p(\mathbb{R}^n)$.

To conclude the proof we will prove that there exists an universal constant $C > 0$ such that

$$(7) \quad \|T_{\alpha,m} a\|_{L_{w^q}^q} \leq C, \quad \text{for all } w^p - (p, p_0, d) \text{ atom } a(\cdot).$$

To prove (7), let $B_i^* = B(A_i x_0, 2Mr)$ for $i = 1, \dots, m$, where $B = B(x_0, r)$ is the ball containing the support of the atom $a(\cdot)$. So

$$\begin{aligned} \int_{\mathbb{R}^n} |T_{\alpha,m} a(x)|^q w^q(x) dx &= \int_{\bigcup_{i=1}^m B_i^*} |T_{\alpha,m} a(x)|^q w^q(x) dx \\ &+ \int_{\mathbb{R}^n \setminus (\bigcup_{i=1}^m B_i^*)} |T_{\alpha,m} a(x)|^q w^q(x) dx = J_1 + J_2 \end{aligned}$$

To estimate J_1 , we apply Hölder's inequality with $\frac{q_0}{q}$ and use that $w^q \in RH_{\frac{q_0}{q}}$, ($p_0 > p \frac{r_{w^p}}{r_{w^p}-1}$ and Lemma 11 imply that $q_0 > q \frac{r_{w^q}}{r_{w^q}-1}$), thus

$$\begin{aligned} J_1 &\lesssim \|T_{\alpha,m} a\|_{L^{q_0}^q}^q \sum_{i=1}^m \left(\int_{B_i^*} [w^q(x)]^{(\frac{q_0}{q})'} dx \right)^{1/(\frac{q_0}{q})'} \\ &\lesssim |B|^{q/p_0} (w^p(B))^{-q/p} \sum_{i=1}^m |B_i^*|^{-q/q_0} w^q(B_i^*) \end{aligned}$$

Lemma 5 implies

$$\lesssim |B|^{q\alpha/n} (w^p(B))^{-q/p} w^q(B).$$

Then, Lemma 9 gives

$$(8) \quad J_1 \leq C.$$

Now we estimate J_2 . By Lemma 14, considering there $d = \lfloor n(\frac{1}{p} - 1) \rfloor$ and w^p instead of w , we have

$$|T_{\alpha,m}a(x)|^q \lesssim (w^p(B))^{-q/p} \sum_{k=1}^m \left[M_{\frac{\alpha n}{n+d+1}}(\chi_B)(A_k^{-1}x) \right]^{q \frac{n+d+1}{n}}, \text{ for all } x \notin \bigcup_{i=1}^m B_i^*,$$

by integrating this pointwise estimate over $\mathbb{R}^n \setminus (\bigcup_{i=1}^m B_i^*)$ with respect to $w^q(x)dx$ and since $w(A_j x) \lesssim w(x)$ a.e. $x \in \mathbb{R}^n$, $1 \leq j \leq m$, we obtain that

$$(9) \quad J_2 \lesssim (w^p(B))^{-q/p} \int_{\mathbb{R}^n} \left[M_{\frac{\alpha n}{n+d+1}}(\chi_B)(x) \right]^{q \frac{n+d+1}{n}} w^q(x) dx.$$

Being $d = \lfloor n(\frac{1}{p} - 1) \rfloor$, we have $q \frac{n+d+1}{n} > p \frac{n+d+1}{n} > 1$. We write $\tilde{q} = q \frac{n+d+1}{n}$ and let $\frac{1}{p} = \frac{1}{\tilde{q}} + \frac{\alpha}{n+d+1}$, so $\frac{\tilde{q}}{q} = \frac{p}{q}$ and $w^{q/\tilde{q}} \in \mathcal{A}_{\tilde{p}, \tilde{q}}$ (see Remark 7). From Theorem 8 we obtain

$$\int_{\mathbb{R}^n} \left[M_{\frac{\alpha n}{n+d+1}}(\chi_B)(x) \right]^{q \frac{n+d+1}{n}} w^q(x) dx \lesssim \left(\int_{\mathbb{R}^n} \chi_B(x) w^p(x) dx \right)^{q/p} = (w^p(B))^{q/p}.$$

This inequality and (9) give

$$(10) \quad J_2 \leq C.$$

Finally, (8) and (10) allow us to obtain (7). This completes the proof. \square

In the following corollary we recover Theorem 4.10 obtained in [16].

Corollary 19. *For $0 < \alpha < n$, let I_α be the Riesz potential on \mathbb{R}^n . If $w^{\frac{n}{(n-\alpha)s}} \in \mathcal{A}_1$ with $0 < s < 1$, and $\frac{r_w}{r_w-1} < \frac{n}{\alpha}$, then I_α can be extended to an $H_{w^p}^p(\mathbb{R}^n)$ - $L_{w^q}^q(\mathbb{R}^n)$ bounded operator for each $s \leq p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$.*

Proof. To apply Theorem 18 with $A_j = id$ for all $j = 1, \dots, m$. \square

REFERENCES

- [1] M. Bownik, Boundedness of operators on Hardy spaces via atomic decompositions. *Proc. Amer. Math. Soc.*, 133, 3535-3542, (2005).
- [2] J. García-Cuerva, Weighted H^p spaces, *Dissertations Math.*, 162, 1-63, (1979).
- [3] T. Godoy and M. Urciuolo, *About the L^p -boundedness of some integral operators*, Rev. de la Un. Mat. Arg., 38 (3), 192-195, (1993).
- [4] T. Godoy and M. Urciuolo, *On certain integral operators of fractional type*, Acta Math. Hungar. 82, 99-105, (1999).
- [5] L. Grafakos, Classical Fourier Analysis, 3rd edition, Graduate Texts in Mathematics, 249, Springer New York, (2014).
- [6] S. Krantz, *Fractional integration on Hardy spaces*, Studia Mathematica, vol 73 (2), 87-94, (1982).
- [7] M. -Y. Lee, C. -C. Lin, The molecular characterization of weighted Hardy spaces, *Journal of Funct. Analysis*, 188, 442 - 460, (2002).
- [8] X. Li and L. Peng, The molecular characterization of weighted Hardy spaces, *Science in China (Series A)*, vol. 44 (2), 201-211, (2001).
- [9] B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal function, *Trans. of the Amer. Math. Soc.*, Vol 165, 207-226, (1972).
- [10] B. Muckenhoupt and R. L. Wheeden, Weighted norm inequalities for fractional integrals, *Trans. of the Amer. Math. Soc.*, Vol 192, 261-274, (1974).
- [11] F. Ricci and P. Sjögren, *Two-parameter maximal functions in the Heisenberg group*, Math. Z. 199 (4), 565-575, (1998).
- [12] M. S. Riveros and M. Urciuolo, *Weighted inequalities for integral operators with some homogeneous kernels*, Czechoslovak Math. J., 55 (130), 423-432, (2005).
- [13] P. Rocha, A note on Hardy spaces and bounded linear operators, *Georgian Math. J.*, 25(1), 73-76, (2018).
- [14] P. Rocha, *A remark on certain integral operators of fractional type*, Canad. Math. Bull., Vol. 61 (2), 370-375, (2018).

- [15] P. Rocha, *Boundedness of generalized Riesz potentials on the variable Hardy spaces*, Journal of the Australian Math. Soc. Vol 104, Issue 2, 255-273, (2018).
- [16] P. Rocha, *On the atomic and molecular decomposition of weighted Hardy spaces*, Rev. de la Un. Mat. Arg., 61 (2), 229-247, (2020).
- [17] P. Rocha and M. Urciuolo, *On the $H^p - L^p$ boundedness of some integral operators*, Georgian Math. Journal, vol 18 (4), 801-808, (2011).
- [18] P. Rocha and M. Urciuolo, *On the $H^p - L^q$ boundedness of some fractional integral operators*, Czech. Math. Journal, 62 (137), 625-635, (2012).
- [19] P. Rocha and M. Urciuolo, *Fractional type integral operators on variable Hardy spaces*, Acta Math. Hungar., 143 (2), 502-514, (2014).
- [20] P. Rocha and M. Urciuolo, *Fractional type integral operators of variable order*, Rev. de la Un. Mat. Arg., 58 (2), 281-296, (2017).
- [21] J. O. Strömberg and R. L. Wheeden, *Fractional integrals on weighted H^p and L^p spaces*, Trans. Amer. Math. Soc., 287, 293-321, (1985).
- [22] J. O. Strömberg and A. Torchinsky, *Weighted Hardy spaces*, Lecture Notes in Mathematics, vol 131, Springer-Verlag, Berlín, (1989).
- [23] M. H. Taibleson and G. Weiss, *The molecular characterization of certain Hardy spaces*, Asterisque 77, 67-149, (1980).
- [24] J. Tan and J. Zhao, *Fractional Integrals on Variable Hardy-Morrey Spaces*, Acta Math. Hungar., 148 (1), 174-190, (2016).
- [25] D. Yang and Y. Zhou, *A boundedness Criterion via atoms for linear operators in Hardy spaces*, Constr. Approx., 29, 207-218, (2009).

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD NACIONAL DEL SUR, BAHÍA BLANCA, 8000 BUENOS AIRES, ARGENTINA.

Email address: `pablo.rocha@uns.edu.ar`