

Rapid mixing of the hardcore Glauber dynamics and other Markov chains in bounded-treewidth graphs

David Eppstein*

Daniel Frishberg* 

November 3, 2021

Abstract

We give a new rapid mixing result for a natural random walk on the independent sets of an input graph G . Rapid mixing is of interest in approximately sampling a structure, over some underlying set or graph, from some target distribution. The structures are typically exponential in number with respect to G . In the case of independent sets, we show that when G has bounded treewidth, this random walk—known as the *hardcore Glauber dynamics*—mixes rapidly for all values of the standard parameter $\lambda > 0$, giving a simple alternative to existing sampling algorithms for these structures.

We also show rapid mixing for analogous Markov chains on dominating sets and b -edge covers (for fixed $b \geq 1$ and $\lambda > 0$) in the case where treewidth is bounded, and for Markov chains on the b -matchings (for fixed $b \geq 1$ and $\lambda > 0$), the maximal independent sets, and the maximal b -matchings of a graph (for fixed $b \geq 1$), in the case where carving width is bounded. Our results imply simpler alternatives to known algorithms for the sampling and approximate counting problems in these graphs.

We prove our results by developing a divide-and-conquer framework using the well-known *multicommodity flows* technique. Using this technique, we additionally show that a similar dynamics on the k -angulations of a convex set of n points mixes in quasipolynomial time for all $k \geq 3$. (McShine and Tetali gave a stronger result in the special case $k = 3$ [41].)

Our technique also allows us to strengthen existing results by Dyer, Goldberg, and Jerrum [16] and by Heinrich [27] for the Glauber dynamics on the q -colorings of G on graphs of bounded carving width, when $q \geq \Delta + 2$ is bounded. Specifically, our technique yields an improvement in the dependence on treewidth when $\Delta < 2t$ or when $q < 4t$ and $\Delta < t^2$.

We additionally show that the Glauber dynamics on the *partial* q -colorings of G mix rapidly for all $\lambda > 0$ when $q \geq \Delta + 2$ is bounded.

1 Introduction

The *Glauber dynamics* on independent sets in a graph—motivated in part by modeling systems in statistical physics—is a Markov chain in which one starts at an arbitrary independent set, then

*Department of Computer Science, University of California, Irvine, {[eppstein](mailto:eppstein@uci.edu), [dfrishbe](mailto:dfrishbe@uci.edu)}@uci.edu

repeatedly chooses a vertex at random and, with probability that depends on a fixed parameter $\lambda > 0$, either removes the vertex from the set (if it is in the set), or adds it to the set (if it is not in the set and has no neighbor in the set). This chain, also known as the *hardcore model*, has seen recent rapid mixing results under various conditions.

Prior work [8, 22] has shown that related chains, including the *softcore model*—in which the sampled sets need not be independent—mix rapidly on graphs of bounded treewidth. However, as Bordewich and Kang [8] note, the hardcore model, which we consider in this paper, is different, and their technique does not apply to this model.

Bezáková and Sun showed [5] that the hardcore model mixes rapidly in chordal graphs with bounded-size separators. Anari, Liu, and Gharan showed [3] that a biased version of the chain mixes rapidly for certain values of λ ; Chen, Galanis, Štefankovič, and Vigoda [14] and Feng, Guo, Yin, and Zhang [19] generalized this work and applied it to graph colorings.

1.1 Our contribution

We prove that the hardcore Glauber dynamics mixes rapidly on graphs of bounded treewidth for all fixed $\lambda > 0$, and that the Glauber dynamics on partial q -colorings (for all $\lambda > 0$) and on q -colorings of a graph of bounded carving width mix rapidly on graphs of bounded carving width. Marc Heinrich proved the latter result, namely for q -colorings, in a 2020 preprint [27]. Heinrich’s result applies to all graphs of bounded treewidth; however, for graphs of bounded carving width whose degree is less than quadratic in their treewidth, we improve on Heinrich’s upper bound—provided that q is fixed.

We also prove that the analogous dynamics on the b -edge covers (when b is bounded) and the dominating sets of a graph of bounded treewidth mix rapidly for all $\lambda > 0$.

In a similar vein, we prove that three additional chains—on b -matchings (when $\lambda > 0$), on maximal independent sets, and on maximal b -matchings—mix rapidly in graphs where carving width is bounded. (For the latter two chains we consider only the unbiased version.)

To prove our results, we introduce a framework that uses the well-established *multicommodity flow* technique (essentially the same as the *canonical paths* technique) for bounding mixing times. The framework consists of a set of conditions that guarantee rapid mixing.

Finally, we prove that the flip chain on the k -angulations of a convex n -point set mixes in time quasipolynomial in n for all fixed $k \geq 3$, although the special case $k = 3$ is well known already to mix rapidly [41].

1.2 Prior work

Sly [56] showed that, except for restricted values of $\lambda < 1$, the hardcore Glauber dynamics does not mix rapidly on general graphs unless $\text{RP} = \text{NP}$, and in fact showed that approximately sampling from the corresponding distribution is hard unless $\text{RP} = \text{NP}$. However, Anari, Liu, and Gharan [3] used a technique known as *spectral independence* to obtain rapid mixing for the hardcore Glauber dynamics when λ is below the so-called *uniqueness threshold* that depends on the maximum degree of the input graph. They showed, by exhibiting an infinite family of examples, that the technique they used could not be further improved (namely beyond the uniqueness threshold) even for trees. By contrast, we show that rapid mixing, for all fixed values of λ , indeed holds not only for trees but for all graphs of bounded treewidth.

Other results exist for trees beyond the uniqueness threshold, however: Martinelli, Sinclair, and Weitz [40] showed that the Glauber dynamics on the hardcore model mixes in $O(n \log n)$ time on the complete $\Delta - 1$ -ary tree with n nodes. They also showed that the dynamics on q -colorings ($q \geq \Delta + 2$) mixes in $O(n \log n)$ time on the same trees. Lucier, Molloy, and Peres [38] showed that the dynamics mixes rapidly on general trees of bounded degree, namely in time $O(n^{O(1+\Delta/(q \log \Delta))})$.

Prior work also exists for q -colorings of bounded-treewidth graphs: Vardi [58] showed that the so-called *single-flaw* dynamics—a variation on the Glauber dynamics in which at most one monochromatic edge is permitted in a valid state—mixes rapidly on bounded-treewidth graphs when $q \geq (1 + \varepsilon)\Delta$, for any fixed parameter $\varepsilon > 0$. The proof used the vertex separation properties of bounded-treewidth graphs to construct a multicommodity flow with bounded congestion, although the construction is substantially different from our divide-and-conquer approach. Dyer, Goldberg, and Jerrum [16] showed rapid mixing when the degree of the graph is at least $2t$ and $q \geq 4t$, where t is the treewidth. On the other hand, Heinrich [27] showed that the Glauber dynamics on q -colorings of a bounded-treewidth graph mixes rapidly when $q \geq \Delta + 2$. Our construction, as we will discuss in more detail in Section 1.4.1, bears some similarity to Heinrich’s. We also require that q (and therefore Δ) be bounded. However, due to a more general analysis of the state spaces of Glauber dynamics as graphs, we obtain a more general framework that holds for a greater variety of chains.

Planar graphs have unbounded but sublinear treewidth. For planar graphs, Hayes [25] showed that the Glauber dynamics on q -colorings of a planar graph of maximum degree Δ mixes rapidly when $q \geq \Delta + O(\sqrt{\Delta})$. Later, Hayes, Vera, and Vigoda [26] proved rapid mixing for q -colorings of planar graphs when $q = \Omega(\Delta / \log \Delta)$, generalizing further to a spectral condition on the adjacency matrix of the graph.

Lastly, Chen, Galanis, Štefankovič, and Vigoda applied the spectral independence technique to prove that the Glauber dynamics on the q -colorings of a triangle-free graph with degree Δ mixes rapidly provided that $q \geq \alpha\Delta + 1$, where α is greater than a threshold approximately equal to 1.763. We show that when the carving width of G is bounded, G need not be triangle free, and it suffices that $q \geq \Delta + 2$ be bounded. We prove a similar result for the natural Glauber dynamics on *partial* q -colorings.

Although our mixing results are new, Wan, Tu, Zhang, and Li showed [59] that exact counting of independent sets is fixed-parameter tractable in treewidth. Furthermore, our result does not technically constitute a proof of fixed-parameter tractability, as the treewidth appears in the exponent of the polynomial we obtain. For this problem and all the other problems we consider, the problem of exact counting—and therefore also uniform sampling—has already been solved on the graphs we consider by an extension of Courcelle’s theorem [48]. In fact, the standard reduction from approximate sampling to approximate counting [54] gives a somewhat different rapidly mixing Markov chain on a larger state space. Nonetheless, our result does settle the question of rapid mixing for a natural chain, and it implies a simpler scheme for approximately sampling independent sets than one would obtain via this reduction.

Such a scheme is known as a *fully polynomial randomized approximation scheme (FPRAS)*. Huang, Lu, and Zhang provided an FPRAS for sampling b -edge covers in general graphs when $b \leq 2$, and for sampling b -matchings when $b \leq 7$ [29]. This FPRAS relied on a rapid mixing argument for a somewhat different Markov chain than ours. Existing dominating set results for certain regular graphs are also known [6].

Exact counting of maximal independent sets—which would give an FPRAS by the equivalence of

counting and sampling—was shown in [47] to be hard for chordal graphs but is known [13] to be tractable in graphs of bounded treewidth. However, again our result improves on the simplicity of existing algorithms.

1.3 Main results

Our main results are the following (see Section 2 for relevant definitions). When $\hat{\lambda}$ appears in a formula, we take $\hat{\lambda} = \max\{\lambda, 1/\lambda\}$, for all fixed values of the standard Glauber dynamics parameter $\lambda > 0$.

Theorem 1.1. *The hardcore Glauber dynamics mixes in time*

$$O(((1 + \hat{\lambda})\hat{\lambda})^2(1 + \log \hat{\lambda})n^{2(t+2)(1+\log \hat{\lambda})+5})$$

on graphs of treewidth t for all $\lambda > 0$.

Theorem 1.2. *The (unbiased) Glauber dynamics on q -colorings (when $q \geq \Delta + 2$) mixes in time*

$$O((q - 1)^2 \log q \cdot n^{4(t+1)\Delta \log q + 7})$$

on graphs of treewidth t and degree Δ . The Glauber dynamics on partial q -colorings (when $q \geq \Delta + 2$) mixes in time

$$O(((1 + \hat{\lambda})\hat{\lambda})^2 q^2 (\log(q + 1) + \log \hat{\lambda}) \cdot n^{2(t+2)(\log(q+1)+\log \hat{\lambda})+5})$$

on graphs of treewidth t for all $\lambda > 0$.

Theorem 1.3. *The Glauber dynamics on b -edge covers mixes in time*

$$O\left(\left((2\hat{\lambda})^{36(t+1)^2}\right)^2 \left((1 + \hat{\lambda})\hat{\lambda}\right)^2 m^3 (\log \hat{\lambda} + 1) n^{\frac{2(3+\log(\frac{1+\hat{\lambda}}{\lambda})+(t+1)\log(b+1)+((t+1)(b+t/2)+1)(1+\log \hat{\lambda}))}{\log(6/5)}}\right)$$

on graphs of treewidth t , for all $\lambda > 0$. The Glauber dynamics on dominating sets mixes in time

$$O\left(\left((2\hat{\lambda})^{12(t+1)}\right)\left((1 + \hat{\lambda})\hat{\lambda}\right)^2 (1 + \log \hat{\lambda}) \cdot n^{\frac{2(3+\log(\frac{1+\hat{\lambda}}{\lambda})+(t+2)(3+\log \hat{\lambda}))}{\log(6/5)}+3}\right)$$

on graphs of treewidth t for all $\lambda > 0$. The Glauber dynamics on b -matchings mixes in time

$$O(((1 + \hat{\lambda})\hat{\lambda})^2(1 + \log \hat{\lambda})m^3 n^{2\Delta(t+2)(1+\log \hat{\lambda})+2})$$

on graphs of treewidth t and degree Δ for all $\lambda > 0$.

Theorem 1.4. *There exist flip chains on maximal independent sets and maximal b -matchings, whose stationary distributions are uniform, that mix respectively in time*

$$O(2^{6\Delta^2} n^{2((t+1)(7\Delta^6+\Delta+1)+4)/\log(3/2)+3})$$

and time

$$O(2^{12\Delta^2} m^3 n^{2((t+1)(8\Delta^7+3\Delta^2))/\log(3/2)+4})$$

on graphs of treewidth t and degree Δ .

Theorem 1.5. *For every fixed $k \geq 3$, the k -angulation flip chain on the convex $(k - 2)n + 2$ -point set mixes in time*

$$O((k - 1)^3 (\log(k - 1)) \cdot n^{2(3k \log(k-1)+k(1+\log \pi)+3k \log n+k)+7}).$$

1.4 The framework: divide and conquer

A *multicommodity flow* (see Section 2.3 for a full definition) in a graph \mathcal{M} is a set of $|V(\mathcal{M})|^2$ flows, one flow for each ordered pair of vertices (u, v) , where each flow sends one unit of a commodity from u to v . If a flow exists in \mathcal{M} with small *congestion*—i.e. one in which no edge carries too much flow—then the natural Markov chain whose states are the vertices of \mathcal{M} mixes rapidly. (We make this precise in Section 2.2.)

All of the chains we analyze are such natural random walks on a “Glauber graph” $\mathcal{M}(G)$ whose vertices are subsets of an underlying set. (For our purposes, this underlying set is either the vertex set or the edge set of the input graph G .) Thus each of these random walks is performed on a graph that may be exponentially large with respect to the size of the input graph. We show that when all of a certain set of conditions hold, we can construct a multicommodity flow in $\mathcal{M}(G)$ with congestion polynomial in n , implying that the unbiased random walk on $\mathcal{M}(G)$ mixes rapidly.

The conditions specify that $\mathcal{M}(G)$ can be partitioned into a small number of induced subgraphs, all of which are approximately the same size, with large numbers of edges between pairs of classes. The conditions also require that each of these induced subgraphs have a special structure, allowing for the decomposition of each induced subgraph into smaller Glauber graphs that are similar in structure to $\mathcal{M}(G)$. This self similarity allows for the inductive construction of a multicommodity flow, by assembling flows on smaller Glauber graphs together into a flow in $\mathcal{M}(G)$ with small congestion.

1.4.1 Prior work and similar approaches

Prior work on rapid mixing of Markov chains on subset systems includes the special case of matroid polytopes. For this case, recent results [1, 2] have partly solved a 30-year-old conjecture of Mihail and Vazirani [42]. Other prior work uses multicommodity flows (and the essentially equivalent *canonical paths* technique) to obtain polynomial mixing upper bounds on structures of exponential size, including matchings and 0/1 knapsack solutions [44, 24]. Madras and Randall [39] used a decomposition of the hardcore model state space to prove rapid mixing under different conditions. We also decompose the state space, but our approach is different from that of Madras and Randall and is more similar to Heinrich’s [27] application of the *projection/restriction* technique pioneered by Jerrum, Son, Tetali, and Vigoda [31]. Essentially, the projection/restriction technique involves partitioning the state space of a chain into a collection of sub-state spaces, each of which internally mixes rapidly, and all of which are well connected to one another. Heinrich used the vertex separation properties of bounded-treewidth graphs to obtain an inductive argument: the resulting sub-spaces are themselves Cartesian products of chains on smaller graphs, and thus mix rapidly. (See Lemma 2.17.)

We partition the state space recursively using the same vertex separation properties, but our analysis is more general, allowing us to prove results for a number of other chains beyond q -colorings. Our framework also solves a key problem that arises in applying the projection/restriction technique: to apply that technique in Heinrich’s recursive fashion, one needs each of the state spaces in the partition to be a Cartesian product of chains on smaller spaces. For four of our nine chains, the sub-spaces obtained in the decomposition are not Cartesian products but may be non-disjoint unions of Cartesian products. In some cases, the sub-spaces may not even be mutually disjoint and may induce non-ergodic chains. We solve this problem by using the structure of the state spaces of Glauber dynamics as graphs to obtain multicommodity flows with bounded congestion. We will discuss this further in Section 8.

Our approach is also inspired by Kaibel’s [32] construction of a flow with bounded congestion in any graph whose vertices are hypercube vertices and whose edges can be partitioned into bipartite graphs in a hierarchical fashion.

1.5 Paper organization

In Section 2, we give relevant definitions and background. This includes defining the flip chains under consideration, as well as the notions of expansion and treewidth. In Section 3.4, we use the chain on independent sets to illustrate what we call a “non-hierarchical” version of the framework. This non-hierarchical version works on this chain when carving width is bounded, and in Sections 3.5 and Section 3.6 we describe how to apply it respectively to q -colorings, and to b -edge covers and b -matchings. We defer some details of this proof to Appendix A.

To fully prove Theorem 1.1 and Theorem 1.3, we need to deal with unbounded-degree graphs. In Section 4, we discuss how to modify the framework to accomplish this, proving Theorem 1.1 for $\lambda = 1$. We defer some of the details of this proof to Appendix B, in which we also finish the proof of Theorem 1.2 for $\lambda = 1$.

We prove the general case $\lambda > 0$ of Theorems 1.1 and 1.2 in Appendix C. We finish the proofs of Theorems 1.3 and 1.4 in Appendix D: applying the framework to the relevant chains requires a refinement of the framework.

In Section 6 we describe the application of our framework to the k -angulation flip chain, and in Appendix E we give detailed proofs and derive the mixing upper bound claimed in Theorem 1.5.

In all of the above, we prove rapid mixing but defer derivation of specific upper bounds to Section 7.

2 Preliminaries

2.1 Rapid mixing and Glauber dynamics

Rapid mixing is of interest in the random generation of certain combinatorial objects, including such subset systems as the set of matchings in a graph. To generate, approximately uniformly at random, an object of a given class—say, an independent set in a given graph—it suffices to conduct a random walk on a graph whose vertices are the objects of interest, and whose edges are *flips* between the objects, under some suitable definition of a flip. (For technical reasons, self loops need to be added to the graph in a standard fashion.) Basic spectral graph theory shows that, under mild conditions, the walk converges to the uniform distribution in the limit. It is of interest for efficient sampling algorithms to determine how rapid the convergence is. In the case of subset systems such as those we consider, the walk takes place over an exponentially large number of subsets defined over an underlying set of size n . If the convergence, or *mixing time*, of the walk is polynomial in n , then the random walk is said to be *rapidly mixing*.

The mixing time is typically denoted as a function $\tau(\varepsilon)$, where ε denotes the desired precision of convergence to the uniform distribution, and the value of τ at ε is the minimum number of steps in the random walk before convergence is guaranteed. Convergence is measured via the *total variation distance* [53] between the distribution over states induced by the walk at a given time step, and the uniform distribution. One can obtain convergence to other distributions by adding weights to the vertices and edges of the graph—see Appendix C.1.

See Levin, Peres, and Wilmer [36] for a comprehensive treatment of rapid mixing.

Of interest for our mixing results is the hardcore Glauber dynamics, defined as follows:

Definition 2.1. *The hardcore Glauber dynamics on the independent sets of a graph G is the following chain, defined with respect to a fixed real parameter $\lambda > 0$:*

1. Let X_0 be an arbitrary independent set in G .
2. For $t \geq 0$, select a vertex $v \in V(G)$ uniformly at random.
3. If $v \notin X_t$ and $X_t \cup \{v\}$ is not a valid independent set, do nothing.
4. Otherwise:

Let $X_{t+1} = X_t \cup \{v\}$ with probability $\lambda/(\lambda + 1)$.

Let $X_{t+1} = X_t \setminus \{v\}$ with probability $1/(\lambda + 1)$.

For dominating sets, b -edge covers, and b -matchings, we consider an almost identical chain, except that in the case of b -edge covers and b -matchings, we are of course selecting edges instead of vertices. Also, in the case of dominating sets and b -edge covers, instead of verifying independence before adding a vertex (or edge), we verify validity of a set (e.g. domination) before dropping a vertex (or edge).

We also consider the Glauber dynamics on q -colorings:

Definition 2.2. *A q -coloring of a graph G is an assignment of a color from the list $[q] = \{1, 2, \dots, q\}$ to each vertex of G , such that no two adjacent vertices have the same color.*

Definition 2.3. *A partial q -coloring of a graph G is an assignment of a color from $[q]$ to each of a subset of the vertices of G , such that no two adjacent vertices have the same color.*

The Glauber dynamics on the partial q -colorings of G is as follows:

Definition 2.4. *Let the Glauber dynamics on the partial q -colorings of a graph G be the following chain defined with respect to $\lambda > 0$:*

1. Let X_0 be an arbitrary partial q -coloring of G .
2. For $t \geq 0$, select a vertex $v \in V(G)$ uniformly at random, and select a color $c \in [q + 1]$ uniformly at random.
3. If $c = q + 1$, then:

If v is already colored in X_t , remove the coloring of v with probability $1/(\lambda + 1)$.

Otherwise, let $X_{t+1} = X_t$.

4. If $c \leq q$, then:

If v is not already colored with c in X_t , set the color of v to c with probability $\lambda/(\lambda + 1)$.

Otherwise, let $X_{t+1} = X_t$.

Finally, the Glauber dynamics on the (complete) q -colorings of G is as follows (for this chain we do not define a biased version):

Definition 2.5. *Let the Glauber dynamics on the q -colorings of a graph G be the following chain:*

1. Let X_0 be an arbitrary q -coloring of G .
2. For $t \geq 0$, select a vertex $v \in V(G)$ uniformly at random, and select a color $c \in [q]$ —other than the color of v —uniformly at random.
3. If v has no neighbor with color c , then change the color of v to c with probability $1/2$ to obtain X_{t+1} .
4. Otherwise, do nothing, i.e. let $X_{t+1} = X_t$.

We define a graph whose vertices are the maximal independent sets of an underlying graph G , and then define the flip chain as a random walk on this graph:

Definition 2.6. *Given a graph $G = (V, E)$, let the maximal independent set Glauber graph be the graph $\mathcal{M}_{\text{MIS}}(G)$ whose vertices are the maximal independent sets of G , and whose edges are the pairs of maximal independent sets that differ by one flip, where a flip is defined as:*

1. adding one vertex v to a given independent set $S \subseteq V$,
2. removing every $u \in S$ such that $(u, v) \in E$, and
3. adding a subset of the vertices at distance two in G from v .

Since $\mathcal{M}_{\text{MIS}}(G)$ is undirected, we also define the reversal of a flip as a flip. See Figure 8 for an example of a flip.

Lemma 2.7. *The graph $\mathcal{M}_{\text{MIS}}(G)$ in Definition 2.6 is connected.*

Proof. The proof relies on an easy greedy transformation argument and is in Appendix F. □

For maximal b -matchings, we define a Glauber graph similar to the maximal independent set Glauber graph, except that we are of course selecting edges instead of vertices in our sets. A flip consists of adding some edge $e = (u, v)$ to the b -matching, then removing edges incident to u and v as needed until a valid b -matching is obtained, then adding edges incident to neighbors of u and v as needed to obtain maximality.

2.2 Mixing and expansion

Formally, the Glauber dynamics is defined as follows:

Definition 2.8. *The Glauber dynamics is a Markov chain, parameterized by $\lambda > 0$, with state space $\Omega = V(\mathcal{M}(G))$ and probability matrix P , where for $S, S' \in V(\mathcal{M}(G))$ with $S \neq S'$,*

$$P(S, S') = \lambda / (\Delta_{\mathcal{M}}(\lambda + 1))$$

when $|S' \setminus S| = 1$, and

$$P(S, S') = 1 / (\Delta_{\mathcal{M}}(\lambda + 1))$$

when $|S \setminus S'| = 1$. If $S = S'$, then $P(S, S') = 1 - \sum_{S'' \neq S} P(S, S'')$.

Here $\Delta_{\mathcal{M}}$ is the maximum degree of the Glauber graph.

The two cases described in Definition 2.8 for $S \neq S'$ are exhaustive for all of the chains that we have parameterized by λ . For all of the chains for which λ is not defined, the transition probability is in every case $1/(2\Delta_{\mathcal{M}})$.

Definition 2.9. The total variation distance between two probability distributions π and π^* over the same set Ω is defined as

$$d(\pi, \pi^*) = \frac{1}{2} \sum_{S \in \Omega} |\pi(S) - \pi^*(S)|.$$

A Markov chain, given a starting state $S \in \Omega$, induces a probability distribution π_t at each time step t . The Glauber dynamics is well known, regardless of starting state, to converge in the limit to a *stationary* distribution $\pi^*(S) = \lambda^{|S|}/Z(\mathcal{M}(G))$, where the term $Z(\mathcal{M}(G))$ is simply a normalizing value. When λ is not defined, one may assume $\lambda = 1$, i.e. the stationary distribution is uniform.

The *mixing time* is defined as follows:

Definition 2.10. Given an arbitrary $\varepsilon > 0$, the mixing time, $\tau(\varepsilon)$, of a Markov chain with state space Ω and stationary distribution π^* is the minimum time t such that, regardless of starting state, we always have

$$d(\pi_t, \pi^*) < \varepsilon.$$

If $\tau(\varepsilon)$ is upper bounded by a function that is polynomial in $\log(1/\varepsilon)$ and in $n = |\Omega|$, the chain is said to be rapidly mixing.

For our chains, we omit the parameter ε , assuming its value to be an arbitrary constant, say $1/4$.

Closely related to mixing is the notion of *expansion*.

Definition 2.11. The edge expansion (or simply expansion), $h(G)$, of a graph $G = (V, E)$ is the quantity

$$\min_{S \subseteq V: 1 \leq |S| \leq |V|/2} |\partial S|/|S|,$$

where

$$\partial S = \{(s, t) | s \in S, t \notin S\}$$

is the set of edges across the $(S, V \setminus S)$ cut.

It is known [53] that a lower bound on expansion leads to an upper bound on mixing when the degree of the Glauber graph is bounded (as is the case for all of our chains):

Lemma 2.12. The mixing time of the Markov chain whose transition matrix is the normalized adjacency matrix of a graph G with N vertices and maximum degree Δ is at most

$$\tau(\varepsilon) = O\left(\frac{\Delta^2}{(h(G))^2} \cdot \ln \frac{N}{\varepsilon}\right).$$

(Tighter bounds are available via Sinclair [53] under some conditions.)

2.3 Cartesian products, expansion, and multicommodity flows

Definition 2.13. The Cartesian product $G \square H$ of the graphs G and H is the graph with vertex set $V(G) \times V(H)$ and the edge set

$$\begin{aligned} & \{((u, v), (u', v)) | (u, u') \in E(G), v \in V(H)\} \\ & \cup \{((u, v), (u, v')) | (v, v') \in E(H), u \in V(G)\}. \end{aligned}$$

Definition 2.14. A multicommodity flow ρ [53, 32] in a graph $\mathcal{M} = (\mathcal{V}, \mathcal{E})$ is a collection of functions $\{\rho_{st} : \mathcal{A} \rightarrow \mathbb{R} \mid s, t \in \mathcal{V}\}$, where

$$\mathcal{A} = \bigcup_{\{u,v\} \in \mathcal{E}} \{(u,v), (v,u)\},$$

and each ρ_{st} is a flow sending a single unit of a commodity from vertex u to vertex v through the edges of G . We consider the capacities of all edges to be infinite. Let $\rho_{st}(u,v)$ be the amount of flow sent by ρ_{st} across (u,v) , divided by $N = |\mathcal{V}|$. That is,

$$\sum_{p \in P(s,t)} \rho_{st}(p) \cdot N = 1,$$

where $P(s,t)$ is the set of simple paths in G from s to t , and where

$$\rho_{st}(u,v) = \sum_{p:(u,v) \in p} \rho_{st}(p)$$

for all $(u,v) \in \mathcal{A}$.

Note that $\rho_{st}(u,v)$ need not be equal to $\rho_{st}(v,u)$. Let

$$\rho(u,v) = \sum_{s,t \in \mathcal{V} \times \mathcal{V}} \rho_{st}(u,v),$$

and let

$$\rho_{max} = \max_{(u,v) \in \mathcal{A}} \rho(u,v).$$

The following is well known and enables the use of multicommodity flows as a powerful lower-bounding technique for expansion. The technique is closely related to *canonical paths* [53].

Lemma 2.15. Given a multicommodity flow ρ in a graph $G = (V, E)$, the expansion $h(G)$ is at least $1/(2\rho_{max})$.

The following is known [23]:

Lemma 2.16. Given graphs G_1, G_2, \dots, G_k , Cartesian product $G_1 \square G_2 \square \dots \square G_k$ satisfies

$$h(G_1 \square G_2 \square \dots \square G_k) \geq \frac{1}{2} \min_i h(G_i).$$

It is easy to prove a similar result for multicommodity flows:

Lemma 2.17. Let $J = G \square H$. Given multicommodity flows ρ_G and ρ_H for G and H respectively, there exists a multicommodity flow ρ_J for J with

$$\rho_{Jmax} \leq \max\{\rho_{Gmax}, \rho_{Hmax}\}.$$

Proof. We defer the proof to Appendix F. □

2.4 Separators and treewidth

The *treewidth* of a graph G is a density parameter that is weaker than expansion, in the sense that a high expansion implies a high treewidth, but not vice versa. Treewidth is defined in terms of a so-called *tree decomposition*:

Definition 2.18. [50] *A tree decomposition of a graph $G = (V, E)$ is a collection of sets $\{X_i\}, i = 1, \dots, k$, called bags, together with a tree T , whose nodes are identified with the bags $\{X_i\}$, such that all of the following hold:*

1. *Every vertex in V lies in some bag, i.e. $\bigcup_{i=1}^k X_i = V$.*
2. *For every $(u, v) \in E$, the vertices u and v belong to at least one bag X_i together, i.e. for some i , $u \in X_i$ and $v \in X_i$.*
3. *The collection of all bags containing any given vertex $v \in V$, i.e. $\{X_i \mid v \in X_i\}$ forms a (connected) subtree of T .*

Definition 2.19. [50] *The width of a tree decomposition is one less than the size of the largest bag in the decomposition. The treewidth of a graph G is the minimum t such that a tree decomposition of G exists with width t .*

Intuitively, treewidth measures how far away a graph is from being a tree. For example, trees have treewidth one; a graph consisting of a single cycle of size at least three has treewidth two. Treewidth is of interest in large part because many NP-hard problems become tractable on graphs of bounded treewidth. For a full definition of treewidth and a survey of this phenomenon, known as *fixed-parameter tractability*, see [7].

For our purposes, treewidth is of interest due to its relationship to *vertex separators*:

Definition 2.20. *A vertex separator for a graph G is a subset $X \subseteq V(G)$ of the vertices of G such that $G \setminus X$ is disconnected. X is a balanced separator if $G[V \setminus X]$ consists of two subgraphs, A and B , such that no edge exists between A and B , and such that $|V(G)|/3 \leq |V(A)| \leq |V(B)| \leq 2|V(G)|/3$. We also say, if $|X| \leq s$ for a given $s \geq 1$, that X is an s -separator.*

Definition 2.21. *With respect to an integer $s \geq 1$, a graph G is recursively s -separable if either $|V(G)| \leq 1$, or G has a balanced s -separator X such that the two mutually disconnected subgraphs induced by removing X from G are both recursively s -separable.*

The following is well known and easy to prove [18]:

Lemma 2.22. *For every $t \geq 1$, every graph with treewidth at most t is recursively $t + 1$ -separable.*

2.5 Carving width

The *carving width* of a graph is a density parameter that is weaker than treewidth, in the sense that high treewidth implies high carving width, but the converse is not true. Carving width is defined with respect to a so-called *carving decomposition* [17] of a given graph G —in short, a binary tree T whose leaves are identified with the vertices of G . Each node $X \in T$ is identified with the subgraph of G induced by the vertices of G (leaves of T) having X as an ancestor in T . Each edge of T induces a cut in T ; this cut induces a partition of the leaves of T (vertices of G) into two sets. This partition is naturally identified with a cut in G .

The *width* of a carving decomposition is the maximum number of edges of G across any such cut, where the maximum is taken over all edges in T . The *carving width* of G is the minimum width of a carving decomposition of G . See Seymour and Thomas [52] for a detailed treatment. For our purposes, carving width is of interest due to its relationship to the treewidth and degree of a graph. Specifically, Eppstein [17] observed the following fact that follows from results of Nestoridis and Thilikos [46] and of Robertson and Seymour [51]:

Lemma 2.23. *Given a graph G with maximum degree Δ , let $\text{tw}(G)$ denote the treewidth of G , and let $\text{cw}(G)$ denote the carving width of G . For every graph G , $(2/3)(\text{tw}(G)+1) \leq \text{cw}(G) \leq \Delta(\text{tw}(G)-1)$.*

It follows from the definition of carving width that every graph with bounded carving width also has bounded degree. Combining this fact with Lemma 2.23 implies the following:

Corollary 2.24. *A graph has bounded degree and treewidth if and only if it has bounded carving width.*

2.6 Dominating sets, b -matchings, and b -edge covers

Definition 2.25. *A dominating set in a graph $G = (V, E)$ is a set $S \subseteq V$ of vertices such that for every vertex $v \in V$, either $v \in S$ or there exists some vertex $u \in S$ such that $(u, v) \in E$.*

b -matchings [34] and b -edge covers [20, 33] generalize the definitions of matchings and edge covers respectively:

Definition 2.26. *Let $G = (V, E)$ be a graph. Let $b : V \rightarrow \mathbb{Z}^{\geq 0}$ be any function assigning a nonnegative integer to each vertex. A b -matching in a graph $G = (V, E)$ is a set $S \subseteq E$ of edges such that every $v \in V$ has at most $b(v)$ incident edges in S .*

Definition 2.27. *Let $G = (V, E)$ be a graph. Let $b : V \rightarrow \mathbb{Z}^{\geq 0}$ be any function assigning a nonnegative integer to each vertex. A b -edge cover in a graph $G = (V, E)$ is a set $S \subseteq E$ of edges such that every $v \in V$ has at least $b(v)$ incident edges in S .*

Sometimes, as in the result by Huang, Lu, and Zhang [29], b -edge covers and b -matchings are defined so that b is a constant, i.e. $b(u) = b(v)$ for all $u, v \in V$.

2.7 k -angulations of convex point sets

2.7.1 Triangulations

A *triangulation* of a point set is a maximal set of non-crossing edges connecting pairs of points. Let P_n be the regular polygon with $n + 1$ vertices. Every triangulation t of P_n has $n - 2$ edges (henceforth *diagonals*), and every diagonal can be *flipped*: that is, every diagonal D belongs to two triangles forming a quadrilateral, such that D can be removed and replaced with the diagonal D' lying in the same quadrilateral and crossing D .

The set of all triangulations of P_n is the vertex set of a graph, denoted $K_{3,n}$, whose edges are the flips described above. This graph is known to be realizable as an $n - 3$ -dimensional polytope [37] called the *associahedron*. It is also known to be isomorphic to the rotation graph on the set of all binary plane trees with n leaves [55], and equivalently the set of all parenthesizations of an algebraic expression with n terms, with “flips” defined as applications of the associative property of multiplication.




Figure 1: A triangulation of the regular octagon.

The structure of this graph depends only on the convexity and the number of vertices of the polygon, and not on its precise geometry. That is, P_n need not be regular for $K_{3,n}$ to be well defined.

McShine and Tetali [41] showed that the *mixing time* of $K_{3,n}$ is $\tau(\varepsilon) = O(n^5 \log(n/\varepsilon))$, and Molloy et al. [43] showed a lower bound of $\tau(\varepsilon) = \Omega(n^{3/2} \log(1/\varepsilon))$.

2.7.2 Quadrangulations

A *quadrangulation* of a point set is a maximal subdivision of the point set into quadrilaterals, where each quadrilateral has all of its vertices in the point set. Consider P_{2n+2} , the regular polygon with $2n + 2$ vertices. We denote by $K_{4,2n+2}$ the graph whose vertex set is the set of all quadrangulations of P_{2n+2} , and whose edges are the flips between quadrilaterals. Here, a flip is defined as follows: each diagonal belongs to two quadrilaterals, which together form a hexagon. Replace the diagonal with one of the other two diagonals in the hexagon. (Thus each diagonal in a quadrangulation can be flipped in two possible ways [12].)

There is a polytope, analogous to the associahedron, known as the *accordiohedron* [30, 4], whose vertices and edges are those of a *subgraph* of $K_{4,2n+2}$. However, we ignore this polytope and just consider the graph $K_{4,n}$.

2.7.3 Counting k -angulations

We refer to a *k -angulation* of a point set as a maximal subdivision of the point set into k -gons, each of whose vertices all belong to the point set. A bijection exists [28] between the k -angulations of $P_{(k-2)n+2}$ and the set of all $k - 1$ -ary plane trees with n internal nodes. We will use the following fact in proving that the random walk on k -angulations fits the framework:

Lemma 2.28. [35, 28] *The number of k -angulations of the convex $(k - 2)n + 2$ -gon is counted by the Fuss-Catalan number*

$$C_{k,n} = \frac{1}{(k-2)n+1} \binom{(k-1)n}{n}.$$

One can show using Stirling's formula, and in particular a result by Robbins [49], that:

Lemma 2.29. For all $k \geq 3$ and $n \geq 1$, $e^{-1/6 \frac{k-2}{k-1}} f(k, n) \leq C_{k,n} \leq e^{1/12} \cdot f(k, n)$, where

$$f(k, n) = \frac{\sqrt{k-1}}{\sqrt{2\pi}((k-2)n)^{3/2}} \cdot \frac{(k-1)^{(k-1)n}}{(k-2)^{(k-2)n}}.$$

2.7.4 k -angulation flip chain

We generalize the associahedron graph K_n as follows:

Definition 2.30. Define the k -angulation flip graph $K_{k,(k-2)n+2}$ as the graph whose vertices are the k -angulations of $P_{(k-2)n+2}$, and whose edges are identified with the flips between k -angulations.

Definition 2.31. Define the k -angulation flip chain as the natural Markov chain whose state space is $K_{k,(k-2)n+2}$.

Theorem 1.5 states that the chain in Definition 2.31 mixes in $O(2^{O(\log^{O(1)} n)})$ time, for all fixed k .

As with the other chains we consider, if one is interested merely in sampling a k -angulation uniformly at random, the feasibility of doing so in polynomial time follows from the existence of the exact formula in Lemma 2.28. However, as previously noted, considerable effort has been devoted to tightening known bounds on the natural Markov chain in the case where $k = 3$ [41, 43], as well as a biased version of the analogous chain on lattice point sets [10, 9]; several recent papers have considered quadrangulations of planar maps [12] and closely related structures [11].

3 $\lambda = 1$: Bounded carving width

To build up to the proof of Theorem 1.1, we first show a weaker result: that the unbiased Glauber dynamics on independent sets mixes rapidly in graphs of bounded carving width. The full proof of Theorem 1.1, even in the unbiased case, requires the non-hierarchical framework.

Consider the independent set flip chain (the hardcore model) as in Definition 2.1. This chain is the natural random walk on what we will call the *independent set Glauber graph*:

Definition 3.1. Given a graph G , let the independent set Glauber graph $\mathcal{M}_{\text{IS}}(G)$ be the graph whose vertices are the independent sets of G , and whose edges are the pairs of independent sets S, S' such that $|S \oplus S'| = 1$.

The following is known, but we give an easy proof:

Lemma 3.2. The independent set Glauber graph in Definition 3.1 is connected.

Proof. Consider the empty independent set \emptyset . Every independent set $S \in V(\mathcal{M}_{\text{IS}}(G))$ has a path of length $|S|$ to \emptyset , formed by removing each vertex in S in arbitrary order. \square

3.1 Partitioning the vertices of $\mathcal{M}_{\text{IS}}(G)$ into classes

The vertices of the Glauber graph $\mathcal{M}_{\text{IS}}(G)$ are subsets of the vertices of an underlying graph G . When G has bounded treewidth, we can choose a small separator X that partitions $V(G) \setminus X$ into two mutually disconnected vertex subsets, A and B , neither of which is too large. Consider the problem of sampling an independent set S from G . Given a separator X for G , partition the independent sets in G into equivalence classes as follows:




Figure 2: Two independent sets in a graph G , belonging to the same class, induced by the restriction of the sets to X .

Definition 3.3. Let $G = (V, E)$ be a graph. Let $\mathcal{M}_{\text{IS}}(G)$ be as in Definition 3.1. Let $X \subseteq V$ be a vertex separator for G . Let $\mathcal{S}_{\text{IS}}(G)$ be the set of equivalence classes of $V(\mathcal{M}_{\text{IS}}(G))$ in which two independent sets S and S' are in the same class if $S \cap X = S' \cap X$. Let $T = S \cap X$, and call the corresponding class $\mathcal{C}_{\text{IS}}(T)$.

See Figure 2 for an example of a partitioning and a class.

Let A and B be the mutually disconnected vertex subsets into which the removal of X partitions $G[V \setminus X]$. Given a fixed independent subset $T \subseteq X$, identify the independent sets in $\mathcal{C}_{\text{IS}}(T)$ with the pairs of the form (S_A, S_B) , where S_A is an independent set in $A \setminus N_A(T)$, and S_B is an independent set in $B \setminus N_B(T)$, where $N_A(T)$ and $N_B(T)$ denote the union of the neighborhoods of vertices in T , in A and B respectively. That is, identify each independent set in $\mathcal{C}_{\text{IS}}(T)$ with a pair of an independent set in A that avoids neighbors of vertices in T , and a similar independent set in B . Consider the two Glauber graphs $\mathcal{M}_{\text{IS}}(A \setminus N_A(T))$ and $\mathcal{M}_{\text{IS}}(B \setminus N_B(T))$, whose vertices are respectively the independent sets in $G[A \setminus N_A(T)]$, and those in $G[B \setminus N_B(T)]$. If two independent sets $S = (S_A, S_B)$ and $S' = (S'_A, S'_B)$ belong to the same class, then a flip exists between S and S' in $\mathcal{M}_{\text{IS}}(G)$ precisely when a flip exists between the restrictions of S and S' to either $\mathcal{M}_{\text{IS}}(A \setminus N_A(T))$ or $\mathcal{M}_{\text{IS}}(B \setminus N_B(T))$ (but not both). See Figure 2. Therefore, each class induces, in $\mathcal{M}_{\text{IS}}(G)$, a subgraph that is isomorphic to a Cartesian product of two smaller Glauber graphs:

Lemma 3.4. Given a graph G and a vertex separator X that partitions $V(G)$ into subgraphs A and B , for every class $T \in \mathcal{S}_{\text{IS}}(G)$,

$$\mathcal{C}_{\text{IS}}(T) \cong \mathcal{M}_{\text{IS}}(A \setminus N_A(T)) \square \mathcal{M}_{\text{IS}}(B \setminus N_B(T)).$$

(Here we identify the class $\mathcal{C}_{\text{IS}}(T)$ with the subgraph it induces in $\mathcal{M}_{\text{IS}}(G)$.)

3.2 Inductive flow construction

As described in Section 3.1, we use a small vertex separator X in G to give a decomposition of $\mathcal{M}_{\text{IS}}(G)$ into subgraphs, each of which has a Cartesian product structure—in which both factor graphs in the product are themselves Glauber graphs. Since Cartesian products preserve flow congestion upper bounds (see Lemma 2.17), this decomposition provides a crucial inductive structure. Namely, we build a multicommodity flow in $\mathcal{M}_{\text{IS}}(G)$ inductively: we assume, for the inductive hypothesis, that a flow with low congestion exists in the Glauber graphs on A and B , then use Lemma 2.17 to give a good flow in each class. Thus for every pair of independent sets S and S' within a given




Figure 3: A schematic view of three classes in the independent set Glauber graph $\mathcal{M}_{\text{IS}}(G)$. The large circles denote classes under the partition described in Section 3.1. The curved arrows illustrate the construction of a flow in $\mathcal{M}_{\text{IS}}(G)$ from an independent set $S \in \mathcal{C}_{\text{IS}}(T)$ to another independent set $S'' \in \mathcal{C}_{\text{IS}}(T)$ —and also to an independent set $S' \in \mathcal{C}_{\text{IS}}(T')$. Here, $\mathcal{C}_{\text{IS}}(T)$ and $\mathcal{C}_{\text{IS}}(T'')$ are adjacent classes in $\mathcal{M}_{\text{IS}}(G)$, connected by a large number of edges, and similarly $\mathcal{C}_{\text{IS}}(T')$ and $\mathcal{C}_{\text{IS}}(T'')$ are adjacent. In Section 3.3 we formalize this flow.




Figure 4: Two independent sets in a graph G : S (left) and S' (right), belonging to distinct classes. S and S' differ by a flip, with the separator X inducing the classes to which the sets belong. S' results from adding v to S . Note that $|S'| < |S|$, since S' excludes those independent sets that contain the vertex u .

class $\mathcal{C}_{\text{IS}}(T)$ of $\mathcal{M}_{\text{IS}}(G)$, we simply use the resulting good internal flow within $\mathcal{C}_{\text{IS}}(T)$ to route the flow between S and S' in our flow for $\mathcal{M}_{\text{IS}}(G)$.

Having constructed a “good” flow (i.e. one with bounded congestion) in each class-induced subgraph $\mathcal{C}_{\text{IS}}(T)$, we then show how to combine these flows with a scheme for routing flow between classes in $\mathcal{M}_{\text{IS}}(G)$. We do so by showing that the edge sets connecting pairs of classes in $\mathcal{M}_{\text{IS}}(G)$ have a useful structure. In particular, we show that every pair of subgraphs sharing at least one edge (flip) is connected by a large number of mutually vertex-disjoint edges—so we can route $S - S'$ flow through $\mathcal{M}_{\text{IS}}(G)$ without sending too much flow through any one edge between a pair of classes. This in turn allows us to route flow throughout the graph without creating too much congestion across any single edge—including across edges within classes. See Figure 3 for a schematic view of the flow we construct.

3.3 Rapid mixing of the independent set flip chain when G has bounded carving width

We establish some properties of $\mathcal{M}_{\text{IS}}(G)$, then sketch a proof that these properties imply bounded congestion. The full proof is in the appendix.

Lemma 3.5. *Let G be a graph with bounded treewidth t , let $\mathcal{M}_{\text{IS}}(G)$ be as in Definition 3.1, and let $\mathcal{S}_{\text{IS}}(G)$ be as in Definition 3.3 with respect to a small balanced separator X with $|X| \leq t + 1$. The*

number of classes in $\mathcal{S}_{\text{IS}}(G)$ is $O(1)$.

Proof. The lemma follows from the fact that $|\mathcal{S}_{\text{IS}}(G)| \leq 2^{|X|} \leq 2^{t+1} = O(1)$, where the first inequality is true because each class is identified with a subset of the vertices in X . \square

Lemma 3.6. *Let G be a graph with bounded treewidth t and bounded degree Δ , let $\mathcal{M}_{\text{IS}}(G)$ be as in Definition 3.1, and let $\mathcal{S}_{\text{IS}}(G)$ be as in Definition 3.3 with respect to a small balanced separator X with $|X| \leq t + 1$. For every pair of classes $\mathcal{C}_{\text{IS}}(T), \mathcal{C}_{\text{IS}}(T') \in \mathcal{S}_{\text{IS}}(G)$, $|\mathcal{C}_{\text{IS}}(T)| = \Theta(1)|\mathcal{C}_{\text{IS}}(T')|$.*

Proof. Consider the class $\mathcal{C}_{\text{IS}}(T_r)$ whose vertex set in X is the empty set, and consider any class $\mathcal{C}_{\text{IS}}(T) \neq \mathcal{C}_{\text{IS}}(T_r)$. $\mathcal{C}_{\text{IS}}(T_r)$ consists of the set of all pairs (S_A, S_B) , where S_A is an independent set in A , and S_B is an independent set in B . $\mathcal{C}_{\text{IS}}(T)$ consists of the set of all pairs (S'_A, S'_B) , where S'_A is an independent set in $A \setminus N_A(T)$, and S'_B is an independent set in $B \setminus N_B(T)$.

Clearly every independent set S'_A in $A \setminus N_A(T)$ is also an independent set in A (and the situation is the same for S'_B), so a trivial injective mapping exists from the sets in $\mathcal{C}_{\text{IS}}(T)$ to the sets in $\mathcal{C}_{\text{IS}}(T_r)$. For the other direction, consider the mapping $f : \mathcal{P}(A) \rightarrow \mathcal{P}(A \setminus N_A(T))$ that sends every independent set $S_A \subseteq A$ to its restriction $S'_A = S_A \setminus N_A(T)$. Because the degree Δ of G is bounded, $|N_A(T)| \leq t\Delta = O(1)$, and thus f is at worst a $2^{t\Delta} = O(1)$ -to-one mapping. This shows that the classes differ in size by a factor of $O(1)$, proving the lemma. \square

Lemma 3.7. *Let G be a graph, let $\mathcal{M}_{\text{IS}}(G)$ be as in Definition 3.1, and let $\mathcal{S}_{\text{IS}}(G)$ be as in Definition 3.3 with respect to a separator X . Let $\mathcal{C}_{\text{IS}}(T), \mathcal{C}_{\text{IS}}(T') \in \mathcal{S}_{\text{IS}}(G)$ be two classes. No independent set in $\mathcal{C}_{\text{IS}}(T)$ has more than $O(1)$ flips to independent sets in $\mathcal{C}_{\text{IS}}(T')$.*

Proof. Each edge (S, S') between independent sets $S \in \mathcal{C}_{\text{IS}}(T)$ and $S' \in \mathcal{C}_{\text{IS}}(T') \neq \mathcal{C}_{\text{IS}}(T)$ consists of flipping a single vertex $v \in X$. It is clear that S has no other flips to independent sets in $\mathcal{C}_{\text{IS}}(T')$. \square

Lemma 3.8. *Let G be a graph with bounded treewidth t and bounded degree Δ , let $\mathcal{M}_{\text{IS}}(G)$ be as in Definition 3.1, and let $\mathcal{S}_{\text{IS}}(G)$ be as in Definition 3.3 with respect to a small balanced separator X with $|X| \leq t + 1$. Let $\mathcal{C}_{\text{IS}}(T), \mathcal{C}_{\text{IS}}(T') \in \mathcal{S}_{\text{IS}}(G)$ be two classes. Suppose there exists at least one flip between an independent set in $\mathcal{C}_{\text{IS}}(T)$ and an independent set in $\mathcal{C}_{\text{IS}}(T')$. Then there exist at least $\Omega(1)|\mathcal{C}_{\text{IS}}(T)|$ flips between independent sets in $\mathcal{C}_{\text{IS}}(T)$ and independent sets in $\mathcal{C}_{\text{IS}}(T')$.*

Proof. T and T' differ by exactly one vertex; call it v . (Or else no flip could exist between $\mathcal{C}_{\text{IS}}(T)$ and $\mathcal{C}_{\text{IS}}(T')$.) Suppose $v \in T$ and $v \notin T'$; then every independent set in $\mathcal{C}_{\text{IS}}(T)$ has a flip to some independent set in $\mathcal{C}_{\text{IS}}(T')$. (See Figure 4.) Thus the number of edges from $\mathcal{C}_{\text{IS}}(T)$ to $\mathcal{C}_{\text{IS}}(T')$ is $|\mathcal{C}_{\text{IS}}(T)|$; the lemma now follows from Lemma 3.6. \square

We are ready to construct our inductive flow.

Lemma 3.9. *Given a graph G with bounded carving width, the natural random walk on the independent set Glauber graph $\mathcal{M}_{\text{IS}}(G)$ has mixing time $\tau(\varepsilon) = O(n^c \log 1/\varepsilon)$, where $c = O(1)$.*

Proof Sketch. The idea of the proof is first to partition $\mathcal{M}_{\text{IS}}(G)$ into classes as described in Definition 3.3. By Lemma 3.4, each class $\mathcal{C}_{\text{IS}}(T) \in \mathcal{S}_{\text{IS}}(G)$ is isomorphic to the Cartesian product

$\mathcal{M}_{\text{IS}}(A \setminus N_A(T)) \square \mathcal{M}_{\text{IS}}(B \setminus N_B(T))$. We make an inductive argument, in which the inductive hypothesis assumes that for each such Cartesian product, the graphs $\mathcal{M}_{\text{IS}}(A \setminus N_A(T))$ and $\mathcal{M}_{\text{IS}}(B \setminus N_B(T))$ have multicommodity flows ρ_A and ρ_B with congestion

$$\rho_{A \max} \leq c^{\log |V(G)| - 1}$$

and

$$\rho_{B \max} \leq c^{\log |V(G)| - 1}$$

respectively, for an appropriate constant c . Lemma 2.17 then implies that $\mathcal{C}_{\text{IS}}(T)$ has a flow ρ_T with congestion

$$\rho_{T \max} \leq c^{\log |V(G)| - 1}.$$

The inductive step is then to combine the ρ_T flows for all of the classes, constructing a multicommodity flow ρ in $\mathcal{M}_{\text{IS}}(G)$ with small congestion. We need to route flow between every pair of independent sets S and S' in $\mathcal{M}_{\text{IS}}(G)$. If S and S' belong to the same class $\mathcal{C}_{\text{IS}}(T)$, this is easy: use the same flow that S uses to send its unit to S' in ρ_T . If $S \in \mathcal{C}_{\text{IS}}(T)$ and $S' \in \mathcal{C}_{\text{IS}}(T') \neq \mathcal{C}_{\text{IS}}(T)$ belong to different classes, we do the following (see Figure 3):

1. Find a path from $\mathcal{C}_{\text{IS}}(T)$ to $\mathcal{C}_{\text{IS}}(T')$, where each pair of consecutive classes on the path share at least one edge. Let this path be

$$\mathcal{C}_{\text{IS}}(T) = \mathcal{C}_{\text{IS}}(T_1), \mathcal{C}_{\text{IS}}(T_2), \dots, \mathcal{C}_{\text{IS}}(T_k) = \mathcal{C}_{\text{IS}}(T').$$

2. Let S send an equal fraction of the S - S' unit (through paths in $\mathcal{C}_{\text{IS}}(T)$) to each independent set $Z \in \mathcal{C}_{\text{IS}}(T)$ that has a neighbor $Y \in \mathcal{C}_{\text{IS}}(T_2)$.
3. For $i = 2, \dots, k - 1$, within $\mathcal{C}_{\text{IS}}(T_i)$, for every independent set Y that receives flow from a neighbor in $\mathcal{C}_{\text{IS}}(T_{i-1})$, let Y send an equal fraction of its S - S' unit (using paths in $\mathcal{C}_{\text{IS}}(T_i)$) to every independent set $Z \in \mathcal{C}_{\text{IS}}(T_i)$ with a neighbor in $\mathcal{C}_{\text{IS}}(T_{i+1})$.
4. For $i = 1, \dots, k - 1$, let each independent set $Z \in \mathcal{C}_{\text{IS}}(T_i)$ having a neighbor $Y \in \mathcal{C}_{\text{IS}}(T_{i+1})$ send its fraction of the $S - S'$ unit to Y across the edge (Z, Y) .
5. Let S' receive an equal fraction of the S - S' unit (through paths in $\mathcal{C}_{\text{IS}}(T')$) from each independent set $Y \in \mathcal{C}_{\text{IS}}(T')$ that has a neighbor in $\mathcal{C}_{\text{IS}}(T_{k-1})$.

We specify in the full proof (Appendix A) how to route the flow at each intermediate step, making use of the existing flows within each class guaranteed by the inductive hypothesis. We then derive upper bounds on the amount of flow resulting from this routing across any given edge within a class, as well as on the amount of flow across each boundary edge e between classes. We show that the latter is $O(N)$, where $N = |V(\mathcal{M}_{\text{IS}}(G))|$, i.e. $\rho(e) = \frac{1}{N}O(N) = O(1)$; we show that the former is at most an $O(1)$ factor times the existing congestion in ρ_T .

This leads to a total congestion of $O(1)^l$, where l is the number of levels of induction. The fact that X is a balanced separator implies that $l = O(\log n)$; the lemma now follows from Lemma 2.12. \square

To prove Theorem 1.1, however, we need to get rid of the assumption that degree is bounded. We address this issue in Section 4. First, however, we distill the observations in the present section into a set of conditions sufficing for rapid mixing—which we then apply to natural flip chains other than the Glauber dynamics on independent sets.

3.4 Abstraction into framework conditions

The observations in Lemmas 3.5 through 3.8 enabled us to construct a flow with polynomially bounded congestion in $\mathcal{M}_{\text{IS}}(G)$. We now abstract these observations into a set of conditions that suffice for rapid mixing. We will then show that the other Glauber graphs we consider satisfy these conditions.

The conditions are, given a connected Glauber graph $\mathcal{M}(G)$, on some set of combinatorial structures over an underlying graph G with n vertices:

1. The vertices of $\mathcal{M}(G)$ can be partitioned into a set \mathcal{S} of classes, where $|\mathcal{S}| = O(1)$.
2. The ratio of the sizes of any two classes in \mathcal{S} is $\Theta(1)$.
3. Given two classes $\mathcal{C}(T), \mathcal{C}(T') \in \mathcal{S}$, no vertex in $\mathcal{C}(T)$ has more than $O(1)$ edges to vertices in $\mathcal{C}(T')$.
4. For every pair of classes that share at least one edge, the number of edges between the two classes is $\Theta(1)$ times the size of each of the two classes.
5. Each class in \mathcal{S} is the Cartesian product of two Glauber graphs $\mathcal{M}(G_1)$ and $\mathcal{M}(G_2)$, each of which can be recursively partitioned in the same way as $\mathcal{M}(G)$.
6. The recursive partitioning mentioned in Condition 5 reaches the base case (graphs with one or zero vertices) in $O(\log n)$ steps.

Conditions 1 through 4 correspond respectively to Lemmas 3.5 through 3.8; Condition 5 corresponds to Lemma 3.4. Condition 6 corresponds to the observation at the end of the proof sketch of Lemma 3.9. Since we only used these conditions in proving the lemma, the following now follows:

Lemma 3.10. *Given a graph $\mathcal{M}(G)$ satisfying the conditions in Section 3.4, the expansion of $\mathcal{M}(G)$ is $\Omega(1/n^c)$, where $c = O(1)$.*

We will use the phrase “non-hierarchical framework” to describe this set of conditions—which apply to the chains we study when the underlying graph G has bounded carving width. In Section 4, we will present an alternative set of conditions—which we will call our “hierarchical framework”—that allows us to handle underlying graphs of unbounded degree, and thus complete the proofs of Theorems 1.1, 1.2, and 1.3.

3.5 q -colorings

We now apply the non-hierarchical framework to q -colorings in graphs of bounded carving width. For reasons that will soon become apparent, we need to generalize to *list colorings*:

Definition 3.11. *A list coloring of a graph $G = (V, E)$, given a function $L : V \rightarrow 2^{[q]}$ assigning a list of colors to each vertex in V , is a coloring of G consistent with L . A partial list coloring is a coloring of some of the vertices of G consistent with L .*

We consider the Glauber graph $\mathcal{M}_{\text{COL}}(G, L)$, defined as follows:

Definition 3.12. *Let the Glauber graph $\mathcal{M}_{\text{COL}}(G, L)$, given an input graph G and a set of colors $[q]$ and a function L as in Definition 3.11, be the graph whose vertices are the list colorings of G consistent with L , and whose edges are the pairs of list colorings C, C' that differ by a color assignment to exactly one vertex $v \in V(G)$.*

The Glauber dynamics in Definition 2.5 is clearly the natural random walk on $\mathcal{M}_{\text{COL}}(G, L)$, with self-loops added in the standard fashion. The following lemma therefore suffices to prove the first claim in Theorem 1.2:

Lemma 3.13. *$\mathcal{M}_{\text{COL}}(G, L)$, defined over a graph G and a list $L : V(G) \rightarrow 2^{[q]}$, with $L(v) \geq \delta(v) + 2$ for every $v \in V(G)$, satisfies the conditions of the non-hierarchical framework whenever G has bounded carving width and q is fixed.*

Proof. We partition $V(\mathcal{M}_{\text{COL}}(G, L))$ into classes induced by a small balanced separator X , where each class is identified with a list coloring T of $G[X]$. This partitioning satisfies Condition 5 since each class $\mathcal{C}_{\text{COL}}(T)$ consists of the tuples of the form (C_A, C_B) , where C_A is a valid list coloring of $G[A]$, and C_B is a valid list coloring of $G[B]$ —with A and B being the mutually disconnected subsets of $V(G)$ resulting from the removal of X . Here, we adjust the list $L(u)$ for each $u \in N_A(X) \cup N_B(X)$, removing from $L(u)$ every color that is assigned to a neighbor of u in X under the coloring C .

It is easy to see that the subproblems on A and B are independent, and that a flip within $\mathcal{C}_{\text{COL}}(T)$ corresponds to a flip within either A or B but not both. Furthermore, the condition that $L(u) \geq \delta(u) + 2$ is preserved even after L is modified, since every color removed from $L(u)$ corresponds to a neighbor of u in X —i.e. a neighbor that is not part of the subproblem on A or B . Condition 5 follows.

Condition 1 follows from the fact that $|X|$ and q are bounded. Condition 2 can be seen from the bounded carving width of G by considering the following mapping $f : V(\mathcal{M}_{\text{COL}}(G, L)) \rightarrow \mathcal{C}_{\text{COL}}(T)$ for any T : given a list coloring $C \in V(\mathcal{M}_{\text{COL}}(G, L))$, let $C' = f(C)$ be the list coloring that (i) agrees with T on its restriction to X , (ii) agrees with C on its assignment of colors to all vertices having no neighbor in X , and (iii) is consistent with both (i) and (ii) on its assignment of colors to neighbors of vertices in X .

(Resolve ambiguity in this mapping via an arbitrary lexicographic ordering on the list colorings of G .) We can always satisfy (iii) because for each $u \in N_A(X) \cup N_B(X)$, we have $|L(u)| \geq \delta(u) + 2$. Condition 4 follows from a similar mapping.

Condition 3 is easy to see from the definition of a flip; Condition 6 follows from the bounded carving width of G . \square

3.6 b -edge covers and b -matchings

For b -edge covers and b -matchings, we now apply the non-hierarchical framework in graphs of bounded carving width. As with independent sets, dealing with unbounded degree in b -edge covers requires the hierarchical framework.

Lemma 3.14. *Given an input graph G of bounded carving width, the Glauber dynamics on b -matchings and on b -edge covers satisfy the conditions of the non-hierarchical framework, when the maximum value of the function b is bounded.*

Proof. The proof for b -edge covers is similar to the proof of Lemma 3.9, with the following modifications.

In defining a b -edge cover, we are selecting subsets of edges instead of vertices. Thus, to define our flip chain on b -edge covers, we modify the flip chain on independent sets in the natural way: dropping or adding edges instead of vertices. The corresponding Glauber graph $\mathcal{M}_{\text{BEC}}(G)$ is connected, since

every b -edge cover has a path in $\mathcal{M}_{\text{BEC}}(G)$ to the trivial b -edge cover (where every edge is selected). We identify each class $\mathcal{C}_{\text{BEC}}(T)$ with the set T of edges chosen incident to vertices in X . Since degree is bounded and $|X| \leq t$, there are $O(1)$ classes, satisfying Condition 1.

Given a class $\mathcal{C}_{\text{BEC}}(T)$, we pass recursively to subproblems on A and B , where we update $b(v)$ for each $v \in A \cup B$ according to the number of edges in T incident to v . That is, for each vertex u selected in T , and for each edge (u, v) with $v \in A$ (similarly $v \in B$), decrement $b(v)$ when passing to the subproblem on A (similarly B). The choices made in the A subproblem and the B subproblem are independent, giving the required Cartesian product structure for Condition 5, and there are still $O(\log n)$ levels of recursion, satisfying Condition 6. For Condition 3, the proof is the same as for independent sets. Conditions 2 and 4 follow from a similar mapping argument to that in the proof of Lemma 3.6.

The proof for b -matchings is similar to that for b -edge covers. □

3.7 Maximal independent sets and maximal b -matchings

The main idea of applying the framework to maximal independent sets and maximal b -matchings is similar to that for independent sets, b -matchings, and b -edge covers, but some adaptation is required: the definition of a flip is somewhat different, and the proof that classes have the required Cartesian product structure has a few more details. We thus defer dealing with these chains to Appendix D.6.2.

4 $\lambda = 1$: Unbounded degree

4.1 Hierarchical framework

We now sketch a set of “hierarchical” framework conditions that guarantee rapid mixing in the case of unbounded degree (when treewidth is bounded). Several of the chains we consider satisfy these conditions so long as the treewidth of the underlying graph is bounded.

In the original framework, we assumed that the classes were approximately the same size. Although all of the Glauber graphs to which we apply this hierarchical framework satisfy this condition in graphs with bounded carving width, this is not the case when the degree is unbounded. Fortunately, in the case of independent sets, partial q -colorings, dominating sets, and b -edge covers, we can solve this problem with some modifications to the framework.

4.2 Independent sets

In the proof of Lemma 3.10, the assumption that the classes were approximately the same size allowed us to argue that even in the worst case, any given class $\mathcal{C}_{\text{IS}}(T)$ can route flow for all pairs of vertices without being too congested, because $\mathcal{C}_{\text{IS}}(T)$ is sufficiently large. Once we discard this assumption, we need to be more explicit in specifying the path through which any given $\mathcal{C}_{\text{IS}}(T)$ routes flow to any given $\mathcal{C}_{\text{IS}}(T')$. Namely, we show that one can engineer the flow so that for any such $\mathcal{C}_{\text{IS}}(T), \mathcal{C}_{\text{IS}}(T')$ pair, every intermediate class $\mathcal{C}_{\text{IS}}(T'')$ that handles flow between sets $S \in \mathcal{C}_{\text{IS}}(T)$ and $S' \in \mathcal{C}_{\text{IS}}(T')$ has a larger cardinality than one of $\mathcal{C}_{\text{IS}}(T)$ or $\mathcal{C}_{\text{IS}}(T')$. This allows us to bound the number of pairs of sets, relative to $|\mathcal{C}_{\text{IS}}(T'')|$, for which $\mathcal{C}_{\text{IS}}(T'')$ carries flow.

To accomplish this, we observe that for any pair of classes $\mathcal{C}_{\text{IS}}(T)$ and $\mathcal{C}_{\text{IS}}(T')$, if there exists *one* flip between an independent set in $\mathcal{C}_{\text{IS}}(T')$ and an independent set in $\mathcal{C}_{\text{IS}}(T)$, then without loss of




Figure 5: Left: a schematic representation of the classes in the independent set Glauber graph and edges between them when degree is unbounded. Right: a class $\mathcal{C}_{\text{IS}}(T)$, with two parents, $\mathcal{C}_{\text{IS}}(P)$ and $\mathcal{C}_{\text{IS}}(P')$, and two children, $\mathcal{C}_{\text{IS}}(C)$ and $\mathcal{C}_{\text{IS}}(C')$. (Classes with larger cardinality are drawn larger.) The parallel edges depict the fact that a child class always has every one of its vertices adjacent to a vertex in a given parent class, and that the edges between any given pair of classes are vertex-disjoint.

generality *every* independent set in $\mathcal{C}_{\text{IS}}(T')$ has a flip to some independent set in $\mathcal{C}_{\text{IS}}(T)$. Namely, this flip consists of dropping some vertex v from $T' \subseteq X$ to obtain T . In this case we call $\mathcal{C}_{\text{IS}}(T)$ a *parent* of $\mathcal{C}_{\text{IS}}(T')$, and $\mathcal{C}_{\text{IS}}(T')$ a *child* of $\mathcal{C}_{\text{IS}}(T)$. See Figure 5. Since the set of these edges is vertex disjoint, this implies that $|\mathcal{C}_{\text{IS}}(T)| \geq |\mathcal{C}_{\text{IS}}(T')|$. In fact, whenever $T \subseteq T'$, we have $|\mathcal{C}_{\text{IS}}(T)| \geq |\mathcal{C}_{\text{IS}}(T')|$.

Thus for any pair of classes $\mathcal{C}_{\text{IS}}(T)$ and $\mathcal{C}_{\text{IS}}(T')$, one can find paths from the two classes to a “common ancestor”, and route flow along these paths, through the common ancestor. Since for every class $\mathcal{C}_{\text{IS}}(T'')$ on this path, either $|\mathcal{C}_{\text{IS}}(T'')| \geq |\mathcal{C}_{\text{IS}}(T)|$ or $|\mathcal{C}_{\text{IS}}(T'')| \geq |\mathcal{C}_{\text{IS}}(T')|$, we are still able to bound the congestion in a fashion similar to the non-hierarchical framework. We make this precise and derive the resulting congestion bounds in Appendix B.

We defer the full description of this hierarchical framework to Appendix B.

5 Dealing with non-independence

The Glauber dynamics on independent sets induces a Glauber graph, $\mathcal{M}_{\text{IS}}(G)$, that behaves well when partitioned into classes. That is, each class $\mathcal{C}_{\text{IS}}(T)$ is isomorphic to the Cartesian product of two Glauber graphs on subgraphs of G . As we will see in Appendix B, the Glauber dynamics on partial q -colorings is similarly well-behaved. Unfortunately, as we will discuss in Appendix D, this does not hold for dominating sets or, in the unbounded-degree case, for b -edge covers. In these problems, the selection $T \subseteq X$ of vertices (or edges) in the separator X with which the class $\mathcal{C}(T)$ is identified imposes constraints on what vertices (or edges) can be chosen in the two subgraphs $A \cup B = G \setminus X$ —and choices in A may invalidate those in B .

In the case of maximal independent sets and maximal b -matchings, the situation is worse: the classes induced by selection of $T \subseteq X$ may not even be internally connected.

We address both of these problems by relaxing the framework condition that each class $\mathcal{C}(T)$ be a Cartesian product of Glauber graphs, and instead require that each class $\mathcal{C}(T)$ be the (not necessarily disjoint) union of Cartesian products of Glauber graphs, satisfying certain conditions. We fully specify this condition, and show that the remaining chains satisfy it, in Appendix D.




Figure 6: Left: a triangular class. Right: a diagonal class. The polygon is depicted as a circle for simplicity of visualization.




Figure 7: Left: the set of edges between two triangular classes. Center: the set of edges between a triangular class and a diagonal class. Right: the set of edges between two diagonal classes.

6 k -angulations

We now return to the k -angulation flip chain discussed in Section 2.7. We will show that a certain partition of the flip graph $K_{k,(k-2)n+2}$ meets the conditions of the non-hierarchical framework. When $k = 3$, this is in fact the same as the partition given by Molloy, Reed, and Steiger [43], which they used to give a *lower* bound on mixing. We generalize this partition to all $k \geq 3$:

6.1 Partition into classes

Definition 6.1. Given a k -gon (or diagonal) T of the regular $(k-2)n+2$ -gon containing the center, identify T with the class $\mathcal{C}_k(T)$ of k -angulations $v \in V(K_{k,(k-2)n+2})$ such that v contains T . Let $\mathcal{S}_{k,(k-2)n+2}$ be the set of all such T .

Remark 6.2. The set $\mathcal{S}_{k,(k-2)n+2}$ is a partition of $V(K_{k,(k-2)n+2})$, because no pair of diagonals or triangles (or diagonal and a triangle) whose endpoints are polygon vertices can contain the origin without crossing.

Definition 6.3. Given classes $\mathcal{C}_k(T), \mathcal{C}_k(T') \in \mathcal{S}_{k,n}$, let $\mathcal{E}(T, T')$ be the set of $K_{k,(k-2)n+2}$ edges with one endpoint in $\mathcal{C}_k(T)$ and one endpoint in $\mathcal{C}_k(T')$.

See Figure 7.

Remark 6.4. *The set of edge sets of the form $\mathcal{E}(T, T')$ is a partition of all flips between pairs of k -angulations in different classes.*

We will show the following in Appendix E:

Lemma 6.5. *The flip graph $K_{k, (k-2)n+2}$, along with the partition $\mathcal{S}_{k, (k-2)n+2}$, meets the conditions of the non-hierarchical framework, except that the $O(1)$ terms described in each condition are replaced with $O(n^{O(1)})$.*

We will prove Theorem 1.5 in Section E. We will use the observation that when there are at most $\log n$ levels of induction, and each application of the inductive step incurs a polynomial congestion factor, the overall resulting congestion is quasipolynomial.

7 Specific upper bounds and discussion of method

7.1 Derivation of upper bounds in main theorems

We now analyze the specific polynomial upper bounds that we obtain from each version of the framework.

In the following bounds, we consider all logarithms to be base two, unless otherwise stated. The $\log n$ terms in the exponents of these bounds come from the balanced separators guaranteed by bounded treewidth. Technically, as we have defined balanced separators in Definition 2.20, one of the two mutually disconnected subgraphs obtained by removing a balanced separator may have size greater than $n/2$. However, one can show [18] that no *connected component* of the resulting disconnected graph has size greater than $n/2$. It is straightforward to modify many of our proofs to account for Cartesian products with multiple factor graphs, by iterating Lemma 2.17. When this is not possible, we will explicitly state the base we use.

We defer the proofs of the following to Appendix A and Appendix B.2 respectively:

Lemma 7.1. *Suppose a Glauber graph $\mathcal{M}(G)$ satisfies the conditions of the non-hierarchical framework. Then the mixing time of the corresponding Glauber dynamics is*

$$O((4N/\mathcal{E}_{\min})^{2 \log n} \cdot \Delta_{\mathcal{M}}^2 \log N),$$

where \mathcal{E}_{\min} is the size of the smallest edge set between adjacent classes.

In the case of q -colorings, tracing the constant factors in the proof of Lemma 3.13, we see that $N/\mathcal{E}_{\min} \leq q^{2\Delta(t+1)}$, that $\Delta_{\mathcal{M}} \leq (q-1)n$, and that $N \leq q^n$. Combining this with Lemma 2.12 and Lemma 2.15 gives the bound claimed in Theorem 1.2.

Lemma 7.2. *Suppose a Glauber graph $\mathcal{M}(G)$ satisfies the conditions of the hierarchical framework. Then the mixing time of the corresponding Glauber dynamics is*

$$O(((2(K+1))^{2 \log n}) \cdot \Delta_{\mathcal{M}}^2 \log N),$$

where $\Delta_{\mathcal{M}}$ is the maximum degree of the Glauber graph $\mathcal{M}(G)$, $n = |V(G)|$, K is the number of classes in the partition, and $N = |V(\mathcal{M}(G))|$.

The unbiased case of the bound in Theorem 1.1 now follows from combining Lemma 7.2 with the observation that for this chain, $\log N \leq n$, $\Delta_{\mathcal{M}} = n$, and $K \leq 2^{t+1}$. Similarly, as we will see, the

bound for the unbiased case of partial q -colorings in Theorem 1.2 will follow from the fact that $K \leq (q+1)^{t+1}$, $\Delta_{\mathcal{M}} \leq qn$, and $N \leq (q+1)^n$, so that

$$\begin{aligned} (2(K+1))^{2 \log n} \Delta_{\mathcal{M}}^2 \log N &= n^{2 \log 2(K+1)} \Delta_{\mathcal{M}}^2 \log N \\ &\leq q^2 \log(q+1) \cdot n^{2(t+2) \log(q+1)+5}. \end{aligned}$$

7.2 Comparison with the projection/restriction technique

In the special cases of the hardcore model, q -colorings, and partial q -colorings—as well as the bounded-carving-width case of b -edge covers and b -matchings, and the case of k -angulations, one could reframe our inductive step in terms of the projection/restriction technique of Jerrum, Son, Tetali, and Vigoda [31]. As we have noted, Heinrich [27] used the projection/restriction technique for q -colorings. The idea is to first partition the Glauber graph into classes, each of which is a Cartesian product of smaller Glauber graphs over an underlying graph at most half the size of the original graph—just as we have done. One then defines a *projection* Markov chain whose states are identified with the classes, where each class has probability mass proportional to its cardinality. The transition probabilities between classes are then proportional to the numbers of edges between classes (up to normalization by the degree of each boundary vertex).

If one then finds a good upper bound on the mixing times of both the projection chain and each of the individual class chains, one can derive a good upper bound on the mixing time of the original chain. One then needs either to incur an $O(1)$ congestion factor at each inductive step of the decomposition, combined with an $O(\log n)$ induction depth—or alternatively to bound a particular quantity γ that describes, essentially, the number of edges between any given class and the rest of the graph.

To find such an upper bound, one could view our “classes” as individual states in the projection chain, with probability masses proportional to their cardinalities. Our construction of flow between classes would become, under this view, a multicommodity flow in the projection chain.

However, constructing the flow in the projection chain still requires showing that each chain satisfies the framework conditions—and in the case of the hardcore model, one needs the hierarchical framework conditions. Furthermore, as formulated by Jerrum, Son, Tetali, and Vigoda [31], the projection/restriction technique requires that the decomposition be a partition of the state space. By contrast, we deal with several chains in which the “classes” overlap or are internally not Cartesian products: namely dominating sets, b -edge covers (when degree is unbounded), maximal independent sets, and maximal b -matchings. For these chains, our flow analysis provides a finer mechanism for dealing with these structural challenges.

Thus we present our construction purely in combinatorial terms—as opposed to considering a flow in a projection state space—for two key reasons: (i) to deal with non-independence as in the third and fourth of our main theorems, and (ii) because we believe our construction makes the graphical analysis of these chains more intuitive.

8 Conclusion and Open Questions

We have developed a framework whose application shows rapid mixing for several natural flip chains on combinatorial structures in graphs of bounded treewidth. However, some work is required in showing that each of the structures satisfies the conditions of the framework. We hope that a more

robust version of the framework can be developed that requires less adaptation for each individual problem.

In particular, all of the structures we have analyzed satisfy the conditions of Courcelle’s theorem, as noted previously. It would be interesting to determine whether the framework can be extended to all structures satisfying these conditions.

The fact that our results hold for all values of $\lambda > 0$ is not especially surprising, as Ioannis Panageas has observed, since the limiting case $\lambda = \infty$ corresponds to the optimization version of each problem, and the case $\lambda = 1$ corresponds to uniform sampling; as stated in the introduction, both of these problems are already known to be fixed-parameter tractable in treewidth. (In fact, as we noted in the introduction, the extension of Courcelle’s theorem, combined with the reduction from sampling to counting, applies to all values of $\lambda > 0$.) Nonetheless, our result does settle a missing case of the mixing question in some generality, through purely combinatorial methods.

Our divide-and-conquer method relies crucially on tree decompositions; it would be interesting to see whether similar methods can be used when treewidth is not bounded but the underlying structure admits some other sparse decomposition—given, say, by the geometry of the structure. For example:

- Similar rapid mixing results exist for *domino tilings* in two dimensions, in which a rectangular region is tiled with dominos of shape 2×1 . The question is open for higher dimensions, although a natural Glauber graph has been examined [21]; could our framework be adapted to the case of thin prisms—identifying classes with, say, all possible choices of a partial tiling through the center of the prism—to obtain mixing results in three dimensions?
- We have used the geometry of convex polygon k -angulations to obtain quasipolynomial mixing (Theorem 1.5). However, as we have noted, McShine and Tetali [41] proved rapid mixing when $k = 3$ using a different analysis. Could a tighter adaptation of our framework yield rapid mixing for all $k \geq 3$?
- As we noted in Section 6, a recent line of research has shown ([10], [57]) that a certain Glauber dynamics on *integer lattice* point set triangulations mixes rapidly for some values of λ , but the case $\lambda = 1$ is still open. Could some variation on our technique be applied to settle the question?

9 Acknowledgements

The authors wish to acknowledge a number of helpful conversations on this topic with Hadi Khodabande and with Milena Mihail. We also acknowledge several similar conversations with Ioannis Panageas, and with Karthik Gajulapalli and Pedro Matias. We acknowledge also some helpful discussion about k -angulations from personal correspondence with Alexandre Stauffer.

A Full proof that non-hierarchical framework conditions imply rapid mixing

Lemma 3.10. *Given a graph $\mathcal{M}(G)$ satisfying the conditions in Section 3.4, the expansion of $\mathcal{M}(G)$ is $\Omega(1/n^c)$, where $c = O(1)$.*

Proof. We fill in the details of the proof sketch given in Section 3.3. First, we state the inductive claim precisely: as stated previously, each class $\mathcal{C}(T)$ is a Cartesian product $\mathcal{M}(G_1) \square \mathcal{M}(G_2)$. The inductive hypothesis is that there exists a constant $c > 0$ such that, for each such $\mathcal{M}(G_1)$, there exists a multicommodity flow ρ_1 in $\mathcal{M}(G_1)$, where $\rho_{1 \max} \leq c^{\log n-1}$, and the same holds for a flow ρ_2 in $\mathcal{M}(G_2)$. Here, $\log n$ is the number of recursive decomposition steps required to reach the base case from $\mathcal{M}(G)$ as in Condition 6; for our problems, n is the number of vertices in the underlying graph.

In the sketch, we described our scheme for routing flow from $u \in \mathcal{C}(T)$ to $v \in \mathcal{C}(T') \neq \mathcal{C}(T)$ (Figure 3). Here we specify how the routing of this flow occurs in further detail. For $i = 2, \dots, k$, let

$$\mathcal{Y}_i = \{Y \in \mathcal{C}(T_i) \mid \exists Z \in \mathcal{C}(T_{i-1}) \text{ s.t. } (Y, Z) \in E(\mathcal{M}(G))\}.$$

For $i = 1, \dots, k-1$, let $\mathcal{Z}_i = N_{\mathcal{C}(T_i)}(\mathcal{Y}_{i+1})$. That is, \mathcal{Z}_i and \mathcal{Y}_{i+1} are the sets of Glauber graph vertices in $\mathcal{C}(T_i)$ and $\mathcal{C}(T_{i+1})$ respectively that lie on the boundary between $\mathcal{C}(T_i)$ and $\mathcal{C}(T_{i+1})$.

As stated in the sketch, by the inductive hypothesis and Lemma 2.17, $\mathcal{C}(T)$ has a flow ρ_T with congestion at most $c^{\log n-1}$. Under our new flow ρ as outlined in the sketch, each Glauber graph vertex S sends a $1/|\mathcal{Z}_1|$ fraction of the $S - S'$ unit to each $Z \in \mathcal{C}(T_1)$; to do so, let S use the same paths it uses to send the $S - Z$ unit in the ρ_T flow. That is, for each edge $e \in E(T_1)$ that carries $\alpha > 0$ units of $S - Z$ flow in ρ_T , let e carry $\alpha/|\mathcal{Z}_1|$ units of $S - S'$ flow in ρ . There are at most N vertices S' to which S must send flow, so the total amount of such flow S sends to S' is at most $N/|\mathcal{Z}_1|$.

Therefore, since the existing flow across each edge in ρ_T is at most $c^{\log n-1}|\mathcal{C}(T)|$, each $e \in E(T_1)$ carries at most an additional $N/|\mathcal{Z}_1| \cdot c^{\log n-1}|\mathcal{C}(T)|$ units. That is, since ρ_T can route one unit of flow from S to every $Z \in \mathcal{Z}_1$ while incurring congestion at most $c^{\log n-1}$, it follows that ρ can route an additional $N/|\mathcal{Z}_1|$ units of flow from S to each $Z \in \mathcal{Z}_1$ while incurring additional congestion at most $N/|\mathcal{Z}_1| \cdot c^{\log n-1}$.

By Conditions 1 through 4, $|\mathcal{Z}_1| = \Omega(N)$, so the total congestion is at most $c^{\log n}$, where c depends only on the constants in Conditions 1 through 4.

The congestion bound for edges in $\mathcal{C}(T')$ is symmetric. Thus it remains to bound the congestion, under ρ , along edges in $\mathcal{C}(T_i)$, for $i = 2, \dots, k-1$. Each $Y \in \mathcal{Y}_i$ needs to send $1/(|\mathcal{Y}_i||\mathcal{Z}_i|)$ $S - S'$ units to each $Z \in \mathcal{Z}_i$, and there are at most N^2 possible such $S - S'$ pairs. Using ρ_{T_i} allows us to route this flow while incurring congestion at most $N^2/(|\mathcal{Y}_i||\mathcal{Z}_i|) \cdot c^{\log n-1}$. Since $|\mathcal{Y}_i| = \Omega(N)$ and $|\mathcal{Z}_i| = \Omega(N)$, this gives congestion $c^{\log n}$.

Finally, we need to specify how each $Z \in \mathcal{Z}_{i-1}$ sends its $S - S'$ fraction to each of its neighbors $Y \in \mathcal{Y}_i$. For simplicity, assume that each Z has only one such neighbor, and that $|\mathcal{Z}_{i-1}| = |\mathcal{Y}_i|$. Then let Z send to its neighbor $Y \in \mathcal{Y}_i$ all of the $S - S'$ flow it has received from vertices in \mathcal{Y}_{i-1} . Summing over all $S - S'$ pairs, this results in each of these boundary edges carrying at most $N^2/|\mathcal{Z}_{i-1}| = O(N)$ flow, for congestion $O(1)$.

Now if some $Z \in \mathcal{Z}_{i-1}$ has multiple neighbors in \mathcal{Y}_i , let Z send an equal fraction of its flow to each such neighbor. To ensure that every $Y \in \mathcal{Y}_i$ carries an equal amount of flow, we must ensure that each $Z \in \mathcal{Z}_{i-1}$ carries an amount of flow proportional to its degree in \mathcal{Y}_{i+1} . To achieve this, let each $Y \in \mathcal{Y}_{i-1}$ send a fraction of its $S - S'$ flow to each $Z \in \mathcal{Z}_{i-1}$ proportional to the degree of Z in \mathcal{Y}_{i+1} .

These modifications clearly produce a valid flow, and they also do not change the congestion incurred at each inductive step by more than an $O(1)$ factor. The lemma follows. \square

We now trace the analysis in the proof of Lemma 3.10 to obtain the bound claimed in Section 7 for the non-hierarchical framework:

Lemma 7.1. *Suppose a Glauber graph $\mathcal{M}(G)$ satisfies the conditions of the non-hierarchical framework. Then the mixing time of the corresponding Glauber dynamics is*

$$O((4N/\mathcal{E}_{\min}))^{2 \log n} \cdot \Delta_{\mathcal{M}}^2 \log N),$$

where \mathcal{E}_{\min} is the size of the smallest edge set between adjacent classes.

Proof. In the construction in the proof of Lemma 3.10, we need to consider the total amount of flow, in the inductive step, that must be handled internally within a class. This includes (i) outbound flow, (ii) inbound flow, and (iii) “through” flow.

For outbound flow, let $\mathcal{E}(T, T')$ denote the set of edges between Glauber graph vertices in $\mathcal{C}(T)$ and those in $\mathcal{C}(T')$. Edges in this edge set carry at most $N|\mathcal{C}(T)|$ units of outbound flow from $\mathcal{C}(T)$ to the rest of the Glauber graph; inbound flow is symmetric. Thus each boundary vertex in $\mathcal{E}(T, T')$ may need to receive (or distribute) at most $N|\mathcal{C}(T)|/(|\mathcal{C}(T)||\mathcal{E}(T, T')|)$ outbound units from (or N inbound units to) each of the rest of the vertices in $\mathcal{C}(T)$. Accounting for both inbound and outbound flow, this increases the amount of flow across each edge in each direction by a factor of at most $2N/(|\mathcal{E}(T, T')|)$.

For “through” flow—that is, flow carried by a class $\mathcal{C}(T)$ originating in some class $\mathcal{C}(T'') \neq \mathcal{C}(T)$ and destined for some class $\mathcal{C}(T''') \neq \mathcal{C}(T)$ —each boundary vertex $S \in \mathcal{E}(T, T')$ may bring in at most $N^2/(|\mathcal{E}(T, T')|)$ units, which it must distribute to the rest of $\mathcal{C}(T)$. Similarly, each boundary vertex may need to carry out the same number of units. This contributes an additional factor of flow at most $2N^2/(|\mathcal{C}(T)||\mathcal{E}(T, T')|)$ across each edge. In the latter expression, we can eliminate an $N/|\mathcal{C}(T)|$ factor using the observation that this term becomes a telescoping product in the induction.

Combining outbound, inbound, and through flow, and taking the product over $\log n$ levels of induction, gives congestion $O((4N/\mathcal{E}_{\min})^{\log n})$. Applying Lemma 2.15 and Lemma 2.12 with the additional $\Delta_{\mathcal{M}}^2 \log N$ factor gives the result claimed. \square

B Hierarchical framework

In this section we complete the proof of the unbiased case of Theorem 1.1 and Theorem 1.2, by fully specifying the hierarchical framework, and showing that the chain on independent sets satisfies the conditions. Fully proving Theorem 1.3 and Theorem 1.4 requires some adaptation of the framework, which we defer to Appendix D.

Recall that in the proof sketch of Lemma 3.10 (Section 3.3), for every pair of Glauber graph vertices $S \in \mathcal{C}(T), S' \in \mathcal{C}(T') \neq \mathcal{C}(T)$, we found a sequence of classes $\mathcal{C}(T) = \mathcal{C}(T_1), \mathcal{C}(T_2), \dots, \mathcal{C}(T_{k-1}), \mathcal{C}(T_k) = \mathcal{C}(T')$, through which to route the $S - S'$ flow. As discussed in Section 4, when degree is unbounded, the classes are no longer nearly the same size, and thus if this sequence is chosen carelessly, some $\mathcal{C}(T_i)$ may carry flow for too many $S - S'$ pairs.

The solution is to choose the sequences carefully. This is possible provided that there exists a partial order $<$ on the classes with a unique maximal element, where $\mathcal{C}(T) > \mathcal{C}(T')$ implies $|\mathcal{C}(T)| \geq |\mathcal{C}(T')|$. Under this condition, we can simply choose our sequence of classes so that for some i with $1 \leq i \leq k$, $|\mathcal{C}(T_1)| \leq |\mathcal{C}(T_2)| \leq \dots \leq |\mathcal{C}(T_i)| \geq |\mathcal{C}(T_{i+1})| \geq \dots \geq |\mathcal{C}(T_{k-1})| \geq |\mathcal{C}(T_k)|$.

B.1 Conditions

The conditions are as follows. Conditions 2 through 4 are new and concern the partial order described above; Condition 1 and Conditions 5 through 7 are as in the non-hierarchical framework.

1. The vertices of $\mathcal{M}(G)$ can be partitioned into a set \mathcal{S} of classes, where $|\mathcal{S}| = O(1)$.
2. There exists a partial order $<$ on the classes in \mathcal{S} , such that whenever $\mathcal{C}(T), \mathcal{C}(T') \in \mathcal{S}$ and $\mathcal{C}(T) > \mathcal{C}(T')$, we have $|\mathcal{C}(T)| \geq |\mathcal{C}(T')|$.
3. The partial order $<$ has a unique maximal element.
4. Whenever an edge exists between vertices in $\mathcal{C}(T)$ and $\mathcal{C}(T')$ with $\mathcal{C}(T) > \mathcal{C}(T')$, the number of such edges is $|\mathcal{C}(T')|$.
5. For every pair of classes $\mathcal{C}(T)$ and $\mathcal{C}(T')$ that share an edge, the maximum degree, in $\mathcal{C}(T)$, of a vertex in $\mathcal{C}(T')$, is $O(1)$, and the maximum degree, in $\mathcal{C}(T')$, of a vertex in $\mathcal{C}(T)$, is $O(1)$.
6. Each class in \mathcal{S} is the Cartesian product of two Glauber graphs $\mathcal{M}(G_1)$ and $\mathcal{M}(G_2)$, each of which can be recursively partitioned in the same way as $\mathcal{M}(G)$.
7. The recursive partitioning mentioned in Condition 6 reaches the base case (graphs with one or zero vertices) in $O(\log n)$ levels of recursion.

B.2 Proof that conditions of the hierarchical framework imply rapid mixing

We are ready to prove the counterpart of Lemma 3.10 for the hierarchical framework, from which the unbiased case of Theorem 1.1 will follow.

Lemma B.1. *Given a graph $\mathcal{M}(G)$ satisfying the conditions in Appendix B.1, the expansion of $\mathcal{M}(G)$ is $\Omega(1/n^c)$, where $c = O(1)$.*

Proof. We use the scheme in the proof of Lemma 3.10, with the following specification: when routing flow from $S \in \mathcal{C}(T)$ to $S' \in \mathcal{C}(T') \neq \mathcal{C}(T)$, we find a sequence of classes $\mathcal{C}(T) = \mathcal{C}(T_1), \mathcal{C}(T_2), \dots, \mathcal{C}(T_{k-1}), \mathcal{C}(T_k) = \mathcal{C}(T')$ as before, where each consecutive pair of classes in the sequence shares an edge in $\mathcal{M}(G)$. In the proof of Lemma 3.10, this sequence was arbitrary; we now require that, under the partial order $<$ in Condition 2, for some $1 \leq i \leq k$, $\mathcal{C}(T_1) < \dots < \mathcal{C}(T_i) > \mathcal{C}(T_{i+1}) > \dots > \mathcal{C}(T_k)$; Condition 3 guarantees that this requirement can be satisfied.

We now bound the resulting congestion. As in the proof of Lemma 3.10 (Appendix A), for $i = 2, \dots, k-1$, the congestion added to edges in $\mathcal{C}(T_i)$ in the inductive step is at most $N^2/(|Y_i||Z_i|) \cdot c^{\log n-1}$. Unfortunately, without assuming that the classes are approximately the same size, we can no longer say that $|Y_i| = \Omega(N)$ or $|Z_i| = \Omega(N)$. Instead, we argue as follows: thanks to the choice of our sequence, for every pair of classes $\mathcal{C}(T)$ and $\mathcal{C}(T')$ that use a given class T_i to route flow, either $|\mathcal{C}(T_i)| \geq |\mathcal{C}(T)|$ (and $|\mathcal{C}(T_i)| \geq |\mathcal{C}(T_{i-1})|$) or $|\mathcal{C}(T_i)| \geq |\mathcal{C}(T')|$ (and $|\mathcal{C}(T_i)| \geq |\mathcal{C}(T_{i+1})|$). Assume the former case without loss of generality. For every pair of classes $\mathcal{C}(T)$ and $\mathcal{C}(T')$ that use the edges between $\mathcal{C}(T_{i-1})$ and $\mathcal{C}(T_i)$, $|\mathcal{C}(T)| \leq |\mathcal{C}(T_{i-1})|$, and therefore the number of pairs S, S' of Glauber graph vertices that use these edges is at most

$$\sum_{T, T': |\mathcal{C}(T)| \leq |\mathcal{C}(T_{i-1})|} |\mathcal{C}(T)||\mathcal{C}(T')| \leq N|\mathcal{S}||\mathcal{C}(T_{i-1})| = O(1)N|\mathcal{C}(T_{i-1})|.$$

Therefore, since there are $|\mathcal{C}(T_{i-1})|$ edges between $\mathcal{C}(T_{i-1})$ and $\mathcal{C}(T_i)$ (by Condition 4), each such edge carries at most $N|\mathcal{S}||\mathcal{C}(T_{i-1})|/|\mathcal{C}(T_{i-1})| = O(N)$ units of flow, giving $O(1)$ congestion.

To bound congestion within $\mathcal{C}(T_i)$, we specify the routing of flow from \mathcal{Y}_i (the set of vertices on the $\mathcal{C}(T_{i-1}), \mathcal{C}(T_i)$ boundary) to \mathcal{Z}_i (the set of vertices on the $\mathcal{C}(T_i), \mathcal{C}(T_{i+1})$ boundary) as follows: first let each $Y \in \mathcal{Y}_i$ send an equal fraction of its flow—of which it receives $O(N)$ units from each of $O(1)$ edges—to *every vertex in $\mathcal{C}(T_i)$* , using ρ_{T_i} to route the flow. Then let each $Z \in \mathcal{Z}_i$ receive its flow similarly from all vertices in $\mathcal{C}(T_i)$. The resulting congestion across each edge is at most

$$(2O(N)/|\mathcal{C}(T_i)|) \cdot c^{\log n - 1} |\mathcal{C}(T_i)| / N \leq c^{\log n},$$

for a constant c . This gives the desired congestion bound, proving the lemma. \square

We now prove Lemma 7.2 by tracing the polynomial factors in the proof of Lemma B.1:

Lemma 7.2. *Suppose a Glauber graph $\mathcal{M}(G)$ satisfies the conditions of the hierarchical framework. Then the mixing time of the corresponding Glauber dynamics is*

$$O(((2(K+1))^{2 \log n}) \cdot \Delta_{\mathcal{M}}^2 \log N),$$

where $\Delta_{\mathcal{M}}$ is the maximum degree of the Glauber graph $\mathcal{M}(G)$, $n = |V(G)|$, K is the number of classes in the partition, and $N = |V(\mathcal{M}(G))|$.

Proof. The analysis is similar to the proof of Lemma 7.1, with the following modifications: each edge set $\mathcal{E}(T, T')$ from $\mathcal{C}(T)$ to a parent $\mathcal{C}(T')$ has $|\mathcal{E}(T, T')| = |\mathcal{C}(T)|$. Therefore, outbound flow along each edge in such an edge set is at most $N|\mathcal{C}(T)|/|\mathcal{E}(T, T')| = N$: each vertex (all vertices in $\mathcal{C}(T)$ are boundary vertices) then receives from each other vertex at most $N/|\mathcal{C}(T)|$ units. As we will show shortly (see analysis of through flow below), edges to children each carry at most $K|\mathcal{C}(T)|$. Thus we will count the flow resulting from edges to children with through flow.

Inbound flow is symmetric. The result is to scale the amount of flow across each edge internal to $\mathcal{C}(T)$ by a factor of $2N/|\mathcal{C}(T)|$.

For through flow (including the outbound flow to children as described above), each boundary vertex in $\mathcal{C}(T)$ carrying flow from (or to) a set of child classes $\{\mathcal{C}(T'_1), \dots, \mathcal{C}(T'_k)\}$ carries at most $\sum_{i=1}^k N|\mathcal{C}(T'_i)|K_i/(|\mathcal{C}(T'_i)|)$ units, where K_i is the number of classes descendent from $\mathcal{C}(T'_i)$, including $\mathcal{C}(T'_i)$ itself. This sum is at most NK . Each boundary vertex carrying flow from (or to) an ancestor similarly carries at most NK units. Thus through flow contributes a factor of $2NK/|\mathcal{C}(T)|$.

The resulting overall congestion is therefore at most

$$(2(K+1))^{\log n},$$

and applying Lemma 2.15 and Lemma 2.12 gives the resulting mixing bound. \square

B.3 Independent sets

We now finish the proof of the unbiased case of Theorem 1.1.

B.3.1 Verification of conditions

To show that the chain on independent sets satisfies the conditions of the hierarchical framework when treewidth is bounded (but degree is unbounded), we first define a partial order $<$ on the classes in $\mathcal{S}_{\text{IS}}(G)$. Recall (Definition 3.3) that these are the classes induced by the separator X in the underlying graph G .

Definition B.2. For $\mathcal{C}_{\text{IS}}(T), \mathcal{C}_{\text{IS}}(T') \in \mathcal{S}_{\text{IS}}(G)$, let $\mathcal{C}_{\text{IS}}(T) < \mathcal{C}_{\text{IS}}(T')$ if $T \subseteq T'$ and $T \neq T'$. Call $\mathcal{C}_{\text{IS}}(T)$ an ancestor of $\mathcal{C}_{\text{IS}}(T')$, and $\mathcal{C}_{\text{IS}}(T')$ a descendant of $\mathcal{C}_{\text{IS}}(T)$. If $\mathcal{C}_{\text{IS}}(T)$ covers $\mathcal{C}_{\text{IS}}(T')$ in this relation, call $\mathcal{C}_{\text{IS}}(T)$ a parent of $\mathcal{C}_{\text{IS}}(T')$, and $\mathcal{C}_{\text{IS}}(T')$ a child of $\mathcal{C}_{\text{IS}}(T)$.

We now prove that the chain on independent sets satisfies the conditions of the hierarchical framework on graphs of bounded treewidth.

Lemma B.3. Given a graph G with fixed treewidth $t - 1$, the hardcore Glauber dynamics on the independent sets of G satisfies the conditions of the hierarchical framework.

Proof. Let $\mathcal{M}_{\text{IS}}(G)$, X , and $\mathcal{S}_{\text{IS}}(G)$ be as previously defined. We have already proven Condition 1 and Conditions 4 through 7 in Lemmas 3.5 through 3.8.

The partial order in Definition B.2 satisfies Condition 2 because for every parent class $\mathcal{C}_{\text{IS}}(T)$ and child class $\mathcal{C}_{\text{IS}}(T')$, the recursive subproblems in the Cartesian product comprising $\mathcal{C}_{\text{IS}}(T')$ are at least as constrained as the subproblems in the product comprising $\mathcal{C}_{\text{IS}}(T)$. That is, $\mathcal{C}_{\text{IS}}(T)$ and $\mathcal{C}_{\text{IS}}(T')$ are each a Cartesian product of two smaller Glauber graphs on the independent sets in subgraphs A_T and B_T of G , and subgraphs $A_{T'}$ and $B_{T'}$ of G respectively. We have $V(A_{T'}) \subseteq V(A_T)$ and $V(B_{T'}) \subseteq V(B_T)$, where the set $V(A_T) \setminus V(A_{T'})$ consists of the vertices in A that have a neighbor in T' but not in T .

Condition 3 follows from the fact that the empty independent set is the unique set that is an ancestor of all other independent sets. □

It now follows by Lemma 2.15 that $\mathcal{M}(G)$ has expansion $\Omega(1/n^{O(1)})$, and Theorem 1.1 follows from this fact and from Lemma 2.12. More precisely, observing that the number of classes in the partition is at most 2^{t+1} and applying Lemma 7.2 gives the bound claimed in Theorem 1.1.

B.4 Partial q -colorings

We now prove the unbiased case of the claim about partial colorings in Theorem 1.2:

Definition B.4. Let $\mathcal{M}_{\text{PCOL}}(G, L)$, given an input graph G and function $L : V(G) \rightarrow 2^{[q]}$, be the graph whose vertices are the partial list colorings of G , and whose edges are the pairs of partial list colorings that differ by the removal or addition of a color assignment to a single vertex.

We show that this Glauber graph satisfies the conditions of the hierarchical framework:

Lemma B.5. Given a graph G with bounded carving width and list function $L : V(G) \rightarrow 2^{[q]}$, where $q \geq \Delta + 2$ is fixed and $L(v) \geq \delta(v) + 2$ for all $v \in V(G)$, the Glauber graph $\mathcal{M}_{\text{PCOL}}(G, L)$ has expansion $\Omega(1/n^c)$, where $c = O(1)$.

Proof. The partitioning is the same as in the proof of Lemma 3.13, except that we allow each class to be identified with a partial list coloring of X . Condition 1, Condition 5, Condition 6, and

Condition 7 can be easily seen as before. For Conditions 2 and 3, the partial order is analogous to the partial order for independent sets: given partial list colorings C and C' of X , let C be a parent of C' if C and C' agree except for a single vertex to which C' assigns a coloring and C does not. Condition 4 is easy to see from this definition. The lemma follows. \square

We obtain the bound in Theorem 1.2 via the observations in Section 7.

C All $\lambda > 0$

Until now, we have only considered the unbiased versions of our chains. In this section we complete the proof of Theorem 1.1, for arbitrary fixed $\lambda > 0$. To do so, we need to introduce the standard notion of *conductance* [53], which extends the definition of expansion in the natural way to the setting of a weighted graph.

C.1 Preliminaries revisited

The conductance is defined with respect to a *stationary* distribution π induced by a random walk. The stationary distribution is the distribution to which the random walk converges in the limit. The convergence requires mild conditions: (i) that walk be *ergodic*, meaning that the Glauber graph is connected; (ii) that the walk be *reversible*; and (iii) that the walk be *lazy*.

Laziness means that with constant probability the walk stays at the current vertex at any step; reversibility means that for every pair of sets $S, S' \in \mathcal{M}_{\text{IS}}(G)$, we have

$$\pi(S)P(S, S') = \pi(S')P(S', S),$$

where $P(S, S')$ denotes the probability that $X_{t+1} = S'$, given that $X_t = S$.

The Glauber dynamics on independent sets is known to satisfy these conditions, and it is easy to show that our other Glauber dynamics satisfy them as well.

In the case of the Glauber dynamics on independent sets, the stationary distribution π evaluates to

$$\pi(S) = \lambda^{|S|} / Z(\mathcal{M}_{\text{IS}}(G)),$$

where for each of our Glauber graphs $\mathcal{M}(G)$,

$$Z(\mathcal{M}(G)) = \sum_{S \subseteq V(\mathcal{M}(G))} \lambda^{|S|}$$

is the normalizing constant. For all independent sets S in G , and for all S' such that $|S \setminus S'| = 1$, it is easy to see that

$$\pi(S)P(S, S') = \pi(S')P(S', S) = \left(\frac{1}{n(1+\lambda)}\right) \left(\frac{\lambda^{|S|}}{Z(\mathcal{M}_{\text{IS}}(G))}\right),$$

where $n = |V(G)|$.

For dominating sets and partial q -colorings, we define the same distribution; for b -edge covers we define the analogous distribution over edges.

Remark C.1. For each of our Glauber graphs $\mathcal{M}(G)$, the probability transition function $P(S, S')$, viewed as a matrix, is in fact the adjacency matrix of an edge-weighted version of $\mathcal{M}(G)$, ignoring self loops.

That is:

Definition C.2. Given a Glauber graph $\mathcal{M}(G)$ and a Markov chain on $\mathcal{M}(G)$ with stationary distribution π and probability transition function P , assign the weight $\pi(S)$ to each vertex S of $\mathcal{M}(G)$, and assign the weight $Q(S, S') = \pi(S)P(S, S')$ to each edge (S, S') .

Definition C.3. Extend the definition of a Cartesian graph product given in Definition 2.13 to the weighted graphs described in this section, so that for vertices $g \in V(G), h \in V(H)$, the weight of the tuple $(g, h) \in V(G \square H)$ is $\pi(g, h) = \pi_G(g)\pi_H(h)$, where π_G and π_H are the vertex weight functions for G and H respectively. Let the weight of each edge e between (g, h) and (g', h') be

$$Q(e) = \pi_H(h)(\Delta_G Q_G(g, g')) / (\Delta_G + \Delta_H),$$

if $g \neq g'$ and $h = h'$, and

$$Q(e) = \pi_G(g)(\Delta_H Q_H(h, h')) / (\Delta_G + \Delta_H),$$

if $g = g'$ and $h \neq h'$, where Q_G and Q_H are the edge weight functions for G and H , and Δ_G and Δ_H are the maximum degrees of G and H .

For the self loop $e = ((g, h), (g, h))$, let

$$Q(e) = \pi(g, h) - \sum_{(g'', h): g'' \neq g} Q((g, h), (g'', h)) - \sum_{(g, h''): h'' \neq h} Q((g, h), (g, h'')),$$

Lemma C.4. Given Definition C.3, the stationary distribution π in the discussion leading to Remark C.1, and the resulting vertex and edge weights as in Definition C.2, for each of our Glauber graphs $\mathcal{M}(G)$ and for each class $\mathcal{C}(T) \cong \mathcal{M}(A) \square \mathcal{M}(B)$, and for each $S \in V(\mathcal{M}(A)), S' \in V(\mathcal{M}(B))$, the following facts hold:

1. The vertex weight of $S \cup S' \cup T$ in $\mathcal{M}(G)$ is equal to

$$\pi_{\mathcal{M}(G)}(S \cup S' \cup T) = \pi_{\mathcal{C}(T)}(S, S') \pi_{\mathcal{M}(G)}(\mathcal{C}(T)),$$

where $\pi_{\mathcal{M}(G)}(\mathcal{C}(T))$ is defined as $\sum_{U \in \mathcal{C}(T)} \pi_{\mathcal{M}(G)}(U)$, and

2. For all S'' with $|S \setminus S''| = 1$, the weight in $\mathcal{M}(G)$ of the edge e between $S \cup S' \cup T$ and $S'' \cup S' \cup T$ is

$$Q_{\mathcal{M}(G)}(e) = Q_{\mathcal{C}(T)}(e) \pi_{\mathcal{M}(G)}(\mathcal{C}(T)) \frac{|V(A)| + |V(B)|}{|V(G)|}.$$

Proof. We have

$$\frac{\lambda^{|S|+|S'|+|T|}}{Z(\mathcal{M}(G))} = \pi_{\mathcal{M}(A)}(S) \pi_{\mathcal{M}(B)}(S') \lambda^{|T|} \cdot \frac{Z(\mathcal{M}(A))Z(\mathcal{M}(B))}{Z(\mathcal{M}(G))} = \pi_{\mathcal{C}(T)}(S, S') \pi_{\mathcal{M}(G)}(\mathcal{C}(T)),$$

and

$$Q(e) = \frac{1}{|V(G)|(\lambda + 1)} \cdot \frac{\lambda^{|S|+|S'|+|T|}}{Z(\mathcal{M}(G))} = Q_{\mathcal{M}(A)}(S, S'') \pi_{\mathcal{M}(B)}(S') \lambda^{|T|} \cdot \frac{Z(\mathcal{M}(A))Z(\mathcal{M}(B))}{Z(\mathcal{M}(G))} \cdot \frac{|V(A)|}{|V(G)|}$$

$$\begin{aligned}
&= Q_{\mathcal{C}(T)}(e)\lambda^{|\mathcal{T}|} \cdot \frac{Z(\mathcal{M}(A))Z(\mathcal{M}(B))}{Z(\mathcal{M}(G))} \cdot \frac{|V(A)| + |V(B)|}{|V(G)|} \\
&= Q_{\mathcal{C}(T)}(e)\pi_{\mathcal{M}(G)}(\mathcal{C}(T)) \frac{|V(A)| + |V(B)|}{|V(G)|}.
\end{aligned}$$

□

Definition C.5. Given a lazy, reversible, ergodic random walk on a weighted graph $\mathcal{M} = (\mathcal{V}, \mathcal{E})$ with stationary distribution π and probability matrix $P : \mathcal{V} \times \mathcal{V} \rightarrow [0, 1]$, the conductance is the quantity

$$\phi(\mathcal{M}) = \min_{\mathcal{S} \subseteq \mathcal{V}: 0 < \pi(\mathcal{S}) \leq 1/2} \frac{Q(\mathcal{S}, \mathcal{V} \setminus \mathcal{S})}{\pi(\mathcal{S})},$$

where for sets $\mathcal{S}, \mathcal{S}' \subseteq \mathcal{V}$,

$$\pi(\mathcal{S}) = \sum_{S \in \mathcal{S}} \pi(S),$$

and

$$Q(\mathcal{S}, \mathcal{S}') = \sum_{S \in \mathcal{S}, S' \in \mathcal{S}'} Q(S, S'),$$

and where $Q(S, S') = \pi(S)P(S, S')$ given $S, S' \in \mathcal{V}$.

We now extend the definitions of multicommodity flows and congestion:

Definition C.6. Let a multicommodity flow ρ in a graph $\mathcal{M} = (\mathcal{V}, \mathcal{E})$ be defined as in Definition 2.14, except that each pair of vertices $S, S' \in \mathcal{V}$ exchanges $\pi(S)\pi(S')$ units of flow in each direction. Let

$$\rho_{\max} = \sum_{e \in \mathcal{E}} \frac{\rho(e)}{Q(e)}.$$

The following generalizations of Lemma 2.15 and Lemma 2.12 relate the conductance, congestion, and mixing time [53]:

Theorem C.7. Given a multicommodity flow ρ in a graph \mathcal{M} , the conductance ϕ satisfies $\phi \geq 1/(2\rho)$.

Theorem C.8. The mixing time of a Markov chain with state space Ω , stationary distribution π^* , and conductance at least ϕ is at most

$$\tau(\varepsilon) = O(\phi^{-2} \log(1/(\pi_{\min}^* \cdot \varepsilon))),$$

where

$$\pi_{\min}^* = \min_{u \in \Omega} \pi^*(u).$$

C.2 Analysis of flow construction

We now complete the proof of Theorem 1.1. It suffices to show the following lemma:

Lemma C.9. The flow ρ constructed in $\mathcal{M}_{\text{IS}}(G)$ in the proof of Lemma B.1, adjusted so that $\mathcal{M}_{\text{IS}}(G)$ is weighted according to the parameter $\lambda > 0$, and so that each pair of sets S, S' exchanges $\pi(S)\pi(S')$ units of flow as in Definition C.6, results in a congestion factor gain of at most $\rho_{\max} = O(1)$ at each of the $O(\log n)$ levels of induction, resulting in at most polynomial overall congestion. The same holds for the flip chain on partial q -colorings.

Proof. We use the same inductive argument, with the following adjustment: if $\mathcal{C}(T)$ is a descendant of $\mathcal{C}(T_{i-1})$, and $\mathcal{C}(T_{i-1})$ is a child of $\mathcal{C}(T_i)$, where $\mathcal{C}(T)$ uses the edges between $\mathcal{C}(T_{i-1})$ and $\mathcal{C}(T_i)$ to send flow to $\mathcal{C}(T)$, then distribute this flow as before across these edges, but now let each edge carry flow in proportion to its weight. We have $\pi(\mathcal{C}(T)) = O(1)\pi(\mathcal{C}(T_{i-1}))$ —because for every independent set $S \in \mathcal{C}(T)$ there exists a distinct independent set $S' \in \mathcal{C}(T_{i-1})$ with $\pi(S') = (1/\lambda^{|\mathcal{C}(T) \setminus T_{i-1}|})\pi(S)$, namely $S' = S \setminus (T \setminus T_{i-1})$.

Each edge (S, S') with $S \in \mathcal{C}(T_i)$, $S' \in \mathcal{C}(T_{i-1})$, satisfies $Q(S, S') = \Omega(\pi(S)/n)$ (where the constant-factor difference depends on λ). Thus the congestion along these edges is still $O(n)$. We then allow each vertex $S \in \mathcal{C}(T_i)$, having received at most $O(\pi(S))$ units along each of $O(1)$ incoming edges from child classes, to distribute these units to all other vertices in $\mathcal{C}(T_i)$ according to their weight. That is, let S send $O(\pi(S)\pi(S'')/\pi(\mathcal{C}(T_i)))$ units to each $S'' \in \mathcal{C}(T_i)$. By the inductive hypothesis and Lemma 2.17, a flow ρ_{T_i} already exists under which S sends $\pi(S)\pi(S'')/(\pi(\mathcal{C}(T_i)))^2$ units to S'' at a congestion cost of $O(nc^{\log n-1})$, for appropriate constant c . Thus letting S send $O(\pi(S)\pi(S'')/\pi(\mathcal{C}(T_i)))$ units to S'' reduces the amount sent across each edge by at least a factor of $1/\pi(\mathcal{C}(T_i))$, while the weight of each edge increases when passing from the factor graphs of $\mathcal{C}(T_i)$ to $\mathcal{M}(G)$ by at most the same factor—up to the change in degree of the flip graph—by Lemma C.4. This gives congestion cost at most $O(nc^{\log n})$. \square

C.2.1 Specific polynomial bounds

We now revisit the discussion in Section 7. In Theorem 1.1, the $1 + \log \hat{\lambda}$ term in the exponent comes from observing that, in Lemma 7.2, we can replace the $K + 1$ term with $K\lambda^{t+1} + 1$ —since in the proof of Lemma 7.2, this is the factor by which the flow carried into a class from a child class increases when adjusting for the weights induced by the parameter λ . A similar analysis gives the result for partial q -colorings in Theorem 1.2.

D Dealing with non-independence

In this section we complete the proofs of Theorem 1.3 and Theorem 1.4. Showing that the framework applies to these chains requires relaxing one of the framework conditions, as described in Section 5. For both theorems, the principal problem is that when attempting to partition the Glauber graph into classes as we did for independent sets, the resulting classes are not isomorphic to Cartesian products of Glauber graphs. For instance, in the case of b -edge covers, we wish to identify a class of b -edge covers with the set T of edges selected within the subgraph of G induced by the separator X . Unfortunately, the resulting subproblems on A and B (where, as before, $A \cup B = V(G) \setminus X$) are not independent. This is because for each vertex $x \in X$, the sum of the number of incident edges selected in A and those in B must be at least $b(x)$, so the choices made in A depend on those made in B , and vice versa.

The solution is as follows: we divide each class into smaller (not necessarily disjoint) “subclasses”, each of which is a Cartesian product of smaller Glauber graphs on b -edge covers. We detail this in Appendix D.2.

We encounter a similar problem in the case of dominating sets, with an additional challenge that will require us to generalize the definition of a dominating set into what we call the “constrained Steiner dominating set” problem. We give the full details in Appendix D.3.

For maximal independent sets and maximal b -matchings (Appendix D.6), the non-hierarchical framework is more natural, as we require bounded degree. The challenge is twofold: first, we need

to define the Glauber graphs and show that they are connected. Secondly, we need to deal with non-independence as with b -edge covers and dominating sets—with the additional challenge, as we will see, that the classes are not necessarily internally connected.

D.1 Framework relaxation to allow non-independence

Lemma D.1. *Suppose a Glauber graph $\mathcal{M}(G)$ satisfies the conditions of the hierarchical framework in Appendix B.1, except for Condition 6. Suppose further that each class $\mathcal{C}(T) \in \mathcal{S}$ is the union of at most $O(1)$ subclasses $\mathcal{C}(T) = \mathcal{C}(T_1) \cup \mathcal{C}(T_2) \cup \dots \cup \mathcal{C}(T_k)$, where for $i = 1, \dots, k-1$:*

1. $|\mathcal{C}(T_i)| = \Theta(1)|\mathcal{C}(T_{i+1})|$, and
2. $\mathcal{C}(T_i)$ and $\mathcal{C}(T_{i+1})$ share at least $\Omega(1)|\mathcal{C}(T_i)| = \Omega(1)|\mathcal{C}(T_{i+1})|$ vertices.

Suppose further that for $i = 1, \dots, k$, $\mathcal{C}(T_i)$ is isomorphic to the Cartesian product of two Glauber graphs $\mathcal{M}(G_1)$ and $\mathcal{M}(G_2)$, each of which can be recursively partitioned in the same way as $\mathcal{M}(G)$.

Then the expansion of $\mathcal{M}(G)$ is $\Omega(1/n^c)$, where $c = O(1)$.

Proof. It suffices to construct a multicommodity flow among the subclasses in $\mathcal{S}(T)$ and bound its congestion. By the inductive hypothesis and Lemma 2.17, each subclass $\mathcal{C}(T_i)$ has an internal flow ρ_{T_i} with congestion $\rho_{T_i \max} \leq c^{\log n-1}$. We would like to derive a flow ρ_T with congestion $\rho_{T \max} \leq c^{\log n-1}$; this will allow the rest of the proof of Lemma B.1 to be applied.

The solution is to follow the proof sketch of Lemma 3.9: for Glauber graph vertices $S \in \mathcal{C}(T_i), S' \in \mathcal{C}(T_j) \neq \mathcal{C}(T_i)$, send the $S-S'$ flow through the classes $\mathcal{C}(T_{i+1}), \mathcal{C}(T_{i+1}), \dots, \mathcal{C}(T_{j-1})$. For $l = i, \dots, j$, let \mathcal{Y}_l and \mathcal{Z}_l be as in the proof of Lemma 3.9, except that $Z_l = Y_{l+1}$. That is, the boundary vertices in consecutive pairs of classes on the path are shared between the two classes. The routing of flow within each class on the path is the same as in Lemma 3.9.

The resulting congestion bound is the same as in Lemma 3.9. The only concern is that since the subclasses may not be disjoint, each edge within a subclass may incur congestion from multiple steps on the path. However, because the number of classes is $O(1)$, there are $O(1)$ such steps, and thus the factor by which this increases congestion is $O(1)$. \square

D.2 b -edge covers in the relaxed hierarchical framework

To finish the proof of Theorem 1.3, it now suffices to show that the chains on dominating sets and b -edge covers satisfy the conditions of the hierarchical framework when treewidth is bounded. We begin with b -edge covers.

Let $\mathcal{C}_{\text{BEC}}(T)$ be defined as in Section 3.6, with the following modification: define each class $\mathcal{C}_{\text{BEC}}(T)$ so that T is identified with the set of selected edges *both of whose endpoints* are in X , instead of including all edges incident to vertices in X .

We now divide each class $\mathcal{C}_{\text{BEC}}(T)$ into subclasses. For each $x \in X$, let $\delta_T(x)$ be the number of edges incident to x (from neighbors in X) that are selected in T . Let $b'(x) = b(x) - \delta_T(x)$, i.e. decrease $b(x)$ by the number of edges incident to x selected in X . Clearly for each $x \in X$, every valid b -edge cover in T includes numbers of edges from neighbors in A and B that sum to at least $b'(x)$.

We will define a subclass of $\mathcal{C}_{\text{BEC}}(T)$ for each possible assignment of b -values to the vertices in X in the subproblems on A and B . (The number of these subclasses, since $|X| = O(1)$ and b is bounded, is still $O(1)$.)

Formally:

Definition D.2. Define functions β and $\bar{\beta}$ as any assignments of b -values, in the subproblems on A and B respectively, to all vertices $x \in X$, such that the β and $\bar{\beta}$ values sum to $b'(x)$ for each x .

There are many degrees of freedom in defining β . Consider each possible choice of β and $\bar{\beta}$.

Definition D.3. Define the subclass $\mathcal{C}_{\text{BEC}}(T_\beta)$ as the set of all b -edge covers that consist of a β -edge cover in A and a $\bar{\beta}$ -edge cover in B .

That is, in class $\mathcal{C}_{\text{BEC}}(T_\beta)$, for each x , the number of incident edges selected in A is at least $\beta(x)$, and the number of incident edges in B is at least $\bar{\beta}(x)$.

Each of these subclasses $\mathcal{C}_{\text{BEC}}(T_\beta)$ is a Cartesian product of b -edge cover Glauber graphs, over subgraphs A and B of G , and thus internally has a good flow ρ_β ; thus it suffices to combine flows within these subclasses together to design a flow ρ_T in T . We can then apply the hierarchical framework to obtain the desired flow in $\mathcal{M}_{\text{BEC}}(G)$.

Lemma D.4. Given a graph G and corresponding Glauber graph $\mathcal{M}_{\text{BEC}}(G)$, each class $\mathcal{C}_{\text{BEC}}(T)$ of b -edge covers in $\mathcal{M}_{\text{BEC}}(G)$ satisfies the conditions of Lemma D.1.

Proof. The number of subclasses is clearly $O(1)$. The subclasses are also all within an $O(1)$ size factor of one another. To see this, compare $|\mathcal{C}_{\text{BEC}}(T)|$ and $|\mathcal{C}_{\text{BEC}}(T_\beta)|$, for any β . Fix some lexicographic ordering of the edges of G . For every b -edge cover $S \in \mathcal{C}_{\text{BEC}}(T)$, there exists a b -edge cover $S' \in \mathcal{C}_{\text{BEC}}(T_\beta)$ that includes the lexicographically first $\beta(x)$ edges in A incident to x , for each $x \in X$, and also includes the first $\bar{\beta}(x)$ edges in B incident to x . (Let S' agree with S on all other edges.) Clearly this is a 2^b -to-1 mapping, i.e. an $O(1)$ -to-1 mapping.

Finally, every pair of subclasses overlaps in at least $\Omega(1)|\mathcal{C}_{\text{BEC}}(T)|$ vertices: consider the set of all b -edge covers in $\mathcal{C}_{\text{BEC}}(T)$ in which for each $x \in X$, x has $\min\{b'(x), \delta_A(x)\}$ incident edges selected in A , and $\min\{b'(x), \delta_B(x)\}$ incident edges selected in B . The number of such b -edge covers is $\Omega(1)|\mathcal{C}_{\text{BEC}}(T)|$, by similar reasoning to the above; furthermore, every pair of subclasses of $\mathcal{C}_{\text{BEC}}(T)$ clearly contains this set of b -edge covers. The lemma follows. \square

The rest of the hierarchical framework conditions are easy to verify, and thus the result in Theorem 1.3 for the unbiased case of b -edge covers follows. The specific bound follows from the following observations: first, constructing the flow within a class $\mathcal{C}_{\text{BEC}}(T)$ incurs a factor of

$$4L|\mathcal{C}_{\text{BEC}}(T)|/(\nu(\min_{\beta} |\mathcal{C}_{\text{BEC}}(T_\beta)|)),$$

where ν is the minimum fraction of vertices shared by a pair of adjacent classes whose intersection is used in the flow, L is the maximum number of subclasses that a vertex can belong to, and $\min_{\beta} |\mathcal{C}_{\text{BEC}}(T_\beta)|$ is the smallest subclass of $\mathcal{C}_{\text{BEC}}(T)$. L is at most the number of subclasses, which is upper-bounded by $(b+1)^{t+1}$; the smallest subclass has size at least $|\mathcal{C}_{\text{BEC}}(T)|/2^{b(t+1)}$; and $\nu \geq 1/2$. (The latter two facts follow readily from observations made in the proof of Lemma D.4.)

Thus the construction of the flow within $\mathcal{C}_{\text{BEC}}(T)$ incurs a factor of at most $8(b+1)^{t+1} \cdot 2^{b(t+1)}$. In the biased chain the overlap between adjacent subclasses is at least $\lambda/(1+\lambda)$ instead of $1/2$, and

we need to adjust the ratio $|\mathcal{C}_{\text{BEC}}(T)|/|\mathcal{C}_{\text{BEC}}(T_\beta)|$ by a factor of $\hat{\lambda}^{b(t+1)}$. Therefore this expression becomes

$$4 \frac{1+\lambda}{\lambda} (b+1)^{t+1} \cdot (2\hat{\lambda})^{b(t+1)}.$$

The rest of the inductive step is as in the case of independent sets, i.e. we apply Lemma 7.2, using $K \leq 2^{t(t+1)/2}$, $\Delta_{\mathcal{M}} \leq m$, and $N \leq 2^m$. (In the weighted case, $\pi_{\min}^* \leq (2\hat{\lambda})^m$. When considering K in the application of Lemma 7.2, we need to weight K by a factor of $\hat{\lambda}^{t(t+1)/2}$.)

Thus we obtain an additional factor of

$$2(K\hat{\lambda}^{t(t+1)/2} + 1) \leq 2((2\hat{\lambda})^{t(t+1)/2+1}).$$

Altogether, these two flow constructions combined, in each iteration, result in a factor of at most

$$8 \left(\frac{1+\lambda}{\lambda} \right) (b+1)^{t+1} \cdot (2\hat{\lambda})^{(t+1)(b+t/2)+1}.$$

The resulting mixing time is therefore at most

$$O(((1+\hat{\lambda})\hat{\lambda})^2 m^3 (\log \hat{\lambda} + 1) n^{2(3+\log(1+\lambda)) - \log(\lambda) + (t+1)\log(b+1) + ((t+1)(b+t/2)+1)(1+\log \hat{\lambda})}).$$

We have ignored one detail: technically the number of levels of induction is greater than $\log n$, because the $t+1$ vertices in the separator are included in the independent subproblems within each subclass. Furthermore, we cannot assume that we have two connected components of size at most $|V(G)|/2$ at each level of decomposition, so the base of the log is $3/2$ and not 2 .

However, for every $\varepsilon < 1/2$, we have for all $n \geq (t+1)/\varepsilon$ that $2n/3 + t + 1 \leq n(2/3 + \varepsilon)$. Thus at the cost of a base case for the induction of $(2\hat{\lambda})^{((t+1)/\varepsilon)^2}$, we adjust the $\log n$ exponent in the congestion term to $\log_{1/(2/3+\varepsilon)} n$. Letting $\varepsilon = 1/6$, we obtain the mixing bound claimed in Theorem 1.3, namely

$$O \left(((2\hat{\lambda})^{36(t+1)^2})^2 \left(\frac{\hat{\lambda}}{(1+\hat{\lambda})} \right)^2 m^3 (\log \hat{\lambda} + 1) n^{\frac{2(3+\log(1+\lambda)) - \log(\lambda) + (t+1)\log(b+1) + ((t+1)(b+t/2)+1)(1+\log \hat{\lambda})}{\log(6/5)}} \right).$$

D.3 Dominating sets in the relaxed hierarchical framework

To finish the proof of Theorem 1.3 in the unbiased case, we now deal with dominating sets.

As with b -edge covers, defining classes in the same way as in the case of independent sets does not result in Cartesian products of dominating set Glauber graphs, because it may be that some vertices in X are not dominated by vertices in $T \cap X$; these vertices must then be dominated by vertices in A or in B .

Therefore, to preserve the recursive structure of the problem and thus complete the proof of Theorem 1.3, we define the *constrained Steiner dominating set problem* as a generalization of the dominating set problem, in which there are three types of vertices:

1. “normal” vertices, which must be dominated and may be selected in a dominating set,

2. “Steiner” vertices, which need not be dominated and may be selected, and
3. “forbidden” vertices, which must be dominated and must not be selected.

Now, we let X be a balanced vertex separator in G as before. We would like to define each class $\mathcal{C}_{\text{DOM}}(T)$ by a choice of vertices in X . In the resulting subproblem in A (similarly B), we then designate each vertex $v \in A \cup B$ having a neighbor selected in $T \subseteq X$ as Steiner. However, there may be vertices in X that are not selected or dominated by any vertex in T . To obtain a valid dominating set, some neighbor of each such vertex w must be chosen in either A or B . Thus we have non-independent subproblems, which ruins the Cartesian product structure needed for the divide-and-conquer argument. To resolve this non-independence, we divide $\mathcal{C}_{\text{DOM}}(T)$ into subclasses as follows:

Definition D.5. *Given a graph G , separator X , and CSDS Glauber graph $\mathcal{M}_{\text{DOM}}(G)$, and class $\mathcal{C}_{\text{DOM}}(T)$ of CSDS’s in $\mathcal{M}_{\text{DOM}}(G)$, let U be the set of undominated vertices in X in class $\mathcal{C}_{\text{DOM}}(T)$. For each subset $W \subseteq U$, let the subclass $\mathcal{C}_{\text{DOM}}(T_W)$ be the set of all CSDS’s that consist of a union of a CSDS on $A \cup W$, and a CSDS on $B \cup U \setminus W$ —in which each $w \in W$ is a forbidden vertex in the A subproblem, and each $\bar{w} \in U \setminus W$ is a forbidden vertex in the B subproblem.*

There are at most $2^t = O(1)$ such subclasses. Technically, as with b -edge covers, these are not equivalence classes, as some CSDS solutions may belong to multiple classes. We will address this shortly, but first we specify how to route flow among the subclasses within $\mathcal{C}_{\text{DOM}}(T)$. Once we have specified this flow, we can simply apply the flow described in the proof of Lemma B.1 to route flow among the “main” classes.

Lemma D.6. *Given a graph G , corresponding CSDS Glauber graph $\mathcal{M}_{\text{DOM}}(G)$, and a class $\mathcal{C}_{\text{DOM}}(T)$ of CSDS’s in $\mathcal{M}_{\text{DOM}}(G)$, the partition into subclasses given in Definition D.5 satisfies the conditions of Lemma D.1.*

Proof. Clearly there are $O(1)$ classes. We observe that the subclasses are all within an $O(1)$ size factor of one another. To see this, compare the sizes of $\mathcal{C}_{\text{DOM}}(T)$ and $\mathcal{C}_{\text{DOM}}(T_W)$. Since $\mathcal{C}_{\text{DOM}}(T_W) \subseteq \mathcal{C}_{\text{DOM}}(T)$, $|\mathcal{C}_{\text{DOM}}(T_W)| \leq |\mathcal{C}_{\text{DOM}}(T)|$. On the other hand, consider the mapping that sends every CSDS $s \in \mathcal{C}_{\text{DOM}}(T)$ to a CSDS $s' \in \mathcal{C}_{\text{DOM}}(T_W)$ in which at least one neighbor (say, the first lexicographically) of each $w \in W$ is selected in A , and in which the first neighbor of each $\bar{w} \in U \setminus W$ is selected in B . This mapping is clearly $2^{|U|}$ -to-1, and thus the size factor difference is $O(1)$.

Now, since the subclasses are not equivalence classes, many pairs of subclasses overlap. In particular, let T and U be as before, and suppose for some $u \in U$, $W' = W \cup \{u\}$. Then $\mathcal{C}_{\text{DOM}}(T_W)$ and $\mathcal{C}_{\text{DOM}}(T_{W'})$ overlap at those CSDS’s in which some neighbor of u in A is selected, and some neighbor of u in B is selected. It is easy to see that for every such pair of subclasses $\mathcal{C}_{\text{DOM}}(T_W)$ and $\mathcal{C}_{\text{DOM}}(T_{W'})$, at least half of the CSDS’s in $\mathcal{C}_{\text{DOM}}(T_W)$ and at least half of those in $\mathcal{C}_{\text{DOM}}(T_{W'})$ are part of the overlap.

It is clear that $\mathcal{C}_{\text{DOM}}(T)$ is internally connected via these overlaps: every CSDS in $\mathcal{C}_{\text{DOM}}(T)$ has a path to the trivial CSDS in which every non-forbidden vertex of $A \cup B$ is selected. Thus the conditions of Lemma D.1 are satisfied. \square

As in the discussion following Lemma D.4, we derive the bound in Theorem 1.3 as follows: the flow

within a class incurs a congestion factor of

$$4 \frac{1+\lambda}{\lambda} \cdot (2\hat{\lambda})^{t+1} |\mathcal{C}_{\text{DOM}}(T)| / (\min_W |\mathcal{C}_{\text{DOM}}(T_W)|) \leq 4 \frac{1+\lambda}{\lambda} (4\hat{\lambda})^{t+1}.$$

The application of Lemma 7.2 contributes a $2(K\hat{\lambda}^{t+1} + 1)$ factor to the inductive step, with $K = 2^{t+1}$.

Thus the factor for one iteration is at most

$$8 \frac{1+\lambda}{\lambda} (8\hat{\lambda}^2)^{t+2}.$$

The inclusion of the $t+1$ separator vertices in the subproblems, as with b -edge covers, increases the induction depth, and an analogous analysis gives a base case of $(2\hat{\lambda})^{6(t+1)}$ using $\varepsilon = 1/6$. Putting all this together with the fact that $\Delta_{\mathcal{M}} \leq n$ and $1/\pi_{\min}^* \leq (2\hat{\lambda})^n$ gives a mixing bound of

$$\begin{aligned} & O(((2\hat{\lambda})^{6(t+1)})^2 \left(\frac{\hat{\lambda}}{1+\hat{\lambda}}\right)^2 (1+\log \hat{\lambda}) \cdot n^{2\log(8\frac{1+\lambda}{\lambda})(8\hat{\lambda}^2)^{t+2}/\log(6/5)+3}) \\ &= O(((2\hat{\lambda})^{12(t+1)}) \left(\frac{\hat{\lambda}}{1+\hat{\lambda}}\right)^2 (1+\log \hat{\lambda}) \cdot n^{2(3+\log(\frac{1+\lambda}{\lambda}))+ (t+2)(3+\log \hat{\lambda})/\log(6/5)+3}). \end{aligned}$$

D.4 Rapid mixing in the relaxed hierarchical framework for all $\lambda > 0$

We now generalize Lemma D.1 to all $\lambda > 0$, finishing the proof of Theorems 1.2 and 1.3.

Lemma D.7. *For the Glauber graphs $\mathcal{M}_{\text{BEC}}(G)$ and $\mathcal{M}_{\text{DOM}}(G)$, with classes defined as in Lemma D.1, and with stationary distribution π induced by parameter λ as in the discussion in Section C.1, the flow construction in Lemma D.1 results in a congestion factor gain of at most $\rho_{\max} = O(1)$ at each of the $O(\log n)$ levels of induction, resulting in at most polynomial overall congestion.*

Proof. We need to show that the flow construction within a class $\mathcal{C}(T)$ in Lemma D.1 produces at most an $O(1)$ -factor increase in congestion; the rest of the argument is similar to the proof of Lemma C.9. For the case of dominating sets, consider a pair of $\mathcal{C}_{\text{DOM}}(T_W) \subseteq \mathcal{C}_{\text{DOM}}(T)$ and $\mathcal{C}_{\text{DOM}}(T_{W'}) \subseteq \mathcal{C}_{\text{DOM}}(T)$. For every such pair, consider the intersection \mathcal{I} of the two subclasses, namely the set of dominating sets in which for every input graph vertex $v \in W \cup W'$, some neighbor of v is selected in A , and for every vertex $w \in U \setminus (W \cap W')$, some neighbor of w is selected in B . There exists an $O(1)$ -to-1 mapping from $\mathcal{C}_{\text{DOM}}(T_W)$ to \mathcal{I} —found by adding $O(1)$ neighbors of vertices in U as described above to each dominating set $S \in \mathcal{C}_{\text{DOM}}(T_W)$ —under which the image S' of S has $|S' \setminus S| = O(1)$, and therefore $\pi(S') = \lambda^{|S' \setminus S|} \pi(S) = \Theta(1)\pi(S)$. This shows that $\pi(\mathcal{I}) = \Theta(\pi(\mathcal{C}_{\text{DOM}}(T_W))) = \Theta(\pi(\mathcal{C}_{\text{DOM}}(T_{W'})))$.

Thus we use the overlaps between classes to route flow along a path of classes as in the proof of Lemma D.1. As before, at each class in the path, the internal routing produces an $O(1)$ factor increase in the congestion within the class. The concern, again, is that due to overlap, there may be edges belonging to multiple classes that thus incur congestion multiple times in the routing of the flow; as before, this is not a problem as there are $O(1)$ pairs of classes for which this occurs.

The argument for $\mathcal{M}_{\text{BEC}}(G)$ is similar, with the intersection \mathcal{I} found by selecting sufficiently many edges incident to each vertex $x \in X$ to satisfy membership in both subclasses $\mathcal{C}_{\text{BEC}}(T_\beta)$ and $\mathcal{C}_{\text{BEC}}(T_{\beta'})$. \square

D.5 Rapid mixing of the Glauber dynamics on b -matchings for all $\lambda > 0$

For the claim about b -matchings in Theorem 1.3, we do not need the relaxed framework; in fact it suffices to combine Lemma 3.14 with the following lemma:

Lemma D.8. *It is easy to adapt the proof of Lemma C.9 to the hierarchical framework, proving the claim about the Glauber dynamics on b -matchings in Theorem 1.3 for all $\lambda > 0$.*

Proof. The proof of Lemma C.9 uses a simple mapping argument to show that for every ancestor $\mathcal{C}(T_i)$ of a class $\mathcal{C}(T)$, $\pi(\mathcal{C}(T_i)) = \Theta(1)\pi(\mathcal{C}(T))$, then allows each boundary edge e between classes to carry $O(1)Q(e)$ units of flow across the boundary, by ensuring that each boundary vertex S carries flow in proportion to its weight $\pi(S)$. Since all pairs of classes $\mathcal{C}(T), \mathcal{C}(T')$ have a common ancestor in the case of b -matchings, we in fact have $\pi(\mathcal{C}(T)) = \Theta(1)\pi(\mathcal{C}(T'))$ for every pair of classes $\mathcal{C}(T), \mathcal{C}(T')$. The bound on flow across the boundary therefore still holds; the argument for bounding congestion factor increase within a class is the same as in the proof of Lemma C.9. \square

For the specific mixing upper bound for b -matchings, we use Lemma 7.2, and observe that $K \leq 2^{\Delta(t+1)}$; the parameter λ contributes at most a $\hat{\lambda}^{\Delta(t+1)}$ factor, $1/\pi_{\min}^* \leq (2\hat{\lambda})^m$, and $\Delta_{\mathcal{M}} \leq m$, so we have a mixing upper bound of

$$\begin{aligned} & O(((1 + \hat{\lambda})\hat{\lambda})^2 (2((2\hat{\lambda})^{\Delta(t+1)} + 1))^{2 \log n} m^3 (1 + \log \hat{\lambda})) \\ & = O(((1 + \hat{\lambda})\hat{\lambda})^2 (1 + \log \hat{\lambda}) m^3 n^{2\Delta(t+2)(1 + \log \hat{\lambda}) + 2}). \end{aligned}$$

D.6 Maximal independent sets and maximal b -matchings in the non-hierarchical framework

D.6.1 Dealing with internally disconnected classes

As noted in Appendix D, we use the non-hierarchical framework and assume bounded carving width for the chains on maximal independent sets and maximal b -matchings. Once we have defined these chains, we will see that partitioning the Glauber graph for each chain in the natural way will result in classes that are not necessarily internally connected. The solution will be to relax the framework conditions so that the classes need not be disjoint—but then require that every pair of overlapping classes must overlap in a large number of vertices. More precisely:

Lemma D.9. *Suppose a Glauber graph $\mathcal{M}(G)$ satisfies the conditions of the non-hierarchical framework in Section 3.4, except that:*

1. *The $O(1)$ classes are not necessarily disjoint.*
2. *Each pair of classes $\mathcal{C}(T)$ and $\mathcal{C}(T')$ sharing at least one vertex shares $\Theta(1)|\mathcal{C}(T)| = \Theta(1)|\mathcal{C}(T')|$ vertices.*

Then the expansion of $\mathcal{M}(G)$ is $\Omega(1/n^c)$, where $c = O(1)$.

Proof. The multicommodity flow construction is as in the proof of Lemma 3.9, except that when sending flow from $S \in \mathcal{C}(T)$ to $S' \in \mathcal{C}(T') \neq \mathcal{C}(T)$ via a path through intermediate classes, we now have some pairs of intermediate classes that share boundary vertices, instead of sharing boundary edges. The flow is the same as before, except that there is no need to send flow across a boundary in these cases.

The congestion analysis is the same as in the proof of Lemma D.1. \square




Figure 8: Two maximal independent sets in a graph G : S (left) and S' (right). S and S' differ by a flip, with the separator X inducing the classes to which the sets belong. S' results from adding v to S , removing the neighbors of v , and adding some of the neighbors of the removed vertices.

D.6.2 Maximal independent sets

We now apply the non-hierarchical framework to the flip chain on maximal independent sets. We first define the maximal independent set mixing problem, then show that it meets the criteria of the framework, up to the conditions in Lemma D.9.

We return to the Glauber graph defined in Definition 2.6.

Lemma D.10. *The maximum degree of $\mathcal{M}_{\text{MIS}}(G)$ is at most $n \cdot 2^{\Delta^2 + \Delta}$, where $n = |V(G)|$ and Δ is the maximum degree of any vertex in G .*

Proof. The number of neighbors of a given maximal independent set S is the same as the number of ways to choose a vertex v to add to or drop from S , along with a subset of the vertices at distance at most two from v to add or drop. \square

Definition D.11. *Define the maximal independent set flip chain on a graph G with Glauber graph $\mathcal{M}_{\text{MIS}}(G)$ as the following Markov chain (let Δ_M be the maximum degree of $\mathcal{M}_{\text{MIS}}(G)$):*

1. Let X_0 be an arbitrary maximal independent set in $V(\mathcal{M}_{\text{MIS}}(G))$.
2. For $t \geq 0$, define X_{t+1} as follows:

With probability $(1/2)(\delta(X_t)/\Delta_M)$, let X_{t+1} be a neighbor in $\mathcal{M}_{\text{MIS}}(G)$ of X_t , selected uniformly at random from the neighbors of X_t .

With probability $1 - (1/2)(\delta(X_t)/\Delta_M)$ let $X_{t+1} = X_t$.

For technical reasons, the following observation is necessary for obtaining a rapid mixing bound from an expansion bound on $\mathcal{M}_{\text{MIS}}(G)$.

Remark D.12. *It is easy to see that this is the standard Markov chain on $\mathcal{M}_{\text{MIS}}(G)$, with appropriate self loops added in the standard way. Furthermore, by Lemma D.10, if G has bounded degree, then the degree-based weighting in Definition D.11 does not cause the spectral expansion of the chain to differ by more than a polynomial factor from the edge (or vertex) expansion of $\mathcal{M}_{\text{MIS}}(G)$.*

D.6.3 Verification of conditions for maximal independent sets

We show how to apply the non-hierarchical version of the framework when the carving width of G is bounded. First, to satisfy Condition 1, we would *like* to use a partition analogous to that defined in Definition 3.3: each class $\mathcal{C}_{\text{MIS}}(T)$ is the set of maximal independent sets that agree on their restriction to the vertex separator X for G . However, a subtlety arises when considering the Cartesian product structure of the Glauber graphs on A and B within a class T : in the independent set Glauber graph, $\mathcal{C}_{\text{IS}}(T)$ was a Cartesian product of two independent set Glauber graphs $\mathcal{M}_{\text{MIS}}(G_A)$ and $\mathcal{M}_{\text{MIS}}(G_B)$, respectively defined on the independent sets in $A \setminus N_A(T)$ and on those in $B \setminus N_B(T)$. Here, however, the independent sets chosen in A and in B need to give a *maximal* independent set when their union is taken with the set chosen in X .

It may be that the independent set in X identified with $\mathcal{C}_{\text{MIS}}(T)$ is not maximal. For a simple example, suppose X is a path of length three, consisting of vertices u, v , and w and edges (u, v) and (v, w) , with u having neighbors only in A , w having neighbors only in B , and v having only u and w as neighbors. Suppose $\mathcal{C}_{\text{MIS}}(T)$ is identified with the independent set $\{u\} \subseteq X$. Then every maximal independent set $S \in T$ has some neighbor of w in B chosen, or else S would not be maximal. Furthermore, it is easy to show that defining classes in this way would result in internally disconnected classes. Thus we cannot simply eliminate vertices in $N_A(T)$ from A and $N_B(T)$ from B and define smaller maximal independent set Glauber graphs. Instead, we define the classes—which, per the conditions of Lemma D.9, need not be disjoint—as follows:

Definition D.13. *Given a graph G with maximal independent set Glauber graph $\mathcal{M}_{\text{MIS}}(G)$ and a class $\mathcal{C}_{\text{MIS}}(T)$, let $U \subseteq X$ be the set of all unselected vertices in T that have no neighbor selected in T . For each independent subset of the vertices in $N_A(U) \cup N_B(U)$ that covers all of U —that is, for each independent subset $C \subseteq N_A(U) \cup N_B(U)$ such that every $x \in U$ has some neighbor $z \in C$, let $\mathcal{C}_{\text{MIS}}(T_C)$ be the class of all independent sets in G that agree with T on X , and that include all of the vertices in C .*

The smaller Glauber graphs on A and B are now independent for a given class $\mathcal{C}_{\text{MIS}}(T_C)$: for each $z \in C$, remove z and all neighbors in $N_A(z) \cup N_B(z)$ from the graph, and consider the resulting maximal independent set Glauber graphs on $A \setminus (C \cup N_A(C))$ and on $B \setminus (C \cup N_B(C))$. Each class $\mathcal{C}_{\text{MIS}}(T_C)$ is a Cartesian product of two such graphs. It suffices to show that this definition obeys the conditions of Lemma D.9:

Lemma D.14. *Given a graph G with bounded carving width and corresponding maximal independent set Glauber graph $\mathcal{M}_{\text{MIS}}(G)$, the definition of classes in Definition D.13 satisfies the conditions of Lemma D.9.*

Proof. The Cartesian product structure of $\mathcal{C}_{\text{MIS}}(T_C)$ and the fact that X is a balanced separator satisfy Conditions 5 and 6 of the non-hierarchical framework.

The classes do not partition $\mathcal{C}_{\text{MIS}}(T)$. However, clearly there are $O(1)$ classes. The classes are also within an $O(1)$ size factor of one another. To see this, define the following mapping f from the set of all maximal independent sets in $\mathcal{M}_{\text{MIS}}(G)$ to the set of maximal independent sets in a class $\mathcal{C}_{\text{MIS}}(T_C)$. For each maximal independent set $S \in V(\mathcal{M}_{\text{MIS}}(G))$, let $S' = f(S) \in \mathcal{C}_{\text{MIS}}(T_C)$ be the following maximal independent set: (i) let S' agree with T on all vertices in X ; (ii) let S' agree with T_C on all vertices in $N_A(T) \cup N_B(T)$; (iii) let S' agree with S on all vertices not in $X \cup N_A(T) \cup N_B(T)$ and having no neighbor in $X \cup N_A(T) \cup N_B(T)$; (iv) add vertices to S , if needed, to obtain maximality.

The symmetric difference $f(S) \oplus S$ is of size at most $\Delta^2|X| \leq \Delta^2(t+1) = O(1)$, where t is the (bounded) treewidth of G and Δ is the (bounded) degree; thus f is an $O(1)$ -to-1 mapping. By similar reasoning, the number of shared maximal independent sets between any two overlapping classes $\mathcal{C}_{\text{MIS}}(T_C)$ and $\mathcal{C}_{\text{MIS}}(T_{C'})$ is at least $\Omega(1)|\mathcal{C}_{\text{MIS}}(T_C)|$, and the number of edges between any two adjacent classes $\mathcal{C}_{\text{MIS}}(T_C)$ and $\mathcal{C}_{\text{MIS}}(T'_{C'})$ is at least $\Omega(1)|\mathcal{C}_{\text{MIS}}(T_C)|$. The lemma follows. \square

D.6.4 Maximal b -matchings in the non-hierarchical framework

We return to the maximal b -matching Glauber graph defined in Section 2.

The argument that the graph is connected is similar to the proof of Lemma 2.7.

It suffices to define a partition and verify the conditions. We would *like* to identify each class $\mathcal{C}_{\text{MBM}}(T)$ with the chosen subset of the edges that have at least one endpoint in the small balanced separator X . However, as with maximal independent sets, the maximality requirement introduces non-independent subproblems. To modify the definition of the classes, we first need to introduce the notion of a *saturated* vertex:

Definition D.15. *Given a b -matching in a graph G , consider a vertex v saturated if $b(v)$ edges incident to v are selected in the matching.*

It may be that a vertex $v \in X$ is not saturated in a maximal b -matching, and thus the choice of edges inducing $\mathcal{C}_{\text{MBM}}(T)$ does not saturate v . In this case, we have a constraint on the subproblems in A and B . Namely, it must be that some neighbor of v , $u \in N(X)$, is saturated, or else the edge (u, v) could be added to the matching. We use this fact to define the subclasses of a class $\mathcal{C}_{\text{MBM}}(T)$:

Definition D.16. *Given a graph G with separator X , maximal b -matching Glauber graph $\mathcal{M}_{\text{MBM}}(G)$, and a set T of edges selected whose endpoints all lie in X , let $U \subseteq X$ be the set of unsaturated vertices in X induced by T ; let C be a minimal set of edges such that, after adding C to T , some neighbor of v is saturated for every $v \in U$. Define the class $\mathcal{C}_{\text{MBM}}(T_C)$ as the set of all maximal b -matchings in T that contain all edges in C .*

As in Section D.6.2, these classes are not equivalence classes, because they overlap. Again, however, each b -matching belongs to at most $O(1)$ subclasses, and thus this overlap does not interfere with the proof.

We now verify that this definition of classes satisfies Lemma D.9:

Lemma D.17. *Given a graph G with bounded carving width, corresponding maximal b -matching Glauber graph $\mathcal{M}_{\text{MBM}}(G)$, the division into classes as in Definition D.16 satisfies the conditions of Lemma D.9.*

Proof. The argument is similar to the proof of Lemma D.14: again we have a Cartesian product structure in each class $\mathcal{C}_{\text{MBM}}(T_C)$ —where the resulting maximal b -matching subproblems on A and B result from (i) removing each edge $(u, w) \in C$ from $G[A]$ and $G[B]$, and (ii) decreasing $b(u)$ and $b(w)$ accordingly.

Clearly the number of classes is $O(1)$, due to the bounded carving width of G . The classes differ by an $O(1)$ size factor, and the overlaps are large; the argument, along with the resulting flow, is similar to that in the proof of Lemma D.14. \square

D.6.5 Specific mixing upper bounds for maximal independent sets and maximal b -matchings

For the derivation of the specific bounds stated in Theorem 1.4, we apply Lemma 7.1, with the modification that the \mathcal{E}_{\min} term must be replaced by the term $\min\{\mathcal{E}_{\min}, \mathcal{O}_{\min}\}$, where \mathcal{O}_{\min} is the size of the smallest overlap between a pair of classes that share at least one vertex. For maximal independent sets, it is easy to show that $\mathcal{E}_{\min} \geq N/2^{7\Delta^6(t+1)}$ and $\mathcal{O}_{\min} \geq N/2^{3\Delta^2(t+1)}$, so $\min\{\mathcal{E}_{\min}, \mathcal{O}_{\min}\} \geq N/2^{7\Delta^6(t+1)}$. We also gain at each level of induction an additional factor of $K \leq 2^{(\Delta+1)(t+1)}$ due to overlaps. Combining this with the fact that $\Delta_{\mathcal{M}} \leq 2^{3\Delta^2}n$ and $N \leq 2^n$ gives a total mixing bound of

$$\begin{aligned} & O((4 \cdot 2^{7\Delta^6(t+1)} \cdot 2^{(\Delta+1)(t+1)2 \log n} \cdot 2^{6\Delta^2} n^3) \\ & = O(2^{6\Delta^2} n^{2(t+1)(7\Delta^6 + \Delta + 1) + 7}), \end{aligned}$$

as claimed. (The $\log(3/2)$ term in the theorem statement comes from the fact that the base of the log in the induction is $3/2$.)

A similar argument for maximal b -matchings gives the result claimed in Theorem 1.4, with $\Delta_{\mathcal{M}} \leq 2^{6\Delta^2}m$, $N \leq 2^m$, $K \leq 2^{3\Delta^2(t+1)}$, $\mathcal{E}_{\min} \geq N/2^{8\Delta^7(t+1)}$, and $\mathcal{O}_{\min} \geq N/2^{4\Delta^3(t+1)}$.

I.e., we have

$$\begin{aligned} & O((4 \cdot 2^{8\Delta^7(t+1)} \cdot 2^{3\Delta^2(t+1)2 \log n} \cdot 2^{12\Delta^2} m^3) \\ & = O(2^{12\Delta^2} m^3 n^{2(t+1)(8\Delta^7 + 3\Delta^2) + 4}). \end{aligned}$$

E Full proof that k -angulations satisfy the framework

E.1 Cardinalities of classes and of edge sets

We make some observations about the nature and cardinalities of the classes in $\mathcal{S}_{k,(k-2)n+2}$, and of the sets and numbers of edges between the classes.

Lemma E.1. *Each k -gonal class in $\mathcal{S}_{k,(k-2)n+2}$ induces a subgraph of $K_{k,(k-2)n+2}$ that is isomorphic to the Cartesian product $K_{k,(k-2)i_1+2} \square K_{k,(k-2)i_2+2} \square \cdots \square K_{k,(k-2)i_k+2}$, for some $1 \leq i_1 \leq \cdots \leq i_k \leq n/2$, $i_1 + \cdots + i_k = n - 1$. Each diagonal class is of the form $K_{i_1} \square K_{i_2}$, $i_1 \leq i_2 \leq n/2$, $i_1 + i_2 = n$.*

Proof. Each k -gon T partitions the regular $(k-2)n+2$ -gon into smaller convex polygons with side lengths $(k-2)i_1+2, (k-2)i_2+2, \dots, (k-2)i_k+2$. Thus each k -angulation in $\mathcal{C}_k(T)$ can be identified with a tuple of k -angulations of these smaller polygons. The Cartesian product structure then follows from the fact that every flip between two k -angulations in $\mathcal{C}_k(T)$ can be identified with a flip in one of the smaller polygons. \square

Lemma E.2. *Each set of edges between classes in $\mathcal{S}_{k,(k-2)n+2}$ is at least equal in cardinality to the Cartesian product $K_{(k-2)i_1+2} \square K_{(k-2)i_2+2} \square \cdots \square K_{(k-2)i_{2k-2}+2}$, for $i_1 \leq \cdots \leq i_{2k-2} \leq n/2$, $i_1 + \cdots + i_{2k-2} = n - 2$. Furthermore, no two edges in any such edge set share a vertex.*

Proof. Each flip between k -angulations in adjacent classes $\mathcal{C}_k(T)$ involves flipping a diagonal of the k -gon T to transform k -angulation $t \in \mathcal{C}_k(T)$ into k -angulation $t' \in \mathcal{C}_k(T')$. Whenever this is possible, there must exist a $2k-2$ -gon Q , sharing $k-1$ sides with T (the $k-1$ sides that are

not flipped), such that both t and t' contain Q . Furthermore, every $t \in \mathcal{C}_k(T)$ containing Q has a flip to a distinct $t' \in \mathcal{C}_k(T')$. The claim follows from the fact that the set of all such boundary vertices $t \in \mathcal{C}_k(T)$ can be identified with the Cartesian product described. (There may be many such $2k - 2$ -gons for a given pair of classes, but the claim holds as a lower bound.) \square

Corollary E.3. *Each k -gonal class in $\mathcal{S}_{k,(k-2)n+2}$ has cardinality $C_{k,i_1}C_{k,i_2} \cdots C_{k,i_k}$, each diagonal class has cardinality $C_{k,i_1}C_{k,i_2}$, and each edge set between classes has cardinality $C_{k,i_1}C_{k,i_2} \cdots C_{k,i_{2k-2}}$. Here, i_1, \dots, i_{2k-2} are as in Lemmas E.1 and E.2.*

E.2 Applying the framework

We are almost ready to prove that $K_{k,(k-2)n+2}$ satisfies the conditions of the framework in Lemma 3.10 (with $O(1)$ factors replaced by polynomial factors), but first we note the following result by Nakamoto, Kawatani, Matsumoto, and Urrutia [45]:

Lemma E.4. *$K_{4,n}$ is connected.*

It is also well known and easy to show that $K_{k,(k-2)n+2}$ is connected. One such way is via the isomorphism [28] between flips on k -angulations and rotations on $k - 1$ -ary plane trees. One can prove that the rotation graph on $k - 1$ -ary plane trees is connected as follows: find a path from any given tree to a “spine,” where all internal nodes belong to a simple path via left children from the root to the leftmost leaf [15]. (This path consists of repeated left rotations.) Every non-spine tree has some internal node at which a left rotation can be performed. Furthermore, when no such operation is still possible, one has a spine.

Sleator, Tarjan, and Thurston also proved [55] that the diameter of $K_{3,n+2}$ is at most $2n - 6$ for $n \geq 11$.

We now prove Lemma 6.5:

Lemma 6.5. *The flip graph $K_{k,(k-2)n+2}$, along with the partition $\mathcal{S}_{k,(k-2)n+2}$, meets the conditions of the non-hierarchical framework, except that the $O(1)$ terms described in each condition are replaced with $O(n^{O(1)})$.*

Proof. By Lemma E.1 and the easy observation that there are at most $\binom{(k-2)n+2}{k}$ classes, the partition $\mathcal{S}_{k,(k-2)n+2}$ meets Conditions 1 and 5 of the framework, with the modification to Condition 1 that the $O(1)$ term is replaced with $O(n^{O(1)})$, and Condition 6 easily follows from the identification of each class with a k -gon containing the center of the $(k - 2)n + 2$ -gon.

Conditions 2, 3, and 4 follow, with the modification that the $O(1)$ terms are replaced with $O(n^{O(1)})$ terms, from Lemma 2.29 and Corollary E.3. \square

The specific bound in Theorem 1.5 comes from combining Lemma 7.1 with the following observation:

Remark E.5. *The smallest edge set between classes in $\mathcal{S}_{k,(k-2)n+2}$ has size at least*

$$\begin{aligned} C_{k,i_1} \cdots C_{k,i_{2k-2}} &\geq N \cdot \frac{f(k, i_1) \cdots f(k, i_{2k-2})}{e^{(2k-2)/6+1/12} ((k-1)/(k-2))^{2k-2} f(k, n)} \\ &\geq N e^{(3-4k)/12} \cdot \frac{(k-2)^{k-3/2}}{(k-1)^{3k-5/2}} \cdot \frac{1}{(2\pi)^{k-3/2}} \cdot \frac{1}{n^{3k}}. \end{aligned}$$

Thus applying Lemma 7.1 (and using the fact that $K_{k,(k-2)n+2}$ is a $(k-2)O(n)$ -regular graph with $\log N \leq (k-1)n \log(k-1)$, gives

$$\begin{aligned} & O((4N/\mathcal{E}_{\min})^{2 \log n} (k-1)^3 (\log(k-1)) n^3) \\ &= O((k-1)^3 (\log(k-1)) n^3 (4e^{(4k-3)/12} \cdot \frac{(k-1)^{3k-5/2}}{(k-2)^{k-3/2}} \cdot (2\pi)^{k-3/2} \cdot n^{3k})^{2 \log n}) \\ &= O((k-1)^3 (\log(k-1)) n^3 (4e^{(4k)/12} \cdot (k-1)^{3k} \cdot (2\pi)^k \cdot n^{3k})^{2 \log n}) \\ &= O((k-1)^3 (\log(k-1)) \cdot n^{2(3k \log(k-1) + k(1+\log \pi) + 3k \log n + k) + 7}). \end{aligned}$$

F Deferred Proof Details

Lemma 2.17. *Let $J = G \square H$. Given multicommodity flows ρ_G and ρ_H for G and H respectively, there exists a multicommodity flow ρ_J for J with*

$$\rho_{Jmax} \leq \max\{\rho_{Gmax}, \rho_{Hmax}\}.$$

Proof. Let ρ_G and ρ_H be as stated; we construct ρ_J as follows:

1. Within each copy of H in J , construct the flow internally according to ρ_H . Similarly, use ρ_G internal to each G copy for each pair of vertices within the G copy.
2. Order the copies of H arbitrarily $H_1, \dots, H_{|V(G)|}$. For each pair of H copies H_r and H_s , $s < r$, and for each vertex $h_r \in H_r, h_s \in H_s$, let the flow from h_r to h_s go through (i) the ρ_H flow in H_r from h_r to the counterpart vertex $u \in H_r$ of h_s , then through (ii) the ρ_G flow that goes from u to h_s (in the G copy that h_s and u both belong to).

Part 1 generates no additional flow. Part 2 generates at most $|V(H)|$ extra flow through each existing ρ_G flow, and at most $|V(G)|$ extra flow through each existing ρ_H flow. This results in scaling the amount of ρ_G flow using any given edge in a G copy by a factor of $|V(H)|$ —while replacing the $\frac{1}{|V(G)|}$ term in the congestion definition by $\frac{1}{|V(J)|} = \frac{1}{|V(G)||V(H)|}$ —and similarly scaling the amount of ρ_H flow using an edge in an H copy by $|V(G)|$. The result follows.

We also need to handle the generalized definition of congestion in the case of weighted graphs for the results in Appendix C. It suffices to use the construction above. In the new flow, h_r needs to send $\pi(h_r)\pi(h_s)$ units to h_s via u . Let $\pi(H)$ denote the combined weight of the H copy in which h_r and u lie. Since h_r already sends $\pi(h_r)\pi(u)/(\pi(H)^2)$ units, under ρ_H , to u —resulting in each edge e in H carrying at most $Q(e)\rho_{Hmax}$ units— h_r can send $\pi(h_r)\pi(u)$ units to u —and $\pi(h_r)\pi(G_s) = \pi(h_r)\pi(u)/\pi(H)$ units to the copies of h_s lying in the G copy G_s containing u and h_s . This is because

$$\pi(u) = \pi_G(u)\pi_H(u) = \pi(G_s)\pi(H),$$

where the latter equality follows from the fact that

$$\pi(G_s) = \sum_{(g,h) \in G_s} \pi(g,h) = \sum_{(g,h) \in G_s} \pi_G(g)\pi_H(h) = \pi_H(u) \sum_{(g,h) \in G_s} \pi_G(g) = \pi_H(u),$$

and from the fact that similarly $\pi(H) = \pi_G(u)$.

Each edge e in H then carries $Q(e)\rho_{Hmax}/\pi(H)$ units. Thus the amount carried across each edge $e = (x,y)$ increases by a factor of at most $1/\pi(H)$, while the weight of each edge decreases by a factor of $1/(\pi(H)(|\Delta_H|/|\Delta_J|))$. \square

Lemma 2.7. *The graph $\mathcal{M}_{\text{MIS}}(G)$ in Definition 2.6 is connected.*

Proof. Let $S \neq S'$ be maximal independent sets, and consider the symmetric difference $S \oplus S'$: if $|S \oplus S'| > 0$, choose some $v \in S' \setminus S$. Obtain a new set S'' by adding v to S and removing all neighbors of v from S , then greedily adding neighbors of neighbors of v until an maximal independent set is obtained. Repeat this process with a new vertex $v' \in S' \setminus S''$, and so on, for every vertex in $S' \setminus S$, obtaining a sequence of sets $S_1 = S, S_2 = S'', S_3, \dots, S_k$. Crucially, once a vertex v is selected from S' in this process, giving set S_i , we have $v \in S_j$ for all $i \leq j \leq k$. This is because the only way for a vertex to be removed in the process is for one of its neighbors to be selected from S' . However, since S' is an independent set, no neighbor of v belongs to S' .

Thus we have $S_k = S'$, proving that there is a path in $\mathcal{M}_{\text{MIS}}(G)$ between every pair of maximal independent sets. \square

References

- [1] Nima Anari, Shayan Oveis Gharan, and Cynthia Vinzant. Log-concave polynomials, entropy, and a deterministic approximation algorithm for counting bases of matroids. In *2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 35–46, 2018. doi: 10.1109/FOCS.2018.00013.
- [2] Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant. Log-concave polynomials II: High-dimensional walks and an FPRAS for counting bases of a matroid. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing (STOC 2019)*, New York, NY, USA, 2019. Association for Computing Machinery. ISBN 9781450367059.
- [3] Nima Anari, Kuikui Liu, and Shayan Oveis Gharan. Spectral independence in high-dimensional expanders and applications to the hardcore model. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1319–1330, 2020. doi: 10.1109/FOCS46700.2020.00125.
- [4] Pinaki Banerjee. Scattering amplitudes from positive geometries. In Prafulla Kumar Behera, Vipin Bhatnagar, Prashant Shukla, and Rahul Sinha, editors, *XXIII DAE High Energy Physics Symposium*, pages 421–427, Singapore, 2021. Springer Singapore. ISBN 978-981-33-4408-2.
- [5] Ivona Bezáková and Wenbo Sun. Mixing of Markov chains for independent sets on chordal graphs with bounded separators. In *Computing and Combinatorics*, pages 664–676. Springer International Publishing, 2020. ISBN 978-3-030-58150-3.
- [6] Ivona Bezáková, Andreas Galanis, Leslie Ann Goldberg, Heng Guo, and Daniel Štefankovič. Approximation via correlation decay when strong spatial mixing fails. *SIAM Journal on Computing*, 48(2):279–349, 2019.
- [7] Hans L. Bodlaender. Fixed-parameter tractability of treewidth and pathwidth. volume 7370 of *Lecture Notes in Computer Science*, pages 196–227. Springer, 2012. doi: 10.1007/978-3-642-30891-8_12.
- [8] Magnus Bordewich and Ross J. Kang. Subset Glauber dynamics on graphs, hypergraphs and matroids of bounded tree-width. *Electronic Journal of Combinatorics*, 21(4), 2014. ISSN 1077-8926. doi: <https://doi.org/10.37236/4195>.

- [9] Pietro Caputo, Fabio Martinelli, Alistair Sinclair, and Alexandre Stauffer. Dynamics of lattice triangulations on thin rectangles. *Electronic Journal of Probability*, 21, 05 2015. doi: 10.1214/16-EJP4321.
- [10] Pietro Caputo, Fabio Martinelli, Alistair Sinclair, and Alexandre Stauffer. Random lattice triangulations: structure and algorithms. *Annals of Applied Probability*, 25:1650–1685, 2015.
- [11] Alessandra Caraceni. A polynomial upper bound for the mixing time of edge rotations on planar maps. *Electronic Journal of Probability*, 25(none):1 – 30, 2020. doi: 10.1214/20-EJP519. URL <https://doi.org/10.1214/20-EJP519>.
- [12] Alessandra Caraceni and Alexandre Stauffer. Polynomial mixing time of edge flips on quadrangulations. *Probability Theory and Related Fields*, 176(1):35–76, Feb 2020. ISSN 1432-2064. doi: 10.1007/s00440-019-00913-5.
- [13] Yijia Chen and Jörg Flum. The parameterized complexity of maximality and minimality problems. *Annals of Pure and Applied Logic*, 151(1):22 – 61, 2008. ISSN 0168-0072. doi: <https://doi.org/10.1016/j.apal.2007.09.003>.
- [14] Zongchen Chen, Andreas Galanis, Daniel Štefankovič, and Eric Vigoda. Rapid mixing for colorings via spectral independence. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1548–1557. doi: 10.1137/1.9781611976465.94.
- [15] Karel Culik II and Derick Wood. A note on some tree similarity measures. *Information Processing Letters*, 15(1):39–42, 1982.
- [16] Martin Dyer, Leslie Ann Goldberg, and Mark Jerrum. Matrix norms and rapid mixing for spin systems. *The Annals of Applied Probability*, 19(1):71–107, 2021/10/28/ 2009. ISSN 10505164. URL <http://www.jstor.org/stable/30243572>. Full publication date: Feb., 2009.
- [17] David Eppstein. The effect of planarization on width. In *Graph Drawing and Network Visualization*, pages 560–572. Springer International Publishing, 2018. ISBN 978-3-319-73915-1.
- [18] Jeff Erickson. Computational topology: Treewidth. Lecture Notes, 2009. URL <http://jeffe.cs.illinois.edu/teaching/comptop/2009/notes/treewidth.pdf>.
- [19] Weiming Feng, Heng Guo, Yitong Yin, and Chihao Zhang. Rapid mixing from spectral independence beyond the boolean domain. In *Proceedings of the Thirty-Second Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2021*, page 1558–1577, USA, 2021. Society for Industrial and Applied Mathematics. ISBN 9781611976465.
- [20] S. M. Ferdous, A. Khan, and A. Pothen. Parallel algorithms through approximation: B-edge cover. In *2018 IEEE International Parallel and Distributed Processing Symposium (IPDPS)*, pages 22–33. IEEE Computer Society, May 2018. doi: 10.1109/IPDPS.2018.00013.
- [21] Juliana Freire, Caroline J Klivans, Pedro H Milet, and Nicolau C Saldanha. On the connectivity of spaces of three-dimensional tilings. *arXiv preprint arXiv:1702.00798*, 2017.
- [22] Q Ge and D Štefankovič. A graph polynomial for independent sets of bipartite graphs. *Combinatorics, Probability and Computing*, 21(5):695–714, 2012.
- [23] F. Graham and P. Tetali. Isoperimetric inequalities for cartesian products of graphs. *Comb. Probab. Comput.*, 7:141–148, 1998.

- [24] Venkatesan Guruswami. Rapidly mixing markov chains: A comparison of techniques (a survey). *ArXiv*, 2016. URL <https://arxiv.org/abs/1603.01512>.
- [25] Thomas P Hayes. A simple condition implying rapid mixing of single-site dynamics on spin systems. In *2006 47th Annual IEEE Symposium on Foundations of Computer Science (FOCS'06)*, pages 39–46. IEEE, 2006.
- [26] Thomas P. Hayes, Juan C. Vera, and Eric Vigoda. Randomly coloring planar graphs with fewer colors than the maximum degree. *Random Structures & Algorithms*, 47(4):731–759, 2015. doi: <https://doi.org/10.1002/rsa.20560>.
- [27] Marc Heinrich. Glauber dynamics for colourings of chordal graphs and graphs of bounded treewidth, 2020. URL <https://arxiv.org/abs/2010.16158>.
- [28] Peter J. Hilton and Jean J. Pedersen. Catalan numbers, their generalization, and their uses. *The Mathematical Intelligencer*, 13:64–75, 1991.
- [29] Lingxiao Huang, Pinyan Lu, and Chihao Zhang. Canonical paths for MCMC: from art to science. In *Proceedings of the twenty-seventh annual ACM-SIAM symposium on Discrete algorithms*, pages 514–527. SIAM, 2016.
- [30] Mrunmay Jagadale and Alok Laddha. Towards positive geometry of multi scalar field amplitudes : Accordiohedron and effective field theory. 2021. URL <https://arxiv.org/abs/2104.04915>.
- [31] Mark Jerrum, Jung-Bae Son, Prasad Tetali, and Eric Vigoda. Elementary bounds on Poincaré and log-Sobolev constants for decomposable Markov chains. *The Annals of Applied Probability*, 14(4):1741–1765, 2004. ISSN 10505164. URL <http://www.jstor.org/stable/4140446>.
- [32] Volker Kaibel. On the expansion of graphs of 0/1-polytopes. In *The Sharpest Cut: The Impact of Manfred Padberg and His Work*, pages 199–216. SIAM, 2004.
- [33] Arif Khan and Alex Pothen. A new 3/2-approximation algorithm for the b-edge cover problem. In *2016 Proceedings of the Seventh SIAM Workshop on Combinatorial Scientific Computing*, pages 52–61. SIAM, 2016.
- [34] Arif Khan, Alex Pothen, Md. Mostofa Ali Patwary, Nadathur Rajagopalan Satish, Narayanan Sundaram, Fredrik Manne, Mahantesh Halappanavar, and Pradeep Dubey. Efficient approximation algorithms for weighted b -matching. *SIAM Journal on Scientific Computing*, 38(5): S593–S619, 2016. doi: 10.1137/15M1026304.
- [35] David A. Klarner. Correspondences between plane trees and binary sequences. *Journal of Combinatorial Theory*, 9(4):401–411, 1970. ISSN 0021-9800. doi: [https://doi.org/10.1016/S0021-9800\(70\)80093-X](https://doi.org/10.1016/S0021-9800(70)80093-X). URL <https://www.sciencedirect.com/science/article/pii/S002198007080093X>.
- [36] David A Levin, Elizabeth Wilmer, and Yuval Peres. *Markov chains and mixing times*, volume 107. American Mathematical Society, 2009. URL <https://bookstore.ams.org/mbk-58>.
- [37] J. Loday. The multiple facets of the associahedron. In *Proc. 2005 Academy Coll. Series*, 2005.
- [38] Brendan Lucier, Michael Molloy, and Yuval Peres. The Glauber dynamics for colourings of bounded degree trees. In Irit Dinur, Klaus Jansen, Joseph Naor, and José Rolim, editors, *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, pages 631–645, Berlin, Heidelberg, 2009. Springer Berlin Heidelberg. ISBN 978-3-642-03685-9.

- [39] Neal Madras and Dana Randall. Markov chain decomposition for convergence rate analysis. *The Annals of Applied Probability*, 12(2):581 – 606, 2002. doi: 10.1214/aoap/1026915617.
- [40] Fabio Martinelli, Alistair Sinclair, and Dror Weitz. Fast mixing for independent sets, colorings, and other models on trees. *Random Structures & Algorithms*, 31(2):134–172, 2007. doi: <https://doi.org/10.1002/rsa.20132>.
- [41] Lisa McShine and P. Tetali. On the mixing time of the triangulation walk and other catalan structures. In *Randomization Methods in Algorithm Design*, 1997.
- [42] Milena Mihail and Umesh Vazirani. On the expansion of 0-1 polytopes. *Journal of Combinatorial Theory, Series B*, 1989.
- [43] Michael Molloy, Bruce Reed, and William Steiger. On the mixing rate of the triangulation walk. *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, 43, 09 2001.
- [44] Ben Morris and Alistair Sinclair. Random walks on truncated cubes and sampling 0-1 knapsack solutions. *SIAM journal on computing*, 34(1):195–226, 2004.
- [45] Atsuhiko Nakamoto, Gen Kawatani, Naoki Matsumoto, and Jorge Urrutia. Geometric quadrangulations of a polygon. *Electronic Notes in Discrete Mathematics*, 68:59–64, 2018. ISSN 1571-0653. doi: <https://doi.org/10.1016/j.endm.2018.06.011>. Discrete Mathematics Days 2018.
- [46] Nestor V. Nestoridis and Dimitrios M. Thilikos. Square roots of minor closed graph classes. *Discrete Applied Mathematics*, 168:34–39, 2014. ISSN 0166-218X. doi: <https://doi.org/10.1016/j.dam.2013.05.026>.
- [47] Yoshio Okamoto, Takeaki Uno, and Ryuhei Uehara. Linear-time counting algorithms for independent sets in chordal graphs. In *Graph-Theoretic Concepts in Computer Science*, pages 433–444. Springer Berlin Heidelberg, 2005. ISBN 978-3-540-31468-4.
- [48] Reinhard Pichler, Stefan Rümmele, and Stefan Woltran. Counting and enumeration problems with bounded treewidth. In *Logic for Programming, Artificial Intelligence, and Reasoning*, pages 387–404. Springer Berlin Heidelberg, 2010. ISBN 978-3-642-17511-4.
- [49] Herbert E. Robbins. A remark on stirling’s formula. *American Mathematical Monthly*, 62: 402–405, 1955.
- [50] Neil Robertson and P.D Seymour. Graph minors. ii. algorithmic aspects of tree-width. *Journal of Algorithms*, 7(3):309–322, 1986. ISSN 0196-6774. doi: [https://doi.org/10.1016/0196-6774\(86\)90023-4](https://doi.org/10.1016/0196-6774(86)90023-4).
- [51] Neil Robertson and P.D Seymour. Graph minors. X. obstructions to tree-decomposition. *Journal of Combinatorial Theory, Series B*, 52(2):153–190, 1991. ISSN 0095-8956. doi: [https://doi.org/10.1016/0095-8956\(91\)90061-N](https://doi.org/10.1016/0095-8956(91)90061-N).
- [52] P. D. Seymour and R. Thomas. Call routing and the ratcatcher. *Combinatorica*, 14(2):217–241, Jun 1994. ISSN 1439-6912. doi: 10.1007/BF01215352.
- [53] Alistair Sinclair. Improved bounds for mixing rates of Markov chains and multicommodity flow. *Combinatorics, Probability and Computing*, 1(4):351–370, 1992. doi: 10.1017/S0963548300000390.

- [54] Alistair Sinclair and Mark Jerrum. Approximate counting, uniform generation and rapidly mixing Markov chains. *Information and Computation*, 82(1):93–133, 1989. ISSN 0890-5401. doi: [https://doi.org/10.1016/0890-5401\(89\)90067-9](https://doi.org/10.1016/0890-5401(89)90067-9).
- [55] Daniel D Sleator, Robert E Tarjan, and William P Thurston. Rotation distance, triangulations, and hyperbolic geometry. *Journal of the American Mathematical Society*, 1(3):647–681, 1988.
- [56] Allan Sly. Computational transition at the uniqueness threshold. In *2010 IEEE 51st Annual Symposium on Foundations of Computer Science*, pages 287–296, 2010. doi: 10.1109/FOCS.2010.34.
- [57] Alexandre Stauffer. A Lyapunov function for Glauber dynamics on lattice triangulations. *Probability Theory and Related Fields*, 169:469–521, 2015.
- [58] Shai Vardi. Randomly coloring graphs of logarithmically bounded pathwidth. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2018)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2018.
- [59] Pengfei Wan, Jianhua Tu, Shenggui Zhang, and Binlong Li. Computing the numbers of independent sets and matchings of all sizes for graphs with bounded treewidth. *Applied Mathematics and Computation*, 332:42 – 47, 2018. ISSN 0096-3003. doi: <https://doi.org/10.1016/j.amc.2018.03.017>.