

Investigating quark confinement from the viewpoint of lattice gauge-scalar models

Ryu Ikeda^{a,*} and Kei-Ichi Kondo^{a,b}

^a*Department of Physics, Graduate School of Science and Engineering, Chiba University,
Chiba 263-8522, Japan*

^b*Department of Physics, Graduate School of Science, Chiba University,
Chiba 263-8522, Japan*

E-mail: cdna0955@chiba-u.jp, kondok@faculty.chiba-u.jp

In this talk, first, we show that the color N -dependent area law falloffs of the double-winding Wilson loop averages for the $SU(N)$ lattice gauge model are reproduced from the Z_N lattice Abelian gauge model due to the center group dominance in quark confinement. Next, we discuss lattice gauge-scalar models which allow analytic continuation for gauge invariant operators between confinement region and Higgs region. Applying the cluster expansion, we try to understand non-trivial contribution from scalar field in quark confinement mechanism. In order to understand quark confinement further, moreover, we study double-winding Wilson loop averages in the analytical region on the phase diagram.

*The 38th International Symposium on Lattice Field Theory, LATTICE2021 26th-30th July, 2021
Zoom/Gather@Massachusetts Institute of Technology*

*Speaker

1. Introduction

In the lattice gauge theory, a double-winding Wilson loop operator $W(C_1 \cup C_2)$ has been introduced in [1] to examine the possible mechanisms for quark confinement. The double-winding Wilson loop operator is defined as a trace of the path-ordered product of gauge link variables U_ℓ along a closed loop C composed of two loops C_1 and C_2 :

$$W(C_1 \cup C_2) \equiv \text{tr} \left[\prod_{\ell \in C_1 \cup C_2} U_\ell \right]. \quad (1)$$

The double-winding Wilson loop is called *coplanar* if the two loops C_1 and C_2 lie in the same plane, while it is called *shifted* if the two loops C_1 and C_2 lie in planes parallel to the x - t plane, but are displaced from one another in the transverse z -direction by distance R , and are connected by lines running parallel to the z -axis to keep the gauge invariance. See Fig.1. Note that the double-winding Wilson loop operators are defined as a gauge invariant operator.




Figure 1: (a) a “coplanar” double-winding Wilson loop, (b) a “shifted” double-winding Wilson loop.

The area dependence of the expectation value $\langle W(C_1 \cup C_2) \rangle$ has been first investigated in [1] to show that the coplanar double-winding Wilson loop average obeys the “difference-of-areas law” in the lattice $SU(2)$ Yang-Mills model by using the strong coupling expansion and the numerical simulations:

$$\langle W(C_1 \cup C_2) \rangle_{R=0} \simeq \exp[-\sigma ||S_1| - |S_2||], \quad (2)$$

where S_1 and S_2 are respectively the minimal areas bounded by loops C_1 and C_2 .

In the continuum $SU(N)$ Yang-Mills model, general multiple-winding Wilson loops have been investigated in [2] to show that there is a novel “max-of-areas law” which is neither difference-of-areas law nor sum-of-areas law for multiple-winding Wilson loop average, provided that the string tension obeys the Casimir scaling for quarks in the higher representations.

In the lattice $SU(N)$ Yang-Mills model, it has been shown in [3] that the coplanar double-winding Wilson loop average has the N -dependent area law falloff in the strong coupling region: “difference-of-areas law” for $N = 2$, “max-of-areas law” for $N = 3$ and “sum-of-areas law” for

$N \geq 4$:

$$\langle W(C_1 \cup C_2) \rangle_{R=0} \simeq \begin{cases} \exp[-\sigma(|S_1| - |S_2|)] & (N = 2) \\ \exp[-\sigma \max(|S_1|, |S_2|)] & (N = 3) \\ \exp[-\sigma(|S_1| + |S_2|)] & (N \geq 4) \end{cases} . \quad (3)$$

Moreover, a shifted double-winding Wilson loop average as a function of the distance R in a transverse direction has the long distance behavior which does not depend on N , while the short distance behavior depends on N .

In our investigation in [4], we examine the center group dominance for a double winding Wilson loop average. It has been shown in [5] that the ordinary single-winding Wilson loop average in the non-Abelian lattice gauge theory with the gauge group G is bounded from above by the same Wilson loop average in the Abelian lattice gauge theory with the center gauge group $Z(G)$:

$$|\langle W_{R(G)}(C) \rangle_G(\beta)| \leq 2\text{tr}(\mathbf{1}) \langle W_{R(Z(G))}(C) \rangle_{Z(G)}(2\dim(G)\beta) . \quad (4)$$

We have extended the above statement to the double winding Wilson loop average, beyond the case of the ordinary single-winding Wilson loop average:

$$|\langle W_{R(G)}(C_1 \cup C_2) \rangle_G(\beta)| \leq 2\text{tr}(\mathbf{1}) \langle W_{R(Z(G))}(C_1 \cup C_2) \rangle_{Z(G)}(2\dim(G)\beta) . \quad (5)$$

From this point of view, we introduce the *character expansion* to the weight $e^{S_G[U]}$ coming from the action and perform the group integration, in order to estimate the expectation value in the Z_N lattice gauge model. We evaluate the double-winding Wilson loop average up to the leading contribution to show that the N -dependent area law falloff in the $SU(N)$ lattice gauge model can be reproduced by using the (Abelian) Z_N lattice gauge model. By taking the limit $N \rightarrow \infty$, we show the center group dominance for a double-winding Wilson loop average in the $U(N)$ lattice gauge model through the $U(1)$ lattice gauge model.

Finally, we extend the above arguments for the lattice gauge-scalar model on the “analytic region”. For this purpose, we estimate the area law falloff, the string tension, and the mass gap by using the *cluster expansion*.

2. Lattice Z_N gauge model

First, we consider the lattice Z_N gauge model with the coupling constant defined by $\beta := 1/g^2$ on a D -dimensional lattice Λ with unit lattice spacing, which is specified by the action

$$S_G[U] = \beta \sum_{p \in \Lambda} \text{Re } U_p , \quad U_p := \prod_{\ell \in \partial p} U_\ell , \quad (6)$$

where ℓ labels a link, p labels an elementary plaquette. To examine this Z_N gauge model analytically, we introduce the *character expansion* to the weight $e^{S_G[U]}$ to obtain the expanded form of the expectation value of an operator \mathcal{F} :

$$\langle \mathcal{F} \rangle_\Lambda := Z_\Lambda^{-1} \int \prod_{\ell \in \Lambda} dU_\ell e^{S_G[U]} \mathcal{F} = Z_\Lambda^{-1} \int \prod_{\ell \in \Lambda} dU_\ell \prod_{p \in \Lambda} \sum_{n=0}^{N-1} b_n(\beta) U_p^n \mathcal{F} , \quad (7)$$

$$Z_\Lambda := \int \prod_{\ell \in \Lambda} dU_\ell e^{S_G[U]} , \quad (8)$$

where the coefficients $b_n(\beta)$ is defined by

$$b_n(\beta) := \frac{1}{N} \sum_{\zeta \in Z_N} \zeta^{-n} e^{\beta \operatorname{Re} \zeta} . \quad (9)$$

We define $c_n(\beta) := b_n(\beta)/b_0(\beta)$. For $N = 2, 3, 4$ and ∞ , $c_1(\beta)$ and $c_2(\beta)$ are written in the form

$$\begin{aligned} c_1(\beta) &= \frac{e^\beta - e^{-\beta}}{e^\beta + e^{-\beta}} \quad (N = 2) , & c_1(\beta) &= \frac{e^\beta - e^{-\beta/2}}{e^\beta + 2e^{-\beta/2}} = c_2(\beta) \quad (N = 3) , \\ c_1(\beta) &= \frac{e^\beta - e^{-\beta}}{e^\beta + 2 + e^{-\beta}} , & c_2(\beta) &= \frac{e^\beta - 2 + e^{-\beta}}{e^\beta + 2 + e^{-\beta}} \quad (N = 4) , \\ c_1(\beta) &= \frac{I_1(\beta)}{I_0(\beta)} , & c_2(\beta) &= \frac{I_2(\beta)}{I_0(\beta)} \quad (N = \infty) . \end{aligned} \quad (10)$$

Note that $b_{N-n}(\beta) = b_n(\beta)$ and $0 \leq c_n(\beta) < 1$ for $0 \leq \beta < \infty$. For $N = 2, 3, 4$ and ∞ , the behavior of $c_1(\beta)$ and $c_2(\beta)$ as functions of β are indicated in Fig.2. We find that $c_1(\beta) \sim O(\beta)$ ($N \geq 2$) and $c_2(\beta) \sim O(\beta^2)$ ($N \geq 4$) for $\beta \ll 1$.




Figure 2: The character expansion coefficient as a function of β , (a) $c_1(\beta)$, (b) $c_2(\beta)$

Next, we evaluate the expectation value of a coplanar double-winding Wilson loop in the lattice Z_N pure gauge model. The leading contribution to a coplanar double-winding Wilson loop average is given by the tiling of a planar set of plaquettes, as shown in the Fig.3. (These result are exact for all β when $D = 2$, while valid for $\beta \ll 1$ when $D > 2$.)

The result of the coplanar double-winding Wilson loop average up to the leading contribution is given by

$$\langle W(C_1 \cup C_2) \rangle_{R=0} \simeq \begin{cases} c_1(\beta)^{|S_1| - |S_2|} & (N = 2) \\ c_1(\beta)^{|S_1|} & (N = 3) \\ c_2(\beta)^{|S_2|} c_1(\beta)^{|S_1| - |S_2|} & (N \geq 4) \end{cases} . \quad (11)$$

Then we obtain the (non-zero) string tension from this result:

$$\sigma(\beta) \simeq \ln \frac{1}{c_1(\beta)} > 0 . \quad (12)$$




Figure 3: A coplanar double-winding Wilson loop, (a) $N = 2$, (b) $N \geq 3$

In the strong coupling region, this result reproduces the area law falloff in the $SU(N)$ lattice gauge model obtained in [3]. Moreover, by taking the continuous group limit $N \rightarrow \infty$, we find that the area law for $N \geq 4$ persists in the $U(1)$ lattice gauge model.

Furthermore, we also evaluate the expectation value of a shifted double-winding Wilson loop in the lattice Z_N pure gauge model. The leading contribution to a shifted double-winding Wilson loop average can be given by the 2 types of tiling by a set of plaquettes, as shown in the Fig.4.




Figure 4: A shifted double-winding Wilson loop, (a) R -independent contribution, (b) R -dependent contribution

The result of the shifted double-winding Wilson loop average up to the leading contribution is given by

$$\langle W(C_1 \cup C_2) \rangle_{R \neq 0} \simeq \begin{cases} c_1(\beta)^{|S_1|+|S_2|} + c_1(\beta)^{2R(L_2+T)} \cdot c_1(\beta)^{|S_1|-|S_2|} & (N = 2) \\ c_1(\beta)^{|S_1|+|S_2|} + c_1(\beta)^{2R(L_2+T)} \cdot c_1(\beta)^{|S_1|} & (N = 3) \\ c_1(\beta)^{|S_1|+|S_2|} + c_1(\beta)^{2R(L_2+T)} \cdot c_2(\beta)^{|S_2|} c_1(\beta)^{|S_1|-|S_2|} & (N \geq 4) \end{cases} \quad (13)$$

This result reproduces the R -dependent behavior of the shifted double-winding Wilson loop average in [3]. In particular, we obtain the (non-zero) mass gap from the case of $S_1 = S_2 = 1$ and $R \gg 1$ in the above result:

$$\Delta(\beta) = 4 \ln \frac{1}{c_1(\beta)} > 0. \quad (14)$$

3. Lattice Z_N gauge-scalar theory

Next, we consider the lattice Z_N gauge-scalar model with the frozen scalar field norm R for simplicity. The action of this model with the coupling constants defined by $\beta := 1/g^2$ and $K := R^2$ on a D -dimensional lattice Λ with unit lattice spacing is given by

$$S[U, \varphi] = \beta \sum_{p \in \Lambda} \text{Re } U_p + K \sum_{\ell \in \Lambda} \text{Re} (\varphi_x U_\ell \varphi_{x+\ell}^*) , \quad (15)$$

where ℓ labels a link, and p labels an elementary plaquette. U_ℓ is a Z_N link variable on link ℓ and φ_x is a Z_N scalar field at site x which transforms according to the fundamental representation of the gauge group Z_N .

In this model, the expectation value of an operator \mathcal{F} has the form

$$\begin{aligned} \langle \mathcal{F} \rangle_\Lambda &:= Z_\Lambda^{-1} \int \prod_{\ell \in \Lambda} dU_\ell \prod_{x \in \Lambda} d\varphi_x e^{S[U, \varphi]} \mathcal{F} = Z_\Lambda^{-1} \int \prod_{\ell \in \Lambda} dU_\ell h[U] e^{\beta \sum_{p \in \Lambda} \text{Re } U_p} \mathcal{F} , \\ Z_\Lambda &:= \int \prod_{\ell \in \Lambda} dU_\ell \prod_{x \in \Lambda} d\varphi_x e^{S[U, \varphi]} , \quad h[U] := \int \prod_{x \in \Lambda} d\varphi_x e^{K \sum_{\ell \in \Lambda} \text{Re} (\varphi_x U_\ell \varphi_{x+\ell}^*)} . \end{aligned} \quad (16)$$

According to [6], we can perform the *cluster expansion* by introducing the new variable ρ_p and the new measure $d\mu_\Lambda$ which absorbs the scalar part $h[U]$:

$$\langle \mathcal{F} \rangle_\Lambda = \frac{\int d\mu_\Lambda \prod_{p \in \Lambda} (1 + \rho_p) \mathcal{F}}{\int d\mu_\Lambda \prod_{p \in \Lambda} (1 + \rho_p)} = \sum_{Q(Q_0) \subset \Lambda} \int d\mu_\Lambda \mathcal{F} \prod_{p \in Q(Q_0)} \rho_p \cdot \frac{Z_{[Q(Q_0) \cup Q_0]^c}}{Z_\Lambda} , \quad (17)$$

$$d\mu_\Lambda := \frac{\prod_{\ell \in \Lambda} dU_\ell h[U]}{\int \prod_{\ell \in \Lambda} dU_\ell h[U]} , \quad \rho_p := e^{\beta \text{Re } U_p} - 1 , \quad (18)$$

where Q_0 is the set of plaquettes which is the support of \mathcal{F} and $Q(Q_0)$ is the set of plaquettes which is connected to Q_0 . For the general set of plaquettes Q , Q^c represents the complement of Q . Here, Z_Q is defined by

$$Z_Q := \sum_{Q' \subset Q} \int d\mu_\Lambda \prod_{p \in Q'} \rho_p . \quad (19)$$

Note that $\rho_p \sim O(\beta)$ for $\beta \ll 1$. It has been showed in [7] that the confinement region ($0 \leq \beta \ll 1, K \ll 1$) and the Higgs region ($\beta \gg 1, K_c \leq K < \infty$) are analytically continued in a single “analytic region”, where the cluster expansion converges uniformly. See Fig.5.

To evaluate $h[U]$, we apply the character expansion and perform the group integration. Ignoring the contributions from multiple plaquettes, then we obtain the expression which is valid up to the lowest plaquettes order:

$$\begin{aligned} h[U] &= \int \prod_{x \in \Lambda} d\varphi_x \prod_{\ell \in \Lambda} \left[b_0(K) + b_1(K) \varphi_x U_\ell \varphi_{x+\ell}^* + \cdots + b_{N-1}(K) (\varphi_x U_\ell \varphi_{x+\ell}^*)^{N-1} \right] \\ &= N^{|\Lambda|} b_0(K)^{D|\Lambda|} \prod_{p \in \Lambda} \sum_{n=0}^{N-1} c_n(K)^4 U_p^n + \cdots . \end{aligned} \quad (20)$$

We estimate the leading contribution to the double-winding Wilson loop average with the above $h[U]$, we also apply the character expansion for ρ_p and evaluate the upper bound of the cluster




Figure 5: The analytic region on the β - K plane

expansion by using the binominal expansion. We find that there is an correspondence between the evaluation for the Z_N lattice gauge model and for the estimated upper bound for the Z_N lattice gauge-scalar model:

$$c_n(\beta) \mapsto a_n(\beta, K) := \frac{[b_0(\beta) - e^\beta] c_n(K)^4 + b_1(\beta) c_{n+1}(K)^4 + \cdots + b_{N-1}(\beta) c_{N+n-1}(K)^4}{b_0(\beta) + b_1(\beta) c_1(K)^4 + \cdots + b_{N-1}(\beta) c_{N-1}(K)^4} + c_n(K)^4 .$$

$$(\text{mod } N, n = 1, \dots, N-1) \quad (21)$$

Note that $a_n(\beta, 0) = c_n(\beta)$ and $a_n(\beta, \infty) = 1$. The above estimation is valid only for the values of parameter β and K on the analytic region in the range where the string breaking does not occur.

By applying the same method as the above, we obtain the estimation for the coplanar double-winding Wilson loop average:

$$\langle W(C_1 \cup C_2) \rangle_{R=0} \lesssim \begin{cases} a_1(\beta, K)^{|S_1| - |S_2|} & (N = 2) \\ a_1(\beta, K)^{|S_1|} & (N = 3) \\ a_2(\beta, K)^{|S_2|} a_1(\beta, K)^{|S_1| - |S_2|} & (N \geq 4) \end{cases} \quad (22)$$

and we obtain the (non-zero) string tension from the above result:

$$\sigma(\beta, K) \gtrsim \ln \frac{1}{a_1(\beta, K)} > 0 . \quad (23)$$

This estimation suggests that the area law falloff in the Z_N lattice gauge model persists in the Z_N lattice gauge-scalar model and the $K \rightarrow 0$ limit agrees with the pure gauge case. Moreover, for $\sigma(\beta, K)$, the $K \rightarrow 0$ limit agrees with $\sigma(\beta)$ in the Z_N lattice gauge model, and $K \rightarrow \infty$ limit converges to 0 uniformly in β . In other words, the string tension is non-zero on the analytic region.

Additionally, we also estimate the shifted double-winding Wilson loop average:

$$\langle W(C_1 \cup C_2) \rangle_{R \neq 0} \lesssim \begin{cases} a_1(\beta, K)^{|S_1| + |S_2|} + a_1(\beta, K)^{2R(L_2+T)} \cdot a_1(\beta, K)^{|S_1| - |S_2|} & (N = 2) \\ a_1(\beta, K)^{|S_1| + |S_2|} + a_1(\beta, K)^{2R(L_2+T)} \cdot a_1(\beta, K)^{|S_1|} & (N = 3) \\ a_1(\beta, K)^{|S_1| + |S_2|} + a_1(\beta, K)^{2R(L_2+T)} \cdot a_2(\beta, K)^{|S_2|} a_1(\beta, K)^{|S_1| - |S_2|} & (N \geq 4) \end{cases} \quad (24)$$

and we obtain the (non-zero) mass gap from the case of $S_1 = S_2 = 1$ and $R \gg 1$ in the above result:

$$\Delta(\beta, K) \gtrsim 4 \ln \frac{1}{a_1(\beta, K)} > 0. \quad (25)$$

For $\Delta(\beta, K)$, the $K \rightarrow 0$ limit agrees with $\Delta(\beta)$ in the Z_N lattice gauge model, and $K \rightarrow \infty$ limit converges to 0 uniformly in β . In other words, the mass gap is non-zero on the analytic region.

4. Conclusion

We investigated the area law falloff of the double-winding Wilson loops in the Z_N lattice gauge model and Z_N lattice gauge-scalar model, where the gauge group is the center group of the original $SU(N)$. First, we evaluated the N -dependent area law falloff for the coplanar double-winding Wilson loop average up to the leading contribution. We found the N -dependence of the area law falloff in the Z_N lattice gauge model, which reproduces the area law falloff in the $SU(N)$ lattice gauge model obtained in [3]. Secondly, we also checked the limit $N \rightarrow \infty$, the area law falloff for $N \geq 4$ persists in the $U(1)$ lattice gauge model. This result implies that the coplanar double-winding Wilson loop average in the $U(N)$ lattice gauge model and the $SU(N)$ ($N \geq 4$) lattice gauge model obeys the same area law up to the leading contribution. Furthermore, we also considered the shifted double-winding Wilson loop average up to the leading contributions. This result reproduces the R -dependent behavior in the $SU(N)$ lattice gauge model obtained in [3]. We obtained the (non-zero) mass gap $\Delta(\beta)$ from this result. Finally, we extended the above study for the Z_N lattice gauge-scalar model on the analytic region. We found that the area law falloff in the Z_N lattice gauge model persists in the Z_N lattice gauge-scalar model. We discovered that the string tension and the mass gap are non-zero on the analytic region from this estimation.

References

- [1] J. Greensite and R. Höllwieser, Phys. Rev. **D91**, 054509 (2015) [arXiv:1411.5091 [hep-lat]]
- [2] R. Matsudo and K.-I. Kondo, Phys. Rev. **D96**, 105011 (2017) [arXiv:1706.05665 [hep-th]]
- [3] S. Kato, A. Shibata and K.-I. Kondo, Phys.Rev. **D102**, 094521 (2020) [arXiv:2008.03684 [hep-lat]]
- [4] R. Ikeda and K.-I. Kondo, Prog. Theor. Exp. Phys. **2021**, ptab114 (2021) [arXiv:2106.14416 [hep-lat]]
- [5] J. Fröhlich, Phys. Lett. **83B**, 195 (1979).
- [6] K. Osterwalder and E. Seiler, Annals Phys. **110**, 440 (1978).
- [7] E.H. Fradkin and S.H. Shenker, Phys.Rev. **D19**, 3682–3697 (1979).