

Extremal Problem for Matchings and Rainbow Matchings on Direct Products

Jian Wang¹, Jie You²

¹Department of Mathematics
Taiyuan University of Technology
Taiyuan 030024, P. R. China

²Center for Applied Mathematics
Tianjin University
Tianjin 300072, P. R. China

E-mail: ¹wangjian01@tyut.edu.cn, ²yj_math@tju.edu.cn

Abstract

Let $n_1, \dots, n_\ell, k_1, \dots, k_\ell$ be integers and let V_1, \dots, V_ℓ be disjoint sets with $|V_i| = n_i$ for $i = 1, \dots, \ell$. Define $\sqcup_{i=1}^\ell \binom{V_i}{k_i}$ as the collection of all subsets F of $\cup_{i=1}^\ell V_i$ with $|F \cap V_i| = k_i$ for each $i = 1, \dots, \ell$. In this paper, we show that if the matching number of $\mathcal{F} \subseteq \sqcup_{i=1}^\ell \binom{V_i}{k_i}$ is at most s and $n_i \geq 4\ell^2 k_i^2 s$ for all i , then $|\mathcal{F}| \leq \max_{1 \leq i \leq \ell} \left[\binom{n_i}{k_i} - \binom{n_i-s}{k_i} \right] \prod_{j \neq i} \binom{n_j}{k_j}$. Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s \subseteq \sqcup_{i=1}^\ell \binom{V_i}{k_i}$ with $n_i \geq 8\ell^2 k_i^2 s$ for all i . We also prove that if $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$ are rainbow matching free, then there exists t in $[s]$ such that $|\mathcal{F}_t| \leq \max_{1 \leq i \leq \ell} \left[\binom{n_i}{k_i} - \binom{n_i-s+1}{k_i} \right] \prod_{j \neq i} \binom{n_j}{k_j}$.

1 Introduction

Let $[n] = \{1, 2, \dots, n\}$ and $\binom{[n]}{k}$ be the collection of all its k -subsets. For a k -graph $\mathcal{F} \subset \binom{[n]}{k}$, the matching number of \mathcal{F} , denote by $\nu(\mathcal{F})$, is the maximum number of pairwise disjoint members of \mathcal{F} . Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s \subseteq \binom{[n]}{k}$. We say that $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$ contain a rainbow matching if there exist s pairwise disjoint sets $F_1 \in \mathcal{F}_1, F_2 \in \mathcal{F}_2, \dots, F_s \in \mathcal{F}_s$.

One of the most important open problems in extremal set theory is to determine the maximum of $|\mathcal{F}|$ subject to $\nu(\mathcal{F}) \leq s$, which is known as the celebrated Erdős matching conjecture.

Conjecture 1.1 (Erdős Matching Conjecture [5]). *Let $n \geq (s+1)k$ and $\mathcal{F} \subseteq \binom{[n]}{k}$. If $\nu(\mathcal{F}) \leq s$, then*

$$|\mathcal{F}| \leq \max \left\{ \binom{k(s+1)-1}{k}, \binom{n}{k} - \binom{n-s}{k} \right\}. \quad (1.1)$$

The two simple constructions showing that (1.1) is best possible (if true) are

$$\mathcal{E}(n, k, s) = \left\{ E \in \binom{[n]}{k} : E \cap [s] \neq \emptyset \right\} \text{ and } \binom{[(s+1)k-1]}{k}.$$

The case $s = 1$ is the classical Erdős-Ko-Rado Theorem [15]. Erdős and Gallai [6] proved (1.1) for $k = 2$. The $k = 3$ case is settled in [10]. For general k , Erdős proved that (1.1) is true and up to isomorphic $\mathcal{E}(n, k, s)$ is the only optimal family provided that $n > n_0(k, s)$. The bounds for $n_0(k, s)$ were subsequently improved by Bollobás, Daykin and Erdős [3], Huang, Loh and Sudakov [14]. The current best bounds establish (1.1) for $n > (2s+1)k$ ([9]) and for $s > s_0$, $n > \frac{5}{3}sk$ ([11]).

A rainbow version of Erdős matching conjecture was proposed by Aharoni, Howard and [1] and independently Huang, Loh, Sudakov [14].

Conjecture 1.2 ([1, 14]). *Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s \subseteq \binom{[n]}{k}$. If*

$$|\mathcal{F}_i| > \max \left\{ \binom{ks-1}{k}, \binom{n}{k} - \binom{n-s+1}{k} \right\}, \quad (1.2)$$

for all $1 \leq i \leq s$, then $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$ contain a rainbow matching.

Using a junta method, Keller and Lifshitz [16] showed the validity of (1.2) for $n > f(s)k$, where $f(s)$ is an unspecified and a very quickly growing function of s . In [12], Frankl and Kupavskii improved the bound to $12sk(2 + \log s)$. By using the absorbing method, Lu, Wang and Yu [19] proved (1.2) for $n \geq 2k(s+1)$ and $s > s_0(k)$. By applying the concentration inequality for random matchings developed in [11], Kupavskii [17] proved (1.2) for $n \geq 3e(s+1)k$ and $s > 10^7$.

Let $n_1, \dots, n_\ell, k_1, \dots, k_\ell$ be integers with $n_1/k_1 \leq \dots \leq n_\ell/k_\ell$ and let V_1, \dots, V_ℓ be pairwise disjoint sets with $|V_i| = n_i$. Define the direct product $\sqcup_{i=1}^\ell \binom{V_i}{k_i}$ as the collection of all subsets F of $\cup_{i=1}^\ell V_i$ with $|F \cap V_i| = k_i$ for $i = 1, \dots, \ell$. By the algebraic method, Frankl [8] proved an Erdős-Ko-Rado theorem for direct products as follows:

Theorem 1.1 ([8]). *Suppose that $\mathcal{F} \subseteq \sqcup_{i=1}^\ell \binom{V_i}{k_i}$ is intersecting and $2 \leq n_1/k_1 \leq \dots \leq n_\ell/k_\ell$. Then*

$$|\mathcal{F}| \leq \frac{k_1}{n_1} \prod_{i=1}^\ell \binom{n_i}{k_i}.$$

In this paper, we mainly consider a direct product version of Erdős matching conjecture. We determine the maximum number of edges in a subfamily of the direct product with matching number at most s for each n_i sufficiently large.

Theorem 1.2. *Let $\ell, s, n_1, \dots, n_\ell, k_1, \dots, k_\ell$ be integers with $n_i \geq 4\ell^2 k_i^2 s$ for $i = 1, \dots, \ell$. Let $\mathcal{F} \subseteq \sqcup_{i=1}^\ell \binom{V_i}{k_i}$ with $|V_i| = n_i$. If $\nu(\mathcal{F}) \leq s$, then*

$$|\mathcal{F}| \leq \max_{1 \leq i \leq \ell} \left[\binom{n_i}{k_i} - \binom{n_i - s}{k_i} \right] \prod_{\substack{j \neq i \\ 1 \leq j \leq \ell}} \binom{n_j}{k_j}. \quad (1.3)$$

Let S_i be an s -subset of V_i . Set $\vec{n} = (n_1, n_2, \dots, n_\ell)$ and $\vec{k} = (k_1, k_2, \dots, k_\ell)$. Define

$$\mathcal{E}_i(\vec{n}, \vec{k}, s) = \left\{ E \in \bigsqcup_{j=1}^{\ell} \binom{V_j}{k_j} : E \cap S_i \neq \emptyset \right\}.$$

The constructions $\mathcal{E}_i(\vec{n}, \vec{k}, s)$ show that (1.3) is best possible.

By the random matching technique developed by Frankl and Kupavskii [11], we obtain a condition on the number of edges for a family of direct products to contain a rainbow matching for each n_i sufficiently large.

Theorem 1.3. *Let $\ell, s, n_1, \dots, n_\ell, k_1, \dots, k_\ell$ be integers with $n_i \geq 8\ell^2 k_i^2 s$ for $i = 1, \dots, \ell$. Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s \subseteq \bigsqcup_{i=1}^{\ell} \binom{V_i}{k_i}$ with $|V_i| = n_i$. If $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$ are rainbow matching free, then there exists $1 \leq t \leq s$ such that*

$$|\mathcal{F}_t| \leq \max_{1 \leq i \leq \ell} \left[\binom{n_i}{k_i} - \binom{n_i - s + 1}{k_i} \right] \prod_{\substack{j \neq i \\ 1 \leq j \leq \ell}} \binom{n_j}{k_j}. \quad (1.4)$$

The constructions $\mathcal{F}_1 = \mathcal{F}_2 = \dots = \mathcal{F}_s = \mathcal{E}_i(\vec{n}, \vec{k}, s)$ show that (1.4) is best possible.

2 Extremal problem for matchings on direct products

Frankl and the first author [13] proved an upper bound on the sum of sizes of overlapping families. In this section, by a probabilistic argument we transfer it to an upper bound on the size of $\mathcal{F} \subseteq \bigsqcup_{i=1}^{\ell} \binom{V_i}{k_i}$ with matching number at most s . By using this upper bound, we prove Theorem 1.2.

Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_m \subseteq \binom{[n]}{k}$ and $m \geq s + 1$. If one cannot choose pairwise disjoint edges from $s + 1$ distinct families of $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_m$, we say these families are s -overlapping.

Lemma 2.1 ([13]). *Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_m \subseteq \binom{[n]}{k}$ be s -overlapping families with $n \geq (s + 1)k$ and $m \geq s + 1$. Then*

$$\sum_{i=1}^m |\mathcal{F}_i| \leq \max \left\{ s \binom{n}{k}, ms \binom{n-1}{k-1} \right\}.$$

By Lemma 2.1, we prove the following lemma via a probabilistic argument.

Lemma 2.2. *Let $\mathcal{F} \subseteq \bigsqcup_{i=1}^{\ell} \binom{V_i}{k_i}$ with $|V_i| = n_i$ for $i = 1, \dots, \ell$ and $s+1 \leq n_1/k_1 \leq \dots \leq n_\ell/k_\ell$. If $\nu(\mathcal{F}) \leq s$, then*

$$|\mathcal{F}| \leq \frac{(s+1)k_1}{n_1} \prod_{i=1}^{\ell} \binom{n_i}{k_i}.$$

Proof. Let $m = \lfloor \frac{n_2}{k_2} \rfloor$ and let B_1, B_2, \dots, B_m be an m -matching chosen from $\bigsqcup_{i=2}^{\ell} \binom{V_i}{k_i}$ uniformly at random. Define

$$\mathcal{F}(B_i) = \left\{ A \in \binom{V_1}{k_1} : A \cup B_i \in \mathcal{F} \right\}$$

for $i = 1, \dots, m$. Since $\nu(\mathcal{F}) \leq s$, $\mathcal{F}(B_1), \dots, \mathcal{F}(B_m)$ are s -overlapping. Note that $m \geq s + 1$. By Lemma 2.1, we have

$$\sum_{i=1}^m |\mathcal{F}(B_i)| \leq \max \left\{ s \binom{n_1}{k_1}, ms \binom{n_1 - 1}{k_1 - 1} \right\}.$$

$n_1/k_1 \leq n_2/k_2$ implies that $n_1/k_1 \leq \lfloor n_2/k_2 \rfloor + 1 = m + 1$. It follows that

$$\sum_{i=1}^m |\mathcal{F}(B_i)| \leq (m + 1)s \binom{n_1 - 1}{k_1 - 1}. \quad (2.1)$$

Taking the expectation on both sides of (2.1), we obtain that

$$\sum_{i=1}^m \mathbb{E}|\mathcal{F}(B_i)| \leq (m + 1)s \binom{n_1 - 1}{k_1 - 1}. \quad (2.2)$$

Since B_1, B_2, \dots, B_m are chosen from $\sqcup_{i=2}^\ell \binom{V_i}{k_i}$ uniformly, we have

$$\mathbb{E}|\mathcal{F}(B_i)| = \sum_{B \in \sqcup_{i=2}^\ell \binom{V_i}{k_i}} |\mathcal{F}(B)| \cdot \frac{1}{\prod_{i=2}^\ell \binom{n_i}{k_i}} = \frac{|\mathcal{F}|}{\prod_{i=2}^\ell \binom{n_i}{k_i}}. \quad (2.3)$$

By (2.2) and (2.3), we have

$$|\mathcal{F}| \leq \left(1 + \frac{1}{m}\right) s \binom{n_1 - 1}{k_1 - 1} \prod_{i=2}^\ell \binom{n_i}{k_i}.$$

Since $m = \lfloor \frac{n_2}{k_2} \rfloor > s$, it follows that

$$\left(1 + \frac{1}{m}\right) s < s + 1$$

and the lemma follows. ■

The following operation, called shifting, was invented by Erdős, Ko and Rado [4] and further developed by Frankl [7]. Let $1 \leq i < j \leq n$, $\mathcal{F} \subset \binom{[n]}{k}$. Define

$$S_{ij}(\mathcal{F}) = \{S_{ij}(F) : F \in \mathcal{F}\},$$

where

$$S_{ij}(F) = \begin{cases} (F \setminus \{j\}) \cup \{i\}, & j \in F, i \notin F \text{ and } (F \setminus \{j\}) \cup \{i\} \notin \mathcal{F}; \\ F, & \text{otherwise.} \end{cases}$$

It is well known (cf. [7]) that shifting does not increase the matching number.

Let us recall some properties of the shifting operator. Let $\mathcal{F} \subseteq \sqcup_{i=1}^\ell \binom{V_i}{k_i}$ with $V_i = \{v_{i,1}, v_{i,2}, \dots, v_{i,n_i}\}$ for $i = 1, \dots, \ell$. Define a partial order \prec on $\cup_{i=1}^\ell V_i$ such that

$$v_{i,1} \prec v_{i,2} \prec \dots \prec v_{i,n_i}$$

for each i and vertices from different parts are incomparable. A family $\mathcal{F} \subseteq \sqcup_{i=1}^{\ell} \binom{V_i}{k_i}$ is called shifted if $S_{ab}(\mathcal{F}) = \mathcal{F}$ for all $a, b \in \cup_{i=1}^{\ell} V_i$ with $a \prec b$. Frankl [7] proved that repeated application of shifting to any family eventually produces a shifted family.

Let $k = k_1 + k_2 + \dots + k_{\ell}$. For two different edges $A = \{a_1, a_2, \dots, a_k\}$ and $B = \{b_1, b_2, \dots, b_k\}$ in $\sqcup_{i=1}^{\ell} \binom{V_i}{k_i}$, define $A \prec B$ iff there exists a permutation $\sigma_1 \sigma_2 \dots \sigma_k$ of $[k]$ such that $a_j \prec b_{\sigma_j}$ or $a_j = b_{\sigma_j}$ for $j = 1, \dots, k$. It is shown in [7] that if \mathcal{F} is a shifted family, $A \prec B$ and $B \in \mathcal{F}$ then $A \in \mathcal{F}$ as well.

In the proof of the following lemma, we need the following inequality, which were already used in [3]. For $n \geq (2s+1)k$,

$$\binom{n-s}{k} / \binom{n}{k} \geq \left(1 - \frac{s}{n-k}\right)^k \geq 1 - \frac{ks}{n-k} \geq \frac{1}{2}. \quad (2.4)$$

Lemma 2.3. *Let $\ell, s, n_1, \dots, n_{\ell}, k_1, \dots, k_{\ell}$ be integers with $n_i \geq 4\ell^2 k_i^2 s$ for $i = 1, \dots, \ell$. If $\mathcal{F} \subseteq \sqcup_{i=1}^{\ell} \binom{V_i}{k_i}$ has matching number at most s , then*

$$|\mathcal{F}| \leq \prod_{i=1}^{\ell} \binom{n_i}{k_i} - \min_{x_1 + \dots + x_{\ell} = s} \prod_{i=1}^{\ell} \binom{n_i - x_i}{k_i}$$

Proof. Without loss of generality, we may assume that $n_1/k_1 \leq \dots \leq n_{\ell}/k_{\ell}$ and \mathcal{F} is shifted. We prove the lemma by induction on s . The case $s = 1$ is verified in Theorem 1.1. Assume that the lemma holds for the case $s-1$. Let $V_i = \{v_{i,j} : j \in [n_i]\}$. For each $x \in \cup_{i=1}^{\ell} V_i$, define

$$\mathcal{F}(x) = \{F \setminus \{x\} : x \in F \in \mathcal{F}\} \text{ and } \mathcal{F}(\overline{x}) = \{F : x \notin F \in \mathcal{F}\}.$$

We distinguish two cases.

Case 1. $\nu(\mathcal{F}(\overline{v_{i,1}})) \leq s-1$ for some $i \in [\ell]$.

By the induction hypothesis, we have

$$|\mathcal{F}(\overline{v_{i,1}})| \leq \binom{n_i-1}{k_i} \prod_{j \neq i} \binom{n_j}{k_j} - \min_{y_1 + \dots + y_{\ell} = s-1} \binom{n_i-1-y_i}{k_i} \prod_{j \neq i} \binom{n_j-y_j}{k_j}.$$

Then,

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{F}(\overline{v_{i,1}})| + |\mathcal{F}(v_{i,1})| \\ &\leq \binom{n_i-1}{k_i} \prod_{j \neq i} \binom{n_j}{k_j} - \min_{y_1 + \dots + y_{\ell} = s-1} \binom{n_i-1-y_i}{k_i} \prod_{j \neq i} \binom{n_j-y_j}{k_j} + \binom{n_i-1}{k_i} \prod_{j \neq i} \binom{n_j}{k_j} \\ &= \prod_{j=1}^{\ell} \binom{n_j}{k_j} - \min_{y_1 + \dots + y_{\ell} = s-1} \binom{n_i-1-y_i}{k_i} \prod_{j \neq i} \binom{n_j-y_j}{k_j} \\ &\leq \prod_{j=1}^{\ell} \binom{n_j}{k_j} - \min_{x_1 + x_2 + \dots + x_{\ell} = s} \prod_{j=1}^{\ell} \binom{n_j-x_j}{k_j} \end{aligned}$$

and the lemma holds.

Case 2. $\nu(\mathcal{F}(\overline{v_{i,1}})) = s$ for all $i \in [\ell]$.

Let M be an s -matching in $\mathcal{F}(\overline{v_{i,1}})$ and let U be the set of vertices covered by M . Since $\nu(\mathcal{F}) \leq s$, each edge in $\mathcal{F}(v_{i,1})$ intersects U . It follows that

$$|\mathcal{F}(v_{i,1})| \leq \prod_{j=1}^{\ell} \binom{n'_j}{k'_j} - \prod_{j=1}^{\ell} \binom{n'_j - sk_j}{k'_j},$$

where $n'_j = n_j, k'_j = k_j$ for all $j \neq i$ and $n'_i = n_i - 1, k'_i = k_i - 1$. Then, we have

$$\begin{aligned} |\mathcal{F}(v_{i,1})| &\leq \sum_{p=1}^{\ell} \left[\binom{n'_p}{k'_p} - \binom{n'_p - sk_p}{k'_p} \right] \prod_{j \leq p-1} \binom{n'_j - sk_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &\leq \sum_{p=1}^{\ell} sk_p \binom{n'_p - 1}{k'_p - 1} \prod_{j \leq p-1} \binom{n'_j - sk_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &\leq \sum_{p=1}^{\ell} sk_p \binom{n'_p - 1}{k'_p - 1} \prod_{j \leq p-1} \binom{n'_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &\leq \sum_{p=1}^{\ell} \frac{sk_p k'_p}{n'_p} \binom{n'_p}{k'_p} \prod_{j \leq p-1} \binom{n'_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &= \left(\sum_{p=1}^{\ell} \frac{sk_p k'_p}{n'_p} \right) \prod_{j=1}^{\ell} \binom{n'_j}{k'_j}. \end{aligned}$$

Note that $\frac{k'_i}{n'_i} = \frac{k_i - 1}{n_i - 1} \leq \frac{k_i}{n_i}$ and $\binom{n'_i}{k'_i} = \frac{k_i}{n_i} \binom{n_i}{k_i}$. It follows that

$$|\mathcal{F}(v_{i,1})| \leq \frac{k_i}{n_i} \left(\sum_{p=1}^{\ell} \frac{sk_p^2}{n_p} \right) \prod_{j=1}^{\ell} \binom{n_j}{k_j}. \quad (2.5)$$

Let M' be an s -matching in \mathcal{F} and let U' be the set of vertices covered by M' . Since \mathcal{F} is shifted, we may assume that

$$U' = \bigcup_{i=1}^{\ell} \{v_{i,1}, \dots, v_{i,sk_i}\}.$$

For each $i \in [\ell]$ and $j \in [sk_i]$, define

$$\mathcal{F}[v_{i,j}] = \left\{ T \in \bigsqcup_{p=1}^{\ell} \binom{V'_p}{k'_p} : T \cup \{v_{i,j}\} \in \mathcal{F} \right\},$$

where $V'_p = V_p, k'_p = k_p$ for $p \neq i$ and $V'_i = V_i \setminus \{v_{i,1}, \dots, v_{i,j}\}, k'_i = k_i - 1$. By shiftedness, we have

$$\mathcal{F}[v_{i,1}] \supseteq \mathcal{F}[v_{i,2}] \supseteq \dots \supseteq \mathcal{F}[v_{i,sk_i}] \quad (2.6)$$

for $i = 1, 2, \dots, \ell$.

We claim that $\nu(\mathcal{F}[v_{i,s+1}]) \leq s$ for $i = 1, 2, \dots, \ell$. Otherwise, let E_1, E_2, \dots, E_{s+1} be a matching of size $s+1$ in $\mathcal{F}[v_{i,s+1}]$. By (2.6), $E_1 \cup \{v_{i,1}\}, E_2 \cup \{v_{i,2}\}, \dots, E_{s+1} \cup \{v_{i,s+1}\}$ is a matching of size $s+1$ in \mathcal{F} , a contradiction. Thus $\nu(\mathcal{F}[v_{i,s+1}]) \leq s$ for $i = 1, 2, \dots, \ell$. Note that $n_i \geq (s+1)k_i$ implies that $\frac{n_i-s-1}{k_i-1} \geq \frac{n_i}{k_i} \geq \frac{n_1}{k_1}$. By Lemma 2.2, we get for each i

$$|\mathcal{F}[v_{i,s+1}]| \leq \frac{(s+1)k_1}{n_1} \binom{n_i-s-1}{k_i-1} \prod_{j \neq i} \binom{n_j}{k_j} \leq \frac{(s+1)k_1 k_i}{n_1 n_i} \prod_{j=1}^{\ell} \binom{n_j}{k_j}. \quad (2.7)$$

Since M' is a largest matching in \mathcal{F} , U' is a vertex cover of \mathcal{F} . By (2.6), it follows that

$$|\mathcal{F}| \leq \sum_{i=1}^{\ell} \sum_{j=1}^{sk_i} |\mathcal{F}[v_{i,j}]| \leq \sum_{i=1}^{\ell} (s|\mathcal{F}[v_{i,1}]| + (sk_i - s)|\mathcal{F}[v_{i,s+1}]|). \quad (2.8)$$

Substitute (2.5) and (2.7) into (2.8), we arrive at

$$\begin{aligned} |\mathcal{F}| &\leq \sum_{i=1}^{\ell} \left(s \frac{k_i}{n_i} \left(\sum_{p=1}^{\ell} \frac{sk_p^2}{n_p} \right) \prod_{j=1}^{\ell} \binom{n_j}{k_j} + s(k_i - 1) \frac{(s+1)k_1 k_i}{n_1 n_i} \prod_{j=1}^{\ell} \binom{n_j}{k_j} \right) \\ &\leq s \prod_{j=1}^{\ell} \binom{n_j}{k_j} \sum_{i=1}^{\ell} \left(\frac{k_i}{n_i} \cdot \sum_{p=1}^{\ell} \frac{sk_p^2}{n_p} + \frac{2sk_1 k_i^2}{n_1 n_i} \right). \end{aligned}$$

Since $n_i \geq 4\ell^2 k_i^2 s$, we have

$$\sum_{p=1}^{\ell} \frac{sk_p^2}{n_p} \leq \frac{1}{4\ell}, \text{ and } \frac{2sk_i^2}{n_i} \leq \frac{1}{2\ell^2} \leq \frac{1}{4\ell}.$$

Moreover, $\frac{k_i}{n_i} \leq \frac{k_1}{n_1}$. Hence, by inequality (2.4) we obtain that

$$|\mathcal{F}| \leq \frac{s}{2} \binom{n_1-1}{k_1-1} \prod_{j=2}^{\ell} \binom{n_j}{k_j} \leq s \binom{n_1-s}{k_1-1} \prod_{j=2}^{\ell} \binom{n_j}{k_j} \leq \left(\binom{n_1}{k_1} - \binom{n_1-s}{k_1} \right) \prod_{j=2}^{\ell} \binom{n_j}{k_j}$$

and this completes the proof. ■

Proof of Theorem 1.2. By Lemma 2.3, we are left to show that

$$\min_{x_1 + \dots + x_{\ell} = s} \prod_{i=1}^{\ell} \binom{n_i - x_i}{k_i} = \min_{1 \leq i \leq \ell} \binom{n_i - s}{k_i} \prod_{j \neq i} \binom{n_j}{k_j}.$$

Set

$$f(x) = \binom{n_i - x}{k_i} \binom{n_j - c + x}{k_j}.$$

Computing its derivative, we have

$$f'(x) = \binom{n_i - x}{k_i} \binom{n_j - c + x}{k_j} \left[\sum_{q=0}^{k_j-1} \frac{1}{n_j - c + x - q} - \sum_{p=0}^{k_i-1} \frac{1}{n_i - x - p} \right].$$

Set

$$g(x) = \sum_{q=0}^{k_j-1} \frac{1}{n_j - c + x - q} - \sum_{p=0}^{k_i-1} \frac{1}{n_i - x - p}.$$

It is easy to see that $g(x)$ is a decreasing function in the range $[k_j - n_j + c, n_i - k_i]$. Moreover,

$$g(k_j - n_j + c) = \sum_{q=0}^{k_j-1} \frac{1}{k_j - q} - \sum_{p=0}^{k_i-1} \frac{1}{n_i - p + n_j - k_j - c} > 1 - \frac{k_i}{n_i - k_i} > 0$$

and

$$g(n_i - k_i) = \sum_{q=0}^{k_j-1} \frac{1}{n_j - c + n_i - k_i - q} - \sum_{p=0}^{k_i-1} \frac{1}{k_i - p} < \frac{k_j}{n_j - k_j} - 1 < 0.$$

Let x_0 be the unique zero of $g(x)$ in the range $[k_j - n_j + c, n_i - k_i]$. It follows that $f(x)$ is increasing in the range $[k_j - n_j, x_0]$ and decreasing in the range $[x_0, n_i - k_i]$. Then we have

$$\min_{0 \leq x \leq c} f(x) = \min \left\{ \binom{n_i - c}{k_i} \binom{n_j}{k_j}, \binom{n_i}{k_i} \binom{n_j - c}{k_j} \right\}.$$

Thus,

$$\min_{x_1 + \dots + x_\ell = s} \prod_{i=1}^{\ell} \binom{n_i - x_i}{k_i} = \min_{1 \leq i \leq \ell} \binom{n_i - s}{k_i} \prod_{j \neq i} \binom{n_j}{k_j}$$

and the theorem follows. ■

3 Extremal problem for rainbow matchings on direct products

Frankl and Kupavskii [11] proved a concentration inequality for the intersections of a k -uniform hypergraph and a random matching. In this section, by the concentration property of the bipartite graph defined on two random matchings, we establish a condition on the number of edges for a family of direct products to contain a rainbow matching.

Aharoni and Howard [1] provided a tight condition on the number of edges for a family of bipartite graphs to contain a rainbow matching.

Lemma 3.1 ([1]). *Let G_1, G_2, \dots, G_s be bipartite graphs with the same partite sets L and R such that $|L| = |R| = m \geq s$. If $e(G_i) > (s-1)m$ for every $i \in [s]$, then G_1, G_2, \dots, G_s contain a rainbow matching.*

We recall that the Kneser graph $KG(n, k)$ is the graph on the vertex set $\binom{[n]}{k}$ and with the edge set formed by pairs of disjoint sets. Lovasz [18] determined all the eigenvalues of the adjacent matrix of $KG(n, k)$.

Lemma 3.2 ([18]). *The eigenvalues of the adjacent matrix of $KG(n, k)$ are $(-1)^i \binom{n-k-1}{k-i}$ with multiplicity $\binom{n}{i} - \binom{n}{i-1}$, $i = 0, 1, \dots, k$. (Here $\binom{n}{-1} = 0$.)*

We also need the following result due to Alon and Chung [2].

Lemma 3.3 ([2]). *Let G be a d -regular graph on n vertices and A be its adjacent matrix. Let $d = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq -d$ be the eigenvalue of A and let $\lambda = \max\{|\lambda_2|, |\lambda_n|\}$. For $S \subseteq V(G)$ with $|S| = \alpha n$, we have*

$$\left| e(G[S]) - \frac{d}{n} \cdot \frac{(\alpha n)^2}{2} \right| \leq \frac{1}{2} \lambda \alpha (1 - \alpha) n.$$

By considering the bipartite graph defined on two random matchings, we prove the following lemma, which is key to the proof of Theorem 1.3.

Lemma 3.4. *Let $\ell, s, n_1, \dots, n_\ell, k_1, \dots, k_\ell$ be integers with $3s \leq n_1/k_1 \leq \dots \leq n_\ell/k_\ell$. Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s \subseteq \sqcup_{i=1}^\ell \binom{V_i}{k_i}$ with $|V_i| = n_i$ for $i = 1, \dots, \ell$. If $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$ are rainbow matching free, then there exists $1 \leq t \leq s$ such that*

$$|\mathcal{F}_t| < \frac{6sk_1}{n_1} \prod_{1 \leq j \leq \ell} \binom{n_j}{k_j}.$$

Proof. Suppose that

$$|\mathcal{F}_t| = \frac{6sk_1}{n_1} \prod_{1 \leq j \leq \ell} \binom{n_j}{k_j}$$

for all $1 \leq t \leq s$. Set $m = \lfloor n_1/k_1 \rfloor$. Let $\mathcal{A} = \{A_1, \dots, A_m\}$ be an m -matching chosen from $\binom{V_1}{k_1}$ uniformly at random and $\mathcal{B} = \{B_1, \dots, B_m\}$ be an m -matching chosen from $\sqcup_{i=2}^\ell \binom{V_i}{k_i}$ uniformly at random. For each $t = 1, \dots, s$, we construct a bipartite graph G_t with partite sets \mathcal{A}, \mathcal{B} where we have an edge (A_i, B_j) iff $A_i \cup B_j \in \mathcal{F}_t$. Note that a rainbow matching in G_1, G_2, \dots, G_s gives a rainbow matching in $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$. Thus, by Lemma 3.1 and the union bound, it suffices to show that

$$\Pr(e(G_t) \leq (s-1)m) < \frac{1}{s}$$

for each $1 \leq t \leq s$. Let $X = e(G_t)$ and X_{ij} be the indicator function of the event that $A_i \cup B_j \in \mathcal{F}_t$. Clearly, we have

$$X = \sum_{1 \leq i, j \leq m} X_{ij}.$$

Set $\alpha = |\mathcal{F}_t| / \prod_{j=1}^\ell \binom{n_j}{k_j}$. Take expectation on both sides, we obtain that

$$\mathbb{E}(X) = \sum_{1 \leq i, j \leq m} \Pr[A_i \cup B_j \in \mathcal{F}_t] = \sum_{1 \leq i, j \leq m} \frac{|\mathcal{F}_t|}{\prod_{j=1}^\ell \binom{n_j}{k_j}} = \alpha m^2.$$

Since $n_1/k_1 \geq 3s$, we have

$$\alpha m = \frac{6sk_1}{n_1} \cdot \left\lfloor \frac{n_1}{k_1} \right\rfloor \geq \frac{6sk_1}{n_1} \cdot \frac{n_1 - k_1}{k_1} \geq 6s \left(1 - \frac{k_1}{n_1}\right) \geq 6s - 2. \quad (3.1)$$

By the Chebyshev inequality, it follows that

$$\Pr(X \leq (s-1)m) \leq \Pr\left(X \leq \frac{\mathbb{E}(X)}{6}\right) \leq \Pr\left(|X - \mathbb{E}(X)| \geq \frac{5}{6}\mathbb{E}(X)\right) \leq \frac{36}{25} \cdot \frac{\text{Var}(X)}{\alpha^2 m^4}. \quad (3.2)$$

Now we need to give an upper bound for $\text{Var}(X)$. For $1 \leq i, i', j, j' \leq m$, define

$$\theta(i, j, i', j') = \Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) - \alpha^2.$$

Then,

$$\begin{aligned} \text{Var}(X) &= \mathbb{E}(X^2) - \mathbb{E}(X)^2 \\ &= \sum_{i, j, i', j' \in [m]} (\mathbb{E}[X_{ij} X_{i'j'}] - \mathbb{E}[X_{ij}] \mathbb{E}[X_{i'j'}]) \\ &= \sum_{i, j, i', j' \in [m]} (\Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) - \Pr(A_i \cup B_j \in \mathcal{F}_t) \Pr(A_{i'} \cup B_{j'} \in \mathcal{F}_t)) \\ &= \sum_{i, j, i', j' \in [m]} \theta(i, j, i', j'). \end{aligned} \tag{3.3}$$

We derive upper bounds on $\theta(i, j, i', j')$ for $i, j, i', j' \in [m]$ by distinguishing four cases.

Case 1. $i \neq i'$ and $j \neq j'$.

Let H be a graph on the vertex set $\sqcup_{i=1}^\ell \binom{V_i}{k_i}$ and with the edge set formed by pairs of disjoint sets in $\sqcup_{i=1}^\ell \binom{V_i}{k_i}$. Let λ be the second largest absolute eigenvalue of adjacent matrix of H . Frankl [8] observed that the adjacency matrix of H is the Kronecker product of the adjacency matrices of the Kneser graphs. By Lemma 3.2, it follows that

$$\lambda = \frac{k_1}{n_1 - k_1} \prod_{1 \leq p \leq \ell} \binom{n_p - k_p}{k_p}.$$

Since $i \neq i'$ and $j \neq j'$, we have $(A_i \cup B_j) \cap (A_{i'} \cup B_{j'}) = \emptyset$. Note that A_1, \dots, A_m and B_1, \dots, B_m are chosen uniformly at random. The probability that $A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t$ equals to the probability that a uniform chosen edge from $E(H)$ is an edge of the induce subgraph $H[\mathcal{F}_t]$, that is,

$$\Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) = \frac{e(H[\mathcal{F}_t])}{e(H)}.$$

Let D be the degree of H and N be the number of vertices of H . It is easy to see that $D = \prod_{1 \leq p \leq \ell} \binom{n_p - k_p}{k_p}$ and $N = \prod_{1 \leq p \leq \ell} \binom{n_p}{k_p}$. Since H is D -regular, we have $e(H) = \frac{DN}{2}$. By Lemma 3.3, we have

$$\theta(i, j, i', j') \leq \left| \frac{e(H[\mathcal{F}_t])}{e(H)} - \alpha^2 \right| \leq \frac{\lambda}{D} \alpha(1 - \alpha) \leq \frac{k_1}{n_1 - k_1} \alpha. \tag{3.4}$$

Case 2. $i \neq i'$ and $j = j'$.

For $B \in \sqcup_{i=2}^\ell \binom{V_i}{k_i}$, define $\mathcal{F}_t(B) = \left\{ K \in \binom{V_1}{k_1} : K \cup B \in \mathcal{F}_t \right\}$. Let H be the Kneser graph on the vertex set $\binom{V_1}{k_1}$. Since $i \neq i'$ and $j = j'$, the event that $A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t$ is equivalent to the event that a uniform chosen edge of H is an edge in $H[\mathcal{F}_t(B)]$. Thus,

$$\Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) = \sum_{B \in \sqcup_{i=2}^\ell \binom{V_i}{k_i}} \Pr(A_i, A_{i'} \in \mathcal{F}_t(B) | B_j = B) \Pr(B_j = B)$$

$$= \sum_{B \in \sqcup_{i=2}^{\ell} \binom{V_i}{k_i}} \frac{e(H[\mathcal{F}_t(B)])}{e(H)} \cdot \Pr(B_j = B). \quad (3.5)$$

Set $\alpha(B) = |\mathcal{F}_t(B)| / \binom{n_1}{k_1}$. Let D be the degree of H and N be the number of vertices in H . Clearly, $D = \binom{n_1 - k_1}{k_1}$ and $N = \binom{n_1}{k_1}$. Let λ be the second largest absolute eigenvalue of adjacent matrix of H . By Lemma 3.2, we have $\lambda = \binom{n_1 - k_1 - 1}{k_1 - 1}$. By lemma 3.3, it follows that

$$\left| \frac{e(H[\mathcal{F}_t(B)])}{e(H)} - \alpha^2(B) \right| \leq \frac{\lambda}{D} \alpha(B)(1 - \alpha(B)) \leq \alpha(B)(1 - \alpha(B)). \quad (3.6)$$

By (3.5) and (3.6), we obtain that

$$\begin{aligned} \theta(i, j, i', j') &= \Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) - \alpha^2 \\ &\leq \frac{1}{\prod_{2 \leq p \leq \ell} \binom{n_p}{k_p}} \sum_{B \in \sqcup_{i=2}^{\ell} \binom{V_i}{k_i}} (\alpha(B)(1 - \alpha(B)) + \alpha^2(B)) - \alpha^2 \\ &\leq \frac{1}{\prod_{2 \leq p \leq \ell} \binom{n_p}{k_p}} \sum_{B \in \sqcup_{i=2}^{\ell} \binom{V_i}{k_i}} \alpha(B) \\ &\leq \frac{1}{\prod_{2 \leq p \leq \ell} \binom{n_p}{k_p}} \sum_{B \in \sqcup_{i=2}^{\ell} \binom{V_i}{k_i}} \frac{|\mathcal{F}_t(B)|}{\binom{n_1}{k_1}} \\ &\leq \frac{1}{\prod_{1 \leq p \leq \ell} \binom{n_p}{k_p}} \sum_{B \in \sqcup_{i=2}^{\ell} \binom{V_i}{k_i}} |\mathcal{F}_t(B)| \\ &\leq \frac{|\mathcal{F}_t|}{\prod_{1 \leq p \leq \ell} \binom{n_p}{k_p}} \\ &= \alpha \end{aligned} \quad (3.7)$$

Case 3. $i = i'$ and $j \neq j'$.

Similarly, let H be a graph on the vertex set $\sqcup_{i=2}^{\ell} \binom{V_i}{k_i}$ and with the edge set formed by pairs of disjoint sets in $\sqcup_{i=2}^{\ell} \binom{V_i}{k_i}$. Let λ be the second largest absolute eigenvalue of adjacent matrix of H . Since the adjacency matrix of H is the Kronecker product of the adjacency matrices of the Kneser graphs. By Lemma 3.2, it follows that

$$\lambda = \frac{k_2}{n_2 - k_2} \prod_{2 \leq p \leq \ell} \binom{n_p - k_p}{k_p}.$$

For each $A \in \binom{V_1}{k_1}$, define $\mathcal{F}_t(A) = \left\{ K \in \sqcup_{i=2}^{\ell} \binom{V_i}{k_i} : A \cup K \in \mathcal{F}_t \right\}$. Set $\alpha(A) = |\mathcal{F}_t(A)| / \prod_{2 \leq p \leq \ell} \binom{n_p}{k_p}$. Let D be the degree of H and let N be the number of vertices in H . Clearly, $D = \prod_{2 \leq p \leq \ell} \binom{n_p - k_p}{k_p}$ and $N = \prod_{2 \leq p \leq \ell} \binom{n_p}{k_p}$. Then,

$$\Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) = \sum_{A \in \binom{V_1}{k_1}} \frac{e(H[\mathcal{F}_t(A)])}{e(H)} \cdot \Pr(A_i = A).$$

By Lemma 3.3, we have for $A \in \binom{V_1}{k_1}$

$$\left| \frac{e(H[\mathcal{F}_t(A)])}{e(H)} - \alpha^2(A) \right| \leq \frac{\lambda}{D} \alpha(A)(1 - \alpha(A)) \leq \alpha(A)(1 - \alpha(A)).$$

Thus,

$$\begin{aligned} \theta(i, j, i', j') &\leq \Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) - \alpha^2 \\ &\leq \frac{1}{\binom{n_1}{k_1}} \sum_{A \in \binom{V_1}{k_1}} (\alpha(A)(1 - \alpha(A)) + \alpha^2(A)) \\ &\leq \frac{1}{\binom{n_1}{k_1}} \cdot \sum_{A \in \binom{V_1}{k_1}} \alpha(A) \\ &\leq \frac{1}{\binom{n_1}{k_1}} \sum_{A \in \binom{V_1}{k_1}} \frac{|\mathcal{F}_t(A)|}{\prod_{2 \leq p \leq \ell} \binom{n_p}{k_p}} \\ &= \alpha. \end{aligned} \tag{3.8}$$

Case 4. $i = i'$ and $j = j'$.

In this case, $\Pr(A_i \cup B_j, A_{i'} \cup B_{j'} \in \mathcal{F}_t) = \Pr(A_i \cup B_j \in \mathcal{F}_t) = \alpha$. Hence,

$$\theta(i, j, i', j') = \alpha - \alpha^2 \leq \alpha. \tag{3.9}$$

Substitute (3.4), (3.7), (3.8) and (3.9) into (3.3), we have

$$\text{Var}(X) = \sum_{i, j, i', j' \in [m]} \theta(i, j, i', j') \leq \frac{k_1}{n_1 - k_1} \alpha m^2 (m - 1)^2 + 2\alpha m^3 + \alpha m^2.$$

Note that $m = \lfloor \frac{n_1}{k_1} \rfloor \leq \frac{n_1}{k_1}$ implies $\frac{k_1}{n_1 - k_1} \leq \frac{1}{m - 1}$. It follows that

$$\text{Var}(X) \leq \frac{1}{m - 1} \alpha m^2 (m - 1)^2 + 2\alpha m^3 + \alpha m^2 = 3\alpha m^3. \tag{3.10}$$

Substitute (3.10) into (3.2), we arrive at

$$\Pr(X \leq (s - 1)m) \leq \frac{36}{25} \cdot \frac{3\alpha m^3}{\alpha^2 m^4} = \frac{36 \cdot 3}{25\alpha m}.$$

Moreover, (3.1) implies that $\alpha m \geq 6s - 2 \geq 5s$. Thus, we obtain that

$$\Pr(X \leq (s - 1)m) \leq \frac{36 \times 3}{25\alpha m} \leq \frac{36 \times 3}{25 \times 5} \cdot \frac{1}{s} < \frac{1}{s},$$

and the lemma follows. ■

The following lemma shows that the shifting operator preserves the rainbow matching free property of a family of uniform hypergraphs, which is due to Huang, Loh and Sudakov [14].

Lemma 3.5 (Huang-Loh-Sudakov [14]). *If the families $\mathcal{F}_1, \dots, \mathcal{F}_s \subseteq \binom{[n]}{k}$ are rainbow matching free and $i, j \in [n]$ with $i < j$, then $S_{ij}(\mathcal{F}_1), \dots, S_{ij}(\mathcal{F}_s)$ are still rainbow matching free.*

Lemma 3.6. *Let $\ell, s, n_1, \dots, n_\ell, k_1, \dots, k_\ell$ be integers with $n_i \geq 8\ell^2 k_i^2 s$ for $i = 1, \dots, \ell$. Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s \subseteq \sqcup_{i=1}^\ell \binom{V_i}{k_i}$ with $|V_i| = n_i$. If $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$ are rainbow matching free, then there exists $1 \leq t \leq s$ such that*

$$|\mathcal{F}_t| \leq \prod_{i=1}^\ell \binom{n_i}{k_i} - \min_{x_1 + \dots + x_\ell = s-1} \prod_{i=1}^\ell \binom{n_i - x_i}{k_i}.$$

Proof. Without loss of generality, we may assume that $n_1/k_1 \leq n_2/k_2 \leq \dots \leq n_\ell/k_\ell$. By Lemma 3.5, we may assume that \mathcal{F}_i is shifted for all $i = 1, \dots, s$. We prove the theorem by induction on s . The case $s = 1$ is trivial. Assume that the lemma holds for the case $s - 1$ and we have to show the lemma holds for the case s .

Let $V_i = \{v_{i,j} : j \in [n_i]\}$. For every $x \in \cup_{i=1}^\ell V_i$ and $1 \leq t \leq s$, define

$$\mathcal{F}_t(x) = \{F \setminus \{x\} : x \in F \in \mathcal{F}_t\} \text{ and } \mathcal{F}_t(\bar{x}) = \{F : x \notin F \in \mathcal{F}_t\}.$$

We distinguish into two cases.

Case 1. There exist $r \in [s]$ and $i \in [\ell]$ such that $\mathcal{F}_1(\overline{v_{i,1}}), \dots, \mathcal{F}_{r-1}(\overline{v_{i,1}}), \mathcal{F}_{r+1}(\overline{v_{i,1}}), \dots, \mathcal{F}_s(\overline{v_{i,1}})$ are rainbow matching free.

By the induction hypothesis, there exists $t \in [s] \setminus \{r\}$ such that

$$|\mathcal{F}_t(\overline{v_{i,1}})| \leq \binom{n_i - 1}{k_i} \prod_{j \neq i} \binom{n_j}{k_j} - \min_{y_1 + \dots + y_\ell = s-1} \binom{n_i - 1 - y_i}{k_i} \prod_{j \neq i} \binom{n_j - y_j}{k_j}.$$

Then,

$$\begin{aligned} |\mathcal{F}_t| &= |\mathcal{F}_t(\overline{v_{i,1}})| + |\mathcal{F}_t(v_{i,1})| \\ &\leq \binom{n_i - 1}{k_i} \prod_{j \neq i} \binom{n_j}{k_j} - \min_{y_1 + \dots + y_\ell = s-1} \binom{n_i - 1 - y_i}{k_i} \prod_{j \neq i} \binom{n_j - y_j}{k_j} + \binom{n_i - 1}{k_i} \prod_{j \neq i} \binom{n_j}{k_j} \\ &= \prod_{j=1}^\ell \binom{n_j}{k_j} - \min_{y_1 + \dots + y_\ell = s-1} \binom{n_i - 1 - y_i}{k_i} \prod_{j \neq i} \binom{n_j - y_j}{k_j} \\ &\leq \prod_{j=1}^\ell \binom{n_j}{k_j} - \min_{x_1 + x_2 + \dots + x_\ell = s} \prod_{j=1}^\ell \binom{n_j - x_j}{k_j} \end{aligned}$$

and the lemma holds.

Case 2. $\mathcal{F}_1(\overline{v_{i,1}}), \dots, \mathcal{F}_{r-1}(\overline{v_{i,1}}), \mathcal{F}_{r+1}(\overline{v_{i,1}}), \dots, \mathcal{F}_s(\overline{v_{i,1}})$ contain a rainbow matching for all $r \in [s]$ and $i \in [\ell]$.

Let M be a rainbow matching in $\{\mathcal{F}_1(\overline{v_{i,1}}), \dots, \mathcal{F}_{r-1}(\overline{v_{i,1}}), \mathcal{F}_{r+1}(\overline{v_{i,1}}), \dots, \mathcal{F}_s(\overline{v_{i,1}})\}$ and let U be the set of vertices covered by M . Since $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_s$ are rainbow matching free, each

edge of $\mathcal{F}_r(v_{i,1})$ intersects U . It follows that

$$|\mathcal{F}_r(v_{i,1})| \leq \prod_{j=1}^{\ell} \binom{n'_j}{k'_j} - \prod_{j=1}^{\ell} \binom{n'_j - (s-1)k_j}{k'_j},$$

where $n'_j = n_j, k'_j = k_j$ for all $j \neq i$ and $n'_i = n_i - 1, k'_i = k_i - 1$. Then,

$$\begin{aligned} |\mathcal{F}_r(v_{i,1})| &\leq \sum_{p=1}^{\ell} \left[\binom{n'_p}{k'_p} - \binom{n'_p - (s-1)k_p}{k'_p} \right] \prod_{j \leq p-1} \binom{n'_j - (s-1)k_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &\leq \sum_{p=1}^{\ell} (s-1)k_p \binom{n'_p - 1}{k'_p - 1} \prod_{j \leq p-1} \binom{n'_j - (s-1)k_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &\leq \sum_{p=1}^{\ell} (s-1)k_p \binom{n'_p - 1}{k'_p - 1} \prod_{j \leq p-1} \binom{n'_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &\leq \sum_{p=1}^{\ell} \frac{(s-1)k_p k'_p}{n'_p} \binom{n'_p}{k'_p} \prod_{j \leq p-1} \binom{n'_j}{k'_j} \prod_{j \geq p+1} \binom{n'_j}{k'_j} \\ &= \left(\sum_{p=1}^{\ell} \frac{(s-1)k_p k'_p}{n'_p} \right) \prod_{j=1}^{\ell} \binom{n'_j}{k'_j}. \end{aligned}$$

Note that $\frac{k'_i}{n'_i} = \frac{k_i - 1}{n_i - 1} \leq \frac{k_i}{n_i}$ and $\binom{n'_i}{k'_i} = \frac{k_i}{n_i} \binom{n_i}{k_i}$. It follows that for every $i \in [\ell]$ and $r \in [s]$

$$|\mathcal{F}_r(v_{i,1})| \leq \frac{(s-1)k_i}{n_i} \left(\sum_{p=1}^{\ell} \frac{k_p^2}{n_p} \right) \prod_{j=1}^{\ell} \binom{n_j}{k_j}. \quad (3.11)$$

For $j \in [\ell]$, let $S_j = \{v_{j,1}, \dots, v_{j,s}\}$ and let $V'_j = V_j \setminus S_j$. For every $r \in [s]$ and $i \in [\ell]$, define

$$\mathcal{F}_r[v_{i,s}] = \left\{ E \in \bigsqcup_{j=1}^{\ell} \binom{V'_j}{k'_j} : E \cup \{v_{i,s}\} \in \mathcal{F}_r \right\},$$

where $k'_j = k_j$ for all $j \neq i$ and $k'_i = k_i - 1$. Define

$$\mathcal{F}_r^*[v_{i,s}] = \{E \cup \{u\} : E \in \mathcal{F}_r[v_{i,s}], u \in V'_i \setminus E\}.$$

Claim 1. There exists $t \in [s]$ such that for all $i \in [\ell]$

$$|\mathcal{F}_t^*[v_{i,s}]| \leq \frac{6sk_1}{n_1} \prod_{1 \leq j \leq \ell} \binom{n_j - s}{k_j}. \quad (3.12)$$

Proof. Suppose to the contrary that for every $t \in [s]$ there is an $i_t \in [\ell]$ such that

$$|\mathcal{F}_t^*[v_{i_t,s}]| > \frac{6sk_1}{n_1} \prod_{1 \leq j \leq \ell} \binom{n_j - s}{k_j}.$$

Note that

$$\mathcal{F}_t^*[v_{i_t,s}] \subseteq \bigsqcup_{j=1}^{\ell} \binom{V'_j}{k_j}.$$

By Lemma 3.4, we conclude that $\mathcal{F}_1^*[v_{i_1,s}], \dots, \mathcal{F}_s^*[v_{i_s,s}]$ contain a rainbow matching. Let $F_1 \in \mathcal{F}_1^*[v_{i_1,s}], \dots, F_s \in \mathcal{F}_s^*[v_{i_s,s}]$ be such a matching. By the definition of $\mathcal{F}_t^*[v_{i_t,s}]$, we see that there exist $E_1 \subset F_1, \dots, E_s \subset F_s$ such that $E_1 \cup \{v_{i_1,s}\} \in \mathcal{F}_1, E_2 \cup \{v_{i_2,s}\} \in \mathcal{F}_2, \dots, E_s \cup \{v_{i_s,s}\} \in \mathcal{F}_s$. Note that $\mathcal{F}_1, \dots, \mathcal{F}_s$ are all shifted. It follows that $E_1 \cup \{v_{i_1,1}\} \in \mathcal{F}_1, E_2 \cup \{v_{i_2,2}\} \in \mathcal{F}_2, \dots, E_s \cup \{v_{i_s,s}\} \in \mathcal{F}_s$ is a rainbow matching, which contradicts the fact that $\mathcal{F}_1, \dots, \mathcal{F}_s$ are rainbow matching free. \blacksquare

By Claim 1, there exists $t \in [s]$ such that (3.12) holds for all $i \in [\ell]$. By the definition of $\mathcal{F}_t^*[v_{i,s}]$, we see that for each $E \in \mathcal{F}_t[v_{i,s}]$ there are $|V'_i \setminus E| = (n_i - s) - (k_i - 1)$ choices of u such that $E \cup \{u\} \in \mathcal{F}_t^*[v_{i,s}]$. Moreover, for each $F \in \mathcal{F}_t^*[v_{i,s}]$ we have $|F \cap V'_i| = k_i$. It follows that at most k_i sets in $\mathcal{F}_t[v_{i,s}]$ are contained in F . Thus, we have

$$|\mathcal{F}_t^*[v_{i,s}]| \geq \frac{n_i - s - k_i + 1}{k_i} \cdot |\mathcal{F}_t[v_{i,s}]|. \quad (3.13)$$

By (3.12) and (3.13), we obtain that for all $i = 1, 2, \dots, \ell$,

$$|\mathcal{F}_t[v_{i,s}]| \leq \frac{6sk_1k_i}{n_1(n_i - s - k_i + 1)} \prod_{1 \leq j \leq \ell} \binom{n_j}{k_j}. \quad (3.14)$$

Note that $\mathcal{F}_1, \dots, \mathcal{F}_{t-1}, \mathcal{F}_{t+1}, \dots, \mathcal{F}_s$ contain a rainbow matching. Let M' be such a matching and let U' be the set of vertices covered by M' . By shiftedness, we may assume that

$$U' = \bigcup_{i=1}^{\ell} \{v_{i,1}, \dots, v_{i,(s-1)k_i}\}.$$

Since $\mathcal{F}_1, \dots, \mathcal{F}_s$ are rainbow matching free, it follows that every edge of \mathcal{F}_t intersects U' . Thus,

$$\mathcal{F}_t = \sum_{i=1}^{\ell} \sum_{j=1}^{(s-1)k_i} \mathcal{F}_t[v_{i,j}].$$

By shiftedness, we have

$$\mathcal{F}_t[v_{i,1}] \supseteq \mathcal{F}_t[v_{i,2}] \supseteq \dots \supseteq \mathcal{F}_t[v_{i,(s-1)k_i}]$$

for $i = 1, 2, \dots, \ell$. Then

$$|\mathcal{F}_t| \leq \sum_{i=1}^{\ell} (s-1) |\mathcal{F}_t[v_{i,1}]| + \sum_{i=1}^{\ell} (k_i - 1)(s-1) |\mathcal{F}_t[v_{i,s}]|. \quad (3.15)$$

Substitute (3.11) and (3.14) into (3.15), we arrive at

$$|\mathcal{F}_t| \leq \sum_{i=1}^{\ell} \left((s-1)^2 \left(\sum_{p=1}^{\ell} \frac{k_p^2}{n_p} \right) \frac{k_i}{n_i} \prod_{j=1}^{\ell} \binom{n_j}{k_j} + 6s(s-1) \frac{k_i(k_i - 1)}{n_i - s - k_i + 1} \frac{k_1}{n_1} \prod_{j=1}^{\ell} \binom{n_j}{k_j} \right).$$

Since $\frac{k_i}{n_i} \leq \frac{k_1}{n_1}$, we have

$$|\mathcal{F}_t| \leq \frac{k_1}{n_1} \prod_{j=1}^{\ell} \binom{n_j}{k_j} \sum_{i=1}^{\ell} \left((s-1)^2 \sum_{p=1}^{\ell} \frac{k_p^2}{n_p} + 6s(s-1) \frac{k_i(k_i-1)}{n_i - s - k_i + 1} \right).$$

Since $n_i \geq 8\ell^2 k_i^2 s$ and $\ell \geq 2$, we have

$$\sum_{i=1}^{\ell} \left((s-1)^2 \sum_{p=1}^{\ell} \frac{k_p^2}{n_p} \right) = \frac{(s-1)^2 \ell^2}{8\ell^2 s} \leq \frac{s-1}{8},$$

and

$$\sum_{i=1}^{\ell} \frac{6s(s-1)k_i(k_i-1)}{n_i - s - k_i + 1} \leq \sum_{i=1}^{\ell} \frac{6s(s-1)k_i(k_i-1)}{16\ell k_i(k_i-1)s + 16\ell k_i s - s - k_i + 1} \leq \frac{3(s-1)}{8}.$$

Moreover, $\frac{k_i}{n_i} \leq \frac{k_1}{n_1}$. Hence, by inequality (2.4) we obtain that

$$\begin{aligned} |\mathcal{F}_t| &\leq \frac{(s-1)}{2} \binom{n_1-1}{k_1-1} \prod_{j=2}^{\ell} \binom{n_j}{k_j} \\ &\leq (s-1) \binom{n_1-s+1}{k_1-1} \prod_{j=2}^{\ell} \binom{n_j}{k_j} \\ &\leq \left(\binom{n_1}{k_1} - \binom{n_1-s+1}{k_1} \right) \prod_{j=2}^{\ell} \binom{n_j}{k_j}, \end{aligned}$$

and this completes the proof. ■

Proof of Theorem 1.3. As the same to the proof of Theorem 1.2, we have

$$\min_{x_1 + \dots + x_{\ell} = s} \prod_{i=1}^{\ell} \binom{n_i - x_i}{k_i} = \min_{1 \leq i \leq \ell} \binom{n_i - s}{k_i} \prod_{j \neq i} \binom{n_j}{k_j}.$$

Thus, Lemma 3.6 implies the theorem. ■

References

- [1] R. Aharoni and D. Howard. A rainbow r -partite version of the Erdős–Ko–Rado theorem. *Combinatorics, Probability and Computing*, 26(3):321–337, 2017.
- [2] N. Alon and F. R. Chung. Explicit construction of linear sized tolerant networks. *Discrete Mathematics*, 72(1-3):15–19, 1988.
- [3] B. Bollobás, D. E. Daykin, and P. Erdős. Sets of independent edges of a hypergraph. *The Quarterly Journal of Mathematics*, 27(1):25–32, 1976.

- [4] P. Erdős, C. Ko, and R. Rado. Intersection theorems for systems of finite sets. *The Quarterly Journal of Mathematics*, 12:313–320, 1961.
- [5] P. Erdős. A problem on independent r -tuples. *Annales Universitatis Budapestinensis*, 8:93–95, 1965.
- [6] P. Erdős and T. Gallai. On maximal paths and circuits of graphs. *Acta Mathematica Academiae Scientiarum Hungarica*, 10(3-4):337–356, 1959.
- [7] P. Frankl. The shifting technique in extremal set theory. *Surveys in combinatorics*, 123:81–110, 1987.
- [8] P. Frankl. An Erdős–Ko–Rado theorem for direct products. *European Journal of Combinatorics*, 17(8):727–730, 1996.
- [9] P. Frankl. Improved bounds for Erdős matching conjecture. *Journal of Combinatorial Theory, Series A*, 120(5):1068–1072, 2013.
- [10] P. Frankl. On the maximum number of edges in a hypergraph with given matching number. *Discrete Applied Mathematics*, 216:562–581, 2017.
- [11] P. Frankl and A. Kupavskii. The Erdős matching conjecture and concentration inequalities. *arXiv preprint arXiv:1806.08855*, 2018.
- [12] P. Frankl and A. Kupavskii. Simple juntas for shifted families. *Discrete Analysis*, (14):18 pp, 2020.
- [13] P. Frankl and J. Wang. On the sum of sizes of overlapping families. *arXiv preprint arXiv:2105.00481*, 2021.
- [14] H. Huang, P.-S. Loh, and B. Sudakov. The size of a hypergraph and its matching number. *Combinatorics, Probability and Computing*, 21(3):442–450, 2012.
- [15] G. Katona. Intersection theorems for systems of finite sets. *Acta Mathematica Academiae Scientiarum Hungaricae*, 15(3-4):329–337, 1964.
- [16] N. Keller and N. Lifshitz. The junta method for hypergraphs and the Erdős–Chvátal simplex conjecture. *Advances in Mathematics*, 392:107991, 2021.
- [17] A. Kupavskii. Rainbow version of the Erdős matching conjecture via concentration. *arXiv preprint arXiv:2104.08083*, 2021.
- [18] L. Lovász. On the shannon capacity of a graph. *IEEE Transactions on Information theory*, 25(1):1–7, 1979.
- [19] H. Lu, Y. Wang, and X. Yu. A better bound on the size of rainbow matchings. *arXiv preprint arXiv:2004.12561*, 2020.