

Treewidth versus clique number. II. Tree-independence number

Clément Dallard^{1,2}, Martin Milanič^{1,2}, Kenny Štorgel^{1,3}

¹ FAMNIT, University of Primorska, Koper, Slovenia

² IAM, University of Primorska, Koper, Slovenia

³ Faculty of Information Studies in Novo mesto, Slovenia

clement.dallard@famnit.upr.si martin.milanic@upr.si kennystorgel.research@gmail.com

Abstract

In 2020, we initiated a systematic study of graph classes in which the treewidth can only be large due to the presence of a large clique, which we call (tw, ω) -bounded. The family of (tw, ω) -bounded graph classes provides a unifying framework for a variety of very different families of graph classes, including graph classes of bounded treewidth, graph classes of bounded independence number, intersection graphs of connected subgraphs of graphs with bounded treewidth, and graphs in which all minimal separators are of bounded size. While Chaplick and Zeman showed in 2017 that (tw, ω) -bounded graph classes enjoy some good algorithmic properties related to clique and coloring problems, it is an interesting open problem to which extent (tw, ω) -boundedness has useful algorithmic implications for problems related to independent sets. We provide a partial answer to this question by identifying a sufficient condition for (tw, ω) -bounded graph classes to admit a polynomial-time algorithm for the Maximum Weight Independent Packing problem and, as a consequence, for the weighted variants of the Independent Set and Induced Matching problems.

Our approach is based on a new min-max graph parameter related to tree decompositions. We define the *independence number* of a tree decomposition \mathcal{T} of a graph as the maximum independence number over all subgraphs of G induced by some bag of \mathcal{T} . The *tree-independence number* of a graph G is then defined as the minimum independence number over all tree decompositions of G . Boundedness of the tree-independence number is a refinement of (tw, ω) -boundedness that is still general enough to hold for all the aforementioned families of graph classes. Generalizing a result on chordal graphs due to Cameron and Hell from 2006, we show that if a graph is given together with a tree decomposition with bounded independence number, then the Maximum Weight Independent Packing problem can be solved in polynomial time. Applications of our general algorithmic result to specific graph classes are given in the third paper of the series [Dallard, Milanič, and Štorgel, Treewidth versus clique number. III. Tree-independence number of graphs with a forbidden structure, 2022].

Contents

1	Introduction	3
1.1	Width parameters of graphs	3
1.2	Chordal graphs	3
1.3	Treewidth versus clique number	4
1.4	Our results	6
1.5	Overview of our methodology	8
1.6	Relation to previous work	9
1.7	Organization of the paper	10
2	Preliminaries	10
3	Definition and basic properties	13
4	A refinement of the tree-independence number	17
5	Weighted independent sets	19
6	Cameron–Hell construction	23
7	Weighted independent packings	25
8	Open questions	27

1 Introduction

1.1 Width parameters of graphs

One of the approaches for dealing with an algorithmically hard problem is to identify restrictions on the input instances under which the problem becomes solvable in polynomial time. In the case of graph problems, various measures of structural complexity of graphs, commonly known as *graph width parameters*, have become one of the central tools for this task. When a width parameter of the input graph is bounded by a fixed constant, this often leads to the development of efficient dynamic programming algorithms, typically applied along some kind of tree-like decomposition of the graph into simpler parts, which is well-behaved according to the specifics of the parameter (see, e.g., [4, 5, 17, 18, 21, 23, 24, 43, 44, 47, 72]). For a concrete example, consider the *treewidth* of a graph G . This parameter measures, roughly speaking, how similar the graph is to a tree. A formal definition is given using the concept of a *tree decomposition* of G , which is a collection of bags (subsets of the vertex set of G) arranged into a tree structure and satisfying certain conditions [86]. The treewidth of a graph is defined as the smallest possible size, over all tree decompositions of the graph, of a largest bag in the decomposition minus one. The key algorithmic result in the context of treewidth is the celebrated metatheorem due to Courcelle [43], which states that any decision problem expressible in monadic second order logic, with quantifiers over vertex and edge subsets, can be solved in linear time if the input graph is equipped with a tree decomposition of bounded width. A similar theorem for optimization problems was developed by Arnborg, Lagergren, and Seese [4].

Graphs of bounded treewidth are necessarily sparse. In 2000, Courcelle and Olariu introduced a more general parameter called *clique-width* [45], which is bounded whenever the treewidth is bounded (see also [42]) but whose boundedness is not restricted to sparse graph families. An algorithmic metatheorem for graph classes with bounded clique-width was developed by Courcelle, Makowsky, and Rotics [44]. Similarly to the algorithmic metatheorems for treewidth, this theorem also applies to decision or optimization problems expressible in monadic second order logic; however, it only allows for quantifiers over vertex subsets. More recently, several other width parameters capable of handling dense graphs were also introduced and studied, including rank-width [81], Boolean-width [23, 24], mim-width [21, 72, 74, 97], and twin-width [16–18]. Bounded clique-width is equivalent to bounded rank-width or bounded Boolean-width, but implies bounded mim-width as well as bounded twin-width. Each of these width parameters has some useful algorithmic features. To mention one more example, the so-called *locally checkable vertex subset* and *vertex partitioning* problems are solvable in polynomial time on any class of bounded mim-width, provided the input graph is given with a branch decomposition of constant width [6, 24]. This includes the INDEPENDENT SET problem, the INDUCED MATCHING problem, and various variants of the DOMINATING SET problem, as well as the k -COLORING and the H -HOMOMORPHISM problems.

1.2 Chordal graphs

Despite a growing variety of graph width parameters and associated algorithmic results, there exist some well-structured graph classes containing dense graphs in which all of the above width parameters remain unbounded. In such a case, different algorithmic approaches are needed. This is the case, for example, for the well-known class of *chordal graphs* (also known as *triangulated graphs*), defined as graphs in which every cycle of length at least four has a chord. Chordal graphs correspond to sparse symmetric positive-definite systems of linear equations that admit a particularly efficient Gaussian elimination (see [88]) and have applications in many other areas, including probabilistic graphical models and continuous optimization (see, e.g., the survey [96]).

They have been extensively studied in the literature from both structural and algorithmic points of view (see, e.g., [9, 19, 67, 93]). However, the class of chordal graphs has unbounded mim-width, as follows from the analogous result for the class of strongly chordal split graphs [79]. Also, the twin-width of chordal graphs can be arbitrarily large, as follows from the fact that this parameter is unbounded in the class of interval graphs, that is, intersection graphs of closed intervals on the real line [16]. Split graphs, strongly chordal graphs, and interval graphs are all subclasses of the class of chordal graphs (see, e.g., [19]).

Thus, one cannot obtain efficient algorithms for the class of chordal graphs by applying any of the algorithmic metatheorems for the aforementioned width parameters. In fact, this is not surprising, as several problems remain NP-complete in the class of chordal graphs, including the DOMINATING SET [41] and the WEIGHTED INDEPENDENT DOMINATING SET problems [29]. Nevertheless, many NP-complete problems are known to be efficiently solvable for chordal graphs, including the INDEPENDENT DOMINATING SET problem [57], as well as the INDEPENDENT SET, the CLIQUE, and the CHROMATIC NUMBER problems, as a consequence of the fact that these problems are solvable in polynomial time in the more general class of perfect graphs [68]. Furthermore, since chordal graphs have only polynomially many minimal separators (see, e.g., [94]), a metatheorem for such graph classes due to Fomin, Todinca, and Villanger [62] applies, establishing polynomial-time solvability of certain problems expressible in counting monadic second order logic. Besides the already mentioned INDEPENDENT SET problem, this approach captures the FEEDBACK VERTEX SET problem, the INDUCED MATCHING problem, and various packing problems.

1.3 Treewidth versus clique number

Chordal graphs have an interesting property regarding treewidth. While graphs with large cliques necessarily have large treewidth, in chordal graphs the converse holds, too: the treewidth of a chordal graph can only be large due to the presence of a large clique (see, e.g., [12]). Recently, we initiated in [48, 50] a systematic study of graph classes in which this sufficient condition for large treewidth—the presence of a large clique—is also necessary, which we call (tw, ω) -bounded. A graph class \mathcal{G} is said to be (tw, ω) -bounded if it admits a (tw, ω) -binding function, that is, a function f such that the treewidth of any graph $G \in \mathcal{G}$ is at most $f(\omega(G))$, where $\omega(G)$ is the clique number of G , and the same holds for all induced subgraphs of G . The family of (tw, ω) -bounded graph classes provides a unifying framework for a variety of very different families of graph classes studied in the literature. Besides graph classes of bounded treewidth, (tw, ω) -boundedness holds for graph classes of bounded independence number (as a consequence of Ramsey’s theorem), intersection graphs of connected subgraphs of graphs with treewidth at most t , for any fixed positive integer t , and classes of graphs in which all minimal separators are of bounded size. Intersection graphs of connected subgraphs of graphs with treewidth at most t were studied in 1990 by Scheffler [89] and in 1998 by Bodlaender, Gustedt, and Telle [10]. They include chordal graphs (for which $t = 1$, since every chordal graph is the intersection graph of subtrees in a tree [25, 66, 98]) and circular-arc graphs, that is, intersection graphs of circular arcs on a circle (for which $t = 2$), as well as H -graphs, that is, the intersection graphs of connected subgraphs of a subdivision of a fixed multigraph H , introduced in 1992 by Bíró, Hujter, and Tuza [8] and studied more recently in a number of papers [30–33, 61]. Classes of graphs in which all minimal separators are of bounded size were studied in 1999 by Skodinis [92]. In [48, 50], we characterized, for each of six well-known graph containment relations (the minor, topological subgraph, subgraph, and their induced variants), the graphs H such that the class of graphs excluding H with respect to the relation is (tw, ω) -bounded.

In view of this richness of the family of (tw, ω) -bounded graph classes, one should not expect algorithmic metatheorems for (tw, ω) -bounded graph classes of similar generality as those available, for example, for graphs of bounded treewidth or bounded clique-width. In fact, as already exemplified via the class of chordal graphs, several NP-complete graph problems remain NP-complete when restricted to some (tw, ω) -bounded graph classes. This is the case not only for the DOMINATING SET and WEIGHTED INDEPENDENT DOMINATING SET problems, but also for the CHROMATIC NUMBER problem, which is NP-complete in the class of circular-arc graphs [63], and for the unweighted INDEPENDENT DOMINATING SET problem, which is NP-complete in the (tw, ω) -bounded class of graphs that have a clique such that each vertex is either in the clique or is adjacent to at most one vertex not in the clique [103]. Nevertheless, (tw, ω) -bounded graph classes do enjoy some good algorithmic properties related to clique and coloring problems. They can be summarized as follows.

1. Chaplick and Zeman showed in [33] (see also [32]) that for every positive integer k , any (tw, ω) -bounded graph class \mathcal{G} having a computable binding function admits linear-time algorithms for the k -CLIQUE and the LIST k -COLORING problems. Furthermore, as we showed in [48, 50], any such class admits a polynomial-time algorithm for the LIST k -COLORING problem that is *robust* in the sense of Raghavan and Spinrad [84]: it either solves the problem or determines that the input graph is not in \mathcal{G} .
2. Similarly, for every positive integer k , there exists a robust quadratic-time algorithm for the LIST k -EDGE-COLORING problem in any (tw, ω) -bounded graph class having a computable binding function f . Indeed, if the input graph G has a vertex with degree more than k , then it cannot be k -edge-colorable with respect to the given lists of colors on the edges, while if the maximum degree is bounded by k , then the clique number is bounded by $k + 1$ and therefore the treewidth is bounded by $c_k = \max\{f(j) : 1 \leq j \leq k + 1\}$. In this case, we can use the linear-time algorithm of Bodlaender [11] to test whether $\text{tw}(G) \leq c_k$. If $\text{tw}(G) > c_k$, then the graph is not in the class. If $\text{tw}(G) \leq c_k$, then the clique-width of G is at most ℓ where $\ell = 3 \cdot 2^{c_k - 1}$ (see Corneil and Rotics [42]), and using a $(2^{2\ell + 1} - 1)$ -expression of G that can be computed in time $\mathcal{O}(|V(G)|^2)$ with an algorithm due to Fomin and Korhonen [59], the LIST k -EDGE-COLORING problem can be solved in time $\mathcal{O}(|V(G)|^2)$ by invoking a result of Kobler and Rotics [76].
3. As also shown in [48, 50], known approximation algorithms for treewidth (see, e.g., [58]) lead to improved approximations for the clique number of a graph from a (tw, ω) -bounded graph class having a computable exponential binding function. The approximation can be improved further if the binding function is computable and either linear or polynomial. For example, in the case of a polynomial (tw, ω) -binding function $f(k) = \mathcal{O}(k^c)$ for some constant c , the clique number can be approximated in polynomial time to within a factor of $\text{opt}^{1 - 1/(c + \varepsilon)}$ for all $\varepsilon > 0$, where opt denotes the maximum size of a clique of the input graph G . For general graphs, this problem is notoriously difficult to approximate: for every $\varepsilon > 0$, there is no polynomial-time algorithm for approximating the maximum clique in an n -vertex graph to within a factor of $n^{1 - \varepsilon}$ unless $\text{P} = \text{NP}$ [102].

An interesting open problem is whether (tw, ω) -boundedness has any further algorithmic implications, for example for problems related to independent sets. An *independent set*—or *stable set*—in a (simple, finite, and undirected) graph G is a set of pairwise non-adjacent vertices. The *independence number* of a graph G is the maximum size of an independent set in G . More generally, given a graph G and a weight function $w : V(G) \rightarrow \mathbb{Q}_+$, the MAX WEIGHT INDEPENDENT

SET problem asks to find an independent set I in G of maximum possible weight $w(I)$, where $w(I) = \sum_{x \in I} w(x)$. The MAX WEIGHT INDEPENDENT SET problem is one of the most studied graph optimization problems. It is not only (strongly) NP-hard in general [75] but also NP-hard to approximate on n -vertex graphs to within a factor of $n^{1-\epsilon}$ for every $\epsilon > 0$ [102]. Understanding what restrictions on the input graph make the problem more tractable has been a subject of investigation for decades, with several recent developments [2, 20, 28, 39, 40, 64, 65, 69, 83].

We observed in [50] that in all the (tw, ω) -bounded graph classes defined by excluding a single graph H with respect to any of the six considered graph containment relations (the minor, topological subgraph, subgraph, and their induced variants), the MAX WEIGHT INDEPENDENT SET problem can be solved in polynomial time, except possibly for the case when H is excluded with respect to the induced minor relation and H is isomorphic to either K_5^- (the complete 5-vertex graph minus an edge), W_4 (a wheel with four spokes, that is, the graph obtained from the 4-vertex cycle by adding to it a universal vertex), or the complete bipartite graph $K_{2,q}$ where $q \geq 3$. (See Section 2 for precise definitions.) We asked whether there is a (tw, ω) -bounded graph class in which the MAX WEIGHT INDEPENDENT SET problem is NP-hard. Unless $\text{P} = \text{NP}$, the question can be equivalently formulated as follows.

Question 1.1. Is the MAX WEIGHT INDEPENDENT SET problem solvable in polynomial time in every (tw, ω) -bounded graph class?

1.4 Our results

Motivated by the quest to increase our understanding of algorithmic implications of (tw, ω) -boundedness, we provide in this work a partial answer to Question 1.1 by identifying a sufficient condition for polynomial-time solvability of the MAX WEIGHT INDEPENDENT SET problem in a (tw, ω) -bounded graph class. In fact, we do this in the more general context of the MAX WEIGHT INDEPENDENT PACKING problem. This problem takes as input a graph G and a family \mathcal{H} of connected subgraphs of G , each equipped with a nonnegative weight, and the task is to compute a maximum-weight subfamily of subgraphs from \mathcal{H} such that no two of them share a vertex or are connected by an edge.

In order to explain the corresponding result, we need some additional definitions. In particular, we introduce a new min-max graph parameter related to tree decompositions. Given a tree decomposition \mathcal{T} of a graph G , we define the *independence number* of \mathcal{T} as the maximum independence number over all subgraphs of G induced by some bag of \mathcal{T} . The *tree-independence number* of a graph G is then defined as the minimum independence number over all tree decompositions of G . We say that a graph class \mathcal{G} has *bounded tree-independence number* if there exists an integer k such that every graph in \mathcal{G} has tree-independence number at most k . Furthermore, tree decompositions with bounded independence number are said to be α -*bounded* and we say that the (boundedness of the) tree-independence number in a graph class \mathcal{G} is *efficiently witnessed* if there exists a polynomial-time algorithm to compute an α -bounded tree decomposition for every graph in \mathcal{G} .

We show that graph classes with bounded tree-independence number satisfy the following properties:

- (1) In every graph class with bounded tree-independence number, the treewidth is bounded by an explicit polynomial function of the clique number (see Lemma 3.2); the degree of the polynomial equals the bound on the tree-independence number. Consequently, every graph class with bounded tree-independence number is (tw, ω) -bounded, admits linear-time

algorithms for the k -CLIQUE and the LIST k -COLORING problems for every positive integer k , as well as a polynomial-time robust algorithm for the LIST k -COLORING and LIST k -EDGE-COLORING problems, and an improved approximation for the clique number.

- (2) In every graph class with bounded tree-independence number, the MAX WEIGHT INDEPENDENT PACKING problem can be solved in polynomial time provided that the input graph G is given along with an α -bounded tree decomposition. (See Theorem 7.2.) This implies analogous results for several problems studied in the literature, including the INDEPENDENT \mathcal{F} -PACKING, the MAX WEIGHT INDEPENDENT SET, the MAX WEIGHT INDUCED MATCHING, the DISSOCIATION SET, and the k -SEPARATOR problems. We refer to Section 7 for more details and references.

As part of our approach related to solving the MAX WEIGHT INDEPENDENT PACKING problem in graph classes with efficiently witnessed tree-independence numbers, we show that, given any graph G and a family \mathcal{H} of connected subgraphs of G , the tree-independence number of a certain derived graph $G(\mathcal{H})$ cannot exceed the tree-independence number of G (see Theorem 6.2). Here, the graph $G(\mathcal{H})$ has the subgraphs in \mathcal{H} as vertices, with two adjacent if and only if they share a vertex or there is an edge connecting them. Cameron and Hell showed in [27] that the following graph classes are closed under the transformation $G \mapsto G(\mathcal{H})$ (for any \mathcal{H}): chordal graphs, interval graphs, circular-arc graphs, polygon-circle graphs, interval filament graphs, cocomparability graphs, AT-free graphs, and weakly chordal graphs. Our result adds the classes of graphs with tree-independence number at most k (for any positive integer k) to the list of graph classes that are known to be closed under this transformation, generalizing the result for chordal graphs, which, as we will see, are exactly the graphs with tree-independence number at most one.

The original motivation for introducing the tree-independence number stems from the fact that in a tree decomposition of constant width each bag interacts with an optimal solution to the problem under consideration in only a bounded number of ways, which can be enumerated efficiently. This often leads to efficient dynamic programming algorithms. It is natural to relax the condition on the width to the requirement that for each bag there are only *polynomially many* ways in which an optimal solution can interact with the bag, and that these interactions can be enumerated efficiently. In particular, this happens whenever each bag can interact with an optimal solution in a bounded number of vertices. (This approach was independently suggested by Maria Chudnovsky [34–37].) The definition of the independence number of a tree decomposition is in line with this discussion: it gives an upper bound on the number of vertices that an optimal solution to the MAX WEIGHT INDEPENDENT SET problem can contain from any given bag. Thus, we are interested in tree decompositions of G that are not necessarily of smallest possible width, but that have the smallest possible independence number. This immediately leads to the definition of the tree-independence number.

The algorithmic result about the MAX WEIGHT INDEPENDENT PACKING problem raises two natural questions:

Question 1.2. Which graph classes have bounded tree-independence number?

Question 1.3. For which graph classes can an α -bounded tree decomposition be computed in polynomial time?

The wide applicability of our algorithmic result is indicated by the following two partial answers to Questions 1.2 and 1.3.

First, the following families of graph classes all have bounded tree-independence number: graph classes of bounded treewidth, graph classes of bounded independence number, classes of intersection graphs of connected subgraphs of graphs with bounded treewidth, and classes of graphs in which all minimal separators are of bounded size (see Theorem 3.12).

Second, when a single graph H is excluded with respect to any of six well-known graph containment relations (the minor, topological subgraph, subgraph, and their induced variants), the resulting graph class is (tw, ω) -bounded if and only if it has bounded tree-independence number (see the third paper of the series [49]). In each such case there is a polynomial-time algorithm for computing an α -bounded tree decomposition. These results are obtained via characterizations, for each of six graph containment relations, of the graphs H such that the class of graphs excluding H has bounded tree-independence number. For completeness, we recall those characterizations below.

A graph is *subcubic* if every vertex of it has degree at most three. We denote by \mathcal{S} the class of forests in which every connected component has at most three leaves (vertices of degree one).

Theorem 1.4 (Dallard, Milanič, and Štorgel [49]). *For every graph H , the following statements hold.*

1. *The class of H -subgraph-free graphs has bounded tree-independence number if and only if $H \in \mathcal{S}$.*
2. *The class of H -topological-minor-free graphs has bounded tree-independence number if and only if H is subcubic and planar.*
3. *The class of H -minor-free graphs has bounded tree-independence number if and only if H is planar.*
4. *The class of H -free graphs has bounded tree-independence number if and only if H is either an induced subgraph of P_3 or an edgeless graph.*
5. *The class of H -induced-topological-minor-free graphs has bounded tree-independence number if and only if H is either an induced topological minor of C_4 or K_4^- , or H is edgeless.*
6. *The class of H -induced-minor-free graphs has bounded tree-independence number if and only if H is an induced minor of W_4 , K_5^- , or $K_{2,q}$ for some $q \in \mathbb{Z}_+$, or H is edgeless.*

In particular, this result leads to a new infinite family of graph classes in which the MAX WEIGHT INDEPENDENT SET problem is solvable in polynomial time. We refer to [49] for details.

Since the tree-independence number is a newly introduced graph invariant, we also provide a set of results about its basic properties (see Section 3). These include a characterization of graphs with tree-independence number at most one (as exactly the chordal graphs), the NP-hardness of the problem of computing the tree-independence number, two sharp upper bounds in terms of the independence number and the treewidth, the monotonicity under vertex deletions and edge contractions, and the behavior under clique cutsets.

1.5 Overview of our methodology

We now summarize the key ideas leading to our main results.

We develop polynomial-time algorithms for the MAX WEIGHT INDEPENDENT PACKING problem in graph classes with bounded and efficiently witnessed tree-independence number in two steps. First, we develop a dynamic programming algorithm for the MAX WEIGHT INDEPENDENT SET

problem, assuming that the input graph is equipped with a tree decomposition of bounded independence number (Corollary 5.3). We do this by adapting the standard dynamic programming approach for the MAX WEIGHT INDEPENDENT SET problem over tree decompositions of bounded width to the more general setting of tree decompositions with bounded independence number. We in fact derive an improved version of this algorithmic result by introducing the concept of an ℓ -refined tree decomposition, which is a tree decomposition in which each bag comes equipped with a subset of at most ℓ vertices (see Section 4). Ignoring such vertices when computing the independence number of the subgraph induced by the bag may lead to better running times (Theorem 5.2). Two particular examples of such improvements are given in [49]. Second, we show that the MAX WEIGHT INDEPENDENT PACKING problem can be solved in polynomial time if the input graph is equipped with an α -bounded tree decomposition (Theorem 7.2). We do this by reducing the problem to the MAX WEIGHT INDEPENDENT SET problem in the derived graph $G(\mathcal{H})$. More specifically, we show that in this case one can compute in polynomial time both the graph $G(\mathcal{H})$ as well as a tree decomposition \mathcal{T}' of $G(\mathcal{H})$ such that the independence number of \mathcal{T}' is at most k , where k is a fixed upper bound on the independence number of the tree decomposition \mathcal{T} of the input graph G (Corollary 6.4).

1.6 Relation to previous work

We asked in [48, 50] whether every (tw, ω) -bounded graph class is in fact *polynomially* (tw, ω) -bounded, in the sense that the treewidth of graphs in the class and all their induced subgraphs is bounded by a fixed *polynomial* function of the clique number. This is indeed the case for graph classes with bounded tree-independence number (Lemma 3.2). In general, the notion of the tree-independence number may give some insight into the properties of a possible counterexample to the question about the equality of (tw, ω) -boundedness and polynomial (tw, ω) -boundedness: any (tw, ω) -bounded graph class that is not polynomially (tw, ω) -bounded has to contain graphs with arbitrarily large tree-independence number.

The concept of tree-independence number is also related to χ -boundedness. In the late 1980s, Gyárfás introduced the concept of χ -bounded graph classes in order to generalize perfection [70]. The *chromatic number* of G is the smallest positive integer k such that G is k -colorable (that is, its vertex set can be expressed as the union of k independent sets) and a graph class \mathcal{G} is said to be χ -bounded if the chromatic number of the graphs in the class and all their induced subgraphs is bounded from above by some function of the clique number. The χ -bounded graph classes were studied extensively in the literature (see [90] for a survey). Every (tw, ω) -bounded graph class is χ -bounded (see, e.g., [50]). Esperet asked whether every χ -bounded graph class is *polynomially* χ -bounded, that is, the chromatic number of the graphs in the class and all their induced subgraphs is bounded by a polynomial function of the clique number (see [56]). For general classes, Esperet's question was recently answered in the negative by Briański, Davies, and Walczak [22]. Nonetheless, since the chromatic number of any graph is bounded from above by its treewidth plus one, we obtain the following.

Fact. *Graph classes with bounded tree-independence number are polynomially χ -bounded.*

Our result on the polynomial-time solvability of the MAX WEIGHT INDEPENDENT SET problem in classes of graphs equipped with a tree decomposition of bounded independence number (Corollary 5.3) generalizes a result of Krause who showed in [77] that the MAX INDEPENDENT SET problem is solvable in polynomial time on intersection graphs of connected subgraphs of graphs of bounded treewidth (assuming that the input graph is given with an intersection model). In some sense, our algorithmic result for the MAX WEIGHT INDEPENDENT PACKING problem (Theorem 7.2)

generalizes the polynomial-time solvability of the MAX INDUCED MATCHING problem in the class of chordal graphs [26], of the k -SEPARATOR problem (for fixed k) in the classes of interval graphs and graphs of bounded treewidth [7], and of the INDEPENDENT \mathcal{F} -PACKING problem in the classes of chordal and circular-arc graphs [27].

On a more general note, let us remark that the tree-independence number shares the property of several graph invariants studied in the literature in the sense that they are defined by the existence of a tree decomposition with certain properties of the bags. These include connectivity properties (connected treewidth [51], also known as bag-connected treewidth [73]), metric properties (tree-length [53] and tree-breadth [54]), and chromatic properties (tree-chromatic number [91]).

1.7 Organization of the paper

In Section 2, we provide the necessary preliminaries. In Section 3, we derive several basic properties of the tree-independence number, including the NP-hardness of the problem of computing it. In Section 4, we introduce the concepts of ℓ -refined tree decompositions and their residual independence numbers. In Section 5, we prove that if \mathcal{G} is a graph class with bounded and efficiently witnessed tree-independence number, then there exists a polynomial-time algorithm for the MAX WEIGHT INDEPENDENT SET problem when the input graph is restricted to belong to \mathcal{G} . We show in Section 6 that for any graph G and any family \mathcal{H} of connected subgraphs of G , the derived graph $G(\mathcal{H})$ has tree-independence number bounded by that of G . This allows us to generalize the algorithmic result of Section 5 by developing, in Section 7, an efficient algorithm for the MAX WEIGHT INDEPENDENT PACKING problem, provided that the input graph G is given along with an α -bounded tree decomposition. We conclude the paper with some open questions in Section 8.

2 Preliminaries

Throughout the paper, we denote by \mathbb{Z}_+ the set of nonnegative integers and by \mathbb{Q}_+ the set of nonnegative rational numbers. We assume familiarity with the basic concepts in graph theory as used, e.g., by West [99]. We denote the vertex set and the edge set of a graph G by $V(G)$ and $E(G)$, respectively. A graph is *null* if it has no vertices. The *neighborhood* of a vertex v in G , which corresponds to the set of vertices adjacent to v in G , is denoted by $N_G(v)$. The *closed neighborhood* of v is the set $N_G[v] = N_G(v) \cup \{v\}$. These concepts are extended to sets $X \subseteq V(G)$ so that $N_G[X]$ is defined as the union of all closed neighborhoods of vertices in X , and $N_G(X)$ is defined as the set $N_G[X] \setminus X$. The *degree* of v , denoted by $d_G(v)$, is the cardinality of the set $N_G(v)$. When there is no ambiguity, we may omit the subscript G in the notations of the degree, and open and closed neighborhoods, and thus simply write $d(v)$, $N(v)$, and $N[v]$, respectively. Given a set $X \subseteq V(G)$, we denote by $G - X$ the graph obtained from G after deleting all the vertices in X , and by $G[X]$ the subgraph of G induced by X , that is, the graph $G - (V(G) \setminus X)$. Similarly, given a vertex $v \in V(G)$, we denote by $G - v$ the graph obtained from G after deleting v . The fact that two graphs G and H are isomorphic to each other is denoted by $G \cong H$. For a positive integer n , we denote the n -vertex complete graph, path, and cycle by K_n , P_n , and C_n , respectively. Similarly, for $n \geq 2$ we denote by K_n^- the graph obtained from the complete graph K_n by deleting an edge, and for positive integers m and n we denote by $K_{m,n}$ the complete bipartite graph with parts of sizes m and n . The graph K_4^- is also called the *diamond*. For $n \geq 4$, we denote by W_n the graph obtained from a cycle C_n by adding to it a universal vertex (that is, a vertex adjacent to all vertices of the C_n).

An *independent set* in a graph G is a set of pairwise non-adjacent vertices, and a *clique* is a set of pairwise adjacent vertices. The *independence number* of G , denoted by $\alpha(G)$, is the maximum size of an independent set in G . The *clique number* of G , denoted by $\omega(G)$, is the maximum size of a clique in G . Given two non-adjacent vertices u and v in G , a *u,v -separator* is a set $S \subseteq V(G) \setminus \{u, v\}$ such that u and v belong to different components of $G - S$. A *u,v -separator* S is *minimal* if no proper subset of S is a *u,v -separator*. A *minimal separator* in G is a set $S \subseteq V(G)$ that is a minimal *u,v -separator* for some non-adjacent vertex pair u, v .

A graph H is said to be an *induced subgraph* of G if H can be obtained from G by deleting vertices. If H is obtained from G by deleting vertices and edges, then H is a *subgraph* of G . The *subdivision of an edge uv* of a graph is the operation that deletes the edge uv and adds a new vertex w and two edges uw and wv . A *subdivision of a graph H* is a graph obtained from H by a sequence of edge subdivisions. A graph H is said to be a *topological minor* of a graph G if G contains a subdivision of H as a subgraph. Similarly, H is an *induced topological minor* of G if some subdivision of H is isomorphic to an induced subgraph of G . An *edge contraction* is the operation of deleting a pair of adjacent vertices and replacing them with a new vertex whose neighborhood is the union of the neighborhoods of the two original vertices. If H can be obtained from G by a sequence of vertex deletions, edge deletions, and edge contractions, then H is said to be a *minor* of G . Finally, we say that G contains H as an *induced minor* if H can be obtained from G by a sequence of vertex deletions and edge contractions. If G does not contain an induced subgraph isomorphic to H , then we say that G is *H -free*. Analogously, we also say that G is *H -subgraph-free*, *H -topological-minor-free*, *H -induced-topological-minor-free*, *H -minor-free*, or *H -induced-minor-free*, respectively, for the other five graph containment relations.

A *tree decomposition* of a graph G is a pair $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ where T is a tree and every node t of T is assigned a vertex subset $X_t \subseteq V(G)$ called a bag such that the following conditions are satisfied: every vertex is in at least one bag, for every edge $uv \in E(G)$ there exists a node $t \in V(T)$ such that X_t contains both u and v , and for every vertex $u \in V(G)$ the subgraph T_u of T induced by the set $\{t \in V(T) : u \in X_t\}$ is connected (that is, a tree). The *width* of \mathcal{T} , denoted by $\text{width}(\mathcal{T})$, is the maximum value of $|X_t| - 1$ over all $t \in V(T)$. The *treewidth* of a graph G , denoted by $\text{tw}(G)$, is defined as the minimum width of a tree decomposition of G .

We next state two useful properties of tree decompositions.

Lemma 2.1 (Scheffler [89], Bodlaender and Möhring [14]). *Let G be a graph, let $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ be a tree decomposition of G , and let C be a clique in G . Then there exists a bag X_t such that $C \subseteq X_t$.*

Corollary 2.2. *Every graph G satisfies $\text{tw}(G) \geq \omega(G) - 1$.*

Lemma 2.3. *Let $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ be a tree decomposition of a graph G . Then for every set $S \subseteq V(G)$ such that every pair of vertices in S is contained in a bag of \mathcal{T} , there exists a bag X_t such that $S \subseteq X_t$.*

Proof. Suppose that every pair of vertices of S is contained in a bag of \mathcal{T} . For each $v \in S$, the set of nodes of T labeled by bags containing v induces a subtree T_v of T . Since every two vertices in S are contained in a bag of \mathcal{T} , every two of the subtrees in $\{T_v : v \in S\}$ have a node in common. It is known (and easy to see) that any family of node sets of subtrees of a tree has the Helly property, and thus there exists a node $t \in V(T)$ common to all the trees T_v , $v \in S$. We infer that $v \in X_t$ for all $v \in S$, that is, $S \subseteq X_t$. \square

There are many equivalent characterizations of treewidth, including one in terms of chordal graphs. A graph G is *chordal* if it has no induced cycles of length at least four. Given a graph $G = (V, E)$, a *triangulation* of G is a chordal graph of the form $(V, E \cup F)$ where F is a subset of the set of non-edges of G . A triangulation $(V, E \cup F)$ of G is a *minimal triangulation* of G if G has no triangulation $(V, E \cup F')$ such that F' is a proper subset of F .

Theorem 2.4 (Robertson and Seymour [87]). *For every graph G , we have*

$$\text{tw}(G) = \min\{\omega(H) - 1 : H \text{ is a triangulation of } G\}.$$

Our approach in proving the NP-hardness of computing the tree-independence number in Section 3 relies on the fact that every tree decomposition has a bag containing the closed neighborhood of some vertex (see Lemma 2.6). This property of tree decompositions implies the inequality $\text{tw}(G) \geq \delta(G)$, where $\delta(G)$ denotes the minimum degree of a vertex in G , valid for any graph G (see, e.g., [13]).

In order to prove Lemma 2.6, we first recall some of the many characterizations of chordal graphs. A *clique tree* of a graph G is a tree decomposition of G such that the bags are exactly the maximal cliques of G . Given a collection $\{T_1, \dots, T_n\}$ of subtrees in a tree T , the *intersection graph* of $\{T_1, \dots, T_n\}$ is the graph with vertex set $\{1, \dots, n\}$, in which two distinct vertices i and j are adjacent if and only if T_i and T_j have a vertex in common. Given a graph G , a *simplicial vertex* in G is a vertex whose neighborhood is a clique. The following theorem combines results from several sources [9, 25, 52, 66, 98].

Theorem 2.5. *Let G be a graph. Then, the following conditions are equivalent:*

1. G is a chordal graph.
2. G has a clique tree.
3. G is the intersection graph of subtrees in a tree.
4. Every nonnull induced subgraph of G has a simplicial vertex.

Lemma 2.6. *Let G be a graph and let $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ be a tree decomposition of G . Then there exists a vertex $v \in V(G)$ and a node $t \in V(T)$ such that $N[v] \subseteq X_t$.*

Proof. Let G' be the graph with vertex set $V(G)$ such that two distinct vertices u and v are adjacent in G' if and only if there exists a bag X_t of \mathcal{T} with $u, v \in X_t$. Note that for every vertex $v \in V(G)$ it holds that $N_G[v] \subseteq N_{G'}[v]$. Recall that for each vertex $v \in V(G')$, the set of nodes $t \in V(T)$ such that $v \in X_t$ induces a subtree T_v of T . Thus, two distinct vertices u and v of G' are adjacent if and only if the corresponding trees T_u and T_v have a node in common. This means that G' is the intersection graph of subtrees in a tree and hence, by Theorem 2.5, G' is a chordal graph. Observe that \mathcal{T} is also a tree decomposition of G' . By Theorem 2.5, G' has a simplicial vertex v , and hence by Lemma 2.1, there must exist a node $t \in V(T)$ such that $N_{G'}[v] \subseteq X_t$. Therefore, $N_G[v] \subseteq X_t$, which concludes the proof. \square

Let us remark that Lemma 2.6 can also be derived from the proofs of Lemmas 2 and 4 in [13] (which do not rely on connections with chordal graphs).

An (integer) *graph invariant* is a mapping from the class of all graphs to the set of nonnegative integers that does not distinguish between isomorphic graphs. For a graph invariant ρ , we say that a graph class \mathcal{G} has *bounded ρ* if there exists an integer k such that $\rho(G) \leq k$ for every graph G

in the class. A graph class \mathcal{G} is said to be (tw, ω) -bounded if it admits a (tw, ω) -binding function, that is, a function f such that for every graph G in the class and any induced subgraph G' of G , the treewidth of G' is at most $f(\omega(G'))$. Ramsey's theorem [85] states that for every two positive integers p and q there exists an integer $N(p, q)$ such that every graph with at least $N(p, q)$ vertices contains either a clique of size p or an independent set of size q . The least such positive integer is denoted by $R(p, q)$.

3 Definition and basic properties

We now define the two main concepts studied in this paper.

Definition 3.1. Consider a graph G and a tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ of G . The *independence number* of \mathcal{T} , denoted by $\alpha(\mathcal{T})$, is defined as follows:

$$\alpha(\mathcal{T}) = \max_{t \in V(T)} \alpha(G[X_t]).$$

The *tree-independence number* of G , denoted by $\text{tree-}\alpha(G)$, is the minimum independence number among all possible tree decompositions of G .

In this section we derive several basic properties of the tree-independence number of graphs. First, we observe that Ramsey's theorem implies that graph classes with bounded tree-independence number are (tw, ω) -bounded with a polynomial binding function.

Lemma 3.2. *For every positive integer k , the class of graphs with tree-independence number at most k is (tw, ω) -bounded, with a binding function $f(p) = R(p+1, k+1) - 2$, which is a polynomial of degree k .*

Proof. The standard proof of Ramsey's theorem is based on the inequality $R(p, q) \leq R(p-1, q) + R(p, q-1)$ for all $p, q \geq 2$, which implies that $R(p, q) \leq \binom{p+q-2}{q-1}$ for all positive integers p and q . For fixed q , this is a polynomial in p of degree $q-1$.

Let us now fix $p \in \mathbb{Z}_+$ and let G be a graph such that $\omega(G) = p$ and $\text{tree-}\alpha(G) \leq k$. Fix a tree decomposition \mathcal{T} of G with independence number at most k . Note that every bag of \mathcal{T} induces a subgraph of G with independence number at most k and clique number at most p . Thus, for every bag X of \mathcal{T} , Ramsey's theorem implies that $|X| \leq R(p+1, k+1) - 1$. It follows that $\text{tw}(G) \leq R(p+1, k+1) - 2$, as claimed. \square

Next, we show that the tree-independence number can be used to characterize chordal graphs.

Theorem 3.3. *Let G be a graph. Then $\text{tree-}\alpha(G) \leq 1$ if and only if G is chordal.*

Proof. If G is chordal, then, by Theorem 2.5, G has a clique tree and thus $\text{tree-}\alpha(G) \leq 1$. Conversely, if G has a tree decomposition \mathcal{T} with independence number at most one, then every bag of \mathcal{T} is a clique in G . Since \mathcal{T} is a tree decomposition, for every vertex u of G the subgraph T_u of T induced by the set $\{t \in V(T) : u \in X_t\}$ is a tree. Furthermore, since each bag of \mathcal{T} is a clique, two distinct vertices u and v of G are adjacent if and only if they belong to a same bag, which is in turn equivalent to the condition that T_u and T_v have a vertex in common. Thus, G is the intersection graph of the collection of subtrees $\{T_u : u \in V(G)\}$. Applying Theorem 2.5, we conclude that G is chordal. \square

A tree decomposition of a graph G is said to be *trivial* if it has a single bag $X_t = V(G)$. Next, we establish the NP-hardness of the problem of computing the tree-independence number of a graph. Our reduction is based on the following lemma, which we derive from Lemma 2.6.

Lemma 3.4. *Let G be a graph and let G' be the graph obtained from two disjoint copies of G by adding all possible edges between them. Then $\text{tree-}\alpha(G') = \alpha(G)$.*

Proof. Let us denote by G_1 and G_2 the two disjoint copies of G such that $V(G') = V(G_1) \cup V(G_2)$. Observe that every independent set in G' is entirely contained in either G_1 or G_2 and hence, we have $\alpha(G') = \alpha(G)$. Thus, the trivial tree decomposition of G' has independence number equal to $\alpha(G)$ and consequently $\text{tree-}\alpha(G') \leq \alpha(G)$.

For the converse direction, consider an arbitrary tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ of G' . By Lemma 2.6, there exists a vertex $v \in V(G)$ and a node $t \in V(T)$ such that $N[v] \subseteq X_t$. Assume without loss of generality that $v \in V(G_1)$. Then $V(G_2) \subseteq N(v) \subseteq X_t$, and therefore $\alpha(G'[X_t]) \geq \alpha(G_2) = \alpha(G)$. Thus, every tree decomposition of G' contains a bag inducing a subgraph with independence number at least $\alpha(G)$. This shows that $\text{tree-}\alpha(G') \geq \alpha(G)$. Therefore, equality must hold. \square

Since computing the independence number of a graph is NP-hard [75], Lemma 3.4 implies the following.

Theorem 3.5. *Computing the tree-independence number of a given graph is NP-hard.*

For later use we also record the following consequence of Lemma 3.4.

Corollary 3.6. *For every positive integer n , we have $\text{tree-}\alpha(K_{n,n}) = n$.*

Given any graph G , the trivial tree decomposition of G has independence number equal to $\alpha(G)$. This immediately implies the following.

Observation 3.7. *For every graph G we have $\text{tree-}\alpha(G) \leq \alpha(G)$.*

As shown by Corollary 3.6, this bound is sharp. Another easily obtained upper bound on the tree-independence number of a graph is related to the treewidth of the graph. This bound is sharp as well.

Theorem 3.8. *For every graph G , $\text{tree-}\alpha(G) \leq \text{tw}(G) + 1$, and this bound is sharp: for every integer $k \neq 2$, there exists a graph G such that $\text{tree-}\alpha(G) = k$ and $\text{tw}(G) = k - 1$.*

Proof. It follows directly from the definitions that the independence number of any tree decomposition is at most its width plus one. Thus, taking \mathcal{T} to be a tree decomposition of G with minimum possible width, we obtain $\text{tree-}\alpha(G) \leq \alpha(\mathcal{T}) \leq \text{width}(\mathcal{T}) + 1 = \text{tw}(G) + 1$.

For $k = 1$, the graph K_1 satisfies $\text{tree-}\alpha(K_1) = k$ and $\text{tw}(K_1) = k - 1$.

Fix $k \geq 3$, let $S = \{1, \dots, k\}$, and let G be the graph obtained from a complete graph with vertex set S by replacing each of its edges ij with k paths of length two connecting i and j . We claim that $\text{tw}(G) = k - 1$ and $\text{tree-}\alpha(G) = k$. Since $\text{tree-}\alpha(G) \leq \text{tw}(G) + 1$, it suffices to show that $\text{tw}(G) \leq k - 1$ and that $\text{tree-}\alpha(G) \geq k$.

The inequality $\text{tw}(G) \leq k - 1$ follows from Theorem 2.4 and the observation that the graph G' obtained from G by adding to it all edges between vertices in the set S is a chordal graph with clique number k .

It remains to show that $\text{tree-}\alpha(G) \geq k$. Consider an arbitrary tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ of G . We claim that there exists a bag of \mathcal{T} having independence number at least k . This is clearly the case if there exists a bag X_t such that $S \subseteq X_t$. Thus, we may assume that no bag of \mathcal{T} contains S . By Lemma 2.3, there exist two distinct vertices $i, j \in S$ that are not contained in the same bag of \mathcal{T} . Since i and j are not contained in a same bag of \mathcal{T} , the subtrees T_i

and T_j are disjoint. Note that, by construction of G , the vertices i and j have k common neighbors. Let $u \in N(i) \cap N(j)$ be such a vertex. Then u belongs to a bag in \mathcal{T} containing i , and similarly for j . In other words, T_u intersects T_i and T_j , which implies that $T_i \cup T_u \cup T_j$ is connected. Let P be the path in T connecting T_i and T_j . Clearly, P is a subgraph of T_u , and thus for any node $t \in P$, the bag X_t of \mathcal{T} contains u . Since this holds for every common neighbor of i and j , every such bag contains $N(i) \cap N(j)$, which is an independent set of size k . Thus, as \mathcal{T} can be any tree decomposition of G , we conclude that $\text{tree-}\alpha(G) \geq k$. \square

Note that there are no graphs G with $\text{tree-}\alpha(G) = \text{tw}(G) + 1 = 2$. Any such graph would satisfy $\text{tw}(G) = 1$ and would therefore be acyclic, and hence chordal. Theorem 3.3 would then imply that $\text{tree-}\alpha(G) \leq 1$, a contradiction.

The treewidth is well-known to be monotone under minors, in the sense that $\text{tw}(G) \leq \text{tw}(G')$ for any graph G and any minor G' of G . This is not the case for the tree-independence number, as it can increase under edge deletions. This fact can be easily observed, e.g., using the families of complete graphs and complete bipartite graphs (cf. Theorem 3.3 and Corollary 3.6). On the other hand, the tree-independence number cannot be increased by vertex deletions and edge contractions.

Proposition 3.9. *Let G be a graph and G' an induced minor of G . Then $\text{tree-}\alpha(G') \leq \text{tree-}\alpha(G)$.*

Proof. Let $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ be an arbitrary tree decomposition of G . First, we show that the deletion of a vertex does not increase the tree-independence number. Let v be a vertex of G . Let \mathcal{T}' be the tree decomposition obtained from \mathcal{T} by removing v from all of the bags that contain it. Observe that \mathcal{T}' is a tree decomposition of $G - v$. Clearly, $\alpha(\mathcal{T}') \leq \alpha(\mathcal{T})$, and hence $\text{tree-}\alpha(G - v) \leq \text{tree-}\alpha(G)$.

Now, we show that the contraction of an edge does not increase the tree-independence number. Let $e = uv$ be an edge of G and G/e denote the graph obtained from G after contracting the edge e . We denote by w the vertex of G/e that corresponds to the contracted edge. We construct a tree decomposition $\mathcal{T}' = (T, \{X'_t\}_{t \in V(T)})$ (where T is the same tree as in \mathcal{T}) of G/e as follows. For each node t of \mathcal{T}' , we have two cases: if X_t contains neither u nor v , then we set $X'_t = X_t$; otherwise, we set $X'_t = (X_t \setminus \{u, v\}) \cup \{w\}$. We claim that \mathcal{T}' is a tree decomposition of G/e such that $\alpha(\mathcal{T}') \leq \alpha(\mathcal{T})$. First, observe that \mathcal{T}' is a tree decomposition of G/e , as it satisfies all the defining conditions of a tree decomposition. To verify that $\alpha(\mathcal{T}') \leq \alpha(\mathcal{T})$, fix a bag X'_t of \mathcal{T}' and let $I \subseteq X'_t$ be an independent set in G/e . If $w \notin I$, then I also corresponds to an independent set in $G[X_t]$, and hence $|I| \leq \alpha(G[X_t])$. On the other hand, if $w \in I$, then either $(I \setminus \{w\}) \cup \{u\}$ or $(I \setminus \{w\}) \cup \{v\}$ is an independent set in $G[X_t]$. In particular, we again get that $|I| \leq \alpha(G[X_t])$. Thus, we have in both cases that $|I| \leq \alpha(\mathcal{T})$. It follows that $\alpha(\mathcal{T}') \leq \alpha(\mathcal{T})$, and hence $\text{tree-}\alpha(G/e) \leq \text{tree-}\alpha(G)$.

If G' is an induced minor of G , then G' can be obtained from G by a sequence of vertex deletions and edge contractions, which implies that $\text{tree-}\alpha(G') \leq \text{tree-}\alpha(G)$. \square

We now show a result that we will use in the proof of Theorem 3.12, and which can find further applications when combined with Tarjan's decomposition of a graph along clique cutsets [95]. A *cut-partition* in a graph G is a triple (A, B, C) such that $A \cup B \cup C = V(G)$, $A \neq \emptyset$, $B \neq \emptyset$, and G has no edges with one endpoint in A and one in B . A *clique cutset* in G is a clique C such that G has a cut-partition (A, B, C) for some A and B .

Proposition 3.10. *Let C be a clique cutset in a graph G and let (A, B, C) be a cut-partition of G . Let $G_A = G[A \cup C]$ and $G_B = G[B \cup C]$, and let \mathcal{T}_A and \mathcal{T}_B be tree decompositions of G_A and G_B , respectively. Then we can compute in time $\mathcal{O}(|\mathcal{T}_A| + |\mathcal{T}_B|)$ a tree decomposition \mathcal{T} of G such that $\alpha(\mathcal{T}) = \max\{\alpha(\mathcal{T}_A), \alpha(\mathcal{T}_B)\}$.*

Proof. Since C is a clique in G_A and in G_B , by Lemma 2.1 there exists a bag X_A of \mathcal{T}_A such that $C \subseteq X_A$, and a bag X_B in \mathcal{T}_B such that $C \subseteq X_B$. Take the disjoint union of the tree T_A of \mathcal{T}_A with the tree T_B of \mathcal{T}_B , and add an edge connecting the nodes corresponding to X_A and X_B . This results in a tree decomposition $\mathcal{T} = (T, \{(X_t, U_t)\}_{t \in V(T)})$ of G . Indeed, since every vertex of G is a vertex of G_A or G_B , every vertex of G is in at least one bag of \mathcal{T} . A similar argument shows that for every edge of G , both endpoints belong to a common bag. Finally, let us verify that for every vertex $u \in V(G)$, the subgraph of T induced by the set of bags containing u is connected. If $u \in A \cup B$, then this follows from the corresponding properties of \mathcal{T}_A and \mathcal{T}_B . Suppose now that $u \in C$. Then the subgraph $T_A(u)$ of T_A induced by the set of bags of \mathcal{T}_A containing u is connected and contains the node corresponding to X_A . Similarly, the subgraph $T_B(u)$ of T_B induced by the set of bags of \mathcal{T}_B containing u is connected and contains the node corresponding to X_B . Thus, the subgraph $T(u)$ of T induced by the set of bags of \mathcal{T} containing u is isomorphic to the graph obtained from the disjoint union of $T_A(u)$ and $T_B(u)$ together with the edge between X_A and X_B , and hence $T(u)$ is connected. By construction, the independence number of \mathcal{T} satisfies $\alpha(\mathcal{T}) = \max\{\alpha(\mathcal{T}_A), \alpha(\mathcal{T}_B)\}$. Finding a bag X_A such that $C \subseteq X_A$ can be done in time $\mathcal{O}(|\mathcal{T}_A|)$, and similarly for X_B . Thus, the time complexity of the above procedure is $\mathcal{O}(|\mathcal{T}_A| + |\mathcal{T}_B|)$. \square

Corollary 3.11. *Let C be a clique cutset in a graph G , let (A, B, C) be a cut-partition of G , and let $G_A = G[A \cup C]$ and $G_B = G[B \cup C]$. Then $\text{tree-}\alpha(G) = \max\{\text{tree-}\alpha(G_A), \text{tree-}\alpha(G_B)\}$.*

Proof. To justify that $\text{tree-}\alpha(G) = \max\{\text{tree-}\alpha(G_A), \text{tree-}\alpha(G_B)\}$, note first that G_A and G_B are induced subgraphs of G and hence $\max\{\text{tree-}\alpha(G_A), \text{tree-}\alpha(G_B)\} \leq \text{tree-}\alpha(G)$ by Proposition 3.9. For the converse inequality, let \mathcal{T}_A and \mathcal{T}_B be tree decompositions of G_A and G_B , respectively, such that $\alpha(\mathcal{T}_A) = \text{tree-}\alpha(G_A)$ and $\alpha(\mathcal{T}_B) = \text{tree-}\alpha(G_B)$. Proposition 3.10 implies the existence of a tree decomposition \mathcal{T} of G such that $\alpha(\mathcal{T}) = \max\{\alpha(\mathcal{T}_A), \alpha(\mathcal{T}_B)\}$. It follows that $\text{tree-}\alpha(G) \leq \alpha(\mathcal{T}) = \max\{\alpha(\mathcal{T}_A), \alpha(\mathcal{T}_B)\} = \max\{\text{tree-}\alpha(G_A), \text{tree-}\alpha(G_B)\}$. \square

We conclude this section by summarizing some sufficient conditions for a graph class to have bounded tree-independence number. To state the result we need to introduce names for two further graph invariants. Given a graph H and connected subgraphs H_1, \dots, H_p of H , the *intersection graph* of H_1, \dots, H_p is the graph with vertex set $\{1, \dots, p\}$ in which two distinct vertices i and j are adjacent if and only if H_i and H_j have a vertex in common. We say that a graph G is *representable* in a graph H if there exist connected subgraphs H_1, \dots, H_p of H such that G is isomorphic to their intersection graph. Note that every graph G is representable in the graph H obtained by subdividing each edge of G exactly once: each vertex v of G corresponds to the subgraph of H induced by the closed neighborhood of v in H . In particular, this means that the following graph invariant is well-defined. The *intersection treewidth* of a graph G , denoted by $\text{itw}(G)$, is the smallest nonnegative integer k such that there exists a graph H with treewidth at most k such that G is representable in H . Note that if G is a chordal graph, then $\text{itw}(G) \leq 1$, and if G is a circular-arc graph, then $\text{itw}(G) \leq 2$.

Furthermore, we define the *maximum minimal separator size* of a graph G , denoted by $\text{mms}(G)$, as the smallest nonnegative integer k such that every minimal separator of G has cardinality at most k . If G is a complete graph, then G does not have any minimal separators. We extend the definition to this case by setting $\text{mms}(G) = 0$ if G is complete.

Theorem 3.12. *Let \mathcal{G} be a graph class that has either bounded treewidth, bounded independence number, bounded intersection treewidth, or bounded maximum minimal separator size. Then \mathcal{G} has bounded tree-independence number.*

Proof. Fix a positive integer k , a graph invariant $\rho \in \{\text{tw}, \alpha, \text{itw}, \text{mms}\}$ and a graph G such that $\rho(G) \leq k$. It suffices to show that $\text{tree-}\alpha(G) \leq f(k)$ for some function $f : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$. For $\rho = \text{tw}$, Theorem 3.8 applies and we can take $f(k) = k + 1$. For $\rho = \alpha$, Observation 3.7 applies and we can take f to be the identity function. Assume now that $\rho = \text{itw}$, that is, $\text{itw}(G) \leq k$. As shown by Bodlaender, Gustedt, and Telle in the proof of [10, Lemma 2.4], G has a tree decomposition \mathcal{T} in which each bag is the union of at most $k + 1$ cliques in G . This implies that \mathcal{T} has independence number at most $k + 1$ and hence $\text{tree-}\alpha(G) \leq k + 1$. Finally, let $\rho = \text{mms}$, that is, $\text{mms}(G) \leq k$. We claim that $\text{tree-}\alpha(G) \leq 2k - 1$. It follows from a result by Skodinis [92] that G is either a complete graph, or G has a clique cutset, or $\text{tw}(G) \leq 2k - 1$. If G is a complete graph, then $\text{tree-}\alpha(G) \leq 1$. If G has treewidth at most $2k - 1$, then $\text{tree-}\alpha(G) \leq 2k - 1$. Assume now that G has a clique cutset. For every induced subgraph G' of G and every minimal separator S' in G' there exists a minimal separator S in G such that $S' \subseteq S$. Indeed, if S' is a minimal u, v -separator in G' , then any minimal u, v -separator in G contained in the u, v -separator $S' \cup (V(G) \setminus V(G'))$ in G satisfies $S' \subseteq S$. Therefore, every induced subgraph G' of G has $\text{mms}(G') \leq \text{mms}(G) \leq k$, and we can use induction on $|V(G)|$ and Proposition 3.10 to infer that $\text{tree-}\alpha(G) \leq 2k - 1$. \square

In the third paper of the series [49], we derive the fact the class of graphs of bounded maximum minimal separator size has bounded tree-independence number also as a consequence of a more general result, namely, that boundedness of the tree-independence number is achieved whenever the independence number of subgraphs induced by minimal separators is bounded.

4 A refinement of the tree-independence number

As explained in the introduction, one of the main motivations for introducing the tree-independence number of a graph is algorithmic: tree decompositions with small independence number allow for the development of efficient dynamic programming algorithms for the MAX WEIGHT INDEPENDENT SET and related problems (see Sections 5 and 7). For the MAX WEIGHT INDEPENDENT SET problem, given a graph G equipped with a tree decomposition \mathcal{T} with independence number at most k , the algorithm takes time $\mathcal{O}(|V(G)|^k)$ for each bag of \mathcal{T} , in order to consider all subsets of at most k vertices from the bag. In some cases, this approach can be improved. Suppose that in each bag X_t we can identify a set $U_t \subseteq X_t$ of at most a constant number ℓ of vertices such that the independence number of the subgraph of G induced by the remaining vertices, $X_t \setminus U_t$, is at most k . Then, even though we can only bound the independence number of the subgraph of G induced by X_t by $k + \ell$, we do not need to examine all $\mathcal{O}(|X_t|^{k+\ell}) = \mathcal{O}(|V(G)|^{k+\ell})$ subsets of the bag of size at most $k + \ell$, but only $\mathcal{O}(2^\ell \cdot |X_t \setminus U_t|^k)$ of them, since every independent set of $G[X_t]$ consists of a subset of U_t along with at most k of the remaining vertices. For constant values of ℓ , the algorithm will thus take time $\mathcal{O}(|V(G)|^k)$ instead of $\mathcal{O}(|V(G)|^{k+\ell})$ for each bag, which may lead to a significantly improved running time. This idea is formalized in the concepts of ℓ -refined tree decompositions and their residual independence numbers.

Definition 4.1. Given a nonnegative integer ℓ , an ℓ -refined tree decomposition of a graph G is a pair $\widehat{\mathcal{T}} = (T, \{(X_t, U_t)_{t \in V(T)}\})$ such that $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ is a tree decomposition of G , and for every $t \in V(T)$ we have $U_t \subseteq X_t$ and $|U_t| \leq \ell$. We will refer to \mathcal{T} as the *underlying tree decomposition* of $\widehat{\mathcal{T}}$. The *width* of $\widehat{\mathcal{T}}$, denoted by $\text{width}(\widehat{\mathcal{T}})$, is defined as the width of \mathcal{T} . More generally, any concept defined for tree decompositions can be naturally extended to ℓ -refined tree decompositions, simply by considering it on the underlying tree decomposition.

Definition 4.2. Given a nonnegative integer ℓ , the *residual independence number* of an ℓ -refined tree decomposition $\widehat{\mathcal{T}}$ of a graph G is defined as $\max_{t \in V(T)} \alpha(G[X_t \setminus U_t])$ and denoted by $\widehat{\alpha}(\widehat{\mathcal{T}})$. The *ℓ -refined tree-independence number* of a graph G is defined as the minimum residual independence number of an ℓ -refined tree decomposition of G , and denoted by $\ell\text{-tree-}\alpha(G)$.

We showcase the applicability of these notions in the third paper of the series [49], by showing that such tree decompositions exist, and can be computed efficiently, for any graph excluding either a W_4 or a K_5^- as an induced minor.

The notion of an ℓ -refined tree decomposition with bounded residual independence number generalizes the notion of (k, ℓ) -semi clique tree decompositions studied by Jacob et al. in [71]. Given two nonnegative integers k and ℓ , a (k, ℓ) -semi clique tree decomposition of G is a tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ of G such that in each bag X_t there exists a subset $U_t \subseteq X_t$ and k (possibly empty) cliques C_1, \dots, C_k in G such that $|U_t| \leq \ell$ and $\cup_{i=1}^k C_i = X_t \setminus U_t$. Since the independence number of any graph that can be covered by k cliques is at most k , the ℓ -refined tree-independence number of any graph G admitting a (k, ℓ) -semi clique tree decomposition is at most k .

Let us say that a graph $G = (V, E)$ is ℓ -vertex-almost chordal if G contains a set $S \subseteq V$ of at most ℓ vertices such that $G - S$ is a chordal graph. Note that it follows from Theorem 3.3 that for any nonnegative integer ℓ , any ℓ -vertex-almost chordal graph admits an ℓ -refined tree decomposition with residual independence number at most 1 (or, equivalently, a $(1, \ell)$ -semi clique tree decomposition). Jacob et al. showed in [71] that for any fixed ℓ and any ℓ -vertex-almost chordal graph G a $(4, 7\ell + 5)$ -semi clique tree decomposition of G can be computed in polynomial time.

A similar but different distance to chordality can be obtained by considering edge additions. Let us say that a graph $G = (V, E)$ is ℓ -edge-almost chordal if G can be turned into a chordal graph by adding at most ℓ edges. In this respect, Fomin and Golovach defined the concept of an ℓ -edge-almost chordal tree decomposition of the graph as a tree decomposition where each bag can be made a clique by adding at most ℓ edges [60] (they called such tree decompositions ℓ -almost chordal tree decompositions). It follows from the definitions that every ℓ -edge-almost chordal graph admits an ℓ -edge-almost chordal tree decomposition, and that the ℓ -refined tree-independence number of any graph G admitting an ℓ -edge-almost chordal tree decomposition is at most 1. Given an ℓ -almost chordal tree decomposition, Fomin and Golovach construct dynamic programming algorithms that are subexponential in ℓ for various problems. In particular, they showed that for any fixed ℓ and any ℓ -edge-almost chordal graph G , an ℓ -almost chordal tree decomposition of G can be computed in polynomial time.

Note that each ℓ -refined tree decomposition $\widehat{\mathcal{T}} = (T, \{(X_t, U_t)_{t \in V(T)}\})$ of a graph G is also an $(\ell + 1)$ -refined tree decomposition, and hence $(\ell + 1)\text{-tree-}\alpha(G) \leq \ell\text{-tree-}\alpha(G)$ for all $\ell \geq 0$ and all graphs G . Furthermore, for each $\ell \geq 0$, every tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ of a graph G naturally corresponds to an ℓ -refined tree decomposition $\widehat{\mathcal{T}} = (T, \{(X_t, U_t)_{t \in V(T)}\})$ of G such that $\widehat{\alpha}(\widehat{\mathcal{T}}) = \alpha(\mathcal{T})$, obtained by setting $U_t = \emptyset$ for all $t \in V(T)$. On the other hand, for any ℓ -refined tree decomposition $\widehat{\mathcal{T}}$ of a graph G , the independence number of the underlying tree decomposition of $\widehat{\mathcal{T}}$ is at most $\widehat{\alpha}(\widehat{\mathcal{T}}) + \ell$. We thus have:

Observation 4.3. For every graph G and every integer $\ell \geq 0$, we have

$$\ell\text{-tree-}\alpha(G) \leq \text{tree-}\alpha(G) \leq \ell\text{-tree-}\alpha(G) + \ell.$$

In particular, equalities hold when $\ell = 0$, that is, $0\text{-tree-}\alpha(G) = \text{tree-}\alpha(G)$.

Using ℓ -refined tree decompositions, we can refine Lemma 3.2 as follows.

Lemma 4.4. *For every two non-negative integers k and ℓ , the class of graphs with ℓ -refined tree-independence number at most k is (tw, ω) -bounded, with a binding function $f(p) = R(p+1, k+1) + \ell - 2$, which is a polynomial of degree k .*

Proof. Recall that $R(p, q) \leq \binom{p+q-2}{q-1}$ for all positive integers p and q , which, for fixed q , is a polynomial in p of degree $q - 1$.

Let us now fix $p \in \mathbb{Z}_+$ and let G be a graph such that $\omega(G) = p$ and ℓ -tree- $\alpha(G) \leq k$. Fix an ℓ -refined tree decomposition $\widehat{\mathcal{T}} = (T, \{(X_t, U_t)_{t \in V(T)}\})$ of G with residual independence number at most k . For every node $t \in V(T)$, the set $X_t \setminus U_t$ induces a subgraph of G with independence number at most k and clique number at most p . Thus, for every bag X_t of $\widehat{\mathcal{T}}$, Ramsey's theorem implies that $|X_t| \leq |U_t| + R(p+1, k+1) - 1 \leq R(p+1, k+1) + \ell - 1$. It follows that $\text{tw}(G) \leq R(p+1, k+1) + \ell - 2$, as claimed. \square

By Theorem 3.3, a graph G has a 0-refined tree decomposition with (residual) independence number at most 1 if and only if G is chordal. In this case Theorem 2.4 implies that $\text{tw}(G) = \omega(G) - 1$. This result generalizes to graphs admitting an ℓ -refined tree decomposition with residual independence number at most 1, for a nonnegative integer ℓ . Such a tree decomposition yields an additive approximation for treewidth to within ℓ .

Proposition 4.5. *Let $\ell \in \mathbb{Z}_+$ and let G be a graph admitting an ℓ -refined tree decomposition $\widehat{\mathcal{T}}$ with residual independence number at most 1. Then $\text{width}(\widehat{\mathcal{T}}) \leq \text{tw}(G) + \ell$ and $\text{tw}(G) \leq \omega(G) - 1 + \ell$.*

Proof. Let us write $\widehat{\mathcal{T}} = (T, \{(X_t, U_t)_{t \in V(T)}\})$ and let X_t be a largest bag of the underlying tree decomposition. Since $X_t \setminus U_t$ induces a subgraph with independence number at most one, it is a clique in G . We thus obtain, using also Corollary 2.2, that

$$\text{width}(\widehat{\mathcal{T}}) = |X_t| - 1 \leq |U_t| + |X_t \setminus U_t| - 1 \leq \ell + \omega(G) - 1 \leq \ell + \text{tw}(G). \quad \square$$

A result analogous to Proposition 3.10 (and with the same proof) also holds for ℓ -refined tree decompositions. This result is used in the third paper of the series [49].

Proposition 4.6. *Let C be a clique cutset in a graph G and let (A, B, C) be a cut-partition of G . Let $G_A = G[A \cup C]$ and $G_B = G[B \cup C]$, and let $\widehat{\mathcal{T}}_A$ and $\widehat{\mathcal{T}}_B$ be ℓ -refined tree decompositions of G_A and G_B , respectively. Then we can compute in time $\mathcal{O}(|\widehat{\mathcal{T}}_A| + |\widehat{\mathcal{T}}_B|)$ an ℓ -refined tree decomposition $\widehat{\mathcal{T}}$ of G such that $\widehat{\alpha}(\widehat{\mathcal{T}}) = \max\{\widehat{\alpha}(\widehat{\mathcal{T}}_A), \widehat{\alpha}(\widehat{\mathcal{T}}_B)\}$.*

5 Weighted independent sets

In this section, we prove that every graph class with bounded and efficiently witnessed tree-independence number admits a polynomial-time algorithm for the MAX WEIGHT INDEPENDENT SET problem.

A tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ of a graph G is said to be *rooted* if we distinguish one node r of T which we take as the root of T . This introduces natural parent-child and ancestor-descendant relations in the tree T . A *leaf* of T is a node with no children. Following [46], we say that a tree decomposition $(T, \{X_t\}_{t \in V(T)})$ is *nice* if it is rooted and the following conditions are satisfied:

- (a) If $t \in V(T)$ is the root or a leaf of T , then $X_t = \emptyset$;
- (b) Every non-leaf node t of T is one of the following three types:

- **Introduce node:** a node t with exactly one child t' such that $X_t = X_{t'} \cup \{v\}$ for some vertex $v \in V(G) \setminus X_{t'}$;
- **Forget node:** a node t with exactly one child t' such that $X_t = X_{t'} \setminus \{v\}$ for some vertex $v \in X_{t'}$;
- **Join node:** a node t with exactly two children t_1 and t_2 such that $X_t = X_{t_1} = X_{t_2}$.

We assume that introduce and forget nodes are also labeled with the vertex v which is introduced or forgotten.

It is well known that, given a graph G and a tree decomposition \mathcal{T} of G with width at most k , one can compute a nice tree decomposition of G with width at most k in polynomial time (see, e.g., [46]). The standard approach for doing so can in fact be used to prove the following lemma.

Lemma 5.1. *Given an ℓ -refined tree decomposition $\widehat{\mathcal{T}} = (T, \{(X_t, U_t)\}_{t \in V(T)})$ with width k of a graph G , one can compute in time $\mathcal{O}(k^2 \cdot |V(T)|)$ a nice ℓ -refined tree decomposition $\widehat{\mathcal{T}}' = (T', \{(X_{t'}, U_{t'})\}_{t' \in V(T')})$ of G that has at most $\mathcal{O}(k \cdot |V(T)|)$ nodes and such that for every node $t' \in V(T')$ there exists a node $t \in V(T)$ such that $X_{t'} \subseteq X_t$ and $U_{t'} = U_t \cap X_{t'}$. In particular, $\widehat{\alpha}(\widehat{\mathcal{T}}') \leq \widehat{\alpha}(\widehat{\mathcal{T}})$.*

Proof. Consider the following algorithm.

1. We traverse the tree T and check for every two adjacent nodes t and t' in T if the bags X_t and $X_{t'}$ are comparable (i.e., if $X_t \subseteq X_{t'}$ or $X_{t'} \subseteq X_t$). If, say, $X_t \subseteq X_{t'}$, then we contract the edge tt' and label the resulting node with the pair $(X_{t'}, U_{t'})$. Hence, we now assume that no two adjacent nodes of T have comparable bags.
2. We choose a node of T with degree at most one as its root r and compute the corresponding parent-child relationship in T .
3. We assure that each node has at most two children, by replacing each node t of T with $d \geq 3$ children c_1, \dots, c_d with a path consisting of d nodes t_1, \dots, t_d , each associated with the same pair (X_t, U_t) , making t_1 a child of the parent of t , and for all $j \in \{1, \dots, d\}$, making c_j a child of t_j (note that if $j < d$, then t_{j+1} is also a child of t_j).
4. For every node t of T with exactly two children t_1 and t_2 , we label t as a join node. For $i \in \{1, 2\}$, if $X_t \neq X_{t_i}$, then we subdivide the edge tt_i and associate the pair (X_t, U_t) to the new node.
5. For each leaf node t of T , we add a new node t' associated with the pair (\emptyset, \emptyset) and make t' a child of t .
6. We add a new node r' associated with the pair (\emptyset, \emptyset) and make r a child of r' (that is, r' becomes the new root).
7. For every node t of T that is not already labeled as a join node there is a unique child t' of t . We replace the edge $t't$ with a path $(t' = t_0, \dots, t_p, t_{p+1}, \dots, t_{p+q} = t)$ of length $p + q$ where $p = |X_{t'} \setminus X_t|$ and $q = |X_t \setminus X_{t'}|$. For all $i \in \{1, \dots, p\}$, the node t_i is a forget node labeled with (X_{t_i}, U_{t_i}) where X_{t_i} forgets one vertex from $X_{t_{i-1}} \setminus X_t$ and $U_{t_i} = U_{t'} \cap X_{t_i}$. Similarly, for all $j \in \{1, \dots, q-1\}$, the node t_{p+j} is an introduce node labeled with $(X_{t_{p+j}}, U_{t_{p+j}})$ where $X_{t_{p+j}}$ introduces one vertex from $X_t \setminus X_{t_{p+j-1}}$, and $U_{t_{p+j}} = U_t \cap X_{t_{p+j}}$. The last node of the path is the node t , which is an introduce node.

Note that each of the above steps modifies an ℓ -refined tree decomposition into another one. Let us denote by $\widehat{T}' = (T', \{(X_{t'}, U_{t'})\}_{t' \in V(T')}$ the final ℓ -refined tree decomposition. By construction, every non-leaf node of T' has a unique label (an introduce node, a forget node, or a join node) and \widehat{T}' is a nice ℓ -refined tree decomposition of G such that for every node t' of T' , there exists a node t in T such that $X_{t'} \subseteq X_t$ and $U_{t'} = U_t \cap X_{t'}$. Consider a node t' of T' such that $\widehat{\alpha}(\widehat{T}') = \alpha(G[X_{t'} \setminus U_{t'}])$ and let t be a node of T such that $X_{t'} \subseteq X_t$ and $U_{t'} = U_t \cap X_{t'}$. Then, $X_{t'} \setminus U_{t'} = X_{t'} \setminus (U_t \cap X_{t'}) = X_{t'} \setminus U_t \subseteq X_t \setminus U_t$ and hence $\widehat{\alpha}(\widehat{T}') = \alpha(G[X_{t'} \setminus U_{t'}]) \leq \alpha(G[X_t \setminus U_t]) \leq \widehat{\alpha}(\widehat{T})$.

We now reason about the complexity of obtaining \widehat{T}' by considering the complexity of each aforementioned step. Step 1 takes $\mathcal{O}(k^2)$ time for every edge of T , and thus $\mathcal{O}(k^2 \cdot |V(T)|)$ overall; the resulting tree T' has $\mathcal{O}(|V(T)|)$ nodes. Step 2 can be done in time $\mathcal{O}(|V(T)|)$. Step 3 can be done in time $\mathcal{O}(|V(T)|)$ and results in a tree T' with $\mathcal{O}(|V(T)|)$ nodes. Step 4 can be done in time $\mathcal{O}(k^2 \cdot |V(T)|)$ and Step 5 in time $\mathcal{O}(|V(T)|)$; both steps yield a tree with $\mathcal{O}(|V(T)|)$ nodes. Step 6 can be done in constant time. Finally, Step 7 can be done in time $\mathcal{O}(k^2 \cdot |V(T)|)$ and results in an ℓ -refined tree decomposition with $\mathcal{O}(k \cdot |V(T)|)$ nodes. \square

In the next theorem, we adapt the standard dynamic programming approach (see, e.g., [46]) for solving the MAX WEIGHT INDEPENDENT SET problem in graphs of bounded treewidth to graphs of bounded ℓ -refined tree independence number, for some integer $\ell \geq 0$. In fact, the algorithm runs in polynomial time even if $\ell = \mathcal{O}(\log |V(G)|)$, which is the case, for example, for graphs with logarithmic treewidth (see, e.g., [1, 15]).

Theorem 5.2. *For every integer $k \geq 1$, MAX WEIGHT INDEPENDENT SET is solvable in time $\mathcal{O}(2^\ell \cdot |V(G)|^{k+1} \cdot |V(T)|)$ if the input vertex-weighted graph G is given with an ℓ -refined tree decomposition $\widehat{T} = (T, \{(X_t, U_t)\}_{t \in V(T)})$ with residual independence number at most k .*

Proof. Let $n = |V(G)|$ and $w : V(G) \rightarrow \mathbb{Q}_+$ be the weight function. We first apply Lemma 5.1 and compute in time $\mathcal{O}(n^2 \cdot |V(T)|)$ a nice ℓ -refined tree decomposition $\widehat{T}' = (T', \{(X_{t'}, U_{t'})\}_{t' \in V(T')}$ of G with $\mathcal{O}(n \cdot |V(T)|)$ bags and such that $\widehat{\alpha}(\widehat{T}') \leq \widehat{\alpha}(\widehat{T}) \leq k$.

Recall that, by definition, \widehat{T}' is rooted at some node r of T' . For every node $t \in V(T')$, we denote by V_t the union of all bags $X_{t'}$ such that $t' \in V(T')$ is a (not necessarily proper) descendant of t in T' .

For each node $t \in V(T')$, we compute the family \mathcal{S}_t of all sets $S \subseteq X_t$ that are independent in G . Note that each set $S \in \mathcal{S}_t$ is the disjoint union of sets S_1 and S_2 where $S_1 = S \cap U_t$ and $S_2 = S \cap (X_t \setminus U_t)$. Since \widehat{T}' is an ℓ -refined tree decomposition with residual independence number at most k , we have that $|S_1| \leq |U_t| \leq \ell$ and $|S_2| \leq \alpha(G[X_t \setminus U_t]) \leq k$. It follows that the family \mathcal{S}_t can be computed in time $\mathcal{O}(2^{|U_t|} \cdot |X_t \setminus U_t|^k) = \mathcal{O}(2^\ell \cdot n^k)$ by enumerating all $\mathcal{O}(2^\ell)$ candidate sets for S_1 , all $\mathcal{O}(n^k)$ candidate sets for S_2 , and verifying if the union $S_1 \cup S_2$ is independent in G .¹ We traverse the tree T' bottom-up and use a dynamic programming approach to compute, for every node $t \in V(T')$ and every set $S \in \mathcal{S}_t$, the value of $c[t, S]$, defined as the maximum weight of an independent set I in the graph $G[V_t]$ such that $I \cap X_t = S$.

Since \widehat{T}' is nice, we have $X_r = \emptyset$; in particular, the only independent set S with $S \subseteq X_r$ is the empty set. Furthermore, $V_r = V(G)$, and hence $c[r, \emptyset]$ corresponds to the maximum weight of an independent set in G , which is what we want to compute.

We consider various cases depending on the type of a node $t \in V(T')$. For each type we give a formula for computing the value $c[t, S]$ for all $S \in \mathcal{S}_t$ from the already computed values of $c[t', S']$ where t' is a child of t in T' and $S' \in \mathcal{S}_{t'}$.

¹We can check in constant time if a set S with $|S| \leq k + \ell$ is independent in G , since $k + \ell$ is constant and we assume that G is represented with an adjacency matrix—if not, we can first compute such a representation from the adjacency lists in time $\mathcal{O}(n^2)$.

Leaf node. By the definition of a nice tree decomposition it follows that $X_t = \emptyset$. Thus, we have $\mathcal{S}_t = \{\emptyset\}$, $V_t = \emptyset$, and $c[t, \emptyset] = 0$.

Introduce node. By definition, t has exactly one child t' and $X_t = X_{t'} \cup \{v\}$ holds for some vertex $v \in V(G) \setminus X_{t'}$. For an arbitrary set $S \in \mathcal{S}_t$, we have

$$c[t, S] = \begin{cases} c[t', S] & \text{if } v \notin S, \\ c[t', S \setminus \{v\}] + w(v) & \text{otherwise.} \end{cases}$$

Forget node. By definition, t has exactly one child t' in T' and $X_t = X_{t'} \setminus \{v\}$ holds for some vertex $v \in X_{t'}$. Note that for a set $S \in \mathcal{S}_t$, the set $S \cup \{v\}$ belongs to $\mathcal{S}_{t'}$ if and only if it is independent in G , that is, if no vertex in S is adjacent to v . For an arbitrary set $S \in \mathcal{S}_t$, we have

$$c[t, S] = \begin{cases} \max\{c[t', S], c[t', S \cup \{v\}]\} & \text{if no vertex in } S \text{ is adjacent to } v, \\ c[t', S] & \text{otherwise.} \end{cases}$$

Join node. By definition, t has exactly two children t_1 and t_2 in T' and it holds that $X_t = X_{t_1} = X_{t_2}$. For an arbitrary set $S \in \mathcal{S}_t$, we have

$$c[t, S] = c[t_1, S] + c[t_2, S] - w(S).$$

The only way in which our algorithm differs from the standard one (see [46]) is that we compute $c[t, S]$ only for sets S in the family \mathcal{S}_t , and not for all subsets of the bag X_t . We therefore omit the description of the recurrence relations leading to the dynamic programming algorithm and the proof of correctness.

It remains to estimate the time complexity. We need time $\mathcal{O}(n^2 \cdot |V(T)|)$ to compute $\widehat{\mathcal{T}}'$. At each of the $\mathcal{O}(n \cdot |V(T)|)$ nodes $t \in V(T')$, we perform a constant-time computation for each set $S \in \mathcal{S}_t$, resulting in an overall time complexity of $\mathcal{O}(2^\ell \cdot n^k)$ per node. Thus, the total time complexity of the algorithm is $\mathcal{O}(n^2 \cdot |V(T)|) + \mathcal{O}(n \cdot |V(T)| \cdot 2^\ell \cdot n^k) = \mathcal{O}(2^\ell \cdot n^{k+1} \cdot |V(T)|)$, as claimed. \square

Since every tree decomposition \mathcal{T} naturally corresponds to 0-refined tree decomposition $\widehat{\mathcal{T}}$ with the same independence number (more precisely, with $\widehat{\alpha}(\widehat{\mathcal{T}}) = \alpha(\widehat{\mathcal{T}}) = \alpha(\mathcal{T})$), Theorem 5.2 implies the following.

Corollary 5.3. *For every $k \geq 1$, MAX WEIGHT INDEPENDENT SET is solvable in time $\mathcal{O}(|V(G)|^{k+1} \cdot |V(T)|)$ if the input vertex-weighted graph G is given with a tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ with independence number at most k .*

Corollary 5.3 generalizes a result of Krause [77] stating that the MAX INDEPENDENT SET problem is solvable in polynomial time on intersection graphs of connected subgraphs of graphs of bounded treewidth (assuming that the input graph is given with an intersection model). Recall that every graph class with bounded intersection treewidth has bounded tree-independence number (cf. Theorem 3.12). More precisely, consider a fixed nonnegative integer k , a graph H with treewidth at most k , and connected subgraphs H_1, \dots, H_p of H . A tree decomposition \mathcal{T} of H with width k can be computed in linear time using, e.g., the algorithm of Bodlaender [11]. Then, as shown by Bodlaender, Gustedt, and Telle in the proof of [10, Lemma 2.4], the tree decomposition \mathcal{T} can be transformed efficiently into a tree decomposition \mathcal{T}' of the intersection graph G of H_1, \dots, H_p such that each bag of \mathcal{T}' is the union of at most $k + 1$ cliques in G . In particular, \mathcal{T}' has independence number at most $k + 1$ and thus Corollary 5.3 applies, implying the result of Krause.

In the third paper of the series [49], we present two particular examples of where the complexity of solving the MAXIMUM WEIGHT INDEPENDENT SET problem is significantly improved when using the result of Theorem 5.2 as compared to using Corollary 5.3.

6 Cameron–Hell construction

Given a graph G and a family $\mathcal{H} = \{H_j\}_{j \in J}$ of connected subgraphs of G , we denote by $G(\mathcal{H})$ the graph with vertex set J , in which two distinct elements $i, j \in J$ are adjacent if and only if H_i and H_j either have a vertex in common or there is an edge in G connecting them. This construction was considered by Cameron and Hell in [27], who focused on the following particular case. Given a graph G and a (finite or infinite) set \mathcal{F} of connected graphs, we denote by $\mathcal{H}(G, \mathcal{F})$ the family of all subgraphs of G isomorphic to a member of \mathcal{F} . In particular:

- for $\mathcal{H} = \mathcal{H}(G, \{K_1\})$, the construction of the graph $G(\mathcal{H})$ is trivial, with $G(\mathcal{H}) \cong G$, and
- for $\mathcal{H} = \mathcal{H}(G, \{K_2\})$, the graph $G(\mathcal{H})$ is isomorphic to the square of the line graph of G .

A construction similar to $G(\mathcal{H})$ was studied by Duchet [55]. More recently, Gartland et al. [65] considered the special case when \mathcal{H} consists of all connected induced subgraphs of G , referring in this case to the derived graph $G(\mathcal{H})$ as the *blob graph* of G .

Cameron and Hell proved in [27] that for any chordal graph G , any set \mathcal{F} of connected graphs, and $\mathcal{H} = \mathcal{H}(G, \mathcal{F})$, the graph $G(\mathcal{H})$ is chordal, generalizing an analogous result due to Cameron in the case when $\mathcal{F} = \{K_2\}$ [26]. Using Theorem 3.3, the result of Cameron and Hell states that $\text{tree-}\alpha(G(\mathcal{H})) \leq 1$ whenever $\text{tree-}\alpha(G) \leq 1$. We now generalize this result by showing that mapping any graph G to the graph $G(\mathcal{H})$, where \mathcal{H} is an arbitrary collection of nonnull connected subgraphs of G , cannot increase the tree-independence number. In particular, this shows that for any graph class \mathcal{G} with bounded tree-independence number and any set \mathcal{F} of connected nonnull graphs, the class $\{G(\mathcal{H}) : G \in \mathcal{G}, \mathcal{H} = \mathcal{H}(G, \mathcal{F})\}$ also has bounded tree-independence number.

Lemma 6.1. *Let G be a graph, let $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ be a tree decomposition of G , and let $\mathcal{H} = \{H_j\}_{j \in J}$ be a finite family of connected nonnull subgraphs of G . Then $\mathcal{T}' = (T, \{X'_t\}_{t \in V(T)})$ with $X'_t = \{j \in J : V(H_j) \cap X_t \neq \emptyset\}$ for all $t \in V(T)$ is a tree decomposition of $G(\mathcal{H})$ such that $\alpha(\mathcal{T}') \leq \alpha(\mathcal{T})$.*

Proof. Let us first show that \mathcal{T}' is a tree decomposition of $G(\mathcal{H})$. First, note that since $V(G(\mathcal{H})) = J$, for each $t \in V(T)$ the set X'_t is indeed a subset of $V(G(\mathcal{H}))$.

Let $j \in J$ be a vertex of $G(\mathcal{H})$. Fix a vertex $v \in V(H_j)$ and consider any bag X_t of \mathcal{T} such that $v \in X_t$. Then $v \in V(H_j) \cap X_t$ and hence $j \in X'_t$. Thus, every vertex of $G(\mathcal{H})$ belongs to a bag of \mathcal{T}' .

Let $\{i, j\}$ be an edge of $G(\mathcal{H})$. Assume first that the subgraphs H_i and H_j have a vertex in common, say v . Since \mathcal{T} is a tree decomposition of G , there exists some $t \in V(T)$ such that $v \in X_t$. The fact that $v \in V(H_i) \cap X_t$ implies that $i \in X'_t$, and similarly, $j \in X'_t$. Assume now that there exist vertices $u \in V(H_i)$ and $v \in V(H_j)$ such that $uv \in E(G)$. Since \mathcal{T} is a tree decomposition of G , there exists some $t \in V(T)$ such that $\{u, v\} \subseteq X_t$. The fact that $u \in V(H_i) \cap X_t$ implies that $i \in X'_t$. Similarly, the fact that $v \in V(H_j) \cap X_t$ implies that $j \in X'_t$. Hence, for every edge of $G(\mathcal{H})$ there exists a bag of \mathcal{T}' containing both endpoints of the edge.

Next, consider an arbitrary vertex $j \in J$ of $G(\mathcal{H})$. We need to show that the set of nodes $t \in V(T)$ such that $j \in X'_t$ induces a connected subgraph of T . Let us denote for each vertex $v \in V(H_j)$ by T_v the subgraph of T induced by the nodes $t \in V(T)$ such that $v \in X_t$. Since \mathcal{T} is a tree decomposition of G , each T_v is a connected subgraph of T , that is, a subtree. For a node $t \in V(T)$, the condition $j \in X'_t$ is equivalent to the condition $V(H_j) \cap X_t \neq \emptyset$, that is, there exists a vertex $v \in V(H_j)$ such that $v \in X_t$. Therefore, $j \in X'_t$ if and only if there exists a vertex $v \in V(H_j)$ such that t belongs to the tree T_v . It thus suffices to show that the union T_j of the trees T_v over all vertices $v \in V(H_j)$ forms a connected graph. Suppose for a contradiction that T_j

is not connected and fix a component C of T_j . Let us denote by U the set of vertices $u \in V(H_j)$ such that $V(T_u) \subseteq V(C)$. Since $V(C) \neq V(T_j)$, we have $U \neq V(H_j)$. By the connectedness of H_j , there is an edge $uv \in E(H_j)$ such that $u \in U$ and $v \in V(H_j) \setminus U$. Let t be a node of T such that $\{u, v\} \subseteq X_t$. The trees T_u and T_v both contain node t , and hence the connected component C of T_j contains both T_u and T_v . This implies that $v \in U$, a contradiction.

It remains to show that $\alpha(\mathcal{T}) \geq \alpha(\mathcal{T}')$. Let t be a node of T maximizing the independence number of the subgraph of $G(\mathcal{H})$ induced by X'_t . Let k be this independence number and let $I \subseteq J$ be an independent set of cardinality k in the subgraph of $G(\mathcal{H})$ induced by X'_t . Then for any two distinct elements $i, j \in I$ the graphs H_i and H_j are vertex-disjoint subgraphs in \mathcal{H} such that no edge of G has one endpoint in H_i and the other one in H_j . Each element $i \in I$ belongs to the bag X'_t , which implies that $V(H_i) \cap X_t \neq \emptyset$, and hence there exists a vertex u_i in $V(H_i) \cap X_t$. Since any two distinct vertices u_i and u_j belong to the subgraphs H_i and H_j of G , which are vertex-disjoint and with no edges between them, the set $\{u_i : i \in I\}$ is an independent set of cardinality k in the subgraph of G induced by X_t . This shows that $\alpha(\mathcal{T}) \geq k = \alpha(\mathcal{T}')$, as claimed. \square

Lemma 6.1 immediately implies the following.

Theorem 6.2. *Let G be a graph and let \mathcal{H} be a finite family of connected nonnull subgraphs of G . Then $\text{tree-}\alpha(G(\mathcal{H})) \leq \text{tree-}\alpha(G)$.*

Note in particular that Theorem 6.2 implies that for any graph G , the tree-independence number of its blob graph is bounded by $\text{tree-}\alpha(G)$.

We will also make use of an algorithmic version of Lemma 6.1. In the analysis of the time complexity we will make use of the following standard lemma (see, e.g., [67] for a similar argument for sorting adjacency lists of graphs).

Lemma 6.3. *Let $V = \{v_1, \dots, v_n\}$ be a set and let $\mathcal{S} = \{S_i\}_{i \in I}$ be a finite family of subsets of V . Then there exists an algorithm running in time $\mathcal{O}(|V| + |I| + \sum_{i \in I} |S_i|)$ that sorts each set S_i with respect to the ordering v_1, \dots, v_n of V .*

Proof. Let B be the bipartite incidence graph of the family \mathcal{S} , that is, B has vertex set $V \cup I$, and edge set $\{\{v, i\} : v \in V, i \in I, v \in S_i\}$. We can compute the adjacency lists of the graph B in time $\mathcal{O}(|V| + |I| + \sum_{i \in I} |S_i|)$ as follows. We fix an ordering of the set I , say $I = \{i_1, \dots, i_m\}$. For each $i \in I$, the set S_i already gives the adjacency list of i . We initialize the adjacency lists for each $v \in V$ to the empty lists. For all $j = 1, \dots, m$, we iterate over the elements v of S_{i_j} and add i_j to the end of the adjacency list of vertex v . We now have the adjacency lists of all vertices of B ; those for $v \in V$ are already sorted, while those for $i \in I$ need not be.

To sort the adjacency lists for vertices $i \in I$, we iterate over the adjacency lists of vertices $v \in V$ in a similar way as we did above for $i \in I$. We reset the adjacency lists for all $i \in I$ to the empty lists. For all $i = 1, \dots, n$, we iterate over the elements $j \in I$ of the adjacency list of v_i and add v_i to the end of the adjacency list for vertex $j \in I$. At the end of this procedure, the adjacency list of each $i \in I$ will contain exactly the elements of S_i , sorted with respect to the ordering v_1, \dots, v_n of V . The total time complexity of the procedure is proportional to the number of vertices and edges of the graph B , that is, $\mathcal{O}(|V| + |I| + \sum_{i \in I} |S_i|)$. \square

Corollary 6.4. *There exists an algorithm that takes as input a graph G , a finite family $\mathcal{H} = \{H_j\}_{j \in J}$ of connected nonnull subgraphs of G , and a tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ of G , and computes in time $\mathcal{O}(|J| \cdot (|J| + |V(T)|) \cdot |V(G)| + |E(G)|)$ the graph $G(\mathcal{H})$ and a tree decomposition $\mathcal{T}' = (T, \{X'_t\}_{t \in V(T)})$ of $G(\mathcal{H})$ such that $\alpha(\mathcal{T}') \leq \alpha(\mathcal{T})$.*

Proof. Fix an arbitrary ordering of the vertex set of G . Using Lemma 6.3, we first sort the vertex set of each of the graphs H_j , $j \in J$, as well as each bag X_t , $t \in V(T)$, with respect to the fixed ordering of $V(G)$, in time

$$\mathcal{O}\left(|V(G)| + |J| + \sum_{j \in J} |V(H_j)|\right) + \mathcal{O}\left(|V(G)| + |V(T)| + \sum_{t \in V(T)} |X_t|\right) = \mathcal{O}\left((|J| + |V(T)|) \cdot |V(G)|\right).$$

Note that using this sortedness assumption, we can compute the union and the intersection of any two sorted sets $X, Y \subseteq V(G)$ in time $\mathcal{O}(|V(G)|)$. To compute the graph $G(\mathcal{H})$, we only need to explain how to compute its edge set, since the vertex set is J . For each $j \in J$, we perform a BFS traversal up to distance two from a new vertex v_j added to G which we make adjacent to all the vertices of H_j . Let R_j be the set of vertices of G reached this way. Then, for all $i \in J \setminus \{j\}$, the graph H_i is adjacent to H_j in $G(\mathcal{H})$ if and only if at least one vertex of H_i belongs to R_j . This can be tested in time $\mathcal{O}(|V(G)|)$ by first sorting the set R_j and then computing the intersection $V(H_i) \cap R_j$. Hence, this procedure over all $j \in J$ can be carried out in time $\mathcal{O}(|J| \cdot (|V(G)| + |E(G)|)) + \mathcal{O}(|J|^2 \cdot |V(G)|) = \mathcal{O}(|J|^2 \cdot |V(G)| + |J| \cdot |E(G)|)$.

To compute \mathcal{T}' , we need to compute for each $t \in V(T)$ the bag X'_t consisting of all vertices $j \in J$ of $G(\mathcal{H})$ such that $V(H_j) \cap X_t \neq \emptyset$. All the intersections $V(H_j) \cap X_t$ can be computed in time $\mathcal{O}(|J| \cdot |V(T)| \cdot |V(G)|)$. Thus, the total time complexity of the algorithm is $\mathcal{O}(|J|^2 \cdot |V(G)| + |J| \cdot |E(G)| + |J| \cdot |V(T)| \cdot |V(G)|) = \mathcal{O}(|J| \cdot (|J| + |V(T)|) \cdot |V(G)| + |E(G)|)$. \square

7 Weighted independent packings

Consider again a graph G and a family $\mathcal{H} = \{H_j\}_{j \in J}$ of connected subgraphs of G . A subfamily \mathcal{H}' of \mathcal{H} is said to be an *independent \mathcal{H} -packing* in G if every two graphs in \mathcal{H}' are vertex-disjoint and there is no edge between them, that is, \mathcal{H}' is an independent set in the graph $G(\mathcal{H})$. Assume now that the subgraphs in \mathcal{H} are equipped with a weight function $w : J \rightarrow \mathbb{Q}_+$ assigning weight w_j to each subgraph H_j . For any set $I \subseteq J$, we define the *weight* of the family $\mathcal{H}' = \{H_i\}_{i \in I}$ as the sum $\sum_{i \in I} w_i$. Given a graph G , a finite family $\mathcal{H} = \{H_j\}_{j \in J}$ of connected nonnull subgraphs of G , and a weight function $w : J \rightarrow \mathbb{Q}_+$ on the subgraphs in \mathcal{H} , the MAX WEIGHT INDEPENDENT PACKING problem asks to find an independent \mathcal{H} -packing in G of maximum weight. In the special case when \mathcal{F} is a fixed finite family of connected graphs and $\mathcal{H} = \mathcal{H}(G, \mathcal{F})$ is the set of all subgraphs of G isomorphic to a member of \mathcal{F} , we obtain the MAX WEIGHT INDEPENDENT \mathcal{F} -PACKING problem. This is a common generalization of several problems studied in the literature, including:

- the INDEPENDENT \mathcal{F} -PACKING problem (see [27]), which corresponds to the unweighted case,
- the MAX WEIGHT INDEPENDENT SET problem, which corresponds to the case $\mathcal{F} = \{K_1\}$,
- the MAX WEIGHT INDUCED MATCHING problem (see, e.g., [3, 82]), which corresponds to the case $\mathcal{F} = \{K_2\}$,
- the DISSOCIATION SET problem (see, e.g., [80, 100, 101]), which corresponds to the case when $\mathcal{F} = \{K_1, K_2\}$ and the weight function assigns to each subgraph H_j , $j \in J$, the weight equal to $|V(H_j)|$, and
- the k -SEPARATOR problem (see, e.g., [7, 78]), which corresponds to the case when \mathcal{F} contains all connected graphs with at most k vertices, the graph G is equipped with a vertex weight function $w : V(G) \rightarrow \mathbb{Q}_+$, and the weight function on \mathcal{H} assigns to each subgraph H_j , $j \in J$, the weight equal to $\sum_{x \in V(H_j)} w(x)$.

The MAX WEIGHT INDEPENDENT PACKING problem can be reduced in polynomial time to the MAX WEIGHT INDEPENDENT SET problem, by means of the following.

Observation 7.1. *Let G be a graph, let $\mathcal{H} = \{H_j\}_{j \in J}$ be a finite family of connected nonnull subgraphs of G , and let $w : J \rightarrow \mathbb{Q}_+$ be a weight function on the subgraphs in \mathcal{H} . Let I be an independent set in $G(\mathcal{H})$ of maximum weight with respect to the weight function w . Then I is an independent \mathcal{H} -packing in G of maximum weight.*

Using Corollaries 5.3 and 6.4 and Observation 7.1, we can now obtain an analogous result to Corollary 5.3 for the MAX WEIGHT INDEPENDENT PACKING problem.

Theorem 7.2. *Let k be a positive integer. Then, given a graph G and a finite family $\mathcal{H} = \{H_j\}_{j \in J}$ of connected nonnull subgraphs of G , the MAX WEIGHT INDEPENDENT PACKING problem can be solved in time $\mathcal{O}(|J| \cdot (|J| + |V(T)|) \cdot |V(G)| + |E(G)| + |J|^k \cdot |V(T)|)$ if G is given together with a tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ with independence number at most k .*

Proof. Let G be a graph, let $\mathcal{H} = \{H_j\}_{j \in J}$ be a finite family of connected nonnull subgraphs of G , let $w : J \rightarrow \mathbb{Q}_+$ be a weight function on the subgraphs in \mathcal{H} , and let $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ be a tree decomposition of G with independence number at most k . By Corollary 6.4, we can compute in time $\mathcal{O}(|J| \cdot (|J| + |V(T)|) \cdot |V(G)| + |E(G)|)$ the graph $G(\mathcal{H})$ and a tree decomposition $\mathcal{T}' = (T, \{X'_t\}_{t \in V(T)})$ of $G(\mathcal{H})$ with independence number at most k . Using Corollary 5.3, we now compute in time $\mathcal{O}(|J|^{k+1} \cdot |V(T)|)$ an independent set I in $G(\mathcal{H})$ of maximum weight with respect to the weight function w . By Observation 7.1, I is a maximum-weight independent \mathcal{H} -packing in G . The claimed running time follows. \square

The case when the subgraphs in \mathcal{H} have bounded order is of particular interest, as it generalizes the MAX WEIGHT INDEPENDENT \mathcal{F} -PACKING problem. For this case, the time complexity can be slightly improved compared to what would be obtained by a direct application of Theorem 7.2.

Theorem 7.3. *Let k and r be two positive integers. Then, given a graph G and a finite family $\mathcal{H} = \{H_j\}_{j \in J}$ of connected nonnull subgraphs of G such that $|V(H_j)| \leq r$ for all $j \in J$, the MAX WEIGHT INDEPENDENT PACKING problem can be solved in time $\mathcal{O}(|V(G)|^{r(k+1)} \cdot |V(T)|)$ if G is given together with a tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ with independence number at most k .*

Proof. We assume that G is represented with an adjacency matrix, since otherwise we can first compute such a representation from the adjacency lists in time $\mathcal{O}(|V(G)|^2)$. Note that $|V(G(\mathcal{H}))| = |J| = \mathcal{O}(|V(G)|^r)$, since by assumption each graph in \mathcal{H} has at most r vertices, and for any such set of vertices we have at most $2^{r(r-1)/2} = \mathcal{O}(1)$ choices for the edge set. We compute the edge set of $G(\mathcal{H})$ in time $\mathcal{O}(|V(G(\mathcal{H}))|^2) = \mathcal{O}(|V(G)|^{2r})$, as follows. For every two distinct $i, j \in J$, we check in time $\mathcal{O}(\max\{|V(H_i)|, |V(H_j)|\}) = \mathcal{O}(1)$ if H_i and H_j have a vertex in common. If this is the case, then we add $\{i, j\}$ to the edge set of $G(\mathcal{H})$. If this is not the case, then we check in time $\mathcal{O}(|V(H_i)| \cdot |V(H_j)|) = \mathcal{O}(1)$ if there is an edge in G connecting a vertex of H_i with a vertex of H_j . If this is the case, then we add $\{i, j\}$ to the edge set of $G(\mathcal{H})$.

For the rest of the proof, we use the same approach as in the proof of Theorem 7.2. In particular, we compute the tree decomposition \mathcal{T}' of $G(\mathcal{H})$ in time $\mathcal{O}(|J| \cdot |V(T)| \cdot |V(G)|) = \mathcal{O}(|V(G)|^{r+1} \cdot |V(T)|)$, and a maximum-weight independent set I in $G(\mathcal{H})$ in time $\mathcal{O}(|J|^{k+1} \cdot |V(T)|) = \mathcal{O}(|V(G)|^{r(k+1)} \cdot |V(T)|)$. The total time complexity of the algorithm is $\mathcal{O}(|V(G)|^{2r} + |V(G)|^{r+1} \cdot |V(T)| + |V(G)|^{r(k+1)} \cdot |V(T)|)$, which simplifies to $\mathcal{O}(|V(G)|^{r(k+1)} \cdot |V(T)|)$, as claimed. \square

Corollary 7.4. *Let \mathcal{F} be a nonempty finite set of connected nonnull graphs and let r be the maximum number of vertices of a graph in \mathcal{F} . Then, for every $k \geq 1$, the MAX WEIGHT INDEPENDENT \mathcal{F} -PACKING problem is solvable in time $\mathcal{O}\left(|V(G)|^{r(k+1)} \cdot |V(T)|\right)$ if the input graph G is given with a tree decomposition $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ with independence number at most k .*

Remark 7.5. The reader may wonder why we did not derive a result generalizing Corollary 7.4 by using the notion of ℓ -refined tree decompositions of bounded residual independence number, as we did in Theorem 5.2 for the MAX WEIGHT INDEPENDENT SET problem. The reason is that Lemma 6.1 does not seem to generalize to ℓ -refined tree decompositions, at least not so that the residual independence number would be preserved.

Consider the following example. Fix two positive integers ℓ and p and let G be the tree consisting of a vertex a adjacent to ℓ other vertices forming a set $B = \{b_1, \dots, b_\ell\}$ such that each b_i is also adjacent to p vertices of degree one, forming a set C_i . Let $\mathcal{T} = (T, \{X_t\}_{t \in V(T)})$ be a tree decomposition of G such that T is the graph $K_{1,p\ell}$, the high-degree node of T is labeled with the bag $\{a\} \cup B$, and the $p\ell$ leaves of T are labeled with bags corresponding to the edges in G containing a vertex of degree one. To make this tree decomposition ℓ -refined, we set $U_t = B$ for the high-degree node t of T and $U_t = \emptyset$ for all the other nodes. The residual independence number of this ℓ -refined tree decomposition is 1. Let $\mathcal{H} = \mathcal{H}(G, \{K_2\})$ be the family of all connected subgraphs of G of order two. Then, the graph $G(\mathcal{H})$ is isomorphic to the graph obtained from the graph $K_{1,\ell}$ by substituting a clique of size $\ell + 1$ into the vertex of degree ℓ and a clique of size p into each vertex of degree one (see, e.g., [38] for the definition of substitution). Let $\mathcal{T}' = (T, \{X'_t\}_{t \in V(T)})$ be the tree decomposition of $G(\mathcal{H})$ as defined in Lemma 6.1, that is, $X'_t = \{H \in V(G(\mathcal{H})) : V(H) \cap X_t \neq \emptyset\}$ for all $t \in V(T)$. Then, there is no way to turn \mathcal{T}' into an $f(\ell)$ -refined tree decomposition of $G(\mathcal{H})$ with residual independence number 1. Indeed, consider the bag $X_t = \{a\} \cup B$ of \mathcal{T} labeling the high-degree node of T . Since every edge of G has an endpoint in X_t , the bag of \mathcal{T}' corresponding to node t is $X'_t = V(G(\mathcal{H}))$. Using the structure of the graph $G(\mathcal{H})$, we see that the smallest subset U'_t of X'_t such that the independence number of the subgraph of $G(\mathcal{H})$ induced by $X'_t \setminus U'_t$ is 1 has size $(\ell - 1)p$, which cannot be bounded from above by any function depending only on ℓ .

8 Open questions

Our main motivation for the study of graph classes with bounded tree-independence number is the fact that for any finite family \mathcal{H} of connected graphs, the MAX WEIGHT INDEPENDENT SUBGRAPH PACKING problem is solvable in polynomial time in any class of graphs with bounded and efficiently witnessed tree-independence number (cf. Theorem 7.2). This motivates the following question.

Question 8.1. Is there a computable function $f : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ such that for every positive integer k there exists a polynomial-time algorithm that takes as input a graph G with tree-independence number at most k and computes a tree decomposition of G with independence number at most $f(k)$?

A positive answer to Question 8.1 would imply a positive answer to the following question.

Question 8.2. Is the MAX WEIGHT INDEPENDENT SET problem polynomial-time solvable in any class of graphs with bounded tree-independence number?

While we showed that the problem of computing the tree-independence number is NP-hard, it remains an open question whether the problem can be solved in polynomial time for fixed values of this parameter.

Question 8.3. For a fixed integer $k \geq 2$, what is the complexity of recognizing graphs with tree-independence number at most k ?

For every graph G , it holds that $|V(G)| \leq \alpha(G) \cdot \chi(G)$. Applying this inequality to every bag of a tree decomposition \mathcal{T} of G with $\alpha(\mathcal{T}) = \text{tree-}\alpha(G)$, we obtain that $\text{tw}(G) + 1 \leq \text{tree-}\alpha(G) \cdot \chi(G)$. Seymour [91] introduced the *tree-chromatic number* of a graph G , denoted $\text{tree-}\chi(G)$, as the smallest nonnegative integer k such that G admits a tree decomposition, each bag of which induces a k -colorable subgraph. With a similar reasoning as above, we also obtain that $\text{tw}(G) + 1 \leq \alpha(G) \cdot \text{tree-}\chi(G)$. Do these two inequalities admit the following common strengthening?

Question 8.4. Does every graph G satisfy $\text{tw}(G) + 1 \leq \text{tree-}\alpha(G) \cdot \text{tree-}\chi(G)$?

Question 8.4 has a positive answer in any class of graphs in which the tree-chromatic number coincides with the chromatic number. In particular, this is the case for any class of graphs in which the chromatic number coincides with the clique number (such as the class of perfect graphs), since in this case $\omega(G) \leq \text{tree-}\chi(G) \leq \chi(G)$, where the first inequality holds for any graph (see [91]).

Acknowledgements

This work is supported in part by the Slovenian Research Agency (I0-0035, research programs P1-0285 and P1-0383, research projects J1-3001, J1-3002, J1-3003, J1-9110, N1-0102, N1-0160, and a Young Researchers Grant).

References

- [1] T. Abrishami, M. Chudnovsky, S. Hajebi, and S. Spirkl. Induced subgraphs and tree decompositions III. Three-path-configurations and logarithmic treewidth, 2021. arXiv:2109.01310.
- [2] T. Abrishami, M. Chudnovsky, M. Pilipczuk, P. Rzażewski, and P. Seymour. Induced subgraphs of bounded treewidth and the container method. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1948–1964. [Society for Industrial and Applied Mathematics (SIAM)], Philadelphia, PA, 2021.
- [3] B. Ahat, T. Ekim, and Z. C. Taşkın. Integer programming formulations and benders decomposition for the maximum induced matching problem. *INFORMS J. Comput.*, 30(1):43–56, 2018.
- [4] S. Arnborg, J. Lagergren, and D. Seese. Easy problems for tree-decomposable graphs. *J. Algorithms*, 12(2):308–340, 1991.
- [5] S. Arnborg and A. Proskurowski. Linear time algorithms for NP-hard problems restricted to partial k -trees. *Discrete Appl. Math.*, 23(1):11–24, 1989.
- [6] R. Belmonte and M. Vatshelle. Graph classes with structured neighborhoods and algorithmic applications. *Theoret. Comput. Sci.*, 511:54–65, 2013.
- [7] W. Ben-Ameur, M.-A. Mohamed-Sidi, and J. Neto. The k -separator problem: polyhedra, complexity and approximation results. *J. Comb. Optim.*, 29(1):276–307, 2015.
- [8] M. Bíró, M. Hujter, and Z. Tuza. Precoloring extension. I. Interval graphs. *Discrete Math.*, 100(1-3):267–279, 1992.
- [9] J. R. S. Blair and B. Peyton. An introduction to chordal graphs and clique trees. In *Graph theory and sparse matrix computation*, volume 56 of *IMA Vol. Math. Appl.*, pages 1–29. Springer, New York, 1993.

- [10] H. Bodlaender, J. Gustedt, and J. A. Telle. Linear-time register allocation for a fixed number of registers. In *Proceedings of the Ninth Annual ACM-SIAM Symposium on Discrete Algorithms (San Francisco, CA, 1998)*, pages 574–583. ACM, New York, 1998.
- [11] H. L. Bodlaender. A linear-time algorithm for finding tree-decompositions of small treewidth. *SIAM J. Comput.*, 25(6):1305–1317, 1996.
- [12] H. L. Bodlaender. A partial k -arboretum of graphs with bounded treewidth. *Theoret. Comput. Sci.*, 209(1-2):1–45, 1998.
- [13] H. L. Bodlaender and A. M. C. A. Koster. Treewidth computations II. Lower bounds. *Inform. and Comput.*, 209(7):1103–1119, 2011.
- [14] H. L. Bodlaender and R. H. Möhring. The pathwidth and treewidth of cographs. *SIAM J. Discrete Math.*, 6(2):181–188, 1993.
- [15] M. Bonamy, E. Bonnet, H. Déprés, L. Esperet, C. Geniet, C. Hilaire, S. Thomassé, and A. Wesolek. Sparse graphs with bounded induced cycle packing number have logarithmic treewidth, 2022. arxiv:2206.00594.
- [16] E. Bonnet, C. Geniet, E. J. Kim, S. Thomassé, and R. Watrigant. Twin-width II: small classes. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1977–1996. [Society for Industrial and Applied Mathematics (SIAM)], Philadelphia, PA, 2021.
- [17] E. Bonnet, C. Geniet, E. J. Kim, S. Thomassé, and R. Watrigant. Twin-width III: Max Independent Set, Min Dominating Set, and coloring. In *48th International Colloquium on Automata, Languages, and Programming*, volume 198 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. 35, 20. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2021.
- [18] E. Bonnet, E. J. Kim, S. Thomassé, and R. Watrigant. Twin-width I: tractable FO model checking. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science—FOCS 2020*, pages 601–612. IEEE Computer Soc., Los Alamitos, CA, 2020.
- [19] A. Brandstädt, V. B. Le, and J. P. Spinrad. *Graph classes: a survey*. SIAM Monographs on Discrete Mathematics and Applications. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999.
- [20] A. Brandstädt and R. Mosca. Maximum weight independent set for ℓ claw-free graphs in polynomial time. *Discrete Appl. Math.*, 237:57–64, 2018.
- [21] N. Brettell, J. Horsfield, A. Munaro, G. Paesani, and D. Paulusma. Bounding the mim-width of hereditary graph classes. In *15th International Symposium on Parameterized and Exact Computation*, volume 180 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. 6, 18. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2020.
- [22] M. Brianiński, J. Davies, and B. Walczak. Separating polynomial χ -boundedness from χ -boundedness. 2022. arXiv:2201.08814.
- [23] B.-M. Bui-Xuan, J. A. Telle, and M. Vatshelle. Boolean-width of graphs. *Theoret. Comput. Sci.*, 412(39):5187–5204, 2011.
- [24] B.-M. Bui-Xuan, J. A. Telle, and M. Vatshelle. Fast dynamic programming for locally checkable vertex subset and vertex partitioning problems. *Theoret. Comput. Sci.*, 511:66–76, 2013.
- [25] P. Buneman. A characterisation of rigid circuit graphs. *Discrete Math.*, 9:205–212, 1974.
- [26] K. Cameron. Induced matchings. *Discrete Appl. Math.*, 24(1-3):97–102, 1989.

- [27] K. Cameron and P. Hell. Independent packings in structured graphs. *Math. Program.*, 105(2-3, Ser. B):201–213, 2006.
- [28] P. Chalermsook and B. Walczak. Coloring and maximum weight independent set of rectangles. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 860–868. [Society for Industrial and Applied Mathematics (SIAM)], Philadelphia, PA, 2021.
- [29] G. J. Chang. The weighted independent domination problem is NP-complete for chordal graphs. *Discrete Appl. Math.*, 143(1-3):351–352, 2004.
- [30] S. Chaplick, F. V. Fomin, P. A. Golovach, D. Knop, and P. Zeman. Kernelization of graph Hamiltonicity: proper H -graphs. *SIAM J. Discrete Math.*, 35(2):840–892, 2021.
- [31] S. Chaplick, M. Töpfer, J. Voborník, and P. Zeman. On H -topological intersection graphs. In *Graph-theoretic concepts in computer science*, volume 10520 of *Lecture Notes in Comput. Sci.*, pages 167–179. Springer, Cham, 2017.
- [32] S. Chaplick, M. Töpfer, J. Voborník, and P. Zeman. On H -topological intersection graphs. *Algorithmica*, 83(11):3281–3318, 2021.
- [33] S. Chaplick and P. Zeman. Combinatorial problems on H -graphs. *Electron. Notes Discret. Math.*, 61:223–229, 2017.
- [34] M. Chudnovsky. Induced subgraphs and tree decompositions. Talk at the Stony Brook Mathematics Colloquium, October 22, 2020 (online).
- [35] M. Chudnovsky. Induced subgraphs and tree decompositions. Talk at the Berlin Mathematical School, MATH+ Friday Colloquium, April 16, 2021 (online).
- [36] M. Chudnovsky. Induced subgraphs and tree decompositions. Talk at the Charles University, Faculty of Mathematics and Physics, Department of Applied Mathematics, Noon seminar, May 14, 2021 (online).
- [37] M. Chudnovsky. Induced subgraphs and tree decompositions. Invited talk at IWOCA 2021: 32nd International Workshop on Combinatorial Algorithms, 5-7 July 2021, Ottawa, Canada (online).
- [38] M. Chudnovsky, I. Penev, A. Scott, and N. Trotignon. Substitution and χ -boundedness. *J. Combin. Theory Ser. B*, 103(5):567–586, 2013.
- [39] M. Chudnovsky, M. Pilipczuk, M. Pilipczuk, and S. Thomassé. On the maximum weight independent set problem in graphs without induced cycles of length at least five. *SIAM J. Discrete Math.*, 34(2):1472–1483, 2020.
- [40] M. Chudnovsky, M. Pilipczuk, M. Pilipczuk, and S. Thomassé. Quasi-polynomial time approximation schemes for the maximum weight independent set problem in H -free graphs. In *Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms*, pages 2260–2278. SIAM, Philadelphia, PA, 2020.
- [41] D. G. Corneil and Y. Perl. Clustering and domination in perfect graphs. *Discrete Appl. Math.*, 9(1):27–39, 1984.
- [42] D. G. Corneil and U. Rotics. On the relationship between clique-width and treewidth. *SIAM J. Comput.*, 34(4):825–847, 2005.
- [43] B. Courcelle. The monadic second-order logic of graphs. I. Recognizable sets of finite graphs. *Inform. and Comput.*, 85(1):12–75, 1990.
- [44] B. Courcelle, J. A. Makowsky, and U. Rotics. Linear time solvable optimization problems on graphs of bounded clique-width. *Theory Comput. Syst.*, 33(2):125–150, 2000.

- [45] B. Courcelle and S. Olariu. Upper bounds to the clique width of graphs. *Discrete Appl. Math.*, 101(1-3):77–114, 2000.
- [46] M. Cygan, F. V. Fomin, Ł. Kowalik, D. Lokshantov, D. Marx, M. Pilipczuk, M. Pilipczuk, and S. Saurabh. *Parameterized algorithms*. Springer, Cham, 2015.
- [47] K. K. Dabrowski, M. Johnson, and D. Paulusma. Clique-width for hereditary graph classes. In *Surveys in combinatorics 2019*, volume 456 of *London Math. Soc. Lecture Note Ser.*, pages 1–56. Cambridge Univ. Press, Cambridge, 2019.
- [48] C. Dallard, M. Milanič, and K. Štorgel. Treewidth versus clique number in graph classes with a forbidden structure. In I. Adler and H. Müller, editors, *Graph-Theoretic Concepts in Computer Science - 46th International Workshop, WG 2020, Leeds, UK, June 24-26, 2020, Revised Selected Papers*, volume 12301 of *Lecture Notes in Computer Science*, pages 92–105. Springer, 2020.
- [49] C. Dallard, M. Milanič, and K. Štorgel. Treewidth versus clique number. III. Tree-independence number of graphs with a forbidden structure, 2022. arXiv:2206.15092.
- [50] C. Dallard, M. Milanič, and K. Štorgel. Treewidth versus clique number. I. Graph classes with a forbidden structure. *SIAM J. Discrete Math.*, 35(4):2618–2646, 2021.
- [51] R. Diestel and M. Müller. Connected tree-width. *Combinatorica*, 38(2):381–398, 2018.
- [52] G. A. Dirac. On rigid circuit graphs. *Abh. Math. Sem. Univ. Hamburg*, 25:71–76, 1961.
- [53] Y. Dourisboure and C. Gavaille. Tree-decompositions with bags of small diameter. *Discrete Math.*, 307(16):2008–2029, 2007.
- [54] F. F. Dragan and E. Köhler. An approximation algorithm for the tree t -spanner problem on unweighted graphs via generalized chordal graphs. *Algorithmica*, 69(4):884–905, 2014.
- [55] P. Duchet. Classical perfect graphs: an introduction with emphasis on triangulated and interval graphs. In *Topics on perfect graphs*, volume 88 of *North-Holland Math. Stud.*, pages 67–96. North-Holland, Amsterdam, 1984.
- [56] L. Esperet. *Graph colorings, flows and perfect matchings*. Habilitation Thesis, Université Grenoble Alpes, 2017.
- [57] M. Farber. Independent domination in chordal graphs. *Oper. Res. Lett.*, 1(4):134–138, 1981/82.
- [58] U. Feige, M. Hajiaghayi, and J. R. Lee. Improved approximation algorithms for minimum weight vertex separators. *SIAM J. Comput.*, 38(2):629–657, 2008.
- [59] F. Fomin and T. Korhonen. Fast FPT-approximation of branchwidth, 2021. arXiv:2111.03492.
- [60] F. V. Fomin and P. A. Golovach. Subexponential parameterized algorithms and kernelization on almost chordal graphs. *Algorithmica*, 83(7):2170–2214, 2021.
- [61] F. V. Fomin, P. A. Golovach, and J.-F. Raymond. On the tractability of optimization problems on H -graphs. *Algorithmica*, 82(9):2432–2473, 2020.
- [62] F. V. Fomin, I. Todinca, and Y. Villanger. Large induced subgraphs via triangulations and CMSO. *SIAM J. Comput.*, 44(1):54–87, 2015.
- [63] M. R. Garey, D. S. Johnson, G. L. Miller, and C. H. Papadimitriou. The complexity of coloring circular arcs and chords. *SIAM J. Algebraic Discrete Methods*, 1(2):216–227, 1980.
- [64] P. Gartland and D. Lokshantov. Independent set on P_k -free graphs in quasi-polynomial time. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science—FOCS 2020*, pages 613–624. IEEE Computer Soc., Los Alamitos, CA, 2020.

- [65] P. Gartland, D. Lokshantov, M. Pilipczuk, M. Pilipczuk, and P. Rzażewski. Finding large induced sparse subgraphs in $C_{>t}$ -free graphs in quasipolynomial time. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2021, page 330–341, New York, NY, USA, 2021. Association for Computing Machinery.
- [66] F. Gavril. The intersection graphs of subtrees in trees are exactly the chordal graphs. *J. Combinatorial Theory Ser. B*, 16:47–56, 1974.
- [67] M. C. Golumbic. *Algorithmic Graph Theory and Perfect Graphs*, volume 57 of *Annals of Discrete Mathematics*. Elsevier Science B.V., Amsterdam, second edition, 2004.
- [68] M. Grötschel, L. Lovász, and A. Schrijver. *Geometric algorithms and combinatorial optimization*, volume 2 of *Algorithms and Combinatorics: Study and Research Texts*. Springer-Verlag, Berlin, 1988.
- [69] A. Grzesik, T. Klímová, M. Pilipczuk, and M. Pilipczuk. Polynomial-time algorithm for maximum weight independent set on P_6 -free graphs. *ACM Trans. Algorithms*, 18(1), jan 2022.
- [70] A. Gyárfás. Problems from the world surrounding perfect graphs. *Zastos. Mat.*, 19(3-4):413–441, 1987.
- [71] A. Jacob, F. Panolan, V. Raman, and V. Sahlhot. Structural parameterizations with modulator oblivion. In *15th International Symposium on Parameterized and Exact Computation*, volume 180 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. No. 19, 18. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2020.
- [72] L. Jaffke, O.-j. Kwon, T. J. F. Strømme, and J. A. Telle. Mim-width III. Graph powers and generalized distance domination problems. *Theoret. Comput. Sci.*, 796:216–236, 2019.
- [73] P. Jégou and C. Terrioux. Tree-decompositions with connected clusters for solving constraint networks. In *International Conference on Principles and Practice of Constraint Programming*, pages 407–423. Springer, 2014.
- [74] D. Y. Kang, O.-j. Kwon, T. J. F. Strømme, and J. A. Telle. A width parameter useful for chordal and co-comparability graphs. *Theoret. Comput. Sci.*, 704:1–17, 2017.
- [75] R. M. Karp. Reducibility among combinatorial problems. In *Complexity of computer computations (Proc. Sympos., IBM Thomas J. Watson Res. Center, Yorktown Heights, N.Y., 1972)*, pages 85–103, 1972.
- [76] D. Kobler and U. Rotics. Edge dominating set and colorings on graphs with fixed clique-width. *Discrete Appl. Math.*, 126(2-3):197–221, 2003.
- [77] P. K. Krause. Optimal register allocation in polynomial time. In R. Jhala and K. D. Bosschere, editors, *Compiler Construction - 22nd International Conference, CC 2013, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2013, Rome, Italy, March 16-24, 2013. Proceedings*, volume 7791 of *Lecture Notes in Computer Science*, pages 1–20. Springer, 2013.
- [78] E. Lee. Partitioning a graph into small pieces with applications to path transversal. *Math. Program.*, 177(1-2, Ser. A):1–19, 2019.
- [79] S. Mengel. Lower bounds on the mim-width of some graph classes. *Discrete Appl. Math.*, 248:28–32, 2018.
- [80] Y. Orlovich, A. Dolgui, G. Finke, V. Gordon, and F. Werner. The complexity of dissociation set problems in graphs. *Discrete Appl. Math.*, 159(13):1352–1366, 2011.
- [81] S.-i. Oum and P. Seymour. Approximating clique-width and branch-width. *J. Combin. Theory Ser. B*, 96(4):514–528, 2006.

- [82] B. S. Panda, A. Pandey, J. Chaudhary, P. Dane, and M. Kashyap. Maximum weight induced matching in some subclasses of bipartite graphs. *J. Comb. Optim.*, 40(3):713–732, 2020.
- [83] M. Pilipczuk, M. Pilipczuk, and P. Rzażewski. Quasi-polynomial-time algorithm for independent set in P_t -free graphs via shrinking the space of induced paths. In H. V. Le and V. King, editors, *4th Symposium on Simplicity in Algorithms, SOSA 2021, Virtual Conference, January 11-12, 2021*, pages 204–209. [Society for Industrial and Applied Mathematics (SIAM)], Philadelphia, PA, 2021.
- [84] V. Raghavan and J. Spinrad. Robust algorithms for restricted domains. *J. Algorithms*, 48(1):160–172, 2003.
- [85] F. P. Ramsey. On a Problem of Formal Logic. *Proc. London Math. Soc. (2)*, 30(4):264–286, 1929.
- [86] N. Robertson and P. D. Seymour. Graph minors. III. Planar tree-width. *J. Combin. Theory Ser. B*, 36(1):49–64, 1984.
- [87] N. Robertson and P. D. Seymour. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms*, 7(3):309–322, 1986.
- [88] D. J. Rose, R. E. Tarjan, and G. S. Lueker. Algorithmic aspects of vertex elimination on graphs. *SIAM J. Comput.*, 5(2):266–283, 1976.
- [89] P. Scheffler. What graphs have bounded tree-width? In *Proceedings of the 7th Fischland Colloquium, III (Wustrow, 1988)*, number 41, pages 31–38, 1990.
- [90] A. Scott and P. Seymour. A survey of χ -boundedness. *J. Graph Theory*, 95(3):473–504, 2020.
- [91] P. Seymour. Tree-chromatic number. *J. Combin. Theory Ser. B*, 116:229–237, 2016.
- [92] K. Skodinis. Efficient analysis of graphs with small minimal separators. In *Graph-theoretic concepts in computer science (Ascona, 1999)*, volume 1665 of *Lecture Notes in Comput. Sci.*, pages 155–166. Springer, Berlin, 1999.
- [93] J. P. Spinrad. *Efficient graph representations*, volume 19 of *Fields Institute Monographs*. American Mathematical Society, Providence, RI, 2003.
- [94] L. Sunil Chandran. A linear time algorithm for enumerating all the minimum and minimal separators of a chordal graph. In *Computing and combinatorics (Guilin, 2001)*, volume 2108 of *Lecture Notes in Comput. Sci.*, pages 308–317. Springer, Berlin, 2001.
- [95] R. E. Tarjan. Decomposition by clique separators. *Discrete Math.*, 55(2):221–232, 1985.
- [96] L. Vandenberghe and M. S. Andersen. Chordal graphs and semidefinite optimization. *Found. Trends Optim.*, 1(4):241–433, 2015.
- [97] M. Vatshelle. *New width parameters of graphs*. PhD thesis, University of Bergen, 2012.
- [98] J. R. Walter. Representations of chordal graphs as subtrees of a tree. *J. Graph Theory*, 2(3):265–267, 1978.
- [99] D. B. West. *Introduction to Graph Theory*. Prentice Hall, Inc., Upper Saddle River, NJ, 1996.
- [100] M. Yannakakis. Node-deletion problems on bipartite graphs. *SIAM J. Comput.*, 10(2):310–327, 1981.
- [101] J. You, J. Wang, and Y. Cao. Approximate association via dissociation. In *Graph-theoretic concepts in computer science*, volume 9941 of *Lecture Notes in Comput. Sci.*, pages 13–24. Springer, Berlin, 2016.
- [102] D. Zuckerman. Linear degree extractors and the inapproximability of max clique and chromatic number. *Theory Comput.*, 3:103–128, 2007.
- [103] I. E. Zverovich. Satgraphs and independent domination. I. *Theoret. Comput. Sci.*, 352(1-3):47–56, 2006.