Noname manuscript No.
(will be inserted by the editor)

Extremal K_4-minor-free graphs without short cycles

János Barát

arXiv:2111.05712v1 [math.CO] 10 Nov 2021

Received: date / Accepted: date

Abstract We determine the maximum number of edges in a K_4-minor-free n-vertex graph of girth g, when $g = 5$ or g is even. We argue that there are many different n-vertex extremal graphs, if n is even and g is odd.

Keywords extremal graph · K_4-minor-free · girth · Euler's formula · graph decomposition

1 Introduction

We assume the reader is familiar with basic graph theory, see e.g. [3]. In particular, let n denote the number of vertices, e the number of edges, f the number of faces of a planar graph. A face with precisely i edges on its boundary is an i-face. The *girth* of a graph is the length of the shortest cycle. We denote the distance[1] between vertices u and v in a graph H by $d_H(u, v)$.

Let u and v be adjacent vertices. The *contraction* of an edge uv is the following operation. We delete the edge uv and merge the vertices u and v into a new vertex w such that edges incident to either u or v correspond to edges incident to w. For our purposes, we only keep single edges if multiple edges are created. A graph H is a *minor* of a graph G if we can get H from G by contracting and deleting edges. A graph G contains a complete graph K_t as a minor if G contains t disjoint connected subgraphs such that any two such subgraphs are connected by an edge of G. A graph G is *K_t-minor-free* if G does not contain K_t as a minor.

Let C be a cycle of a graph G, and let F be a connected component of $G - C$. The edges between vertices of F and C are the *leg*s of F. A *bridge* B of C in G is

Supported by ERC Advanced Grant "GeoScape", National Research, Development and Innovation Office, NKFIH, K-131529. and the grant of the Hungarian Ministry for Innovation and Technology (Grant Number: NKFIH-1158-6/2019).

János Barát
Alfréd Rényi Institute of Mathematics, Eötvös Loránd Research Network, Budapest, Hungary
and Department of Mathematics, University of Pannonia, Hungary
E-mail: barat@mik.uni-pannon.hu

[1] length of the shortest path

a connected component of $G - C$ together with its legs. The vertices in $B \cap C$ are the *attachment* vertices.

In an *extremal* problem, we would like to determine the maximum number of edges of an n-vertex graph G that belongs to a graph class $\mathcal{C}$. The graphs that have this maximum number of edges are the *extremal graph*s. In the classical example, Turán's theorem, there is precisely one extremal graph. In our focus, the graph class is the K_t-minor-free graphs of girth g, where both t and g are small. In some cases there is only one extremal graph and in some other cases there are many.

Our motivation comes from graph colorings. An easy list-coloring argument is the following. Assume every vertex has a list of colors of size k. Consider a class $\mathcal{C}$ of graphs, where every graph G in the class has minimum degree $\delta(G) \leq k-1$. Now every graph in the class can be k-list-colored by the greedy algorithm, always coloring a vertex of minimum degree.

Our distant goal is to study K_6-minor-free graphs without short cycles. The preliminary results show that proving an extremal result might be enough to use the simple algorithm given above to improve on the present upper bounds. To find the necessary tools for the problem, we first study K_4- and K_5-minor-free graphs with a condition on the girth. See also the separate paper on K_5-minor-free graphs [1].

Any K_4-minor-free graph is a subgraph of a 3-tree. That is, a graph built from triangles pasted along edges. The K_4-minor-free graphs are also called *series-parallel* graphs. Observe that any K_4-minor-free graph is planar.

2 Even girth

Let G be a K_4-minor-free graph and C a cycle of G. Studying planar graphs, the relative position of bridges of a cycle plays a crucial role [2]. Two bridges B_1 and B_2 of C are *crossing*, if there are attachment vertices u_1, u_2 of B_1 and v_1, v_2 of B_2 such that u_1, v_1, u_2, v_2 appears on C in this order. We use the following simple facts.

Proposition 1 *Let C be any cycle of a K_4-minor-free graph G.*
(i) Every bridge of C can have at most 2 legs.
(ii) The bridges of C are non-crossing.

Since G is K_4-minor-free, it has a planar drawing. Consider G^* the dual of any drawing of G, where we ignore the unbounded region. Any cycle in G^* would imply a K_4-minor in G. Therefore, G^* must be a tree.

Let G be a K_4-minor-free graph of girth $2k$, where $k \geq 2$.

Proposition 2 *A K_4-minor-free graph G on n vertices of girth $2k$ can have at most $\frac{k}{k-1}(n-2)$ edges, where $k \geq 2$. This bound is tight, if $n = s(k-1)+2$, where $s \geq 2$.*

Proof The graph G satisfies Euler's formula: $e + 2 = f + n$. Also, by the girth condition $2e = \sum i f_i \geq 2kf$, where f_i denotes the number of i-faces of G. Therefore, $2ke + 4k = 2kf + 2kn \leq 2e + 2kn$, which implies $e \leq \frac{2kn-4k}{2k-2} = \frac{k}{k-1}(n-2)$.

Next, we construct a K_4-minor-free graph G with $n = s(k-1)+2$ vertices and $e = ks$ edges and of girth $2k$, where $s \geq 2$. This gives us a sharp example as $\frac{k\,(s(k-1)+2)-2k}{k-1} = ks$. The graph G is the following: let u and v be two vertices

and connect them by s edge-disjoint paths with k edges. Now every cycle must go through u and v, hence the girth is indeed $2k$. The graph G is also K_4-minor-free since removing any cycle leaves only bridges with 2 legs.

We show that actually these are all the extremal examples, when $n = s(k-1)+2$. Notice that the application of Euler's formula shows us that in any drawing of an extremal graph G all faces must have $2k$ edges. Let C be a face of length $2k$ in G. Consider all bridges of C in G. Any such bridge B can have at most two legs (actually must have 2 legs). Also bridges must be non-crossing. Finally, any two bridges must have the same attachment vertices. Otherwise we get a face with length greater than $2k$. Now we know that there are two vertices u and v of C and all bridges are attached to these two vertices in G.

Taking into account that every face has length $2k$, we get that a bridge B cannot contain a path longer than the path between u and v in C. This implies that indeed B cannot be other than a copy of the path between u and v. □

3 Girth 5

In this section G denotes a K_4-minor-free graph of girth 5. We start with an important observation.

Lemma 1 *We may assume that an extremal graph G is 2-connected.*

Proof Suppose that a K_4-minor-free graph G' of girth 5 has a cutvertex x such that $G_1 \cup G_2 = G'$ and $G_1 \cap G_2 = x$. Let v_1 be a neighbor of x in G_1 and v_2 a neighbor in G_2. We modify the graph G' by deleting the edge v_2x and adding the edge v_1v_2. Notice the girth condition is still satisfied. How could there be a K_4-minor now? Only if there is a cycle C through x and v_1v_2 and a bridge B of C with at least three legs. However, the attachment vertices of B must be entirely in G_1 or G_2. If they are in G_1, then we contract the path of C between x and v_1 through v_2 to find a K_4-minor already in G_1. If the attachment vertices are in G_2, then we contract the path of C between x and v_2 through v_1 to find a K_4-minor already in G_2. It shows that the modified graph is still K_4-minor-free. Repeating the above process we can transform the original graph to one without cutvertices, but still having the same parameters. □

In what follows, we assume G is 2-connected. As it often happens, the odd girth case is different and more complicated than even girth. We can use Euler's formula as before to set an upper bound. However, there is no matching lower bound construction.

Theorem 1 *Any n-vertex K_4-minor-free graph G of girth 5 can have at most $\lceil \frac{3}{2}n-3 \rceil$ edges, where $n \geq 5$.*

Proof We prove the statement by induction on the number of vertices n. For the base of induction, we notice that on 5 vertices, G can have at most 5 edges. We get $5 \leq \lceil \frac{3}{2}5 - 3 \rceil = 5$ and the statement holds. On 6 vertices, the 6-cycle is the only possible graph and the claim holds again. Finally on 7 vertices, there are two possible graphs and one extremal example. The inequality holds for these too.

For the induction step, let G be a 2-connected extremal K_4-minor-free graph of girth 5. Let $\{u,v\}$ be a 2-cut[2] such that $G_1 \cup G_2 = G$ and $G_1 \cap G_2 = \{u,v\}$. Let n_i and m_i denote the number of vertices in G_i for $i \in \{1,2\}$. Observe here that we can always select u and v such that G_1 has at least 4 vertices[3], when $n \geq 8$.

First suppose that $d_{G_1}(u,v) = 1$ and $d_{G_2}(u,v) \geq 4$. We add an edge between u and v in G_2 to create G_2^+. By the induction hypothesis for G_1 and G_2^+, we deduce the following:
$m_1 \leq \lceil \frac{3}{2}n_1 - 3 \rceil \leq \frac{3}{2}n_1 - \frac{5}{2}$.
$m_2 + 1 \leq \lceil \frac{3}{2}n_2 - 3 \rceil \leq \frac{3}{2}n_2 - \frac{5}{2}$.

Since $m = m_1 + m_2$ and $n = n_1 + n_2 - 2$, we get:
$m = m_1 + m_2 \leq \frac{3}{2}n_1 - \frac{5}{2} + \frac{3}{2}n_2 - \frac{7}{2} = \frac{3}{2}(n_1 + n_2) - 6 = \frac{3}{2}n - 3 \leq \lceil \frac{3}{2}n - 3 \rceil$.

Now suppose that $d_{G_1}(u,v) \geq 2$ and $d_{G_2}(u,v) \geq 3$.

Case 1. n_1 and n_2 are even. Now we directly apply the induction hypothesis on G_1 and G_2:
$m_1 \leq \lceil \frac{3}{2}n_1 - 3 \rceil = \frac{3}{2}n_1 - 3$.

$m_2 \leq \lceil \frac{3}{2}n_2 - 3 \rceil = \frac{3}{2}n_2 - 3$.

Since $m = m_1 + m_2$ and $n = n_1 + n_2 - 2$, we get:
$m = m_1 + m_2 \leq \frac{3}{2}n_1 - 3 + \frac{3}{2}n_2 - 3 = \frac{3}{2}n - 3 = \lceil \frac{3}{2}n - 3 \rceil$, since n is even.

Case 2. n_1 and n_2 are odd. Now we add a path with 2 edges between u and v in G_2 to create G_2^+. By the induction hypothesis for G_1 and G_2^+, we deduce the following:
$m_1 \leq \lceil \frac{3}{2}n_1 - 3 \rceil = \frac{3}{2}n_1 - \frac{5}{2}$.
$m_2 + 2 \leq \lceil \frac{3}{2}(n_2 + 1) - 3 \rceil = \frac{3}{2}(n_2 + 1) - 3$.

Since $m = m_1 + m_2$ and $n = n_1 + n_2 - 2$, we get:
$m = m_1 + m_2 \leq \frac{3}{2}n_1 - \frac{5}{2} + \frac{3}{2}(n_2 + 1) - 5 = \frac{3}{2}(n_1 + n_2) - 6 = \frac{3}{2}n - 3 \leq \lceil \frac{3}{2}n - 3 \rceil$.

Case 3. n_1 is even and n_2 is odd. Again we add a path with 2 edges between u and v in G_2 to create G_2^+. By the induction hypothesis for G_1 and G_2^+, we deduce the following:
$m_1 \leq \lceil \frac{3}{2}n_1 - 3 \rceil = \frac{3}{2}n_1 - 3$.
$m_2 + 2 \leq \lceil \frac{3}{2}(n_2 + 1) - 3 \rceil = \frac{3}{2}(n_2 + 1) - 3$.

Since $m = m_1 + m_2$ and $n = n_1 + n_2 - 2$, we get:
$m = m_1 + m_2 \leq \frac{3}{2}n_1 - 3 + \frac{3}{2}(n_2 + 1) - 5 = \frac{3}{2}(n_1 + n_2) - 8 + \frac{3}{2} = \frac{3}{2}n - \frac{7}{2} < \frac{3}{2}n - 3 \leq \lceil \frac{3}{2}n - 3 \rceil$.

Case 4. n_1 is odd and n_2 is even. We add a path with 2 edges between u and v in G_2 to create G_2^+. By the induction hypothesis for G_1 and G_2^+, we deduce the following:
$m_1 \leq \lceil \frac{3}{2}n_1 - 3 \rceil = \frac{3}{2}n_1 - \frac{5}{2}$.
$m_2 + 2 \leq \lceil \frac{3}{2}(n_2 + 1) - 3 \rceil = \frac{3}{2}(n_2 + 1) - \frac{5}{2}$.

Since $m = m_1 + m_2$ and $n = n_1 + n_2 - 2$, we get:
$m = m_1 + m_2 \leq \frac{3}{2}n_1 - \frac{5}{2} + \frac{3}{2}(n_2 + 1) - \frac{5}{2} - 2 = \frac{3}{2}(n_1 + n_2) - 7 + \frac{3}{2} = \frac{3}{2}n - 4 + \frac{3}{2} = \frac{3}{2}n - \frac{5}{2} = \lceil \frac{3}{2}n - 3 \rceil$, since n is odd. □

[2] There is always a 2-cut since G is the subgraph of a 3-tree.
[3] That is, the graph G_1 is larger than a path with 2 edges.

4 Lower bound construction

Next, we show a K_4-minor-free graph G_s with $n = 2s+1$ vertices and $e = 3s-1 = \lceil \frac{3}{2}n - 3 \rceil$ edges and of girth 5 for every $s \geq 2$. This shows that Theorem 1 is tight for every odd n, where $n \geq 5$. The construction is similar to the one for even girth. Let u and v be two vertices and connect them by $s-1$ edge-disjoint paths with 3 edges and one path with 2 edges. See Fig 1.




Fig. 1 Extremal K_4-minor-free graphs of girth 5.

If we subdivide an edge[4] of G_{s-1} not in the 2-edge path between u and v, then we get a K_4-minor-free graph of girth 5 on $n = 2s$ vertices and $e = 3s-3 = \frac{3}{2}n-3$ edges. Next, we indicate that if the number of vertices is even, there are many other graphs with the same parameters.

We determine all K_4-minor-free graphs on 10 vertices of girth 5 and 12 edges. Let us consider the planar drawings of these graphs. We distinguish them by the length l of their outer cycle C.

If $l = 10$, then there must be two more edges connecting vertices of C. Since G is K_4-minor-free, these two edges form non-crossing bridges. So there are 3 faces inside of C. Now two edges can make the total length of these 3 faces to 14. There must be a face with at most 4 edges, a contradiction.

If $l = 9$, then there must be a vertex x in C such that x is adjacent to y in C forming a 5-face and x is connected to z in C by a subdivided edge forming two more 5-faces. This gives us the first example: H_1, see Fig 2.

If $l = 8$, then the remaining 2 vertices and 4 edges can form an edge and a 3-edge-path or two 2-edge-paths connecting vertices of C. The total length of the faces inside C must be 16. If an edge is connecting two vertices x and y of C, it must form two 5-cycles with C. We may assume that the remaining 3-path is also incident to x. There are two different ways of connecting x to z in C by a 3-path. In the drawing, either the length of the faces are 5,6,5 or 6,5,5. Now we have to check whether these examples are isomorhic to H_1. We find that indeed the 5,6,5 example is H_1, so we found a new graph H_2, see Fig 2.

If there are two 2-edge-paths, then they either have a vertex x in common or not. If they are independent, then they form faces of length 5,6,5 inside C. This is a new graph H_3 by its degree sequence. If x is connected to y and z in C by two 2-paths, then the faces have length either 5,6,5 or 5,5,6 in order. These are also new examples: H_4 and H_5, see Fig 2.

If $l = 7$, then there are 3 remaining vertices and 5 edges, so the total length of the faces inside C must be 17. Any chord of C would create a short cycle and a 4-path would create a long cycle tracking us back to a previous case. So there can

[4] replace an edge by a 2-edge path

be a 2-path Q and a 3-path P connecting vertices of C. The 2-path must connect two vertices at distance 3 on C creating a 5-cycle and a 6-cycle. If the 3-path P has attachments different from the 2-path, then P connects two vertices at distance 2 on the created 6-cycle. However, this is isomorphic to H_3. Otherwise a common vertex x is incident to both Q and P. Principally there are four possibilities for the other endvertex of P.

1. The other endvertex is on the 5-cycle at distance 2 from x. This creates faces of length 5,6,6 inside C. However, this is isomorphic to H_5.

2. The other endvertex coincides with the other endvertex of Q. Now there are two vertices of degree 4, and the length of the faces inside C are 5,5,7. This is a new example H_6, see Fig 2.

3. The other endvertex of P is at distance 3 from x on C and is different from the other endvertex of Q. This creates faces of length 5,6,6 inside C. This is again a new example H_7, see Fig 2.

4. The other endvertex of P is at distance 2 from x on C. This creates faces of length 5,7,5 inside C. However, this example is isomorphic to H_4.

If $l = 6$, then there are 4 remaining vertices and 6 edges. If they form a path with at least 4 edges, then a cycle of length at least 7 is created. That was covered in the previous cases. Therefore the remaining bridges must be two 3-paths. If we connect two vertices at distance 2 on C by a 3-path, then we create a cycle of length 7. Therefore the 3-paths must connect vertices at distance 3. Since the bridges are non-crossing, the attachments must be the same for the two 3-paths. This is graph H_8, see Fig 2.




Fig. 2 Extremal K_4-minor-free graphs of girth 5 on 10 vertices.

We emphasize again, that thereby we proved that Theorem 1 is tight for $n = 10$, and we have determined the complete list of extremal graphs as shown in Fig 2.

Acknowledgements We thank an anonymous referee of [1], who suggested this project. We thank another anonymous referee for suggesting structural changes that improved the presentation.

References

1. János Barát. On the number of edges in a K_5-minor-free graph of given girth. *submitted*
2. J.Adrian Bondy and U.S.Ram Murty *Graph theory*, vol. 244. of *Graduate Texts in Mathematics*. Springer, 1st corrected edition, 2008.
3. Reinhard Diestel. *Graph theory*, vol. 173. of *Graduate Texts in Mathematics*. Springer, 5th edition, 2016.
4. Carsten Thomassen. A theorem on paths in planar graphs. *Journal of Graph Theory*, **7** 169–176, 1983.