

# WEAK DEGENERACY OF GRAPHS

ANTON BERNSHTEYN AND EUGENE LEE

**ABSTRACT.** Motivated by the study of greedy algorithms for graph coloring, we introduce a new graph parameter, which we call *weak degeneracy*. By definition, every  $d$ -degenerate graph is also weakly  $d$ -degenerate. On the other hand, if  $G$  is weakly  $d$ -degenerate, then  $\chi(G) \leq d + 1$  (and, moreover, the same bound holds for the list-chromatic and even the DP-chromatic number of  $G$ ). It turns out that several upper bounds in graph coloring theory can be phrased in terms of weak degeneracy. For example, we show that planar graphs are weakly 4-degenerate, which implies Thomassen's famous theorem that planar graphs are 5-list-colorable. We also prove a version of Brooks's theorem for weak degeneracy: a connected graph  $G$  of maximum degree  $d \geq 3$  is weakly  $(d - 1)$ -degenerate unless  $G \cong K_{d+1}$ . (By contrast, all  $d$ -regular graphs have degeneracy  $d$ .) We actually prove an even stronger result, namely that for every  $d \geq 3$ , there is  $\varepsilon > 0$  such that if  $G$  is a graph of weak degeneracy at least  $d$ , then either  $G$  contains a  $(d + 1)$ -clique or the maximum average degree of  $G$  is at least  $d + \varepsilon$ . Finally, we show that graphs of maximum degree  $d$  and either of girth at least 5 or of bounded chromatic number are weakly  $(d - \Omega(\sqrt{d}))$ -degenerate, which is best possible up to the value of the implied constant.

## 1. Introduction

All graphs in this paper are finite and simple. Recall that for a graph  $G$ ,  $\chi(G)$  denotes its **chromatic number**, i.e., the minimum number of colors necessary to color the vertices of  $G$  so that adjacent vertices are colored differently. A well-studied generalization of graph coloring is **list coloring**, which was introduced independently by Vizing [Viz76] and Erdős, Rubin, and Taylor [ERT79]. In the setting of list coloring, each vertex  $u \in V(G)$  is given a set  $L(u)$ , called its **list of available colors**. A **proper  $L$ -coloring** is then a function  $\varphi$  defined on  $V(G)$  such that:

- $\varphi(u) \in L(u)$  for all  $u \in V(G)$ ; and
- $\varphi(u) \neq \varphi(v)$  for all  $uv \in E(G)$ .

The **list-chromatic number** of  $G$ , denoted by  $\chi_\ell(G)$ , is the minimum  $k$  such that  $G$  admits a proper  $L$ -coloring whenever  $|L(u)| \geq k$  for all  $u \in V(G)$ . Clearly,  $\chi_\ell(G) \geq \chi(G)$  for all graphs  $G$ .

A further generalization of list coloring is **DP-coloring** (also known as **correspondence coloring**), which was recently introduced by Dvořák and Postle [DP18]. A related notion of *local conflict coloring* was studied independently from the algorithmic standpoint by Fraigniaud, Heinrich, and Kosowski [FHK16]. Just as in list coloring, we assume that every vertex  $u \in V(G)$  of a graph  $G$  is given a list  $L(u)$  of colors to choose from. In contrast to list coloring though, the identifications between the colors in the lists are allowed to vary from edge to edge. That is, each edge  $uv \in E(G)$  is assigned a matching  $C_{uv}$  (not necessarily perfect and possibly empty) from  $L(u)$  to  $L(v)$ . If  $\alpha\beta \in C_{uv}$ , we say that  $\alpha$  **corresponds** to  $\beta$  (under the correspondence  $C$ ). A **proper  $(L, C)$ -coloring** of  $G$  is a function  $\varphi$  defined on  $V(G)$  such that:

- $\varphi(u) \in L(u)$  for all  $u \in V(G)$ ; and
- $\varphi(u)\varphi(v) \notin C_{uv}$  for all  $uv \in E(G)$ .

---

(Anton Bernshteyn) School of Mathematics, Georgia Institute of Technology, Atlanta, GA, USA  
 (Eugene Lee) Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh, PA, USA  
*E-mail addresses:* bahtoh@gatech.edu, eleeuhaj@andrew.cmu.edu.

Research of the first named author is partially supported by the NSF grant DMS-2045412.

List coloring is a special case of this framework where  $\alpha \in L(u)$  corresponds to  $\beta \in L(v)$  if and only if  $\alpha = \beta$ . The **DP-chromatic number** of  $G$ , denoted by  $\chi_{DP}(G)$ , is the minimum  $k$  such that  $G$  admits a proper  $(L, C)$ -coloring whenever  $|L(u)| \geq k$  for all  $u \in V(G)$ . Again, it is clear from the definition that  $\chi_{DP}(G) \geq \chi_\ell(G)$ .

In this paper we are interested in greedy algorithms for graph coloring. The basic greedy algorithm considers the vertices of  $G$  one at a time. When we get to consider a vertex  $u$ , we assign to it an arbitrary color, say  $\alpha$ , from  $L(u)$ . At this point, to ensure that the coloring is proper, we have to remove the colors corresponding to  $\alpha$  from the lists of colors available to the neighbors of  $u$ . Thus, the list size for every neighbor of  $u$  may decrease by 1, while all the other lists remain unchanged. If throughout this process no list size reduces to 0 (i.e., if every uncolored vertex always has at least one available color), then we successfully obtain a proper (DP-)coloring of  $G$ . This idea is formally captured in the notion of *graph degeneracy*:

**Definition 1.1** (Degeneracy). Let  $G$  be a graph and let  $f: V(G) \rightarrow \mathbb{N}$  be a function.<sup>1</sup> For a vertex  $u \in V(G)$ , the operation  $\text{DELETE}(G, f, u)$  outputs the graph  $G' := G - u$  and the function  $f': V(G') \rightarrow \mathbb{Z}$  given by the formula

$$f'(v) := \begin{cases} f(v) - 1 & \text{if } uv \in E(G); \\ f(v) & \text{otherwise.} \end{cases}$$

An application of the operation  $\text{DELETE}$  is **legal** if the resulting function  $f'$  is non-negative, i.e., if  $f'(v) \geq 0$  for all  $v \in V(G')$ . A graph  $G$  is  **$f$ -degenerate** if it is possible to remove all vertices from  $G$  by a sequence of legal applications of the operation  $\text{DELETE}$ . Given  $d \in \mathbb{N}$ , we say that  $G$  is  **$d$ -degenerate** if it is degenerate with respect to the constant  $d$  function. The **degeneracy** of  $G$ , denoted by  $\mathbf{d}(G)$ , is the minimum  $d$  such that  $G$  is  $d$ -degenerate.

It follows from the above discussion that  $\chi_{DP}(G) \leq \mathbf{d}(G) + 1$  for every graph  $G$ ; because of this, the quantity  $\mathbf{d}(G) + 1$  is sometimes referred to as the *coloring number* of  $G$  [EH66]. It is not hard to see that a graph  $G$  is  $d$ -degenerate if and only if every nonempty subgraph of  $G$  has a vertex of degree at most  $d$  [Die17, Proposition 5.2.2].

The upper bound  $\chi_{DP}(G) \leq \mathbf{d}(G) + 1$  is usually not sharp. For instance, if  $G$  is a  $d$ -regular graph, then  $\mathbf{d}(G) = d$ , which implies that  $\chi_{DP}(G) \leq d + 1$ . However, the only connected  $d$ -regular graphs  $G$  with  $\chi_{DP}(G) = d + 1$  are the complete graph  $K_{d+1}$  and—if  $d = 2$ —cycles [BKP17]. (A curious distinction between DP-coloring and list coloring is that  $\chi_\ell(C_n)$  is 2 if  $n$  is even and 3 if  $n$  is odd, while  $\chi_{DP}(C_n) = 3$  for all  $n \geq 3$  [DP18, §1.1].) It is therefore interesting to see if we can modify the greedy coloring procedure to “save” some of the colors and get a better bound on  $\chi_{DP}(G)$ . Here we investigate a particularly simple (but, as we shall see, already quite powerful) way of doing so.

To motivate our main definition, consider a vertex  $u \in V(G)$  and let  $w$  be its neighbor. In general, if we assign a color to  $u$ , then  $w$  may lose one of its colors. However, suppose that  $|L(u)| > |L(w)|$ , i.e., that  $u$  has *strictly more* available colors than  $w$ . In this case, there must be a color in  $L(u)$  that does not correspond to any color in  $L(w)$ , and assigning such a color to  $u$  does not affect  $L(w)$  (of course, the other neighbors of  $u$  may still lose a color). In this way, we “save” an extra color for  $w$ . This idea naturally leads to the notion that we call *weak degeneracy*:

**Definition 1.2** (Weak degeneracy). Let  $G$  be a graph and let  $f: V(G) \rightarrow \mathbb{N}$  be a function. For a pair of adjacent vertices  $u, w \in V(G)$ , the operation  $\text{DELSAVE}(G, f, u, w)$  outputs the graph  $G' := G - u$  and the function  $f': V(G') \rightarrow \mathbb{Z}$  given by the formula

$$f'(v) := \begin{cases} f(v) - 1 & \text{if } uv \in E(G) \text{ and } v \neq w; \\ f(v) & \text{otherwise.} \end{cases}$$

<sup>1</sup>In this paper  $\mathbb{N} = \{0, 1, 2, \dots\}$  denotes the set of all non-negative integers.

An application of the operation `DELSAVE` is **legal** if  $f(u) > f(w)$  and the resulting function  $f'$  is non-negative. A graph  $G$  is **weakly  $f$ -degenerate** if it is possible to remove all vertices from  $G$  by a sequence of legal applications of the operations `DELETE` and `DELSAVE`. Given  $d \in \mathbb{N}$ , we say that  $G$  is **weakly  $d$ -degenerate** if it is weakly degenerate with respect to the constant  $d$  function. The **weak degeneracy** of  $G$ , denoted by  $\text{wd}(G)$ , is the minimum  $d$  such that  $G$  is weakly  $d$ -degenerate.

Again, the above discussion shows that  $\chi_{DP}(G) \leq \text{wd}(G) + 1$  for every graph  $G$ . Actually, the same bound holds even for the on-line version of DP-chromatic number called *DP-paint number*, which was introduced by Kim, Kostochka, Li, and Zhu [Kim+20] (see §3 for the definition):

**Proposition 1.3.** *For every graph  $G$ ,*

$$\chi(G) \leq \chi_\ell(G) \leq \chi_{DP}(G) \leq \chi_{DPP}(G) \leq \text{wd}(G) + 1,$$

where  $\chi_{DPP}(G)$  is the DP-paint number of  $G$ .

It turns out that the simple way of “saving” colors using the `DELSAVE` operation is sufficient for several non-trivial upper bounds. For example, consider the case of planar graphs. It follows from Euler’s formula that planar graphs are 5-degenerate, which gives a simple proof of their 6-colorability (and even 6-DP-colorability). On the other hand, Thomassen [Tho94] proved that every planar graph is 5-list-colorable, and this result was extended to DP-coloring by Dvořák and Postle [DP18]. The value 5 here is optimal as Voigt [Voi93] constructed planar graphs of list-chromatic number exactly 5. While degeneracy is not sufficient to establish Thomassen’s theorem, we show in §4 that weak degeneracy is:

**Theorem 1.4.** *Every planar graph is weakly 4-degenerate.*

Next we consider Brooks-type theorems for weak degeneracy. As mentioned earlier, if  $d \geq 3$ , then the only connected graph  $G$  of maximum degree  $d \geq 3$  with  $\chi_{DP}(G) = d + 1$  is the complete graph  $K_{d+1}$ . We show that there is a corresponding bound on weak degeneracy:

**Theorem 1.5.** *If  $G$  is a connected graph of maximum degree  $d \geq 3$ , then either  $G \cong K_{d+1}$  or  $G$  is weakly  $(d - 1)$ -degenerate.*

More generally, suppose that  $G$  is a connected graph and  $|L(u)| \geq \deg_G(u)$  for every vertex  $u \in V(G)$  (that is, the lower bound on the list size varies depending on the degree of the vertex). In the list-coloring framework, Borodin [Bor79] and, independently, Erdős, Rubin, and Taylor [ERT79] showed that  $G$  is  $L$ -colorable unless it is a **Gallai tree**, i.e., a connected graph in which every block is either a clique or an odd cycle. In the DP-coloring setting the same result holds, except that the graphs that need to be excluded are the **GDP trees**, i.e., connected graphs in which every block is either a clique or a cycle (not necessarily odd) [BKP17]. We again establish the corresponding result for weak degeneracy:

**Theorem 1.6.** *Let  $G$  be a connected graph. The following statements are equivalent:*

- (1)  $G$  is weakly  $f$ -degenerate, where  $f(u) = \deg_G(u) - 1$  for all  $u \in V(G)$ ;
- (2)  $G$  is not a GDP-tree.

Theorem 1.5 is an immediate corollary of Theorem 1.6. We prove Theorem 1.6 in §5.

Recall that the **average degree** of a nonempty graph  $G$ , denoted by  $\text{ad}(G)$ , is the average of the degrees of the vertices of  $G$ . Equivalently, we have  $\text{ad}(G) = 2|E(G)|/|V(G)|$ . The **maximum average degree** of  $G$ , denoted by  $\text{mad}(G)$ , is defined by  $\text{mad}(G) := \max_H \text{ad}(H)$ , where the maximum is taken over all nonempty subgraphs  $H$  of  $G$ . The maximum average degree of a graph is a natural measure of its local density. There is a close relationship between a graph’s maximum average degree and its degeneracy; namely, we have

$$2d(G) \geq \text{mad}(G) \geq d(G).$$

For  $d$ -regular graphs  $G$ ,  $\text{mad}(G) = d(G) = d$ . By contrast, we show that if  $\text{wd}(G) \geq 3$  and  $G$  contains no  $(\text{wd}(G) + 1)$ -clique, then  $\text{mad}(G) \geq \text{wd}(G) + \varepsilon$ , where  $\varepsilon > 0$  only depends on  $\text{wd}(G)$ :

**Theorem 1.7.** *Let  $G$  be a nonempty graph. If the weak degeneracy of  $G$  is at least  $d \geq 3$ , then either  $G$  contains a  $(d + 1)$ -clique or*

$$\text{mad}(G) \geq d + \frac{d - 2}{d^2 + 2d - 2}.$$

Note that Theorem 1.7 is a strengthening of Theorem 1.5, since  $\text{mad}(G)$  is at most the maximum degree of  $G$ . Our proof of Theorem 1.7, which we present in §5.2, relies on Theorem 1.6 and follows an approach similar to the one used by Gallai [Gal63] to establish a lower bound on the average degree of critical graphs.

As far as lower bounds on weak degeneracy are concerned, a fairly straightforward double counting argument gives the following:

**Proposition 1.8.** *Let  $G$  be a  $d$ -regular graph with  $n \geq 2$  vertices. Then  $\text{wd}(G) \geq d - \sqrt{2n}$ .*

In particular, if  $n = O(d)$ , then  $\text{wd}(G) \geq d - O(\sqrt{d})$ . For example, Proposition 1.8 yields the bound  $\text{wd}(K_{d,d}) \geq d - 2\sqrt{d}$  for  $d \geq 2$ . Actually, this can be improved to  $d - \sqrt{2d} - 1$ :

**Proposition 1.9.** *If  $G$  is a triangle-free  $d$ -regular graph with  $n \geq 4$  vertices, then  $\text{wd}(G) > d - \sqrt{n} - 1$ . In particular, the complete bipartite graph  $K_{d,d}$  with  $d \geq 2$  satisfies  $\text{wd}(K_{d,d}) > d - \sqrt{2d} - 1$ .*

This should be contrasted with the fact that  $\chi(K_{d,d}) = 2$ ,  $\chi_\ell(K_{d,d}) = (1 + o(1)) \log_2 d$  [ERT79], and  $\chi_{DP}(K_{d,d}) = \Theta(d/\log d)$  [Ber16]. We prove Propositions 1.8 and 1.9 in §6.

It seems plausible that *every*  $d$ -regular graph has weak degeneracy at least  $d - O(\sqrt{d})$ ; we leave verifying or refuting this supposition as an open problem:

**Conjecture 1.10.** *Every  $d$ -regular graph  $G$  satisfies  $\text{wd}(G) \geq d - O(\sqrt{d})$ .*

In view of the above lower bounds, it makes sense to ask, for what classes of graphs  $G$  does the upper bound  $\text{wd}(G) \leq d - \Omega(\sqrt{d})$  hold, where  $d$  is the maximum degree of  $G$ ? Along these lines, we establish the following results:

**Theorem 1.11.** *For each integer  $k \geq 1$ , there exist  $c > 0$  and  $d_0 \in \mathbb{N}$  such that if  $G$  is a graph of maximum degree  $d \geq d_0$  with  $\chi(G) \leq k$ , then  $\text{wd}(G) \leq d - c\sqrt{d}$ .*

**Theorem 1.12.** *There exist  $c > 0$  and  $d_0 \in \mathbb{N}$  such that if  $G$  is a graph of maximum degree  $d \geq d_0$  and girth at least 5, then  $\text{wd}(G) \leq d - c\sqrt{d}$ .*

Theorems 1.11 and 1.12 are proved in §7 using probabilistic arguments.

We finish the introduction with another conjecture that implies both Theorems 1.11 and 1.12:

**Conjecture 1.13.** *For each integer  $k \geq 1$ , there exist  $c > 0$  and  $d_0 \in \mathbb{N}$  such that if  $G$  is a graph of maximum degree  $d \geq d_0$  and without a  $k$ -clique, then  $\text{wd}(G) \leq d - c\sqrt{d}$ .*

At present, we do not even know if Conjecture 1.13 holds for  $k = 3$ , i.e., whether  $\text{wd}(G) \leq d - \Omega(\sqrt{d})$  for triangle-free graphs of maximum degree  $d$ .

*Acknowledgments.*—We are very grateful to the anonymous referees for their comments.

## 2. Preliminary results

In this section we establish several basic results about weak degeneracy that will be used throughout the rest of this article.

**Lemma 2.1** (Weak degeneracy is monotone). *Let  $G$  be a weakly  $f$ -degenerate graph. If  $g: V(G) \rightarrow \mathbb{N}$  is a function such that  $g(u) \geq f(u)$  for all  $u \in V(G)$ , then  $G$  is weakly  $g$ -degenerate.*

PROOF. We wish to show that  $G$  is weakly  $g$ -degenerate by removing its vertices via the same sequence of operations that witnesses that  $G$  is weakly  $f$ -degenerate. The only possible issue is that an application of DELSAVE may become illegal when  $f$  is replaced by  $g$ . Namely, it can happen that  $\text{DELSAVE}(G, f, u, w)$  is legal, while  $\text{DELSAVE}(G, g, u, w)$  is not due to the fact that  $g(u) \leq g(w)$ . However, since  $f(u) > f(w)$ , this means that  $g(w) > f(w)$ , so instead of  $\text{DELSAVE}(G, g, u, w)$  we can simply use  $\text{DELETE}(G, g, u)$ : this replaces  $g(w)$  by  $g(w) - 1$ , which is still at least  $f(w)$ . ■

**Lemma 2.2** (Weak degeneracy and DP-coloring). *Let  $G$  be a weakly  $f$ -degenerate graph. Suppose that every vertex  $u \in V(G)$  is given a list  $L(u)$  of available colors and that for each edge  $uv \in E(G)$ , there is a matching  $C_{uv}$  from  $L(u)$  to  $L(v)$ . If  $|L(u)| \geq f(u) + 1$  for all  $u \in V(G)$ , then  $G$  admits a proper  $(L, C)$ -coloring.*

PROOF. This statement was essentially established in the introduction (in the discussion preceding Definition 1.2). We give a more detailed proof here for completeness. Since  $G$  is weakly  $f$ -degenerate, it is possible to remove all vertices from  $G$  by a sequence of legal applications of the operations DELETE and DELSAVE. Fix any such sequence  $\mathcal{S} = (\mathcal{O}_0, \dots, \mathcal{O}_{n-1})$ . Set  $(G_0, f_0) := (G, f)$  and for each  $0 \leq i \leq n-1$ , let  $(G_{i+1}, f_{i+1})$  be the result of applying the operation  $\mathcal{O}_i$  to  $(G_i, f_i)$ . We color the vertices of  $G$  one by one, in the order in which they are removed by  $\mathcal{S}$ . Each time a vertex  $u$  is assigned a color  $\alpha$ , we remove the colors corresponding to  $\alpha$  from the lists of colors available to the neighbors of  $u$ , thus ensuring that the resulting coloring is proper. Let  $L_i(u)$  be the list of colors available to a vertex  $u \in V(G_i)$  at the start of step  $i$  (in particular,  $L_0(u) := L(u)$ ). Throughout our coloring procedure, we will maintain the following property:

$$(P_i) \quad |L_i(u)| \geq f_i(u) + 1 \text{ for all } u \in V(G_i).$$

If we can achieve this, then we will successfully color the entire graph, since no uncolored vertex will ever run out of available colors. Now, property  $(P_0)$  holds by assumption. On step  $i$ , we assume that  $(P_i)$  holds and consider two cases.

**Case 1:**  $\mathcal{O}_i = \text{DELETE}(G_i, f_i, u_i)$ .

In this case we assign to  $u_i$  an arbitrary available color. It is clear that property  $(P_{i+1})$  holds regardless of what color is assigned to  $u_i$ .

**Case 2:**  $\mathcal{O}_i = \text{DELSAVE}(G_i, f_i, u_i, w_i)$ .

If  $|L(w_i)| > f_i(w_i) + 1$ , we can, as in Case 1, assign an arbitrary available color to  $u_i$ . Now suppose that  $|L(w_i)| = f_i(w_i) + 1$ . Then, by  $(P_i)$  and since this application of DELSAVE is legal, we have  $|L_i(u_i)| \geq f_i(u_i) + 1 > f_i(w_i) + 1 = |L_i(w_i)|$ . This means that  $u_i$  must have an available color  $\alpha_i \in L_i(u_i)$  that does not correspond to any color in  $L_i(w_i)$ . If we assign  $\alpha_i$  to  $u_i$ , then the list of available colors for  $w_i$  will not change, and thus  $(P_{i+1})$  will not be violated, as desired. ■

**Lemma 2.3** (Partitioning lemma). *Let  $G$  be a weakly  $f$ -degenerate graph. Suppose that functions  $f_1, f_2: V(G) \rightarrow \mathbb{Z}$  satisfy  $f_1(u) + f_2(u) = f(u) - 1$  for all  $u \in V(G)$ . Then there is a partition  $V(G) = V_1 \sqcup V_2$  such that the subgraph  $G[V_i]$  is weakly  $f_i$ -degenerate for each  $i \in \{1, 2\}$ .*

PROOF. The proof is by induction on  $|V(G)|$ . If  $V(G) = \emptyset$ , the statement holds vacuously. Now suppose that  $|V(G)| \geq 1$  and the claim holds for all graphs with  $|V(G)| - 1$  vertices. Since  $G$  is weakly  $f$ -degenerate, there is a legal application of an operation  $\mathcal{O} \in \{\text{DELETE}, \text{DELSAVE}\}$  that produces a pair  $(G', f')$  in which the graph  $G'$  is weakly  $f'$ -degenerate. We consider the two cases depending on whether  $\mathcal{O}$  is DELETE or DELSAVE.

**Case 1:**  $\mathcal{O} = \text{DELETE}(G, f, u)$ .

Then  $G' = G - u$ . Since  $f_1(u) + f_2(u) = f(u) - 1 \geq -1$ , we have  $f_1(u) \geq 0$  or  $f_2(u) \geq 0$ . For concreteness, say  $f_1(u) \geq 0$ . Define a function  $f'_1: V(G') \rightarrow \mathbb{Z}$  by

$$f'_1(v) := \begin{cases} f_1(v) - 1 & \text{if } uv \in E(G); \\ f_1(v) & \text{otherwise.} \end{cases}$$

Then  $f'_1 + f_2 = f' - 1$ , so, by the inductive hypothesis, there is a partition  $V(G') = V'_1 \sqcup V_2$  such that  $G[V'_1]$  is weakly  $f'_1$ -degenerate and  $G[V_2]$  is weakly  $f_2$ -degenerate. Set  $V_1 := V'_1 \sqcup \{u\}$ . We claim that the partition  $V(G) = V_1 \sqcup V_2$  is as desired. Since  $G[V_2]$  is weakly  $f_2$ -degenerate by assumption, we just need to argue that  $G[V_1]$  is weakly  $f_1$ -degenerate. As  $f_1(u) \geq 0$ , the function  $f_1$  is non-negative on  $V_1$ . Now we are done since  $\text{DELETE}(G[V_1], f_1, u) = (G[V'_1], f'_1)$  and  $G[V'_1]$  is weakly  $f'_1$ -degenerate.

**Case 2:**  $\mathcal{O} = \text{DELSAVE}(G, f, u, w)$ .

Again we have  $G' = G - u$ . It will be convenient to assume that  $f_1(w), f_2(w) \geq -1$ . If this is not the case and, say,  $f_1(w) < -1$ , then we replace  $f_1$  and  $f_2$  by the functions  $f_1^*, f_2^*: V(G) \rightarrow \mathbb{Z}$  given by  $f_1^*(w) := -1$ ,  $f_2^*(w) := f(w)$ , and  $f_i^*(v) := f_i(v)$  for all  $i \in \{1, 2\}$  and  $v \neq w$ . We can do this because every weakly  $f_i^*$ -degenerate subgraph of  $G$  is also weakly  $f_i$ -degenerate. For  $i = 2$  this follows from Lemma 2.1 since  $f_2 \geq f_2^*$  by definition. On the other hand, if a subgraph  $H$  of  $G$  is weakly  $f_1^*$ -degenerate, then  $w \notin V(H)$  since  $f_1^*(w) < 0$ . As  $f_1^*$  and  $f_1$  agree on all vertices except  $w$ ,  $H$  must be weakly  $f_1$ -degenerate as well.

Since this application of  $\text{DELSAVE}$  is legal, we have  $f(u) > f(w)$ , which implies that  $f_1(u) > f_1(w)$  or  $f_2(u) > f_2(w)$ . For concreteness, say  $f_1(u) > f_1(w)$ . Define  $f'_1: V(G') \rightarrow \mathbb{Z}$  by

$$f'_1(v) := \begin{cases} f_1(v) - 1 & \text{if } uv \in E(G) \text{ and } v \neq w; \\ f_1(v) & \text{otherwise.} \end{cases}$$

Then  $f'_1 + f_2 = f' - 1$ , so, by the inductive hypothesis, there is a partition  $V(G') = V'_1 \sqcup V_2$  such that  $G[V'_1]$  is weakly  $f'_1$ -degenerate and  $G[V_2]$  is weakly  $f_2$ -degenerate. Set  $V_1 := V'_1 \sqcup \{u\}$ . We claim that the partition  $V(G) = V_1 \sqcup V_2$  is as desired. We just need to argue that  $G[V_1]$  is weakly  $f_1$ -degenerate. Since  $f_1(u) > f_1(w) \geq -1$ , we have  $f_1(u) \geq 0$ . Hence,  $f_1$  is non-negative on  $V_1$ . It remains to observe that by a legal application of one of the operations  $\text{DELETE}$ ,  $\text{DELSAVE}$  it is possible to reduce the pair  $(G[V_1], f_1)$  to  $(G[V'_1], f'_1)$ . Indeed, if  $w \notin V_1$ , then  $(G[V'_1], f'_1) = \text{DELETE}(G[V_1], f_1, u)$ , while if  $w \in V_1$ , then  $(G[V'_1], f'_1) = \text{DELSAVE}(G[V_1], f_1, u, w)$ . ■

### 3. On-line DP-coloring and weak degeneracy

As mentioned in the introduction, DP-paint number is an on-line version of DP-chromatic number introduced by Kim, Kostochka, Li, and Zhu in [Kim+20]. It is defined by means of a certain game on a graph  $G$ :

**Definition 3.1** (DP-painting game). Let  $G$  be a graph and let  $g: V(G) \rightarrow \mathbb{N}$  be a function. The **DP-painting game on**  $(G, g)$  is played between two players—**Lister** and **Painter**—as follows. The game proceeds in **rounds**, starting with Round 0. At the start of Round  $i$ , we have a graph  $G_i$ , where we initially set  $G_0 := G$ . **Lister** then picks a list  $L_i(u)$  of colors for each vertex  $u \in V(G_i)$  and assigns to every edge  $uv \in E(G_i)$  a matching  $C_{i,uv}$  from  $L_i(u)$  to  $L_i(v)$  (the matching  $C_{i,uv}$  need not be perfect and, in particular, may be empty). In response, **Painter** picks a function  $\varphi_i$  defined on some subset  $U_i \subseteq V(G_i)$  with the following properties:

- $\varphi_i(u) \in L_i(u)$  for all  $u \in U_i$  (in particular,  $L_i(u) \neq \emptyset$  for all  $u \in U_i$ ); and
- $\varphi_i(u)\varphi_i(v) \notin C_{i,uv}$  for all  $u, v \in U_i$  that are adjacent in  $G_i$ .

Then we set  $G_{i+1} := G_i - U_i$  and proceed to Round  $i + 1$ . **Lister** wins the game if for some  $i \in \mathbb{N}$ , there is a vertex  $u \in V(G_i)$  with  $\sum_{j < i} |L_j(u)| \geq g(u)$ ; otherwise, **Painter** wins.

A graph  $G$  is  $g$ -**DP-paintable** if **Painter** has a winning strategy in the DP-painting game on  $(G, g)$ . Given  $k \in \mathbb{N}$ , we say that  $G$  is  $k$ -**DP-paintable** if it is DP-paintable with respect to the constant  $k$  function. The **DP-paint number**  $\chi_{\text{DPP}}(G)$  of  $G$  is the least  $k$  such that  $G$  is  $k$ -DP-paintable.

Take  $k \in \mathbb{N}$  and consider the DP-painting game on  $(G, \mathbf{k})$ , where  $\mathbf{k}$  is the constant  $k$  function. On Round 0, **Lister** may decide to give each vertex  $u \in V(G)$  a list  $L_0(u)$  of colors of size  $|L_0(u)| = k$ .

Then **Painter** must immediately assign a color to every vertex. Therefore, **Painter** can win only if  $\chi_{DP}(G) \leq k$ , which shows that  $\chi_{DP}(G) \leq \chi_{DPP}(G)$  for all  $G$ . On the other hand, if **Lister** always plays so that  $|L_i(u)| \leq 1$  for all  $i$  and  $u \in V(G_i)$ , then **Painter** can win if and only if  $\chi_P(G) \leq k$ , where  $\chi_P(G)$  is the classical *paint number* of  $G$ , i.e., the on-line analog of list-chromatic number (see [Kim+20, §2] for details). Thus,  $\chi_P(G) \leq \chi_{DPP}(G)$  as well, so  $\chi_{DPP}(G)$  provides a common upper bound on  $\chi_{DP}(G)$  and  $\chi_P(G)$ . It is shown in [Kim+20] that either inequality  $\chi_{DP}(G) \leq \chi_{DPP}(G)$  and  $\chi_P(G) \leq \chi_{DPP}(G)$  can be strict; however, it is unknown if both of them can be strict at the same time. It is also not known if the difference  $\chi_{DPP}(G) - \chi_{DP}(G)$  can be arbitrarily large.

The goal of this section is to prove Proposition 1.3, which says that the DP-paint number is bounded above by weak degeneracy plus 1. We prove it in the following stronger form:

**Proposition 1.3.** *If  $G$  is a weakly  $f$ -degenerate graph, then  $G$  is  $(f + 1)$ -DP-paintable.*

PROOF. The strategy for **Painter** is to pick functions  $\varphi_i$  so as to maintain the following property:

(P<sub>*i*</sub>)  $G_i$  is weakly  $f_i$ -degenerate, where  $f_i(u) := f(u) - \sum_{j < i} |L_j(u)|$  for all  $u \in V(G_i)$ .

If this can be achieved, then **Painter** will never lose, since for all  $u \in V(G_i)$ , we will have  $f_i(u) \geq 0$ , or, equivalently,  $\sum_{j < i} |L_j(u)| < f(u) + 1$ . Since  $f_0 = f$ , property (P<sub>0</sub>) holds by assumption, so it remains to argue that if (P<sub>*i*</sub>) holds at the start of Round  $i$ , then **Painter** will be able to pick  $\varphi_i$  so that (P<sub>*i+1*</sub>) holds.

Suppose **Lister** assigned a list  $L_i(u)$  of colors and a matching  $C_{i,uv}$  to each vertex  $u \in V(G_i)$  and edge  $uv \in E(G_i)$  respectively. For all  $u \in V(G_i)$ , let

$$f_{i,1}(u) := |L_i(u)| - 1 \quad \text{and} \quad f_{i,2}(u) := f_i(u) - |L_i(u)|.$$

Since  $f_{i,1}(u) + f_{i,2}(u) = f_i(u) - 1$  for all  $u \in V(G_i)$ , Lemma 2.3 yields a partition  $V(G_i) = U_i \sqcup W_i$  such that  $G[U_i]$  is weakly  $f_{i,1}$ -degenerate, while  $G[W_i]$  is weakly  $f_{i,2}$ -degenerate. By Lemma 2.2,  $G[U_i]$  admits a proper  $(L_i, C_i)$ -coloring  $\varphi_i$ . **Painter** plays this coloring  $\varphi_i$ . Then  $G_{i+1} = G[W_i]$ , so, to establish (P<sub>*i+1*</sub>), we need to show that  $G[W_i]$  is weakly  $f_{i+1}$ -degenerate. To this end, note that for each  $u \in W_i$ ,

$$f_{i+1}(u) = f(u) - \sum_{j \leq i} |L_j(u)| = f_i(u) - |L_i(u)| = f_{i,2}(u),$$

and  $G[W_i]$  is indeed weakly  $f_{i,2}$ -degenerate by construction. ■

## 4. Planar graphs are weakly 4-degenerate

In this section we prove the analog of Thomassen's theorem [Tho94] on 5-list-colorability of planar graphs in the context of weak degeneracy:

**Theorem 1.4.** *Every planar graph is weakly 4-degenerate.*

As in the proof of Thomassen's theorem, we use induction to establish a technical lemma, which then easily yields Theorem 1.4. First, we need a definition. Let  $G$  be a graph and let  $f: V(G) \rightarrow \mathbb{N}$  be a function. Given a subset  $U \subseteq V(G)$ , we say that  $G$  is  **$U$ -safely weakly  $f$ -degenerate** if, starting with  $(G, f)$ , it is possible to remove all vertices from  $G$  by a sequence of legal applications of the operations DELETE and DELSAVE, so that every vertex in  $U$  is removed using the DELETE operation. In particular,  $G$  is  $V(G)$ -safely weakly  $f$ -degenerate if and only if  $G$  is  $f$ -degenerate.

**Lemma 4.1.** *Let  $G$  be a planar graph on at least 3 vertices where every non-outer face is triangular and the outer face is a cycle  $C$  of length  $k$ . Let the vertices of  $C$  in the natural order be  $v_1, \dots, v_k$ . Define  $f: V(G) \setminus \{v_1, v_2\} \rightarrow \mathbb{Z}$  by*

$$f(u) := \begin{cases} 2 - |N_G(u) \cap \{v_1, v_2\}| & \text{if } u \in V(C); \\ 4 - |N_G(u) \cap \{v_1, v_2\}| & \text{otherwise.} \end{cases}$$

Then  $G - v_1 - v_2$  is  $(V(C) \setminus \{v_1, v_2\})$ -safely weakly  $f$ -degenerate.

PROOF. We proceed by induction on  $|V(G)|$ . If  $|V(G)| = 3$ , then  $G - v_1 - v_2$  comprises a single vertex, which is 0-degenerate, as desired. Now suppose that  $|V(G)| \geq 4$  and that the induction hypothesis holds for smaller graphs. We consider two cases.

**Case 1:**  $C$  has a chord  $v_a v_b$ .

Then  $C + v_a v_b$  is the union of two cycles  $C_1, C_2$  with  $E(C_1) \cap E(C_2) = \{v_a v_b\}$ . Without loss of generality, suppose  $v_1 v_2 \in E(C_1)$  (and so  $v_1 v_2 \notin E(C_2)$ ). Let  $G_1, G_2$  be the respective induced subgraphs of  $G$  on the vertices of each  $C_i$  along with the vertices on the interiors of each cycle. Let  $f_1 := f|_{V(G_1) \setminus \{v_1, v_2\}}$  and define  $f_2: V(G_2) \setminus \{v_1, v_2\} \rightarrow \mathbb{Z}$  by

$$f_2(u) := \begin{cases} 2 - |N_G(u) \cap \{v_a, v_b\}| & \text{if } u \in V(C_2); \\ 4 - |N_G(u) \cap \{v_a, v_b\}| & \text{otherwise.} \end{cases}$$

By the induction hypothesis, starting with  $(G_1 - v_1 - v_2, f_1)$ , we can remove all vertices from  $V(G_1) \setminus \{v_1, v_2\}$  via legal applications of the operations DELETE and DELSAVE, where each vertex in  $V(C_1) \setminus \{v_1, v_2\}$  is removed using DELETE. Applying the same sequence of operations but starting with  $(G - v_1 - v_2, f)$  yields the pair  $(G_2 - v_a - v_b, f_2)$ . By the inductive hypothesis again, we can now remove every remaining vertex via a sequence of legal applications of DELETE and DELSAVE, with every vertex in  $V(C_2) \setminus \{v_a, v_b\}$  removed using DELETE, as desired.

**Case 2:**  $C$  has no chord.

Since every non-outer face of  $G$  is a triangle, the neighbors of  $v_k$  form a path  $u_1 \dots u_\ell$ , where  $u_1 = v_1$  and  $u_\ell = v_{k-1}$ . The assumption that  $C$  has no chord implies that  $u_2, \dots, u_{\ell-1}$  belong to the interior of  $C$ . Then the cycle  $C' := u_1 \dots u_\ell v_{k-2} \dots v_1$  bounds the outer face of  $G' := G - v_k$ . Applying the induction hypothesis to  $G'$  shows that  $G'' := G - v_1 - v_2 - v_k$  is  $(V(C') \setminus \{v_1, v_2\})$ -safely weakly  $f'$ -degenerate, where  $f': V(G'') \rightarrow \mathbb{Z}$  is defined by

$$f'(u) := \begin{cases} 2 - |N_{G'}(u) \cap \{v_1, v_2\}| & \text{if } u \in V(C'); \\ 4 - |N_{G'}(u) \cap \{v_1, v_2\}| & \text{otherwise.} \end{cases}$$

In other words, starting with  $(G'', f')$ , we can remove every vertex by a sequence of legal applications of DELETE and DELSAVE, where each vertex in  $V(C') \setminus \{v_1, v_2\}$  is removed using DELETE. Since  $f' \leq f$ , we may apply the same sequence of operations starting with  $(G - v_1 - v_2, f)$  instead (see Lemma 2.1). Moreover, we can accrue some extra savings for the vertex  $v_k$ , as follows. Consider any  $u_i$  with  $2 \leq i \leq \ell - 1$ . By assumption,  $u_i$  is removed from  $G''$  using the DELETE operation, but since  $u_i \notin V(C)$ , we are now allowed to remove  $u_i$  using DELSAVE. Notice that  $f'(u_i) = f(u_i) - 2$ , because  $u_i$  is in  $V(C')$  but not in  $V(C)$ . When  $u_i$  was removed from  $G''$ , the value of the function at  $u_i$  was at least 0, which means that at the same stage of the process on  $G - v_1 - v_2$ , the value of the function at  $u_i$  is at least 2. On the other hand, since  $v_k \in V(C)$  and  $v_1 \in N_G(v_k)$ , we have  $f(v_k) \leq 1 < 2$ . This means that instead of using the operation DELETE, we can legally remove  $u_i$  using DELSAVE( $\cdot, \cdot, u_i, v_k$ ). Upon performing this modified sequence of operations, we only have  $v_k$  left to remove, so we just need to check that the value of the function at  $v_k$  is at least 0. To this end, note that  $f(v_k)$  is 1 if  $v_{k-1} \neq v_2$  and 0 otherwise. Since the only neighbor of  $v_k$  that may be removed without saving  $v_k$  is  $v_{k-1}$ , and that can only happen when  $v_{k-1} \neq v_2$ , it follows that the value at  $v_k$  cannot drop below 0, as desired. ■

We now complete the proof of the theorem.

PROOF of Theorem 1.4. Since adding vertices or edges cannot decrease the weak degeneracy of a graph, it suffices to prove the theorem for maximal planar graphs  $G$  on at least 3 vertices. Then  $G$  is a planar triangulation. Let  $v_1, v_2$  be adjacent vertices on its outer face. Removing  $v_1$  and  $v_2$  using DELETE and then applying Lemma 4.1, we see that  $G$  is weakly 4-degenerate, as desired. ■



## 5. Brooks-type results

### 5.1. Weakly $(\deg - 1)$ -degenerate graphs

We say that a graph  $G$  is **weakly  $(\deg - 1)$ -degenerate** if it is weakly degenerate with respect to the function  $u \mapsto \deg_G(u) - 1$ . Recall that a **GDP tree** is a connected graph in which every block is either a clique or a cycle. The main result of this section is the following characterization of connected weakly  $(\deg - 1)$ -degenerate graphs:

**Theorem 1.6.** *Let  $G$  be a connected graph. The following statements are equivalent:*

- (1)  $G$  is weakly  $(\deg - 1)$ -degenerate;
- (2)  $G$  is not a GDP-tree.

To begin with, we need the following standard fact:

**Lemma 5.1.** *Let  $G$  be a connected graph and let  $f: V(G) \rightarrow \mathbb{N}$  be a function. Suppose that:*

- (a)  $f(u) \geq \deg_G(u) - 1$  for all  $u \in V(G)$ ; and
- (b)  $f(x) \geq \deg_G(x)$  for some  $x \in V(G)$ .

*Then  $G$  is  $f$ -degenerate.*

PROOF. Fix a vertex  $x$  witnessing (b) and list the vertices of  $G$  as  $u_1, u_2, \dots, u_n$  in order of decreasing distance to  $x$ , resolving ties arbitrarily. Then  $u_n = x$  and, for each  $1 \leq i \leq n - 1$ , the vertex  $u_i$  has at least one neighbor among  $u_{i+1}, \dots, u_n$ . We can now remove all vertices from  $G$  by applying the operation DELETE to them in this order. ■

The next lemma contains the central part of our argument:

**Lemma 5.2.** *Let  $G$  be a connected graph that is not weakly  $(\deg - 1)$ -degenerate. Then every connected induced subgraph of  $G$  without cut vertices is regular.*

PROOF. Take a subset  $A \subseteq V(G)$  such that the subgraph  $G[A]$  has no cut vertices and suppose, toward a contradiction, that  $G[A]$  is not regular. Define  $f: V(G) \rightarrow \mathbb{N}$  by  $f(u) := \deg_G(u) - 1$  for all  $u \in V(G)$ . Our goal is to show that  $G$  is weakly  $f$ -degenerate. Note that every connected component of  $G - A$  contains at least one vertex  $v$  that has a neighbor in  $A$  and hence satisfies  $f(v) \geq \deg_{G-A}(v)$ . Therefore, by Lemma 5.1, we can remove all vertices from  $G - A$  using only the operation DELETE. After this, the graph  $G$  will be replaced by  $G' := G[A]$  and the function  $f$  by  $f': A \rightarrow \mathbb{N}: u \mapsto \deg_{G[A]}(u) - 1$ . Since the graph  $G[A]$  is connected and not regular, we can pick two adjacent vertices  $x, y \in A$  with  $\deg_{G[A]}(x) < \deg_{G[A]}(y)$  and hence  $f'(x) < f'(y)$ . Now we let

$$(G'', f'') := \text{DELSAVE}(G[A], f', y, x).$$

Since  $f'(y) > f'(x)$ , this is a legal application of DELSAVE. As the graph  $G[A]$  has no cut vertices, the graph  $G'' = G[A] - y$  is connected. It remains to observe that  $G''$  is  $f''$ -degenerate by Lemma 5.1, where condition (b) is witnessed by the vertex  $x$ . ■

It remains to characterize the graphs satisfying the conclusion of Lemma 5.2:

**Lemma 5.3.** *Let  $G$  be a connected graph such that every connected induced subgraph of  $G$  without cut vertices is regular. Then  $G$  is a GDP-tree.*

PROOF. Suppose, toward a contradiction, that  $G$  is a counterexample with the fewest vertices. Note that  $|V(G)| \geq 4$ , since all connected graphs on at most 3 vertices are GDP-trees. By the minimality of  $|V(G)|$ , every proper connected induced subgraph of  $G$  must be a GDP-tree.

We claim that  $G$  is 2-connected. Otherwise, every block in  $G$  would be a proper connected induced subgraph of  $G$ , hence a GDP-tree. The only GDP-trees without cut vertices are cliques and cycles, so this implies that every block in  $G$  is a clique or a cycle, i.e.,  $G$  is a GDP-tree.

Since  $G$  is 2-connected, it must be regular. Let  $d$  be the common degree of the vertices of  $G$ . Then  $d \geq 2$  by 2-connectedness. Furthermore, if  $d$  were equal to 2, then  $G$  would be a cycle and hence a GDP-tree. Therefore,  $d \geq 3$ .

Pick an arbitrary vertex  $x \in V(G)$  and consider the graph  $G' := G - x$ . Then  $G'$  is connected, so it is a GDP-tree. Since  $G$  is regular and not a clique, not every vertex in  $V(G')$  is adjacent to  $x$ . This implies that  $G'$  is not regular, so it must have a cut vertex and at least two blocks.

Let  $B$  be an arbitrary leaf block in  $G'$  and let  $c \in V(B)$  be the cut vertex of  $G'$  in  $B$ . The graph  $B$  is regular, so let  $k$  be the common degree of every vertex of  $B$ . The degree of a vertex  $u \in V(B) \setminus \{c\}$  in  $G$  is either  $k + 1$  or  $k$ , depending on whether  $u$  is adjacent to  $x$  or not. Since  $G$  is 2-connected,  $x$  must be adjacent to at least one vertex in  $V(B) \setminus \{c\}$ , which, since  $G$  is  $d$ -regular, implies that  $k + 1 = d$  and  $x$  is in fact adjacent to every vertex in  $V(B) \setminus \{c\}$ . Hence,  $x$  has at least  $|V(B)| - 1 \geq k = d - 1$  neighbors in  $B$ .

Finally, as there are at least 2 distinct leaf blocks in  $G'$ , we conclude that  $x$  has at least  $2(d - 1)$  neighbors. Therefore,  $d \geq 2(d - 2)$ , i.e.,  $d \leq 2$ , which is a contradiction. ■

Theorem 1.6 now follows easily:

**PROOF of Theorem 1.6.** The implication (2)  $\implies$  (1) is a combination of Lemmas 5.2 and 5.3. The implication (1)  $\implies$  (2) follows since GDP-trees are not DP-degree-colorable [BKP17, Theorem 9]. That is, if  $G$  is a GDP-tree, then it is possible to give each vertex  $u \in V(G)$  a list  $L(u)$  of available colors of size  $|L(u)| \geq \deg_G(u)$  and assign to each edge  $uv \in E(G)$  a matching  $C_{uv}$  from  $L(u)$  to  $L(v)$  so that  $G$  does not admit a proper  $(L, C)$ -coloring. By Lemma 2.2, this implies that  $G$  is not weakly  $(\deg - 1)$ -degenerate. ■

## 5.2. Weak degeneracy and maximum average degree

Here we establish a lower bound on the maximum average degree of a graph in terms of its weak degeneracy:

**Theorem 1.7.** *Let  $G$  be a nonempty graph. If the weak degeneracy of  $G$  is at least  $d \geq 3$ , then either  $G$  contains a  $(d + 1)$ -clique or*

$$\text{mad}(G) \geq d + \frac{d - 2}{d^2 + 2d - 2}.$$

We derive Theorem 1.7 from Theorem 1.6. Our argument is closely analogous to the proof of the lower bound on the average degree of DP-critical graphs due to Kostochka, Pron, and the first named author [BKP17, Corollary 10], which in turn is based on earlier work of Gallai [Gal63].

We need the following result, essentially established by Gallai in [Gal63] (Gallai's paper is in German; see [BKP17, Appendix] for a proof in English):

**Lemma 5.4** ([BKP17, Lemma 20]). *Let  $T$  be a GDP-tree of maximum degree at most  $d \geq 3$  and without a  $(d + 1)$ -clique. Then  $\text{ad}(T) \leq d - 1 + 2/d$ .*

We say that  $G$  is a **minimal** graph of weak degeneracy  $d$  if  $\text{wd}(G) = d$  and  $\text{wd}(H) < d$  for every proper subgraph  $H$  of  $G$ .

**Lemma 5.5.** *Let  $G$  be a minimal graph of weak degeneracy  $d \geq 3$ .*

- (a) *The minimum degree of  $G$  is at least  $d$ .*
- (b) *Let  $U := \{u \in V(G) : \deg_G(u) = d\}$ . Then every component of  $G[U]$  is a GDP-tree.*

**PROOF.** (a) Suppose that there is a vertex  $u \in V(G)$  with  $\deg_G(u) \leq d - 1$ . We will show that  $G$  is weakly  $(d - 1)$ -degenerate. Let  $f$  be the constant  $d - 1$  function on  $V(G)$ . By the minimality of  $G$ , we may remove every vertex from  $(G - u, f)$  via a sequence of legal applications of the operations DELETE and DELSAVE. Since  $\deg_G(u) \leq d - 1$ , we may use the same sequence of operations to

remove every vertex except  $u$  from  $(G, f)$  (at which point the function  $f$  will be replaced by the map sending  $u$  to  $d - 1 - \deg_G(u)$ ) and then remove  $u$  using the operation **DELETE**.

(b) Let  $C$  be a connected component of  $G[U]$  and let  $f$  be the constant  $d - 1$  function on  $V(G)$ . By the minimality of  $G$ , we may remove every vertex from  $(G - V(C), f)$  via a sequence of legal applications of the operations **DELETE** and **DELSAVE**. If we perform the same sequence of operations on  $(G, f)$ , then the graph  $G$  will be replaced by  $C$ , while the function  $f$  will be replaced by the map sending each  $u \in V(C)$  to  $d - 1 - \deg_{G-V(C)}(u) = \deg_C(u) - 1$ . Since  $G$  is not weakly  $(d - 1)$ -degenerate, this implies that  $C$  is not weakly  $(\deg - 1)$ -degenerate. Hence, by Theorem 1.6,  $C$  is a GDP-tree.  $\blacksquare$

**PROOF of Theorem 1.7.** Fix  $d \geq 3$ . It suffices to argue that every minimal graph  $G$  of weak degeneracy  $d$  and without a  $(d + 1)$ -clique satisfies

$$\text{ad}(G) \geq d + \frac{d - 2}{d^2 + 2d - 2}.$$

To this end, we use discharging. Let the initial charge of each vertex  $u \in V(G)$  be  $\text{ch}(u) := \deg_G(u)$ . The only discharging rule is: Every vertex  $u \in V(G)$  with  $\deg_G(u) > d$  sends to each neighbor the charge  $c := d/(d^2 + 2d - 2)$ . Let the new charge of each vertex  $u$  be  $\text{ch}^*(u)$ . Note that

$$\text{ad}(G)|V(G)| = \sum_{u \in V(G)} \text{ch}(u) = \sum_{u \in V(G)} \text{ch}^*(u).$$

For any vertex  $u$  with  $\deg_G(u) > d$ , we have

$$\text{ch}^*(u) \geq \deg_G(u) - c \deg_G(u) \geq \left(1 - \frac{d}{d^2 + 2d - 2}\right)(d + 1) = d + \frac{d - 2}{d^2 + 2d - 2}.$$

Let  $C$  be any connected component of  $G[U]$ , where  $U$  is the set of all vertices of degree  $d$  in  $G$ . By Lemma 5.5(b),  $C$  is a GDP-tree. Hence, by Lemma 5.4,  $\text{ad}(C) \leq d - 1 + 2/d$ . Therefore,

$$\begin{aligned} \sum_{u \in V(C)} \text{ch}^*(u) &= d|V(C)| + c \sum_{u \in V(C)} (d - \deg_C(u)) \\ &\geq \left(d + \frac{d}{d^2 + 2d - 2} \left(1 - \frac{2}{d}\right)\right) |V(C)| = \left(d + \frac{d - 2}{d^2 + 2d - 2}\right) |V(C)|. \end{aligned}$$

The above bounds imply that

$$\sum_{u \in V(G)} \text{ch}^*(u) \geq \left(d + \frac{d - 2}{d^2 + 2d - 2}\right) |V(G)|,$$

which yields the desired result.  $\blacksquare$

## 6. Lower bounds for regular graphs

In this section we establish lower bounds on weak degeneracy for regular graphs.

**Proposition 1.8.** *Let  $G$  be a  $d$ -regular graph with  $n \geq 2$  vertices. Then  $\text{wd}(G) \geq d - \sqrt{2n}$ .*

**PROOF.** Let  $k := \text{wd}(G)$ . Set  $G_0 := G$  and let  $f_0$  be the constant  $k$  function on  $V(G)$ . By definition, starting with  $(G_0, f_0)$ , it is possible to remove all vertices from  $G$  via a sequence of legal applications of the operations **DELETE** and **DELSAVE**. Fix any such sequence  $\mathcal{S} = (\mathcal{O}_0, \dots, \mathcal{O}_{n-1})$ . For each  $0 \leq i \leq n - 1$ , let  $(G_{i+1}, f_{i+1})$  be the result of applying  $\mathcal{O}_i$  to  $(G_i, f_i)$ . Then we can write

$$\mathcal{O}_i = \text{DELETE}(G_i, f_i, u_i) \quad \text{or} \quad \mathcal{O}_i = \text{DELSAVE}(G_i, f_i, u_i, w_i).$$

For each  $0 \leq i \leq n - 1$ , define

$$d_i := |\{j < i : u_j u_i \in E(G)\}| \quad \text{and} \quad \sigma_i := |\{j < i : \mathcal{O}_j = \text{DELSAVE}(G_j, f_j, u_j, u_i)\}|.$$

(So  $\sigma_i$  is the number of vertices that “save”  $u_i$ .) Then  $0 \leq f_i(u_i) = k - d_i + \sigma_i$  and thus

$$k \geq d_i - \sigma_i. \quad (6.1)$$

Adding (6.1) up over the interval  $n - t \leq i \leq n - 1$  for some integer  $1 \leq t \leq n$  yields

$$kt \geq \left( \sum_{i=n-t}^{n-1} d_i \right) - \left( \sum_{i=n-t}^{n-1} \sigma_i \right). \quad (6.2)$$

Each index  $j$  contributes to  $\sigma_i$  for at most one  $i$ , so  $\sum_{i=0}^{n-1} \sigma_i \leq n$ . Also, since  $G$  is  $d$ -regular,

$$\sum_{i=n-t}^{n-1} d_i = dt - |E(G[u_{n-t}, \dots, u_{n-1}])| \geq dt - \binom{t}{2}. \quad (6.3)$$

Therefore, (6.2) implies that

$$k \geq d - \frac{t-1}{2} - \frac{n}{t}.$$

Finally, taking  $t := \lceil \sqrt{2n} \rceil$  gives

$$k \geq d - \frac{\lceil \sqrt{2n} \rceil - 1}{2} - \frac{n}{\lceil \sqrt{2n} \rceil} \geq d - \frac{\sqrt{2n}}{2} - \frac{n}{\sqrt{2n}} = d - \sqrt{2n},$$

as desired. ■

**Proposition 1.9.** *If  $G$  is a triangle-free  $d$ -regular graph with  $n \geq 4$  vertices, then  $\text{wd}(G) > d - \sqrt{n} - 1$ .*

PROOF. The argument is virtually the same as in the proof of Proposition 1.8, except that we use Mantel’s theorem to replace the bound (6.3) by  $\sum_{i=n-t}^{n-1} d_i \geq dt - t^2/4$ . This yields

$$\text{wd}(G) \geq d - \frac{t}{4} - \frac{n}{t}.$$

Now we set  $t := \lceil 2\sqrt{n} \rceil$  and conclude that

$$\text{wd}(G) \geq d - \frac{\lceil 2\sqrt{n} \rceil}{4} - \frac{n}{\lceil 2\sqrt{n} \rceil} \geq d - \frac{2\sqrt{n} + 1}{4} - \frac{n}{2\sqrt{n}} \geq d - \sqrt{n} - \frac{1}{4},$$

as desired. ■

## 7. Going below the maximum degree

### 7.1. Preliminaries

In this section we review some necessary background facts. First, we will need the Lovász Local Lemma, in the following form:

**Theorem 7.1** (Lovász Local Lemma [AS16, Lemma 5.1.1]). *Let  $\mathcal{X}$  be a finite family of random events such that each  $X \in \mathcal{X}$  has probability at most  $p$  and is mutually independent from all but  $\Delta$  other events in  $\mathcal{X}$ . If  $ep(\Delta + 1) \leq 1$ , then the probability that no event in  $\mathcal{X}$  happens is positive.*

We shall also use the Chernoff bound for binomial random variables:

**Theorem 7.2** (Chernoff bound [MR02, p. 43]). *If  $X \sim \text{Bin}(n, p)$  is a binomial random variable, then for all  $0 \leq t \leq np$ ,*

$$\mathbb{P}[|X - np| > t] < 2 \exp\left(-\frac{t^2}{3np}\right).$$

Next, we need a quantitative version of the Central Limit Theorem due to Berry and Esseen:

**Theorem 7.3** (Berry–Esseen [Fel72, §XVI.5]). *There is a universal constant  $A > 0$  with the following property. Let  $X_1, \dots, X_n$  be independent identically distributed random variables such that  $\mathbb{E}[X_i] = 0$ ,  $\mathbb{E}[X_i^2] = \sigma^2 > 0$ , and  $\mathbb{E}[|X_i|^3] = \rho < \infty$ . Then for all  $t \in \mathbb{R}$ ,*

$$\left| \mathbb{P} \left[ \frac{\sum_{i=1}^n X_i}{\sigma \sqrt{n}} \leq t \right] - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-x^2/2} dx \right| \leq \frac{A\rho}{\sigma^3 \sqrt{n}}.$$

In particular, if  $X \sim \text{Bin}(n, p)$  is a binomial random variable, then for any  $\beta > 0$ ,

$$\mathbb{P}[X \leq np - \beta \sqrt{n}] = \int_{-\infty}^{\beta/\sqrt{p(1-p)}} e^{-x^2/2} dx + O \left( \frac{1 - 2p(1-p)}{\sqrt{p(1-p)n}} \right) \quad (7.4)$$

This means that for large  $n$ ,  $\mathbb{P}[X \leq np - \beta \sqrt{n}]$  is separated from 0. By applying this result to the random variable  $n - X \sim \text{Bin}(n, 1 - p)$ , we see that  $\mathbb{P}[X \geq np + \beta \sqrt{n}]$  is separated from 0 as well.

The following is a standard consequence of Hall’s theorem:

**Lemma 7.5.** *Let  $G$  be a graph and let  $A, B \subseteq V(G)$  be disjoint sets. Suppose that each vertex in  $A$  has at most  $d_1$  neighbors in  $B$ , while each vertex in  $B$  has at least  $d_2$  neighbors in  $A$ . Let  $t \in \mathbb{N}$  satisfy  $td_1 \leq d_2$ . Then there exists a partial function  $s : A \dashrightarrow B$  such that:*

- for all  $u \in A$ , if  $s(u)$  is defined, then  $s(u)$  is a neighbor of  $u$ ;
- the preimage of every vertex  $w \in B$  under  $s$  has cardinality exactly  $t$ .

**PROOF.** Let  $H$  be the maximal bipartite subgraph of  $G$  with parts  $A$  and  $B$ , and let  $H^*$  be obtained from  $H$  by replacing every vertex  $w \in B$  by  $t$  copies, denoted  $w_1, \dots, w_t$ . By construction,  $H^*$  is a bipartite graph with parts  $A$  and  $B^* := \{w_j : w \in B, 1 \leq j \leq t\}$ . For all  $u \in A$ ,  $\deg_{H^*}(u) \leq td_1 \leq d_2$ . On the other hand, every vertex  $w_j \in B^*$  satisfies  $\deg_{H^*}(w_j) \geq d_2$ . These inequalities, together with Hall’s theorem [Die17, Theorem 2.1.2], imply that  $H^*$  has a matching  $M$  that saturates  $B^*$ . We can now define the desired function  $s : A \dashrightarrow B$  by mapping each  $u \in A$  that is covered by  $M$  to the unique  $w \in B$  such that  $uw_j \in M$  for some  $j$ . ■

It will be convenient for us to work with  $d$ -regular graphs rather than graphs of maximum degree  $d$ . To this end, we shall employ the following facts:

**Lemma 7.6** (Chartrand–Wall [CW75]). *If  $G$  is a graph of maximum degree  $d$  and chromatic number at most  $k$ , then  $G$  can be embedded into a  $d$ -regular graph  $G^*$  of chromatic number at most  $k$ .*

**Lemma 7.7** ([MR02, Exercise 12.4]). *If  $G$  is a graph of maximum degree  $d$  and girth at least  $g$ , then  $G$  can be embedded into a  $d$ -regular graph  $G^*$  of girth at least  $g$ .*

**PROOF.** This fact is well-known, but we include a proof for completeness. We use a simplified version of the construction from [ABD21, Proposition 4.1]. Set

$$N := \sum_{u \in V(G)} (d - \deg_G(u))$$

and let  $\Gamma$  be an  $N$ -regular graph of girth at least  $g$ , which exists by [Imr84; Mar82]. We may assume that  $V(\Gamma) = \{1, \dots, q\}$ , where  $q := |V(\Gamma)|$ . Take  $q$  vertex-disjoint copies of  $G$ , say  $G_1, \dots, G_q$  and define  $S_i := \{u \in V(G_i) : \deg_{G_i}(u) < d\}$  for every  $1 \leq i \leq q$ . The graph  $G^*$  is obtained from the disjoint union of  $G_1, \dots, G_q$  by performing the following sequence of operations once for each edge  $ij \in E(\Gamma)$ , one edge at a time:

- (1) Pick arbitrary vertices  $u \in S_i$  and  $v \in S_j$ .
- (2) Add the edge  $uv$  to  $E(G^*)$ .
- (3) If  $\deg_{G^*}(u) = d$ , remove  $u$  from  $S_i$ .
- (4) If  $\deg_{G^*}(v) = d$ , remove  $v$  from  $S_j$ .

It is clear that the resulting graph  $G^*$  is as desired. ■

## 7.2. Removal schemes

In the next definition we introduce the technical notion of a *removal scheme* on a graph  $G$ . Roughly speaking, a removal scheme records the order in which we attempt to remove the vertices from  $G$ . Additionally, it indicates whether each vertex is removed using a `DELETE` or a `DELSAVE` operation, and in the latter case, what other vertex we “save” an extra color for.

**Definition 7.8** (Removal schemes). Fix a graph  $G$ . A **removal scheme** on  $G$  is a pair  $(<, \text{save})$ , where  $<$  is a linear ordering on  $V(G)$  and  $\text{save}: V(G) \dashrightarrow V(G)$  is a partial function such that for every vertex  $u \in V(G)$ , if  $\text{save}(u)$  is defined, then it is a neighbor of  $u$  and  $u < \text{save}(u)$ . For convenience, we write  $\text{save}(u) = \text{blank}$  if  $\text{save}(u)$  is undefined. Given a removal scheme  $(<, \text{save})$ , we call  $<$  the **removal order** and say that a vertex  $u$  **saves** the vertex  $\text{save}(u)$ . A removal scheme  $(<, \text{save})$  is **legal** if for all  $u, w \in V(G)$  such that  $w = \text{save}(u)$ , we have

$$|\{v \in N_G(u) : v < u \text{ and } \text{save}(v) \neq u\}| < |\{v \in N_G(w) : v < u \text{ and } \text{save}(v) \neq w\}|. \quad (7.9)$$

The **gap** of a vertex  $u$  with respect to a removal scheme  $(<, \text{save})$  is the quantity

$$\text{gap}(u; <, \text{save}) := |\{v \in N_G(u) : v > u\}| + |\{v \in N_G(u) : \text{save}(v) = u\}|.$$

We also let  $\text{gap}(<, \text{save}) := \min\{\text{gap}(u; <, \text{save}) : u \in V(G)\}$ .

**Lemma 7.10.** *Let  $G$  be a graph of maximum degree at most  $d$  and let  $(<, \text{save})$  be a legal removal scheme on  $G$ . Then  $G$  is weakly  $(d - \text{gap}(<, \text{save}))$ -degenerate.*

PROOF. For brevity, let  $g := \text{gap}(<, \text{save})$ . Let  $u_0, \dots, u_{n-1}$  be the vertices of  $G$  listed in the order given by  $<$ . Define a sequence  $(G_i, f_i)$ ,  $0 \leq i \leq n-1$  by setting  $(G_0, f_0) := (G, d - g)$  and

$$(G_{i+1}, f_{i+1}) := \begin{cases} \text{DELETE}(G_i, f_i, u_i) & \text{if } \text{save}(u_i) = \text{blank}; \\ \text{DELSAVE}(G_i, f_i, u_i, \text{save}(u_i)) & \text{otherwise.} \end{cases}$$

We claim that this construction yields a sequence of legal applications of `DELETE` and `DELSAVE` that removes every vertex from  $G$ . Indeed, consider any vertex  $u_i$ . Note that

$$f_i(u_i) = d - g - |\{v \in N_G(u_i) : v < u_i \text{ and } \text{save}(v) \neq u_i\}| \geq \text{gap}(u_i; <, \text{save}) - g \geq 0.$$

This shows that the functions  $f_i$  are non-negative. Now suppose that  $\text{save}(u_i) = w \in V(G)$ . Then, by definition,  $w$  is a neighbor of  $u_i$  that appears after  $u_i$  in the ordering  $<$ , and thus the operation  $\text{DELSAVE}(G_i, f_i, u_i, w)$  may be applied. Furthermore,  $f_i(u_i) > f_i(w)$  by (7.9), so this application of `DELSAVE` is legal, as desired. ■

## 7.3. Regular sets

Let  $G$  be a graph. Given a vertex  $u \in V(G)$  and a set  $A \subseteq V(G)$ , we let  $N_A(u) := N_G(u) \cap A$  denote the set of all neighbors of  $u$  in  $A$  and write  $\deg_A(u) := |N_A(u)|$ . Several times in our arguments, we will need to perform the following operation: given a set  $A$  and a number  $p \in [0, 1]$ , we will need to pick a subset  $A' \subseteq A$  such that every vertex  $u \in V(G)$  has roughly  $p \deg_A(u)$  neighbors in  $A'$ . Formally, we introduce the following definition:

**Definition 7.11** (Regular sets). Fix a graph  $G$  of maximum degree  $d$  and a subset  $A \subseteq V(G)$ . Given  $p, \varepsilon \in (0, 1]$ , a  $(p, \varepsilon)$ -**regular subset** of  $A$  is a set  $A' \subseteq A$  such that every vertex  $u \in V(G)$  satisfies one of the following conditions:

- either  $\deg_A(u) < 9 \log d / (\varepsilon^2 p)$  (i.e.,  $u$  has very few neighbors in  $A$ ),
- or  $(1 - \varepsilon)p \deg_A(u) \leq \deg_{A'}(u) \leq (1 + \varepsilon)p \deg_A(u)$  (i.e.,  $\deg_{A'}(u) \approx p \deg_A(u)$ ).

Using the Lovász Local Lemma, it is not hard to prove that  $(p, \varepsilon)$ -regular subsets exist:

**Lemma 7.12.** *Let  $G$  be a graph of maximum degree  $d$ . Fix  $p, \varepsilon \in (0, 1]$ . Then every set  $A \subseteq V(G)$  has a  $(p, \varepsilon)$ -regular subset.*

PROOF. We may assume  $d > 8$ , as otherwise  $\deg_A(u) \leq d < 9 \log d$  for all  $u \in V(G)$ , so any subset  $A' \subseteq A$  is  $(p, \varepsilon)$ -regular. Form a random set  $A' \subseteq A$  by picking each vertex independently with probability  $p$ . We shall use the Lovász Local Lemma (Theorem 7.1) to argue that  $A'$  is  $(p, \varepsilon)$ -regular with positive probability. Let  $U \subseteq V(G)$  be the set of all vertices  $u \in V(G)$  with  $\deg_A(u) \geq 9 \log d / (\varepsilon^2 p)$ . For each  $u \in U$ , let  $X_u$  be the random event that

$$\deg_{A'}(u) \notin [(1 - \varepsilon)p \deg_A(u), (1 + \varepsilon)p \deg_A(u)].$$

We need to argue that with positive probability, none of the events  $X_u$  happen. By the Chernoff bound (Theorem 7.2), for each  $u \in U$  we have

$$\mathbb{P}[X_u] < 2 \exp\left(-\frac{\varepsilon^2 p \deg_A(u)}{3}\right) \leq 2 \exp(-3 \log d) = 3d^{-3}.$$

Each event  $X_u$  is mutually independent from all the events  $X_v$  corresponding to the vertices  $v$  that do not share a neighbor with  $u$ . Since there are at most  $d(d-1)$  vertices that share a neighbor with  $u$  (not including  $u$  itself), the Lovász Local Lemma shows that with positive probability none of the events  $X_u$ ,  $u \in U$  happen provided that

$$e \cdot 3d^{-3} \cdot (d(d-1) + 1) < 1.$$

This inequality holds for all  $d > 8$ , and the proof is complete. ■

#### 7.4. Graphs of bounded chromatic number

**Theorem 1.11.** *For each integer  $k \geq 1$ , there exist  $c > 0$  and  $d_0 \in \mathbb{N}$  such that if  $G$  is a graph of maximum degree  $d \geq d_0$  with  $\chi(G) \leq k$ , then  $\text{wd}(G) \leq d - c\sqrt{d}$ .*

Let  $G$  be a graph of maximum degree  $d$  and chromatic number at most  $k$ , where we assume that  $d$  is sufficiently large in terms of  $k$ . Upon replacing  $G$  with a supergraph if necessary, we may assume that  $G$  is  $d$ -regular (Lemma 7.6). Let  $c$  be a sufficiently small positive quantity depending on  $k$  (but not on  $d$ ). We will construct a legal removal scheme  $(\prec, \text{save})$  on  $G$  such that  $\text{gap}(\prec, \text{save}) \geq c\sqrt{d}$ . By Lemma 7.10, this will yield the desired result.

We start by applying Lemma 7.12 to obtain a  $(2/\sqrt{d}, 1/2)$ -regular subset  $B$  of  $V(G)$ . Since  $G$  is  $d$ -regular and  $d > 18\sqrt{d} \log d$ , every vertex  $u \in V(G)$  satisfies

$$\sqrt{d} \leq \deg_B(u) \leq 3\sqrt{d}. \tag{7.13}$$

Set  $A := V(G) \setminus B$ . We will find a legal removal scheme  $(\prec, \text{save})$  on  $G$  such that:

- (a) In the ordering  $\prec$ , every vertex in  $A$  comes before every vertex in  $B$ .
- (b) Every vertex in  $B$  is saved at least  $c\sqrt{d}$  times.

Notice that if  $(\prec, \text{save})$  satisfies conditions (a) and (b), then  $\text{gap}(\prec, \text{save}) \geq c\sqrt{d}$ , which is the property we want. Indeed, take any vertex  $u \in V(G)$ . If  $u \in A$ , then, by (a) and (7.13),

$$\text{gap}(u; \prec, \text{save}) \geq |\{v \in N_G(u) : v \succ u\}| \geq \deg_B(u) \geq \sqrt{d}.$$

On the other hand, if  $u \in B$ , then, by (b),

$$\text{gap}(u; \prec, \text{save}) \geq |\{v \in N_G(u) : \text{save}(v) = u\}| \geq c\sqrt{d}.$$

Assuming  $c < 1$ , we have  $\text{gap}(u; \prec, \text{save}) \geq c\sqrt{d}$  in both cases, as desired.

A legal removal scheme  $(\prec, \text{save})$  satisfying (a) and (b) is constructed as follows. For  $1 \leq i \leq k$ , we recursively define the following numerical parameters:

$$N_1 := 1 \quad \text{and} \quad N_i := 20k \sum_{j=1}^{i-1} N_j \text{ for } i \geq 2.$$

Set  $p_i := N_i/(6kN_k)$ . Note that  $p_1 < p_2 < \dots < p_k = 1/(6k)$ . We shall assume  $c$  is so small that

$$32kc < p_1. \quad (7.14)$$

Since  $\chi(G) \leq k$ , we can partition  $A$  into  $k$  independent sets  $A_1, \dots, A_k$ . Let  $C_i$  be a  $(p_i, 1/2)$ -regular subset of  $A_i$  and let  $D_i$  be a  $(p_i, 1/2)$ -regular subset of  $A_i \setminus C_i$ . The ordering  $<$  is defined by listing the elements of  $V(G)$  in the following order:

$$C_1, D_1, C_2, D_2, \dots, C_k, D_k, A \setminus \bigcup_{i=1}^k (C_i \cup D_i), B.$$

(The order of the elements in each set in this list is arbitrary.) Since the elements of  $B$  appear last in this ordering, condition (a) is fulfilled.

Now we need to define the function **save** so that condition (b) holds. We start by recording the following observation:

**Claim 7.15.** *Every vertex  $u \in V(G)$  satisfies*

$$\deg_{C_i}(u) \leq \frac{3p_i}{2}d \quad \text{and} \quad \deg_{D_i}(u) \leq \frac{3p_i}{2}d.$$

PROOF. Immediate from the definitions of  $C_i$  and  $D_i$  and since the maximum degree of  $G$  is  $d$ . ■

By (7.13), each vertex  $u \in V(G)$  has at least  $d - 3\sqrt{d} \geq d/2$  neighbors in  $A$ . Therefore, we may partition  $B$  into  $k$  sets  $B_1, \dots, B_k$  so that each vertex in  $B_i$  has at least  $d/(2k)$  neighbors in  $A_i$ . This implies that every vertex in  $B_i$  has many neighbors in  $C_i$  and  $D_i$ .

**Claim 7.16.** *Every vertex  $w \in B_i$  satisfies*

$$\deg_{C_i}(w) \geq \frac{p_i}{4k}d \quad \text{and} \quad \deg_{D_i}(w) \geq \frac{p_i}{8k}d.$$

PROOF. The first inequality holds since  $C_i$  is a  $(p_i, 1/2)$ -regular subset of  $A_i$  and  $\deg_{A_i}(w) \geq d/(2k)$ . The second inequality follows similarly since, by Claim 7.15,

$$\deg_{A_i \setminus C_i}(w) \geq \frac{d}{2k} - \frac{3p_i}{2}d \geq \frac{d}{4k}. \quad \blacksquare$$

Note that, by (7.13), each vertex in  $D_i$  has at most  $3\sqrt{d}$  neighbors in  $B_i$ . On the other hand, by Claim 7.16, every vertex in  $B_i$  has at least  $p_i d/(8k) \geq p_1 d/(8k)$  neighbors in  $D_i$ . Since, by (7.14),

$$\lceil c\sqrt{d} \rceil \cdot 3\sqrt{d} < 4cd < \frac{p_1}{8k}d,$$

we can apply Lemma 7.5 to find a partial function  $s_i: D_i \dashrightarrow B_i$  such that:

- for all  $u \in D_i$ , if  $s_i(u)$  is defined, then  $s_i(u)$  is a neighbor of  $u$ ;
- the preimage of every vertex  $w \in B_i$  under  $s_i$  has cardinality  $\lceil c\sqrt{d} \rceil$ .

Now we can define **save**:  $V(G) \dashrightarrow V(G)$  by

$$\text{save}(u) := \begin{cases} s_i(u) & \text{if } u \in D_i \text{ and } s_i(u) \text{ is defined;} \\ \text{blank} & \text{otherwise.} \end{cases}$$

By the choice of  $s_i$ ,  $(<, \text{save})$  is a removal scheme that satisfies (b). It remains to verify that this removal scheme is legal. To this end, take any  $u, w \in V(G)$  such that  $\text{save}(u) = w$ . By construction, this means that  $u \in D_i$  and  $w \in B_i$  for some  $i$ . The vertices that precede  $u$  in the ordering  $<$  are the ones in  $C_1, D_1, \dots, C_{i-1}, D_{i-1}, C_i$ , plus possibly some vertices in  $D_i$ . Since the set  $A_i$  is independent,  $u$  has no neighbors in  $C_i \cup D_i$ , and hence, by Claim 7.15,

$$|\{v \in N_G(u) : v < u\}| = \sum_{j=1}^{i-1} (\deg_{C_j}(u) + \deg_{D_j}(u)) \leq \sum_{j=1}^{i-1} 3p_j d = \frac{3p_i}{20k}d < \frac{p_i}{4k}d.$$



On the other hand, since no vertex in  $C_i$  saves  $w$ , Claim 7.16 yields

$$|\{v \in N_G(w) : v < u \text{ and } \text{save}(v) \neq w\}| \geq \deg_{C_i}(w) \geq \frac{p_i}{4k}d.$$

Therefore, inequality (7.9) holds, and the proof of Theorem 1.11 is complete.

## 7.5. Graphs of girth at least 5

**Theorem 1.12.** *There exist  $c > 0$  and  $d_0 \in \mathbb{N}$  such that if  $G$  is a graph of maximum degree  $d \geq d_0$  and girth at least 5, then  $\text{wd}(G) \leq d - c\sqrt{d}$ .*

Let  $G$  be a graph of maximum degree  $d$  and girth at least 5, where  $d$  is sufficiently large. Upon replacing  $G$  with a supergraph if necessary, we may assume that  $G$  is  $d$ -regular (Lemma 7.7). Let  $c$  be a sufficiently small positive constant. As in the proof of Theorem 1.11, we will construct a legal removal scheme  $(<, \text{save})$  on  $G$  such that  $\text{gap}(<, \text{save}) \geq c\sqrt{d}$ . By Lemma 7.10, this will yield the desired result.

By Lemma 7.12, there is a  $(2/\sqrt{d}, 1/2)$ -regular subset  $B$  of  $V(G)$ . Then for every vertex  $u \in V(G)$ ,

$$\sqrt{d} \leq \deg_B(u) \leq 3\sqrt{d}. \quad (7.17)$$

Set  $A := V(G) \setminus B$ . Every vertex in  $A$  has at most  $3\sqrt{d}$  neighbors in  $B$ , while every vertex in  $B$  has at least  $d - 3\sqrt{d} \geq d/2$  neighbors in  $A$ . Since  $\lceil \sqrt{d}/8 \rceil \cdot 3\sqrt{d} < d/2$ , Lemma 7.5 gives a partial function  $s: A \dashrightarrow B$  such that:

- for all  $u \in A$ , if  $s(u)$  is defined, then  $s(u)$  is a neighbor of  $u$ ;
- the preimage of each  $w \in B$  under  $s$  has cardinality  $\lceil \sqrt{d}/8 \rceil$ .

For each  $w \in B$ , let  $S_w$  denote the preimage of  $w$  under  $s$ ; for  $u \in A$ , set  $S_u := \emptyset$ .

Now we assemble a removal scheme  $(<, \text{save})$  using a randomized procedure. Pick a linear ordering  $\triangleleft$  of  $A$  uniformly at random. The ordering  $<$  will start with the vertices of  $A$  listed according to  $\triangleleft$ , followed by the vertices of  $B$  in some order (to be specified shortly). Intuitively, we imagine that every vertex  $u \in A$  attempts to save the vertex  $s(u)$ . This attempt only succeeds if condition (7.9) is satisfied. Formally, we say that  $u \in A$  with  $s(u) = w \in B$  is **successful** if

$$|\{v \in N_A(u) : v \triangleleft u\}| < |\{v \in N_A(w) \setminus S_w : v \triangleleft u\}|.$$

If  $u \in A$  is successful, then we write  $\text{save}(u) := s(u)$ ; for all other vertices  $u$  we set  $\text{save}(u) := \text{blank}$ .

Say that a vertex  $w \in B$  is **happy** if its preimage under the function  $\text{save}$  has cardinality at least  $c\sqrt{d}$ . Let  $H \subseteq B$  be the set of all happy vertices. The ordering  $<$  consists of  $A$  listed according to  $\triangleleft$ , followed by  $B \setminus H$  in an arbitrary order, and then by  $H$  in an arbitrary order. By construction,  $(<, \text{save})$  is a legal removal scheme, and we claim that  $\text{gap}(<, \text{save}) \geq c\sqrt{d}$  with positive probability. The key fact we need to establish is the following:

**Claim 7.18.** *With positive probability, every vertex of  $G$  has at least  $c\sqrt{d}$  neighbors in  $H$ .*

Let us see why Claim 7.18 implies the desired result. Suppose that every vertex of  $G$  has at least  $c\sqrt{d}$  neighbors in  $H$ . Take any  $u \in V(G)$ . If  $u \notin H$ , then

$$\text{gap}(u; <, \text{save}) \geq |\{v \in N_G(u) : v > u\}| \geq \deg_H(u) \geq c\sqrt{d}.$$

On the other hand, if  $u \in H$ , then, by the definition of  $H$ ,

$$\text{gap}(u; <, \text{save}) \geq |\{v \in N_G(u) : \text{save}(v) = u\}| \geq c\sqrt{d}.$$

In either case,  $\text{gap}(u; <, \text{save}) \geq c\sqrt{d}$ , as desired.

In the remainder of this section we prove Claim 7.18. It will be convenient to assume that the random ordering  $\triangleleft$  is sampled according to the following procedure: each vertex  $u \in A$  picks a real number  $\vartheta(u) \in [0, 1]$  uniformly at random, and then we set  $u_1 \triangleleft u_2$  if and only if  $\vartheta(u_1) < \vartheta(u_2)$  (note that  $\vartheta(u_1) \neq \vartheta(u_2)$  with probability 1). For each  $u \in V(G)$ , let  $X_u$  be the random event that

$\deg_H(u) < c\sqrt{d}$ . It is clear that  $X_u$  only depends on the values of the function  $\vartheta$  on the vertices at distance at most 3 from  $u$ . Therefore,  $X_u$  is mutually independent from the events  $X_v$  corresponding to the vertices  $v$  at distance more than 6 from  $u$ . Hence, by the Lovász Local Lemma, to prove that with positive probability none of the events  $X_u$  happen it suffices to show that

$$\mathbb{P}[X_u] = o(d^{-6}). \quad (7.19)$$

The proof of (7.19) is somewhat technical, so before getting into its details, let us briefly explain the intuition behind our approach. Assuming  $c$  is small enough, it is possible to show that for each  $w \in B$ ,  $\mathbb{P}[w \text{ is happy}] = \Omega(1)$ . Since every vertex  $u \in V(G)$  has at least  $\sqrt{d}$  neighbors in  $B$ , we have  $\mathbb{E}[\deg_H(u)] = \Omega(\sqrt{d})$ . Ideally, we would now argue that the random variable  $\deg_H(u)$  is close to its expected value with very high probability. One way to achieve this would be to show that the random events “ $w$  is happy” for  $w \in N_B(u)$  are close to being mutually independent and then apply the Chernoff bound or some other similar result. This strategy indeed works in the case when  $G$  has girth at least 7. This is because for each  $w \in N_B(u)$ , the event “ $w$  is happy” is determined by the values of  $\vartheta$  in the radius-2 ball around  $w$ , and the girth-7 assumption implies that the radius-2 balls around the vertices in  $N_B(u)$  do not overlap too much.

It turns out that, with a more clever argument, we can reduce the girth requirement from 7 to 5. The idea is to define a certain property of vertices  $w \in B$ , which we call being *powerful* (or, more accurately,  $\varepsilon$ -*powerful* for some  $\varepsilon > 0$ ), so that the following statements hold:

- (a) the event “ $w$  is powerful” is determined by the values of  $\vartheta$  on the neighbors of  $w$ ;
- (b) the probability that  $w$  is powerful is at least  $\Omega(1)$  (Claim 7.22);
- (c) if  $w$  is powerful, then  $w$  is happy with very high probability (Claim 7.24).

Thanks to (b), the expected number of powerful neighbors for each vertex  $u \in V(G)$  is  $\Omega(\sqrt{d})$ . Using (a) and the girth-5 assumption, we can show that in fact  $u$  has  $\Omega(\sqrt{d})$  powerful neighbors with very high probability. Finally, according to (c), once  $u$  has  $\Omega(\sqrt{d})$  powerful neighbors, it is extremely likely that it has  $\Omega(\sqrt{d})$  happy neighbors as well.

Let us now begin the formal proof. We start by associating to each vertex of  $G$  a (random) vector with entries in  $[0, 1]$  by setting, for every  $u \in V(G)$ ,

$$x_u := (\vartheta(v) : v \in N_A(u) \setminus S_u).$$

(Recall that  $S_u = \emptyset$  for  $u \in A$ .) Now we introduce the following definitions:

**Definition 7.20** (Powerful vectors and vertices). Given a vector  $x = (x_1, \dots, x_k) \in [0, 1]^k$  and a real number  $\alpha \in [0, 1]$ , let the  $\alpha$ -**power** of  $x$  be the quantity

$$\pi(x, \alpha) := |\{i : x_i < \alpha\}|.$$

For  $\varepsilon > 0$ , we say that a vector  $x \in [0, 1]^k$  is  $\varepsilon$ -**powerful** if the following statement holds: If we pick a real number  $\alpha \in [0, 1]$  and a vector  $y \in [0, 1]^d$  uniformly at random, then

$$\mathbb{P}[\pi(y, \alpha) < \pi(x, \alpha)] \geq \varepsilon. \quad (7.21)$$

A vertex  $w \in B$  is  $\varepsilon$ -**powerful** if the vector  $x_w$  is  $\varepsilon$ -powerful.

We remark that if  $x \in [0, 1]^k$  is  $\varepsilon$ -powerful, then (7.21) also holds for  $y$  drawn uniformly at random from  $[0, 1]^\ell$  for any  $\ell \leq d$ . Similarly, if an  $\varepsilon$ -powerful vector  $x$  is obtained from another vector  $x'$  by removing some of the coordinates, then  $x'$  is  $\varepsilon$ -powerful as well, since  $\pi(x', \alpha) \geq \pi(x, \alpha)$  for all  $\alpha$ .

Using this notation, we can say that a vertex  $u \in A$  with  $s(u) = w$  is successful if and only if

$$\pi(x_u, \vartheta(u)) < \pi(x_w, \vartheta(u)).$$

**Claim 7.22.** *There exists a constant  $\varepsilon > 0$  such that if  $k \geq d - 5\sqrt{d}$ , then the probability that a uniformly random vector  $x \in [0, 1]^k$  is  $\varepsilon$ -powerful is at least  $\varepsilon$ .*

PROOF. For  $\varepsilon > 0$ , let  $p_\varepsilon$  denote the probability that a uniformly random vector  $x \in [0, 1]^k$  is  $\varepsilon$ -powerful. If we sample  $x \in [0, 1]^k$ ,  $\alpha \in [0, 1]$ , and  $y \in [0, 1]^d$  uniformly at random, then

$$\begin{aligned} \mathbb{P}[\pi(y, \alpha) < \pi(x, \alpha)] &= \mathbb{P}[x \text{ is } \varepsilon\text{-powerful}] \mathbb{P}[\pi(y, \alpha) < \pi(x, \alpha) \mid x \text{ is } \varepsilon\text{-powerful}] \\ &\quad + \mathbb{P}[x \text{ is not } \varepsilon\text{-powerful}] \mathbb{P}[\pi(y, \alpha) < \pi(x, \alpha) \mid x \text{ is not } \varepsilon\text{-powerful}] \\ &\leq p_\varepsilon + (1 - p_\varepsilon)\varepsilon \leq p_\varepsilon + \varepsilon. \end{aligned} \tag{7.23}$$

We now prove a lower bound on the left-hand side of (7.23). We sample  $\alpha \in [0, 1]$  first. Note that with probability  $1/3$ , we get  $1/3 \leq \alpha \leq 2/3$ . Now  $\pi(x, \alpha)$  and  $\pi(y, \alpha)$  are independent random variables sampled from the binomial distributions  $\text{Bin}(k, \alpha)$  and  $\text{Bin}(d, \alpha)$  respectively. It follows from the Berry–Esseen theorem (specifically from equation (7.4)) that there exists a constant  $\gamma > 0$  such that, assuming  $d$  is large enough and  $1/3 \leq \alpha \leq 2/3$ , we have

$$\mathbb{P}[\pi(x, \alpha) > \alpha k] \geq \gamma \quad \text{and} \quad \mathbb{P}[\pi(y, \alpha) < \alpha(d - 5\sqrt{d})] \geq \gamma.$$

Since  $\alpha(d - 5\sqrt{d}) \leq \alpha k$ , we conclude that

$$\mathbb{P}[\pi(y, \alpha) < \pi(x, \alpha)] \geq \frac{\gamma^2}{3}.$$

By (7.23), setting  $\varepsilon := \gamma^2/6$  finishes the proof. ■

In the remainder of the proof we fix a constant  $\varepsilon$  satisfying the conclusion of Claim 7.22. We shall assume that the ratio  $c/\varepsilon$  is sufficiently small, say  $c < \varepsilon/10$ . To simplify the presentation, we will use the asymptotic notation  $O(\cdot)$  to hide positive constant factors (which may be computed as functions of  $\varepsilon$  and  $c$ ).

**Claim 7.24.** *For every vertex  $w \in B$ , we have*

$$\mathbb{P}[w \text{ is happy} \mid w \text{ is } \varepsilon\text{-powerful}] \geq 1 - \exp\left(-O(\sqrt{d})\right).$$

PROOF. Let us fix the values  $\vartheta(v)$  for  $v \in N_A(w) \setminus S_w$  so that the vector  $x_w$  is  $\varepsilon$ -powerful. Now consider any  $u \in S_w$ . The value  $\vartheta(u)$  is chosen uniformly at random from  $[0, 1]$ . Moreover, since  $G$  is triangle-free,  $u$  and  $w$  have no common neighbors, which means that the values  $\vartheta(v)$  for  $v \in N_A(u)$  have not yet been determined. In other words,  $x_u$  is a uniformly random vector from  $[0, 1]^{\deg_A(u)}$ . Since  $x_w$  is  $\varepsilon$ -powerful and  $\deg_A(u) \leq d$ ,

$$\mathbb{P}[u \text{ is successful}] = \mathbb{P}[\pi(x_u, \vartheta(u)) < \pi(x_w, \vartheta(u))] \geq \varepsilon.$$

Since  $G$  has girth at least 5, the vertices in  $S_w$  have no common neighbors except  $w$ , and thus the random events “ $\pi(x_u, \vartheta(u)) < \pi(x_w, \vartheta(u))$ ” for  $u \in S_w$  are mutually independent. Therefore, the random variable  $\xi$  equal to the cardinality of the preimage of  $w$  under the function `save` is bounded below by a binomial random variable with distribution  $\text{Bin}(|S_w|, \varepsilon)$ . Hence, we may apply the Chernoff bound (Theorem 7.2) and the inequality  $|S_w| \geq \sqrt{d}/8$  to conclude that

$$\mathbb{P}[w \text{ is not happy}] = \mathbb{P}[\xi < c\sqrt{d}] < 2 \exp\left(-\left(\frac{\varepsilon}{8} - c\right)^2 \frac{8\sqrt{d}}{3\varepsilon}\right) \leq \exp\left(-O(\sqrt{d})\right). \quad \blacksquare$$

For a vertex  $u \in V(G)$ , define

$$P_\varepsilon(u) := \{w \in N_B(u) : w \text{ is } \varepsilon\text{-powerful}\}.$$

**Claim 7.25.** *For every vertex  $u \in V(G)$ , we have*

$$\mathbb{P}[|P_\varepsilon(u)| \geq c\sqrt{d}] \geq 1 - \exp\left(-O(\sqrt{d})\right).$$

PROOF. A slight technical issue here arises from the fact that the vectors  $x_w$  for  $w \in N_B(u)$  may not be probabilistically independent from each other, since each of them may include  $\vartheta(u)$  as one of the coordinates. To remedy this, we define for every  $w \in N_B(u)$  a vector  $x'_w$  as follows:

$$x'_w := (\vartheta(v) : v \in N_A(w) \setminus (S_w \cup \{u\})).$$

That is,  $x'_w$  is obtained from  $x_w$  by deleting the coordinate corresponding to  $u$ . Let

$$P'_\varepsilon(u) := \{w \in N_B(u) : x'_w \text{ is } \varepsilon\text{-powerful}\}.$$

Then  $P'_\varepsilon(u) \subseteq P_\varepsilon(u)$ , so it suffices to argue that

$$\mathbb{P}[|P'_\varepsilon(u)| \geq c\sqrt{d}] \geq 1 - \exp(-O(\sqrt{d})).$$

For  $w \in N_B(u)$ , let  $k(w) := |N_A(w) \setminus (S_w \cup \{u\})|$ . Then, by (7.17) and since  $|S_w| = \lceil \sqrt{d}/8 \rceil$ , we have

$$k(w) \geq d - 3\sqrt{d} - \lceil \sqrt{d}/8 \rceil - 1 \geq d - 5\sqrt{d}.$$

By the choice of  $\varepsilon$  and since  $x'_w$  is drawn uniformly at random from  $[0, 1]^{k(w)}$ , we conclude that

$$\mathbb{P}[x'_w \text{ is } \varepsilon\text{-powerful}] \geq \varepsilon.$$

As  $G$  has girth at least 5, the vertices in  $N_B(u)$  have no common neighbors except  $u$ , so we can apply the Chernoff bound and the inequality  $\deg_B(u) \geq \sqrt{d}$  to obtain the desired bound

$$\mathbb{P}[|P'_\varepsilon(u)| < c\sqrt{d}] < 2 \exp\left(-\frac{(\varepsilon - c)^2 \sqrt{d}}{3\varepsilon}\right) \leq \exp(-O(\sqrt{d})). \quad \blacksquare$$

Finally, we can bound the probability of each event  $X_u$ :

**Claim 7.26.** *Let  $u \in V(G)$ . Recall that  $X_u$  is the event that  $\deg_H(u) < c\sqrt{d}$ . Then*

$$\mathbb{P}[X_u] \leq \exp(-O(\sqrt{d})).$$

PROOF. By Claim 7.24, for each  $w \in N_B(u)$ , we have

$$\mathbb{P}[w \in P_\varepsilon(u) \setminus H] = \mathbb{P}[w \text{ is } \varepsilon\text{-powerful but not happy}] \leq \exp(-O(\sqrt{d})).$$

Therefore, by the union bound,

$$\mathbb{P}[P_\varepsilon(u) \setminus H \neq \emptyset] \leq \deg_B(u) \cdot \exp(-O(\sqrt{d})) \leq \exp(-O(\sqrt{d})).$$

Hence, by Claim 7.25,

$$\mathbb{P}[X_u] \leq \mathbb{P}[|P_\varepsilon(u)| < c\sqrt{d}] + \mathbb{P}[P_\varepsilon(u) \setminus H \neq \emptyset] \leq \exp(-O(\sqrt{d})). \quad \blacksquare$$

The upper bound on  $\mathbb{P}[X_u]$  given by Claim 7.26 implies the asymptotic bound (7.19). As discussed earlier, this yields Claim 7.18 and completes the proof of Theorem 1.12.

## References

- [AS16] N. ALON and J.H. SPENCER. *The Probabilistic Method*. 4th. Wiley Publishing, 2016. ISBN: 1119061954 (cit. on p. 12)
- [ABD21] J. ANDERSON, A. BERNSHTEYN, and A. DHAWAN. *Coloring graphs with forbidden bipartite subgraphs*, <https://arxiv.org/pdf/2107.05595> (preprint), 2021 (cit. on p. 13)
- [Ber16] A. BERNSHTEYN. *The asymptotic behavior of the correspondence chromatic number*, Disc. Math., **339** (2016), 2680–2692 (cit. on p. 4)
- [BKP17] A. BERNSHTEYN, A. KOSTOCHKA, and S. PRON. *О DP-раскраске графов и мультиграфов* (Russian) [On DP-coloring of graphs and multigraphs], Sib. Mat. Zhurnal, **58** (1) (2017). English version: <https://arxiv.org/abs/1609.00763>, 36–47 (cit. on pp. 2, 3, 10)

- [Bor79] O.V. BORODIN. *Задачи раскраски и покрытия вершин графов индуцированными подграфами* (Russian) [Problems of coloring and of covering the vertex set of a graph by induced subgraphs], Ph.D. Thesis. Novosibirsk: Novosibirsk State University, 1979 (cit. on p. 3)
- [CW75] G. CHARTRAND and C.E. WALL. *On regular bipartite-preserving supergraphs*, Aeq. Math., **13** (1975), 97–101 (cit. on p. 13)
- [Die17] R. DIESTEL. *Graph Theory*. 5th ed. Berlin Heidelberg: Springer-Verlag, 2017 (cit. on pp. 2, 13)
- [DP18] Z. DVOŘÁK and L. POSTLE. *Correspondence coloring and its application to list-coloring planar graphs without cycles of lengths 4 to 8*, J. Combin. Theory, Ser. B, **129** (2018), 38–54 (cit. on pp. 1–3)
- [EH66] P. ERDŐS and A. HAJNAL. *On chromatic number of graphs and set-systems*, Acta Math. Hung., **17** (1966), 61–99 (cit. on p. 2)
- [ERT79] P. ERDŐS, A.L. RUBIN, and H. TAYLOR. *Choosability in graphs*, Proc. West Coast Conf. on Combinatorics, Graph Theory and Computing, Congressus Numerantium XXVI, (1979), 125–157 (cit. on pp. 1, 3, 4)
- [Fel72] W. FELLER. *An Introduction to Probability Theory and Its Applications*. 2nd ed. Vol. II. New York: John Wiley & Sons, 1972 (cit. on p. 13)
- [FHK16] P. FRAIGNIAUD, M. HEINRICH, and A. KOSOWSKI. *Local conflict coloring*, IEEE Symposium on Foundations of Computer Science (FOCS), (2016), 625–634. Full version: <https://arxiv.org/abs/1511.01287> (cit. on p. 1)
- [Gal63] T. GALLAI. *Kritische Graphen I* (German) [Critical graphs I], Publ. Math. Inst. Hungar. Acad. Sci., **8** (1963), 165–192 (cit. on pp. 4, 10)
- [Imr84] W. IMRICH. *Explicit construction of regular graphs without small cycles*, Combinatorica, **4** (1984), 53–59 (cit. on p. 13)
- [Kim+20] S.J. KIM, A. KOSTOCHKA, X. LI, and X. ZHU. *On-line DP-coloring of graphs*, Discret. Appl. Math., **285** (2020), 443–453 (cit. on pp. 3, 6, 7)
- [Mar82] G. A. MARGULIS. *Explicit constructions of graphs without short cycles and low density codes*, Combinatorica, **2** (1982), 71–78 (cit. on p. 13)
- [MR02] M. MOLLOY and B. REED. *Graph Colourings and the Probabilistic Method*. Springer, 2002 (cit. on pp. 12, 13)
- [Tho94] C. THOMASSEN. *Every planar graph is 5-choosable*, J. Combin. Theory, Ser. B, **62** (1994), 180–181 (cit. on pp. 3, 7)
- [Viz76] V.G. VIZING. *Раскраска вершин графа в предписанные цвета* (Russian) [Vertex colorings with given colors], Metody Diskret. Analiz., **29** (1976), 3–10 (cit. on p. 1)
- [Voi93] M. VOIGT. *List colourings of planar graphs*, Disc. Math., **120** (1–3) (1993), 215–219 (cit. on p. 3)