

A Few Identities of the Takagi Function on Dyadic Rationals

Laura Monroe
Los Alamos National Laboratory
Los Alamos, NM 87501, USA
lmonroe@lanl.gov

Abstract

The number of imbalanced interior nodes of divide-and-conquer trees on n leaves is known to form a sequence of dilations of the Takagi function on dyadic rationals. We use this fact to derive identities on the Takagi function, and on the Hamming weight of an integer in terms of the Takagi function.

Mathematics Subject Classification: 68R05 (Primary), 26A27 (Secondary), 05C05, 28A80
Keywords: Takagi function, full binary trees, divide-and-conquer trees, Hamming weight.

Contents

1	Introduction	1
2	Notation	2
3	The Takagi function and divide-and-conquer trees	2
4	D-nodes in divide-and-conquer trees	4
4.1	Counting D-nodes in a divide-and-conquer tree	4
4.2	Counting D-nodes on a divide-and-conquer tree in terms of Hamming weight	6
5	From D-nodes to the Takagi function	9
6	Acknowledgements	11

1 Introduction

The Takagi function is a widely-studied self-similar continuous nowhere-differentiable function on $[0, 1]$, identified by Takagi in 1901 [Tak01]. It has connections to many areas of mathematics, including number theory, combinatorics, probability theory and analysis, and many people have studied this function over the past century. Two recent comprehensive surveys on the subject have

This work was supported by Triad National Security, LLC, operator of the Los Alamos National Laboratory under Contract No.89233218CNA000001 with the U.S. Department of Energy, and by LANL's Ultrascale Systems Research Center at the New Mexico Consortium (Contract No. DE-FC02-06ER25750). The United States Government retains and the publisher, by accepting this work for publication, acknowledges that the United States Government retains a nonexclusive, paid-up, irrevocable, world-wide license to publish or reproduce this work, or allow others to do so for United States Government purposes. This paper has been assigned the LANL identification number LA-UR-21-31244.

been published, both in 2011: one by Lagarias [Lag11], and one by Allaart and Kawamura [AK11]. OEIS [A268289](#) has also been related to the Takagi function; there are a number of identities linking the two [Lag11, AK11, Bar19].

There is an interesting connection between binary trees representing divide-and-conquer calculations and the Takagi function, shown in [CFH⁺20]: the number of imbalanced interior nodes of a divide-and-conquer tree is a sequence of dilations of the Takagi function on dyadic rationals.

In this paper we derive several formulas for counting such nodes on a divide-and-conquer tree, and use the dilations to extend them to identities on the Takagi function on dyadic rationals. These identities are in Section 5, and include:

- Recursive and closed formulas for the Takagi function on dyadic rationals. The recursive formulas are in Theorems 5.0.1 and 5.0.3, and the closed are in Theorems 5.0.2 and 5.0.4.
- A formula for the Hamming weight of an integer in terms of the Takagi function on dyadic rationals, in Theorem 5.0.5, derived from the formulas for counting D -nodes. As a consequence, we provide another proof of Trollope’s theorem on cumulative binary digit sums [Tro68] in Corollary 5.0.7, and give three succinct forms of the cumulative binary digit sum in Corollary 5.0.8.

2 Notation

We assume throughout the paper that $n > 0$ is an integer, and use the following representations. If additional assumptions are made in individual theorems on these variables, we state them explicitly; otherwise, the definitions in use are as stated in Notation 2.0.1.

Notation 2.0.1. The non-negative integer n may be represented as:

- (i) $n = 2^k + r$, with $k \geq 0$ and $0 \leq r < 2^k$.
- (ii) $n = 2^k \cdot (1 + x)$, with $k \geq 0$ and $0 \leq x < 1$.
- (iii) $n = \sum_{i=0}^k n_i \cdot 2^i$, where $k = \log_2(n)$ (binary decomposition).

Lemma 2.0.2. Let n, k, r, x be as in Notation 2.0.1. Then $x = \frac{r}{2^k}$, and $1 + x = \frac{n}{2^k}$.

3 The Takagi function and divide-and-conquer trees

The Takagi function, shown in Fig. 1(d), is a continuous nowhere-differentiable function on $[0, 1]$. We recast the original definition of the Takagi function on dyadic rationals from [Tak01] to be consistent with notation and numbering convenient to the study of binary trees.

Definition 3.0.1 (Takagi function). [Tak01] Let n be an integer. The *Takagi function* on dyadic rationals is defined as

$$\tau\left(\frac{n}{2^{k+1}}\right) = \sum_{i=1}^{\infty} \frac{\ell_i(n)}{2^i}, \text{ where } \ell_{i+1}(n) = \begin{cases} \sum_{j=0}^{i-1} n_{k-j}, & \text{if } n_{k-i} = 0, \\ i - \sum_{j=0}^{i-1} n_{k-j}, & \text{if } n_{k-i} = 1. \end{cases}$$

Theorem 3.0.6 gives a simple formula for the dilations of the Takagi function on dyadic rationals in $[0, 1]$, in terms of certain interior nodes of full binary trees, called D -nodes.

Definition 3.0.2 (S-nodes and D-nodes). [Mon21] An interior node of a rooted full binary tree is called an *S-node* if its two subtrees have the **S**ame number of leaves, and a *D-node* if its subtrees have **D**ifferent numbers of leaves.

Such a labeling gives an *SD-tree*. The SD-tree was defined to express the notion of balance in a binary commutative non-associative product [Mon21].

The next lemma is true because an n -leaf binary tree has $(n - 1)$ interior nodes [Knu97], and these are partitioned into *S*- and *D*-nodes.

Lemma 3.0.3. *Let $a(n)$ be the number of S-nodes in a binary tree having n leaves. Then the number of D-nodes is $((n - 1) - a(n))$.*

Definition 3.0.4 (Divide-and-conquer tree). A *divide-and-conquer tree* is a full binary tree in which the number of leaf descendants of the left and right children of any node differ by at most 1.

The binary tree generated is thus as balanced as it can be. At every level, the leaf descendants of an interior node are evenly or almost evenly divided between the left and the right branches. There is only one divide-and-conquer tree on n leaves, up to tree isomorphism.

Notation 3.0.5. The number of *D*-nodes in a divide-and-conquer tree is denoted by $\delta(n)$.

Theorem 3.0.6 is the same as Corollary 4 in [CFH⁺20] but is stated here in terms of *D*-nodes and Takagi’s original definition. It is also closely related to 3.3 in [Bar19]. We show several examples of these dilations in Fig. 1.

Theorem 3.0.6. [CFH⁺20] *Let $\delta(n)$ be the number of D-nodes in a divide-and-conquer tree on $n = 2^k + r$ leaves, with $0 \leq r \leq 2^k$ and with the binary expansion of r $r_{k-1}r_{k-2}\dots r_0$. Let $\tau(x)$ be the Takagi function on x , with $0 \leq x \leq 1$. Then*

$$\tau\left(\frac{r}{2^k}\right) = \frac{\delta(2^k + r)}{2^k}$$




Figure 1: **Divide-and-conquer dilations of the Takagi function on the dyadic rationals.** Subfigures (a), (b) and (c) show examples of the dilations $y = \frac{\delta(2^k+x)}{2^k} = \tau\left(\frac{x}{2^k}\right)$ from Theorem 3.0.6, where $\delta(n)$ is the number of *D*-nodes on a divide-and-conquer tree on n leaves. Here, $k = 4, 6,$ and $8,$ and x is an integer with $0 \leq x \leq 2^k$. These may be visually compared to subfigure (d), showing the continuous, self-similar, nowhere-differentiable Takagi (blancmange) curve on $[0, 1]$. (The blancmange curve image in subfigure (d) is taken from Wiki Commons.)

4 D-nodes in divide-and-conquer trees

In this section, we develop several identities on $\delta(n)$, the number of D -nodes in a divide-and-conquer tree. The first subsection is general, and the second develops identities in terms of $s_1(n)$, the Hamming weight of n . We will then apply Theorem 3.0.6 to these in Section 5, for some quick identities on the Takagi function.

4.1 Counting D -nodes in a divide-and-conquer tree

We state here formulas for the number of D -nodes in a divide-and-conquer form with n leaves. The number of D -nodes corresponds to sequence OEIS [A296062](#) [rOFI20], as was noted in [CFH⁺20].

Theorem 4.1.1. [CFH⁺20] *The number of D -nodes in a divide-and-conquer SD-tree with n leaf nodes is*

$$\delta(n) = \begin{cases} 2\delta(m), & \text{if } n = 2m; \\ \delta(m) + \delta(m+1) + 1, & \text{if } n = 2m + 1; \\ 0 & \text{if } n = 1. \end{cases}$$

The following corollary applies to n when n is not a power of 2.

Corollary 4.1.2. *Let $n = 2^k + r$, where $0 < r < 2^k$, and let ρ_1 be the position of the smallest non-0 bit in r . The number of D -nodes in a divide-and-conquer SD-tree with n leaf nodes is*

$$\delta(n) = \frac{1}{2}[\delta(n-1) + \delta(n+1)] + 1 - \rho_1$$

Proof. If $r = 2d + 1$ is odd:

$$\begin{aligned} \delta(n) &= \delta(2(2^{k-1} + d) + 1) \\ &= \delta(2^{k-1} + d) + \delta(2^{k-1} + d + 1) + 1 && \text{by Theorem 4.1.1} \\ &= \frac{1}{2}(2 \cdot \delta(2^{k-1} + d) + 2 \cdot \delta(2^{k-1} + d + 1)) + 1 \\ &= \frac{1}{2}(\delta(2^k + 2d) + \delta(2^k + 2d + 2)) + 1 && \text{by Theorem 4.1.1} \\ &= \frac{1}{2}(\delta(n-1) + \delta(n+1)) + 1 - 0 \\ &= \frac{1}{2}(\delta(n-1) + \delta(n+1)) + 1 - \rho_1 && \text{since } n \text{ is odd} \end{aligned}$$

Now let $r = 2d$ be even. First, we use what has been proved for n odd:

$$\begin{aligned} \delta(n-1) &= \frac{1}{2}(\delta(n-2) + \delta(n)) + 1 = \frac{1}{2}\delta(n-2) + \frac{1}{2}\delta(n) + 1 \\ \delta(n+1) &= \frac{1}{2}(\delta(n) + \delta(n+2)) + 1 = \frac{1}{2}\delta(n+2) + \frac{1}{2}\delta(n) + 1 \end{aligned}$$

Thus

$$\delta(n-1) + \delta(n+1) = \delta(n) + \frac{1}{2}(\delta(n-2) + \delta(n+2)) + 2$$

and

$$\delta(n-1) + \delta(n+1) - 2 - \delta(n) = \frac{1}{2}(\delta(n-2) + \delta(n+2)) \tag{1}$$

Now we apply induction on k , noting that the smallest non-0 bit in $\frac{n}{2} = (2^{k-1} + d)$ is $(\rho_1 - 1)$:

$$\begin{aligned}
\delta(n) &= \delta(2(2^{k-1} + d)) \\
&= 2\delta(2^{k-1} + d) && \text{by Theorem 4.1.1} \\
&= 2 \cdot \left(\frac{1}{2}[\delta(2^{k-1} + d - 1) + \delta(2^{k-1} + d + 1)] + 1 - (\rho_1 - 1)\right) && \text{by induction} \\
&= \frac{1}{2}[2 \cdot \delta(2^{k-1} + d - 1) + 2 \cdot \delta(2^{k-1} + d + 1)] + 2 - (2 \cdot \rho_1 - 2) \\
&= \frac{1}{2}[\delta(2^k + 2d - 2) + \delta(2^k + 2d + 2)] + 2 - (2 \cdot \rho_1 - 2) && \text{by Theorem 4.1.1} \\
&= \frac{1}{2}(\delta(n - 2) + \delta(n + 2)) + 4 - 2 \cdot \rho_1 \\
&= \delta(n - 1) + \delta(n + 1) - 2 - \delta(n) + 4 - 2 \cdot \rho_1 && \text{by Equation (1)} \\
&= \delta(n - 1) + \delta(n + 1) - \delta(n) + 2 - 2 \cdot \rho_1
\end{aligned}$$

So

$$2 \cdot \delta(n) = \delta(n - 1) + \delta(n + 1) + 2 - 2 \cdot \rho_1$$

and

$$\delta(n) = \frac{1}{2}(\delta(n - 1) + \delta(n + 1)) + 1 - \rho_1$$

□

Theorem 4.1.3. *The number of D -nodes in a divide-and-conquer SD-tree with n leaf nodes is*

$$\delta(n) = \sum_{i=0}^{\lfloor \log_2(n) \rfloor - 1} \lambda_i(n), \text{ where } \lambda_i(n) = \begin{cases} (n \bmod 2^i), & \text{if } n_i = 0; \\ 2^i - (n \bmod 2^i), & \text{if } n_i = 1. \end{cases} \quad (2)$$

An explicit form is

$$\delta(n) = \sum_{i=0}^{\lfloor \log_2(n) \rfloor - 1} [n_i \cdot 2^i + (-1)^{n_i} \cdot (n \bmod 2^i)] \quad (3)$$

Proof. Because every SD-tree is a full complete binary tree, there are 2^i nodes at level i except for the bottom level. At each level, the descendant leaf nodes are almost evenly divided between the nodes: at the i^{th} level, $2^i - (n \bmod 2^i)$ nodes have $\lfloor \frac{n}{2^i} \rfloor$ leaf descendants, and $(n \bmod 2^i)$ of these nodes have $\lfloor \frac{n}{2^i} \rfloor + 1$ leaf descendants.

Consider the binary representation of the number n where the bits are ordered from least significant to most significant. A node in a divide-and-conquer tree is an D -node if and only if it has an odd number of leaf descendants. If n_i is 0, then $\lfloor \frac{n}{2^i} \rfloor$ is even and $(n \bmod 2^i)$ nodes have an odd number of descendants. If n_i is 1, then $\lfloor \frac{n}{2^i} \rfloor$ is odd, and $2^i - (n \bmod 2^i)$ nodes have an odd number of descendants. So $\beta_i(n)$, the number of D -nodes at level i , is

$$\begin{cases} (n \bmod 2^i), & \text{if } n_i \text{ is } 0; \\ 2^i - (n \bmod 2^i), & \text{if } n_i \text{ is } 1. \end{cases}$$

Equation (2) is obtained by summing across the levels.

So the number of D -nodes at level i is

$$n_i \cdot 2^i + (-1)^{n_i} \cdot (n \bmod 2^i)$$

Equation (3) is obtained by summing across the levels. □

An extensive analysis of solutions to divide-and-conquer recurrences is provided in [HJT17]. Theorem 4.1.3 might be obtained from Theorem 4.1.1 using the methods in that paper. Instead we prove Theorem 4.1.3 directly by analyzing D -nodes.

4.2 Counting D -nodes on a divide-and-conquer tree in terms of Hamming weight

We begin this section with some simple lemmas on Hamming weight used throughout this section that appear in many papers. They can be seen by inspection.

Definition 4.2.1 (Hamming weight). The *Hamming weight* of n , $\sum_{i=0}^k n_i$, is the number of 1s in the binary expansion of n , and is denoted in this paper by $s_1(n)$.

Lemma 4.2.2. $s_1(n) = s_1(r) + 1$.

Lemma 4.2.3. If n is even, then $s_1(n+1) = s_1(n) + 1$.

Lemma 4.2.4. If n is odd, then $s_1(n-1) = s_1(n) - 1$.

Lemma 4.2.5. If $r < 2^{k-1}$ is odd, then $s_1(2^k - r) = 1 + k - s_1(r)$.

Proof. The binary representation of 2^k is 1 0...0, where there are k 0s following the leading 1. Let the binary representation of r be 0 0 $r_{k-2} \dots r_1 1$, since $r < 2^{k-1}$ is odd. Subtracting r from 2^k gives 0 1 $\overline{r_{k-2}} \dots \overline{r_1} 1$, where $\overline{r_i}$ is the negation of r_i . The lemma follows. \square

We use the above lemmas to prove a series of assertions on the D -nodes of a tree with n leaves.

Lemma 4.2.6. Let $n = 2^k + r$, with r odd and $0 < r < 2^k$. Let λ be as in Theorem 4.1.3, with $0 < i < k$. Then

$$\lambda_i(n) - \lambda_i(n-1) = \begin{cases} 1, & \text{if } n_i = 0; \\ -1, & \text{if } n_i = 1. \end{cases}$$

Proof. For $0 < i < k$, all bits of r and $(r-1)$ besides the 0^{th} are the same, since r is odd. Thus,

$$\lambda_i(n) - \lambda_i(n-1) = \begin{cases} (n \bmod 2^i) - ((n-1) \bmod 2^i) = 1, & \text{if } n_i = 0; \\ (2^i - (n \bmod 2^i)) - (2^i - ((n-1) \bmod 2^i)) = -1, & \text{if } n_i = 1. \end{cases}$$

\square

The next lemma follows from Theorem 4.1.1 by induction.

Lemma 4.2.7. $\delta(2^{k+1} - r) = \delta(2^k + r)$, when $0 \leq r \leq 2^k$.

Lemma 4.2.8. Let $n > 0$ be even. Then

$$\delta(n+1) = \delta(n) + (\lfloor \log_2(n) \rfloor - 2 \cdot s_1(n) + 2).$$

Proof. Let $n = 2^k + r$, with $0 \leq r < 2^k$, and let n have as its binary expansion $n_k n_{k-1} \dots n_0$. n is even, so $\lfloor \log_2(n) \rfloor = \lfloor \log_2(n+1) \rfloor$.

$$\begin{aligned} \delta(n+1) - \delta(n) &= \sum_{i=0}^{k-1} \lambda_i(n+1) - \sum_{i=0}^{k-1} \lambda_i(n) && \text{by Theorem 4.1.3} \\ &= \sum_{i=0}^{k-1} (\lambda_i(n+1) - \lambda_i(n)) \\ &= (\lambda_0(n+1) - \lambda_0(n)) + \sum_{i=1}^{k-1} (\lambda_i(n+1) - \lambda_i(n)) \end{aligned}$$

$n_0 = 0$, so $\lambda_0(n+1) - \lambda_0(n) = (2^0 - 0) - 0 = 1$.

There are $s_1(n) - 1 = (s_1(n+1) - 2)$ 1s in the 1st through $(k-1)$ th bits of $(n+1)$.

Thus there are $((k-1) - (s_1(n+1) - 2))$ 0s in 1st through $(k-1)$ th bits of $(n+1)$. So

$$\begin{aligned}
\delta(n+1) - \delta(n) &= 1 - 1 \cdot [s_1(n+1) - 2] + 1 \cdot [(k-1) - (s_1(n+1) - 2)] && \text{by Lemma 4.2.6} \\
&= k - 2 \cdot s_1(n+1) + 4 \\
&= k - 2 \cdot (s_1(n) + 1) + 4 && \text{by Lemma 4.2.3} \\
&= k - 2 \cdot s_1(n) + 2 \\
&= \lfloor \log_2(n) \rfloor - 2 \cdot s_1(n) + 2
\end{aligned}$$

□

Lemma 4.2.9. *Let $n = 2^k + r > 0$ be odd. Then*

$$\delta(n+1) = \delta(n) + (\lfloor \log_2(n) \rfloor - 2 \cdot s_1(n) + 2).$$

Proof. Let $n = 2^k + r$, with $0 < r < 2^k$. We proceed by moving from n and $n+1$ to the symmetric $m = 2^{k+1} - (r+1)$ (which is even) and $m+1 = 2^{k+1} - r$ (which is odd), by Lemma 4.2.7. We then apply Lemma 4.2.8, and move back to n and $n+1$, again by Lemma 4.2.7.

$$\begin{aligned}
\delta(n+1) &= \delta(m) && \text{by Lemma 4.2.7} \\
&= \delta(m+1) - (\lfloor \log_2(m) \rfloor - 2 \cdot s_1(m) + 2) && \text{by Lemma 4.2.8} \\
&= \delta(m+1) - (k - 2 \cdot s_1(m) + 2) \\
&= \delta(n) - (k - 2 \cdot s_1(m) + 2) && \text{by Lemma 4.2.5} \\
&= \delta(n) - (k - 2 \cdot (s_1(m+1) - 1) + 2) && \text{by Lemma 4.2.3} \\
&= \delta(n) - (k - 2 \cdot s_1(2^k - r) + 4) \\
&= \delta(n) - (k - 2 \cdot (2 + k - s_1(r)) + 4) && \text{by Lemma 4.2.5} \\
&= \delta(n) - (k - 2 \cdot (2 + k - (s_1(n) - 1)) + 4) && \text{by Lemma 4.2.2} \\
&= \delta(n) + (k - 2 \cdot s_1(n) + 2) \\
&= \delta(n) + (\lfloor \log_2(n) \rfloor - 2 \cdot s_1(n) + 2)
\end{aligned}$$

□

We then have a general recurrence relation for $\delta(n)$, following from Lemmas 4.2.8 and 4.2.9.

Theorem 4.2.10.

$$\delta(n+1) = \delta(n) + (\lfloor \log_2(n) \rfloor - 2 \cdot s_1(n) + 2).$$

Repeated application of Theorem 4.1.1 (for even sub-sums) and Lemma 4.2.8 (for odd) gives rise to Theorem 4.2.11, another explicit formula for the number of D -nodes in a divide-and-conquer tree.

Theorem 4.2.11.

$$\delta(n) = \sum_{i=0}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right]$$

or alternatively

$$\delta(n) = \sum_{i=0}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot s_1 \left(\left\lfloor \frac{n}{2^i} \right\rfloor \right) + 4 \right]$$

Proof. Proof by induction, noting that this is true for $n = 1$ and $n = 2$.

If n is even, $n/2$ has as its binary expansion $n_k n_{k-1} \dots n_1$, so

$$\begin{aligned} \delta(n/2) &= \sum_{i=1}^{k-1} 2^{i-1} \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] && \text{by induction} \\ &= \frac{1}{2} \sum_{i=1}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] \end{aligned}$$

$$\begin{aligned} \delta(n) &= 2\delta(n/2), && \text{by Theorem 4.1.1} \\ &= 2 \cdot \frac{1}{2} \sum_{i=1}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] \\ &= \sum_{i=0}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] && \text{since } n_0 = 0 \end{aligned}$$

If n is odd, n has as its binary expansion $n_k n_{k-1} \dots n_1 1$, and $n - 1$ has as its binary expansion $n_k n_{k-1} \dots n_1 0$.

$$\begin{aligned} \delta(n) &= \delta(n-1) + (\lfloor \log_2(n-1) \rfloor - 2 \cdot s_1(n-1) + 2) && \text{by Theorem 4.2.10} \\ &= \delta(n-1) + (k-2 \cdot s_1(n-1) + 2) \\ &= \sum_{i=0}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] + [k-2 \cdot s_1(n-1) + 2] && \text{by induction} \\ &= \sum_{i=1}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] + [k-2 \cdot s_1(n-1) + 2] \\ &= \sum_{i=1}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] + [k-2 \cdot (s_1(n) - 1) + 2] && \text{by Lemma 4.2.4} \\ &= \sum_{i=1}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] + \left[k-2 \cdot \sum_{j=0}^k n_j + 4 \right] \\ &= \sum_{i=1}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] + 2^0 \cdot 1 \cdot \left[(k-0) - 2 \cdot \sum_{j=0}^k n_j + 4 \right] \\ &= \sum_{i=1}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] + 2^0 \cdot n_0 \cdot \left[(k-0) - 2 \cdot \sum_{j=0}^k n_j + 4 \right] \\ &= \sum_{i=0}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right] \end{aligned}$$

□

Theorem 4.2.10 also gives a corollary on the Hamming weight of n .

Corollary 4.2.12.

$$s_1(n) = \frac{\delta(n) - \delta(n+1) + \lfloor \log_2(n) \rfloor}{2} + 1.$$

5 From D-nodes to the Takagi function

In this section, we take the identities on D -nodes on a divide-and-conquer tree from Section 4 and apply Theorem 3.0.6 to obtain identities on the Takagi function.

Theorem 5.0.1. *Let $1 \leq r \leq 2^k - 1$. Let ρ_1 be the position of the smallest non-0 bit in r . Then*

$$\tau\left(\frac{r}{2^k}\right) = \frac{1}{2} \left[\tau\left(\frac{r-1}{2^k}\right) + \tau\left(\frac{r+1}{2^k}\right) \right] + \frac{1 - \rho_1}{2^k}$$

Proof. Follows from Theorem 4.1.2. □

We observe that Theorem 5.0.1 is a special case of Boros' inequality [Bor08]

$$\tau\left(\frac{x+y}{2}\right) \leq \frac{1}{2} [\tau(x) + \tau(y)] + \frac{|x-y|}{2}, \quad (4)$$

where

$$x = \frac{r-1}{2^k} \text{ and } y = \frac{r+1}{2^k}.$$

When r is odd, $\rho_1 = 0$, so equality holds in (4). When r is even, $\rho_1 > 0$ and (4) is a strict inequality.

Theorem 5.0.2.

$$\tau\left(\frac{r}{2^k}\right) = \frac{1}{2^k} \cdot \sum_{i=0}^{k-1} \lambda_i(n), \text{ where } \lambda_i(n) = \begin{cases} (n \bmod 2^i), & \text{if } n_i = 0; \\ 2^i - (n \bmod 2^i), & \text{if } n_i = 1. \end{cases}$$

An explicit form is

$$\tau\left(\frac{r}{2^k}\right) = \frac{1}{2^k} \cdot \sum_{i=0}^{k-1} [n_i \cdot 2^i + (-1)^{n_i} \cdot (n \bmod 2^i)]$$

Proof. Follows from Theorem 4.1.3. □

Theorem 5.0.3.

$$\tau\left(\frac{r+1}{2^k}\right) = \tau\left(\frac{r}{2^k}\right) + \frac{1}{2^k} \cdot (k - 2 \cdot s_1(n) + 2).$$

Proof. Follows from Theorem 4.2.10. □

Theorem 5.0.3 may also be derived from Krüppel's Proposition 2.1 (Formula (2.2) in [Krü07], by substituting 1 for x , r for k , and k for ℓ in the statement of the formula.

Theorem 5.0.4.

$$\tau\left(\frac{r}{2^k}\right) = \frac{1}{2^k} \cdot \sum_{i=0}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot \sum_{j=i}^k n_j + 4 \right]$$

or alternatively

$$\tau\left(\frac{r}{2^k}\right) = \frac{1}{2^k} \cdot \sum_{i=0}^{k-1} 2^i \cdot n_i \cdot \left[(k-i) - 2 \cdot s_1\left(\left\lfloor \frac{n}{2^i} \right\rfloor\right) + 4 \right]$$

Proof. Follows from Theorem 4.2.11. □

We end the paper with Theorem 5.0.5 on the Hamming weight of binary integers and its Corollary 5.0.7 on the cumulative sum of the weights of integers. Corollary 5.0.7 was shown by Trollope in 1968 [Tro68], but also follows directly from the weight result, so we provide that proof.

Theorem 5.0.5.

$$s_1(n) = 2^{k-1} \left[\tau \left(\frac{r}{2^k} \right) - \tau \left(\frac{r+1}{2^k} \right) \right] + \frac{k+2}{2}.$$

Proof. Follows from Theorem 5.0.3. □

Lemma 5.0.6. *Let $S_1(n) = \sum_{i=0}^{n-1} s_1(i)$. If n is a power of 2, then $S_1(n) = \frac{1}{2} \cdot n \cdot \log_2(n)$.*

Proof. Exactly half of the bits in the integers between 1 and $n-1$ have value 1. □

Corollary 5.0.7. [Tro68] *Let $S_1(n) = \sum_{i=0}^{n-1} s_1(i)$. Then*

$$S_1(n) = \frac{n \cdot \log_2(n)}{2} + 2^{k-1} \cdot [2x - \tau(x) - (1+x) \cdot \log_2(1+x)].$$

Proof.

$$\begin{aligned} S_1(n) &= \sum_{i=1}^{2^k-1} s_1(i) + \sum_{i=2^k}^{2^k+r-1} s_1(i) \\ &= \frac{2^k \cdot k}{2} + \sum_{i=2^k}^{2^k+r-1} s_1(i) && \text{by Lemma 5.0.6} \\ &= \frac{2^k \cdot k}{2} + 2^{k-1} \cdot \left(\tau \left(\frac{0}{2^k} \right) - \tau \left(\frac{r}{2^k} \right) \right) + \sum_{i=2^k}^{2^k+r-1} \frac{k+2}{2} && \text{by Theorem 5.0.5} \\ &= \frac{2^k \cdot k}{2} - 2^{k-1} \cdot \tau \left(\frac{r}{2^k} \right) + r \cdot \frac{k+2}{2} \\ &= \frac{(2^k+r) \cdot k}{2} - 2^{k-1} \cdot \tau \left(\frac{r}{2^k} \right) + r \\ &= \frac{(2^k+r) \cdot k}{2} + 2^{k-1} \cdot \left[2 \cdot \frac{r}{2^k} - \tau \left(\frac{r}{2^k} \right) \right] \\ &= \frac{n \cdot k}{2} + 2^{k-1} \cdot [2x - \tau(x)] && \text{by Lemma 2.0.2 (5)} \\ &= \frac{n \cdot k}{2} + \frac{n}{2} \cdot \log_2 \left(\frac{n}{2^k} \right) + 2^{k-1} \cdot \left[2x - \tau(x) - \frac{n}{2^k} \cdot \log_2 \left(\frac{n}{2^k} \right) \right] \\ &= \frac{n \cdot k}{2} + \frac{n}{2} \cdot (\log_2 n - k) + 2^{k-1} \cdot \left[2x - \tau(x) - \frac{n}{2^k} \cdot \log_2 \left(\frac{n}{2^k} \right) \right] \\ &= \frac{n \cdot \log_2 n}{2} + 2^{k-1} \cdot \left[2x - \tau(x) - \frac{n}{2^k} \cdot \log_2 \left(\frac{n}{2^k} \right) \right] \\ &= \frac{n \cdot \log_2 n}{2} + 2^{k-1} \cdot [2x - \tau(x) - (1-x) \cdot \log_2(1-x)] && \text{by Lemma 2.0.2} \end{aligned}$$

□

Finally, Corollary 5.0.8 gives three succinct formulas of the cumulative binary digit sum in terms of the Takagi function and in terms of the number of D -nodes.

Corollary 5.0.8. *Let n be as in Notation 2.0.1, and let $S_1(n) = \sum_{i=0}^{n-1} s_1(i)$. Then*

$$(a) S_1(n) = \frac{1}{2} \cdot [nk + 2^k \cdot (2x - \tau(x))].$$

$$(b) S_1(n) = \frac{1}{2} \cdot (nk + 2r - 2^k \cdot \tau(\frac{r}{2^k})).$$

$$(c) S_1(n) = \frac{1}{2} \cdot [nk + 2r - \delta(n)].$$

Proof. Follows from Equation (5) in the proof of Corollary 5.0.7, and Theorem 3.0.6. □

Corollaries 5.0.8 (a) and (b) may also be derived from Krüppel’s Proposition 2.1 (Formula (2.4) in [Krü07], by substituting r for n and ℓ for k in the statement of the formula and using Lemma 2.0.2 to obtain $S_1(n)$. Corollary 5.0.8 (c) would then follow by Theorem 3.0.6.

6 Acknowledgements

We thank Jean-Paul Allouche for calling our attention to Krüppel’s results on Hamming weights.

References

- [AK11] Pieter Allaart and Kiko Kawamura. The Takagi function: a survey. *Real Analysis Exchange*, 37, 2011.
- [Bar19] Thomas Baruchel. Properties of the cumulated deficient binary digit sum. <https://arxiv.org/abs/1908.02250>, 2019.
- [Bor08] Zoltán Boros. An inequality for the takagi function. *Math. Inequal. Appl.*, 11:757–765, 2008.
- [CFH⁺20] Tomás M. Coronado, Mareike Fischer, Lina Herbst, Francesc Rosselló, and Kristina Wicke. On the minimum value of the Colless index and the bifurcating trees that achieve it. *J. Math. Biol.*, 80:1993–2054, 2020.
- [HJT17] Hsien-Kuei Hwang, Svante Janson, and Tsung-Hsi Tsai. Exact and asymptotic solutions of a divide-and-conquer recurrence dividing at half: Theory and applications. *ACM Trans. Algorithms*, 13(4), 2017.
- [Knu97] D.E. Knuth. *The Art of Computer Programming: Volume 1: Fundamental Algorithms*. Pearson Education, 1997.
- [Krü07] M. Krüppel. On the extrema and the improper derivatives of Takagi’s continuous nowhere differentiable function. *Rostocker Mathematisches Kolloquium*, 62:41–59, 2007.
- [Lag11] Jeffrey Lagarias. The Takagi function and its properties. *Functions in Number Theory and Their Probabilistic Aspects*, 2011.
- [Mon21] Laura Monroe. A class of trees having near-best balance. <https://arxiv.org/abs/2108.11496>, 2021.

- [rOFI20] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences. <http://oeis.org>, 2020. References sequences A268289 and A296062.
- [Tak01] Teiji Takagi. A simple example of the continuous function without derivative. *Tokyo Sugaku-Butsurigakkwai Hokoku*, 1:176–177, 1901.
- [Tro68] J. R. Trollope. An explicit expression for binary digital sums. *Mathematics Magazine*, 41(1):21–25, 1968.