

# Deep inelastic scattering with proton target in the presence of gluon condensation using holography

Sara Tahery,<sup>a 1</sup> Xiaopeng Wang,<sup>b 2</sup> Xurong Chen<sup>c 3</sup>

<sup>a,b,c</sup>*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

<sup>b</sup>*Lanzhou University, Lanzhou 730000, China*

<sup>b,c</sup>*University of Chinese Academy of Sciences, Beijing 100049, China*

<sup>c</sup>*Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter,*

<sup>c</sup>*South China Normal University, Guangzhou 510006, China*

## Abstract

We study the deep inelastic scattering (DIS) of a proton-targeted lepton in the presence of gluon condensation using gauge/gravity duality. We use a modified  $AdS_5$  background where the modification parameter  $c$  corresponds to the gluon condensation in the boundary theory. Firstly, when examining the electromagnetic field, we find that non-zero  $c$  can increase the magnitude of the field. Our goal is to find the acceptable value of  $c$  for this scattering and our method is based on setting the mass of the proton as an eigenvalue of the baryonic state equations of the DIS to find the acceptable value of the parameter  $c$  on the other side of the equations. Therefore in the second step, we calculate wave function equations for the baryonic states where the mass of the proton target requires a value contribution of  $c$  as  $c = 0.0120 \text{ GeV}^4$ . Proceeding by the electromagnetic field and the baryonic states, we derive the holographic interaction action related to the amplitude of the scattering. Finally, we compute the corresponding structure functions numerically as functions of  $x$  and  $q$ , which are Björken variables and the lepton momentum transfers, respectively. Comparing the Jlab Hall C data with our theoretical calculations, our results are acceptable.

---

<sup>1</sup>E-mail: s.tahery@impcas.ac.cn

<sup>2</sup>E-mail: wangxiaopeng@impcas.ac.cn

<sup>3</sup>E-mail: xchen@impcas.ac.cn

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>DIS parametrization and holography</b>	<b>4</b>
<b>3</b>	<b>Electromagnetic interactions in deep inelastic scattering</b>	<b>6</b>
<b>4</b>	<b>Baryonic state equations in deep inelastic scattering</b>	<b>7</b>
<b>5</b>	<b>Modified geometry and the action of deep inelastic scattering</b>	<b>11</b>
<b>6</b>	<b>Conclusions</b>	<b>15</b>

## 1 Introduction

The Deep inelastic scattering (DIS) process is used to study internal structure of proton [1] which is a puzzle in quantum chromodynamic (QCD). The main difficulty arises from the fact that massless gluons and nearly massless quarks give rise to the mass of the proton  $M \sim 1 \text{ GeV}$ . One can say that from the point of view of special relativity the missing mass could be considered as initiation by kinetic energy of quarks and gluons inside the proton, but it turns out that the mentioned kinetic energy is not enough to describe the total vanishing mass. To learn more about this topic, breaking the conformal symmetry in QCD is another reason that should be considered. It leads to anomalous dimension contributions in the DIS. Recall that QCD coupling is large at low energies, so perturbation calculations cannot be used to study many properties of hadrons. The alternative way for this purpose, is to use a holographic approach to study DIS.

In AdS/CFT description a strongly coupled field theory at the boundary of the AdS space could be described by a weakly coupled gravitational theory in the bulk of the AdS [2, 3]. In other words a ten dimensional geometry at the corresponding boundary is an exact dual of a supersymmetric  $SU(N)$  gauge theory with large  $N$  in the bulk. In particular, the string theory in the space  $AdS_5 \times S_5$  is a dual of a four dimensional gauge theory. AdS/CFT duality is for conformal theories originally, but fortunately it has been generalized to non-conformal theories like QCD, so in any case of interest one can study a QCD problem by assuming an appropriate AdS background. In the current holographic study a DIS process with a proton target is considered using a modified  $AdS_5$  background. In general in a phenomenological holographic approach, AdS/QCD tries to adapt a five-dimensional effective field theory to QCD as much as possible. Therefore mass gap, confinement and supersymmetry breaking are obtained by considering some modifications in gravitational duals. To break the conformal symmetry, one can modify the radial coordinate, the 5th dimension of spacetime, as was done in the reference [4]. In AdS/QCD approach, the anomalous dimension

associated with DIS, is replaced by the introducing of the modification parameter in gravity dual. Because of this, a holographic modified model can deal with phenomenological aspects of DIS. Introducing an anomalous dimension into the holographic context is applied to fit the masses for mesons, baryons and glueballs [5–7]. The correct value of the modification parameter in different models, can be determined using phenomenology, by setting values of known physical quantities. For example, one may adopt a modified AdS that introduces the gluon condensation on the QCD side of duality.

Originally, gluon condensation was a measure of the non-perturbative physics in zero-temperature QCD [8]. It was later identified as an order parameter for confinement to study some non-perturbative phenomena [9–12]. The gluon condensate  $G_2$  is the vacuum expectation value of the operator  $\frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{a,\mu\nu}$  where  $G_{\mu\nu}^a$  is the gluon field strength tensor. A non-zero trace of the energy-momentum tensor appears in a full quantum theory of QCD. The anomaly implies a non-zero gluon condensate which can be calculated as [13–15],

$$\Delta G_2(T) = G_2(T) - G_2(0) = -(\varepsilon(T) - 3P(T)), \quad (1)$$

where  $G_2(T)$  denotes the thermal gluon condensate,  $G_2(0)$ , being equal to the condensate value at the deconfinement transition temperature, is the zero temperature condensate value,  $\varepsilon(T)$  is the energy density,  $P(T)$  is the pressure of the QGP system.

A well-known modified holographic model introducing gluon condensation in the boundary theory is given by the following background metric in *Minkowski* spacetime [16],

$$ds^2 = g_{mn} dx^m dx^n = \frac{L^2}{z^2} (\sqrt{1 - c^2 z^8} (-dt^2 + \sum_i dx_i^2) + dz^2), \quad (2)$$

$$\varphi(z) = \sqrt{\frac{3}{2}} \ln \frac{1 + cz^4}{1 - cz^4} + \varphi_0. \quad (3)$$

In the above dilaton-wall solution,  $i = 1, 2, 3$  are orthogonal spatial boundary coordinates,  $z$  denotes the 5th dimension, radial coordinate and  $z = 0$  sets the boundary.  $\varphi_0$  is a constant,  $c = \frac{1}{z_c^4}$  and  $z_c$  denotes the IR cutoff. It shows that  $z$  is defined from zero to the IR limit as usual. To clarify, parameter  $c$  does not bound upper limit of  $z$  to values less than cutoff, in fact it should be interpreted as  $c < \frac{1}{z_c^4}$ . In other words, for a very large value of  $z$ , the parameter  $c$  tends to an infinitesimal value based on the range  $1 - c^2 z^8 > 0$ . Also we calculate in the unit where  $L = 1$ . To investigate how the forth correction of radial coordinate (with the coefficient denoted  $c$ ) appears in the metric, let us mention that the dilaton field is dual to a scalar operator and the metric is dual to the energy-momentum tensor of the dual field theory [17] (for more discussions see [18, 19]). Expanding the dilaton profile near  $z = 0$  will give,

$$\varphi(z) = \varphi_0 + \sqrt{6}cz^4 + \dots \quad (4)$$

According to the holographic dictionary  $\varphi$  and  $c$  are the source and the parameter associated with the confinement respectively. We expect the constant piece to correspond to the source for the operator  $TrG_2$  and coefficient of the  $z_4$  to give the gluon condensate [20]. Obviously  $c$  in the background metric breaks the conformal symmetry so the gluon condensation appears in the boundary theory. The relevant phenomenological information show its value generally lie in the range  $0 < c \leq 0.9 \text{ GeV}^4$  [21–23], and recent lattice calculations based on a QCD sum rule method show that the gluon condensate behaves a rapid change around  $T_c$  [24, 25]. We are interested in finding the exact value of  $c$  in a proton-targeted DIS process.

The holographic description of the gluon condensation allows many physical quantities to be studied in this context. Firstly in order to get familiar with its phenomenological aspects notice that the dilaton wall solution represented by (2), (3) is related to the zero temperature case, hence this is suitable for studying DIS and its physics. As it must be, one can readily check that in the limit  $c \mapsto 0$ , (2) reduces to  $AdS_5$  which does not present mass gap, while modification of radial coordinate would yield more phenomenological results. In fact, it has become an approach to discuss more phenomenological aspects using modified AdS [26–28].

In the most related work to our case in [29], such a scattering has been investigated by using a deformed AdS. Since models with anomalous dimension in AdS/QCD lead to mass-scale fermionic field generation, many works have used them to deal with DIS [29–61]. In view of all the above motivations, we will use a holographic model of gluon condensation (2) and (3) in the current work to study DIS with proton target.

This paper is structured as follows, after a brief overview of the DIS properties using holography in section 2 we will study electromagnetic interactions and baryonic states in deep inelastic scattering in sections 3 and 4 respectively. Based on these results, Section 5 provides the interaction action, then we will study structure functions according to the relationship between such action and scattering amplitude. In section 6 we briefly review and discuss our results.

## 2 DIS parametrization and holography

We begin this section with a brief overview of deep inelastic scattering to clarify our motivation and goals. The main application of DIS in particle physics is the study of internal hadronic structure and strong interactions. Consider a DIS process in which a lepton scattered off a proton target, as it is shown in figure 1. During this scattering a virtual photon is exchanged. Proton fragmentation creates a lepton and some final hadronic states. It should be noted that production of final hadronic states depends on four momentum the initial lepton transfers. Therefore, the four momentum causes the inner quarks and gluons of proton expelled out, finally in the next step quark anti-quark pairs are hadronized. According to [62] DIS is parametrized by Bjorken dynamical variable which is




Figure 1: Deep inelastic scattering process of the lepton on the proton by exchange of the virtual photon.

defined as,

$$x = -\frac{q^2}{2P \cdot q}, \quad (5)$$

where  $q$  is the momentum that lepton transfers to the proton target via a virtual photon and  $P$  is the initial momentum of the proton. We adopt the method has been explained in [62] and rederived in [5, 29]. Doing so, the hadronic transition amplitude is given as,

$$W^{\mu\nu} = F_1(\eta^{\mu\nu} - \frac{q^\mu q^\nu}{q^2}) + \frac{2x}{q^2} F_2(P^\mu + \frac{q^\mu}{2x})(P^\nu + \frac{q^\nu}{2x}), \quad (6)$$

where  $F_{1,2} = F_{1,2}(x, q^2)$  are some structure functions.

Now let's relate the matrix above to holography. From the AdS/QCD dictionary, elements of (6) on the QCD side are associated with the interaction action on the AdS side as [30],

$$\eta_\mu \langle P + q, s_X | J^\mu(0) | P, s_i \rangle = \mathcal{K}_{eff} S_{int}, \quad (7)$$

where  $\eta_\mu$  is polarization of virtual photon,  $|P, s_i \rangle$  represents a normalizable proton state with spin  $s_i$ ,  $J^\mu$  is the electromagnetic quark current and  $s_X$  denotes the final state. It is worth to mention that  $\mathcal{K}_{eff}$  is an effective factor that adjusts the bulk supergravity quantities to the boundary phenomenologically. This is based on a different perspective in reference [5] that bulk/boundary quantities of (7) are proportional, and not necessarily equal.

The interaction action is written as,

$$S_{int} = g_V \int dz d^4 y e^{-\varphi} \sqrt{-g} \phi^\mu \bar{\Psi}_X \Gamma_\mu \Psi_i, \quad (8)$$

and  $g_V$  is a coupling constant related to the electric charge of the baryon,  $\varphi$  is the dilaton field and  $\sqrt{-g}$  is given by the metric,  $\phi^\mu$  is the electromagnetic gauge field,  $\Psi_i$  and  $\Psi_X$  are the initial and final state spinors for the baryon, respectively and  $\Gamma_\mu$  are Dirac gamma matrices in the curved

space. By computing all above quantities according to (2) and (3), we study interaction action of DIS.

### 3 Electromagnetic interactions in deep inelastic scattering

A photon is exchanged during scattering, so we study the electromagnetic interactions in the bulk. It can be described as the presence of photon in the modified AdS. The action for a five dimensional massless gauge field  $\phi^n$  is given by,

$$S = -\frac{1}{4} \int d^5x e^{-\varphi} \sqrt{-g} F^{mn} F_{mn}, \quad (9)$$

where  $F^{mn} = \partial^m \phi^n - \partial^n \phi^m$ , and  $m, n$  refer to the 5-dimensional space includes Minkowski spacetime coordinates,  $\mu, \nu$  and  $z$ , and  $\varphi$  is the dilaton field given by (3). Note that  $\varphi$  and  $\phi$  should be differentiated. In fact (9) is an action showing the gauge field  $\phi$  on a background coupled to a dilaton field  $\varphi$ . From (9) the equation of motion of such an electromagnetic field is derived as,

$$\partial_m [e^{-\varphi} \sqrt{-g} F^{mn}] = 0. \quad (10)$$

Considering  $m, n \equiv \mu, \nu, z$  the relation (10) leads to,

$$\begin{aligned} \partial_\mu \left[ \frac{1}{z} (1 + cz^4)^{1-\sqrt{\frac{3}{2}}} (1 - cz^4)^{1+\sqrt{\frac{3}{2}}} F^{\mu z} \right] &= 0, \\ \partial_z \left[ \frac{1}{z} (1 + cz^4)^{1-\sqrt{\frac{3}{2}}} (1 - cz^4)^{1+\sqrt{\frac{3}{2}}} F^{z\mu} \right] &= 0. \end{aligned} \quad (11)$$

In order to solve the equations of motion of the gauge field in (11), we should first fix the gauge. Suppose there is an electromagnetic field in the bulk defined with the metric (2). This obeys the 5-dimensional Maxwell equation supplemented by a gauge condition which we take to be,

$$e^{-\varphi} \sqrt{-g} \partial_\mu \phi^\mu + \partial_z (e^{-\varphi} \sqrt{-g} \phi_z) = 0. \quad (12)$$

From (12) one can write,

$$\partial_\mu \phi^\mu + \frac{z}{(1 + cz^4)^{1-\sqrt{\frac{3}{2}}} (1 - cz^4)^{1+\sqrt{\frac{3}{2}}}} \partial_z \left( \frac{(1 + cz^4)^{1-\sqrt{\frac{3}{2}}} (1 - cz^4)^{1+\sqrt{\frac{3}{2}}}}{z} \phi_z \right) = 0, \quad (13)$$

so,

$$\square \phi_\mu + \partial_\mu \partial_z \phi_z - \frac{1 + 4\sqrt{6}cz^4 + 7c^2z^8}{z(1 - c^2z^8)} \partial_\mu \phi_z = 0. \quad (14)$$

Using the gauge (14) together with (11) leads to the following equations,

$$\square \phi_z - \partial_\mu \partial_z \phi^\mu = 0, \quad (15)$$

$$\square\phi_\mu + \partial_z^2\phi_\mu - \frac{1 + 4\sqrt{6}cz^4 + 7c^2z^8}{z(1 - c^2z^8)}\partial_z\phi_\mu = 0. \quad (16)$$

At this point one could consider a photon with a particular polarization as  $\eta_\mu q^\mu = 0$  for simplicity, hence only the  $\phi^\mu$  component contributes in the scattering [5, 30, 33]. In the latter case we need to solve only (16). This equation cannot be solved analytically and we have to use numerical methods. Let us consider  $\phi_\mu(z, q, y) = \eta_\mu e^{iq \cdot y} \phi_1(z, q)$ , and the condition  $\phi_\mu(z, q, y)|_{z=0} = \eta_\mu e^{iq \cdot y}$  in (16). We use reference [5] for the initial condition, because we should get same results at the boundary  $z = 0$ . At the IR limit we take the Neumann boundary condition. So  $\phi_1(z, q)$  should be convergent and the assumption is taken as follows,  $\frac{\partial\phi_1}{\partial z}|_{z=z_c} = 0$ . Then we can get  $\phi_1(z, q)$  versus  $z$  that describes the behaviour of the electromagnetic field in the bulk. Figure. 2 shows  $\phi_1(z, q)$  for different values




Figure 2: Electromagnetic field in the bulk with a) large value of  $q^2 = 10 \text{ GeV}^2$  and two different values of  $c = 0.01 \text{ GeV}^4$  and  $c = 0.001 \text{ GeV}^4$ , b) small value of  $q^2 = 1 \text{ GeV}^2$  and two different values of  $c = 0.01 \text{ GeV}^4$  and  $c = 0.001 \text{ GeV}^4$ , c) two different values of parameter  $q^2 = 1 \text{ GeV}^2$  and  $q^2 = 10 \text{ GeV}^2$  at fixed value of  $c = 0.01 \text{ GeV}^4$ .

of  $q^2$  and  $c$ . Plot a) shows that for large values of  $q^2$  increasing the parameter  $c$  does not affect  $\phi_1(z, q)$  significantly. Also  $\phi_1$  has its maximum value near the boundary ( $z = 0$ ). In plot b) for small values of  $q^2$ , increasing the parameter  $c$ , increases magnitude of  $\phi_1$  which is more visible with large values of  $z$ . Plot c) is a comparison of  $\phi_1$  at different values of  $q^2$  and a fixed  $c$ . Obviously  $\phi_1$  is stronger, the smaller  $q^2$  is. In other words the magnitude of the electromagnetic field is larger for small values of  $q^2$ .

## 4 Baryonic state equations in deep inelastic scattering

In this section, we study the baryonic initial and final states for further requirements of the interaction action (8). The action for the fermionic fields is written as,

$$S = \int dx^5 e^{-\varphi} \sqrt{g} \bar{\Psi} (\not{D} - m_5) \Psi. \quad (17)$$

where  $m_5$  is the baryon bulk mass and the operator  $\mathcal{D}$  is defined as,

$$\mathcal{D} = g^{mn} e_n^a \gamma_a (\partial_m + \frac{1}{2} \omega_m^{bc} \Sigma_{bc}), \quad (18)$$

in which  $g^{mn}$  is given by metric (2),  $\gamma_\alpha = (\gamma_\mu, \gamma_5)$ ,  $\{\gamma_a, \gamma_b\} = 2\eta_{ab}$  and  $\Sigma_{\mu 5} = \frac{1}{4}[\gamma_\mu, \gamma_5]$  [63–67].  $\gamma_\mu$  are Dirac's gamma matrices.  $a, b, c$  are flat space and,  $m, n, p, q$  are AdS space indices respectively. As before  $\mu, \nu$  represent the Minkowski space. The equations of motion of fermionic states are,

$$(\mathcal{D} - m_5)\Psi = 0. \quad (19)$$

With the metric (2) Vielbein are computed as,

$$\begin{aligned} e_n^a &= \frac{(1 - c^2 z^8)^{\frac{1}{4}}}{z} \delta_n^a, \\ a &= t, x_1, x_2, x_3, \\ e_n^b &= \frac{1}{z} \delta_n^b, \\ b &= z. \end{aligned} \quad (20)$$

The above terms give us first term of (18). Now we should calculate the second term of that. Spin connection is given by,

$$\omega_m^{ab} = e_n^a \partial_m e^{nb} + e_n^a e^{pb} \Gamma_{pm}^n, \quad (21)$$

where the Christoffel symbols are,

$$\Gamma_{mn}^p = \frac{1}{2} g^{pq} (\partial_n g_{mq} + \partial_m g_{nq} - \partial_q g_{mn}). \quad (22)$$

From the metric (2), one may write,  $g_{\mu\nu} = \frac{\sqrt{1-c^2 z^8}}{z^2} \eta_{\mu\nu}$  and  $g_{zz} = \frac{1}{z^2}$ . So the only non vanishing terms are,  $\Gamma_{mn}^p \Rightarrow \Gamma_{\mu\nu}^z, \Gamma_{zz}^z, \Gamma_{\nu z}^\mu$ . After computation they are written as,

$$\begin{aligned} \Gamma_{\mu\nu}^z &= -\frac{(1 + c^2 z^8)}{z(1 - c^2 z^8)} \eta_{\mu\nu} \\ \Gamma_{zz}^z &= \frac{1}{z} \\ \Gamma_{\nu z}^\mu &= \frac{(1 + c^2 z^8)}{z(1 - c^2 z^8)} \delta_\nu^\mu. \end{aligned} \quad (23)$$

Also from (2) together with (20) and (22) the relation (21) turns to,

$$\omega_\mu^{z\nu} = -\omega_\mu^{\nu z} = -\frac{(1 + c^2 z^8)}{z(1 - c^2 z^8)^{\frac{3}{4}}} \delta_\mu^\nu, \quad (24)$$

hence other components of  $\omega_m^{ab}$  are zero. Using these solutions, (18) is given by,

$$\mathcal{D} = z\gamma^5 \partial_z + \frac{z}{(1 - c^2 z^8)^{\frac{1}{4}}} \gamma^\mu \partial_\mu - 2 \frac{(1 + c^2 z^8)}{z(1 - c^2 z^8)^{\frac{3}{4}}} \gamma^5, \quad (25)$$



and (19) is written as,

$$[z\gamma^5\partial_z + \frac{z}{(1-c^2z^8)^{\frac{1}{4}}}\gamma^\mu\partial_\mu - 2\frac{(1+c^2z^8)}{z(1-c^2z^8)^{\frac{3}{4}}}\gamma^5 - m_5]\Psi = 0. \quad (26)$$

According to the fact that spinor is either left-handed or right-handed, and since Kaluza-Klein modes are dual to the chirality spinors we decompose these components and expand as,

$$\Psi_{L/R}(x^\mu, z) = \sum_n f_{L/R}^n(x^\mu) \chi_{L/R}^n(z), \quad (27)$$

by applying (27) in the equation of motion (26) we find the coupled equations as,

$$(\partial_z - 2\frac{(1+c^2z^8)}{z^2(1-c^2z^8)^{\frac{3}{4}}} + \frac{m_5}{z})\chi_L(z) = \frac{M_n}{(1-c^2z^8)^{\frac{1}{4}}}\chi_R(z), \quad (28)$$

$$(\partial_z - 2\frac{(1+c^2z^8)}{z^2(1-c^2z^8)^{\frac{3}{4}}} - \frac{m_5}{z})\chi_R(z) = \frac{-M_n}{(1-c^2z^8)^{\frac{1}{4}}}\chi_L(z). \quad (29)$$

Decoupling (28) and (29) leads to the following equation which describes both left-handed and right-handed sectors as,

$$\begin{aligned} -(1-c^2z^8)^{\frac{1}{4}}\left(\partial_z - 2\frac{(1+c^2z^8)}{z^2(1-c^2z^8)^{\frac{3}{4}}} \pm \frac{m_5}{z}\right)(1-c^2z^8)^{\frac{1}{4}}\left(\partial_z - 2\frac{(1+c^2z^8)}{z^2(1-c^2z^8)^{\frac{3}{4}}} \mp \frac{m_5}{z}\right)\chi_{R/L}(z) \\ = M_n^2\chi_{R/L}(z). \end{aligned} \quad (30)$$

Below we create a Schrödinger-like equation by applying a transformation like this,

$$\chi_{R/L}(z) = \frac{e^{-\frac{2(1-c^2z^8)^{\frac{1}{4}}}{z}}}{(1-c^2z^8)^{\frac{1}{8}}}\psi_{R/L}(z), \quad (31)$$

so the equation (30) is written as,

$$\sqrt{1-c^2z^8}\left(-\psi_{R/L}''(z) + \frac{m_5(m_5 \mp 1) - c^2z^8(2m_5^2 + 7) + c^4z^{16}m_5(m_5 \pm 1)}{z^2(1-c^2z^8)^2}\psi_{R/L}(z)\right) = M_n^2\psi_{R/L}(z) \quad (32)$$

In (32),  $m_5$  is a parameter on the AdS side of gauge/ gravity duality and is related to the baryon mass on the gauge side, so the normalizable solutions of the equations above are dual to the states in the boundary theory. In pure AdS space, the bulk mass is related to the canonical conformal dimension  $\Delta_{can}$  of a boundary operator as,

$$|m_5^{AdS}| = \Delta_{can} - 2. \quad (33)$$

Recall that  $QCD$  is not a conformal field theory since it has a mass gap. So the gravity side should be modified somehow and then it is not pure AdS any more. If one modifies AdS, the

canonical dimension  $\Delta_{can}$  of an operator has an anomalous contribution  $\gamma$  implying an effective scaling dimension.

$$|m_5| = \Delta_{can} + \gamma - 2. \quad (34)$$

The contribution of the anomaly is related to how one modifies the theory. For example in [5] modification of the scale introduces the mass gap in the theory. Therefore the anomalous contribution represents the energy scale in the theory and leads to the mass spectra. So, the main task is to find the value of the bulk mass in (34). In AdS/CFT dictionary, the bulk mass is related to the dimension, means the energy scale of the boundary theory is holographically related to the localization in the  $z$ -coordinate, therefore we have  $z$ -dependent mass in the bulk. Let us focus on  $m_5$ . One can fit  $m_5$  numerically as the equations (32) have normalizable solutions. By fixing  $M$  as proton mass, we should find suitable values for  $c$  and  $m_5$  which give us well defined answers.

Figures 3 and 4 show initial ( $n=1$ ) and final (for two excited states as  $n=2,3$ ) chiral components of the wave function respectively. To solve the equations (32) numerically, we fix proton mass  $M$  as eigenvalue of equation, therefore  $m_5$  and  $c$  are found as  $c = 0.0120 \text{ GeV}^4$  and  $m_5 = 0.081 \text{ GeV}$ . Interestingly, the value of parameter  $c$  on the AdS side is very close to the phenomenological GC value of QCD as found  $G_2 = 0.010 \pm 0.0023 \text{ GeV}^4$  in the reference [26]. Another consequence of the presence of  $c$  is that the anomaly  $\gamma$  in (34) affects the bulk mass intensely. After finding both




Figure 3: Left-handed (dashed) and right-handed (solid) sectors of wave function from (32) for the initial state (target proton), by considering  $c = 0.0120 \text{ GeV}^4$  and  $m_5 = 0.081 \text{ GeV}$ .

left-handed and right-handed modes from (32) we have,

$$\Psi_i = \frac{e^{-\frac{2(1-c^2z^8)^{\frac{1}{4}}}{z}}}{(1-c^2z^8)^{\frac{1}{8}}} e^{iP \cdot y} \left[ \left( \frac{1+\gamma_5}{2} \right) \psi_L^i + \left( \frac{1-\gamma_5}{2} \right) \psi_R^i \right] u_{s_i}(p), \quad (35)$$

as the initial wave function for the target proton and,

$$\Psi_X = \frac{e^{-\frac{2(1-c^2z^8)^{\frac{1}{4}}}{z}}}{(1-c^2z^8)^{\frac{1}{8}}} e^{iP_X \cdot y} \left[ \left( \frac{1+\gamma_5}{2} \right) \psi_L^X + \left( \frac{1-\gamma_5}{2} \right) \psi_R^X \right] u_{s_X}(p), \quad (36)$$




Figure 4: Left-handed (dashed) and right-handed (solid) sectors of wave function from (32) for the final state a)  $n=2$  and b)  $n=3$ , by considering  $c = 0.0120 \text{ GeV}^4$  and  $m_5 = 0.081 \text{ GeV}$ .

as the final wave function for the hadronic state. These will be used later.

## 5 Modified geometry and the action of deep inelastic scattering

According to (7) and (8) we find the interaction action with the electromagnetic field and the baryonic states obtained from (16) and (35) - (36) respectively. The interaction action (8) is written as,

$$\begin{aligned}
 S_{int} &= g_V \int dz d^4 y e^{-\varphi} \sqrt{-g} \phi^\mu \bar{\Psi}_X e_\mu^\alpha \gamma_\alpha \Psi_i \\
 &= g_V \int dz d^4 y \frac{1}{z} (1 - cz^4)^{1+\sqrt{\frac{3}{2}}} (1 + cz^4)^{1-\sqrt{\frac{3}{2}}} \phi^\mu \bar{\Psi}_X \frac{1}{z} (1 - c^2 z^8)^{\frac{1}{4}} \delta_\mu^\alpha \gamma_\alpha \Psi_i \\
 &= g_V \int dz d^4 y \frac{1}{z^2} (1 - cz^4)^{\frac{5}{4}+\sqrt{\frac{3}{2}}} (1 + cz^4)^{\frac{5}{4}-\sqrt{\frac{3}{2}}} \phi^\mu \bar{\Psi}_X \gamma_\mu \Psi_i,
 \end{aligned} \tag{37}$$

and from (36) one writes,

$$\bar{\Psi}_X = \frac{e^{-\frac{2(1-c^2 z^8)^{\frac{1}{4}}}{z}}}{(1 - c^2 z^8)^{\frac{1}{8}}} e^{-i P_X \cdot y} \bar{u}_{s_X}(p) \left[ \left( \frac{1 + \gamma_5}{2} \right) \psi_L^X + \left( \frac{1 - \gamma_5}{2} \right) \psi_R^X \right]. \tag{38}$$

Therefore (37) is given by,

$$\begin{aligned}
S_{int} &= \frac{g_V}{2} \int dz d^4 y e^{-i(P_X - P - q) \cdot y} \eta^\mu \phi_1 \frac{1}{z^2} (1 - cz^4)^{\frac{5}{4} + \sqrt{\frac{3}{2}}} (1 + cz^4)^{\frac{5}{4} - \sqrt{\frac{3}{2}}} \\
&\quad \frac{e^{-\frac{4(1-c^2 z^8)^{\frac{1}{4}}}{z}}}{(1 - c^2 z^8)^{\frac{1}{4}}} \left[ \bar{u}_{s_X} (\hat{P}_L \psi_L^X + \hat{P}_R \psi_R^X) \gamma_\mu (\hat{P}_L \psi_L^i + \hat{P}_R \psi_R^i) u_{s_i} \right] \\
&= \frac{g_V}{2} (2\pi)^4 \delta^4(P_X - P - q) \eta^\mu \int dz \frac{1}{z^2} (1 - cz^4)^{\frac{5}{4} + \sqrt{\frac{3}{2}}} (1 + cz^4)^{\frac{5}{4} - \sqrt{\frac{3}{2}}} \\
&\quad \frac{e^{-\frac{4(1-c^2 z^8)^{\frac{1}{4}}}{z}}}{(1 - c^2 z^8)^{\frac{1}{4}}} \phi_1 \left[ \bar{u}_{s_X} \gamma_\mu \hat{P}_R u_{s_i} \psi_L^X \psi_L^i + \bar{u}_{s_X} \gamma_\mu \hat{P}_L u_{s_i} \psi_R^X \psi_R^i \right].
\end{aligned} \tag{39}$$

By defining the following integral,

$$\mathcal{B}_{R,L} = \int dz \frac{e^{-\frac{4(1-c^2 z^8)^{\frac{1}{4}}}{z}}}{(1 - c^2 z^8)^{\frac{1}{4}}} \frac{1}{z^2} (1 - cz^4)^{\frac{5}{4} + \sqrt{\frac{3}{2}}} (1 + cz^4)^{\frac{5}{4} - \sqrt{\frac{3}{2}}} \phi_1 \psi_{R,L}^X \psi_{R,L}^i, \tag{40}$$

(39) is written as,

$$S_{int} = \frac{g_V}{2} (2\pi)^4 \delta^4(P_X - P - q) \eta^\mu [\bar{u}_{s_X} \gamma_\mu \hat{P}_R u_{s_i} \mathcal{B}_L + \bar{u}_{s_X} \gamma_\mu \hat{P}_L u_{s_i} \mathcal{B}_R], \tag{41}$$

and (7) is written as,

$$\begin{aligned}
\eta_\mu < P_X | J_\mu(q) | P_i > &= \frac{g_{eff}}{2} \delta^4(P_X - P - q) \eta_\mu [\bar{u}_{s_X} \gamma_\mu \hat{P}_R u_{s_i} \mathcal{B}_L + \bar{u}_{s_X} \gamma_\mu \hat{P}_L u_{s_i} \mathcal{B}_R] \\
\eta_\nu < P_i | J_\mu(q) | P_X > &= \frac{g_{eff}}{2} \delta^4(P_X - P - q) \eta_\nu [\bar{u}_{s_i} \gamma_\mu \hat{P}_R u_{s_X} \mathcal{B}_L + \bar{u}_{s_i} \gamma_\mu \hat{P}_L u_{s_X} \mathcal{B}_R],
\end{aligned} \tag{42}$$

where  $g_{eff}^2 = \mathcal{K}_{eff}^2 g_V^2 (2\pi)^8$ , and  $g_V^2 = \frac{1}{137}$ .  $\mathcal{K}_{eff}^2$  should be fitted numerically as shown in table 1. With (6) we get,

$$\begin{aligned}
\eta_\mu \eta_\nu W^{\mu\nu} &= \frac{\eta_{\mu\nu}}{4} \sum_{M_x^2} \sum_{s_i, s_X} \frac{g_{eff}^2}{4} \delta(M_X^2 - (P + q)^2) \left[ \bar{u}_{s_X} \gamma^\mu \hat{P}_R u_{s_i} \bar{u}_{s_i} \gamma^\nu \hat{P}_R u_{s_X} \mathcal{B}_L^2 \right. \\
&\quad + \bar{u}_{s_X} \gamma^\mu \hat{P}_R u_{s_i} \bar{u}_{s_i} \gamma^\nu \hat{P}_L u_{s_X} \mathcal{B}_L \mathcal{B}_R + \bar{u}_{s_X} \gamma^\mu \hat{P}_L u_{s_i} \bar{u}_{s_i} \gamma^\nu \hat{P}_R u_{s_X} \mathcal{B}_R \mathcal{B}_L \\
&\quad \left. + \bar{u}_{s_X} \gamma^\mu \hat{P}_L u_{s_i} \bar{u}_{s_i} \gamma^\nu \hat{P}_L u_{s_X} \mathcal{B}_R^2 \right]
\end{aligned} \tag{43}$$

Since our calculation is spin independent, we write,

$$\sum_s (u_s)_\alpha(p) (\bar{u}_s)_\beta(p) = (\gamma^\mu p_\mu + M)_{\alpha\beta}, \tag{44}$$

with the summation over the initial and final spin states, and then applying trace engineering we arrive at,

$$\eta_\mu \eta_\nu W^{\mu\nu} = \frac{g_{\text{eff}}^2}{4} \sum_{M_X^2} \delta(M_X^2 - (P+q)^2) \left\{ (\mathcal{B}_L^2 + \mathcal{B}_R^2) \left[ (P \cdot \eta)^2 - \frac{1}{2} \eta \cdot \eta (P^2 + P \cdot q) \right] \right. \\ \left. + \mathcal{B}_L \mathcal{B}_R M_X^2 M_0^2 \eta \cdot \eta \right\}, \quad (45)$$

where  $\not{p} = \gamma^\mu p_\mu$ ,  $\{\gamma_5, \gamma_\mu\} = 0$ , and  $P_{R/L} \gamma^\mu = \gamma^\mu P_{L/R}$ . Summing over the outgoing states  $P_X$  and carrying it on to the continuum limit we have the invariant mass delta function which is related to the functional form of the mass spectrum of the produced particles with the excitation number  $n$  [30],

$$\delta(M_X^2 - (P+q)^2) \propto \left( \frac{\partial M_n^2}{\partial n} \right)^{-1}. \quad (46)$$

We consider the lowest state produced at the collision, since the spectrum is linear with  $n$ , and the delta will account for  $1/M_X^2$  [30,33]. With the transversal polarization ( $\eta \cdot q = 0$ ), the hadronic tensor (6) is obtained as,

$$\eta_\mu \eta_\nu W^{\mu\nu} = \eta^2 F_1(q^2, x) + \frac{2x}{q^2} (\eta \cdot P)^2 F_2(q^2, x), \quad (47)$$

where  $F_1$  and  $F_2$  are,

$$F_1(q^2, x) = \frac{g_{\text{eff}}^2}{4} \left[ M_0 M_X \mathcal{B}_L \mathcal{B}_R + (\mathcal{B}_L^2 + \mathcal{B}_R^2) \left( \frac{q^2}{4x} + \frac{M_0^2}{2} \right) \right] \frac{1}{M_X^2}, \quad (48)$$

and

$$F_2(q^2, x) = \frac{g_{\text{eff}}^2}{8} \frac{q^2}{x} (\mathcal{B}_L^2 + \mathcal{B}_R^2) \frac{1}{M_X^2}, \quad (49)$$

respectively. Also  $M_0$  is the mass of the initial hadron and  $M_X$  is the mass of the final hadron as,

$$M_X = \sqrt{M_0^2 + q^2 \left( \frac{1-x}{x} \right)}. \quad (50)$$

## Numerical strategy

As we mentioned in section 4, in the equation of states the eigenvalue of the ground state should be close to the square of the proton mass. So, we consider the ranges  $0 < m_5 < 1 \text{ GeV}$  and  $0.001 < c < 1 \text{ GeV}^4$ , while the eigenvalue of the ground state equation, is in the range from  $0.876 \text{ GeV}$  to  $1 \text{ GeV}$  ( $M_{\text{proton}} = 0.938 \text{ GeV}$ ). Accordingly we get a set of suitable values of the parameters  $m_5$  and  $c$ . Their approximate ranges are  $0.001 < m_5 < 0.2 \text{ GeV}$  and  $0.006 < c < 0.02 \text{ GeV}^4$ . In the next step we look for the appropriate values of  $\mathcal{K}_{\text{eff}}^2$  as  $m_5$  and  $c$  satisfy their ranges and our theoretical calculations can be fitted with the experimental data for  $F_2$ .

As it is shown in table 1, in accordance with the acceptable ranges for  $\mathcal{K}_{eff}^2$ ,  $m_5$  and  $c$  we need to continue by considering 0.01 – 0.04 order of  $x$  and small  $q^2$  which our model works well. In sweep spectrum form, we determine a set of  $m_5$  and  $c$  and then for each set, we fit the experimental data for  $F_2$  to get  $\mathcal{K}_{eff}^2$  by Least squares method. What needs to be mentioned here is that for  $m_5$ , the scanning step is 0.01, and for  $c$ , the scanning step is 0.001. The optimal parameter values with the smallest uncertainty are  $m_5 = 0.081$  GeV,  $c = 0.0120$  GeV<sup>4</sup>,  $\mathcal{K}_{eff}^2 = 37.3259$ . The uncertainty of  $c$  comes from the size of scanning step. Using this parameters set, we will get the proton structure function  $F_2$  as a function of  $q^2$ .

$x$	$m_5/\text{GeV}$	$c/\text{GeV}^4$	$k_{eff}^2$
0.015	0.081	0.012	37.3259
0.025			
0.04			

Table 1: Adjustment of parameter  $k_{eff}^2$  at different  $x$  with  $c = 0.0120$  GeV<sup>4</sup>. Remind that the value of parameter  $c$  is demanded by phenomenological value of proton mass.




Figure 5: a), b), c) Comparison between Jlab Hall C data [68] and our theoretical results. Dashed lines are theoretical results and square dots are experimental data.

Figure 5 is a comparison between Jlab Hall C data [68] and our theoretical results. In plots a), b), c) our results have good agreements with experimental data at  $x = 0.015$ ,  $x = 0.025$  and  $x = 0.04$ . Note that the electromagnetic wave function and the hadron function are independent of the Bjorken variable  $x$ . In our calculation this variable only appears in the  $F_2$  formula (49) which introduces some uncertainty that is more visible in diagram (a). In practice, the GC model is suitable for such scattering for these values of  $x$ . Applying the background (2) and (3) on the one hand and setting the proton as the DIS target on the other, results in some specific small values of the  $c$  parameter being acceptable, then our model turned out to work better for the mentioned

region of  $x$  values (which are not large), than large values, as an alternative to [29] discussing large  $x$  values in the soft- wall model.

There are differences between the applications of the SW model that was used in the same DIS in the [29] and our current study. The modification parameter in [29] appears in an exponential function having dimension mass squared ( $GeV^2$ ) and is associated with a QCD mass scale. The background metric is not coupled to a dilaton field. Considering such physics with the modified exponential function and due to the kinematical region in this reference their model is expected to produce better results for large  $x$ . With the goal of discussing other areas of  $x$ , we considered a different model. In our case the modification parameter introduced as  $c$ , has the dimension of ( $GeV^4$ ), also there is a coupled dilaton field in the action. The background structure and the proton target, strongly require the small value of the  $c$  parameter and non-large  $x$ .

## 6 Conclusions

In a holographic description of DIS we found effects of the parameter  $c$  appearing in the background metric representing gluon condensation in the boundary theory. Since there is a proton target in the scattering, the mass of the proton and the value of  $c$  parameter both play an important role in this study. One of our main aim was to determine the value of  $c$  from experimental data. First, we found the behaviour of the electromagnetic field with respect to  $z$  for different values of  $c$ . We have shown that the  $c$  parameter can increase the magnitude of this field, especially for small values of  $q^2$ . Then, we solved the equation of baryonic wave function numerically and set the proton mass as the ground state eigenvalue to find the best bulk mass,  $c$  parameter and the  $k_{eff}^2$  values. Hence, only small values of  $c$  lead to a well-defined answer of the equation, or proton target requires a small value of  $c$ . It could be suggested that since the  $c$  parameter breaks the conformal symmetry, its value represents the confinement. So in our study case confinement is not strong.

Based on the above results, we discussed the structure functions in the scattering versus Jlab Hall C data (with order  $0.01 - 0.04$  of  $x$  and small  $q^2$ ). Numerically, we used an appropriate set of bulk mass and  $c$  in the form of a sweep spectrum which were already determined to fit experimental data and the structure function. With this formalism at finite  $x$  [30] our results are useful for understanding QCD and proton structure.

## Acknowledgement

Authors would like to thank Danning Li, Zi-qiang Zhang and Wei kou for useful discussions. This work was supported by Strategic Priority Research Program of Chinese Academy of Sciences (XDB34030301). ST is supported by the PIFI (2021PM0065).

## References

- [1] Sayipjamal Dulat, Tie Jiun Hou, Jun Gao, Marco Guzzi, Joey Huston, Pavel Nadolsky, Jon Pumplin, Carl Schmidt, Daniel Stump, C. P. Yuan , “*New parton distribution functions from a global analysis of quantum chromodynamics*”, *Phys. Rev. D* **93**, 033006 (2016) [arXiv: 1506.07443 [hep-ph]].
- [2] J. M. Maldacena, “*The Large  $N$  limit of superconformal field theories and supergravity*”, *Adv. Theor. Math. Phys* **2**, 231 (1998) [arXiv: 9711200 [hep-th]].
- [3] E. Witten, “*Anti-de Sitter space and holography*”, *Adv. Theor. Math. Phys* **2**, 253 (1998) [arXiv: 9802150 [hep-th]].
- [4] Joseph Polchinski and Matthew J. Strassler, “*Hard Scattering and Gauge/String Duality*”, *Phys. Rev. Lett* **88**, 031601 (2002) [arXiv: 0109174 [hep-th]].
- [5] N. R. F. Braga and A. Vega, “*Deep inelastic scattering of baryons in a modified soft wall model*”, *Eur. Phys. J. C* **72**, 2236 (2012) [arXiv: 1110.2548 [hep-ph]].
- [6] H. Boschi-Filho, N. Braga, F. Jugeau and M. Torres , “*Anomalous dimensions and scalar glueball spectroscopy in AdS/QCD*”, *Eur. Phys. J. C* **73**, 2540 (2013) [arXiv: 1208.2291[hep-ph]].
- [7] E. Folco Capossoli, D. Li and H. Boschi-Filho, “*Dynamical corrections to the anomalous holographic soft-wall model: the pomeron and the odderon*”, *Eur. Phys. J. C* **76**, no.6, 320 (2016) [arXiv:1604.01647 [hep-ph]].
- [8] M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, , “*QCD and resonance physics. theoretical foundations*”, *Nucl. Phys. B* **147**, 385 (1979).
- [9] R. Carcasses Quevedo, J. L. Goity, R. Trinchero, “*QCD condensates and holographic Wilson loops for asymptotically AdS spaces*”, *Phys. Rev.D* **89**, 036004 (2014)[arXiv: 1311.1175[hep-ph]].
- [10] M.D. Elia,A.D.Giacomo, E.Meggiolaro, “*Gauge-invariant field-strength correlators in pure Yang-Mills theory and full QCD at finite temperature*”, *Phys. Rev.D* **67** 114504 (2003) [arXiv: 0205018 [hep-lat]].
- [11] D.E. Miller, “*Lattice QCD Calculation for the Physical Equation of State*”, *Phys. Rept.* **443**, 55 (2007)[arXiv: 0608234[hep-ph]].
- [12] S.H. Lee, “*Gluon condensates above  $T_c$* ”, *Phys. Rev. D* **40**, 2484 (1989).



- [13] P. Colangelo, F. Giannuzzi, S. Nicotri, F. Zuo , “*Temperature and chemical potential dependence of the gluon condensate: a holographic study*”, *Phys. Rev. D* **88**, 115011 (2013) [arXiv: 1308.0489 [hep-ph]].
- [14] H. Leutwyler, “*in QCD 20 Years Later: Proceedings, Workshop, Aachen, Germany*”, 693 (1992).
- [15] P. Castorina, M. Mannarelli, “*Effective degrees of freedom and gluon condensation in the high temperature deconfined phase*”, *Phys.Rev.C* **75**, 054901 (2007) [arXiv: hep-ph/0701206].
- [16] Shin’ichi Nojiri, Sergei D. Odintsov, “*Two-Boundaries AdS/CFT Correspondence in Dilatonic Gravity*”, *Phys.Lett.B* **449** 39 (1999) [arXiv: 9812017[hep-th]].
- [17] Luis A. H. Mamani, “*Conformal symmetry breaking in holographic QCD*”, *Phys. Rev. D* **100** 106009 (2019) [arXiv: 1910.00026[hep-th]].
- [18] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “*Gauge Theory Correlators from Non-Critical String Theory*”, *Phys. Lett. B* **428** 105 (1998) [arXiv: hep-th/9802109].
- [19] Alfonso Ballon-Bayona, Henrique Boschi-Filho, Luis A. H. Mamani, Alex S. Miranda, Vilson T. Zanchin, “*An effective holographic approach to QCD*”, *Phys. Rev. D* **97** 4 046001 (2018) [arXiv: 1804.01579 [hep-th]].
- [20] Csaba Csaki, Matthew Reece , “*Toward a Systematic Holographic QCD: A Braneless Approach*”, *JHEP* **0705** 062 (2007) [arXiv: hep-ph/0608266].
- [21] Steven S. Gubser, “*Dilaton-driven confinement*”, *HUPT-A010* [arXiv: hep-th/9902155].
- [22] Y. Kim, B.H. Lee, C. Park, S.J. Sin “*The Effect of gluon condensate on holographic heavy quark potential*”, *Phys. Rev. D* **80** 105016 (2009) [arXiv: 0808.1143 [hep-th]].
- [23] Z.q. Zhang, X. Zhu, “*Effect of gluon condensate on jet quenching parameter and drag force*”, *Eur. Phys. J. C* **107** 79 (2019).
- [24] Kenji Morita, Su Houn Lee , “*Critical behavior of charmonia across the phase transition: A QCD sum rule approach*”, *Phys.Rev.C* **77** , 064904 (2008) [arXiv: 0711.3998 [hep-ph]]
- [25] G. Boyd, D. E. Miller, , “*The Temperature Dependence of the SU(N) Gluon Condensate from Lattice Gauge Theory*”, [arXiv: hep-ph/9608482].
- [26] Oleg Andreev, Valentin I. Zakharov , “*Gluon Condensate, Wilson Loops and Gauge/String Duality*”, *PhysRevD* **76** 047705 (2007) [arXiv: hep-ph/0703010].
- [27] Yumi Ko, Bum-Hoon Lee, Chanyong Park, “*Meson spectra in a gluon condensate background*”, *JHEP* **04** 037 (2010) [arXiv: 0912.5274 [hep-ph]].

- [28] Oleg Andreev, “Nonlocal and Mixed Condensates Tests of Gauge/String Duality”, *Phys.Rev.D* **82** 086012 (2010) [arXiv: 1008.4738 [hep-ph]].
- [29] Eduardo Folco Capossoli, Miguel Angel Martín Contreras, Danning Li, Alfredo Vega, Henrique Boschi-Filho , “Proton Structure Functions from an AdS/QCD model with a deformed background”, *Phys. Rev. D* **102**, 086004 (2020) [arXiv: 2007.09283 [hep-ph]].
- [30] J. Polchinski and M. J. Strassler, “Deep inelastic scattering and gauge / string duality”, *JHEP* **0305**, 012 (2003) [arXiv: 0209211 [hep-th]].
- [31] Y.Hatta, E.Iancu and A.H.Mueller, “Deep inelastic scattering at strong coupling from gauge/string duality: The Saturation line”, *JHEP* **0801**, 026 (2008) [arXiv: 0710.2148 [hep-th]].
- [32] C. A. Ballon Bayona, H. Boschi-Filho and N. R. F. Braga, “Deep inelastic structure functions from supergravity at small  $x$ ”, *JHEP* **0810**, 088 (2008) [arXiv: 0712.3530[hep-th]].
- [33] C. A. Ballon Bayona, H. Boschi-Filho and N. R. F. Braga, *Deep inelastic scattering from gauge string duality in the soft wall model*, *JHEP* **0803**, 064 (2008) [arXiv: 0711.0221 [hep-th]].
- [34] L. Cornalba and M. S. Costa, “Saturation in Deep Inelastic Scattering from AdS/CFT”, *Phys. Rev. D* **78**, 096010 (2008) [arXiv: 0804.1562 [hep-ph]].
- [35] B. Pire, C. Roiesnel, L. Szymanowski and S. Wallon, “On AdS/QCD correspondence and the partonic picture of deep inelastic scattering”, *Phys. Lett. B* **670**, 84 (2008) [arXiv: 0805.4346 [hep-ph]].
- [36] J. L. Albacete, Y. V. Kovchegov and A. Taliotis, “DIS on a Large Nucleus in AdS/CFT”, *JHEP* **0807**, 074 (2008) [arXiv: 0806.1484 [hep-th]].
- [37] C. A. Ballon Bayona, H. Boschi-Filho and N. R. F. Braga, “Deep inelastic scattering from gauge string duality in D3-D7 brane model”, *JHEP* **0809**, 114 (2008) [arXiv: 0807.1917 [hep-th]].
- [38] J. H. Gao and B. W. Xiao, “Polarized Deep Inelastic and Elastic Scattering From Gauge/String Duality”, *Phys. Rev. D* **80**, 015025 (2009) [arXiv: 0904.2870 [hep-ph]].
- [39] A. Taliotis, “DIS from the AdS/CFT correspondence”, *Nucl. Phys. A* **830**, 299C (2009) [arXiv: 0907.4204 [hep-th]].
- [40] Y. Yoshida, “The Virtual Photon Structure Functions and AdS/QCD Correspondence”, *Prog. Theor. Phys* **123**, 79 (2010) [arXiv: 0902.1015 [hep-th]].

- [41] Y. Hatta, T. Ueda and B. W. Xiao, “Polarized DIS in  $N=4$  SYM: Where is spin at strong coupling?”, *JHEP* **0908**, 007 (2009) [arXiv: 0905.2493 [hep-ph]].
- [42] E. Avsar, E. Iancu, L. McLerran and D. N. Triantafyllopoulos, “Shockwaves and deep inelastic scattering within the gauge/gravity duality”, *JHEP* **0911**, 105 (2009) [arXiv: 0907.4604 [hep-th]].
- [43] L. Cornalba, M. S. Costa and J. Penedones, “Deep Inelastic Scattering in Conformal QCD”, *JHEP* **1003**, 133 (2010) [arXiv: 0911.0043 [hep-th]].
- [44] C. A. B. Bayona, H. Boschi-Filho and N. R. F. Braga, “Deep inelastic scattering off a plasma with flavour from  $D3$ - $D7$  brane model”, *Phys. Rev. D* **81**, 086003 (2010) [arXiv: 0912.0231 [hep-th]].
- [45] L. Cornalba, M. S. Costa and J. Penedones, “AdS black disk model for small- $x$  DIS”, *Phys. Rev. Lett* **105**, 072003 (2010) [arXiv: 1001.1157 [hep-ph]].
- [46] R. C. Brower, M. Djuric, I. Sarcevic and C. I. Tan, “String-Gauge Dual Description of Deep Inelastic Scattering at Small- $x$ ”, *JHEP* **1011**, 051 (2010) [arXiv: 1007.2259 [hep-ph]].
- [47] J. H. Gao and Z. G. Mou, “Polarized Deep Inelastic Scattering Off the Neutron From Gauge/String Duality”, *Phys. Rev. D* **81**, 096006 (2010) [arXiv: 1003.3066 [hep-ph]].
- [48] C. A. Ballon Bayona, H. Boschi-Filho, N. R. F. Braga and M. A. C. Torres, “Deep inelastic scattering for vector mesons in holographic  $D4$ - $D8$  model”, *JHEP* **1010**, 055 (2010) [arXiv: 1007.2448 [hep-th]].
- [49] E. Koile, S. Macaluso and M. Schvellinger, “Deep inelastic scattering structure functions of holographic spin-1 hadrons with  $N_f \geq 1$ ”, *JHEP* **1401**, 166 (2014) [arXiv: 1311.2601 [hep-th]].
- [50] E. Koile, N. Kovensky and M. Schvellinger, “Hadron structure functions at small  $x$  from string theory”, *JHEP* **1505**, 001 (2015) [arXiv: 1412.6509 [hep-th]].
- [51] J. H. Gao and Z. G. Mou, “Structure functions in deep inelastic scattering from gauge/string duality beyond single-hadron final states”, *Phys. Rev. D* **90**, 075018 (2014) [arXiv: 1406.7576 [hep-ph]].
- [52] E. Folco Capossoli and H. Boschi-Filho, “Deep Inelastic Scattering in the Exponentially Small Bjorken Parameter Regime from the Holographic Softwall Model”, *Phys. Rev. D* **92**, 126012 (2015) [arXiv: 1509.01761 [hep-th]].
- [53] E. Koile, N. Kovensky and M. Schvellinger, “Deep inelastic scattering cross sections from the gauge/string duality”, *JHEP* **1512**, 009 (2015) [arXiv: 1507.07942 [hep-th]].

- [54] D. Jorin, N. Kovensky and M. Schvellinger, “Towards  $1/N$  corrections to deep inelastic scattering from the gauge/gravity duality”, *JHEP* **1604**, 113 (2016) [arXiv: 1601.01627 [hep-th]].
- [55] D. Jorin, M. Schvellinger and N. Kovensky, “Deep inelastic scattering off scalar mesons in the  $1/N$  expansion from the  $D3D7$ -brane system”, *JHEP* **1612**, 003 (2016) [arXiv: 1609.01202 [hep-th]].
- [56] N. Kovensky, G. Michalski and M. Schvellinger, “DIS off glueballs from string theory: the role of the chiral anomaly and the Chern-Simons term”, *JHEP* **1804**, 118 (2018) [arXiv: 1711.06171 [hep-th]].
- [57] A. Amorim, R. Carcassés Quevedo and M. S. Costa, “Nonminimal coupling contribution to DIS at low  $x$  in Holographic QCD”, *Phys. Rev. D* **98**, 026016 (2018) [arXiv: 1804.07778 [hep-ph]].
- [58] G. F. de Teramond *et al.* “Universality of Generalized Parton Distributions in Light-Front Holographic QCD”, *Phys. Rev. Lett* **120**, 182001 (2018) [arXiv: 1801.09154 [hep-ph]].
- [59] T. Liu, R. S. Sufian, G. F. de Téramond, H. G. Dosch, S. J. Brodsky and A. Deur, “Unified Description of Polarized and Unpolarized Quark Distributions in the Proton”, *Phys. Rev. Lett* **124**, 082003 (2020) [arXiv: 1909.13818 [hep-ph]].
- [60] A. Watanabe, T. Sawada and M. Huang, “Extraction of gluon distributions from structure functions at small  $x$  in holographic QCD”, *Phys. Lett. B* **805**, 135470 (2020) [arXiv: 1910.10008 [hep-ph]].
- [61] D. Jorin, G. Michalski and M. Schvellinger, “Spin-1/2 fermionic operators of  $\mathcal{N} = 4$  SYM theory and DIS from type IIB supergravity”, *JHEP* **06**, 063 (2020) [arXiv: 2004.02909 [hep-th]].
- [62] Aneesh V. Manohar, “An Introduction to Spin Dependent Deep Inelastic Scattering”, *Contribution to: Lake Louise Winter Institute: Symmetry and Spin in the Standard Model*, [arXiv: 9204208 [hep-ph]].
- [63] Mans Henningson, Konstadinos Sfetsos, “Spinors and the AdS/CFT correspondence”, *Phys. Lett. B* **431**, 63 (1998) [arXiv: 9803251 [hep-th]].
- [64] W. Mück, K. S. Viswanathan, “Conformal Field Theory Correlators from Classical Field Theory on Anti-de Sitter Space II. Vector and Spinor Fields”, *Phys. Rev. D* **58**, 106006 (1998) [arXiv: 9805145 [hep-th]].
- [65] Ingo Kirsch, “Spectroscopy of fermionic operators in AdS/CFT”, *JHEP* **0609**, 052 (2006) [arXiv: 0607205 [hep-th]].
- [66] Theodoros Nakas, Konstantinos S. Rigatos, “Fermions and baryons as open-string states from brane junctions”, *JHEP* **12**, (2020) 157 [arXiv: 2010.00025 [hep-th]].

- [67] Zainul Abidin, Carl E. Carlson, “*Nucleon electromagnetic and gravitational form factors from holography*”, *Phys. Rev. D* **79**, 115003 (2009) [arXiv: 0903.4818 [hep-ph]].
- [68] V. Tvaskis, et al , *The proton and deuteron  $F_2$  structure function at low  $Q^2$* , *Phys.Rev.C* **81** 055207 (2010) [arXiv: 1002.1669 [nucl-ex]].