

On the phase diagram of a three-dimensional dipolar model.

V. Russier^{1,*} and Juan J. Alonso^{2,3,†}

¹*ICMPE, UMR 7182 CNRS and UPE 2-8 rue Henri Dunant 94320 Thiais, France.*

²*Física Aplicada I, Universidad de Málaga, 29071 Málaga, Spain*

³*Instituto Carlos I de Física Teórica y Computacional, Universidad de Málaga, 29071 Málaga, Spain*

(Dated: January 17, 2022)

The magnetic phase diagram at zero external field of an ensemble of dipoles with uniaxial anisotropy on a FCC lattice has been investigated from tempered Monte Carlo simulations. The uniaxial anisotropy is characterized by a random distribution of easy axes and its magnitude λ_u is the driving force of disorder and consequently frustration. The phase diagram, separating the paramagnetic, ferromagnetic and spin-glass regions, was thus considered in the temperature, λ_u plane. Here we interpret this phase diagram in terms of the more convenient variables namely the bare dipolar interaction and anisotropy energies ϵ_d and ϵ_u on the one hand and the volume fraction Φ on the other hand and compare the result with that corresponding to the random distribution of particles in the absence of anisotropy. We also display the nature of the ordered phase reached at low temperature by the ensemble of dipoles on the FCC lattice in terms of both the dipolar coupling and the texturation of the easy axes distribution when the latter is no more random. This system is aimed at modeling the magnetic phase diagram of supracrystals of magnetic nanoparticles.

In Ref.¹ we investigated the phase diagram of an ensemble of dipolar hard spheres located on the nodes of a perfect FCC lattice. The phase diagram was investigated in the reduced temperature, disorder control parameter, (T^*, λ_u) plane. Given the $1/r^3$ dependence of the dipole-dipole interaction (DDI), it has been found convenient to define the reduced temperature, $T^* = T/T_r$ through the reference temperature $T_r = (\Phi/\Phi_m)(\epsilon_d/k_B)$ where Φ , Φ_m and ϵ_d are the volume fraction, its maximum value of the FCC lattice and the dipole-dipole energy for particles at contact. The disorder control parameter is nothing but the anisotropy energy (MAE) over DDI ratio, namely $\lambda_u = (\epsilon_u/\epsilon_d)(\Phi_m/\Phi)$. Then the (T^*, λ_u) phase diagram, displayed on figure (1), is qualitatively similar to those corresponding to the $\pm J$ Ising model^{2,3}. The essential feature is the succession at low temperature of long range order FM, quasi long range FM (QLRO-FM) plus transverse SG and SG phases with increasing values of λ_u . Now, when considering the system as a model for the magnetic phase diagram of supracrystals of bare magnetic nanoparticles characterized by given values of ϵ_d and ϵ_u where the volume fraction ϕ can be tuned through the thickness of the coating layer, it is convenient to translate the phase diagram in terms of $(k_B T/\epsilon_u)$ and $(\Phi/\Phi_m)(\epsilon_d/\epsilon_u)$, the latter being understood as the increasing dipolar coupling through the increase of concentration at constant ϵ_d and ϵ_u . This is shown on figure (2). In this representation, the dipolar lattice phase diagram appears qualitatively similar to the one deduced from experiments on the DMIM samples⁴. Furthermore one must limit the reachable part of this phase diagram to the region defined by $\Phi/\Phi_m \leq 1$. Doing this we see, as displayed on figure (3), that according to the value of (ϵ_d/ϵ_u) , namely the relative importance of the dipolar coupling to the anisotropy energy, the FM-QLRO and the FM phases may be not reachable. In the limit $\epsilon_d/\epsilon_u \rightarrow \infty$, one gets only the PM/FM transition of the pure dipolar system on the FCC lattice free of anisotropy ($\epsilon_u = 0$)^{1,5,6} and all the non trivial features of the phase diagram are shrunk on the $\Phi = 0$ line. Moreover, the $((k_B T/\epsilon_u), (\Phi/\Phi_m)(\epsilon_d/\epsilon_u))$ phase diagram for particles located on the perfect FCC lattice can be directly compared to the one we have got⁷ for the system of dipolar spheres frozen in a random hard sphere like distribution, in the absence of anisotropy, where the increasing disorder is quantified by $1 - (\phi/\phi_{RCP})$, the volume fraction being then limited to the so-called RCP value, $\Phi_{RCP} \approx 0.64$, given on figure (4). It is also qualitatively close

* e-mail address: russier@icmpe.cnrs.fr

† e-mail address: jjalonso@uma.es

to the one of dipolar Ising model on the SC lattice in terms of dilution⁸, where however the ordered phase is anti-FM due to the lattice symmetry. Finally, we present on figure (5) the nature of the ordered phase at low temperature in terms of the dipolar coupling and the texturation of the easy axes distribution as was studied in^{9,10} in the limit of infinitely strong anisotropy energy leading to the dipolar Ising model. As is the case above, the condition $\Phi/\Phi_m \leq 1$ restricts the diagram to the left hand part defined by the abscissa $\leq (\epsilon_d/\epsilon_u)$ and has been displayed for two particular cases corresponding to a weak and a strong dipolar coupling respectively.




FIG. 1. $(T_c/T_r, \lambda_u)$ phase diagram of the dipolar plus uniaxial anisotropy with random distribution of easy axes on a FCC lattice. λ_u is the natural disorder control parameter.




FIG. 2. Phase diagram of figure (1), shown as T_c in terms of increasing dipolar coupling for ϵ_d and ϵ_u considered as constant.




FIG. 3. Same as figure (2) where the constraint $\Phi / \Phi_m \leq 1$ is explicated for three characteristic values of ϵ_d / ϵ_u as indicated.




FIG. 4. Phase diagram of the dipolar hard spheres distributed according to a frozen hard-sphere like distribution with $\epsilon_u = 0$. The limiting value of the volume fraction, Φ_{RCP} is indicated.




FIG. 5. Nature of the ordered phase at low temperature for the FCC lattice in terms of the texturation quantified by the variance σ of the easy axes distribution (see Ref.^{9,10}) and the dipolar coupling $(\Phi/\Phi_m)\epsilon_d/\epsilon_u$. Open circle and open triangle are the FM/SG transition in the dipolar Ising limit¹⁰, and $(\Phi/\Phi_m)\epsilon_d/\epsilon_u = 0.25$ ¹¹ respectively. The dotted line is only indicative. The upper horizontal red line, $\sigma = \pi/2$ indicates the lower bound of random distribution of easy axes. The SG region is separated in the true dipolar SG (3D) and the limiting dipolar Ising SG (1D) obtained for $\lambda_u > 30$ ¹. The reachable region is located on the left hand side of the line $(\Phi/\Phi_m)\epsilon_d/\epsilon_u = \epsilon_d/\epsilon_u$ which is shown for a weak and a strong dipolar couplings as particular cases.

-
- ¹ V. Russier, J. J. Alonso, I. Lisiecki, A. T. Ngo, C. Salzemann, S. Nakamae, and C. Raepsaet, Phys. Rev. B **102**, 174410 (2020).
- ² M. Hasenbusch, F. P. Toldin, A. Pelissetto, and E. Vicari, Phys. Rev. B **76**, 184202 (2007).
- ³ T. Papakonstantinou and A. Malakis, Phys. Rev. E **87**, 012132 (2013).
- ⁴ O. Petracic, Superlattices and Microstructures **47**, 569 (2010).
- ⁵ J. P. Bouchaud and P. G. Zerah, Phys. Rev. B **47**, 9095 (1993).
- ⁶ V. Russier and E. Ngo, Condensed Matter Physics **20**, 33703 (2017).
- ⁷ J. J. Alonso, B. Allés, and V. Russier, Phys. Rev. B **102**, 184423 (2020).
- ⁸ J.-J. Alonso and J.-F. Fernández, Phys. Rev. B **81**, 064408 (2010).
- ⁹ J.-J. Alonso, B. Allés, and V. Russier, Phys. Rev. B **100**, 134409 (2019), arXiv:1909.13573.
- ¹⁰ V. Russier and J. J. Alonso, J. Phys.-Cond. Matt. **32**, 135804 (2020).
- ¹¹ V. Russier, (2020), unpublished.