

# ON TWO CONGRUENCE CONJECTURES OF Z.-W. SUN INVOLVING FRANEL NUMBERS

GUO-SHUAI MAO AND YAN LIU

**ABSTRACT.** In this paper, we mainly prove the following conjectures of Z.-W. Sun [17]: Let  $p > 2$  be a prime. If  $p = x^2 + 3y^2$  with  $x, y \in \mathbb{Z}$  and  $x \equiv 1 \pmod{3}$ , then

$$x \equiv \frac{1}{4} \sum_{k=0}^{p-1} (3k+4) \frac{f_k}{2^k} \equiv \frac{1}{2} \sum_{k=0}^{p-1} (3k+2) \frac{f_k}{(-4)^k} \pmod{p^2},$$

and if  $p \equiv 1 \pmod{3}$ , then

$$\sum_{k=0}^{p-1} \frac{f_k}{2^k} \equiv \sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} \pmod{p^3},$$

where  $f_n = \sum_{k=0}^n \binom{n}{k}^3$  stands for the  $n$ th Franel number.

## 1. INTRODUCTION

In 1894, Franel [2] found that the numbers

$$f_n = \sum_{k=0}^n \binom{n}{k}^3 \quad (n = 0, 1, 2, \dots)$$

satisfy the recurrence relation (cf. [12, A000172]):

$$(n+1)^2 f_{n+1} = (7n^2 + 7n + 2) f_n + 8n^2 f_{n-1} \quad (n = 1, 2, 3, \dots).$$

These numbers are now called Franel numbers. Callan [1] found a combinatorial interpretation of the Franel numbers. The Franel numbers play important roles in combinatorics and number theory. The sequence  $(f_n)_{n \geq 0}$  is one of the five sporadic sequences (cf. [21, Section 4]) which are integral solutions of certain Apéry-like recurrence equations and closely related to the theory of modular forms. In 2013, Sun [17] revealed some unexpected connections between the numbers

---

*Key words and phrases.* Congruences; franel numbers;  $p$ -adic Gamma function; Gamma function.

*Mathematics Subject Classification.* Primary 11A07; Secondary 05A10; 05A19; 33B15.

The first author is the corresponding author. This research was supported by the Natural Science Foundation of China (grant 12001288).

$f_n$  and representations of primes  $p \equiv 1 \pmod{3}$  in the form  $x^2 + 3y^2$  with  $x, y \in \mathbb{Z}$ , for example, Z.-W. Sun [17, (1.2)] showed that

$$\sum_{k=0}^{p-1} \frac{f_k}{2^k} \equiv \sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} \equiv 2x - \frac{p}{2x} \pmod{p^2}, \quad (1.1)$$

and in the same paper, Sun proposed some conjectures involving Franel numbers, one of which is

**Conjecture 1.1.** *Let  $p > 2$  be a prime. If  $p = x^2 + 3y^2$  with  $x, y \in \mathbb{Z}$  and  $x \equiv 1 \pmod{3}$ , then*

$$x \equiv \frac{1}{4} \sum_{k=0}^{p-1} (3k+4) \frac{f_k}{2^k} \equiv \frac{1}{2} \sum_{k=0}^{p-1} (3k+2) \frac{f_k}{(-4)^k} \pmod{p^2}.$$

For more studies on Franel numbers, we refer the readers to ([3, 4, 6, 7, 8, 15, 18] and so on).

In this paper, our first goal is to prove the above conjecture.

**Theorem 1.1.** *Conjecture 1.1 is true.*

Z.-W. Sun [17] also gave the following conjecture which is much difficult and complex.

**Conjecture 1.2.** *Let  $p > 2$  be a prime. If  $p \equiv 1 \pmod{3}$ , then*

$$\sum_{k=0}^{p-1} \frac{f_k}{2^k} \equiv \sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} \pmod{p^3}.$$

Our last goal is to prove this conjecture.

**Theorem 1.2.** *Conjecture 1.2 is true.*

We are going to prove Theorem 1.1 in Section 2. Section 3 is devoted to proving Theorem 1.2. Our proofs make use of some combinatorial identities which were found by the package **Sigma** [11] via the software **Mathematica** and the  $p$ -adic Gamma function. The proof of Theorem 1.2 is somewhat difficult and complex because it is rather convoluted. Throughout this paper, prime  $p$  always  $\equiv 1 \pmod{3}$ , so in the following Lemmas  $p > 5$  or  $p > 3$  or  $p > 2$  is the same, we mention it here first.

## 2. PROOF OF THEOREM 1.1

For a prime  $p$ , let  $\mathbb{Z}_p$  denote the ring of all  $p$ -adic integers and let  $\mathbb{Z}_p^\times := \{a \in \mathbb{Z}_p : a \text{ is prime to } p\}$ . For each  $\alpha \in \mathbb{Z}_p$ , define the  $p$ -adic

order  $\nu_p(\alpha) := \max\{n \in \mathbb{N} : p^n \mid \alpha\}$  and the  $p$ -adic norm  $|\alpha|_p := p^{-\nu_p(\alpha)}$ . Define the  $p$ -adic gamma function  $\Gamma_p(\cdot)$  by

$$\Gamma_p(n) = (-1)^n \prod_{\substack{1 \leq k < n \\ (k, p) = 1}} k, \quad n = 1, 2, 3, \dots,$$

and

$$\Gamma_p(\alpha) = \lim_{\substack{|\alpha - n|_p \rightarrow 0 \\ n \in \mathbb{N}}} \Gamma_p(n), \quad \alpha \in \mathbb{Z}_p.$$

In particular, we set  $\Gamma_p(0) = 1$ . Following, we need to use the most basic properties of  $\Gamma_p$ , and all of them can be found in [9, 10]. For example, we know that

$$\frac{\Gamma_p(x+1)}{\Gamma_p(x)} = \begin{cases} -x, & \text{if } |x|_p = 1, \\ -1, & \text{if } |x|_p > 1. \end{cases} \quad (2.1)$$

$$\Gamma_p(1-x)\Gamma_p(x) = (-1)^{a_0(x)}, \quad (2.2)$$

where  $a_0(x) \in \{1, 2, \dots, p\}$  such that  $x \equiv a_0(x) \pmod{p}$ . And a property we need here is the fact that for any positive integer  $n$ ,

$$z_1 \equiv z_2 \pmod{p^n} \quad \text{implies} \quad \Gamma_p(z_1) \equiv \Gamma_p(z_2) \pmod{p^n}. \quad (2.3)$$

**Lemma 2.1.** ([17, Lemma 2.2]) *For any  $n \in \mathbb{N}$  we have*

$$\sum_{k=0}^n \binom{n}{k}^3 z^k = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n+k}{3k} \binom{2k}{k} \binom{3k}{k} z^k (1+z)^{n-2k} \quad (2.4)$$

and

$$f_n = \sum_{k=0}^n \binom{n+2k}{3k} \binom{2k}{k} \binom{3k}{k} (-4)^{n-k}. \quad (2.5)$$

For  $n, m \in \{1, 2, 3, \dots\}$ , define

$$H_n^{(m)} = \sum_{1 \leq k \leq n} \frac{1}{k^m},$$

these numbers with  $m = 1$  are often called the classic harmonic numbers.

Recall that the Bernoulli polynomials are given by

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_k x^{n-k} \quad (n = 0, 1, 2, \dots).$$

**Lemma 2.2.** ([13, 14]) *Let  $p > 5$  be a prime. Then*

$$\begin{aligned}
H_{p-1}^{(2)} &\equiv 0 \pmod{p}, \quad H_{\frac{p-1}{2}}^{(2)} \equiv 0 \pmod{p}, \quad H_{p-1} \equiv 0 \pmod{p^2}, \\
\frac{1}{5}H_{\lfloor \frac{p}{6} \rfloor}^{(2)} &\equiv H_{\lfloor \frac{p}{3} \rfloor}^{(2)} \equiv \frac{1}{2} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) \pmod{p}, \\
H_{\lfloor \frac{p}{6} \rfloor} &\equiv -2q_p(2) - \frac{3}{2}q_p(3) + pq_p^2(2) + \frac{3p}{4}q_p^2(3) - \frac{5p}{12} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) \pmod{p^2}, \\
H_{\lfloor \frac{p}{3} \rfloor} &\equiv -\frac{3}{2}q_p(3) + \frac{3p}{4}q_p^2(3) - \frac{p}{6} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) \pmod{p^2}, \\
H_{\frac{p-1}{2}} &\equiv -2q_p(2) + pq_p^2(2) \pmod{p^2}, \quad H_{\lfloor \frac{p}{4} \rfloor}^{(2)} \equiv (-1)^{\frac{p-1}{2}} 4E_{p-3} \pmod{p}, \\
H_{\lfloor \frac{2p}{3} \rfloor} &\equiv -\frac{3}{2}q_p(3) + \frac{3p}{4}q_p^2(3) + \frac{p}{3} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) \pmod{p^2},
\end{aligned}$$

where  $q_p(a) = (a^{p-1} - 1)/p$  stands for the Fermat quotient.

**Lemma 2.3.** *Let  $p > 2$  be a prime and  $p \equiv 1 \pmod{3}$ . If  $0 \leq j \leq (p-1)/2$ , then we have*

$$\binom{3j}{j} \binom{p+j}{3j+1} \equiv \frac{p}{3j+1} (1 - pH_{2j} + pH_j) \pmod{p^3}.$$

*Proof.* If  $0 \leq j \leq (p-1)/2$  and  $j \neq (p-1)/3$ , then we have

$$\begin{aligned}
\binom{3j}{j} \binom{p+j}{3j+1} &= \frac{(p+j) \cdots (p+1)p(p-1) \cdots (p-2j)}{j!(2j)!(3j+1)} \\
&\equiv \frac{pj!(1 + pH_j)(-1)^{2j}(2j)!(1 - pH_{2j})}{j!(2j)!(3j+1)} \\
&\equiv \frac{p}{3j+1} (1 - pH_{2j} + pH_j) \pmod{p^3}.
\end{aligned}$$

If  $j = (p-1)/3$ , then by Lemma 2.2, we have

$$\begin{aligned}
&\binom{p-1}{\frac{p-1}{3}} \binom{p+\frac{p-1}{3}}{\frac{p-1}{3}} \\
&\equiv \left( 1 - pH_{\frac{p-1}{3}} + \frac{p^2}{2}(H_{\frac{p-1}{3}}^2 - H_{\frac{p-1}{3}}^{(2)}) \right) \left( 1 + pH_{\frac{p-1}{3}} + \frac{p^2}{2}(H_{\frac{p-1}{3}}^2 - H_{\frac{p-1}{3}}^{(2)}) \right) \\
&\equiv 1 - p^2 H_{\frac{p-1}{3}}^{(2)} \equiv 1 - \frac{p^2}{2} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) \pmod{p^3}
\end{aligned}$$

and

$$1 - pH_{\frac{2p-2}{3}} + pH_{\frac{p-1}{3}} \equiv 1 - \frac{p^2}{2} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) \pmod{p^3}.$$

Now the proof of Lemma 2.3 is complete.  $\square$

*Proof of Theorem 1.1.* With the help of (2.4), we have

$$\begin{aligned} \sum_{k=0}^{p-1} (3k+4) \frac{f_k}{2^k} &= \sum_{k=0}^{p-1} \frac{3k+4}{2^k} \sum_{j=0}^{\lfloor k/2 \rfloor} \binom{k+j}{3j} \binom{2j}{j} \binom{3j}{j} 2^{k-2j} \\ &= \sum_{j=0}^{(p-1)/2} \frac{\binom{2j}{j} \binom{3j}{j}}{4^j} \sum_{k=2j}^{p-1} (3k+4) \binom{k+j}{3j}. \end{aligned} \quad (2.6)$$

By loading the package **Sigma** in the software **Mathematica**, we find the following identity:

$$\sum_{k=2j}^{n-1} (3k+4) \binom{k+j}{3j} = \frac{9nj+3n+9j+5}{3j+2} \binom{n+j}{3j+1}.$$

Thus, replacing  $n$  by  $p$  in the above identity and then substitute it into (2.6), we have

$$\sum_{k=0}^{p-1} (3k+4) \frac{f_k}{2^k} = \sum_{j=0}^{(p-1)/2} \frac{\binom{2j}{j} \binom{3j}{j}}{4^j} \frac{9pj+3p+9j+5}{3j+2} \binom{p+j}{3j+1}.$$

Hence we immediately obtain the following result by Lemma 2.3,

$$\sum_{k=0}^{p-1} (3k+4) \frac{f_k}{2^k} \equiv p \sum_{j=0}^{(p-1)/2} \frac{\binom{2j}{j}}{4^j} \frac{9j+5}{(3j+1)(3j+2)} \pmod{p^2}. \quad (2.7)$$

It is easy to verify that

$$p \sum_{j=0}^{(p-1)/2} \frac{\binom{2j}{j}}{4^j} \frac{9j+5}{(3j+1)(3j+2)} \equiv S_1 + S_2 \pmod{p^2}, \quad (2.8)$$

where

$$S_1 = p \sum_{j=0}^{(p-1)/2} \binom{(p-1)/2}{j} (-1)^j \left( \frac{2}{3j+1} + \frac{1}{3j+2} \right) \quad (2.9)$$

and

$$S_2 = \frac{3p+2}{p+1} \left( \binom{(2p-2)/3}{(p-1)/3} 4^{(1-p)/3} - \binom{(p-1)/2}{(p-1)/3} \right).$$

Applying the famous identity

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{k+x} = \frac{n!}{x(x+1) \cdots (x+n)} \quad (2.10)$$

with  $x = 1/3, n = (p-1)/2$  and  $x = 2/3, n = (p-1)/2$ , we may simplify (2.9) as

$$S_1 = \frac{4p}{3p-1} \frac{(1)_{(p-1)/2}}{(1/3)_{(p-1)/2}} + \frac{2p}{3p+1} \frac{(1)_{(p-1)/2}}{(2/3)_{(p-1)/2}},$$

where  $(a)_n = a(a+1)\cdots(a+n-1)$  is the rising factorial or the Pochhammer symbol.

In view of (2.2), we have

$$\begin{aligned} \frac{4p}{3p-1} \frac{(1)_{\frac{p-1}{2}}}{(1/3)_{\frac{p-1}{2}}} &= \frac{4p}{3p-1} \frac{\Gamma(\frac{p+1}{2})\Gamma(\frac{1}{3})}{\Gamma(\frac{1}{3} + \frac{p-1}{2})} = \frac{4p}{3p-1} \frac{(-1)^{\frac{p+1}{2}}\Gamma_p(\frac{p+1}{2})\Gamma_p(\frac{1}{3})}{(-1)^{\frac{p-1}{2}}\frac{p}{3}\Gamma_p(\frac{1}{3} + \frac{p-1}{2})} \\ &= \frac{12}{1-3p} \frac{\Gamma_p(\frac{p+1}{2})\Gamma_p(\frac{1}{3})}{\Gamma_p(\frac{p}{2} - \frac{1}{6})} = \frac{12(-1)^{\frac{p-1}{6}}}{1-3p} \Gamma_p\left(\frac{p+1}{2}\right) \Gamma_p\left(\frac{1}{3}\right) \Gamma_p\left(\frac{7}{6} - \frac{p}{2}\right), \end{aligned}$$

where  $\Gamma(\cdot)$  is the Gamma function. It is known that for  $\alpha, s \in \mathbb{Z}_p$ , we have

$$\Gamma_p(\alpha + ps) \equiv \Gamma_p(\alpha) + ps\Gamma'_p(\alpha) \pmod{p^2} \quad (2.11)$$

and

$$\frac{\Gamma'_p(\alpha)}{\Gamma_p(\alpha)} \equiv 1 + H_{p-\langle-\alpha\rangle_{p-1}} \pmod{p}, \quad (2.12)$$

where  $\Gamma'_p(x)$  denotes the  $p$ -adic derivative of  $\Gamma_p(x)$ ,  $\langle\alpha\rangle_n$  denotes the least non-negative residue of  $\alpha$  modulo  $n$ , i.e., the integer lying in  $\{0, 1, \dots, n-1\}$  such that  $\langle\alpha\rangle_n \equiv \alpha \pmod{n}$ .

Therefore modulo  $p^2$ , we have

$$\frac{4p}{3p-1} \frac{(1)_{\frac{p-1}{2}}}{(\frac{1}{3})_{\frac{p-1}{2}}} \equiv \frac{12(-1)^{\frac{p-1}{6}}\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{3})\Gamma_p(\frac{7}{6})}{1-3p} \left(1 + \frac{p}{2}(H_{\frac{p-1}{2}} - H_{\frac{p-7}{6}})\right).$$

In view of (2.1) and (2.2), we have

$$\frac{4p}{3p-1} \frac{(1)_{\frac{p-1}{2}}}{(\frac{1}{3})_{\frac{p-1}{2}}} \equiv \frac{2(1+3p)\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{3})}{\Gamma_p(\frac{5}{6})} \left(1 + \frac{p}{2}(H_{\frac{p-1}{2}} - H_{\frac{p-7}{6}})\right) \pmod{p^2}.$$

And then by using [20, Proposition 4.1], we have

$$\frac{\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{3})}{\Gamma_p(\frac{5}{6})} \equiv \frac{\binom{(5p-5)/6}{(p-1)/3}}{\left(1 + \frac{p}{6}(5H_{(5p-5)/6} - 3H_{(p-1)/2} - 2H_{(p-1)/3})\right)} \pmod{p^2}.$$

Then with the help of [20, Theorem 4.12] and Lemma 2.2, we have

$$\frac{4p}{3p-1} \frac{(1)_{\frac{p-1}{2}}}{(1/3)_{\frac{p-1}{2}}} \equiv 4x + 3pxq_p(3) - \frac{p}{x} \pmod{p^2} \quad (2.13)$$

and

$$\frac{2p}{3p+1} \frac{(1)_{(p-1)/2}}{(2/3)_{(p-1)/2}} \equiv \frac{p}{x} \pmod{p^2}. \quad (2.14)$$

Hence

$$S_1 \equiv 4x + 3pxq_p(3) \pmod{p^2}. \quad (2.15)$$

**Lemma 2.4.** *Let  $p > 3$  be a prime. For any  $p$ -adic integer  $t$ , we have*

$$\binom{\frac{p-1}{2} + pt}{\frac{p-1}{3}} \equiv \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \left( 1 + pt \left( H_{\frac{p-1}{2}} - H_{\frac{p-1}{6}} \right) \right) \pmod{p^2}. \quad (2.16)$$

*Proof.* Set  $m = (p-1)/2$ . It is easy to check that

$$\begin{aligned} \binom{m+pt}{(p-1)/3} &= \frac{(m+pt) \cdots (m+pt - (p-1)/3 + 1)}{((p-1)/3)!} \\ &\equiv \frac{m \cdots (m - (p-1)/3 + 1)}{((p-1)/3)!} (1 + pt(H_m - H_{m-(p-1)/3})) \\ &= \binom{m}{(p-1)/3} (1 + pt(H_m - H_{m-(p-1)/3}) \pmod{p^2}). \end{aligned}$$

So Lemma 2.4 is finished.  $\square$

Now we evaluate  $S_2$  modulo  $p^2$ . It is easy to obtain that

$$\begin{aligned} S_2 &\equiv 2 \left( \binom{-\frac{1}{2}}{(p-1)/3} - \binom{(p-1)/2}{(p-1)/3} \right) \\ &\equiv -p \binom{(p-1)/2}{(p-1)/3} (H_{(p-1)/2} - H_{(p-1)/6}) \\ &\equiv -3pxq_p(3) \pmod{p^2} \end{aligned} \quad (2.17)$$

with the help of Lemma 2.2, Lemma 2.4 and [20, Theorem 4.12].

Therefore, in view of (2.7), (2.8), (2.15) and (2.17), we immediately get the desired result

$$\frac{1}{4} \sum_{k=0}^{p-1} (3k+4) \frac{f_k}{2^k} \equiv x \pmod{p^2}.$$

On the other hand, we use the equation (2.5) to obtain that

$$\begin{aligned} \sum_{k=0}^{p-1} (3k+2) \frac{f_k}{(-4)^k} &= \sum_{k=0}^{p-1} \frac{3k+2}{(-4)^k} \sum_{j=0}^k \binom{k+2j}{3j} \binom{2j}{j} \binom{3j}{j} (-4)^{k-j} \\ &= \sum_{j=0}^{p-1} \frac{\binom{2j}{j} \binom{3j}{j}}{(-4)^j} \sum_{k=j}^{p-1} (3k+2) \binom{k+2j}{3j}. \end{aligned}$$

By using the package **Sigma** again, we find the following identity:

$$\sum_{k=j}^{n-1} (3k+2) \binom{k+2j}{3j} = \frac{9nj+3n+1}{3j+2} \binom{n+2j}{3j+1}.$$

Thus,

$$\sum_{k=0}^{p-1} (3k+2) \frac{f_k}{(-4)^k} = \sum_{j=0}^{p-1} \frac{\binom{2j}{j} \binom{3j}{j} \binom{p+2j}{3j+1}}{(-4)^j} \frac{9pj+3p+1}{3j+2}. \quad (2.18)$$

**Lemma 2.5.** *Let  $p > 3$  be a prime and  $p \equiv 1 \pmod{3}$ . If  $0 \leq j \leq (p-1)/2$  and  $j \neq (p-1)/3$ , then*

$$\binom{3j}{j} \binom{p+2j}{3j+1} \equiv \frac{p(-1)^j}{3j+1} (1 + pH_{2j} - pH_j) \pmod{p^3}.$$

If  $(p+1)/2 \leq j \leq p-1$ , then

$$\binom{3j}{j} \binom{p+2j}{3j+1} \equiv \frac{2p(-1)^j}{3j+1} \pmod{p^2}.$$

*Proof.* If  $0 \leq j \leq (p-1)/2$  and  $j \neq (p-1)/3$ , then we have

$$\begin{aligned} \binom{3j}{j} \binom{p+2j}{3j+1} &= \frac{(p+2j) \cdots (p+1)p(p-1) \cdots (p-j)}{j!(2j)!(3j+1)} \\ &\equiv \frac{p(2j)!(1 + pH_{2j})(-1)^j(j)!(1 - pH_j)}{j!(2j)!(3j+1)} \\ &\equiv \frac{p(-1)^j}{3j+1} (1 + pH_{2j} - pH_j) \pmod{p^3}. \end{aligned}$$

If  $(p+1)/2 \leq j \leq p-1$ , then

$$\begin{aligned} &\binom{3j}{j} \binom{p+2j}{3j+1} \\ &= \frac{(p+2j) \cdots (2p+1)(2p)(2p-1) \cdots (p+1)p(p-1) \cdots (p-j)}{j!(2j)!(3j+1)} \\ &\equiv \frac{2p^2(2j) \cdots (p+1)(p-1)!(-1)^j(j)!}{j!(2j)!(3j+1)} = \frac{2p(-1)^j}{3j+1} \pmod{p^2}. \end{aligned}$$

Now the proof of Lemma 2.5 is complete.  $\square$

It is known that  $\binom{2k}{k} \equiv 0 \pmod{p}$  for each  $(p+1)/2 \leq k \leq p-1$ , and it is easy to check that for each  $0 \leq j \leq (p-1)/2$ ,

$$\binom{3j}{j} \binom{p+2j}{3j+1} \equiv \frac{p(-1)^j}{3j+1} \pmod{p^2}.$$



These, with (2.18) yield that

$$\begin{aligned}
\sum_{k=0}^{p-1} (3k+2) \frac{f_k}{(-4)^k} &\equiv \sum_{j=0}^{\frac{p-1}{2}} \frac{\binom{2j}{j}}{(-4)^j} \frac{p(-1)^j}{3j+1} \frac{9pj+3p+1}{3j+2} + \sum_{j=\frac{p+1}{2}}^{p-1} \frac{\binom{2j}{j}}{(-4)^j} \frac{2p(-1)^j}{3j+1} \frac{1}{3j+2} \\
&\equiv \sum_{j=0}^{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{j} \frac{p(-1)^j}{3j+1} \frac{1}{3j+2} + S_3 \\
&= p \sum_{j=0}^{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{j} (-1)^j \left( \frac{1}{3j+1} - \frac{1}{3j+2} \right) + S_3 \pmod{p^2}, \tag{2.19}
\end{aligned}$$

where

$$\begin{aligned}
S_3 &= \binom{\frac{2p-2}{3}}{\frac{p-1}{3}} \frac{1}{(p+1)4^{\frac{p-1}{3}}} - \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \frac{1}{p+1} - \binom{\frac{4p-4}{3}}{\frac{2p-2}{3}} \frac{1}{4^{\frac{2p-2}{3}}} \\
&= \frac{1}{p+1} \left( \binom{-1/2}{(p-1)/3} - \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \right) - \binom{-1/2}{(2p-2)/3}.
\end{aligned}$$

In the same way of above, with (2.10), (2.13), (2.14), Lemma 2.2 and [20, Theorem 4.12], we have the following congruence modulo  $p^2$

$$p \sum_{j=0}^{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{j} (-1)^j \left( \frac{1}{3j+1} - \frac{1}{3j+2} \right) \equiv 2x + \frac{3px}{2} q_p(3) - \frac{3p}{2x}. \tag{2.20}$$

Now we evaluate  $S_3$ . It is easy to see that

$$\begin{aligned}
\binom{-1/2}{(2p-2)/3} &= \frac{(-\frac{1}{2})(-\frac{1}{2}-1)\cdots(-\frac{1}{2}-\frac{2p-2}{3}+1)}{(\frac{2p-2}{3})!} \\
&= \frac{(\frac{1}{2})(\frac{3}{2})\cdots(\frac{p}{2}-1)\frac{p}{2}(\frac{p}{2}+1)\cdots(\frac{p}{2}+\frac{p-7}{6})}{(\frac{2p-2}{3})!} \\
&= \frac{(\frac{p}{2}-\frac{p-1}{2})\cdots(\frac{p}{2}-1)\frac{p}{2}(\frac{p}{2}+1)\cdots(\frac{p}{2}+\frac{p-7}{6})}{(\frac{2p-2}{3})!} \\
&\equiv \frac{(-1)^{\frac{p-1}{2}} \frac{p}{2} (\frac{p-1}{2})! (\frac{p-7}{6})!}{(\frac{2p-2}{3})!} = \frac{(-1)^{\frac{p-1}{2}} 3p}{p-1} \frac{1}{\binom{\frac{2p-2}{3}}{\frac{p-1}{2}}} \\
&\equiv \frac{-3p(-1)^{(p-1)/2}}{\binom{\frac{2p-2}{3}}{\frac{p-1}{2}}} \pmod{p^2}.
\end{aligned}$$

In view of (2.17) and [20, Theorem 4.12], we immediately obtain that

$$S_3 \equiv -\frac{3px}{2}q_p(3) + \frac{3p}{2x} \pmod{p^2}.$$

This, with (2.19) and (2.20) yields that

$$\frac{1}{2} \sum_{k=0}^{p-1} (3k+2) \frac{f_k}{(-4)^k} \equiv x \pmod{p^2}$$

Now the proof of Theorem 1.1 is complete.  $\square$

### 3. PROOF OF THEOREM 1.2

*Proof of Theorem 1.2.* With the help of (2.4), we have

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{f_k}{2^k} &= \sum_{k=0}^{p-1} \frac{1}{2^k} \sum_{j=0}^{\lfloor k/2 \rfloor} \binom{k+j}{3j} \binom{2j}{j} \binom{3j}{j} 2^{k-2j} \\ &= \sum_{j=0}^{(p-1)/2} \frac{\binom{2j}{j} \binom{3j}{j}}{4^j} \sum_{k=2j}^{p-1} \binom{k+j}{3j}. \end{aligned} \quad (3.1)$$

By loading the package **Sigma** in the software **Mathematica**, we have the following identity:

$$\sum_{k=2j}^{n-1} \binom{k+j}{3j} = \binom{n+j}{3j+1}.$$

Thus, replace  $n$  by  $p$  in the above identity and then substitute it into (3.1), we have

$$\sum_{k=0}^{p-1} \frac{f_k}{2^k} = \sum_{j=0}^{(p-1)/2} \frac{\binom{2j}{j} \binom{3j}{j}}{4^j} \binom{p+j}{3j+1}.$$

Hence we immediately obtain the following result by Lemma 2.3,

$$\sum_{k=0}^{p-1} \frac{f_k}{2^k} \equiv p \sum_{j=0, j \neq \frac{p-1}{3}}^{(p-1)/2} \frac{\binom{2j}{j}}{4^j} \frac{1 - pH_{2k} + pH_k}{(3j+1)} + S_1 \pmod{p^3}, \quad (3.2)$$

where

$$S_1 = \frac{\binom{\frac{2p-2}{3}}{\frac{p-1}{3}} \binom{p-1}{\frac{p-1}{3}} \binom{p+\frac{p-1}{3}}{p}}{4^{\frac{p-1}{3}}} = \binom{-\frac{1}{2}}{\frac{p-1}{3}} \binom{p-1}{\frac{p-1}{3}} \binom{p+\frac{p-1}{3}}{p}.$$

It is easy to verify that

$$\begin{aligned}
& p \sum_{j=0, j \neq \frac{p-1}{3}}^{(p-1)/2} \frac{\binom{2j}{j}}{4^j} \frac{1 - pH_{2k} + pH_k}{(3j+1)} \\
& \equiv p \sum_{j=0, j \neq \frac{p-1}{3}}^{(p-1)/2} \frac{\binom{\frac{p-1}{2}}{j} (-1)^j (1 - pH_{2k} + pH_k)}{(3j+1) \left(1 - p \sum_{r=1}^j \frac{1}{2r-1}\right)} \\
& \equiv p \sum_{j=0}^{(p-1)/2} \frac{\binom{\frac{p-1}{2}}{j} (-1)^j \left(1 + \frac{p}{2} H_k\right)}{(3j+1)} - S_2 \pmod{p^3},
\end{aligned}$$

where

$$S_2 = \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \left(1 + \frac{p}{2} H_{\frac{p-1}{3}}\right).$$

So

$$\sum_{k=0}^{p-1} \frac{f_k}{2^k} \equiv p \sum_{j=0}^{(p-1)/2} \frac{\binom{\frac{p-1}{2}}{j} (-1)^j \left(1 + \frac{p}{2} H_k\right)}{(3j+1)} + S_1 - S_2 \pmod{p^3}. \quad (3.3)$$

It is easy to see that

$$\frac{2p}{3p-1} \frac{(1)_{\frac{p-1}{2}}}{(\frac{1}{3})_{\frac{p-1}{2}}} = \frac{(\frac{p-1}{2})!}{\frac{1}{3} \cdots (\frac{p}{3}-1)(\frac{p}{3}+1) \cdots (\frac{p}{3} + \frac{p-1}{6})} \equiv \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \pmod{p}. \quad (3.4)$$

On the other hand. We have

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} &= \sum_{k=0}^{p-1} \frac{1}{(-4)^k} \sum_{j=0}^k \binom{k+2j}{3j} \binom{2j}{j} \binom{3j}{j} (-4)^{k-j} \\
&= \sum_{j=0}^{p-1} \frac{\binom{2j}{j} \binom{3j}{j}}{(-4)^j} \sum_{k=j}^{p-1} \binom{k+2j}{3j} = \sum_{j=0}^{p-1} \frac{\binom{2j}{j} \binom{3j}{j}}{(-4)^j} \binom{p+2j}{3j+1}.
\end{aligned}$$

So by Lemma 2.5 and the fact that for each  $0 \leq k \leq (p-1)/2$ ,

$$\frac{\binom{2k}{k}}{(-4)^k} \equiv \frac{\binom{(p-1)/2}{k}}{(1 - p \sum_{j=1}^k \frac{1}{2j-1})} \pmod{p^2}, \quad j \binom{2j}{j} \binom{2p-2j}{p-j} \equiv 2p \pmod{p^2}$$

for each  $(p+1)/2 \leq j \leq p-1$ , we have the following congruence modulo  $p^3$ ,

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} - S_3 &\equiv p \sum_{\substack{j=0 \\ j \neq \frac{p-1}{3}}}^{\frac{p-1}{2}} \frac{\binom{2j}{j} (1 + pH_{2j} - pH_j)}{(3j+1)4^j} + 2p \sum_{j=\frac{p+1}{2}}^{p-1} \frac{\binom{2j}{j}}{(3j+1)4^j} \\
&\equiv \sum_{\substack{j=0 \\ j \neq \frac{p-1}{3}}}^{\frac{p-1}{2}} \frac{p(-1)^j \binom{\frac{p-1}{2}}{j} (1 + 2pH_{2j} - \frac{3}{2}pH_j)}{3j+1} + \sum_{j=\frac{p+1}{2}}^{p-1} \frac{4p^2}{4^j(3j+1)j \binom{2p-2j}{p-j}} \\
&\equiv \sum_{j=0}^{\frac{p-1}{2}} \frac{p(-1)^j \binom{\frac{p-1}{2}}{j} (1 + 2pH_{2j} - \frac{3}{2}pH_j)}{3j+1} + \sum_{j=1}^{\frac{p-1}{2}} \frac{p^2 4^j}{(3j-1)j \binom{2j}{j}} - S_4,
\end{aligned}$$

where

$$\begin{aligned}
S_3 &= \frac{\binom{\frac{2p-2}{3}}{\frac{p-1}{3}} \binom{p-1}{\frac{p-1}{3}} \binom{p+\frac{2p-2}{3}}{p}}{(-4)^{\frac{p-1}{3}}} = \binom{-\frac{1}{2}}{\frac{p-1}{3}} \binom{p-1}{\frac{p-1}{3}} \binom{p+\frac{2p-2}{3}}{p}, \\
S_4 &= \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \left( 1 + 2pH_{\frac{2p-2}{3}} - \frac{3}{2}pH_{\frac{p-1}{3}} \right).
\end{aligned}$$

Hence we have

$$\begin{aligned}
&\sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} - \sum_{k=0}^{p-1} \frac{f_k}{2^k} \\
&\equiv 2p^2 \sum_{j=0}^{\frac{p-1}{2}} \frac{\binom{\frac{p-1}{2}}{j} (-1)^j (H_{2j} - H_j)}{3j+1} + S_5 + \sum_{j=1}^{\frac{p-1}{2}} \frac{p^2 4^j}{(3j-1)j \binom{2j}{j}} \pmod{p^3},
\end{aligned} \tag{3.5}$$

where

$$S_5 = S_3 - S_4 + S_2 - S_1.$$

By **Sigma**, we can find and prove the following identity:

$$\begin{aligned}
&\sum_{j=0}^n \frac{2 \binom{n}{j} (-1)^j (H_{2j} - H_j)}{3j+1} \\
&= \frac{1}{3n+1} \prod_{k=1}^n \frac{3k}{3k-2} \left( \sum_{k=1}^n \frac{1}{k} \prod_{j=1}^k \frac{3j-2}{3j} - \sum_{k=1}^n \frac{1}{k} \prod_{j=1}^k \frac{2(3j-2)}{3(2j-1)} \right) \\
&= \frac{(1)_n}{(3n+1) \left(\frac{1}{3}\right)_n} \left( \sum_{k=1}^n \frac{\left(\frac{1}{3}\right)_k}{k(1)_k} - \sum_{k=1}^n \frac{\left(\frac{1}{3}\right)_k}{k\left(\frac{1}{2}\right)_k} \right).
\end{aligned} \tag{3.6}$$

In view of [16, Lemma 3.1] and Lemma 2.2, we have

$$\sum_{k=1}^{\frac{p-1}{2}} \frac{\left(\frac{1}{3}\right)_k}{k(1)_k} = \sum_{k=1}^{\frac{p-1}{2}} \frac{\left(-1/3\right)_k}{k\binom{-1}{k}} \equiv \frac{3}{2}q_p(3) - \frac{3p}{4}q_p^2(3) - \frac{p}{3} \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k^2\binom{2k}{k}} \pmod{p^2}. \quad (3.7)$$

$$\begin{aligned} \sum_{k=1}^{\frac{p-1}{2}} \frac{\left(\frac{1}{3}\right)_k}{k\left(\frac{1}{2}\right)_k} &= \sum_{k=1}^{\frac{p-1}{2}} \frac{\left(-1/3\right)_k}{k\binom{-1/2}{k}} \equiv \frac{4p}{3}(-1)^{\frac{p-1}{2}}E_{p-3} + \frac{3}{2}q_p(3) - \frac{3p}{4}q_p^2(3) \\ &\quad - \frac{2p}{3}(-1)^{\frac{p-1}{2}} \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{(2k-1)k\binom{2k}{k}} \pmod{p^2}. \end{aligned} \quad (3.8)$$

It is easy to check that

$$\sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{(2k-1)k\binom{2k}{k}} = 2 \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{(2k-1)\binom{2k}{k}} - \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k\binom{2k}{k}}. \quad (3.9)$$

And by [19, (6)], we have

$$\frac{1}{\binom{n+1+k}{k}} = (n+1) \sum_{r=0}^n \binom{n}{r} (-1)^r \frac{1}{k+r+1}. \quad (3.10)$$

$$\begin{aligned} 2 \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{(2k-1)\binom{2k}{k}} &\equiv 2 \sum_{k=1}^{\frac{p-1}{3}} \frac{(-1)^k}{(2k-1)\binom{\frac{p-1}{2}}{k}} \equiv (-1)^{\frac{p+1}{2}} \sum_{k=\frac{p-1}{6}}^{\frac{p-3}{2}} \frac{(-1)^k}{(k+1)\binom{\frac{p-1}{2}}{k}} \\ &= (-1)^{\frac{p+1}{2}} \left( \sum_{k=0}^{\frac{p-3}{2}} \frac{(-1)^k}{(k+1)\binom{\frac{p-1}{2}}{k}} - \sum_{k=0}^{\frac{p-7}{6}} \frac{(-1)^k}{(k+1)\binom{\frac{p-1}{2}}{k}} \right) \pmod{p}. \end{aligned} \quad (3.11)$$

By **Sigma**, we find the following identity which can be proved by induction on  $n$ :

$$\sum_{k=0}^n \frac{(-1)^k}{(k+1)\binom{n}{k}} = \frac{2(-1)^n - 1}{n+1} - (n+1)H_n^{(2)} - 2(n+1) \sum_{k=1}^n \frac{(-1)^k}{k^2}.$$

So by setting  $n = (p-1)/2$  in the above identity, we have

$$\sum_{k=0}^{\frac{p-3}{2}} \frac{(-1)^k}{(k+1)\binom{\frac{p-1}{2}}{k}} \equiv 2 \left( (-1)^{\frac{p-1}{2}} - 1 \right) - (-1)^{\frac{p-1}{2}} 2E_{p-3} \pmod{p}. \quad (3.12)$$

And by (3.10), we have

$$\begin{aligned}
& \sum_{k=0}^{\frac{p-7}{6}} \frac{(-1)^k}{(k+1)\binom{\frac{p-1}{2}}{k}} \equiv \sum_{k=0}^{\frac{p-7}{6}} \frac{1}{(k+1)\binom{\frac{p-1}{2}+k}{k}} \\
&= \sum_{k=0}^{\frac{p-7}{6}} \frac{1}{k+1} \frac{p-1}{2} \sum_{r=0}^{\frac{p-3}{2}} \binom{\frac{p-3}{2}}{r} (-1)^r \frac{1}{k+r+1} \\
&\equiv -\frac{1}{2} \sum_{k=1}^{\frac{p-1}{6}} \frac{1}{k} \sum_{r=0}^{\frac{p-3}{2}} \binom{\frac{p-3}{2}}{r} (-1)^r \frac{1}{k+r} \\
&= -\frac{1}{2} H_{\frac{p-1}{6}}^{(2)} - \frac{1}{2} \sum_{r=1}^{\frac{p-3}{2}} \frac{(-1)^r}{r} \binom{\frac{p-3}{2}}{r} \sum_{k=1}^{\frac{p-1}{6}} \left( \frac{1}{k} - \frac{1}{k+r} \right) \pmod{p}.
\end{aligned}$$

It is easy to check that

$$H_{\frac{p-1}{6}} - \sum_{k=1}^{\frac{p-1}{6}} \frac{1}{k+r} \equiv - \sum_{k=1}^r \frac{1}{k(6k-1)} \pmod{p}.$$

And by Sigma again, we have

$$\sum_{r=1}^n \frac{(-1)^r}{r} \binom{n}{r} \sum_{k=1}^r \frac{1}{k(6k-1)} = H_n^{(2)} - \sum_{k=1}^n \frac{(1)_k}{k \binom{\frac{5}{6}}{k}}.$$

So by Lemma (2.2) and [20], we have

$$\begin{aligned}
& \sum_{k=0}^{\frac{p-7}{6}} \frac{(-1)^k}{(k+1)\binom{\frac{p-1}{2}}{k}} \\
&\equiv \frac{(-1)^{\frac{p-1}{2}}}{x} - 2 - \frac{5}{4} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) - \frac{1}{2} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-1)^k}{k^2 \binom{-\frac{5}{6}}{k}} \pmod{p}.
\end{aligned}$$

Again, by (3.10), we have

$$\begin{aligned}
\sum_{k=1}^{\frac{p-1}{2}} \frac{(-1)^k}{k^2 \binom{-\frac{5}{6}}{k}} &= -\frac{6}{5} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-1)^k}{k \binom{-\frac{11}{6}}{k-1}} \equiv \frac{6}{5} \sum_{k=0}^{\frac{p-3}{2}} \frac{(-1)^k}{(k+1) \binom{\frac{5p-11}{6}}{k}} \\
&= \frac{6}{5} \sum_{k=0}^{\frac{p-3}{2}} \frac{1}{(k+1) \binom{\frac{p+5}{6}+k}{k}} = \frac{6}{5} \sum_{k=0}^{\frac{p-3}{2}} \frac{1}{k+1} \frac{p+5}{6} \sum_{r=0}^{\frac{p-1}{6}} (-1)^r \binom{\frac{p-1}{6}}{r} \frac{1}{k+1+r} \\
&\equiv \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k} \sum_{r=0}^{\frac{p-1}{6}} (-1)^r \binom{\frac{p-1}{6}}{r} \frac{1}{k+r} \\
&= H_{\frac{p-1}{2}}^{(2)} + \sum_{r=1}^{\frac{p-1}{6}} \frac{(-1)^r}{r} \binom{\frac{p-1}{6}}{r} \sum_{k=1}^{\frac{p-1}{2}} \left( \frac{1}{k} - \frac{1}{k+r} \right) \pmod{p}.
\end{aligned}$$

Also it is easy to see that

$$H_{\frac{p-1}{2}} - \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k+r} \equiv - \sum_{k=1}^r \frac{1}{k(2k-1)} \pmod{p}.$$

And by Sigma, we have

$$\sum_{r=1}^n \frac{(-1)^r}{r} \binom{n}{r} \sum_{k=1}^r \frac{1}{k(2k-1)} = H_n^{(2)} - \sum_{k=1}^n \frac{4^k}{k^2 \binom{2k}{k}}.$$

So by Lemma 2.2, we have

$$\sum_{k=1}^{\frac{p-1}{2}} \frac{(-1)^k}{k^2 \binom{-\frac{5}{6}}{k}} \equiv \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} - \frac{5}{2} \left( \frac{p}{3} \right) B_{p-2} \left( \frac{1}{3} \right) \pmod{p}.$$

Hence

$$\sum_{k=0}^{\frac{p-7}{6}} \frac{(-1)^k}{(k+1) \binom{\frac{p-1}{2}}{k}} \equiv \frac{(-1)^{\frac{p-1}{2}}}{x} - 2 - \frac{1}{2} \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \pmod{p}.$$

This, with (3.11) and (3.12) yields that

$$\begin{aligned}
&2 \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{(2k-1)} \\
&\equiv -2 + \frac{1}{x} + 2E_{p-3} - \frac{1}{2} (-1)^{\frac{p-1}{2}} \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \pmod{p}. \tag{3.13}
\end{aligned}$$

By Sigma, we have

$$\sum_{k=1}^n \frac{4^k}{k \binom{2k}{k}} = -2 + 2 \frac{4^n}{\binom{2n}{n}}. \quad (3.14)$$

So by [20], we have

$$\sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k \binom{2k}{k}} \equiv -2 + \frac{2}{\binom{\frac{p-1}{2}}{\frac{p-1}{3}}} \equiv -2 + \frac{1}{x} \pmod{p}.$$

This, with (3.9) and (3.13) yields that

$$\sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{(2k-1)k \binom{2k}{k}} \equiv 2E_{p-3} - \frac{1}{2}(-1)^{\frac{p-1}{2}} \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \pmod{p}.$$

This with (3.8) yields that

$$\sum_{k=1}^{\frac{p-1}{2}} \frac{\left(\frac{1}{3}\right)_k}{k \left(\frac{1}{2}\right)_k} \equiv \frac{3}{2}q_p(3) - \frac{3p}{4}q_p^2(3) + \frac{p}{3} \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \pmod{p^2}. \quad (3.15)$$

So by (3.7), we have

$$\sum_{k=1}^{\frac{p-1}{2}} \frac{\left(\frac{1}{3}\right)_k}{k \left(1\right)_k} - \sum_{k=1}^{\frac{p-1}{2}} \frac{\left(\frac{1}{3}\right)_k}{k \left(\frac{1}{2}\right)_k} \equiv -\frac{p}{3} \left( \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k^2 \binom{2k}{k}} + \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \right) \pmod{p^2}.$$

Therefore, by (3.6) and (3.4), we have

$$\begin{aligned} & 2p^2 \sum_{j=0}^{\frac{p-1}{2}} \frac{\left(\frac{p-1}{2}\right)_j (-1)^j (H_{2j} - H_j)}{3j+1} \\ & \equiv -\frac{p^2}{3} \left(\frac{p-1}{3}\right) \left( \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k^2 \binom{2k}{k}} + \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \right) \pmod{p^3}. \end{aligned} \quad (3.16)$$

Now we evaluate the second sum in the right side of (3.5). It is easy to see that

$$\sum_{j=1}^{\frac{p-1}{2}} \frac{4^j}{(3j-1)j \binom{2j}{j}} = 3 \sum_{j=1}^{\frac{p-1}{2}} \frac{4^j}{(3j-1) \binom{2j}{j}} - \sum_{j=1}^{\frac{p-1}{2}} \frac{4^j}{j \binom{2j}{j}}. \quad (3.17)$$

It is easy to see from (3.14) that

$$\sum_{j=1}^{\frac{p-1}{2}} \frac{4^j}{j \binom{2j}{j}} \equiv -2 + 2(-1)^{\frac{p-1}{2}} \pmod{p}. \quad (3.18)$$



Now we consider the first sum of the right side in (3.17).

$$\sum_{j=1}^{\frac{p-1}{2}} \frac{4^j}{(3j-1)\binom{2j}{j}} = \sum_{j=1}^{\frac{p-1}{3}} \frac{4^j}{(3j-1)\binom{2j}{j}} + \sum_{j=\frac{p+2}{3}}^{\frac{p-1}{2}} \frac{4^j}{(3j-1)\binom{2j}{j}}.$$

The following identity is very import to us:

$$\sum_{k=1}^n \frac{4^k}{(k+n)\binom{2k}{k}} = -2 + 2 \frac{4^n}{\binom{2n}{n}} - \frac{n\binom{2n}{n}}{4^n} \sum_{k=1}^n \frac{4^k}{k^2 \binom{2k}{k}}. \quad (3.19)$$

This, with [20] yields that

$$\begin{aligned} 3 \sum_{j=1}^{\frac{p-1}{3}} \frac{4^j}{(3j-1)\binom{2j}{j}} &\equiv \sum_{j=1}^{\frac{p-1}{3}} \frac{4^j}{(j + \frac{p-1}{3}) \binom{2j}{j}} \\ &\equiv -2 + \frac{2}{\binom{-1/2}{\frac{p-1}{3}}} + \frac{1}{3} \binom{-1/2}{\frac{p-1}{3}} \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k^2 \binom{2k}{k}} \\ &\equiv -2 + \frac{1}{x} + \frac{1}{3} \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k^2 \binom{2k}{k}} \pmod{p}. \end{aligned} \quad (3.20)$$

It is easy to check that And by (3.19), we have

$$\begin{aligned} 3 \sum_{j=\frac{p+2}{3}}^{\frac{p-1}{2}} \frac{4^j}{(3j-1)\binom{2j}{j}} &\equiv 3 \sum_{j=0}^{\frac{p-7}{6}} \frac{(-1)^{\frac{p-1}{2}-j}}{(3(\frac{p-1}{2}-j)-1)\binom{\frac{p-1}{2}}{j}} \\ &\equiv 6(-1)^{\frac{p+1}{2}} \sum_{j=0}^{\frac{p-7}{6}} \frac{4^j}{(6j+5)\binom{2j}{j}} \equiv (-1)^{\frac{p+1}{2}} \sum_{j=0}^{\frac{p-7}{6}} \frac{(-1)^j}{(j + \frac{p+5}{6})\binom{\frac{p-1}{2}}{j}} \\ &\equiv \frac{6}{5}(-1)^{\frac{p+1}{2}} + (-1)^{\frac{p+1}{2}} \sum_{j=1}^{\frac{p+5}{6}} \frac{4^j}{(j + \frac{p+5}{6})\binom{2j}{j}} + \frac{3}{\binom{\frac{p-1}{2}}{\frac{p-1}{3}}} \pmod{p}. \end{aligned} \quad (3.21)$$

By (3.19) and [20], we have

$$\sum_{j=1}^{\frac{p+5}{6}} \frac{4^j}{(j + \frac{p+5}{6})\binom{2j}{j}} \equiv -\frac{16}{5} + \frac{5(-1)^{\frac{p-1}{6}}}{2x} - \frac{(-1)^{\frac{p-1}{6}}}{6x} \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \pmod{p}.$$

This, with (3.21) yields that

$$3 \sum_{j=\frac{p+2}{3}}^{\frac{p-1}{2}} \frac{4^j}{(3j-1)\binom{2j}{j}} \equiv 2(-1)^{\frac{p-1}{2}} - \frac{1}{x} + \frac{1}{3} \left( \frac{p-1}{2} \right) \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \pmod{p}.$$

Combining this with (3.20), we have

$$\begin{aligned} & 3 \sum_{j=1}^{\frac{p-1}{2}} \frac{4^j}{(3j-1)\binom{2j}{j}} \\ & \equiv -2 + 2(-1)^{\frac{p-1}{2}} + \frac{1}{3} \left( \frac{p-1}{2} \right) \left( \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k^2 \binom{2k}{k}} + \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \right) \pmod{p}. \end{aligned}$$

Thus, by (3.17) and (3.18), we have

$$\sum_{j=1}^{\frac{p-1}{2}} \frac{4^j}{(3j-1)j\binom{2j}{j}} \equiv \frac{1}{3} \left( \frac{p-1}{2} \right) \left( \sum_{k=1}^{\frac{p-1}{3}} \frac{4^k}{k^2 \binom{2k}{k}} + \sum_{k=1}^{\frac{p-1}{6}} \frac{4^k}{k^2 \binom{2k}{k}} \right) \pmod{p}.$$

This, with (3.5) and (3.16) yields that

$$\sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} - \sum_{k=0}^{p-1} \frac{f_k}{2^k} \equiv S_5 \pmod{p^3}. \quad (3.22)$$

While

$$\begin{aligned} S_5 &= \binom{-\frac{1}{2}}{\frac{p-1}{3}} \binom{p-1}{\frac{p-1}{3}} \left( \binom{p+\frac{2p-2}{3}}{\frac{p-1}{3}} - \binom{p+\frac{p-1}{3}}{\frac{p-1}{3}} \right) \\ &+ 2p \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \left( H_{\frac{p-1}{3}} - H_{\frac{2p-2}{3}} \right). \end{aligned}$$

It is easy to check that

$$\binom{p+\frac{2p-2}{3}}{\frac{p-1}{3}} \equiv 1 + pH_{\frac{2p-2}{3}} + \frac{p^2}{2} \left( H_{\frac{2p-2}{3}}^2 - H_{\frac{2p-2}{3}}^{(2)} \right) \pmod{p^3}$$

and

$$\binom{p+\frac{p-1}{3}}{\frac{p-1}{3}} \equiv 1 + pH_{\frac{p-1}{3}} + \frac{p^2}{2} \left( H_{\frac{p-1}{3}}^2 - H_{\frac{p-1}{3}}^{(2)} \right) \pmod{p^3}.$$

So by Lemma 2.2 and the fact that  $H_{p-1-k}^{(2)} \equiv -H_k^{(2)} \pmod{p}$  for each  $0 \leq k \leq p-1$ , we have

$$\begin{aligned} \binom{p + \frac{2p-2}{3}}{\frac{p-1}{3}} - \binom{p + \frac{p-1}{3}}{\frac{p-1}{3}} &\equiv p(H_{\frac{2p-2}{3}} - H_{\frac{p-1}{3}}) + \frac{p^2}{2}(H_{\frac{p-1}{3}}^{(2)} - H_{\frac{2p-2}{3}}^{(2)}) \\ &\equiv p^2 \left(\frac{p}{3}\right) B_{p-2} \left(\frac{1}{3}\right) \pmod{p^3} \end{aligned}$$

and

$$2p \left(H_{\frac{p-1}{3}} - H_{\frac{2p-2}{3}}\right) \equiv -p^2 \left(\frac{p}{3}\right) B_{p-2} \left(\frac{1}{3}\right) \pmod{p^3}.$$

So by  $\binom{-\frac{1}{2}}{\frac{p-1}{3}} \equiv \binom{\frac{p-1}{2}}{\frac{p-1}{3}} \pmod{p}$  and  $\binom{p-1}{\frac{p-1}{3}} \equiv (-1)^{\frac{p-1}{3}} = 1 \pmod{p}$ , we can immediately obtain that

$$S_5 \equiv 0 \pmod{p^3}.$$

This, with (3.22) yields that

$$\sum_{k=0}^{p-1} \frac{f_k}{(-4)^k} \equiv \sum_{k=0}^{p-1} \frac{f_k}{2^k} \pmod{p^3}.$$

Now the proof of Theorem 1.2 is complete.  $\square$

**Acknowledgment.** The first author is funded by the National Natural Science Foundation of China (12001288) and China Scholarship Council (202008320187).

## REFERENCES

- [1] D. Callan, *A combinatorial interpretation for an identity of Barrucand*, J. Integer Seq. **11** (2008), Article 08.3.4, 3pp (electronic).
- [2] J. Franel, *On a question of Laisant*, L'Intermédiaire des Mathématiciens **1** (1894), 45–47.
- [3] V.J.W. Guo, *Proof of two conjectures of Sun on congruences for Franel numbers*, Integral Transforms Spec. Funct. **24** (2013), 532–539.
- [4] V.J.W. Guo, *Proof of a superconjecture conjectured by Z.-H. Sun*, Integral Transforms Spec. Funct. **25** (2014), 1009–1015.
- [5] E. Lehmer, *On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson*, Ann. Math. **39**(1938), 350–360.
- [6] J.-C. Liu, *On two congruences involving Franel numbers*, Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Math. **114** (2020), Art.201.
- [7] G.-S. Mao, *On two congruences involving Apéry and Franel numbers*, Results Math. **75** (2020), Art 159.
- [8] G.-S. Mao, *Proof of some congruence conjectures of Z.-H. Sun involving Apéry-like numbers*, preprint, arXiv:2111.08778.
- [9] M. R. Murty, *Introduction to p-adic analytic number theory*, AMS/IP Studies in Advanced Mathematics, 27, American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2002.

- [10] A. M. Robert, *A course in p-adic analysis*, Graduate Texts in Mathematics, 198. Springer-Verlag, New York, 2000.
- [11] C. Schneider, *Symbolic summation assists combinatorics*, Sémin. Lothar. Combin. **56** (2007), Article B56b.
- [12] N.J.A. Sloane, Sequence A000172 in OEIS (On-Line Encyclopedia of Integer Sequences), <http://oeis.org/A000172>.
- [13] Z.-H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105** (2000), no.1-3, 193–223.
- [14] Z.-H. Sun, *Congruences involving Bernoulli and Euler numbers*, J. Number Theory **128** (2008), no.2, 280–312.
- [15] Z.-H. Sun, *Congruences for sums involving Franel numbers*, Int. J. Number Theory, **14** (2018), no.1, 123–142.
- [16] Z.-H. Sun, *Super congruences concerning Bernoulli polynomials*, Int. J. Number Theory **11** (2015), no.8, 2393–2404.
- [17] Z.-W. Sun, *Connections between  $p = x^2 + 3y^2$  and Franel numbers*, J. Number Theory **133** (2013), 2914–2928.
- [18] Z.-W. Sun, *Congruences for Franel numbers*, Adv. in Appl. Math. **51** (2013), no.4, 524–535.
- [19] B. Sury, T.-M. Wang and F.-Z. Zhao, *Identities involving reciprocals of binomial coefficients*, J. Integer Seq. **7** (2004), Article 04.2.8.
- [20] K. M. Yeung, *On congruences for Binomial Coefficients*, J. Number Theory **33** (1989), 1–17.
- [21] D. Zagier, *Integral solutions of Apéry-like recurrence equations*, in: Groups and Symmetries: from Neolithic Scots to John McKay, CRM Proc. Lecture Notes 47, Amer. Math. Soc., Providence, RI, 2009, pp. 349–366.

(GUO-SHUAI MAO) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY OF INFORMATION SCIENCE AND TECHNOLOGY, NANJING 210044, PEOPLE’S REPUBLIC OF CHINA, [maogsmath@163.com](mailto:maogsmath@163.com)

(YAN LIU) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY OF INFORMATION SCIENCE AND TECHNOLOGY, NANJING 210044, PEOPLE’S REPUBLIC OF CHINA, [1325507759@qq.com](mailto:1325507759@qq.com)