

Analytic extensions of Starobinsky model of inflation

Vsevolod R. Ivanov,^a Sergei V. Ketov,^{b,c,d} Ekaterina O. Pozdeeva,^e
Sergey Yu. Vernov^{e,1}

^aPhysics Department, Lomonosov Moscow State University
Leninskie Gory 1, 119991 Moscow, Russia

^bInstitut für Theoretische Physik and Riemann Center for Geometry and Physics
Leibniz Universität Hannover, Appelstrasse 2, 30167 Hannover, Germany

^cDepartment of Physics, Tokyo Metropolitan University
1-1 Minami-ohsawa, Hachioji-shi, Tokyo 192-0397, Japan

^dKavli Institute for the Physics and Mathematics of the Universe (WPI)
The University of Tokyo Institutes for Advanced Study, Kashiwa 277-8583, Japan

^eSkobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University,
Leninskiye Gory 1, 119991 Moscow, Russia

E-mail: vsvd.ivanov@gmail.com, ketov@tmu.ac.jp,
pozdeeva@www-hep.sinp.msu.ru, svernov@theory.sinp.msu.ru

Abstract. We study several extensions of the Starobinsky model of inflation, which obey all observational constraints on the inflationary parameters, by demanding that both the inflaton scalar potential in the Einstein frame and the $F(R)$ gravity function in the Jordan frame have the explicit dependence upon fields *and* parameters in terms of elementary functions. Our models are continuously connected to the original Starobinsky model via changing the parameters. We modify the Starobinsky $(R + R^2)$ model by adding an R^3 -term, an R^4 -term, and an $R^{3/2}$ -term, respectively, and calculate the scalar potentials, the inflationary observables and the allowed limits on the deformation parameters by using the latest observational bounds. We find that the tensor-to-scalar ratio in the Starobinsky model modified by the $R^{3/2}$ -term significantly increases with raising the parameter in front of that term. On the other side, we deform the scalar potential of the Starobinsky model in the Einstein frame in powers of $y = \exp\left(-\sqrt{\frac{2}{3}}\phi/M_{Pl}\right)$, where ϕ is the canonical inflaton (scalon) field, calculate the corresponding $F(R)$ gravity functions in the two new cases, and find the restrictions on the deformation parameters in the lowest orders with respect to the variable y that is physically small during slow-roll inflation.

Keywords: inflation, modified gravity, cosmology

ArXiv ePrint: [2111.09058](https://arxiv.org/abs/2111.09058)

¹The corresponding author.

Contents

1	Introduction	1
2	Setup	4
2.1	Duality transformations	4
2.2	Equations of motion	6
2.3	Slow-roll approximation and inflation observables	7
3	Polynomial modifications of Starobinsky model	8
3.1	The $(R + R^2 + R^3)$ gravity models of inflation	8
3.2	The $(R + R^2 + R^4)$ gravity model of inflation	12
4	The $(R + R^{3/2} + R^2)$ gravity model of inflation	15
4.1	The inflaton potential	15
4.2	The inflationary parameters	16
5	Deforming the scalar potential in the Starobinsky model with analytic F-functions	20
5.1	New case I	21
5.2	New case II	22
6	Conclusion	25

1 Introduction

The *duality* relation between modified $F(R)$ gravity theories and scalar-tensor gravity theories is the standard tool in modern cosmology, see Refs. [1–5] for the original papers about the duality transformation and Refs. [6–9] for a review of $F(R)$ gravity theories and their physical applications. In the literature, the duality relation is usually used only in one direction, from an $F(R)$ gravity model to the equivalent scalar-tensor (or quintessence) model in the Einstein frame with a propagating scalar field. In the context of inflationary models [10–19], the scalar field is identified with the inflaton having the clear gravitational origin as a physical excitation of the higher-derivative gravity (called scalaron).

A well-known example of the correspondence is given by the celebrated Starobinsky model of inflation [10], whose action is given by

$$S_{\text{Star.}}[g_{\mu\nu}^J] = \frac{M_{Pl}^2}{2} \int d^4x \sqrt{-g_J} \left(R_J + \frac{1}{6m^2} R_J^2 \right), \quad (1.1)$$

where we have introduced the reduced Planck mass M_{Pl} and the inflaton mass m , in terms of metric $g_{\mu\nu}^J$ having the Ricci scalar curvature R_J , with the spacetime signature $(-, +, +, +)$. The action (1.1) is dual to the quintessence (or scalar-tensor gravity) action

$$S_{\text{quint.}}[g_{\mu\nu}, \phi] = \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V_{\text{Star.}}(\phi) \right] \quad (1.2)$$

in terms of the canonical scalar ϕ and another metric $g_{\mu\nu}$ in the Einstein frame, related to $g_{\mu\nu}^J$ (in the Jordan frame) by a Weyl transform (see Sec. 2 for details), and having the Ricci scalar R . The induced scalar potential is given by

$$V_{\text{Star.}}(\phi) = \frac{3}{4} M_{Pl}^2 m^2 \left[1 - \exp \left(-\sqrt{\frac{2}{3}} \phi / M_{Pl} \right) \right]^2. \quad (1.3)$$

The Starobinsky model is known as the excellent model of large-field slow-roll cosmological inflation with very good agreement to the Planck measurements of the Cosmic Microwave Background (CMB) radiation [20]. In particular, the observable CMB amplitude fixes the only parameter of the Starobinsky model as $m \sim 10^{-5} M_{Pl}$. It is also remarkable that the Lagrangian (1.1) and the corresponding scalar potential (1.3) are very simple.

The Starobinsky model (1.1) is the particular case of the modified $F(R)$ gravity theories having the action

$$S_F[g_{\mu\nu}^J] = \int d^4x \sqrt{-g^J} F(R_J) \quad (1.4)$$

with a differentiable function F . In contrast to Starobinsky's Lagrangian quadratic in R_J , the $F(R_J)$ theories with generic functions F do not lead to a simple scalar potential, and the duality transformation itself is non-trivial. This is the reason why the induced scalar potentials are often obtained and studied in the $F(R)$ gravity literature by numerical methods. Moreover, when starting from a generic F -function, one often arrives at a multi-valued scalar potential with a number of branching cuts and points, see e.g., Ref. [11] for some examples. Therefore, mathematically and physically well-defined $F(R)$ gravity functions have to be carefully chosen.

One also has to avoid graviton as a ghost and scalaron (inflaton) as a tachyon. It leads to further restrictions

$$\frac{dF}{dR_J} > 0 \quad \text{and} \quad \frac{d^2F}{dR_J^2} > 0 \quad (1.5)$$

that restrict possible values of parameters and R_J [21, 22]. For example, in the Starobinsky model, the first condition in (1.5) is violated for the large *negative* $R_J < -3m^2$, where the duality transformation does not exist.¹

When considering an inflationary model in modified gravity as the effective gravitational theory, its Hubble function H must be negligible against the Planck scale M_{Pl} and the UV-cutoff Λ_{UV} . This guarantees decoupling of heavy modes (like Kaluza-Klein modes and string theory massive modes), which is required for consistency [25]. In the Starobinsky model, both conditions are satisfied because $H \sim \mathcal{O}(m)$ and $\Lambda_{UV} = M_{Pl}$, while the latter easily follows from expanding the scalar potential (1.3) in power series with respect to the inflaton field ϕ . In this paper, we confine ourselves to non-negative values of the Ricci scalar curvature well below $\mathcal{O}(M_{Pl}^2)$.

When using an $F(R)$ gravity model for describing inflation, an agreement with the CMB observables (the amplitude of fluctuations A_s , the scalar power spectrum index n_s and the tensor-to-scalar ratio r) is also required. As regards the scalar potential, it is reasonable to demand a Minkowski (or de Sitter) vacuum and the boundedness of the potential from below.

¹An absence of the duality transformation does not imply non-existence of smooth solutions on which the function dF/dR_J changes its sign. In the Friedmann-Lemaître-Robertson-Walker (FLRW) universe such solutions do exist [23] but in more general spacetimes, e.g., with a Bianchi I metric, anisotropic instabilities arise and isotropic solutions are not stable [24].

Taken together, all these restrictions also significantly restrict possible choices of F function and scalar potential.

In this paper, we consider analytic deformations of the Starobinsky model (1.1), which obey all the above mentioned restrictions, at least for small values of the deformation parameters. The inverse duality transformation relating a generic scalar potential $V(\phi)$ in the action (1.2) to the corresponding $F(R)$ gravity function in Eq. (1.4) is known in the parametric form [17, 26, 27]. However, it is often impossible to get an explicit analytic solution to the F -function in terms of elementary functions for a given scalar potential. For example, a simple quadratic scalar potential (a mass term) leads to the complicated F -function in terms of the special (Lambert) function. It is therefore of interest to find other cases (generalizing the Starobinsky model) where *both* functions $V(\phi)$ and $F(R)$ exist in terms of elementary functions. It is certainly relevant for studying the parameter spaces of cosmological models because analytical methods are superior to numerical methods there.

The Starobinsky model is just the simplest model of inflation with sharp predictions for the inflationary observables and no free parameters, i.e. it has the maximal predictive power. However, its viable extensions may be required by future experiments, should the observed values of the scalar perturbation index n_s and the tensor-to-scalar ratio r deviate from their values in the Starobinsky model of inflation. Having more freedom in the choice of new parameters consistent with inflation is also useful for other purposes, such as studies of reheating and astrophysical constraints [6–9], or formation of primordial black holes [28]. In the context of inflationary models, we are only interested in the physically viable theories consistent with CMB measurements. In this paper, we find some continuous deformations of the Starobinsky model under the additional condition that both functions $V(\phi)$ and $F(R)$ can be explicitly given in a finite form in terms of elementary functions, and focus on the deformations with only one free parameter.

Our motivation can thus be summarized as follows. The Starobinsky model gives sharp predictions for observables, so that any new viable extension of the model is worth investigating because future measurements may deviate from its predictions. Also, the Starobinsky model has no free parameters, whereas its extensions have new parameters. A full investigation of the parameter space requires both analytic and numeric methods because numerical calculations alone are often possible only for specific values of the parameters. Given extra parameters, it is important to provide specific constraints on their possible values. On the technical side, it is desirable to identify a "small field" in the Starobinsky model, which can serve for an expansion with respect to that field in possible extensions.

Our paper is organized as follows. In Sec. 2 we review the duality transformations between $F(R)$ gravity theories and scalar-tensor theories in *both* directions, formulate the equations of motion, and define the slow-roll approximation for inflation. The polynomial deformations of the Starobinsky model in $F(R)$ gravity by adding the R^3 term or the R^4 term in the context of slow-roll inflation are studied in Sec. 3. These terms may arise as the quantum gravity corrections from Planck scale physics. We estimate their contribution to inflation by finding the upper limits of their coefficients, via demanding consistency of slow-roll inflation with CMB observations [20, 29]. The scalar potentials in these models are drastically different from that in Eq. (1.3) at very large values of the scalar curvature, independently upon the smallness of the deformation parameters. In Sec. 4 we study the impact of adding the $R^{3/2}$ term to the Starobinsky model of inflation, where our methods also apply. In Sec. 5 we begin with the scalar potential (1.3), give the two new examples of its one-parametric deformation, which are also consistent with observations, and find

the corresponding $F(R)$ gravity functions in the explicit analytic form. Section 6 is our conclusion.

2 Setup

2.1 Duality transformations

The $F(R)$ gravity action (1.4) can be rewritten as

$$S_J[g_{\mu\nu}^J, \sigma] = \int d^4x \sqrt{-g^J} [F_{,\sigma}(R_J - \sigma) + F] , \quad (2.1)$$

where the new scalar field σ has been introduced, and $F_{,\sigma}(\sigma) = \frac{dF(\sigma)}{d\sigma}$. Eliminating σ via its algebraic equation of motion, $R_J = \sigma$, yields back the action (1.4) when assuming that $F_{,\sigma\sigma}(\sigma) = d^2F/d\sigma^2 \neq 0$. After the Weyl transformation of the metric

$$g_{\mu\nu} = \frac{2F_{,\sigma}(\sigma)}{M_{Pl}^2} g_{\mu\nu}^J \quad (2.2)$$

one gets the following action in the Einstein frame [3]:

$$S_E[g_{\mu\nu}, \sigma] = \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{2} R - \frac{h(\sigma)}{2} g^{\mu\nu} \partial_\mu \sigma \partial_\nu \sigma - V \right] , \quad (2.3)$$

where we have introduced the functions

$$h(\sigma) = \frac{3M_{Pl}^2}{2F_{,\sigma}^2} F_{,\sigma\sigma}^2 \quad \text{and} \quad V(\sigma) = M_{Pl}^4 \frac{F_{,\sigma}\sigma - F}{4F_{,\sigma}^2} . \quad (2.4)$$

Introducing the canonical scalar field ϕ instead of σ as

$$\phi = \sqrt{\frac{3}{2}} M_{Pl} \ln \left[\frac{2}{M_{Pl}^2} F_{,\sigma} \right] \quad (2.5)$$

allows one to rewrite the action S_E to the standard (quintessence or scalar-tensor) form:

$$S_E[g_{\mu\nu}, \phi] = \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right] . \quad (2.6)$$

The inverse transformation reads as follows [17, 26, 27]:

$$R_J = \left[\frac{\sqrt{6}}{M_{Pl}} V_{,\phi} + \frac{4V}{M_{Pl}^2} \right] \exp \left(\sqrt{\frac{2}{3}} \frac{\phi}{M_{Pl}} \right) , \quad (2.7)$$

$$F = \frac{M_{Pl}^2}{2} \left[\frac{\sqrt{6}}{M_{Pl}} V_{,\phi} + \frac{2V}{M_{Pl}^2} \right] \exp \left(2\sqrt{\frac{2}{3}} \frac{\phi}{M_{Pl}} \right) , \quad (2.8)$$

where $V_{,\phi} = \frac{dV}{d\phi}$, defining the function $F(R_J)$ in the parametric form with the parameter ϕ .

Being motivated by the potential (1.3), we find useful to introduce the non-canonical dimensionless field

$$y \equiv \exp \left(-\sqrt{\frac{2}{3}} \frac{\phi}{M_{Pl}} \right) = \frac{M_{Pl}^2}{2F_{,\sigma}} > 0 \quad (2.9)$$

because it is (physically) *small* during slow-roll inflation. Defining $\tilde{V}(y) = V(\phi)$ and using

$$\frac{dV}{d\phi} = -\sqrt{\frac{2}{3}} \frac{y}{M_{Pl}} \frac{d\tilde{V}}{dy} ,$$

we simplify Eqs. (2.7) and (2.8) as follows:

$$R_J = \frac{2}{M_{Pl}^2} \left(2 \frac{\tilde{V}}{y} - \tilde{V}_{,y} \right) , \quad (2.10)$$

$$F = \frac{\tilde{V}}{y^2} - \frac{\tilde{V}_{,y}}{y} , \quad (2.11)$$

respectively.

Equation (2.9) can be obtained as a consequence of Eqs. (2.10) and (2.11). Using Eqs. (2.9) and (2.10), we get

$$F_{,\sigma\sigma} = \frac{M_{Pl}^4}{4 \left(2y\tilde{V}_{,y} - 2\tilde{V} - y^2\tilde{V}_{,yy} \right)} . \quad (2.12)$$

In the Starobinsky model we have

$$\tilde{V}_{\text{Star.}}(y) = V_0(1-y)^2 , \quad \text{where} \quad V_0 = \frac{3}{4}m^2M_{Pl}^2 . \quad (2.13)$$

The general equations in this Section should be supplemented by demanding the existence of real solutions, choosing appropriate branches and imposing the physical no-ghost and no-tachyon conditions, which restrict choices of the allowed functions $F(R_J)$ and $V(\phi)$.

The simple extensions of the Starobinsky model in the form (1.3) are given by the so-called *T-models* [30] or the $\alpha = 1$ *attractors* [31] with the canonical potential

$$V(\phi) = f^2 \left(\tanh \frac{\phi}{\sqrt{6}M_{Pl}} \right) \quad (2.14)$$

in terms of a regular (monotonic) function $f(z)$, where we have introduced the new dimensionless variable

$$z = \tanh \frac{\phi}{\sqrt{6}M_{Pl}} \geq 0 . \quad (2.15)$$

In terms of this variable, the Starobinsky potential (1.3) takes the simple form

$$V_{\text{Star.}}(z) = 4V_0 \left(\frac{z}{z+1} \right)^2 . \quad (2.16)$$

The simplest T-model of inflation is defined by the even simpler function

$$V(z) = 4V_0 z^2 . \quad (2.17)$$

All these models have the same values of the inflationary observables n_s and r because for large values of the inflaton field ϕ we have $z \approx 1 - 2 \exp \left(-\sqrt{2/3} \phi / M_{Pl} \right)$. The inverse transformation (2.7) and (2.8) in terms of the new variable (2.15) takes the form

$$R_J = \frac{1}{M_{Pl}^2} (1+z) \left[(1+z) \frac{dV}{dz} + \frac{4V}{1-z} \right] \quad (2.18)$$

and

$$F = \frac{(1+z)^2}{2} \left[\left(\frac{1+z}{1-z} \right) \frac{dV}{dz} + \frac{2V}{(1-z)^2} \right]. \quad (2.19)$$

These equations are suitable in the framework of the *pole* inflation near $z \approx 1_-$ [32–34]. For example, in the Starobinsky model we find

$$R_J = 6m^2 \frac{z}{1-z} \quad \text{and} \quad F_{\text{Star.}} = 4V_0 \frac{z}{(z-1)^2}. \quad (2.20)$$

The variable z is simply connected to the variable y of Eq. (2.9) as

$$z = \frac{1-y}{1+y} \quad \text{and} \quad y = \frac{1-z}{1+z}. \quad (2.21)$$

It is to be compared to the map $z(\tilde{z})$ between a disc and a half-plane in complex analysis,

$$z = \frac{\tilde{z}-1}{\tilde{z}+1} \quad \text{and} \quad \tilde{z} = \frac{1+z}{1-z}, \quad (2.22)$$

with $0 < z < 1$ and $1 < t < \infty$ in our (real) case. It gives the mathematical origin of the variable y because $y = \tilde{z}^{-1}$, and allows us to rewrite Eqs. (2.18), (2.19) and (2.20) in the regular form (without poles) in terms of the real variable $\tilde{z}$ that is large during slow-roll inflation.

2.2 Equations of motion

In the spatially flat FLRW universe with the metric

$$ds^2 = -dt^2 + a^2(t) (dx^2 + dy^2 + dz^2),$$

the action (2.6) leads to the standard system of evolution equations:

$$6M_{Pl}^2 H^2 = \dot{\phi}^2 + 2V, \quad (2.23)$$

$$2M_{Pl}^2 \dot{H} = -\dot{\phi}^2, \quad (2.24)$$

$$\ddot{\phi} + 3H\dot{\phi} + V_{,\phi} = 0, \quad (2.25)$$

where $H = \dot{a}/a$ is the Hubble parameter, $a(t)$ is the scale factor, and the dots denote the derivatives with respect to the cosmic time t . Equation (2.23) is the Friedmann equation. In the inflationary model building, the e-foldings number

$$N_e = \ln \left(\frac{a_{\text{end}}}{a} \right), \quad (2.26)$$

where a_{end} is the value of a at the end of inflation, is considered instead of the time variable. Using the relation $d/dt = -H d/dN_e$, one can rewrite Eq. (2.23) as follows:

$$Q = \frac{2V}{6M_{Pl}^2 - \chi^2}, \quad (2.27)$$

where $Q \equiv H^2$ and $\chi = \phi' = -\dot{\phi}/H$, and the primes (here and below) denote the derivatives with respect to N_e . Equations (2.24) and (2.25) yield the dynamical system of equations:

$$Q' = \frac{1}{M_{Pl}^2} Q \chi^2, \quad \phi' = \chi, \quad \chi' = 3\chi - \frac{1}{2M_{Pl}^2} \chi^3 - \frac{1}{Q} \frac{dV}{d\phi}. \quad (2.28)$$

Using Eq. (2.27), we rewrite the last equation as

$$\chi' = 3\chi - \frac{1}{2M_{Pl}^2} \chi^3 - \frac{6M_{Pl}^2 - \chi^2}{2V} \frac{dV}{d\phi}. \quad (2.29)$$

2.3 Slow-roll approximation and inflation observables

We associate the spacetime of our Universe with the Einstein frame. The slow-roll parameters are defined by [35]

$$\epsilon = \frac{M_{Pl}^2}{2} \left(\frac{V_{,\phi}}{V} \right)^2 = \frac{y^2}{3} \left(\frac{\tilde{V}_{,y}}{\tilde{V}} \right)^2, \quad (2.30)$$

$$\eta = M_{Pl}^2 \left(\frac{V_{,\phi\phi}}{V} \right) = \frac{2y}{3\tilde{V}} \left(\tilde{V}_{,y} + y\tilde{V}_{,yy} \right). \quad (2.31)$$

The scalar spectral index n_s and the tensor-to-scalar ratio r in terms of the slow-roll parameters are given by [35]

$$n_s = 1 - 6\epsilon + 2\eta, \quad r = 16\epsilon. \quad (2.32)$$

In the slow-roll approximation, we have $\chi' \ll \chi$ and $\chi \ll M_{Pl}$. Hence, the function $\phi(N_e)$ can be found as a solution of the differential equation

$$\chi \equiv \phi' \simeq \frac{M_{Pl}^2}{V} V_{,\phi} \quad (2.33)$$

when demanding that $\epsilon = 1$ corresponds to the end of inflation with $a = a_{\text{end}}$.

Equation (2.33) is equivalent to

$$y' = \frac{2y^2 \tilde{V}_{,y}}{3\tilde{V}}. \quad (2.34)$$

Using Eq. (2.33), we connect n_s and V' as follows:²

$$n_s = 1 + \frac{d}{dN_e} \left[\ln \frac{d}{dN_e} \left(-\frac{1}{V} \right) \right] \quad (2.35)$$

that allows us to reconstruct $V(N_e)$ for a given $n_s(N_e)$.

The main cosmological parameters of inflation are given by the scalar tilt n_s and the tensor-to-scalar ratio r , whose values are constrained by the combined Planck, WMAP and BICEP/Keck observations of CMB as [20, 29]

$$n_s = 0.9649 \pm 0.0042 \quad (68\% \text{CL}) \quad \text{and} \quad r < 0.036 \quad (95\% \text{CL}). \quad (2.36)$$

The theoretical values of these observables are sensitive to the duration of inflation and the initial value of the inflaton field, ϕ_i . For instance, in the case of the Starobinsky inflation, we find

ϕ_i/M_{Pl}	5.2262	5.4971
n_s	0.961	0.969
r	0.0043	0.0027
N_e	49.258	62.335

²When using the notion of the effective potential [36], it is possible to get an analogue of Eq. (2.35) in more general models with the scalar field non-minimally coupled to the Ricci scalar and the Gauss-Bonnet term [37, 38].

The values of n_s and r do not depend upon the scalaron mass m . The amplitude of scalar perturbations is given by

$$A_s = \frac{2V}{3\pi^2 M_{Pl}^4 r} \quad , \quad (2.37)$$

while its observed value (Planck) is $A_s = 2.1 \times 10^{-9}$. Therefore, Eq. (2.37) relates the height of the inflationary potential to the tensor-to-scalar ratio r .

In the Starobinsky model, we have

$$A_s = \frac{N_e^2 m^2}{24\pi^2 M_{Pl}^2} \quad (2.38)$$

that determines the value of $m/M_{Pl} \sim \mathcal{O}(10^{-5})$.

The primordial power spectra of scalar and tensor perturbations in the inflationary models based on $f(R)$ gravity were first quantitatively derived in Refs. [39, 40].

3 Polynomial modifications of Starobinsky model

3.1 The $(R + R^2 + R^3)$ gravity models of inflation

To the best of our knowledge, adding the higher-order terms in R was first proposed in Ref. [4]. The slow-roll large-field inflation models, continuously connected to the Starobinsky model, were also studied in Refs. [11, 13–16, 18, 19, 41–43]. We revisit only those models that allow a fully analytic treatment in our approach by using the non-canonical field y defined by Eq. (2.9). We do not change the coefficient in front of the R^2 term but add a single term with a higher power in R and a dimensionless parameter in front of it. The most obvious option is a modification of the Starobinsky model by adding an R^3 term, while the corresponding scalar potential can be derived in the analytic form. It is natural to interpret the higher-order terms in R as the quantum gravity corrections to the Starobinsky model. It is our purpose to evaluate the size of those corrections that are consistent with the recent CMB measurements [20, 29] by using our methods.

A generic $(R + R^2 + R^3)$ gravity action is given by

$$S_{3\text{-gen.}} = \frac{M_{Pl}^2}{2} \int d^4x \sqrt{-g_J} \left[(1 + \delta_1) R_J + \frac{(1 + \delta_2)}{6m^2} R_J^2 + \frac{\delta_3}{36m^4} R_J^3 \right] \quad , \quad (3.1)$$

where we have introduced the three dimensionless parameters δ_i , $i = 1, 2, 3$. Similarly to the Starobinsky case, inflation is supposed to be mainly driven by the R^2 term with the dimensionless coefficient in the action, while the coefficient in front of the additional R^3 term has the negative (mass) dimension. The latter is usually exploited in the inflation literature via the standard argument that the higher-order curvature terms (beyond the quadratic order) are suppressed by powers of the Planck mass and, therefore, are irrelevant. We use the Starobinsky mass in Eq. (3.1) instead, while the variable R_J/m^2 is not small during inflation (our parameters δ_i do not have to be small). We also assume that all the coefficients in $S_{3\text{-gen.}}$ are non-negative in order to avoid problems with ghosts and negative values of the scalar potential.

The corresponding inflaton scalar potential (2.4) is given by

$$V(\sigma) = \frac{M_{Pl}^2 \sigma^2 \left(1 + \delta_2 + \frac{\delta_3}{3m^2} \sigma \right)}{12m^2 \left[1 + \delta_1 + \left(\frac{1+\delta_2}{3} \right) \frac{\sigma}{m^2} + \frac{\delta_3}{12m^4} \sigma^2 \right]^2} = \frac{16V_0 \tilde{\sigma}^2 [3(1 + \delta_2) + \delta_3 \tilde{\sigma}]}{3 [12(1 + \delta_1) + 4(1 + \delta_2) \tilde{\sigma} + \delta_3 \tilde{\sigma}^2]^2} \quad , \quad (3.2)$$

where the dimensionless variable $\tilde{\sigma} = \sigma/m^2$ has been introduced.

It is easy to see that $V(0) = 0$, $V(\tilde{\sigma}) > 0$ at $\tilde{\sigma} > 0$, and V tends to zero at $\tilde{\sigma} \rightarrow +\infty$, while the potential has a maximum at some positive value of $\tilde{\sigma}$. The extreme equation $V' = 0$ has only one positive root given by

$$\tilde{\sigma}_{\max.} = 6\sqrt{\frac{1+\delta_1}{\delta_3}}. \quad (3.3)$$

This case is qualitatively different from the pure R^2 -gravity inflation, because at $\delta_3 = 0$ the potential (3.2) is a monotonically increasing function of $\tilde{\sigma}$, approaching a positive constant at $\tilde{\sigma} \rightarrow +\infty$, whereas when $\delta_3 > 0$ one must have $\tilde{\sigma} < \tilde{\sigma}_{\max.}$ during and after (hilltop) inflation.

To study the impact of the R^3 -term on inflation in more detail, let us consider the simplest non-trivial case with $\delta_1 = \delta_2 = 0$, in which Eq. (2.9) implies

$$\frac{1}{y} = 1 + \frac{1}{3}\tilde{\sigma} + \frac{\delta_3}{12}\tilde{\sigma}^2. \quad (3.4)$$

Equation (3.4) is a quadratic equation on $\tilde{\sigma}$ as a function of y . The only positive root of this equation is given by

$$\tilde{\sigma} = \frac{2}{\delta_3} \left[\sqrt{1 + 3\delta_3(y^{-1} - 1)} - 1 \right] = \frac{2}{\delta_3} \left[\sqrt{1 + 3\delta_3 \left(e^{\sqrt{\frac{2}{3}}\phi/M_{Pl}} - 1 \right)} - 1 \right]. \quad (3.5)$$

Using Eqs. (2.4) and (2.9), we find the scalar potential in terms of y or the inflaton field ϕ as follows:

$$\begin{aligned} \tilde{V}(y) &= \frac{4V_0}{27\delta_3^2 y} \left[y + 2\sqrt{y(y + 3\delta_3(1 - y))} \right] \left(y - \sqrt{y(y + 3\delta_3(1 - y))} \right)^2, \\ V(\phi) &= \frac{4V_0}{27\delta_3^2} e^{-2\sqrt{2/3}\phi/M_{Pl}} \left(\sqrt{1 + 3\delta_3 \left(e^{\sqrt{2/3}\phi/M_{Pl}} - 1 \right)} - 1 \right)^2 \times \\ &\quad \times \left[1 + 2\sqrt{1 + 3\delta_3 \left(e^{\sqrt{2/3}\phi/M_{Pl}} - 1 \right)} \right]. \end{aligned} \quad (3.6)$$

It is worth noticing that $\tilde{V}_{\text{Star.}}(y)$ is reproduced in the limit $\delta_3 \rightarrow 0$.

When $\delta_3 = 1/3$, the potential (3.6) is greatly simplified to

$$V(\phi) = \frac{4}{3}V_0 [y^2 - 3y + 2\sqrt{y}] = \frac{4}{3}V_0 e^{-2\sqrt{2/3}\phi/M_{Pl}} \left(1 + 2e^{\sqrt{3/2}\phi/M_{Pl}} - 3e^{\sqrt{2/3}\phi/M_{Pl}} \right). \quad (3.7)$$

In terms of the variable z defined by Eq. (2.15), the potential reads

$$V(z) = \frac{2V_0 \left[(1 + (18\delta_3 - 1)z)(z - 1)^2 - [(z - 1)^2 - 12\delta_3(z - 1)z]^{3/2} \right]}{27\delta_3^2(1 + z)^2(z - 1)}. \quad (3.8)$$

We also find in this case

$$R_J = \frac{m^2 \left(z - 1 + \sqrt{(12\delta_3 z + 1 - z)(1 - z)} \right)}{\delta_3(1 - z)} \quad (3.9)$$

and

$$z = \frac{R_J (\delta_3 R_J + 2m^2)}{12m^4 + \delta_3 R_J^2 + 2m^2 R_J} = \frac{\tilde{\sigma} (\delta_3 \tilde{\sigma} + 2)}{12 + \delta_3 \tilde{\sigma}^2 + 2\tilde{\sigma}} \quad . \quad (3.10)$$

The profile of the potential $V(\phi)$ is given on Fig. 1. The maximum occurs at

$$\phi_{\max.} = \sqrt{\frac{3}{2}} M_{Pl} \ln \left[2 \left(2 + \delta_3^{-1/2} \right) \right] \quad , \quad (3.11)$$

while this value does not depend upon m .




Figure 1: The normalized potential $V(\phi)/V_0$ of Eq. (3.6) for $\delta_3 = 0.000001$ (red), $\delta_3 = 0.000247$ (blue), and $\delta_3 = 1/3$ (green).

As is clear from Fig. 1, the plateau of the potential (on the left-hand-side from the maximum) becomes longer, as well as the duration of slow-roll, with decreasing δ_3 .

The key discriminator for viable inflation in the given class of models is the value of the scalar perturbation index n_s . For example, in the case of the potential (3.7) with $\delta_3 = 1/3$, the n_s never exceeds 0.9 for any value of the canonical inflaton field ϕ , so that it is not suitable for inflation.

The condition $\phi_i < \phi_{\max.}$ yields the additional restriction on the possible initial values of ϕ . Equation (3.11) can be written to the following form:

$$\delta_3 = \frac{4}{\left(e^{\sqrt{2/3} \phi_{\max.}/M_{Pl}} - 4 \right)^2} \quad , \quad (3.12)$$

being represented by the blue curve on the left-hand-side of Fig. 2.

The upper bound on the parameter δ_3 can be estimated by assuming the observable value of n_s to be calculated at the maximum of the potential. Then we find

$$n_s(\phi_{\max.}) = 1 - \frac{8\sqrt{\delta_3} (1 + 4\sqrt{\delta_3} + 4\delta_3)}{3 (3\sqrt{\delta_3} + 1) (2\sqrt{\delta_3} + 1)^2} \quad . \quad (3.13)$$

Since observations require $n_s > 0.960$, we get $\delta_3 < 0.0002467$. The dependence of n_s upon δ_3 is given on the right-hand-side of Fig. 2. Therefore, the domain of allowed values of δ_3 and ϕ is highly restricted.




Figure 2: The allowed range of δ_3 and ϕ from the observational constraints (Planck): $0.961 < n_s < 0.969$ (left), and the dependence of n_s upon δ_3 (right), under the assumption that inflation started at the maximum of the potential.

Since viable inflation requires $\delta_3 \ll 1$, it allows us to consider δ_3 as a truly small parameter and expand the potential in power series of δ_3 as follows:

$$\tilde{V}(y) = V_0(y-1)^2 \left[1 + \frac{y-1}{2y}\delta_3 + \frac{9(y-1)^2}{16y^2}\delta_3^2 + \mathcal{O}(\delta_3^3) \right]. \quad (3.14)$$

In this approximation, using Eqs. (2.30) and (2.31), we get

$$\begin{aligned} \epsilon &\simeq \frac{4y^2}{3(y-1)^2} + \frac{2}{3(y-1)}\delta_3 + \frac{14y+1}{12y^2}\delta_3^2, \\ \eta &\simeq \frac{4y(2y-1)}{3(y-1)^2} + \frac{3y+1}{3y(y-1)}\delta_3 + \frac{21y+16}{12y^2}\delta_3^2, \end{aligned}$$

and, hence,

$$n_s \simeq 1 - \frac{8y(y+1)}{3(y-1)^2} - \frac{2(3y-1)}{3y(y-1)}\delta_3 - \frac{21y-13}{6y^2}\delta_3^2. \quad (3.15)$$

The conditions $\epsilon(y_{end}) = 1$ and $y_{end} = 2\sqrt{3} - 3$ at $\delta_3 = 0$ give

$$y_{end} = 2\sqrt{3} - 3 + \frac{45 - 26\sqrt{3}}{12\sqrt{3} - 21}\delta_3 + \frac{2146\sqrt{3} - 3717}{9(4\sqrt{3} - 7)^2}\delta_3^2, \quad (3.16)$$

Solving Eq. (2.34), we get for small δ_3 :

$$N_e = \frac{3(y \ln(y) + 1)}{4y} + \frac{3y^2 - 3y + 1}{16y^3}\delta_3 + \frac{3(70y^4 - 110y^3 + 80y^2 - 25y + 2)}{640y^5}\delta_3^2 - N_0, \quad (3.17)$$

where N_0 can be obtained by the condition $N_e(y_{end}) = 0$:

$$\begin{aligned} N_0 &= \frac{(1896102\sqrt{3} - 3284145) \ln(2\sqrt{3} - 3) - 293328\sqrt{3} + 508059}{4(2\sqrt{3} - 3)^5(4\sqrt{3} - 7)^2} + \\ &\quad \frac{2578194\sqrt{3} - 4465563}{16(2\sqrt{3} - 3)^5(4\sqrt{3} - 7)^2}\delta_3 + \frac{187776774\sqrt{3} - 325238913}{640(2\sqrt{3} - 3)^5(4\sqrt{3} - 7)^2}\delta_3^2. \end{aligned} \quad (3.18)$$

With a value of δ_3 within $0 \leq \delta_3 \leq 0.00025$ we obtain $N_0 \simeq 1.040$.

In Fig. 2, one can see that suitable values of ϕ are not less than in the case of the Starobinsky inflation. So, to calculate inflationary parameters we should take $\phi \geq 5.226 M_{Pl}$ that corresponds to $y \leq 0.0140$. For these values, N_e is an increasing function of δ_3 , so the number of e-folding during inflation is always more than 49.26 that corresponds to the Starobinsky model. The request $N_e < 65$ gives the additional restriction of the maximal value of δ_3 . Namely, for $\delta_3 = 0.00012$, the condition $n_s > 0.961$ gives $\phi > 5.438 M_{Pl}$. The corresponding $N_e > 65.1$ that is not suitable for inflationary scenario.

We come to the conclusion that the model under investigation gives the inflationary parameters that do not contradict observations only if $\delta_3 < 0.00012$ and the inflation started in the narrow domain of the scalar field ϕ values (a part of the marked green domain in the left picture of Fig. 2), which implies that this inflationary scenario is rather unrealistic. The same model was also studied in detail in Ref. [18, 19]. A similar inflationary model in the framework of the unimodular gravity was proposed in Ref. [43].

It is no surprise that the dimensionless parameter δ_3 in front of the R^3 -term must be very small³ because slow roll inflation in $f(R)$ gravity only applies for the range of R (in the large curvature regime) where $f(R)/R^2$ is a slowly-changing function of R , together with its first and second derivatives with respect to $\ln R$ [22]. Our considerations above provide the quantitative estimates for the δ_3 and n_s .

3.2 The $(R + R^2 + R^4)$ gravity model of inflation

In this Subsection, we consider another model defined by

$$F(R_J) = \frac{M_{Pl}^2}{2} \left[R_J + \frac{1}{6m^2} R_J^2 + \frac{\delta_4 R_J^4}{48m^6} \right] \quad (3.19)$$

with the dimensionless parameter $\delta_4 > 0$, as the natural alternative to the previous model.

We compute the inflaton scalar potential from Eq. (9) as follows:

$$V = V_0 \frac{\tilde{\sigma}^2(8 + 3\tilde{\sigma}^2\delta_4)}{72y^{-2}} = V_0 \frac{2\tilde{\sigma}^2(8 + 3\delta_4\tilde{\sigma}^2)}{(12 + 4\tilde{\sigma} + \delta_4\tilde{\sigma}^3)^2} \quad , \quad (3.20)$$

where y is related to F' by Eq. (2.9). In the case under consideration we have

$$y^{-1} = 1 + \frac{\tilde{\sigma}}{3} + \frac{\delta_4}{12} \tilde{\sigma}^3 \quad . \quad (3.21)$$

Solving the cubic equation (3.21) yields $\tilde{\sigma}$ in terms of y^{-1} ,

$$\tilde{\sigma} = \frac{Z}{3\delta_4} - \frac{4}{Z} \quad , \quad \text{where} \quad Z = \sqrt[3]{\left(162(y^{-1} - 1) + 6\sqrt{3}\sqrt{243(y^{-1} - 1)^2 + 16\delta_4^{-1}}\right)} \delta_4^2 \quad . \quad (3.22)$$

Assuming $\delta_4 \ll 1$ and using Eq. (3.22), we can expand $\tilde{\sigma}^2$ in power series of δ_4 near zero as follows:

$$\tilde{\sigma}^2 = 9(y^{-1} - 1)^2 - \frac{81(y^{-1} - 1)^4}{2} \delta_4 + \frac{5103(y^{-1} - 1)^6}{16} \delta_4^2 + \mathcal{O}(\delta_4^3) \quad . \quad (3.23)$$

³It also applies to the parameter δ_4 in front of the R^4 -term studied in the next Subsection 3.2.

Substituting Eq. (3.23) into Eq. (3.20), we get the scalar potential with the same accuracy,

$$V(y) \approx V_0 \left[\frac{(y^{-1} - 1)^2}{y^{-2}} - \frac{9}{8} \frac{(y^{-1} - 1)^4}{y^{-2}} \delta_4 + \frac{81}{16} \frac{(y^{-1} - 1)^6}{y^{-2}} \delta_4^2 \right] \\ \approx \tilde{V}_{\text{Star.}}(y) \left[1 - \frac{9}{8} (y^{-1} - 1)^2 \delta_4 + \frac{81}{16} (y^{-1} - 1)^4 \delta_4^2 \right], \quad (3.24)$$

where the potential $\tilde{V}_{\text{Star.}}(y)$ has been defined by Eq. (2.13).

The profile of the scalar potential in the $(R + R^2 + R^4)$ model versus the scalar potential of the Starobinsky $(R + R^2)$ model is given in Fig. 3.




Figure 3: The scalar potential in the $(R + R^2 + R^4)$ model (red) versus the scalar potential in the Starobinsky $(R + R^2)$ model (green). The parameter value is $\delta_4 = 10^{-7}$. The approximated potential (3.24) in the $(R + R^2 + R^4)$ model is given by the blue line with the same value of $\delta_4 = 10^{-7}$.

The slow-roll parameters are

$$\epsilon \approx \frac{4}{3(y^{-1} - 1)^2} - 3y^{-1}\delta_4 + \frac{27(y^{-1} + 14)(y^{-1} - 1)^2 y^{-1}}{16} \delta_4^2 \quad (3.25)$$

and

$$\eta \approx -\frac{4(y^{-1} - 2)}{3(y^{-1} - 1)^2} - \frac{3(2y^{-1} + 3)y^{-1}}{2} \delta_4 - \frac{81y^{-1}(-10y^{-3} + 13y^{-2} + 4y^{-1} - 7)}{16} \delta_4^2. \quad (3.26)$$

Accordingly, the inflationary observables $n_s = 1 - 6\epsilon + 2\eta$ and $r = 16\epsilon$ are given by

$$n_s \approx \frac{3y^{-2} - 14y^{-1} - 5}{3(y^{-1} - 1)^2} - 3y^{-1}(2y^{-1} - 3)\delta_4 - \frac{81y^{-1}(-9y^{-3} + 25y^{-2} - 23y^{-1} + 7)}{8} \delta_4^2$$

and

$$r \approx \frac{64}{3(y^{-1} - 1)^2} - 48y^{-1}\delta_4 + 27y^{-1}(y^{-1} + 14)(y^{-1} - 1)^2\delta_4^2 \quad .$$

The e-foldings number N_e as a function of the inflaton field is

$$\begin{aligned} N_e \approx & \frac{3}{4}(y^{-1} + \ln y) + \frac{27}{128}(y^{-1} - 1)^4\delta_4 \\ & + \frac{243}{256}(7^{-1}y^{-7} - 2y^{-6} + 9y^{-5} - 20y^{-4} + 25y^{-3} - 18y^{-2} + 7y^{-1})\delta_4^2 - N_0 \quad , \end{aligned} \quad (3.27)$$

where the integration constant N_0 is fixed by the condition $\epsilon(y_{end}) = 1$. A numerical solution to the condition $\epsilon = 1$ with $\delta_4 \leq 10^{-6}$ yields the approximate values of the field and the e-foldings number at the end of inflation as $\phi_{end} \approx 0.9402M_{Pl}$ and $N_0 \approx 1.040$. The relevant values of the canonical inflaton field ϕ_i at the beginning of inflation versus the values of δ_4 are given in Fig. 4.




Figure 4: The green area on these pictures describes the values of the parameter δ_4 and $\phi = \phi_i$ that are in agreement with the observed values of the spectral index n_s and the tensor-to-scalar ratio r . The upper boundary of the green area (dark blue curve) corresponds to $n_s = 0.961$ and the lower boundary (blue curve) corresponds to $n_s = 0.969$. In the left picture, the blue area gives the values of δ_4 and ϕ_i leading to the observed values of r , the red curve corresponds to the maximum of the potential V , and the grey curve shows the boundary of applicability of our approximation. In the right picture, the magenta area shows the values of the parameter δ_4 and ϕ_i required for inflation with a decreasing scalar field. The yellow area corresponds to the e-foldings number N_e of the Starobinsky inflation for the allowed interval $0.961 \leq n_s \leq 0.969$, the left black dotted line corresponds to $N_e \approx 49.258$, the right orange dotted line corresponds to $N_e \approx 62.335$, and the red line corresponds to $r = 0.0027$.

The red curve in the middle of the left picture in Fig. 4 gives the function $\delta(\phi_{\max})$, where $\phi_{\max}$ is the point of the maximum potential of V . This function can be presented in

an analytical form

$$\delta_4(\phi_{max}) = \frac{y_{max}^{-1} + 1 - \sqrt{y_{max}^{-2} - 30y_{max}^{-1} - 15}}{9(2y_{max}^{-1} + 1)(y_{max}^{-1} - 1)^2},$$

where $y_{max} = y(\phi_{max})$. The scalar field ϕ tends to zero during inflation if its initial value is less than ϕ_{max} .

As shown in Fig. 4, the maximum allowed value of δ_4 is about 2×10^{-7} . In the case of the Starobinsky inflation, $\delta_4 = 0$, limiting the maximum value of the scalar spectral index n_s leads to a minimum value of the ratio of the tensor to the scalar $r = 0.0027$. When $\delta_4 > 0$, the minimum value of r decreases. In the right picture of Fig. 4, the red line corresponds to the value $r = 0.0027$ for $\delta_4 \neq 0$. To the left of the red line we have $r > 0.0027$, and $r < 0.0027$ to the right.

Our consideration provides quantitative estimation for value of δ_4 for which the parameters n_s and r are consistent with current observations.

4 The $(R + R^{3/2} + R^2)$ gravity model of inflation

4.1 The inflaton potential

Our methods apply to yet another model of inflation based on the modified gravity with the $(R + R^{3/2} + R^2)$ terms,

$$F(R_J) = \frac{M_{Pl}^2}{2} \left[R_J + \frac{1}{6m^2} R_J^2 + \frac{\delta}{m} R_J^{3/2} \right], \quad (4.1)$$

where we have introduced the dimensionless parameter δ .

The $R^{3/2}$ -term in $F(R)$ gravity (or in the Jordan frame) arises in an approximate description of the Higgs field with a small cubic term in its scalar potential and a large non-minimal coupling to R [44]. The $R^{3/2}$ term also appears in the (chiral) modified supergravity [12, 45].

The possible values of δ are restricted by the condition (1.5). Given $\tilde{\sigma} > 0$, we find

$$F_{,\sigma} = \frac{M_{Pl}^2}{6} \left(\sqrt{\tilde{\sigma}} + \frac{9\delta}{4} \right)^2 - \frac{3}{16} (27\delta^2 - 16) > 0 \quad (4.2)$$

when $\delta > -4\sqrt{3}/9$, and

$$F_{,\sigma\sigma} = \frac{M_{Pl}^2}{24m^2} \left(4 + \frac{9\delta}{\sqrt{\tilde{\sigma}}} \right) > 0 \quad (4.3)$$

only when $\delta > 0$. Hence, the condition $\delta > 0$ is necessary to get a stable $F(R)$ gravity model for all $R_J > 0$.

The corresponding scalar potential (2.4) is given by

$$V = \frac{4V_0\tilde{\sigma}(3\delta\sqrt{\tilde{\sigma}} + \tilde{\sigma})}{(6 + 9\delta\sqrt{\tilde{\sigma}} + 2\tilde{\sigma})^2}. \quad (4.4)$$

Equation (2.9) in this case is a quadratic equation on $\sqrt{\tilde{\sigma}}$, and its only real solution is

$$\tilde{\sigma} = \frac{3(1-y)}{y} + \frac{9\delta}{8y} \left[9\delta y - \sqrt{3y(27\delta^2 y - 16y + 16)} \right]. \quad (4.5)$$

The potential (4.4) can be rewritten as

$$\begin{aligned}\tilde{V} &= \frac{V_0}{2304y^2} (s + 3\delta y) (s - 9\delta y)^3 \\ &= \frac{243V_0\delta^4 y^2}{256} \left(3\sqrt{1 + \frac{16(1-y)}{27\delta^2 y}} + 1 \right) \left(\sqrt{1 + \frac{16(1-y)}{27\delta^2 y}} - 1 \right)^3,\end{aligned}\quad (4.6)$$

where we have introduced $s = \sqrt{3y(27\delta^2 y - 16y + 16)}$.

When $\delta = 4\sqrt{3}/9$, the $F_{,\sigma}$ function is a perfect square, and the potential simplifies as

$$\tilde{V}_{\text{special}}(y) = \frac{V_0}{3} (3 + \sqrt{y}) (1 - \sqrt{y})^3, \quad (4.7)$$

or

$$V_{\text{special}}(\phi) = \frac{V_0}{3} \left(e^{\phi/(\sqrt{6}M_{Pl})} - 1 \right)^3 \left(1 + 3e^{\phi/(\sqrt{6}M_{Pl})} \right) e^{-2\sqrt{2/3}\phi/M_{Pl}}.$$




Figure 5: The potential $V(\phi)$ for $\delta = 0$ (red), $\delta = 1/5$ (blue), $\delta = 4\sqrt{3}/9$ (green), and $\delta = 5$ (black).

The profile of the potential $V(\phi)$ as a function of the canonical inflaton ϕ is given in Fig. 5.

It is possible to further generalize our model by replacing the $R^{3/2}$ -term by the $(R + R_0)^{3/2}$ -term with a positive constant R_0 [12, 13].

4.2 The inflationary parameters

Equations (2.30) and (2.31) give the slow-roll parameters in our model (4.1) as

$$\epsilon = \frac{48y^2 [s\delta + (8 - 9\delta^2)y]^2}{(s - 9\delta y)^2 (s + 3\delta y)^2} \quad (4.8)$$

and

$$\eta = \frac{4y [24(3\delta + 2)(3\delta - 2)(27\delta^2 - 16)y^3 - (27\delta(21s\delta^2 - 16s - 80\delta) + 768)y^2 - s^3\delta]}{(3\delta y + s)(s - 9\delta y)^2 s} . \quad (4.9)$$

Therefore, according to Eq. (2.32), the inflationary parameters are given by

$$\begin{aligned} n_s = & 1 + \frac{8y (3s(3\delta(9\delta^2 - 16)s + 720\delta^2 - 256)y^2 - s^3\delta(39\delta y + s))}{(-9\delta y + s)^2 (3\delta y + s)^2 s} \\ & - \frac{8y [72\delta(4 - 9\delta^2)(27\delta^2 - 16)y^4 + ((-1215\delta^4 - 432\delta^2 + 768)s - 144\delta(45\delta^2 - 16))y^3]}{(-9\delta y + s)^2 (3\delta y + s)^2 s} \end{aligned} \quad (4.10)$$

and

$$r = \frac{768 y^2 (-9\delta^2 y + s\delta + 8y)^2}{(-9\delta y + s)^2 (3\delta y + s)^2} . \quad (4.11)$$

The slow-roll evolution equation (2.34) allows us to relate N_e with y at the end of inflation,

$$\begin{aligned} N_e = & \left(\frac{9}{8} - \delta^{-2} \right) \ln [9\delta^3 (9\delta y - s) + 24(1 - 4y)\delta^2 + 8(\delta s + 4y)] \\ & + \left(\delta^{-2} - \frac{3}{8} \right) \ln y + \frac{s}{4\delta y} - N_0 , \end{aligned} \quad (4.12)$$

where the integration constant N_0 is fixed by the condition $N_e(y_{end}) = 0$. The analytic formula for $N_0(\delta)$ is obtained by substituting $N_e = 0$ and $y = y_{end}$ (see below). It results in a long equation that is not very illuminating, so we do not present it here.

It follows from Eq. (4.8) that the condition $\epsilon = 1$ with arbitrary positive parameter δ leads to a quadratic equation on $y = y_{end}$ with the only solution as

$$y_{end} = \frac{3(4 - 3\delta^2 + \sqrt{3}\delta^2) - \sqrt{9(4 - 3\delta^2 + \sqrt{3}\delta^2)^2 - 72(2 - 3\delta^2)}}{2(2 - 3\delta^2)(3 + 2\sqrt{3})} . \quad (4.13)$$

It is worth noticing that this solution has no singularity at $\delta = \sqrt{2/3}$, while $y_{end}(\delta)$ is a smooth monotonically decreasing function.

The slow-roll parameters ϵ and η remain finite in the limit $\delta \rightarrow +\infty$ at fixed y ,

$$\epsilon_\infty(y) = \frac{(2y + 1)^2}{3(1 - y)^2} , \quad \eta_\infty(y) = \frac{2(4y^2 + y + 1)}{3(1 - y)^2} . \quad (4.14)$$

Since the value of y at the end of inflation is determined by the condition $\epsilon(y_{end}) = 1$, y_{end} also approaches a finite limit as $\delta \rightarrow +\infty$, which is given by a solution to the equation

$$\epsilon_\infty = \frac{(2y + 1)^2}{3(1 - y)^2} = 1. \quad (4.15)$$

This equation has only one positive solution $y_{end}|_{\delta \rightarrow +\infty} = 3\sqrt{3} - 5 \approx 0.196$.

The amplitude of scalar perturbations (2.37) is given by

$$A_s = \frac{(-9\delta y + s)^5 (3\delta y + s)^3 m^2}{3538944y^4\pi^2 (-9\delta^2 y + s\delta + 8y)^2} . \quad (4.16)$$

The observed value of A_s determines the value of the parameter m .

We summarize our results for various values of the parameter δ in Table 1 and Fig. 6. The values of the tensor-to-scalar ratio r at $\delta = 5$ are about four times more the corresponding values in the Starobinsky model. When $\delta > 5$, the r is almost constant. The value of y_{end} at $\delta = 25$ is close to its value at $\delta \rightarrow +\infty$. Unlike the models with the R^3 and R^4 terms, the potential has no extremum at $\phi < 0$, so there is no fine-tuning of initial conditions. The number of e-foldings before the end of inflation monotonically increases for $\delta > 1$. Restricting this number to be less or equal to 65, we get an estimation for the maximal value of the parameter as $\delta = 100$ at which the model still does not contradict observations.

Table 1: The values of y , N_e and r corresponding to $n_s = 0.961$ and $n_s = 0.969$, respectively, and the values of y_{end} for some values of the parameter δ .

δ	y_{end}	$y_{in, n_s=0.961}$	$y_{in, n_s=0.969}$	$N_{e, n_s=0.961}$	$N_{e, n_s=0.969}$	$r_{n_s=0.961}$	$r_{n_s=0.969}$
0	0.464	0.0140	0.0112	49.3	62.3	0.0043	0.0027
0.2	0.395	0.00682	0.00505	45.0	56.8	0.0096	0.0065
$\frac{4\sqrt{3}}{9}$	0.299	0.00146	0.000968	48.1	60.9	0.0152	0.0099
1	0.279	0.000939	0.000616	49.4	62.4	0.0157	0.0102
5	0.205	$4.32 \cdot 10^{-5}$	$2.75 \cdot 10^{-5}$	56.3	69.7	0.0168	0.0108
10	0.199	$1.08 \cdot 10^{-5}$	$6.91 \cdot 10^{-6}$	58.7	72.0	0.0168	0.0108
25	0.197	$1.74 \cdot 10^{-6}$	$1.11 \cdot 10^{-6}$	61.4	74.8	0.0169	0.0108
50	0.196	$4.34 \cdot 10^{-7}$	$2.77 \cdot 10^{-7}$	63.5	76.9	0.0169	0.0108
100	0.196	$1.09 \cdot 10^{-7}$	$6.92 \cdot 10^{-8}$	65.5	79.1	0.0169	0.0108

When $\delta = 4\sqrt{3}/9$, the function $F_{,\sigma}$ simplifies as

$$F_{,\sigma} = \frac{M_{Pl}^2}{6} \left(\sqrt{\sigma} + \sqrt{3} \right)^2 .$$

Accordingly, the slow-roll parameters are also simplified as

$$\epsilon = \frac{4y^2 (2\sqrt{y} + y)^2}{3 (\sqrt{y} - y)^2 (3\sqrt{y} + y)^2} \quad (4.17)$$

and

$$\eta = \frac{4y^2 (2\sqrt{y} + 2y - 1)}{3 (\sqrt{y} - y)^2 (3\sqrt{y} + y)} . \quad (4.18)$$

In this special case we find

$$n_s = 1 - \frac{8y^2 (7y + 4y\sqrt{y} + y^2 + 3\sqrt{y})}{3 (\sqrt{y} - y)^2 (3\sqrt{y} + y)^2} \quad (4.19)$$




Figure 6: The inflaton field values against the values of the parameter δ . The green area corresponds to the observational restrictions on n_s and r . The blue area is defined by the restrictions on r only. When $\delta > 5$, the allowed domain is restricted by the lines $\delta y^2 = \text{const.}$

and

$$r = \frac{64y^2 (2\sqrt{y} + y)^2}{3(\sqrt{y} - y)^2 (3\sqrt{y} + y)^2} . \quad (4.20)$$

Equation (4.12) also simplifies to

$$N_e = \frac{9}{4\sqrt{y}} + \frac{21}{16} \ln y - \frac{9}{8} \ln(\sqrt{y} + 2) - N_0 , \quad (4.21)$$

so that $\epsilon = 1$ is at $y_{\text{end}} = (\sqrt{6\sqrt{3} - 8} - 1)^2$, and the integration constant N_0 is given by

$$N_0 = -\frac{1}{8} \left[21 \ln(\sqrt{6\sqrt{3} - 8} - 1) - 9 \ln(1 + \sqrt{6\sqrt{3} - 8}) + (6 + 4\sqrt{3}) \left(1 + \sqrt{6\sqrt{3} - 8} \right) \right] \\ \approx -1.48 . \quad (4.22)$$

The inflationary parameters in the special case are also given in Table 1. It is worth noticing that the values of the tensor-to-scalar ratio r are significantly higher than those in the Starobinsky model.

The amplitude of scalar perturbations at the special value of δ is given by

$$A_s \approx \frac{2\tilde{V}}{3\pi^2 M_{Pl}^4 r} \approx \frac{m^2 (1 - \sqrt{y})^5 (3 + \sqrt{y})^3}{128\pi^2 M_{Pl}^2 (2 + \sqrt{y})^2} . \quad (4.23)$$

Its observed value $A_s \approx 2.1 \cdot 10^{-9}$ implies $m \approx 2 \cdot 10^{-5}$.

5 Deforming the scalar potential in the Starobinsky model with analytic F -functions

The Starobinsky inflation based on the inflaton potential (1.3) is a *large* ϕ -field inflation, with the relevant values of ϕ around the Planck scale. Hence, the field y defined by (2.9) is *small* during slow-roll inflation. The inflaton potential (1.3) as a function of y is

$$V(\phi) = \tilde{V}(y) = V_0 [1 - 2y + \mathcal{O}(y^2)] , \quad (5.1)$$

where only the first two terms are essential for the CMB observables [46]. The inflaton potential (1.3) can therefore be modified as

$$\tilde{V}(y) = V_0 [1 - 2y + y^2 \omega(y)] , \quad (5.2)$$

with arbitrary analytic function $\omega(y)$ *without* changing the CMB observables predicted by the Starobinsky model, at least for those values of ω that are not very large. The Starobinsky model appears at $\omega = 1$ with inflation taking place for positive values of R_J .

The physical conditions (1.5) are satisfied with the potential (5.2). In particular, we find the second derivative $F_{,\sigma\sigma}(y)$ in the form

$$F_{,\sigma\sigma}(y) = \frac{M_{Pl}^2}{3m^2 \left(2 + 2y^3 \frac{d\omega}{dy} + y^4 \frac{d^2\omega}{dy^2} \right)} . \quad (5.3)$$

Equation (2.10) reads

$$\tilde{\sigma} \equiv \frac{R_J}{m^2} = 3 \left(\frac{1}{y} - 1 - \frac{1}{2} y^2 \frac{d\omega}{dy} \right) , \quad (5.4)$$

and Eq. (2.11) is given by

$$F = V_0 \left(\frac{1}{y^2} - \omega - y \frac{d\omega}{dy} \right) . \quad (5.5)$$

As a check, in the Starobinsky case, $\omega = 1$ and $V = V_0(1 - y)^2$, and Eq. (2.10) gives

$$y = \left(1 + \frac{R_J}{3m^2} \right)^{-1} . \quad (5.6)$$

Substituting it into Eq. (2.11), we get

$$F_{Star.}(R_J) = \frac{M_{Pl}^2}{2} \left(R_J + \frac{R_J^2}{6m^2} \right) , \quad (5.7)$$

as it should be. Moreover, when ω is an arbitrary constant, we find

$$F(R_J) = F_{Star.}(R_J) - \Lambda , \quad (5.8)$$

where $\Lambda = V_0(1 - \omega)$ is a cosmological constant.

5.1 New case I

As a new example, let us now consider the non-trivial case with

$$\omega(y) = \omega_0 + \omega_1 y , \quad (5.9)$$

where $\omega_0 \leq 1$ and $\omega_1 > 0$ are constants. The constant ω_1 should be positive for the potential V bounded from below. The inequality $\omega_0 \leq 1$ is needed for positivity of a cosmological constant, see Eq. (5.8).

Equation (5.4) leads to the depressed cubic equation

$$y^3 + \frac{2}{\omega_1} \left(1 + \frac{R_J}{3m^2}\right) y - \frac{2}{\omega_1} \equiv y^3 + py + q = 0 \quad (5.10)$$

with the negative discriminant

$$\Delta = -(4p^3 + 27q^2) = -\frac{32}{\omega_1^3} \left(1 + \frac{R_J}{3m^2}\right)^3 - \frac{108}{\omega_1^2} < 0 , \quad (5.11)$$

so that it has only one real root given by the Cardano formula

$$\begin{aligned} y &= \left(-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}\right)^{1/3} + \left(-\frac{q}{2} - \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}\right)^{1/3} \\ &= \frac{1}{\omega_1^{1/3}} \left(1 + \sqrt{1 + \frac{8}{27\omega_1} \left(1 + \frac{R_J}{3m^2}\right)^3}\right)^{1/3} + \frac{1}{\omega_1^{1/3}} \left(1 - \sqrt{1 + \frac{8}{27\omega_1} \left(1 + \frac{R_J}{3m^2}\right)^3}\right)^{1/3} . \end{aligned} \quad (5.12)$$

Equation (5.5) yields the explicit F -function in this case as follows:

$$\begin{aligned} \frac{F}{V_0} &= \frac{1}{y^2} - \omega_0 - 2\omega_1 y \\ &= \omega_1^{2/3} \left[\left(1 + \sqrt{1 + \frac{8}{27\omega_1} \left(1 + \frac{R_J}{3m^2}\right)^3}\right)^{1/3} + \left(1 - \sqrt{1 + \frac{8}{27\omega_1} \left(1 + \frac{R_J}{3m^2}\right)^3}\right)^{1/3} \right]^{-2} \\ &\quad - 2\omega_1^{2/3} \left[\left(1 + \sqrt{1 + \frac{8}{27\omega_1} \left(1 + \frac{R_J}{3m^2}\right)^3}\right)^{1/3} + \left(1 - \sqrt{1 + \frac{8}{27\omega_1} \left(1 + \frac{R_J}{3m^2}\right)^3}\right)^{1/3} \right] \\ &\quad - \omega_0 , \end{aligned} \quad (5.13)$$

where

$$\begin{aligned} \omega_0 &= \omega_1^{2/3} \left[\left(1 + \sqrt{1 + \frac{8}{27\omega_1}}\right)^{1/3} + \left(1 - \sqrt{1 + \frac{8}{27\omega_1}}\right)^{1/3} \right]^{-2} \\ &\quad - 2\omega_1^{2/3} \left[\left(1 + \sqrt{1 + \frac{8}{27\omega_1}}\right)^{1/3} + \left(1 - \sqrt{1 + \frac{8}{27\omega_1}}\right)^{1/3} \right] . \end{aligned} \quad (5.14)$$

The limit $\omega_1 \rightarrow 0$ is smooth and gives back the case (5.8). The shape of the scalar potential of the canonical inflaton field ϕ is shown on the left-hand-side of Fig. 7: it remains largely

unchanged for the values of the deformation parameter up to $\omega_1 \approx 10$, though the "waterfall" down to the minimum becomes steeper and the non-vanishing vacuum expectation value $\langle \phi \rangle \neq 0$ appears for $\omega_1 > 0$. Since a vacuum expectation value should be less than M_{Pl} , we find that the deformation parameter $\omega_1 < 2 \left(1 - e^{-\sqrt{2/3}}\right) e^3 \sqrt{2/3} \approx 12.9$.

The corresponding $F(R)$ gravity functions are shown on the right-hand-side of Fig. 7. Compared to the Starobinsky curve ($\omega_1 = 0$), they have larger values in the given range of R_J/m^2 . For larger values of R_J/m^2 , all those F -curves converge to the Starobinsky curve.




Figure 7: The scalar potential of the canonical inflaton field ϕ (left) and the related $F(R)$ -function (right) in the case of the deformation of the Starobinsky model for some values of the parameter ω_1 : 0, 1, 5, and 10.

The results of our numerical calculation for the index n_s of scalar perturbations and the tensor-to-scalar ratio r , as the functions of e-folds N_e , are given in Fig. 8 for the values of the parameter ω_1 equal to 0, 1, 5, and 10. As is clear from Fig. 8, there are (small) changes in the duration of inflation and its initial and final moments.

From Eq. (5.3) we get a simple formula for the second derivative $F_{,\sigma\sigma}$ as

$$F_{,\sigma\sigma}(y) = \frac{M_{Pl}^2}{6m^2 (1 + \omega_1 y^3)} \quad , \quad (5.15)$$

so that the $F(R)$ gravity model (5.13) satisfies the conditions (1.5) with $\omega_1 \geq 0$.

5.2 New case II

Our approach suggests another one-parametric deformation of the Starobinsky potential (1.3) as follows:

$$V(\phi) = V_0 \left[1 - e^{-\sqrt{2/3}\phi/M_{Pl}} - \zeta e^{-2\sqrt{2/3}\phi/M_{Pl}} \right]^2 = V_0 (1 - y - \zeta y^2)^2 \quad , \quad (5.16)$$

where we assume the parameter $\zeta \geq 0$. This potential can be realized in supergravity [47], while the potential (1.3) is recovered at $\zeta = 0$. The Minkowski minimum of the potential (5.16) is realized by demanding $V(y) = V'(y) = 0$ that yields a quadratic equation whose solution is given by

$$y_{\min.} = \frac{\sqrt{1 + 4\zeta} - 1}{2\zeta} \quad . \quad (5.17)$$




Figure 8: The index n_s (left) of scalar perturbations and the tensor-to-scalar ratio r (right) as the functions of e-folds N_e in the case I of the deformation of the Starobinsky model for some values of the parameter ω_1 : 0, 1, 5, and 10.

The potential (5.16) can also be rewritten to the form (5.2) with the $\omega(y)$ -function

$$\omega = 1 - 2\zeta + 2\zeta y + \zeta^2 y^2 . \quad (5.18)$$

Accordingly, Eq. (5.4) takes the form

$$\zeta^2 y^4 + \zeta y^3 + \left(1 + \frac{R_J}{3m^2}\right) y - 1 \equiv ay^4 + by^3 + cy^2 + dy + e = 0 \quad (5.19)$$

with

$$a = \zeta^2, \quad b = \zeta, \quad c = 0, \quad d = 1 + \frac{R_J}{3m^2}, \quad e = -1 . \quad (5.20)$$

The discriminant of the quartic equation (5.19),

$$\begin{aligned} \Delta = & -256\zeta^6 - 192\zeta^5 \left(1 + \frac{R_J}{3m^2}\right) - 3\zeta^4 \left[9 \left(1 + \frac{R_J}{3m^2}\right)^4 - 2 \left(1 + \frac{R_J}{3m^2}\right)^2 + 9\right] \\ & - 4\zeta^3 \left(1 + \frac{R_J}{3m^2}\right)^3 , \end{aligned} \quad (5.21)$$

is negative, so that there are two distinct real roots and two conjugated complex roots. The equivalent depressed quartic equation

$$\tilde{y}^4 + p\tilde{y}^2 + q\tilde{y} + r = 0, \quad \text{where} \quad \tilde{y} = y + \frac{b}{4a} = y + \frac{\zeta^{-1}}{4} , \quad (5.22)$$

has the following parameters:

$$p = -\frac{3b^2}{8a^2} = -\frac{3}{8\zeta^2} < 0, \quad q = \frac{b^3 + 8a^2d}{8a^3} = \frac{1 + 8\zeta \left(1 + \frac{R_J}{3m^2}\right)}{8\zeta^3} > 0, \quad (5.23)$$

and

$$r = -\frac{3 + 256\zeta^2 + 64\zeta \left(1 + \frac{R_J}{3m^2}\right)}{256\zeta^4} < 0 . \quad (5.24)$$

The Ferrari formula gives the roots of the depressed quartic equation in terms of the quantities

$$\begin{aligned}\Delta_0 &= -3bd + 12ae = -3\zeta \left(1 + \frac{R_J}{3m^2}\right) - 12\zeta^2 < 0, \\ \Delta_1 &= 27b^2e + 27ad^2 = 27\zeta^2 \left[\left(1 + \frac{R_J}{3m^2}\right)^2 - 1 \right] > 0, \\ Q &= \left(\frac{\Delta_1 + \sqrt{\Delta_1^2 - 4\Delta_0^3}}{2} \right)^{1/3}, \quad S = \frac{1}{2} \sqrt{-\frac{2}{3}p + \frac{1}{3a} \left(Q + \frac{\Delta_0}{Q}\right)},\end{aligned}\tag{5.25}$$

where in our case $Q > 0$, $S > 0$ and $27\Delta = 4\Delta_0^3 - \Delta_1^2 < 0$. The real roots are given by

$$y_{1,2} = -\frac{b}{4a} - S \pm \frac{1}{2} \sqrt{-4S^2 - 2p + \frac{q}{S}} = -\frac{1}{4\zeta} - S \pm \frac{1}{2} \sqrt{-4S^2 + \frac{3}{4\zeta^2} + \frac{1 + 8\zeta \left(1 + \frac{R_J}{3m^2}\right)}{8S\zeta^3}}.\tag{5.26}$$

In the limit $\zeta \rightarrow 0_+$ we find

$$p \rightarrow -\frac{3}{8\zeta^2}, \quad q \rightarrow \frac{1}{8\zeta^3}, \quad \Delta_1 \rightarrow 0, \quad \Delta_0 \rightarrow -3\zeta \left(1 + \frac{R_J}{3m^2}\right), \quad Q \rightarrow \sqrt{|\Delta_0|}, \quad S \rightarrow \frac{1}{4\zeta}.$$

Actually, only the root with the *upper sign* choice in Eq. (5.26) leads to the physical solution connected to the Starobinsky model because the lower sign choice implies the negative value of y and, therefore, should be discarded. The scalar potential and the F function are given in Fig. 9 for some values of the deformation parameter ζ . Since a vacuum expectation value should be less than M_{Pl} , we find a restriction on the deformation parameter,

$$\zeta < \left(1 - e^{-\sqrt{2/3}}\right) e^{2\sqrt{2/3}} \approx 2.9.\tag{5.27}$$




Figure 9: The scalar potential of the canonical inflaton field ϕ , and the related $F(R)$ function in case II.

The results of our numerical calculation for the index n_s of scalar perturbations and the tensor-to-scalar ratio r , as the functions of e-folds N_e , in case II are given in Fig. 10 for the values of the parameter ζ equal to 0, 1/2, 1, and 2. There are (small) changes in the duration of inflation and its initial and final moments, similarly to case I.




Figure 10: The index n_s (left) of scalar perturbations and the tensor-to-scalar ratio r (right) as the functions of e-folds N_e in case II for some values of the parameter ζ : 0, 1/2, 1, and 2.

From Eq. (2.12), we find

$$F_{,\sigma\sigma}(y) = \frac{M_{Pl}^2}{6m^2 (1 + 2\zeta y^3 + 3\zeta^2 y^4)} > 0 \quad .$$

To the end of this Section, we comment on case II with a negative parameter $\zeta = -|\zeta|$. The critical points of the potential (5.16) correspond to three solutions,

$$y_{1,2} = \frac{1 \mp \sqrt{1 - 4|\zeta|}}{2|\zeta|} \quad , \quad y_3 = \frac{1}{2|\zeta|} \quad . \quad (5.28)$$

To get slow-roll inflation, we need $|\zeta| < 1/4$. It leads to the scalar potential with two minima and one maximum, while only one minimum has $y < 1$, thus leading to hilltop inflation with restricted initial conditions.

6 Conclusion

In this paper, we studied several extensions of the Starobinsky inflation model of the $(R + R^2)$ gravity in the context of $F(R)$ gravity and scalar-tensor gravity, including the known models (Sec. 3) and the new ones (Secs. 4 and 5) by using our methods. These models are consistent with cosmological observations and have no ghosts, while *both* the F -function and the inflaton (scalaron) potential $V(\phi)$ are available in the explicit analytic form including their dependence upon the parameters. The last feature appears to be very restrictive. We focused on the models with only one free parameter for simplicity. Of course, the list of such models can be extended but not much. Besides the physical and observational constraints, on the one side, it is must be possible to invert the dependence of a polynomial in R upon the inflaton field ϕ as an analytic function $R(\phi)$, see Sec. 3. On the other side, it must be possible to analytically invert a function $R(y)$, see Sec. 5. It is only possible when there is an analytic formula for the roots of the polynomials, which restricts their order to 4 or less.

In Secs. 3 and 4 we deformed the Starobinsky $(R + R^2)$ gravity model of inflation and found the inflaton potential in the analytic form in the three specific cases, by adding an R^3 term, an R^4 -term and an $R^{3/2}$ term, respectively. In Sec. 5 we started from the opposite side,

by deforming the scalar potential of the Starobinsky model, and derived the corresponding F -function in the analytic form in the two different models with a single parameter, and found the lower and upper bounds on the values of the parameters. The new models in Sec. 5 are very close to the original Starobinsky model of inflation, as regards their cosmological parameters. However, unlike the Starobinsky model, the inflaton (scalaron) acquires a non-vanishing vacuum expectation value in those models.

The asymptotic de Sitter solution in the pure $(R+R^2)$ model at very large positive values of R turns out to be unstable against any correction of the higher order in R in the F -function that drastically changes the behavior of the scalar potential before inflation. Even though those values of R are apparently beyond the scope of applicability of the modified $F(R)$ -gravity as the effective theory of gravity, it introduces a dependence upon initial values of the inflaton field that must roll down to the left from the maximum of the potential (see Figs. 7 and 9) for viable hilltop inflation. In other words, the higher-order (in R) quantum corrections imply tuning the initial conditions for inflation.⁴ This feature is apparently universal because there is no fundamental reason for the absence of quantum corrections proportional to the higher powers of R . This observation is, however, limited to the perturbative treatment with respect to the R -dependence of F -function.

Consistency with CMB observations in our models demands the higher order terms in R to be negligible against the R^2 term during inflation. We achieved it via demanding the dimensionless coefficients in front of the higher-order terms to be small enough because the variable $\tilde{\sigma} = R/m^2$ is not small during inflation. We found the upper limits on the values of the coefficients at the R^3 and R^4 terms. However, we are not aware of any fundamental reason demanding those coefficients to be small.⁵

Unlike the R^3 and R^4 terms, the modification of the Starobinsky model by the $R^{3/2}$ term does not lead to significant constraints on its coefficient in slow-roll inflation, at least for $0 < \delta < 100$. However, the $R^{3/2}$ term has a significant impact on the value of the tensor-to-scalar ratio r (see Table 1 in Sec. 4)⁶ and, therefore, implies higher production rate of primordial gravitational waves caused by inflation. There is another source of primordial gravitational waves caused by possible formation of primordial black holes; in the framework of $F(R)$ gravity it was studied in Ref. [50].

More restrictions of the F -function and the inflaton scalar potential arise when demanding their minimal embedding into supergravity, along the lines of Refs. [45, 51, 52]. For instance, the R^3 term is excluded in supergravity, whereas only case II in Subsec. 5.2 is extendable in the minimal supergravity framework that requires the scalar potential to be a real function squared. The $R^{3/2}$ term also arises in certain versions of the chiral $F(\mathcal{R})$ supergravity [45].

Acknowledgments

The authors are grateful to Gia Dvali, Olaf Lechtenfeld, Kei-ichi Maeda and Alexei Starobinsky for discussions and correspondence.

V.R.I. was supported by the Theoretical Physics and Mathematics Advancement Foundation “BASIS”. S.V.K. was supported by Tokyo Metropolitan University, the World Premier

⁴The similar results were obtained in the $(R + R^2 + R^3)$ model [19].

⁵It might be possible to get a flat potential for very large inflaton field values via a resummation of all higher-curvature corrections, e.g., in the framework of asymptotically-safe quantum gravity [48, 49].

⁶It also has a significant impact on reheating after inflation [12].

International Research Center Initiative (MEXT) and Yamada Foundation in Japan. S.V.K. is grateful to Leibniz Universität Hannover in Germany for pleasant hospitality during this investigation. E.O.P. and S.Yu.V. were partially supported by the Russian Foundation for Basic Research grant No. 20-02-00411.

References

- [1] B. Whitt, *Fourth Order Gravity as General Relativity Plus Matter*, *Phys. Lett. B* **145** (1984) 176.
- [2] K.-i. Maeda, *Inflation as a Transient Attractor in R^{**2} Cosmology*, *Phys. Rev. D* **37** (1988) 858.
- [3] K.-i. Maeda, *Towards the Einstein-Hilbert Action via Conformal Transformation*, *Phys. Rev. D* **39** (1989) 3159.
- [4] J.D. Barrow and S. Cotsakis, *Inflation and the Conformal Structure of Higher Order Gravity Theories*, *Phys. Lett. B* **214** (1988) 515.
- [5] D.C. Rodrigues, F. de O. Salles, I.L. Shapiro and A.A. Starobinsky, *Auxiliary fields representation for modified gravity models*, *Phys. Rev. D* **83** (2011) 084028 [[1101.5028](#)].
- [6] T.P. Sotiriou and V. Faraoni, *$f(R)$ Theories Of Gravity*, *Rev. Mod. Phys.* **82** (2010) 451 [[0805.1726](#)].
- [7] A. De Felice and S. Tsujikawa, *$f(R)$ theories*, *Living Rev. Rel.* **13** (2010) 3 [[1002.4928](#)].
- [8] S. Capozziello and M. De Laurentis, *Extended Theories of Gravity*, *Phys. Rept.* **509** (2011) 167 [[1108.6266](#)].
- [9] S.V. Ketov, *Supergravity and Early Universe: the Meeting Point of Cosmology and High-Energy Physics*, *Int. J. Mod. Phys. A* **28** (2013) 1330021 [[1201.2239](#)].
- [10] A.A. Starobinsky, *A new type of isotropic cosmological models without singularity*, *Phys. Lett. B* **91** (1980) 99 .
- [11] T. Saidov and A. Zhuk, *Bouncing inflation in nonlinear $R^2 + R^4$ gravitational model*, *Phys. Rev. D* **81** (2010) 124002 [[1002.4138](#)].
- [12] S.V. Ketov and S. Tsujikawa, *Consistency of inflation and preheating in $F(R)$ supergravity*, *Phys. Rev. D* **86** (2012) 023529 [[1205.2918](#)].
- [13] L. Sebastiani, G. Cognola, R. Myrzakulov, S.D. Odintsov and S. Zerbini, *Nearly Starobinsky inflation from modified gravity*, *Phys. Rev. D* **89** (2014) 023518 [[1311.0744](#)].
- [14] H. Motohashi, *Consistency relation for R^p inflation*, *Phys. Rev. D* **91** (2015) 064016 [[1411.2972](#)].
- [15] K. Bamba and S.D. Odintsov, *Inflationary cosmology in modified gravity theories*, *Symmetry* **7** (2015) 220 [[1503.00442](#)].
- [16] T. Miranda, J.C. Fabris and O.F. Piattella, *Reconstructing a $f(R)$ theory from the α -Attractors*, *JCAP* **09** (2017) 041 [[1707.06457](#)].
- [17] H. Motohashi and A.A. Starobinsky, *$f(R)$ constant-roll inflation*, *Eur. Phys. J. C* **77** (2017) 538 [[1704.08188](#)].
- [18] D.Y. Cheong, H.M. Lee and S.C. Park, *Beyond the Starobinsky model for inflation*, *Phys. Lett. B* **805** (2020) 135453 [[2002.07981](#)].
- [19] G. Rodrigues-da Silva, J. Bezerra-Sobrinho and L.G. Medeiros, *A higher-order extension of Starobinsky inflation: initial conditions, slow-roll regime and reheating phase*, [2110.15502](#).

- [20] PLANCK collaboration, *Planck 2018 results. X. Constraints on inflation*, *Astron. Astrophys.* **641** (2020) A10 [[1807.06211](#)].
- [21] A.A. Starobinsky, *Disappearing cosmological constant in $f(R)$ gravity*, *JETP Lett.* **86** (2007) 157 [[0706.2041](#)].
- [22] S.A. Appleby, R.A. Battye and A.A. Starobinsky, *Curing singularities in cosmological evolution of $F(R)$ gravity*, *JCAP* **06** (2010) 005 [[0909.1737](#)].
- [23] V.R. Ivanov and S.Y. Vernov, *Integrable modified gravity cosmological models with an additional scalar field*, *Eur. Phys. J. C* **81** (2021) 985 [[2108.10276](#)].
- [24] M.F. Figueiro and A. Saa, *Anisotropic singularities in modified gravity models*, *Phys. Rev. D* **80** (2009) 063504 [[0906.2588](#)].
- [25] G. Dvali, A. Kehagias and A. Riotto, *Inflation and Decoupling*, [2005.05146](#).
- [26] S.V. Ketov and N. Watanabe, *On the Higgs-like Quintessence for Dark Energy*, *Mod. Phys. Lett. A* **29** (2014) 1450117 [[1401.7756](#)].
- [27] S.Y. Vernov, V.R. Ivanov and E.O. Pozdeeva, *Superpotential Method for $F(R)$ Cosmological Models*, *Phys. Part. Nucl.* **51** (2020) 744 [[1912.07049](#)].
- [28] J. Garcia-Bellido and E. Ruiz Morales, *Primordial black holes from single field models of inflation*, *Phys. Dark Univ.* **18** (2017) 47 [[1702.03901](#)].
- [29] BICEP/KECK collaboration, *Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season*, *Phys. Rev. Lett.* **127** (2021) 151301 [[2110.00483](#)].
- [30] D.H. Lyth and A. Riotto, *Particle physics models of inflation and the cosmological density perturbation*, *Phys. Rept.* **314** (1999) 1 [[hep-ph/9807278](#)].
- [31] R. Kallosh, A. Linde and D. Roest, *Superconformal Inflationary α -Attractors*, *JHEP* **11** (2013) 198 [[1311.0472](#)].
- [32] B.J. Broy, M. Galante, D. Roest and A. Westphal, *Pole inflation — Shift symmetry and universal corrections*, *JHEP* **12** (2015) 149 [[1507.02277](#)].
- [33] T. Terada, *Generalized Pole Inflation: Hilltop, Natural, and Chaotic Inflationary Attractors*, *Phys. Lett. B* **760** (2016) 674 [[1602.07867](#)].
- [34] C. Pallis, *Pole-induced Higgs inflation with hyperbolic Kähler geometries*, *JCAP* **05** (2021) 043 [[2103.05534](#)].
- [35] A.R. Liddle, P. Parsons and J.D. Barrow, *Formalizing the slow roll approximation in inflation*, *Phys. Rev. D* **50** (1994) 7222 [[astro-ph/9408015](#)].
- [36] E.O. Pozdeeva, M. Sami, A.V. Toporensky and S.Y. Vernov, *Stability analysis of de Sitter solutions in models with the Gauss-Bonnet term*, *Phys. Rev. D* **100** (2019) 083527 [[1905.05085](#)].
- [37] E.O. Pozdeeva, M.R. Gangopadhyay, M. Sami, A.V. Toporensky and S.Y. Vernov, *Inflation with a quartic potential in the framework of Einstein-Gauss-Bonnet gravity*, *Phys. Rev. D* **102** (2020) 043525 [[2006.08027](#)].
- [38] E.O. Pozdeeva and S.Y. Vernov, *Construction of inflationary scenarios with the Gauss-Bonnet term and nonminimal coupling*, *Eur. Phys. J. C* **81** (2021) 633 [[2104.04995](#)].
- [39] A.A. Starobinsky, *The Perturbation Spectrum Evolving from a Nonsingular Initially De-Sitter Cosmology and the Microwave Background Anisotropy*, *Sov. Astron. Lett.* **9** (1983) 302.
- [40] J.-c. Hwang and H. Noh, *Cosmological perturbations in generalized gravity theories*, *Phys. Rev. D* **54** (1996) 1460.

- [41] A.L. Berkin and K.-i. Maeda, *Effects of R^{**3} and R box R terms on R^{**2} inflation*, *Phys. Lett. B* **245** (1990) 348.
- [42] Q.-G. Huang, *A polynomial $f(R)$ inflation model*, *JCAP* **02** (2014) 035 [[1309.3514](#)].
- [43] S.D. Odintsov and V.K. Oikonomou, *Unimodular Mimetic $F(R)$ Inflation*, *Astrophys. Space Sci.* **361** (2016) 236 [[1602.05645](#)].
- [44] J.S. Martins, O.F. Piattella, I.L. Shapiro and A.A. Starobinsky, *Inflation with sterile scalar coupled to massive fermions and to gravity*, [2010.14639](#).
- [45] S.V. Ketov and A.A. Starobinsky, *Embedding $(R + R^2)$ -Inflation into Supergravity*, *Phys. Rev. D* **83** (2011) 063512 [[1011.0240](#)].
- [46] H. Nakada and S.V. Ketov, *Inflation from higher dimensions*, *Phys. Rev. D* **96** (2017) 123530 [[1710.02259](#)].
- [47] S.V. Ketov, *On the equivalence of Starobinsky and Higgs inflationary models in gravity and supergravity*, *J. Phys. A* **53** (2020) 084001 [[1911.01008](#)].
- [48] L.-H. Liu, T. Prokopec and A.A. Starobinsky, *Inflation in an effective gravitational model and asymptotic safety*, *Phys. Rev. D* **98** (2018) 043505 [[1806.05407](#)].
- [49] J. Chojnacki, J. Krajecka, J.H. Kwapisz, O. Slowik and A. Strag, *Is asymptotically safe inflation eternal?*, *JCAP* **04** (2021) 076 [[2101.00866](#)].
- [50] T. Papanikolaou, C. Tzerefos, S. Basilakos and E.N. Saridakis, *Scalar induced gravitational waves from primordial black hole Poisson fluctuations in Starobinsky inflation*, [2112.15059](#).
- [51] F. Farakos, A. Kehagias and A. Riotto, *On the Starobinsky Model of Inflation from Supergravity*, *Nucl. Phys. B* **876** (2013) 187 [[1307.1137](#)].
- [52] S. Ferrara, R. Kallosh, A. Linde and M. Porrati, *Minimal Supergravity Models of Inflation*, *Phys. Rev. D* **88** (2013) 085038 [[1307.7696](#)].