Nonlinear electromagnetic fields in strictly stationary spacetimes

A. Bokulić* and I. Smolić[†] Department of Physics, Faculty of Science, University of Zagreb, 10000 Zagreb, Croatia

T. Jurić[‡] Rudjer Bošković Institute, Bijenička cesta 54, HR-10002 Zagreb, Croatia

We prove two theorems which imply that any nonlinear electromagnetic field obeying a dominant energy condition in a strictly stationary, everywhere regular, asymptotically flat spacetime, must be either trivial or a stealth field. First theorem holds in static spacetimes and is independent of gravitational part of the action, as long as the coupling of electromagnetic field to the gravitational field is minimal. Second theorem assumes Einstein–Hilbert gravitational action and relies on the positive energy theorem, but does not assume that the spacetime metric is static. In addition, we discuss possible generalizations of these results, to theories with charged matter, as well as higher dimensional nonlinear electromagnetic fields.

I. INTRODUCTION

Interaction of the gravitational and the electromagnetic field, governed by the gravitational-gauge field equations, is highly nonlinear. It is quite optimistic to hope that we might reach a complete classification of all solutions, even under the constraints of some regularity and boundary conditions. Indeed, a slightly less ambitious goal, understanding of time-independent solutions, is still a formidable task, but one worth taking as stationary solutions serve as models of the equilibrium field configurations.

For example, an important class of the *stationary* black hole spacetimes is heavily narrowed by the series of black hole uniqueness and no-hair theorems [1, 2], distilled and polished over the past several decades. These solutions, however, are not *strictly* stationary, as the Killing vector field corresponding to stationary isometry, timelike on some domain of the black hole exterior, may change its causal character in the black hole interior and ergoregions surrounding rotating black holes. Also, black holes may harbor a singularity, in which case they are not globally regular spacetimes.

This begs a question whether it is possible to have a strictly stationary, everywhere regular, asymptotically flat solution with nonvanishing electromagnetic field. Such spacetime would represent an instance of Wheeler's gravitational-electromagnetic geon [3], at least up to a nontrivial question of stability. Negative answer in the case of Einstein–Maxwell theory is a canonical, well-known result, sometimes referred to as the absence of self-gravitating electromagnetic solitons [1]. Setting aside

a delicate historical question of primacy, basic strategy of proofs can be traced back to the seminal work of Lichnerowicz [4]: construct a convenient nonnegative quantity, whose integral over the spacetime domain in problem is nonpositive, implying that this quantity has to be identically zero. This was masterfully utilized in foundational uniqueness theorems obtained by Carter [5] (cf. republished, corrected paper [6]) and, more recently, by Heusler [2, 7]. Several generalizations of the "no-soliton theorem" for the Einstein–Maxwell theory in the presence of various scalar fields was obtained by Shiromizu, Ohashi and Suzuki [8], and Herdeiro and Oliveira [9, 10].

One step further is to ask what happens in the theories where classical Maxwell's electrodynamics is replaced by its nonlinear modifications. Nonlinear electrodynamics (NLE) has its roots at the dawn of quantum field theory, back in the 1930s, sprouting over the following decades with innumerable NLE Lagrangians. Born-Infeld theory [11, 12] was constructed with specific aim to cure the inconsistencies of the Maxwell's electrodynamics associated with the infinite self-energy of the point charges and, remarkably, reappeared much later in low energy limits of the string theory [13]. Another prominent NLE theory is defined by the Euler-Heisenberg's one-loop QED correction to Maxwell's Lagrangian [14, 15]. Novel Mod-Max electrodynamics [16, 17] is a one-parameter class of NLE theories which is both conformally invariant and invariant with respect to electromagnetic duality rotations [18]. Nonlinearities in the electromagnetic interaction are being tested by the ATLAS Collaboration [19–22] and new generations of the ultraintense lasers at the Extreme Light Infrastructure [23].

Intriguing feature of NLE theories is that they may admit a resolution of the black hole singularities, up to delicate constraints [24–27]. An example of regular black hole spacetime, originally proposed *ad hoc* by Bardeen [28], was later interpreted by Ayón-Beato and García

^{*} abokulic@phy.hr

[†] ismolic@phy.hr

[‡] tjuric@irb.hr

[29, 30] as a solution of Einstein-NLE Maxwell field equations for a particular NLE theory. Nevertheless, these examples still leave the original question open, whether a self-gravitating electromagnetic field may settle in a nontrivial, regular configuration which is not a black hole? First extension of the "no-soliton" theorem (referred to by the authors as the "Lichnerowicz-type theorem") for theories with NLE was given in [31] for the truncated Born–Infeld theory and the power-Maxwell theory. Our aim is to provide much broader generalization of this result, for NLE Lagrangians which are general smooth functions of both electromagnetic invariants, $F_{ab}F^{ab}$ and $F_{ab} \star F^{ab}$. Also, we shall present first steps in generalization of these results for theories with charged matter or theories in different number of spacetime dimensions.

The paper is organized as follows. In Sec. II we briefly overview fundamentals of gravitational theories with nonlinear electromagnetic fields. Main results of the paper, theorems 1 and 2, are stated in Sec. III and their proofs presented in Sec. IV. We discuss various generalizations of these theorems in Sec. V and review remaining open questions in Sec. VI. Several basic identities from differential geometry are stated in the Appendix.

Conventions and notation. Interior of a set S is denoted by S° , boundary of S by ∂S and closure of S by \overline{S} . Difference between sets A and B is denoted by A-B. We shall use the "mostly plus" metric signature and natural system of units with $G=c=4\pi\epsilon_0=1$. Differential forms are denoted by bolded indexless letters, abstract index notation or combination of both. Volume 4-form is denoted by ϵ . Contraction of a symmetric tensor S_{ab} with vector X^a is a 1-form denoted by $\mathbf{S}(X)$. Following the reference [1], we write $f=O(r^{-k})$ when f is of order $O(r^{-k})$ as $r\to\infty$ and $f=O_{\infty}(r^{-k})$ when $\partial_{i_1}\ldots\partial_{i_\ell}f=O(r^{-k-\ell})$ for arbitrary set of coordinate indices $\{i_1,\ldots,i_\ell\}$.

II. BRIEF OVERVIEW OF NLE

Let us, before stating the central theorems of the paper, briefly introduce the nonlinear electrodynamics. Ubiquitous elements are two electromagnetic invariants

$$\mathcal{F} := F_{ab}F^{ab} \quad \text{and} \quad \mathcal{G} := F_{ab} \star F^{ab} .$$
 (1)

We follow the nomenclature from [32], by sorting NLE theories into the \mathcal{F} -class, with a Lagrangian density \mathcal{L} depending only on invariant \mathcal{F} , and the $\mathcal{F}\mathcal{G}$ -class, with Lagrangian density \mathcal{L} depending on both invariants. In this paper the main focus is on the broader, $\mathcal{F}\mathcal{G}$ -class of NLE theories, with NLE Lagrangian density \mathcal{L} which is a C^2 function on some neighbourhood of the origin of the \mathcal{F} - \mathcal{G} plane. We can always choose Lagrangian density, by adding an appropriate constant, such that $\mathcal{L}(0,0) = 0$. Partial derivatives of the Lagrangian density \mathcal{L} are denoted by abbreviations such as $\mathcal{L}_{\mathcal{F}} := \partial_{\mathcal{F}} \mathcal{L}$, $\mathcal{L}_{\mathcal{G}} := \partial_{\mathcal{G}} \mathcal{L}$, and so on. We say that a NLE

Lagrangian density \mathscr{L} obeys the Maxwell's weak field limit if $\mathscr{L}_{\mathscr{F}}(0,0) = -1/4$ and $\mathscr{L}_{\mathscr{G}}(0,0) = 0$.

The Lagrangian 4-form, defined with some (diffeomorphism covariant) gravitational Lagrangian density $\mathcal{L}^{(g)}$, is

$$\mathbf{L} = \frac{1}{16\pi} \left(\mathcal{L}^{(g)} + 4\mathcal{L} \right) \boldsymbol{\epsilon} . \tag{2}$$

Corresponding gravitational field equation is of form

$$E_{ab} = 8\pi T_{ab} , \qquad (3)$$

where the gravitational tensor E_{ab} is divergence free, $\nabla^a E_{ab} = 0$, and the NLE energy-momentum tensor may be conveniently written as

$$T_{ab} = -4\mathcal{L}_{\mathcal{F}} T_{ab}^{(\text{Max})} + \frac{1}{4} T g_{ab} \tag{4}$$

with Maxwell's electromagnetic energy-momentum tensor

$$T_{ab}^{(\text{Max})} := \frac{1}{4\pi} \left(F_{ac} F_b^{\ c} - \frac{1}{4} g_{ab} \mathcal{F} \right)$$
 (5)

and the trace

$$T := g^{ab} T_{ab} = \frac{1}{\pi} \left(\mathscr{L} - \mathscr{L}_{\mathfrak{F}} \mathfrak{F} - \mathscr{L}_{\mathfrak{G}} \mathfrak{G} \right) . \tag{6}$$

For Einstein-Hilbert Lagrangian density $\mathcal{L}^{(g)} = R$ we have $E_{ab} = G_{ab}$, the Einstein tensor.

Using an auxiliary 2-form

$$\mathbf{Z} := -4\left(\mathcal{L}_{\mathcal{F}}\mathbf{F} + \mathcal{L}_{\mathcal{G}}\star\mathbf{F}\right) , \qquad (7)$$

the NLE Maxwell's equations may be written in a form

$$d\mathbf{F} = 0 , \quad d \star \mathbf{Z} = 0 . \tag{8}$$

We shall invoke two energy conditions, null energy condition (NEC) and dominant energy condition (DEC). It can be shown [32, 33] that NEC holds if and only if $\mathcal{L}_{\mathcal{F}} \leq 0$, while DEC holds if and only if $\mathcal{L}_{\mathcal{F}} \leq 0$ and $T \leq 0$.

III. TWO THEOREMS

We shall present two "no-soliton" theorems, each of which has its strengths and limitations. Both theorems assume that the spacetime is strictly stationary, so that we do not consider spacetimes with either black hole or cosmological horizons. First result rests upon a stronger assumption, that spacetime is static, which admits a simpler proof that does not depend on details of gravitational Lagrangian of the theory, as long as the coupling of the electromagnetic field to gravitation is minimal. Second result does not rely on this assumption, so that it may be applied to "rotating" solutions. However, this comes at a price: proof rests upon a highly nontrivial, celebrated positive energy theorem [34–40] and, correspondingly, works only in those gravitational theories for which this theorem has been proven.

Let us, before the statement of main results, list technical assumptions necessary for the theorems.

- (1) Spacetime consists of a 4-dimensional smooth, simply connected manifold \mathcal{M} , with smooth Lorentzian metric g_{ab} and a smooth electromagnetic 2-form F_{ab} , which are solutions of the gravitational-NLE fields equations (3) and (8), with NLE Lagrangian density \mathcal{L} obeying the Maxwell's weak field limit.
- (2) Spacetime admits a strictly timelike Killing vector field k^a (namely, $k^a k_a < 0$ on whole \mathscr{M}) and the electromagnetic field inherits this symmetry, $\mathscr{L}_k F_{ab} = 0$ [41–43].
- (3) Through each point $p \in \mathcal{M}$ passes at least one complete oriented spacelike hypersurface Σ with induced metric h_{ij} and associated second fundamental form (extrinsic curvature) K_{ij} , Euclidean at infinity¹ and asymptotically flat in the sense [1, 39, 40] that on each of its "ends" the following fall-off conditions, written in Cartesian coordinates, are met: $1 + k^{\alpha}k_{\alpha} = O_{\infty}(r^{-1})$, $k^{\alpha}g_{\alpha i} = O_{\infty}(r^{-1})$, $\gamma_{ij} = O_{\infty}(r^{-1})$ and $K_{ij} = O_{\infty}(r^{-2})$, while the electromagnetic 2-form F_{ab} satisfies $k^{\alpha}F_{\alpha i} = O_{\infty}(r^{-2})$ and $k^{\alpha} \star F_{\alpha i} = O_{\infty}(r^{-2})$, and the associated potentials (defined below) are of order $O_{\infty}(r^{-1})$.

We shall refer to the assumptions (1)–(3) as the basic assumptions. It is quite possible that some of the assumptions above may be slightly relaxed without any significant effect on the further conclusions, but we shall not pursue such nuances here. Furthermore, we introduce the following notion.

Definition. We say that an electromagnetic field is *stealth* at a point $p \in \mathcal{M}$ if at that point the corresponding energy-momentum tensor T_{ab} is zero, but the electromagnetic 2-form F_{ab} is non-zero.

In other words, stealth fields do not affect the spacetime metric, as their contribution to the energy-momentum tensor is vanishing. Such configurations are absent in the classical, Maxwell's electrodynamics, but appear in NLE theories if and only if $F_{ab} \neq 0$, $\mathcal{L}_{\mathcal{F}} = 0$ and T = 0 hold at a given point [44]. Note that in this paper, for clarity, we keep trivial fields, $F_{ab} = 0$, apart from the stealth fields. Prominent class of stealth solution examples [44] may be found among the null electromagnetic fields in power-Maxwell theory [45, 46], which also belong to a family of so-called universal electromagnetic fields [47–49].

The two central results of the paper are as follows.

Theorem 1. Suppose that spacetime with electromagnetic field satisfies basic assumptions, with electromagnetic energy-momentum tensor obeying the null energy

condition, and Killing vector field k^a is hypersurface orthogonal. Then the electromagnetic field is at each point of the spacetime either trivial, $F_{ab} = 0$, or stealth.

Theorem 2. Suppose that spacetime with electromagnetic field satisfies basic assumptions and the gravitational part of the action is the Einstein-Hilbert's with the electromagnetic energy-momentum tensor obeying the dominant energy condition. Then the spacetime is isometric to the Minkowski spacetime (\mathbb{R}^4 , η_{ab}) and the electromagnetic field is at each point of the spacetime either trivial, $F_{ab} = 0$, or stealth.

We stress that the theorem 1 relies on a weaker, null energy condition.

IV. PROOFS OF THEOREMS

In both theorems we are looking at a spacetime admitting a strictly timelike Killing vector field k^a . It is convenient to introduce function $V := -k_a k^a > 0$ and the associated twist 1-form

$$\boldsymbol{\omega} := -\star (\mathbf{k} \wedge d\mathbf{k}) \ . \tag{9}$$

One should, however, beware of variations in definition of the twist 1-form throughout the literature (e.g. Heusler [2] introduces twist 1-form $\widetilde{\omega}$, such that $\omega = -2\widetilde{\omega}$). Our choice is mainly motivated by the fact that, in the abstract index notation, it corresponds simply to $\omega_a = \epsilon_a^{bcd} k_b \nabla_c k_d$, without additional factors.

Vector field k^a allows us to define two electric 1-forms,

$$\mathbf{E} := -i_k \mathbf{F} , \quad \mathbf{D} := -i_k \mathbf{Z} , \tag{10}$$

and two magnetic 1-forms,

$$\mathbf{B} := i_k \star \mathbf{F} , \quad \mathbf{H} := i_k \star \mathbf{Z} . \tag{11}$$

As a consequence of the symmetry inheritance and generalized Maxwell's equations we know that ${\bf E}$ and ${\bf H}$ are closed forms,

$$d\mathbf{E} = -di_k \mathbf{F} = (-\pounds_k + i_k d) \mathbf{F} = 0 , \qquad (12)$$

$$d\mathbf{H} = di_k \star \mathbf{Z} = (\pounds_k - i_k d) \star \mathbf{Z} = 0.$$
 (13)

As manifold \mathcal{M} is, by assumption, simply connected, we can globally define scalar potentials, electric Φ and magnetic Ψ , such that $\mathbf{E} = -\mathrm{d}\Phi$ and $\mathbf{H} = -\mathrm{d}\Psi$. Also, directly from the definition we know that $\pounds_k \Phi = -i_k \mathbf{E} = 0$ and $\pounds_k \Psi = -i_k \mathbf{H} = 0$.

Backbone of proofs are divergence identities which have to be carefully chosen. First, using

$$d\left(\frac{\mathbf{k}}{V}\right) = \frac{1}{V^2} \left(V d\mathbf{k} - dV \wedge \mathbf{k}\right) = \frac{1}{V^2} \star (\boldsymbol{\omega} \wedge \mathbf{k}) \qquad (14)$$

and

$$- \star i_k \mathbf{Z} = \star i_k \star \star \mathbf{Z} = \mathbf{k} \wedge \star \mathbf{Z} \tag{15}$$

¹ We say that a n-smooth manifold S is Euclidean at infinity if there is a compact set $\mathcal{C} \subseteq S$, such that $S - \mathcal{C}$ is a disjoint union of a finite number of sets ("ends"), each of which is diffeomorphic to the complement of a contractible compact set in \mathbb{R}^n .

we have

$$\nabla^{a} \left(\frac{D_{a}}{V} \right) = -\star d \star \left(-\frac{1}{V} i_{k} \mathbf{Z} \right)$$

$$= -\star d \left(\frac{1}{V} \mathbf{k} \wedge \star \mathbf{Z} \right)$$

$$= -\star \left(\frac{1}{V^{2}} \star (\boldsymbol{\omega} \wedge \mathbf{k}) \wedge \star \mathbf{Z} \right)$$

$$= \frac{1}{2V^{2}} (\boldsymbol{\omega} \wedge \mathbf{k})_{ab} \star \mathbf{Z}^{ab} . \tag{16}$$

Therefore,

$$\nabla^a \left(\frac{D_a}{V} \right) = -\frac{\omega_a H^a}{V^2} \tag{17}$$

and, analogously,

$$\nabla^a \left(\frac{B_a}{V} \right) = \frac{\omega_a E^a}{V^2} \ . \tag{18}$$

 $Proof\ of\ theorem\ 1.$ Let us introduce an auxiliary open set

$$O := \{ x \in \mathcal{M} \mid \mathcal{L}_{\mathcal{F}}(x) \neq 0 \} . \tag{19}$$

In other words, due to assumed NEC, $\mathcal{L}_{\mathcal{F}}(x) < 0$ for all $x \in O$ and $\mathcal{L}_{\mathcal{F}}(y) = 0$ for all $y \in \mathcal{M} - O$. As the electromagnetic field decays along each "end" and Lagrangian density obeys Maxwellian weak field limit, we know that O is nonempty.

At each point of the complement $\mathcal{M} - O$, the gravitational field equation is reduced to $E_{ab} = 2\pi T g_{ab}$. Thus, divergence $\nabla^a E_{ab} = 0$ implies that trace T is constant on each connected component of the interior $(\mathcal{M} - O)^{\circ}$. Furthermore, by the assumption of the theorem $\omega = 0$, so that k^a is a hypersurface orthogonal vector field. Let Σ be an arbitrary spacelike hypersurface from the basic assumption (3). Inserting, respectfully, $\alpha = \mathbf{D}/V$ and $\alpha = \mathbf{B}/V$ in the equation (A7), both of which satisfy $\mathcal{L}_k \alpha = 0$, we get

$$\int_{\partial \Sigma} \frac{1}{V} \star (\mathbf{k} \wedge \mathbf{D}) = 0 , \quad \int_{\partial \Sigma} \frac{1}{V} \star (\mathbf{k} \wedge \mathbf{B}) = 0 . \quad (20)$$

Formally, the integral over $\partial \Sigma$ may denote the limit for the integral over the "sphere at infinity". Furthermore, for $\alpha = \Phi \mathbf{D}/V$ and $\alpha = \Psi \mathbf{B}/V$ we get, respectfully,

$$\nabla^a \left(\frac{\Phi}{V} D_a \right) = \frac{4}{V} \left(\mathscr{L}_{\mathcal{F}} E_a E^a - \mathscr{L}_{\mathcal{G}} E_a B^a \right) \tag{21}$$

and

$$\nabla^a \left(\frac{\Psi}{V} B_a \right) = \frac{4}{V} \left(\mathcal{L}_{\mathcal{F}} B_a B^a + \mathcal{L}_{\mathcal{G}} E_a B^a \right) . \tag{22}$$

Sum of these two equations,

$$\nabla^a \left(\frac{\Phi}{V} D_a + \frac{\Psi}{V} B_a \right) = \frac{4}{V} \left(\mathcal{L}_{\mathcal{F}} E_a E^a + \mathcal{L}_{\mathcal{F}} B_a B^a \right) \tag{23}$$

integrated over Σ , with help of (20) and the fall-off conditions on potentials Φ and Ψ , leads to

$$\int_{\Sigma} \frac{\mathscr{L}_{\mathcal{F}}}{V} \left(E_a E^a + B_a B^a \right) \hat{\epsilon} = 0 , \qquad (24)$$

with induced volume 3-form $\hat{\epsilon}$. Now, as \mathbf{k} is strictly timelike, V>0, neither E^a nor B^a can be causal (as $k_aE^a=0$ and $k_aB^a=0$) and the integrand above is strictly nonnegative on $O\cap\Sigma$ and zero on $(\mathscr{M}-O)\cap\Sigma$. As the integral is zero, it follows that $E^a=0=B^a$ and, as $\mathscr{L}(0,0)=0$, consequently T=0 on $O\cap\Sigma$. By continuity this implies that T=0 on $\overline{O}\cap\Sigma$, thus T=0 on the whole Σ . In conclusion, on each point of the set $O\cap\Sigma$ we have $F_{ab}=0$, while on each point of the set $(\mathscr{M}-O)\cap\Sigma$ the electromagnetic field F_{ab} is either zero or stealth.

We stress that the argument works irrespectively of gravitational part of the equations of motion, as long as the coupling is minimal and tensor E_{ab} is divergence free. If $\mathcal{L}_{\mathcal{F}} = 0$ we can find simple counterexamples, such as take stealth field on static background [44].

Proof of theorem 2. Here we turn to the Einstein-Hilbert case, $E_{ab} = G_{ab}$. Exterior derivative of the twist 1-form, with help of the Killing lemma $d \star d \mathbf{k} = 2 \star \mathbf{R}(k)$, may be written as

$$d\boldsymbol{\omega} = -2\star(\mathbf{k} \wedge \mathbf{R}(k)) =$$

$$= 64\pi \mathcal{L}_{\mathcal{F}} \star(\mathbf{k} \wedge \mathbf{T}^{(\text{Max})}(k)) =$$

$$= 4\mathbf{E} \wedge \mathbf{H} . \tag{25}$$

Using electromagnetic scalar potentials,

$$d\boldsymbol{\omega} = -4(d\Phi \wedge \mathbf{H}) = -4(\mathbf{E} \wedge d\Psi) \tag{26}$$

we see that both $\omega + 4\Phi \mathbf{H}$ and $\omega - 4\Psi \mathbf{E}$ are closed 1-forms. Thus, as \mathscr{M} is by assumption, simply connected, we can globally define new scalar potentials U_E and U_H , such that

$$\omega = -4\Phi \mathbf{H} + dU_H \tag{27}$$

$$=4\Psi\mathbf{E}+dU_E. \qquad (28)$$

As with the electromagnetic potentials Φ and Ψ , we see directly from the definition that $\pounds_k U_E = 0$ and $\pounds_k U_H = 0$. Note that

$$\omega_a \omega^a = -4\Phi \omega_a H^a + \omega^a \nabla_a U_H \tag{29}$$

$$=4\Psi\omega_a E^a + \omega^a \nabla_a U_E \ . \tag{30}$$

Now, using a basic formula

$$\nabla^a \left(\frac{\omega_a}{V^2} \right) = 0 \tag{31}$$

and relations above, we have four divergence identities,

$$\nabla^a \left(U_E \frac{\omega_a}{V^2} \right) = \frac{\omega^a \nabla_a U_E}{V^2} \,, \tag{32}$$

$$\nabla^a \left(U_H \frac{\omega_a}{V^2} \right) = \frac{\omega^a \nabla_a U_H}{V^2} \,, \tag{33}$$

$$\nabla^a \left(\frac{\Phi}{V} D_a \right) = \frac{4}{V} \left(\mathcal{L}_{\mathcal{F}} E_a E^a - \mathcal{L}_{\mathcal{G}} E_a B^a \right) - \Phi \frac{\omega_a H^a}{V^2} ,$$
(34)

$$\nabla^a \left(\frac{\Psi}{V} B_a \right) = \frac{4}{V} \left(\mathcal{L}_{\mathcal{F}} B_a B^a + \mathcal{L}_{\mathcal{G}} E_a B^a \right) + \Psi \frac{\omega_a E^a}{V^2} . \tag{35}$$

A brief inspection reveals that an auxiliary 1-form

$$\mathbf{W} := \frac{U_E + U_H}{V^2} \,\boldsymbol{\omega} + \frac{4}{V} \left(\Phi \mathbf{D} + \Psi \mathbf{B} \right) \tag{36}$$

has a rather simple covariant divergence

$$\nabla_a W^a = \frac{16}{V} \mathcal{L}_{\mathcal{F}} \left(E_a E^a + B_a B^a \right) + 2 \frac{\omega_a \omega^a}{V^2} \ . \tag{37}$$

Taking into account Einstein's field equation,

$$R_{ab} = 8\pi \left(T_{ab} - \frac{1}{2} T g_{ab} \right) \tag{38}$$

and

$$8\pi T_{ab}^{(\text{Max})} k^a k^b = E_a E^a + B_a B^a , \qquad (39)$$

we get

$$\frac{4}{V} R_{ab} k^a k^b - \frac{2}{V^2} \omega_a \omega^a = -\nabla_a W^a + 8\pi T \ . \tag{40}$$

Now we turn to Heusler's mass formula [7]. If we contract

$$\star d\omega = 2 \mathbf{k} \wedge \mathbf{R}(k) \tag{41}$$

with k^a and take Hodge dual, we obtain

$$-\star \mathbf{R}(k) = \frac{R_{ab}k^ak^b}{V} \star \mathbf{k} + \frac{1}{2V}\mathbf{k} \wedge d\boldsymbol{\omega} . \qquad (42)$$

Also, using

$$-d\left(\frac{1}{V}\mathbf{k}\wedge\boldsymbol{\omega}\right) = \frac{\omega_a\omega^a}{V^2}\star\mathbf{k} + \frac{1}{V}\mathbf{k}\wedge d\boldsymbol{\omega}$$
(43)

we have

$$-\star \mathbf{R}(k) = \left(\frac{R_{ab}k^ak^b}{V} - \frac{\omega_a\omega^a}{2V^2}\right)\star\mathbf{k} - d\left(\frac{1}{2V}\,\mathbf{k}\wedge\boldsymbol{\omega}\right). \tag{44}$$

Thus, relying on the fall-off properties of the twist 1-form ω inferred from the basic assumptions, Komar's mass

$$M = -\frac{1}{4\pi} \int_{\Sigma} \star \mathbf{R}(k) \tag{45}$$

may be written in a form

$$M = \frac{1}{4\pi} \int_{\Sigma} \left(\frac{R_{ab} k^a k^b}{V} - \frac{\omega_a \omega^a}{2V^2} \right) \star \mathbf{k} , \qquad (46)$$

which, in combination with (40), becomes

$$M = -\frac{1}{16\pi} \int_{\Sigma} \nabla_a W^a \star \mathbf{k} + \frac{1}{2} \int_{\Sigma} T \star \mathbf{k} . \tag{47}$$

We note in passing that the formula obtained here is consistent with the generalized Smarr formula [32, 50, 51]. The W-term vanishes at infinity, while the T-term is nonpositive, given that DEC holds. Finally, positive energy theorem implies that $M \geq 0$ and M = 0 if and only if the spacetime is Minkowski. As we have proven that $M \leq 0$, it follows that M = 0. Therefore $T_{ab} = 0$, implying that any nontrivial electromagnetic field must be zero or stealth.

V. FURTHER GENERALIZATIONS

We now turn to possible generalizations of the main results, which we shall sort into two directions.

Theories with charged matter. As a simplest model of matter we may choose a complex scalar field ϕ , with the total Lagrangian

$$\mathscr{L}^{(\text{tot})} = \mathscr{L}(\mathcal{F}, \mathcal{G}) + (\mathcal{D}_a \phi)^* (\mathcal{D}^a \phi) - \mathscr{U}(\phi^* \phi) , \quad (48)$$

constructed with the covariant gauge derivative $\mathcal{D}_a = \nabla_a + iqA_a$, gauge 1-form **A** which defines the electromagnetic 2-form $\mathbf{F} = \mathrm{d}\mathbf{A}$, and scalar (self-interaction) potential \mathscr{U} . In this theory generalized Maxwell's equations have a form

$$d \star \mathbf{Z} = 4\pi \star \mathbf{J} \tag{49}$$

with the current 1-form

$$J_a = \frac{iq}{4\pi} \left(\phi^* \mathcal{D}_a \phi - \phi (\mathcal{D}_a \phi)^* \right) . \tag{50}$$

Electric 1-form **E** is again closed and we have associated electric scalar potential Φ . Let us, for simplicity, focus on the strictly static case, $\omega = 0$. Then

$$\nabla^a \left(\frac{1}{V} D_a \right) = -\frac{4\pi}{V} k^a J_a , \quad \nabla^a \left(\frac{1}{V} B_a \right) = 0 \quad (51)$$

and

$$\nabla^a \left(\frac{\Phi}{V} D_a \right) = \frac{4}{V} \left(\mathcal{L}_{\mathcal{F}} E_a E^a - \mathcal{L}_{\mathcal{G}} E_a B^a \right) - \frac{4\pi\Phi}{V} k^a J_a .$$
(52)

First technical obstacle is a treatment of the term $\Phi k^a J_a$ which, without additional assumptions, is in general neither positive nor negative definite. Setting aside spacetimes with symmetry noninheriting scalar fields [52–54], let us for simplicity assume that $\mathcal{L}_k \phi = 0$. Also, taking

into account remarks from [32, 42], we assume that the gauge choice is made such that $\mathcal{L}_k \mathbf{A} = 0$. Now, as

$$d(\Phi + i_k \mathbf{A}) = -\mathbf{E} + (\pounds_k - i_k d)\mathbf{A} = 0 , \qquad (53)$$

given that both Φ and k^aA_a vanish at infinity, we may set $\Phi = -k^aA_a$. This leads us to simplification

$$\Phi k^a J_a = \frac{(q\Phi)^2}{2\pi} \,\phi^* \phi \ge 0 \ . \tag{54}$$

Furthermore, magnetic 1-form \mathbf{H} is no longer necessary a closed form as

$$d\mathbf{H} = 4\pi \star (\mathbf{k} \wedge \mathbf{J}) . \tag{55}$$

This is a familiar obstacle to introduction of magnetic scalar potential on domains which contain nonvanishing electric currents. There are several subcases in which we can proceed with similar strategy of proof as above:

(a) if $\mathbf{k} \wedge \mathbf{J} = 0$, which allows us to introduce magnetic scalar potential Ψ and deduce

$$\nabla^a \left(\frac{\Psi}{V} B_a \right) = \frac{4}{V} \left(\mathcal{L}_{\mathcal{F}} B_a B^a + \mathcal{L}_{\mathcal{G}} E_a B^a \right) ; \qquad (56)$$

(b) if we have a strictly electric system, in sense that $\mathbf{B} = 0$, so that the equation (52) may suffice for the proof.

Given that any of two conditions above, (a) or (b), are met, repetition of the argument from the previous section leads to the conclusion that the electromagnetic field is trivial on the set O where $\mathcal{L}_{\mathcal{F}} \neq 0$. We note in passing that equation (52) may be also used if we have an \mathcal{F} -class theory, but only to deduce that $\mathbf{E} = 0$ on the set O, without any control of the magnetic field \mathbf{B} (unless, again, we invoke condition (a) or (b)).

Furthermore, divergence of the gravitational field equation on the interior $(\mathcal{M} - O)^{\circ}$ leads to $\nabla_a T = 4J^b F_{ba}$. If, in addition, we assume either (a) or (b) from above, decomposition $V\mathbf{F} = \mathbf{k} \wedge \mathbf{E} + \star (\mathbf{k} \wedge \mathbf{B})$ and $k^a \nabla_a T = 0$ allow us to deduce $V \nabla_a T = 4(k^b J_b) E_a$. In particular, $\mathbf{B} = 0$ on the set $(\mathcal{M} - O)^{\circ}$ also implies $\mathbf{D} = 0$ and, via divergence identities, $k^a J_a = 0$, leading to the conclusion that trace T is constant on each connected component of the domain $(\mathcal{M} - O)^{\circ}$ (it is not clear if this necessarily holds in the (a) subcase).

In conclusion, at least under some additional assumptions, the initial problem can be reduced to the question of existence of self-gravitating scalar solitons [2, 55]. It is important to stress that spacetimes with charged boson stars evade the partial no-go result from above due to symmetry noninheriting scalar field. Namely, typical ansatz for such solutions feature scalar field of the form $\phi(t,r) = f(r)e^{i\omega t}$, so that $\pounds_k \phi = i\omega \phi$ and definiteness of the term $\Phi k^a J_a$ is in general lost.

Higher dimensional theories. Let us look at spacetimes of dimension $m \geq 5$. As \mathcal{G} invariant is a scalar only in four

spacetime dimensions, where **F** and its Hodge dual \star **F** are both 2-forms, here we treat only \mathcal{F} -class theories. We may define twist (m-3)-form as $\boldsymbol{\omega} := (-1)^{m+1} \star (\mathbf{k} \wedge d\mathbf{k})$, in order to preserve its form in abstract indices,

$$\omega_{a_1...a_{m-3}} = \epsilon_{a_1...a_{m-3}}{}^{bcd}k_b \nabla_c k_d , \qquad (57)$$

and 1-form $\mathbf{D} := -i_k \mathbf{Z}$ as above. Taking into account that

$$i_k \mathbf{Z} = (-1)^{m+1} \star (\mathbf{k} \wedge \star \mathbf{Z}) \tag{58}$$

we have

$$\nabla^a \left(\frac{D_a}{V} \right) = \frac{1}{(m-2)!V^2} \left(\boldsymbol{\omega} \wedge \mathbf{k} \right)_{a_1 \dots a_{m-2}} \star \mathbf{Z}^{a_1 \dots a_{m-2}} .$$
(59)

Basic divergence identity for strictly static spacetime, with $\omega = 0$, is

$$\nabla^a \left(\frac{\Phi}{V} D_a \right) = \frac{4}{V} \mathcal{L}_{\mathcal{F}} E_a E^a . \tag{60}$$

Assuming the appropriate fall-off conditions, namely $\Phi = O(r^{-(m-3)})$ and $\mathbf{D} = O(r^{-(m-2)})$ (cf. also [56]), we may repeat the previous argument to conclude that $\mathbf{E} = 0$ on the set O. Further technical obstacle is that in a m-dimensional spacetime magnetic fields $\mathbf{B} = i_k \star \mathbf{F}$ and $\mathbf{H} = i_k \star \mathbf{Z}$ are (m-3)-forms, so that we less control on sign of their squares, such as $B_{a_1...a_{m-3}}B^{a_1...a_{m-3}}$.

VI. DISCUSSION

We have proved that, up to exotic stealth solutions, NLE theories on a simply connected four dimensional spacetimes do not admit globally regular, stationary solitonic solutions. These results admit only partial generalizations in the presence of the charged matter and in higher dimensional theories. Limitations in both directions do not come as a surprise, due to known solutions with charged bosonic stars and increase of number of electromagnetic degrees of freedom with number of spacetime dimensions.

Remaining open questions may be grouped as follows:

- 1. Are conclusions altered if the spacetime manifold \$\mathcal{M}\$ is not simply connected?
- 2. How to treat a NLE theory which does not obey the Maxwellian weak field limit?
- 3. Can constrains on 4-dimensional theories with complex scalar fields and higher dimensional theories be strengthened?

Simple connectedness of the manifold \mathcal{M} was invoked in order to guarantee the existence of scalar potentials Φ , Ψ , U_E and U_H . This assumption is not necessary, but on a non-simply connected manifold one needs to either (a) impose some boundary conditions which imply existence of scalar potentials, or (b) construct divergence identities which do not involve scalar potentials and which allow one to prove theorems analogous to the ones treated in this paper.

Maxwellian weak field limit or, slightly weaker, mere assumption that partial derivatives $\mathcal{L}_{\mathcal{F}}$ and $\mathcal{L}_{\mathcal{G}}$ are well-defined and finite at the origin of the \mathcal{F} - \mathcal{G} plane, plays a role in elimination of the boundary terms at asymptotic ends. However, there are NLE theories, such as the power-Maxwell [45, 46] (for powers less than 1) and ModMax theory [16–18, 57], which do not behave well in this sense. Here one must first find a proper way to incorporate some notion of asymptotic flatness, with the appropriate fall-off conditions for the fields.

We feel that conditions under which a NLE theory with complex scalar field (or some other form of the charged matter) does not admit strictly stationary solitonic solutions should be mapped more carefully. Also, it is not clear how to sensibly choose additional assumptions which might lead to generalizations in higher dimensional theories. Finally, we remark that in (1+2)-dimensional spacetimes one might again rely on the divergence relation (60), as well as the lower dimensional positive energy theorem [58], but the problem is that natural fall-off condition for the scalar potential $\Phi = O(\ln r)$ does not seem to be sufficient to get rid of the boundary terms.

Appendix A: Menagerie of identities

Let us first, for generality, assume that (\mathcal{M}, g_{ab}) is a smooth m-dimensional Lorentzian manifold. The Hodge dual of a p-form α , defined as

$$(\star \alpha)_{a_{p+1}...a_m} := \frac{1}{p!} \alpha_{a_1...a_p} \epsilon^{a_1...a_p}{}_{a_{p+1}...a_m} , \qquad (A1)$$

twice applied produces a sign according to

$$\star\star\boldsymbol{\alpha}=(-1)^{p(m-p)+1}\boldsymbol{\alpha}\;. \tag{A2}$$

A simple useful rule, so-called "flipping over the Hodge", reads

$$i_X \star \boldsymbol{\alpha} = \star (\boldsymbol{\alpha} \wedge \mathbf{X}) ,$$
 (A3)

where **X** is the associated 1-form, $X_a = g_{ab}X^b$.

We shall introduce an auxiliary coderivative operator, acting on p-form α as

$$\delta \boldsymbol{\alpha} := (-1)^{m(p+1)+1} \star d \star \boldsymbol{\alpha} , \qquad (A4)$$

which in the abstract index notation is simply

$$\delta \alpha_{a_1 \dots a_{p-1}} = \nabla^b \alpha_{ba_1 \dots a_{p-1}} . \tag{A5}$$

Note that for even m we have $\delta \alpha = -\star d \star \alpha$. If K^a is a Killing vector field and $K_a = g_{ab}K^b$ associated 1-form, then the following identity holds,

$$\pounds_K \alpha = \delta(\mathbf{K} \wedge \alpha) + \mathbf{K} \wedge \delta \alpha . \tag{A6}$$

We are mostly interested in the case where α is a 1-form such that $\pounds_K \alpha = 0$. Whence, equation (A6) integrated over a smooth hypersurface Σ , with application of the generalized Stokes' theorem, leads to

$$\int_{\Sigma} (\delta \boldsymbol{\alpha}) \star \mathbf{K} = \int_{\partial \Sigma} \star (\mathbf{K} \wedge \boldsymbol{\alpha}) , \qquad (A7)$$

where we have, for simplicity, suppressed pullback symbol

ACKNOWLEDGMENTS

The research was supported by Croatian Science Foundation Project No. IP-2020-02-9614.

P. Chruściel, J. Costa, and M. Heusler, Living Rev.Rel. 15, 7 (2012), arXiv:1205.6112 [gr-qc].

^[2] M. Heusler, *Black Hole Uniqueness Theorems* (Cambridge University Press, Cambridge New York, 1996).

^[3] J. A. Wheeler, Phys. Rev. 97, 511 (1955).

^[4] A. Lichnerowicz, Théories Relativistes de la Gravitation et de l'Électromagnétisme (Masson et Cie, Paris, 1955).

^[5] B. Carter, in Black Holes (Gordon and Breach, New York, 1973).

^[6] B. Carter, Gen. Relativ. Gravit. 42, 653 (2010).

^[7] M. Heusler, Class. Quantum Grav. 12, 779 (1995), arXiv:gr-qc/9411054.

 ^[8] T. Shiromizu, S. Ohashi, and R. Suzuki, Phys. Rev. D 86, 064041 (2012),

arXiv:1207.7250 [gr-qc].

^[9] C. Herdeiro and J. Oliveira, Class. Quantum Grav. 36, 105015 (2019), arXiv:1902.07721 [gr-qc].

^[10] C. Herdeiro and J. Oliveira, Phys. Lett. B 800, 135076 (2020), arXiv:1909.08915 [gr-qc].

^[11] M. Born, Proc. R. Soc. A 143, 410 (1934).

^[12] M. Born and L. Infeld, Proc. R. Soc. A 144, 425 (1934).

^[13] E. Fradkin and A. Tseytlin, Phys. Lett. **163B**, 123 (1985).

^[14] W. Heisenberg and H. Euler, Z. Phys. 98, 714 (1936), arXiv:physics/0605038 [physics].

^[15] G. Dunne, "Heisenberg-Euler effective La-

- grangians: Basics extensions," and in From fields to strings: Circumnavigating theoretical physics. [39] Togan mParken collected (3 vQume set). (World Scientific, 2004) 445-522. pp. arXiv:hep-th/0406216.
- [16] I. Bandos, K. Lechner, D. Sorokin, and Ρ. Townsend, Phys. Rev. D **102**, 121703 (2020), arXiv:2007.09092 [hep-th].
- Kosyakov, Phys. Lett. B 810, 135840 (2020), [17] B. arXiv:2007.13878 [hep-th].
- [18] G. W. Gibbons D. Α. Rasheed. and Nucl. Phys. B 454, 185 (1995), arXiv:hep-th/9506035.
- [19] M. Aaboud etal.(ATLAS), Nature Phys. 13, 852 (2017), arXiv:1702.01625 [hep-ex].
- [20] J. Ellis, N. Mavromatos, and Phys. Rev. Lett. 118, 261802 (2017), arXiv:1703.08450 [hep-ph].
- Niau Akmansoy Medeiros, and Eur. Phys. J. C 78, 143 (2018), arXiv:1712.05486 [hep-ph].
- [22] P. Niau Akmansoy and G. Medeiros, Phys. Rev. **D** 99, 115005 (2019), arXiv:1809.01296 [hep-ph].
- Α. Tanaka Matter Radiat. at Extremes 5, 024402 (2020).
- [24] K. Bronnikov, V. Melnikov, G. Shikin, and K. Staniukowicz, Annals Phys. 118, 84 (1979).
- Bronnikov, Phys. Rev. D 63, 044005 (2001), [25] K. arXiv:gr-qc/0006014 [gr-qc].
- [26] A. Burinskii S. Hildebrandt, and Phys. Rev. D 65, 104017 (2002), arXiv:hep-th/0202066 [hep-th].
- [27] K. A. Bronnikov, Phys. Rev. D **96**, 128501 (2017), arXiv:1712.04342 [gr-qc].
- [28] J. Bardeen, in Proceeding of the International Conference GR5 (Tbilisi, 1968) p. 174.
- [29] E. Avón-Beato García, and Α. Phys. Rev. Lett. 80, 5056 (1998), arXiv:gr-qc/9911046 [gr-qc].
- Ayón-Beato A. García, and Phys. Lett. **B493**, 149 (2000), arXiv:gr-qc/0009077 [gr-qc].
- [31] L.-M. Cao, Υ. Peng, Xu, J. Phys. Rev. D **90**, 024046 (2014), arXiv:1404.6639 [gr-qc].
- [32] A. Bokulić, Т. Jurić, and I. Smolić, Phys. Rev. D **103**, 124059 (2021), arXiv:2102.06213 [gr-qc].
- [33] J. Plebański, (1970), Nordita.
- [34] R. Schoen and S.-T. Yau, Phys. Rev. Lett. 43, 1457 (1979).
- [35] R. Schoen S.-T. and Yau, Commun. Math. Phys. **65**, 45 (1979)
- [36] R. Schoen S.-T. and Yau, Commun. Math. Phys. **79**, 47 (1981)
- Schoen S.-T. [37] R. and Yau, Commun. Math. Phys. **79**, 231 (1981).

- [38] E. Witten, Commun. Math. Phys. 80, 381 (1981).
- Taubes. Commun. Math. Phys. 84, 223 (1982).
- [40] S. Dain, "Positive energy theorems in General Relativity," in Springer Handbook of Spacetime, edited by A. Ashtekar and V. Petkov (2014) pp. 363–380, arXiv:1302.3405 [gr-qc].
- [41] P. Tod. Gen. Relativ. Gravit. 39, 111 (2007), arXiv:gr-qc/0611035 [gr-qc].
- [42] M. Cvitan, P. Dominis Prester, and I. Smolić, Class. Quantum Grav. 33, 077001 (2016), arXiv:1508.03343 [gr-qc].
- [43] I. Barjašić, L. Gulin, Smolić, Phys. Rev. **D** 95, 124037 (2017), I. arXiv:1705.00628 [gr-qc].
- [44] I. Smolić, Phys. Rev. D 97, 084041 (2018), arXiv:1711.07490 [gr-qc].
- [45] M. Hassaïne Martínez, and Phys. Rev. D **75**, 027502 (2007), arXiv:hep-th/0701058 [hep-th].
- C. [46] M. Hassaïne and Martínez, Class. Quantum Grav. 25, 195023 (2008), arXiv:0803.2946 [hep-th].
- V[47] M. Ortaggio and Pravda, Class. Quantum Grav. 33, 115010 (2016), arXiv:1506.04538 [gr-qc].
- V. [48] M. Pravda, Ortaggio and Phys. Lett. B 779, 393 (2018), arXiv:1708.08017 [hep-th].
- [49] S. Hervik, M. Ortaggio, and Pravda, Class. Quant. Grav. 35, 175017 (2018), arXiv:1806.05835 [gr-qc].
- [50] L. Gulin Smolić, and Class. Quantum Grav. 35, 025015 (2018), arXiv:1710.04660 [gr-qc].
- [51] Y. Zhang Gao, and Class. Quantum Grav. 35, 145007 (2018), arXiv:1610.01237 [gr-qc].
- [52] I. Smolić, Class. Quantum Grav. 32, 145010 (2015), arXiv:1501.04967 [gr-qc].
- Phys. Rev. 95, 024016 (2017), Smolić, arXiv:1609.04013 [gr-qc].
- Franzin and Smolić, Class. Quantum Grav. 38, 115004 (2021), arXiv:2101.05816 [gr-qc].
- [55] C. Herdeiro Radu, and Int. J. Mod. Phys. D 24, 1542014 (2015), arXiv:1504.08209 [gr-qc].
- Ortaggio, Phys. Rev. D 90, 124020 (2014), arXiv:1406.3186 [gr-qc].
- [57] D. Flores-Alfonso, B. González-Morales, R. Linares, and M. Maceda, Phys. Lett. B 812, 136011 (2021), arXiv:2011.10836 [gr-qc].
- [58] W. W.-Y. Wong, (2012), arXiv:1202.6279 [gr-qc].