# Wormhole inducing exponential expansion in $R^2$ gravity

B Modak\*and Gargi Biswas<sup>†</sup>

November 23, 2021

Department of Physics, University of Kalyani, Kalyani, India-741235

#### Abstract

We present a few exact solutions of the Euclidean field equations in  $R^2$  gravity in spatial homogeneous and isotropic background in 4-dimension. Wormhole configuration, as well as oscillatory scale factors are obtained as a function of Euclidean time  $\tau$  for  $\kappa = 0, \pm 1$ . Each of the oscillatory solution in  $\tau$  reduces to the Lorentzian wormhole using analytic continuation from Euclidean space to Lorentz spacetime by  $\tau = it$ . The Lorentzian wormhole leads to an exponential expanding era asymptotically. An Euclidean wormhole in the very early universe could evolve to an inflationary era at later epoch for  $\kappa = 1$ .

**PACS:** 98.80.-k, 04.50.+h **Keywords:** Wormhole, Analytic continuation, Inflation and  $R^2$  gravity

## 1 Introduction

Inflationary expansion is necessary [1–6] in the early vacuum dominated universe to resolve some of the problems of standard cosmology, like horizon problem, structure formation problem, etc. However, the initial singularity problem prevails in the cosmological model before the Planck era of strong curvature regime. In the strong curvature regime quantum gravity is necessary, however it revealed that Einstein's gravity is not renormalizable.

In this endeavour the quadratic curvature (i.e.  $R^2$ ) correction was introduced from the fundamental of Physics to improve the renormalizability of general relativity [7]. Subsequently, other higher order curvature terms appear naturally as renormalization terms in quantum theory in curved space time [8]. In this venture modified theory of gravity [9–11] appears as an alternative theory of gravitation. Starobinsky [6] first proposed inflationary model based on  $R^2$  term. The modified theory of gravitation can accommodate not only inflation, but it can explain late time acceleration of the universe [12–14]. On due course of development low energy effective Heterotic string theory [15, 16], Kaluza Klein gravity [17–19], etc., appear as an alternative theory of gravitation.

The problem of cosmological singularity prevails irrespective of the success of aforesaid gravity theories, whose resolution has not yet been settled. A few proposals have been advocated to alleviate the singularity problem [20, 21]. First of all, the wave function of the universe [22] in quantum cosmology is finite even at classical singularity. Further, the Euclidean wormhole that appears in the context of quantum gravity, can alleviate cosmic singularity problem by introducing tunnelling of the universe through the classical singularity. Hawking and Page [23] proposed that the wave function is finite even at classical singularity and die away at large three space for the existence of quantum wormhole.

Wormhole solutions [24–31] and their consequences [32–39] are explored with different fields after the discovery of wormhole [40] as a solution of the Euclidean field equations invoking the axionic field in Einstein's gravity. In this context we specify the wormhole as a configuration governed by the Euclidean field equations, wherein two asymptotic regions of Robertson walker spacetimes are connected by a tube of minimum radius of scale factor  $a(\tau_0)$  at some time  $\tau_0$ . The axionic field in [40] acts as an exotic field, which gives rise to violation of null energy condition. The wormhole solution demands violation of null energy condition [41]. The violation can be minimised by introducing some geometric contribution instead of exotic fields in some theories, like the modified theory of gravity with higher order curvature terms [9–11], the low energy heterotic string effective theory [15, 16], Kaluza Klein gravity [17–19], etc.

<sup>\*</sup>E-mail: bijanmodak@yahoo.co.in

<sup>&</sup>lt;sup>†</sup>E-mail: biswasgb@gmail.com

Physics onward GUT era is well understood and evolution of the universe from the inflationary era to the radiation dominated era, subsequent re-ionization, structure formation, etc are well known in the literature. Physics of the pre-inflationary era, or the Planck era is not well understood in the literature and the chronological scenario of subsequent physical configuration is not known; however, the wormhole, if it exits at the Planck era, subsequent cosmic evolution should reproduce a regime of inflation at later stage. Though the cosmological wormholes are considered [42], [43], yet the viability of an inflationary era from an early wormhole configuration has not yet been clear in the literature. The evolution from an initial wormhole configuration at the Planck era to the subsequent inflationary scenario at later era may not be a smooth process, since the Planck era is determined by the quantum gravity, while the inflationary era can be described by the field theory and the Physics at these eras are completely different from the Planck era. Specifically, the term "evolution" in cosmology is specified with respect to proper time t and it is unique in the classical allowed domain, while the "time" in the quantum field theory in the preinflationary era is specified by the Euclidean time  $\tau$  and the corresponding background is the Euclidean space. The Lorentz time or the cosmic time "t" in the classically allowed domain cannot probe the events or evolutionary scenario in the classical forbidden domain in general, as well as the time " $\tau$ " in the Euclidean domain cannot probe the Lorentz domain. So to probe the scenario outside its domain we need analytic continuation of the function from the Euclidean domain to the Lorentzian domain, vice versa (or the change of geometry) and the Wicks rotation  $\tau = it$  serves its purpose. Thus to study evolution from an Euclidean wormhole to an inflationary scenario the transformation  $\tau = it$  acts as an important tool.

The work [44] is worthy to achieve an evolution of the Euclidean wormhole to a deSitter universe through a tube with oscillating radius invoking R,  $R^2$  and  $\Lambda$  terms in the action in Robertson Walker metric for closed three space section, however it depends on approximation. A reconstruction of the scalar tensor f(R) function corresponding to evolving wormhole is considered in [45] in an inhomogeneous spacetime generalizing Morris-Thorn metric, while in [46] Darabi presents the form of  $f(\hat{R})$  in Palatini  $f(\tilde{R})$  cosmology as  $f(\tilde{R}) \sim \tilde{R} - c_1 \tilde{R}^2 + c$  for small  $\tilde{R}$  from an Euclidean wormhole in O'Hanlon theory. Though the works [45], [46] are reconstruction of f(R) for evolving wormhole, however, they could not reproduce inflation from a wormhole configuration. Kehagias et al [47] outlined the solutions of pure  $R^2$  gravity in two distinct classes of solutions, namely class (i), the solution satisfying R = 0, but  $R_{\mu\nu} \neq 0$  and class (ii), the solution satisfying  $R_{\mu\nu} = 3\lambda g_{\mu\nu}$ , where  $\lambda$  is an arbitrary constant. The solution corresponding to class (i) is one of our Euclidean wormholes. Further, the solutions for the "00"-component of  $R_{\mu\nu} = 3\lambda g_{\mu\nu}$  are relevant to our solutions both for positive and negative values of  $\lambda$ , but the "space-space" components of the equations of class (ii) do not belong to our class of solutions. Interestingly, power law inflation [48] is obtained asymptotically from an Euclidean wormhole in gravity with  $R^{\frac{3}{2}}$  term using Noether symmetry. In a recent work [49] it is shown that an Euclidean wormhole in the early universe evolved to an inflationary era away from the throat of the wormhole using  $\tau = it$  in 5-dimensional Kaluza Klein spacetime minimally coupled with a scalar field.

Our motivation is to study the viability of transition of an Euclidean wormhole configuration in the early era to the inflationary scenario invoking analytic continuation by  $\tau = it$  in  $R^2$  gravity. It is important to note that all the gravitational theories contain at least linear *R*-term apart from higher order curvature contribution. However, pure  $R^2$  gravity in the context of supergravity and superstring theory possess two important features [50], namely it is the only ghost free theory in quadratic curvature tensor and one can show it is a scale invariant gravitational theory. Under conformal transformation, [47] the action of  $R^2$  gravity is equivalent to Einstein action coupled minimally with a mass-less scalar field and a non-zero cosmological constant. Interestingly, traversable Lorentzian wormhole [51] solutions have been considered in the framework of  $R^2$  gravity in time dependent inhomogeneous spacetime assuming constant Ricci scalar and trace-less energy momentum tensor. However, our work on wormhole is based on Robertson Walker background.

We investigate the Euclidean field equations with a view to obtain possible wormhole solution in the very early era and consequence of them in the later epoch in Lorentz time t using the modified theory of gravity in the spatially homogeneous and isotropic Euclidean background in 4-dimension. In a toy model, we consider  $R^2$  term in the Einstein-Hilbert action since such term alone (without any matter field) admits inflationary solution [5] in cosmic time t. The field equations contains fourth derivative terms in the standard variational principle. Thus to get an alternative form of the field equations in second order we consider Boulware et al [52] proposal. In this proposal one can introduce a new variable, which is the first derivative of the action with respect to the highest derivative of the field variable that appears in the action after removing the total derivative terms in the action following the works [52–55].

Some exact solutions of the Euclidean field equations are obtained in  $R^2$  gravity. Euclidean wormholes, as well as oscillatory solution of  $a^2(\tau)$ , are obtained as a function of Euclidean time  $\tau$  for all 3-space

curvature parameter  $\kappa$   $(0, \pm 1)$ . Oscillatory solution of  $a^2(\tau)$  reduces to a Lorentzian wormhole under analytic continuation  $\tau = it$  and asymptotically it leads to an exponential expansion. A polynomial solution of  $a^2(\tau)$  leads to a Lorentzian wormhole with  $\tau = it$  and asymptotically it reduces to power law inflation. Further, an Euclidean wormhole is allowed in the very early universe for  $\kappa = 1$ , while at later era (i.e. asymptotic region) inflationary expansion is realized naturally using  $\tau = it$ . All the solutions in this work satisfy the null energy condition, which are being discussed in the Appendix.

We present the field equations considering  $R^2$  term in the Einstein-Hilbert action in section 2. In section 3 we present the solution of the field equations for  $R^2$  case. Finally, we present a brief discussion in section 4 and an Appendix in section 5.

# 2 The Field equations in $R^2$ gravity

Let the Einstein Hilbert action including quadratic curvature term as

$$S = \int d^4x \sqrt{g} \,\left[\alpha R + \frac{\beta}{6}R^2\right],\tag{1}$$

where  $\alpha$  and  $\beta$  are constants. The Ricci scalar in the Euclidean background of Robertson-Walker spacetime

$$ds^{2} = d\tau^{2} + a^{2}(\tau) \left[ \frac{dr^{2}}{1 - \kappa r^{2}} + r^{2} \left( d\theta^{2} + \sin^{2} \theta d\phi^{2} \right) \right]$$

is  $R = -6(\frac{a''}{a} + \frac{a'^2}{a^2} - \frac{\kappa}{a^2})$ , where the Euclidean time  $\tau$  and the Lorentz time t are related by  $\tau = it$ . The scale factor is  $a(\tau)$ , an overhead prime stands derivative with respect to  $\tau$  and  $\kappa$  is the three space curvature parameter and it can take  $0, \pm 1$ . Action (1) leads to

$$S = m \int d\tau \Big[ \alpha (aa'^2 + \kappa a) + \beta \{ aa''^2 + \frac{1}{a} (-a'^2 + \kappa)^2 \} \Big] + S_{m_1},$$
(2)

where  $m = 12\pi^2$  is a constant and  $S_{m_1} = -m \left[ \int d\tau \left\{ \alpha a^2 a' + 2\beta a'(\kappa - a'^2) + \frac{4\beta}{3} a'^3 \right\}' \right]$ . The integrand in the action S in (2) contains a term quadratic in a'', so the Lagrangian cannot be obtained in terms of usual canonical variables. To remove this problem and to reduce the fourth order gravity theory to a second order theory without changing degrees of freedom of the system we introduce new variable Qfollowing [52] as

$$m\beta \ Q = -\frac{\partial S}{\partial a''}, \text{ which gives } Q = -2aa''.$$
 (3)

Now replacing  $aa''^2$  by Q using (3) as  $aa''^2 = -a''Q - \frac{Q^2}{4a} = -(a'Q)' + a'Q' - \frac{Q^2}{4a}$  in equation (2) we can reduce following [52,54] as

$$S = m \int d\tau \Big[ \alpha \Big( aa'^2 + \kappa a \Big) + \beta \Big\{ a'Q' - \frac{Q^2}{4a} + \frac{1}{a} (-a'^2 + \kappa)^2 \Big\} \Big] + S_m,$$

where  $S_m = S_{m_1} - m\beta \int d\tau (Qa')'$ . The term  $S_m$  is a surface term (or boundary term ) and extraction of the surface term from the action is necessary to get well defined action principle. This term is analogous to the Gibbons-Hawking-York surface term of general relativity. Now with proper choice  $S_m$  the Lagrangian can be written as

$$L = m \Big[ \alpha \Big( aa'^2 + \kappa a \Big) + \beta \Big( a'Q' - \frac{Q^2}{4a} + \frac{1}{a} (-a'^2 + \kappa)^2 \Big) \Big].$$
(4)

The field equations are

$$\alpha \left(\frac{{a'}^2}{a^2} - \frac{\kappa}{a^2}\right) + \beta \left(\frac{a'Q'}{a^3} + \frac{Q^2}{4a^4} - \frac{(\kappa - {a'}^2)(\kappa + 3{a'}^2)}{a^4}\right) = 0,$$
(5)

$$\alpha(2aa'' + {a'}^2 - \kappa) + \beta \Big[ Q'' + 12\frac{{a'}^2 a''}{a} - 4\kappa \frac{a''}{a} + 4(\kappa - {a'}^2)\frac{{a'}^2}{a^2} + \frac{(\kappa - {a'}^2)^2}{a^2} - \frac{Q^2}{4a^2} \Big] = 0.$$
(6)

Now Q variation in (4) gives Q = -2aa'', which is just the definition of Q in (3). Now using above Q in (5) and (6) we have usual fourth order gravity theory in terms of a, however the field equations in terms of a and Q represent the second order equations. Solution of the field equations needs special care as they are coupled non linear equations. In [44] evolution of wormhole is considered assuming

 $a'^2 = 1 + f(a)$  in (5) for  $\kappa = 1$ , where f(a) is arbitrary function of a and the authors [44] extract solution when  $(f(a) - f_0)$  is very small, where  $f(a) = f_0$  at the extreme regime of a. In our presentation, we consider exact solution of evolving wormhole using all the field equations in  $R^2$  gravity. It is important to mention that a wormhole has two asymptotic regions connected by a tube with a minimum radius, so  $a(\tau)$  should have a' = 0 and a'' > 0 at some  $\tau = \tau_0$  at the minimum, otherwise  $a(\tau)$  is finite.

## **3** Solution of the field equations:

Now multiplying (5) by  $a^3$  and (6) by a and taking sum of them gives

$$2\alpha(a^2a'' + aa'^2 - \kappa a) + \beta \left(aQ' + 4a'^3 - 4\kappa a'\right)' = 0.$$
(7)

The solution of  $a(\tau)$  from (7) is not possible in closed form for  $\alpha \neq 0$  and  $\beta \neq 0$ . Thus considering  $R^2$  term only in the action ( $\alpha = 0$  and  $\beta \neq 0$ ) with Q = -2aa'' in (7) we have

$$aQ' + 4a'^3 - 4\kappa a' = M = -2a' \Big[ a^2 \frac{d}{da} (a'') + aa'' - 2a'^2 + 2\kappa \Big],$$
(8)

where M is an integration constant, further (8) can be simplified as

$$(a'^2)_{xx} - 4a'^2 + 4\kappa = -\frac{M}{a'},\tag{9}$$

where  $x = \ln a$  and subscript x denotes derivative with respect to x. The constants are usually determined by the initial condition on the variable and its derivative, however, one may have a prior information depending on the configuration of the physical system. Since we are seeking wormhole configuration, one should have a' = 0 and a'' > 0 at some  $\tau$  for existence of non-vanishing minimum of  $a(\tau)$  at the throat of the wormhole. Now in view of the wormhole configuration a' = 0, a'' > 0 at the throat, so equation (8) yields M = 0 without loss of generality. So integration of (9) for M = 0 yields

$$a'^2 = \kappa + c \ a^2 + \frac{c_1}{a^2},\tag{10}$$

where c and  $c_1$  are constants. It is important to note that  $\frac{c_1}{a^2}$  term in (10) is dominant in the very early universe, while the term  $ca^2$  has a large contribution in the late epoch when  $a(\tau)$  is sufficiently large. Further the signature of c and  $c_1$  are also important in determining the solution. So the different terms lead to distinct configurations at different epoch in the cosmic evolution. Without specifying any of them *a priori*, we consider solution of (10). Now an integration of (10) leads to

$$a^{2}(\tau) = -\frac{\kappa}{2c} + \frac{r}{4c} e^{2\sqrt{c}\tau} + \frac{(\kappa^{2} - 4cc_{1})}{4cr} e^{-2\sqrt{c}\tau},$$
(11)

where r is an integration constant. The constant r can be obtained imposing  $a'(\tau = 0) = 0$  as one of the conditions applied for obtaining minimum of wormhole at the throat. The equation (11) allows class of solutions depending on the sign of  $\kappa$ , c, r and  $c_1$ . In the forthcoming section, we classify them according to the values of  $\kappa$  (0 and  $\pm 1$ ), further, we specify the details of the solutions depending on c,  $c_1$  and r. To get a clear view and difference among the solutions depending on different signs of above constants we invoke separate symbols for c,  $c_1$  and r in each case. So we consider the piece-wise solutions of (10) according to the weightage of individual terms in the right hand side of (10). Finally, we consider interpretation of the solutions relevant to cosmic evolution invoking idea of the change of geometry from the Euclidean space to the Lorentz space by using  $\tau = it$ .

## **3.1** Possible solutions and cosmic scenario for $\kappa = 0$ :

Now we consider possible solution from equation (11) with a few restriction on c,  $c_1$  and r for  $\kappa = 0$ .

#### **3.1.1** Solution with $c_1 < 0, c > 0$ and r > 0:

The term  $\frac{c_1}{a^2}$  in (10) dominates over other terms in the very early universe, however it does not yield an interesting solution for  $\kappa = 0$ . A wormhole configuration follows from (11) for  $\kappa = 0$  when the constants are restricted to satisfy  $c_1 = -H_0^2 < 0$ ,  $c = h_0^2 > 0$  and r > 0. The extremum of  $a(\tau)$  (i.e. a' = 0) occurs at  $\tau = 0$  for  $r = 2h_0H_0$ . The scale factor is  $a^2(\tau) = a_0^2 \cosh(2h_0\tau)$ , where the radius of the wormhole at the throat is  $a_0 = (\frac{H_0}{h_0})^{\frac{1}{2}}$ . Now analytic continuation with  $\tau = it$  does not yield a valid  $a^2(t)$  for all t.

### **3.1.2** Solution with $c_1 > 0$ , c < 0 and r < 0:

The scale factor (11) satisfying restriction  $c_1 = H_1^2 > 0$ ,  $c = -h_1^2 < 0$  and  $r = -r_1 < 0$  leads to

$$a^{2}(\tau) = \frac{r_{1}}{4h_{1}^{2}} e^{2ih_{1}\tau} + \frac{H_{1}^{2}}{r_{1}} e^{-2ih_{1}\tau},$$

which is an oscillating function in Euclidean time  $\tau$ , and the  $a^2(\tau)$  is not acceptable for some values of  $\tau$ . One may avoid the problem by introducing analytic continuation with  $\tau = it$  which gives

$$a^{2}(t) = \frac{r_{1}}{4h_{1}^{2}} e^{-2h_{1}t} + \frac{H_{1}^{2}}{r_{1}} e^{2h_{1}t},$$
(12)

so it can be interpreted as a wormhole solution, not in Euclidean time, rather a wormhole configuration in Lorentz time t for  $\kappa = 0$ . The non-vanishing minimum of a(t) occurs at t = 0 for  $r_1 = 2h_1H_1$  and the scale factor reduces to  $a^2(t) = \sqrt{\frac{h_1^2}{H_1^2}} \cosh(2h_1t)$ . In this configuration with time t the radius at the throat is  $a_0 = (\frac{h_1}{H_1})^{1/2}$ . Further, the solution also shows an exponential expanding era asymptotically at  $2h_1t >> 1$  emanating from an early Lorentzian wormhole configuration, so a Lorentzian wormhole evolves to an inflationary scenario.

### 3.2 Possible solutions and cosmic scenario for $\kappa = -1$ :

Now we consider possible solutions of (10) and other solutions from (11) with a few restriction on c,  $c_1$  and r for  $\kappa = -1$ .

## **3.2.1** Solution of (10) when $\frac{c_1}{a^2} >> ca^2$ for $c_1 > 0$ :

The term  $\frac{c_1}{a^2}$  in (10) is dominating compared to the term  $ca^2$  in the very early universe, thus neglecting it in the early era we can find approximate solution of (10) for  $\frac{c_1}{a^2} >> ca^2$  as

$$a^{2}(\tau) = H_{1}^{2} - \tau^{2}$$
, under analytic continuation  $\tau = it \implies a^{2}(t) = H_{1}^{2} + t^{2}$ , (13)

where  $c_1 = H_1^2$ . In (13), the scale factor  $a(\tau)$  encounter singularity at  $\tau = \pm H_1$  and the Euclidean solution is valid in the domain  $-H_1 < \tau < H_1$ . Thus considering analytic continuation  $\tau = it$ , we get a wormhole solution, not in the Euclidean time, rather in the Lorentz time. The solution (13) is symmetric with t and the radius of the wormhole at the throat is  $H_1$  at t = 0. Further  $a(t) \approx t$ , at  $t >> H_1$ , so we have a power law expansion. Thus a Lorentzian wormhole evolves to power law expanding phase at  $t >> H_1$ . Further the effect of  $ca^2$  term in (10) will be more pronounced with expansion and one should consider the contribution of other terms in (10) at later epoch.

#### **3.2.2** Solution with c < 0, r < 0 and $c_1 > 0$ :

The equation (11) allows extrema of  $a(\tau)$  at  $\tau = 0$ , wherein a' = 0 with restriction  $c = -h_1^2 < 0$ , r < 0, but  $c_1 = H_1^2 > 0$ , which gives  $r_3 = \sqrt{1 + 4h_1^2H_1^2} > 1$ . The scale factor than reduces to

$$a^{2}(\tau) = \frac{1}{2h_{1}^{2}} \Big[ -1 + r_{3}\cos(2h_{1}\tau) \Big],$$
(14)

where  $r = -r_3 < 0$ . Apparently it shows an oscillating universe in  $\tau$ , but it faces problem with  $a^2(\tau) < 0$ at the minima of  $a^2(\tau)$ . Thus to avoid it we introduce  $\tau = it$ , then (14) gives

$$a^{2}(t) = \frac{1}{2h_{1}^{2}} \Big[ -1 + r_{3} \cosh(2h_{1}t) \Big].$$
(15)

The scale factor (15) represents a Lorentzian wormhole with the radius  $\sqrt{\frac{r_3-1}{2h_1^2}}$  at the throat. In the very early universe when t is small  $(2h_1t < 1)$  (15) reduces to a form similar to (13) as  $a^2(t) \approx \frac{r_3-1}{2h_1^2} + r_3t^2$  with a lowest order contribution and with a re-definition t. Further in the asymptotic regime  $2h_1t >> 1$ , (15) gives an exponential expansion. So one can obtain inflation from an early Lorentzian wormhole. Further continuation of (15) from (14) using analytic continuation  $\tau = it$  can be interpreted as an early oscillating universe (except  $a^2(\tau) < 0$  domain) with Euclidean time  $\tau$  which eventually evolves to an exponential expanding era.

### **3.2.3** Solution with c > 0, r > 0 and $4cc_1 < 1$

The equation (11) also allows a minimum of  $a(\tau)$  at  $\tau = 0$ , wherein a' = 0 with restriction  $c = h_0^2 > 0$ , r > 0 and  $4h_0^2c_1 < 1$ , which gives  $r = \sqrt{1 - 4h_0^2c_1} < 1$ . The scale factor then reduces to

$$a^{2}(\tau) = \frac{1}{2h_{0}^{2}} \Big[ 1 + r \cosh(2h_{0}\tau) \Big],$$
(16)

which gives a wormhole with a radius  $a_0 = \sqrt{\frac{1+r}{2h_0^2}}$  at the throat. The scale factor is quite large at greater  $\tau$  and  $a(\tau)$  increases exponentially with  $\tau$  at  $2h_0\tau >> 1$ , eventually the term  $ca^2$  in (10) dominates over  $\frac{c_1}{a^2}$  at later era. Further analytic continuation of the solution with  $\tau = it$  also yields an oscillating universe in time t with the period  $\frac{1}{2h_0}$ , where non-vanishing lower bound and upper maximum of a(t) are respectively  $\sqrt{\frac{1-r}{2h_0^2}}$  and  $\sqrt{\frac{1+r}{2h_0^2}}$ , provided r < 1. Thus an oscillatory universe in time 't' for  $\kappa = -1$  may be obtained from an Euclidean wormhole with suitable choice of constants.

## **3.2.4** Solution of (10) when $ca^2 >> \frac{c_1}{a^2}$ :

The solution of (10) when the scale factor is sufficiently large, or with the condition  $ca^2 >> \frac{c_1}{a^2}$  in (10) for  $\kappa = -1$  gives

$$a(\tau) = \frac{r_2}{2h_0^2} e^{h_0 \tau} + \frac{1}{2r_2} e^{-h_0 \tau},$$
(17)

where  $r_2$  is constant and  $c = h_0^2$ . Now the extremum of  $a(\tau)$  in (17) at  $\tau = 0$  gives  $r_2 = h_0$ . So (17) leads to

$$a(\tau) = \frac{1}{h_0} \cosh(h_0 \tau),\tag{18}$$

which is also a wormhole solution. So all the solutions (13)-(17) for  $\kappa = -1$  yield wormhole configuration. The solution (13) and (15) yield wormhole in the early era with respect to Lorentz time t, while the wormhole solutions (16) and (18) are with respect to Euclidean time $\tau$ , however, evolution from an Euclidean wormhole to an exponential expanding era has not been observed for  $\kappa = -1$ .

## **3.3** Possible solutions and cosmic scenario for $\kappa = 1$ :

In this section we consider all possible solutions of (10) and other results from (11) with a few restriction on c,  $c_1$  and r for  $\kappa = 1$ .

## **3.3.1** Solution of (10) with c = 0, but $c_1 = -H_0^2$ :

In the early universe when  $a(\tau)$  is very small the contribution of  $ca^2$  term is not significant compared to  $\frac{c_1}{a^2}$  in (10), so we can neglect the contribution of  $ca^2$  in the early epoch of evolution. The equation (10) with  $c_1 = -H_0^2$  leads to

$$a^2(\tau) = \tau^2 + H_0^2,\tag{19}$$

which represents a wormhole solution and the radius at the throat of the wormhole is  $H_0$ . The scale factor is symmetric with  $\tau$  and it is increasing with  $\tau$ . Now at sufficiently large scale factor the contribution of  $ca^2$  also dominates in (10). However, at some intermediate stage the contribution of all the terms in (10) play an important role in evolution.

#### **3.3.2** Solution with c > 0, r > 0 and $c_1 < 0$ :

The solution (11) with the restrictions  $c = h_0^2 > 0$ , r > 0 and  $c_1 = -H_0^2 < 0$  shows an extremum of  $a(\tau)$  at  $\tau = 0$  for  $r = \sqrt{1 + 4h_0^2H_0^2} > 1$  and the value of  $a(\tau)$  at the extremum at  $\tau = o$  is  $a_0 = \sqrt{\frac{r-1}{2h_0^2}}$ . Then the scale factor reads as

$$a^{2}(\tau) = \frac{1}{2h_{0}^{2}} \left[ -1 + r \cosh(2h_{0}\tau) \right],$$
(20)

so we have a wormhole solution for  $H_0^2 > 0$ . Again the scale factor is large enough with  $\tau$ , and  $a(\tau)$  increases exponentially with  $\tau$  at  $2h_0\tau >> 1$ , however, at small  $\tau$ , (20) leads  $a^2(\tau) \approx \frac{r-1}{2h_0^2} + r\tau^2$  for  $2h_0\tau < 1$  with lowest order contribution, which agrees with the scenario of the early universe given in (19) with re-definition of  $\tau$ . At large scale factor the contribution  $\frac{c_1}{a^2}$  is very small and  $ca^2$  plays a crucial role in evolution. Before considering it we present another solution with different sign of c,  $c_1$  and r.

### **3.3.3** Solution with c < 0, r < 0 and $c_1 < 0$ :

The equation (11) with the conditions c < 0, r < 0 and  $(1 - 4cc_1) > 0$  or  $c_1 < 0$  reduces to

$$a^{2}(\tau) = \frac{1}{2h_{1}^{2}} + \frac{r_{0}}{4h_{1}^{2}} e^{2ih_{1}\tau} + \frac{(1 - 4h_{1}^{2}H_{0}^{2})}{4h_{1}^{2}r_{0}}e^{-2ih_{1}\tau}.$$
(21)

The solution (21) leads to an oscillating universe in  $\tau$  with a period  $\frac{1}{2h_1}$ , where  $c = -h_1^2 < 0$ ,  $c_1 = -H_0^2 < 0$  and  $r = -r_0 < 0$ . The function  $a^2(\tau)$  attains one of the extrema at  $\tau = 0$ , which yields  $r_0 = \sqrt{1 - 4h_1^2 H_0^2} < 1$ , and it leads to

$$a^{2}(\tau) = \frac{1}{2h_{1}^{2}} \left[ 1 + r_{0} \cos(2h_{1}\tau) \right],$$
(22)

the solution (22) represents an oscillating universe with  $\tau$  satisfying  $a^2(\tau) > 0$  at the minima of the function  $a^2(\tau)$  as  $r_0 < 1$ . So introducing analytic continuation  $\tau = it$  in (22) we get

$$a^{2}(t) = \frac{1}{2h_{1}^{2}} \left[ 1 + r_{0} \cosh(2h_{1}t) \right].$$
(23)

The solution (23) represents a Lorentzian wormhole like earlier (15), and the radius at the throat is  $\sqrt{\frac{r_0+1}{2h_1^2}}$ . This solution also leads to an exponential expanding universe with t at an era  $2h_1t >> 1$ . So an early oscillating wormhole (22) in Euclidean time  $\tau$  evolves to an exponential expansion from (23)

## **3.3.4** Solution of (10) with $c_1 = 0$ , but $c = -h_1^2$ :

In course of evolution when the scale factor is sufficiently large at later epoch of  $\tau$ , we can assume  $\frac{c_1}{a^2} \simeq 0$ , or we can put  $c_1 = 0$ . The scale factor from (10) reduces to

$$a^{2}(\tau) = \frac{1}{h_{1}^{2}} \sin^{2}\left(h_{1}\tau\right),$$
(24)

apart from integration constant. The solution of the scale factor is valid as long as  $a(\tau)h_1 < 1$ , which represents an oscillating universe with  $\tau$  within a maxima  $1/h_1$  (deSitter radius) and vanishing minima radii. Now in the era when  $a(\tau)h_1 > 1$  the solution from (10) reduces to

$$a(\tau) = \frac{1}{h_1} \cosh\left(ih_1\tau\right), \text{ and it reduces to } a(t) = \frac{1}{h_1} \cosh\left(h_1t\right), \tag{25}$$

under analytic continuation with  $\tau = it$ , which yields an expanding universe with the Lorentz time. Further in the asymptotic domain  $t >> \frac{1}{h_1}$ , the universe expands exponentially with t. In this case at large t we have an exponential expansion. The universe before crossing deSitter radius is oscillating with Euclidean time  $\tau$  and after crossing the deSitter radius it enters an era of exponential expansion. This agrees with the idea of [44], however, our results are independent of approximation. Thus to sum up the results for  $\kappa = 1$  we have both the configuration of Euclidean and Lorentzian wormholes in the very early universe, eventually the cosmic scenario evolves to an inflationary era in the asymptotic regime of Lorentz time t.

## **3.4** Interpretation of the solutions:

Interpretation of the piece-wise solutions  $\{(19), (20), (22), (24), (25)\}$  and  $\{(13), (15), (16), (18)\}$  respectively for  $\kappa = 1$  and  $\kappa = -1$  are given in the following way. In a nutshell, we can interpret and bring together these piece-wise solutions of (10) for  $\kappa = 1$  representing cosmic scenario of the universe. Beginning with the solution (19) in the very early era, the subsequent evolution passes through an era described by (20) and approaches to an oscillating universe with  $\tau$  in solution (22) and then (23). Finally, the universe after crossing the deSitter radius enters to an era of faster expansion, given by (25) and asymptotically one can realize an exponential expansion with Lorentz time.

Similarly, we can bring together the piece-wise solutions of (10) for  $\kappa = -1$  as representing the cosmic scenario of an universe beginning with (13) as a Lorentzian wormhole evolving towards the solution (15), which could initiate an exponential expansion in Lorentz time depending on constant  $h_1$ . Thereafter, the evolution pushes in the form of (16) to an Euclidean wormhole depending on the parameter  $h_0$  and finally the evolution continues with (18) at very large scale factor. So the evolution from an Euclidean wormhole to an exponential expanding era can be achieved in  $\kappa = 1$ , which is different from the scenario obtained with  $\kappa = -1$ . The new feature of our solution is that we can achieve exponential expansion naturally in the later epoch beginning from a Euclidean wormhole in the very early universe for  $\kappa = 1$ .

## 4 Discussion

Physics of the pre-inflationary universe is not well understood, as the regime is beyond the realm of Einstein's gravity, rather it is determined by the quantum gravity. The initial cosmic singularity, or other relevant configuration of the universe or the wormhole in the Planck era is thus determined by the quantum theory, wherein the spacetime is assumed to be Euclidean. However, the field theory in the Hyperbolic spacetime with time t can explain the inflationary scenario at the GUT era. Thus to explore Physics one may seek viability of evolution from a wormhole configuration to the inflationary scenario. So we consider a toy model of  $R^2$  gravity to study such evolution or possible transition. We consider  $R^2$  term since the term has a huge contribution in the very early universe as well as it reveals inflation.

We simplify the fourth order Euclidean field equations to a second order field equations introducing a new variable following [52, 54] proposal. Exact solution of the field equations in  $R^2$  gravity allows Euclidean wormhole as well as oscillatory solution of  $a^2(\tau)$  depending on three constants and the curvature parameter  $\kappa$ . The oscillatory solutions of  $a^2(\tau)$  are singular with  $\tau$ , however by analytic continuation  $\tau = it$ , the  $a^2(\tau)$ -function reduces to the Lorentzian wormhole with time t. The Lorentzian wormhole gives rise to an exponential expansion asymptotically. A polynomial solution of  $a^2(\tau)$  also reduces to Lorentzian wormhole for  $\kappa = -1$  with use of  $\tau = it$  and expands as a power law inflation in the asymptotic domain. The Euclidean wormhole does not evolve to an era of inflationary solution for all  $\kappa$  using  $\tau = it$ . However, the piece-wise solutions can be brought together to trace the evolution in chronological sequence depending on the relative values of the scale factor. In this context an interesting case appears, wherein an Euclidean wormhole (19) in the very early universe could evolve to an inflationary solution (25) with t at large scale factor only for  $\kappa = 1$ .

Finally, we can state that an Euclidean wormhole (given in (19)) or an oscillating Euclidean wormhole (given in (22) and (24)) in the very early era can evolve to an inflationary era at later epoch under analytic continuation with  $\tau = it$  for  $\kappa = 1$ . In fact the Lorentzian wormhole solutions are able to produce exponential expansion asymptotically in some cases, while in another case power law inflation is followed asymptotically. Thus introduction of wormhole configuration in the very early universe is well justified with onset of inflation at later epoch.

## 5 Appendix

## Null energy condition in our solutions:

In general relativity, wormholes are supported by the exotic matter, which leads to a stress energy tensor (i.e.  $T_{\mu\nu}$ ) that violates null energy condition (NEC). The NEC is given by  $T_{\mu\nu}k^{\mu}k^{\nu} \ge 0$ , where  $k^{\mu}$ is any null vector. The violation of NEC can be minimised by introducing some geometric contribution [9–11], [15,16], etc. instead of exotic field. In our study of  $R^2$  gravity, no such energy momentum tensor exist, however the condition  $R_{\mu\nu}k^{\mu}k^{\nu} \ge 0$  acts as an alternative to the NEC from the Raychaudhuri equation, which is also consistent following the Hawking-Penrose theorem "a space-time is not timelike and null geodesically complete if  $R_{\mu\nu}X^{\mu}X^{\nu} \ge 0$  for every non-spacelike vector  $X^{\mu}$ ". So we evaluate  $R_{\mu\nu}k^{\mu}k^{\nu}$  by contracting the field equation of  $R^2$  gravity with the null vector  $k^{\mu}$  as

$$\left[RR_{\mu\nu} - \frac{R^2}{4}g_{\mu\nu} - R_{;\nu;\mu} + R^{;\alpha}_{;\alpha} g_{\mu\nu}\right]k^{\mu}k^{\nu} = 0,$$

where  $k^{\mu}k_{\mu} = 0$  and a semi-colon stands for covariant derivative, which gives  $R_{\mu\nu}k^{\mu}k^{\nu} = \frac{1}{R}R_{;\nu;\mu}k^{\nu}k^{\mu} = \frac{1}{R}(R_{;\nu;\mu}k^{-\mu}$ 

$$R_{\mu\nu}k^{\mu}k^{\nu} = \frac{k^{0^2}}{R}(R'' - \frac{a'}{a}R'),$$

where we assume  $k^{\mu} = (k^0, k^1, 0, 0)$  satisfying  $(k^0)^2 + g_{11}(k^1)^2 = 0$ ,  $g_{00} = 1$  and  $g_{11} = \frac{a^2(\tau)}{1 - \kappa r^2}$ . Now using (10) and the expression of Ricci scalar *R* we get  $a^2 R'' - aa' R' = 0$ , which yields

$$R_{\mu\nu}k^{\mu}k^{\nu} = 0.$$

So our solutions satisfies NEC, hence the spacetime is not null geodesically complete, i.e., we cannot extend null lines arbitrarily and focusing of null geodesics are possible.

# References

- [1] A Guth, Phys. Rev. **D23** 347 (1981)
- [2] A D Linde, Phys. Lett. **B108** 389 (1982)
- [3] A Albrecht and P J Steinhardt, Phys. Rev. Lett. 48 1220 (1982)
- [4] A Guth and S Y Pi, Phys. Rev. Lett. 49 1110 (1982)
- [5] A Starobinsky, Phys. Lett. **B117** 175 (1982)
- [6] A Starobinsky, Phys. Lett **B** 91, 99 (1980)
- [7] K S Stell, Phys. Rev. **D16** 953 (1977)
- [8] N D Birrel and P C W Davis, Quantum Fields in curved Space (Cambridge University Press, Cambridge, England, 1982)
- [9] S Capozziello, T Harko, T S Koivisto, F S N Lobo and G J Olmo, Phys. Rev. D86 127504 (2012)
- [10] T Harko, F S N Lobo, M K Mak and S V Sushkov, Phys. Rev. D87 067504 (2013)
- [11] F S N Lobo, M A Oliveira, Phys Rev D 80 104012 (2009)
- [12] S Capozziello and R Garattini, Class. Quantum Gravity 24 1627 (2007)
- [13] S Capozziello and M Francaviglia, Gen. Rel.Grav. 40 357 (2008)
- [14] B Modak, K Sarkar and A K Sanyal, Astrophys space Sci 353 707 (2014)
- [15] D J Gross and J H Sloan, Nucl Phys **B291** 41 (1987)
- [16] R R Metsaev and A A Tseytlin, Nucl Phys **B293** 385 (1987)
- [17] D Sahdev, Phys. Lett. **B137** 155 (1984)
- [18] P S Wesson, Astrophysical Journal, 394 19 (1992)
- [19] V Dzhunushaliev and D Singleton, Phys. Rev. **D59** 064018 (1999)
- [20] P Kanti, J Rizos and K Tamvakis, Phys. Rev D 59 083512 (1999)
- [21] I Antoniadis, J Rizos and K Tamvakis, Nucl. Phys. B415 497 (1994)
- [22] J B Hartle and S W Hawking, Phys. Rev. **D28** 2960 (1083)
- [23] S W Hawking and D N Page, Phys. Rev. D 42 2665 (1990)
- [24] M S Morris, K S Thorne, Am J Phys 56 (1988) 395
- [25] M S Morris, K S Thorne and U Yurtsever, Phys Rev Lett.61 1446 (1988)
- [26] K Lee, Phys. Rev. Lett. **61** 263 (1988)
- [27] A Hosoya and W Ogura, Phys. Lett. B 225 117 (1989)
- [28] J Halliwell and R Laflamme, Class. Quantum Grav. 6 1839 (1989)
- [29] D H Coule and K I Maeda, Class. Quantum Grav. 7 955 (1990)
- [30] F S N Lobo and M Oliviera, Phy. Rev D 81 067501 (2010)
- [31] K K Nandi, B Bhattacharjee, S M K Alam and J Evans, Phys Rev D 57 823 (1998)
- [32] S W Hawking, Phys. Rev. D 37 904 (1988)
- [33] S Coleman, Nucl. Phys. B 307, 867 (1988)
- [34] S Coleman, Nucl. Phys. B 310 643 (1988)

- [35] W Fischler and L Susskind, Phys. Lett. B 217 48 (1989)
- [36] J Polchinski, Phys. Lett. B 219 251 (1989)
- [37] W J Unruh, Phys. Rev. D 40 1053 (1989)
- [38] S W Hawking, Nucl. Phys. B 335 155 (1990)
- [39] I Klebanov, L Susskind and T Banks Nucl. Phys. B 317 665 (1989)
- [40] S B Giddings and A Strominger, Nucl. Phys. B 306 890 (1988)
- [41] M Visser, S Kar and N Dadhich, Phys. Rev. Lett. 90 201102 (2003)
- [42] H Maeda, T Harada and B J Carr, Phys. Rev. D79 044034 (2009)
- [43] M Cataldo and S del Campo, Phys. Rev. **D79** 024005 (2009)
- [44] H Fukutaka, K Ghoroku and K Tanaka, Phys. Lett. B 222 191 (1989)
- [45] S Bhattacharya and S Chakroborty, Eur P Journal C77 558 (2017)
- [46] F Darabi, Theor. Math. Phys. 173 1734 (2012)
- [47] A Kehagias, C Kounnas, D Lüst and A Riotto, "Black Hole solutions in  $R^2$  gravity", arXiv:1502.04192 [hep-th], (2015)
- [48] K Sarkar, G Biswas and B Modak, Gen. Rel. Grav 50 157 (2018)
- [49] G Biswas, K Sarkar and B Modak, Mod. Phys. Lett A. 30 16, 2150107 (2021)
- [50] C Kounnas, D Lüst and N Toumbas, " $R^2$  inflation from scale invariant supergravity and anomaly free superstrings with fluxes", arXiv: 1409:7076 [hep-th], (2014)
- [51] S N Sajadi and N Riazi, Prog theo Phys 126 753 (2011)
- [52] D Boulware, A Strominger and E T Tomboulis, in Quantum Theory of Gravity, edited by S Christensen (Adam Hilger, Bristol, 1984)
- [53] G T Horowitz, Phys. Rev. **D31** 1169 (1985)
- [54] A K Sanyal and B Modak, Phys. Rev. D 63 064021 (2001)
- [55] A K Sanyal and B Modak, Class Quantum Grav 19 515 (2002)