

No-Regret Dynamics in the Fenchel Game: A Unified Framework for Algorithmic Convex Optimization

Jun-Kun Wang¹, Jacob Abernethy² and Kfir Y. Levy³

¹Department of Computer Science, Yale University, New Haven, USA.

²School of Computer Science, Georgia Institute of Technology, Atlanta, USA.

³Department of Electrical & Computer Engineering, Technion - Israel Institute of Technology, Haifa, Israel.

Contributing authors: jun-kun.wang@yale.edu; prof@gatech.edu; kfirylevy@technion.ac.il;

Abstract

We develop an algorithmic framework for solving convex optimization problems using no-regret game dynamics. By converting the problem of minimizing a convex function into an auxiliary problem of solving a min-max game in a sequential fashion, we can consider a range of strategies for each of the two-players who must select their actions one after the other. A common choice for these strategies are so-called no-regret learning algorithms, and we describe a number of such and prove bounds on their regret. We then show that many classical first-order methods for convex optimization—including average-iterate gradient descent, the Frank-Wolfe algorithm, the Heavy Ball algorithm, and Nesterov’s acceleration methods—can be interpreted as special cases of our framework as long as each player makes the correct choice of no-regret strategy. Proving convergence rates in this framework becomes very straightforward, as they follow from plugging in the appropriate known regret bounds. Our framework also gives rise to a number of new first-order methods for special cases of convex optimization that were not previously known.

Keywords: Online learning, No-regret learning, Zero-sum game, Convex optimization, Frank-Wolfe method, Nesterov’s accelerated gradient methods, Momentum methods

1 Introduction

The main goal of this work is to develop a framework for solving convex optimization problems using iterative methods. Given a convex function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, domain $K \subset \mathbb{R}^d$, and some tolerance $\epsilon > 0$, we want to find an approximate minimizer $x \in \mathcal{K}$ so that $f(x) - \min_{x' \in \mathcal{K}} f(x') \leq \epsilon$, using a sequence of oracle calls to f and its derivatives. This foundational problem has received attention for decades, and researchers have designed numerous methods for this problem under a range of oracle query models and structural assumptions on $f(\cdot)$. What we aim to show in this paper is that a surprisingly large number of these methods—including those of Nesterov [1–5], Frank and Wolfe [6], Polyak [7], and Beck and Teboulle [8]—can all be described and analyzed through a single unified algorithmic framework, which we call the *Fenchel game no-regret dynamics* (FGNRD). We show that several novel methods, with fast rates, emerge from FGNRD as well.

Let us give a short overview before laying out the FGNRD framework more precisely. A family of tools, largely developed by researchers in theoretical machine learning, consider the problem of sequential prediction and decision making in non-stochastic environments, often called *adversarial online learning*. This online learning setting has found numerous applications in several fields beyond machine learning—finance, for example, as well as statistics—but it has also emerged as a surprisingly useful tool in game theory. What we call *no-regret online learning algorithms* are particularly well-suited for computing equilibria in two-player zero-sum games, as well as solving saddle point problems more broadly. If each agent employs a no-regret online learning algorithm to choose their action at each of a sequence of rounds, it can be shown that the agents’ choices will converge to a saddle point, and at a rate that depends on their choice of learning algorithm. Thus, if we are able to simulate the two agents’ sequential strategies, where each aims to minimize the “regret” of their chosen actions, then what emerges from the resulting *no-regret dynamics* (NRD) can be implemented explicitly as an algorithm for solving min-max problems.

How does NRD help us to develop and analyze methods for minimizing a convex f , i.e. solving $\min_{x \in \mathcal{K}} f(x)$.

$$g(x, y) := \langle x, y \rangle - f^*(y).$$

We view this as a game in the sense that if one player selects an action x and a second player selects action y , then $g(x, y)$ is the former’s “cost” and the latter’s “gain” associated to their decisions. If the two players continue to update their decisions sequentially, first choosing x_1 and y_1 then x_2 and y_2 , etc., and each player relies on some no-regret algorithm for this purpose, then one can show that the time-averaged iterates \bar{x}, \bar{y} form an approximate equilibrium of the Fenchel game—that is, $g(\bar{x}, y') - \epsilon \leq g(\bar{x}, \bar{y}) \leq g(x', \bar{y}) + \epsilon$ for any alternative x', y' . But indeed, this approximate equilibrium brings us right back to where we started, since using the construction of the Fenchel

game it is easy to show that \bar{x} then satisfies $f(\bar{x}) - \min_{x \in \mathcal{K}} f(x) \leq \epsilon$. The approximation factor ϵ is important, and we will see that it depends upon the number of iterations of the dynamic and the players' strategies.

What FGNRD gives us is a recipe book for constructing and analyzing iterative algorithms for convex optimization. To simulate a dynamic we still need to make particular choices as for both players' strategies and analyze their performance. We begin in Section 3 by giving a brief overview of tools from adversarial online learning, and we introduce a handful of simple online learning algorithms, including variants of FollowTheLeader and OnlineMirrorDescent, and prove bounds on the *weighted* regret—we generalize slightly the notion of regret by introducing weights $\alpha_t > 0$ for each round. We prove a key result that relates the error ϵ of the approximate equilibrium pair \bar{x}, \bar{y} , which are the weighted-average of the iterates of the two players, to the weighted regret of the players' strategies. In Section 4 we show how several algorithms, including the Heavy Ball method [7], Frank-Wolfe's method [6], and several variants of Nesterov Accelerated Gradient Descent [1–5, 8], are all special cases of the FGNRD framework, all with special choices of the learning algorithms for the x - and y -players, and the weights α_t ; see Table 1 for a summary of these recipes. In addition we provide several new algorithms using FGNRD in Section 5, summarized in Table 2.

The work presented in this manuscript combines results from our previous publications, [9–11], with a significantly cleaner and more general analysis. But we also include a number of new algorithms and convergence guarantees, described in Algorithm 8, Algorithm 9, Algorithm 18, Algorithm 19, and Algorithm 20, and Theorem 6, Theorem 7, Theorem 20, Theorem 21, Theorem 22. Moreover, for a family of online learning strategies listed in Algorithm 4, we develop a meta lemma in this work, Lemma 13, such that the guarantee of each online algorithm is obtained by an easy application of the meta lemma.

2 Preliminaries

2.1 Convex Analysis

We summarize some results in convex analysis that will be used in this paper. We also refer the readers to some excellent textbooks (e.g. [5, 14–18]).

Smoothness and strong convexity

A function $f(\cdot)$ on \mathbb{R}^d is L -smooth with respect to a norm $\|\cdot\|$ if $f(\cdot)$ is everywhere differentiable and it has Lipschitz continuous gradient $\|\nabla f(x) - \nabla f(z)\|_* \leq L\|x - z\|$, where $\|\cdot\|_*$ denotes the dual norm. A function $f(\cdot)$ is μ -strongly convex w.r.t. a norm $\|\cdot\|$ if the domain of $f(\cdot)$ is convex and that $f(\theta x + (1 - \theta)z) \leq \theta f(x) + (1 - \theta)f(z) - \frac{\mu}{2}\theta(1 - \theta)\|x - z\|^2$ for all $x, z \in \text{dom}(f)$ and $\theta \in [0, 1]$. If a function is μ -strongly convex, then $f(z) \geq f(x) + \partial f(x)^\top(z - x) + \frac{\mu}{2}\|z - x\|^2$ for all $x, z \in \text{dom}(f)$, where $\partial f(x)$ denotes a subgradient of f at x .

Table 1 Summary of recovering existing optimization algorithms from *Fenchel Game*. Here T denotes the total number of iterations, α_t are the weights which set the emphasis on iteration t , the last two columns on the table indicate the specific strategies of the players in the FGNRD.

Non-smooth convex optimization: $\min_w f(w)$ ($G := \max_{w \in \mathcal{K}} \ \partial f(w)\ ^2$ and $R := \ w_0 - w^*\ $) as a game $g(x, y) := \langle x, y \rangle - f^*(y)$.				
Algorithm	rate	weights	y-player	x-player
Gradient Descent with averaging	Thm. 7 $O\left(\frac{GR}{\sqrt{T}}\right)$	$\alpha_t = 1$	Sec. 3.3.1 BESTRESP ⁺	Sec. 3.3.7 OMD
<i>L</i> -smooth convex optimization: $\min_w f(w)$ as a game $g(x, y) := \langle x, y \rangle - f^*(y)$.				
Algorithm	rate	weights	y-player	x-player
Frank-Wolfe method [6]	Prop. 3 and 4 $O\left(\frac{L}{T}\right)$	$\alpha_t = t$	Sec. 3.3.2 FTL	Sec. 3.3.1 BESTRESP ⁺
Linear rate FW [12]	Thm. 5 $O\left(\exp\left(-\frac{\lambda T}{L}\right)\right)$	$\alpha_t = \frac{1}{\ \ell_t(x_t)\ ^2}$	Sec. 3.3.2 FTL	Sec. 3.3.1 BESTRESP ⁺
Nesterov's 1-memory method [4]	Thm. 11 $O\left(\frac{L}{T^2}\right)$	$\alpha_t = t$	Sec. 3.3.6 OPTIMISTICFTL	Sec. 3.3.8 OMD ⁺
Nesterov's ∞ -memory method [3]	Thm. 12 $O\left(\frac{L}{T^2}\right)$	$\alpha_t = t$	Sec. 3.3.6 OPTIMISTICFTL	Sec. 3.3.5 FTRL ⁺
Nesterov's first acceleration method [2]	Thm. 11 $O\left(\frac{L}{T^2}\right)$	$\alpha_t = t$	Sec. 3.3.6 OPTIMISTICFTL	Sec. 3.3.8 OMD ⁺ with $\phi(x) = \frac{1}{2}\ x\ _2^2$
Heavy Ball method [7]	Thm. 9 $O\left(\frac{L}{T}\right)$	$\alpha_t = t$	Sec. 3.3.2 FTL	Sec. 3.3.5 FTRL ⁺
Composite optimization: $\min_w f(w) + \psi(w)$, where $\psi(\cdot)$ is possibly non-differentiable, as a game $g(x, y) := \langle x, y \rangle - f^*(y) + \psi(x)$.				
Algorithm	rate	weights	y-player	x-player
Accelerated proximal method [8]	Thm. 13 $O\left(\frac{L}{T^2}\right)$	$\alpha_t = t$	Sec. 3.3.6 OPTIMISTICFTL	Sec. 3.3.8 OMD ⁺
<i>L</i> -smooth and μ strongly convex optimization: $\min_w f(w)$ as a game $g(x, y) := \langle x, y \rangle - \tilde{f}^*(y) + \frac{\mu\ x\ ^2}{2}$, where $\tilde{f}(\cdot) := f(\cdot) - \frac{\mu}{2}\ \cdot\ ^2$.				
Algorithm	rate	α_t	y-player	x-player
Nesterov's method [5]	Thm. 15 $O\left(\exp\left(-\sqrt{\frac{\mu}{L}}T\right)\right)$	$\alpha_t \propto \exp(t)$	Sec. 3.3.6 OPTIMISTICFTL	Sec. 3.3.5 FTRL ⁺
<i>L</i> -smooth convex finite-sum optimization: $\min_{w \in \mathcal{K}} f(w) := \sum_{i=1}^n f_i(w)$ as a game $g(x, y) := \langle x, y \rangle - f^*(y)$				
Algorithm	rate	weights	y-player's alg.	x-player's alg.
Incremental FW [13]	Thm. 6 $O\left(\frac{n}{T}\right)$	$\alpha_t = t$	Appendix. C Lazy FTL	Sec. 3.3.1 BESTRESP ⁺

Convex function and conjugate

For any convex function $f(\cdot)$, its Fenchel conjugate is

$$f^*(y) := \sup_{x \in \text{dom}(f)} \langle x, y \rangle - f(x). \quad (1)$$

If a function $f(\cdot)$ is convex, then its conjugate $f^*(\cdot)$ is also convex, as it is a supremum over linear functions. Furthermore, if the function $f(\cdot)$ is closed and

Table 2 Summary of *new* optimization algorithms from *Fenchel Game*. Here T denotes the total number of iterations, α_t are the weights which set the emphasis on iteration t , the last two columns on the table indicate the specific strategies of the players in the FGNRD.

Non-smooth convex optimization: $\min_{w \in \mathcal{K}} f(w)$, where \mathcal{K} is a λ -strongly convex set as a game $g(x, y) := \langle x, y \rangle - f^*(y)$,				
Assume that the norm of cumulative gradient does not vanish, $\ \frac{1}{T} \sum_{s=1}^T \partial f(x_s)\ \geq \rho$.				
Algorithm	rate	weights	y-player's alg.	x-player's alg.
Boundary FW	Thm. 17 $O\left(\frac{1}{\lambda \rho T}\right)$	$\alpha_t = 1$	Sec. 3.3.2 FTL	Sec. 3.3.1 BESTRESP ⁺
L -smooth convex optimization: $\min_{w \in \mathcal{K}} f(w)$, where \mathcal{K} is a λ -strongly convex set that is centrally symmetric and contains the origin, as a game $g(x, y) := \langle x, y \rangle - f^*(y)$				
Algorithm	rate	weights	y-player's alg.	x-player's alg.
Gauge FW	Thm. 19 $O\left(\frac{L}{\lambda T^2}\right)$	$\alpha_t = t$	Sec. 3.3.6 OPTIMISTICFTL	Sec. 3.3.5 FTRL ⁺ with gauge function
L -smooth and μ -strongly convex optimization: $\min_{w \in \mathcal{K}} f(w)$, where \mathcal{K} is a λ -strongly convex set that is centrally symmetric and contains the origin, as a game $g(x, y) := \langle x, y \rangle - \tilde{f}^*(y) + \frac{\mu}{\gamma_{\mathcal{K}}} \gamma_{\mathcal{K}}^2(x)$, where $\tilde{f}(\cdot) := f(\cdot) - \frac{\mu}{\gamma_{\mathcal{K}}} \gamma_{\mathcal{K}}^2(x)$				
Algorithm	rate	weights	y-player's alg.	x-player's alg.
Gauge FW	Thm. 20 $O\left(\exp\left(-\sqrt{\frac{\mu}{L}} T\right)\right)$	$\alpha_t \propto \exp(t)$	Sec. 3.3.6 OPTIMISTICFTL	Sec. 3.3.5 FTRL ⁺ with gauge function
Non-smooth convex optimization: $\min_w f(w)$ ($G := \max_{w \in \mathcal{K}} \ \partial f(w)\ ^2$ and $R := \ w_0 - w^*\ $) as a game $g(x, y) := \langle x, y \rangle - f^*(y)$.				
Algorithm	rate	weights	y-player	x-player
Cumulative GD	Thm. 21 $O\left(\frac{GR}{\sqrt{T}}\right)$	$\alpha_t = 1$	Sec. 3.3.3 FTL ⁺	Sec. 3.3.7 OMD
L -smooth convex optimization: $\min_{w \in \mathcal{K}} f(w)$ as a game $g(x, y) := \langle x, y \rangle - f^*(y)$				
Algorithm	rate	weights	y-player's alg.	x-player's alg.
Optimistic Mirror Descent with averaging	Thm. 22 $O\left(\frac{L}{T^2}\right)$	$\alpha_t = t$	Sec. 3.3.3 FTL ⁺	Sec. 3.3.9 OPTIMISTICMD

convex, the following are equivalent: (I) $y \in \partial f(x)$, (II) $x \in \partial f^*(y)$, and (III)

$$\langle x, y \rangle = f(x) + f^*(y), \quad (2)$$

which also implies that the bi-conjugate is equal to the original function, i.e. $f^{**}(\cdot) = f(\cdot)$. Moreover, when the function $f(\cdot)$ is differentiable, we have $\nabla f(x) = \sup_y \langle x, y \rangle - f^*(y)$. We refer to the readers to [19, 20] for more details of Fenchel conjugate. Throughout this paper, unless specifically mentioned, we assume that the underlying convex function is proper, closed, and differentiable.

An important property of a closed and convex function is that $f(\cdot)$ is L -smooth w.r.t. some norm $\|\cdot\|$ if and only if its conjugate $f^*(\cdot)$ is $1/L$ -strongly convex w.r.t. the dual norm $\|\cdot\|_*$ (e.g. Theorem 6 in [20]).

Bregman Divergence.

We will denote the Bregman divergence $D_z^\phi(\cdot)$ centered at a point z with respect to a β -strongly convex distance generating function $\phi(\cdot)$ as

$$D_z^\phi(x) := \phi(x) - \langle \nabla \phi(z), x - z \rangle - \phi(z). \quad (3)$$

Strongly convex sets.

A convex set $\mathcal{K} \subseteq \mathbb{R}^m$ is an λ -strongly convex set w.r.t. a norm $\|\cdot\|$ if for any $x, z \in \mathcal{K}$, any $\theta \in [0, 1]$, the $\|\cdot\|$ ball centered at $\theta x + (1 - \theta)z$ with radius $\theta(1 - \theta)\frac{\lambda}{2}\|x - z\|^2$ is included in \mathcal{K} [21]. Examples of strongly convex sets include ℓ_p balls: $\|x\|_p \leq r, \forall p \in (1, 2]$, Schatten p balls: $\|\sigma(X)\|_p \leq r$ for $p \in (1, 2]$, where $\sigma(X)$ is the vector consisting of singular values of the matrix X , and Group (s,p) balls: $\|X\|_{s,p} = \|(\|X_1\|_s, \|X_2\|_s, \dots, \|X_m\|_s)\|_p \leq r, \forall s, p \in (1, 2]$. In Appendix F, we discuss more about strongly-convex sets.

2.2 Min-max Problems and the Fenchel Game**Min-max problems and (approximate) Nash equilibrium**

Given a zero-sum game with *payoff function* $g(x, y)$ which is convex in x and concave in y , define $V^* = \inf_{x \in \mathcal{K}} \sup_y g(x, y)$. An ϵ -equilibrium of $g(\cdot, \cdot)$ is a pair \hat{x}, \hat{y} such that

$$V^* - \epsilon \leq \inf_{x \in \mathcal{K}} g(x, \hat{y}) \leq V^* \leq \sup_y g(\hat{x}, y) \leq V^* + \epsilon. \quad (4)$$

The Fenchel Game.

One of the core tools of this paper is as follows. In order to solve the problem

$$\min_{x \in \mathcal{K}} f(x) \quad (5)$$

we instead construct a saddle-point problem which we call the *Fenchel Game*. We define $g : \mathcal{K} \times \mathbb{R}^d$ as follows:

$$g(x, y) := \langle x, y \rangle - f^*(y). \quad (6)$$

This payoff function is useful for solving the original optimization problem, since an equilibrium of this game provides us with a solution to $\min_{x \in \mathcal{K}} f(x)$. Let \hat{x}, \hat{y} be any equilibrium pair of g , with $\hat{x} \in \mathcal{K}$. that is, where $V^* = \sup_y g(\hat{x}, y)$. Then we have

$$\begin{aligned} \inf_{x \in \mathcal{K}} f(x) &= \inf_{x \in \mathcal{K}} \sup_y \{\langle x, y \rangle - f^*(y)\} = \inf_{x \in \mathcal{K}} \sup_y g(x, y) \\ &= \sup_y g(\hat{x}, y) = \sup_y \{\langle \hat{x}, y \rangle - f^*(y)\} = f(\hat{x}). \end{aligned}$$

In other words, given an equilibrium pair \hat{x}, \hat{y} of $g(\cdot, \cdot)$, we immediately have a minimizer of $f(\cdot)$. This simple observation can be extended to approximate equilibria as well.

Lemma 1 *If (\hat{x}, \hat{y}) is an ϵ -equilibrium of the Fenchel Game (6), then $f(\hat{x}) - \min_x f(x) \leq \epsilon$.*

Lemma 1 sets us up for the remainder of the paper. The framework, which we lay out precisely in Section 3.2, will consider two players sequentially playing the Fenchel game, where the y -player sequentially outputs iterates y_1, y_2, \dots , while alongside the x -player returns iterates x_1, x_2, \dots . Each player may use the previous sequence of actions of their opponent in order to choose their next point x_t or y_t , and we will rely heavily on the use of no-regret online learning algorithms described in Section 3. In addition, we need to select a sequence of weights $\alpha_1, \alpha_2, \dots > 0$ which determine the “strength” of each round, and can affect the players’ update rules. What we will be able to show is that the α -weighted average iterate pair, defined as

$$(\hat{x}, \hat{y}) := \left(\frac{\alpha_1 x_1 + \dots + \alpha_T x_T}{\alpha_1 + \dots + \alpha_T}, \frac{\alpha_1 y_1 + \dots + \alpha_T y_T}{\alpha_1 + \dots + \alpha_T} \right),$$

is indeed an ϵ -equilibrium of $g(\cdot, \cdot)$, and thus via Lemma 1 we have that \hat{x} approximately minimizes f . To get a precise estimate of ϵ requires us to prove a family of regret bounds, which is the focus of the following section.

3 No-regret learning algorithms

An algorithmic framework, often referred to as *no-regret learning* or *online convex optimization*, has been developed mostly within the machine learning research community, has grown quite popular as it can be used in a broad class of sequential decision problems. As we explain in Section 3.1, one imagines an algorithm making repeated decisions by selecting a vector of parameters in a convex set, and on each round is charged according to a varying convex loss function. The algorithm’s goal is to minimize an objective known as regret. In Section 3.2 we describe how online convex optimization algorithms with vanishing regret can be implemented in a two-player protocol which sequentially computes an approximate equilibria for a convex-concave payoff function. This is the core tool that allows us to describe a range of known and novel algorithms for convex optimization, by modularly combining pairs of OCO strategies. In Section 3.3 we provide several such OCO algorithms, most of which have been proposed and analyzed over the past 10-20 years.

Protocol 1 Weighted Online Convex Optimization

```

1: Input: decision set  $\mathcal{Z} \subset \mathbb{R}^n$ 
2: Input: number of rounds  $T$ 
3: Input: weights  $\alpha_1, \alpha_2, \dots, \alpha_T > 0$  // Weights determined in advance
4: Input: algorithm OAlg // This implements the learner's update strategy
5: for  $t = 1, 2, \dots$ , do
6:   Return:  $z_t \leftarrow$  OAlg // Alg returns a point  $z_t$ 
7:   Receive:  $\alpha_t, \ell_t(\cdot) \rightarrow$  OAlg // Alg receives loss fn. and round weight
8:   Evaluate: Loss  $\leftarrow$  Loss +  $\alpha_t \ell_t(z_t)$  // Alg suffers weighted loss for choice of  $z_t$ 
9: end for

```

3.1 Online Convex Optimization and Regret

Here we describe the framework, given precisely in Protocol 1, for online convex optimization. We assume we have some learning algorithm known as OAlg that is tasked with selecting “actions” from a compact and convex *decision set* $\mathcal{Z} \subset \mathbb{R}^d$. On each round $t = 1, \dots, T$, OAlg returns a point $z_t \in \mathcal{Z}$, and is then presented with the pair α_t, ℓ_t , where $\alpha_t > 0$ is a weight for the current round and $\ell_t : \mathcal{Z} \rightarrow \mathbb{R}$ is a convex loss function that evaluates the choice z_t . While OAlg is essentially forced to “pay” the cost $\alpha_t \ell_t(z_t)$, it can then update its state to provide better choices in future rounds.

On each round t , the learner must select a point $z_t \in \mathcal{Z}$, and is then “charged” a loss of $\alpha_t \ell_t(z_t)$ for this choice. Typically it is assumed that, when the learner selects z_t on round t , she has observed all loss functions $\alpha_1 \ell_1(\cdot), \dots, \alpha_{t-1} \ell_{t-1}(\cdot)$ up to, but not including, time t . However, we will also consider learners that are *prescient*, i.e. that can choose z_t with knowledge of the loss functions up to *and including* time t . The objective of interest in most of the online learning literature is the learner’s *regret*, defined as

$$\alpha\text{-REG}^z := \sum_{t=1}^T \alpha_t \ell_t(z_t) - \min_{z \in \mathcal{Z}} \sum_{t=1}^T \alpha_t \ell_t(z). \quad (7)$$

Oftentimes we will want to refer to the *average regret*, or the regret normalized by the time weight $A_T := \sum_{t=1}^T \alpha_t$, which we will denote $\bar{\alpha}\text{-REG}^z := \frac{\alpha\text{-REG}^z}{A_T}$. Note that in online learning literature, What has become a cornerstone of online learning research has been the existence of *no-regret algorithms*, i.e. learning strategies that guarantee $\bar{\alpha}\text{-REG}^x \rightarrow 0$ as $A_T \rightarrow \infty$.

Let us consider some very simple learning strategies that will be used in this paper, and we note the available guarantees for each. We also refer the readers to some tutorial of online learning for more online learning algorithms. (see e.g. [22–25])

3.2 Framework: optimization as *Fenchel Game*

In this paper, we consider Fenchel game (6) with weighted losses depicted in Algorithm 2. In this game, the y -player plays before the x -player plays and the x -player sees what the y -player plays before choosing its action. The y -player receives loss functions $\alpha_t \ell_t(\cdot)$ in round t , in which $\ell_t(y) := f^*(y) - \langle x_t, y \rangle$, while the x -player see its loss functions $\alpha_t h_t(\cdot)$ in round t , in which $h_t(x) := \langle x, y_t \rangle - f^*(y_t)$. Consequently, we can define the *weighted regret* of the x and y players as

$$\alpha\text{-REG}^y := \sum_{t=1}^T \alpha_t \ell_t(y_t) - \min_y \sum_{t=1}^T \alpha_t \ell_t(y) \quad (8)$$

$$\alpha\text{-REG}^x := \sum_{t=1}^T \alpha_t h_t(x_t) - \min_x \sum_{t=1}^T \alpha_t \ell_t(x). \quad (9)$$

Protocol 2 Fenchel Game No-Regret Dynamics

```

1: Input: number of rounds  $T$ 
2: Input: Convex decision sets  $\mathcal{X}, \mathcal{Y} \subset \mathbb{R}^d$ 
3: Input: Convex-concave payoff function  $g : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ 
4: Input: weights  $\alpha_1, \alpha_2, \dots, \alpha_T > 0$  // Weights determined in advance
5: Input: algorithms  $\text{OAlg}^Y, \text{OAlg}^X$  // Learning algorithms for both players
6: for  $t = 1, 2, \dots, T$  do
7:   Return:  $y_t \leftarrow \text{OAlg}^Y$  //  $y$ -player returns a point  $y_t$ 
8:   Update:  $\alpha_t, h_t(\cdot) \rightarrow \text{OAlg}^X$  //  $x$ -player updates with  $\alpha_t$  and loss  $-g(\cdot, y_t)$ 
9:     where  $h_t(\cdot) := -g(\cdot, y_t)$ 
10:  Return:  $x_t \leftarrow \text{OAlg}^X$  //  $x$ -player returns a point  $x_t$ 
11:  Update:  $\alpha_t, \ell_t(\cdot) \rightarrow \text{OAlg}^Y$  //  $y$ -player updates with  $\alpha_t$  and loss  $g(x_t, \cdot)$ 
12:    where  $\ell_t(\cdot) := g(x_t, \cdot)$ 
13: end for
14: Output  $(\bar{x}_T, \bar{y}_T) := \left( \frac{\sum_{s=1}^T \alpha_s x_s}{A_T}, \frac{\sum_{s=1}^T \alpha_s y_s}{A_T} \right)$ .
```

At times when we want to refer to the regret on another sequence y'_1, \dots, y'_T we may refer to this as $\alpha\text{-REG}(y'_1, \dots, y'_T)$. We also denote A_t as the cumulative sum of the weights $A_t := \sum_{s=1}^t \alpha_s$ and the weighted average regret $\overline{\alpha\text{-REG}} := \frac{\alpha\text{-REG}}{A_T}$. Finally, for offline constrained optimization (i.e. $\min_{x \in \mathcal{K}} f(x)$), we let the decision space of the benchmark/comparator in the weighted regret definition to be $\mathcal{X} = \mathcal{K}$; for offline unconstrained optimization, we let the decision space of the benchmark/comparator to be a norm ball that contains the optimum solution of the offline problem (i.e. contains $\operatorname{argmin}_{x \in \mathbb{R}^n} f(x)$), which means that \mathcal{X} of the comparator is a norm ball.

Theorem 1 Assume a T -length sequence α are given. Suppose in Protocol 2 the online learning algorithms $OAlg^x$ and $OAlg^y$ have the α -weighted average regret $\overline{\alpha\text{-REG}}^x$ and $\overline{\alpha\text{-REG}}^y$ respectively. Then the output (\bar{x}_T, \bar{y}_T) is an ϵ -equilibrium for $g(\cdot, \cdot)$, with $\epsilon = \overline{\alpha\text{-REG}}^x + \overline{\alpha\text{-REG}}^y$. Moreover, if $f(x) := \sup_{y \in \mathcal{Y}} g(x, y)$, then it follows that

$$f(\bar{x}_T) - \min_{x \in \mathcal{X}} f(x) \leq \overline{\alpha\text{-REG}}^x + \overline{\alpha\text{-REG}}^y.$$

Proof Suppose that the loss function of the x -player in round t is $\alpha_t h_t(\cdot) : \mathcal{X} \rightarrow \mathbb{R}$, where $h_t(\cdot) := g(\cdot, y_t)$. The y -player, on the other hand, observes her own sequence of loss functions $\alpha_t \ell_t(\cdot) : \mathcal{Y} \rightarrow \mathbb{R}$, where $\ell_t(\cdot) := -g(x_t, \cdot)$.

$$\begin{aligned} \frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T \alpha_t g(x_t, y_t) &= \frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T -\alpha_t \ell_t(y_t) \\ &\geq -\frac{1}{\sum_{s=1}^T \alpha_s} \inf_{y \in \mathcal{Y}} \left(\sum_{t=1}^T \alpha_t \ell_t(y) \right) - \frac{\alpha\text{-REG}^y}{\sum_{s=1}^T \alpha_s} \\ &= \sup_{y \in \mathcal{Y}} \left(\frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T \alpha_t g(x_t, y) \right) - \overline{\alpha\text{-REG}}^y \\ \text{(Jensen)} &\geq \sup_{y \in \mathcal{Y}} g \left(\frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T \alpha_t x_t, y \right) - \overline{\alpha\text{-REG}}^y \\ &= \sup_{y \in \mathcal{Y}} g(\bar{x}_T, y) - \overline{\alpha\text{-REG}}^y \end{aligned} \tag{10}$$

$$\geq \inf_{x \in \mathcal{X}} \sup_{y \in \mathcal{Y}} g(x, y) - \overline{\alpha\text{-REG}}^y. \tag{11}$$

Let us now apply the same argument on the right hand side, where we use the x -player's regret guarantee.

$$\begin{aligned} \frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T \alpha_t g(x_t, y_t) &= \frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T \alpha_t h_t(x_t) \\ &\leq \inf_{x \in \mathcal{X}} \left(\sum_{t=1}^T \frac{1}{\sum_{s=1}^T \alpha_s} \alpha_t g(x, y_t) \right) + \frac{\alpha\text{-REG}^x}{\sum_{s=1}^T \alpha_s} \\ &= \inf_{x \in \mathcal{X}} \left(\sum_{t=1}^T \frac{1}{\sum_{s=1}^T \alpha_s} \alpha_t g(x, y_t) \right) + \overline{\alpha\text{-REG}}^x \\ \text{(Jensen)} &\leq \inf_x g \left(x, \sum_{t=1}^T \frac{1}{\sum_{s=1}^T \alpha_s} \alpha_t y_t \right) + \overline{\alpha\text{-REG}}^x \\ &= \inf_x g(x, \bar{y}_T) + \overline{\alpha\text{-REG}}^x \end{aligned} \tag{12}$$

$$\leq \sup_{y \in \mathcal{Y}} \inf_{x \in \mathcal{X}} g(x, y) + \overline{\alpha\text{-REG}}^x. \tag{13}$$

By (13) and (10), we have $\sup_{y \in \mathcal{Y}} g(\bar{x}_T, y) \leq V^* + \overline{\alpha\text{-REG}}^x + \overline{\alpha\text{-REG}}^y$. By (12) and (11), we have $V^* - \overline{\alpha\text{-REG}}^x - \overline{\alpha\text{-REG}}^y \leq \inf_{x \in \mathcal{X}} g(x, \bar{y}_T)$. Thus (\bar{x}_T, \bar{y}_T) is an $\epsilon = \overline{\alpha\text{-REG}}^x + \overline{\alpha\text{-REG}}^y$ equilibrium.

We also have

$$\begin{aligned}
 \frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T \alpha_t g(x_t, y_t) &= \frac{1}{\sum_{s=1}^T \alpha_s} \sum_{t=1}^T \alpha_t h_t(x_t) \\
 &\leq \left(\sum_{t=1}^T \frac{1}{\sum_{s=1}^T \alpha_s} \alpha_t h_t(x^*) \right) + \frac{\alpha\text{-REG}^x}{\sum_{s=1}^T \alpha_s} \quad (14) \\
 &= \left(\sum_{t=1}^T \frac{1}{\sum_{s=1}^T \alpha_s} \alpha_t g(x^*, y_t) \right) + \overline{\alpha\text{-REG}}^x \\
 (\text{Jensen}) &\leq g \left(x^*, \sum_{t=1}^T \frac{1}{\sum_{s=1}^T \alpha_s} \alpha_t y_t \right) + \overline{\alpha\text{-REG}}^x \\
 &= g(x^*, \bar{y}_T) + \overline{\alpha\text{-REG}}^x \\
 &\leq \sup_{y \in \mathcal{Y}} g(x^*, y) + \overline{\alpha\text{-REG}}^x, \quad (15)
 \end{aligned}$$

where (14) is because $\sum_{t=1}^T \alpha_t h_t(x_t) - \sum_{t=1}^T \alpha_t h_t(x^*) \leq \sum_{t=1}^T \alpha_t h_t(x_t) - \min_{x \in \mathcal{X}} \sum_{t=1}^T \alpha_t h_t(x) := \alpha\text{-REG}^x$. Combining (10) and (15), we see that $f(\bar{x}_T) - \min_{x \in \mathcal{X}} f(x) \leq \overline{\alpha\text{-REG}}^x + \overline{\alpha\text{-REG}}^y$. \square

In order to utilize minimax duality, we have to define decision sets for two players, and we must produce a convex-concave payoff function. First we will assume, for convenience, that $f(x) := \infty$ for any $x \notin \mathcal{X}$. That is, it takes the value ∞ outside of the convex/compact set \mathcal{X} , which ensures that $f(\cdot)$ is lower semi-continuous and convex. We let the y -player be given the set $\mathcal{Y} := \{\partial f(x) : x \in \mathcal{X}\}$. One can check that the closure of the set \mathcal{Y} is a convex set. Appendix A describes the proof.

Theorem 2 *The closure of (sub-)gradient space $\{\partial f(x) : x \in \mathcal{X}\}$ is a convex set.*

Faster rates for min-max problems.

We conclude this section by noting that many of the accelerated rates that we prove in this work emerge from a new set of tools that have been developed within the past decade. The use of no-regret learning to solve zero-sum games, as described in Theorem 1, was popularized by Freund and Schapire [26] in 1996, although the idea goes back to Blackwell [27] and Hannan [28] in the 1950s. While this basic tool has been used in many settings and applications, a key trick was discovered by Rakhlin and Sridharan [29] in 2013, showing that the rate for solving a simple matrix game could be improved from $O(1/\sqrt{T})$ to $O(1/T)$ via a more sophisticated type of algorithm which we herein refer to as *optimistic*; this will be explained precisely in Section 3.3. The first known presentation of such an algorithm was given by [30], who showed that a better

regret bound is achievable if the sequence of loss functions changes slowly. As explained in [29], this is very useful when used in Protocol 2. This new trick is a core piece of many of the tools presented in this paper, and we emphasize that all of the *accelerated* optimization algorithms we describe through the FGNRD framework rely on one of the players using an optimistic online learning algorithm.

3.3 Online Convex Optimization: An Algorithmic Menu

In this section we introduce and analyze several core online learning algorithms. Later in Sections 4 & 5, we will show how composing different online learning algorithm within the Fenchel Game No-Regret Dynamics (Protocol 2) enables to easily recover known results and methods for convex optimization (Section 4), as well as to design new algorithm with novel guarantees (Section 5).

We present two families of online learning strategies: the “batch-style” family of algorithms, and the “update” family, although we note that one can relate these two. Within each family, we have algorithms that require a regularization function $R(\cdot)$, or a “link” or “distance-generating” function $\phi(\cdot)$. In some cases, the algorithm may peek at the current loss function ℓ_t before selecting z_t (these are denoted with a superscript +), or they may only have access to $\ell_1, \dots, \ell_{t-1}$. We also present “optimistic” variants, where the algorithm can not see ℓ_t but can try to “guess” this function with some m_t , and use this guess in it’s optimization to compute z_t . We note, importantly, that we have given different names to these algorithms, keeping up with historical precedents in the online learning community, but we observe that with the appropriate parameters every one of the batch-style algorithms is in fact a special case of OPTIMISTICFTRL, which is in some sense the Master Algorithm. We utilize this fact when proving regret bounds, as a Master Bound is provided in Lemma 13 of Subsection 3.4, and nearly all of the remaining bounds of the “batch-style” family of algorithms in Algorithm 4 follow as an easy corollary. Therefore, in the following, we describe the online learning strategies with their regret bounds first, and we defer the proofs in Subsection 3.4.

Before diving into the two families, we begin with one simple algorithm BESTRESP⁺ that stands as something of an outlier.

Algorithm 3 BESTRESP⁺ - Best Response

Parameters: convex set \mathcal{Z}
 Receives: $\alpha_1, \dots, \alpha_T > 0, \ell_1, \dots, \ell_T : \mathcal{Z} \rightarrow \mathbb{R}$
 Action: $z_t \leftarrow \underset{z \in \mathcal{Z}}{\operatorname{argmin}} \{ \ell_t(z) \}$

Algorithm 4 A family of batch-style online learning strategies

Parameters: convex set \mathcal{Z} , initial point $z_{\text{init}} \in \mathcal{Z}$

Receives: $\alpha_1, \dots, \alpha_T > 0, \ell_1, \dots, \ell_T : \mathcal{Z} \rightarrow \mathbb{R}$

FTRL: $z_t \leftarrow z_{\text{init}}$ if $t = 1$ otherwise (16)

$$z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\sum_{s=1}^{t-1} \alpha_s \ell_s(z) \right)$$

FTRL⁺: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\sum_{s=1}^t \alpha_s \ell_s(z) \right)$ (17)

FTRL[$R(\cdot), \eta$]: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\sum_{s=1}^{t-1} \alpha_s \ell_s(z) + \frac{1}{\eta} R(z) \right)$ (18)

FTRL⁺[$R(\cdot), \eta$]: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\sum_{s=1}^t \alpha_s \ell_s(z) + \frac{1}{\eta} R(z) \right)$ (19)

OPTIMISTICFTRL: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_t m_t(z) + \sum_{s=1}^{t-1} \alpha_s \ell_s(z) \right)$ (20)

Needs “guesses” $m_1, \dots, m_T : \mathcal{Z} \rightarrow \mathbb{R}$

OPTIMISTICFTRL[$R(\cdot), \eta$]: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_t m_t(z) + \sum_{s=1}^{t-1} \alpha_s \ell_s(z) + \frac{R(z)}{\eta} \right)$ (21)

Requires “guesses” $m_1, \dots, m_T : \mathcal{Z} \rightarrow \mathbb{R}$

Algorithm 5 A family of update-style online learning strategies

Parameters: convex set \mathcal{Z} , initial point $z_0 = z_{-\frac{1}{2}} \in \mathcal{Z}$

Receives: $\alpha_1, \dots, \alpha_T > 0, \ell_1, \dots, \ell_T : \mathcal{Z} \rightarrow \mathbb{R}$

Set: $\alpha_0 := 0, \ell_0(\cdot) := 0$

OMD[$\phi(\cdot), z_0, \gamma$]: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_{t-1} \ell_{t-1}(z) + \frac{1}{\gamma} D_{z_{t-1}}^\phi(z) \right)$ (22)

OMD⁺[$\phi(\cdot), z_0, \gamma$]: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_t \ell_t(z) + \frac{1}{\gamma} D_{z_{t-1}}^\phi(z) \right)$ (23)

OPTIMISTICMD[$\phi(\cdot), z_0, \gamma$]: $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_t \langle z, m_t \rangle + \frac{1}{\gamma} D_{z_{t-\frac{1}{2}}}^\phi(z) \right)$

$$z_{t+\frac{1}{2}} \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_t \langle z, \nabla \ell_t(z_t) \rangle + \frac{1}{\gamma} D_{z_{t-\frac{1}{2}}}^\phi(z) \right)$$

Requires “guesses” $m_1, \dots, m_T : \mathcal{Z} \rightarrow \mathbb{R}$

3.3.1 Best Response – BESTRESP⁺

Perhaps the most trivial strategy for a prescient learner is to ignore the history of the ℓ_s 's, and simply play the best choice of z_t on the current round. We call this algorithm BESTRESP⁺. The ability to see the cost of your decisions in advance guarantees, quite naturally, that you will not suffer positive regret. We note that algorithms with a + superscript may “peek” at the loss ℓ_t before choosing z_t .

Lemma 2 (BESTRESP⁺) *For any sequence of loss functions $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$, if z_1, \dots, z_T is chosen according to BESTRESP⁺ ensures,*

$$\overline{\alpha\text{-REG}}^z := \sum_{t=1}^T \alpha_t \ell_t(z_t) - \min_{z \in \mathcal{Z}} \sum_{t=1}^T \alpha_t \ell_t(z) \leq 0. \quad (24)$$

Proof Since $z_t = \operatorname{argmin}_{z \in \mathcal{Z}} \ell_t(z)$, we have $\ell_t(z_t) \leq \ell_t(z)$ for any $z \in \mathcal{Z}$. The result follows by summing the inequalities from $t = 1, \dots, T$, and recalling that the α_t 's are non-negative. \square

3.3.2 Follow The Leader – FTL

FTL ((16) in Algorithm 4) is perhaps the simplest strategy in online learning, it plays the best fixed action for the cumulative (weighted) loss seen in each previous round. It has been shown in various places (e.g. [22, 25]) that FTL exhibits logarithmic regret, at least when the loss functions are strongly convex. Our proof will ultimately be a consequence of a larger meta-result which we lay out in Section 3.4.

Lemma 3 (FTL[z_{init}]) *Let $\alpha_1, \dots, \alpha_T > 0$ be arbitrary, and assume we have a sequence of μ -strongly convex loss functions $\ell_1(\cdot), \dots, \ell_T(\cdot)$ for some $\mu \geq 0$. Then for any initial point $z_{\text{init}} \in \mathcal{Z}$ the OCO procedure FTL[z_{init}] satisfies*

$$\alpha\text{-REG} \leq \sum_{t=1}^T \frac{2\alpha_t^2}{\left(\sum_{s=1}^t \alpha_s \mu\right)} \|\nabla \ell_t(z_t)\|_*^2. \quad (25)$$

By Lemma 3, when we set the weights uniformly, i.e. $\alpha_t = 1 \forall t$, and assume a bound on the gradient norms, i.e. $\|\nabla \ell_t(z_t)\|_*^2 \leq G$, the uniform regret is

$$\text{REG} := \sum_{t=1}^T \ell_t(z_t) - \min_z \sum_{t=1}^T \ell_t(z) \leq \frac{G \log(T+1)}{2\mu}. \quad (26)$$

It has been observed [31] that following the leader does not always lead to vanishing regret, for example when the loss function is linear, $\ell_t(\cdot) := \langle \theta_t, \cdot \rangle$, the regret can grow linearly with T (see e.g. Example 2.2 [25]). On the other hand, if the constraint set \mathcal{Z} is “suitably round”, then logarithmic regret is achievable even for linear losses.

Lemma 4 (Theorem 3.3 in [32]) *Let $\{\ell_t(\cdot) := \langle \theta_t, \cdot \rangle\}_{t=1}^T$ be any sequence of linear loss functions. Denote $G := \max_{t \leq T} \|\theta_t\|$ and assume that the support function $\Phi(\cdot) := \max_{z \in \mathcal{Z}} \langle z, \cdot \rangle$ has a unique maximizer for each cumulative loss vector $L_t := \sum_{s=1}^t \theta_s$ at round t . Define $\nu_T := \min_{1 \leq t \leq T} \|L_t\|$. Let $\mathcal{Z} \subset \mathbb{R}^d$ be a λ -strongly convex set. Choose $z_{\text{init}} \in \text{boundary}(\mathcal{Z})$. Then $\text{FTL}[z_{\text{init}}]$ ensures,*

$$\text{REG} := \sum_{t=1}^T \ell_t(z_t) - \min_{z \in \mathcal{Z}} \sum_{t=1}^T \ell_t(z) \leq \frac{2G^2}{\lambda \nu_T} (1 + \log(T)). \quad (27)$$

3.3.3 Be The Leader – FTL^+

FTL^+ (a.k.a. Be The Leader), defined in (17), is similar to FTL, except that the learner plays the best fixed action for the weighted cumulative loss seen thus far *including the current round*.¹ Of course, in practical scenarios, we are unable to see the the loss on the current round before we select our action, but it is nevertheless useful for analytical purposes to consider such “one-step lookahead” algorithms; in each such case we superscript the algorithm’s name with a + symbol. In the the FGNRD framework, we can design dynamics in where the player that acts second in the protocol can indeed observe the loss function prior to choosing its action, and thus it is quite natural to consider such one-step lookahead OCO algorithms.

Lemma 5 (FTL^+) *Let $\alpha_1, \dots, \alpha_T > 0$ be arbitrary, and assume we have a sequence of μ -strongly convex loss functions $\ell_1(\cdot), \dots, \ell_T(\cdot)$ for some $\mu \geq 0$. Then the regret of FTL^+ satisfies*

$$\alpha\text{-REG} \leq - \sum_{t=1}^T \frac{\mu A_{t-1}}{2} \|z_{t-1} - z_t\|^2 \leq 0. \quad (28)$$

3.3.4 Follow The Regularized Leader – FTRL

FTRL, also called dual averaging in optimization literature [33] is a classic algorithm in online learning (see e.g. [22, 24]). Looking at Equation (18) in Algorithm 4 one can notice that FTRL is similar to FTL with an additional *Regularization term* $R(\cdot)$ scaled by parameter $1/\eta$. The regularization term induces *stability* into the decisions of the player, ensuring that consecutive decisions are close to each other, and this property is often crucial in order to ensure regret guarantees. For example, in the case of linear loss functions, FTRL, with appropriate choices of η and $R(\cdot)$, can guarantee sublinear regret guarantees, while FTL cannot. In what follows we assume that $R(\cdot)$ is a β -strongly-convex function over \mathcal{Z} .

Lemma 6 ($\text{FTRL}[R(\cdot), \eta]$) *Given $\eta > 0$ and a β -strongly convex $R : \mathcal{Z} \rightarrow \mathbb{R}$, assume we have $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$ be a sequence of loss functions such that each $\ell_t(\cdot)$ is μ -strongly convex for some $\mu \geq 0$. Then $\text{FTRL}[R(\cdot), \eta]$, which generates the sequence*

¹The term “Be The Leader” was coined by [31], who also showed it exhibits non-positive regret for linear loss functions.

$z_1, \dots, z_T \in \mathcal{Z}$, satisfies the following regret bound for any comparator $z^* \in \mathcal{Z}$,

$$\alpha\text{-REG}^z \leq \sum_{t=1}^T \frac{2\alpha_t^2}{\left(\sum_{s=1}^t \alpha_s \mu\right) + \beta} \|\nabla \ell_t(z_t)\|_*^2 + \frac{1}{\eta} (R(z^*) - R(z_1)). \quad (29)$$

3.3.5 Be The Regularized Leader – FTRL⁺

FTRL⁺ ((19)) is very similar to FTRL, with the difference that the former has an access to all past loss functions up to and *including* the current round. Recall that in our FGNRD template (protocol 2) the x -player is allowed to view the current loss prior to playing, and can therefore apply FTRL⁺.

Lemma 7 (FTRL⁺[$R(\cdot)$, $1/\eta$]) *Given $\eta > 0$ and a β -strongly convex $R : \mathcal{Z} \rightarrow \mathbb{R}$, assume we have $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$ be a sequence of loss functions such that each $\ell_t(\cdot)$ is μ -strongly convex for some $\mu \geq 0$. Then FTRL⁺[$R(\cdot)$, η] satisfies the following regret bound for any comparator $z^* \in \mathcal{Z}$,*

$$\alpha\text{-REG}^z \leq \frac{R(z^*) - R(z_0)}{\eta} - \sum_{t=1}^T \left(\frac{\mu A_{t-1}}{2} + \frac{\beta}{2\eta} \right) \|z_{t-1} - z_t\|^2, \quad (30)$$

where $z_0 = \min_{z \in \mathcal{Z}} R(z)$.

3.3.6 Optimistic Follow the Leader – OPTIMISTICFTL

In the previous subsection, we have seen that FTL⁺ uses the knowledge of the loss function at rounds t in order to ensure negative regret. While this knowledge is oftentimes unavailable, one can often access a “hint” or “guess” function $m_t(\cdot)$ that approximates $\ell_t(\cdot)$ prior to choosing an action z_t . As can be seen from Equation (20) in Algorithm 4 and Lemma 8, OPTIMISTICFTL makes use of the availability of such hints in order to provide better guarantees. The next statement shows that when we have “good” hints, in the sense that $m_t(\cdot) \approx \ell_t(\cdot)$, then OPTIMISTICFTL obtains improved guarantees compared to standard FTL.

Lemma 8 (OPTIMISTICFTL) *Let $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$ be a sequence of loss functions such that each $\ell_t(\cdot)$ is μ_t -strongly convex, let $m_1(\cdot), \dots, m_T(\cdot) : \mathcal{Z} \rightarrow \mathbb{R}$ be an arbitrary sequence of convex functions. Assume that we select our sequence of actions z_1, \dots, z_T according to OCO algorithm OPTIMISTICFTL, then we have the following regret bound*

$$\alpha\text{-REG}^z \leq \sum_{t=1}^T \alpha_t (\ell_t(z_t) - \ell_t(w_{t+1})) - \alpha_t (m_t(z_t) - m_t(w_{t+1})), \quad (31)$$

where we assume the sequence w_2, \dots, w_T satisfies $w_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \sum_{s=1}^{t-1} \ell_s(z)$.

3.3.7 Mirror Descent – OMD

OMD ((22) in Algorithm 5) is another popular algorithm in online learning (see e.g. [22, 24]). Compared to those in Algorithm 4, OMD updates the

current iterate using only the latest loss function instead of the cumulative loss functions seen so far. Yet, it still has a nice regret guarantee.

Lemma 9 (OMD $[\phi(\cdot), z_0, \gamma]$) *Assume that $\phi(\cdot)$ is β -strongly convex w.r.t $\|\cdot\|$ and $D := \sup_{z_0, z^* \in \mathcal{X}} D_{z_0}^\phi(z^*) < \infty$. For any sequence of proper lower semi-continuous convex loss functions $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$, OMD $[\phi(\cdot), z_0, \gamma]$ satisfies the following regret bound for any comparator $z^* \in \mathcal{Z}$,*

$$\alpha\text{-REG}^z \leq \frac{1}{\gamma} D_{z_0}^\phi(z^*) + \frac{\gamma}{2} \sum_{t=1}^T \|\alpha_t \nabla \ell_t(z_t)\|_*^2. \quad (32)$$

Proof The per-round regret with respect to any comparator $z^* \in \mathcal{Z}$ can be bounded as

$$\begin{aligned} \alpha_t \ell_t(z_t) - \alpha_t \ell_t(z^*) &\leq \langle \alpha_t \nabla \ell_t(z_t), z_t - z^* \rangle \\ &= \langle \frac{1}{\gamma} (\nabla \phi(z_{t+1}) - \nabla \phi(z_t)), z^* - z_{t+1} \rangle + \langle \alpha_t \nabla \ell_t(z_t), z_t - z_{t+1} \rangle \\ &\quad + \langle \frac{1}{\gamma} (\nabla \phi(z_t) - \nabla \phi(z_{t+1})) - \alpha_t \nabla \ell_t(z_t), z^* - z_{t+1} \rangle \\ &\leq \langle \frac{1}{\gamma} (\nabla \phi(z_{t+1}) - \nabla \phi(z_t)), z^* - z_{t+1} \rangle + \langle \alpha_t \nabla \ell_t(z_t), z_t - z_{t+1} \rangle \\ &= \frac{1}{\gamma} (D_{z_t}^\phi(z^*) - D_{z_{t+1}}^\phi(z^*) - D_{z_{t+1}}^\phi(z_t)) + \langle \alpha_t \nabla \ell_t(z_t), z_t - z_{t+1} \rangle \\ &\leq \frac{1}{\gamma} (D_{z_t}^\phi(z^*) - D_{z_{t+1}}^\phi(z^*)) + \frac{\gamma}{2} \|\alpha_t \nabla \ell_t(z_t)\|_*^2. \end{aligned} \quad (33)$$

where the second inequality uses the optimality of z_{t+1} for $\min_{z \in \mathcal{Z}} \alpha_t \ell_t(z) + \frac{1}{\gamma} D_{z_t}^\phi(z)$ and the last one uses that $\langle \alpha_t \nabla \ell_t(z_t), z_t - z_{t+1} \rangle \leq \frac{\gamma}{2\beta} \|\alpha_t \nabla \ell_t(z_t)\|_*^2 + \frac{\beta}{2\gamma} \|z_t - z_{t+1}\|^2$ and that $\phi(\cdot)$ is β strongly convex.

Summing (33) from $t = 1$ to T leads to the bounds. \square

3.3.8 Prescient Mirror Descent – OMD⁺

OMD⁺ ((23) in Algorithm 5) assumes that the online player is *prescient*, i.e., it knows the loss functions ℓ_t prior to choosing z_t . Recall that in FGNRD (protocol 2) the x -player is allowed to view the current loss prior to playing, and can therefore apply OMD⁺.

Lemma 10 (OMD⁺ $[\phi(\cdot), z_0, \gamma]$) *Assume that $\phi(\cdot)$ is β -strongly convex and $D := \sup_{z_0, z^* \in \mathcal{X}} D_{z_0}^\phi(z^*) < \infty$. For any sequence of proper lower semi-continuous convex loss functions $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$, OMD⁺ $[\phi(\cdot), z_0, \gamma]$ satisfies the following regret bound for any comparator $z^* \in \mathcal{Z}$,*

$$\alpha\text{-REG}^z \leq \frac{1}{\gamma} D_{z_0}^\phi(z^*) - \sum_{t=1}^T \frac{\beta}{2\gamma} \|z_{t-1} - z_t\|^2 \leq \frac{D}{\gamma} - \sum_{t=1}^T \frac{\beta}{2\gamma} \|z_{t-1} - z_t\|^2.$$

Proof The key inequality we need is Lemma 11; using the lemma with $\theta(z) = \gamma\alpha_t\ell_t(z)$, $z^+ = z_t$ and $c = z_{t-1}$ we have

$$\gamma\alpha_t\ell_t(z_t) - \gamma\alpha_t\ell_t(z^*) = \theta(z_t) - \theta(z^*) \leq D_{z_{t-1}}^\phi(z^*) - D_{z_t}^\phi(z^*) - D_{z_{t-1}}^\phi(z_t). \quad (34)$$

Therefore, the regret with respect to any comparator $z^* \in \mathcal{Z}$ can be bounded as

$$\begin{aligned} \alpha\text{-REG}^z &:= \sum_{t=1}^T \alpha_t \ell_t(z_t) - \sum_{t=1}^T \alpha_t \ell_t(z^*) \\ &\stackrel{(34)}{\leq} \sum_{t=1}^T \frac{1}{\gamma} (D_{z_{t-1}}^\phi(z^*) - D_{z_t}^\phi(z^*) - D_{z_{t-1}}^\phi(z_t)) \\ &= \frac{1}{\gamma} D_{z_0}^\phi(z^*) - \frac{1}{\gamma} D_{z_T}^\phi(z^*) + \sum_{t=1}^{T-1} \left(\frac{1}{\gamma} - \frac{1}{\gamma} \right) D_{z_t}^\phi(z^*) - \frac{1}{\gamma} D_{z_{t-1}}^\phi(z_t) \\ &\leq \frac{1}{\gamma} D_{z_0}^\phi(z^*) - \sum_{t=1}^T \frac{\beta}{2\gamma} \|z_{t-1} - z_t\|^2, \end{aligned}$$

where the last inequality uses the definition of D and the strong convexity of ϕ , which grants $D_{z_{t-1}}^\phi(z_t) \geq \frac{\beta}{2} \|z_t - z_{t-1}\|^2$. \square

Lemma 11 (Property 1 in [34]) *For any proper lower semi-continuous convex function $\theta(z)$, let $z^+ = \operatorname{argmin}_{z \in \mathcal{Z}} \theta(z) + D_c^\phi(z)$. Then, it satisfies that for any $z^* \in \mathcal{Z}$,*

$$\theta(z^+) - \theta(z^*) \leq D_c^\phi(z^*) - D_{z^+}^\phi(z^*) - D_c^\phi(z^+). \quad (35)$$

3.3.9 Optimistic Mirror Descent – OPTIMISTICMD

When a hint of the loss function at the beginning of each round is available, the online learner can exploit the hint in the hope to get a better regret. Denote m_t a vector that represents the hint before the learner outputs a point at t . OPTIMISTICMD in Algorithm 5 has the following regret guarantee.

Lemma 12 (OPTIMISTICMD $[\phi(\cdot), z_0, \gamma]$) *Assume that $\phi(\cdot)$ is β -strongly convex w.r.t $\|\cdot\|$ and $D := \sup_{z_0, z^* \in \mathcal{X}} D_{z_0}^\phi(z^*) < \infty$. For any sequence of proper lower semi-continuous convex loss functions $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$, OPTIMISTICMD $[\phi(\cdot), z_0, \gamma]$ satisfies the following regret bound for any comparator $z^* \in \mathcal{Z}$,*

$$\alpha\text{-REG}^z \leq \frac{1}{\gamma} D_{z_0}^\phi(z^*) + \frac{\gamma}{2\beta} \sum_{t=1}^T \alpha_t^2 \|\nabla \ell_t(z_t) - m_t\|_*^2. \quad (36)$$

Proof The proof in the following is based on the proof in [29]. By applying Lemma 11 to the update $z_t \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_t \langle z, m_t \rangle + \frac{1}{\gamma} D_{z_{t-\frac{1}{2}}}^\phi(z) \right)$ we have

$$\alpha_t \langle m_t, z_t - z_{t+\frac{1}{2}} \rangle \leq \frac{1}{\gamma} \left(D_{z_{t-\frac{1}{2}}}^\phi(z_{t+\frac{1}{2}}) - D_{z_t}^\phi(z_{t+\frac{1}{2}}) - D_{z_{t-\frac{1}{2}}}^\phi(z_t) \right). \quad (37)$$

Also, by applying Lemma 11 to $z_{t+\frac{1}{2}} \leftarrow \operatorname{argmin}_{z \in \mathcal{Z}} \left(\alpha_t \langle z, \nabla \ell_t(z_t) \rangle + \frac{1}{\gamma} D_{z_{t-\frac{1}{2}}}^\phi(z) \right)$, we get

$$\alpha_t \langle \nabla \ell_t(z_t), z_{t+\frac{1}{2}} - z^* \rangle \leq \frac{1}{\gamma} \left(D_{z_{t-\frac{1}{2}}}^\phi(z^*) - D_{z_{t+\frac{1}{2}}}^\phi(z^*) - D_{z_{t-\frac{1}{2}}}^\phi(z_{t+\frac{1}{2}}) \right). \quad (38)$$

Then, we can bound the per-round regret with respect to the comparator z^* as follows:

$$\begin{aligned} \alpha_t \ell_t(z_t) - \alpha_t \ell_t(z^*) &\leq \alpha_t \langle \nabla \ell_t(z_t), z_t - z^* \rangle \\ &= \alpha_t \langle \nabla \ell_t(z_t), z_{t+\frac{1}{2}} - z^* \rangle + \alpha_t \langle m_t, z_t - z_{t+\frac{1}{2}} \rangle + \alpha_t \langle \nabla \ell_t(z_t) - m_t, z_t - z_{t+\frac{1}{2}} \rangle \\ &\leq \frac{1}{\gamma} \left(D_{z_{t-\frac{1}{2}}}^\phi(z^*) - D_{z_{t+\frac{1}{2}}}^\phi(z^*) - D_{z_t}^\phi(z_{t+\frac{1}{2}}) - D_{z_{t-\frac{1}{2}}}^\phi(z_t) \right) + \alpha_t \|\nabla \ell_t(z_t) - m_t\|_* \|z_t - z_{t+\frac{1}{2}}\| \\ &\leq \frac{1}{\gamma} \left(D_{z_{t-\frac{1}{2}}}^\phi(z^*) - D_{z_{t+\frac{1}{2}}}^\phi(z^*) - \frac{\beta}{2} \|z_t - z_{t+\frac{1}{2}}\|^2 - \frac{\beta}{2} \|z_t - z_{t-\frac{1}{2}}\|^2 \right) \\ &\quad + \frac{\alpha_t^2 \gamma}{2\beta} \|\nabla \ell_t(z_t) - m_t\|_*^2 + \frac{\beta}{2\gamma} \|z_t - z_{t+\frac{1}{2}}\|^2, \end{aligned}$$

where the second inequality is due to (37) and (38) and the last one is by the β -strong convexity of $\phi(\cdot)$.

Summing the above from $t = 1$ to $t = T$ and noting that $z_{-\frac{1}{2}} = z_0 \in \mathcal{K}$, we obtain the bound. \square

3.4 A meta online learning algorithm: OPTIMISTICFTRL

Here we describe OPTIMISTICFTRL, a Meta-algorithm that captures all previously mentioned methods as special cases. As can be seen from Equation (21) in Algorithm 4, OPTIMISTICFTRL employs a regularization term (similarly to FTRL and FTRL⁺), and makes use of a hint sequence $m_t(\cdot)$ (similarly to OPTIMISTICFTL).

In Lemma 13 we state the regret guarantees of OPTIMISTICFTRL, and then show how does the guarantees of FTL, FTL⁺, OPTIMISTICFTL, FTRL, and FTRL⁺ follow as corollaries of this Lemma. The proof of Lemma 13 is provided in Subsection 3.4.

Lemma 13 (OPTIMISTICFTRL[$R(\cdot), \eta$]) *Given $\eta > 0$ and a β -strongly convex $R : \mathcal{Z} \rightarrow \mathbb{R}$, assume we have $\{\alpha_t \ell_t(\cdot)\}_{t=1}^T$ a sequence of weights and loss functions such that each $\ell_t(\cdot)$ is μ_t -strongly convex for some $\mu_t \geq 0$. Let $m_1, \dots, m_T : \mathcal{Z} \rightarrow \mathbb{R}$ be the sequence of hint functions given to OPTIMISTICFTRL, where each $m_t(\cdot)$ is $\hat{\mu}_t$ -strongly convex function over \mathcal{Z} . Then OPTIMISTICFTRL[$R(\cdot), \eta$] satisfies*

$$\begin{aligned} \alpha\text{-REG}^z &\leq & (39) \\ &\sum_{t=1}^T \alpha_t (\ell_t(z_t) - \ell_t(w_{t+1}) - m_t(z_t) + m_t(w_{t+1})) & \text{(term (A))} \\ &+ \frac{1}{\eta} (R(z^*) - R(w_1)) & \text{(term (B))} \end{aligned}$$

$$-\frac{1}{2} \sum_{t=1}^T \left(\frac{\beta}{\eta} + \sum_{s=1}^{t-1} \alpha_s \mu_s \right) \|z_t - w_t\|^2 \quad (\text{term (C)})$$

$$-\frac{1}{2} \sum_{t=1}^T \left(\frac{\beta}{\eta} + \alpha_t \hat{\mu}_t + \sum_{s=1}^{t-1} \alpha_s \mu_s \right) \|z_t - w_{t+1}\|^2, \quad (\text{term (D)})$$

where w_1, \dots, w_T is the alternative sequence chosen according to $\text{FTRL}[R(\cdot), \eta]$ $w_t := \arg\min_{z \in \mathcal{Z}} \sum_{s=1}^{t-1} \alpha_s \ell_s(z) + \frac{1}{\eta} R(z)$, and $z^* \in \mathcal{Z}$ is arbitrary.

We prove Lemma 13 at the end of this section. But let us first show how several of the aforementioned guarantees—for algorithms FTL , FTL^+ , OPTIMISTICFTL , FTRL , and FTRL^+ —follow from the above Lemma.

Proof of Lemma 8 on OPTIMISTICFTL Observe that OPTIMISTICFTL is actually OPTIMISTICFTRL when $R(\cdot)$ is a zero function. Therefore, let $R(\cdot) = 0$ in Lemma 13 and drop term (C) and (D) in (39) as they are non-positive, we obtain the result. \square

Proof of Lemma 7 on FTRL^+ Observe that the FTRL^+ update is exactly equivalent to OPTIMISTICFTRL with $m_t(\cdot) = \ell_t(\cdot)$. Furthermore, w_{t+1} in Lemma 13 is actually z_t of FTRL^+ shown on (19). So term (A) and term (D) on (39) in Lemma 13 is 0, Therefore, FTRL^+ regret satisfies

$$\alpha\text{-REG}^z \leq \frac{R(z^*) - R(z_0)}{\eta} - \sum_{t=1}^T \left(\frac{\mu A_{t-1}}{2} + \frac{\beta}{2\eta} \right) \|z_{t-1} - z_t\|^2. \quad (40)$$

\square

Proof of Lemma 5 on FTL^+ Observe that FTL^+ is actually FTRL^+ with $R(\cdot) = 0$. Therefore, let $R(\cdot) = 0$ and $\beta = 0$ in Equation (40), we obtain the regret of FTL^+

$$\alpha\text{-REG}^z \leq - \sum_{t=1}^T \frac{\mu A_{t-1}}{2} \|z_{t-1} - z_t\|^2 \leq 0.$$

\square

Proof of Lemma 6 on FTRL Observe that FTRL is actually OPTIMISTICFTRL where $m_t(\cdot) = 0 \forall t$. Therefore, let $m_t(\cdot) = 0$ in Lemma 13, we obtain the regret of FTRL ,

$$\alpha\text{-REG}^z \leq \sum_{t=1}^T \alpha_t (\ell_t(z_t) - \ell_t(z_{t+1})) + \frac{1}{\eta} (R(z^*) - R(z_1)), \quad (41)$$

where we have dropped term (C) and term (D) on (39) since they are non-positive, and we also note that w_t in Lemma 13 is the same as z_t here. To continue, we use Lemma 14. Specifically, in Lemma 14, we let $\psi_1(\cdot) \leftarrow \sum_{s=1}^{t-1} \alpha_s \ell_s(\cdot) + \frac{1}{\eta} R(\cdot)$ and

$\psi_2(\cdot) \leftarrow \sum_{s=1}^t \alpha_s \ell_s(\cdot) + \frac{1}{\eta} R(\cdot)$. Then, we have $\phi(\cdot) = \alpha_t \ell_t(\cdot)$, $u_1 = z_t$, $u_2 = z_{t+1}$ and that $\sigma = \sum_{s=1}^t \alpha_s \mu + \beta$. So by Lemma 14 below, we have

$$\alpha_t (\ell_t(z_t) - \ell_t(z_{t+1})) \leq \frac{2\alpha_t^2}{(\sum_{s=1}^t \alpha_s \mu) + \beta} \|\nabla \ell_t(z_t)\|_*^2. \quad (42)$$

Combining (41) and (42) leads to the result.

Lemma 14 (Lemma 5 in [35]) *Let $\psi_1(\cdot), \psi_2(\cdot) : \mathcal{Z} \rightarrow \mathbb{R}$ be two convex functions defined over a closed and convex domain. Denote $u_1 := \operatorname{argmin}_{z \in \mathcal{Z}} \psi_1(z)$ and $u_2 := \operatorname{argmin}_{z \in \mathcal{Z}} \psi_2(z)$. Assume that ψ_2 is σ -strongly convex with respect to a norm $\|\cdot\|$. Define $\phi(\cdot) := \psi_2(\cdot) - \psi_1(\cdot)$. Then,*

$$\|u_1 - u_2\| \leq \frac{2}{\sigma} \|\nabla \phi(u_1)\|_*. \quad (43)$$

Furthermore, if $\phi(\cdot)$ is convex, then,

$$0 \leq \phi(u_1) - \phi(u_2) \leq \frac{2}{\sigma} \|\nabla \phi(u_1)\|_*^2. \quad (44)$$

□

Proof of Lemma 3 on FTL Observe that FTL is actually FTRL with $R(\cdot) = 0$. Therefore, let $R(\cdot) = 0$ and $\beta = 0$ in Lemma 6, we obtain the result. □

We now complete this section by proving the meta result for OPTIMISTICFTRL.

of Lemma 13 We can re-write the regret as

$$\begin{aligned} \alpha\text{-REG}^z &:= \sum_{t=1}^T \alpha_t \ell_t(z_t) - \sum_{t=1}^T \alpha_t \ell_t(z^*) \\ &= \underbrace{\sum_{t=1}^T \alpha_t (\ell_t(z_t) - \ell_t(w_{t+1})) - \alpha_t (m_t(z_t) - m_t(w_{t+1}))}_{\text{first term}} \\ &\quad + \underbrace{\sum_{t=1}^T \alpha_t (m_t(z_t) - m_t(w_{t+1}))}_{\text{second term}} + \underbrace{\sum_{t=1}^T \alpha_t (\ell_t(w_{t+1}) - \ell_t(z^*))}_{\text{third term}}. \end{aligned} \quad (45)$$

In the following, we will denote

$$D_T := \frac{1}{2} \left(\sum_{t=1}^T \left(\frac{\beta}{\eta} + \sum_{s=1}^t \alpha_s \mu_s \right) \|z_t - w_t\|^2 + \sum_{t=1}^T \left(\frac{\beta}{\eta} + \sum_{s=1}^t \alpha_s \mu_s + \alpha_t \hat{\mu}_t \right) \|z_t - w_{t+1}\|^2 \right)$$

for brevity. Let us first deal with the second term and the third term on (45). We will use induction to show that

$$\begin{aligned} &\underbrace{\sum_{t=1}^T \alpha_t (m_t(z_t) - m_t(w_{t+1}))}_{\text{second term}} + \underbrace{\sum_{t=1}^T \alpha_t (\ell_t(w_{t+1}) - \ell_t(z^*))}_{\text{third term}} \\ &\leq \frac{1}{\eta} (R(z^*) - R(w_1)) - D_T \end{aligned} \quad (46)$$

for any point $z^* \in \mathcal{Z}$. For the base case $T = 0$, we have

$$\begin{aligned} & \sum_{t=1}^0 \alpha_t (m_t(z_t) - m_t(w_{t+1})) + \sum_{t=1}^0 \alpha_t (\ell_t(w_{t+1}) - \ell_t(z^*)) \\ & = 0 \leq \frac{1}{\eta} (R(z^*) - R(w_1)) - 0, \end{aligned}$$

as $w_1 := \operatorname{argmin}_{z \in \mathcal{Z}} R(z)$. So the base case trivially holds.

Let us assume that the inequality (46) holds for $t = 0, 1, \dots, T-1$. Now consider round T . We have

$$\begin{aligned} & \sum_{t=1}^T \alpha_t (m_t(z_t) - m_t(w_{t+1}) + \ell_t(w_{t+1})) \\ & \stackrel{(a)}{\leq} \alpha_T (m_T(z_T) - m_T(w_{T+1}) + \ell_T(w_{T+1})) + \frac{1}{\eta} (R(w_T) - R(w_1)) - D_{T-1} \\ & \quad + \sum_{t=1}^{T-1} \alpha_t \ell_t(w_T). \\ & \stackrel{(b)}{\leq} \alpha_T (m_T(z_T) - m_T(w_{T+1}) + \ell_T(w_{T+1})) + \frac{1}{\eta} (R(z_T) - R(w_1)) - D_{T-1} \\ & \quad - \frac{1}{2} \left(\frac{\beta}{\eta} + \sum_{s=1}^{T-1} \alpha_s \mu_s \right) \|z_T - w_T\|^2 + \sum_{t=1}^{T-1} \alpha_t \ell_t(z_T) \\ & = \alpha_T (\ell_T(w_{T+1}) - m_T(w_{T+1})) + \frac{1}{\eta} (R(z_T) - R(w_1)) - D_{T-1} \\ & \quad - \frac{1}{2} \left(\frac{\beta}{\eta} + \sum_{s=1}^{T-1} \alpha_s \mu_s \right) \|z_T - w_T\|^2 + \sum_{t=1}^{T-1} \alpha_t \ell_t(z_T) + \alpha_T m_T(z_T) \\ & \stackrel{(c)}{\leq} \alpha_T (\ell_T(w_{T+1}) - m_T(w_{T+1})) + \frac{1}{\eta} (R(w_{T+1}) - R(w_1)) - D_{T-1} \\ & \quad - \frac{1}{2} \left(\frac{\beta}{\eta} + \sum_{s=1}^{T-1} \alpha_s \mu_s \right) \|z_T - w_T\|^2 - \frac{1}{2} \left(\frac{\beta}{\eta} + \sum_{s=1}^{T-1} \alpha_s \mu_s + \alpha_T \hat{\mu}_T \right) \|z_T - w_{T+1}\|^2 \\ & \quad + \sum_{t=1}^{T-1} \alpha_t \ell_t(w_{T+1}) + \alpha_T m_T(w_{T+1}) \\ & = \sum_{t=1}^T \alpha_t \ell_t(w_{T+1}) + \frac{1}{\eta} (R(w_{T+1}) - R(w_1)) - D_T \\ & \stackrel{(d)}{\leq} \sum_{t=1}^T \alpha_t \ell_t(z^*) + \frac{1}{\eta} (R(z^*) - R(w_1)) - D_T, \end{aligned}$$

where (a) we use the induction such that the inequality (46) holds for any $z^* \in \mathcal{Z}$ including $z^* = w_T$, and (b) is because

$$\sum_{t=1}^{T-1} \alpha_t \ell_t(w_T) + \frac{1}{\eta} R(w_T) \leq \sum_{t=1}^{T-1} \alpha_t \ell_t(z_T) + \frac{1}{\eta} R(z_T) - \frac{1}{2} \left(\frac{\beta}{\eta} + \sum_{s=1}^{T-1} \alpha_s \mu_s \right) \|z_T - w_T\|^2,$$

as w_T is the minimizer of a $\frac{\beta}{\eta} + \sum_{t=1}^{T-1} \alpha_t \mu_t$ strongly convex function since $w_T := \operatorname{argmin}_{z \in \mathcal{Z}} \sum_{s=1}^{T-1} \alpha_s \ell_s(z) + \frac{1}{\eta} R(z)$, and (c) is because

$$\begin{aligned} \sum_{t=1}^{T-1} \alpha_t \ell_t(z_T) + \alpha_T m_T(z_T) + \frac{1}{\eta} R(z_T) &\leq \sum_{t=1}^{T-1} \alpha_t \ell_t(w_{T+1}) + \alpha_T m_T(w_{T+1}) + \frac{1}{\eta} R(w_{T+1}) \\ &\quad - \frac{1}{2} \left(\frac{\beta}{\eta} + \sum_{s=1}^{T-1} \alpha_s \mu_s + \alpha_T \hat{\mu}_T \right) \|z_T - w_{T+1}\|^2, \end{aligned}$$

as z_T is the minimizer of a $\frac{\beta}{\eta} + \sum_{t=1}^{T-1} \alpha_t \mu_t + \alpha_T \hat{\mu}_T$ strongly convex function since $z_T := \operatorname{argmin}_{z \in \mathcal{Z}} \left(\sum_{s=1}^{T-1} \alpha_s \ell_s(z) \right) + \alpha_T m_T(z) + \frac{1}{\eta} R(z)$, and (d) is due to $w_{T+1} := \operatorname{argmin}_{z \in \mathcal{Z}} \sum_{t=1}^T \alpha_t \ell_t(z) + \frac{1}{\eta} R(z)$. □

4 Recovery of existing algorithms

What we are now able to establish, using the tools developed above, is that several iterative first order methods to minimize a convex function can be cast as simple instantiations of the Fenchel game no-regret dynamics. But more importantly, using this framework and the various regret bounds stated above, we able to establish a convergence rate for each via a unified analysis.

For every one of the optimization methods we explore below we provide the following:

1. We state the update method described in its standard iterative form, alongside an equivalent formulation given as a no-regret dynamic. To provide the FGNRD form, we must specify the payoff function $g(\cdot, \cdot)$ —typically the Fenchel game, with some variants—as well as the sequence of weights α_t , and the no-regret algorithms $\text{OAlg}^Y, \text{OAlg}^X$ the two players.
2. We provide a proof of this equivalence, showing that the FGNRD formulation does indeed produce the same sequence of iterates as the iterative form; this is often deferred to the appendix.
3. Leaning on Theorem 1, we prove a convergence rate for the method.

4.1 Frank-Wolfe method and its variants

The *Frank-Wolfe method* (FW) [6], also known as *conditional gradient*, is known for solving constrained optimization problems. FW is entirely first-order, while requiring access to a linear optimization oracle. Specifically, given a compact and convex constraint set $\mathcal{K} \subset \mathbb{R}^d$, FW relies on the ability to (quickly) answer queries of the form $\operatorname{argmin}_{x \in \mathcal{K}} \langle x, v \rangle$, for any vector $v \in \mathbb{R}^d$. In many cases this linear optimization problem is much faster for well-behaved constraint sets; e.g. simple convex polytopes, the PSD cone, and various balls defined by vector and matrix norms [36, 37]. When the constraint set is the nuclear norm ball, which arises in matrix completion problems, then the linear optimization oracle corresponds to computing a top singular vector, which requires time roughly linear in the size of the matrix [37].

Algorithm 6 Frank-Wolfe [6]

Given: L -smooth $f(\cdot)$, convex domain \mathcal{K} , arbitrary w_0 , iterations T .

$$\begin{array}{l} \gamma_t \leftarrow \frac{2}{t+1} \\ v_t \leftarrow \underset{v \in \mathcal{K}}{\operatorname{argmin}} \langle v, \nabla f(w_{t-1}) \rangle \\ w_t \leftarrow (1 - \gamma_t)w_{t-1} + \gamma_t v_t \end{array}$$

Iterative Description

$$\begin{array}{l} g(x, y) := \langle x, y \rangle - f^*(y) \\ \alpha_t \leftarrow t \\ \text{OAlg}^Y := \text{FTL}[\nabla f(w_0)] \\ \text{OAlg}^X := \text{BESTRESP}^+ \end{array}$$

FGNRD Equivalence

Output: $w_T = \bar{x}_T$

We describe the Frank-Wolfe method precisely in Algorithm 6, in both its iterative form and its FGNRD interpretation. We begin by showing that these two representations are equivalent.

Proposition 3 *The two interpretations of Frank-Wolfe, as described in Algorithm 6, are equivalent. That is, for every t , the iterate w_t computed iteratively on the left hand side is identically the weighted-average point \bar{x}_t produced by the dynamic on the right hand side.*

Proof We show, via induction, that the following three equalities are maintained for every t . Note that three objects on the left correspond to the iterative description given in Algorithm 6 whereas the three on the right correspond to the FGNRD description.

$$\nabla f(w_{t-1}) = y_t \tag{47}$$

$$v_t = x_t \tag{48}$$

$$w_t = \bar{x}_t. \tag{49}$$

To start, we observe that since the OAlg^Y is set as $\text{FTL}[\nabla f(w_0)]$, we have the base case for (47), $y_1 = \nabla f(w_0)$, holds by definition. Furthermore, we observe that for any t we have (47) \implies (48). This is because, if $y_t = \nabla f(w_{t-1})$, the definition of BESTRESP^+ implies that

$$x_t = \underset{x \in \mathcal{X}}{\operatorname{argmin}} \alpha_t (\langle x, y_t \rangle - f^*(y_t)) = \underset{x \in \mathcal{X}}{\operatorname{argmin}} \langle x, \nabla f(w_{t-1}) \rangle = v_t \tag{50}$$

Next, we can show that (48) \implies (49) for any t as well using induction. Assuming that $w_{t-1} = \frac{\sum_{s=1}^{t-1} \alpha_s x_s}{\sum_{s=1}^{t-1} \alpha_s} = \frac{\sum_{s=1}^{t-1} s v_s}{\sum_{s=1}^{t-1} s}$, a bit of algebra verifies

$$\begin{aligned} w_t &:= (1 - \gamma_t)w_{t-1} + \gamma_t v_t = \left(\frac{t-1}{t+1}\right) \frac{\sum_{s=1}^{t-1} s v_s}{\sum_{s=1}^{t-1} s} + \left(\frac{2}{t+1}\right) v_t \\ &= \frac{\sum_{s=1}^t s v_s}{\sum_{s=1}^t s} = \frac{\sum_{s=1}^t \alpha_s x_s}{\sum_{s=1}^t \alpha_s} =: \bar{x}_t \end{aligned}$$

Finally, we show that (47) holds for $t > 1$ via induction. Recall that y_t is selected via FTL against the sequence of loss functions $\alpha_t \ell_t(\cdot) := -\alpha_t g(x_t, \cdot)$. Precisely this means that, for $t > 1$,

$$y_t := \underset{y \in \mathcal{Y}}{\operatorname{argmin}} \left\{ \frac{1}{A_{t-1}} \sum_{s=1}^{t-1} \alpha_s \ell_s(y) \right\}$$

$$\begin{aligned}
 &= \operatorname{argmin}_{y \in \mathcal{Y}} \left\{ \frac{1}{A_{t-1}} \sum_{s=1}^{t-1} (\alpha_s \langle -x_s, y \rangle + f^*(y)) \right\} \\
 &= \operatorname{argmax}_{y \in \mathcal{Y}} \{ \langle \bar{x}_{t-1}, y \rangle - f^*(y) \} = \nabla f(\bar{x}_{t-1}).
 \end{aligned}$$

The final line follows as a result of the Legendre transform [17]. Finally, by induction, we have $\bar{x}_{t-1} = w_{t-1}$, and hence we have established (47). This completes the proof. \square

Now that we have established Frank-Wolfe as an instance of Protocol 2, we can now prove a bound on convergence using the tools established in Section 3.

Theorem 4 *Let w_T be the output of Algorithm 6. Let f be L -smooth and let \mathcal{K} have square ℓ_2 diameter no more than D . Then we have*

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) \leq \frac{8LD}{T+1}.$$

Proof Now that we have established that Algorithm 6 is an instance of Protocol 2, we can appeal directly to Theorem 1 to see that

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) \leq \overline{\alpha\text{-REG}}^x[\text{BESTRESP}^+] + \overline{\alpha\text{-REG}}^y[\text{FTL}].$$

Recall that, by Lemma 2, we have $\overline{\alpha\text{-REG}}^x[\text{BESTRESP}^+] \leq 0$. Let us then turn our attention to the regret of OAlg^Y .

First note that, since $f(\cdot)$ is L -smooth, its conjugate $f^*(\cdot)$ is $\frac{1}{L}$ -strongly convex, and thus the function $-g(x, \cdot)$ is also $\frac{1}{L}$ -strongly convex in its second argument. Next, if we define $\ell_t(\cdot) := -g(x_t, \cdot)$, then we can bound the norm of the gradient as

$$\|\nabla \ell_t(y_t)\|^2 = \|x_t - \nabla f^*(y_t)\|^2 = \|x_t - \bar{x}_{t-1}\|^2 \leq D.$$

Combining with Lemma 3 we see that

$$\overline{\alpha\text{-REG}}^y[\text{FTL}] \leq \frac{1}{A_T} \sum_{t=1}^T \frac{2\alpha_t^2 \|\nabla \ell_t(y_t)\|^2}{\sum_{s=1}^t \alpha_s(1/L)} = \frac{8L}{T(T+1)} \sum_{t=1}^T \frac{t^2 D}{t(t+1)} \leq \frac{8LD}{T+1}.$$

This completes the proof. \square

4.1.1 Variant 1: a linear rate Frank-Wolfe over strongly convex set

[12, 38, 39] show that under certain conditions, Frank-Wolfe for smooth convex function (*not necessarily a strongly convex function*) for strongly convex sets has linear rate under certain conditions. Algorithm 7 shows that a similar result can be derived from the game framework, in which the y -player uses a variant of FTL called AFTL (defined in Algorithm 21 of Appendix B).

Algorithm 7 Adaptive Frank-Wolfe

Given: L -smooth convex $f(\cdot)$, arbitrary $\bar{x}_0 = x_0 = w_0 \in \mathcal{K}$, iterations T , $A_0 = 0$.

$\begin{aligned} x_t &\leftarrow \operatorname{argmin}_{x \in \mathcal{K}} \langle x, \nabla f(w_{t-1}) \rangle \\ \alpha_t &\leftarrow \frac{1}{\ x_t - \bar{x}_{t-1}\ } \\ A_t &\leftarrow A_{t-1} + \alpha_t \\ w_t &\leftarrow \frac{1}{A_t} \sum_{s=1}^t \alpha_s x_s \end{aligned}$	$\begin{aligned} g(x, y) &:= \langle x, y \rangle - f^*(y) \\ \text{OAlg}^Y &:= \text{AFTL}[\nabla f(x_0)] \\ \text{OAlg}^X &:= \text{BESTRESP}^+ \end{aligned}$
Iterative Description	FGNRD Equivalence

Output: $w_T = \bar{x}_T$

Theorem 5 *Suppose that $f(\cdot)$ is L -smooth convex. and that \mathcal{K} is a λ -strongly convex set. Also assume that the gradients of the $f(\cdot)$ in \mathcal{K} are bounded away from 0, i.e., $\max_{w \in \mathcal{K}} \|\nabla f(w)\| \geq B$. Then, Algorithm 7 has*

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) = O\left(\exp\left(-\frac{\lambda B}{L} T\right)\right).$$

Note that the weights α_t are not predefined but rather depend on the queries of the algorithm. The proof of Theorem 5 is described in full detail in Appendix B.

4.1.2 Variant 2: an incremental Frank-Wolfe

Recently, [13] and [40] propose stochastic Frank-Wolfe algorithms for optimizing smooth convex finite-sum functions, i.e. $\min_{x \in \mathcal{K}} f(x) := \frac{1}{n} \sum_{i=1}^n f_i(x)$, where each $f_i(x) := \phi(x^\top z_i)$ represents a loss function $\phi(\cdot)$ associated with sample z_i . In each iteration the algorithms only require a gradient computation of a single component. [13] show that their algorithm has $O(\frac{c_\kappa}{T})$ expected convergence rate, where c_κ is a number that depends on the underlying data matrix z and in worst case is bounded by the number of components n . Algorithm 8 shows a similar algorithm which picks a sample in each iteration by cycling through the data points. It is obtained when the y-player uses a variant of FTL called LAZYFTL defined in Appendix C. We have the following theorem that shows an $\tilde{O}(\frac{n}{T})$ *deterministic* convergence rate. Its proof is in Appendix C.

Algorithm 8 Stochastic Frank-Wolfe

 Given: L -smooth $f(\cdot)$, convex domain \mathcal{K} , arbitrary $w_0 \in \mathcal{K}$, iterations T .

 Init: For each sample i , compute $g_{i,0} := \frac{1}{n} \nabla f_i(w_0) \in \mathbb{R}^d$ so that $\nabla f(w_0) = \sum_{i=1}^n g_{i,0}$.

Select i_t via by cycling through $[n]$

$$g_{i_t,t} \leftarrow \frac{1}{n} \nabla f_{i_t}(w_t) \text{ and}$$

$$g_{j,t} \leftarrow g_{j,t-1}, \text{ for } j \neq i_t$$

$$g_t \leftarrow \sum_{i=1}^n g_{i,t}$$

$$v_t \leftarrow \operatorname{argmin}_{x \in \mathcal{K}} \langle x, g_t \rangle$$

$$w_t \leftarrow (1 - \gamma_t)w_{t-1} + \gamma_t v_t$$

Iterative Description

$$g(x, y) := \langle x, y \rangle - f^*(y)$$

$$\alpha_t \leftarrow 1$$

$$\text{OAlg}^Y := \text{LAZYFTL}[\nabla f(w_0)]$$

$$\text{OAlg}^X := \text{BESTRESP}^+$$

FGNRD Equivalence

 Output: $w_T = \bar{x}_T$

Theorem 6 Assume that $f(\cdot)$ is L -smooth convex and that its conjugate is L_0 -Lipschitz. Algorithm 8 has

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) = O\left(\frac{\max\{LR, L(L_0 + r)nr\} \log T}{T}\right),$$

where r is a bound of the length of any point x in the constraint set \mathcal{K} , i.e. $\max_{x \in \mathcal{K}} \|x\| \leq r$, and R is the square of the diameter of \mathcal{K} .

4.1.3 More related works

There has been growing interest in Frank-Wolfe in recent years, e.g. [41, 42]. For the constraint set that belongs to a certain class of convex polytopes, there are FW-like algorithms achieving linear convergence rate, see e.g. [36, 43–47]. There are also works that study Frank-Wolfe on various aspects, e.g. minimizing some structural norms [48], reducing the number of gradient evaluations [49]. Bach [50] shows that for certain types of objectives, subgradient descent applied to the primal domain is equivalent to FW applied to the dual domain.

4.2 Vanilla gradient descent, with averaging

In Algorithm 9 we describe a variant of gradient descent. It outputs a weighted average of iterates of the gradient descent steps. The algorithm is obtained by swapping the ordering of the players. The x -player plays first according to OMD and then the y -player plays BESTRESP^+ . Theorem 7 shows that the algorithm has a guarantee even for non-smooth convex functions.

Theorem 7 Denote the constant $G = \max_{w \in \mathcal{K}} \|\partial f(w)\|^2$, and let R be an upper bound on $\|w_0 - w^*\|$, where $w_* := \operatorname{argmin}_{w \in \mathcal{K}} f(w)$. The output $\bar{w}_T = \bar{x}_T$ of Algorithm 9 satisfies

$$f(\bar{w}_T) - \min_{w \in \mathcal{K}} f(w) = O\left(\frac{GR}{\sqrt{T}}\right).$$

Algorithm 9 Vanilla gradient descent, with averagingGiven: Convex $f(\cdot)$ and iterations T .Init: $w_0 = x_0 \in \mathcal{K}$.

$$\begin{aligned} \eta &\leftarrow \frac{R}{G\sqrt{T}} \\ w_t &\leftarrow w_{t-1} - \eta \partial f(w_{t-1}) \\ \bar{w}_t &\leftarrow \frac{1}{t} \sum_{s=1}^t w_s \end{aligned}$$

Iterative Description

$$\begin{aligned} g(x, y) &:= \langle x, y \rangle - f^*(y) \\ \alpha_t &:= 1 \text{ for } t = 1, \dots, T \\ \text{OAlg}^X &:= \text{OMD}[\frac{1}{2}\|\cdot\|_2^2, x_0, \frac{R}{G\sqrt{T}}] \\ \text{OAlg}^Y &:= \text{BESTRESP}^+ \end{aligned}$$

FGNRD Equivalence

Output: $\bar{w}_T = \bar{x}_T$

Proof The equivalence $w_t = x_t$ between the two displays can be easily shown by noting that the update of OMD is $x_t = x_{t-1} - \alpha_t \gamma y_{t-1}$, where $\gamma = \frac{R}{G\sqrt{T}}$ and $\alpha_t = 1$, and the update of BESTRESP^+ is $y_t = \text{argmin}_y f^*(y) - \langle x_t, y \rangle = \partial f(x_t)$.

By Lemma 9, the regret of the x-player is, $\alpha\text{-REG}^x \leq \frac{1}{\gamma} D_{x_0}^\phi(x^*) + \frac{\gamma}{2} \sum_{t=1}^T \|\alpha_t y_t\|_2^2$. On the other hand, by Lemma 2, the regret of the y-player is 0. By setting $\alpha_t = 1$, adding average regrets of both players, and plugging in $\gamma = \frac{R}{G\sqrt{T}}$,

$$\begin{aligned} \overline{\alpha\text{-REG}}^x[\text{OMD}] + \overline{\alpha\text{-REG}}^y[\text{BESTRESP}^+] &\leq \frac{1}{A_t} \left(\frac{1}{\gamma} D_{x_0}^\phi(x^*) + \sum_{t=1}^T \frac{\gamma}{2} \|\alpha_t y_t\|_2^2 \right) \\ &\leq \frac{1}{T} \left(\frac{R^2}{\gamma} + \frac{\gamma T G^2}{2} \right) \\ &= O\left(\frac{GR}{\sqrt{T}}\right). \end{aligned}$$

□

We should note that BESTRESP^+ is not necessarily a well-defined algorithm, when the required argmin doesn't have a unique solution. This is relevant in the case when $f(\cdot)$ is not a differentiable function, which is why we have set $y_t = \partial f(x_t)$, any subgradient of f at x_t . On the other hand, the convergence in Theorem 7 holds for any choice of subgradients.

4.3 Accelerated methods for smooth convex optimization

In this subsection, we are going to introduce several *accelerated* algorithms. An optimization procedure is generally referred to as accelerated if it achieves a faster convergence rate relative to a vanilla method, e.g. gradient descent. The convergence of gradient descent on smooth convex optimization problems was known to be $O(1/T)$, and it was believed by many up until the early 1980s that this was the fastest rate achievable despite the only known lower bound of $\Omega(1/T^2)$. It was shown by Nesterov [1] that indeed an upper bound of $O(1/T^2)$ is indeed possible with a slightly more complex update scheme.

We will discuss *Nesterov Accelerated Gradient Descent*, and its variants, in the context of our FGNRD framework. Nesterov's first acceleration method [1, 2] (see also [51]) for the unconstrained setting is frequently described as follows. First initialize z_0 and w_0 arbitrarily, and then iterate as follows:

Algorithm 10 Unconstrained Nesterov Accelerated Gradient Descent

```

1: Input:  $w_0, z_0$  arbitrary
2: for  $t = 1, 2, \dots$ , do
3:    $\theta_t \leftarrow \frac{t}{2(t+1)L}$ ,  $\beta_t \leftarrow \frac{t-1}{t+2}$ 
4:    $w_t \leftarrow z_{t-1} - \theta_t \nabla f(z_{t-1})$ 
5:    $z_t \leftarrow w_t + \beta_t(w_t - w_{t-1})$ 
6: end for
7: return  $w_T$ 
    
```

But to get the ball rolling let us first consider a related method of Polyak [7].

4.3.1 The Heavy Ball Method

In Algorithm 11 we describe the classical iterative version of the Heavy Ball algorithm in the box on the left. One observes that the update looks quite similar to vanilla gradient descent, but there's the addition of a so-called *momentum* term v_t which is the difference of the previous two iterates.

Algorithm 11 Heavy Ball

Given: L -smooth $f(\cdot)$, arbitrary $x_0 = w_0 = w_{-1} \in \mathcal{K}$, iterations T .

$\eta_t \leftarrow \frac{t}{4(t+1)L}, \quad \beta_t \leftarrow \frac{t-1}{t+2}$ $v_t \leftarrow w_{t-1} - w_{t-2}$ $w_t \leftarrow w_{t-1} - \eta_t \nabla f(w_{t-1}) + \beta_t v_t$	$g(x, y) := \langle x, y \rangle - f^*(y)$ $\alpha_t := t \text{ for } t = 1, \dots, T$ $\text{OAlg}^Y := \text{FTL}[\nabla f(x_0)]$ $\text{OAlg}^X := \text{OMD}^+[\frac{1}{2}\ \cdot\ _2^2, x_0, \frac{1}{8L}]$
Iterative Description	FGNRD Equivalence

Output: $w_T = \bar{x}_T$

We provide a formulation of Heavy Ball in the right box of Algorithm 11, and let us begin by stating the equivalence. The proof is deferred to Appendix D.

Proposition 8 *The two interpretations of Heavy Ball described in Algorithm 6 are equivalent when $K = \mathbb{R}^d$.*

Sketch Similar to the proof of Proposition 3, we need to show that the following three equalities are maintained throughout each iteration:

$$\begin{aligned} \nabla f(w_{t-1}) &= y_t \\ \bar{x}_t &= \bar{x}_{t-1} - \frac{\gamma_t \alpha_t^2}{A_t} \nabla f(\bar{x}_{t-1}) + (\bar{x}_{t-1} - \bar{x}_{t-2}) \left(\frac{\alpha_t A_{t-2}}{A_t \alpha_{t-1}} \right), \\ w_t &= \bar{x}_t \end{aligned} \quad (51)$$

The first and last lines follow along the same lines as the proof of Proposition 3. The expression (51) can be established when we see that the $\text{OMD}^+[\frac{1}{2}\|\cdot\|_2^2, x_0, \frac{1}{8L}]$ update algorithm can be written as

$$x_t = x_{t-1} - \frac{\alpha_t}{4L} y_t.$$

Plugging this expression into the definition of \bar{x}_t establishes the update (51). \square

Theorem 9 *The output \bar{x}_T of Algorithm 11 satisfies*

$$f(\bar{x}_T) - \min_{w \in \mathcal{K}} f(w) = O\left(\frac{4L\|x_0 - x^*\|^2 + \sum_{t=1}^T 4L(\|\bar{x}_{t-1} - x_t\|^2 - \|x_{t-1} - x_t\|^2)}{T^2}\right).$$

Furthermore, given a convex constraint set $\mathcal{K} \subset \mathbb{R}^d$ with diameter R , we have $f(\bar{x}_T) - \min_{w \in \mathcal{K}} f(w) = O(\frac{LR}{T})$.

4.3.2 Nesterov's Methods

Starting from 1983, Nesterov has proposed various accelerated methods for smooth convex problems (i.e. [1–4]), all of which are often described as accelerated gradient descent. Our goal now will be to try to understand Nesterov's various algorithms. We will show that all the methods can be generated from *Fenchel game* with some simple tweaks.

Algorithm 12 Nesterov's 1-memory method [4, 34]

Given: L -smooth $f(\cdot)$, convex domain \mathcal{K} , arbitrary $x_0 = v_0 \in \mathcal{K}$, 1-strongly convex distance generating function $\phi(\cdot)$, iterations T .

$\begin{aligned} \beta_t &\leftarrow \frac{2}{t+1}, \gamma_t \leftarrow \frac{t}{4L} \\ z_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_tv_{t-1} \\ v_t &\leftarrow \operatorname{argmin}_{x \in \mathcal{K}} \gamma_t \langle \nabla f(z_t), x \rangle + D_{v_{t-1}}^\phi(x) \\ w_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_tv_t \end{aligned}$
--

Iterative Description

$\begin{aligned} g(x, y) &:= \langle x, y \rangle - f^*(y) \\ \alpha_t &:= t \text{ for } t = 1, \dots, T \\ \text{OAlg}^Y &:= \text{OPTIMISTICFTL} \\ \text{OAlg}^X &:= \text{OMD}^+[\phi(\cdot), x_0, \frac{1}{4L}] \end{aligned}$
--

FGNRD Equivalence

Output: $w_T = \bar{x}_T$,

Let us begin with the version known as the *1-memory method*. To be precise, we adopt the presentation of Nesterov's algorithm given in Algorithm 1 of [34].

Proposition 10 *The two interpretations of Nesterov's 1-memory method in Algorithm 12 are equivalent. That is, the iteratively generated points w_t coincide with the sequence \bar{x}_t produced by the FGNRD dynamic.*

Proof Let us recall the notations (57), $\bar{x}_t := \frac{1}{A_t} \sum_{s=1}^t \alpha_s x_s$ and $\tilde{x}_t := \frac{1}{A_t} \left(\alpha_t x_{t-1} + \sum_{s=1}^{t-1} \alpha_s x_s \right)$. We show, via induction, that the following three equalities are maintained for every t . Note that three objects on the left correspond to the iterative description given in Algorithm 15 whereas the three on the right correspond to the FGNRD description.

$$\nabla f(z_t) = y_t \quad (52)$$

$$v_t = x_t \quad (53)$$

$$w_t = \bar{x}_t. \quad (54)$$

We first note that the initialization ensures that (52) holds for $t = 1$. Also, observe that we have $\beta_t = \frac{\alpha_t}{A_t}$. Therefore,

$$w_t = (1 - \beta_t) w_{t-1} + \beta_t v_t = \left(1 - \frac{\alpha_t}{A_t}\right) w_{t-1} + \frac{\alpha_t}{A_t} v_t = \frac{A_{t-1}}{A_t} w_{t-1} + \frac{\alpha_t}{A_t} v_t \quad (55)$$

From (55), we see that (53) and $w_{t-1} = \bar{x}_{t-1}$ implies (54), as w_t is always an average of the updates v_t . It remains to establish (52) and (53) via induction.

Let us first show (52). But before that, we show $y_t = \nabla f(\tilde{x}_t)$ as follows.

In Algorithm 12 the y -player is using the OPTIMISTICFTL algorithm to choose the sequence of y_t 's, which is given by

$$y_t := \operatorname{argmin}_{y \in \mathcal{Y}} \left\{ \alpha_t \ell_{t-1}(\cdot) + \sum_{s=1}^{t-1} \alpha_s \ell_s(y) \right\}, \quad (56)$$

where the guess $m_t(\cdot)$ to be $m_t(\cdot) \leftarrow \ell_{t-1}(\cdot)$, $\ell_s(y) := f^*(y) - \langle x_s, y \rangle$, and $\ell_0(x_0) := f^*(y) - \langle x_0, y \rangle$ so that $y_0 = \nabla f(x_0)$. We need to consider two parallel sequence of points for both the x - and y -players. For convenience, let

$$\begin{aligned} \bar{x}_t &:= \frac{1}{A_t} \sum_{s=1}^t \alpha_s x_s \\ \tilde{x}_t &:= \frac{1}{A_t} \left(\alpha_t x_{t-1} + \sum_{s=1}^{t-1} \alpha_s x_s \right), \end{aligned} \quad (57)$$

$$\hat{y}_{t+1} := \operatorname{argmin}_y \sum_{s=1}^t \alpha_s (f^*(y) - \langle x_s, y \rangle)$$

Let us now make a few observations about these sequences.

$$\hat{y}_{t+1} = \operatorname{argmax}_y \langle \bar{x}_t, y \rangle - f^*(y) = \nabla f(\bar{x}_t) \quad (58)$$

$$y_t = \operatorname{argmin}_y \{A_T (f^*(y) - \langle \tilde{x}_t, y \rangle)\} = \nabla f(\tilde{x}_t), \quad (59)$$

$$\tilde{x}_t - \bar{x}_t = \frac{\alpha_t}{A_t} (x_{t-1} - x_t). \quad (60)$$

Equations (58) and (59) follow from simple arithmetic as well as elementary properties of Fenchel conjugation and the Legendre transform [16]. Equation (60) is by a simple algebraic calculation.

Given that $y_t = \nabla f(\tilde{x}_t)$ shown in (59), it suffices to show that $\tilde{x}_t = z_t$ for establishing (52). We have

$$\begin{aligned} z_t &= (1 - \beta_t)w_{t-1} + \beta_tv_{t-1} = \left(1 - \frac{\alpha_t}{A_t}\right)\tilde{x}_{t-1} + \frac{\alpha_t}{A_t}x_{t-1} \\ &= \frac{A_{t-1}}{A_t} \frac{\sum_{s=1}^{t-1} \alpha_s x_s}{A_{t-1}} + \frac{\alpha_t}{A_t} x_{t-1} = \frac{\sum_{s=1}^{t-1} \alpha_s x_s + \alpha_t x_{t-1}}{A_t} = \tilde{x}_t. \end{aligned}$$

Hence, (52) holds.

To show (53), observe that the update of v_t is exactly equivalent to OMD^+ for which $\gamma \leftarrow \frac{1}{4L}$, $\alpha_t \leftarrow t$, and $\ell_t(\cdot) \leftarrow \langle y_t, \cdot \rangle$. We thus have completed proof. \square

We remark that when the problem is unconstrained, i.e. $\mathcal{K} = \mathbb{R}^d$, Algorithm 12 with the distance generating function $\phi(\cdot) = \frac{1}{2}\|\cdot\|_2^2$ can be rewritten as Algorithm 10 — the Nesterov's first acceleration method.

Lemma 15 *Consider the FGNRD implementation in Algorithm 12, where the function $f(\cdot)$ is convex and L -smooth with respect to the norm $\|\cdot\|$, whose dual norm is $\|\cdot\|_*$. Regardless of the sequence of points x_1, \dots, x_T generated from the protocol, we have*

$$\alpha\text{-REG}^y \leq L \sum_{t=1}^T \frac{\alpha_t^2}{A_t} \|x_{t-1} - x_t\|^2. \quad (61)$$

Proof By using Lemma 8 with the guess $m_t(\cdot) \leftarrow \ell_{t-1}(\cdot)$, $w_t \leftarrow \hat{y}_t$ of (58), and $z_t \leftarrow y_t$ of (59), and that $\alpha_t(\ell_t(y) - \ell_{t-1}(y)) = \alpha_t \langle x_{t-1} - x_t, y \rangle$ in Fenchel Game, we have

$$\begin{aligned} \sum_{t=1}^T \alpha_t \ell_t(y_t) - \alpha_t \ell_t(y^*) &\leq \sum_{t=1}^T \alpha_t (\ell_t(y_t) - \ell_{t-1}(y_t) - (\ell_t(\hat{y}_{t+1}) - \ell_{t-1}(\hat{y}_{t+1}))) \\ \text{(Eqns. 58, 59)} &= \sum_{t=1}^T \alpha_t \langle x_{t-1} - x_t, \nabla f(\tilde{x}_t) - \nabla f(\bar{x}_t) \rangle \\ \text{(Hölder's Ineq.)} &\leq \sum_{t=1}^T \alpha_t \|x_{t-1} - x_t\| \|\nabla f(\tilde{x}_t) - \nabla f(\bar{x}_t)\|_* \\ \text{(L-smoothness of } f) &\leq L \sum_{t=1}^T \alpha_t \|x_{t-1} - x_t\| \|\tilde{x}_t - \bar{x}_t\| \\ \text{(Eqn. 60)} &= L \sum_{t=1}^T \frac{\alpha_t^2}{A_t} \|x_{t-1} - x_t\| \|x_{t-1} - x_t\| \end{aligned}$$

as desired, where the first inequality is because that $m_t(\cdot) = \ell_{t-1}(\cdot)$. \square

Theorem 11 *Let us consider the output $\bar{x}_T \equiv w_T$ of Algorithm 12. Given that f is L -smooth, and we have $D := \sup_{w \in \mathcal{K}} D_{w_0}^\phi(w)$, then w_T satisfies*

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) \leq \frac{8LD}{T^2}.$$

Proof The equivalence $\bar{x}_T \equiv w_T$ was established in Proposition 10. We may now appeal to Theorem 1 to prove this result, which gives us that

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) \leq \frac{1}{A_T} (\alpha\text{-REG}^x[\text{OPTIMISTICFTL}] + \alpha\text{-REG}^y[\text{OMD}^+]).$$

We have a bound for $\alpha\text{-REG}^x$ from Lemma 10, with parameters $\gamma = \frac{1}{4L}$ and $\beta = 1$, and a bound on $\alpha\text{-REG}^y$ in Lemma 15. Combining these we obtain

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) \leq \frac{1}{A_T} \left(4LD + \sum_{t=1}^T \left(\frac{\alpha_t^2}{A_t} L - 2L \right) \|x_{t-1} - x_t\|^2 \right). \quad (62)$$

Given the choice of weights $\alpha_t = t$, we have $A_t = \frac{t(t+1)}{2}$ and therefore $\frac{\alpha_t^2}{A_t} = \frac{2t^2}{t(t+1)} \leq 2$. With this in mind, the sum on the right hand side of (62) is non-positive. Noticing that $\frac{1}{A_T} \leq \frac{2}{T^2}$ completes the proof. \square

It is worth dwelling on exactly how we obtained the above result. A less refined analysis of the OMD⁺ algorithm would have simply ignored the negative summation term in Lemma 10, and simply upper bounded this by 0. But the negative terms $\|x_t - x_{t-1}\|^2$ in this sum happen to correspond *exactly* to the positive terms one obtains in the regret bound for the y -player, but this is true *only as a result of* using the OPTIMISTICFTL algorithm. To obtain a cancellation of these terms, we need a γ_t which is roughly constant, and hence we need to ensure that $\frac{\alpha_t^2}{A_t} = O(1)$. The final bound, of course, is determined by the inverse quantity $\frac{1}{A_T}$, and a quick inspection reveals that the best choice of $\alpha_t = \theta(t)$. This is not the only choice that could work, and we conjecture that there are scenarios in which better bounds are achievable for different α_t tuning. We show in Subsection 4.4 that a *linear rate* is achievable when $f(\cdot)$ is also strongly convex, and there we tune α_t to grow exponentially in t rather than linearly.

Infinite memory method.

We finish this section by mentioning one additional algorithm. Nesterov has proposed another first-order method with $O(1/T^2)$ convergence, although the technique has the slight downside in that it requires maintaining the full sequence of gradients observed along the algorithm’s path. But this method also has a natural and simple interpretation in our FGNRD framework, and we include it here for completeness.

We leave it to the reader to establish the equivalence described in Algorithm 13, and here we state the convergence rate whose proof is deferred to Appendix E.

Theorem 12 *Assume that $f(\cdot)$ is L -smooth convex. Nesterov’s ∞ -memory method (Algorithm 13) satisfies*

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) \leq O \left(\frac{L(R(x^*) - \min_{x \in \mathcal{K}} R(x))}{T^2} \right).$$

Algorithm 13 Nesterov's ∞ -memory method [3, 34]

Given: L -smooth $f(\cdot)$, convex domain \mathcal{K} , arbitrary $v_0 = x_0 \in \mathcal{K}$, 1-strongly convex regularizer $R(\cdot)$, iterations T .

$$\begin{aligned} \beta_t &\leftarrow \frac{2}{t+1}, \gamma_t \leftarrow \frac{t}{4L} \\ z_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_t v_{t-1} \\ v_t &\leftarrow \operatorname{argmin}_{x \in \mathcal{K}} \sum_{s=1}^t \gamma_s \langle \nabla f(z_s), x \rangle + R(x) \\ w_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_t v_t \end{aligned}$$

Iterative Description

$$\begin{aligned} g(x, y) &:= \langle x, y \rangle - f^*(y) \\ \alpha_t &:= t \text{ for } t = 1, \dots, T \\ \text{OAlg}^Y &:= \text{OPTIMISTICFTL} \\ \text{OAlg}^X &:= \text{FTRL}^+ [R(\cdot), \frac{1}{4L}] \end{aligned}$$

FGNRD Equivalence

Output: $w_T = \bar{x}_T$ **4.3.3 Accelerated proximal methods****Algorithm 14** Accelerated proximal method.

Given: L -smooth $f(\cdot)$, arbitrary $v_0 = x_0 \in \mathbb{R}^d$, iterations T .

$$\begin{aligned} \beta_t &\leftarrow \frac{2}{t+1}, \gamma_t \leftarrow \frac{t}{4L} \\ z_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_t v_{t-1} \\ v_t &\leftarrow \mathbf{prox}_{t\gamma\psi}(x_{t-1} - t\gamma\nabla f(z_t)) \\ w_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_t v_t \end{aligned}$$

Iterative Description

$$\begin{aligned} g(x, y) &:= \langle x, y \rangle - f^*(y) + \psi(x) \\ \alpha_t &:= t \text{ for } t = 1, \dots, T \\ \text{OAlg}^Y &:= \text{OPTIMISTICFTL} \\ \text{OAlg}^X &:= \text{OMD}^+ [\phi(\cdot) = \frac{1}{2}\|\cdot\|_2^2, \frac{1}{4L}] \end{aligned}$$

FGNRD Equivalence

Output: $w_T = \bar{x}_T$

In this section, we consider solving composite optimization problems

$$\min_{x \in \mathbb{R}^d} f(x) + \psi(x), \quad (63)$$

where $f(\cdot)$ is smooth convex but $\psi(\cdot)$ is possibly non-differentiable convex. Examples of $\psi(\cdot)$ include ℓ_1 norm ($\|\cdot\|_1$), ℓ_∞ norm ($\|\cdot\|_\infty$), and elastic net ($\|\cdot\|_1 + \gamma\|\cdot\|^2$ for $\gamma > 0$). We want to show that the game analysis still applies to this problem. We just need to change the payoff function g to account for $\psi(x)$. Specifically, we consider the following two-player zero-sum game,

$$\min_x \max_y \{g(x, y) := \langle x, y \rangle - f^*(y) + \psi(x)\}. \quad (64)$$

Notice that the minimax value of the game is $\min_x f(x) + \psi(x)$, which is exactly the optimum value of the composite optimization problem. Let us denote the proximal operator as $\mathbf{prox}_{\lambda\psi}(v) := \operatorname{argmin}_x (\psi(x) + \frac{1}{2\lambda}\|x - v\|_2^2)$.²

²It is known that for some $\psi(\cdot)$, their corresponding proximal operations have closed-form solutions (see e.g. [52] for details).

We remark that the FGNRD interpretation in Algorithm 14 is essentially the same as Algorithm 12, except here the payoff function g is defined differently (64). The weighting scheme α_t and the players' strategies remain the same, and we thus omit the equivalence between the two interpretations of Algorithm 14.

In this new game, the x -player plays OMD⁺ with the distance generating function $\phi(x) = \frac{1}{2}\|x\|_2^2$ and receives the loss functions $\alpha_t h_t(x) := \alpha_t \{\langle x, y_t \rangle + \psi(x)\}$, which leads to the following update,

$$\begin{aligned} x_t &= \underset{x}{\operatorname{argmin}} \gamma(\alpha_t h_t(x)) + D_{x_{t-1}}^\phi(x) = \underset{x}{\operatorname{argmin}} \gamma(\alpha_t \{\langle x, y_t \rangle + \psi(x)\}) + D_{x_{t-1}}^\phi(x) \\ &= \underset{x}{\operatorname{argmin}} \psi(x) + \frac{1}{2\alpha_t \gamma} (\|x\|_2^2 + 2\langle \alpha_t \gamma y_t - x_{t-1}, x \rangle) \\ &= \mathbf{prox}_{\alpha_t \gamma \psi}(x_{t-1} - \alpha_t \gamma \nabla f(\tilde{x}_t)), \end{aligned}$$

where the last equality follows that $y_t = \nabla f(\tilde{x}_t)$ and the definition of the operator $\mathbf{prox}(\cdot)$. One can view Algorithm 14 as a variant of the so called ‘‘Accelerated Proximal Gradient’’ in [8]. Yet, the design and analysis of our algorithm is simpler than that of [8].

Theorem 13 *Suppose that $D_{x_0}^\phi(x^*) \leq D$. The weighted average of \bar{x}_T in Algorithm 14 satisfies*

$$f(\bar{x}_T) - \min_x f(x) \leq O\left(\frac{LD}{T^2}\right).$$

Proof Even though the payoff function $g(\cdot, \cdot)$ is a bit different, the proof still essentially follows the same line as Theorem 11, as y -player plays OPTIMISTICFTL and the x -player plays OMD⁺. □

4.3.4 Related works

In recent years, there are growing interest in giving new interpretations of Nesterov’s accelerated algorithms or proposing new variants. For example, [34] gives a unified analysis for some Nesterov’s accelerated algorithms [3–5], using the standard techniques and analysis in optimization literature. [53, 54] connects the design of accelerated algorithms with dynamical systems and control theory. [55, 56] gives a geometric interpretation of the Nesterov’s method for unconstrained optimization, inspired by the ellipsoid method. [57] studies the Nesterov’s methods and the HEAVYBALL method for quadratic non-strongly convex problems by analyzing the eigen-values of some linear dynamical systems. [58] proposes a variant of accelerated algorithms by mixing the updates of gradient descent and mirror descent and showing the updates are complementary. [59, 60] propose a primal-dual view that recovers several first-order algorithms with careful discretizations of a continuous-time dynamic, which also leads to a new accelerated extra-gradient descent method. [61] show a

simple acceleration proof of mirror prox [62] and dual extrapolation [63] based on solving the Fenchel game. [51, 64–69] analyze the acceleration algorithms via the lens of ordinary differential equations and the numerical analysis. See also a monograph [70] for various methods of acceleration. Finally, we note an independent work [71, 72] provide a game interpretation of Nesterov’s accelerated method but does not have the idea of the regret analysis. In our work, we show a deeper connection between online learning and optimization, and propose a modular framework that is not limited to Nesterov’s method.

4.4 Proving Accelerated Linear Rates

Algorithm 15 Accelerated Gradient with Linear Convergence

Given: L -smooth, μ -strongly convex $f(\cdot)$, convex domain \mathcal{K} , iterations T , and a distance generating function $\phi(\cdot)$ that is 1-strongly convex, L_ϕ -smooth, and differentiable. Finally, let $R(x) = \phi(x)$.

Init: $w_0 = v_0 = x_0 = \operatorname{argmin}_{x \in \mathcal{K}} \phi(x)$.

$\beta \leftarrow \frac{1}{2} \sqrt{\frac{\mu}{L(1+L_\phi)}}, \gamma_t \leftarrow \alpha_t$ $z_t \leftarrow (1 - \beta)w_{t-1} + \beta v_{t-1}$ $\Phi_t(x) := \sum_{s=1}^t \gamma_s (\langle \nabla \tilde{f}(z_s), x \rangle + \mu \phi(x))$ $v_t \leftarrow \operatorname{argmin}_{x \in \mathcal{K}} R(x) + \Phi_t(x)$ $w_t \leftarrow (1 - \beta)w_{t-1} + \beta v_t$	$\tilde{f}(x) := f(x) - \mu \phi(x)$ $g(x, y) := \langle x, y \rangle - \tilde{f}^*(y) + \mu \phi(x)$ $\alpha_1 = \frac{1}{2L(1+L_\phi)}$ $\frac{\alpha_t}{A_t} = \frac{1}{2} \sqrt{\frac{\mu}{L(1+L_\phi)}}$ <p style="text-align: center;">for $t = 2, \dots, T$</p> $\text{OAlg}^Y := \text{OPTIMISTICFTL}$ $\text{OAlg}^X := \text{FTRL}^+[\phi, 1]$
Iterative Description	FGNRD Equivalence

Output: $w_T = \bar{x}_T$

Nesterov observed that for strongly convex smooth functions, one can achieve an accelerated linear rate (e.g. page 71-81 of [5]). It is natural to ask if the zero-sum game and regret analysis in the present work also recovers this faster rate in the same fashion. We answer this in the affirmative. Assume $f(\cdot)$ is μ -strongly convex in the following sense (see also Section 3.3 of [72]),

$$f(z) \geq f(x) + \langle \nabla f(x), z - x \rangle + \mu D_x^\phi(z), \quad (65)$$

for all $z, x \in \mathcal{K}$, where $D_x^\phi(z)$ is the Bregman divergence. When the distance generating function is $\phi(x) := \frac{1}{2} \|x\|_2^2$, the strong convexity condition (65) becomes

$$f(z) \geq f(x) + \langle \nabla f(x), z - x \rangle + \frac{1}{2} \|z - x\|_2^2. \quad (66)$$

It is known that the function $\tilde{f}(x) := f(x) - \mu \phi(x)$ is a convex function for all $x \in \mathcal{K}$ (see e.g. [73]). Based on this property, we consider a new game

$$\tilde{g}(x, y) := \langle x, y \rangle - \tilde{f}^*(y) + \mu \phi(x), \quad (67)$$

where the minimax value of the game is $V^* := \min_x \max_y \tilde{g}(x, y) = \min_x \tilde{f}(x) + \mu\phi(x) = \min_x f(x)$. We begin by showing the following in this new game.

Proposition 14 *The two interpretations of Algorithm 15 are equivalent. That is, the iteratively generated points w_t coincide with the sequence \bar{x}_t produced by the FGNRD dynamic.*

Proof The proof follows the same lines as that of Proposition 10 until the last line by setting $\beta_t = \beta = \frac{1}{2} \sqrt{\frac{\mu}{L(1+L_\phi)}} = \frac{\alpha_t}{A_t}$. We only need to replace the last line of the proof. Specifically, the update of v_t is exactly equivalent to FTRL^+ on the loss sequences $\alpha_t \ell_t(\cdot) \leftarrow \alpha_t (\langle \nabla \tilde{f}(z_t), \cdot \rangle + \mu\phi(\cdot))$. Hence, the proof is completed. \square

We now prove the accelerated linear convergence rate of Algorithm 15. First, we denote the *condition number* $\kappa := \frac{L}{\mu}$.

Theorem 15 *Assume we are given a norm $\|\cdot\|$, and that we have a distance-generating function $\phi(\cdot)$ that is differentiable, L_ϕ -smooth, and 1-strongly convex with respect to $\|\cdot\|$. Assume we also have a function $f(\cdot)$ that is L -smooth with respect to $\|\cdot\|$ and μ -strongly convex in the sense of (65). Then the output of Algorithm 15 satisfies*

$$f(w_T) - \min_w f(w) \leq c \exp\left(-\frac{1}{2\sqrt{\kappa}\sqrt{1+L_\phi}}T\right) (\phi(w_*) - \phi(w_0))$$

for some constant $c := 2L(1+L_\phi) > 0$.

Proof As the proof of Lemma 15, we first bound the regret of the y-player as follows.

$$\begin{aligned} \sum_{t=1}^T \alpha_t \ell_t(y_t) - \alpha_t \ell_t(y^*) &\leq \sum_{t=1}^T \alpha_t \langle x_{t-1} - x_t, y_t - \hat{y}_{t+1} \rangle \\ \text{(Eqns. 58, 59)} \quad &= \sum_{t=1}^T \alpha_t \langle x_{t-1} - x_t, \nabla \tilde{f}(\tilde{x}_t) - \nabla \tilde{f}(\bar{x}_t) \rangle \\ \text{(Hölder's Ineq.)} \quad &\leq \sum_{t=1}^T \alpha_t \|x_{t-1} - x_t\| \|\nabla \tilde{f}(\tilde{x}_t) - \nabla \tilde{f}(\bar{x}_t)\|_* \\ &= \sum_{t=1}^T \alpha_t \|x_{t-1} - x_t\| \|\nabla f(\tilde{x}_t) - \mu \nabla \phi(\tilde{x}_t) - \nabla f(\bar{x}_t) + \mu \nabla \phi(\bar{x}_t)\|_* \\ \text{(Triangle Ineq.)} \quad &\leq \sum_{t=1}^T \alpha_t \|x_{t-1} - x_t\| (\|\nabla f(\tilde{x}_t) - \nabla f(\bar{x}_t)\|_* + \mu L_\phi \|\bar{x}_t - \tilde{x}_t\|) \\ \text{(smoothness)} \quad &\leq L(1+L_\phi) \sum_{t=1}^T \alpha_t \|x_{t-1} - x_t\| \|\tilde{x}_t - \bar{x}_t\| \end{aligned}$$

$$\text{(Eqn. 60)} \quad = L(1 + L_\phi) \sum_{t=1}^T \frac{\alpha_t^2}{A_t} \|x_{t-1} - x_t\| \|x_{t-1} - x_t\|.$$

So the regret satisfies

$$\alpha\text{-REG}^y \leq L(1 + L_\phi) \sum_{t=1}^T \frac{\alpha_t^2}{A_t} \|x_{t-1} - x_t\|^2. \quad (68)$$

For the x-player, according to Lemma 7, its regret is

$$\alpha\text{-REG}^x \leq R(x^*) - R(x_0) - \sum_{t=1}^T \left(\frac{\mu A_{t-1}}{2} + \frac{1}{2} \right) \|x_{t-1} - x_t\|_2^2, \quad (69)$$

where $x_0 = \operatorname{argmin}_x R(x)$. Summing (68) and (69), we have

$$\begin{aligned} & \alpha\text{-REG}^y[\text{OPTIMISTICFTL}] + \alpha\text{-REG}^x[\text{FTRL}^+] \\ & \leq R(x^*) - R(x_0) + \sum_{t=1}^T \left(\frac{L(1 + L_\phi)\alpha_t^2}{A_t} - \frac{\mu A_{t-1} + 1}{2} \right) \|x_{t-1} - x_t\|_2^2. \end{aligned}$$

By choosing the weight $\{\alpha_t\}$ to satisfy $\alpha_1 = \frac{1}{2L(1+L_\phi)}$ and that for $t \geq 2$, $\frac{\alpha_t}{A_t} = \frac{1}{2} \sqrt{\frac{\mu}{L(1+L_\phi)}}$, the coefficient of the distance terms will be non-positive for all t , i.e. $\left(\frac{L(1+L_\phi)\alpha_t^2}{A_t} - \frac{\mu A_{t-1}}{2} \right) \leq 0$, which means that the distance terms will cancel out. Therefore, the optimization error ϵ after T iterations satisfies that

$$\begin{aligned} \epsilon & \leq \frac{\alpha\text{-REG}^y + \alpha\text{-REG}^x}{A_T} \leq \frac{1}{A_1} \frac{A_1}{A_2} \dots \frac{A_{T-1}}{A_T} (R(x^*) - R(x_0)) \\ & = \frac{1}{A_1} \left(1 - \frac{\alpha_2}{A_2} \right) \dots \left(1 - \frac{\alpha_T}{A_T} \right) (R(x^*) - R(x_0)) \\ & \leq \frac{1}{A_1} \left(1 - \frac{\alpha_2}{\bar{A}_2} \right) \dots \left(1 - \frac{\alpha_T}{\bar{A}_T} \right) (R(x^*) - R(x_0)) \\ & \leq \left(1 - \frac{1}{2\sqrt{1 + L_\phi\sqrt{\kappa}}} \right)^{T-1} \frac{R(x^*) - R(x_0)}{A_1} = O \left(\exp \left(-\frac{1}{2\sqrt{1 + L_\phi\sqrt{\kappa}}} T \right) \right). \end{aligned}$$

□

5 New algorithms

In Section 4 we established that the FGNRD framework can be used to describe a set of known algorithms for convex optimization under various assumptions. But we have argued that the framework is indeed quite generic, given that we have flexibility to choose different no-regret update algorithms, to change the game's payoff function, and to modify the protocol. In the present section we use FGNRD to describe two novel algorithms, and to provide strong convergence rates using the same methodology.

5.1 Boundary Frank-Wolfe

In original formulation of the Fenchel Game No-Regret Dynamics, Protocol 2, we specified that the two players act in a particular order, first the y -player

followed by the x -player. This ordering is important, as it allows us to consider scenarios in which x_t can be chosen in response to y_t , as was the case in nearly all of the algorithms proposed in Section 4. But we can instead reverse the order of the two players, allowing the x -player to act before the y -player, while otherwise following the same protocol. This will give us a new algorithm for constrained convex optimization, *Boundary Frank-Wolfe*, which has the unusual property that it only computes gradients of f on the boundary of the constraint set \mathcal{K} . We will show the algorithm has a $O(\log T/T)$ convergence rate for any convex Lipschitz f , without any smoothness nor strong-convexity requirement for f , yet under two conditions: (1) the true minimizer of f in \mathcal{K} must exist on the boundary, and \mathcal{K} must be a *strongly convex set*. The result relies on recent work in online convex optimization under strongly convex constraints [32]. (Notably, there is prior work that develops a Frank-Wolfe-like algorithm that strongly convex decision sets [21] with $O(1/T^2)$ convergence, but this result requires that f is both smooth and strongly convex as well.)

Algorithm 16 Boundary Frank-Wolfe

Given: possibly non-smooth convex $f(\cdot)$, convex domain \mathcal{K} , arbitrary $w_0 = z_0$, iterations T .

$$\begin{aligned} z_t &\leftarrow \operatorname{argmin}_{z \in \mathcal{K}} \frac{1}{t-1} \sum_{s=1}^{t-1} \langle z, \partial f(z_s) \rangle \\ w_t &\leftarrow \frac{(t-1)w_{t-1} + z_t}{t} \end{aligned}$$

Iterative Description

$$\begin{aligned} g(x, y) &:= \langle x, y \rangle - f^*(y) \\ \alpha_t &\leftarrow 1 \\ \text{OAlg}^X &:= \text{FTL}[z_0] \\ \text{OAlg}^Y &:= \text{BESTRESP}^+ \end{aligned}$$

FGNRD Equivalence

Output: $w_T := \bar{x}_T$.

The equivalence between the two forms of Algorithm 16 can be proven in the usual fashion, and so we omit the proof of the following proposition.

Proposition 16 *The two interpretations of Algorithm 16 are equivalent. That is, the iteratively generated points w_t coincide with the sequence \bar{x}_t produced by the FGNRD dynamic.*

Theorem 17 *Assume that the constraint set \mathcal{K} is λ -strongly convex, and that $\sum_{s=1}^t \partial f(x_s)$ has non-zero norm, then Boundary Frank-Wolfe (Algorithm 16) has*

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) = O\left(\frac{M \log T}{\lambda L_T T}\right),$$

where $M := \sup_{x \in \mathcal{K}} \|\partial f(x)\|$, $\Theta_t := \sum_{s=1}^t \frac{1}{t} \partial f(x_s)$, and $L_T := \min_{1 \leq t \leq T} \|\Theta_t\|$.

Proof Since y -player plays BESTRESP^+ , its regret ≤ 0 . For the x -player, we use Lemma 4, which grants $\alpha\text{-REG}^x \leq O(\frac{M \log T}{\lambda L_T})$. Using Theorem 1 we have

$$f(\bar{x}_T) - \min_{x \in \mathcal{K}} f(x) \leq \frac{1}{T} (\alpha\text{-REG}^x[\text{FTL}] + \alpha\text{-REG}^y[\text{BESTRESP}^+])$$

$$= O\left(\frac{M \log T}{\lambda L_T T}\right)$$

as desired. \square

Note that the rate depends crucially on L_T , which is the smallest averaged-gradient norm computed during the optimization. According to Theorem 2, the closure of the gradient space is a convex set. This implies the following lemma that ensures L_T is bounded away from 0.

Lemma 16 *If $\mathbf{0} \notin \text{closure}(\{\nabla f(x) : x \in \mathcal{K}\})$, then the average of gradient norm L_T satisfies $L_T > 0$.*

Hence, the proposed algorithm may only be useful when it is known, a priori, that the solution x^* will occur not in the interior but on the boundary of \mathcal{K} . It is indeed an odd condition, but it does hold in many typical scenarios. One may add a perturbed vector to the gradient and show that with high probability, L_T is a non-zero number. The downside of this approach is that it would generally grant a slower convergence rate.

5.2 Gauge Frank-Wolfe

Using the FGNRD framework further, we propose a new algorithm that is “FRANK-WOLFE-like” in the following sense: we can optimize a smooth objective function using only linear optimization oracle queries. This method requires the constraint set to be “suitably round” but achieves a rate of $O(1/T^2)$ without additional assumptions on f . The main trick is to utilize a regularization function that is adapted to the constraint set \mathcal{K} . With this in mind, define the “gauge function” of \mathcal{K} [74, 75] as

$$\gamma_{\mathcal{K}}(x) := \inf\{c \geq 0 : \frac{x}{c} \in \mathcal{K}\}. \quad (70)$$

Notice that, for a closed convex \mathcal{K} that contains the origin, one has $\mathcal{K} = \{x \in \mathbb{R}^d : \gamma_{\mathcal{K}}(x) \leq 1\}$. Furthermore, the boundary points on \mathcal{K} satisfy $\gamma_{\mathcal{K}}(x) = 1$.

Theorem 18 (λ -Gauge set) *(Theorem 4 in [76] and Theorem 2 in [77]) Let \mathcal{K} be a $\frac{\lambda}{2}$ -strongly convex set that contains the origin in its interior. Then, its squared gauge function, i.e. $\gamma_{\mathcal{K}}^2(\cdot)$, is λ -strongly-convex.*

Examples of the λ -Gauge set are given in Appendix F.

Algorithmically, in our two-player game framework, we will utilize FTRL⁺ for the x -player with regularization $\gamma_{\mathcal{K}}^2(\cdot)$. Naturally, this requires us to solve problems of the form

$$x = \operatorname{argmin}_{x \in \mathcal{K}} \langle L, x \rangle + \gamma_{\mathcal{K}}^2(x), \quad (71)$$

for arbitrary vectors L . While at first glance this does not appear to be a linear optimization problem, let us reparametrize the problem as follows. Denote $\text{bndry}(\mathcal{K})$ as the boundary of the constraint set \mathcal{K} , and notice that (71) is equivalent to solving

$$\min_{\rho \in [0,1], z \in \text{bndry}(\mathcal{K})} \langle L, \rho z \rangle + \gamma_{\mathcal{K}}^2(\rho z) = \min_{\rho \in [0,1]} \left[\left(\min_{z \in \text{bndry}(\mathcal{K})} \langle L, z \rangle \right) \rho + \rho^2 \right]. \quad (72)$$

Notice that the inner minimization is now a linear problem, and the outer problem is a 1-parameter constrained quadratic problem which can be solved directly. In summary, we have

$$\underset{x \in \mathcal{K}}{\text{argmin}} \langle L, x \rangle + \gamma_{\mathcal{K}}^2(x) = \rho^* z^* \quad \text{where} \quad \begin{cases} z^* = \underset{z \in \text{bndry}(\mathcal{K})}{\text{argmin}} \langle L, z \rangle, \\ \rho^* = \max(0, \min(1, -\langle L, z^* \rangle / 2)). \end{cases} \quad (73)$$

Algorithm 17 Gauge Frank-Wolfe (smooth convex $f(\cdot)$)

Given: L -smooth convex $f(\cdot)$, arbitrary $v_0 = x_0 \in \mathcal{K}$, iteration T .

$\begin{aligned} \beta_t &\leftarrow \frac{2}{t+1}, \gamma_t \leftarrow \frac{t}{4L} \\ z_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_t v_{t-1} \\ z_t^* &\leftarrow \underset{z \in \text{bndry}(\mathcal{K})}{\text{argmin}} \langle \sum_{s=1}^t \gamma_s \nabla f(z_s), z \rangle \\ \gamma_t &\leftarrow \frac{1}{2} \langle \sum_{s=1}^t \gamma_s \nabla f(z_s), z^* \rangle \\ \rho_t^* &\leftarrow \max(0, \min(1, -\gamma_t)) \\ v_t &\leftarrow \rho_t^* z_t^* \\ w_t &\leftarrow (1 - \beta_t)w_{t-1} + \beta_t v_t \end{aligned}$	$\begin{aligned} g(x, y) &:= \langle x, y \rangle - f^*(y) \\ \alpha_t &:= t \text{ for } t = 1, \dots, T \\ \text{OAlg}^Y &:= \text{OPTIMISTICFTL} \\ \text{OAlg}^X &:= \text{FTRL}^+[\gamma_{\mathcal{K}}^2(\rho x), \frac{1}{4L}] \end{aligned}$
--	---

Iterative Description

FGNRD Equivalence

Output: $w_T = \bar{x}_T$,

Theorem 19 *Suppose the constraint set \mathcal{K} is a λ -Gauge set. Assume that the function $f(\cdot)$ is L -smooth convex with respect to the induced gauge norm $\gamma_{\mathcal{K}}(\cdot)$. Suppose that the step size η satisfies $\frac{1}{CL} \leq \frac{\eta}{\lambda} \leq \frac{1}{4L}$ for some constant $C \geq 4$. Then, the output \bar{x}_T of Algorithm 17 satisfies*

$$f(\bar{x}_T) - \min_{x \in \mathcal{K}} f(x) \leq \frac{2CL}{\lambda T^2}.$$

Proof We have just shown that line 4-5 is due to that the x-player plays FTRL^+ with the square of the gauge function as the regularizer. So Algorithm 17 is an instance of Algorithm 13, and we can invoke Theorem 12 to obtain the convergence rate, noting that $\gamma_{\mathcal{K}}^2(x^*) \leq 1$. \square

We want to emphasize again that our analysis does not need the function $f(\cdot)$ to be strongly convex to show $O(1/T^2)$ rate. On the other hand, [21] shows

the $O(1/T^2)$ rate under the additional assumption that the function is strongly convex. [21] raises an open question if an accelerated linear rate is possible under the assumption that the function is smooth and strongly convex. We partially answer the open question through Algorithm 18 and Theorem 20.

Algorithm 18 Gauge Frank-Wolfe (smooth and strongly convex $f(\cdot)$)

Given: L -smooth and μ -strongly convex $f(\cdot)$, arbitrary $w_0 = v_0 = x_0 \in \mathcal{K}$, iterations T .

$$\begin{aligned}
 \beta &\leftarrow \frac{1}{2} \sqrt{\frac{\mu}{L(1+L_\phi/\lambda)}}, \gamma_t \leftarrow \alpha_t \\
 z_t &\leftarrow (1-\beta)w_{t-1} + \beta v_{t-1} \\
 z_t^* &\leftarrow \operatorname{argmin}_{z \in \operatorname{bdry}(\mathcal{K})} \langle \sum_{s=1}^t \gamma_s \nabla \tilde{f}(z_s), z \rangle \\
 \gamma_t &\leftarrow \frac{\langle \sum_{s=1}^t \gamma_s \nabla \tilde{f}(z_s), z_t^* \rangle}{2(\frac{1}{\mu} + \sum_{s=1}^t \gamma_s) \frac{\mu}{\lambda}} \\
 \rho_t^* &\leftarrow \max(0, \min(1, -\gamma_t)) \\
 v_t &\leftarrow \rho_t^* z_t^* \\
 w_t &\leftarrow (1-\beta)w_{t-1} + \beta v_t
 \end{aligned}$$

Iterative Description

$$\begin{aligned}
 g(x, y) &:= \langle x, y \rangle - \tilde{f}^*(y) + \frac{\mu}{\lambda} \gamma_{\mathcal{K}}^2(\cdot) \\
 &\quad \text{where } \tilde{f}(x) := f(x) - \frac{1}{\lambda} \gamma_{\mathcal{K}}^2(\cdot) \\
 \alpha_1 &:= \frac{1}{2L(1+L_\phi/\lambda)} \\
 \frac{\alpha_t}{A_t} &:= \frac{1}{2} \sqrt{\frac{\mu}{L(1+L_\phi/\lambda)}}, \\
 &\quad \text{for } t = 2, \dots, T \\
 \text{OAlg}^Y &:= \text{OPTIMISTICFTL} \\
 \text{OAlg}^X &:= \text{FTRL}^+[\frac{1}{\lambda} \gamma_{\mathcal{K}}^2(\cdot), 1]
 \end{aligned}$$

FGNRD Equivalence

Output: $w_T = \bar{x}_T$,

Theorem 20 *Suppose the constraint set \mathcal{K} is a λ -Gauge set and the induced square of the gauge function $\gamma_{\mathcal{K}}^2(\cdot)$ is differentiable and L_ϕ -smooth. Assume that the function $f(\cdot)$ is L -smooth with respect to the induced gauge norm $\gamma_{\mathcal{K}}(\cdot)$ and $f(\cdot)$ is μ -strongly convex satisfying (65) with $\frac{1}{\lambda} \gamma_{\mathcal{K}}^2(\cdot)$ being the distance generating function. Then, the output \bar{x}_T of Algorithm 18 satisfies*

$$f(\bar{x}_T) - \min_{x \in \mathcal{K}} f(x) = O\left(\exp\left(-\frac{T}{2\sqrt{1+L_\phi/\lambda\sqrt{\kappa}}}\right)\right).$$

Lemma 2.2 and Theorem A.2 in [78] implies that if a strongly convex set \mathcal{K} which is centrally-symmetric at the origin satisfies the notion of a smooth set ³, then the induced square of the gauge function $\gamma_{\mathcal{K}}^2(\cdot)$ is L_ϕ -smooth and differentiable for some constant $L_\phi > 0$; consequently, the condition of Theorem 20 holds. Examples include l_p balls and Schatten p balls for $1 < p \leq 2$ [78].

Proof The equivalence of the two displays in Algorithm 18 can be shown by following that of the proof of Proposition 14, where the update FTRL^+ :

$$x_t \leftarrow \operatorname{argmin}_{x \in \mathcal{K}} \sum_{s=1}^t \alpha_s h_s(x) + R(x) = \operatorname{argmin}_{x \in \mathcal{K}} \sum_{s=1}^t \alpha_s \left(\langle y_s, x \rangle + \frac{\mu}{\lambda} \gamma_{\mathcal{K}}^2(x) \right) + \frac{1}{\lambda} \gamma_{\mathcal{K}}^2(x)$$

³For a compact convex set \mathcal{K} , it is smooth if and only if $\#\{N_{\mathcal{K}}(z) \cap \partial \mathcal{K}^\circ\} = 1$ for any point z on the boundary of \mathcal{K} , i.e. $z \in \partial \mathcal{K}$, where $N_{\mathcal{K}}(z) := \{w \in \mathbb{R}^d : \langle z - x, w \rangle \geq 0, \forall x \in \mathcal{K}\}$ is the normal cone at z and $\mathcal{K}^\circ := \{w \in \mathbb{R}^d : \langle w, x \rangle \leq 1, \forall x \in \mathcal{K}\}$ is the polar of \mathcal{K} , see e.g. Definition 2.1 in [78].

is equivalent to solving

$$(\rho_t^*, z_t^*) \leftarrow \operatorname{argmin}_{\rho \in [0,1]} \left(\operatorname{argmin}_{z \in \operatorname{bndry}(\mathcal{K})} \left\langle \sum_{s=1}^t \alpha_s \nabla \tilde{f}(z_s), z \right\rangle \rho + \left(\frac{1}{\mu} + \sum_{s=1}^t \alpha_s \right) \frac{\mu}{\lambda} \rho^2 \right),$$

and setting $v_t = \rho_t^* z_t^*$. Specifically, Algorithm 18 is an instance of Algorithm 15 with the regularization of FTRL^+ being the square of the gauge function, i.e. $\phi(\cdot) = \frac{1}{\lambda} \gamma_{\mathcal{K}}^2(\cdot)$. Hence, the convergence rate can be shown by following the proof of Theorem 15, where we bound the Lipschitzness of $\nabla \tilde{f}(\cdot)$ as $\|\nabla \tilde{f}(\tilde{x}_t) - \nabla \tilde{f}(\tilde{x}_t)\| \leq \|\nabla f(\tilde{x}_t) - \frac{\mu}{\lambda} \nabla \phi(\tilde{x}_t) - \nabla f(\tilde{x}_t) + \frac{\mu}{\lambda} \nabla \phi(\tilde{x}_t)\|_* \leq \|\nabla f(\tilde{x}_t) - \nabla f(\tilde{x}_t)\|_* + \mu \frac{L_\phi}{\lambda} \|\tilde{x}_t - \tilde{x}_t\| \leq L \left(1 + \frac{L_\phi}{\lambda} \right) \|\tilde{x}_t - \tilde{x}_t\|$. \square

5.3 Cumulative gradient descent

We describe yet another algorithm from the game for non-smooth optimization. Algorithm 19 below has the last-iteration guarantee, while subgradient descent method, i.e. $w_t = w_{t-1} - \partial f(w_{t-1})$, does not enjoy the last-iteration guarantee, see e.g. Chapter 2 of [79] or [80]. This highlights the benefit of the proposed algorithm.

Algorithm 19 Cumulative gradient descent

Given: Convex $f(\cdot)$ and iterations T .

Init: $w_0 = x_0 \in \mathcal{K}$.

$\begin{aligned} \eta &\leftarrow \frac{(R/G)}{\sqrt{T}} \\ w_t &= \left(1 - \frac{1}{t}\right)w_{t-1} - \frac{1}{t}\eta \sum_{s=1}^{t-1} \partial f(w_s) \\ &= \left(1 - \frac{1}{t}\right)w_{t-1} - \frac{1}{t}\eta \partial f(w_{t-1}) \\ &\quad - \frac{t-1}{t}\eta \left(w_{t-1} - \left(1 - \frac{1}{t-1}\right)w_{t-2}\right) \end{aligned}$	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px;">$g(x, y) := \langle x, y \rangle - f^*(y)$</td> </tr> <tr> <td style="padding: 5px;">$\alpha_t := 1$ for $t = 1, \dots, T$</td> </tr> <tr> <td style="padding: 5px;">$\text{OAlg}^X := \text{OMD}[\frac{1}{2}\ \cdot\ _2^2, x_0, \frac{(R/G)}{\sqrt{T}}]$</td> </tr> <tr> <td style="padding: 5px;">$\text{OAlg}^Y := \text{FTL}^+$</td> </tr> </table>	$g(x, y) := \langle x, y \rangle - f^*(y)$	$\alpha_t := 1$ for $t = 1, \dots, T$	$\text{OAlg}^X := \text{OMD}[\frac{1}{2}\ \cdot\ _2^2, x_0, \frac{(R/G)}{\sqrt{T}}]$	$\text{OAlg}^Y := \text{FTL}^+$
$g(x, y) := \langle x, y \rangle - f^*(y)$					
$\alpha_t := 1$ for $t = 1, \dots, T$					
$\text{OAlg}^X := \text{OMD}[\frac{1}{2}\ \cdot\ _2^2, x_0, \frac{(R/G)}{\sqrt{T}}]$					
$\text{OAlg}^Y := \text{FTL}^+$					

Iterative Description

FGNRD Equivalence

Output: $w_T = \tilde{x}_T$

Theorem 21 Denote the constant $G = \max_{w \in \mathcal{K}} \|\partial f(w)\|^2$, and let R be an upper bound on $\|w_0 - w^*\|$. The output $\tilde{w}_T = \tilde{x}_T$ of Algorithm 19 satisfies

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) = O\left(\frac{GR}{\sqrt{T}}\right),$$

where $w_* := \operatorname{argmin}_{w \in \mathcal{K}} f(w)$.

Proof Let $\gamma = \frac{(R/G)}{\sqrt{T}}$ and $\alpha_t = 1$. The update of OMD is $x_t = x_{t-1} - \gamma y_{t-1}$, while the update of FTL^+ is $y_t = \operatorname{argmax}_y \sum_{t=1}^T (f^*(y) - \langle x_t, y \rangle) = \partial f(\tilde{x}_t)$. Recursively

expanding \bar{x}_t leads to

$$\begin{aligned}\bar{x}_t &= \frac{(t-1)\bar{x}_{t-1} + x_t}{t} = \frac{(t-1)\bar{x}_{t-1} + x_{t-1} - \gamma \partial f(\bar{x}_{t-1})}{t} \\ &= \left(1 - \frac{1}{t}\right) \bar{x}_{t-1} - \frac{1}{t} \gamma \sum_{s=0}^{t-1} \partial f(\bar{x}_s).\end{aligned}$$

We now see the equivalence of the two displays in Algorithm 19.

By Lemma 9, the regret of the x-player is, $\alpha\text{-REG}^x \leq \frac{1}{\gamma} D_{x_0}^\phi(x^*) + \frac{\gamma}{2} \sum_{t=1}^T \|\alpha_t y_t\|_2^2$. On the other hand, by Lemma 2, the regret of the y-player is not greater than 0. Adding average regrets of both players, we get

$$\begin{aligned}\overline{\alpha\text{-REG}}^x[\text{OMD}] + \overline{\alpha\text{-REG}}^y[\text{FTL}^+] &\leq \frac{1}{A_t} \left\{ \frac{1}{\gamma} D_{x_0}^\phi(x^*) + \sum_{t=1}^T \frac{\gamma}{2} \|\alpha_t y_t\|_2^2 \right\} \\ &= O\left(\frac{GR}{\sqrt{T}}\right).\end{aligned}$$

□

Let us comment that though the same algorithm can be produced from Algorithm 1 in [81] with the online gradient descent strategy, it is not shown in the paper. Therefore, we consider cumulative gradient descent to be new.

5.4 Optimistic mirror descent, with weighted averaging

We give another accelerated algorithm for smooth convex optimization in the following. Algorithm 20 is obtained when we swap the ordering of the player and let the x-player plays first using OPTIMISTICMD and then the y-player plays FTL⁺. It is noted that the algorithm here only requires a single gradient evaluation in each iteration, while the extra-gradient methods in the literature, e.g. [59, 62, 82] need computing two gradients at different points in every iteration.

Algorithm 20 Optimistic mirror descent, with weighted averaging

Given: L -smooth $f(\cdot)$, a 1-strongly convex $\phi(\cdot)$, iterations T

Init: arbitrary $w_{-\frac{1}{2}} = w_0 = x_0 = x_{-\frac{1}{2}} \in \mathcal{K}$.

$$\begin{aligned}\gamma &\leftarrow \frac{1}{L} \\ w_t &\leftarrow \operatorname{argmin}_{w \in \mathcal{K}} \left(\alpha_t \langle w, \nabla f(\bar{w}_{t-1}) \rangle \right. \\ &\quad \left. + \frac{1}{\gamma} D_{w_{t-\frac{1}{2}}}^\phi(w) \right) \\ w_{t+\frac{1}{2}} &\leftarrow \operatorname{argmin}_{w \in \mathcal{K}} \left(\alpha_t \langle w, \nabla f(\bar{w}_t) \rangle \right. \\ &\quad \left. + \frac{1}{\gamma} D_{w_{t-\frac{1}{2}}}^\phi(w) \right)\end{aligned}$$

Iterative Description

$$\begin{aligned}g(x, y) &:= \langle x, y \rangle - f^*(y) \\ \alpha_t &:= t \text{ for } t = 1, \dots, T \\ \text{OAlg}^X &:= \text{OPTIMISTICMD}[\phi(\cdot), x_0, \frac{1}{L}] \\ \text{OAlg}^Y &:= \text{FTL}^+\end{aligned}$$

FGNRD Equivalence

Output: $\bar{w}_T = \bar{x}_T$

Theorem 22 *The output $\bar{w}_T = \bar{x}_T$ of Algorithm 20 satisfies*

$$f(\bar{w}_T) - \min_{w \in \mathcal{K}} f(w) = \frac{2LD_{w_0}^\phi(w^*) + L\|\bar{w}_0 - \bar{w}_1\|^2}{T^2},$$

where $w_* := \operatorname{argmin}_{w \in \mathcal{K}} f(w)$.

Proof The equivalence of the two displays, namely, for all t ,

$$\begin{aligned} w_t &= x_t \\ w_{t-\frac{1}{2}} &= x_{t-\frac{1}{2}} \\ \nabla f(\bar{w}_t) &= y_t \end{aligned}$$

can be trivially shown by induction. Specifically, the first two relations hold in the beginning by letting the initial point $w_0 = w_{-\frac{1}{2}} = x_0 = x_{-\frac{1}{2}} \in \mathcal{K}$. To show the last one, we apply the definition of FTL⁺,

$$y_t \leftarrow \operatorname{argmax}_y \sum_{s=1}^T \ell_s(y) = \sum_{s=1}^T (f^*(y) - \alpha_s \langle x_s, y \rangle) = \nabla f(\bar{x}_t).$$

By induction, we have $w_t = x_t$. So we have $\bar{w}_t = \bar{x}_t$ and consequently $y_t = \nabla f(\bar{w}_t)$.

By summing the regret bound of each player, i.e. Lemma 12 and Lemma 5, we get

$$\begin{aligned} & \overline{\alpha\text{-REG}}^x[\text{OPTIMISTICMD}] + \overline{\alpha\text{-REG}}^y[\text{FTL}^+] \\ & \stackrel{(a)}{\leq} \frac{1}{A_t} \left\{ \frac{1}{\gamma} D_{x_0}^\phi(x^*) + \frac{\gamma}{2} \sum_{t=1}^T \alpha_t^2 \|\nabla f(\bar{w}_t) - \nabla f(\bar{w}_{t-1})\|_*^2 - \sum_{t=1}^T \frac{A_{t-1}}{L} \|\nabla f(\bar{w}_t) - \nabla f(\bar{w}_{t-1})\|_*^2 \right\} \\ & \stackrel{(b)}{\leq} \frac{LD_{x_0}^\phi(x^*) + L\|\bar{w}_0 - \bar{w}_1\|^2}{A_T} \leq \frac{2LD_{x_0}^\phi(x^*) + L\|\bar{w}_0 - \bar{w}_1\|^2}{T^2}, \end{aligned}$$

where in (a) we used the relation $m_t = y_{t-1} = \nabla f(\bar{w}_{t-1})$ when we applied Lemma 12 and the fact that the y-player's loss function $\ell_t(y) := f^*(y) - \langle x, y_t \rangle$ is $\frac{1}{L}$ -strongly convex w.r.t the dual norm $\|\cdot\|_*$ so that we have $\mu = \frac{1}{L}$ when we invoked Lemma 5, and in (b) we used that $\frac{\gamma\alpha_t^2}{2} \leq \frac{A_{t-1}}{L}$ for $\gamma \leq \frac{1}{L}$ so that the distance terms cancel out except the one at $t = 1$. □

Appendix A Proof of Theorem 2

Proof This is a result of the following lemmas.

Definition: [Definition 12.1 in [83]] *A mapping $T : \mathcal{R}^n \rightarrow \mathcal{R}^n$ is called monotone if it has the property that*

$$\langle v_1 - v_0, x_1 - x_0 \rangle \geq 0 \text{ whenever } v_0 \in T(x_0), v_1 \in T(x_1).$$

Moreover, T is maximal monotone if there is no monotone operator that properly contains it.

Lemma 2: [Theorem 12.17 in [83]] *For a proper, lsc, convex function f , ∂f is a maximum monotone operator.*

Lemma 3: [Theorem 12.41 in [83]] *For any maximal monotone mapping T , the set*

“domain of T “ is nearly convex, in the sense that there is a convex set C such that $C \subset \text{domain of } T \subset \text{cl}(C)$. The same applies to the range of T .

Therefore, the closure of $\{\partial f(x) : x \in \mathcal{X}\}$ is also convex, because we can define another proper, lsc, convex function $\hat{f}(x)$ such that it is $\hat{f}(x) = f(x)$ if $x \in \mathcal{X}$; otherwise, $\hat{f}(x) = \infty$. Then, the sub-differential of $\hat{f}(x)$ is equal to $\{\partial f(x) : x \in \mathcal{X}\}$. So, we can apply the the lemmas to get the result. \square

Appendix B Proof of Theorem 5

Note that in the algorithm that we describe below the weights α_t are not pre-defined but rather depend on the queries of the algorithm. These adaptive weights are explicitly defined in Algorithm 21 which is used by the y -player. Note that Algorithm 21 is equivalent to performing FTL updates over the following loss sequence: $\{\tilde{\ell}_t(y) := \alpha_t \ell_t(y)\}_{t=1}^T$. The x -player plays best response, which only involves the linear optimization oracle.

Algorithm 21 Adaptive Follow-the-Leader (AFTL)

- 1: **for** $t = 1, 2, \dots, T$ **do**
 - 2: Play $y_t \in \mathcal{Y}$
 - 3: Receive a strongly convex loss function $\alpha_t \ell_t(\cdot)$ with $\alpha_t = \frac{1}{\|\nabla \ell_t(y_t)\|^2}$.
 - 4: Update $y_{t+1} = \min_{y \in \mathcal{Y}} \sum_{s=1}^t \alpha_y \ell_s(y)$
 - 5: **end for**
-

Proof We can easily show the equivalence of the two displays shown on Algorithm 7 by using the induction and noting that $\nabla \ell_t(y_t) = x_t - \nabla f^*(y_t) = x_t - \bar{x}_{t-1}$, where in the last equality we used that $y_t = \nabla f(\bar{x}_{t-1})$ if and only if $\bar{x}_{t-1} = \nabla f^*(y_t)$.

Now we switch to show the convergence rate. Since the x -player plays best response, $\overline{\alpha\text{-REG}}^x[\text{BESTRESP}^+] = 0$, we only need to show that $\overline{\alpha\text{-REG}}^y[\text{SC-AFTL}] = O(\exp(-\frac{\lambda B}{L}T))$, which we do next.

We start by defining a function $s(y) := \max_{x \in \mathcal{X}} -x^\top y + f^*(y)$, which is a strongly convex function. We are going to show that $s(\cdot)$ is also smooth. We have

$$\begin{aligned} \|\nabla_w s(\cdot) - \nabla_z s(\cdot)\| &= \left\| \arg \max_{x \in \mathcal{X}} (-w^\top x + f^*(x)) - \arg \max_{x \in \mathcal{X}} (-z^\top x + f^*(x)) \right\| \\ &= \left\| \arg \max_{x \in \mathcal{X}} (-w^\top x) - \left(\arg \max_{x' \in \mathcal{X}} -z^\top x' \right) \right\| \leq \frac{2\|w - z\|}{\lambda(\|w\| + \|z\|)} \leq \frac{\|w - z\|}{\lambda B}, \end{aligned}$$

where the second to last inequality uses Lemma 18 regarding λ -strongly convex sets, and the last inequality is by assuming the gradient of $\|\nabla f(\cdot)\| \geq B$ and the fact that $w, z \in \mathcal{Y}$ are gradients of f . This shows that $s(\cdot)$ is a smooth function with smoothness constant $L' := \frac{1}{\lambda B}$.

$$\begin{aligned} T &= \sum_{t=1}^T \frac{\|\nabla \ell_t(y_t)\|^2}{\|\nabla \ell_t(y_t)\|^2} \stackrel{\text{Proposition 23}}{=} \sum_{t=1}^T \frac{\|\nabla s(y_t)\|^2}{\|\nabla \ell_t(y_t)\|^2} \stackrel{\text{Lemma 17}}{\leq} \sum_{t=1}^T \frac{L'}{\|\nabla \ell_t(y_t)\|^2} (s(y_t) - s(y^*)) \\ &\leq \sum_{t=1}^T \frac{L'}{\|\nabla \ell_t(x_t)\|^2} (\ell_t(y_t) - \ell_t(y^*)), \end{aligned}$$

where we denote $y^* := \operatorname{argmin}_y s(y)$ and the last inequality follows from the fact that $s(y_t) := \ell_t(y_t)$ and $\ell_t(y) = -g(x_t, y) \leq -g(x_y, y) = s(y)$ for any y .

In the following, we will denote c a constant such that $\|\nabla \ell_t(y_t)\| = \|x_t - \nabla f^*(y_t)\| = \|x_t - \bar{x}_{t-1}\| \leq c$. We have

$$\begin{aligned} T &\leq \sum_{t=1}^T \frac{L'}{\|\ell_t(y_t)\|^2} (\ell_t(y_t) - \ell_t(y^*)) \\ &\stackrel{(a)}{=} \sum_{t=1}^T L' (\tilde{\ell}_t(y_t) - \tilde{\ell}_t(y^*)) \\ &\stackrel{(b)}{\leq} \frac{L \cdot L'}{2} \sum_{t=1}^T \frac{\|\nabla \ell_t(y_t)\|^{-2}}{\sum_{s=1}^t \|\nabla \ell_s(y_t)\|^{-2}} \\ &\stackrel{(c)}{\leq} \frac{L \cdot L'}{2} \left(1 + \log \left(c^2 \sum_{t=1}^T \|\nabla \ell_t(y_t)\|^{-2} \right) \right), \end{aligned}$$

where (a) is by the definition of $\tilde{\ell}_t(\cdot)$, and (b) is shown using Lemma 3 with strong convexity parameter of $\ell_t(\cdot)$ being $\frac{1}{L}$, and (c) is by Lemma 19 so that

$$\sum_{t=1}^T \frac{\|\ell_t(y_t)\|^{-2}}{\sum_{s=1}^t \|\ell_s(y_s)\|^{-2}} = \sum_{t=1}^T \frac{c^2 \|\ell_t(y_t)\|^{-2}}{\sum_{s=1}^t c^2 \|\ell_s(y_s)\|^{-2}} \leq 1 + \log \left(c^2 \sum_{t=1}^T \|\ell_t(y_t)\|^{-2} \right).$$

Thus, we get

$$c^2 \sum_{t=1}^T \|\nabla \ell_t(y_t)\|^{-2} = O(e^{\frac{1}{L \cdot L'} T}) = O(e^{\frac{\lambda B}{L} T}). \quad (\text{B1})$$

So

$$\begin{aligned} \frac{\alpha\text{-REG}^y}{A_T} &:= \frac{\sum_{t=1}^T \alpha_t (\ell_t(y_t) - \ell_t(y^*))}{A_T} \leq \frac{L}{2A_T} \sum_{t=1}^T \frac{\|\nabla \ell_t(y_t)\|^{-2}}{\sum_{\tau=1}^t \|\nabla \ell_t(y_\tau)\|^{-2}} \\ &\stackrel{(a)}{\leq} \frac{Lc^2 \left(1 + \log \left(c^2 \sum_{t=1}^T \|\nabla \ell_t(y_t)\|^{-2} \right) \right)}{2c^2 \sum_{t=1}^T \|\nabla \ell_t(y_t)\|^{-2}} \stackrel{(b)}{\leq} O \left(\frac{Lc^2 \left(1 + \left(\frac{\lambda B T}{L} \right) \right)}{e^{\frac{\lambda B T}{L}}} \right) = O \left(Lc^2 e^{-\frac{\lambda B}{L} T} \right), \end{aligned}$$

where (a) is by Lemma 19, (b) is by (B1) and the fact that $\frac{1+\log z}{z}$ is monotonically decreasing for $z \geq 1$. Thus, we have

$$f(w_T) - \min_{w \in \mathcal{K}} f(w) \leq \overline{\alpha\text{-REG}}^x[\text{BESTRESP}^+] + \overline{\alpha\text{-REG}}^y[\text{AFTL}] = O \left(Lc^2 e^{-\frac{\lambda B}{L} T} \right).$$

This completes the proof. \square

Proposition 23 *For arbitrary y , let $\ell(\cdot) := -g(x_y, \cdot)$. Then $-\nabla_y \ell(\cdot) \in \partial_y s(\cdot)$, where x_y means that the x -player plays x by BESTRESP^+ after observing the y -player plays y .*

Proof Consider any point $w \in \mathcal{Y}$,

$$\begin{aligned} s(w) - s(y) &= g(x_y, y) - g(x_w, w) \\ &= g(x_y, y) - g(x_y, w) + g(x_y, w) - g(x_w, w) \geq g(x_y, y) - g(x_y, w) + 0 \\ &\geq \langle \partial_y g(x_y, y), w - y \rangle = \langle -\nabla_y \ell(y), w - y \rangle, \end{aligned}$$

where the first inequality is because that x_w is the best response to w , the second inequality is due to the concavity of $g(x_y, \cdot)$. The overall statement implies that $-\nabla_y \ell(y)$ is a subgradient of s at y . \square

Lemma 17 For any L -smooth convex function $\ell(\cdot) : \mathbb{R}^d \mapsto \mathbb{R}$, if $x^* = \operatorname{argmin}_{x \in \mathbb{R}^d} \ell(x)$, then

$$\|\nabla \ell(x)\|^2 \leq 2L (\ell(x) - \ell(x^*)), \quad \forall x \in \mathbb{R}^d.$$

Lemma 18⁴ Denote $x_p = \operatorname{argmax}_{x \in \mathcal{K}} \langle p, x \rangle$ and $x_q = \operatorname{argmax}_{x \in \mathcal{K}} \langle q, x \rangle$, where $p, q \in \mathbb{R}^d$ are any nonzero vectors. If a compact set \mathcal{K} is a λ -strongly convex set, then

$$\|x_p - x_q\| \leq \frac{2\|p - q\|}{\lambda(\|p\| + \|q\|)}.$$

Proof [76] show that a strongly convex set \mathcal{K} can be written as intersection of some Euclidean balls. Namely,

$$\mathcal{K} = \bigcap_{u: \|u\|_2=1} B_{\frac{1}{\lambda}} \left(x_u - \frac{u}{\lambda} \right),$$

where x_u is defined as $x_u = \operatorname{argmax}_{x \in \mathcal{K}} \langle \frac{u}{\|u\|}, x \rangle$.

Let $x_p = \operatorname{argmax}_{x \in \mathcal{K}} \langle \frac{p}{\|p\|}, x \rangle$ and $x_q = \operatorname{argmax}_{x \in \mathcal{K}} \langle \frac{q}{\|q\|}, x \rangle$. Based on the definition of strongly convex sets, we can see that $x_q \in B_{\frac{1}{\lambda}}(x_p - \frac{p}{\lambda\|p\|})$ and $x_p \in B_{\frac{1}{\lambda}}(x_q - \frac{q}{\lambda\|q\|})$. Therefore,

$$\|x_q - x_p - \frac{p}{\lambda\|p\|}\|^2 \leq \frac{1}{\lambda^2},$$

which leads to

$$\|p\| \cdot \|x_p - x_q\|^2 \leq \frac{2}{\lambda} \langle x_p - x_q, p \rangle. \quad (\text{B2})$$

Similarly,

$$\|x_p - x_q - \frac{q}{\lambda\|q\|}\|^2 \leq \frac{1}{\lambda^2},$$

which results in

$$\|q\| \cdot \|x_p - x_q\|^2 \leq \frac{2}{\lambda} \langle x_q - x_p, q \rangle. \quad (\text{B3})$$

Summing (B2) and (B3), one gets $(\|p\| + \|q\|)\|x_p - x_q\|^2 \leq \frac{2}{\lambda} \langle x_p - x_q, p - q \rangle$. Applying the Cauchy-Schwarz inequality completes the proof. \square

Lemma 19 ([84]) For any non-negative real numbers $a_1, \dots, a_n \geq 1$,

$$\sum_{i=1}^n \frac{a_i}{\sum_{j=1}^i a_j} \leq 1 + \log \left(\sum_{i=1}^n a_i \right).$$

⁴[76] discuss the smoothness of the support function on strongly convex sets. Here, we state a more general result.

Appendix C Proof of Theorem 6

Proof The equivalence of the updates follows the proof of Theorem 3. Specifically, we have that the objects on the left in the following equalities correspond to Alg. 2 and those on the right to Alg. 8.

$$\begin{aligned} x_t &= v_t \\ \bar{x}_t &= w_t. \end{aligned}$$

To analyze the regret of the y -player, we define $\{\hat{y}_t\}$ as the points if the y -player would have played FTL, which is

$$\begin{aligned} \hat{y}_t &:= \operatorname{argmin}_y \frac{1}{t-1} \sum_{s=1}^{t-1} \ell_t(y) = \operatorname{argmax}_y \frac{1}{t-1} \sum_{s=1}^{t-1} \langle x_s, y \rangle - f^*(y) \\ &= \nabla f(\bar{x}_t) = \frac{1}{n} \sum_{i=1}^n \nabla f_i(\bar{x}_t). \end{aligned}$$

On the other hand, we call the actual strategy used by the y -player LAZYFTL as it only compute the gradient of a single component. We can show the average regret of LAZYFTL as follows.

$$\begin{aligned} \overline{\alpha\text{-REG}}^y &= \frac{1}{T} \sum_{t=1}^T (\ell_t(\hat{y}_t) - \ell_t(y_*)) + \frac{1}{T} \sum_{t=1}^T (\ell_t(y_t) - \ell_t(\hat{y}_t)) \\ &\stackrel{(a)}{\leq} \frac{4LR \log T}{T} + \frac{1}{T} \sum_{t=1}^T (\ell_t(y_t) - \ell_t(\hat{y}_t)) \\ &= \frac{4LR \log T}{T} + \frac{1}{T} \sum_{t=1}^T (f^*(y_t) - f^*(\hat{y}_t) + \langle x_t, \hat{y}_t - y_t \rangle) \\ &\stackrel{(b)}{\leq} \frac{4LR \log T}{T} + \sum_{t=1}^T \frac{1}{T} (L_0 + r) \|y_t - \hat{y}_t\| \\ &= \frac{4LR \log T}{T} + \frac{1}{T} (L_0 + r) \sum_{t=1}^T \left\| \frac{1}{n} \sum_{i=1}^n g_{i,t} - \frac{1}{n} \sum_{i=1}^n \nabla f_i(\bar{x}_t) \right\| \\ &= \frac{4LR \log T}{T} + \frac{1}{T} (L_0 + r) \sum_{t=1}^T \left\| \frac{1}{n} \sum_{i \neq i_t}^n (g_{i,t} - \nabla f_i(\bar{x}_t)) \right\| \\ &\stackrel{(c)}{\leq} \frac{4LR \log T}{T} + \frac{L(L_0 + r)}{Tn} \sum_{t=1}^T \sum_{i \neq i_t}^n \|\bar{x}_{\tau_t(i)} - \bar{x}_t\| \\ &\stackrel{(d)}{\leq} \frac{4LR \log T}{T} + \frac{L(L_0 + r)}{Tn} \sum_{t=1}^T \sum_{i \neq i_t}^n \frac{2nr}{t} \\ &= O\left(\frac{\max\{LR, L(L_0 + r)nr\} \log T}{T}\right), \end{aligned}$$

where (a) is due to the regret of FTL (Lemma 3),

$$\frac{1}{T} \left(\sum_{t=1}^T \ell_t(\hat{y}_t) - \ell_t(y_*) \right) \leq \frac{1}{T} \sum_{t=1}^T \frac{2\|\nabla \ell_t(\hat{y}_t)\|^2}{\sum_{s=1}^t (1/L)} = \frac{4LR \log T}{T},$$

and that $\|\nabla\ell_t(\hat{y}_t)\|^2 = \|x_t - \nabla f^*(\hat{y}_t)\|^2 = \|x_t - \bar{x}_{t-1}\|^2 \leq R$, (b) we assume that the conjugate is L_0 -Lipschitz and that $\max_{x \in \mathcal{K}} \|x\| \leq r$, (c) we denote $\tau_t(i) \in [T]$ as the last iteration that i_{th} sample's gradient is computed at t , and (d) is because that

$$\begin{aligned} \|\bar{x}_{\tau_t(i)} - \bar{x}_t\| &= \left\| \frac{1}{\tau_t(i)} \sum_{s=1}^{\tau_t(i)} x_s - \frac{1}{t} \sum_{s=1}^t x_s \right\| \leq \left\| \sum_{s=1}^{\tau_t(i)} x_s \left(\frac{1}{\tau_t(i)} - \frac{1}{t} \right) \right\| + \left\| \frac{1}{t} \sum_{s=\tau_t(i)+1}^t x_s \right\| \\ &= \frac{t - \tau_t(i)}{t} \|\bar{x}_{\tau_t(i)}\| + \left\| \frac{1}{t} \sum_{s=\tau_t(i)+1}^t x_s \right\| \\ &\leq \frac{nr}{t} + \left\| \frac{1}{t} \sum_{s=\tau_t(i)+1}^t x_s \right\| = \frac{nr}{t} + \frac{t - \tau_t(i)}{t} \left\| \frac{1}{t - \tau_t(i)} \sum_{s=\tau_t(i)+1}^t x_s \right\| \\ &\leq \frac{2nr}{t}. \end{aligned}$$

For the x -player, since it plays BESTRESP^+ , the regret is non-positive, i.e. $\overline{\alpha\text{-REG}}^x[\text{BESTRESP}^+] = 0$. By combining the regrets of both players, we have

$$\begin{aligned} f(w_T) - \min_{w \in \mathcal{K}} f(w) &\leq \overline{\alpha\text{-REG}}^x[\text{BESTRESP}^+] + \overline{\alpha\text{-REG}}^y[\text{LAZYFTL}] \\ &= O\left(\frac{\max\{LR, L(L_0 + r)nr\} \log T}{T}\right). \end{aligned}$$

□

Appendix D Proof of Theorem 9

Proof First, we can bound the norm of the gradient as

$$\|\nabla\ell_t(y_t)\|^2 = \|x_t - \nabla f^*(y_t)\|^2 = \|x_t - \bar{x}_{t-1}\|^2.$$

Combining this with Lemma 3 we see that

$$\overline{\alpha\text{-REG}}^y[\text{FTL}] \leq \frac{1}{A_T} \sum_{t=1}^T \frac{2\alpha_t^2 \|\nabla\ell_t(y_t)\|^2}{\sum_{s=1}^t \alpha_s(1/L)} = \frac{1}{A_T} \sum_{t=1}^T \frac{2\alpha_t^2 \|x_t - \bar{x}_{t-1}\|^2}{\sum_{s=1}^t \alpha_s(1/L)}.$$

On the other hand, the x -player plays OMD^+ , according to Lemma 10, its regret satisfies

$$\overline{\alpha\text{-REG}}^x \leq \frac{\frac{1}{\gamma} D_{x_0}^\phi(x^*) - \sum_{t=1}^T 4L \|x_{t-1} - x_t\|^2}{A_T}.$$

By adding the average regret of both players, we get

$$\overline{\alpha\text{-REG}}^y[\text{FTL}] + \overline{\alpha\text{-REG}}^x[\text{OMD}^+] \leq \frac{4L \|x_0 - x^*\|^2 + \sum_{t=1}^T 4L \left(\|\bar{x}_{t-1} - x_t\|^2 - \|x_{t-1} - x_t\|^2 \right)}{A_T},$$

where we used that $D_{x_0}^\phi(x^*) = \frac{1}{2} \|x_0 - x^*\|^2$. Since the distance terms may not cancel out, one can only bound the differences of the distance terms by a constant, which leads to the non-accelerated $O(1/T)$ rate. □

Appendix E Proof of Theorem 12

Proof The proof is similar to that of Theorem 11. The only difference is that the x-player is FTRL⁺ instead of OMD⁺. By using a bound on α -REG^y in Lemma 15 and the bound in Lemma 7 of FTRL⁺ with $\mu = 0$ (as the x-player sees linear loss functions), we have

$$\begin{aligned} f(w_T) - \min_{w \in \mathcal{K}} f(w) &\leq \frac{1}{A_T} (\alpha\text{-REG}^x[\text{OPTIMISTICFTL}] + \alpha\text{-REG}^y[\text{FTRL}^+]) \\ &\leq \frac{1}{A_T} \left(\frac{R(x^*) - \min_{x \in \mathcal{K}} R(x)}{\eta} + \sum_{t=1}^T \left(\frac{\alpha_t^2}{A_t} L - \frac{\beta}{2\eta} \right) \|x_{t-1} - x_t\|^2 \right). \end{aligned} \quad (\text{E4})$$

Choosing $\eta = \frac{1}{4L}$, $\beta = 1$, and the weight $\alpha_t = t$, we have $A_t := \frac{t(t+1)}{2}$ and therefore $\frac{\alpha_t^2}{A_t} = \frac{2t^2}{t(t+1)} \leq 2$. With this in mind, the sum on the right hand side of (E4) is non-positive. Noticing that $\frac{1}{A_T} \leq \frac{2}{T^2}$ completes the proof. \square

Appendix F Examples of strongly convex sets

[21] lists three known classes of λ -strongly convex sets as follows, which are all gauge sets.

1. ℓ_p balls: $\|x\|_p \leq r, \forall p \in (1, 2]$. The strong convexity of the set is $\lambda = \frac{p-1}{r}$ and its square of gauge function is $\frac{1}{r^2} \|x\|_p^2$, which is a $\frac{2(p-1)}{r^2}$ -strongly convex function with respect to norm $\|\cdot\|_p$ by Lemma 4 in [21].
2. Schatten p balls: $\|\sigma(X)\|_p \leq r$ for $p \in (1, 2]$, where $\sigma(X)$ is the vector consisting of singular values of the matrix X . The strong convexity of the set is $\lambda = \frac{p-1}{r}$ and the square gauge function is $\frac{1}{r^2} \|\sigma(X)\|_p^2$, which is a $\frac{2(p-1)}{r^2}$ -strongly convex function with respect to norm $\|\sigma(\cdot)\|_p$ by Lemma 6 in [21].
3. Group (s,p) balls: $\|X\|_{s,p} = \|(\|X_1\|_s, \|X_2\|_s, \dots, \|X_m\|_s)\|_p \leq r$ where $X \in \mathbb{R}^{m \times n}$, X_j represents the j -th row of X , and $s, p \in (1, 2]$. The strong convexity of the set is $\lambda = \frac{2(s-1)(p-1)}{r((s-1)+(p-1))}$ and its square gauge function is $\frac{1}{r^2} \|X\|_{s,p}^2$, which is a $\frac{2(s-1)(p-1)}{r^2((s-1)+(p-1))}$ -strongly convex function with respect to norm $\|\cdot\|_{s,p}$ by Lemma 8 in [21].

References

- [1] Nesterov, Y. A method of solving a convex programming problem with convergence rate $o(1/k^2)$. *Soviet Mathematics Doklady* **27**, 372–376 (1983).
- [2] Nesterov, Y. A method for unconstrained convex minimization problem with the rate of convergence $o(1/k^2)$. *Doklady AN USSR* (1983).

- [3] Nesterov, Y. Smooth minimization of nonsmooth functions. *Mathematical programming* (2005) .
- [4] Nesterov, Y. On an approach to the construction of optimal methods of minimization of smooth convex functions. *Ekonom. i. Mat. Metody* **24**, 509–517 (1988) .
- [5] Nesterov, Y. Introductory lectures on convex optimization: A basic course. *Springer* (2004) .
- [6] Frank, M. & Wolfe, P. An algorithm for quadratic programming. *Naval research logistics quarterly* **3** (1-2), 95–110 (1956) .
- [7] Polyak, B. Some methods of speeding up the convergence of iteration methods. *USSR Computational Mathematics and Mathematical Physics* (1964) .
- [8] Beck, A. & Teboulle, M. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J. on Imaging Sciences* (2009) .
- [9] Abernethy, J. & Wang, J.-K. Frank-wolfe and equilibrium computation. *NeurIPS* (2017) .
- [10] Abernethy, J., Levy, K., Lai, K. & Wang, J.-K. Faster rates for convex-concave games. *COLT* (2018) .
- [11] Wang, J.-K. & Abernethy, J. Acceleration through optimistic no-regret dynamics. *NeurIPS* (2018) .
- [12] Levitin, E. S. & Polyak, B. T. Constrained minimization methods. *USSR Computational mathematics and mathematical physics* (1966) .
- [13] Négiar, G. *et al.* Stochastic frank-wolfe for constrained finite-sum minimization. *ICML* (2020) .
- [14] Bertsekas, D. P., Nedic, A. & Ozdaglar, A. E. Convex analysis and optimization. *Athena Scientific* (2003) .
- [15] Hiriart-Urruty, J.-B. & Lemarechal, C. Convex analysis and minimization algorithms. *Springer* (1993) .
- [16] Rockafellar, T. Convex analysis. *Princeton University Press* (1996) .
- [17] Boyd, S. & Vandenberghe, L. Convex optimization. *Cambridge University Press* (2004) .
- [18] Borwein, J. & Lewis, A. S. Convex analysis and nonlinear optimization theory and examples. *Springer* (2006) .

- [19] Bauschke, H. H. & Lucet, Y. What is a fenchel conjugate? *Notices of the AMS* (2012) .
- [20] Kakade, S. M., Shalev-shwartz, S. & Tewari, A. On the duality of strong convexity and strong smoothness: Learning applications and matrix regularization. *SIAM Journal on Control and Optimization* (2009) .
- [21] Garber, D. & Hazan, E. Faster rates for the frank-wolfe method over strongly-convex sets. *ICML* (2015) .
- [22] Orabona, F. A modern introduction to online learning. *arXiv:1912.13213* (2019) .
- [23] Rakhlin, A. & Sridharan, K. Statistical learning theory and sequential prediction (2016) .
- [24] Hazan, E. *et al.* Introduction to online convex optimization. *Foundations and Trends® in Optimization* **2** (3-4), 157–325 (2016) .
- [25] Shalev-Shwartz, S. *et al.* Online learning and online convex optimization. *Foundations and Trends® in Machine Learning* **4** (2), 107–194 (2012) .
- [26] Freund, Y. & Schapire, R. E. Game theory, on-line prediction and boosting. *COLT* 325–332 (1996) .
- [27] Blackwell, D. An analog of the minimax theorem for vector payoffs. *Pacific Journal of Mathematics* **6** (1), 1–8 (1956) .
- [28] Hannan, J. Approximation to bayes risk in repeated play. *Contributions to the Theory of Games* **39**, 97 (1957) .
- [29] Rakhlin, A. & Sridharan, K. Optimization, learning, and games with predictable sequences. *NeurIPS* (2013) .
- [30] Chiang, C.-K. *et al.* Online optimization with gradual variations. (2012) .
- [31] Kalai, A. & Vempala, S. Efficient algorithms for online decision problems. *Journal of Computer and System Sciences* **71** (3), 291–307 (2005) .
- [32] Huang, R., Lattimore, T., György, A. & Szepesvari, C. Following the leader and fast rates in linear prediction: Curved constraint sets and other regularities. *NeurIPS* (2016) .
- [33] Xiao, L. Dual averaging methods for regularized stochastic learning and online optimization. *JMLR* (2010) .
- [34] Tseng, P. On accelerated proximal gradient methods for convex-concave optimization (2008) .

- [35] Koren, T. & Levy, K. Fast rates for exp-concave empirical risk minimization. *NeurIPS* (2015) .
- [36] Garber, D. & Meshi, O. Linear-memory and decomposition-invariant linearly convergent conditional gradient algorithm for structured polytopes. *NeurIPS* (2016) .
- [37] Combettes, C. W. & Pokutta, S. Complexity of linear minimization and projection on some sets. *arXiv:2101.10040* (2021) .
- [38] Demyanov, V. F. & Rubinov, A. M. Approximate methods in optimization problems. *Elsevier Publishing Company*, (1970) .
- [39] Dunn, J. C. Rates of convergence for conditional gradient algorithms near singular and nonsingular extremals. *SIAM Journal on Control and Optimization* (1979) .
- [40] Lu, H. & Freund, R. M. Generalized stochastic frankwolfe algorithm with stochastic “substitute” gradient for structured convex optimization. *Mathematical Programming* (2020) .
- [41] Jaggi, M. Revisiting frank-wolfe: Projection-free sparse convex optimization. *ICML* (2013) .
- [42] Clarkson, K. Coresets, sparse greedy approximation, and the frank-wolfe algorithm. *SODA* (2008) .
- [43] Garber, D. & Hazan, E. A linearly convergent conditional gradient algorithm with applications to online and stochastic optimization. *SIAM Journal on Optimization* (2016) .
- [44] Wolf., P. Convergence theory in nonlinear programming. *Integer and Nonlinear Programming* (1970) .
- [45] Lacoste-Julien, S. & Jaggi, M. On the global linear convergence of frank-wolfe optimization variants. *NeurIPS* (2015) .
- [46] Freund, R. M. & Grigas, P. New analysis and results for the frank-wolfe method. *Mathematical Programming* (2016) .
- [47] Braun, G., Pokutta, S., Tu, D. & Wright, S. Blended conditional gradients. *ICML* (2019) .
- [48] Harchaoui, Z., Juditsky, A. & Nemirovski, A. Conditional gradient algorithms for norm-regularized smooth convex optimization. *Math. Prog., Series A* (2013) .

- [49] Lan, G. & Zhou, Y. Conditional gradient sliding for convex optimization. *SIAM Journal on Optimization* (2016) .
- [50] Bubeck, S. Convex optimization: Algorithms and complexity. *Foundations and Trends in Machine Learning* (2015) .
- [51] Su, W., Boyd, S. & Candes, E. A differential equation for modeling nesterov’s accelerated gradient method: Theory and insights. *NeurIPS* (2014) .
- [52] Parikh, N. & Boyd, S. Proximal algorithms. *Foundations and Trends in Optimization* (2014) .
- [53] Lessard, L., Recht, B. & Packard, A. Analysis and design of optimization algorithms via integral quadratic constraints. *SIAM Journal on Optimization* (2016) .
- [54] Hu, B. & Lessard, L. Dissipativity theory for nesterov’s accelerated method. *ICML* (2017) .
- [55] Bubeck, S., Lee, Y. T. & Singh, M. A geometric alternative to nesterov’s accelerated gradient descent. *arXiv:1506.08187* (2015) .
- [56] Drusvyatskiy, D., Fazel, M. & Roy, S. An optimal first order method based on optimal quadratic averaging. *SIAM Journal on Optimization* (2018) .
- [57] Flammarion, N. & Bach, F. From averaging to acceleration, there is only a step-size. *COLT* (2015) .
- [58] Allen-Zhu, Z. & Orecchia, L. Linear coupling: An ultimate unification of gradient and mirror descent. *ITCS* (2017) .
- [59] Diakonikolas¹, J. & Orecchia, L. Accelerated extra-gradient descent: A novel accelerated first-order method. *ITCS* (2018) .
- [60] Diakonikolas, J. & Orecchia, L. The approximate duality gap technique: A unified theory of first-order methods. *SIAM Journal on Optimization* (2019) .
- [61] Cohen, M. B., Sidford, A. & Tian, K. Relative lipschitzness in extra-gradient methods and a direct recipe for acceleration. *ITCS* (2021) .
- [62] Nemirovski, A. Prox-method with rate of convergence $o(1/t)$ for variational inequalities with lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization* (2004) .

- [63] Nesterov, Y. Dual extrapolation and its applications to solving variational inequalities and related problems. *mathematical programming* (2007) .
- [64] Wibisono, A., Wilson, A. C. & Jordan, M. I. A variational perspective on accelerated methods in optimization. *Proceedings of the National Academy of Sciences* **113** (47), E7351–E7358 (2016) .
- [65] Shi, B., Du, S. S., Jordan, M. I. & Su, W. J. Understanding the acceleration phenomenon via high-resolution differential equations. *Mathematical Programming* (2021) .
- [66] Krichene, W., Bayen, A. & Bartlett, P. L. Accelerated mirror descent in continuous and discrete time. *NeurIPS* (2015) .
- [67] Scieur, D., Roulet, V., Bach, F. & d’Aspremont, A. Integration methods and optimization algorithms. *NeurIPS* (2017) .
- [68] Wilson, A. C., Recht, B. & Jordan, M. I. A lyapunov analysis of accelerated methods in optimization. *JMLR* (2021) .
- [69] Attouch, H., Chbani, Z., Peypouquet, J. & Redont, P. Fast convergence of inertial dynamics and algorithms with asymptotic vanishing viscosity. *Mathematical Programming* (2018) .
- [70] d’Aspremont, A., Scieur, D. & Taylor, A. Acceleration methods. *arXiv:2101.09545* (2021) .
- [71] Lan, G. & Zhou, Y. An optimal randomized incremental gradient method. *Mathematical Programming* (2017) .
- [72] Lan, G. First-order and stochastic optimization methods for machine learning. *Springer* (2020) .
- [73] Lu, H., Freund, R. M. & Nesterov, Y. Relatively-smooth convex optimization by first-order methods and applications. *SIAM Journal on Optimization* (2018) .
- [74] Freund, R. M. Dual gauge programs, with applications to quadratic programming and the minimum-norm problem. *Mathematical Programming* (1987) .
- [75] Friedlander, M. P., Macedo, I. & Pong, T. K. Gauge optimization and duality. *SIAM Journal on Optimization* (2014) .
- [76] Polovinkin, E. S. Strongly convex analysis. *Sbornik: Mathematics* *187(2):259* (1996) .

- [77] Molinaro, M. Curvature of feasible sets in offline and online optimization. *arXiv:2002.03213* (2020) .
- [78] Kerdreux, T., Roux, C., d'Aspremont, A. & Pokutta, S. Linear bandits on uniformly convex sets. *arXiv:2103.05907* (2021) .
- [79] Shor, N. Z. Minimization methods for non-differentiable functions. *Springer* (1985) .
- [80] Boyd, S. Subgradient methods. *Notes for EE364b* (2014) .
- [81] Cutkosky, A. Anytime online-to-batch, optimism and acceleration. *ICML* (2019) .
- [82] Kavis, A., Levy, K. Y., Bach, F. & Cevher, V. Unixgrad: A universal, adaptive algorithm with optimal guarantees for constrained optimization. *arXiv:1910.13857* (2019) .
- [83] Rockafellar, R. T. & Wets, R. J.-B. Variational analysis. *Springer* (1998) .
- [84] Levy, K. Online to offline conversions, universality and adaptive minibatch sizes. *NeurIPS* (2017) .