

Private and polynomial time algorithms for learning Gaussians and beyond

Hassan Ashtiani* Christopher Liaw†

November 23, 2021

Abstract

We present a fairly general framework for reducing (ϵ, δ) differentially private (DP) statistical estimation to its non-private counterpart. As the main application of this framework, we give a polynomial time and (ϵ, δ) -DP algorithm for learning (unrestricted) Gaussian distributions in \mathbb{R}^d . The sample complexity of our approach for learning the Gaussian up to total variation distance α is $\tilde{O}\left(\frac{d^2}{\alpha^2} + \frac{d^2\sqrt{\ln 1/\delta}}{\alpha\epsilon}\right)$, matching (up to logarithmic factors) the best known information-theoretic (non-efficient) sample complexity upper bound (Aden-Ali, Ashtiani, Kamath [1]). In an independent work, Kamath, Mouzakis, Singhal, Steinke, and Ullman [22] proved a similar result using a different approach and with $O(d^{5/2})$ sample complexity dependence on d .

As another application of our framework, we provide the first polynomial time (ϵ, δ) -DP algorithm for robust learning of (unrestricted) Gaussians.

1 Introduction

Learning a multivariate Gaussian distribution with respect to the total variation (TV) distance is one of the most basic tasks in statistical estimation. It is folklore that simply taking the empirical mean and covariance matrix achieves the optimal sample complexity of $\Theta(d^2/\alpha^2)$ for learning *unbounded* Gaussians in \mathbb{R}^d up to TV distance α (see, for example, [3, Theorem C.1]). However, this simple approach can compromise the privacy of the individuals/entities whose data has been used in the estimation procedure. Differential privacy [16] offers a formal framework to study and guarantee privacy in such estimation tasks. The central question that we aim to address is whether unbounded Gaussians can be learned *privately* and *efficiently*.

For the strong notion of *pure* $(\epsilon, 0)$ -differential privacy, learning unbounded Gaussians is impossible. In fact, even learning univariate Gaussians requires some boundedness assumptions on the parameters of the Gaussian [26, Theorem 1.4]. This motivates the study of learning Gaussians under the *approximate* (ϵ, δ) -differential privacy model [14] (or the approximate-DP for short).

*McMaster University, zokaaiam@mcmaster.ca. Hassan Ashtiani was supported by an NSERC Discovery Grant

†Google, cvliaw@google.com

Previous work has established the polynomial-time learnability of univariate Gaussians (with unbounded mean and variance) in the approximate-DP model. The seminal work of Karwa and Vadhan [26] uses stability-based histograms [7] to find a bound on the parameters; once the problem is reduced to the bounded case, Laplace mechanism [16] offers a private way to release a noisy—but sufficiently accurate—estimate for the parameters of the Gaussian¹.

The problem is more difficult in the multivariate setting. For example, simply estimating the mean and variance along each axis does not necessarily yield an accurate overall solution. While this approach does provide an acceptable solution for learning the special case of axis-aligned Gaussians² [26], in general it requires the number of samples to depend on the condition number of the covariance matrix. This dependence is also present in the other private Gaussian estimation approaches such as the CoinPress method [4] and its predecessor [21]³.

More recently, Aden-Ali, Ashtiani, and Kamath [1] established a sharp upper bound on the sample complexity of learning Gaussians in the approximate-DP model, demonstrating that the dependence on the condition number (or the size of the parameters) can be totally avoided. Unfortunately, the approach of [1] is non-constructive and does not yield a computationally efficient method. In fact, it only shows the *existence* of a “locally small” cover for Gaussians in TV distance, which yields an upper bound on the sample complexity through the use of the private hypothesis selection method [6]. This raises the following concrete question.

Is there an (ε, δ) -DP and polynomial time⁴ algorithm for learning unbounded Gaussian distributions over \mathbb{R}^d ?

A different but conceptually related question is whether there is an efficient algorithms for learning high-dimensional Gaussians *robustly*, where a small fraction of samples are arbitrarily corrupted by an adversary. The breakthrough of [11, 28] answered this question positively. In particular, the work of [11] offers a polynomial time algorithm for robustly learning Gaussians with respect to the TV distance. This raises the following natural question.

Is there an (ε, δ) -DP, *robust*, and polynomial time algorithm for learning unbounded Gaussian distributions over \mathbb{R}^d ?

2 Main Results

Theorem 2.1 (Informal). *There is an (ε, δ) -DP algorithm that takes $m = \tilde{O}\left(\frac{d^2}{\alpha^2} + \frac{d^2\sqrt{\ln 1/\delta}}{\alpha\varepsilon}\right)$ i.i.d. samples from an arbitrary Gaussian distributions over \mathbb{R}^d , runs in polynomial time in m , and with high probability outputs a Gaussian whose TV distance to the true distribution is smaller than α .*

¹Although the method of [26] works with a histogram with infinite number of bins, it can be easily implemented efficiently by only constructing the bins that are non-empty (i.e., lazy initialization).

²An axis-aligned Gaussian distribution is a Gaussian whose covariance matrix is diagonal.

³These two papers study a notion of differential privacy known as concentrated-DP [8, 18], which is stronger than pure DP but weaker than approximate DP.

⁴We are looking for an algorithm whose time complexity is polynomial in terms of the optimal sample complexity of the problem.

See [Theorem 9.7](#) for the formal version. The sample complexity of this approach matches the best (non-efficient) learning method [\[1\]](#) up to logarithmic factors. Moreover, [\[21, Theorem 56\]](#) and [\[26, Theorem 1.4\]](#) show sample complexity lower bounds of $\Omega(d/\alpha\varepsilon)$ and $\Omega(\log(1/\delta)/\varepsilon)$ for private learning of spherical and univariate Gaussians respectively. Finally, a lower bound of $\tilde{\Omega}(d^2/\alpha^2)$ is known for the non-private setting [\[3, Theorem 1.2\]](#).

The main ingredient of proving [Theorem 2.1](#) is a polynomial-time preprocessing algorithm which brings the condition number of the covariance matrix down to a constant. This algorithm is extremely simple: it compute a number of candidate covariance matrices from disjoint samples, averages the “good” candidates together, and then adds noise to the resulting covariance. Here, a candidate covariance matrix is “good” if it is close to a large majority of the other candidate covariance matrices; intuitively, this should also mean that it is also close to the true covariance matrix. The first two steps come directly from the framework discussed in [Section 7](#) and the last step is discussed in [Section 9](#).

In order to handle Gaussians with degenerate covariance matrices, we need a private and efficient method for learning (exactly) the subspace of a Gaussian. The recent work of Singhal and Steinke [\[33\]](#) offers a differentially private but inefficient method for this problem. We develop an efficient method for this problem which is of independent interest (see [Theorem 8.1](#) for a formal statement). We note that our result is somewhat incomparable to that of Singhal and Steinke [\[33\]](#) since the assumptions are different.

Theorem 2.2 (Informal). *There is an (ε, δ) -DP algorithm that given an i.i.d. samples from an arbitrary Gaussian distributions over \mathbb{R}^d , outputs the subspace that the Gaussian is supported on almost surely. Furthermore, the sample complexity and time complexities of this algorithm are $m = O\left(\frac{d \log(1/\delta)}{\varepsilon}\right)$ and $O(md)$ respectively.*

In this paper, we prove [Theorem 2.2](#) as an example of our technique from [Section 7](#). However, we remark that this is not strictly necessary and the same theorem can be proved using standard techniques such as private lazy histogram or propose-test-release.

Finally, we show how to learn Gaussians privately, robustly, and efficiently. An α -corrupted sample is a sample that is corrupted by an adversary in the following way: the adversary chooses an α -fraction of the points in the sample and modifies them arbitrarily. See [Corollary 11.5](#) for more details.

Theorem 2.3 (Informal). *There is a robust and (ε, δ) -DP algorithm which given an α -corrupted sample from an arbitrary but nonsingular⁵ Gaussian distributions over \mathbb{R}^d , outputs a distribution whose TV distance to the true distribution is $O(\alpha \log(1/\alpha))$ with probability at least $1 - \beta$. Furthermore, the sample complexity and time complexity of this algorithm is polynomial in $\left(\frac{d \log(1/\beta\delta)}{\alpha\varepsilon}\right)$.*

This is the first polynomial time algorithm for private and robust learning of Gaussian distributions. An important ingredient of the above theorem is a subroutine for learning the mean robustly. The following theorem settles the polynomial learnability of private and robust mean estimation (see [Theorem 11.4](#)).

⁵Our robust learning guarantee holds when the covariance matrix satisfies $\Sigma \succ 0$.

Theorem 2.4 (Informal). *There is a robust and (ε, δ) -DP algorithm which given an α -corrupted sample of size $\tilde{\Omega}\left(\frac{d^{3/2}\ln(1/\delta)}{\varepsilon\alpha^2}\right)$ from $\mathcal{N}(\mu, I_d)$ with unknown $\mu \in \mathbb{R}^d$, outputs $\tilde{\mu}$ such that $\|\mu - \tilde{\mu}\|_2 = O(\alpha\sqrt{\log(1/\alpha)})$ with probability at least $1 - \beta$. Furthermore, the time complexity of this algorithm is polynomial in m and d .*

We note that the sample complexity of the above result matches that of Theorem 7 in [30]; however, unlike [30] we do not need any boundedness assumptions on the mean.

3 Techniques

We present a simple, efficient, and fairly general reduction from private statistical estimation to its non-private counterpart. For the reduction, we run the non-private estimator on a number of disjoint subsets of the data set (in the spirit of Subsample-And-Aggregate framework [31]). We then privately check whether most of the (non-private) outcomes are “close” to each other (basically using the Propose-Test-Release framework [15]). If this is the case, then we aggregate the “good” outcomes by simple averaging, and output a “noisy” version of this average.

In Definition 7.1, we define the notion of a *convex semimetric space*, which generalizes metric spaces (e.g., by requiring only approximate triangle inequality). Our reduction works when the outcomes of the non-private estimators belong to a convex semimetric space. This allows us use our reduction for non-metric spaces. A key motivation of this generalization is handling covariance matrices of a Gaussian distribution. As such, one of our technical results is Lemma 7.2 which shows that covariance matrices can be worked with in this framework.

The final part of the reduction is to output a “noisy” version of the average. The notion of a *masking mechanism* abstracts the requirements we need from such a mechanism (see Definition 7.3). It is well known that the Laplace and Gaussian mechanisms work well for ℓ_1 and ℓ_2 spaces. Masking mechanisms generalize this idea for convex semimetric spaces. Another technical ingredient of learning covariance matrices is designing an efficient masking mechanism for them (see Lemma 9.1).

4 Discussion of a concurrent result

In an independent work, Kamath, Mouzakis, Singhal, Steinke, and Ullman [22] propose a different polynomial time algorithm for privately learning unbounded high-dimensional Gaussian distributions. At a high-level, they devise an iterative preconditioning method for the covariance matrix (in the spirit of [4, 21]). To resolve the challenge of handling covariance matrices with unbounded condition numbers, they propose a subspace learning method inspired by the approach of [33]. The iterative approach to bound the ratio of consecutive eigenvalues incurs additional cost of DP composition, resulting in a term in the sample complexity of order $O(d^{5/2}/\varepsilon)$ rather than $O(d^2/\varepsilon)$.

In contrast, our approach for preconditioning the covariance matrix is not iterative, and in its core relies on a masking/noising mechanism for covariance matrices. In particular, we are able to compute a preconditioner with roughly $\tilde{O}(d^2/\varepsilon)$ samples. Non-privately, $O(d)$ samples are needed to compute a preconditioner. It is an interesting open problem to see if $O(d)$ samples suffices for

the private setting as well. Our general reduction also allows us to apply the result in other settings such as *robust* learning of Gaussians in polynomial time.

5 More on related work

Most relevant to our work are the results on learning Gaussians with respect to the TV distance under the (ε, δ) -DP. Karwa and Vadhan [26] established a polynomial-time and sample-efficient method for learning unbounded univariate Gaussians under (ε, δ) -DP. One can use this method for learning multivariate Gaussians as well, by simply learning the one dimensional projection of the Gaussians along each axis (and invoking the composition for DP). While this approach works well for axis-aligned Gaussians, for general Gaussians its sample complexity depends on the condition number of the covariance matrix. Kamath, Li, Singhal, and Ullman [21] consider the problem of privately learning high-dimensional distributions. They provide a polynomial time algorithm for learning high-dimensional Gaussians under the (stronger notion of) concentrated DP [8, 18]. The sample complexity of this approach improves over the multivariate version of [26]; however, it still depends (logarithmically) on the condition number of covariance matrix. On a follow up work, Biswas, Dong, Kamath, and Ullman [4] propose the more practical method of CoinPress whose sample complexity matches that of [21] for learning multivariate Gaussians. CoinPress is shown to empirically work better than its counterparts—namely the multivariate versions of [13, 26]—in high dimensions.

The best known lower bounds for learning d -dimensional Gaussians up to TV distance α under (ε, δ) -DP is $\tilde{\Omega}(d/\alpha\varepsilon)$ [21]. In contrast, the sample complexity of all the methods described above depends on the condition number of the covariance matrix. Utilizing the private hypothesis selection framework proposed in [6], Aden-Ali, Ashtiani, and Kamath [1] proved the first sample complexity bound for privately learning general Gaussians that does not depend on the condition number or the size of the parameters. The proof is, however, non-constructive, as it invokes Zorn’s lemma to prove the existence of a locally small cover for covariance matrices.

In the non-private setting, an α -accurate estimation of the mean of a d -dimensional Gaussian distribution with respect to the Mahalanobis distance can be found using $O(d/\alpha^2)$ samples. Achieving linear dependence on d , however, is not easy in the private setting. Brown, Gaboardi, Smith, Ullman, and Zakyntinou [5] propose a method for solving this problem under (ε, δ) -DP with linear dependence on d . Their method is, however, not computationally efficient. One interesting aspect of [5] is the use of a variant of the Gaussian mechanism that is scaled with the empirical covariance. We also make use of a similar idea to noise the covariance matrix.

A related line of work is the problem of private mean estimation with respect to the Euclidean distance. It is noteworthy that these results do not translate to learning Gaussians with respect to the TV distance. Kamath, Singhal, and Ullman [24] prove tight minimax rates for mean estimation for distributions with bounded k -th moment. Liu, Kong, Kakade, and Oh [30] proposed a robust, efficient, and private mean estimation method for sub-Gaussian distributions with bounded mean.

Other related work include private Principal Component Analysis [9, 38] and private learning of Gaussian mixture models [2, 23]. We refer the readers to [17, 35] for an introduction to differential privacy, and to [25] for a survey on private statistical learning.

6 Preliminaries

For a vector $v \in \mathbb{R}^d$, we use $\|v\|_2$ to denote the Euclidean norm of the vector v . For a matrix $A \in \mathbb{R}^{d \times d}$, we use $\|A\|_F$ to denote the Frobenius norm of A and $\|A\|$ to denote the operator norm of A . The matrix A is positive semi-definite (resp. positive definite), denoted $A \succeq 0$ (resp. $A \succ 0$) if A is symmetric and all its eigenvalues are non-negative (resp. positive). If $A \succeq 0$ then $A^{1/2}$ denotes the unique positive semi-definite matrix satisfying $A = (A^{1/2})^2$. Further, if $A \succ 0$ then $A^{-1/2} = (A^{1/2})^{-1}$. In this paper, we also write \mathcal{S}^d to denote the positive-definite cone in $\mathbb{R}^{d \times d}$.

6.1 Probability Facts

In this paper, we use $d_{\text{TV}}(\mathcal{D}_1, \mathcal{D}_2)$ and $D_{\text{KL}}(\mathcal{D}_1 \parallel \mathcal{D}_2)$ to denote the total variation distance and KL divergence between two distributions $\mathcal{D}_1, \mathcal{D}_2$

Fact 6.1. *Let $\mu_1, \mu_2 \in \mathbb{R}^d$ and $\Sigma_1, \Sigma_2 \succ 0$. Then*

$$D_{\text{KL}}(\mathcal{N}(\mu_1, \Sigma_1) \parallel \mathcal{N}(\mu_2, \Sigma_2)) = \frac{1}{2} [\text{tr}(\Sigma_2^{-1}\Sigma_1 - I) + (\mu_2 - \mu_1)^\top \Sigma_2^{-1}(\mu_2 - \mu_1) - \ln \det(\Sigma_2^{-1}\Sigma_1)].$$

Moreover, suppose that all the eigenvalues of $\Sigma_2^{-1}\Sigma_1$ are at least $\frac{1}{2}$. Then

$$D_{\text{KL}}(\mathcal{N}(\mu_1, \Sigma_1) \parallel \mathcal{N}(\mu_2, \Sigma_2)) \leq \frac{1}{2} [\|\Sigma_2^{-1/2}\Sigma_1\Sigma_2^{-1/2} - I\|_F^2 + (\mu_2 - \mu_1)^\top \Sigma_2^{-1}(\mu_2 - \mu_1)]$$

The first assertion of [Fact 6.1](#) is a well-known formula for the KL divergence between Normal distributions. The proof of the second assertion can be found in [Appendix B.1](#).

Lemma 6.2 (Pinsker's Inequality [[34](#), Lemma 2.5]). *For any two distributions $\mathcal{D}_1, \mathcal{D}_2$, $d_{\text{TV}}(\mathcal{D}_1, \mathcal{D}_2)^2 \leq \frac{1}{2}D_{\text{KL}}(\mathcal{D}_1 \parallel \mathcal{D}_2)$.*

The following lemma is a standard bound on the (upper) tails of a Gaussian. (For example, it is a special case of [[37](#), Proposition 2.5] for $n = 1$).

Lemma 6.3. *Let $g \sim \mathcal{N}(0, \sigma^2)$. Then for all $x > 0$, $\mathbb{P}[g \geq \sigma x] \leq e^{-x^2/2}$.*

We require the following concentration inequality which can be easily derived from well-known techniques (see [[37](#), §2.1.3]). For completeness, we provide a proof in [Appendix B.2](#).

Lemma 6.4. *Let $g \sim \mathcal{N}(0, I_d)$ and A be a symmetric matrix. Then for all $x > 0$,*

$$\mathbb{P} \left[g^\top A g \geq \text{tr}(A) + 4\|A\|_F \cdot \sqrt{x} + 4\|A\| \cdot x \right] \leq \exp(-x).$$

6.2 Differential Privacy

In this section, we let \mathcal{X}, \mathcal{Y} denote sets and $n \in \mathbb{N}$. Two datasets $D = (X_1, \dots, X_n), D' = (X'_1, \dots, X'_n) \in \mathcal{X}^n$ are said to be *neighbouring* if $d_H(D, D') \leq 1$ where d_H denotes Hamming distance, i.e. $d_H(D, D') = |\{i \in [d] : X_i \neq X'_i\}|$.

Definition 6.5 ([14, 16]). A mechanism $\mathcal{M}: \mathcal{X}^n \rightarrow \mathcal{Y}$ is said to be (ε, δ) -differentially private if for all neighbouring datasets $D, D' \in \mathcal{X}^n$ and all $Y \subseteq \mathcal{Y}$, we have

$$\mathbb{P}[\mathcal{M}(D) \in Y] \leq e^\varepsilon \cdot \mathbb{P}[\mathcal{M}(D') \in Y] + \delta.$$

In this paper, we use DP as shorthand for differentially private or differential privacy, depending on context.

Remark 6.6. *The astute reading may notice that the algorithms we present in this paper are not invariant under permutation of the input. Given this, it may seem more desirable to define differential privacy where we view datasets as unordered sets instead of ordered tuples. However, one can show that if one has an (ε, δ) -DP algorithm for ordered tuples than one also has a (ε, δ) -DP for unordered sets by randomly permuting the input [5, Proposition D.6].*

Definition 6.7. Let $\mathcal{D}_1, \mathcal{D}_2$ be two continuous distributions defined on \mathbb{R}^d and let f_1, f_2 be the respective density functions. We use $\mathcal{L}_{\mathcal{D}_1 \parallel \mathcal{D}_2}: \mathbb{R}^d \rightarrow \mathbb{R}$ to denote the logarithm of the likelihood ratio, i.e. for any $x \in \mathbb{R}^d$,

$$\mathcal{L}_{\mathcal{D}_1 \parallel \mathcal{D}_2}(x) := \ln \frac{f_1(x)}{f_2(x)}. \quad (6.1)$$

Remark 6.8. *Note that $\mathbb{E}_{Y \sim \mathcal{D}_1} [\mathcal{L}_{\mathcal{D}_1 \parallel \mathcal{D}_2}(Y)] = D_{\text{KL}}(\mathcal{D}_1 \parallel \mathcal{D}_2)$.*

Definition 6.9. Let $\mathcal{D}_1, \mathcal{D}_2$ be two distributions defined on a set \mathcal{X} . Then $\mathcal{D}_1, \mathcal{D}_2$ are said to be (ε, δ) -indistinguishable if for all measurable $S \subseteq \mathcal{X}$, we have

$$\mathbb{P}_{Y \sim \mathcal{D}_1} [Y \in S] \leq e^\varepsilon \mathbb{P}_{Y \sim \mathcal{D}_2} [Y \in S] + \delta \quad \text{and} \quad \mathbb{P}_{Y \sim \mathcal{D}_2} [Y \in S] \leq e^\varepsilon \mathbb{P}_{Y \sim \mathcal{D}_1} [Y \in S] + \delta.$$

In light of Definition 6.9, a mechanism \mathcal{M} is (ε, δ) -DP if and only if for every neighbouring datasets D, D' , the distributions given by $\mathcal{M}(D)$ and $\mathcal{M}(D')$ are (ε, δ) -indistinguishable.

In this paper, we make use of the following sufficient condition for indistinguishability; a proof can be found in Appendix B.3.

Lemma 6.10. *Let $\mathcal{D}_1, \mathcal{D}_2$ be continuous distributions defined on \mathbb{R}^d . If*

$$\mathbb{P}_{Y \sim \mathcal{D}_1} [\mathcal{L}_{\mathcal{D}_1 \parallel \mathcal{D}_2}(Y) \geq \varepsilon] \leq \delta \quad \text{and} \quad \mathbb{P}_{Y \sim \mathcal{D}_2} [\mathcal{L}_{\mathcal{D}_2 \parallel \mathcal{D}_1}(Y) \geq \varepsilon] \leq \delta$$

then $\mathcal{D}_1, \mathcal{D}_2$ are (ε, δ) -indistinguishable.

Claim 6.11. *Let $\mu_1, \mu_2 \in \mathbb{R}^d$ and $\Sigma_1, \Sigma_2 \succ 0$. Then*

$$\mathcal{L}_{\mathcal{N}(\mu_1, \Sigma_1) \parallel \mathcal{N}(\mu_2, \Sigma_2)}(x) = \ln \left(\frac{\det(\Sigma_1)^{-1/2} \exp\left(-\frac{1}{2}(x - \mu_1)^\top \Sigma_1^{-1}(x - \mu_1)\right)}{\det(\Sigma_2)^{-1/2} \exp\left(-\frac{1}{2}(x - \mu_2)^\top \Sigma_2^{-1}(x - \mu_2)\right)} \right). \quad (6.2)$$

Proof. The density of a normal distribution $\mathcal{N}(\mu, \Sigma)$ in \mathbb{R}^d is given by

$$f(x) = (2\pi)^{-d/2} \det(\Sigma)^{-1/2} \exp\left(-\frac{1}{2}(x - \mu)^\top \Sigma^{-1}(x - \mu)\right),$$

so the claim follows by a direct calculation. □

In this paper, we make use of the truncated Laplace distribution, denoted by $\text{TLap}(\Delta, \varepsilon, \delta)$ whose p.d.f. is given by

$$f_{\text{TLap}(\Delta, \varepsilon, \delta)}(x) := \begin{cases} B e^{-|x|/\lambda} & x \in [-A, A] \\ 0 & x \notin [-A, A] \end{cases},$$

where $\lambda = \frac{\Delta}{\varepsilon}$, $A = \frac{\Delta}{\varepsilon} \ln \left(1 + \frac{e^\varepsilon - 1}{2\delta}\right)$, $B = \frac{1}{2\lambda(1 - e^{-A/\lambda})}$.

Theorem 6.12 ([19, Theorem 1]). *Suppose that $q: \mathcal{X} \rightarrow \mathbb{R}$ is a function with sensitivity Δ . Then the mechanism $q(x) + Y$ where $Y \sim \text{TLap}(\Delta, \varepsilon, \delta)$ is (ε, δ) -DP.*

7 A framework for private statistical estimation

In this section, we describe an efficient and fairly general reduction from private statistical estimation to the non-private counterpart. Our reduction is inspired by the Propose-Test-Release framework [15] and the Subsample-And-Aggregate framework [31]. At a very high level, we partition the data set into multiple disjoint subsets and run the non-private estimation method on each subset. We then privately check whether most of these solutions are “close” to each-other. If this is not the case then the method fails. Otherwise, we compute a weighted average of the “concentrated” solutions. Roughly speaking, solutions which are “close” to many other candidate solutions (say 0.7 fraction of them) are given weight 1 while solutions that are “close” to fewer candidate solutions (say less than 0.6 fraction of them) are given weight 0. To help with sensitivity, we then linearly interpolate the weight for solutions that are in the middle. Finally, we output a “noisy/masked” version of this average. In order for the reduction to work in rather general scenarios (such as learning the high dimensional Gaussians with respect the TV distance), we first define the notion of convex semimetric spaces. This helps to formalizes the notion of “closeness” between solutions as well as the weighted average over a number of solutions.

7.1 Convex semimetric spaces and masking mechanisms

In this paper, we work with the following notion of a convex semimetric space. The key property to keep in mind is that for semimetric spaces, we only have an *approximate* triangle inequality, as long as the points are significantly close together.

Definition 7.1 (Convex semimetric space). Let \mathcal{Y} be a convex set and let $\text{dist}: \mathcal{Y} \times \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$. We say $(\mathcal{Y}, \text{dist})$ is a convex semimetric space if there exists absolute constants $t \geq 1$, $\phi \geq 0$, and $r > 0$ such that for every $k \in \mathbb{N}$ and every $Y, Y_1, Y_2, \dots, Y_k \in \mathcal{Y}$, the following conditions hold.

1. $\text{dist}(Y, Y) = 0$ and $\text{dist}(Y_1, Y_2) \geq 0$.
2. **Symmetry.** $\text{dist}(Y_1, Y_2) = \text{dist}(Y_2, Y_1)$.
3. **t -approximate r -restricted triangle inequality.** If $\text{dist}(Y_1, Y_2), \text{dist}(Y_2, Y_3) \leq r$ then $\text{dist}(Y_1, Y_3) \leq t \cdot (\text{dist}(Y_1, Y_2) + \text{dist}(Y_2, Y_3))$.
4. **Convexity.** For all $\alpha \in \Delta_k$, $\text{dist}(\sum \alpha_i Y_i, Y) \leq \sum \alpha_i \text{dist}(Y_i, Y)$.
5. **ϕ -Locality.** For all $\alpha, \alpha' \in \Delta_k$, $\text{dist}(\sum_i \alpha_i Y_i, \sum_i \alpha'_i Y_i) \leq \sum_i |\alpha_i - \alpha'_i| (\phi + \max_{i,j} (\text{dist}(Y_i, Y_j)))$

where Δ_k denotes the k -dimensional probability simplex. When r is unspecified we take it to mean that $r = \infty$ and refer to it as a t -approximate triangle inequality.

The following technical lemma (whose proof appears in Appendix C) is helpful for learning covariance matrices.

Lemma 7.2. *Let \mathcal{S}^d be the set of all $d \times d$ positive definite matrices. For $A, B \in \mathcal{S}^d$ let $\text{dist}(A, B) = \max\{\|A^{-1/2}BA^{-1/2} - I\|, \|B^{-1/2}AB^{-1/2} - I\|\}$. Then $(\mathcal{S}^d, \text{dist})$ is a convex semimetric which satisfies a $(3/2)$ -approximate 1-restricted triangle inequality and 1-locality.*

Definition 7.3 (Masking mechanism). Let $(\mathcal{Y}, \text{dist})$ be a convex semimetric space. A $(\gamma, \varepsilon, \delta)$ -masking mechanism for $(\mathcal{Y}, \text{dist})$, denoted by $\mathcal{B}_{\gamma, \varepsilon, \delta} : \mathcal{Y} \rightarrow \mathcal{Y}$, is a randomized algorithm such that for all $Y_1, Y_2 \in \mathcal{Y}$, if $\text{dist}(Y_1, Y_2) \leq \gamma$ then $\mathcal{B}_{\gamma, \varepsilon, \delta}(Y_1)$ is (ε, δ) -indistinguishable from $\mathcal{B}_{\gamma, \varepsilon, \delta}(Y_2)$.

Remark 7.4. *It follows from standard analysis that for $p \geq 1$, the vector space \mathbb{R}^d equipped with $\text{dist}(Y_1, Y_2) = \|Y_1 - Y_2\|_p$ forms a convex semimetric space (see Proposition 11.1). Interestingly, the Laplace mechanism [16] can be viewed as a masking mechanism for (\mathbb{R}^d, ℓ_1) . Also, Gaussian mechanism can be viewed as a masking mechanism for (\mathbb{R}^d, ℓ_2) . More challenging is to establish a masking mechanism for the set of covariance matrices with respect to $\text{dist}(A, B) = \max\{\|A^{-1/2}BA^{-1/2} - I\|, \|B^{-1/2}AB^{-1/2} - I\|\}$ which we do in Section 9.*

Definition 7.5 (Concentration of a masking mechanism). Let $(\mathcal{Y}, \text{dist})$ be a convex semimetric space and $\mathcal{B}_{\gamma, \varepsilon, \delta}$ be $(\gamma, \varepsilon, \delta)$ -masking mechanism for it. We say $\mathcal{B}_{\gamma, \varepsilon, \delta}$ is (α, β) concentrated if for all $Y \in \mathcal{Y}$, $\mathbb{P}[\text{dist}(\mathcal{B}_{\gamma, \varepsilon, \delta}(Y), Y) > \alpha] \leq \beta$.

7.2 The Private Populous Mean Estimator

In this section we outline an efficient and fairly general reduction from private statistical estimation to the non-private counterpart. We use $\mathcal{A} : \mathcal{X}^* \rightarrow \mathcal{Y}$ to denote a generic (efficient) non-private mechanism which takes a collection of samples as inputs and outputs an element in \mathcal{Y} . We assume that $(\mathcal{Y}, \text{dist})$ is a convex semimetric space, and we have access to (an efficient) masking mechanism for $(\mathcal{Y}, \text{dist})$. Intuitively, the error of the estimator is measured by $\text{dist}(\hat{Y}, Y^*)$, where \hat{Y} is our estimation while Y^* is the ground truth.

At a high-level, the algorithm works as follows. We split the dataset into k disjoint groups and run the non-private algorithm \mathcal{A} on each group i to obtain an output Y_i . We then consider a ball of radius r/t around Y_i with respect to dist ⁶ and compute q_i , the fraction of Y_j 's that are inside this ball. We then look at the average value of the q_i , i.e., $Q = \frac{1}{k} \sum_{i \in [k]} q_i$. Intuitively, Q efficiently measures the “stability” of \mathcal{A} across the data set partitions. Namely, Q is large when most of the Y_i 's are concentrated in a small “popular region”, and small when Y_i 's are scattered around \mathcal{Y} . If Q is small, then \mathcal{A} has not worked well and we can announce failure privately using a noisy threshold. If Q is large, then we take a weighted average of Y_i 's that are in the popular region and then output a “masked” version of this average. The following lemma establishes the privacy and accuracy of Algorithm 1 and its proof can be found in Appendix C.2.

Lemma 7.6. *Let $(\mathcal{Y}, \text{dist})$ be a convex semimetric space that satisfies r -restricted t -approximate triangle inequality and ϕ -locality for some $r > 0, t \geq 1, \phi \geq 0$. Let $\mathcal{B}_{400(r+\phi)/k, \varepsilon, \delta}$ be a $(400(r+\phi)/k, \varepsilon, \delta)$ -masking mechanism with respect to $(\mathcal{Y}, \text{dist})$. Then*

1. **Privacy.** For $k \geq 30$, Algorithm 1 is $(2\varepsilon, 3e^\varepsilon\delta)$ -DP.

⁶ r and t are chosen in a way that $(\mathcal{Y}, \text{dist})$ satisfies r -restricted t -approximate triangle inequality.

Algorithm 1 Private to non-private reduction via populous mean estimation

Input: Dataset $D = (X_1, \dots, X_m)$; parameters $\varepsilon, \delta, r, \phi > 0$; $k \leq m$; $t \geq 2$.

- 1: Let $s \leftarrow \lfloor m/k \rfloor$.
 - 2: For $i \in [k]$, let $Y_i \leftarrow \mathcal{A}(\{X_\ell\}_{\ell=(i-1)s+1}^{is})$.
 - 3: For $i \in [k]$, let $q_i \leftarrow \frac{1}{k+3} |\{j \in [k] : \text{dist}(Y_i, Y_j) \leq r/t\}|$.
 - 4: Let $Q \leftarrow \frac{1}{k} \sum_{i \in [k]} q_i$.
 - 5: Let $Z \sim \text{TLap}(2/k, \varepsilon, \delta)$.
 - 6: Let $\widehat{Q} \leftarrow Q + Z$.
 - 7: If $\widehat{Q} < 0.8 + \frac{2}{k\varepsilon} \ln(1 + \frac{e^\varepsilon - 1}{2\delta})$, fail and return \perp .
 - 8: For $i \in [k]$, let $w_i = \min(1, 10 \max(0, q_i - 0.6))$.
 - 9: $\mu = (\sum_{i=1}^k w_i Y_i) / (\sum_{i=1}^k w_i)$.
 - 10: Return $\mathcal{B}_{400(r+\phi)/k, \varepsilon, \delta}(\mu)$.
-

2. **Utility.** Assume $k \geq \max\{30, \frac{20}{\varepsilon} \ln(1 + \frac{e^\varepsilon - 1}{2\delta})\}$. For $\alpha_1 \leq r/t$ assume that there exist $Y^* \in \mathcal{Y}$ such that $\text{dist}(Y_i, Y^*) \leq \alpha_1/t$ for all $i \in [k]$. If \mathcal{B} is $(\alpha_2/t, \beta)$ -concentrated then the output \widehat{Y} of Algorithm 1 satisfies

$$\mathbb{P} \left[\text{dist}(\widehat{Y}, Y^*) > \alpha_1 + \alpha_2 \right] \leq \beta$$

Remark 7.7. For privacy, Lemma 7.6 requires that k is large enough (at least 30). If this is not the case, we can always add “dummy” points into the dataset.

Remark 7.8. The computational complexity of Algorithm 1 is dominated by these computations:

- k queries to the oracle \mathcal{A} (with inputs of size m/k).
- k^2 computations of the $\text{dist}(\cdot, \cdot)$ function.
- One query to the masking mechanism \mathcal{B} .

In particular, if \mathcal{A} , \mathcal{B} , and the algorithm for computing dist all run in polynomial time then Algorithm 1 runs in polynomial time.

8 Learning the subspace

As a simple application of our technique, we consider the problem of privately learning a subspace. More specifically, if the samples are drawn from $\mathcal{N}(0, \Sigma)$ then the goal is to learn the range of Σ . Recently, Singhal and Steinke [33] also studied the problem of privately learning a subspace albeit under slightly different assumption than ours. For learning a subspace *exactly*, they gave an algorithm that was sample-optimal but not computationally-efficient. For learning a subspace *approximately*, they provided an algorithm that was sample-efficient. The latter was also used (with some modifications) in [22] to efficiently learn the covariance matrix of a Gaussian distribution. In this section, we give a computationally-efficient algorithm for learning a subspace *exactly* with a slightly different sample complexity (albeit with different assumptions). Formally, we prove the following theorem.

Theorem 8.1. *There exists a mechanism \mathcal{M} satisfying the following. For all $m, d \in \mathbb{N}$, $\varepsilon, \delta \in (0, 1)$, and dataset $D = (X_1, \dots, X_m)$:*

1. Given ε, δ, D as input, \mathcal{M} is $(2\varepsilon, 3e^\varepsilon\delta)$ -DP and runs in time $O(md)$.
2. If $\Sigma \succeq 0$, $X_1, \dots, X_m \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \Sigma)$, and $m \geq \max\{30, \lceil \frac{20}{\varepsilon} \ln(1 + \frac{e^\varepsilon - 1}{2\delta}) \rceil\} \cdot d = O(d \log(1/\delta)/\varepsilon)$ then, almost surely, \mathcal{M} outputs the orthogonal projection matrix onto the range of Σ .

Remark 8.2. We remark that the result we prove in this section can also be obtained using standard techniques such as propose-test-release or private lazy histograms. However, we choose to apply our technique to give the reader an example of how it may be used.

The mechanism that achieves [Theorem 8.1](#) appears in [Algorithm 2](#) and is designed using the template given in [Algorithm 1](#). In this section, we define

$$\text{dist}(A, B) = \begin{cases} 0 & \text{if } A = B \text{ are orthogonal projection matrices} \\ \infty & \text{otherwise} \end{cases},$$

where A, B are two matrices in $\mathbb{R}^{d \times d}$. Note that this makes the choice of r, t, ϕ moot since the distance between any two matrices is either 0 or ∞ .

Algorithm 2 A private algorithm for learning the subspace.

Input: Dataset $D = (X_1, \dots, X_m)$; parameters $\varepsilon, \delta \in (0, 1)$.

▷ Parameter settings.

1: $r \leftarrow 1, t \leftarrow 1, \phi \leftarrow 1$.

▷ Not used in [Algorithm 1](#).

2: $k \leftarrow \max\{30, \lceil \frac{20}{\varepsilon} \ln(1 + \frac{e^\varepsilon - 1}{2\delta}) \rceil\}$

3: **function** $\mathcal{A}(X_1, \dots, X_s)$

4: Let P be orthogonal projection matrix onto $\text{span}\{X_1, \dots, X_s\}$.

5: **Return** P .

6: **end function**

7: **function** $\mathcal{B}(P)$

8: **Return** P .

9: **end function**

Proof of [Theorem 8.1](#). Using the convention $\infty \leq \infty$, it is easy to check that the dist function we defined above forms a convex semimetric space for matrices (see [Definition 7.1](#)). Also, the identity function is a $(\gamma, \varepsilon, \delta)$ -masking mechanism for all $\gamma, \varepsilon, \delta \in (0, 1)$ since for all A, B , $\text{dist}(A, B) < \infty$ only if A, B are both orthogonal projection matrices and $A = B$. Therefore, the privacy statement follows from [Lemma 7.6](#).

For the utility, note that the masking mechanism is the identity function, so it is $(0, 0)$ -concentrated. Since we set $k = \max\{30, \lceil \frac{20}{\varepsilon} \ln(1 + \frac{e^\varepsilon - 1}{2\delta}) \rceil\}$, our assumption on m implies that $m \geq kd$. In particular, $s \geq d$ (see [Algorithm 1](#)). Therefore, with probability 1, the output of $\mathcal{A}(\{X_\ell\}_{\ell=(i-1)s+1}^{is})$ is the orthogonal projection matrix onto the range of Σ for all $i \in [k]$ almost surely. Therefore, the utility conditions of [Lemma 7.6](#) are satisfied with $\alpha_1 = \alpha_2 = \beta = 0$. \square

9 An efficient algorithm for learning the covariance matrix

In this section we show how to use the general reduction to learn high-dimensional Gaussian distributions with respect to the total variation distance. Here, the main hurdle to find an approximation to the covariance matrix for the zero-mean case. Based on [Lemma 7.2](#), the following dist function forms a semimetric space for positive definite matrices.

$$\text{dist}(\Sigma_1, \Sigma_2) = \begin{cases} \max \left(\|\Sigma_2^{-1/2} \Sigma_1 \Sigma_2^{-1/2} - I_d\|, \|\Sigma_1^{-1/2} \Sigma_2 \Sigma_1^{-1/2} - I_d\| \right) & \text{if } \text{rank } \Sigma_1 = \text{rank } \Sigma_2 = d \\ \infty & \text{otherwise} \end{cases} \quad (9.1)$$

The main steps to use the reduction are to define a masking mechanism and show that it is concentrated. We start by stating the main algorithmic idea of this section, which is a masking mechanism with respect to the dist function in Eq. (9.1).

Lemma 9.1. *For a matrix $\Sigma \succ 0$, define $\mathcal{B}(\Sigma) = \Sigma^{1/2}(I + \eta G)(I + \eta G)^\top \Sigma^{1/2}$ where $G \in \mathbb{R}^{d \times d}$ is a $d \times d$ matrix with independent $\mathcal{N}(0, 1)$ entries.*

Then for every $\eta > 0$ and $\varepsilon, \delta \in (0, 1]$, \mathcal{B} is a $(\gamma, \varepsilon, \delta)$ -masking mechanism with respect to dist for

$$\gamma = \min \left\{ \sqrt{\frac{\varepsilon}{2d(d + 1/\eta^2)}}, \frac{\varepsilon}{8d\sqrt{\ln(2/\delta)}}, \frac{\varepsilon}{8\ln(2/\delta)}, \frac{\varepsilon\eta}{12\sqrt{d}\sqrt{\ln(2/\delta)}} \right\}.$$

The proof of [Lemma 9.1](#) can be found in [Appendix D.1](#). Next, we show that the masking mechanism in [Lemma 9.1](#) is concentrated.

Lemma 9.2. *Let $\Sigma \succ 0$ and set $\eta = \frac{1}{C_1(\sqrt{d} + \sqrt{\ln(4/\beta)})}$ for a sufficiently large (universal) constant $C_1 > 0$. Then the masking mechanism $\mathcal{B}(\Sigma)$ defined in [Lemma 9.1](#) is $(1/100, \beta/2)$ -concentrated.*

Proof. Let $\widehat{\Sigma} = \mathcal{B}(\Sigma) = \Sigma^{1/2}(I + \eta G)(I + \eta G)^\top \Sigma^{1/2}$. First, we prove that

$$\|\Sigma^{-1/2} \widehat{\Sigma} \Sigma^{-1/2} - I_d\| \leq 1/400,$$

with probability $1 - \beta/2$. Indeed, we have that, for a sufficiently large constant $C' > 0$,

$$\begin{aligned} \|\Sigma^{-1/2} \widehat{\Sigma} \Sigma^{-1/2} - I_d\| &= \|(I_d + \eta G)(I + \eta G)^\top - I_d\| \\ &\leq 2\eta G + \eta^2 \|GG^\top\| \\ &\leq 2\eta C'(\sqrt{d} + \sqrt{\ln(4/\beta)}) + \eta^2 (C'(\sqrt{d} + \sqrt{\ln(4/\beta)}))^2, \end{aligned}$$

where we used the fact that $\|G\| \leq C(\sqrt{d} + \sqrt{\ln(4/\beta)})$ with probability $1 - \beta/2$ (see [Theorem A.5](#)). Taking $\eta \leq \frac{1}{1600C(\sqrt{d} + \sqrt{\ln(4/\beta)})}$ gives that

$$\|\Sigma^{-1/2} \widehat{\Sigma} \Sigma^{-1/2} - I_d\| \leq 1/400,$$

as desired.

In order to show that $\text{dist}(\Sigma, \widehat{\Sigma})$, it remains to show that

$$\|\widehat{\Sigma}^{-1/2}\Sigma\widehat{\Sigma}^{-1/2} - I_d\| \leq 1/100.$$

This follows directly from [Lemma A.6](#). \square

Algorithm 3 summarizes the modules needed for [Lemma 7.6](#) to work, including the definitions of the mechanism \mathcal{A}, \mathcal{B} as well as the required parameters. Note that \mathcal{A} returns just the empirical covariance matrix (although we could easily replace it by any algorithm which tries to approximate the covariance matrix). One last piece is to show that the output of the non-private \mathcal{A} is accurate enough.

Lemma 9.3. *There is a universal constant C_2 such that the following holds. Suppose $s \geq C_2(d + \ln(4/\beta))$. Let $X_1, \dots, X_s \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \Sigma)$ and $\widehat{\Sigma} = \sum_{\ell=1}^s X_\ell X_\ell^\top$. Then $\text{dist}(\Sigma, \widehat{\Sigma}) \leq 1/100$ with probability $1 - \beta/2$.*

Proof. Applying a standard concentration inequality (see [Lemma A.4](#)), we have $\|\Sigma^{-1/2}\widehat{\Sigma}\Sigma^{-1/2} - I_d\| \leq C \left(\sqrt{\frac{d + \ln(4/\beta)}{s}} + \frac{d + \ln(4/\beta)}{s} \right)$. Choosing $s \geq 800^2 C^2 (d + \ln(4/\beta))$ gives that $\|\Sigma^{-1/2}\widehat{\Sigma}\Sigma^{-1/2} - I_d\| \leq 1/400$. Applying [Lemma A.6](#) gives that $\text{dist}(\widehat{\Sigma}, \Sigma) \leq 1/100$ with probability $1 - \beta/2$ as required. \square

Algorithm 3 A private algorithm for learning the covariance matrix up to a constant factor with respect to the spectral distance

Input: Dataset $D = (X_1, \dots, X_m)$; parameters $\varepsilon, \delta, \alpha, \beta \in (0, 1)$

▷ Some parameter settings.

- 1: $\eta \leftarrow \frac{1}{C_1(\sqrt{d} + \sqrt{\ln(4/\beta)})}$ ▷ C_1 from [Lemma 9.2](#)
- 2: $\gamma \leftarrow \min \left\{ \sqrt{\frac{\varepsilon}{2d(d+1/\eta^2)}}, \frac{\varepsilon}{8d\sqrt{\ln(2/\delta)}}, \frac{\varepsilon}{8\ln(2/\delta)}, \frac{\varepsilon\eta}{12\sqrt{d\ln(2/\delta)}} \right\}$.
- 3: $r \leftarrow 1, \phi \leftarrow 1$
- 4: $k \leftarrow \max \left\{ \frac{400(r+\phi)}{\gamma}, \frac{20}{\varepsilon} \ln \left(1 + \frac{e^\varepsilon - 1}{2\delta} \right) \right\}$
- 5: $t \leftarrow 3/2$
- 6: **function** $\mathcal{A}(X_1, \dots, X_s)$
- 7: $\widehat{\Sigma} = \frac{1}{s} \sum_{\ell=1}^s X_\ell X_\ell^\top$
- 8: **Return** $\widehat{\Sigma}$.
- 9: **end function**
- 10: **function** $\mathcal{B}(\widehat{\Sigma})$
- 11: Let G be $d \times d$ matrix with independent $\mathcal{N}(0, 1)$ entries.
- 12: **Return** $\widehat{\Sigma}^{1/2}(I + \eta G)(I + \eta G)^\top \widehat{\Sigma}^{1/2}$.
- 13: **end function**

Now we are ready to apply [Lemma 7.6](#). The following theorem shows that we can learn the covariance matrix up to a constant distance with respect to the spectral distance using more or less $O(d^2/\varepsilon)$ samples.

Theorem 9.4. Applying Lemma 7.6 with parameters specified in Algorithm 3 gives a mechanism \mathcal{M} satisfying the following. For all $m, d \in \mathbb{N}$, $\varepsilon, \delta, \beta \in (0, 1)$, and dataset $D = (X_1, \dots, X_m)$:

1. Given $\varepsilon, \delta, \beta, D$ as input, \mathcal{M} is $(2\varepsilon, 3e^\varepsilon \delta)$ -DP and runs in time $O(md^2 + kd^3)$.
2. If $\Sigma \succ 0$, $X_1, \dots, X_m \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \Sigma)$, and $m = \tilde{\Omega}\left(\frac{d^2 \ln(1/\beta)^{3/2} \ln(1/\delta)^{1/2}}{\varepsilon}\right)$ then with probability at least $1 - \beta$, \mathcal{M} outputs a matrix $\hat{\Sigma}$ such that $\text{dist}(\Sigma, \hat{\Sigma}) \leq 1/10$.

Proof. Lemma 7.2 shows that $(\mathcal{S}^d, \text{dist})$ satisfies the assumptions of Definition 7.1. Lemma 9.1 (with $r = \phi = 1, t = 3/2$) shows that \mathcal{B} (as defined in Lemma 9.1 and Algorithm 3) as a $(\gamma, \varepsilon, \delta)$ -masking mechanism. Since $k \geq 400(r + \phi)/\gamma$ (so $400(r + \phi)/k \leq \gamma$) it follows that \mathcal{B} is also a $(400(r + \phi)/k, \varepsilon, \delta)$ -masking mechanism. Therefore, privacy immediately follows from the first assertion of Lemma 7.2.

Next, if $m \geq C_2(d + \ln(4k/\beta)) \cdot k = \tilde{O}\left(\frac{d(d\sqrt{\ln(1/\delta)} + \ln(1/\delta)) \ln(1/\beta)^{3/2}}{\varepsilon}\right)$ then $\hat{\Sigma}_i = \mathcal{A}(\{X_\ell\}_{\ell=(i-1)s+1}^{is})$ satisfies $\text{dist}(\Sigma, \hat{\Sigma}_i) \leq 1/10 \leq r/t$ for all $i \in [k]$ with probability $1 - \beta/2$ by Lemma 9.3 and a union bound over k . In addition, \mathcal{B} is $(1/100, \beta/2)$ -concentrated.

To conclude, let $Y_i = \hat{\Sigma}_i$, $Y^* = \Sigma$, and $\hat{Y} = \hat{\Sigma}$ be the output of Algorithm 1 with the modules specified in Algorithm 3. Then, taking $\alpha_1 = \alpha_2 = 1/25$ in Lemma 7.2 gives that $\text{dist}(\hat{\Sigma}, \Sigma) \leq 1/10$ with probability $1 - \beta$. \square

Assuming $\Sigma \succ 0$, Theorem 9.4 gives us a covariance matrix $\hat{\Sigma}$ such that $0.9I_d \preceq \hat{\Sigma}^{-1/2} \Sigma \hat{\Sigma}^{-1/2} \preceq 1.1I_d$. Note that if $X \sim \mathcal{N}(0, \Sigma)$ then $\hat{\Sigma}^{-1/2} X \sim \mathcal{N}(0, \hat{\Sigma}^{-1/2} \Sigma \hat{\Sigma}^{-1/2})$. We can now make use of off-the-shelf private covariance methods which are efficient when the covariance matrix is well-conditioned [4, 21]. For concreteness, we use the following theorem which is paraphrased from [21, Theorem 1.1], [4, Theorem 3.3].

Theorem 9.5 ([4, 21]). *There is a polynomial time (ε, δ) -DP algorithm that given*

$$m = \tilde{\Omega}\left(\left(\frac{d^2}{\alpha^2} + \frac{d^2 \sqrt{\ln(1/\delta)}}{\alpha \varepsilon}\right) \cdot \ln(1/\beta)\right)$$

samples from a Gaussian $\mathcal{N}(0, \Sigma)$ with unknown covariance matrix Σ such that $0.9I \preceq \Sigma \preceq 1.1I$ and outputs $\hat{\Sigma}$ satisfying $\|\hat{\Sigma}^{-1/2} \Sigma \hat{\Sigma}^{-1/2} - I_d\|_F \leq \alpha$. In particular, $d_{\text{TV}}(\mathcal{N}(0, \Sigma), \mathcal{N}(0, \hat{\Sigma})) \leq \alpha$.

Finally, we prove our result for general, full-rank covariance matrices. Note that one can use Theorem 8.1 to easily extend this to the case where Σ is not necessarily full-rank. The details can be found in Subsection 9.1.

Theorem 9.6. *There exists a mechanism \mathcal{M} satisfying the following. For all $m, d \in \mathbb{N}$, $\varepsilon, \delta, \alpha, \beta \in (0, 1)$, and dataset $D = (X_1, \dots, X_m)$:*

1. Given $\varepsilon, \delta, \alpha, \beta, D$ as input, \mathcal{M} is $(3\varepsilon, (3e^\varepsilon + 1)\delta)$ -DP and runs in polynomial time.
2. If $\Sigma \in \mathbb{S}^d$, $X_1, \dots, X_m \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \Sigma)$, and

$$m = \tilde{\Omega}\left(\frac{(d^2 \sqrt{\ln(1/\delta)} + d \ln(1/\delta)) \ln(1/\beta)^{3/2}}{\varepsilon} + \left(\frac{d^2}{\alpha^2} + \frac{d^2 \sqrt{\ln(1/\delta)}}{\alpha \varepsilon}\right) \cdot \ln(1/\beta)\right)$$

then with probability $1 - \beta$, \mathcal{M} outputs a matrix $\widehat{\Sigma}$ such that $d_{\text{TV}}\left(\mathcal{N}(0, \Sigma), \mathcal{N}(0, \widehat{\Sigma})\right) \leq \alpha$.

Proof. Combining [Theorem 9.4](#) and [Theorem 9.5](#) concludes the result. The privacy follows from basic composition since the algorithm has only three steps. The utility directly follows from the adopted theorems. \square

9.1 An efficient algorithm for learning a Gaussian

In this subsection, we sketch the ideas needed to learn a Gaussian $\mathcal{N}(\mu, \Sigma)$ under (ε, δ) -DP with no assumptions on μ, Σ . Suppose we draw $2m_1 + 2m_2 + m_3 + m_4$ samples from a Gaussian distribution $\mathcal{N}(\mu, \Sigma)$, where m_1 is the sample complexity for learning a subspace using [Algorithm 2](#) and m_2 is the sample complexity for learning a full-rank covariance matrix using [Algorithm 3](#) and [Theorem 9.5](#). The values of m_3 and m_4 are discussed below. We now show how to privately and efficiently learn a Gaussian in four steps.

1. Let $Z_i = \frac{X_i - X_{m_1+i}}{\sqrt{2}}$ for $i \in [m_1]$. We use Z_1, \dots, Z_{m_1} as input into the private subspace learning algorithm [Algorithm 2](#). As long as m_1 is sufficiently large, with probability $1 - \beta$, it will output a matrix P which is the projection onto the column space of Σ .
2. For $i \in [m_2]$, let $Z_i = \frac{X_{2m_1+i} - X_{2m_1+m_2+i}}{\sqrt{2}}$. Also, write $P = U_r U_r^\top$ where U_r is a $d \times r$ matrix whose rows form an orthonormal basis for the column space of Σ (equivalently, P). Note that $U_r^\top Z_i \sim \mathcal{N}(0, U_r^\top \Sigma U_r)$ and that $U_r^\top \Sigma U_r$ is a full rank $r \times r$ covariance matrix. We use [Algorithm 3](#) to learn an $r \times r$ covariance matrix $\widehat{\Sigma}_r$ such that $\|\widehat{\Sigma}_r^{-1/2} U_r^\top \Sigma U_r \widehat{\Sigma}_r^{-1/2} - I_r\|_F \leq \alpha$. In particular, $d_{\text{TV}}\left(\mathcal{N}(0, \widehat{\Sigma}_r), \mathcal{N}(0, U_r^\top \Sigma U_r)\right) \leq O(\alpha)$ (see [Theorem 9.6](#)).

Next, we learn the mean μ in two steps.

3. For $i \in [m_3]$, let $Z_i = X_{2m_1+2m_2+i}$. Then $\widehat{\Sigma}_r^{-1/2} U_r^\top Z_i \sim \mathcal{N}(\widehat{\Sigma}_r^{-1/2} U_r^\top \mu, \widehat{\Sigma}_r^{-1/2} U_r^\top \Sigma U_r \widehat{\Sigma}_r^{-1/2})$. Note that (for $\alpha \leq 0.5$), we have $0.5I \preceq \widehat{\Sigma}_r^{-1/2} U_r^\top \Sigma U_r \widehat{\Sigma}_r^{-1/2} \preceq 1.5I$. In particular, each coordinate of $\widehat{\Sigma}_r^{-1/2} U_r^\top Z_i$ has a variance which is bounded below by $1/2$ and above by 1.5 . Hence, applying [\[26, Theorem 1.3\]](#), we can obtain an estimate $\widehat{\mu}_r \in \mathbb{R}^d$ such that $\|\widehat{\mu}_r - \widehat{\Sigma}_r^{-1/2} U_r^\top \mu\|_2 \leq \alpha$ provided $m_3 = \widetilde{\Omega}\left(\frac{d}{\alpha^2} + \frac{d \ln(1/\delta)}{\alpha \varepsilon}\right)$ where $\widetilde{\Omega}$ hides $\text{polylog}(1/\beta, d, \ln(1/\delta), 1/\varepsilon, 1/\alpha)$. We remark that this step can be done in polynomial time.

We now have that

$$\begin{aligned}
d_{\text{TV}}\left(\mathcal{N}(U_r^\top \mu, U_r^\top \Sigma U_r), \mathcal{N}(\widehat{\Sigma}_r^{1/2} \widehat{\mu}_r, \widehat{\Sigma}_r)\right) &\leq d_{\text{TV}}\left(\mathcal{N}(U_r^\top \mu, U_r^\top \Sigma U_r), \mathcal{N}(U_r^\top \mu, \widehat{\Sigma}_r)\right) \\
&\quad + d_{\text{TV}}\left(\mathcal{N}(U_r^\top \mu, \widehat{\Sigma}_r), \mathcal{N}(\widehat{\Sigma}_r^{1/2} \widehat{\mu}_r, \widehat{\Sigma}_r)\right) \\
&= d_{\text{TV}}\left(\mathcal{N}(0, U_r^\top \Sigma U_r), \mathcal{N}(0, \widehat{\Sigma}_r)\right) \\
&\quad + d_{\text{TV}}\left(\mathcal{N}(\widehat{\Sigma}_r^{-1/2} U_r^\top \mu, I_r), \mathcal{N}(\widehat{\mu}_r, I_r)\right) \\
&\leq O(\alpha),
\end{aligned}$$

where the first line uses the triangle inequality, the second line uses the fact that total variation distance is invariant under bijective transformations, and the last line uses steps 2 and 3 above as

well as [Fact 6.1](#) and [Lemma 6.2](#) to assert that the total variation distance is $O(\alpha)$. Using the fact that bijective transformations are invariant once again, that $P = U_r U_r^\top$, and that $P\Sigma P = \Sigma$, we have

$$d_{\text{TV}}\left(\mathcal{N}(P\mu, \Sigma), \mathcal{N}(U_r \widehat{\Sigma}_r^{1/2} \widehat{\mu}_r, U_r \widehat{\Sigma}_r U_r^\top)\right) = d_{\text{TV}}\left(\mathcal{N}(U_r^\top \mu, U_r^\top \Sigma U_r), \mathcal{N}(\widehat{\Sigma}_r^{1/2} \widehat{\mu}_r, \widehat{\Sigma}_r)\right) \leq O(\alpha).$$

All that remains is to learn $(I - P)\mu$.

4. Now let $Z_i = X_{2m_1+2m_2+m_3+i}$ for $i \in [m_4]$. Note that $(I - P)$ is the orthogonal projection matrix onto the orthogonal complement of the column space of Σ . In particular $(I - P)Z_i = (I - P)\mu$ with probability 1 if $Z_i \sim \mathcal{N}(0, \Sigma)$. Learning $(I - P)\mu$ can now be done using standard techniques (for example, lazy private histograms or propose-test-release) with sample complexity $O(\log(1/\delta)/\varepsilon)$. Let us call the output of this step $\widehat{\mu}_\perp$.

With the last step (and again, using that bijective maps does not affect the total variation distance to add in $\widehat{\mu}_\perp$), we have

$$d_{\text{TV}}\left(\mathcal{N}(\mu, \Sigma), \mathcal{N}(U_r \widehat{\Sigma}_r^{1/2} \widehat{\mu}_r + \widehat{\mu}_\perp, U_r \widehat{\Sigma}_r U_r^\top)\right) \leq O(\alpha)$$

To conclude, we have the following theorem. We remind the reader that Σ is not assumed to be full-rank.

Theorem 9.7. *There exists a mechanism \mathcal{M} satisfying the following. For all $m, d \in \mathbb{N}$, $\varepsilon, \delta, \alpha, \beta \in (0, 1)$, and dataset $D = (X_1, \dots, X_m)$:*

1. *Given $\varepsilon, \delta, \alpha, \beta, D$ as input, \mathcal{M} is $(O(\varepsilon), O(e^\varepsilon \delta))$ -DP and runs in polynomial time.*
2. *If $X_1, \dots, X_m \sim \mathcal{N}(\mu, \Sigma)$ and*

$$m = \widetilde{\Omega}\left(\frac{d^2}{\alpha^2} + \frac{d^2 \sqrt{\ln(1/\delta)}}{\alpha \varepsilon} + \frac{d \ln(1/\delta)}{\alpha \varepsilon}\right)$$

then with probability $1 - \beta$, \mathcal{M} outputs a matrix $\widehat{\Sigma} \in \mathbb{R}^{d \times d}$ and a vector $\widehat{\mu} \in \mathbb{R}^d$ such that $d_{\text{TV}}(\mathcal{N}(\mu, \Sigma), \mathcal{N}(\widehat{\mu}, \widehat{\Sigma})) \leq \alpha$. Here, the $\widetilde{\Omega}$ hides $\text{polylog}(d, 1/\alpha, 1/\beta, 1/\varepsilon, \ln(1/\delta))$.

10 Efficient and robust learning of the covariance matrix

From the previous sections we know that the spectral dist function forms a convex semimetric space for positive definite matrices. Furthermore, we already have a masking mechanism for covariance matrices (see [Lemma 9.1](#)) which is shown to be concentrated (see [Lemma 9.2](#)). In this section we show how one can replace the non-robust subroutines of [Section 9](#)—namely [Lemma 9.3](#) and [Theorem 9.5](#)—with robust ones to get a robust and efficient learning method for Gaussians.

Following [\[10, 28\]](#), we use the notion of α -corrupted sample from a distribution \mathcal{D} , where an adversary receives an i.i.d. sample from \mathcal{D} , alters at most an α -fraction of them arbitrarily, and outputs the corrupted sample. The following result is based on [Theorem 4.35](#) in [\[10\]](#) and is our main tool for robust learning of the covariance matrix.

Theorem 10.1 ([10]). *There are constants $C_4 > 0, 0 < \alpha_0 < 1/2$ such that the following holds. There exist an efficient algorithm that receives $0 < \alpha < \alpha_0, \beta > 0$, and X_1, \dots, X_m as input, and outputs $\widehat{\Sigma}$ with the following guarantee: if X_1, \dots, X_m are α -corrupted samples from $\mathcal{N}(0, \Sigma)$ for some $\Sigma \in \mathbb{S}_{>0}^d$, and if $m = \widetilde{\Omega}\left(\frac{d^2 \log^5 1/\beta}{\alpha^2}\right)$, then $\left\| \Sigma^{1/2} \widehat{\Sigma}^{-1} \Sigma^{1/2} - I \right\|_F \leq C_4 \alpha \ln(1/\alpha) < 1/9$.*

Our plan is use a very similar algorithm to Algorithm 3. The key differences are that we use Theorem 10.1 for \mathcal{A} and a smaller value for η . The proof of the following proposition is very similar to that of Lemma 9.2 and is omitted for brevity.

Proposition 10.2. *Let $\Sigma \succ 0$ and $\alpha \in (0, 1)$. Set $\eta = \frac{\alpha}{C_1(d + \sqrt{d \ln(4/\beta)})}$ for a sufficiently large (universal) constant $C_1 > 0$. Then the masking mechanism $\mathcal{B}(\Sigma)$ defined in Lemma 9.1 is $(\alpha/\sqrt{d}, \beta/2)$ -concentrated.*

Algorithm 4 A private and robust algorithm for learning the covariance matrix

Input: Dataset $D = (X_1, \dots, X_m)$; parameters $\varepsilon, \delta, \alpha, \beta \in (0, 1)$

- ▷ Some parameter settings.
 - 1: $\phi \leftarrow 1, t \leftarrow 3/2, r \leftarrow 1$
 - 2: $\eta \leftarrow \frac{\alpha}{C_1(d + \sqrt{d \ln(4/\beta)})}$ ▷ C_1 from Proposition 10.2
 - 3: $\gamma \leftarrow \min \left\{ \sqrt{\frac{\varepsilon}{2d(d+1/\eta^2)}}, \frac{\varepsilon}{8d\sqrt{\ln(2/\delta)}}, \frac{\varepsilon}{8\ln(2/\delta)}, \frac{\varepsilon\eta}{12\sqrt{d}\sqrt{\ln(2/\delta)}} \right\}$.
 - 4: $k \leftarrow \max \left\{ \frac{400(r+\phi)}{\gamma}, \frac{20}{\varepsilon} \ln \left(1 + \frac{e^\varepsilon - 1}{2\delta} \right) \right\}$
 - 5: **function** $\mathcal{A}(X_1, \dots, X_s, \alpha, \beta)$
 - 6: $\widehat{\Sigma} = \text{LearnCovariance}(\alpha, \beta/k, X_1, \dots, X_s)$ ▷ Algorithm from [10]
 - 7: **Return** $\widehat{\Sigma}$.
 - 8: **end function**
 - 9: **function** $\mathcal{B}(\widehat{\Sigma})$
 - 10: Let G be $d \times d$ matrix with independent $\mathcal{N}(0, 1)$ entries.
 - 11: **Return** $\widehat{\Sigma}^{1/2}(I + \eta G)(I + \eta G)^\top \widehat{\Sigma}^{1/2}$.
 - 12: **end function**
-

Theorem 10.3. *Applying Lemma 7.6 with parameters specified in Algorithm 4 gives a mechanism \mathcal{M} satisfying the following. For all $m, d \in \mathbb{N}, \varepsilon, \delta, \alpha, \beta \in (0, 1)$, and dataset $D = (X_1, \dots, X_m)$:*

1. *Given $\varepsilon, \delta, \beta, \alpha, D$ as input, \mathcal{M} is $(2\varepsilon, 3e^\varepsilon \delta)$ -DP and runs in $\text{poly}(m, d)$ time.*
2. *If D is an α -corrupted i.i.d. sample of size m from $\mathcal{N}(0, \Sigma)$, for $\Sigma \succ 0, \alpha$ is sufficiently small, and $m = \widetilde{\Omega}\left(\frac{d^{3.5} \ln^{5.5}(1/\beta) \ln(1/\delta)}{\varepsilon \alpha^3}\right)$, then with probability at least $1 - \beta$, \mathcal{M} outputs a matrix $\widehat{\Sigma}$ such that $\left\| \widehat{\Sigma}^{-1/2} \Sigma \widehat{\Sigma}^{-1/2} - I_d \right\|_F = O(\alpha \ln(1/\alpha))$.*

Proof. The privacy claim follows from the exact same argument as in the proof of Theorem 9.4 (still we have $400(r + \phi)/k \leq \gamma$). Also, the time complexity of the algorithm remains polynomial since the non-private robust covariance learning method runs in polynomial time.

We now prove utility although the proof does not directly make use of the utility statement in [Lemma 7.6](#). For $i \in [k]$, let $\widehat{\Sigma}_i$ be the candidate covariance matrix computed by \mathcal{A} with inputs $\{X_\ell\}_{\ell=(i-1)s+1}^{is}$. Provided that $s = \widetilde{\Omega}\left(\frac{d^2 \log^5(k/\beta)}{\alpha^2}\right)$, we have that $\|\Sigma^{-1/2}\widehat{\Sigma}_i\Sigma^{-1/2} - I_d\|_F \leq C_4\alpha \ln(1/\alpha) < 1/9$ for $i \in [k]$ with probability $1 - \beta/2$. Note that a sufficient condition for this is that $m \geq k \cdot s = \widetilde{\Omega}\left(k \cdot \frac{d^2 \log^5(k/\beta)}{\alpha^2}\right) = \widetilde{\Omega}\left(\frac{d^{3.5} \ln(1/\delta) \ln(1/\beta)^{5.5}}{\varepsilon \alpha^3}\right)$. Next, using the trivial fact that Frobenius norm is an upper bound on spectral norm, we also have $\|\Sigma^{-1/2}\widehat{\Sigma}_i\Sigma^{-1/2} - I_d\| < 1/9$. By [Lemma A.6](#), we also have $\|\widehat{\Sigma}_i^{-1/2}\Sigma^{1/2}\widehat{\Sigma}_i^{-1/2} - I_d\| < 4/9$. Thus $\text{dist}(\Sigma, \widehat{\Sigma}_i) < 4/9$, where dist is as defined in [Eq. \(9.1\)](#). Now, using the fact that $\text{dist}(\cdot, \cdot)$ satisfies a 3/2-approximate triangle inequality, we have $\text{dist}(\widehat{\Sigma}_i, \widehat{\Sigma}_j) < 2/3$ for all $i, j \in [k]$. In particular, since k is sufficiently large, the algorithm does not fail with probability 1 (conditioned on the event that $\text{dist}(\widehat{\Sigma}_j, \widehat{\Sigma}_j) < 2/3$ for all $i, j \in [k]$).

Now, let $\widehat{\Sigma}_\mu$ be the weighted average computed in [Line 9](#) of [Algorithm 1](#). Note that if $\text{dist}(\widehat{\Sigma}_i, \widehat{\Sigma}_j) < 2/3$ for all $i, j \in [k]$ then $\widehat{\Sigma}_\mu = \frac{1}{k} \sum_{i \in [k]} \widehat{\Sigma}_i$. It is easy to check that since $\|\Sigma^{-1/2}\widehat{\Sigma}_i\Sigma^{-1/2} - I_d\|_F = O(\alpha \ln(1/\alpha))$, we also have $\|\Sigma^{-1/2}\widehat{\Sigma}_\mu\Sigma^{-1/2} - I_d\|_F = O(\alpha \ln(1/\alpha))$.

Next, note that by our choice of η in [Algorithm 4](#), [Proposition 10.2](#) shows that \mathcal{B} is $(\alpha/\sqrt{d}, \beta/2)$ -concentrated. In other words, if $\widehat{\Sigma} = \mathcal{B}(\widehat{\Sigma}_\mu)$ then $\|\widehat{\Sigma}^{-1/2}\widehat{\Sigma}_\mu\widehat{\Sigma}^{-1/2} - I_d\| = O(\alpha/\sqrt{d})$. In particular, $\|\widehat{\Sigma}^{-1/2}\widehat{\Sigma}_\mu\widehat{\Sigma}^{-1/2} - I_d\|_F = O(\alpha)$.

We are now ready to complete the proof. We have that $d_{\text{TV}}(\mathcal{N}(0, \Sigma), \mathcal{N}(0, \widehat{\Sigma}_\mu)) \leq O(\alpha \ln(1/\alpha))$ and $d_{\text{TV}}(\mathcal{N}(0, \widehat{\Sigma}_\mu), \mathcal{N}(0, \widehat{\Sigma})) \leq O(\alpha)$ by [Fact 6.1](#) and [Lemma 6.2](#). By the triangle inequality, we conclude that $d_{\text{TV}}(\mathcal{N}(0, \Sigma), \mathcal{N}(0, \widehat{\Sigma})) \leq O(\alpha \ln(1/\alpha))$. \square

11 Efficient and robust mean estimation

As another application of our general framework, we show how to efficiently and robustly learn the mean of a Gaussian distribution with identity covariance matrix. The following proposition indicates that every normed vector space induces a convex semimetric space, which means the Euclidean space is a convex semimetric space too.

Proposition 11.1. *Every normed vector space equipped with its induced metric forms a convex semimetric space. In particular, it satisfies triangle inequality ($r = \infty, t = 1$) and 0-locality.*

Proof. The first four properties of [Definition 7.1](#) hold for norm-induced metrics by standard arguments. Locality holds because

$$\left\| \sum_{i \in [k]} \alpha_i Y_i - \sum_{i \in [k]} \alpha'_i Y_i \right\| = \left\| \sum_{i \in [k]} (\alpha_i - \alpha'_i)(Y_i - Y_1) \right\| \leq \sum_{i \in [k]} |\alpha_i - \alpha'_i| \|Y_i - Y_1\| \leq \sum_{i \in [k]} |\alpha_i - \alpha'_i| \max_{i,j} \|Y_i - Y_j\|,$$

where in the first equality we used that $\sum_{i \in [k]} \alpha_i = \sum_{i \in [k]} \alpha'_i = 1$. \square

The Gaussian mechanism is known to be a private mechanism for functions with low ℓ_2 -sensitivity. The following theorem presents this observation in the terminology of the masking mechanisms.

Lemma 11.2. *For $\mu \in \mathbb{R}^d$, define $\mathcal{B}(\mu) = \mu + \eta X$ where $X \sim \mathcal{N}(0, I)$ and $\eta = \frac{2\gamma \ln(1.25/\delta)}{\varepsilon}$. Then \mathcal{B} is a $(\gamma, \varepsilon, \delta)$ -masking mechanism. Moreover, for all $\beta > 0$, \mathcal{B} is $(\eta\sqrt{2d \ln(2/\beta)}, \beta/2)$ concentrated.*

Proof. The privacy result is standard and is identical to that of the Gaussian mechanism; see Theorem A.1. in [17] for details. The concentration result is also standard, and follows directly from the Hoeffding's inequality (see [37, Proposition 2.5]). \square

Now we have the ingredients to reduce the problem of privately learning the mean to the non-private one. The following result is about the polynomial learnability of the mean in the robust (non-private) setting and is a direct consequence of Theorem 1.1 in [12].

Theorem 11.3 ([12]). *There an absolute constant C_3 and an algorithm that receives $0 < \alpha < 1/2$, $\beta > 0$, and X_1, \dots, X_m as input, runs in $\text{poly}(m, d)$, and outputs $\hat{\mu}$ with the following guarantee: if X_1, \dots, X_m are α -corrupted samples from $\mathcal{N}(\mu, \Sigma)$ for some $(1/2)I_d \preceq \Sigma \preceq 2I_d$, and if $m = \Omega\left(\frac{d + \log 1/\beta}{\alpha^2}\right)$, then $\|\mu - \hat{\mu}\|_2 \leq C_3 \alpha \sqrt{\log(1/\alpha)}$.*

Finally, we are ready to apply the Lemma 7.6. Algorithm 5 summarizes the parameters that we use for this reduction.

Algorithm 5 A private and robust algorithm for learning the mean

Input: Dataset $D = (X_1, \dots, X_m)$; parameters $\varepsilon, \delta, \alpha, \beta \in (0, 1)$

▷ Some parameter settings.

- 1: $t \leftarrow 1, \phi \leftarrow 0$ ▷ See Proposition 11.1.
- 2: $r \leftarrow C_3 \alpha \log(1/\alpha)$ ▷ C_3 is from Theorem 11.3.
- 3: $\eta \leftarrow \frac{\alpha}{\sqrt{2d \ln(2/\beta)}}$
- 4: $\gamma \leftarrow \frac{\eta \varepsilon}{2 \ln(1.25/\delta)}$
- 5: $k \leftarrow \max \left\{ \frac{400r}{\gamma}, \frac{20}{\varepsilon} \ln \left(1 + \frac{e^\varepsilon - 1}{2\delta} \right) \right\}$
- 6: **function** $\mathcal{A}(X_1, \dots, X_s, \alpha, \beta)$
- 7: $\hat{\mu} = \text{QUEScoreFilter}(\alpha, \beta/k, X_1, \dots, X_s)$ ▷ Algorithm from [12]
- 8: **Return** $\hat{\mu}$.
- 9: **end function**
- 10: **function** $\mathcal{B}(\hat{\mu})$
- 11: Let $X \sim \mathcal{N}(0, I_d)$
- 12: **Return** $\hat{\mu} + \eta X$.
- 13: **end function**

Theorem 11.4. *Applying Lemma 7.6 with parameters specified in Algorithm 5 gives a mechanism \mathcal{M} satisfying the following. For all $m, d \in \mathbb{N}$, $\varepsilon, \delta, \alpha, \beta \in (0, 1)$, and dataset $D = (X_1, \dots, X_m)$:*

1. Given $\varepsilon, \delta, \beta, \alpha, D$ as input, \mathcal{M} is $(2\varepsilon, 3e^\varepsilon \delta)$ -DP and runs in $\text{poly}(m, d)$ time.

2. If D is an α -corrupted i.i.d. sample of size m from $\mathcal{N}(\mu, \Sigma)$, for $(1/2)I \preceq \Sigma \preceq 2I$, $\alpha \in (0, 0.5)$, and $m = \tilde{\Omega}\left(\frac{d^{3/2} \ln(1/\delta)}{\varepsilon \alpha^2}\right)$, then with probability at least $1 - \beta$, \mathcal{M} outputs a vector $\tilde{\mu}$ such that $\|\tilde{\mu} - \mu\|_2 = O(\alpha \sqrt{\ln(1/\alpha)})$.

Proof. Proposition 11.1 indicates that $(\mathbb{R}^d, \|\cdot\|_2)$ form a convex semimetric space with non-restricted triangle inequality and 0-locality. We take $r = C_3 \alpha \log(1/\alpha) < \infty$. By Lemma 11.2, we know that \mathcal{B} is a $(400r/\gamma, \varepsilon, \delta)$ -masking mechanism. Therefore, the privacy of the algorithm follows from Lemma 7.6. Also, the time complexity of the algorithm remains polynomial since the non-private robust mean estimation runs in polynomial time.

Lemma 11.2 shows that \mathcal{B} is $(\eta \sqrt{2d \ln(2/\beta)}, \beta/2)$ -concentrated. By our choice of η in Algorithm 5, we conclude that \mathcal{B} is $(\alpha, \beta/2)$ -concentrated.

Let $\mu_i = \mathcal{A}(\{X_\ell\}_{\ell=(i-1)s+1}^{is}, \alpha, \beta)$ be the outputs of the non-private mean estimation method. Since $m = \tilde{\Omega}\left(k \cdot \frac{d + \log(k/\beta)}{\alpha^2}\right)$, we conclude that $\|\mu_i - \mu\|_2 = O(\alpha \log(1/\alpha))$ for all $i \in [k]$ with probability $1 - \beta/2$ by Theorem 11.3 and a union bound over k .

Finally, setting $Y_i = \mu_i$, $Y^* = \mu$, and $\hat{Y} = \tilde{\mu}$ for Algorithm 1, with the modules specified in Algorithm 4. Then, taking $\alpha_1 = \alpha_2 = O(\alpha \log(1/\alpha))$ in Lemma 7.6 gives that $\text{dist}(\tilde{\mu}, \mu) = O(\alpha \log(1/\alpha))$ with probability $1 - \beta$. Moreover, it is enough for the number of samples to be $m = \tilde{\Omega}\left(k \cdot \frac{d + \log(k/\beta)}{\alpha^2}\right) = \tilde{\Omega}\left(\frac{d^{3/2} \ln(1/\delta)}{\varepsilon \alpha^2}\right)$, which concludes the proof. \square

Finally, we can combine the mean estimation approach of Theorem 11.4 with the covariance estimation of Theorem 10.3 to have a complete private algorithm for robust learning of Gaussians. The sample complexity bottleneck will be that of covariance estimation. The proof of the following corollary is very similar to that of Subsection 9.1 and is omitted for brevity.

Corollary 11.5. *There exist an (ε, δ) -DP algorithm that receives $0 < \alpha < 1/2$, $\beta > 0$, and X_1, \dots, X_m as input, and outputs $\hat{\Sigma}, \hat{\mu}$ with the following guarantee: if X_1, \dots, X_m are α -corrupted samples from $\mathcal{N}(\mu, \Sigma)$ for $\Sigma \in \mathbb{S}_{>0}^d$, $\mu \in \mathbb{R}^d$ then $d_{TV}(\mathcal{N}(\mu, \Sigma), \mathcal{N}(\hat{\mu}, \hat{\Sigma})) = O(\alpha \log(1/\alpha))$ as long as $m = \Omega\left(\frac{d^{4.5} \ln(1/\delta) \ln^5(d/\varepsilon \alpha \beta)}{\varepsilon \alpha^3}\right)$.*

Remark 11.6. *Unlike Theorem 9.7 to for the non-robust case, here the utility guarantee is conditioned on the assumption that $\Sigma \succ 0$. In order to address this, one needs to come up with either a variant of the robust covariance estimation approach of [10] which works for singular covariance matrices, or an efficient and robust subspace recovery method which can be a hard task [20].*

Remark 11.7. *The sample complexities of Theorem 10.3 (and therefore Corollary 11.5) as well as Theorem 11.4 are not optimal. It seems possible to improve these sample complexities by a more careful analysis. We leave this as a future research direction.*

References

- [1] Ishaq Aden-Ali, Hassan Ashtiani, and Gautam Kamath. On the sample complexity of privately learning unbounded high-dimensional gaussians. In *Algorithmic Learning Theory*, pages 185–216. PMLR, 2021.

- [2] Ishaq Aden-Ali, Hassan Ashtiani, and Christopher Liaw. Privately learning mixtures of axis-aligned gaussians. *Advances in Neural and Information Processing Systems*, 2021.
- [3] Hassan Ashtiani, Shai Ben-David, Nicholas JA Harvey, Christopher Liaw, Abbas Mehrabian, and Yaniv Plan. Near-optimal sample complexity bounds for robust learning of gaussian mixtures via compression schemes. *Journal of the ACM (JACM)*, 67(6):1–42, 2020.
- [4] Sourav Biswas, Yihe Dong, Gautam Kamath, and Jonathan Ullman. Coinpress: Practical private mean and covariance estimation. *Advances in Neural and Information Processing Systems*, 2020.
- [5] Gavin Brown, Marco Gaboardi, Adam Smith, Jonathan Ullman, and Lydia Zakynthinou. Covariance-aware private mean estimation without private covariance estimation. *arXiv preprint arXiv:2106.13329*, 2021.
- [6] Mark Bun, Gautam Kamath, Thomas Steinke, and Steven Z Wu. Private hypothesis selection. *Advances in Neural Information Processing Systems*, 32:156–167, 2019.
- [7] Mark Bun, Kobbi Nissim, and Uri Stemmer. Simultaneous private learning of multiple concepts. *Journal of Machine Learning Research*, 20(94):1–34, 2019.
- [8] Mark Bun and Thomas Steinke. Concentrated differential privacy: Simplifications, extensions, and lower bounds. In *Theory of Cryptography Conference*, pages 635–658. Springer, 2016.
- [9] Kamalika Chaudhuri, Anand D Sarwate, and Kaushik Sinha. A near-optimal algorithm for differentially-private principal components. *Journal of Machine Learning Research*, 14, 2013.
- [10] Ilias Diakonikolas, Gautam Kamath, Daniel Kane, Jerry Li, Ankur Moitra, and Alistair Stewart. Robust estimators in high-dimensions without the computational intractability. *SIAM Journal on Computing*, 48(2):742–864, 2019.
- [11] Ilias Diakonikolas, Gautam Kamath, Daniel M Kane, Jerry Li, Ankur Moitra, and Alistair Stewart. Robust estimators in high dimensions without the computational intractability. In *2016 IEEE 57th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 655–664. IEEE, 2016.
- [12] Yihe Dong, Samuel Hopkins, and Jerry Li. Quantum entropy scoring for fast robust mean estimation and improved outlier detection. *Advances in Neural Information Processing Systems*, 32:6067–6077, 2019.
- [13] Wenxin Du, Canyon Foot, Monica Moniot, Andrew Bray, and Adam Groce. Differentially private confidence intervals. *arXiv preprint arXiv:2001.02285*, 2020.
- [14] Cynthia Dwork, Krishnaram Kenthapadi, Frank McSherry, Ilya Mironov, and Moni Naor. Our data, ourselves: Privacy via distributed noise generation. In *Annual International Conference on the Theory and Applications of Cryptographic Techniques*, pages 486–503. Springer, 2006.
- [15] Cynthia Dwork and Jing Lei. Differential privacy and robust statistics. In *Proceedings of the forty-first annual ACM symposium on Theory of computing*, pages 371–380, 2009.

- [16] Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam Smith. Calibrating noise to sensitivity in private data analysis. *Journal of Privacy and Confidentiality*, 7(3):17–51, 2017.
- [17] Cynthia Dwork and Aaron Roth. The algorithmic foundations of differential privacy. *Found. Trends Theor. Comput. Sci.*, 9(3-4):211–407, 2014.
- [18] Cynthia Dwork and Guy N Rothblum. Concentrated differential privacy. *arXiv preprint arXiv:1603.01887*, 2016.
- [19] Quan Geng, Wei Ding, Ruiqi Guo, and Sanjiv Kumar. Tight analysis of privacy and utility tradeoff in approximate differential privacy. *Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics*, 108:89–99, 2020.
- [20] Moritz Hardt and Ankur Moitra. Algorithms and hardness for robust subspace recovery. In *Conference on Learning Theory*, pages 354–375. PMLR, 2013.
- [21] Gautam Kamath, Jerry Li, Vikrant Singhal, and Jonathan Ullman. Privately learning high-dimensional distributions. In *Conference on Learning Theory*, pages 1853–1902. PMLR, 2019.
- [22] Gautam Kamath, Argyris Mouzakis, Vikrant Singhal, Thomas Steinke, and Jonathan Ullman. A private and computationally-efficient estimator for unbounded gaussians. *arXiv preprint arXiv:2111.04609*, 2021.
- [23] Gautam Kamath, Or Sheffet, Vikrant Singhal, and Jonathan Ullman. Differentially private algorithms for learning mixtures of separated gaussians. In *2020 Information Theory and Applications Workshop (ITA)*, pages 1–62. IEEE, 2020.
- [24] Gautam Kamath, Vikrant Singhal, and Jonathan Ullman. Private mean estimation of heavy-tailed distributions. In *Conference on Learning Theory*, pages 2204–2235. PMLR, 2020.
- [25] Gautam Kamath and Jonathan Ullman. A primer on private statistics. *arXiv preprint arXiv:2005.00010*, 2020.
- [26] Vishesh Karwa and Salil Vadhan. Finite sample differentially private confidence intervals. In *9th Innovations in Theoretical Computer Science Conference (ITCS 2018)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2018.
- [27] Solomon Kullback. *Information Theory and Statistics*. Dover, 1968.
- [28] Kevin A Lai, Anup B Rao, and Santosh Vempala. Agnostic estimation of mean and covariance. In *2016 IEEE 57th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 665–674. IEEE, 2016.
- [29] B. Laurent and P. Massart. Adaptive estimation of a quadratic functional by model selection. *The Annals of Statistics*, 28(5):1302–1338, 2000.
- [30] Xiyang Liu, Weihao Kong, Sham Kakade, and Sewoong Oh. Robust and differentially private mean estimation. *arXiv preprint arXiv:2102.09159*, 2021.
- [31] Kobbi Nissim, Sofya Raskhodnikova, and Adam Smith. Smooth sensitivity and sampling in private data analysis. In *Proceedings of the thirty-ninth annual ACM symposium on Theory of computing*, pages 75–84, 2007.

- [32] Carl Edward Rasmussen and Christopher K. I. Williams. *Gaussian Processes for Machine Learning*. The MIT Press, 2006.
- [33] Vikrant Singhal and Thomas Steinke. Privately learning subspaces. *arXiv preprint arXiv:2106.00001*, 2021.
- [34] Alexandre B Tsybakov. *Introduction to Nonparametric Estimation*. Springer, 2009.
- [35] Salil Vadhan. The complexity of differential privacy. In *Tutorials on the Foundations of Cryptography*, pages 347–450. Springer, 2017.
- [36] Roman Vershynin. *High-Dimensional Probability: An Introduction with Applications in Data Science*. Cambridge University Press, 2018.
- [37] Martin J. Wainwright. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press, 2019.
- [38] Lu Wei, Anand D Sarwate, Jukka Corander, Alfred Hero, and Vahid Tarokh. Analysis of a privacy-preserving pca algorithm using random matrix theory. In *2016 IEEE Global Conference on Signal and Information Processing (GlobalSIP)*, pages 1335–1339. IEEE, 2016.

A Standard Facts

Claim A.1. *If $x \leq 1/2$ then $\frac{1+x}{1-x} \leq 1 + 4x$.*

Proof. Let $f(x) = 1 + 4x - \frac{1+x}{1-x}$. Then $f''(x) = -\frac{2(1+x)}{(1-x)^3} - \frac{2}{(1-x)^2}$. So f is concave on the interval $[0, 1)$. Since $f(0) = f(1/2) = 0$, we have that $f(x) \geq 0$ for $x \in [0, 1/2]$ as desired. \square

Claim A.2. *For all $x \in \mathbb{R}$, $\frac{1-x}{1+x} \geq 1 - 2x$.*

Proof. We have $\frac{1-x}{1+x} \geq (1-x)^2 = 1 - 2x + x^2 \geq 1 - 2x$. \square

Lemma A.3 ([29, Lemma 1]). *Let g_1, \dots, g_k be i.i.d. $\mathcal{N}(0, 1)$ random variables. Then*

$$\mathbb{P} \left[\sum_{i=1}^k g_i^2 - k \geq 2\sqrt{kt} + 2t \right] \leq e^{-t}.$$

Lemma A.4 ([36, Exercise 4.7.3]). *There is an absolute constant $C > 0$ such that the following holds. Let $\Sigma \succ 0$ and $\beta > 0$. Let $X_1, \dots, X_m \sim \mathcal{N}(0, \Sigma)$ and $\widehat{\Sigma} = \frac{1}{m} \sum_{i=1}^m X_i X_i^\top$. Then $\|\Sigma^{-1/2} \widehat{\Sigma} \Sigma^{-1/2} - I_d\| \leq C \left(\sqrt{\frac{d + \ln(2/\beta)}{m}} + \frac{d + \ln(2/\beta)}{m} \right)$ with probability $1 - \beta$.*

Theorem A.5 ([36, Theorem 4.4.5]). *There is a universal constant $C > 0$ such that the following holds. Let G be a $d \times d$ random matrix where each entry is an independent $\mathcal{N}(0, 1)$ random variable. Then $\|G\| \leq C(\sqrt{d} + \sqrt{\ln(2/\beta)})$ with probability $1 - \beta$.*

Lemma A.6. Let A, B be $d \times d$ matrices and suppose that $\|A^{-1/2}BA^{-1/2} - I_d\| \leq \gamma \leq 1/2$. Then $\|B^{-1/2}AB^{-1/2} - I_d\| \leq 4\gamma$.

Proof. The assumption that $\|A^{-1/2}BA^{-1/2} - I_d\| \leq \gamma$ means that $(1-\gamma)A \preceq B \preceq (1+\gamma)A$. Hence, we also have that $\frac{1}{1+\gamma}B \preceq A \preceq \frac{1}{1-\gamma}B$. The asserted inequality is then proved using the bounds in Claim A.7. \square

Claim A.7. Suppose that $\gamma \in [0, 1/2]$. Then $\frac{1}{1+\gamma} \geq 1 - 4\gamma$ and $\frac{1}{1-\gamma} \leq 1 + 4\gamma$.

Proof. To prove the first inequality, let $f(\gamma) = 1 - 4\gamma - \frac{1}{1+\gamma}$. We need to show that $f(\gamma) \geq 0$ for all $\gamma \in [0, 1/2]$. Differentiating, we have $f'(\gamma) = 4 - \frac{1}{1+\gamma^2}$ so f is increasing. Since $f(0) = 0$, the first inequality follows.

For the second inequality, let $f(\gamma) = 1 + 4\gamma - \frac{1}{1-\gamma}$. The derivative $f'(\gamma) = 4 - \frac{1}{(1-\gamma)^2} \geq 0$ for $\gamma \in [0, 1/2]$. So f is increasing on $[0, 1/2]$ and $f(0) = 0$ so the second inequality is proved. \square

B Missing Proofs From Section 6

B.1 Proof of Fact 6.1

Proof of Fact 6.1. The first assertion is a well-known formula for the KL divergence between two normal distributions (see [27, §9.1] or [32, Eq. A.23]).

Next, let $\lambda_1, \dots, \lambda_d$ be the eigenvalues of $\Sigma_2^{-1}\Sigma_1$. Then

$$\text{tr}(\Sigma_2^{-1}\Sigma_1 - I) - \ln \det(\Sigma_2^{-1}\Sigma_1) = \sum_{i=1}^d \lambda_i - 1 - \ln \lambda_i \leq \sum_{i=1}^d (\lambda_i - 1)^2 = \|\Sigma_2^{-1}\Sigma_1 - I\|_F^2,$$

where the inequality follows from Claim B.1. The lemma now follows because the two matrices $\Sigma_2^{-1}\Sigma_1$ and $\Sigma_2^{-1/2}\Sigma_1\Sigma_2^{-1/2}$ have the same spectrum. \square

Claim B.1. Suppose that $x \geq 1/2$. Then $x - 1 - \ln(x) \leq (x - 1)^2$.

Proof. Let $f(x) = (x - 1)^2 - (x - 1) + \ln(x)$. Then $f'(x) = 2(x - 1) - 1 + \frac{1}{x} = \frac{2x^2 - 3x + 1}{x} = \frac{(2x-1)(x-1)}{x}$. So $f'(x) \leq 0$ on $[1/2, 1]$ and $f'(x) \geq 0$ on $[1, \infty)$. Hence, on the interval $[1/2, \infty)$, $f(x)$ is minimized at $x = 1$ and $f(x) \geq f(1) = 0$ for all $x \in [1/2, \infty)$. \square

Claim B.2. Suppose $x \leq 1/2$. Then $\sqrt{1-x} \geq 1 - x/2 - x^2/2$.

Proof. Let $f(x) = \sqrt{1-x} - 1 + x/2 + x^2/2$. Then $f'(x) = x - \frac{1}{2\sqrt{1-x}} + \frac{1}{2}$ and $f''(x) = 1 - \frac{1}{4(1-x)^{3/2}}$. Note that $f'(0) = 0$ so $x = 0$ is a critical point of f . Moreover, $f''(x) \geq 0$ for all $x \leq 1/2$ so f is convex on $(-\infty, 1/2)$. Hence, the minimizer of f on the interval $(-\infty, 1/2]$ is at $x = 0$ and $f(x) \geq f(0) = 0$ for all $x \in (-\infty, 1/2]$. \square

B.2 Proof of Lemma 6.4

Claim B.3 (see [37, Example 2.8]). *Let $X \sim \mathcal{N}(0, 1)$ and $|u| \leq 1/4$. Then $\mathbb{E} \left[e^{u(X^2-1)} \right] \leq e^{2u^2}$.*

We first prove Lemma 6.4 for the special case where the matrix is diagonal. The general case follows easily from rotation invariance.

Lemma B.4. *Let Λ be a $d \times d$ diagonal matrix and $g \sim \mathcal{N}(0, I_d)$. Then for all $x > 0$,*

$$\mathbb{P} \left[g^\top \Lambda g \geq \text{tr}(\Lambda) + 4\|\Lambda\|_F \cdot \sqrt{x} + 4\|\Lambda\| \cdot x \right] \leq e^{-x}.$$

Proof. Note that $g^\top \Lambda g = \sum_{i=1}^d \lambda_i (g_i^2 - 1)$ where $\lambda_1, \dots, \lambda_d$ are the diagonal entries of Λ . From Claim B.3, we have $\mathbb{E} \left[e^{u\lambda_i(g_i^2-1)} \right] \leq e^{2u^2\lambda_i^2}$ provided $|u| \leq 1/4\|\Lambda\|$. Hence, $\mathbb{E} \left[e^{u(g^\top \Lambda g - \text{tr}(\Lambda))} \right] = \prod_{i=1}^d \mathbb{E} \left[e^{u\lambda_i(g_i^2-1)} \right] \leq e^{2u^2\|\Lambda\|_F^2}$. Applying Markov's Inequality, we thus have, for $0 \leq u \leq 1/4\|\Lambda\|$,

$$\mathbb{P} \left[g^\top \Lambda g - \text{tr}(\Lambda) \geq 4\|\Lambda\|_F \cdot \sqrt{x} + 4\|\Lambda\| \cdot x \right] \leq e^{-u(4\|\Lambda\|_F \cdot \sqrt{x} + 4\|\Lambda\| \cdot x) + 2u^2\|\Lambda\|_F^2}.$$

It is a straightforward calculation to verify that choosing $u = \min \left\{ \frac{1}{4\|\Lambda\|}, \frac{\sqrt{x}}{\|\Lambda\|_F} \right\}$ makes the RHS at most e^{-x} which proves the lemma. \square

Proof of Lemma 6.4. Let $A = U\Lambda U^\top$ be the eigendecomposition of A . Then $g^\top A g = g^\top U\Lambda U^\top g \stackrel{d}{=} g^\top \Lambda g$ where the last equality in distribution is by rotation invariance (Fact B.5). The claim now follows from Lemma B.4 since A and Λ have the same spectrum. \square

Fact B.5. *Let $g \sim \mathcal{N}(0, I_d)$ and U be a $d \times d$ orthogonal matrix. Then $Ug \sim \mathcal{N}(0, I_d)$.*

B.3 Proof of Lemma 6.10

Proof of Lemma 6.10. Let f_1, f_2 be the densities of $\mathcal{D}_1, \mathcal{D}_2$, respectively. Let $S \subseteq \mathbb{R}^d$ and let $T = \{y \in \mathbb{R}^d : f_1(y) \geq e^\varepsilon f_2(y)\}$. The assumption $\mathbb{P}_{Y \sim \mathcal{D}_1} [\mathcal{L}_{\mathcal{D}_1 \| \mathcal{D}_2}(Y) \geq \varepsilon] \leq \delta$ means that $\mathbb{P}_{Y \sim \mathcal{D}_1} [Y \in T] \leq \delta$. Thus, we have

$$\begin{aligned} \mathbb{P}_{Y \sim \mathcal{D}_1} [Y \in S] &= \int_S f_1(y) \, dy \\ &= \int_{S \setminus T} f_1(y) \, dy + \int_{S \cap T} f_1(y) \, dy \\ &\leq \int_{S \setminus T} e^\varepsilon f_2(y) \, dy + \delta \\ &\leq e^\varepsilon \mathbb{P}_{Y \sim \mathcal{D}_2} [S] + \delta. \end{aligned}$$

An identical argument shows that $\mathbb{P}_{Y \sim \mathcal{D}_2} [Y \in S] \leq e^\varepsilon \mathbb{P}_{Y \sim \mathcal{D}_1} [S] + \delta$, which proves that $\mathcal{D}_1, \mathcal{D}_2$ are (ε, δ) -indistinguishable. \square

C Missing proofs from Section 7

C.1 Proof of Lemma 7.2

Proof. The first two conditions in Definition 7.1 are trivial.

Triangle Inequality.

Next, we prove that dist satisfies a $(3/2)$ -approximate 1-restricted triangle inequality. Let $A, B, C \in \mathcal{S}^d$ and let $\rho_{ab} = \text{dist}(A, B)$, $\rho_{bc} = \text{dist}(B, C)$ and $\rho_{ac} = \text{dist}(A, C)$. We need to show that if $\rho_{ab}, \rho_{bc} \leq 1$ then $\rho_{ac} \leq \frac{3}{2}(\rho_{ab} + \rho_{bc})$. Indeed, we have

$$\begin{aligned} A &\preceq (1 + \rho_{ab})B \preceq (1 + \rho_{ab})(1 + \rho_{bc})C = (1 + \rho_{ab} + \rho_{bc} + \rho_{ab}\rho_{bc})C \\ &\preceq (1 + \rho_{ab} + \rho_{bc} + \rho_{ab}^2/2 + \rho_{bc}^2/2)C \preceq (1 + \frac{3}{2}(\rho_{ab} + \rho_{bc}))C, \end{aligned}$$

where the first two inequalities used that $\text{dist}(A, B) = \rho_{ab}$ and $\text{dist}(B, C) = \rho_{bc}$, respectively, the third inequality used that $xy \leq x^2/2 + y^2/2$ for any $x, y \in \mathbb{R}$ and the last inequality used that $\rho_{ab}, \rho_{bc} \leq 1$. Similarly, we can show that

$$A \succeq (1 - \frac{3}{2}(\rho_{ab} + \rho_{bc}))C,$$

which imply, with the previous quality, that $\|C^{-1/2}AC^{-1/2} - I\| \leq \frac{3}{2}(\rho_{ab} + \rho_{bc})$. We can similarly show that $\|A^{-1/2}CA^{-1/2} - I\| \leq \frac{3}{2}(\rho_{ab} + \rho_{bc})$ so $\rho_{ac} \leq \frac{3}{2}(\rho_{ab} + \rho_{bc})$, as desired.

Convexity.

For this, it would suffice to show that for all $\alpha \in [0, 1]$, $\text{dist}(\alpha A + (1 - \alpha)B, C) \leq \alpha \text{dist}(A, C) + (1 - \alpha) \text{dist}(B, C)$, and the rest follows from induction over k . Let $\rho_a = \text{dist}(A, C)$, $\rho_b = \text{dist}(B, C)$. We need to show the following four inequalities:

1. $C \preceq (1 + \alpha\rho_a + (1 - \alpha)\rho_b)(\alpha A + (1 - \alpha)B)$;
2. $C \succeq (1 - \alpha\rho_a - (1 - \alpha)\rho_b)(\alpha A + (1 - \alpha)B)$;
3. $\alpha A + (1 - \alpha)B \preceq (1 + \alpha\rho_a + (1 - \alpha)\rho_b)C$; and
4. $\alpha A + (1 - \alpha)B \succeq (1 - \alpha\rho_a - (1 - \alpha)\rho_b)C$.

The third and fourth items are easy to check. Indeed, for the third item, we have $A \preceq (1 + \rho_a)C$ and $B \preceq (1 + \rho_b)C$ so

$$\alpha A + (1 - \alpha)B \preceq \alpha(1 + \rho_a)C + (1 - \alpha)(1 + \rho_b)C = (1 + \alpha\rho_a + (1 - \alpha)\rho_b)C.$$

The fourth item follows similarly.

It remains to prove the first two inequalities. Here, we prove only the first inequality since the second follows by an analogous argument. Let

$$K = \left(\frac{\alpha(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_a} + \frac{(1 - \alpha)(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_b} \right)^{-1}$$

and set

$$w = K \frac{\alpha(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_a}.$$

Our choice of K means that

$$1 - w = K \frac{(1 - \alpha)(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_b}.$$

Note that $w \in [0, 1]$. Since $C \preceq (1 + \rho_a)A$ and $C \preceq (1 + \rho_b)B$, we have

$$\begin{aligned} C &= wC + (1 - w)C \\ &\preceq w(1 + \rho_a)C + (1 - w)(1 + \rho_b)B \\ &= K [\alpha(1 + \alpha\rho_a + (1 - \alpha)\rho_b)A + (1 - \alpha)(1 + \alpha\rho_a + (1 - \alpha)\rho_b)B] \\ &= K [(1 + \alpha\rho_a + (1 - \alpha)\rho_b)(\alpha A + (1 - \alpha)B)]. \end{aligned}$$

Claim C.1 shows that $K \leq 1$ so we have the desired inequality.

Claim C.1. *Let $\alpha \in [0, 1]$ and let $\rho_a, \rho_b \in \mathbb{R}_{\geq 0}$. Then*

$$\frac{\alpha(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_a} + \frac{(1 - \alpha)(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_b} \geq 1.$$

Proof. Let

$$f(\alpha) := \frac{\alpha(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_a} + \frac{(1 - \alpha)(1 + \alpha\rho_a + (1 - \alpha)\rho_b)}{1 + \rho_b}.$$

Observe that $f(0) = f(1) = 1$. Thus, it suffices to prove that f is concave. Indeed, we have

$$f'(\alpha) = \frac{1 + \alpha\rho_a + (1 - \alpha)\rho_b}{1 + \rho_a} + \frac{\alpha(\rho_a - \rho_b)}{1 + \rho_a} - \frac{1 + \alpha\rho_a + (1 - \alpha)\rho_b}{1 + \rho_b} + \frac{(1 - \alpha)(\rho_a - \rho_b)}{1 + \rho_b}.$$

and

$$f''(\alpha) = 2\frac{\rho_a - \rho_b}{1 + \rho_a} + 2\frac{\rho_b - \rho_a}{1 + \rho_b} = 2(\rho_a - \rho_b) \cdot \left(\frac{1}{1 + \rho_a} - \frac{1}{1 + \rho_b} \right).$$

Finally, we check that $f'' \leq 0$. Indeed, if $\rho_a = \rho_b$ then $f'' = 0$ and if $\rho_a \neq \rho_b$ then $\rho_a - \rho_b$ and $1/(1 + \rho_a) - 1/(1 + \rho_b)$ have opposite signs so $f'' < 0$. \square

Locality.

Let $\widehat{\Sigma} = \sum_{i \in [k]} w_i \widehat{\Sigma}_i$ and $\widehat{\Sigma}' = \sum_{i \in [k]} w'_i \widehat{\Sigma}_i$. Let $\rho = \max_{i, j \in [k]} \text{dist}(\widehat{\Sigma}_i, \widehat{\Sigma}_j)$. Then

$$\begin{aligned} \sum_{i \in [k]} w_i \widehat{\Sigma}_i &\preceq \sum_{i \in [k]} w'_i \widehat{\Sigma}_i + \sum_{i \in [k]} |w_i - w'_i| \widehat{\Sigma}_i \\ &\preceq \widehat{\Sigma}' + \sum_{i \in [k]} |w_i - w'_i| (1 + \rho) \widehat{\Sigma}' \\ &= \left(1 + (1 + \rho) \sum_{i \in [k]} |w_i - w'_i| \right) \widehat{\Sigma}' \end{aligned}$$

For the second inequality, we used $\widehat{\Sigma}_i \preceq (1 + \rho) \widehat{\Sigma}_j$ for all $i, j \in [k]$, which imply $\widehat{\Sigma}_i \preceq (1 + \rho) \widehat{\Sigma}'$ for all $i \in [k]$. \square

C.2 Proof of Lemma 7.6

Proof. We first prove the utility and then the privacy result.

Utility. We first show that the algorithm will not return \perp if the assumptions of the utility guarantee hold. Based on the assumption, $\text{dist}(Y_i, Y^*) \leq \alpha_1/t$ for all $i \in [k]$. By the t -approximate r -restricted triangle inequality we have $\text{dist}(Y_i, Y_j) \leq \alpha_1 \leq r/t$ for all $i, j \in [k]$. Therefore, for all $i \in [k]$, $q_i = k/(k+3) \geq 30/33 > 0.9$. Finally, $Q = q_i > 0.9 > 0.8 + \frac{2}{k\varepsilon} \ln\left(1 + \frac{e^\varepsilon - 1}{2\delta}\right)$ and therefore the algorithm does not fail in Line 7.

Let $W = \sum_{i \in [k]} w_i$. To show that \hat{Y} concentrates around Y^* we have

$$\begin{aligned} \mathbb{P}\left[\text{dist}(\hat{Y}, Y^*) > \alpha_1 + \alpha_2\right] &\leq \mathbb{P}\left[\text{dist}(\hat{Y}, \mu) + \text{dist}(\mu, Y^*) > \alpha_1/t + \alpha_2/t\right] \leq \\ \mathbb{P}\left[\text{dist}(\hat{Y}, \mu) + \sum_{i \in [k]} \frac{w_i}{W} \text{dist}(Y_i, Y^*) > \alpha_1/t + \alpha_2/t\right] &\leq \\ \mathbb{P}\left[\text{dist}(\hat{Y}, \mu) + \alpha_1/t > \alpha_1/t + \alpha_2/t\right] &\leq \mathbb{P}\left[\text{dist}(\hat{Y}, \mu) > \alpha_2/t\right] \leq \beta \end{aligned}$$

where the first inequality follows from the t -approximate triangle inequality, and the second one follows from the convexity of the dist function and the fact that μ is a convex combination of Y_i 's. The third inequality uses the assumption that $\text{dist}(Y_i, Y^*) \leq \alpha_1/t$ for all $i \in [k]$. Finally, the last inequality follows from the assumption that the masking mechanism \mathcal{B} is $(\alpha_2/t, \beta)$ concentrated.

Privacy. Let D, D' be two neighbouring datasets and let \mathcal{M} denote the mechanism specified by Algorithm 1. Note that the statistic Q computed in Line 4 has sensitivity less than $2/k$. This is because if we modify a single datapoint, there is at most a single i such that Y_i changes. Hence, the value of q_i changes by at most 1 and for $j \neq i$, the value of q_j changes by less than $1/k$. Thus, Q has sensitivity at most $2/k$. In particular, since we apply the Truncated Laplace mechanism in Line 7 (see Theorem 6.12), we have

$$\mathbb{P}[\mathcal{M}(D) = \perp] \leq e^\varepsilon \mathbb{P}[\mathcal{M}(D') = \perp] + \delta \quad (\text{C.1})$$

We now show that for any $T \subseteq \mathcal{Y}$, we have

$$\mathbb{P}[\mathcal{M}(D) \in T] \leq e^{2\varepsilon} \mathbb{P}[\mathcal{M}(D') \in T] + 3e^\varepsilon \delta \quad \text{and} \quad (\text{C.2})$$

$$\mathbb{P}[\mathcal{M}(D) \in T \cup \{\perp\}] \leq e^{2\varepsilon} \mathbb{P}[\mathcal{M}(D') \in T \cup \{\perp\}] + 3e^\varepsilon \delta \quad (\text{C.3})$$

which establishes that Algorithm 1 is (ε, δ) -DP. To this end, we consider two different cases.

Case 1: $Q < 0.8$. In this case, $\hat{Q} < 0.8 + \frac{2}{k\varepsilon} \ln\left(1 + \frac{e^\varepsilon - 1}{2\delta}\right)$ with probability 1 so $\mathbb{P}[\mathcal{M}(D) = \perp] = 1$. Now, we verify that Eq. (C.2) and Eq. (C.3) hold. For any $T \subseteq \mathcal{Y}$, we have $\mathbb{P}[\mathcal{M}(D) \in T] = 0$ so Eq. (C.2) is trivially satisfied. To check Eq. (C.3) holds, we apply Eq. (C.1) to see that

$$\mathbb{P}[\mathcal{M}(D) \in T \cup \{\perp\}] = \mathbb{P}[\mathcal{M}(D) = \perp] \leq e^\varepsilon \mathbb{P}[\mathcal{M}(D') = \perp] + \delta \leq e^\varepsilon \mathbb{P}[\mathcal{M}(D') \in T \cup \{\perp\}] + \delta.$$

Case 2: $Q \geq 0.8$. Let Y_1, \dots, Y_k (resp. Y'_1, \dots, Y'_k) be the output of the non-private algorithm \mathcal{A} in Line 2 when the dataset is D (resp. D'). Further, define let q_i, w_i, μ be as defined in Algorithm 1 when the input is D and q'_i, w'_i, μ' be the analogous quantities when the input is D' . For notation, let $W = \sum_{i=1}^k w_i$ and $W' = \sum_{i=1}^k w'_i$. Our goal is to show that $\text{dist}(\mu, \mu') \leq 400(r + \phi)/k$ so that the $(400(r + \phi)/k, \varepsilon, \delta)$ -masking mechanism guarantees the output is (ε, δ) -indistinguishable.

We proceed by proving a series of claims. The first claim gives a lower bound on W .

Claim C.2. $W \geq k/3$.

Proof. Let $\alpha = |\{i \in [k] : q_i \geq 0.7\}|/k$. Then $0.8 \leq Q \leq \alpha + 0.7(1 - \alpha)$ so $\alpha \geq 1/3$. Finally, $W \geq \alpha k$ because if $q_i \geq 0.7$ then $w_i = 1$. We conclude that $W \geq k/3$. \square

Let us make one simple observation about the sequence Y_1, \dots, Y_k and Y'_1, \dots, Y'_k . Since only one datapoint has been changed, it must be the case that $|\{i \in [k] : Y_i \neq Y'_i\}| \leq 1$. Without loss of generality (perhaps by reindexing), we assume that $Y_i = Y'_i$ for $2 \leq i \leq k$. In addition, we have the following straightforward claim which follows from the aforementioned observation.

Claim C.3. For $2 \leq i \leq k$, we have $|w_i - w'_i| \leq 10/k$. In addition, $|W - W'| \leq 11$.

Proof. That $|w_i - w'_i| \leq 10/k$ for $2 \leq i \leq k$ follows from the fact that $Y_i = Y'_i$ for $2 \leq i \leq k$. Next, $|W - W'| \leq |w_1 - w'_1| + \sum_{2 \leq i \leq k} |w_i - w'_i| \leq 1 + (k - 1) \cdot 10/k < 11$. \square

Now, let us proceed to bound $\text{dist}(\mu, \mu')$. Using repeated applications of convexity, we have

$$\begin{aligned} \text{dist}(\mu, \mu') &= \text{dist} \left(\frac{w_1}{W} Y_1 + \sum_{i=2}^k \frac{w_i}{W} Y_i, \frac{w'_1}{W'} Y'_1 + \sum_{j=2}^k \frac{w'_j}{W'} Y_j \right) \\ &\leq \frac{w_1}{W} \text{dist} \left(Y_1, \frac{w'_1}{W'} Y'_1 + \sum_{j=2}^k \frac{w'_j}{W'} Y_j \right) \\ &\quad + \left(1 - \frac{w_1}{W}\right) \text{dist} \left(\frac{1}{1 - w_1/W} \sum_{i=2}^k \frac{w_i}{W} Y_i, \frac{w'_1}{W'} Y'_1 + \sum_{j=2}^k \frac{w'_j}{W'} Y_j \right) \\ &\leq \frac{w_1}{W} \frac{w'_1}{W'} \text{dist}(Y_1, Y'_1) \end{aligned} \tag{C.4}$$

$$+ \frac{w_1}{W} \left(1 - \frac{w'_1}{W'}\right) \text{dist} \left(Y_1, \frac{1}{1 - w'_1/W'} \sum_{j=2}^k \frac{w'_j}{W'} Y_j \right) \tag{C.5}$$

$$+ \left(1 - \frac{w_1}{W}\right) \frac{w'_1}{W'} \text{dist} \left(\frac{1}{1 - w_1/W} \sum_{i=2}^k \frac{w_i}{W} Y_i, Y'_1 \right) \tag{C.6}$$

$$+ \left(1 - \frac{w_1}{W}\right) \left(1 - \frac{w'_1}{W'}\right) \text{dist} \left(\frac{1}{1 - w_1/W} \sum_{i=2}^k \frac{w_i}{W} Y_i, \frac{1}{1 - w'_1/W'} \sum_{j=2}^k \frac{w'_j}{W'} Y_j \right), \tag{C.7}$$

where we used convexity in the first argument for the first inequality and convexity in the second argument for the second inequality. We now bound each Eq. (C.4), Eq. (C.5), Eq. (C.6), and Eq. (C.7) separately. To do so, we make use of the following claim.

Claim C.4. *Suppose that $w_i, w'_j > 0$. Then $\text{dist}(Y_i, Y'_j) \leq 2r$.*

Proof. Let $S_i = \{s \in [k] : \text{dist}(Y_i, Y_s) \leq r/t\}$ and $S'_j = \{s \in [k] : \text{dist}(Y'_j, Y'_s) \leq r/t\}$. Since $w_i, w'_j > 0$, we must have $|S_i| \geq 0.6(k+3)$ and $|S'_j| \geq 0.6(k+3)$. Hence, $|S_i \cap S'_j| = |S_i| + |S'_j| - |S_i \cup S'_j| \geq 0.2k + 3.6 \geq 2$. So there exists $s \in S_i \cap S'_j$ such that $2 \leq s \leq k$ which implies that $Y_s = Y'_s$. In particular, for this s , we have $\text{dist}(Y_i, Y_s) \leq r/t$ and $\text{dist}(Y'_j, Y_s) \leq r/t$. Recalling that dist satisfies a t -approximate r -restricted triangle inequality, we have $\text{dist}(Y_i, Y'_j) \leq t(\text{dist}(Y_i, Y_s) + \text{dist}(Y'_j, Y_s)) \leq 2r$. \square

By Claim C.4, we have that Eq. (C.4) is bounded by

$$\frac{w_1 w'_1}{W W'} \text{dist}(Y_1, Y'_1) \leq \frac{w_1 w'_1}{W W'} 2r \quad (\text{C.8})$$

We now bound Eq. (C.5) which is very similar to the bound of Eq. (C.4) we just obtained. Using convexity, we have

$$\begin{aligned} \frac{w_1}{W} \left(1 - \frac{w'_1}{W'}\right) \text{dist} \left(Y_1, \frac{1}{1 - w'_1/W'} \sum_{j=2}^k \frac{w'_j}{W'} Y_j \right) &\leq \frac{w_1}{W} \sum_{j=2}^k \frac{w'_j}{W'} \text{dist}(Y_1, Y_j) \\ &\leq \frac{w_1}{W} \sum_{j=2}^k \frac{w'_j}{W'} \cdot 2r, \end{aligned} \quad (\text{C.9})$$

where we applied Claim C.4 for the last inequality. Similarly, we have

$$\left(1 - \frac{w_1}{W}\right) \frac{w'_1}{W'} \text{dist} \left(\frac{1}{1 - w_1/W} \sum_{i=2}^k \frac{w_i}{W} Y_i, Y'_1 \right) \leq \frac{w'_1}{W'} \sum_{i=2}^k \frac{w_i}{W} 2r. \quad (\text{C.10})$$

Note that Eq. (C.8), Eq. (C.9), Eq. (C.10) shows that the sum of Eq. (C.4), Eq. (C.5), Eq. (C.6) is bounded above by $\frac{2r}{W} + \frac{2r}{W'}$.

Finally, we bound Eq. (C.7). We require the following claim whose proof is similar to that of Claim C.4.

Claim C.5. *Let $S = \{2 \leq i \leq k : \max(w_i, w'_i) > 0\}$. Then for all $i, j \in S$, we have $\text{dist}(Y_i, Y_j) \leq 2r$.*

Proof. Let $S_i = \{s \in [k] : \text{dist}(Y_i, Y_s) \leq r/t\}$ and $S_j = \{s \in [k] : \text{dist}(Y_j, Y_s) \leq r/t\}$. Since $\max(w_i, w'_i), \max(w_j, w'_j) > 0$, we must have $|S_i| \geq 0.6(k+3) - 1$ and $|S_j| \geq 0.6(k+3) - 1$. Hence, $|S_i \cap S_j| = |S_i| + |S_j| - |S_i \cup S_j| \geq 0.2k + 1.6 > 1$ so $|S_i \cap S_j| \geq 2$. So there exists $s \in S_i \cap S_j$ such that $s \geq 2$. In particular, for this s , we have $\text{dist}(Y_i, Y_s) \leq r/t$ and $\text{dist}(Y_j, Y_s) \leq r/t$. Recalling that dist satisfies a t -approximate r -restricted triangle inequality, we have $\text{dist}(Y_i, Y_j) \leq t(\text{dist}(Y_i, Y_s) + \text{dist}(Y_j, Y_s)) \leq 2r$. \square

Note that $\sum_{i=2}^k \frac{w_i}{W} Y_i = \sum_{i \in S} \frac{w_i}{W} Y_i$ and $\sum_{i=2}^k \frac{w'_i}{W'} Y_i = \sum_{i \in S} \frac{w'_i}{W'} Y_i$. Now, we have

$$\begin{aligned} & \text{dist} \left(\frac{1}{1 - w_1/W} \sum_{i \in S} \frac{w_i}{W} Y_i, \frac{1}{1 - w'_1/W'} \sum_{i \in S} \frac{w'_i}{W'} Y_i \right) \\ & \leq \left(\phi + \max_{i,j \in S} \text{dist}(Y_i, Y_j) \right) \sum_{i \in S} \left| \frac{w_i/W}{1 - w_1/W} - \frac{w'_i/W'}{1 - w'_1/W'} \right|. \end{aligned}$$

Where we used the fact that dist satisfies locality in Definition 7.1. Note that $\text{dist}(Y_i, Y_j) \leq 2r$ for all $i, j \in S$. Also, by the Claim C.6 we see that the right hand side is bounded by $100/k$. Therefore, the term in Eq. (C.7) is upper bounded by $100(2r + \phi)/k$. We now state and prove Claim C.6.

Claim C.6. $\sum_{i \in S} \left| \frac{w_i/W}{1 - w_1/W} - \frac{w'_i/W'}{1 - w'_1/W'} \right| \leq 100/k$.

Proof.

$$\begin{aligned} & \sum_{i \in S} \left| \frac{w_i/W}{1 - w_1/W} - \frac{w'_i/W'}{1 - w'_1/W'} \right| = \sum_{i \in S} \left| \frac{w_i}{W - w_1} - \frac{w'_i}{W' - w'_1} \right| \\ & \leq \sum_{i \in S} \left(\left| \frac{w_i}{W - w_1} - \frac{w_i}{W' - w'_1} \right| + \left| \frac{w_i}{W' - w'_1} - \frac{w'_i}{W' - w'_1} \right| \right) \\ & \leq \sum_{i \in S} \left(\frac{w_i}{W} \left| \frac{(W' - W) + (w_1 - w'_1)}{(1 - w_1/W)(W' - w'_1)} \right| + |w_i - w'_i| \left| \frac{1}{W' - w'_1} \right| \right) \\ & \leq \sum_{i \in S} \left(\frac{w_i}{W} \left| \frac{11 + 2}{(1 - 3/k)(k/3 - 1)} \right| + |10/k| \frac{1}{k/3 - 1} \right) \\ & \leq \left(\frac{39}{0.9(k - 3)} \right) + \frac{30/k}{k - 3} \leq 100/k \end{aligned}$$

where in the last line we used the fact that $k \geq 30$ and in the second to the last line we use Claim C.2 to lower bound W and W' and Claim C.3 to upper bound $|W - W'|$ and $|w_i - w'_i|$.

We have now bounded Eq. (C.8), Eq. (C.9), Eq. (C.10), and Eq. (C.7). The sum of these terms is at most $\frac{2r}{W} + \frac{2r}{W'} + 100(2r + \phi)/k$. Using the Claim C.2 again we can upper bound this sum by $(212r + 100\phi)/k < 400(r + \phi)/k$. \square

We now verify that the algorithm is $(2\varepsilon, 2\delta)$ -DP in the present case. Let $T \subseteq \mathcal{Y}$. Then

$$\begin{aligned} \mathbb{P}[\mathcal{M}(D) \in T] &= \mathbb{P}[\mathcal{M}(D) \neq \perp] \mathbb{P}[\mathcal{B}_{400(r+\phi)/k, \varepsilon, \delta}(\mu) \in T] \\ &\leq (e^\varepsilon \mathbb{P}[\mathcal{M}(D') \neq \perp] + \delta) (e^\varepsilon \mathbb{P}[\mathcal{B}_{400(r+\phi)/k, \varepsilon, \delta}(\mu') \in T] + \delta) \\ &= e^{2\varepsilon} \mathbb{P}[\mathcal{M}(D') \neq \perp] \mathbb{P}[\mathcal{B}_{400(r+\phi)/k, \varepsilon, \delta}(\mu') \in T] \\ &\quad + e^\varepsilon \delta (\mathbb{P}[\mathcal{M}(D') \neq \perp] + \mathbb{P}[\mathcal{B}_{400(r+\phi)/k, \varepsilon, \delta}(\mu') \in T]) + \delta^2 \\ &\leq e^{2\varepsilon} \mathbb{P}[\mathcal{M}(D') \in T] + 3e^\varepsilon \delta, \end{aligned}$$

which establishes Eq. (C.2). Finally,

$$\begin{aligned}\mathbb{P}[\mathcal{M}(D) \in T \cup \{\perp\}] &= \mathbb{P}[\mathcal{M}(D) = \perp] + \mathbb{P}[\mathcal{M}(D) \in T] \\ &\leq (e^\varepsilon \mathbb{P}[\mathcal{M}(D') = \perp] + \delta) + (e^{2\varepsilon} \mathbb{P}[\mathcal{M}(D') \in T] + 3e^\varepsilon \delta) \\ &\leq e^{2\varepsilon} \mathbb{P}[\mathcal{M}(D') \in T \cup \{\perp\}] + 3e^\varepsilon \delta,\end{aligned}$$

which establishes Eq. (C.3). \square

D Missing proofs from Section 9

D.1 Proof of Lemma 9.1

Lemma 9.1 follows easily from the following lemma.

Lemma D.1. *Suppose that*

$$\max\{\|\Sigma_1^{-1/2}\Sigma_2\Sigma_1^{-1/2} - I_d\|, \|\Sigma_2^{-1/2}\Sigma_1\Sigma_2^{-1/2} - I_d\|\} \leq \gamma \leq 1/2.$$

Then for every $\delta_0 > 0$, $\Sigma_1^{1/2}(I + \eta G)$ and $\Sigma_2^{1/2}(I + \eta G)$ are $(\varepsilon_0(\delta_0), \delta_0)$ -indistinguishable where

$$\varepsilon_0(\delta_0) = \frac{\gamma^2 d}{2}(d + 1/\eta^2) + 2d\gamma\sqrt{\ln(2/\delta_0)} + 2\gamma \ln(2/\delta_0) + 3\gamma\sqrt{d}\sqrt{\ln(2/\delta_0)}/\eta. \quad (\text{D.1})$$

Indeed, given Lemma D.1, Lemma 9.1 follows by setting γ appropriately so that each term in Eq. (D.1) is at most $\varepsilon/4$ and post-processing, i.e. since $\Sigma_1^{1/2}(I + \eta G)$ and $\Sigma_2^{1/2}(I + \eta G)$ are (ε, δ) -indistinguishable then so are $\Sigma_1^{1/2}(I + \eta G)(I + \eta G)^\top \Sigma_1^{1/2}$ and $\Sigma_2^{1/2}(I + \eta G)(I + \eta G)^\top \Sigma_2^{1/2}$.

First, we require the following technical lemma which asserts that for Gaussians $\mathcal{N}(\mu, \Sigma), \mathcal{N}(\hat{\mu}, \hat{\Sigma})$, if $g \sim \mathcal{N}(0, I_d)$ then the random variable $\mathcal{L}_{\mathcal{N}(\mu, \Sigma) \parallel \mathcal{N}(\hat{\mu}, \hat{\Sigma})}(\Sigma^{1/2}g + \mu)$ concentrates tightly around $D_{\text{KL}}(\mathcal{N}(\mu, \Sigma) \parallel \mathcal{N}(\hat{\mu}, \hat{\Sigma}))$. The plan is to apply the lemma with $\mu \in \mathbb{R}^{d^2}$ (resp. $\hat{\mu}$) is $\Sigma_1^{1/2}$ (resp. $\Sigma_2^{1/2}$) but viewed as a vector of length d^2 and Σ (resp. $\hat{\Sigma}$) is the covariance matrix of (the vector) $\Sigma_1^{1/2}(I + \eta G)$ (resp. $\Sigma_2^{1/2}(I + \eta G)$).

Lemma D.2. *Let $\mu, \hat{\mu} \in \mathbb{R}^d$ and $\Sigma, \hat{\Sigma} \in \mathbb{S}_{>0}^d$. Let $g \sim \mathcal{N}(0, I_d)$. Then*

$$\begin{aligned}\mathcal{L}_{\mathcal{N}(\mu, \Sigma) \parallel \mathcal{N}(\hat{\mu}, \hat{\Sigma})}(\Sigma^{1/2}g + \mu) &\leq D_{\text{KL}}(\mathcal{N}(\mu, \Sigma) \parallel \mathcal{N}(\hat{\mu}, \hat{\Sigma})) \\ &\quad + 2\|\Sigma^{1/2}\hat{\Sigma}^{-1}\Sigma^{1/2} - I\|_F\sqrt{\ln(2/\delta)} + 2\|\Sigma^{1/2}\hat{\Sigma}^{-1}\Sigma^{1/2} - I\| \ln(2/\delta) \\ &\quad + \|\Sigma^{1/2}\hat{\Sigma}^{-1}\Sigma^{1/2}\| \cdot \|\Sigma^{-1/2}(\hat{\mu} - \mu)\|_2\sqrt{2\ln(2/\delta)}\end{aligned}$$

with probability at least $1 - \delta$.

The proof of Lemma D.2 can be found in Subsection D.2.

We use Lemma D.2 to show that if $\|\Sigma_1^{-1/2}\Sigma_2\Sigma_1^{-1/2} - I\|$ and $\|\Sigma_2^{-1/2}\Sigma_1\Sigma_2^{-1/2} - I\|$ are small then $\Sigma_1^{1/2}(I_d + \eta G)$ and $\Sigma_2^{1/2}(I_d + \eta G)$ are indistinguishable. As a consequence, the matrices $\Sigma_1^{1/2}(I_d + \eta G)(I_d + \eta G)^\top \Sigma_1^{1/2}$ and $\Sigma_2^{1/2}(I_d + \eta G)(I_d + \eta G)^\top \Sigma_2^{1/2}$ are also indistinguishable.

Before we prove [Lemma D.1](#), it is convenient to set up some notation. For a matrix $B = [b_1 \cdots b_d] \in \mathbb{R}^{d \times d}$, we write $\text{vec}(B)$ to denote a vector of length d^2 with the columns of B stacked together. More precisely,

$$\text{vec}(B) = \begin{bmatrix} b_1 \\ \vdots \\ b_d \end{bmatrix} \in \mathbb{R}^{d^2}.$$

Next, for a matrix $A = [a_1 \cdots a_d] \in \mathbb{R}^{k \times d}$, we let $\text{bd}(A)$ be the block-diagonal matrix with d copies of A along its diagonal, i.e.

$$\text{bd}(A) = \begin{bmatrix} A & 0 & \cdots & 0 \\ 0 & A & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A \end{bmatrix} \in \mathbb{R}^{kd \times kd}.$$

Note that for a size-compatible matrix $\text{vec}(AB) = \text{bd}(A) \text{vec}(B)$. Indeed, $AB = [Ab_1 \cdots Ab_d]$ so

$$\underbrace{\begin{bmatrix} Ab_1 \\ \vdots \\ Ab_d \end{bmatrix}}_{\text{vec}(AB)} = \underbrace{\begin{bmatrix} A & 0 & \cdots & 0 \\ 0 & A & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A \end{bmatrix}}_{\text{bd}(A)} \underbrace{\begin{bmatrix} b_1 \\ \vdots \\ b_d \end{bmatrix}}_{\text{vec}(B)}.$$

We require a few calculations before we can prove [Lemma D.1](#).

Claim D.3. *Suppose that $\|\Sigma_1^{1/2} \Sigma_2^{-1} \Sigma_1^{1/2} - I_d\| \leq \gamma$. Then*

$$\|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2} - I_{d^2}\|_F^2 = d \cdot \|\Sigma_1^{-1/2} \Sigma_2 \Sigma_1^{-1/2} - I_d\|_F^2 \leq d^2 \gamma^2.$$

Proof. The equality is because the matrices $\text{bd}(\Sigma_1), \text{bd}(\Sigma_2), I_{d^2}$ are all block diagonal with copies of Σ_1, Σ_2, I_d along its diagonal.

For the inequality, we have $\|\Sigma_1^{1/2} \Sigma_2^{-1} \Sigma_1^{1/2} - I_d\| \leq \gamma$ which implies $\|\Sigma_1^{1/2} \Sigma_2^{-1} \Sigma_1^{1/2} - I_d\|_F^2 \leq \gamma^2 \cdot d$. Multiplying both sides by d gives the result. \square

Claim D.4. *Suppose that $\|\Sigma_1^{1/2} \Sigma_2^{-1} \Sigma_1^{1/2} - I_d\| \leq \gamma$. Then $\|\eta^{-1} \text{bd}(\Sigma_2)^{-1/2} (\text{vec}(\Sigma_2^{1/2}) - \text{vec}(\Sigma_1^{1/2}))\|_2^2 \leq \gamma^2 d / \eta^2$.*

Proof. Note that $\|\text{bd}(\Sigma_2)^{-1/2} (\text{vec}(\Sigma_2^{1/2}) - \text{vec}(\Sigma_1^{1/2}))\|_2^2 = \|I - \Sigma_2^{-1/2} \Sigma_1^{1/2}\|_F^2$. Let $\lambda_1, \dots, \lambda_d$ be the (positive) eigenvalues of $\Sigma_1^{1/2} \Sigma_2^{-1} \Sigma_1^{1/2}$.⁷ Then $\sqrt{\lambda_1}, \dots, \sqrt{\lambda_d}$ are the singular values of $\Sigma_2^{-1/2} \Sigma_1^{1/2}$

⁷The eigenvalues of $\Sigma_2^{-1/2} \Sigma_1 \Sigma_2^{-1/2} - I_d$ lie in the interval $[-1/2, 1/2]$ so the eigenvalues of $\Sigma_2^{-1/2} \Sigma_1 \Sigma_2^{-1/2}$ lie in the interval $[1/2, 3/2]$. In particular, the eigenvalues are positive.

and $1 - \sqrt{\lambda_1}, \dots, 1 - \sqrt{\lambda_d}$ are the singular values of $I_d - \Sigma_2^{-1/2} \Sigma_1^{1/2}$. Thus,

$$\begin{aligned}
\|I_d - \Sigma_2^{-1/2} \Sigma_1^{1/2}\|_F^2 &= \sum_{i=1}^d (1 - \sqrt{\lambda_i})^2 \\
&= \sum_{i=1}^d 1 - 2\sqrt{\lambda_i} + \lambda_i \\
&\leq \sum_{i=1}^d 1 - 2\sqrt{1 - \gamma} + (1 + \gamma) \\
&\leq \sum_{i=1}^d 1 - 2(1 - \gamma/2 - \gamma^2/2) + (1 + \gamma) \\
&= \sum_{i=1}^d \gamma^2 = \gamma^2 d,
\end{aligned}$$

where in the third line we used that $\lambda_i \in [1 - \gamma, 1 + \gamma]$ for all $i \in [d]$ and in the fourth line, we used the assumption that $\gamma \leq 1/2$ and Claim B.2 to bound $\sqrt{1 - \gamma} \leq 1 - \gamma/2 - \gamma^2/2$. The lemma follows by dividing both sides by η^2 . \square

Claim D.5. *Suppose that $\|\Sigma_1^{1/2} \Sigma_2^{-1} \Sigma_1^{1/2} - I_d\| \leq \gamma \leq 1/2$. Then*

$$D_{\text{KL}} \left(\mathcal{N}(\text{vec}(\Sigma_1^{1/2}), \eta^2 \text{bd}(\Sigma_1)) \parallel \mathcal{N}(\text{vec}(\Sigma_2^{1/2}), \eta^2 \text{bd}(\Sigma_2)) \right) \leq \frac{\gamma^2 d}{2} (d + 1/\eta^2).$$

Proof. Follows by plugging in the bounds from Claim D.3 and Claim D.4 into Fact 6.1. \square

Proof of Lemma D.1. It is convenient to view $\Sigma_i^{1/2}(I_d + \eta G)$ as d^2 -dimensional vectors. Indeed, we have $\text{vec}(\Sigma_i^{1/2}(I_d + \eta G)) = \text{vec}(\Sigma_i^{1/2}) + \eta \text{bd}(\Sigma_i^{1/2}) \text{vec}(G)$ which is a random d^2 -dimensional vector distributed as $\mathcal{N}(\text{vec}(\Sigma_i^{1/2}), \eta^2 \text{bd}(\Sigma_i))$. For notation, let us write $\mathcal{D}_i = \mathcal{N}(\text{vec}(\Sigma_i^{1/2}), \eta^2 \text{bd}(\Sigma_i))$. We will use Lemma D.2 to bound

$$\mathcal{L} := \mathcal{L}_{\mathcal{D}_1 \parallel \mathcal{D}_2}(\eta \text{bd}(\Sigma_1)^{1/2} \text{vec}(G) + \text{vec}(\Sigma_1)^{1/2}).$$

An identical argument holds with Σ_1, Σ_2 reversed and so we may apply Lemma 6.10.

By Lemma D.2, we have that

$$\begin{aligned}
\mathcal{L} &\leq D_{\text{KL}}(\mathcal{D}_1 \parallel \mathcal{D}_2) \\
&\quad + 2\|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2} - I\|_F \sqrt{\ln(2/\delta)} \\
&\quad + 2\|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2} - I_{d^2}\| \ln(2/\delta) \\
&\quad + \|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2}\| \cdot \|\text{bd}(\Sigma_1)^{-1} (\text{vec}(\Sigma_2) - \text{vec}(\Sigma_1))\|_2 \sqrt{2 \ln(2/\delta)},
\end{aligned} \tag{D.2}$$

with probability at least $1 - \delta$. We now bound each term in Eq. (D.2).

The first term in Eq. (D.2) is bounded by Claim D.5 and we have

$$D_{\text{KL}}(\mathcal{D}_1 \parallel \mathcal{D}_2) \leq \frac{\gamma^2 d}{2}(d + 1/\eta^2). \quad (\text{D.3})$$

The second term is bounded by Claim D.3 and we have

$$\|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2} - I_{d^2}\|_F \leq \gamma d. \quad (\text{D.4})$$

For the third term, note that $\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2}$ and $\Sigma_1^{1/2} \Sigma_2^{-1} \Sigma_1$ have the same set of eigenvalues but with different multiplicities (by a factor of d). Hence,

$$\|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2} - I_{d^2}\| \leq \gamma. \quad (\text{D.5})$$

Finally, for the last term, we use that $\|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2}\| \leq 1 + \gamma \leq 3/2$ and Claim D.4 to obtain

$$\|\text{bd}(\Sigma_1)^{1/2} \text{bd}(\Sigma_2)^{-1} \text{bd}(\Sigma_1)^{1/2}\| \cdot \|\text{bd}(\Sigma_1)^{-1}(\text{vec}(\Sigma_2) - \text{vec}(\Sigma_1))\|_2 < 3\gamma\sqrt{d}/2\eta. \quad (\text{D.6})$$

Plugging Eq. (D.3) through Eq. (D.6) into Eq. (D.2) gives

$$\mathcal{L} \leq \frac{\gamma^2 d}{2}(d + 1/\eta^2) + 2d\gamma\sqrt{\ln(2/\delta)} + 2\gamma \ln(2/\delta) + 3\gamma\sqrt{d}\sqrt{\ln(2/\delta)}/\eta,$$

with probability at least $1 - \delta$. \square

D.2 Proof of Lemma D.2

Proof of Lemma D.2. Let $\mathcal{L} = \mathcal{L}_{\mathcal{N}(\mu, \Sigma) \parallel \mathcal{N}(\hat{\mu}, \hat{\Sigma})}(\Sigma^{1/2}g + \mu)$. From Claim 6.11, we have that

$$\mathcal{L} = -\frac{1}{2} \ln(\det(\hat{\Sigma}^{-1}\Sigma)) - \frac{1}{2}g^\top g + \frac{1}{2}(g - \Sigma^{-1/2}(\hat{\mu} - \mu))^\top \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2} (g - \Sigma^{-1/2}(\hat{\mu} - \mu)). \quad (\text{D.7})$$

Expanding the last term, we have

$$\mathcal{L} = -\frac{1}{2} \ln(\det(\hat{\Sigma}^{-1}\Sigma)) + \frac{1}{2}g^\top (I - \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2})g - (\hat{\mu} - \mu)^\top \hat{\Sigma}^{-1} \Sigma^{1/2} g + \frac{1}{2}(\hat{\mu} - \mu)^\top \hat{\Sigma}^{-1} (\hat{\mu} - \mu). \quad (\text{D.8})$$

By Lemma 6.4, we have that

$$\begin{aligned} g^\top (I - \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2})g &\leq \text{tr}(I - \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}) + 4\|I - \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}\|_F \sqrt{\ln(2/\delta)} \\ &\quad + 4\|I - \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}\| \ln(2/\delta), \end{aligned} \quad (\text{D.9})$$

with probability $1 - \delta/2$. Next, $(\hat{\mu} - \mu)^\top \hat{\Sigma}^{-1} \Sigma^{1/2} g$ is a mean-zero Gaussian random variable with variance $\|\Sigma^{1/2} \hat{\Sigma}^{-1} (\hat{\mu} - \mu)\|_2^2 \leq \|\Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}\|^2 \|\Sigma^{-1/2} (\hat{\mu} - \mu)\|_2^2$. Hence, using the tails of a Gaussian random variable (Lemma 6.3), we have

$$(\hat{\mu} - \mu)^\top \hat{\Sigma}^{-1} \Sigma^{1/2} g \leq \|\Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}\| \|\Sigma^{-1/2} (\hat{\mu} - \mu)\|_2 \sqrt{2 \ln(2/\delta)} \quad (\text{D.10})$$

with probability $1 - \delta/2$. Plugging Eq. (D.9) and Eq. (D.10) along with the formula for the KL divergence between two multivariate Normal distributions (Fact 6.1), we have

$$\begin{aligned} \mathcal{L} &\leq D_{\text{KL}}(\mathcal{N}(\mu, \Sigma) \parallel \mathcal{N}(\hat{\mu}, \hat{\Sigma})) + 2\|I - \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}\|_F \sqrt{\ln(2/\delta)} \\ &\quad + 2\|I - \Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}\| \ln(2/\delta) + \|\Sigma^{1/2} \hat{\Sigma}^{-1} \Sigma^{1/2}\| \|\Sigma^{-1/2} (\hat{\mu} - \mu)\|_2 \sqrt{2 \ln(2/\delta)}, \end{aligned} \quad (\text{D.11})$$

with probability $1 - \delta$. \square