

On a class of critical N -Laplacian problems*

Tsz Chung Ho and Kanishka Perera

Department of Mathematical Sciences

Florida Institute of Technology

Melbourne, FL 32901, USA

tho2011@my.fit.edu & kperera@fit.edu

Abstract

We establish some existence results for a class of critical N -Laplacian problems in a bounded domain in \mathbb{R}^N . In the absence of a suitable direct sum decomposition of the underlying Sobolev space to which the classical linking theorem can be applied, we use an abstract linking theorem based on the \mathbb{Z}_2 -cohomological index to obtain a nontrivial critical point.

1 Introduction

In this paper we establish some existence results for the class of critical N -Laplacian problems

$$\begin{cases} -\Delta_N u = h(u) e^{\alpha |u|^{N'}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where Ω is a smooth bounded domain in \mathbb{R}^N , $N \geq 2$, $\alpha > 0$, $N' = N/(N-1)$ is the Hölder conjugate of N , and h is a continuous function such that

$$\lim_{|t| \rightarrow \infty} h(t) = 0 \quad (1.2)$$

and

$$0 < \beta := \liminf_{|t| \rightarrow \infty} th(t) < \infty. \quad (1.3)$$

**MSC2010*: Primary 35J92, Secondary 35B33, 35B38

Key Words and Phrases: critical N -Laplacian problems, existence, critical points, linking, \mathbb{Z}_2 -cohomological index

This problem is motivated by the Trudinger-Moser inequality

$$\sup_{\substack{u \in W_0^{1,N}(\Omega) \\ \|u\| \leq 1}} \int_{\Omega} e^{\alpha_N |u|^{N'}} dx < \infty, \quad (1.4)$$

where $W_0^{1,N}(\Omega)$ is the usual Sobolev space with the norm

$$\|u\| = \left(\int_{\Omega} |\nabla u|^N dx \right)^{1/N},$$

$$\alpha_N = N \omega_{N-1}^{1/(N-1)},$$

and

$$\omega_{N-1} = \frac{2\pi^{N/2}}{\Gamma(N/2)}$$

is the area of the unit sphere in \mathbb{R}^N (see Trudinger [14] and Moser [10]). Problem (1.1) is critical with respect to this inequality and hence lacks compactness. Indeed, the associated variational functional satisfies the Palais-Smale compactness condition only at energy levels below a certain threshold (see Proposition 2.1 in the next section).

In dimension $N = 2$, problem (1.1) is semilinear and has been extensively studied in the literature (see, e.g., [2, 3, 4, 6]). In dimensions $N \geq 3$, this problem is quasilinear and has been studied mainly when

$$G(t) := \int_0^t h(s) e^{\alpha |s|^{N'}} ds \leq \lambda |t|^N \quad \text{for small } t \quad (1.5)$$

for some $\lambda \in (0, \lambda_1)$ (see, e.g., [1, 5, 9]). Here

$$\lambda_1 = \inf_{u \in W_0^{1,N}(\Omega) \setminus \{0\}} \frac{\int_{\Omega} |\nabla u|^N dx}{\int_{\Omega} |u|^N dx} \quad (1.6)$$

is the first eigenvalue of the eigenvalue problem

$$\begin{cases} -\Delta_N u = \lambda |u|^{N-2} u & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases} \quad (1.7)$$

The case $h(t) = \lambda |t|^{N-2} t$ with $\lambda > 0$, for which $\beta = \infty$, was recently studied in Yang and Perera [15]. The remaining case, where $N \geq 3$, $\lambda \geq \lambda_1$, and $\beta < \infty$, does not seem to have been studied in the literature. This case is covered in our results here, which are for large $\beta < \infty$ and allow $N \geq 3$ and $\lambda \geq \lambda_1$ in (1.5).

Let d be the radius of the largest open ball contained in Ω . Our first result is the following theorem.

Theorem 1.1. *Assume that $\alpha > 0$, h satisfies (1.2) and (1.3), and G satisfies*

$$G(t) \geq -\frac{1}{N} \sigma_0 |t|^N \quad \text{for } t \geq 0, \quad (1.8)$$

$$G(t) \leq \frac{1}{N} (\lambda_1 - \sigma_1) |t|^N \quad \text{for } |t| \leq \delta \quad (1.9)$$

for some $\sigma_0 \geq 0$ and $\sigma_1, \delta > 0$. If

$$\beta > \frac{1}{N\alpha^{N-1}} \left(\frac{N}{d} \right)^N e^{\sigma_0/(N-1)\kappa}, \quad (1.10)$$

where $\kappa = \frac{1}{N!} \left(\frac{N}{d} \right)^N$, then problem (1.1) has a nontrivial solution.

In particular, we have the following corollary for $\sigma_0 = 0$.

Corollary 1.2. *Assume that $\alpha > 0$, h satisfies (1.2) and (1.3), and G satisfies*

$$G(t) \geq 0 \quad \text{for } t \geq 0,$$

$$G(t) \leq \frac{1}{N} (\lambda_1 - \sigma_1) |t|^N \quad \text{for } |t| \leq \delta$$

for some $\sigma_1, \delta > 0$. If

$$\beta > \frac{1}{N\alpha^{N-1}} \left(\frac{N}{d} \right)^N,$$

then problem (1.1) has a nontrivial solution.

Corollary 1.2 should be compared with Theorem 1 of do Ó [9], where this result is proved under the stronger assumption $h(t) \geq 0$ for $t \geq 0$.

To state our second result, let (λ_k) be the sequence of eigenvalues of problem (1.7) based on the \mathbb{Z}_2 -cohomological index that was introduced in Perera [11] (see Proposition 2.3 in the next section). We have the following theorem.

Theorem 1.3. *Assume that $\alpha > 0$, h satisfies (1.2) and (1.3), and G satisfies*

$$G(t) \geq \frac{1}{N} (\lambda_{k-1} + \sigma_0) |t|^N \quad \forall t, \quad (1.11)$$

$$G(t) \leq \frac{1}{N} (\lambda_k - \sigma_1) |t|^N \quad \text{for } |t| \leq \delta \quad (1.12)$$

for some $k \geq 2$ and $\sigma_0, \sigma_1, \delta > 0$. Then there exists a constant $c > 0$ depending on Ω , α , and k , but not on σ_0 , σ_1 , or δ , such that if

$$\beta > \frac{1}{\alpha^{N-1}} \left(\frac{N}{d} \right)^N e^{c/\sigma_0^{N-1}},$$

then problem (1.1) has a nontrivial solution.

Theorem 1.3 should be compared with Theorem 1.4 of de Figueiredo et al. [3, 4], where this result is proved in the case $N = 2$ under the additional assumption that $0 < 2G(t) \leq th(t)e^{\alpha t^2}$ for all $t \in \mathbb{R} \setminus \{0\}$. However, the linking argument used in [3, 4] is based on a splitting of $H_0^1(\Omega)$ that involves the eigenspaces of the Laplacian, and this argument does not extend to the case $N \geq 3$ where the N -Laplacian is a nonlinear operator and therefore has no linear eigenspaces. We will prove Theorem 1.3 using an abstract critical point theorem based on the \mathbb{Z}_2 -cohomological index that was proved in Yang and Perera [15] (see Section 2.4).

In the proofs of Theorems 1.1 and 1.3, the inner radius d of Ω comes into play when verifying that certain minimax levels are below the compactness threshold given in Proposition 2.1.

2 Preliminaries

2.1 A compactness result

Weak solutions of problem (1.1) coincide with critical points of the C^1 -functional

$$E(u) = \frac{1}{N} \int_{\Omega} |\nabla u|^N dx - \int_{\Omega} G(u) dx, \quad u \in W_0^{1,N}(\Omega).$$

We recall that a $(PS)_c$ sequence of E is a sequence $(u_j) \subset W_0^{1,N}(\Omega)$ such that $E(u_j) \rightarrow c$ and $E'(u_j) \rightarrow 0$. Proofs of Theorem 1.1 and Theorem 1.3 will be based on the following compactness result.

Proposition 2.1. *Assume that $\alpha > 0$ and h satisfies (1.2) and (1.3). Then for all $c \neq 0$ satisfying*

$$c < \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1},$$

every $(PS)_c$ sequence of E has a subsequence that converges weakly to a nontrivial solution of problem (1.1).

Proof. Let $(u_j) \subset W_0^{1,N}(\Omega)$ be a $(PS)_c$ sequence of E . Then

$$E(u_j) = \frac{1}{N} \|u_j\|^N - \int_{\Omega} G(u_j) dx = c + o(1) \tag{2.1}$$

and

$$E'(u_j) u_j = \|u_j\|^N - \int_{\Omega} u_j h(u_j) e^{\alpha |u_j|^{N'}} dx = o(\|u_j\|). \quad (2.2)$$

First we show that (u_j) is bounded in $W_0^{1,N}(\Omega)$. Multiplying (2.1) by $2N$ and subtracting (2.2) gives

$$\|u_j\|^N + \int_{\Omega} \left(u_j h(u_j) e^{\alpha |u_j|^{N'}} - 2NG(u_j) \right) dx = 2Nc + o(\|u_j\| + 1),$$

so it suffices to show that $th(t) e^{\alpha |t|^{N'}} - 2NG(t)$ is bounded from below. Let $0 < \varepsilon < \beta/(2N+1)$. By (1.2) and (1.3), for some constant $C_\varepsilon > 0$,

$$|G(t)| \leq \varepsilon e^{\alpha |t|^{N'}} + C_\varepsilon \quad (2.3)$$

and

$$th(t) e^{\alpha |t|^{N'}} \geq (\beta - \varepsilon) e^{\alpha |t|^{N'}} - C_\varepsilon \quad (2.4)$$

for all t . So

$$th(t) e^{\alpha |t|^{N'}} - 2NG(t) \geq [\beta - (2N+1)\varepsilon] e^{\alpha |t|^{N'}} - (2N+1)C_\varepsilon,$$

which is bounded from below.

Since (u_j) is bounded in $W_0^{1,N}(\Omega)$, a renamed subsequence converges to some u weakly in $W_0^{1,N}(\Omega)$, strongly in $L^p(\Omega)$ for all $p \in [1, \infty)$, and a.e. in Ω . We have

$$E'(u_j) v = \int_{\Omega} |\nabla u_j|^{N-2} \nabla u_j \cdot \nabla v dx - \int_{\Omega} v h(u_j) e^{\alpha |u_j|^{N'}} dx \rightarrow 0 \quad (2.5)$$

for all $v \in W_0^{1,N}(\Omega)$. By (1.2), given any $\varepsilon > 0$, there exists a constant $C_\varepsilon > 0$ such that

$$|h(t) e^{\alpha |t|^{N'}}| \leq \varepsilon e^{\alpha |t|^{N'}} + C_\varepsilon \quad \forall t. \quad (2.6)$$

By (2.2),

$$\sup_j \int_{\Omega} u_j h(u_j) e^{\alpha |u_j|^{N'}} dx < \infty,$$

which together with (2.4) gives

$$\sup_j \int_{\Omega} e^{\alpha |u_j|^{N'}} dx < \infty. \quad (2.7)$$

For $v \in C_0^\infty(\Omega)$, it follows from (2.6) and (2.7) that the sequence $(v h(u_j) e^{\alpha |u_j|^{N'}})$ is uniformly integrable and hence

$$\int_{\Omega} v h(u_j) e^{\alpha |u_j|^{N'}} dx \rightarrow \int_{\Omega} v h(u) e^{\alpha |u|^{N'}} dx$$

by Vitali's convergence theorem, so it follows from (2.5) that

$$\int_{\Omega} |\nabla u|^{N-2} \nabla u \cdot \nabla v dx - \int_{\Omega} v h(u) e^{\alpha |u|^{N'}} dx = 0.$$

Then this holds for all $v \in W_0^{1,N}(\Omega)$ by density, so the weak limit u is a solution of problem (1.1).

Suppose that $u = 0$. Then

$$\int_{\Omega} G(u_j) dx \rightarrow 0$$

since (2.3) and (2.7) imply that the sequence $(G(u_j))$ is uniformly integrable, so (2.1) gives $c \geq 0$ and

$$\|u_j\| \rightarrow (Nc)^{1/N}. \quad (2.8)$$

Let $Nc < \nu < (\alpha_N/\alpha)^{N-1}$. Then $\|u_j\| \leq \nu^{1/N}$ for all $j \geq j_0$ for some j_0 . Let $q = \alpha_N/\alpha \nu^{1/(N-1)} > 1$. By the Hölder inequality,

$$\left| \int_{\Omega} u_j h(u_j) e^{\alpha |u_j|^{N'}} dx \right| \leq \left(\int_{\Omega} |u_j h(u_j)|^p dx \right)^{1/p} \left(\int_{\Omega} e^{q\alpha |u_j|^{N'}} dx \right)^{1/q},$$

where $1/p + 1/q = 1$. The first integral on the right-hand side converges to zero since h is bounded and $u_j \rightarrow 0$ in $L^p(\Omega)$, and the second integral is bounded by (1.4) since $q\alpha |u_j|^{N'} = \alpha_N |\tilde{u}_j|^{N'}$, where $\tilde{u}_j = u_j/\nu^{1/N}$ satisfies $\|\tilde{u}_j\| \leq 1$ for $j \geq j_0$, so

$$\int_{\Omega} u_j h(u_j) e^{\alpha |u_j|^{N'}} dx \rightarrow 0.$$

Then $u_j \rightarrow 0$ by (2.2) and hence $c = 0$ by (2.8), contrary to assumption. So u is a nontrivial solution. \square

2.2 \mathbb{Z}_2 -cohomological index

The \mathbb{Z}_2 -cohomological index of Fadell and Rabinowitz [8] is defined as follows. Let W be a Banach space and let \mathcal{A} denote the class of symmetric subsets of $W \setminus \{0\}$. For $A \in \mathcal{A}$, let $\overline{A} = A/\mathbb{Z}_2$ be the quotient space of A with each u and $-u$ identified, let $f : \overline{A} \rightarrow \mathbb{R}P^\infty$ be

the classifying map of \overline{A} , and let $f^* : H^*(\mathbb{RP}^\infty) \rightarrow H^*(\overline{A})$ be the induced homomorphism of the Alexander-Spanier cohomology rings. The cohomological index of A is defined by

$$i(A) = \begin{cases} \sup \{m \geq 1 : f^*(\omega^{m-1}) \neq 0\}, & A \neq \emptyset \\ 0, & A = \emptyset, \end{cases}$$

where $\omega \in H^1(\mathbb{RP}^\infty)$ is the generator of the polynomial ring $H^*(\mathbb{RP}^\infty) = \mathbb{Z}_2[\omega]$. For example, the classifying map of the unit sphere S^{m-1} in \mathbb{R}^m , $m \geq 1$ is the inclusion $\mathbb{RP}^{m-1} \subset \mathbb{RP}^\infty$, which induces isomorphisms on H^q for $q \leq m-1$, so $i(S^{m-1}) = m$.

The following proposition summarizes the basic properties of the cohomological index (see Fadell and Rabinowitz [8]).

Proposition 2.2. *The index $i : \mathcal{A} \rightarrow \mathbb{N} \cup \{0, \infty\}$ has the following properties:*

- (i) *Definiteness:* $i(A) = 0$ if and only if $A = \emptyset$.
- (ii) *Monotonicity:* If there is an odd continuous map from A to B (in particular, if $A \subset B$), then $i(A) \leq i(B)$. Thus, equality holds when the map is an odd homeomorphism.
- (iii) *Dimension:* $i(A) \leq \dim W$.
- (iv) *Continuity:* If A is closed, then there is a closed neighborhood $N \in \mathcal{A}$ of A such that $i(N) = i(A)$. When A is compact, N may be chosen to be a δ -neighborhood $N_\delta(A) = \{u \in W : \text{dist}(u, A) \leq \delta\}$.
- (v) *Subadditivity:* If A and B are closed, then $i(A \cup B) \leq i(A) + i(B)$.
- (vi) *Stability:* If SA is the suspension of $A \neq \emptyset$, obtained as the quotient space of $A \times [-1, 1]$ with $A \times \{1\}$ and $A \times \{-1\}$ collapsed to different points, then $i(SA) = i(A) + 1$.
- (vii) *Piercing property:* If A , A_0 and A_1 are closed, and $\varphi : A \times [0, 1] \rightarrow A_0 \cup A_1$ is a continuous map such that $\varphi(-u, t) = -\varphi(u, t)$ for all $(u, t) \in A \times [0, 1]$, $\varphi(A \times [0, 1])$ is closed, $\varphi(A \times \{0\}) \subset A_0$ and $\varphi(A \times \{1\}) \subset A_1$, then $i(\varphi(A \times [0, 1]) \cap A_0 \cap A_1) \geq i(A)$.
- (viii) *Neighborhood of zero:* If U is a bounded closed symmetric neighborhood of 0, then $i(\partial U) = \dim W$.

2.3 Eigenvalues

Eigenvalues of problem (1.7) coincide with critical values of the functional

$$\Psi(u) = \frac{1}{\int_{\Omega} |u|^N dx}, \quad u \in S = \left\{ u \in W_0^{1,N}(\Omega) : \int_{\Omega} |\nabla u|^N dx = 1 \right\}.$$

We have the following proposition (see Perera [11] and Perera et al. [12, Proposition 3.52 and Proposition 3.53]).

Proposition 2.3. *Let \mathcal{F} denote the class of symmetric subsets of S and set*

$$\lambda_k := \inf_{\substack{M \in \mathcal{F} \\ i(M) \geq k}} \sup_{u \in M} \Psi(u), \quad k \in \mathbb{N}.$$

Then $0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \rightarrow +\infty$ is a sequence of eigenvalues of problem (1.7). Moreover, if $\lambda_{k-1} < \lambda_k$, then

$$i(\Psi^{\lambda_{k-1}}) = i(S \setminus \Psi_{\lambda_k}) = k - 1,$$

where $\Psi^a = \{u \in S : \Psi(u) \leq a\}$ and $\Psi_a = \{u \in S : \Psi(u) \geq a\}$ for $a \in \mathbb{R}$.

We will also need the following result of Degiovanni and Lancelotti ([7, Theorem 2.3]).

Proposition 2.4. *If $\lambda_{k-1} < \lambda_k$, then $\Psi^{\lambda_{k-1}}$ contains a compact symmetric set C of index $k - 1$ that is bounded in $C^1(\overline{\Omega})$.*

2.4 An abstract critical point theorem

We will use the following abstract critical point theorem proved in Yang and Perera [15, Theorem 2.2] to prove Theorem 1.3. This result generalizes the linking theorem of Rabinowitz [13].

Theorem 2.5. *Let E be a C^1 -functional defined on a Banach space W and let A_0 and B_0 be disjoint nonempty closed symmetric subsets of the unit sphere $S = \{u \in W : \|u\| = 1\}$ such that*

$$i(A_0) = i(S \setminus B_0) < \infty. \tag{2.9}$$

Assume that there exist $R > \rho > 0$ and $\omega \in S \setminus A_0$ such that

$$\sup E(A) \leq \inf E(B), \quad \sup E(X) < \infty,$$

where

$$A = \{sv : v \in A_0, 0 \leq s \leq R\} \cup \{R\pi((1-t)v + t\omega) : v \in A_0, 0 \leq t \leq 1\},$$

$$B = \{\rho u : u \in B_0\},$$

$$X = \{sv + t\omega : v \in A_0, s, t \geq 0, \|sv + t\omega\| \leq R\},$$

and $\pi : W \setminus \{0\} \rightarrow S, u \mapsto u/\|u\|$ is the radial projection onto S . Let

$$\Gamma = \{\gamma \in C(X, W) : \gamma(X) \text{ is closed and } \gamma|_A = id_A\},$$

and set

$$c := \inf_{\gamma \in \Gamma} \sup_{u \in \gamma(X)} E(u).$$

Then $\inf E(B) \leq c \leq \sup E(X)$, and E has a $(PS)_c$ sequence.

2.5 Moser sequence

For $j \geq 2$, let

$$\omega_j(x) = \frac{1}{\omega_{N-1}^{1/N}} \begin{cases} (\log j)^{(N-1)/N}, & |x| \leq d/j \\ \frac{\log(d/|x|)}{(\log j)^{1/N}}, & d/j < |x| < d \\ 0, & |x| \geq d \end{cases} \quad (2.10)$$

(see Moser [10]).

Proposition 2.6. *We have*

$$\int_{\Omega} \omega_j^m dx = \frac{m! \omega_{N-1}^{1-m/N} d^N}{N^{m+1} (\log j)^{m/N}} \left[1 - \frac{1}{j^N} \sum_{l=1}^m \frac{(N \log j)^{m-l}}{(m-l)!} \right], \quad m = 1, \dots, N \quad (2.11)$$

and

$$\int_{\Omega} |\nabla \omega_j|^m dx = \begin{cases} \frac{\omega_{N-1}^{1-m/N} d^{N-m}}{(N-m) (\log j)^{m/N}} \left(1 - \frac{1}{j^{N-m}} \right), & m = 1, \dots, N-1 \\ 1, & m = N. \end{cases} \quad (2.12)$$

Proof. We have

$$\int_{\Omega} \omega_j^m dx = \frac{\omega_{N-1}^{1-m/N} d^N}{(\log j)^{m/N}} \left[I_m + \frac{(\log j)^m}{N j^N} \right],$$

where

$$I_m = \int_{1/j}^1 (-\log s)^m s^{N-1} ds.$$

We have

$$I_1 = \frac{1}{N^2} \left[1 - \frac{1}{j^N} (N \log j + 1) \right],$$

and integrating by parts gives the recurrence relation

$$I_m = \frac{m}{N} I_{m-1} - \frac{(\log j)^m}{N j^N}, \quad m \geq 2.$$

So

$$I_m = \frac{m!}{N^{m+1}} \left[1 - \frac{1}{j^N} \sum_{l=0}^m \frac{(N \log j)^{m-l}}{(m-l)!} \right],$$

and (2.11) follows. The integral in (2.12) is easily evaluated. \square

2.6 A limit calculation

We will need the following limit in the proof of Theorem 1.1.

Proposition 2.7. *We have*

$$\lim_{n \rightarrow \infty} \int_0^1 n e^{-n(t-t^{N'})} dt = N.$$

Proof. Let $f_n(t) = n e^{-n(t-t^{N'})}$ and set $t_0 = (N')^{-1/(N'-1)}$. For $t \neq t_0$,

$$f_n(t) = g_n(t) - \frac{d}{dt} \left(\frac{e^{-n(t-t^{N'})}}{1 - N' t^{N'-1}} \right), \quad (2.13)$$

where

$$g_n(t) = \frac{N'(N'-1)t^{N'-2}e^{-n(t-t^{N'})}}{(1 - N' t^{N'-1})^2}.$$

Fix δ so small that $0 < \delta < t_0 < 1 - \delta < 1$ and write

$$\int_0^1 f_n(t) dt = \int_0^\delta f_n(t) dt + \int_\delta^{1-\delta} f_n(t) dt + \int_{1-\delta}^1 f_n(t) dt. \quad (2.14)$$

By (2.13),

$$\int_0^\delta f_n(t) dt = \int_0^\delta g_n(t) dt - \frac{e^{-n(\delta-\delta^{N'})}}{1 - N' \delta^{N'-1}} + 1. \quad (2.15)$$

For all $t \in (0, \delta)$, $g_n(t) \rightarrow 0$ as $n \rightarrow \infty$ and $|g_n(t)| \leq N'(N'-1)t^{N'-2}/(1 - N' \delta^{N'-1})^2$, so $\int_0^\delta g_n(t) dt \rightarrow 0$ by the dominated convergence theorem. So $\int_0^1 f_n(t) dt \rightarrow 1$ by (2.15). A similar calculation shows that $\int_{1-\delta}^1 f_n(t) dt \rightarrow N - 1$. On the other hand, it is easily seen that $\int_\delta^{1-\delta} f_n(t) dt \rightarrow 0$. So $\int_0^1 f_n(t) dt \rightarrow N$ by (2.14). \square

3 Proof of Theorem 1.1

In this section we prove Theorem 1.1 by showing that the functional E has the mountain pass geometry with the mountain pass level $c \in (0, (1/N)(\alpha_N/\alpha)^{N-1})$ and applying Proposition 2.1.

Lemma 3.1. *There exists a $\rho > 0$ such that*

$$\inf_{\|u\|=\rho} E(u) > 0.$$

Proof. Since (1.2) implies that h is bounded, there exists a constant $C_\delta > 0$ such that

$$|G(t)| \leq C_\delta |t|^{N+1} e^{\alpha |t|^{N'}} \quad \text{for } |t| > \delta,$$

which together with (1.9) gives

$$\int_{\Omega} G(u) dx \leq \frac{1}{N} (\lambda_1 - \sigma_1) \int_{\Omega} |u|^N dx + C_\delta \int_{\Omega} |u|^{N+1} e^{\alpha |u|^{N'}} dx. \quad (3.1)$$

By (1.6),

$$\int_{\Omega} |u|^N dx \leq \frac{\rho^N}{\lambda_1}, \quad (3.2)$$

where $\rho = \|u\|$. By the Hölder inequality,

$$\int_{\Omega} |u|^{N+1} e^{\alpha |u|^{N'}} dx \leq \left(\int_{\Omega} |u|^{2(N+1)} dx \right)^{1/2} \left(\int_{\Omega} e^{2\alpha |u|^{N'}} dx \right)^{1/2}. \quad (3.3)$$

The first integral on the right-hand side is bounded by $C\rho^{2(N+1)}$ for some constant $C > 0$ by the Sobolev embedding theorem. Since $2\alpha |u|^{N'} = 2\alpha \rho^{N'} |\tilde{u}|^{N'}$, where $\tilde{u} = u/\rho$ satisfies $\|\tilde{u}\| = 1$, the second integral is bounded when $\rho^{N'} \leq \alpha_N/2\alpha$ by (1.4). So combining (3.1)–(3.3) gives

$$\int_{\Omega} G(u) dx \leq \frac{1}{N} \left(1 - \frac{\sigma_1}{\lambda_1} \right) \rho^N + O(\rho^{N+1}) \quad \text{as } \rho \rightarrow 0.$$

Then

$$E(u) \geq \frac{1}{N} \frac{\sigma_1}{\lambda_1} \rho^N + O(\rho^{N+1}),$$

and the desired conclusion follows from this for sufficiently small $\rho > 0$. \square

We may assume without loss of generality that $B_d(0) \subset \Omega$. Let (ω_j) be the sequence of functions defined in (2.10).

Lemma 3.2. *We have*

(i) $E(t\omega_j) \rightarrow -\infty$ as $t \rightarrow \infty$ for all $j \geq 2$,

(ii) $\exists j_0 \geq 2$ such that

$$\sup_{t \geq 0} E(t\omega_{j_0}) < \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1}.$$

Proof. (i) Fix $0 < \varepsilon < \beta$. By (1.3), $\exists M_\varepsilon > 0$ such that

$$th(t) e^{\alpha |t|^{N'}} > (\beta - \varepsilon) e^{\alpha |t|^{N'}} \quad \text{for } |t| > M_\varepsilon. \quad (3.4)$$

Since $e^{\alpha |t|^{N'}} > \alpha^{2N-2} t^{2N} / (2N-2)!$ for all t , then there exists a constant $C_\varepsilon > 0$ such that

$$th(t) e^{\alpha |t|^{N'}} \geq \frac{1}{(2N-2)!} (\beta - \varepsilon) \alpha^{2N-2} t^{2N} - C_\varepsilon |t| \quad (3.5)$$

and

$$G(t) \geq \frac{2N-1}{(2N)!} (\beta - \varepsilon) \alpha^{2N-2} t^{2N} - C_\varepsilon |t| \quad (3.6)$$

for all t . Since $\|\omega_j\| = 1$ and $\omega_j \geq 0$, then

$$E(t\omega_j) \leq \frac{t^N}{N} - \frac{2N-1}{(2N)!} (\beta - \varepsilon) \alpha^{2N-2} t^{2N} \int_{\Omega} \omega_j^{2N} dx + C_\varepsilon t \int_{\Omega} \omega_j dx,$$

and the conclusion follows.

(ii) Set

$$H_j(t) = E(t\omega_j) = \frac{t^N}{N} - \int_{\Omega} G(t\omega_j) dx, \quad t \geq 0.$$

If the conclusion is false, then it follows from (i) that for all $j \geq 2$, $\exists t_j > 0$ such that

$$H_j(t_j) = \frac{t_j^N}{N} - \int_{\Omega} G(t_j\omega_j) dx = \sup_{t \geq 0} H_j(t) \geq \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1}, \quad (3.7)$$

$$H'_j(t_j) = t_j^{N-1} - \int_{\Omega} \omega_j h(t_j\omega_j) e^{\alpha t_j^{N'} \omega_j^{N'}} dx = 0. \quad (3.8)$$

Since $G(t) \geq -C_\varepsilon t$ for all $t \geq 0$ by (3.6), (3.7) gives

$$t_j^N \geq t_0^N - N\delta_j t_j, \quad (3.9)$$

where

$$t_0 = \left(\frac{\alpha_N}{\alpha} \right)^{(N-1)/N}$$

and

$$\delta_j = C_\varepsilon \int_{\Omega} \omega_j dx \rightarrow 0 \quad \text{as } j \rightarrow \infty \quad (3.10)$$

by Proposition 2.6. First we will show that $t_j \rightarrow t_0$.

By (3.9) and the Young's inequality,

$$(1 + \nu) t_j^N \geq t_0^N - \frac{N-1}{\nu^{1/(N-1)}} \delta_j^{N'} \quad \forall \nu > 0,$$

which together with (3.10) gives

$$\liminf_{j \rightarrow \infty} t_j \geq t_0. \quad (3.11)$$

Write (3.8) as

$$t_j^N = \int_{\{t_j \omega_j > M_\varepsilon\}} t_j \omega_j h(t_j \omega_j) e^{\alpha t_j^{N'} \omega_j^{N'}} dx + \int_{\{t_j \omega_j \leq M_\varepsilon\}} t_j \omega_j h(t_j \omega_j) e^{\alpha t_j^{N'} \omega_j^{N'}} dx =: I_1 + I_2. \quad (3.12)$$

Set $r_j = d e^{-M_\varepsilon (\omega_{N-1} \log j)^{1/N} / t_j}$. Since $\liminf t_j > 0$, for all sufficiently large j , $d/j < r_j < d$ and $t_j \omega_j(x) > M_\varepsilon$ if and only if $|x| < r_j$. So (3.4) gives

$$\begin{aligned} I_1 &\geq (\beta - \varepsilon) \int_{\{|x| < r_j\}} e^{\alpha t_j^{N'} \omega_j^{N'}} dx = (\beta - \varepsilon) \left(\int_{\{|x| \leq d/j\}} e^{\alpha t_j^{N'} \omega_j^{N'}} dx \right. \\ &\quad \left. + \int_{\{d/j < |x| < r_j\}} e^{\alpha t_j^{N'} \omega_j^{N'}} dx \right) =: (\beta - \varepsilon) (I_3 + I_4). \end{aligned} \quad (3.13)$$

We have

$$I_3 = \frac{\omega_{N-1}}{N} \left(\frac{d}{j} \right)^N e^{\alpha t_j^{N'} \log j / \omega_{N-1}^{1/(N-1)}} = \frac{\omega_{N-1}}{N} d^N j^{\alpha (t_j^{N'} - t_0^{N'}) / \omega_{N-1}^{1/(N-1)}}. \quad (3.14)$$

Since $th(t) e^{\alpha |t|^{N'}} \geq -C_\varepsilon t$ for all $t \geq 0$ by (3.5),

$$I_2 \geq -C_\varepsilon t_j \int_{\{t_j \omega_j \leq M_\varepsilon\}} \omega_j dx \geq -\delta_j t_j. \quad (3.15)$$

Combining (3.12)–(3.15) and noting that $I_4 \geq 0$ gives

$$t_j^N \geq (\beta - \varepsilon) \frac{\omega_{N-1}}{N} d^N j^{\alpha (t_j^{N'} - t_0^{N'}) / \omega_{N-1}^{1/(N-1)}} - \delta_j t_j.$$

It follows from this that

$$\limsup_{j \rightarrow \infty} t_j \leq t_0,$$

which together with (3.11) shows that $t_j \rightarrow t_0$.

Next we estimate I_4 . We have

$$\begin{aligned}
I_4 &= \int_{\{d/j < |x| < r_j\}} e^{\alpha t_j^{N'} [\log(d/|x|)]^{N'} / (\omega_{N-1} \log j)^{1/(N-1)}} dx \\
&= \omega_{N-1} \left(\int_{d/j}^d e^{\alpha t_j^{N'} [\log(d/r)]^{N'} / (\omega_{N-1} \log j)^{1/(N-1)}} r^{N-1} dr \right. \\
&\quad \left. - \int_{r_j}^d e^{\alpha t_j^{N'} [\log(d/r)]^{N'} / (\omega_{N-1} \log j)^{1/(N-1)}} r^{N-1} dr \right) \\
&= \omega_{N-1} d^N \left(\log j \int_0^1 e^{-Nt [1 - (t_j/t_0)^{N'} t^{1/(N-1)}] \log j} dt \right. \\
&\quad \left. - \int_{s_j}^1 s^{N-1} e^{\alpha t_j^{N'} (-\log s)^{N'} / (\omega_{N-1} \log j)^{1/(N-1)}} ds \right), \tag{3.16}
\end{aligned}$$

where $t = \log(d/r)/\log j$, $s = r/d$, and $s_j = r_j/d = e^{-M_\varepsilon (\omega_{N-1} \log j)^{1/N}/t_j} \rightarrow 0$. For $s_j < s < 1$, $\alpha t_j^{N'} (-\log s)^{N'} / (\omega_{N-1} \log j)^{1/(N-1)}$ is bounded by $\alpha M_\varepsilon^{N'}$ and goes to zero as $j \rightarrow \infty$, so the last integral converges to

$$\int_0^1 s^{N-1} ds = \frac{1}{N}.$$

So combining (3.12)–(3.16) and letting $j \rightarrow \infty$ gives

$$t_0^N \geq (\beta - \varepsilon) \frac{\omega_{N-1}}{N} d^N (L_1 + L_2 - 1),$$

where

$$L_1 = \liminf_{j \rightarrow \infty} e^{-n [1 - (t_j/t_0)^{N'}]},$$

$$L_2 = \liminf_{j \rightarrow \infty} \int_0^1 n e^{-n [t - (t_j/t_0)^{N'} t^{N'}]} dt,$$

and $n = N \log j \rightarrow \infty$. Letting $\varepsilon \rightarrow 0$ in this inequality gives

$$\beta \leq \frac{1}{\alpha^{N-1}} \left(\frac{N}{d} \right)^N \frac{1}{L_1 + L_2 - 1}. \tag{3.17}$$

By (3.7), (1.8), and Proposition 2.6,

$$t_j^N - t_0^N \geq N \int_{\Omega} G(t_j \omega_j) dx \geq -\sigma_0 t_j^N \int_{\Omega} \omega_j^N dx \geq -\frac{\sigma_0 t_j^N}{\kappa n},$$

so

$$\left(\frac{t_j}{t_0}\right)^{N'} \geq \left(1 + \frac{\sigma_0}{\kappa n}\right)^{-1/(N-1)} \geq 1 - \frac{\sigma_0}{(N-1)\kappa n}.$$

This gives

$$L_1 \geq e^{-\sigma_0/(N-1)\kappa}$$

and

$$L_2 \geq \lim_{n \rightarrow \infty} \int_0^1 n e^{-n(t-t^{N'}) - \sigma_0 t^{N'}/(N-1)\kappa} dt \geq N e^{-\sigma_0/(N-1)\kappa}$$

by Proposition 2.7. So (3.17) gives

$$\beta \leq \frac{1}{\alpha^{N-1}} \left(\frac{N}{d}\right)^N \frac{1}{N e^{-\sigma_0/(N-1)\kappa} - (1 - e^{-\sigma_0/(N-1)\kappa})} \leq \frac{1}{N \alpha^{N-1}} \left(\frac{N}{d}\right)^N e^{\sigma_0/(N-1)\kappa},$$

contradicting (1.10). \square

We are now ready to prove Theorem 1.1.

Proof of Theorem 1.1. Let j_0 be as in Lemma 3.2 (ii). By Lemma 3.2 (i), $\exists R > \rho$ such that $E(R\omega_{j_0}) \leq 0$, where ρ is as in Lemma 3.1. Let

$$\Gamma = \left\{ \gamma \in C([0, 1], W_0^{1,N}(\Omega)) : \gamma(0) = 0, \gamma(1) = R\omega_{j_0} \right\}$$

be the class of paths joining the origin to $R\omega_{j_0}$, and set

$$c := \inf_{\gamma \in \Gamma} \max_{u \in \gamma([0,1])} E(u).$$

By Lemma 3.1, $c > 0$. Since the path $\gamma_0(t) = tR\omega_{j_0}$, $t \in [0, 1]$ is in Γ ,

$$c \leq \max_{u \in \gamma_0([0,1])} E(u) \leq \sup_{t \geq 0} E(t\omega_{j_0}) < \frac{1}{N} \left(\frac{\alpha_N}{\alpha}\right)^{N-1}.$$

If there are no $(PS)_c$ sequences of E , then E satisfies the $(PS)_c$ condition vacuously and hence has a critical point u at the level c by the mountain pass theorem. Then u is a solution of problem (1.1) and u is nontrivial since $c > 0$. So we may assume that E has a $(PS)_c$ sequence. Then this sequence has a subsequence that converges weakly to a nontrivial solution of problem (1.1) by Proposition 2.1. \square

4 Proof of Theorem 1.3

In this section we prove Theorem 1.3 using Theorem 2.5. We take A_0 to be the set C in Proposition 2.4 and $B_0 = \Psi_{\lambda_k}$. Since $i(S \setminus B_0) = k - 1$ by Proposition 2.3, (2.9) holds.

Lemma 4.1. *There exists a $\rho > 0$ such that $\inf E(B) > 0$, where $B = \{\rho u : u \in B_0\}$.*

Proof. As in the proof of Lemma 3.1, there exists a constant $C_\delta > 0$ such that

$$|G(t)| \leq C_\delta |t|^{N+1} e^{\alpha |t|^{N'}} \quad \text{for } |t| > \delta,$$

which together with (1.12) gives

$$G(t) \leq \frac{1}{N} (\lambda_k - \sigma_1) |t|^N + C_\delta |t|^{N+1} e^{\alpha |t|^{N'}} \quad \forall t. \quad (4.1)$$

For $u \in B_0$ and $\rho > 0$,

$$\int_{\Omega} |\rho u|^N dx \leq \frac{\rho^N}{\lambda_k} \quad (4.2)$$

and

$$\int_{\Omega} |\rho u|^{N+1} e^{\alpha |\rho u|^{N'}} dx \leq \rho^{N+1} \left(\int_{\Omega} |u|^{2(N+1)} dx \right)^{1/2} \left(\int_{\Omega} e^{2\alpha \rho^{N'} |u|^{N'}} dx \right)^{1/2}. \quad (4.3)$$

The first integral on the right-hand side of (4.3) is bounded by the Sobolev embedding theorem, and the second integral is bounded when $\rho^{N'} \leq \alpha_N/2\alpha$ by (1.4). So combining (4.1)–(4.3) gives

$$\int_{\Omega} G(\rho u) dx \leq \frac{1}{N} \left(1 - \frac{\sigma_1}{\lambda_k} \right) \rho^N + O(\rho^{N+1}) \quad \text{as } \rho \rightarrow 0.$$

Then

$$E(\rho u) \geq \frac{1}{N} \frac{\sigma_1}{\lambda_k} \rho^N + O(\rho^{N+1}),$$

and the desired conclusion follows from this for sufficiently small ρ . \square

We may assume without loss of generality that $B_d(0) \subset \Omega$. Let (ω_j) be the sequence of functions defined in (2.10).

Lemma 4.2. *We have*

$$(i) \quad E(sv) \leq 0 \quad \forall v \in A_0, s \geq 0,$$

(ii) for all $j \geq 2$,

$$\sup \{E(R\pi((1-t)v + t\omega_j)) : v \in A_0, 0 \leq t \leq 1\} \rightarrow -\infty \text{ as } R \rightarrow \infty,$$

(iii) $\exists j_0 \geq 2$ such that

$$\sup \{E(sv + t\omega_{j_0}) : v \in A_0, s, t \geq 0\} < \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1}.$$

Proof. (i) By (1.11),

$$E(u) \leq \frac{1}{N} \left[\int_{\Omega} |\nabla u|^N dx - (\lambda_{k-1} + \sigma_0) \int_{\Omega} |u|^N dx \right]. \quad (4.4)$$

For $v \in A_0$ and $s \geq 0$,

$$\int_{\Omega} |sv|^N dx \geq \frac{s^N}{\lambda_{k-1}}$$

since $A_0 \subset \Psi^{\lambda_{k-1}}$, so (4.4) gives

$$E(sv) \leq -\frac{1}{N} \frac{\sigma_0}{\lambda_{k-1}} s^N \leq 0.$$

(ii) Fix $0 < \varepsilon < \beta$. As in the proof of Lemma 3.2 (i), $\exists M_\varepsilon > 0$ such that

$$th(t) e^{\alpha|t|^{N'}} > (\beta - \varepsilon) e^{\alpha|t|^{N'}} \quad \text{for } |t| > M_\varepsilon \quad (4.5)$$

and there exists a constant $C_\varepsilon > 0$ such that

$$th(t) e^{\alpha|t|^{N'}} \geq \frac{1}{(2N-2)!} (\beta - \varepsilon) \alpha^{2N-2} t^{2N} - C_\varepsilon |t| \quad (4.6)$$

and

$$G(t) \geq \frac{2N-1}{(2N)!} (\beta - \varepsilon) \alpha^{2N-2} t^{2N} - C_\varepsilon |t| \quad (4.7)$$

for all t . Let $A_1 = \{\pi((1-t)v + t\omega_j) : v \in A_0, 0 \leq t \leq 1\}$. For $u \in A_1$ and $R > 0$, (4.7) gives

$$E(Ru) \leq \frac{R^N}{N} - \frac{2N-1}{(2N)!} (\beta - \varepsilon) \alpha^{2N-2} R^{2N} \int_{\Omega} |u|^{2N} dx + C_\varepsilon R \int_{\Omega} |u| dx.$$

The set A_1 is compact since A_0 is compact, so the first integral on the right-hand side is bounded away from zero on A_1 . Since the second integral is bounded, the desired conclusion follows.

(iii) If the conclusion is false, then it follows from (i) and (ii) that for all $j \geq 2$, there exist $v_j \in A_0$, $s_j \geq 0$, $t_j > 0$ such that

$$E(s_j v_j + t_j \omega_j) = \sup \{E(sv + t\omega_j) : v \in A_0, s, t \geq 0\} \geq \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1}.$$

Set $u_j = s_j v_j + t_j \omega_j$. Then

$$E(u_j) = \frac{1}{N} \|u_j\|^N - \int_{\Omega} G(u_j) dx \geq \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1}. \quad (4.8)$$

Moreover, $\tau u_j \in \{sv + t\omega_j : v \in A_0, s, t \geq 0\}$ for all $\tau \geq 0$ and $E(\tau u_j)$ attains its maximum at $\tau = 1$, so

$$\left. \frac{\partial}{\partial \tau} E(\tau u_j) \right|_{\tau=1} = E'(u_j) u_j = \|u_j\|^N - \int_{\Omega} u_j h(u_j) e^{\alpha |u_j|^{N'}} dx = 0. \quad (4.9)$$

Since $\|v_j\| = \|\omega_j\| = 1$ and $G(t) \geq 0$ for all t by (1.11), (4.8) gives

$$s_j + t_j \geq t_0,$$

where

$$t_0 = \left(\frac{\alpha_N}{\alpha} \right)^{(N-1)/N}.$$

First we show that $s_j \rightarrow 0$ and $t_j \rightarrow t_0$ as $j \rightarrow \infty$.

Combining (4.8) with (1.11) gives

$$\|s_j v_j + t_j \omega_j\|^N \geq (\lambda_{k-1} + \sigma_0) \int_{\Omega} |s_j v_j + t_j \omega_j|^N dx + t_0^N.$$

Set $\tau_j = s_j/t_j$. Then

$$\|\tau_j v_j + \omega_j\|^N \geq (\lambda_{k-1} + \sigma_0) \int_{\Omega} |\tau_j v_j + \omega_j|^N dx + \left(\frac{t_0}{t_j} \right)^N. \quad (4.10)$$

Since (v_j) is bounded in $C^1(\overline{\Omega})$, Proposition 2.6 gives

$$\begin{aligned} \|\tau_j v_j + \omega_j\|^N &\leq \int_{\Omega} (\tau_j |\nabla v_j| + |\nabla \omega_j|)^N dx = \tau_j^N \int_{\Omega} |\nabla v_j|^N dx + \int_{\Omega} |\nabla \omega_j|^N dx \\ &\quad + \sum_{m=1}^{N-1} \binom{N}{m} \tau_j^{N-m} \int_{\Omega} |\nabla v_j|^{N-m} |\nabla \omega_j|^m dx \leq \tau_j^N + 1 + c_1 \sum_{m=1}^{N-1} \frac{\tau_j^{N-m}}{(\log j)^{m/N}} \end{aligned}$$

and

$$\begin{aligned} \int_{\Omega} |\tau_j v_j + \omega_j|^N dx &\geq \int_{\Omega} (\tau_j |v_j| - \omega_j)^N dx = \tau_j^N \int_{\Omega} |v_j|^N dx \\ &+ \sum_{m=1}^N (-1)^m \binom{N}{m} \tau_j^{N-m} \int_{\Omega} |v_j|^{N-m} \omega_j^m dx \geq \frac{\tau_j^N}{\lambda_{k-1}} - c_2 \sum_{m=1}^N \frac{\tau_j^{N-m}}{(\log j)^{m/N}} \end{aligned}$$

for some constants $c_1, c_2 > 0$. So (4.10) gives

$$\frac{\sigma_0}{\lambda_{k-1}} \tau_j^N + \left(\frac{t_0}{t_j} \right)^N \leq 1 + c_3 \sum_{m=1}^N \frac{\tau_j^{N-m}}{(\log j)^{m/N}} \quad (4.11)$$

for some constant $c_3 > 0$, which implies that (τ_j) is bounded and

$$\liminf_{j \rightarrow \infty} t_j \geq t_0. \quad (4.12)$$

Next combining (4.9) with (4.5) and (4.6) gives

$$\begin{aligned} \|u_j\|^N &= \int_{\{|u_j| > M_\varepsilon\}} u_j h(u_j) e^{\alpha |u_j|^{N'}} dx + \int_{\{|u_j| \leq M_\varepsilon\}} u_j h(u_j) e^{\alpha |u_j|^{N'}} dx \\ &\geq (\beta - \varepsilon) \int_{\{|u_j| > M_\varepsilon\}} e^{\alpha |u_j|^{N'}} dx - C_\varepsilon \int_{\{|u_j| \leq M_\varepsilon\}} |u_j| dx. \end{aligned} \quad (4.13)$$

For $|x| \leq d/j$,

$$|u_j| \geq t_j \omega_j - s_j |v_j| \geq \frac{t_j}{\omega_{N-1}^{1/N}} \left[(\log j)^{(N-1)/N} - c_4 \tau_j \right]$$

for some constant $c_4 > 0$, and the last expression is greater than M_ε for all sufficiently large j since (τ_j) is bounded and $\liminf t_j > 0$. So

$$\begin{aligned} \int_{\{|u_j| > M_\varepsilon\}} e^{\alpha |u_j|^{N'}} dx &\geq e^{\alpha t_j^{N'} [(\log j)^{(N-1)/N} - c_4 \tau_j]^{N'}} / \omega_{N-1}^{1/(N-1)} \int_{\{|x| \leq d/j\}} dx \\ &= \frac{\omega_{N-1} d^N}{N} j^{\alpha [t_j^{N'} (1 - c_4 \tau_j / (\log j)^{(N-1)/N})^{N'} - t_0^{N'}] / \omega_{N-1}^{1/(N-1)}} \end{aligned}$$

for large j . On the other hand,

$$\int_{\{|u_j| \leq M_\varepsilon\}} |u_j| dx \leq \int_{\Omega} (s_j |v_j| + t_j \omega_j) dx \leq c_5 t_j \left[\tau_j + \frac{1}{(\log j)^{1/N}} \right]$$

for some constant $c_5 > 0$ by Proposition 2.6. So (4.13) gives

$$(\beta - \varepsilon) j^\alpha [t_j^{N'} (1 - c_4 \tau_j / (\log j)^{(N-1)/N})^{N'} - t_0^{N'}] / \omega_{N-1}^{1/(N-1)} \leq \frac{N t_j^N (\tau_j + 1)^N}{\omega_{N-1} d^N} + c_6 t_j \left[\tau_j + \frac{1}{(\log j)^{1/N}} \right] \quad (4.14)$$

for some constant $c_6 > 0$. Since (τ_j) is bounded, it follows from this that

$$\limsup_{j \rightarrow \infty} t_j \leq t_0,$$

which together with (4.12) shows that $t_j \rightarrow t_0$. Then (4.11) implies that $\tau_j \rightarrow 0$, so $s_j = \tau_j t_j \rightarrow 0$.

Now we show that there exists a constant $c > 0$ depending only on Ω , α , and k such that

$$\beta \leq \frac{1}{\alpha^{N-1}} \left(\frac{N}{d} \right)^N e^{c/\sigma_0^{N-1}}. \quad (4.15)$$

The right-hand side of (4.14) goes to $(N/d)^N / \alpha^{N-1}$ as $j \rightarrow \infty$. If $\beta \leq (N/d)^N / \alpha^{N-1}$, then we may take any $c > 0$, so suppose $\beta > (N/d)^N / \alpha^{N-1}$. Then for $\varepsilon < \beta - (N/d)^N / \alpha^{N-1}$ and all sufficiently large j , (4.14) gives $j^\alpha [t_j^{N'} (1 - c_4 \tau_j / (\log j)^{(N-1)/N})^{N'} - t_0^{N'}] / \omega_{N-1}^{1/(N-1)} \leq 1$, so

$$\frac{t_0}{t_j} \geq 1 - \frac{c_4 \tau_j}{(\log j)^{(N-1)/N}}.$$

Combining this with (4.11) gives

$$\frac{\sigma_0}{\lambda_{k-1}} \tau_j^N - \frac{N c_4 \tau_j}{(\log j)^{(N-1)/N}} \leq c_3 \sum_{m=1}^N \frac{\tau_j^{N-m}}{(\log j)^{m/N}},$$

so

$$\sigma_0 \tau_j^N \leq c_7 \sum_{m=1}^N \frac{\tau_j^{N-m}}{(\log j)^{m/N}}$$

for some constant $c_7 > 0$. Set $\tilde{\tau}_j = \tau_j (\log j)^{1/N}$. Then

$$\sigma_0 \tilde{\tau}_j^N \leq c_7 \sum_{m=1}^N \tilde{\tau}_j^{N-m}. \quad (4.16)$$

We claim that

$$\tilde{\tau}_j \leq \frac{c_8}{\sigma_0} \quad (4.17)$$

for some constant $c_8 > 0$. Taking σ_0 smaller in (1.11) if necessary, we may assume that $\sigma_0 \leq 1$. So if $\tilde{\tau}_j < 1$, then (4.17) holds with $c_8 = 1$, so suppose $\tilde{\tau}_j \geq 1$. Then (4.16) gives (4.17) with $c_8 = Nc_7$. Now (4.11) gives

$$\left(\frac{t_0}{t_j}\right)^N \leq 1 + \frac{c_3}{\log j} \sum_{m=1}^N \tilde{\tau}_j^{N-m} \leq 1 + \frac{c_9}{\sigma_0^{N-1} \log j}$$

for some constant $c_9 > 0$, so

$$\left(\frac{t_0}{t_j}\right)^{N'} \leq \left(1 + \frac{c_9}{\sigma_0^{N-1} \log j}\right)^{1/(N-1)} \leq 1 + \frac{c_9}{\sigma_0^{N-1} \log j}.$$

Then

$$\begin{aligned} t_j^{N'} \left[1 - \frac{c_4 \tau_j}{(\log j)^{(N-1)/N}}\right]^{N'} - t_0^{N'} &= t_j^{N'} \left[\left(1 - \frac{c_4 \tilde{\tau}_j}{\log j}\right)^{N'} - \left(\frac{t_0}{t_j}\right)^{N'}\right] \\ &\geq t_j^{N'} \left[\left(1 - \frac{c_{10}}{\sigma_0 \log j}\right)^{N'} - \left(1 + \frac{c_9}{\sigma_0^{N-1} \log j}\right)\right] \geq -t_j^{N'} \left(\frac{N' c_{10}}{\sigma_0 \log j} + \frac{c_9}{\sigma_0^{N-1} \log j}\right) \\ &\geq -\frac{c_{11}}{\sigma_0^{N-1} \log j} \end{aligned}$$

for some constants $c_{10}, c_{11} > 0$, so

$$j^{\alpha} [t_j^{N'} (1 - c_4 \tau_j / (\log j)^{(N-1)/N})^{N'} - t_0^{N'}] / \omega_{N-1}^{1/(N-1)} \geq j^{-c/\sigma_0^{N-1} \log j} = e^{-c/\sigma_0^{N-1}}$$

for some constant $c > 0$. Combining this with (4.14) and passing to the limit gives

$$(\beta - \varepsilon) e^{-c/\sigma_0^{N-1}} \leq \frac{1}{\alpha^{N-1}} \left(\frac{N}{d}\right)^N,$$

and letting $\varepsilon \rightarrow 0$ gives (4.15). □

We are now ready to prove Theorem 1.3.

Proof of Theorem 1.3. Let $j_0 \geq 2$ be as in Lemma 4.2 (iii). By Lemma 4.2 (ii), $\exists R > \rho$ such that

$$\sup \{E(R\pi((1-t)v + t\omega_{j_0})) : v \in A_0, 0 \leq t \leq 1\} \leq 0, \quad (4.18)$$

where $\rho > 0$ is as in Lemma 4.1. Let

$$A = \{sv : v \in A_0, 0 \leq s \leq R\} \cup \{R\pi((1-t)v + t\omega_{j_0}) : v \in A_0, 0 \leq t \leq 1\},$$

$$X = \{sv + t\omega_{j_0} : v \in A_0, s, t \geq 0, \|sv + t\omega_{j_0}\| \leq R\}.$$

Combining Lemma 4.2 (i), (4.18), and Lemma 4.1 gives

$$\sup E(A) \leq 0 < \inf E(B), \quad (4.19)$$

while Lemma 4.2 (iii) gives

$$\sup E(X) \leq \sup \{E(sv + t\omega_{j_0}) : v \in A_0, s, t \geq 0\} < \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1}. \quad (4.20)$$

Let

$$\Gamma = \{\gamma \in C(X, W) : \gamma(X) \text{ is closed and } \gamma|_A = id_A\},$$

and set

$$c := \inf_{\gamma \in \Gamma} \sup_{u \in \gamma(X)} E(u).$$

By Theorem 2.5, $\inf E(B) \leq c \leq \sup E(X)$, and E has a $(PS)_c$ sequence. By (4.19) and (4.20),

$$0 < c < \frac{1}{N} \left(\frac{\alpha_N}{\alpha} \right)^{N-1},$$

so a subsequence of this $(PS)_c$ sequence converges weakly to a nontrivial solution of problem (1.1) by Proposition 2.1. \square

5 Competing interests declaration

The authors declare no competing interests.

References

- [1] Adimurthi. Existence of positive solutions of the semilinear Dirichlet problem with critical growth for the n -Laplacian. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 17(3):393–413, 1990.
- [2] Adimurthi and S. L. Yadava. Bifurcation results for semilinear elliptic problems with critical exponent in \mathbf{R}^2 . *Nonlinear Anal.*, 14(7):607–612, 1990.

- [3] D. G. de Figueiredo, O. H. Miyagaki, and B. Ruf. Elliptic equations in \mathbf{R}^2 with nonlinearities in the critical growth range. *Calc. Var. Partial Differential Equations*, 3(2):139–153, 1995.
- [4] D. G. de Figueiredo, O. H. Miyagaki, and B. Ruf. Corrigendum: “Elliptic equations in \mathbf{R}^2 with nonlinearities in the critical growth range”. *Calc. Var. Partial Differential Equations*, 4(2):203, 1996.
- [5] Djairo G. de Figueiredo, João Marcos do Ó, and Bernhard Ruf. On an inequality by N. Trudinger and J. Moser and related elliptic equations. *Comm. Pure Appl. Math.*, 55(2):135–152, 2002.
- [6] Djairo G. de Figueiredo, João Marcos do Ó, and Bernhard Ruf. Elliptic equations and systems with critical Trudinger-Moser nonlinearities. *Discrete Contin. Dyn. Syst.*, 30(2):455–476, 2011.
- [7] Marco Degiovanni and Sergio Lancelotti. Linking solutions for p -Laplace equations with nonlinearity at critical growth. *J. Funct. Anal.*, 256(11):3643–3659, 2009.
- [8] Edward R. Fadell and Paul H. Rabinowitz. Generalized cohomological index theories for Lie group actions with an application to bifurcation questions for Hamiltonian systems. *Invent. Math.*, 45(2):139–174, 1978.
- [9] João Marcos B. do Ó. Semilinear Dirichlet problems for the N -Laplacian in \mathbb{R}^N with nonlinearities in the critical growth range. *Differential Integral Equations*, 9(5):967–979, 1996.
- [10] J. Moser. A sharp form of an inequality by N. Trudinger. *Indiana Univ. Math. J.*, 20:1077–1092, 1970/71.
- [11] Kanishka Perera. Nontrivial critical groups in p -Laplacian problems via the Yang index. *Topol. Methods Nonlinear Anal.*, 21(2):301–309, 2003.
- [12] Kanishka Perera, Ravi P. Agarwal, and Donal O’Regan. *Morse theoretic aspects of p -Laplacian type operators*, volume 161 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [13] Paul H. Rabinowitz. Some critical point theorems and applications to semilinear elliptic partial differential equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 5(1):215–223, 1978.
- [14] Neil S. Trudinger. On imbeddings into Orlicz spaces and some applications. *J. Math. Mech.*, 17:473–483, 1967.
- [15] Yang Yang and Kanishka Perera. N -Laplacian problems with critical Trudinger-Moser nonlinearities. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 16(4):1123–1138, 2016.