

ARAZY-CWIKEL PROPERTY FOR QUASI-BANACH COUPLES

SERGEY V. ASTASHKIN AND PER G. NILSSON

ABSTRACT. The main result of this paper establishes that the known Arazy-Cwikel property holds for classes of uniformly K -monotone spaces in the quasi-Banach setting provided that the initial couple is mutually closed. As a consequence, we get that the class of all quasi-Banach K -spaces (i.e., interpolation spaces which are described by the real K -method) with respect to an arbitrary mutually closed Banach couple enjoys the Arazy-Cwikel property. Another consequence complements some previous results by Bykov and Ovchinnikov, showing that this property holds also for the class of all interpolation quasi-Banach spaces with respect to a quasi-Banach couple whenever all the couples involved have the uniform Calderón-Mityagin property. We apply these results to some classical families of spaces.

1. INTRODUCTION

According to the classical result of the interpolation theory of operators, which was obtained independently by Calderón [9] and Mityagin [31], a Banach function space X on an arbitrary underlying measure space is an interpolation space with respect to the couple (L^1, L^∞) on that measure space if and only if the following monotonicity property¹ holds: if $f \in X$, $g \in L^1 + L^\infty$ and

$$\int_0^t g^*(s) \, ds \leq \int_0^t f^*(s) \, ds, \quad t > 0$$

(where h^* denotes the nonincreasing left-continuous rearrangement of $|h|$), then $g \in X$ and $\|g\|_X \leq \|f\|_X$ (for all undefined terminology see the next section). Since Peetre [37, 38] had proved (cf. also a similar result due independently to Oklander [34], and cf. also [23, pp. 158–159]) that the functional $t \mapsto \int_0^t f^*(s) \, ds$ is in fact the K -functional of the function $f \in L^1 + L^\infty$ for the couple (L^1, L^∞) , the results of [9, 31] naturally led to the introduction of the following definition. An intermediate Banach space X with respect to a Banach couple $\overline{X} = (X_0, X_1)$

Date: January 21, 2022.

2010 Mathematics Subject Classification. Primary 46B70; Secondary 46E30, 46M35.

Key words and phrases. interpolation space, quasi-Banach space, p -convex space, K -functional, Calderón-Mityagin property, Lions-Peetre K -spaces.

The work of the first author was completed as a part of the implementation of the development program of the Scientific and Educational Mathematical Center Volga Federal District, agreement no. 075-02-2021-1393.

¹Here we are using Calderón's terminology. Mityagin formulates this result somewhat differently.

is said to be *K-monotone* with respect to $\overline{X}$ if whenever elements $x \in X$ and $y \in \Sigma(\overline{X})$ satisfy

$$K(t, y; \overline{X}) \leq K(t, x; \overline{X}), \text{ for all } t > 0,$$

it follows that $y \in X$. Moreover, if each interpolation Banach space with respect to a Banach couple $\overline{X}$ is *K-monotone*, $\overline{X}$ is said to have the Calderón-Mityagin property. As is well known now this property is shared by many Banach couples and, in particular, by each couple (L^p, L^q) , $1 \leq p < q \leq \infty$ (see e.g. [42] or [12]).

Using the Calderón-Mityagin property of couples of L^p -spaces, Arazy and Cwikel proved, in [1], that for all $1 \leq p < q \leq \infty$ and for each underlying measure space a Banach function space X is an interpolation space with respect to the couple (L^p, L^q) if and only if X is such a space with respect to each of the couples (L^1, L^q) and (L^p, L^∞) , or more formally

$$(1.1) \quad \text{Int}(L^p, L^q) = \text{Int}(L^1, L^q) \cap \text{Int}(L^p, L^\infty)$$

($\text{Int}(X_0, X_1)$ denotes the class of all interpolation Banach spaces with respect to a Banach couple $\overline{X} = (X_0, X_1)$). This result somewhat resembles the well-known interpolation Boyd theorem together with its one-sided refinements. Indeed, the latter theorem reads that, for $1 \leq p < q \leq \infty$, the condition $1/q < \alpha(X) \leq \beta(X) < 1/p$ ($\alpha(X)$ and $\beta(X)$ are the Boyd indices of X) ensures that a rearrangement invariant space X belongs to the set $\text{Int}(L^p, L^q)$ [25, Theorem 2.b.11], while in the extreme case when $1 = p < q < \infty$ (resp. $1 < p < q = \infty$) the same result holds for each X such that $X \in \text{Int}(L^1, L^\infty)$ provided that only the one-sided estimate $\alpha(X) > 1/q$ (resp. $\beta(X) < 1/p$) is valid (see [26] and [3]).

Later on, Bykov and Ovchinnikov obtained a result similar to (1.1) for families of interpolation spaces, corresponding to weighted couples of shift-invariant ideal sequence spaces [7]. Moreover, let X_0 and X_1 be classical Lions-Peetre *K*-spaces with respect to a Banach couple (A_0, A_1) , i.e., $X_0 = (A_0, A_1)_{\alpha_0, p_0}$, $X_1 = (A_0, A_1)_{\alpha_1, p_1}$, where $0 < \alpha_i < 1$, $1 \leq p_i \leq \infty$, $i = 0, 1$. Then, from [7, Theorem 2.9], in particular, it follows that for all $0 < \theta < \eta < 1$ and $0 < p, q \leq \infty$

$$(1.2) \quad \text{Int}(\overline{X}_{\theta, p}, \overline{X}_{\eta, q}) = \text{Int}(X_0, \overline{X}_{\eta, q}) \cap \text{Int}(\overline{X}_{\theta, p}, X_1).$$

Recently some results of Arazy-Cwikel type were proved also in the quasi-Banach setting. So, relation (1.1) has been extended to the range $0 \leq p < q \leq \infty$ and the classes of all interpolation quasi-Banach spaces with respect to the couples of L^p -spaces of measurable functions on the semi-axis $(0, \infty)$ with the Lebesgue measure [8] and the couples of sequence l^p -spaces [2] (the definition of the extreme spaces L^0 and l^0 see also in [8] and [2]).

On the other hand, already in [1], it was indicated that formula (1.2) fails to be valid for *all* Banach couples (X_0, X_1) . In contrast to that, the main result of this paper establishes that the Arazy-Cwikel property holds if we consider classes of

uniformly K -monotone (rather than of all interpolation) spaces even in the quasi-Banach setting provided that the initial couple is mutually closed. Since in the Banach case classes of K -monotone and K spaces coincide [6, Theorem 4.1.11], we immediately obtain that the class of all quasi-Banach K -spaces (i.e., interpolation spaces which are described by the real K -method) with respect to an arbitrary mutually closed Banach couple enjoys the Arazy-Cwikel property. Observe that the latter result has been announced, without a proof, in [13] (see also [6, p. 672]).

By the well-known theorem due to Cwikel [12, Theorem 1], for every Banach couple $\overline{X} = (X_0, X_1)$ and all $0 < \theta_0, \theta_1 < 1$, $1 \leq p_0, p_1 \leq \infty$, the couple $(\overline{X}_{\theta_0, p_0}, \overline{X}_{\theta_1, p_1})$ is uniformly K -monotone. Hence, the above-mentioned result of [7] suggests that equality (1.2) holds for any uniformly K -monotone Banach couple. Indeed, this statement is another consequence of our main theorem. Moreover, we show that (1.2) is valid for an arbitrary quasi-Banach couple $\overline{X} = (X_0, X_1)$ and all $0 < \theta_0, \theta_1 < 1$, $0 < p_0, p_1 \leq \infty$ whenever all the couples involved in (1.2) have the uniform Calderón-Mityagin property.

Let us describe briefly the content of the paper. In Section 2, we give preliminaries with basic definitions and notation. We address some properties of quasi-Banach spaces and lattices, which relate to their convexity, and necessary definitions and results from interpolation theory. In the next section we collect some auxiliary results, many of which are apparently to some extent known. Section 4 contains the main results of the paper (Theorem 4.1 and Corollaries 4.2 and 4.4) related to the Arazy-Cwikel property for the Lions-Peetre K -spaces $X_{\theta, p}$, $0 < \theta < 1$, $0 < p \leq \infty$, in the quasi-Banach setting. In conclusion, in Section 5, we apply these results to some classical families of spaces.

The authors would like to thank Professor Michael Cwikel, whose work and insight are behind the results announced in the paper [13].

2. PRELIMINARIES

2.1. Quasi-Banach spaces and lattices. Recall that a (real) quasi-Banach space X is a complete real vector space whose topology is given by a quasi-norm $x \mapsto \|x\|$ which satisfy the conditions: $\|x\| > 0$ if $x \neq 0$, $\|\alpha x\| = |\alpha| \|x\|$, $\alpha \in \mathbb{R}$, $x \in X$, and $\|x_1 + x_2\|_X \leq C(\|x_1\| + \|x_2\|)$ for some $C > 0$ and all $x_1, x_2 \in X$.

If a quasi-Banach space X is additionally a vector lattice such that $\|x\| \leq \|y\|$ whenever $|x| \leq |y|$, we say that X is a quasi-Banach lattice (see [25], [30].)

Let X be a quasi-Banach lattice, $x_k \in X$, $k = 1, 2, \dots, n$. Then, any element of the form $(\sum_{k=1}^n |x_k|^p)^{1/p}$, $0 < p < \infty$, can be defined by means of a "homogeneous functional calculus" in the quasi-Banach setting exactly as in the case of Banach lattices (cf. [25, pp. 40-41], [21], [10]). A quasi-Banach lattice X is said to be (*lattice*) *p -convex*, $0 < p \leq \infty$, if for some constant M and any $x_k \in X$,

$k = 1, 2, \dots, n$ we have

$$(2.1) \quad \left\| \left(\sum_{k=1}^n |x_k|^p \right)^{1/p} \right\|_X \leq M \left(\sum_{k=1}^n \|x_k\|_X^p \right)^{1/p} \quad \text{if } p < \infty,$$

with the usual modification for $p = \infty$ (see e.g. [25] for $p \geq 1$ and [21] for $p > 0$). By $M^{(p)}(X)$ we will denote the minimal value of M satisfying (2.1).

Next, we will use the so-called p -convexification procedure which just an abstract description of the mapping $f \mapsto |f|^p \text{sign } f$ from $L_r(\mu)$, $0 < r < \infty$, into $L_{rp}(\mu)$. Note that in a general lattice X there is no meaning to the symbol x^p that makes us to introduce new algebraic operations in X (see [25, pp. 53-54]).

Let X be a quasi-Banach lattice with the algebraic operations denoted by $+$ and $\cdot$ and let $p > 0$. For every $x, y \in X$ and $\alpha \in \mathbb{R}$ we define

$$x \oplus y := (x^{1/p} + y^{1/p})^p \quad \text{and} \quad \alpha \odot x := \alpha^p \cdot x,$$

where α^p is $|\alpha|^p \text{sign } \alpha$. Then, the set X , endowed with the operations $\oplus$, $\odot$ and the same order as in X is a vector lattice, which denoted by $X^{(p)}$. Moreover, $\|x\|_{X^{(p)}} := \|x\|_X^{1/p}$ is a lattice norm on $X^{(p)}$ and $(X^{(p)}, \|\cdot\|_{X^{(p)}})$ is a quasi-Banach lattice for every $0 < p < \infty$ [10, Proposition 1.2]. One can easily check also that if X is r -convex with the constant $M^{(r)}$, then $X^{(p)}$ is pr -convex with the same constant.

Note that if X is a quasi-Banach function lattice defined on some measure space, $X^{(p)}$ can be identified with the space of functions f such that $f^p := |f|^p \text{sign } f \in X$ equipped with the norm $\|f\|_{X^{(p)}} = \| |f|^p \|_X^{1/p}$.

2.2. Interpolation of quasi-Banach spaces. Let us recall some basic constructions and definitions related to the interpolation theory of operators. For more detailed information we refer to [4, 5, 6, 24, 36].

In this paper we are mainly concerned with interpolation within the class of quasi-Banach spaces, while the linear bounded operators are considered as the corresponding morphisms. A pair $\overline{X} = (X_0, X_1)$ of quasi-Banach spaces is called a *quasi-Banach couple* if X_0 and X_1 are both linearly and continuously embedded in some Hausdorff topological vector space.

For each quasi-Banach couple $\overline{X} = (X_0, X_1)$ we define the *intersection* $\Delta(\overline{X}) = X_0 \cap X_1$ and the *sum* $\Sigma(\overline{X}) = X_0 + X_1$ as the quasi-Banach spaces equipped with the quasi-norms

$$\|x\|_{\Delta(\overline{X})} := \max \{ \|x\|_{X_0}, \|x\|_{X_1} \}$$

and

$$\|x\|_{\Sigma(\overline{X})} := \inf \{ \|x_0\|_{X_0} + \|x_1\|_{X_1} : x = x_0 + x_1, x_i \in X_i, i = 0, 1 \},$$

respectively.

A quasi-Banach space X is called an *intermediate space* for a quasi-Banach couple $\overline{X} = (X_0, X_1)$ if the continuous inclusions $\Delta(\overline{X}) \subset X \subset \Sigma(\overline{X})$ hold. The set of intermediate spaces with respect to $\overline{X}$ will be denoted by $I(\overline{X})$ or $I(X_0, X_1)$.

If $\overline{X} = (X_0, X_1)$ is a quasi-Banach couple, then we let $\mathfrak{L}(\overline{X})$ (or $\mathfrak{L}(X_0, X_1)$) denote the space of all linear operators $T : \Sigma(\overline{X}) \rightarrow \Sigma(\overline{X})$ that are bounded on X_i , $i = 0, 1$, equipped with the quasi-norm

$$\|T\|_{\mathfrak{L}(\overline{X})} := \max_{i=0,1} \|T\|_{X_i \rightarrow X_i}.$$

Let $\overline{X} = (X_0, X_1)$ be a quasi-Banach couple and let $X \in I(\overline{X})$. Then, X is said to be an *interpolation space* with respect to the couple $\overline{X}$, if every operator $T \in \mathfrak{L}(\overline{X})$ is bounded on X . Recall that, by the Aoki-Rolewicz theorem (see e.g. [5, Lemma 3.10.1]), every quasi-Banach space is a F -space (i.e., the topology in that space is generated by a complete invariant metric). In particular, this applies to the space $\mathfrak{L}(\overline{X})$ which is obviously a quasi-Banach space with respect to the quasi-norm $T \mapsto \|T\|_{\mathfrak{L}(\overline{X})}$ and also with respect to the quasi-norm $T \mapsto \max(\|T\|_{\mathfrak{L}(\overline{X})}, \|T\|_{X \rightarrow X})$ whenever the quasi-Banach space X is an interpolation space with respect to the quasi-Banach couple $\overline{X} = (X_0, X_1)$. As is well known (see e.g. [40, Theorem 2.2.15]), the Closed Graph Theorem and the equivalent Bounded Inverse Theorem (see e.g. [40, Corollary 2.2.12]) hold for F -spaces. Therefore, exactly the same reasoning as required for the Banach case (see Theorem 2.4.2 of [5, p. 28]) shows that, if X is an interpolation quasi-Banach space with respect to a quasi-Banach couple $\overline{X} = (X_0, X_1)$, then there exists a constant $C > 0$ such that for every $T \in \mathfrak{L}(\overline{X})$ we have $\|T\|_{X \rightarrow X} \leq C\|T\|_{\mathfrak{L}(\overline{X})}$. The least constant C , satisfying the last inequality for all such T , is called the *interpolation constant* of X with respect to the couple $\overline{X}$. The collection of all interpolation spaces with respect to the couple $\overline{X}$ will be denoted by $Int(\overline{X})$ (or $Int(X_0, X_1)$).

One of the most important ways of constructing interpolation spaces is based on use of the *Peetre K -functional*, which is defined for an arbitrary quasi-Banach couple $\overline{X} = (X_0, X_1)$, for every $x \in \Sigma(\overline{X})$ and each $t > 0$ as follows:

$$(2.2) \quad K(t, x; \overline{X}) := \inf\{\|x_0\|_{X_0} + t\|x_1\|_{X_1} : x = x_0 + x_1, x_i \in X_i, i = 0, 1\}.$$

For each fixed $x \in \Sigma(\overline{X})$ one can easily show that the function $t \mapsto K(t, x; \overline{X})$ is continuous, non-decreasing, concave and non-negative on $(0, \infty)$ [5, Lemma 3.1.1].

Let X be an intermediate quasi-Banach space with respect to a quasi-Banach couple $\overline{X} = (X_0, X_1)$. Then, X is said to be a *K -monotone space* with respect to the couple $\overline{X}$ if whenever elements $x \in X$ and $y \in \Sigma(\overline{X})$ satisfy

$$K(t, y; \overline{X}) \leq K(t, x; \overline{X}), \text{ for all } t > 0,$$

it follows that $y \in X$. If additionally $\|y\|_X \leq C\|x\|_X$, for a constant C which does not depend on x and y , then we say that X is a *uniformly K -monotone space*

with respect to the couple $\overline{X}$. The infimum of all constants C with this property is referred as the *K-monotonicity constant* of X . Clearly, each K -monotone space with respect to the couple $\overline{X}$ is an interpolation space with respect to this couple. As was already defined in Introduction, a couple $\overline{X}$ has the Calderón-Mityagin property whenever every interpolation quasi-Banach space with respect to $\overline{X}$ is K -monotone. The collection of all uniformly K -monotone spaces with respect to the couple $\overline{X}$ will be denoted by $Int^{KM}(\overline{X})$ (or $Int^{KM}(X_0, X_1)$).

Suppose E is a quasi-Banach function lattice on $(0, \infty)$ (with respect to the usual Lebesgue measure and a.e. order) and $w : (0, \infty) \rightarrow \mathbb{R}$ is a nonnegative measurable function. Then, $E(w)$ is the weighted quasi-Banach function lattice with the norm $\|x\|_{E(w)} := \|xw\|_E$. In particular, in what follows, by $\overline{L^\infty}$ we denote the couple $(L^\infty, L^\infty(1/t))$.

In particular, if $E = L^p(t^{-\theta}, \frac{dt}{t})$ (i.e., the weighted space $L^p(t^{-\theta})$ on $(0, \infty)$ equipped with the measure $\frac{dt}{t}$), where $0 < \theta < 1, 0 < p \leq \infty$, we get the classical Lions-Peetre K -spaces $\overline{X}_{\theta,p}$ endowed with the quasi-norms

$$\|x\|_{\overline{X}_{\theta,p}} := \left(\int_0^\infty (K(t, x; \overline{X}) t^{-\theta})^p \frac{dt}{t} \right)^{1/p}$$

(with usual modification if $p = \infty$), see [5].

One of the most important properties of the real K -method is the following reiteration theorem due to Brudnyi-Kruglyak (see e.g. [6, Theorem 3.3.24]). If E_0, E_1 are quasi-Banach function lattices such that $E_0, E_1 \in Int(\overline{L^\infty})$, then for every quasi-Banach couple $\overline{X} = (X_0, X_1)$ we have

$$(2.3) \quad K(t, x; \overline{X}_{E_0:K}, \overline{X}_{E_1:K}) \cong K(t, K(\cdot, x; X_0, X_1); E_0, E_1), \quad t > 0,$$

with some constants independent of $x \in \overline{X}_{E_0:K} + \overline{X}_{E_1:K}$. This implies that for every quasi-Banach function lattice F on $(0, \infty)$

$$(2.4) \quad (\overline{X}_{E_0:K}, \overline{X}_{E_1:K})_{F:K} = \overline{X}_{E:K},$$

where $E := (E_0, E_1)_{F:K}$.

If $\overline{X} = (X_0, X_1)$ is a quasi-Banach couple and X is an intermediate space with respect to $\overline{X}$, then the *relative closure* X^c of X consists of all $x \in \Sigma(\overline{X})$ for which there exists a bounded sequence $\{x_n\} \subset X$, converging to x in $\Sigma(\overline{X})$. The norm in X^c is taken as the infimum of all bounds of such sequences in X . Note that $K(t, x; \overline{X}) = K(t, x; \overline{X}^c)$ for all $x \in \Sigma(\overline{X})$ and $t > 0$, where $\overline{X}^c = (X_0^c, X_1^c)$ (see [11, Lemma 2]). We will say that a space $X \in I(\overline{X})$ is *relatively closed* in $\Sigma(\overline{X})$ whenever $X = X^c$ with equivalence of norms². A quasi-Banach couple is said to be *mutually closed* (or *Gagliardo couple*) if both spaces X_0 and X_1 are relatively closed in $\Sigma(\overline{X})$.

²This differs from the usual definition saying that a space X is relatively closed in $\Sigma(\overline{X})$ if $X^c = X$ isometrically (see, for instance, [24, Definition I.1.4] or [6, Definition 2.2.16]).

Clearly, for every quasi-Banach couple $\overline{X} = (X_0, X_1)$ we have $X_0 \subset \overline{X}_{L^\infty:K}$ and $X_1 \subset \overline{X}_{L^\infty(1/t):K}$ with constant 1. Moreover, a couple $\overline{X} = (X_0, X_1)$ is mutually closed if and only if the opposite embeddings $\overline{X}_{L^\infty:K} \subset X_0$ and $\overline{X}_{L^\infty(1/t):K} \subset X_1$ hold, i.e.,

$$\|x\|_{X_0} \cong \sup_{t>0} K(t, x; X_0, X_1) \quad \text{and} \quad \|x\|_{X_1} \cong \sup_{t>0} \frac{1}{t} K(t, x; X_0, X_1)$$

with constants independent of $x \in (X_0, X_1)_{L^\infty:K}$ and $x \in (X_0, X_1)_{L^\infty(1/t):K}$, respectively (see, for instance, [6, Lemma 2.2.21, p.123] or [36, p. 384]).

Let $\overline{X} = (X_0, X_1)$ be a quasi-Banach couple and let $x \in X_0 + X_1$, $x \neq 0$. The orbit $\text{Orb}(x; \overline{X})$ of x with respect to the class of operators $\mathfrak{L}(\overline{X})$ is the linear space $\{Tx : T \in \mathfrak{L}(\overline{X})\}$, which is equipped with the quasi-norm defined by

$$\|y\|_{\text{Orb}(x)} := \inf \left\{ \|T\|_{\mathfrak{L}(\overline{X})} : y = Tx, T \in \mathfrak{L}(\overline{X}) \right\}.$$

Since any orbit $\text{Orb}(x; \overline{X})$ can be regarded as a quotient of the quasi-Banach space $\mathfrak{L}(\overline{X})$, it is a quasi-Banach space itself. If for every nonzero $x \in X_0 + X_1$ there exists a linear functional $x^* \in (X_0 + X_1)^*$ with $\langle x, x^* \rangle \neq 0$ then $X_0 \cap X_1$ is contained in $\text{Orb}(x; X_0, X_1)$ continuously (see e.g. [36, Section 1.6, p. 368]). It is easy to see that then, moreover, each orbit $\text{Orb}(x; \overline{X})$ is an interpolation space between X_0 and X_1 .

A similar notion may be defined also by using the K -functional. If $\overline{X} = (X_0, X_1)$ is a quasi-Banach couple, then the K -orbit $\text{Orb}^K(x; \overline{X})$ of an element $x \in \Sigma(\overline{X})$, $x \neq 0$, is the space of all $y \in \Sigma(\overline{X})$ such that the quasi-norm

$$\|y\|_{\text{Orb}^K(x)} := \sup_{t>0} \frac{K(t, y; \overline{X})}{K(t, x; \overline{X})}$$

is finite.

It is obvious that for every quasi-Banach couple (X_0, X_1) and each $x \in X_0 + X_1$ we have the continuous embedding $\text{Orb}(x; \overline{X}) \subset \text{Orb}^K(x; \overline{X})$ with constant 1. A quasi-Banach couple $\overline{X} = (X_0, X_1)$ has the Calderón-Mityagin property if and only if the opposite embedding $\text{Orb}^K(x; \overline{X}) \subset \text{Orb}(x; \overline{X})$ holds for each $x \in X_0 + X_1$, i.e., if for every $y \in \text{Orb}^K(x; \overline{X})$ there exists an operator $T \in \mathfrak{L}(\overline{X})$ such that $y = Tx$. Moreover, $\overline{X} = (X_0, X_1)$ has the uniform Calderón-Mityagin property if and only if additionally we can choose $T \in \mathfrak{L}(\overline{X})$ so that $\|T\|_{\mathfrak{L}(\overline{X})} \leq C \|y\|_{\text{Orb}^K(x)}$, where C is independent of x and y .

2.3. Quasi-Banach lattice couples. Suppose X_0 and X_1 are two quasi-Banach lattices. We will say that $\overline{X} = (X_0, X_1)$ is a *quasi-Banach lattice couple* if there exists a Hausdorff topological vector lattice $\mathcal{H}$ such that both X_0 and X_1 are embedded into $\mathcal{H}$ via a continuous, interval preserving, lattice homeomorphism (see e.g. [39]). Then, one can easily check that the intersection $\Delta(\overline{X})$ and the sum $\Sigma(\overline{X})$ are quasi-Banach lattices.

Suppose that a quasi-Banach lattice X is intermediate with respect to a quasi-Banach lattice couple $\overline{X} = (X_0, X_1)$ as a quasi-Banach space. We will say that X is an *intermediate quasi-Banach lattice* with respect to $\overline{X}$ if the canonical embeddings $I_\Delta : \Delta(\overline{X}) \rightarrow X$ and $I_\Sigma : X \rightarrow \Sigma(\overline{X})$ are continuous, interval preserving, lattice homeomorphisms (see [6] and [39]). By $I(\overline{X})$ we will denote in this case the set of all intermediate quasi-Banach lattices with respect to $\overline{X}$.

Observe that every K -monotone quasi-Banach lattice with respect to a quasi-Banach lattice couple is, in fact, uniform K -monotone. This can be proved by a simple modification of the arguments for the Banach lattice case used in [14, Theorem 6.1].

We will say that a quasi-Banach lattice couple $\overline{X} = (X_0, X_1)$ is *p-convex* if both spaces X_0 and X_1 are p -convex.

2.4. The cone $Conv$ and K -functionals. Let $Conv$ denote the cone of all continuous, concave, non-negative functions defined on the half-line $(0, \infty)$. Each $f \in Conv$ is a nondecreasing function such that the function $t \mapsto f(t)/t$ is nonincreasing. Consequently, $f(t) \leq \max(1, t/s) f(s)$, $0 < s, t < \infty$, and thus $Conv$ is a subset of the space $\Sigma(\overline{L^\infty}) = L^\infty(\min(1, 1/s))$ and for any $f \in Conv$ we have $\|f\|_{\Sigma(\overline{L^\infty})} = f(1)$.

Note that, for every quasi-Banach couple $\overline{X}$ and any $x \in \Sigma(\overline{X})$, the function $t \mapsto K(t, x; \overline{X})$ belongs to the cone $Conv$. Moreover, for each function $h \in \Sigma(\overline{L^\infty})$ the function $t \mapsto K(t, h; \overline{L^\infty})$ is the least concave majorant of $|h|$ on $(0, \infty)$ [15] (see also [6, Proposition 3.1.17]). Given a set $\mathcal{U} \subseteq Conv$, a quasi-Banach couple $\overline{X}$ is called $\mathcal{U}$ -*abundant* if for every $f \in \mathcal{U}$ there exists $x \in \Sigma(\overline{X})$ such that

$$K(t, x; \overline{X}) \cong f(t), \quad t > 0,$$

with equivalence constants independent of f (see [6, Definition 4.4.8])³. In particular, if $\mathcal{U} = \{K(\cdot, x; \overline{Y}), x \in \Sigma(\overline{Y})\}$ for some quasi-Banach couple $\overline{Y}$, we will say that $\overline{X}$ is $\overline{Y}$ -*abundant*.

One can check easily that the couple $\overline{L^\infty}$ is $Conv$ -abundant. Therefore, a quasi-Banach couple $\overline{X}$ is $\overline{L^\infty}$ -abundant if and only if it is $Conv$ -abundant.

The notation $A \preceq B$ means that there exists a positive constant C with $A \leq C \cdot B$ for all applicable values of the arguments (parameters) of the functions (expressions) A and B . We will write $A \cong B$ if $A \preceq B$ and $B \preceq A$.

3. AUXILIARY RESULTS

Lemma 3.1. *Let X be a p -convex quasi-Banach lattice, $p \in (0, 1)$. Then the $1/p$ -convexification $X^{(1/p)}$ of X has an equivalent lattice norm and hence $X^{(1/p)}$ is lattice isomorphic to a Banach lattice.*

³This property is referred sometimes as the K -surjectivity of a couple, see e.g. [33, p. 217].

Proof. Let $M^{(p)}$ be the p -convexity constant of X . Hence, if $x_i \in X^{(1/p)}$, $i = 1, 2, \dots, n$, then from definition of the $(1/p)$ -convexification $X^{(1/p)}$ (see Section 2.1) it follows

$$\left\| \sum_{i=1}^n \oplus x_i \right\|_{X^{(1/p)}} \leq (M^{(p)})^p \sum_{i=1}^n \|x_i\|_{X^{(1/p)}}.$$

Now, a straightforward inspection shows that the functional

$$|||x||| := \inf \left\{ \sum_{i=1}^n \|x_i\|_{X^{(1/p)}} : |x| \leq \sum_{i=1}^n \oplus |x_i|, x_i \in X^{(1/p)} \right\}$$

is a lattice norm on the space $X^{(1/p)}$, which is equivalent to the original quasi-norm. $\square$

Some variants of the next result in the Banach case are well known (see e.g. [25, Proposition 2.g.6]).

Proposition 3.2. *Suppose $\overline{X} = (X_0, X_1)$ is a p -convex quasi-Banach lattice couple, $p > 0$. Then $\Sigma(\overline{X})$ is a p -convex lattice and*

$$M^{(p)}(\Sigma(\overline{X})) \leq \max(2^{1-1/p}, 2^{2/p-2}) \max_{i=0,1} M^{(p)}(X_i).$$

Proof. We will assume that $0 < p \leq 1$. The case when $p > 1$ can be treated similarly.

Let $x_i \in \Sigma(\overline{X})$, $i = 1, 2, \dots, n$, and let $x_i = x_i^0 + x_i^1$ be an arbitrary representation, with $x_i^0 \in X_0, x_i^1 \in X_1$. Then, we have

$$\begin{aligned} \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} &\leq \left(\sum_{i=1}^n (|x_i^0|^p + |x_i^1|^p) \right)^{1/p} \\ &\leq 2^{1/p-1} \left(\left(\sum_{i=1}^n |x_i^0|^p \right)^{1/p} + \left(\sum_{i=1}^n |x_i^1|^p \right)^{1/p} \right). \end{aligned}$$

Therefore,

$$\begin{aligned}
\left\| \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \right\|_{\Sigma(\overline{X})}^p &\leq 2^{1-p} \left\| \left(\sum_{i=1}^n |x_i^0|^p \right)^{1/p} + \left(\sum_{i=1}^n |x_i^1|^p \right)^{1/p} \right\|_{\Sigma(\overline{X})}^p \\
&\leq 2^{1-p} \left(\left\| \left(\sum_{i=1}^n |x_i^0|^p \right)^{1/p} \right\|_{X_0} + \left\| \left(\sum_{i=1}^n |x_i^1|^p \right)^{1/p} \right\|_{X_1} \right)^p \\
&\leq 2^{1-p} \max_{i=0,1} M^{(p)}(X_i)^p \left(\left(\sum_{i=1}^n \|x_i^0\|_{X_0}^p \right)^{1/p} + \left(\sum_{i=1}^n \|x_i^1\|_{X_1}^p \right)^{1/p} \right)^p \\
&\leq 2^{1-p} \max_{i=0,1} M^{(p)}(X_i)^p \sum_{i=1}^n \left(\|x_i^0\|_{X_0}^p + \|x_i^1\|_{X_1}^p \right) \\
&\leq 2^{2-2p} \max_{i=0,1} M^{(p)}(X_i)^p \left(\sum_{i=1}^n \left(\|x_i^0\|_{X_0} + \|x_i^1\|_{X_1} \right) \right)^p.
\end{aligned}$$

Now passing in the right-hand side to the infimum over all admissible representations $x_i = x_i^0 + x_i^1$, $i = 1, 2, \dots, n$, we obtain

$$\left\| \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \right\|_{\Sigma(\overline{X})}^p \leq 2^{2-2p} \left(\max_{i=0,1} M^{(p)}(X_i) \right)^p \sum_{i=1}^n \|x_i\|_{\Sigma(\overline{X})}^p,$$

which completes the proof. $\square$

The following simple fact is well-known in the case of Banach function lattices (see e.g. [27], [6, Proposition 3.1.15], [28] and [29, Lemma 1]).

Lemma 3.3. *Let $\overline{X}$ be a quasi-Banach lattice couple. Then for every $x \in \Sigma(\overline{X})$ and all $t > 0$ we have*

$$\begin{aligned}
K(t, x; \overline{X}) &= K(t, |x|; \overline{X}) \\
&= \inf \{ \|x_0\|_{X_0} + t \|x_1\|_{X_1} : |x| = x_0 + x_1, 0 \leq x_0 \in X_0, 0 \leq x_1 \in X_1 \} \\
&= \inf \{ \|x_0\|_{X_0} + t \|x_1\|_{X_1} : |x| \leq x_0 + x_1, 0 \leq x_0 \in X_0, 0 \leq x_1 \in X_1 \}.
\end{aligned}$$

Proof. Denote by $A(t, x)$ (resp. $B(t, x)$) the first (resp. second) infimum in the statement of the lemma.

Let $x \in \Sigma(\overline{X})$ and $|x| \leq x_0 + x_1$, $0 \leq x_i \in X_i$. Then, by the decomposition property (see e.g. [25, p. 2]), we may write $|x| = y_0 + y_1$, where $0 \leq y_i \leq |x_i|$, $i = 0, 1$. Hence, $y_i \in X_i$ and $\|y_i\|_{X_i} \leq \|x_i\|_{X_i}$, $i = 0, 1$. Thus,

$$\|x_0\|_{X_0} + t \|x_1\|_{X_1} \geq \|y_0\|_{X_0} + t \|y_1\|_{X_1} \geq A(t, x).$$

Passing to the infimum in the left-hand side of this inequality, we deduce that $B(t, x) \geq A(t, x)$. Since the opposite inequality is obvious, we get $A(t, x) = B(t, x)$.

Next, if $x \in \Sigma(\overline{X})$ and $|x| = x_0 + x_1, x_i \in X_i$, then $|x| \leq |x_0| + |x_1|$ and hence

$$\|x_0\|_{X_0} + t\|x_1\|_{X_1} \geq B(t, x).$$

Therefore, $K(t, |x|; \overline{X}) \geq B(t, x)$. It is clear also that $K(t, |x|; \overline{X}) \leq A(t, x)$. Summing up, we have

$$(3.1) \quad K(t, |x|; \overline{X}) = A(t, x) = B(t, x).$$

Let now $x \in \Sigma(\overline{X})$ and $x = x_0 + x_1, x_i \in X_i$. Then $|x| \leq |x_0| + |x_1|$, which implies as above that

$$(3.2) \quad K(t, x; \overline{X}) \geq B(t, x).$$

Conversely, assume that $x \in \Sigma(\overline{X})$, $|x| = x_0 + x_1$ and $0 \leq x_i \in X_i$. Denoting $x_+ = x \vee 0, x_- = (-x) \vee 0$, we set $z_0 := x_0 \wedge x_+ - x_0 \wedge x_-$ and $z_1 := x_1 \wedge x_+ - x_1 \wedge x_-$. Clearly, $z_i \in X_i$ and, since $x_i \leq |x|$ (see also [25, Theorem 1.d.1] or [21, p. 142]), we have

$$\|z_i\|_{X_i} \leq \|x_i \wedge x_+ + x_i \wedge x_-\|_{X_i} = \|x_i \wedge |x|\|_{X_i} = \|x_i\|_{X_i}, \quad i = 0, 1.$$

Moreover,

$$\begin{aligned} z_0 + z_1 &= (x_0 \wedge x_+ + x_1 \wedge x_+) - (x_0 \wedge x_- + x_1 \wedge x_-) \\ &= (x_0 + x_1) \wedge x_+ - (x_0 + x_1) \wedge x_- = x_+ - x_- = x. \end{aligned}$$

Thus,

$$\|x_0\|_{X_0} + t\|x_1\|_{X_1} \geq \|z_0\|_{X_0} + t\|z_1\|_{X_1} \geq K(t, x; \overline{X}),$$

whence

$$A(t, x) \geq K(t, x; \overline{X}).$$

Combining this together with (3.1) and (3.2), we complete the proof. $\square$

Proposition 3.4. *Let $p > 0$ and $\overline{X} = (X_0, X_1)$ be a p -convex quasi-Banach lattice couple. Then $\Sigma(\overline{X}^{(1/p)}) = \Sigma(\overline{X})^{(1/p)}$. Moreover, for arbitrary $x \in \Sigma(\overline{X}^{(1/p)})$ and all $t > 0$ we have*

$$(3.3) \quad \min(2^{2p-2}, 2^{1-p}) \min_{i=0,1} (M^{(p)}(X_i))^{-p} K(t^{1/p}, x; \overline{X})^p \leq K(t, x; \overline{X}^{(1/p)})$$

and

$$(3.4) \quad K(t, x; \overline{X}^{(1/p)}) \leq \max(2^{p-1}, 2^{1-p}) K(t^{1/p}, x; \overline{X})^p.$$

Proof. For definiteness, we consider again only the case when $0 < p \leq 1$. Moreover, by Lemma 3.3, we can (and will) assume that $x \geq 0$.

Suppose first $x \in \Sigma(\overline{X}^{(1/p)})$. Take any decomposition $x = x_0 \oplus x_1 = (x_0^p + x_1^p)^{1/p}$ with $0 \leq x_0 \in X_0^{(1/p)}, 0 \leq x_1 \in X_1^{(1/p)}$. Since X_0 and X_1 are p -convex, then, by

Proposition 3.2, the sum $X_0 + t^{1/p}X_1$ is p -convex for every $t > 0$ with the constant $C_p := \max(2^{1-1/p}, 2^{2/p-2}) \max_{i=0,1} M^{(p)}(X_i)$. Consequently,

$$\begin{aligned} K(t^{1/p}, x; \overline{X})^p &= \|x\|_{X_0+t^{1/p}X_1}^p \\ &= \|(x_0^p + x_1^p)^{1/p}\|_{X_0+t^{1/p}X_1}^p \\ &\leq C_p^p (\|x_0\|_{X_0+t^{1/p}X_1}^p + \|x_1\|_{X_0+t^{1/p}X_1}^p) \\ &\leq C_p^p (\|x_0\|_{X_0}^p + (t^{1/p}\|x_1\|_{X_1})^p) \\ &= C_p^p (\|x_0\|_{X_0^{(1/p)}} + t\|x_1\|_{X_1^{(1/p)}}). \end{aligned}$$

Thus, $x \in \Sigma(\overline{X})^{(1/p)}$ and from Lemma 3.3 it follows that

$$K(t^{1/p}, x; \overline{X})^p \leq C_p^p K(t, x; \overline{X^{(1/p)}}),$$

which is equivalent to inequality (3.3).

Conversely, let $x \in \Sigma(\overline{X})^{(1/p)}$ and $x = y_0 + y_1$, with $0 \leq y_0 \in X_0$, $0 \leq y_1 \in X_1$. Then, since $0 < p \leq 1$, we have $x \leq (y_0^p + y_1^p)^{1/p}$. Applying once more the decomposition property to the lattice $\Sigma(\overline{X})^{(1/p)}$ (see [25, p. 2]), we may write $x = (x_0^p + x_1^p)^{1/p}$, where $0 \leq x_i \leq y_i$, $i = 0, 1$. Since this implies that $x_i \in X_i^{(1/p)}$, $i = 0, 1$, it follows that $x \in \Sigma(\overline{X^{(1/p)}})$ and moreover

$$\begin{aligned} K(t, x; \overline{X^{(1/p)}}) &\leq \|x_0\|_{X_0^{(1/p)}} + t\|x_1\|_{X_1^{(1/p)}} = \|x_0\|_{X_0}^p + t\|x_1\|_{X_1}^p \\ &\leq \|y_0\|_{X_0}^p + t\|y_1\|_{X_1}^p \leq 2^{1-p}(\|y_0\|_{X_0} + t^{1/p}\|y_1\|_{X_1})^p. \end{aligned}$$

Applying Lemma 3.3, we see that this inequality implies (3.4). Therefore, the proof is completed. $\square$

Proposition 3.5. *If $\overline{X} = (X_0, X_1)$ is a quasi-Banach couple, then for arbitrary $E_2, E_3 \in \text{Int}(\overline{L^\infty})$ we have*

$$\overline{X}_{E_2;K} + \overline{X}_{E_3;K} = \overline{X}_{E_2+E_3;K}.$$

Proof. Let $x \in \overline{X}_{E_2+E_3;K}$. Then $K(\cdot, x; \overline{X}) \in E_2 + E_3$ and so, by Lemma 3.3, $K(\cdot, x; \overline{X}) = f_2(\cdot) + f_3(\cdot)$, where $0 \leq f_2 \in E_2$, $0 \leq f_3 \in E_3$ and

$$(3.5) \quad \|x\|_{\overline{X}_{E_2+E_3;K}} \succeq \|f_2\|_{E_2} + \|f_3\|_{E_3}$$

with some constant independent of x . Moreover, since $E_2, E_3 \in \text{Int}(\overline{L^\infty})$, we may assume that $f_2, f_3 \in \text{Conv}$. Therefore, by a weak version of K -divisibility property, which holds for quasi-Banach couples [6, Theorem 3.2.12], we can find a decomposition $x = x_2 + x_3$ such that $K(t, x_i; \overline{X}) \leq \gamma f_i(t)$, $i = 2, 3$, for a universal constant γ and all $t > 0$. Combining these inequalities with (3.5), we get

$$\begin{aligned} \|x\|_{\overline{X}_{E_2+E_3;K}} &\succeq \|K(t, x_2; \overline{X})\|_{E_2} + \|K(t, x_3; \overline{X})\|_{E_3} \\ &= \|x_2\|_{\overline{X}_{E_2;K}} + \|x_3\|_{\overline{X}_{E_3;K}} \geq \|x\|_{\overline{X}_{E_2;K} + \overline{X}_{E_3;K}}, \end{aligned}$$

whence

$$\overline{X}_{E_2+E_3;K} \subset \overline{X}_{E_2;K} + \overline{X}_{E_3;K}.$$

To prove the opposite embedding, assume that $x \in \overline{X}_{E_2;K} + \overline{X}_{E_3;K}$. Then, $x = x_2 + x_3$, with $x_i \in \overline{X}_{E_i;K}$, $i = 2, 3$, and

$$\|x\|_{\overline{X}_{E_2;K} + \overline{X}_{E_3;K}} \succeq \|x_2\|_{\overline{X}_{E_2;K}} + \|x_3\|_{\overline{X}_{E_3;K}}$$

with some constant independent of x . Therefore, since

$$K(t, x; \overline{X}) \leq \max(C_0, C_1) (K(t, x_2; \overline{X}) + K(t, x_3; \overline{X})), \quad t > 0,$$

where C_i is the constant in the quasi-triangle inequality for X_i , $i = 0, 1$, we have

$$\begin{aligned} \|x\|_{\overline{X}_{E_2+E_3;K}} &= \|K(\cdot, x; \overline{X})\|_{E_2+E_3} \\ &\preceq \|K(\cdot, x_2; \overline{X})\|_{E_2} + \|K(\cdot, x_3; \overline{X})\|_{E_3} \\ &= \|x_2\|_{\overline{X}_{E_2;K}} + \|x_3\|_{\overline{X}_{E_3;K}} \preceq \|x\|_{\overline{X}_{E_2;K} + \overline{X}_{E_3;K}}, \end{aligned}$$

which implies that

$$\overline{X}_{E_2;K} + \overline{X}_{E_3;K} \subset \overline{X}_{E_2+E_3;K},$$

and the proposition is proved. $\square$

For some versions of the next result see [6, Theorem 3.6.7, p. 415] and [32, Theorem 3.20]. Recall that a quasi-Banach function lattice E on a measure space (T, Σ, μ) has the *Fatou property* if from $x_n \in E$, $n = 1, 2, \dots$, $\sup_{n=1,2,\dots} \|x_n\|_E < \infty$ and $x_n \rightarrow x$ a.e. on T it follows that $x \in E$ and $\|x\|_E \leq \liminf_{n \rightarrow \infty} \|x_n\|_E$. Observe that such a lattice E has the Fatou property whenever from $x_n \in E$, $x_n \geq 0$, $n = 1, 2, \dots$, and $x_n \uparrow x$ a.e. on T it follows that $x \in E$ and $\lim_{n \rightarrow \infty} \|x_n\|_E = \|x\|_E$.

Proposition 3.6. *Let $\overline{X} = (X_0, X_1)$ be a quasi-Banach couple and let $X_2 = (X_0, X_1)_{E_2;K}$, $X_3 = (X_0, X_1)_{E_3;K}$, where $E_2, E_3 \in \text{Int}(\overline{L^\infty})$. Suppose that at least one of the following conditions holds:*

- (a) (E_2, E_3) is a mutually closed quasi-Banach couple;
- (b) E_2 and E_3 are relatively complete spaces with respect to the couple $\overline{L^\infty}$;
- (c) E_2 and E_3 have the Fatou property.

Then, the couple (X_2, X_3) is mutually closed.

Proof. Assuming the condition (a) to be hold, we have

$$\|f\|_{E_2} \cong \sup_{t>0} K(t, f; E_2, E_3) \quad \text{and} \quad \|f\|_{E_3} \cong \sup_{t>0} \frac{1}{t} K(t, f; E_2, E_3),$$

with some constants independent of f (see Section 2.2 or [6, Lemma 2.2.21 p.123] or [36, p. 384]). Moreover, by the reiteration theorem (see equivalence (2.3) or [6, Theorem 3.3.11]),

$$K(t, x; X_2, X_3) \cong K(t, K(\cdot, x; \overline{X}); E_2, E_3), \quad t > 0,$$

with constants independent of $x \in X_2 + X_3$. Therefore,

$$\begin{aligned} \sup_{t>0} K(t, x; X_2, X_3) &\cong \sup_{t>0} K(t, K(\cdot, x; \overline{X}); E_2, E_3) \\ &\cong \|K(\cdot, x; \overline{X})\|_{E_2} = \|x\|_{E_2} \end{aligned}$$

and similarly

$$\begin{aligned} \sup_{t>0} \frac{1}{t} K(t, x; X_2, X_3) &\cong \sup_{t>0} \frac{1}{t} K(t, K(\cdot, x; \overline{X}); E_2, E_3) \\ &\cong \|K(\cdot, x; \overline{X})\|_{E_3} = \|x\|_{E_3}. \end{aligned}$$

This implies that the couple (X_2, X_3) is mutually closed.

Since $E_2, E_3 \in \text{Int}(\overline{L^\infty})$, then the condition (a) is a consequence of (b). Hence, the result follows also in the case (b).

Finally, assume that the condition (c) holds. Suppose that $\{x_n\}$ is a sequence from E_2 such that $C := \sup_{n=1,2,\dots} \|x_n\|_{E_2} < \infty$ and $x_n \rightarrow x$ in $\Sigma(\overline{L^\infty})$. Then, $x_n \rightarrow x$ a.e. on $(0, \infty)$. Therefore, since E_2 has the Fatou property, we obtain that $x \in E_2$ and $\|x\|_{E_2} \leq C$. As a result, E_2 is a relatively complete space with respect to the couple $\overline{L^\infty}$. Since the same result holds for E_3 , we get (b). This completes the proof. $\square$

In particular, since the space $L^p(t^{-\theta}, \frac{dt}{t})$ is relatively complete with respect to the couple $\overline{L^\infty}$ for all $0 < \theta < 1$ and $0 < p \leq \infty$, we have

Corollary 3.7. *For every quasi-Banach couple $\overline{X} = (X_0, X_1)$ and all $0 < \theta_0, \theta_1 < 1$, $0 < p_0, p_1 \leq \infty$ the couple $(\overline{X}_{\theta_0, p_0}, \overline{X}_{\theta_1, p_1})$ is mutually closed.*

Remark 3.8. The result of Corollary 3.7 is well known in the case when $\overline{X} = (X_0, X_1)$ is a Banach couple and $1 \leq p_0, p_1 \leq \infty$. Indeed, then $(\overline{X}_{\theta_0, p_0}, \overline{X}_{\theta_1, p_1})$ is a K -monotone Banach couple [12, Theorem 1] and hence it is mutually closed [11, Lemma 3].

4. MAIN RESULTS

In [13], it was announced a statement, asserting that the class of all K -spaces with respect to a Banach couple possesses the Arazy-Cwikel property (see also [6, p. 672]). Below we give a proof of this result for mutually closed Banach couples (see Corollary 4.4). In fact, by using some ideas due to Bykov-Ovchinnikov [7], we obtain here some more general results of such type in the quasi-Banach setting.

The following main result of this paper establishes the Arazy-Cwikel property for the class of K -monotone spaces with respect to an arbitrary mutually closed quasi-Banach couple.

Theorem 4.1. *Let $\overline{X} = (X_0, X_1)$ be a mutually closed quasi-Banach couple. Then, for all $0 < \theta < \eta < 1$ and $0 < p, q \leq \infty$ we have*

$$(4.1) \quad \text{Int}^{KM}(\overline{X}_{\theta, p}, \overline{X}_{\eta, q}) = \text{Int}^{KM}(X_0, \overline{X}_{\eta, q}) \cap \text{Int}^{KM}(\overline{X}_{\theta, p}, X_1).$$

Since the classes of K -monotone and interpolation spaces for couples with the uniform Calderón-Mityagin property coincide, we immediately get the following result, extending thereby some results due to Bykov-Ovchinnikov [7] (see Introduction) to the quasi-Banach case.

Corollary 4.2. *Suppose $\overline{X} = (X_0, X_1)$ is a mutually closed quasi-Banach couple such that the couples $(\overline{X}_{\theta,p}, \overline{X}_{\eta,q})$, $(X_0, \overline{X}_{\eta,q})$ and $(\overline{X}_{\theta,p}, X_1)$ have the uniform Calderón-Mityagin property for all $0 < \theta < \eta < 1$ and $0 < p, q \leq \infty$. Then,*

$$(4.2) \quad \text{Int}(\overline{X}_{\theta,p}, \overline{X}_{\eta,q}) = \text{Int}(X_0, \overline{X}_{\eta,q}) \cap \text{Int}(\overline{X}_{\theta,p}, X_1).$$

Remark 4.3. In the Banach case the conditions of the last corollary may be relaxed (see, for instance, [12, Theorem 1], [16, Theorem 2], [33, Theorem 4.17] and [6, Theorem 4.4.18]). In particular, (4.2) holds for every Banach couple $\overline{X} = (X_0, X_1)$ with the uniform Calderón-Mityagin property for all $0 < \theta < \eta < 1$ and $1 \leq p, q \leq \infty$.

It seems to be unknown by now whether the classes of K -monotone and K -spaces coincide in the quasi-Banach setting. At the same time, it is well known that this coincidence holds in the Banach case [6, Theorem 4.1.11] (see also [33, Corollary 4.3]). Therefore, from Theorem 4.1 it follows

Corollary 4.4. *If $\overline{X} = (X_0, X_1)$ is an arbitrary mutually closed Banach couple, then*

$$\text{Int}^K(\overline{X}_{\theta,p}, \overline{X}_{\eta,q}) = \text{Int}^K(X_0, \overline{X}_{\eta,q}) \cap \text{Int}^K(\overline{X}_{\theta,p}, X_1)$$

for all $0 < \theta < \eta < 1$ and $0 < p, q \leq \infty$.

To prove the above results we need some additional notions and auxiliary assertions.

Lemma 4.5. *Let E and F be quasi-Banach function lattices such that $E, F \in \text{Int}(\overline{L^\infty})$. Then, the following continuous embeddings hold:*

$$\begin{aligned} (L^\infty + F) \cap (E + L^\infty(1/t)) &\subset E + F \\ (L^\infty \cap F) + (E \cap L^\infty(1/t)) &\supset E \cap F. \end{aligned}$$

Proof. Suppose $x \in (L^\infty + F) \cap (E + L^\infty(1/t))$. Then, on the one hand, $x = x_0 + x_1$, where $x_0 \in L^\infty$ and $x_1 \in F$. Since $F \in \text{Int}(\overline{L^\infty})$, we have $x\chi_{[1,\infty)} = x_0\chi_{[1,\infty)} + x_1\chi_{[1,\infty)} \in F$ and $\|x\chi_{[1,\infty)}\|_F \preceq \|x\|_{L^\infty+F}$. On the other hand, $x = x'_0 + x'_1$, where $x'_0 \in E$ and $x'_1 \in L^\infty(1/t)$. As above, we get $x\chi_{[0,1]} = x'_0\chi_{[0,1]} + x'_1\chi_{[0,1]} \in E$ and $\|x\chi_{[0,1]}\|_E \preceq \|x\|_{E+L^\infty(1/t)}$. Consequently, $x = x\chi_{[0,1]} + x\chi_{[1,\infty)} \in E + F$ and $\|x\|_{E+F} \preceq \|x\|_{(L^\infty+F) \cap (E+L^\infty(1/t))}$.

Since the second embedding can be proved quite similarly, we skip its proof. $\square$

Let $\overline{X} = (X_0, X_1)$ be a mutually closed quasi-Banach couple and let X_2, X_3 be quasi-Banach spaces such that $X_2 \in \text{Int}^K(X_0, X_3)$, $X_3 \in \text{Int}^K(X_2, X_1)$ and both couples (X_0, X_3) and (X_2, X_1) are mutually closed. We will call such a collection $\{\overline{X}, X_2, X_3\}$ *admissible*.

Remark 4.6. Further, the following sufficient condition for the admissibility of a collection $\{\overline{X}, X_2, X_3\}$ will be useful.

Assume that $\overline{X} = (X_0, X_1)$ is a mutually closed quasi-Banach couple and X_2, X_3 are quasi-Banach spaces such that $X_2 \in \text{Int}^K(X_0, X_3)$, $X_3 \in \text{Int}^K(X_2, X_1)$. Then, by the reiteration theorem for the K -method (see (2.4) or [6, Theorem 3.3.11]), $X_2, X_3 \in \text{Int}^K(X_0, X_1)$ and hence there are parameters $E_2, E_3 \in$

$\text{Int}(\overline{L^\infty})$ with $X_2 = (X_0, X_1)_{E_2;K}$, $X_3 = (X_0, X_1)_{E_3;K}$. Moreover, by the hypothesis, $X_0 = (X_0, X_1)_{L^\infty;K}$ and $X_1 = (X_0, X_1)_{L^\infty(1/t);K}$. Then, applying Proposition 3.6 to the couples (X_0, X_3) and (X_2, X_1) , we conclude that they are mutually closed. Hence, the collection $\{\overline{X}, X_2, X_3\}$ is admissible.

Recall that $I(X, Y)$ denotes the set of all intermediate spaces with respect to a quasi-Banach couple (X, Y) .

Lemma 4.7. *Let $\overline{X} = (X_0, X_1)$ be a quasi-Banach couple and $\{\overline{X}, X_2, X_3\}$ be an admissible collection. Then, we have*

$$I(X_2, X_3) = I(X_0, X_3) \cap I(X_2, X_1).$$

Proof. Suppose first $X \in I(X_2, X_3)$, i.e., $X_2 \cap X_3 \subseteq X \subseteq X_2 + X_3$. Since $X_2 \in \text{Int}(X_0, X_3)$, it follows that $X_0 \cap X_3 \subseteq X_2 \subseteq X_0 + X_3$. Hence, $X_0 \cap X_3 \subseteq X \subseteq X_0 + X_3$, which means that $X \in I(X_0, X_3)$. In the same way, $X \in I(X_2, X_1)$.

Conversely, assume that $X \in I(X_0, X_3) \cap I(X_2, X_1)$, i.e., $X_0 \cap X_3 \subseteq X \subseteq X_0 + X_3$ and $X_2 \cap X_1 \subseteq X \subseteq X_2 + X_1$. Then, in particular,

$$X \subseteq (X_0 + X_3) \cap (X_2 + X_1).$$

It is clear also that $X_0 \subseteq \overline{X}_{L^\infty;K}$, $X_1 \subseteq \overline{X}_{L^\infty(1/t);K}$. Combining this with the equalities $X_2 = \overline{X}_{E_2;K}$, $X_3 = \overline{X}_{E_3;K}$, by Proposition 3.5, we conclude

$$(X_0 + X_3) \cap (X_2 + X_1) \subseteq \overline{X}_{L^\infty + E_3;K} \cap \overline{X}_{E_2 + L^\infty(1/t);K} = \overline{X}_{E;K},$$

where $E := (L^\infty + E_3) \cap (E_2 + L^\infty(1/t))$. On the other hand, by Lemma 4.5, $E \subset E_2 + E_3$, and so from Proposition 3.5 it follows

$$\overline{X}_{E;K} \subset \overline{X}_{E_2 + E_3;K} = \overline{X}_{E_2;K} + \overline{X}_{E_3;K} = X_2 + X_3.$$

As a result, the last embeddings yield $X \subseteq X_2 + X_3$.

Similarly,

$$X \supset X_0 \cap X_3 + X_2 \cap X_1 = \overline{X}_{E';K},$$

where $E' := (L^\infty \cap E_3) + (E_2 \cap L^\infty(1/t))$. Again applying Lemma 4.5, we get

$$\overline{X}_{E';K} \supset \overline{X}_{E_2 \cap E_3;K} = \overline{X}_{E_2;K} \cap \overline{X}_{E_3;K} = X_2 \cap X_3$$

and hence $X_2 \cap X_3 \subseteq X$. Thus, $X \in I(X_2, X_3)$, which completes the proof. $\square$

Lemma 4.8. *Let $\{\overline{X}, X_2, X_3\}$ be an admissible collection. The following embeddings hold:*

- (a) $\text{Int}^K(X_2, X_3) \subseteq \text{Int}^K(X_0, X_3) \cap \text{Int}^K(X_2, X_1)$;
- (b) $\text{Int}^{KM}(X_2, X_3) \subseteq \text{Int}^{KM}(X_0, X_3) \cap \text{Int}^{KM}(X_2, X_1)$.

Proof. Observe first that, thanks to Lemma 4.7, either of the conditions $X \in \text{Int}^K(X_2, X_3)$ and $X \in \text{Int}^{KM}(X_2, X_3)$ implies that

$$X \in I(X_0, X_3) \cap I(X_2, X_1).$$

(a) Suppose $X \in \text{Int}^K(X_2, X_3)$, i.e., $X = (X_2, X_3)_{E;K}$ for some $E \in \text{Int}(\overline{L^\infty})$. According to the definition of an admissible collection, $X_3 = (X_0, X_3)_{L^\infty(1/t);K}$ and there are $F_2, F_3 \in \text{Int}(\overline{L^\infty})$ such that $X_2 = (X_0, X_3)_{F_2;K}$, $X_3 = (X_2, X_1)_{F_3;K}$.

Therefore, again by the reiteration theorem (see (2.3) or [6, Theorem 3.3.11]), with constants independent of $x \in X_2 + X_3$, we have

$$(4.3) \quad K(t, x; X_2, X_3) \cong K(t, K(\cdot, x; X_0, X_3); F_2, L^\infty(1/t)), \quad t > 0.$$

Consequently, $X = (X_0, X_3)_{E_X:K}$, where $E_X = (F_2, L^\infty(1/t))_{E:K}$. Hence, $X \in \text{Int}^K(X_0, X_3)$. In the same way, $X \in \text{Int}^K(X_2, X_1)$.

(b) Let $X \in \text{Int}^{KM}(X_2, X_3)$. Suppose that $x \in X$ and $y \in X_0 + X_3$ satisfy

$$K(t, y; X_0, X_3) \leq K(t, x; X_0, X_3), \quad t > 0.$$

Then, by (4.3),

$$K(1, K(\cdot, y; X_0, X_3); F_2, L^\infty(1/t)) \leq CK(1, K(\cdot, x; X_0, X_3); F_2, L^\infty(1/t)) < \infty,$$

that is,

$$K(\cdot, y; X_0, X_3) \in F_2 + L^\infty(1/t).$$

Thus, by Proposition 3.5, we obtain

$$y \in (X_0, X_3)_{F_2 + L^\infty(1/t):K} = (X_0, X_3)_{F_2:K} + (X_0, X_3)_{L^\infty(1/t):K} = X_2 + X_3$$

and hence from (4.3) and similar equivalence for y we infer

$$K(t, y; X_2, X_3) \preceq K(t, x; X_2, X_3), \quad t > 0.$$

Therefore, by the assumption, $y \in X$ and so $X \in \text{Int}^{KM}(X_0, X_3)$. In the same way, $X \in \text{Int}^{KM}(X_2, X_1)$. $\square$

The following definition is inspired by some results due to Bykov and Ovchinnikov [7].

Definition 4.9. We will say that an admissible collection $\{\overline{X}, X_2, X_3\}$, where $\overline{X} = (X_0, X_1)$, has *additive K -orbits* whenever for each $x \in X_2 + X_3$ the equality

$$\text{Orb}^K(x; X_2, X_3) = \text{Orb}^K(x; X_0, X_3) + \text{Orb}^K(x; X_2, X_1)$$

holds uniformly with respect to x .

Lemma 4.10. *If an admissible collection $\{\overline{X}, X_2, X_3\}$ has additive K -orbits, then*

$$\text{Int}^{KM}(X_0, X_3) \cap \text{Int}^{KM}(X_2, X_1) = \text{Int}^{KM}(X_2, X_3).$$

Proof. In view of Lemma 4.8(b), it suffices only to prove the embedding

$$\text{Int}^{KM}(X_0, X_3) \cap \text{Int}^{KM}(X_2, X_1) \subset \text{Int}^{KM}(X_2, X_3).$$

First, if $X \in \text{Int}^{KM}(X_0, X_3) \cap \text{Int}^{KM}(X_2, X_1)$. Then, by Lemma 4.7, $X \in I(X_2, X_3)$. To prove that $X \in \text{Int}^{KM}(X_2, X_3)$, assume that $x \in X$ and $y \in X_2 + X_3$ satisfy

$$K(t, y; X_2, X_3) \leq K(t, x; X_2, X_3), \quad t > 0.$$

Therefore, $y \in \text{Orb}^K(x; X_2, X_3)$ and $\|y\|_{\text{Orb}^K(x; X_2, X_3)} \leq 1$. Thus, by the assumption, $y = y_3 + y_2$, where $y_3 \in \text{Orb}^K(x; X_0, X_3)$, $y_2 \in \text{Orb}^K(x; X_2, X_1)$,

$$K(t, y_3; X_0, X_3) \leq CK(t, x; X_0, X_3), \quad t > 0,$$

and

$$K(t, y_2; X_2, X_1) \leq CK(t, x; X_2, X_1), \quad t > 0,$$

with some constant C independent of x . Since $X \in \text{Int}^{KM}(X_0, X_3) \cap \text{Int}^{KM}(X_2, X_1)$, it follows that $y_2, y_3 \in X$. In consequence, $y \in X$ and $\|y\|_X \leq C'\|x\|_X$ with a constant C' independent of x , which implies that $X \in \text{Int}^{KM}(X_2, X_3)$. $\square$

Proposition 4.11. *Let $\{\overline{X}, X_2, X_3\}$ and $\{\overline{Y}, Y_2, Y_3\}$ be two admissible collections such that $X_2 = \overline{X}_{E_2:K}$, $Y_2 = \overline{Y}_{E_2:K}$, $X_3 = \overline{X}_{E_3:K}$ and $Y_3 = \overline{Y}_{E_3:K}$, where $E_2, E_3 \in \text{Int}(\overline{L^\infty})$. If $\overline{Y}$ is an $\overline{X}$ -abundant couple and the collection $\{\overline{Y}, Y_2, Y_3\}$ has additive K -orbits, then the collection $\{\overline{X}, X_2, X_3\}$ has K -additive orbits as well.*

Proof. Suppose first $x \in X_2 + X_3$ and $y \in \text{Orb}^K(x; X_0, X_3)$. Then, arguing in the same way as in the proof of Lemma 4.8 (b) (see also (4.3)), we conclude that $y \in \text{Orb}^K(x; X_2, X_3)$. Similarly, the fact that $y \in \text{Orb}^K(x; X_2, X_1)$ yields $y \in \text{Orb}^K(x; X_2, X_3)$. Thus,

$$\text{Orb}^K(x; X_0, X_3) + \text{Orb}^K(x; X_2, X_1) \subset \text{Orb}^K(x; X_2, X_3),$$

with constants independent of x . Therefore, it remains only to prove the opposite embedding

$$(4.4) \quad \text{Orb}^K(x; X_2, X_3) \subset \text{Orb}^K(x; X_0, X_3) + \text{Orb}^K(x; X_2, X_1).$$

Let $x \in X_2 + X_3$ and $y \in \text{Orb}^K(x; X_2, X_3)$ be arbitrary. Since $x, y \in \Sigma(\overline{X})$ and $\overline{Y}$ is $\overline{X}$ -abundant, we can select elements $f, g \in \Sigma(\overline{Y})$ such that

$$K(t, x; \overline{X}) \cong K(t, f; \overline{Y}) \quad \text{and} \quad K(t, y; \overline{X}) \cong K(t, g; \overline{Y}), \quad t > 0,$$

with universal constants. Moreover, by the conditions and reiteration arguments,

$$\begin{aligned} K(t, x; X_2, X_3) &= K(t, x; \overline{X}_{E_2:K}, \overline{X}_{E_3:K}) \cong K(t, K(\cdot, x; \overline{X}); E_2, E_3) \\ &\cong K(t, K(\cdot, f; \overline{Y}); E_2, E_3) = K(t, f; Y_2, Y_3), \quad t > 0. \end{aligned}$$

Similarly, we obtain

$$K(t, y; X_2, X_3) \cong K(t, g; Y_2, Y_3), \quad t > 0.$$

Combining these equivalences with the assumption that $y \in \text{Orb}^K(x; X_2, X_3)$ we conclude that $g \in \text{Orb}^K(f; Y_2, Y_3)$. Therefore, since the collection $\{\overline{Y}, Y_2, Y_3\}$ has additive K -orbits, we may write $g = g_0 + g_1$, where $g_0 \in \text{Orb}^K(f; Y_0, Y_3)$, $g_1 \in \text{Orb}^K(f; Y_2, Y_1)$. Since $\overline{Y} = (Y_0, Y_1)$ is a quasi-Banach couple, then the K -functional $K(t, g; \overline{Y})$ is a quasi-norm on the sum $Y_0 + Y_1$ with the constant $C = \max(C_0, C_1)$, where C_i is the quasi-norm constant for Y_i , $i = 0, 1$. Consequently, we have

$$K(t, y; \overline{X}) \cong K(t, g; \overline{Y}) \leq C \cdot (K(t, g_0; \overline{Y}) + K(t, g_1; \overline{Y})), \quad t > 0.$$

Using now once more a weak version of K -divisibility property [6, Theorem 3.2.12], we can find a decomposition $y = y_0 + y_1$ such that $K(t, y_i; \overline{X}) \leq \gamma CK(t, g_i; \overline{Y})$, $i = 0, 1$, for all $t > 0$. Combining these inequalities with the above relations

and the hypothesis that $\{\overline{X}, X_2, X_3\}$ and $\{\overline{Y}, Y_2, Y_3\}$ are admissible collections, by reiteration arguments, we get

$$\begin{aligned} K(t, y_0; X_0, X_3) &= K(t, K(\cdot, y_0; \overline{X}); L_\infty, E_3) \preceq K(t, K(\cdot, g_0; \overline{Y}); L_\infty, E_3) \\ &\cong K(t, g_0; Y_0, Y_3) \preceq K(t, f; Y_0, Y_3) \cong K(t, K(\cdot, f; \overline{Y}); L_\infty, E_3) \\ &\cong K(t, K(\cdot, x; \overline{X}); L_\infty, E_3) = K(t, x; X_0, X_3), \quad t > 0. \end{aligned}$$

Thus, $y_0 \in Orb^K(x; X_0, X_3)$. In the same way, $y_1 \in Orb^K(x; X_2, X_1)$. As a result, we get embedding (4.4) (with a constant independent of $x \in X_2 + X_3$), and therefore the proof is completed. $\square$

Lemma 4.12. *Let $\overline{X} = (X_0, X_1)$ be a p -convex quasi-Banach lattice couple. Then, for each $r \geq 1/p$ and every $x \in \Sigma(\overline{X})$ we have*

$$Orb^K(x; \overline{X})^{(r)} = Orb^K(x; \overline{X}^{(r)}).$$

Proof. Observe that, by Proposition 3.4, the inequalities

$$K(t, y; \overline{X}) \leq CK(t, x; \overline{X}), \quad t > 0,$$

and

$$K(t, y; \overline{X}^{(r)}) \leq C_r \cdot K(t, x; \overline{X}^{(r)}), \quad t > 0,$$

where the constant C_r depends only on $\overline{X}$, C and r , are equivalent. As a result, the spaces $Orb^K(x; \overline{X})^{(r)}$ and $Orb^K(x; \overline{X}^{(r)})$ coincide with equivalent norms. $\square$

Lemma 4.13. *Let $0 < \theta < 1, 0 < p \leq \infty$. Suppose $\overline{X} = (X_0, X_1)$ is a p -convex quasi-Banach lattice couple and r is a positive number such that $r \geq 1/p$. Then, $(\overline{X}_{\theta,p})^{(r)} = \overline{X}^{(r)}_{\theta,rp}$.*

Proof. Assuming that $p < \infty$ and applying once more Proposition 3.4, we obtain

$$\begin{aligned} \|x\|_{(\overline{X}_{\theta,p})^{(r)}}^r &= \left(\int_0^\infty (s^{-\theta} K(s, x; \overline{X}))^p ds/s \right)^{1/p} \\ &\cong \left(\int_0^\infty \left(s^{-\theta} K(s^{1/r}, x; \overline{X}^{(r)})^r \right)^p ds/s \right)^{1/p} \\ &\cong \left(\left(\int_0^\infty (u^{-\theta} K(u, x; \overline{X}^{(r)}))^{rp} du/u \right)^{1/rp} \right)^r \\ &= \|x\|_{\overline{X}^{(r)}_{\theta,rp}}^r. \end{aligned}$$

The case when $p = \infty$ can be handled quite similarly. $\square$

Proof of Theorem 4.1. Let $\overline{X} = (X_0, X_1)$ be a mutually closed quasi-Banach couple, $0 < \theta < \eta < 1, 0 < p, q \leq \infty$. We put $X_2 = \overline{X}_{\theta,p}$ and $X_3 = \overline{X}_{\eta,q}$. Since the space $L_*^r(t^{-s})$ is relatively complete with respect to the couple $\overline{L}^\infty$ for all

$0 < s < 1$ and $0 < r \leq \infty$, by Remark 4.6 (see also Proposition 3.6), it follows that the collection $\{\overline{X}, \overline{X}_{\theta,p}, \overline{X}_{\eta,q}\}$ is admissible. Moreover, by the classical reiteration theorem (see e.g. [5, Theorem 3.5.3]), we have

$$(4.5) \quad \overline{X}_{\theta,p} = (X_0, \overline{X}_{\eta,q})_{\theta/\eta,p} \quad \text{and} \quad \overline{X}_{\eta,q} = (\overline{X}_{\theta,p}, X_1)_{(\eta-\theta)/(1-\eta),q}.$$

Along with the collection $\{\overline{X}, \overline{X}_{\theta,p}, \overline{X}_{\eta,q}\}$ we consider also the (admissible) collection $\{\overline{L}^\infty, L_*^p(t^{-\theta}), L_*^q(t^{-\eta})\}$, where $L_*^r(t^{-s}) = L^r(t^{-s}, \frac{dt}{t})$, $0 < s < 1$, $0 < r \leq \infty$. Since for all $0 < s < 1$ and $0 < r \leq \infty$

$$(4.6) \quad L_*^r(t^{-s}) = \overline{L}^\infty_{s,r},$$

then, applying (4.5) to the couple $\overline{L}^\infty$, we get

$$L_*^p(t^{-\theta}) = (L^\infty, L_*^q(t^{-\eta}))_{\theta/\eta,p}$$

and

$$L_*^q(t^{-\eta}) = (L_*^p(t^{-\theta}), L^\infty(1/t))_{(\eta-\theta)/(1-\eta),q}.$$

In view of Lemma 4.10, we need only to prove that the collection $\{\overline{X}, X_2, X_3\}$ has K -additive orbits. In turn, since the couple $\overline{L}^\infty$ is *Conv*-abundant, according to Proposition 4.11, this would be established, once we prove that the collection $\{\overline{L}^\infty, L_*^p(t^{-\theta}), L_*^q(t^{-\eta})\}$ has K -additive orbits. In other words, it suffices to show that

$$(4.7) \quad \begin{aligned} Orb^K(x; L_*^p(t^{-\theta}), L_*^q(t^{-\eta})) &= Orb^K(x; L^\infty, L_*^q(t^{-\eta})) \\ &+ Orb^K(x; L_*^p(t^{-\theta}), L^\infty(1/t)), \end{aligned}$$

for every $x \in L_*^p(t^{-\theta}) + L_*^q(t^{-\eta})$ and $0 < \theta < \eta < 1$, $0 < p, q \leq \infty$.

Let $r \geq \max(1/p, 1/q, 1)$. Then, from (4.6) and Lemmas 4.12, 4.13 it follows

$$\begin{aligned} Orb^K(x; L_*^p(t^{-\theta}), L_*^q(t^{-\eta}))^{(r)} &= Orb^K\left(x; (\overline{L}^\infty_{\theta,p})^{(r)}, (\overline{L}^\infty_{\eta,q})^{(r)}\right) \\ &= Orb^K\left(x; (\overline{L}^\infty)^{(r)}_{\theta,pr}, (\overline{L}^\infty)^{(r)}_{\eta,qr}\right), \end{aligned}$$

and similarly

$$Orb^K(x; L^\infty, L_*^q(t^{-\eta}))^{(r)} = Orb^K\left(x; (L^\infty)^{(r)}, (\overline{L}^\infty)^{(r)}_{\eta,qr}\right),$$

$$Orb^K(x; L_*^p(t^{-\theta}), L^\infty(1/t))^{(r)} = Orb^K\left(x; (\overline{L}^\infty)^{(r)}_{\theta,pr}, (L^\infty(1/t))^{(r)}\right).$$

Observe that $(L^\infty)^{(r)} = L^\infty$ and $(L^\infty(1/t))^{(r)} = L^\infty(t^{-1/r})$. Therefore, since $pr \geq 1$, $qr \geq 1$, applying [7, Theorem 2.5] to the Banach couple $(L^\infty, L^\infty(t^{-1/r}))$, we obtain

$$\begin{aligned} Orb\left(x; (\overline{L}^\infty)^{(r)}_{\theta,pr}, (\overline{L}^\infty)^{(r)}_{\eta,qr}\right) &= Orb\left(x; (L^\infty)^{(r)}, (\overline{L}^\infty)^{(r)}_{\eta,qr}\right) \\ &+ Orb\left(x; (\overline{L}^\infty)^{(r)}_{\theta,pr}, (L^\infty(1/t))^{(r)}\right). \end{aligned}$$

Note that all the couples involved here have the uniform Calderón-Mityagin property (see, for instance, [12] or [42]). Hence (see also Section 2.2), we can replace in the last equality the orbits with the K -orbits:

$$\begin{aligned} Orb^K \left(x; \left(\overline{L^\infty}^{(r)} \right)_{\theta, pr}, \left(\overline{L^\infty}^{(r)} \right)_{\eta, qr} \right) &= Orb^K \left(x; (L^\infty)^{(r)}, \left(\overline{L^\infty}^{(r)} \right)_{\eta, qr} \right) \\ &\quad + Orb^K \left(x; \left(\overline{L^\infty}^{(r)} \right)_{\theta, pr}, (L^\infty(1/t))^{(r)} \right), \end{aligned}$$

and combining this with the preceding formulae, we get

$$\begin{aligned} (4.8) \quad Orb^K \left(x; L_*^p(t^{-\theta}), L_*^q(t^{-\eta}) \right)^{(r)} &= Orb^K \left(x; L^\infty, L_*^q(t^{-\eta}) \right)^{(r)} \\ &\quad + Orb^K \left(x; L_*^p(t^{-\theta}), L^\infty(1/t) \right)^{(r)}. \end{aligned}$$

Next, since $1/r \leq \min(p, q)$, the quasi-Banach function couples $(L^\infty, L_*^q(t^{-\eta}))$ and $(L^\infty, L_*^q(t^{-\eta}))$ are $1/r$ -convex. Hence,

$$U := Orb^K \left(x; L^\infty, L_*^q(t^{-\eta}) \right) \quad \text{and} \quad V := Orb^K \left(x; L_*^p(t^{-\theta}), L^\infty(1/t) \right)$$

are $1/r$ -convex quasi-Banach function lattices. Indeed, since the spaces L^∞ and $L_*^q(t^{-\eta})$ are $1/r$ -convex with constant 1, in view of Lemma 3.2, for any $y_1, y_2, \dots, y_n \in U$ we have

$$\begin{aligned} \left\| \left(\sum_{k=1}^n |y_k|^{1/r} \right)^r \right\|_U &= \sup_{t>0} \frac{K \left(t, \left(\sum_{k=1}^n |y_k|^{1/r} \right)^r; L^\infty, L_*^q(t^{-\eta}) \right)}{K(t, x; L^\infty, L_*^q(t^{-\eta}))} \\ &\leq 2^{r-1} \sup_{t>0} \left(\frac{\sum_{k=1}^n K(t, |y_k|; L^\infty, L_*^q(t^{-\eta}))^{1/r}}{K(t, x; L^\infty, L_*^q(t^{-\eta}))^{1/r}} \right)^r \\ &\leq 2^{r-1} \left(\sum_{k=1}^n \left(\sup_{t>0} \frac{K(t, |y_k|; L^\infty, L_*^q(t^{-\eta}))}{K(t, x; L^\infty, L_*^q(t^{-\eta}))} \right)^{1/r} \right)^r \\ &= 2^{r-1} \left(\sum_{k=1}^n \|y_k\|_U^{1/r} \right)^r. \end{aligned}$$

Quite similarly, for any $z_1, z_2, \dots, z_n \in V$,

$$\left\| \left(\sum_{k=1}^n |z_k|^{1/r} \right)^r \right\|_V \leq 2^{r-1} \left(\sum_{k=1}^n \|z_k\|_V^{1/r} \right)^r.$$

From this observation and Proposition 3.4 it follows that $(U + V)^{(r)} = U^{(r)} + V^{(r)}$. Combining this together with (4.8) we get (4.7), which completes the proof of the theorem. $\square$

Remark 4.14. In the paper [1], by using some ideas from [35], it is constructed an example of (Banach) rearrangement invariant spaces of measurable functions B_1, B_p, B_q, B_∞ and A such that $B_p = (B_1, B_\infty)_{1/4, p}$, $B_q = (B_1, B_\infty)_{3/4, q}$, A is an interpolation space with respect to both (B_1, B_q) and (B_q, B_∞) , but A fails to be

an interpolation space with respect to (B_p, B_q) . This indicates that the Arazy-Cwikel formula (4.2) cannot be extended to the class of all Banach couples and hence the conditions of Corollary 4.2, in general, cannot be skipped even in the Banach case. At the same time, note that the example of l^p -spaces, $0 < p \leq \infty$, shows that the latter conditions are not necessary to have (4.2) (see Example 5 below or [2]).

Remark 4.15. By the well-known Wolff's theorem [44] (see also [19] and [6, Theorem 4.5.16]), if X_2, X_3 are intermediate spaces with respect to a quasi-Banach couple $\overline{X} = (X_0, X_1)$ such that $(X_0, X_3)_{\mu, p} = X_2, (X_2, X_1)_{\nu, q} = X_3$ for some $0 < \mu, \nu < 1$ and $0 < p, q \leq \infty$, then $X_2 = \overline{X}_{\theta, p}$ and $X_3 = \overline{X}_{\eta, q}$ for some $0 < \theta < \eta < 1$. This allows to relax somewhat the assumptions of Theorem 4.1 and Corollary 4.4. For instance, we obtain that the equality

$$Int^{KM}(X_2, X_3) = Int^{KM}(X_0, X_3) \cap Int^{KM}(X_2, X_1)$$

holds for intermediate spaces X_2, X_3 with respect to a quasi-Banach couple $\overline{X} = (X_0, X_1)$ whenever $(X_0, X_3)_{\mu, p; K} = X_2$ and $(X_2, X_1)_{\nu, q; K} = X_3$ for some $0 < \mu, \nu < 1, 0 < p, q \leq \infty$.

5. SOME APPLICATIONS

5.1. l^p -spaces. As usual, the space l^p , $0 < p \leq \infty$, consists of all sequences $x = (x_k)_{k=1}^\infty$ such that

$$\|x\|_{l^p} := \left(\sum_{k=1}^{\infty} |x_k|^p \right)^{1/p} < \infty$$

(with the usual modification for $p = \infty$).

Let $0 < s < p < q < r \leq \infty$. Since the couple (l^s, l^r) is mutually closed, from Theorem 4.1 it follows that

$$(5.1) \quad Int^{KM}(l^p, l^q) = Int^{KM}(l^s, l^q) \cap Int^{KM}(l^p, l^r).$$

Note that, if $q \geq 1$, then all the couples from (5.1) have the uniform Calderón-Mityagin property [2, Corollary 4.6] and hence the corresponding classes of K -monotone and interpolation spaces coincide. Therefore, in this case we have

$$(5.2) \quad Int(l^p, l^q) = Int(l^s, l^q) \cap Int(l^p, l^r).$$

Moreover, by [2, Theorem 4.1], (5.2) holds for all $0 < s < p < q < r \leq \infty$. At the same time, if $q < 1$, (l^p, l^q) is not a Calderón-Mityagin couple [2, Theorem 5.3], and so $Int(l^p, l^q) \neq Int^{KM}(l^p, l^q)$ and $Int(l^s, l^q) \neq Int^{KM}(l^s, l^q)$. Thus, if $q < 1$, equalities (5.1) and (5.2) are different.

Next, let us consider a more general case.

5.2. Lorentz $L^{p,q}$ -spaces. Let $L^{p,q}$, $0 < p < \infty, 0 < q \leq \infty$, be the Lorentz spaces of measurable functions on an arbitrary underlying σ -finite measure space

(T, Σ, μ) , which can be defined using the quasi-norms

$$\|f\|_{L^{p,q}} := \left(\int_0^\infty f^*(u)^q d(u^{q/p}) \right)^{1/q} \text{ for } q < \infty,$$

and

$$\|f\|_{L^{p,\infty}} := \operatorname{ess\,sup}_{u>0} (f^*(u)u^{1/p}).$$

Here, f^* is the non-increasing rearrangement of the function $|f|$, i.e.,

$$f^*(u) := \inf\{s > 0 : \mu(\{t \in T : |f(t)| > s\}) \leq u\}, \quad 0 < u < \mu(T).$$

Recall that $L^{p,p} = L^p$, $0 < p < \infty$, isometrically.

Let $0 < p_0 < p_2 < p_3 < p_1 < \infty$ and $0 < q_0, q_1, q_2, q_3 \leq \infty$ be arbitrary. Since $L^{p_i, q_i} = (L^r, L^s)_{\theta_i, q_i}$ for $0 < r < p_i < s < \infty$ and $1/p_i = (1 - \theta_i)/r + \theta_i/s$ [5, Theorem 5.3.1], then from Corollary 3.7 it follows that the couple $(L^{p_0, q_0}, L^{p_1, q_1})$ is mutually closed. Moreover, $L^{p_2, q_2} = (L^{p_0, q_0}, L^{p_1, q_1})_{\theta, q_2}$ and $L^{p_3, q_3} = (L^{p_0, q_0}, L^{p_1, q_1})_{\eta, q_3}$, where $1/p_2 = (1 - \theta)/p_0 + \theta/p_1$ and $1/p_3 = (1 - \eta)/p_0 + \eta/p_1$ [5, Theorem 5.3.1]. Since $0 < \theta < \eta < 1$, by Theorem 4.1, we have

$$\operatorname{Int}^{KM}(L^{p_2, q_2}, L^{p_3, q_3}) = \operatorname{Int}^{KM}(L^{p_0, q_0}, L^{p_3, q_3}) \cap \operatorname{Int}^{KM}(L^{p_2, q_2}, L^{p_1, q_1}).$$

Suppose now that (T, Σ, μ) is the half-line $(0, \infty)$ with the Lebesgue measure. Then, arguing similarly as in the paper [8], one can prove that the couple $(L^{p,q}, L^{r,s})$ has the uniformly Calderón-Mityagin property for all $0 < p < r \leq \infty$ and $0 < q, s \leq \infty$. Therefore, from Corollary 4.2 it follows that for all $0 < p_0 < p_2 < p_3 < p_1 \leq \infty$ and $0 < q_0, q_1, q_2, q_3 \leq \infty$

$$\operatorname{Int}(L^{p_2, q_2}, L^{p_3, q_3}) = \operatorname{Int}(L^{p_0, q_0}, L^{p_3, q_3}) \cap \operatorname{Int}(L^{p_2, q_2}, L^{p_1, q_1}).$$

Observe that the last result was obtained in the case of L^p -spaces in [8, Theorem 1.1].

5.3. Weighted L^p -spaces. Let $0 < p \leq \infty$ and $w(t)$ be a nonnegative measurable function on a σ -finite measure space (T, Σ, μ) . Consider a couple the weighted L^p -spaces $(L^{p_0}(w_0), L^{p_1}(w_1))$, $0 < p_0, p_1 \leq \infty$ (in particular, p_0 and p_1 may be equal).

Let $0 < \theta < \eta < 1$. Assuming that $w_2 = w_0^{1-\theta} w_1^\theta$, $w_3 = w_0^{1-\eta} w_1^\eta$, $1/p_2 = (1 - \theta)/p_0 + \theta/p_1$ and $1/p_3 = (1 - \eta)/p_0 + \eta/p_1$, by [5, Theorem 5.5.1], we get

$$(L^{p_0}(w_0), L^{p_1}(w_1))_{\theta, p_2} = L^{p_2}(w_2) \quad \text{and} \quad (L^{p_0}(w_0), L^{p_1}(w_1))_{\eta, p_3} = L^{p_3}(w_3).$$

In consequence, from Theorem 4.1 it follows

$$\operatorname{Int}^{KM}(L^{p_2}(w_2), L^{p_3}(w_3)) = \operatorname{Int}^{KM}(L^{p_0}(w_0), L^{p_3}(w_3)) \cap \operatorname{Int}^{KM}(L^{p_2}(w_2), L^{p_1}(w_1)).$$

In particular, if $p_0, p_1 \geq 1$, by [42] and Corollary 4.4, we obtain

$$\operatorname{Int}(L^{p_2}(w_2), L^{p_3}(w_3)) = \operatorname{Int}(L^{p_0}(w_0), L^{p_3}(w_3)) \cap \operatorname{Int}(L^{p_2}(w_2), L^{p_1}(w_1))$$

(cf. [7, Theorem 2.6]).

5.4. Hardy spaces. Let $0 < p \leq \infty$ and let $H^p := H^p(\mathbb{R})$ be the (real) Hardy space on the real line (see, for instance, [6, § 3.9.3]). Then, as is well known (see, for instance, [4, Corollary 5.6.16, p. 374]), for all $0 < \theta < 1$ and $1 \leq q \leq \infty$ we have

$$(H^1, L^\infty)_{\theta, q} = L^{p, q}$$

where $1/p = 1 - \theta$. Moreover, the couple (H^1, L^∞) has the uniform Calderón-Mityagin property [41]. Consequently, applying Corollary 4.2 (see also Remark 4.3) for all $1 < p_0 < p_1 \leq \infty$ and $1 \leq q_0, q_1 \leq \infty$ we get

$$\text{Int}(L^{p_0, q_0}, L^{p_1, q_1}) = \text{Int}(H^1, L^{p_1, q_1}) \cap \text{Int}(L^{p_0, q_0}, L^\infty).$$

Next, by [17] (see also [22]), for all $0 < p_0 < p_1 \leq \infty$ and $0 < \theta < 1$ we have

$$(H^{p_0}, H^{p_1})_{\theta, p} = H^p,$$

where $1/p = (1 - \theta)/p_0 + \theta/p_1$. Hence, from Corollary 3.7 and Theorem 4.1 it follows that for all $0 < p_0 < p_2 < p_3 < p_1 \leq \infty$

$$\text{Int}^{KM}(H^{p_2}, H^{p_3}) = \text{Int}^{KM}(H^{p_0}, H^{p_3}) \cap \text{Int}^{KM}(H^{p_2}, H^{p_1}).$$

In particular, since the couple (H^1, H^∞) has the uniform Calderón-Mityagin property [20] (see also [6, p. 684]), from Corollary 4.2 and Remark 4.3 it follows

$$\text{Int}(H^p, H^q) = \text{Int}(H^1, H^q) \cap \text{Int}(H^p, H^\infty)$$

for all $1 < p < q < \infty$.

5.5. Besov spaces. Let $B_{p, q}^s := B_{p, q}^s(\mathbb{R}^n)$ be the Besov spaces on $\mathbb{R}^n$ (see, for instance, [43, § 2.3]). Suppose $-\infty < s_0 < s_1 < \infty$, $0 < \theta < 1$, $s = (1 - \theta)s_0 + \theta s_1$, $0 < q_0, q_1, q \leq \infty$. Then, for all $0 < p \leq \infty$

$$(B_{p, q_0}^{s_0}, B_{p, q_1}^{s_1})_{\theta, q} = B_{p, q}^s$$

(see [43, Theorem 2.4.2]). Hence, from Corollary 3.7 and Theorem 4.1 it follows that for all $\infty < s_0 < s_2 < s_3 < s_1 < \infty$ and $0 < q_0, q_1, q_2, q_3 \leq \infty$ we have

$$\text{Int}^{KM}(B_{p, q_2}^{s_2}, B_{p, q_3}^{s_3}) = \text{Int}^{KM}(B_{p, q_0}^{s_0}, B_{p, q_3}^{s_3}) \cap \text{Int}^{KM}(B_{p, q_2}^{s_2}, B_{p, q_1}^{s_1}).$$

5.6. Lorentz quasi-normed ideals of compact operators. Let H be a separable complex Hilbert space. For every compact operator $A : H \rightarrow H$ let $s(A) = \{s_n(A)\}_{n=1}^\infty$ be the sequence of s -numbers of A determined by the Schmidt expansion [18]. For every $p, q > 0$, the class $\mathfrak{S}^{p, q}$ consists of all compact operators $A : H \rightarrow H$ such that

$$\|A\|_{p, q} := \|s(A)\|_{l^{p, q}} < \infty,$$

where $l^{p, q}$, the discrete version of the function space $L^{p, q}$, consists of all sequences $x = (x_n)_{n=1}^\infty$ of real numbers such that the quasi-norm

$$\|x\|_{l^{p, q}} := \left(\sum_{n=1}^\infty (x_n^*)^q n^{\frac{q}{p}-1} \right)^{\frac{1}{q}}$$

is finite. Here, $x^* = (x_n^*)_{n=1}^\infty$ denotes the nonincreasing permutation of the sequence $(|x_n|)_{n=1}^\infty$.

The classes $\mathfrak{S}^{p,q}$, $p, q > 0$, are two-sided symmetrically quasi-normed ideals in the space of all bounded operators in H , which sometimes are referred to as Lorentz ideals. The classical Schatten-von Neumann ideals $\mathfrak{S}^p$ correspond to the case $p = q$, i.e., $\mathfrak{S}^p = \mathfrak{S}^{p,p}$.

Let $0 < p_0 < p_2 < p_3 < p_1 < \infty$ and $0 < q_0, q_1, q_2, q_3 \leq \infty$ be arbitrary. Since $\mathfrak{S}^{p_i, q_i} = (\mathfrak{S}^r, \mathfrak{S}^s)_{\theta_i, q_i}$ for $0 < r < p_i < s < \infty$ and $1/p_i = (1 - \theta_i)/r + \theta_i/s$ (see, for instance, [5, §7.3], then from Corollary 3.7 it follows that the couple $(\mathfrak{S}^{p_0, q_0}, \mathfrak{S}^{p_1, q_1})$ is mutually closed. Moreover, $\mathfrak{S}^{p_2, q_2} = (\mathfrak{S}^{p_0, q_0}, \mathfrak{S}^{p_1, q_1})_{\theta, q_2}$ and $\mathfrak{S}^{p_3, q_3} = (\mathfrak{S}^{p_0, q_0}, \mathfrak{S}^{p_1, q_1})_{\eta, q_3}$, where $1/p_2 = (1 - \theta)/p_0 + \theta/p_1$ and $1/p_3 = (1 - \eta)/p_0 + \eta/p_1$ [5, Theorem 5.3.1]. Since $0 < \theta < \eta < 1$, by Theorem 4.1, we have

$$Int^{KM}(\mathfrak{S}^{p_2, q_2}, \mathfrak{S}^{p_3, q_3}) = Int^{KM}(\mathfrak{S}^{p_0, q_0}, \mathfrak{S}^{p_3, q_3}) \cap Int^{KM}(\mathfrak{S}^{p_2, q_2}, \mathfrak{S}^{p_1, q_1}).$$

REFERENCES

- [1] J. Arazy and M. Cwikel, *A new characterization of the interpolation spaces between L^p and L^q* . Math. Scand. **55** (1984), 253–270. 1, 1, 4.14
- [2] S. V. Astashkin, M. Cwikel, and P. G. Nilsson, *Arazy-Cwikel and Calderón-Mityagin type properties of the couples (l^p, l^q) , $0 \leq p < q \leq \infty$* . arXiv.org 2106.03083v1 Math.FA (2021). 1, 4.14, 5.1, 5.1
- [3] S. V. Astashkin and L. Maligranda) *Interpolation between L_1 and L_p , $1 < p < \infty$* , Proc. Amer. Math. Soc. **132** (2004), no. 10, 2929–2938. 1
- [4] C. Bennett and R. Sharpley, *Interpolation of operators*. London: Academic Press, 1988. 2.2, 5.4
- [5] J. Bergh and J. Löfström, *Interpolation Spaces. An Introduction*. Springer-Verlag, Berlin, 1976. 2.2, 2.2, 4, 5.2, 5.3, 5.6
- [6] Yu. A. Brudnyi and N. Ya. Kruglyak, *Interpolation Functors and Interpolation Spaces 1*, North-Holland, 1991. 1, 2.2, 2.2, 2, 2.2, 2.3, 2.4, 3, 3, 3, 4, 4.3, 4, 4.6, 4, 4, 4.15, 5.4
- [7] Yu. N. Bykov and V.I. Ovchinnikov, *Interpolation properties of scales of Banach spaces*, Math. Notes **80** (2006), no. 6, 761–769. 1, 1, 4, 4, 4, 4, 5.3
- [8] L. Cadilhac, F. Sukochev, and D. Zanin, *Lorentz-Shimogaki-Arazy-Cwikel theorem revisited*. arXiv.org 2009.02145v1 math.FA (2020). 1, 5.2
- [9] A. P. Calderón, *Spaces between L^1 and L^∞ and the theorem of Marcinkiewicz*. Studia Math. **26** (1966), 273–299. 1
- [10] B. Cuartero and M. A. Triana, *(p, q) -Convexity in quasi-Banach lattices and applications*. Studia Math. **84** (1986), 113–124. 2.1, 2.1
- [11] M. Cwikel, *Monotonicity properties of interpolation spaces*. Ark. Mat. **14** (1976), 213–236. 2.2, 3.8
- [12] M. Cwikel, *Monotonicity properties of interpolation spaces II*. Ark. Mat. **19** (1981), 123–136. 1, 1, 3.8, 4.3, 4
- [13] M. Cwikel and P. Nilsson, *On Calderón-Mityagin couples of Banach lattices*. Proc. Conf. Constructive Theory of Functions (1984). 1, 4
- [14] M. Cwikel and P. G. Nilsson, *Interpolation of weighted Banach lattices*. Memoirs Amer. Math. Soc. **165** (2003) no. 787, 1–105. 2.3
- [15] V. I. Dmitriev, *Theorems on parameters of interpolation method of constants*. DAN USSR, **216** (1974), no. 2, 257–258 (in Russian). 2.4
- [16] V.I. Dmitriev and V. I. Ovchinnikov, *On interpolation in real method spaces*. Dokl. Akad. Nauk. SSSR **246** (1979), 794–797. 4.3
- [17] C. Fefferman, N. M. Rivière and Y. Sagher, *Interpolation between H^p spaces: the real method*. Trans. Amer. Math. Soc. **191** (1974), 75–81. 5.4
- [18] I. C. Gohberg and M. G. Krein, *Introduction to the Theory of Linear Nonselfadjoint Operators*, Amer. Math. Soc., Transl. Math. Monographs, Volume 18, Providence 1969. 5.6
- [19] S. Janson, P. Nilsson, and J. Peetre (with Appendix by M. Zafran), *Notes on Wolff’s note on interpolation spaces*. Proc. London Math. Soc. (3), **48** (1984), 283–299. 4.15
- [20] P. Jones, *On interpolation between H^1 and H^∞* . Lect. Notes Math. Springer, **1070** (1984), 143–151. 5.4
- [21] N. J. Kalton, *Convexity conditions for non-locally convex lattices*. Glasgow Math. J. **25** (1984), 141–152. 2.1, 2.1, 3
- [22] S. V. Kislyakov, *Interpolation of H^p -spaces: some recent developments*. Israel Math. Conf. **13** (1999), 102–140. 5.4
- [23] P. Krée, *Interpolation d’espaces qui ne sont ni normés ni complets. Applications*. Ann. Inst. Fourier (Grenoble) **17** (1967), fasc. 2, 137–174 (1968). 1
- [24] S. G. Krein, Yu. I. Petunin, and E. M. Semenov, *Interpolation of linear operators*. Amer. Math. Soc., Providence, RI, 1982. 2.2, 2

- [25] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces vol. II.* (Springer-Verlag, Berlin, 1979). 1, 2.1, 2.1, 3, 3, 3, 3
- [26] L. Maligranda, *A generalization of the Shimogaki theorem.* Studia Math. **71** (1981), 69–83. 1
- [27] L. Maligranda, *The K -functional for function spaces.* Research report, 33 pages. Poznań. 1983. 3
- [28] L. Maligranda, *Orlicz Spaces and Interpolation.* Seminars in Math. 5. Univ. of Campinas, Campinas SP, Brazil, 1989. 3
- [29] L. Maligranda, *The K -functional for p -convexifications.* Positivity **17** (2013), 707–710. 3
- [30] P. Meyer-Nieberg, *Banach lattices.* Universitext xv + 395 (1991). 2.1
- [31] B. S. Mityagin, *An interpolation theorem for modular spaces.* Proc. Conf. "Interpol. Spaces and Allied Topics in Anal." (Lund, 1983), Lecture Notes in Math. **1070** (1984), 10–23 (translated from Mat. Sb. **66** (1965), 473–482 (in Russian)). 1
- [32] P. Nilsson, *Reiteration theorems for real interpolation and approximation spaces.* Ann. Mat. Pura Appl. **132** (1982), 291–330. 3
- [33] P. Nilsson, *Interpolation of Calderón and Ovčinnikov pairs.* Ann. Mat. Pura Appl. **134** (1983), 201–232. 3, 4.3, 4
- [34] E. T. Oklander, *On interpolation of Banach Spaces.* Ph.D. thesis, University of Chicago, 1964. 1
- [35] V. I. Ovchinnikov, *On estimates of interpolation orbits,* Math. USSR-Sb. **43** (1982), 573–583. 4.14
- [36] V. I. Ovchinnikov, *The Method of Orbits in Interpolation Theory,* Mathematical Reports **1** (1984), 349–516. 2.2, 2.2, 3
- [37] J. Peetre, *Nouvelles propriétés d'espaces d'interpolation.* C. R. Acad. Sci. Paris **256** (1963), 1424–1426 (French). 1
- [38] J. Peetre, *A theory of interpolation of normed spaces.* (Lecture notes, Brasilia, 1963.) Notas de mat. **39** (1968), 1–86. 1
- [39] Y. Raynaud and P. Tradacete, *Calderón-Lozanovskii interpolation on quasi-Banach lattices.* Banach J. Math. Anal. **12** (2018), no. 2, 294–313. 2.3
- [40] W. Rudin, *Functional analysis.* Second edition., MacGraw-Hill, New York, 1991. 2.2
- [41] R. Sharpley, *A characterization of the interpolation spaces of H^1 and L^∞ on the line.* Constr. Approx. **4** (1988), 199–209. 5.4
- [42] G. Sparr, *Interpolation of weighted L_p -spaces.* Studia Math. **62** (1978), 229–271. 1, 4, 5.3
- [43] H. Triebel, *Theory of function spaces.* Birkhäuser Verlag, Basel, Boston, Stuttgart, 1983. 5.5
- [44] T. Wolff, *A note on interpolation spaces,* Proceedings Conference on Harmonic Analysis, Minneapolis, 1981, Lecture Notes in Mathematics 908 (Springer, Berlin, 1982), pp. 199–204.

4.15

ASTASHKIN: DEPARTMENT OF MATHEMATICS, SAMARA NATIONAL RESEARCH UNIVERSITY, MOSKOVSKOYE SHOSSE 34, 443086, SAMARA, RUSSIA

Email address: `astash56@mail.ru`

NILSSON: ROSLAGSGATAN 6, 113 55 STOCKHOLM, SWEDEN

Email address: `pgn@plntx.com`