

A SHORT NONCOMPLEX PROOF FOR THE SOLUTION SET OF HOMOGENEOUS SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS WITH CONSTANT COEFFICIENTS

PATRIK LUNDSTRÖM

ABSTRACT. We provide a short proof, not utilizing complex numbers, for the solution set of homogeneous second order linear differential equations with constant coefficients.

We argue that the solution set of second order linear differential equations with constant coefficients should be presented using hyperbolic functions in the distinct real root case:

Theorem. Let $a, b \in \mathbb{R}$. Put $\alpha = a/2$ and $\beta = \sqrt{|a^2/4 - b|}$. All solutions to $y'' + ay' + by = 0$ are given by (1) $y = e^{-\alpha x}(C_1 \cosh(\beta x) + C_2 \sinh(\beta x))$, if $a^2/4 > b$, (2) $y = e^{-\alpha x}(C_1 + C_2 x)$, if $a^2/4 = b$, and (3) $y = e^{-\alpha x}(C_1 \cos(\beta x) + C_2 \sin(\beta x))$, if $a^2/4 < b$, for arbitrary $C_1, C_2 \in \mathbb{R}$.

There are many reasons for doing so. First of all, this presentation highlights the beautiful symmetry between (1) and (3). Secondly, if we in (1) and (3) use the first order approximations $\cosh(\beta x) \approx 1$, $\sinh(\beta x) \approx \beta x$, $\cos(\beta x) \approx 1$ and $\sin(\beta x) \approx \beta x$, for small β , then (2) can be seen, qualitatively, as a limiting case of both (1) and (3), as $\beta \rightarrow 0$. Thirdly, and perhaps most importantly, utilizing a condensed version of a clever trick by Dobos [1], and here expanding it to the real case also, one can construct a short coherent elementary noncomplex proof of (1)-(3).

Namely, it is a straightforward calculation to show that the expressions in (1), (2) and (3) indeed are solutions to the differential equation. Now suppose that y is an arbitrary solution and put $z = e^{\alpha x}y$. Then $z' = \alpha z + e^{\alpha x}y'$ and thus $z'' = \alpha^2 z + 2\alpha e^{\alpha x}y' + e^{\alpha x}y'' = e^{\alpha x}((a^2/4)y + ay' + y'')$.

(1) Let $a^2/4 > b$. Then $z'' - \beta^2 z = e^{\alpha x}((a^2/4)y + ay' + y'' - (a^2/4 - b)y) = e^{\alpha x}(y'' + ay' + by) = 0$ and hence $(-z \sinh(\beta x) + z' \cosh(\beta x)/\beta)' = 0$ and $(-z' \sinh(\beta x)/\beta + z \cosh(\beta x))' = 0$ which implies that $-z \sinh(\beta x) + z' \cosh(\beta x)/\beta = C_2$ and $-z' \sinh(\beta x)/\beta + z \cosh(\beta x) = C_1$ for some $C_1, C_2 \in \mathbb{R}$. Therefore $C_1 \cosh(\beta x) + C_2 \sinh(\beta x) = z \cosh^2(\beta x) - z \sinh^2(\beta x) = z = e^{\alpha x}y$ and thus finally we get that $y = e^{-\alpha x}z = e^{-\alpha x}(C_1 \cosh(\beta x) + C_2 \sinh(\beta x))$.

(2) Suppose that $a^2/4 = b$. Then $z'' = e^{\alpha x}((a^2/4)y + ay' + y'') = e^{\alpha x}(y'' + ay' + by) = 0$. Hence $z = C_1 + C_2 x$ for some $C_1, C_2 \in \mathbb{R}$ and therefore we get that $y = e^{-\alpha x}z = e^{-\alpha x}(C_1 + C_2 x)$.

(3) Let $a^2/4 < b$. Then $z'' + \beta^2 z = e^{\alpha x}((a^2/4)y + ay' + y'' + (b - a^2/4)y) = e^{\alpha x}(y'' + ay' + by) = 0$ and hence $(z \sin(\beta x) + z' \cos(\beta x)/\beta)' = 0$ and $(-z' \sin(\beta x)/\beta + z \cos(\beta x))' = 0$ which implies that $z \sin(\beta x) + z' \cos(\beta x)/\beta = C_2$ and $-z' \sin(\beta x)/\beta + z \cos(\beta x) = C_1$ for some $C_1, C_2 \in \mathbb{R}$. Therefore $C_1 \cos(\beta x) + C_2 \sin(\beta x) = z \cos^2(\beta x) + z \sin^2(\beta x) = z = e^{\alpha x}y$ and thus finally we get that $y = e^{-\alpha x}z = e^{-\alpha x}(C_1 \cos(\beta x) + C_2 \sin(\beta x))$. \square

Note that the solutions in (1) above of course easily can be rewritten in the familiar form $y = e^{-\alpha x}(C_1 \cosh(\beta x) + C_2 \sinh(\beta x)) = D_1 e^{r_1 x} + D_2 e^{r_2 x}$ where $D_1 = (C_1 + C_2)/2$, $D_2 = (C_1 - C_2)/2$ and $r_1 = -\alpha + \beta$ and $r_2 = -\alpha - \beta$ are the two roots of the characteristic equation $r^2 + ar + b = 0$.

REFERENCES

- [1] J. Dobos, ODE without complex numbers: another look. *Teach. Math. Appl.* **26**, 196–200 (2007).

UNIVERSITY WEST, DEPARTMENT OF ENGINEERING SCIENCE, SE-46186 TROLLHÄTTAN, SWEDEN
Email address: patrik.lundstrom@hv.se