

# The intrinsic Toeplitz structure and its applications in algebraic Riccati equations

Zhen-Chen Guo\*      Xin Liang†

December 3, 2021

## Abstract

In this paper we propose a new algorithm for solving large-scale algebraic Riccati equations with low-rank structure. The algorithm is based on the found Toeplitz-structured closed form of the stabilizing solution and the fast Fourier transform. It works without unnecessary assumptions, shift selection strategies, or matrix calculations of the cubic order with respect to the problem scale. Numerical examples are given to illustrate its features. Besides, we show that it is theoretically equivalent to several algorithms existing in the literature in the sense that they all produce the same sequence under the same parameter setting.

**Key words.** Toeplitz matrix, FFT, algebraic Riccati equations, large-scale, low-rank

**AMS subject classifications.** 15A24, 15B05, 65F45, 93B52

## 1 Introduction

Consider a continuous-time algebraic/limiting Riccati equation (CARE)

$$A^T X + X A - X B B^T X + C^T C = 0, \quad (1.1)$$

where  $A \in \mathbb{R}^{n \times n}$ ,  $B \in \mathbb{R}^{n \times m}$ ,  $C \in \mathbb{R}^{l \times n}$ . The CAREs arise in various models related to control theory, such as linear-quadratic optimal regulator design, and  $H^2$  and  $H^\infty$  controller design for linear systems, see, e.g., [35, 3]. They also arise in nonlinear systems, like nonlinear controller design by state-dependent Riccati equations [13], or solving differential Riccati equations by implicit integration schemes [17, 8]. Usually (1.1) has infinite many solutions, but in many applications including those mentioned above only the so-called c-stabilizing solution is hoped to be chased. Here a solution  $X$  is called c-stabilizing if  $A - B B^T X$  is stable, namely all the eigenvalues of  $A - B B^T X$  lies in the open left half complex plane  $\mathbb{C}_-$ . Its existence and uniqueness are guaranteed by the assumption that the pairs  $(A, B B^T)$  and  $(A^T, C^T C)$  are c-stabilizable, or equivalently,  $\text{rank}([A - \lambda I \quad B B^T]) = \text{rank}([A^T - \lambda I \quad C^T C]) = n$  for any  $\lambda \in \mathbb{C} \setminus \mathbb{C}_-$ .

During many years, people have developed many numerical methods to find out the c-stabilizing solution of (1.1). Reader are referred to [12] to obtain an overview. In this paper, we are focusing on a special case that  $A$  is large-scale and sparse, and  $B, C$  are low-rank, namely  $m, l \ll n$ . The existing methods are categorized into four classes:

1. projection methods, including extended Krylov subspace method [23], rational Krylov subspace method [18], tangential rational Krylov subspace method [19], global extended Krylov subspace method [27], etc.;
2. non-projective iterations, including quadratic ADI [44], Cayley transformed Hamiltonian subspace iteration [38], RADI [5], etc.;
3. Newton-type methods, including the Galerkin projected variant of Newton-Kleinman ADI [9] and its inexact line-search variant [6], etc.;
4. methods adopted from those suited for small-scale problems, including structure-preserving doubling algorithm (SDA) [14, 36], and Hamiltonian stable subspace methods [1, 4], etc..

---

\*Department of Mathematics, Nanjing University, Nanjing 210093, China; e-mail: guozhenchen@nju.edu.cn. Supported in part by NSFC-11901290 and Fundamental Research Funds for the Central Universities.

†Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China, and Yanqi Lake Beijing Institute of Mathematical Sciences and Applications, Beijing 101408, China; e-mail: liangxinslm@tsinghua.edu.cn. Supported in part by NSFC-11901340.



For ease,  $\mathcal{L}_{p_1 \times p_2}(A) = \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} A_0 \\ A_1 \\ \vdots \\ A_{m-1} \end{bmatrix} \right)$  if  $A = \begin{bmatrix} A_0 \\ A_1 \\ \vdots \\ A_{m-1} \end{bmatrix}$ , and this notation makes no confusion for the subscript  $p_1 \times p_2$  demonstrates how the matrix is composed. Similarly,

$$\mathcal{U}_{p_1 \times p_2} \left( \begin{bmatrix} A_0 \\ A_1 \\ \vdots \\ A_{m-1} \end{bmatrix} \right) = \begin{bmatrix} A_{m-1} & \cdots & \cdots & A_2 & A_1 & A_0 \\ & A_{m-1} & \ddots & & A_2 & A_1 \\ & & \ddots & \ddots & & A_2 \\ & & & \ddots & \ddots & \vdots \\ & & & & A_{m-1} & \vdots \\ & & & & & A_{m-1} \end{bmatrix} \in \mathbb{R}^{p_1 m \times p_2 m}.$$

Besides,

$$\begin{aligned} \mathcal{L}_{p_1 \times p_2} \left( [A_0 \ A_1 \ \cdots \ A_{m-1}]^T \right)^T &= \mathcal{U}_{p_1 \times p_2} \left( \begin{bmatrix} A_{m-1} \\ \vdots \\ A_1 \\ A_0 \end{bmatrix} \right), \\ \mathcal{U}_{p_1 \times p_2} \left( [A_0 \ A_1 \ \cdots \ A_{m-1}]^T \right)^T &= \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} A_{m-1} \\ \vdots \\ A_1 \\ A_0 \end{bmatrix} \right). \end{aligned}$$

The following lemma will be used several times later.

**Lemma 2.1.** *Given  $Y \in \mathbb{R}^{p_1 \times p_2}$ ,  $Y^L \in \mathbb{R}^{p_2 \times p_1}$ ,  $D_{t-1} \in \mathbb{R}^{p_1(t-1) \times p_2}$ ,  $D_{t-1}^L \in \mathbb{R}^{p_2 \times p_1(t-1)}$ , let*

$$T_t = \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} Y \\ D_{t-1} \end{bmatrix} \right) = \begin{bmatrix} Y & 0 \\ D_{t-1} & T_{t-1} \end{bmatrix} \in \mathbb{R}^{p_1 t \times p_2 t}, \quad T_t^L = \mathcal{L}_{p_1 \times p_2} \left( [Y^L \ D_{t-1}^L]^T \right)^T = \begin{bmatrix} Y^L & D_{t-1}^L \\ 0 & T_{t-1}^L \end{bmatrix} \in \mathbb{R}^{p_2 t \times p_1 t}.$$

If  $I - T_t T_t^L$  is nonsingular, then

$$\begin{aligned} (I - T_t T_t^L)^{-1} &= \mathcal{U}_{p_1 \times p_1} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p_1 \times p_1} \left( [Q_2^L \ Q_{2,b}]^T \right)^T \\ &\quad + \mathcal{U}_{p_1 \times p_2} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes [W + WY^L YW])^{-1} \mathcal{U}_{p_1 \times p_2} \left( [Q_3^L \ 0]^T \right)^T, \end{aligned} \quad (2.1a)$$

where the following equations are solvable and  $Q_{2,b}, W + WY^L YW$  are nonsingular:

$$Q_3^L [I - D_{t-1} D_{t-1}^L - T_{t-1} T_{t-1}^L] = D_{t-1}^L, \quad (2.1b)$$

$$[I - D_{t-1} D_{t-1}^L - T_{t-1} T_{t-1}^L] Q_3 = D_{t-1}, \quad W = -(I + Q_3^L D_{t-1}), \quad (2.1c)$$

$$[Q_2^L \ Q_{2,b}] (I - T_t T_t^L) = [0 \ I_{p_1}], \quad (2.1d)$$

$$(I - T_t T_t^L) \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} = \begin{bmatrix} 0 \\ I_{p_1} \end{bmatrix}, \quad Q_{2,b} \in \mathbb{R}^{p_1 \times p_1}. \quad (2.1e)$$

### 3 DARE

Given a linear time-invariant control system in discrete-time:

$$\begin{aligned} x_0 &\text{ is given,} \\ x_{k+1} &= Ax_k + Bu_k, \quad k = 0, 1, 2, \dots, \\ y_k &= Cx_k, \end{aligned} \quad (3.1)$$

where  $A \in \mathbb{R}^{n \times n}$ ,  $B \in \mathbb{R}^{n \times m}$ ,  $C \in \mathbb{R}^{l \times n}$ . Suppose the following condition holds through out this section:

$$\boxed{(A, B) \text{ is d-stabilizable and } (C, A) \text{ is d-detectable,}}$$

or equivalently,  $\text{rank}([A - \lambda I \quad BB^T]) = \text{rank}([A^T - \lambda I \quad C^T C]) = n$  for any  $\lambda \in \mathbb{C} \setminus \mathbb{D}$ , where  $\mathbb{D}$  is the open unit disk.

Its linear-quadratic optimal control can be expressed as

$$\arg \min_{\{u_k\}} \frac{1}{2} \sum_{k=0}^{\infty} (y_k^T y_k + u_k^T u_k) = \{u_k = -(I + B^T X_* B)^{-1} B^T X_* A x_k\}, \quad (3.2)$$

where  $X_*$  is the unique symmetric positive semi-definite d-stabilizing solution of the DARE [12, 15, 34, 39]:

$$-X + A^T X (I + BB^T X)^{-1} A + C^T C = 0. \quad (3.3)$$

Here a solution  $X$  is called d-stabilizing, if the closed loop matrix  $A_X = (I + BB^T X)^{-1} A$  is d-stable, namely all of its eigenvalues lie in the open unit disk  $\mathbb{D}$ , or equivalently,  $\rho(A_X) < 1$ .

### 3.1 In the operator view

In this section, we briefly state the existence and uniqueness shown by the operator theory, which is based on the monograph [26].

In order to make things simple, first we assume that  $A$  is d-stable. Write  $\mathbf{x} = \{x_k\}_{k \in \mathbb{N}}$ ,  $\mathbf{u} = \{u_k\}_{k \in \mathbb{N}}$ ,  $\mathbf{y} = \{y_k\}_{k \in \mathbb{N}}$ . Let  $\ell_+^{2,n}$  denote the Hilbert space of norm-square summable  $\mathbb{R}^n$ -valued series.

Suppose  $\mathbf{u} \in \ell_+^{2,m}$  and consider the cost functional (aka restricted quadratic index)

$$J(\mathbf{u}) = \sum_{k=0}^{+\infty} \begin{bmatrix} x_k \\ u_k \end{bmatrix}^T \begin{bmatrix} Q & L \\ L^T & R \end{bmatrix} \begin{bmatrix} x_k \\ u_k \end{bmatrix} = \left\langle \begin{bmatrix} \mathbf{x} \\ \mathbf{u} \end{bmatrix}, \begin{bmatrix} Q & L \\ L^T & R \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{u} \end{bmatrix} \right\rangle_{\ell_+^{2,n} \times \ell_+^{2,m}},$$

where  $\mathbf{x}$  is the solution to (3.1). Here for any matrix  $U$  and any series  $\mathbf{z} = \{z_k\}_{k \in \mathbb{N}}$ ,  $U\mathbf{z}$  is understood as  $U\mathbf{z} := \{Uz_k\}_{k \in \mathbb{N}}$ .

In fact  $\mathbf{x} = \mathcal{F}x_0 + \mathcal{L}\mathbf{u} \in \ell_+^{2,n}$ , where  $\mathcal{F}: \mathbb{R}^n \rightarrow \ell_+^{2,n}$ ,  $(\mathcal{F}x_0)_k = A^k x_0$ ,  $k \geq 0$ , and  $\mathcal{L}: \ell_+^{2,m} \rightarrow \ell_+^{2,n}$ ,  $(\mathcal{L}\mathbf{u})_0 = 0$ ,  $(\mathcal{L}\mathbf{u})_k = \sum_{i=0}^{k-1} A^{k-i-1} B u_i$ ,  $k \geq 1$ . Clearly  $\mathcal{F}$  and  $\mathcal{L}$  are bounded linear operators. Also, it is not difficult to find  $\mathcal{L}$  is a Toeplitz operator. Hence

$$\begin{aligned} J(\mathbf{u}) &= \left\langle \begin{bmatrix} \mathbf{x} \\ \mathbf{u} \end{bmatrix}, \begin{bmatrix} Q & L \\ L^T & R \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{u} \end{bmatrix} \right\rangle \\ &= \left\langle \begin{bmatrix} \mathcal{F} & \mathcal{L} \\ I & \end{bmatrix} \begin{bmatrix} x_0 \\ \mathbf{u} \end{bmatrix}, \begin{bmatrix} Q & L \\ L^T & R \end{bmatrix} \begin{bmatrix} \mathcal{F} & \mathcal{L} \\ I & \end{bmatrix} \begin{bmatrix} x_0 \\ \mathbf{u} \end{bmatrix} \right\rangle \\ &= \left\langle \begin{bmatrix} x_0 \\ \mathbf{u} \end{bmatrix}, \begin{bmatrix} \mathcal{P}_o & \mathcal{P} \\ \mathcal{P}^* & \mathcal{R} \end{bmatrix} \begin{bmatrix} x_0 \\ \mathbf{u} \end{bmatrix} \right\rangle \quad (\mathcal{A}^* \text{ is the adjoint of operator } \mathcal{A}) \end{aligned}$$

where  $\mathcal{P}_o = \mathcal{F}^* Q \mathcal{F}$ ,  $\mathcal{P} = \mathcal{F}^* (Q \mathcal{L} + L)$ ,  $\mathcal{R} = R + L^T \mathcal{L} + \mathcal{L}^* L + \mathcal{L}^* Q \mathcal{L}$ . Then the unique symmetric d-stabilizing solution  $X_*$  is given by

$$X_* = \mathcal{P}_o - \mathcal{P} \mathcal{R}^{-1} \mathcal{P}^*. \quad (3.4)$$

Clearly,  $\mathcal{P}_o, \mathcal{P}, \mathcal{R}$  are bounded linear operators. [26, Theorem 4.7.1] tells that the DARE (3.3) has a unique d-stabilizing solution, if and only if the Toeplitz-like operator  $\mathcal{R}$  has a bounded inverse.

Moreover, if  $A$  is not d-stable, a similar result still holds. Since  $(A, B)$  is d-stabilizable, there exists  $F$  such that  $\tilde{A} = A + BF$  is d-stable. Note the system defined by  $A, B, Q = C^T C, L = 0, R = I$  is equivalent to the system defined by  $\tilde{A}, \tilde{B} = B, \tilde{Q} = C^T C + F^T F, \tilde{L} = F^T, \tilde{R} = I$  in the sense that they share the same d-stabilizing solution. Hence

$$X_* = \tilde{\mathcal{P}}_o - \tilde{\mathcal{P}} \tilde{\mathcal{R}}^{-1} \tilde{\mathcal{P}}^*. \quad (3.5)$$

In [26], the authors did not provide the (infinite) matrix representation of (3.4) or (3.5). Let us make it. First consider the case that  $A$  is d-stable. Clearly, the matrix representations of  $\mathcal{F}, \mathcal{L}$ , still denoted by  $\mathcal{F}, \mathcal{L}$ , are

$$\mathcal{F} = \begin{bmatrix} I \\ A \\ A^2 \\ A^3 \\ \vdots \end{bmatrix}, \quad \mathcal{L} = \begin{bmatrix} 0 & & & & \\ B & 0 & & & \\ AB & B & 0 & & \\ A^2 B & AB & B & 0 & \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{bmatrix}.$$

Hence

$$X_* = \mathcal{F}^* Q \mathcal{F} - \mathcal{F}^* (Q \mathcal{L} + L) (R + L^T \mathcal{L} + \mathcal{L}^* L + \mathcal{L}^* Q \mathcal{L})^{-1} (\mathcal{L}^* Q^T + L^T) \mathcal{F}$$

$$\begin{aligned}
&= \mathcal{F}^* C^T C \mathcal{F} - \mathcal{F}^* C^T C \mathcal{L} (I + \mathcal{L}^* C^T C \mathcal{L})^{-1} \mathcal{L}^* C^T C \mathcal{F} \\
&= \mathcal{F}^* C^T [I - C \mathcal{L} (I + \mathcal{L}^* C^T C \mathcal{L})^{-1} \mathcal{L}^* C^T] C \mathcal{F} \\
&\stackrel{(1.3)}{=} \mathcal{F}^* C^T (I + C \mathcal{L} \mathcal{L}^* C^T)^{-1} C \mathcal{F}.
\end{aligned}$$

Write

$$\mathcal{V} = C \mathcal{F} = \begin{bmatrix} C \\ CA \\ CA^2 \\ CA^3 \\ \vdots \end{bmatrix}, \quad \mathcal{T} = C \mathcal{L} = \begin{bmatrix} 0 & & & & \\ CB & 0 & & & \\ CAB & CB & 0 & & \\ CA^2 B & CAB & CB & 0 & \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{bmatrix}, \quad (3.6)$$

and then  $\mathcal{V}: \mathbb{R}^n \rightarrow \ell_+^{2,l}$ ,  $\mathcal{T}: \ell_+^{2,m} \rightarrow \ell_+^{2,l}$ ,  $(I + \mathcal{T} \mathcal{T}^*)^{-1}: \ell_+^{2,l} \rightarrow \ell_+^{2,l}$  are bounded linear operators, and  $\mathcal{T}$  is also a Toeplitz operator. Thus

$$X_\star = \mathcal{V}^* (I + \mathcal{T} \mathcal{T}^*)^{-1} \mathcal{V}, \quad (3.7)$$

which is a closed form of the solution.

Then consider the case  $A$  is not d-stable.

$$\begin{aligned}
X_\star &= \widetilde{\mathcal{F}}^* \widetilde{Q} \widetilde{\mathcal{F}} - \widetilde{\mathcal{F}}^* (\widetilde{Q} \widetilde{\mathcal{L}} + \widetilde{L}) (\widetilde{R} + \widetilde{L}^T \widetilde{\mathcal{L}} + \widetilde{\mathcal{L}}^* \widetilde{L} + \widetilde{\mathcal{L}}^* \widetilde{Q} \widetilde{\mathcal{L}})^{-1} (\widetilde{\mathcal{L}}^* \widetilde{Q}^T + \widetilde{L}^T) \widetilde{\mathcal{F}} \\
&= \widetilde{\mathcal{F}}^* [C^T C + F^T F] \widetilde{\mathcal{F}} \\
&\quad - \widetilde{\mathcal{F}}^* [(C^T C + F^T F) \widetilde{\mathcal{L}} + F^T] (I + F \widetilde{\mathcal{L}} + \widetilde{\mathcal{L}}^* F^T + \widetilde{\mathcal{L}}^* [C^T C + F^T F] \widetilde{\mathcal{L}})^{-1} (\widetilde{\mathcal{L}}^* [C^T C + F^T F] + F) \widetilde{\mathcal{F}} \\
&= \widetilde{\mathcal{F}}^* \begin{bmatrix} F \\ C \end{bmatrix}^T \begin{bmatrix} F \\ C \end{bmatrix} \widetilde{\mathcal{F}} - \widetilde{\mathcal{F}}^* \begin{bmatrix} F \\ C \end{bmatrix}^T \begin{bmatrix} I + F \widetilde{\mathcal{L}} \\ C \widetilde{\mathcal{L}} \end{bmatrix} \left( \begin{bmatrix} I + F \widetilde{\mathcal{L}} \\ C \widetilde{\mathcal{L}} \end{bmatrix}^* \begin{bmatrix} I + F \widetilde{\mathcal{L}} \\ C \widetilde{\mathcal{L}} \end{bmatrix} \right)^{-1} \begin{bmatrix} I + F \widetilde{\mathcal{L}} \\ C \widetilde{\mathcal{L}} \end{bmatrix}^* \begin{bmatrix} F \\ C \end{bmatrix} \widetilde{\mathcal{F}} \\
&\stackrel{(1.3)}{=} \widetilde{\mathcal{F}}^* \begin{bmatrix} F \\ C \end{bmatrix}^T (I - \mathcal{Q} (\mathcal{Q}^* \mathcal{Q})^{-1} \mathcal{Q}^*) \begin{bmatrix} F \\ C \end{bmatrix} \widetilde{\mathcal{F}} \quad \left( \text{write } \mathcal{Q} = \begin{bmatrix} I + F \widetilde{\mathcal{L}} \\ C \widetilde{\mathcal{L}} \end{bmatrix} \right).
\end{aligned}$$

Clearly  $I - \mathcal{Q} (\mathcal{Q}^* \mathcal{Q})^{-1} \mathcal{Q}^*$  is an orthogonal projection operator onto  $\mathcal{R}(\mathcal{Q})^\perp$ , the orthogonal complement of the image set of  $\mathcal{Q}$ . We temporarily forget the convergence of series and the boundedness of operators. Then  $C \widetilde{\mathcal{L}} = C \mathcal{L} (I + F \widetilde{\mathcal{L}})$ , because the three linear operators  $C \widetilde{\mathcal{L}}, C \mathcal{L}, I + F \widetilde{\mathcal{L}}$  are all Toeplitz operators, and the generating polynomials satisfy

$$\begin{aligned}
&(CB\lambda + CAB\lambda^2 + CA^2B\lambda^3 + \dots)(I + FB\lambda + F(A + BF)B\lambda^2 + F(A + BF)^2B\lambda^2 + \dots) \\
&= CB\lambda + (CBFB + CAB)\lambda^2 + (CBF(A + BF)B + CABFB + CA^2B)\lambda^3 + \dots \\
&= CB\lambda + C(A + BF)B\lambda^2 + C(A + BF)^2B\lambda^3 + \dots.
\end{aligned}$$

Hence  $\mathcal{Q} = \begin{bmatrix} I + F \widetilde{\mathcal{L}} \\ C \mathcal{L} (I + F \widetilde{\mathcal{L}}) \end{bmatrix} = \begin{bmatrix} I \\ C \mathcal{L} \end{bmatrix} (I + F \widetilde{\mathcal{L}})$  where  $I + F \widetilde{\mathcal{L}}$  is invertible. Since

$$\begin{bmatrix} I & -\mathcal{L}^* C^T \\ C \mathcal{L} & I \end{bmatrix} = \begin{bmatrix} I & \\ C \mathcal{L} & I \end{bmatrix} \begin{bmatrix} I & \\ & I + C \mathcal{L} \mathcal{L}^* C^T \end{bmatrix} \begin{bmatrix} I & -\mathcal{L}^* C^T \\ & I \end{bmatrix}$$

is invertible, and  $\begin{bmatrix} I \\ C \mathcal{L} \end{bmatrix}^* \begin{bmatrix} -\mathcal{L}^* C^T \\ I \end{bmatrix} = 0$ , we know that  $\mathcal{Q}_\perp := \begin{bmatrix} -\mathcal{L}^* C^T \\ I \end{bmatrix}$  satisfies  $\mathcal{Q}_\perp (\mathcal{Q}_\perp^* \mathcal{Q}_\perp)^{-1} \mathcal{Q}_\perp^* = I - \mathcal{Q} (\mathcal{Q}^* \mathcal{Q})^{-1} \mathcal{Q}^*$ . Similarly we have  $C \widetilde{\mathcal{F}} = C \mathcal{F} + C \mathcal{L} F \widetilde{\mathcal{F}}$ . Thus

$$\begin{aligned}
X_\star &= \begin{bmatrix} F \widetilde{\mathcal{F}} \\ C \mathcal{F} + C \mathcal{L} F \widetilde{\mathcal{F}} \end{bmatrix}^* \begin{bmatrix} -\mathcal{L}^* C^T \\ I \end{bmatrix} (I + C \mathcal{L} \mathcal{L}^* C^T)^{-1} \begin{bmatrix} -\mathcal{L}^* C^T \\ I \end{bmatrix}^* \begin{bmatrix} F \widetilde{\mathcal{F}} \\ C \mathcal{F} + C \mathcal{L} F \widetilde{\mathcal{F}} \end{bmatrix} \\
&= \mathcal{F}^* C^T (I + C \mathcal{L} \mathcal{L}^* C^T)^{-1} C \mathcal{F} \\
&= \mathcal{V}^* (I + \mathcal{T} \mathcal{T}^*)^{-1} \mathcal{V},
\end{aligned}$$

where  $\mathcal{V}, \mathcal{T}$  are as in (3.6), which implies that the solution in the unstable case has the same form as that in the stable case. However, is this true, considering the convergence of series? In the next subsection, we will show the validity of (3.6).

### 3.2 In the matrix view

It is well known that  $X_\star = \lim_{t \rightarrow \infty} X_t$ , where  $X_t$  is generated by the difference Riccati equation (DRE):

$$X_0 = 0, \quad X_{t+1} = \mathcal{D}(X_t) := C^T C + A^T X_t (I + B B^T X_t)^{-1} A. \quad (3.8)$$

which can be recognized as a variant of fixed point iteration for (3.3).

Based on the fixed point iteration, in 1970s, people have developed the doubling algorithm to solve DAREs (3.3) and CAREs. Anderson [2] proposed a variant, which is recently usually called SDA and has three iterative recursions:

$$A_{k+1} = A_k (I_n + G_k H_k)^{-1} A_k, \quad (3.9a)$$

$$G_{k+1} = G_k + A_k (I_n + G_k H_k)^{-1} G_k A_k^T, \quad (3.9b)$$

$$H_{k+1} = H_k + A_k^T H_k (I_n + G_k H_k)^{-1} A_k, \quad (3.9c)$$

provided that all matrix inversions are feasible (i.e.,  $I_n + G_k H_k$  are nonsingular for  $k = 0, 1, \dots$ ). The initial terms are usually set by

$$A_0 = A, \quad G_0 = B B^T, \quad H_0 = C^T C.$$

It has been shown that for those initial terms,  $I_n + G_k H_k$  are nonsingular for  $k \geq 0$ , and  $A_k \rightarrow 0$ ,  $G_k \rightarrow Y_\star$  (the solution to the dual DARE) and  $H_k \rightarrow X_\star$ , all quadratically [39] except for the critical case [25].

In [2], it is stated clearly that  $H_k = X_{2^k}$ , implying that the iteration for  $H_k$  can be treated as an acceleration of (3.8), because it only computes the terms  $X_1, X_2, X_4, \dots, X_{2^k}, \dots$ . Moreover, [2] also argued that (3.8) with any initial  $X_0 \succeq 0$  leads  $X_t \rightarrow X_\star$  in usual situation (but did not mention which situation satisfies).

Questions arise naturally, of which two are:

1. can we even only compute less terms in the sequence  $\{X_t\}$ , namely accelerate (3.8) even further?
2. how things go when arbitrary initial terms are set?

Before we begin the analysis, an easy property of the DRE (3.8) is given.

**Lemma 3.1.** *The operator  $\mathcal{D}$  is monotonic on the set consisting of all positive semi-definite matrices with respect to the partial order “ $\succeq$ ”. In details, if  $Z_1 \succeq Z_2 \succeq 0$ , then*

$$Z_1 \succeq Z_2 \Rightarrow \mathcal{D}(Z_1) \succeq \mathcal{D}(Z_2).$$

*Proof.* First suppose  $Z_2 \succ 0$  and thus  $Z_2$  is nonsingular. Then

$$\begin{aligned} Z_1 \succeq Z_2 &\Leftrightarrow Z_1^{-1} \preceq Z_2^{-1} \Leftrightarrow (Z_1^{-1} + B B^T)^{-1} \succeq (Z_2^{-1} + B B^T)^{-1} \\ &\Leftrightarrow Z_1 (I + B B^T Z_1)^{-1} \succeq Z_2 (I + B B^T Z_2)^{-1} \Rightarrow \mathcal{D}(Z_1) \succeq \mathcal{D}(Z_2). \end{aligned}$$

If  $Z_2$  is singular, then  $Z_2 + \varepsilon I \succ 0$  for any  $\varepsilon > 0$ . Thus, taking limits,

$$Z_1 \succeq Z_2 \Leftrightarrow Z_1 + \varepsilon I \succeq Z_2 + \varepsilon I \Rightarrow \mathcal{D}(Z_1 + \varepsilon I) \succeq \mathcal{D}(Z_2 + \varepsilon I) \Rightarrow \mathcal{D}(Z_1) \succeq \mathcal{D}(Z_2). \quad \square$$

Let us observe the first several iterations of the DRE (3.8). Clearly  $X_0 = 0, X_1 = C^T C$ . Since  $X_1 \succeq X_0 = 0$ , by Lemma 3.1,  $X_2 = \mathcal{D}(X_1) \succeq \mathcal{D}(X_0) = X_1$ . Similarly  $0 = X_0 \preceq X_1 \preceq X_2 \preceq \dots \preceq X_t \preceq \dots$ , namely the sequence  $\{X_t\}$  generated by (3.8) is monotonic. On the other hand, the d-stabilizing solution  $X_\star \succeq 0 = X_0$ , and thus  $X_\star = \mathcal{D}(X_\star) \succeq \mathcal{D}(X_0) = X_1$ . Similarly  $X_\star \succeq X_t$  for any  $t$ , namely the sequence  $\{X_t\}$  is bounded. As a result, it holds that  $X_t \rightarrow X_\star$ .

Using some calculations, we have

$$\begin{aligned} X_2 &= C^T C + A^T C^T C (I + B B^T C^T C)^{-1} A \\ &= C^T C + A^T C^T (I + C B B^T C^T)^{-1} C A \\ &= \begin{bmatrix} C \\ C A \end{bmatrix}^T \begin{bmatrix} I & \\ & I + C B B^T C^T \end{bmatrix}^{-1} \begin{bmatrix} C \\ C A \end{bmatrix} \\ &= \begin{bmatrix} C \\ C A \end{bmatrix}^T \left( I + \begin{bmatrix} 0 & \\ C B & 0 \end{bmatrix} \begin{bmatrix} 0 & \\ C B & 0 \end{bmatrix}^T \right)^{-1} \begin{bmatrix} C \\ C A \end{bmatrix}. \end{aligned}$$

This encourages us to use an ansatz  $X_t = V_t^T (I + T_t T_t^T)^{-1} V_t$ . Note that this ansatz is always valid because  $X_t \succeq 0$ . The following theorem, namely Theorem 3.1, gives one of the detailed forms of the ansatz.

**Theorem 3.1.** Write

$$V_t = \begin{bmatrix} C \\ CA \\ CA^2 \\ \vdots \\ \vdots \\ CA^{t-1} \end{bmatrix}, \quad T_t = \begin{bmatrix} 0 & & & & & \\ CB & 0 & & & & \\ CAB & CB & \ddots & & & \\ \vdots & & \ddots & \ddots & & \\ \vdots & & & & CB & 0 \\ CA^{t-2}B & \dots & \dots & \dots & CAB & CB & 0 \end{bmatrix}, \quad (3.10)$$

$T_1 = 0$ . Then the terms of the sequence  $\{X_t\}$  generated by the DRE (3.8) are

$$X_t = V_t^T (I + T_t T_t^T)^{-1} V_t, \quad t = 1, 2, \dots \quad (3.11)$$

Moreover,  $\{X_t\}$  is monotonically nondecreasing, and  $X_t \rightarrow X_*$ , the solution of DARE (3.3).

*Proof.* The ‘‘moreover’’ part has been illustrated above. Only (3.11) is proved here.

We have already shown (3.11) is correct for  $t = 1, 2$ . Assuming (3.11) is correct for  $t$ , we are going to prove it is also correct for  $t + 1$ . By the DRE (3.8),

$$\begin{aligned} X_{t+1} &= C^T C + A^T V_t^T (I + T_t T_t^T)^{-1} V_t (I + BB^T V_t^T (I + T_t T_t^T)^{-1} V_t)^{-1} A \\ &\stackrel{(1,2)}{=} C^T C + A^T V_t^T (I + T_t T_t^T)^{-1} (I + V_t BB^T V_t^T (I + T_t T_t^T)^{-1})^{-1} V_t A \\ &= C^T C + A^T V_t^T (I + T_t T_t^T + V_t BB^T V_t^T)^{-1} V_t A \\ &= \begin{bmatrix} C \\ V_t A \end{bmatrix}^T \begin{bmatrix} I \\ I + T_t T_t^T + V_t BB^T V_t^T \end{bmatrix}^{-1} \begin{bmatrix} C \\ V_t A \end{bmatrix} \\ &= \begin{bmatrix} C \\ V_t A \end{bmatrix}^T \left( I + \begin{bmatrix} 0 & \\ V_t B & T_t \end{bmatrix} \begin{bmatrix} 0 & \\ V_t B & T_t \end{bmatrix}^T \right)^{-1} \begin{bmatrix} C \\ V_t A \end{bmatrix} \\ &= V_{t+1}^T (I + T_{t+1} T_{t+1}^T)^{-1} V_{t+1}. \quad \square \end{aligned}$$

It is not difficult to discover that Theorem 3.1 coincides with the decoupled formulae of the dSDA for DAREs introduced in [21], which is actually guaranteed by the fact that the sequence  $\{H_k\}$  generated by SDA (3.9c) is a subsequence of  $\{X_t\}$ .

One can easily find (3.11) is the truncated form of (3.7), which is given by the operator theory. Under the assumption that  $A$  is d-stable, obviously  $V_t \rightarrow \mathcal{V}, T_t \rightarrow \mathcal{T}$ , by the fact that  $\mathcal{V}, \mathcal{T}$  are bounded linear operators, we have  $X_t \rightarrow X_*$ . However, for the case that  $A$  is unstable,  $\mathcal{V}, \mathcal{T}$  are no longer bounded, it would be difficult to show  $X_t \rightarrow X_*$  by the operator theory, which is however guaranteed by the matrix analysis.

### 3.3 Efficient method

Now we acquire the non-iterative form (3.11) of  $X_t$ , which allows us to compute the terms  $X_t$  directly for arbitrary  $t$ . In the following, we will work on an efficient method to compute  $X_t$  for any given  $t$ .

Using the notations for Toeplitz matrices in Section 1, we have

$$T_t = \mathcal{L}_{l \times m} \left( \begin{bmatrix} 0 \\ V_{t-1} B \end{bmatrix} \right) = \begin{bmatrix} 0 & 0 \\ \mathcal{L}_{l \times m}(V_{t-1} B) & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ V_{t-1} B & T_{t-1} \end{bmatrix}.$$

Thus,

$$\mathcal{L}_{l \times m}(V_{t-1} B) \mathcal{L}_{l \times m}(V_{t-1} B)^T = T_{t-1} T_{t-1}^T + V_{t-1} B B^T V_{t-1}^T,$$

and

$$X_t = C^T C + A^T V_{t-1}^T [I + \mathcal{L}_{l \times m}(V_{t-1} B) \mathcal{L}_{l \times m}(V_{t-1} B)^T]^{-1} V_{t-1} A. \quad (3.12)$$

Clearly  $\mathcal{L}_{l \times m}(V_{t-1} B)$  is block-Toeplitz. Hence the results in Section 2 can be applied.

By Lemma 2.1,  $X_t$  can be computed by solving only  $l + m$  rather than  $n$  equations.

**Theorem 3.2.** Let

$$[I + \mathcal{L}_{l \times m}(V_{t-1} B) \mathcal{L}_{l \times m}(V_{t-1} B)^T] \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} = \begin{bmatrix} 0 \\ I_l \end{bmatrix}, \quad Q_{2,b} \in \mathbb{R}^{l \times l}, \quad (3.13a)$$

$$[I + \mathcal{L}_{l \times m}(V_{t-1}B) \mathcal{L}_{l \times m}(V_{t-1}B)^T] \begin{bmatrix} Q_{3,t} \\ Q_{3,c} \end{bmatrix} = V_{t-1}B, \quad Q_{3,t} \in \mathbb{R}^{l \times m}, \quad (3.13b)$$

and  $W = I - \begin{bmatrix} Q_{3,t} \\ Q_{3,c} \end{bmatrix}^T V_{t-1}B$ . Then the sequence  $X_t$  defined by (3.11) can be generated by

$$X_t = \begin{bmatrix} C \\ \Xi_1 \\ \Xi_2 \end{bmatrix}^T \begin{bmatrix} I & & \\ & (I \otimes Q_{2,b})^{-1} & \\ & & (I \otimes W)^{-1} \end{bmatrix} \begin{bmatrix} C \\ \Xi_1 \\ \Xi_2 \end{bmatrix}, \quad (3.14)$$

where

$$\Xi_1 = \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T V_{t-1}A, \quad \Xi_2 = \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} \right)^T V_{t-1}A.$$

*Proof.* By Lemma 2.1 with  $Y = -(Y^L)^T \leftarrow 0$ ,  $D_{t-1} = -(D_{t-1}^L)^T \leftarrow V_{t-1}B$ ,

$$(I + T_t T_t^T)^{-1} = \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right)^T + \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes W)^{-1} \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right)^T,$$

where

$$\begin{aligned} [I + \mathcal{L}_{l \times m}(V_{t-1}B) \mathcal{L}_{l \times m}(V_{t-1}B)^T] Q_3 &= V_{t-1}B, & W &= I - Q_3^T V_{t-1}B, \\ (I + T_t T_t^T) \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} &= \begin{bmatrix} 0 \\ I_l \end{bmatrix}, & Q_{2,b} &\in \mathbb{R}^{l \times l}. \end{aligned}$$

Note that

$$(I + T_t T_t^T)^{-1} = \begin{bmatrix} I & \\ & [I + \mathcal{L}_{l \times m}(V_{t-1}B) \mathcal{L}_{l \times m}(V_{t-1}B)^T]^{-1} \end{bmatrix}.$$

Letting  $Q_2 = \begin{bmatrix} Q_{2,t} \\ Q_{2,c} \end{bmatrix}$  where  $Q_{2,t} \in \mathbb{R}^{l \times l}$ , we have  $Q_{2,t} = 0$  and

$$[I + \mathcal{L}_{l \times m}(V_{t-1}B) \mathcal{L}_{l \times m}(V_{t-1}B)^T] \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} = \begin{bmatrix} 0 \\ I_l \end{bmatrix}.$$

Write  $Q_3 = \begin{bmatrix} Q_{3,t} \\ Q_{3,c} \end{bmatrix}$  where  $Q_{3,t} \in \mathbb{R}^{l \times m}$ , and then it follows that

$$\begin{aligned} & [I + \mathcal{L}_{l \times m}(V_{t-1}B) \mathcal{L}_{l \times m}(V_{t-1}B)^T]^{-1} \\ &= \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T + \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} \right) (I \otimes W)^{-1} \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} \right)^T. \end{aligned} \quad (3.15)$$

Then the result is a direct consequence of (3.12).  $\square$

Theorem 3.2 suggests a new algorithm, Algorithm 1, to approximate the solution of DAREs. The key is how to fast compute  $Q_{*,*}$ , or equivalently solve the linear system (3.13), and compute the products of  $\mathcal{U}_*(*)^T V_{t-1}A$ . Both are related to the computation on block Toeplitz matrices. It is well known that Fast Fourier Transform (FFT) can be used in the calculation with Toeplitz matrices involved, see, e.g., [43, 28, 29] and the references therein.

Some remarks are given below to illustrate the algorithm.

### Parameter and output

1. In order to use FFT,  $t$  is usually chosen as powers of 2, namely  $t = 2^k$ . There is no strategy to determine a proper  $t$  in advance. In practice, we may choose a heuristic  $k$ , for example 5–8. If the output is a good approximation of the solution, then we stop here; otherwise, we use the output as a new initial guess, and run another round to achieve a better approximation; the process is repeated until convergence, namely some criterion is satisfied. The details and the validity of this method are discussed in Section 3.4.
2. Note that  $\text{rank}(S) = \text{rank}(V_{2^k})$ . Numerically  $V_{2^k}$  probably has rank much less than  $2^k l$ . An obvious clue is that  $V_{2^k}$  contains a power series of  $A$  performing on  $C$ , and as  $k$  goes larger and larger, the terms in it become more and more likely to be linearly dependent. This implies that (3.14) is not a compact form. To deal with this, some compression technique may be brought in. This idea needs more discussions on the convergence, which is given in Section 3.4.



---

**Algorithm 1** FFT-based Toeplitz-structured Approximation (FTA) for DARE
 

---

**Input:**  $A, B, C$  and  $t$ .

- 1: Compute sequentially  $C \cdot A, CA \cdot A, \dots, CA^{t-3} \cdot A, CA^{t-2} \cdot A$ , and form  $V_{t-1}$  by stacking  $C$  and the first  $t-2$  terms vertically in order, and form  $V_{t-1}A$  by stacking the  $t-1$  terms vertically in order.
- 2: Compute  $V_{t-1}B$ .
- 3: Use Preconditioned Conjugate Gradient method to solve (3.13).
- 4: Compute  $W = I - \begin{bmatrix} Q_{3,t} \\ Q_{3,c} \end{bmatrix}^T V_{t-1}B$  and then the Cholesky factorizations of  $Q_{2,b} = L_Q L_Q^T$  and  $W = L_W L_W^T$ .
- 5: Use fast multiplication to obtain  $S_1 = \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} L_Q^{-T} \right)^T V_{t-1}A$  and  $S_2 = \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} L_W^{-T} \right)^T V_{t-1}A$ , and form  $S = \begin{bmatrix} C \\ S_1 \\ S_2 \end{bmatrix}$ .

**Output:**  $S$  which satisfies  $S^T S \approx X_*$ .

---

3. In the output, we do not give an approximation of  $X_*$  directly but its factor, namely a  $2^{kl} \times n$  matrix  $S$ . If some compression technique is used during the process, an approximation of  $S$  would have relatively small low row rank, say  $r \ll n$ . Then in practice we only need the products of  $X$  and other matrices, for example, in obtaining the optimal control (3.2). The setting  $r < n$  makes multiplication with  $X_*$ 's factor save time and space. This is also considered in many literatures, e.g., [11].

**Time complexity** Complexity for  $X_{k+1}$ :

1. Step 1, compute  $V_{t-1}A$ , namely  $CA, CA^2, \dots, CA^{t-1}$ , in  $(t-1)(2n-1)nl$  flops.
2. Step 2, compute  $V_{t-1}B$  in  $(t-1)(2n-1)lm$  flops.
3. Step 3, use  $M$ -step PCG ( $Ntl \ln l$  for fast multiplication), to compute  $Q_{*,*}$ , in  $O(MNtl[\ln^2 t + \ln(tl)](l+m))$  flops.
4. Step 4, compute  $W$  in  $(2tl-1)\frac{m(m+1)}{2} + m$  flops, and  $L_Q, L_W, \tilde{D}_Q, \tilde{D}_W$  in  $\frac{2}{3}l(l-1)(l+4)$  flops.
5. Step 5, compute  $\begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} L_Q^{-T}, \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} L_W^{-T}$  in  $tl^3 + (t-1)lm^2$  flops; compute  $S_1, S_2$  in  $O(2Nntl \ln l)$  flops.
6. To sum up, assuming  $l \ll n, m \ll n$  and omitting lower order terms, the total complexity is  $2ln^2t + 2lmnt + O(MNl(l+m)t \ln^2 t + 2Nltn \ln l) = O(t(n^2 + \ln^2 t))$  flops.
7. Suppose  $A$  is sparse, and the number of nonzero entries is  $\text{nnz}(A)$ . Only Step 1 is different, and the total complexity is  $O(lt \text{nnz}(A) + lmnt + MNl(l+m)t \ln^2 t + 2Nltn \ln l) = O(t(\text{nnz}(A) + n + \ln^2 t))$  flops.

**Space complexity**

1. Step 1, store  $V_{t-1}A$  in  $(t-1)nl$  units.
2. Step 2, store  $V_{t-1}B$  in  $(t-1)lm$  units.
3. Step 3, store  $Q_{*,*}$  in  $(t-1)l(l+m)$  units.
4. Step 4, store  $W$  and then  $L_W, L_Q$  in part of the storage for  $V_{t-1}B$ . (The storage is enough because the three matrices need in total  $m(m+1)/2 + l(l+1)/2$  units, which is less than  $(t-1)lm$ .)
5. Step 5, store  $S_2$  in the storage for  $V_{t-1}B$  and additional storage, consuming in total  $(t-1)ln$  units; store  $S_1$  in the storage for  $V_{t-1}A$ .
6. to sum up, the total storage is  $(t-1)l(2n+l+m) = O(tn)$  units.

### 3.4 Arbitrary initial term

In this subsection, we consider (3.8) with an arbitrary initial  $X_0 = \Gamma^T \Gamma \succeq 0$ :

$$X_0 = \Gamma^T \Gamma, \quad X_{t+1} = \mathcal{D}(X_t) = C^T C + A^T X_t (I + B B^T X_t)^{-1} A. \quad (3.8')$$

**Theorem 3.3.** Write

$$\Upsilon_t = [A^{t-1} B \quad \dots \quad A B \quad B].$$

Then the sequence  $\{X_t\}$  generated by (3.8') is given by

$$X_t = \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix}^T \left( I + \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix} \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix}^T \right)^{-1} \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix}, \quad (3.16)$$

where  $V_t, T_t$  is defined by (3.10).

**Remark 3.1.** Note that (3.16) coincides with (3.11) at  $\Gamma = 0$ .

*Proof.* First examine the initial term.

$$\begin{aligned} \begin{bmatrix} C \\ \Gamma A \end{bmatrix}^T \left( I + \begin{bmatrix} 0 \\ \Gamma B \end{bmatrix} \begin{bmatrix} 0 \\ \Gamma B \end{bmatrix}^T \right)^{-1} \begin{bmatrix} C \\ \Gamma A \end{bmatrix} &= C^T C + A^T \Gamma^T (I + \Gamma B B^T \Gamma^T)^{-1} \Gamma A \\ &= C^T C + A^T \Gamma^T \Gamma (I + B B^T \Gamma^T \Gamma)^{-1} A = X_1. \end{aligned}$$

Then examine the recursion.

$$\begin{aligned} &C^T C + A^T X_t (I + B B^T X_t)^{-1} A \\ &= C^T C + A^T \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix}^T \left( I + \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix} \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix}^T \right)^{-1} \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix} \left( I + B B^T \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix}^T \left( I + \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix} \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix}^T \right)^{-1} \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix} \right)^{-1} A \\ &\stackrel{(1.2)}{=} C^T C + A^T \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix}^T \left( I + \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix} \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix}^T + \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix} B B^T \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix}^T \right)^{-1} \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix} A \\ &= \begin{bmatrix} C \\ V_t A \\ \Gamma A^{t+1} \end{bmatrix}^T \begin{bmatrix} I \\ I + \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix} \begin{bmatrix} T_t \\ \Gamma \Upsilon_t \end{bmatrix}^T + \begin{bmatrix} V_t B \\ \Gamma A^t B \end{bmatrix} \begin{bmatrix} V_t B \\ \Gamma A^t B \end{bmatrix}^T \end{bmatrix}^{-1} \begin{bmatrix} C \\ V_t A \\ \Gamma A^{t+1} \end{bmatrix} \\ &= \begin{bmatrix} V_{t+1} \\ \Gamma A^{t+1} \end{bmatrix}^T \left( I + \begin{bmatrix} 0 & 0 \\ V_t B & T_t \end{bmatrix} \begin{bmatrix} 0 & 0 \\ \Gamma A^t B & \Gamma \Upsilon_t \end{bmatrix}^T \right)^{-1} \begin{bmatrix} V_{t+1} \\ \Gamma A^{t+1} \end{bmatrix} \\ &= \begin{bmatrix} V_{t+1} \\ \Gamma A^{t+1} \end{bmatrix}^T \left( I + \begin{bmatrix} T_{t+1} \\ \Gamma \Upsilon_{t+1} \end{bmatrix} \begin{bmatrix} T_{t+1} \\ \Gamma \Upsilon_{t+1} \end{bmatrix}^T \right)^{-1} \begin{bmatrix} V_{t+1} \\ \Gamma A^{t+1} \end{bmatrix} \\ &= X_{t+1}. \quad \square \end{aligned}$$

**Theorem 3.4.** Let  $Q_{2,c}, Q_{2,b}, Q_{3,t}, Q_{3,c}, W, \Xi_1, \Xi_2$  as in Theorem 3.2. Then the sequence  $X_t$  defined by (3.16) can be generated by

$$X_t = \begin{bmatrix} C \\ \Xi_1 \\ \Xi_2 \\ \Xi_\Gamma \end{bmatrix}^T \begin{bmatrix} I & & & \\ & (I \otimes Q_{2,b})^{-1} & & \\ & & (I \otimes W)^{-1} & \\ & & & W_\Gamma^{-1} \end{bmatrix} \begin{bmatrix} C \\ \Xi_1 \\ \Xi_2 \\ \Xi_\Gamma \end{bmatrix},$$

where

$$\begin{aligned} W_\Gamma &= I + \Gamma \Upsilon_t \Upsilon_t^T \Gamma^T - \Xi_{1,\Gamma}^T (I \otimes Q_{2,b})^{-1} \Xi_{1,\Gamma} - \Xi_{2,\Gamma}^T (I \otimes W)^{-1} \Xi_{2,\Gamma}, \\ \Xi_\Gamma &= \Gamma A^t - \Xi_{1,\Gamma}^T (I \otimes Q_{2,b})^{-1} \Xi_1 - \Xi_{2,\Gamma}^T (I \otimes W)^{-1} \Xi_2, \end{aligned}$$

and

$$\Xi_{1,\Gamma} = \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T \mathcal{L}_{l \times m} (V_{t-1} B) \Upsilon_{t-1}^T A^T \Gamma^T, \quad \Xi_{2,\Gamma} = \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} \right)^T \mathcal{L}_{l \times m} (V_{t-1} B) \Upsilon_{t-1}^T A^T \Gamma^T.$$

*Proof.* By (3.16),

$$\begin{aligned}
X_t &= \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix}^T \begin{bmatrix} I + T_t T_t^T & T_t \Upsilon_t^T \Gamma^T \\ \Gamma \Upsilon_t T_t^T & I + \Gamma \Upsilon_t \Upsilon_t^T \Gamma^T \end{bmatrix}^{-1} \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix} \\
&= (*)^T \begin{bmatrix} I + T_t T_t^T & \\ & I + \Gamma \Upsilon_t \Upsilon_t^T \Gamma^T - \Gamma \Upsilon_t T_t^T (I + T_t T_t^T)^{-1} T_t \Upsilon_t^T \Gamma^T \end{bmatrix}^{-1} \begin{bmatrix} I & \\ -\Gamma \Upsilon_t T_t^T (I + T_t T_t^T)^{-1} & I \end{bmatrix} \begin{bmatrix} V_t \\ \Gamma A^t \end{bmatrix} \\
&= \begin{bmatrix} V_t \\ \Xi_\Gamma \end{bmatrix}^T \begin{bmatrix} I + T_t T_t^T & \\ & I + \Gamma \Upsilon_t \Upsilon_t^T \Gamma^T - \Gamma \Upsilon_t T_t^T (I + T_t T_t^T)^{-1} T_t \Upsilon_t^T \Gamma^T \end{bmatrix}^{-1} \begin{bmatrix} V_t \\ \Xi_\Gamma \end{bmatrix} \\
&= V_t^T (I + T_t T_t^T)^{-1} V_t + \Xi_\Gamma^T [I + \Gamma \Upsilon_t \Upsilon_t^T \Gamma^T - \Gamma \Upsilon_t T_t^T (I + T_t T_t^T)^{-1} T_t \Upsilon_t^T \Gamma^T]^{-1} \Xi_\Gamma,
\end{aligned}$$

where  $\Xi_\Gamma = \Gamma A^t - \Gamma \Upsilon_t T_t^T (I + T_t T_t^T)^{-1} V_t$  and  $*$  is used to indicate the same part limited by the symmetry. Similarly to (3.12), by (3.15), writing  $\mathcal{L} = \mathcal{L}_{l \times m}(V_{t-1}B)$ , it can be simplified to

$$\begin{aligned}
X_t &= C^T C + A^T V_{t-1}^T [I + \mathcal{L} \mathcal{L}^T]^{-1} V_{t-1} A + \Xi_\Gamma^T [I + \Gamma \Upsilon_t \Upsilon_t^T \Gamma^T - \Gamma A \Upsilon_{t-1} \mathcal{L}^T (I + \mathcal{L} \mathcal{L}^T)^{-1} \mathcal{L} \Upsilon_{t-1}^T A^T \Gamma^T]^{-1} \Xi_\Gamma \\
&= C^T C + \Xi_1^T (I \otimes Q_{2,b})^{-1} \Xi_1 + \Xi_2^T (I \otimes W)^{-1} \Xi_2 \\
&\quad + \Xi_\Gamma^T [I + \Gamma \Upsilon_t \Upsilon_t^T \Gamma^T - \Xi_{1,\Gamma}^T (I \otimes Q_{2,b})^{-1} \Xi_{1,\Gamma} - \Xi_{2,\Gamma}^T (I \otimes W)^{-1} \Xi_{2,\Gamma}]^{-1} \Xi_\Gamma,
\end{aligned}$$

where

$$\begin{aligned}
\Xi_\Gamma &= \Gamma A^t - \Gamma A \Upsilon_{t-1} \mathcal{L}^T (I + \mathcal{L} \mathcal{L}^T)^{-1} V_{t-1} A \\
&= \Gamma A^t - \Xi_{1,\Gamma}^T (I \otimes Q_{2,b})^{-1} \Xi_1 - \Xi_{2,\Gamma}^T (I \otimes W)^{-1} \Xi_2. \quad \square
\end{aligned}$$

Note that the product of two lower triangular block-Toeplitz matrices is still a lower triangular block-Toeplitz matrix. Hence writing  $\Xi_1 = \begin{bmatrix} \Xi_{1,c} \\ \Xi_{1,b} \end{bmatrix}$  where  $\Xi_{1,b} \in \mathbb{R}^{l \times n}$ ,

$$\begin{aligned}
\mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T \mathcal{L}_{l \times m}(V_{t-1}B) &= \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T \left[ I \otimes (CB) + \mathcal{L}_{l \times m} \left( \begin{bmatrix} 0 \\ V_{t-2} A \end{bmatrix} \right) I \otimes B \right] \\
&= \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T I \otimes (CB) + \mathcal{L}_{l \times n} \left( \begin{bmatrix} 0 \\ \Xi_{1,c} \end{bmatrix} \right) I \otimes B, \\
&= \mathcal{U}_{l \times l} \left( (CB)^T \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T + \mathcal{L}_{l \times m} \left( \begin{bmatrix} 0 \\ \Xi_{1,c} \end{bmatrix} B \right).
\end{aligned}$$

Similarly, writing  $\Xi_2 = \begin{bmatrix} \Xi_{2,c} \\ \Xi_{2,b} \end{bmatrix}$  where  $\Xi_{2,b} \in \mathbb{R}^{l \times n}$ ,

$$\mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} \right)^T \mathcal{L}_{l \times m}(V_{t-1}B) = \mathcal{U}_{l \times l} \left( (CB)^T \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} \right)^T + \mathcal{L}_{l \times m} \left( \begin{bmatrix} 0 \\ \Xi_{2,c} \end{bmatrix} B \right).$$

These can be used to reduce calculations.

What kind of the initial  $\Gamma$  makes the iteration (3.8) converge? Lemma 3.2 gives an easy sufficient condition, which will be immediately used in our algorithm.

**Lemma 3.2.** *The unique positive semi-definite  $d$ -stabilizing solution  $X_\star$  of the DARE (3.3) is an attractor (i.e., asymptotically stable fixed point) of the DRE (3.8) or (3.8'). Moreover, any symmetric matrix  $X$  satisfying either following condition lies in its attraction basin:*

1.  $0 \leq X \leq X_\star$ ;
2.  $\rho(B^T(X - X_\star)B) \leq 1 - \rho(A_{X_\star})$ .

As a result, (3.8') with the matrix above as its initial term converges to  $X_\star$ .

*Proof.* First calculate the differentials.

$$\begin{aligned}
d\mathcal{D}(X) &= A^T dX(I + BB^T X)^{-1} A + A^T X d((I + BB^T X)^{-1}) A \\
&= A^T dX(I + BB^T X)^{-1} A - A^T X (I + BB^T X)^{-1} BB^T dX(I + BB^T X)^{-1} A \\
&= A^T [I - X(I + BB^T X)^{-1} BB^T] dX(I + BB^T X)^{-1} A
\end{aligned}$$

$$\begin{aligned}
&= A^T(I + XBB^T)^{-1} dX(I + BB^T X)^{-1} A \\
&= A_X^T dX A_X,
\end{aligned}$$

where  $A_X$  is the closed loop matrix. In order to avoid the appearance of 4th-order tensor, we use vectorization to obtain

$$d(\text{vec } \mathcal{D}(x)) = A_X^T \otimes A_X^T d(\text{vec } X) \quad \text{and} \quad \frac{d(\text{vec } \mathcal{D}(x))}{d(\text{vec } X)} = A_X^T \otimes A_X^T.$$

Since  $X_*$  is d-stabilizing,  $\rho(A_{X_*}) < 1$  and thus the Fréchet derivative at  $X_*$  has norm less than 1, which guarantees  $X_*$  is an attractor.

For the first kind of matrices, by Lemma 3.1,  $X_* = \mathcal{D}^t(X_*) \succeq \mathcal{D}^t(X) \succeq \mathcal{D}^t(0) \rightarrow X_*$ , which forces  $\mathcal{D}^t(X) \rightarrow X_*$ . For the second one, the only thing to be needed is the Fréchet derivative at  $X$  has norm less than 1. Writing  $X - X_* = \Delta$  and using an norm induced by some vector norm,

$$\begin{aligned}
\|(I + BB^T X)^{-1} A\| &= \|[I + (I + BB^T X_*)^{-1} BB^T \Delta]^{-1} (I + BB^T X_*)^{-1} A\| \\
&\leq \|[I + (I + BB^T X_*)^{-1} BB^T \Delta]^{-1}\| \|(I + BB^T X_*)^{-1} A\| \\
&\leq \frac{\|I\|}{1 - \|(I + BB^T X_*)^{-1} BB^T \Delta\|} \|(I + BB^T X_*)^{-1} A\| \\
&\leq \frac{\|A_{X_*}\|}{1 - \|(I + BB^T X_*)^{-1}\| \|BB^T \Delta\|} \\
&\leq \frac{\|A_{X_*}\|}{1 - \|BB^T \Delta\|},
\end{aligned}$$

which is less than 1, as long as  $\rho(B^T \Delta B) = \rho(BB^T \Delta) \leq 1 - \rho(A_{X_*})$ .  $\square$

According to Lemma 3.2, the compression technique can be used in Algorithm 1 without breaking its convergence. We roughly describe the process here: after performing Algorithm 1 for a small/mid  $t$ , a truncation technique (e.g., SVD/QR) is used on  $\Xi$  to produce  $S'_X = \Gamma$ ; then (3.16) is used to generate a new approximation; repeat this process until convergence. To decrease the number of calculations, the same  $t$  is used in each outer iteration.

## 4 CARE

Given a linear time-invariant control system in continuous-time:

$$\begin{aligned}
\dot{x}(t) &= Ax(t) + Bu(t), \\
y(t) &= Cx(t),
\end{aligned}$$

where  $A \in \mathbb{R}^{n \times n}$ ,  $B \in \mathbb{R}^{n \times m}$ ,  $C \in \mathbb{R}^{l \times n}$ . Suppose the following condition holds through out this section:

$$\boxed{(A, B) \text{ is c-stabilizable and } (C, A) \text{ is c-detectable,}}$$

or equivalently,  $\text{rank}([A - \lambda I \quad BB^T]) = \text{rank}([A^T - \lambda I \quad C^T C]) = n$  for any  $\lambda \in \mathbb{C} \setminus \mathbb{C}_-$ , where  $\mathbb{C}_-$  is the open left half complex plane.

Its linear-quadratic optimal control can be expressed as

$$\arg \min_{u(t)} \int_0^\infty [y(t)^T y(t) + u(t)^T u(t)] dt = -B^T X_* x(t),$$

where  $X_*$  is the unique symmetric positive semi-definite c-stabilizing solution  $X$  of the CARE [12, 14, 34, 39]:

$$\mathcal{E}(X) := A^T X + X A - X B B^T X + C^T C = 0. \quad (4.1)$$

Here a solution  $X$  is called c-stabilizing, if the closed loop matrix  $A_X = A - BB^T X$  is c-stable, namely all of its eigenvalues lie in the open left half complex plane  $\mathbb{C}_-$ .

### 4.1 FTA and its equivalence to many other methods

Many numerical methods to solve CAREs are based on performing Cayley transformation on the associated Hamiltonian matrix

$$\mathcal{H} = \begin{bmatrix} A & BB^T \\ C^T C & -A^T \end{bmatrix}.$$



*Proof.* The ‘‘moreover’’ part is the same as that for DAREs in Section 3 and hence omitted. Only (4.4) is proved here.

It is easy to verify that (4.4) is correct for  $t = 1$ . Assuming (4.4) is correct for  $t$ , we are going to prove it is also correct for  $t + 1$ . By the DRE (4.3),

$$\begin{aligned}
X_{t+1} &= \tilde{C}^T(I + Y_\gamma Y_\gamma^T)^{-1} \tilde{C} + (\tilde{A} - \tilde{B} Y_\gamma^T (I + Y_\gamma Y_\gamma^T)^{-1} \tilde{C})^T \tilde{V}_t^T \left( I + \tilde{T}_t \tilde{T}_t^T \right)^{-1} \tilde{V}_t \\
&\quad \cdot \left( I + \tilde{B} (I + Y_\gamma^T Y_\gamma)^{-1} \tilde{B}^T \tilde{V}_t^T \left( I + \tilde{T}_t \tilde{T}_t^T \right)^{-1} \tilde{V}_t \right)^{-1} (\tilde{A} - \tilde{B} Y_\gamma^T (I + Y_\gamma Y_\gamma^T)^{-1} \tilde{C}) \\
&\stackrel{(1.2)}{=} \tilde{C}^T (I + Y_\gamma Y_\gamma^T)^{-1} \tilde{C} + (\tilde{A} - \tilde{B} Y_\gamma^T (I + Y_\gamma Y_\gamma^T)^{-1} \tilde{C})^T \tilde{V}_t^T \\
&\quad \cdot \left( I + \tilde{T}_t \tilde{T}_t^T + \tilde{V}_t \tilde{B} (I + Y_\gamma^T Y_\gamma)^{-1} \tilde{B}^T \tilde{V}_t^T \right)^{-1} \tilde{V}_t (\tilde{A} - \tilde{B} Y_\gamma^T (I + Y_\gamma Y_\gamma^T)^{-1} \tilde{C}) \\
&= (*)^T \begin{bmatrix} I + Y_\gamma Y_\gamma^T & \\ & I + \tilde{T}_t \tilde{T}_t^T + \tilde{V}_t \tilde{B} (I + Y_\gamma^T Y_\gamma)^{-1} \tilde{B}^T \tilde{V}_t^T \end{bmatrix}^{-1} \begin{bmatrix} \tilde{C} \\ \tilde{V}_t \tilde{A} - \tilde{V}_t \tilde{B} Y_\gamma^T (I + Y_\gamma Y_\gamma^T)^{-1} \tilde{C} \end{bmatrix} \\
&= (*)^T \begin{bmatrix} I + Y_\gamma Y_\gamma^T & \\ & I + \tilde{T}_t \tilde{T}_t^T + \tilde{V}_t \tilde{B} \tilde{B}^T \tilde{V}_t^T - \tilde{V}_t \tilde{B} Y_\gamma^T (I + Y_\gamma Y_\gamma^T)^{-1} Y_\gamma \tilde{B}^T \tilde{V}_t^T \end{bmatrix}^{-1} \\
&\quad \cdot \begin{bmatrix} I & \\ -\tilde{V}_t \tilde{B} Y_\gamma^T (I + Y_\gamma Y_\gamma^T)^{-1} & I \end{bmatrix} \begin{bmatrix} \tilde{C} \\ \tilde{V}_t \tilde{A} \end{bmatrix} \\
&= \begin{bmatrix} \tilde{C} \\ \tilde{V}_t \tilde{A} \end{bmatrix}^T \begin{bmatrix} I + Y_\gamma Y_\gamma^T & & & \\ & Y_\gamma \tilde{B}^T \tilde{V}_t^T & & \\ & & I + \tilde{T}_t \tilde{T}_t^T + \tilde{V}_t \tilde{B} \tilde{B}^T \tilde{V}_t^T & \\ & & & \end{bmatrix}^{-1} \begin{bmatrix} \tilde{C} \\ \tilde{V}_t \tilde{A} \end{bmatrix} \\
&= \begin{bmatrix} \tilde{C} \\ \tilde{V}_t \tilde{A} \end{bmatrix}^T \left( I + \begin{bmatrix} Y_\gamma & 0 \\ \tilde{V}_t \tilde{B} & \tilde{T}_t \end{bmatrix} \begin{bmatrix} Y_\gamma & 0 \\ \tilde{V}_t \tilde{B} & \tilde{T}_t \end{bmatrix}^T \right)^{-1} \begin{bmatrix} \tilde{C} \\ \tilde{V}_t \tilde{A} \end{bmatrix} \\
&= \tilde{V}_{t+1}^T (I + \tilde{T}_{t+1} \tilde{T}_{t+1}^T)^{-1} \tilde{V}_{t+1}.
\end{aligned}$$

Here \* is still used to indicate the same part limited by the symmetry. □

Using the notations for Toeplitz matrices in Section 2, we have

$$\tilde{T}_t = \mathcal{L}_{l \times m} \left( \begin{bmatrix} Y_\gamma \\ \tilde{V}_{t-1} \tilde{B} \end{bmatrix} \right) = I_t \otimes Y_\gamma + \begin{bmatrix} 0 & 0 \\ \mathcal{L}_{l \times m}(\tilde{V}_{t-1} \tilde{B}) & 0 \end{bmatrix} = \begin{bmatrix} Y_\gamma & 0 \\ \tilde{V}_{t-1} \tilde{B} & \tilde{T}_{t-1} \end{bmatrix}.$$

**Theorem 4.2.** *Let*

$$\left[ I + \tilde{T}_t \tilde{T}_t^T \right] \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} = \begin{bmatrix} 0 \\ I_l \end{bmatrix}, \quad Q_{2,b} \in \mathbb{R}^{l \times l}, \quad (4.5a)$$

$$\left[ I + \tilde{V}_{t-1} \tilde{B} \tilde{B}^T \tilde{V}_{t-1}^T + \tilde{T}_{t-1} \tilde{T}_{t-1}^T \right] Q_3 = \tilde{V}_{t-1} \tilde{B}, \quad (4.5b)$$

and  $W = I - Q_3^T \tilde{V}_{t-1} \tilde{B}$ . Then the sequence  $X_t$  defined by DRE (4.4) can be generated by

$$X_t = \begin{bmatrix} \Xi_1 \\ \Xi_2 \end{bmatrix}^T \begin{bmatrix} (I \otimes Q_{2,b})^{-1} & \\ & (I \otimes [W + W Y_\gamma^T Y_\gamma W])^{-1} \end{bmatrix} \begin{bmatrix} \Xi_1 \\ \Xi_2 \end{bmatrix}, \quad (4.6)$$

where

$$\Xi_1 = \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right)^T \tilde{V}_t, \quad \Xi_2 = \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right)^T \tilde{V}_t.$$

*Proof.* By Lemma 2.1 with  $Y = -(Y^L)^T \leftarrow Y_\gamma$ ,  $D_{t-1} = -(D_{t-1}^L)^T \leftarrow \tilde{V}_{t-1} \tilde{B}$ ,

$$\begin{aligned}
&(I + \tilde{T}_t \tilde{T}_t^T)^{-1} \\
&= \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right)^T + \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes [W + W Y_\gamma^T Y_\gamma W])^{-1} \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right)^T. \quad (4.7)
\end{aligned}$$

Then the result follows (4.4). □

Theorem 4.2 suggests a similar algorithm to approximate the solution of CARE. Some remarks are given below to illustrate the algorithm.

---

**Algorithm 2** FFT-based Toeplitz-structured Approximation (FTA) for CARE

---

**Input:**  $A, B, C$  and  $\gamma, t$ .

- 1: Compute  $\hat{A}_\gamma = A - \gamma I$  and generate the linear-system solver  $\hat{A}_\gamma^{-1}$  for the sparse  $A$  (or its PLU factorization  $\hat{A}_\gamma = PLU$  for the dense  $A$ ).
- 2: Compute  $(\mathbf{tmp}) = U^{-1}L^{-1}P^{-1}B, \tilde{C} = \sqrt{2\gamma}C\hat{A}_\gamma^{-1}$  by the linear solver or forward/backward substitution, and compute  $\tilde{B} = \sqrt{2\gamma}(\mathbf{tmp}), Y_\gamma = C(\mathbf{tmp})$ .
- 3: Compute sequentially  $\tilde{C} \cdot \tilde{A}, \tilde{C}\tilde{A} \cdot \tilde{A}, \dots, \tilde{C}\tilde{A}^{t-3} \cdot \tilde{A}, \tilde{C}\tilde{A}^{t-2} \cdot \tilde{A}$  by the way  $(\mathbf{tmp})\tilde{A} = (\mathbf{tmp}) + 2\gamma(\mathbf{tmp})\hat{A}_\gamma^{-1}$  by the linear solver or forward/backward substitution, and form  $\tilde{V}_t$  by stacking  $\tilde{C}$  and the  $t-1$  terms vertically in order, where the first  $t-1$  terms consists of  $\tilde{V}_{t-1}$ .
- 4: Compute  $\tilde{V}_{t-1}\tilde{B}$  and form  $\begin{bmatrix} Y_\gamma \\ \tilde{V}_{t-1}\tilde{B} \end{bmatrix}$ .
- 5: Use Preconditioned Conjugate Gradient method to solve (4.5).
- 6: Compute  $W = I - Q_3^T \tilde{V}_{t-1} \tilde{B}, (\mathbf{tmp}) = Y_\gamma W, \tilde{W} = W + (\mathbf{tmp})^T (\mathbf{tmp})$  and then the Cholesky factorizations of  $Q_{2,b} = L_Q L_Q^T$  and  $\tilde{W} = L_W L_W^T$ .
- 7: Use fast multiplication to obtain  $S_1 = \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} L_Q^{-T} \right)^T \tilde{V}_t$  and  $S_2 = \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} L_W^{-T} \right)^T \tilde{V}_t$ , and form  $S = \begin{bmatrix} S_1 \\ S_2 \end{bmatrix}$ .

**Output:**  $S$  which satisfies  $S^T S \approx X_*$ .

---

### Parameter and output

1. considerations similar to Algorithm 1 have to be made. This implies that (4.6) is not a compact form either, and some truncation/reduction/shrinking technique may be brought in.

**Time complexity** Complexity for  $X_{k+1}$ :

1. Step 1, compute  $\hat{A}_\gamma$  and its PLU factorization in  $n + \frac{n(n-1)(4n+1)}{6}$  flops.
2. Step 2, compute  $\tilde{C}, \tilde{B}, Y_\gamma$  in  $2n^2m + 2n^2l + nm + nl + lm(2n-1)$  flops.
3. Step 3, compute  $\tilde{V}_t$ , namely  $\tilde{C}\tilde{A}, \tilde{C}\tilde{A}^2, \dots, \tilde{C}\tilde{A}^{t-1}$ , in  $(t-1)[2n^2l + nl + nl]$  flops.
4. Step 4, compute  $\tilde{V}_{t-1}\tilde{B}$  in  $(t-1)(2n-1)lm$  flops.
5. Step 5, Use  $M$ -step PCG ( $Ntl \ln l$  for fast multiplication), to compute  $Q_{*,*}$ , in  $O(MNtl[\ln^2 t + \ln(t)](l+m))$  flops.
6. Step 6, compute  $W, \tilde{W}$  in  $(2tl-1)\frac{m(m+1)}{2} + m + (2m-1)lm + (2l-1)\frac{m(m+1)}{2}$  flops, and  $L_Q, L_W$  in  $\frac{2}{3}l(l-1)(l+4)$  flops.
7. Step 7, compute  $\begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} L_Q^{-T}, \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} L_W^{-T}$  in  $tl^3 + (t-1)lm^2$  flops; compute  $S_1, S_2$  in  $O(2Nntl \ln l)$  flops.
8. To sum up, assuming  $l \ll n, m \ll n$  and omitting lower order terms, the total complexity is  $\frac{2}{3}n^3 + 2(l+m)n^2 + 2lmn + 2ln^2t + 2lmnt + O(MNl(l+m)t \ln^2 t + 2Nltn \ln l) = \frac{2}{3}n^3 + O(t(n^2 + \ln^2 t))$  flops.
9. Suppose  $A$  is sparse, and the number of nonzero entries is  $\text{nnz}(A)$ . The PLU factorization can be replaced by an iterative solver with at most  $P$  steps, such as GMRES. The total complexity is  $O(P(m+l) \text{nnz}(A) + lmn + lt \text{nnz}(A) + lmnt + MNl(l+m)t \ln^2 t + 2Nltn \ln l) = O(t(\text{nnz}(A) + n + \ln^2 t))$  flops.

### Space complexity

1. the storage is similar to that of Algorithm 1.

## 4.2 Incorporation technique

In the following, we consider the incorporation technique (aka defect correction). The key idea is: once an approximate solution  $\tilde{X}$  is obtained, letting the difference from the exact solution  $X_*$  be  $\Delta$ , namely  $X_* = \tilde{X} + \Delta$ , the difference satisfies  $A^T(\tilde{X} + \Delta) + (\tilde{X} + \Delta)A + C^T C - (\tilde{X} + \Delta)BB^T(\tilde{X} + \Delta) = 0$ , from which an approximation  $\tilde{\Delta}$  can be generated and then  $\tilde{X} + \tilde{\Delta}$  should be an approximate solution to the original equation better than  $\tilde{X}$ . More details can be found in [24, 5]. The following lemma is important as the guarantee of the validity of the incorporation technique.

**Lemma 4.1** ([5, Theorem 1]). *Let  $\tilde{X}$  be an approximation to a solution to (4.1).*

1.  $\Delta = X_* - \tilde{X}$  is a solution to the equation

$$(A - BB^T\tilde{X})^T\Delta + \Delta(A - BB^T\tilde{X}) + \mathcal{C}(\tilde{X}) - \Delta BB^T\Delta = 0. \quad (4.8)$$

2. Conversely, if  $\Delta$  is a solution to (4.8), then  $\tilde{X} + \Delta$  is a solution to (4.1). Moreover, if  $\tilde{X} \succeq 0$  and  $\Delta$  is a stabilizing solution to (4.8), then  $\tilde{X} + \Delta$  is the stabilizing solution to (4.1).

3. If  $\tilde{X} \succeq 0, \mathcal{C}(\tilde{X}) \succeq 0$ , then  $\Delta$  is the unique stabilizing solution to (4.8).

4. If  $\tilde{X} \succeq 0, \mathcal{C}(\tilde{X}) \succeq 0$ , then  $\Delta \preceq X_*$ .

To make the technique useful for Algorithm 2, the fundamental problem we face is the rank factorization of  $\mathcal{C}(\tilde{X})$ . Let  $\tilde{X} = X_t$  that we obtained.

**Theorem 4.3.** *Let  $\mathbf{1}_t \in \mathbb{R}^t$  be a vector with each entry one. Then for  $X_t$  defined by DRE (4.3),  $\mathcal{C}(X_t) = C_t^T C_t$ , where*

$$C_0 = C, \quad C_t = C + \sqrt{2\gamma}(\mathbf{1}_t^T \otimes I_l)(I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t. \quad (4.9)$$

*Proof.* Clearly,

$$\begin{aligned} \mathcal{C}(X_t) &= A^T X_t + X_t A + C^T C - X_t B B^T X_t \\ &= [\gamma I + 2\gamma(\tilde{A} - I)^{-T}] X_t + X_t [\gamma I + 2\gamma(\tilde{A} - I)^{-1}] + 2\gamma(\tilde{A} - I)^{-T} \tilde{C}^T \tilde{C} (\tilde{A} - I)^{-1} - 2\gamma X_t (\tilde{A} - I)^{-1} \tilde{B} \tilde{B}^T (\tilde{A} - I)^{-T} X_t \\ &= 2\gamma \left[ (\tilde{A} - I)^{-T} \tilde{C}^T \tilde{C} (\tilde{A} - I)^{-1} + (\tilde{A} - I)^{-T} X_t + X_t (\tilde{A} - I)^{-1} + X_t - X_t (\tilde{A} - I)^{-1} \tilde{B} \tilde{B}^T (\tilde{A} - I)^{-T} X_t \right]. \end{aligned}$$

Note that by (4.4)

$$\begin{aligned} X_t (\tilde{A} - I)^{-1} \tilde{B} &= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t (\tilde{A} - I)^{-1} \tilde{B} \\ &= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \begin{bmatrix} \tilde{C} \\ \tilde{C} \tilde{A} \\ \vdots \\ \tilde{C} \tilde{A}^{t-1} \end{bmatrix} (\tilde{A} - I)^{-1} \tilde{B} \\ &= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \begin{bmatrix} Y_\gamma \\ \tilde{C} \tilde{B} + Y_\gamma \\ \vdots \\ \tilde{C} \tilde{A}^{t-2} \tilde{B} + \dots + \tilde{C} \tilde{B} + Y_\gamma \end{bmatrix} \\ &= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{T}_t \begin{bmatrix} I_m \\ I_m \\ \vdots \\ I_m \end{bmatrix} = \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{T}_t (\mathbf{1}_t \otimes I_m), \end{aligned}$$

and

$$X_t (\tilde{A} - I)^{-1} = \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \begin{bmatrix} \tilde{C} \\ \tilde{C} \tilde{A} \\ \vdots \\ \tilde{C} \tilde{A}^{t-1} \end{bmatrix} (\tilde{A} - I)^{-1}$$



$$\begin{aligned}
&= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \begin{bmatrix} \tilde{C}(\tilde{A} - I)^{-1} \\ \tilde{C} + \tilde{C}(\tilde{A} - I)^{-1} \\ \vdots \\ \tilde{C}\tilde{A}^{t-2} + \dots + \tilde{C} + \tilde{C}(\tilde{A} - I)^{-1} \end{bmatrix} \\
&= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \left( (\mathbf{1}_t \otimes I_l) \tilde{C}(\tilde{A} - I)^{-1} + \mathcal{L}_{l \times n} \left( \begin{bmatrix} 0 \\ \tilde{V}_{t-1} \end{bmatrix} \right) (\mathbf{1}_t \otimes I_l) \right) \\
&= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \left( (\mathbf{1}_t \otimes I_l) \tilde{C}(\tilde{A} - I)^{-1} + \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right) \tilde{V}_t \right),
\end{aligned}$$

of which the last equality is guaranteed by

$$\mathcal{L}_{l \times n} \left( \begin{bmatrix} 0 \\ \tilde{V}_{t-1} \end{bmatrix} \right) (\mathbf{1}_t \otimes I_l) = \begin{bmatrix} 0 \\ \tilde{C} \\ \tilde{C}\tilde{A} + \tilde{C} \\ \vdots \\ \tilde{C}\tilde{A}^{t-2} + \dots + \tilde{C} \end{bmatrix} = \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right) \tilde{V}_t.$$

Hence

$$\begin{aligned}
\frac{1}{2\gamma} (\mathcal{E}(X_t) - C_t^T C_t) &= \frac{(\tilde{A} - I)^{-T} \tilde{C}^T \tilde{C} (\tilde{A} - I)^{-1}}{2\gamma} \\
&\quad + \frac{(\tilde{A} - I)^{-T} \tilde{C}^T (\mathbf{1}_t^T \otimes I_l) (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t + \tilde{V}_t^T \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right)^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t}{2\gamma} \\
&\quad + \frac{\tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} (\mathbf{1}_t \otimes I_l) \tilde{C}(\tilde{A} - I)^{-1} + \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right) \tilde{V}_t}{2\gamma} \\
&\quad + \frac{\tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t - \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{T}_t (\mathbf{1}_t \otimes I_m) (\mathbf{1}_t^T \otimes I_m) \tilde{T}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t}{2\gamma} \\
&\quad - \frac{(\tilde{A} - I)^{-T} \tilde{C}^T \tilde{C} (\tilde{A} - I)^{-1} - \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} (\mathbf{1}_t \otimes I_l) (\mathbf{1}_t^T \otimes I_l) (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t}{2\gamma} \\
&\quad - \frac{(\tilde{A} - I)^{-T} \tilde{C}^T (\mathbf{1}_t^T \otimes I_l) (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t - \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} (\mathbf{1}_t \otimes I_l) \tilde{C}(\tilde{A} - I)^{-1}}{2\gamma} \\
&= \tilde{V}_t^T \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right)^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t + \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right) \tilde{V}_t \\
&\quad + \frac{\tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t - \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{T}_t (\mathbf{1}_t \otimes I_m) (\mathbf{1}_t^T \otimes I_m) \tilde{T}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t}{2\gamma} \\
&\quad - \frac{\tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} (\mathbf{1}_t \otimes I_l) (\mathbf{1}_t^T \otimes I_l) (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t}{2\gamma} \\
&= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \left[ (I + \tilde{T}_t \tilde{T}_t^T) \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right)^T + \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right) (I + \tilde{T}_t \tilde{T}_t^T) \right. \\
&\quad \left. + (I + \tilde{T}_t \tilde{T}_t^T) - \tilde{T}_t (\mathbf{1}_t \otimes I_m) (\mathbf{1}_t^T \otimes I_m) \tilde{T}_t^T - (\mathbf{1}_t \otimes I_l) (\mathbf{1}_t^T \otimes I_l) \right] (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t \\
&= \tilde{V}_t^T (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \left[ \tilde{T}_t \tilde{T}_t^T \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right)^T + \mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right) \tilde{T}_t \tilde{T}_t^T \right. \\
&\quad \left. + \tilde{T}_t \tilde{T}_t^T - \tilde{T}_t (\mathbf{1}_t \otimes I_m) (\mathbf{1}_t^T \otimes I_m) \tilde{T}_t^T \right] (I + \tilde{T}_t \tilde{T}_t^T)^{-1} \tilde{V}_t \\
&= 0,
\end{aligned}$$

of which the last equality holds for

$$\mathcal{L}_{l \times l} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_l \end{bmatrix} \right) \tilde{T}_t = \tilde{T}_t \mathcal{L}_{m \times m} \left( \begin{bmatrix} 0 \\ \mathbf{1}_{t-1} \otimes I_m \end{bmatrix} \right). \quad \square$$

According to Theorem 4.3, we are able to make incorporation easily and solve (4.8). One thing worth mentioning is that  $C_t$  is not difficult to calculate. Putting (4.7) into (4.9),

$$C_t = C + \sqrt{2\gamma} \begin{bmatrix} \Xi_{1,I} \\ \Xi_{2,I} \end{bmatrix}^T \begin{bmatrix} (I \otimes Q_{2,b})^{-1} & \\ & (I \otimes [W + WY_\gamma^T Y_\gamma W])^{-1} \end{bmatrix} \begin{bmatrix} \Xi_1 \\ \Xi_2 \end{bmatrix},$$

where  $\Xi_1, \Xi_2$  is defined as in (4.6), and

$$\Xi_{1,I} = \mathcal{U}_{l \times l} \left( \begin{bmatrix} Q_{2,c} \\ Q_{2,b} \end{bmatrix} \right)^T (\mathbf{1}_t \otimes I_l), \quad \Xi_{2,I} = \mathcal{U}_{l \times m} \left( \begin{bmatrix} Q_{3,c} \\ 0 \end{bmatrix} \right)^T (\mathbf{1}_t \otimes I_l).$$

Theorem 4.3 gives the detailed form of (4.8) at  $\tilde{X} = X_t$ . Note that the sequence  $\{X_t\}$  can be generated by the RADI method introduced in [5], if the initial approximation is 0 and the same shifts are used, which is actually make incorporation at each  $t = 1$  iteratively. As a direct consequence, we have the following result.

**Theorem 4.4.** For  $X_t$  defined by DRE (4.3) and  $C_t$  defined by (4.9),

$$\Delta_t := X_{t+1} - X_t = 2\gamma(C_t A_{t,\gamma}^{-1})^T (I + C_t A_{t,\gamma}^{-1} B B^T A_{t,\gamma}^{-T} C_t^T)^{-1} C_t A_{t,\gamma}^{-1}, \quad A_{t,\gamma} = A - B B^T X_t - \gamma I,$$

or equivalently,  $\Delta_t$  is the approximate solution to (4.8) at  $\tilde{X} = X_t$  generated by DRE (4.3) on  $t = 1$ .

## 5 Experiments and discussions

In this section, we will provide several examples to illustrate the new algorithm FTA and compare it with some existing methods. As is stated in Section 4.1, many methods solve a CARE through an equivalent DARE. Hence here we only test the CARE of the form

$$A^T X + X A - X B B^T X + C^T C = 0, \quad A \in \mathbb{R}^{n \times n}, B \in \mathbb{R}^{n \times m}, C \in \mathbb{R}^{l \times n},$$

for the performance of the methods on CARE can be recognized as those on DARE. We will use these methods in the tests:

- FTA: our FFT-based Toeplitz-structured approximation with the incorporation technique;
- RKSM: rational Krylov subspace method [22, 18, 42];
- RADI+opt: RADI method [5], with the residual minimizing shifts;
- RADI+proj: RADI method with the residual Hamiltonian shifts;
- NK-ADI+GP: the Galerkin projected variant of Newton-Kleinman ADI method [6, 7, 10, 9];
- iNK-ADI+LS: the inexact variant of Newton-Kleinman ADI method with line search.

All experiments are done in MATLAB 2021a under the Windows 10 Professional 64-bit operating system on a PC with a Intel Core i7-8700 processor at 3.20GHz and 64GB RAM. The implementation of RKSM comes from the source codes from Simoncini's homepage<sup>1</sup> with some modifications; we use corresponding functions in the package M-M.E.S.S. version 2.1 [41] as the implementations of the last four methods.

The methods are intentionally chosen: RADI is the recommended method by the package M-M.E.S.S., and it is usually one of the fastest methods among the non-projective methods, and two different shift selection strategies are used, for there does not exist a definitely good one and these two are good in many tests; the two variants of NK-ADI are Newton-type methods; RKSM is a projection method. On the other hand, FTA, SDA, RADI, quadratic ADI and Cayley transformed Hamiltonian subspace iteration are theoretically equivalent if the shifts are the same; the last three are of the same type, so only one of them, namely RADI, is chosen; SDA is appropriate for small-to-mid scale dense problems, so we give up putting it into comparison.

Since classical performance indices behave very different in different methods, we directly use the accuracy vs. the running time to compare. The accuracy is measured by

$$\text{NRes}(X) := \frac{\|\mathcal{C}(X)\|_F}{\|\mathcal{C}(0)\|_F} = \frac{\|A^T X + X A - X B B^T X + C^T C\|_F}{\|C^T C\|_F}.$$

In the following, three examples are tested, where the results are shown in Figure 5.1.

**Example 5.1 (Rail).** The example is a version of the steel profile cooling model from the Oberwolfach Model Reduction Benchmark Collection, hosted at MORwiki [40]. The data include  $A \preceq 0, E \succeq 0, B, C$  with  $n = 79841, m = 7, l = 6$ . Since we only focus on solving the CARE,  $E$  is simply dropped.

For the parameters, in FTA, we use a heuristic shift  $\gamma = 8 \times 10^{-7}$  and in each incorporation step  $\gamma \leftarrow \gamma/1.01$ . Each of the other five methods has its own way to choose shifts, so we leave the task for their own. The situations in the following examples are the same.

<sup>1</sup><http://www.dm.unibo.it/~simoncin/software.html>




Figure 5.1: accuracy vs. time

In this example  $A \prec 0$  and thus c-stable, which implies the properties of this problem are good. This results in the fact that all methods converge. We can see that the FTA is the slowest one in all the six methods. This phenomenon is reasonable. FTA and RADI are theoretically equivalent, while the only difference is that RADI has much more chances to choose different shifts to accelerate its convergence. Good shifts largely accelerate its convergence, and on the opposite, bad shifts would slow it down. RKSM and NK-ADI also benefit from the choice of shifts.

**Example 5.2** (Lung2+). The example is generated in this way:  $A$  is the matrix `lung2` in the SuiteSparse Matrix Collection [16] (formerly the University of Florida Sparse Matrix Collection), modelling temperature and water vapor transport in the human lung;  $B, C$  are generated by MATLAB function `rand`. Here  $n = 109460, m = 10, l = 10$ .

For the parameters, in FTA, we use a heuristic shift  $\gamma = 5 \times 10^3$  and in each incorporation step  $\gamma \leftarrow \gamma/1.01$ .

In this example  $A$  is nonsymmetric and the eigenvalues of  $A$  lie in the right half plane, namely  $A$  is c-anti-stable, or  $-A$  is c-stable. None of RADI+opt, RADI+proj, and RKSM converges. NK-ADI+GP and iNK-ADI+LS both report that non-stable Ritz values were detected and terminated the process. Only FTA produces a good approximate solution. This tells that the other methods strongly rely on the stability of  $A$ . For example, a sufficient condition for achieving the convergence is that  $A$  is stable and the shifts  $\gamma_k$  satisfy the non-Blaschke condition  $\sum_{k=1}^{+\infty} \frac{\Re(\gamma_k)}{1+|\gamma_k|^2} = -\infty$ . However, the FTA works well even for the case that  $A$  is not stable, which implies that in this sense the FTA is more robust with respect to the spectrum of  $A$ .

**Example 5.3** (Lung2-). The example is almost the same with Example 5.2 except that the matrix `lung2` is used as  $-A$  rather than  $A$ .

For the parameters, in FTA, we use a heuristic shift  $\gamma = 5 \times 10^3$  and in each incorporation step  $\gamma \leftarrow \gamma/1.01$ .

In this example  $A$  is nonsymmetric and c-stable. Note that the only difference of RADI+opt and RADI+proj is the different shift selection strategies. RADI+opt tends to converge fast but finally stays at a low accuracy; RADI+proj becomes convergent very late but soon converges in a very short time. The phenomenon illustrates that the choice of shifts fatally affects its speed of convergence. The iNK-ADI+LS converges very slowly, while the NK-ADI+GP reports that non-stable Ritz values were detected again. The RKSM converges in a fairly good speed but finally slow down. The FTA converges steadily in a predictable speed.

On the other hand, compared with Example 5.2, the running time of the FTA is nearly the same for different  $A$ 's, so the running time is predictable and can be estimated in advance.

Summarizing the numerical results, we see that the FTA has two significant features:

1. FTA is robust in some sense and it converges no matter how the property of  $A$  is;
2. FTA has a steady convergence rate and the execution time is predictable, although in good cases it may converge slowly compared with other methods.

Moreover, it is easy to see that if  $A$  is dense, FTA need the LU/PLU factorization only several times, while the other methods need as many as number of iterations, according to the number of used shifts.

## 6 Conclusion

We have presented our FFT-based Toeplitz-structured approximation method for computing the stabilizing solution of large-scale algebraic Riccati equations with low-rank structure. It is shown that the closed form given by operator theory under good assumptions is also valid for the general case, which is proved by matrix analysis. It is quite natural to ask whether the closed form can be directly produced by the analysis of unbounded linear operators, which would be a difficult task for future work. On the numerical front, our method works robust in some sense and few parameters are needed. As the readers may see, there is still possibility to improve the behavior by adopting more techniques. However, to keep this paper compact and concentrated, we leave it for another work.

## A Displacement rank and Toeplitz matrix

In order to continue our discussion, we first give a few but urgent tips to the displacement rank and Toeplitz matrices, and interested readers are referred to the review paper [33] and the references therein.

For any matrix  $R \in \mathbb{R}^{pn \times pn}$ , its ( $\pm$ )-displacement rank  $\alpha_{\pm}(R, p)$  with respect to block size  $p \times p$ , is defined by

$$\alpha_+(R, p) := \text{rank}_p(R - Z_{n,p} R Z_{n,p}^T), \quad \alpha_-(R, p) := \text{rank}_p(R - Z_{n,p}^T R Z_{n,p}),$$

where  $Z_{n,p} = \begin{bmatrix} 0 & 0 \\ I_{(n-1)p} & 0 \end{bmatrix}_{pn \times pn}$ , and  $\text{rank}_p(\cdot)$  is considered as the rank of the linear transformation on the module  $\mathbb{R}^{np \times p}$  over the ring  $\mathbb{R}^{p \times p}$ . For the case  $p = 1$ ,  $\text{rank}_p(\cdot) = \text{rank}(\cdot)$ , the ordinary rank of matrices in  $\mathbb{R}^{n \times n}$ .

The definition is based on the following result, namely Lemma A.1.

**Lemma A.1** ([32]). Given  $R_1, R_2 \in \mathbb{R}^{pn \times p}$  and  $R \in \mathbb{R}^{pn \times pn}$ , then

$$\begin{aligned} R - Z_{n,p} R Z_{n,p}^T &= R_1 R_2^T \iff R = \mathcal{L}_{p \times p}(R_1) \mathcal{L}_{p \times p}(R_2)^T, \\ R - Z_{n,p}^T R Z_{n,p} &= R_1 R_2^T \iff R = \mathcal{U}_{p \times p}(R_1) \mathcal{U}_{p \times p}(R_2)^T. \end{aligned}$$

Lemma A.1 implies that for a matrix its displacement rank is related to how it can be expressed as a sum of products of block-Toeplitz matrices, as is shown in Lemma A.2.

**Lemma A.2** ([32, 31, 33]). Given a matrix  $R \in \mathbb{R}^{pn \times pn}$ .

1. Its (+)-displacement rank  $\alpha_+(R, p)$  is the smallest integer  $\beta$  such that  $R$  can be written in the form

$$R = \sum_{i=1}^{\beta} \mathcal{L}_{p \times p}(R_i) \mathcal{U}_{p \times p}(\tilde{R}_i), \quad (\text{A.2a})$$

where  $R_i, \tilde{R}_i \in \mathbb{R}^{pn \times p}$ .

2. Its (-)-displacement rank  $\alpha_-(R, p)$  is the smallest integer  $\beta$  such that  $R$  can be written in the form

$$R = \sum_{i=1}^{\beta} \mathcal{U}_{p \times p}(R_i) \mathcal{L}_{p \times p}(\tilde{R}_i), \quad (\text{A.2b})$$

where  $R_i, \tilde{R}_i \in \mathbb{R}^{pn \times p}$ .

3. If  $R$  is symmetric and positive semidefinite, (A.2a) and (A.2b) can be replaced respectively by

$$R = \sum_{i=1}^{\beta} \mathcal{L}_{p \times p}(R_i) \mathcal{L}_{p \times p}(R_i)^T, \quad \text{and} \quad R = \sum_{i=1}^{\beta} \mathcal{U}_{p \times p}(R_i) \mathcal{U}_{p \times p}(R_i)^T.$$

4. If  $R$  is nonsingular, then  $\alpha_+(R, p) = \alpha_-(R^{-1}, p)$ ,  $\alpha_-(R, p) = \alpha_+(R^{-1}, p)$ .

Lemma A.2 tells the relation between the displacement ranks of a matrix and its inverse, which is the theoretical foundation of the fast and superfast algorithms on Toeplitz matrices.

The following result, namely Lemma A.3, gives an expression of the inverse related to the displacement rank.

**Lemma A.3** ([20]). Given  $R \in \mathbb{R}^{pn \times pn}$ , suppose

1.  $R$  is nonsingular, and  $R^{-1} = \begin{bmatrix} Q_{1,t} & Q_1^L \\ Q_1 & * \end{bmatrix} = \begin{bmatrix} * & Q_2 \\ Q_2^L & Q_{2,b} \end{bmatrix}$  where  $Q_{1,t}, Q_{2,b} \in \mathbb{R}^{p \times p}$  are nonsingular;
2.  $R - Z_{n,p} R Z_{n,p}^T = \begin{bmatrix} * & * \\ * & D_1 \Sigma D_2^T \end{bmatrix}$  where  $D_1, D_2 \in \mathbb{R}^{p(n-1) \times p\alpha}$  and  $\Sigma$  is a diagonal matrix whose diagonal entries are  $\pm 1$ ;
3. writing  $R = \begin{bmatrix} * & * \\ * & R_s \end{bmatrix}$  where  $R_s \in \mathbb{R}^{p(n-1) \times p(n-1)}$ , there exists  $Q_3, Q_3^L$  such that  $R_s Q_3 = D_1, Q_3^L R_s = D_2^T$ .

Then

$$\begin{aligned} R^{-1} &= -\mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_1 \\ 0 \end{bmatrix} \right) (I \otimes Q_{1,t})^{-1} \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_1^L & 0 \end{bmatrix}^T \right)^T + \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2^L & Q_{2,b} \end{bmatrix}^T \right)^T \\ &\quad + \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes W)^{-1} \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3^L & 0 \end{bmatrix}^T \right)^T, \end{aligned} \quad (\text{A.3a})$$

or alternatively,

$$\begin{aligned} R^{-1} &= \mathcal{L}_{p \times p} \left( \begin{bmatrix} Q_{1,t} \\ Q_1 \end{bmatrix} \right) (I \otimes Q_{1,t})^{-1} \mathcal{L}_{p \times p} \left( \begin{bmatrix} Q_{1,t} & Q_1^L \end{bmatrix}^T \right)^T - \mathcal{L}_{p \times p} \left( \begin{bmatrix} 0 \\ Q_2 \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{L}_{p \times p} \left( \begin{bmatrix} 0 & Q_2^L \end{bmatrix}^T \right)^T \\ &\quad - \mathcal{L}_{p \times p\alpha} \left( \begin{bmatrix} 0 \\ Q_3 \end{bmatrix} \right) (I \otimes W)^{-1} \mathcal{L}_{p \times p\alpha} \left( \begin{bmatrix} 0 & Q_3^L \end{bmatrix}^T \right)^T, \end{aligned} \quad (\text{A.3b})$$

where  $W = \Sigma - Q_3^L D_1$ .

Moreover, if  $R$  is symmetric, then there exists a factorization to make  $D_1 = D_2$ ; for that case, (A.3) can be rewritten by  $Q_1^L = Q_1^T, Q_2^L = Q_2^T, Q_3^L = Q_3^T$ .

**Remark A.1.** Item 2 of Lemma A.3 implies that  $R - Z_{n,p}^T R Z_{n,p} = \begin{bmatrix} -D_1 \Sigma D_2^T & * \\ * & * \end{bmatrix}$ .

Note that (A.3) presents a sum of  $\alpha + 2$  products of block-Toeplitz matrices, in which the number of terms may not be the smallest one, namely  $\alpha_{\mp}(R, p)$ .

In the following, we will derive a sum of the  $\alpha_+(R, p) = \alpha_-(R^{-1}, p)$  terms, to coincide with Lemma A.2, of which the form is called a *shortest sum*. Using the same way a sum of  $\alpha_-(R, p) = \alpha_+(R^{-1}, p)$  terms can also be derived, so we omit the details.

Write  $R = \begin{bmatrix} R_{11} & R_{12} \\ R_{21} & R_s \end{bmatrix}$ , and then  $R - Z_{n,p} R Z_{n,p}^T = \begin{bmatrix} R_{11} & R_{12} \\ R_{21} & D_1 \Sigma D_2^T \end{bmatrix}$ . Thus,  $\alpha \leq \alpha_+(R, p) \leq \alpha + 2$ .

On the other hand, by (1.3), under sufficient nonsingular conditions, it is easy to have

$$\begin{aligned} \begin{bmatrix} R_{11} & R_{12} \\ R_{21} & R_s \end{bmatrix}^{-1} &= \begin{bmatrix} R_{11}^{-1} + R_{11}^{-1} R_{12} (R_s - R_{21} R_{11}^{-1} R_{12})^{-1} R_{21} R_{11}^{-1} & -R_{11}^{-1} R_{12} (R_s - R_{21} R_{11}^{-1} R_{12})^{-1} \\ -(R_s - R_{21} R_{11}^{-1} R_{12})^{-1} R_{21} R_{11}^{-1} & (R_s - R_{21} R_{11}^{-1} R_{12})^{-1} \end{bmatrix} \\ &= \begin{bmatrix} (R_{11} - R_{12} R_s^{-1} R_{21})^{-1} & -(R_{11} - R_{12} R_s^{-1} R_{21})^{-1} R_{12} R_s^{-1} \\ -R_s^{-1} R_{21} (R_{11} - R_{12} R_s^{-1} R_{21})^{-1} & R_s^{-1} + R_s^{-1} R_{21} (R_{11} - R_{12} R_s^{-1} R_{21})^{-1} R_{12} R_s^{-1} \end{bmatrix}. \end{aligned}$$

Compared with the conditions,

$$Q_1 = -R_s^{-1} R_{21} Q_{1,t}, \quad Q_1^L = -Q_{1,t} R_{12} R_s^{-1}, \quad Q_{1,t} = (R_{11} - R_{12} R_s^{-1} R_{21})^{-1}.$$

If  $\alpha_+(R, p) = \alpha$ , then it has to hold that  $R - Z_{n,p} R Z_{n,p}^T = \begin{bmatrix} S_1^T \Sigma^{-1} S_2 & S_1^T D_2^T \\ D_1 S_2 & D_1 \Sigma D_2^T \end{bmatrix}$  for some  $S_1, S_2 \in \mathbb{R}^{p\alpha \times p}$ .

Clearly  $S_1, S_2$  are of full column rank for  $R$  is nonsingular. Noticing  $\Sigma^{-1} = \Sigma$ , we have

$$\begin{aligned} Q_1 &= -R_s^{-1} D_1 S_2 Q_{1,t} = -Q_3 S_2 Q_{1,t}, \\ Q_1^L &= -Q_{1,t} S_1^T D_2^T R_s^{-1} = -Q_{1,t} S_1^T Q_3^L, \\ Q_{1,t} &= (S_1^T \Sigma^{-1} S_2 - S_1^T D_2^T R_s^{-1} D_1 S_2)^{-1} = (S_1^T \Sigma^{-1} S_2 - S_1^T Q_3^L D_1 S_2)^{-1} = (S_1^T W S_2)^{-1}. \end{aligned}$$

Thus

$$\begin{aligned} & \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_1 \\ 0 \end{bmatrix} \right) (I \otimes Q_{1,t})^{-1} \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_1^L & 0 \end{bmatrix} \right)^T \\ &= \mathcal{U}_{p \times p} \left( \begin{bmatrix} -Q_3 S_2 Q_{1,t} \\ 0 \end{bmatrix} \right) (I \otimes Q_{1,t})^{-1} \mathcal{U}_{p \times p} \left( \begin{bmatrix} -Q_{1,t} S_1^T Q_3^L & 0 \end{bmatrix} \right)^T \\ &= \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t}) (I \otimes Q_{1,t})^{-1} (I \otimes Q_{1,t} S_1^T) \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3^L & 0 \end{bmatrix} \right)^T \\ &= \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t} S_1^T) \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3^L & 0 \end{bmatrix} \right)^T \\ &= \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 (S_1^T W S_2)^{-1} S_1^T) \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3^L & 0 \end{bmatrix} \right)^T. \end{aligned}$$

Note that

$$\left[ W^{-1} - S_2 (S_1^T W S_2)^{-1} S_1^T \right] W S_2 = 0.$$

Complement  $S_2$  to a nonsingular matrix  $\begin{bmatrix} S_2 & S_2^c \end{bmatrix}$ , and then

$$\begin{aligned} \left[ W^{-1} - S_2 (S_1^T W S_2)^{-1} S_1^T \right] W \begin{bmatrix} S_2 & S_2^c \end{bmatrix} &= \begin{bmatrix} S_2 & S_2^c \end{bmatrix} \begin{bmatrix} 0 & -(S_1^T W S_2)^{-1} S_1^T W S_2^c \\ 0 & I \end{bmatrix} \\ &= \begin{bmatrix} S_2 & S_2^c \end{bmatrix} \begin{bmatrix} -(S_1^T W S_2)^{-1} S_1^T W S_2^c \\ I \end{bmatrix} \begin{bmatrix} 0 & I \end{bmatrix}, \end{aligned}$$

whose rank is  $p(\alpha - 1)$ . Write

$$W_1 = \begin{bmatrix} S_2 & S_2^c \end{bmatrix} \begin{bmatrix} -(S_1^T W S_2)^{-1} S_1^T W S_2^c \\ I_{p(\alpha-1)} \end{bmatrix} \in \mathbb{R}^{p\alpha \times p(\alpha-1)}, \quad W_1^L = \begin{bmatrix} 0 & I_{p(\alpha-1)} \end{bmatrix} \begin{bmatrix} S_2 & S_2^c \end{bmatrix}^{-1} W^{-1} \in \mathbb{R}^{p(\alpha-1) \times p\alpha},$$

and then  $W^{-1} - S_2 (S_1^T W S_2)^{-1} S_1^T = W_1 W_1^L$ . Hence

$$\begin{aligned} R^{-1} &= \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p \times p} \left( [Q_2^L \quad Q_{2,b}]^T \right)^T + \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes W_1 W_1^L) \mathcal{U}_{p \times p\alpha} \left( [Q_3^L \quad 0]^T \right)^T \\ &= \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p \times p} \left( [Q_2^L \quad Q_{2,b}]^T \right)^T + \mathcal{U}_{p \times p(\alpha-1)} \left( \begin{bmatrix} Q_3 W_1 \\ 0 \end{bmatrix} \right) \mathcal{U}_{p \times p(\alpha-1)} \left( [W_1^L Q_3^L \quad 0]^T \right)^T. \end{aligned} \quad (\text{A.4})$$

If  $\alpha_+(R, p) = \alpha + 1$ , then it holds that

$$R - Z_{n,p} R Z_{n,p}^T = \begin{bmatrix} S_1^T \Sigma^{-1} S_2 & S_1^T D_2^T \\ D_1 S_2 & D_1 \Sigma D_2^T \end{bmatrix} + \begin{bmatrix} S_3 & D_3^T \\ 0 & 0 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} S_1^T \Sigma^{-1} S_2 & S_1^T D_2^T \\ D_1 S_2 & D_1 \Sigma D_2^T \end{bmatrix} + \begin{bmatrix} S_3 & 0 \\ D_3 & 0 \end{bmatrix}$$

for some  $S_1, S_2 \in \mathbb{R}^{p\alpha \times p}$ ,  $S_3 \in \mathbb{R}^{p \times p}$  and  $D_3 \in \mathbb{R}^{p(n-1) \times p}$ .

Consider the former one. Then,

$$\begin{aligned} Q_1 &= -R_s^{-1} D_1 S_2 Q_{1,t} = -Q_3 S_2 Q_{1,t}, \\ Q_1^L &= -Q_{1,t} (S_1^T D_2^T + D_3^T) R_s^{-1} = -Q_{1,t} S_1^T Q_3^L - Q_{1,t} D_3^T R_s^{-1}, \\ Q_{1,t} &= (S_1^T \Sigma^{-1} S_2 + S_3 - (S_1^T D_2^T + D_3^T) R_s^{-1} D_1 S_2)^{-1} \\ &= (S_1^T \Sigma^{-1} S_2 + S_3 - D_3^T Q_3 S_2 - S_1^T Q_3^L D_1 S_2)^{-1} \\ &= (S_3 - D_3^T Q_3 S_2 + S_1^T W S_2)^{-1}. \end{aligned}$$

Thus,

$$\begin{aligned} &\mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_1 \\ 0 \end{bmatrix} \right) (I \otimes Q_{1,t})^{-1} \mathcal{U}_{p \times p} \left( [Q_1^L \quad 0]^T \right)^T \\ &= \mathcal{U}_{p \times p} \left( \begin{bmatrix} -Q_3 S_2 Q_{1,t} \\ 0 \end{bmatrix} \right) (I \otimes Q_{1,t})^{-1} \mathcal{U}_{p \times p} \left( [-Q_{1,t} S_1^T Q_3^L - Q_{1,t} D_3^T R_s^{-1} \quad 0]^T \right)^T \\ &= \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t}) (I \otimes Q_{1,t})^{-1} (I \otimes Q_{1,t} S_1^T) \mathcal{U}_{p \times p\alpha} \left( [Q_3^L \quad 0]^T \right)^T \\ &\quad + \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t}) (I \otimes Q_{1,t})^{-1} (I \otimes Q_{1,t}) \mathcal{U}_{p \times p} \left( [D_3^T R_s^{-1} \quad 0]^T \right)^T \\ &= \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t} S_1^T) \mathcal{U}_{p \times p\alpha} \left( [Q_3^L \quad 0]^T \right)^T \\ &\quad + \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t}) \mathcal{U}_{p \times p} \left( [D_3^T R_s^{-1} \quad 0]^T \right)^T \\ &= \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 (S_3 - D_3^T Q_3 S_2 + S_1^T W S_2)^{-1} S_1^T) \mathcal{U}_{p \times p\alpha} \left( [Q_3^L \quad 0]^T \right)^T \\ &\quad + \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t}) \mathcal{U}_{p \times p} \left( [D_3^T R_s^{-1} \quad 0]^T \right)^T. \end{aligned}$$

Since

$$\begin{aligned} W^{-1} - S_2 (S_3 - D_3^T Q_3 S_2 + S_1^T W S_2)^{-1} S_1^T &= W^{-1} (W - W S_2 (S_3 - D_3^T Q_3 S_2 + S_1^T W S_2)^{-1} S_1^T W) W^{-1} \\ &\stackrel{(1.3)}{=} W^{-1} (W^{-1} + S_2 (S_3 - D_3^T Q_3 S_2)^{-1} S_1^T)^{-1} W^{-1} \\ &= (W + W S_2 (S_3 - D_3^T Q_3 S_2)^{-1} S_1^T W)^{-1} =: W_1^{-1}, \end{aligned}$$

we have

$$\begin{aligned} R^{-1} &= \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p \times p} \left( [Q_2^L \quad Q_{2,b}]^T \right)^T + \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes W_1)^{-1} \mathcal{U}_{p \times p\alpha} \left( [Q_3^L \quad 0]^T \right)^T \\ &\quad - \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes S_2 Q_{1,t}) \mathcal{U}_{p \times p} \left( [D_3^T R_s^{-1} \quad 0]^T \right)^T \\ &= \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p \times p} \left( [Q_2^L \quad Q_{2,b}]^T \right)^T \\ &\quad + \mathcal{U}_{p \times p\alpha} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes W_1)^{-1} \mathcal{U}_{p \times p\alpha} \left( [Q_3^L - W_1 S_2 Q_{1,t} D_3^T R_s^{-1} \quad 0]^T \right)^T. \end{aligned} \quad (\text{A.5})$$

Similarly, for the latter one,

$$R^{-1} = \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p \times p} \left( [Q_2^L \quad Q_{2,b}]^T \right)^T \quad (\text{A.6})$$

$$+ \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_3 - R_s^{-1} D_3 Q_{1,t} S_1^T W_1 \\ 0 \end{bmatrix} \right) (I \otimes W_1)^{-1} \mathcal{U}_{p \times p} \left( [Q_3^L \quad 0]^T \right)^T,$$

where  $W_1 = W + W S_2 (S_3 - S_1^T Q_3^L D_3)^{-1} S_1^T W$ .

To sum up, we have Lemma A.4.

**Lemma A.4.** *Given  $R \in \mathbb{R}^{p_n \times p_n}$ , suppose the conditions in Lemma A.3 hold. Then  $\alpha \leq \alpha_+(R, p) \leq \alpha + 2$ , and the following statements hold.*

1. if  $\alpha_+(R, p) = \alpha$ , then (A.4) is a shortest sum.
2. if  $\alpha_+(R, p) = \alpha + 1$ , then (A.5) or (A.6) is a shortest sum.
3. if  $\alpha_+(R, p) = \alpha + 2$ , then (A.3a) is a shortest sum.

Moreover, if  $R$  is symmetric, then there exists a factorization to make  $D_1 = D_2$ ; for that case, (A.4) to (A.6) and (A.3a) can be rewritten by  $Q_1^L = Q_1^T, Q_2^L = Q_2^T, Q_3^L = Q_3^T, S_1 = S_2, D_3 = 0$ .

Finally, we prove Lemma 2.1.

*Proof of Lemma 2.1.* First consider the case  $p_1 = p_2 = p$ . Since

$$I - T_t T_t^L = \begin{bmatrix} I - Y Y^L & -Y D_{t-1}^L \\ -D_{t-1} Y^L & I - D_{t-1} D_{t-1}^L - T_{t-1} T_{t-1}^L \end{bmatrix},$$

and

$$\begin{aligned} (I - T_t T_t^L) - Z_{t,p} (I - T_t T_t^L) Z_{t,p}^T &= I - Z_{t,p} Z_{t,p}^T - T_t T_t^L + Z_{t,p} T_t T_t^L Z_{t,p}^T \\ &= \begin{bmatrix} I & \\ & 0 \end{bmatrix} - \begin{bmatrix} Y & 0 \\ D_{t-1} & T_{t-1} \end{bmatrix} \begin{bmatrix} Y^L & D_{t-1}^L \\ 0 & T_{t-1}^L \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ T_{t-1} & 0 \end{bmatrix} \begin{bmatrix} 0 & T_{t-1}^L \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} I - Y Y^L & -Y D_{t-1}^L \\ -D_{t-1} Y^L & -D_{t-1} D_{t-1}^L \end{bmatrix}, \end{aligned}$$

we have  $\alpha_+(I - T_t T_t^L, p) = 2$ . By Lemma A.4, since  $\alpha = 1$ , the case falls in Item 2 with substitutions

$$D_3 \leftarrow 0, S_3 \leftarrow I, D_1 \leftarrow D_{t-1}, \Sigma \leftarrow -I, D_2^T \leftarrow D_{t-1}^L, S_1^T \leftarrow Y, S_2 \leftarrow Y^L.$$

Then (A.5) (or equivalently (A.6)) becomes

$$\begin{aligned} (I - T_t T_t^L)^{-1} &= \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p \times p} \left( [Q_2^L \quad Q_{2,b}]^T \right)^T \\ &\quad + \mathcal{U}_{p \times p} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes [W + W Y^L Y W])^{-1} \mathcal{U}_{p \times p} \left( [Q_3^L \quad 0]^T \right)^T, \end{aligned}$$

where  $Q_2, Q_{2,b}, Q_2^L, Q_3, Q_3^L, W$  is as in (2.1).

Then consider the case  $p_1 > p_2$ . Complement  $Y$  to a  $p_1 \times p_1$  matrix  $\tilde{Y} = [Y \quad 0]$  and similarly for  $\tilde{D}_{t-1} = [D_{t-1} \quad 0], \tilde{Y}^L = \begin{bmatrix} Y^L \\ 0 \end{bmatrix}, \tilde{D}_{t-1}^L = \begin{bmatrix} D_{t-1}^L \\ 0 \end{bmatrix}$ . Immediately we are able to use the result above on the case  $p_1 = p_2$  to

obtain  $\left[ I - \mathcal{L}_{p_1 \times p_1} \left( \begin{bmatrix} \tilde{Y} \\ \tilde{D}_{t-1} \end{bmatrix} \right) \mathcal{L}_{p_1 \times p_1} \left( \begin{bmatrix} \tilde{Y}^L & \tilde{D}_{t-1}^L \end{bmatrix}^T \right)^T \right]^{-1}$ . Note that

$$\mathcal{L}_{p_1 \times p_1} \left( \begin{bmatrix} \tilde{Y} \\ \tilde{D}_{t-1} \end{bmatrix} \right) \mathcal{L}_{p_1 \times p_1} \left( \begin{bmatrix} \tilde{Y}^L & \tilde{D}_{t-1}^L \end{bmatrix}^T \right)^T = \begin{bmatrix} * & 0 & * & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \dots \\ * & 0 & * & 0 & \dots \end{bmatrix} \begin{bmatrix} * & \dots & * \\ 0 & \dots & 0 \\ * & \dots & * \\ 0 & \dots & 0 \\ \vdots & & \vdots \end{bmatrix}$$



$$\begin{aligned}
&= \begin{bmatrix} * & * & \cdots \\ \vdots & \vdots & \\ * & * & \cdots \end{bmatrix} \begin{bmatrix} * & \cdots & * \\ * & \cdots & * \\ \vdots & & \vdots \end{bmatrix} \\
&= \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} Y \\ D_{t-1} \end{bmatrix} \right) \mathcal{L}_{p_1 \times p_2} \left( [Y^L \ D_{t-1}^L]^T \right)^T.
\end{aligned}$$

Thus,  $\tilde{Q}_2 = Q_2, \tilde{Q}_2^L = Q_2^L, \tilde{Q}_{2,b} = Q_{2,b}$ , and  $\tilde{Q}_3 = [Q_3 \ 0], \tilde{Q}_3^L = \begin{bmatrix} Q_3^L \\ 0 \end{bmatrix}$ . Therefore,

$$\begin{aligned}
\tilde{W} &= -I - \begin{bmatrix} Q_3^L \\ 0 \end{bmatrix} [D_{t-1} \ 0] = \begin{bmatrix} -I - Q_3^L D_{t-1} & 0 \\ 0 & -I \end{bmatrix} = \begin{bmatrix} W & \\ & -I \end{bmatrix}, \\
\tilde{W} \tilde{Y}^L \tilde{Y} \tilde{W} &= \begin{bmatrix} W & \\ & -I \end{bmatrix} \begin{bmatrix} Y^L \\ 0 \end{bmatrix} [Y \ 0] \begin{bmatrix} W & \\ & -I \end{bmatrix} = \begin{bmatrix} W Y^L Y W & \\ & 0 \end{bmatrix}.
\end{aligned}$$

Hence

$$\begin{aligned}
&\mathcal{U}_{p_1 \times p_1} \left( \begin{bmatrix} \tilde{Q}_3 \\ 0 \end{bmatrix} \right) (I \otimes [\tilde{W} + \tilde{W} \tilde{Y}^L \tilde{Y} \tilde{W}])^{-1} \mathcal{U}_{p_1 \times p_1} \left( \begin{bmatrix} \tilde{Q}_3^L & 0 \end{bmatrix}^T \right)^T \\
&= \begin{bmatrix} * & 0 & * & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \\ * & 0 & * & 0 & \cdots \end{bmatrix} \begin{bmatrix} * & & & & \\ & -I & & & \\ & & * & & \\ & & & -I & \\ & & & & \ddots \end{bmatrix} \begin{bmatrix} * & \cdots & * \\ 0 & \cdots & 0 \\ * & \cdots & * \\ 0 & \cdots & 0 \\ \vdots & & \vdots \end{bmatrix} \\
&= \begin{bmatrix} * & * & \cdots \\ \vdots & \vdots & \\ * & * & \cdots \end{bmatrix} \begin{bmatrix} * & & \\ & * & \\ & & \ddots \end{bmatrix} \begin{bmatrix} * & \cdots & * \\ * & \cdots & * \\ \vdots & & \vdots \end{bmatrix} \\
&= \mathcal{U}_{p_1 \times p_2} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes [W + W Y^L Y W])^{-1} \mathcal{U}_{p_1 \times p_2} \left( [Q_3^L \ 0]^T \right)^T.
\end{aligned}$$

Finally consider the case  $p_1 < p_2$ . Complement  $Y$  to a  $p_2 \times p_2$  matrix  $\tilde{Y} = \begin{bmatrix} Y \\ 0 \end{bmatrix}$  and similarly for  $\tilde{D}_{t-1}^T = \begin{bmatrix} * & 0 & * & 0 & \cdots \end{bmatrix}$  where  $D_{t-1}^T = \begin{bmatrix} * & * & \cdots \end{bmatrix}$ , and  $\tilde{Y}^L = [Y^L \ 0], \tilde{D}_{t-1}^L = \begin{bmatrix} * & 0 & * & 0 & \cdots \end{bmatrix}$  where  $D_{t-1}^L = \begin{bmatrix} * & * & \cdots \end{bmatrix}$ . To make things clear, two permutations  $P, P_s$  are used to make  $P \begin{bmatrix} \tilde{Y} \\ \tilde{D}_{t-1} \end{bmatrix} = \begin{bmatrix} Y \\ D_{t-1} \\ 0 \end{bmatrix}, P_s \tilde{D}_{t-1} = \begin{bmatrix} D_{t-1} \\ 0 \end{bmatrix}$ . So  $[\tilde{Y}^L \ \tilde{D}_{t-1}^L] P^T = [Y^L \ D_{t-1}^L \ 0], \tilde{D}_{t-1}^L P_s^T = [D_{t-1}^L \ 0]$ , and

$$P \mathcal{L}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Y} \\ \tilde{D}_{t-1} \end{bmatrix} \right) = \left[ \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} Y \\ D_{t-1} \\ 0 \end{bmatrix} \right) \right], \mathcal{L}_{p_2 \times p_2} \left( [\tilde{Y}^L \ \tilde{D}_{t-1}^L]^T \right)^T P^T = \left[ \mathcal{L}_{p_1 \times p_2} \left( [Y^L \ D_{t-1}^L]^T \right)^T \ 0 \right].$$

Then we use the result above on the case  $p_1 = p_2$  to obtain  $\left[ I - \mathcal{L}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Y} \\ \tilde{D}_{t-1} \end{bmatrix} \right) \mathcal{L}_{p_2 \times p_2} \left( [\tilde{Y}^L \ \tilde{D}_{t-1}^L]^T \right)^T \right]^{-1}$ .

Note that

$$P \left[ I - \mathcal{L}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Y} \\ \tilde{D}_{t-1} \end{bmatrix} \right) \mathcal{L}_{p_2 \times p_2} \left( [\tilde{Y}^L \ \tilde{D}_{t-1}^L]^T \right)^T \right] P^T = \left[ I - \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} Y \\ D_{t-1} \\ 0 \end{bmatrix} \right) \mathcal{L}_{p_1 \times p_2} \left( [Y^L \ D_{t-1}^L]^T \right)^T \right] I.$$

Thus,

$$\begin{aligned}
P \begin{bmatrix} \tilde{Q}_2 \\ \tilde{Q}_{2,b} \end{bmatrix} &= P \left[ I - \mathcal{L}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Y} \\ \tilde{D}_{t-1} \end{bmatrix} \right) \mathcal{L}_{p_2 \times p_2} \left( [\tilde{Y}^L \ \tilde{D}_{t-1}^L]^T \right)^T \right]^{-1} P^T P \begin{bmatrix} 0 \\ I_{p_2} \end{bmatrix} \\
&= \left[ \begin{array}{c} I - \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} Y \\ D_{t-1} \\ 0 \end{bmatrix} \right) \mathcal{L}_{p_1 \times p_2} \left( [Y^L \ D_{t-1}^L]^T \right)^T \\ I \end{array} \right]^{-1} \begin{bmatrix} 0 \\ [I_{p_1} \ 0] \\ 0 \\ [0 \ I_{p_2-p_1}] \end{bmatrix}
\end{aligned}$$

$$= \begin{bmatrix} \left[ I - \mathcal{L}_{p_1 \times p_2} \left( \begin{bmatrix} Y \\ D_{t-1}^L \end{bmatrix} \right) \mathcal{L}_{p_1 \times p_2} \left( [Y^L \quad D_{t-1}^L]^T \right)^T \right]^{-1} \begin{bmatrix} 0 \\ I_{p_1} \end{bmatrix} & 0 \\ 0 & \begin{bmatrix} 0 \\ I_{p_2-p_1} \end{bmatrix} \end{bmatrix} = \begin{bmatrix} \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} & 0 \\ 0 & \begin{bmatrix} 0 \\ I_{p_2-p_1} \end{bmatrix} \end{bmatrix},$$

and similarly,  $\begin{bmatrix} \tilde{Q}_2^L & \tilde{Q}_{2,b} \end{bmatrix} P^T = \begin{bmatrix} [Q_2^L \quad Q_{2,b}] & 0 \\ 0 & I_{p_2-p_1} \end{bmatrix}$ . Therefore,  $\tilde{Q}_{2,b} = \begin{bmatrix} Q_{2,b} & 0 \\ 0 & I_{p_2-p_1} \end{bmatrix}$  and

$$\begin{aligned} & P \mathcal{U}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Q}_2 \\ \tilde{Q}_{2,b} \end{bmatrix} \right) P^T P (I \otimes \tilde{Q}_{2,b})^{-1} P^T P \mathcal{U}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Q}_2^L & \tilde{Q}_{2,b} \end{bmatrix}^T \right)^T P^T \\ &= \begin{bmatrix} \mathcal{U}_{p_1 \times p_1} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} (I \otimes Q_{2,b})^{-1} & \\ & I \end{bmatrix} \begin{bmatrix} \mathcal{U}_{p_1 \times p_1} \left( [Q_2^L \quad Q_{2,b}]^T \right) & 0 \\ 0 & I \end{bmatrix}^T \\ &= \begin{bmatrix} \mathcal{U}_{p_1 \times p_1} \left( \begin{bmatrix} Q_2 \\ Q_{2,b} \end{bmatrix} \right) (I \otimes Q_{2,b})^{-1} \mathcal{U}_{p_1 \times p_1} \left( [Q_2^L \quad Q_{2,b}]^T \right)^T & \\ & I \end{bmatrix}. \end{aligned}$$

Similarly,

$$P_s \tilde{Q}_3 = P_s \left[ I - \tilde{D}_{t-1} \tilde{D}_{t-1}^L - \tilde{T}_{t-1} \tilde{T}_{t-1}^L \right]^{-1} P_s^T P_s \tilde{D}_{t-1} = \begin{bmatrix} [I - D_{t-1} D_{t-1}^L - T_{t-1} T_{t-1}^L]^{-1} & \\ & I \end{bmatrix} \begin{bmatrix} D_{t-1} \\ 0 \end{bmatrix} = \begin{bmatrix} Q_3 \\ 0 \end{bmatrix},$$

and similarly,  $\tilde{Q}_3^L P_s^T = \begin{bmatrix} Q_3^L & 0 \end{bmatrix}$ , and then

$$\begin{aligned} \tilde{W} &= -I - \tilde{Q}_3^L P_s^T P_s \tilde{D}_{t-1} = -I - Q_3^L D_{t-1} = W, \\ \tilde{W} \tilde{Y}^L \tilde{Y} \tilde{W} &= W \begin{bmatrix} Y^L & 0 \\ & 0 \end{bmatrix} W = W Y^L Y W. \end{aligned}$$

Hence

$$\begin{aligned} & P \mathcal{U}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Q}_3 \\ 0 \end{bmatrix} \right) (I \otimes [\tilde{W} + \tilde{W} \tilde{Y}^L \tilde{Y} \tilde{W}])^{-1} \mathcal{U}_{p_2 \times p_2} \left( \begin{bmatrix} \tilde{Q}_3^L & 0 \end{bmatrix}^T \right)^T P^T \\ &= \begin{bmatrix} \mathcal{U}_{p_1 \times p_2} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) \\ 0 \end{bmatrix} (I \otimes [W + W Y^L Y W])^{-1} \begin{bmatrix} \mathcal{U}_{p_1 \times p_2} \left( [Q_3^L \quad 0]^T \right) \\ 0 \end{bmatrix}^T \\ &= \begin{bmatrix} \mathcal{U}_{p_1 \times p_2} \left( \begin{bmatrix} Q_3 \\ 0 \end{bmatrix} \right) (I \otimes [W + W Y^L Y W])^{-1} \mathcal{U}_{p_1 \times p_2} \left( [Q_3^L \quad 0]^T \right)^T \\ 0 \end{bmatrix}. \quad \square \end{aligned}$$

## References

- [1] L. Amodei and J.-M. Buchot. An invariant subspace method for large-scale Riccati equation. *Appl. Numer. Math.*, 60:1067–1082, 2010.
- [2] Brian D.O. Anderson. Second-order convergent algorithms for the steady-state Riccati equation. *Internat. J. Control*, 28(2):295–306, 1978.
- [3] Panos J. Antsaklis and Anthony N. Michel. *A Linear Systems Primer*. Birkhäuser, Boston, 2007.
- [4] P. Benner and Z. Bujanović. On the solution of large-scale algebraic Riccati equations by using low-dimensional invariant subspaces. *Linear Algebra Appl.*, 488:430–459, 2016.
- [5] P. Benner, Z. Bujanović, P. Kürschner, and J. Saak. RADI: a low-rank ADI-type algorithm for large-scale algebraic Riccati equations. *Numer. Math.*, 138:301–330, 2018.
- [6] P. Benner, M. Heinkenschloss, J. Saak, and H. K. Weichelt. An inexact low-rank Newton-ADI method for large-scale Riccati equations. *Appl. Numer. Math.*, 108:125–142, 2016.
- [7] P. Benner, J.-R. Li, and T. Penzl. Numerical solution of large Lyapunov equations, Riccati equations, and linear-quadratic control problems. *J. Numer. Lin. Alg. Appl.*, pages 755–777, 2008.
- [8] P. Benner and H. Mena. Numerical solution of the infinite-dimensional LQR-problem and the associated differential Riccati equations. *Numer. Math.*, 26:1–20, 2018.
- [9] P. Benner and J. Saak. A Newton-Galerkin-ADI method for large-scale algebraic Riccati equations. In *Applied Linear Algebra*. GAMM Workshop Applied and Numerical Linear Algebra, May 2010.

- [10] P. Benner and J. Saak. Numerical solution of large and sparse continuous time algebraic matrix Riccati and Lyapunov equations: a state of the art survey. *GAMM-Mitteilungen*, 36:32–52, 2013.
- [11] Peter Benner, Zvonimir Bujanović, Patrick Kürschner, and Jens Saak. A numerical comparison of different solvers for large-scale, continuous-time algebraic Riccati equations and LQR problems. *SIAM J. Sci. Comput.*, 42(2):A957–A996, 2020.
- [12] D. A. Bini, B. Iannazzo, and B. Meini. *Numerical Solution of Algebraic Riccati Equations*, volume 9 of *Fundamentals of Algorithms*. SIAM Publications, Philadelphia, 2012.
- [13] T. Çimen. State-dependent Riccati equations. *IFAC Proc. Vol.*, 41:3761–3775, 2008.
- [14] Eric K.-W. Chu, H. Y. Fan, and W.-W. Lin. A structure-preserving doubling algorithm for continuous-time algebraic Riccati equations. *Linear Algebra Appl.*, 396:55–80, 2005.
- [15] Eric K.-W. Chu, H.-Y. Fan, W.-W. Lin, and C.-S. Wang. A structure-preserving doubling algorithm for periodic discrete-time algebraic Riccati equations. *Internat. J. Control*, 77:767–788, 2004.
- [16] Timothy A. Davis and Yifan Hu. The university of Florida sparse matrix collection. *ACM Trans. Math. Software*, 38(1):Article 1, 2011. 25 pages.
- [17] L. Dieci. Numerical integration of the differential Riccati equation and some related issues. *SIAM J. Numer. Anal.*, 29:781–815, 1992.
- [18] V. Druskin and V. Simoncini. Adaptive rational Krylov subspaces for large-scale dynamical systems. *Sys. Contr. Lett.*, 60:546–560, 2011.
- [19] V. Druskin, V. Simoncini, and M. Zaslavsky. Adaptive tangential interpolation in rational krylov subspaces for MIMO dynamical systems. *SIAM J. Matrix Anal. Appl.*, 35:476–498, 2014.
- [20] B. Friedlander, M. Morf, T. Kailath, and L. Ljung. New inversion formulas for matrices classified in terms of their distance from Toeplitz matrices. *Linear Algebra Appl.*, 27:31–60, 1979.
- [21] Z.-C. Guo, E. K.-W. Chu, X. Liang, and W.-W. Lin. A decoupled form of the structure-preserving doubling algorithm with low-rank structures. *ArXiv e-prints*, may 2020. 18 pages, arXiv: 2005.08288.
- [22] Stefan Güttel. Rational Krylov approximation of matrix functions: numerical methods and optimal pole selection. *GAMM-Mitteilungen*, 36:8–31, 2013.
- [23] M. Heyouni and K. Jbilou. An extended block Arnoldi algorithm for large-scale solutions of continuous-time algebraic Riccati equation. *Electron. Trans. Numer. Anal.*, 33:53–62, 2009.
- [24] T.-M. Huang, R.-C. Li, and W.-W. Lin. *Structure-Preserving Doubling Algorithms for Nonlinear Matrix Equations*. Fundamentals of Algorithms. SIAM, Philadelphia, 2018.
- [25] T.-M. Huang and W.-W. Lin. Structured doubling algorithms for weakly stabilizing Hermitian solutions of algebraic Riccati equations. *Linear Algebra Appl.*, 430:1452–1478, 2009.
- [26] Vlad Ionescu, Cristian Oară, and Martin Weiss. *Generalized Riccati Theory and Robust Control: A Popov Function Approach*. John Wiley & Sons, Chichester, UK, 1999.
- [27] K. Jbilou. An Arnoldi based algorithm for large algebraic Riccati equations. *Appl. Math. Lett.*, 19:437–444, 2006.
- [28] Xiaoqing Jin. *Developments and Applications of Block Toeplitz Iterative Solvers*. Science Press, Beijing, 2002.
- [29] Xiaoqing Jin. *Preconditioning Techniques for Toeplitz Systems*. Higher Education Press, Beijing, 2010.
- [30] Thomas Kailath and Joohwan Chun. Generalized Gohberg-Semencul formulas for matrix inversion. *Oper. Theory: Adv. Appl.*, 40:231–246, 1989.
- [31] Thomas Kailath and Joohwan Chun. Generalized displacement structure for block-Toeplitz, Toeplitz-block, and Toeplitz-derived matrices. *SIAM J. Matrix Anal. Appl.*, 15(1):114–128, 1994.
- [32] Thomas Kailath, Sun-Yuan Kung, and Martin Morf. Displacement ranks of matrices and linear equations. *J. Math. Anal. and Appl.*, 68:395–407, 1979.
- [33] Thomas Kailath and Ali H. Sayed. Displacement structure: theory and applications. *SIAM Rev.*, 37(3):297–386, 1995.
- [34] P. Lancaster and L. Rodman. Solutions of continuous and discrete time algebraic riccati equations: A review. In S. Bittanti, A. J. Laub, and J. C. Willems, editors, *The Riccati Equations*, pages 11–52. Springer-Verlag, Berlin, Heidelberg, 1991.
- [35] P. Lancaster and L. Rodman. *Algebraic Riccati Equations*. The clarendon Press, Oxford Sciece Publications, New York, 1995.
- [36] T. Li, E. K.-W. Chu, W.-W. Lin, and P. C.-Y. Weng. Solving large-scale continuous-time algebraic Riccati equations by doubling. *J. Comput. Appl. Math.*, 237:373–383, 2013.
- [37] W.-W. Lin and S.-F. Xu. Convergence analysis of structure-preserving doubling algorithm for Riccati-type matrix equations. *SIAM J. Matrix Anal. Appl.*, 28:26–39, 2006.
- [38] Y. Lin and V. Simoncini. A new subspace iteration method for the algebraic Riccati equation. *J. Numer. Lin. Alg. Appl.*, 22(1):26–47, 2015.
- [39] V. L. Mehrmann. The autonomous linear quadratic control problems. In *Lecture Notes in Control and Information Sciences*, volume 163. Springer-Verlag, Berlin, 1991.

- [40] Oberwolfach Benchmark Collection. Steel profile. hosted at MORwiki – Model Order Reduction Wiki, 2005.
- [41] J. Saak, M. Köhler, and P. Benner. M-M.E.S.S.-2.1 – the matrix equations sparse solvers library, April 2021. see also: <https://www.mpi-magdeburg.mpg.de/projects/mess>.
- [42] V. Simoncini, D. Szyld, and M. Monsalve. On two numerical methods for the solution of large-scale algebraic Riccati equations. *IMA J. Numer. Anal.*, 34:904–920, 2014.
- [43] Gilbert Strang. A proposal for Toeplitz matrix calculations. *Stud. Appl. Math.*, 74:171–176, 1986.
- [44] N. Wong and V. Balakrishnan. Quadratic alternating direction implicit iteration for the fast solution of algebraic Riccati equations. In *Proceedings of International Symposium on Intelligent Signal Processing and Communication Systems*, pages 373–376, 2005.
- [45] N. Wong and V. Balakrishnan. Fast positive-real balanced truncation via quadratic alternating direction implicit iteration. *IEEE Trans. Computer-Aided Design of Integrated Circuits and Systems*, 26(9):1725–1731, 2007.