

# Embeddings and Integrable Charges for Extended Corner Symmetry

Luca Ciambelli

*Physique Mathématique des Interactions Fondamentales & International Solway Institutes,  
Université Libre de Bruxelles, Campus Plaine - CP 231, 1050 Bruxelles, Belgium*

Robert G. Leigh and Pin-Chun Pai

*Illinois Center for Advanced Studies of the Universe & Department of  
Physics, University of Illinois, 1110 West Green St., Urbana IL 61801, U.S.A.*

We revisit the problem of extending the phase space of diffeomorphism-invariant theories to account for embeddings associated with the boundary of sub-regions. We do so by emphasizing the importance of a careful treatment of embeddings in all aspects of the covariant phase space formalism. In so doing we introduce a new notion of the extension of field space associated with the embeddings which has the important feature that the Noether charges associated with all extended corner symmetries are in fact integrable, but not necessarily conserved. We give an intuitive understanding of this description. We then show that the charges give a representation of the extended corner symmetry via the Poisson bracket, without central extension.

## INTRODUCTION

Classical gravitational theories are gauge theories for which the principal symmetry is diffeomorphism invariance. The study of symmetry in gauge theories has a long history going back to the seminal work of Noether [1]. While the first Noether theorem deals with global symmetries, Noether herself remarked that for a gauge symmetry, defined as a local symmetry acting differently at different spacetime points, the first theorem gives a current associated to the symmetry that necessarily vanishes on-shell up to total derivatives. It then follows that Noether charges for gauge symmetries must be defined as surface integrals of  $d - 2$  forms, where  $d$  is the spacetime dimension, i.e., as surface charges. Thus codimension-2 surfaces play a central role in gravity and other gauge theories. We will refer to them here as *corners* [2–8] for brevity.

The first discussion of the role of gauge symmetries and their surface charges appeared using the Hamiltonian formalism in the seminal work of Regge and Teitelboim [9]. Later, Wald and others [10–12], using the covariant phase-space formalism, and Barnich and Brandt [13], using Anderson’s variational bicomplex [14, 15], formulated the Lagrangian analysis of gauge symmetries and their conserved quantities. The theory of asymptotic symmetries and surface charges following from these works has been shown to give the same results, modulo ambiguities, as reviewed for example in [16].

In any diffeomorphism-invariant classical theory on a manifold  $M$ , in the absence of any extra structure we are free to perform any diffeomorphism, as it corresponds to a gauge redundancy. However, in the presence of some geometric structure, such as a subregion of  $M$  or a corner, some diffeomorphisms will become physical symmetries. This is a familiar feature of other gauge theories as well, where in the corresponding quantum theory, non-trivial physics is involved in the gluing of subregions and is im-

plicated in entanglement properties. Often one addresses such geometric structures through the mathematical construction of embeddings [2, 3, 7]. In particular, in [7] we explored the consequences of carefully treating such embeddings, and found a theory-independent finite sub-algebra of  $\mathfrak{diff}(M)$ , compatible with the presence of an embedded submanifold  $S$ , whose Lie brackets close on itself, which we referred to as the maximal embedding symmetry,  $\mathcal{A}_k = (\mathfrak{diff}(S) \ltimes \mathfrak{gl}(k, \mathbb{R})) \ltimes \mathbb{R}^k$ . Applied to corners ( $k = 2$ ), this algebra includes the so-called extended corner algebra [8], which in turn includes notable sub-cases, such as the BMSW algebra [17], the generalized BMS algebra [18–20], the extended BMS algebra [21, 22], as well as the original BMS algebra [23–25].

In diffeomorphism-invariant theories, a basic construction is the integration of a Noether charge density over a corner. An important issue in this regard is *integrability*, whether or not a diffeomorphism is equivalently associated with a Hamiltonian vector field on the space of fields of the theory and a function on field space. If integrability pertains, then diffeomorphisms are represented by such functions and the algebra of charges reproduces canonically the vector field algebra via the Poisson bracket, possibly with central extension [26]. A separate issue is how the possible non-conservation of charge in a given region is associated with the flow of flux in or out of the region. As advocated in [2], such features can be obtained in general if one extends the field space of the theory to include effects associated with embedded submanifolds. However, in that and subsequent analyses, it was found that diffeomorphisms that do not preserve a corner are on a much different footing than those that do: only the latter were found to correspond to integrable charges. The diffeomorphisms that do not preserve a corner are its normal translations, which appear in the  $\mathbb{R}^2$  factor in the extended corner symmetry  $\mathcal{A}_2 = (\mathfrak{diff}(S) \ltimes \mathfrak{gl}(2, \mathbb{R})) \ltimes \mathbb{R}^2$ . In the context of asymptotic symmetries, it is this factor that gives rise to the so-called supertranslations, and so

it is certainly of central interest to fully understand it.

In [7] by carefully treating embeddings we found that the maximal embedding algebra corresponds precisely to the set of those diffeomorphisms that become physical in the presence of a corner and furthermore that the variation of the charge with respect to any diffeomorphism gives precisely  $\delta_{\eta} H_{\underline{\xi}} = H_{[\underline{\xi}, \eta]}$ , which appears ready-made for an interpretation in terms of Poisson brackets. This result was obtained purely geometrically without reference to the covariant phase space formalism beyond a specification of the Noether charge density.

In this letter, we will show how to extend the field space of diffeomorphism-invariant theories such that the entire extended corner symmetry is realized by integrable charges. This is achieved by introducing the same careful treatment of embeddings into the covariant phase space methodology, which leads to a natural extension of the symplectic structure of the theory which differs from that introduced in [2]. We stress that integrability is achieved on the entire field space (on-shell) without a need for first setting some flux to zero. The existing literature contains many examples where integrability is achieved only in this sense. Here we propose a universal resolution and the consequences are far-reaching: we find that the algebra of charges associated with the full extended corner symmetry (and thus any of its subalgebras appropriate to any particular physical situation) is represented in terms of Poisson brackets on the extended phase space, without central extension. As a further consequence, the flux is accounted for fully by the field-space contraction of the Hamiltonian vector field associated to the symmetry and the extended pre-symplectic potential.

## EMBEDDING MAPS

In this section, we describe in detail embedding maps. This material of course is standard but it is important to carefully establish our notation. We are concerned entirely here with classical diffeomorphism-invariant theories. We thus suppose that we have a spacetime manifold  $M$  of dimension  $d$  that we endow locally with coordinates via the trivialization  $\Phi_i : U_i \rightarrow \mathbb{R}^d$ ,  $U_i \subset M$ , giving  $\Phi_i : p \mapsto y^\mu$ . An embedded submanifold of  $M$  of codimension  $k$  is the image of a (smooth) map  $\phi_k : S_k \rightarrow M$ , where  $S_k$  is a manifold of dimension  $d - k$ . We introduce coordinates  $\{\sigma^\alpha\}$  on  $S_k$  with the embedding described by  $\phi_k : \{\sigma^\alpha\} \mapsto \{y^\mu(\sigma)\}$ , the latter thought of as the locus of points of the embedded manifold in  $M$ . Given the embedding, differential forms  $\alpha \in \Gamma(\wedge^{d-k} T^*M)$  can be pulled back to a top form  $\phi_k^* \alpha \in \Gamma(\wedge^{d-k} T^*S_k)$  on  $S_k$ . Such a form can then be integrated over  $S_k$ .

We will be interested in describing a variation of such an embedding  $\phi_k \rightarrow \phi_k + \delta\phi_k$ , promoted to an arbitrary variation from the perspective of covariant phase space. Given such, we can then express the response of the em-

bedding to a diffeomorphism as a contraction<sup>1</sup> of  $\delta\phi_k$  with a vector field on phase space, i.e.,  $\delta_{\eta}\phi_k = l_{V_{\underline{\xi}}}\delta\phi_k$ . Essentially, this idea was implemented originally in [2] by writing the corresponding change of the embedding coordinates as

$$\{\sigma^\alpha\} \rightarrow \{y^\mu(\sigma^\alpha) + \chi^\mu(\sigma^\alpha)\} \quad (1)$$

which we interpret as defining a vector field on  $M$ ,  $\underline{\chi} \in TM$  defined at all points on the embedded manifold. Thus we have  $\delta\phi_k \leftrightarrow \underline{\chi}$  and we will consequently regard  $\underline{\chi}$  as a 1-form on field space, the details of which we will review below.

Consider a  $(d - k)$ -form  $\alpha[g, y]$  where  $g$  collectively denotes the fields of a theory and  $y$  denotes the coordinates on  $M$  in which we are working. Given an embedding  $\phi_k : S_k \rightarrow M$ , we can pull back  $\alpha$  to  $S_k$  and integrate

$$A[g, \phi] := \int_{S_k} \phi_k^* (\alpha[g, y]). \quad (2)$$

Often in the literature these details are not expressed clearly and instead integrals are expressed directly in the coordinates of  $M$ . We will see that keeping track of such details will have important consequences for the covariant phase space formalism. An important result that we will use repeatedly is Stokes' theorem, which in the notation we are using reads

$$\int_{S_k} \phi_k^* (d\beta[g, y]) = \int_{\partial S_k} \phi_{k-1}^* (\beta[g, y]) \quad (3)$$

where  $\partial S_k$  is the boundary of  $S_k$ , its embedding given by  $\phi_{k-1} : \partial S_k \rightarrow M$ .

In [7], we made use of this formalism in a simple setting, in which we simply integrated the Noether charge density in this way over a corner, corresponding to the case  $k = 2$ ,

$$H_{\underline{\xi}} = \int_{S_2} \phi_2^* (Q_{\underline{\xi}}) = \int_{S_2} \phi_2^* (*d\underline{\xi}g), \quad (4)$$

where  $\underline{\xi} \in TM$  is an arbitrary vector field on  $M$  and  $Q_{\underline{\xi}}$  is the Noether charge density, making no further reference to any other aspects of the covariant phase space formalism. In the second equality we inserted the specific representation of the Noether charge for the Einstein-Hilbert theory. We found that only those vector fields in the maximal embedding symmetry<sup>2</sup>  $\tilde{\mathcal{A}}_k =$

<sup>1</sup> Throughout the paper, we use the notation  $i_{\underline{\xi}}$  for the contraction of a spacetime vector field with a differential form and  $l_{V_{\underline{\xi}}}$  for the corresponding contraction on field space; here  $V_{\underline{\xi}}$  denotes the vector field on field space associated with the spacetime diffeomorphism generated by  $\underline{\xi}$ . Similarly, Lie derivatives are denoted  $\mathcal{L}_{\underline{\xi}} = d i_{\underline{\xi}} + i_{\underline{\xi}} d$  and  $\mathcal{L}_{V_{\underline{\xi}}} = \delta l_{V_{\underline{\xi}}} + l_{V_{\underline{\xi}}} \delta$ .

<sup>2</sup> The tilde denotes the fact that whereas the maximal embedding algebra is the algebra of the group  $(\text{Diff}(S) \times \text{GL}(k, \mathbb{R})) \times \mathbb{R}^k$ , only the group  $(\text{Diff}(S) \times \text{SL}(k, \mathbb{R})) \times \mathbb{R}^k$  is realized in the Einstein-Hilbert theory.

$(\text{diff}(S) \ltimes \mathfrak{sl}(k, \mathbb{R})) \ltimes \mathbb{R}^k$  give rise to non-zero charges. Thus all other diffeomorphisms remain pure gauge, even in the presence of a corner. We then showed that

$$\delta_{\underline{\xi}} H_{\underline{\eta}} = -H_{[\underline{\xi}, \underline{\eta}]}. \quad (5)$$

It is tempting to interpret this result as a bracket

$$\{[H_{\underline{\xi}}, H_{\underline{\eta}}]\} = -H_{[\underline{\xi}, \underline{\eta}]}. \quad (6)$$

If this bracket were to coincide with the Poisson bracket of the covariant phase space formalism for all  $\tilde{\mathcal{A}}_2$  diffeomorphisms, then we would conclude that the symmetry is represented on phase space without central extension.

We will show in the remainder of the paper that precisely this pertains if we correctly enlarge the field space to include the degrees of freedom associated with embedding maps corresponding to subregions of  $M$  and hypersurfaces, and extend the symplectic structure accordingly. Such a correction is in the spirit of [2] but differs in a crucial way. In particular we will show that the correct extension of the symplectic structure yields integrable charges for *all* of  $\tilde{\mathcal{A}}_2$  and furthermore, non-conservation of the charges associated with flux is fully accounted for by the extended symplectic structure.

Consider the variation of (2)

$$\delta A[g, \phi] = \delta \int_{S_k} \phi_k^*(\alpha[g, y]). \quad (7)$$

The variation receives two contributions, one from the variation of  $\alpha$ , and one from the variation of the embedding  $\phi_k$ ,

$$\delta A[g, \phi] = \int_{S_k} \phi_k^*(\delta\alpha[g, y]) + \int_{S_k} \delta\phi_k^*(\alpha[g, y]) \quad (8)$$

where as usual

$$\delta\alpha[g, y] := \alpha[g + \delta g, y] - \alpha[g, y] \quad (9)$$

while the variation of the embedding is given by

$$(\delta\phi_k^*)(\alpha[g, y]) := \phi_k^*(\alpha[g, y + \chi] - \alpha[g, y]) \quad (10)$$

$$= \phi_k^*(\mathcal{L}_{\underline{\chi}}\alpha[g, y]) \quad (11)$$

where we are using the notation introduced in eq. (1). Thus

$$\delta A[g, \phi] = \int_{S_k} \phi_k^*(\delta\alpha[g, y] + \mathcal{L}_{\underline{\chi}}\alpha[g, y]). \quad (12)$$

The variation with respect to a diffeomorphism in particular is given by

$$l_{V_{\underline{\eta}}}\delta g = \mathfrak{L}_{V_{\underline{\eta}}}g = \mathcal{L}_{\underline{\eta}}g, \quad l_{V_{\underline{\eta}}}\underline{\chi} = -\underline{\eta}|_{\phi_k(S_k)} \quad (13)$$

which implies in particular that  $\delta_{\underline{\eta}}\phi_k = l_{V_{\underline{\eta}}}\delta\phi_k$ . Essentially similar relations have also appeared in [2, 27] for

example. Before moving on to covariant phase space formalism, we consider the implication of the nilpotency of  $\delta$ ,

$$0 = \delta^2 A[g, \phi] = \int_{S_k} \phi_k^* \left( \mathcal{L}_{\delta\underline{\chi}}\alpha[g, y] + \frac{1}{2}\mathcal{L}_{[\underline{\chi}, \underline{\chi}]} \alpha[g, y] \right)$$

from which we conclude

$$\delta\underline{\chi} = -\frac{1}{2}[\underline{\chi}, \underline{\chi}]. \quad (14)$$

Thus  $\underline{\chi}$  is not an exact form in field space generally. We note that this result resembles a property of ghosts, which is by no means an accident. Indeed this is another sign that in the presence of an embedding, a ‘vertical’ degree of freedom becomes physical. Related ideas appear in [3, 28].

## COVARIANT PHASE SPACE

In the case where one integrates a Lagrangian density over all of the manifold  $M$ , no embedding is required because the Lagrangian is a top-form on  $M$ . However, suppose we consider integrating the Lagrangian over some subregion  $R$  of  $M$ ; we will regard that subregion as embedded in  $M$  via  $\phi_0 : R \rightarrow M$ , with the boundary of the subregion regarded as an embedded hypersurface in  $M$ . So we write

$$S_R[g, \phi] = \int_R \phi_0^*(L[g, y]). \quad (15)$$

Given this, we then write

$$\begin{aligned} \delta S_R[g, \phi] &= \int_R \phi_0^*(\delta L[g, y]) + \int_R \delta\phi_0^*(L[g, y]) \\ &\hat{=} \int_{\partial R} \phi_1^*(\theta[g, \delta g, y]) + \int_R \phi_0^*(\mathcal{L}_{\underline{\chi}}L[g, y]) \\ &\hat{=} \int_{\partial R} \phi_1^*(\theta[g, \delta g, y] + i_{\underline{\chi}}L[g, y]). \end{aligned} \quad (16)$$

where  $\theta$  is the usual pre-symplectic form satisfying  $\delta L \hat{=} d\theta$ . This result suggests a corresponding extended pre-symplectic potential,

$$\Theta_{\Sigma}^{\text{ext.}} \equiv \int_{\Sigma} \phi_1^*(\theta[g, \delta g, y] + i_{\underline{\chi}}L[g, y]) \quad (17)$$

where  $\phi_1$  now embeds a hypersurface  $\phi_1 : \Sigma \rightarrow M$ , possibly with boundary. This extension is different than the one first introduced in [2]. To understand its properties, we consider the corresponding extended pre-symplectic structure, which can be shown to be of the form

$$\begin{aligned} \Omega_{\Sigma}^{\text{ext.}} &:= \delta\Theta_{\Sigma}^{\text{ext.}} = \int_{\Sigma} \phi_1^*(\delta\theta[g, \delta g, y]) \\ &+ \int_{\partial\Sigma} \phi_2^*(i_{\underline{\chi}}\theta[g, \delta g, y] + \frac{1}{2}i_{\underline{\chi}}i_{\underline{\chi}}L[g, y]). \end{aligned} \quad (18)$$

We then identify

$$\Omega_{\Sigma}^{\text{ext.}}[g, \delta g, \underline{\chi}, y] = \Omega_{\Sigma}[g, \delta g, y] + \Omega_{\partial\Sigma}^{\text{cor.}}[g, \delta g, \underline{\chi}, y].$$

Note that if embeddings are not considered as in the standard treatment, then the result would reduce to the first line of (18). This is only consistent however if we restrict attention to diffeomorphisms that preserve the corner  $\partial\Sigma$ . On the other hand, similar extended structures were given in [2, 3] but the prescription used there differs from ours in an important way, the details of which we will discuss below.

We now turn our attention to the question of integrability, and so we compute

$$l_{V_{\underline{\eta}}}\Omega_{\Sigma}^{\text{ext.}} = l_{V_{\underline{\eta}}}\Omega_{\Sigma} + l_{V_{\underline{\eta}}}\Omega_{\partial\Sigma}^{\text{cor.}}.$$

For the first term, we have the standard result

$$l_{V_{\underline{\eta}}}\Omega_{\Sigma} = \int_{\Sigma} \phi_1^* (l_{V_{\underline{\eta}}}\delta\theta) \hat{=} \int_{\Sigma} \phi_1^* (di_{\underline{\eta}}\theta - \delta dQ_{\underline{\eta}}) \quad (19)$$

where here we have assumed that  $\underline{\eta}$  is an arbitrary field-independent vector field as well as the covariance of  $\theta$ ,  $\mathfrak{L}_{V_{\underline{\eta}}}\theta = \mathcal{L}_{\underline{\eta}}\theta$  [11] (see also [29]). Here  $Q_{\underline{\eta}}$  is the Noether charge density, defined as usual by

$$dQ_{\underline{\eta}} = l_{V_{\underline{\eta}}}\theta - i_{\underline{\eta}}L. \quad (20)$$

Next, we consider the corner contribution,

$$\begin{aligned} l_{V_{\underline{\eta}}}\Omega_{\Sigma}^{\text{cor.}} &= \int_{\partial\Sigma} \phi_2^* \left( l_{V_{\underline{\eta}}}i_{\underline{\chi}}\theta + \frac{1}{2}l_{V_{\underline{\eta}}}i_{\underline{\chi}}i_{\underline{\chi}}L \right) \\ &= - \int_{\partial\Sigma} \phi_2^* \left( i_{\underline{\eta}}\theta + i_{\underline{\chi}}l_{V_{\underline{\eta}}}\theta + \frac{1}{2}i_{\underline{\eta}}i_{\underline{\chi}}L - \frac{1}{2}i_{\underline{\chi}}i_{\underline{\eta}}L \right) \\ &= - \int_{\partial\Sigma} \phi_2^* \left( i_{\underline{\eta}}\theta + i_{\underline{\chi}}(dQ_{\underline{\eta}} + \frac{1}{2}i_{\underline{\eta}}L) + \frac{1}{2}i_{\underline{\eta}}i_{\underline{\chi}}L \right) \\ &= - \int_{\partial\Sigma} \phi_2^* \left( i_{\underline{\eta}}\theta + i_{\underline{\chi}}dQ_{\underline{\eta}} \right) \end{aligned} \quad (21)$$

and so

$$l_{V_{\underline{\eta}}}\Omega_{\Sigma}^{\text{ext.}} \hat{=} - \int_{\Sigma} \phi_1^* (\delta dQ_{\underline{\eta}} + \mathcal{L}_{\underline{\chi}}dQ_{\underline{\eta}}) = -\delta \int_{\partial\Sigma} \phi_2^* (Q_{\underline{\eta}})$$

where we made use of eq. (12) in reverse. Thus the charge is integrable, and we write

$$l_{V_{\underline{\eta}}}\Omega_{\Sigma}^{\text{ext.}} = -\delta \int_{\partial\Sigma} \phi_2^* (Q_{\underline{\eta}}) = -\delta H_{\underline{\eta}} \quad (22)$$

Thus  $V_{\underline{\eta}}$  is the Hamiltonian vector field on field space corresponding to the Hamiltonian function  $H_{\underline{\eta}}$ , the latter coinciding precisely with that appearing in (4). Continuing in the same vein, it can be shown that

$$l_{V_{\underline{\xi}}}l_{V_{\underline{\eta}}}\Omega_{\Sigma}^{\text{ext.}} = -l_{V_{\underline{\xi}}}\delta \int_{\partial\Sigma} \phi_2^* (Q_{\underline{\eta}}) = \int_{\partial\Sigma} \phi_2^* (Q_{[\underline{\xi}, \underline{\eta}]}) = H_{[\underline{\xi}, \underline{\eta}]}$$

which coincides with the result of [7]. Thus we have the Poisson bracket

$$\{H_{\underline{\xi}}, H_{\underline{\eta}}\} = \delta_{\underline{\xi}}H_{\underline{\eta}} = -H_{[\underline{\xi}, \underline{\eta}]}. \quad (23)$$

This proves that the charges are canonically represented, given the choice (17). This is achieved without any restriction on the field space, without boundary conditions being imposed and without involvement of ambiguities that would alter the definition of the Noether charge.

Finally we note that  $\Theta_{\Sigma}^{\text{ext.}}$  correctly accounts for flux. Consider for example a ‘pillbox’ with spacelike boundaries  $\Sigma_{1,2}$  and a timelike boundary  $B$  (whose boundary is  $S_1 \cup S_2$ ). Then

$$H_{\underline{\xi}}|_{S_2} - H_{\underline{\xi}}|_{S_1} = \int_B \phi_1^* (dQ_{\underline{\xi}}) \quad (24)$$

$$= \int_B \phi_1^* (l_{V_{\underline{\xi}}}\theta - i_{\underline{\xi}}L) \quad (25)$$

$$= l_{V_{\underline{\xi}}}\int_B \phi_1^* (\theta + i_{\underline{\chi}}L) \quad (26)$$

$$= l_{V_{\underline{\xi}}}\Theta_B^{\text{ext.}} \quad (27)$$

where we used again our definition (17) applied to  $B$ . Thus the flux of the extended symplectic form precisely accounts for possible non-conservation of the charges. This applies to all vector fields  $\underline{\xi}$ , whether it be tangential or orthogonal.

### Comparison to Previous Treatments

In the usual treatment, where embeddings are not considered at all, one has  $\Theta_{\Sigma} = \int_{\Sigma} \theta$  and  $l_{V_{\underline{\eta}}}\Omega_{\Sigma} = -\int_{\partial\Sigma} (\delta Q_{\underline{\eta}} - i_{\underline{\eta}}\theta)$  as above. Thus, one says that the charge is integrable if  $i_{\underline{\eta}}\theta$  is either zero, or is set to zero by imposing some condition. Note that it is automatically zero if  $\underline{\eta}$  is tangent to  $\partial\Sigma$ , but not if  $\underline{\eta}$  is normal. In fact, this is precisely the problem that we have resolved by including the embedding – the charges are then integrable for all  $\underline{\eta}$ .

Finally we consider the relationship of our treatment to that of [2, 3]. In fact, they chose a different extension by making use of an ambiguity in the pre-symplectic form

$$\Theta_{\Sigma}^{\text{DF}} = \Theta_{\Sigma}^{\text{ext.}} + \int_{\partial\Sigma} \phi_2^* (Q_{\underline{\chi}}) \quad (28)$$

$$= \int_{\Sigma} \phi_1^* (\theta + i_{\underline{\chi}}L + dQ_{\underline{\chi}}). \quad (29)$$

This has the property that  $l_{V_{\underline{\eta}}}\Theta_{\Sigma}^{\text{DF}} = 0$ , and thus all diffeomorphism charges have been made to vanish. However, as we have seen, it is the pre-symplectic structure  $\Theta_{\Sigma}^{\text{ext.}}$  that gives rise to the actual Noether charges that generate the extended corner symmetry.

### CONCLUSIONS

In this letter, we have emphasized the need for an interpretation in which physical symmetries are associated

with corners and we have found the description of field space that gives rise to that interpretation. We plan to follow up this work with a more extensive account in which we study many of the examples of current interest and extend the construction to field-dependent vector fields and the corresponding modified brackets. For example, one may anticipate that our formalism has important implications when applied to the case of black hole horizons.

*Acknowledgements* We thank L. Freidel for communication about this result and for sharing an unpublished note containing a related discussion. The research of LC was partially supported by a Marina Solvay Fellowship, by the ERC Advanced Grant “High-Spin-Grav” and by FNRS-Belgium (convention FRFC PDR T.1025.14 and convention IISN 4.4503.15). The work of RGL was supported by the U.S. Department of Energy under contract DE-SC0015655.

- 
- [1] E. Noether, Invariante variationsprobleme, *Nach. Ges. Wiss. Gött.* **1918**, 235 (1918).
- [2] W. Donnelly and L. Freidel, Local subsystems in gauge theory and gravity, *JHEP* **09**, 102, [arXiv:1601.04744 \[hep-th\]](#).
- [3] A. J. Speranza, Local phase space and edge modes for diffeomorphism-invariant theories, *JHEP* **02**, 021, [arXiv:1706.05061 \[hep-th\]](#).
- [4] L. Freidel, M. Geiller, and D. Pranzetti, Edge modes of gravity. Part III. Corner simplicity constraints, *JHEP* **01**, 100, [arXiv:2007.12635 \[hep-th\]](#).
- [5] L. Freidel, M. Geiller, and D. Pranzetti, Edge modes of gravity. Part II. Corner metric and Lorentz charges, *JHEP* **11**, 027, [arXiv:2007.03563 \[hep-th\]](#).
- [6] L. Freidel, M. Geiller, and D. Pranzetti, Edge modes of gravity. Part I. Corner potentials and charges, *JHEP* **11**, 026, [arXiv:2006.12527 \[hep-th\]](#).
- [7] L. Ciambelli and R. G. Leigh, Isolated surfaces and symmetries of gravity, *Phys. Rev. D* **104**, 046005 (2021), [arXiv:2104.07643 \[hep-th\]](#).
- [8] L. Freidel, R. Oliveri, D. Pranzetti, and S. Speziale, Extended corner symmetry, charge bracket and Einstein’s equations, *JHEP* **09**, 083, [arXiv:2104.12881 \[hep-th\]](#).
- [9] T. Regge and C. Teitelboim, Role of Surface Integrals in the Hamiltonian Formulation of General Relativity, *Annals Phys.* **88**, 286 (1974).
- [10] R. M. Wald, Black hole entropy is the Noether charge, *Phys. Rev. D* **48**, 3427 (1993), [arXiv:gr-qc/9307038](#).
- [11] V. Iyer and R. M. Wald, Some properties of Noether charge and a proposal for dynamical black hole entropy, *Phys. Rev. D* **50**, 846 (1994), [arXiv:gr-qc/9403028](#).
- [12] R. M. Wald and A. Zoupas, A General definition of ‘conserved quantities’ in general relativity and other theories of gravity, *Phys. Rev. D* **61**, 084027 (2000), [arXiv:gr-qc/9911095](#).
- [13] G. Barnich and F. Brandt, Covariant theory of asymptotic symmetries, conservation laws and central charges, *Nucl. Phys. B* **633**, 3 (2002), [arXiv:hep-th/0111246](#).
- [14] I. M. Anderson, Introduction to the variational bicomplex, in *Math. aspects of classical field theory*, Contemp. Math., Vol. 132 (Amer. Math. Soc., Providence, RI, 1992) pp. 51–73.
- [15] I. M. Anderson and C. G. Torre, Asymptotic conservation laws in field theory, *Phys. Rev. Lett.* **77**, 4109 (1996), [arXiv:hep-th/9608008](#).
- [16] G. Compère, *Advanced Lectures on General Relativity*, Lecture Notes in Physics, Vol. 952 (Springer, 2019).
- [17] L. Freidel, R. Oliveri, D. Pranzetti, and S. Speziale, The Weyl BMS group and Einstein’s equations, *JHEP* **07**, 170, [arXiv:2104.05793 \[hep-th\]](#).
- [18] M. Campiglia and A. Laddha, Asymptotic symmetries and subleading soft graviton theorem, *Phys. Rev. D* **90**, 124028 (2014), [arXiv:1408.2228 \[hep-th\]](#).
- [19] G. Compère, A. Fiorucci, and R. Ruzzi, Superboost transitions, refraction memory and super-Lorentz charge algebra, *JHEP* **11**, 200, [Erratum: *JHEP* 04, 172 (2020)], [arXiv:1810.00377 \[hep-th\]](#).
- [20] M. Campiglia and J. Peraza, Generalized BMS charge algebra, *Phys. Rev. D* **101**, 104039 (2020), [arXiv:2002.06691 \[gr-qc\]](#).
- [21] G. Barnich and C. Troessaert, Aspects of the BMS/CFT correspondence, *JHEP* **05**, 062, [arXiv:1001.1541 \[hep-th\]](#).
- [22] G. Barnich and C. Troessaert, BMS charge algebra, *JHEP* **12**, 105, [arXiv:1106.0213 \[hep-th\]](#).
- [23] H. Bondi, M. G. J. Van der Burg, and A. W. K. Metzner, Gravitational waves in general relativity, VII. Waves from axi-symmetric isolated system, *Proc. Roy. Soc. London.* **A269**, 21 (1962).
- [24] R. Sachs and H. Bondi, Gravitational Waves in General Relativity VIII. Waves in Asymptotically Flat Space-time, *Proc. Roy. Soc. London.* **A270**, 103 (1962).
- [25] R. Sachs, Gravitational Waves in General Relativity. 8. Waves in asymptotically flat space-times, *Proc. Roy. Soc. Lond. A* **270**, 103 (1962).
- [26] J. D. Brown and M. Henneaux, Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three-Dimensional Gravity, *Commun. Math. Phys.* **104**, 207 (1986).
- [27] A. J. Speranza, Geometrical tools for embedding fields, submanifolds, and foliations, (2019), [arXiv:1904.08012 \[gr-qc\]](#).
- [28] H. Gomes and A. Riello, The observer’s ghost: notes on a field space connection, *JHEP* **05**, 017, [arXiv:1608.08226 \[math-ph\]](#).
- [29] V. Chandrasekaran and A. J. Speranza, Anomalies in gravitational charge algebras of null boundaries and black hole entropy, *JHEP* **01**, 137, [arXiv:2009.10739 \[hep-th\]](#).