

LOGARITHMIC HARNACK INEQUALITIES FOR TRANSITION SEMIGROUPS IN HILBERT SPACES

LUCIANA ANGIULI, DAVIDE A. BIGNAMINI*, SIMONE FERRARI

ABSTRACT. We consider the stochastic differential equation

$$\begin{cases} dX(t) = [AX(t) + F(X(t))]dt + C^{1/2}dW(t), & t > 0; \\ X(0) = x \in \mathcal{X}; \end{cases}$$

where \mathcal{X} is a Hilbert space, $\{W(t)\}_{t \geq 0}$ is a \mathcal{X} -valued cylindrical Wiener process, A, C are suitable operators on \mathcal{X} and $F : \text{Dom}(F) \subseteq \mathcal{X} \rightarrow \mathcal{X}$ is a smooth enough function. We establish a logarithmic Harnack inequality for the transition semigroup $\{P(t)\}_{t \geq 0}$ associated with the stochastic problem above, under less restrictive conditions than those considered in the literature. Some applications to these inequalities are also shown.

1. INTRODUCTION

The first formulation of the Harnack inequality dates back to 1887 and can be found in his seminal paper [21], and concerns positive harmonic functions. After some partial extensions, the most important contribution is due to J. Moser [29] which proved the Harnack inequality for positive (weak) solutions of uniformly elliptic linear equations with bounded coefficients in variational form. Moser also stresses the usefulness of such kind of estimates to deduce regularity results, such as the local hölderianity of the solutions. The further passage towards non-linear elliptic equations was made first by J. Serrin [39] and then by N.S. Trudinger [40] a few years later, and is based on Moser's approach.

The first parabolic version of the Harnack inequality is proved separately from J. Hadamard [20] and B. Pini [35] for positive solutions of the heat equation. Many years later this kind of estimates have been extended to positive solutions of more general linear parabolic equations by Moser himself [30]. Hence the extension to almost linear parabolic equations was due to D. G. Aronson and J. Serrin [1] and N.S. Trudinger [40]. Differently from the elliptic case, however, the case of operators with non-linear coefficients turned out to be more difficult and remained unresolved for a long time. In this direction we refer to [11, 12] where an *intrinsic* Harnack type inequality was proved for solutions of a large class of nonlinear equations and for operators with nonlinear coefficients. The techniques used in these latter results were inspired by the method of E. De Giorgi and J. Nash (see [10, 32]) to show boundedness and regularity for certain classes of functions (the so-called De Giorgi classes), which contain in particular the solutions of some elliptic equations.

We refer to [24] and the reference therein for a more in-depth analysis of the Harnack inequality. In all the quoted results, the formulation of the Harnack inequality allows to compare the values of a positive solution of some elliptic or parabolic differential equation, at two different points. All these Harnack inequalities are dimension-dependent and thus they cannot pass to infinite dimension. A possibility to get the Harnack-type inequality in an infinite dimensional setting consists in replacing the classical formulation to the dimension-free logarithmic Harnack Inequality (LHI) first introduced by F.-Y. Wang in [41] for the study of diffusion semigroups on

Date: November 29, 2021.

2010 Mathematics Subject Classification. 60H10, 60J60.

Key words and phrases. Infinite dimensional analysis, transition semigroup, logarithmic Harnack inequality.

*Corresponding author.

a Riemannian manifold M . It reads as

$$(P(t)f)^\alpha(x) \leq (P(t)f^\alpha)(x)e^{c(t)\rho(x,y)}, \quad t > 0, x, y \in M \quad (1.1)$$

which holds true for any positive and Borel bounded function f , any $\alpha > 1$ and some continuous function $c(t)$. Here ρ is a Riemannian metric on M . Also in the infinite dimensional setting, this kind of inequality has been used to obtain a lot of results, like some regularizing effects of the semigroup (see, for example, [6, Proposition 4.1], [37, Corollary 1.2] and [43, Corollary 7.3.14]) as well as some hyperboundedness properties for the semigroup $\{P(t)\}_{t \geq 0}$ (see, for example, [37] and [42]). We refer to [43] and the reference therein for a discussion of this inequality and its consequences.

The aim of this paper consists in proving a (LHI) like (1.1), for transition semigroups associated to some stochastic partial differential equations in infinite dimensional separable Hilbert spaces under more less restrictive conditions than those considered in the literature.

We consider $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ a normal filtered probability space, \mathcal{X} a separable Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and associated norm $\|\cdot\|$ and the stochastic partial differential equation

$$\begin{cases} dX(t) = [AX(t) + F(X(t))]dt + C^{1/2}dW(t), & t > 0; \\ X(0) = x \in \mathcal{X}, \end{cases} \quad (1.2)$$

where $\{W(t)\}_{t \geq 0}$ is a \mathcal{X} -valued cylindrical Wiener process, $C \in \mathcal{L}(\mathcal{X})$ is a positive operator, $F : \text{Dom}(F) \subseteq \mathcal{X} \rightarrow \mathcal{X}$ is a regular enough function, and $A : \text{Dom}(A) \subseteq \mathcal{X} \rightarrow \mathcal{X}$ is a possibly unbounded operator. Throughout the paper we will assume some hypotheses on A , F and C to guarantee the existence of a mild solution for (1.2) (see Hypotheses 2.6 and 3.1). Stochastic partial differential equations like (1.2) are widely studied in the literature (see [8, 43] and the references therein) as well as the validity of some (LHI) in Hilbert spaces for the associated semigroup

$$P(t)\varphi(x) := \mathbb{E}[\varphi(X(t, x))], \quad x \in \mathcal{X}, t \geq 0, \varphi \in B_b(\mathcal{X}); \quad (1.3)$$

(see, for example, [6, 38]). Here $\mathbb{E}[\cdot]$ denotes the expectation with respect to \mathbb{P} and $B_b(\mathcal{X})$ is the space of bounded and Borel measurable function on \mathcal{X} . Estimates like (1.1) for the transition semigroup (1.3) can be found for instance in [6, 17, 22, 27, 38, 44]. In all the quoted papers two different sets of assumptions for A , F and C are made in order to get inequalities like (1.1). Concerning the operator C , it is required some sort of invertibility. In [6, 44] it is required that C itself admits continuous and bounded linear inverse. In [22, 27] it is assumed that CC^* is invertible, while in [17] the authors restrict themselves to consider as C the identity operator. On the other hand, for what concerns the function F , it is usually required that it is Lipschitz continuous and that it satisfies the following dissipativity type condition:

$$\langle F(x) - F(y), C^{-1}(x - y) \rangle \leq \zeta \|C^{-1/2}(x - y)\|, \quad x - y \in C(\mathcal{X}). \quad (1.4)$$

for some $\zeta \in \mathbb{R}$. This latter condition can be found in [38] where C actually can depend also on x .

The main results of this paper are stated in Theorems 3.12 and 4.7 where a (LHI) similar to (1.1) is obtained for the transition semigroup (1.3) without any hypotheses of invertibility on C and assuming only one between the Lipschitz and the dissipativity condition appearing in (1.4) (see Section 3 for the Lipschitz continuous case and Section 4 for the dissipative case). To be more precise we will prove a (LHI) type inequality along the direction of the square root of the diffusion operator C , namely

$$|P(t)\varphi(x + h)|^p \leq P(t)|\varphi(x)|^p e^{c(t)\|C^{-1/2}h\|^2}, \quad t > 0, x \in \mathcal{X}, h \in C^{1/2}(\mathcal{X}); \quad (1.5)$$

for any bounded and Borel measurable function $\varphi : \mathcal{X} \rightarrow \mathbb{R}$, any $p > 1$ and some continuous function $c : (0, +\infty) \rightarrow \mathbb{R}$. We point out that if C has a continuous inverse then (1.5) is equivalent to (1.1).

The key tool we use to prove the (LHI) (3.14) and (4.15) in both cases is an approximation method. In the Lipschitz continuous case the approximants which allow us to get our estimate are suitable finite dimensional semigroups which satisfy suitable gradient estimates. On the other

hand, the dissipative case is solved by using a double approximation procedure which consider a finite dimensional approximation of the Yosida approximants.

For our results, we have collected some standard consequences in Section 5. For instance, the (LHI) can be used to prove a strong Feller-type property for $\{P(t)\}_{t \geq 0}$ and, assuming further the existence of an invariant measure μ for $\{P(t)\}_{t \geq 0}$, the occurrence of an entropy-cost inequality (see Corollary 5.1 (iii)). In this last case, another classical consequence of the (LHI) is a hypercontractivity type estimate for the semigroup $\{P(t)\}_{t \geq 0}$ in $L^p(\mathcal{X}, \mu)$. Such estimate relies on the Hölder inequality and some integrability conditions with respect to μ of some exponential functions (see Corollary 5.3). Finally, a considerable set of examples of operators A, C and functions F to which our results can be applied, are collected at the end of Section 5.

2. NOTATIONS AND PRELIMINARIES

In this section we fix the notations and we recall some basic results we will use throughout the paper.

Let H_1 and H_2 be two Hilbert spaces with inner products $\langle \cdot, \cdot \rangle_{H_1}$ and $\langle \cdot, \cdot \rangle_{H_2}$ and associated norms $\|\cdot\|_{H_1}$ and $\|\cdot\|_{H_2}$, respectively. We denote by $B_b(H_1; H_2)$ (resp. $C_b(H_1; H_2)$) the set of functions $f : H_1 \rightarrow H_2$ which are bounded and Borel measurable (resp. continuous). When $H_2 = \mathbb{R}$ we simply write $B_b(H_1)$ (resp. $C_b(H_1)$). For any $k \in \mathbb{N} \cup \{\infty\}$, $C_b^k(H_1; H_2)$ consists of continuous functions $f : H_1 \rightarrow H_2$ which are k -times Fréchet differentiable with bounded and continuous derivatives up to order k . If $H_2 = \mathbb{R}$ we simply write $C_b^k(H_1)$ and for $k = 1$ then, for any $x \in H_1$, we denote by $\mathcal{D}f(x)$ the unique $k \in H_1$ such that

$$\lim_{\|h\|_{H_1} \rightarrow 0} \frac{|f(x+h) - f(x) - \langle h, k \rangle_{H_1}|}{\|h\|_{H_1}} = 0.$$

For any $k \in \mathbb{N} \cup \{\infty\}$, we denote by $\mathcal{F}C_b^{k,n}(H_1)$, the space of cylindrical C_b^k functions depending on n variables, i.e., the set of functions $f : H_1 \rightarrow \mathbb{R}$ such that $f(x) = \varphi(\langle x, h_1 \rangle_{H_1}, \dots, \langle x, h_n \rangle_{H_1})$, $x \in H_1$, for some $\varphi \in C_b^k(\mathbb{R}^n)$, $h_1, \dots, h_n \in H_1$ and $n \in \mathbb{N}$.

Let $G : \text{Dom}(G) \subseteq H_1 \rightarrow H_1$ be a linear operator and let $H_2 \subseteq H_1$. We call *part of G in H_2* the operator $G_{H_2} : \text{Dom}(G_{H_2}) \subseteq H_2 \rightarrow H_2$ defined as

$$\begin{aligned} \text{Dom}(G_{H_2}) &:= \{x \in \text{Dom}(G) \cap H_2 \mid Gx \in H_2\}; \\ G_{H_2}x &:= Gx, \quad x \in \text{Dom}(G_{H_2}). \end{aligned}$$

By $\mathcal{L}(H_1)$ we denote the set of all bounded linear operators from H_1 into itself and by $\text{Id}_{H_1} \in \mathcal{L}(H_1)$ the identity operator on H_1 . We say that $B \in \mathcal{L}(H_1)$ is *non-negative* (resp. *positive*) if for every $x \in H_1 \setminus \{0\}$

$$\langle Bx, x \rangle_{H_1} \geq 0 \quad (> 0).$$

On the other hand, $B \in \mathcal{L}(H_1)$ is a *non-positive* (resp. *negative*) operator, if $-B$ is non-negative (resp. positive). We remind the reader that such operators are self-adjoint (see [36, Section IV.4]). We recall that if a semigroup $\{P(t)\}_{t \geq 0} \subseteq \mathcal{L}(H_1)$ is strongly continuous, then there exists $M_0 \geq 1$ and $\eta_0 \in \mathbb{R}$ such that

$$\|P(t)\|_{\mathcal{L}(H_1)} \leq M_0 e^{\eta_0 t}, \quad t \geq 0.$$

For the general theory concerning linear operators and semigroups we refer to [14, 15, 16].

Now, let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ be a normal filtered probability space, \mathcal{K} be a separable Banach space and by $\mathcal{B}(\mathcal{K})$ we denote the family of the Borel subsets of \mathcal{K} . For any random variable $\xi : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow (\mathcal{K}, \mathcal{B}(\mathcal{K}))$, the *law of ξ* on $(\mathcal{K}, \mathcal{B}(\mathcal{K}))$, denoted by $\mathcal{L}(\xi)$ and the *expectation of ξ* with respect to \mathbb{P} denoted by $\mathbb{E}[\xi]$ are defined by the formulas

$$\mathcal{L}(\xi) := \mathbb{P} \circ \xi^{-1}$$

and

$$\mathbb{E}[\xi] := \int_{\Omega} \xi(w) \, \mathbb{P}(dw) = \int_{\mathcal{K}} x \, \mathcal{L}(\xi)(dx).$$

We need to recall the definition of some Banach spaces often used in the literature (see, for example, [4, Section 6.2]).

Definition 2.1. Let $I \subseteq [0, +\infty)$ be an interval and let $p \geq 1$. We denote by $C_p(I; \mathcal{K})$ the space of \mathcal{K} -valued processes $\{Y(t)\}_{t \in I}$ such that the function $(Y(\cdot))(\omega) : I \rightarrow \mathcal{K}$ is bounded and continuous for \mathbb{P} -a.e. $\omega \in \Omega$. We endow the space $C_p(I; \mathcal{K})$ with the norm

$$\|\{Y(t)\}_{t \in I}\|_{C_p(I; \mathcal{K})}^p := \mathbb{E} \left[\sup_{t \in I} \|Y(t)\|_{\mathcal{K}}^p \right].$$

We denote by $\mathcal{K}^p(I)$ the space of progressively measurable¹ \mathcal{K} -valued process $\{Y(t)\}_{t \in I}$ such that the quantity

$$\|\{Y(t)\}_{t \in I}\|_{\mathcal{K}^p(I)}^p := \sup_{t \in I} \mathbb{E} \left[\|Y(t)\|_{\mathcal{K}}^p \right],$$

is finite. We endow the space $\mathcal{K}^p(I)$ with the norm $\|\cdot\|_{\mathcal{K}^p(I)}$,

A map $f : \text{Dom}(f) \subseteq \mathcal{K} \rightarrow \mathcal{K}$ is said to be *dissipative* if for any $\alpha > 0$ and $x, y \in \text{Dom}(f)$, it holds

$$\|x - y - \alpha(f(x) - f(y))\|_{\mathcal{K}} \geq \|x - y\|_{\mathcal{K}} \quad (2.1)$$

In particular if \mathcal{K} is a Hilbert space (2.1) is equivalent to

$$\langle f(x) - f(y), x - y \rangle_{\mathcal{K}} \leq 0.$$

We say that f is *m-dissipative* if it is dissipative and the range of $\text{Id}_{\mathcal{K}} - f$ is the whole space \mathcal{K} .

Remark 2.2. If $f = A$ is linear, then (2.1) becomes $\|(\lambda \text{Id} - A)x\|_{\mathcal{K}} \geq \lambda \|x\|_{\mathcal{K}}$ for every $x \in \text{Dom}(A)$ and $\lambda > 0$. In addition, if \mathcal{K} is a Hilbert space with inner product $\langle \cdot, \cdot \rangle_{\mathcal{K}}$, then (2.1) is equivalent to say that $\langle Ah, h \rangle_{\mathcal{K}} \leq 0$ for any $h \in \mathcal{K}$.

Now we introduce a Hilbert space that plays a key role in the proof of the (LHI) (see Sections 3 and 4). In what follows, \mathcal{X} will be an Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and associated norm $\|\cdot\|$, and $C \in \mathcal{L}(\mathcal{X})$ will be a positive operator. We set

$$H_C := C^{1/2}(\mathcal{X}), \quad [h_1, h_2]_C := \langle C^{-1/2}h_1, C^{-1/2}h_2 \rangle, \quad h_1, h_2 \in C^{1/2}(\mathcal{X}).$$

It can be proved that $(H_C, [\cdot, \cdot]_C)$ is a Hilbert space continuously embedded in \mathcal{X} (see [3]). We set $\|\cdot\|_C$ the norm on H_C associated to the inner product $[\cdot, \cdot]_C$.

We give the definition of H_C -Fréchet differentiability.

Definition 2.3. A function $\Phi : \mathcal{X} \rightarrow \mathbb{R}$ is H_C -Fréchet differentiable at $x \in \mathcal{X}$, if there exists $h_x \in H_C$ such that

$$\lim_{\|h\|_C \rightarrow 0} \frac{|\Phi(x+h) - \Phi(x) - [h_x, h]_C|}{\|h\|_C} = 0.$$

Clearly, when h_x exists, it is unique and we set $\mathcal{D}_C \Phi(x) := h_x$.

Remark 2.4. Observe that if $\Phi : \mathcal{X} \rightarrow \mathbb{R}$ is a Fréchet differentiable function, then it is H_C -differentiable too. Furthermore, for every $x \in \mathcal{X}$ it holds

$$\mathcal{D}_C \Phi(x) = C \mathcal{D} \Phi(x),$$

This result can be found in [3, Proposition 17].

The definition of H_C -Gateaux differentiability is the Gateaux counterpart of the previous definition. We stress that for our purposes we need a more technical definition in comparison to the “standard” one.

¹We say that a \mathcal{K} -valued process ψ is *progressively measurable* if, as a transformation from $\Omega \times [0, t]$ equipped with the σ -field $\mathcal{F}_t \times \mathcal{B}([0, t])$ into $(\mathcal{K}, \mathcal{B}(\mathcal{K}))$ is measurable for any $[0, t] \subset I$.

Definition 2.5. We say that a function $\Phi : \mathcal{X} \rightarrow \mathcal{X}$ is H_C -Gateaux differentiable if for every $x \in \mathcal{X}$ and $h \in H_C$, there exists $\varepsilon_{x,h} > 0$ such that the function $\varphi_{x,h} : (-\varepsilon_{x,h}, \varepsilon_{x,h}) \rightarrow \mathcal{X}$ defined as

$$\varphi_{x,h}(r) := \Phi(x + rh) - \Phi(x)$$

is H_C -valued and there exists $L_x \in \mathcal{L}(H_C)$ such that for every $h \in H_C$

$$\lim_{r \rightarrow 0} \left\| \frac{1}{r} \varphi_{x,h}(r) - L_x h \right\|_C = 0. \quad (2.2)$$

See [3] for further properties of the space H_C .

We end this section by recalling a result of existence and uniqueness of the mild solution of the problem

$$\begin{cases} dX(t) = [AX(t) + F(X(t))]dt + C^{1/2}dW(t), & t > 0; \\ X(0) = x \in \mathcal{X}. \end{cases} \quad (2.3)$$

Here $\{W(t)\}_{t \geq 0}$ is a \mathcal{X} -valued cylindrical Wiener process (see [8, Section 4.1.2] for a definition) and the operator A, F and C satisfy the following assumptions.

Hypotheses 2.6. Let \mathcal{X} be a Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and associated norm $\|\cdot\|$.

- (i) There exists E a Borel subset of \mathcal{X} which is a Banach space continuously and densely embedded in \mathcal{X} ;
- (ii) $A : \text{Dom}(A) \subseteq \mathcal{X} \rightarrow \mathcal{X}$ generates a strongly continuous semigroup e^{tA} on \mathcal{X} and A_E (the part of A in E) generates an analytic semigroup e^{tA_E} on E . There exists $\zeta_A \in \mathbb{R}$ such that $A - \zeta_A \text{Id}_{\mathcal{X}}$ is dissipative in \mathcal{X} and $A_E - \zeta_A \text{Id}_E$ is dissipative in E . For any $T > 0$ and for \mathbb{P} -a.e. $\omega \in \Omega$ the function $W_A(\cdot)(\omega) : [0, T] \rightarrow E$ defined by

$$W_A(t)(\omega) := \left(\int_0^t e^{(t-s)A} C^{1/2} dW(s) \right) (\omega), \quad t \in [0, T], \quad (2.4)$$

is continuous and

$$\int_0^T \text{Tr}[e^{sA} C e^{sA^*}] ds < +\infty. \quad (2.5)$$

- (iii) $E \subseteq \text{Dom}(F)$ and $F(E) \subseteq E$. There exists $\zeta_F \in \mathbb{R}$ such that $F - \zeta_F \text{Id}_{\mathcal{X}}$ is m -dissipative in \mathcal{X} and $F|_E - \zeta_F \text{Id}_E$ is m -dissipative in E . $F|_E : E \rightarrow E$ is locally Lipschitz on E , namely $F|_E$ is Lipschitz continuous on bounded subsets of E and, there exist $M > 0$ and $m \in \mathbb{N}$ such that

$$\|F|_E(x)\|_E \leq M(1 + \|x\|_E^m), \quad x \in E. \quad (2.6)$$

- (iv) $C \in \mathcal{L}(\mathcal{X})$ is a positive operator.

Remark 2.7. If $E = \mathcal{X}$, by [8, Theorem 5.11], a sufficient condition that guarantees both the continuity of the trajectories of W_A and (2.5) is provided by the following: there exists $\eta \in (0, 1)$ such that

$$\int_0^T s^{-\eta} \text{Tr}[e^{sA} C e^{sA^*}] ds < +\infty. \quad (2.7)$$

If $E \neq \mathcal{X}$, other conditions can be found in [4, Sections 6.1-8.2] and [8, Sections 5.4-5.5].

For a x belonging to \mathcal{X} , a mild solution of (2.3) is a \mathcal{X} -valued adapted stochastic process $\{X(t, x)\}_{t \geq 0}$ satisfying

$$X(t, x) = e^{tA}x + \int_0^t e^{(t-s)A} F(X(s, x)) ds + \int_0^t e^{(t-s)A} C^{1/2} dW(s), \quad t \geq 0.$$

The following is a result of existence and uniqueness of a mild solution of (2.3) whose proof can be found in [2, Section 3].

Theorem 2.8. Assume Hypotheses 2.6 hold true.

- (i) For any $x \in E$, (2.3) has a unique mild solution $\{X(t, x)\}_{t \geq 0}$ belonging to $C_p([0, T], \mathcal{X}) \cap C_p((0, T], E)$, for any $T > 0$ and $p \geq 1$.

- (ii) For any $x \in \mathcal{X}$, (2.3) has a unique generalized mild solution $\{X(t, x)\}_{t \geq 0}$ belonging to $C_p([0, T], \mathcal{X})$, for any $T > 0$ and $p \geq 1$, namely, for any sequence $\{x_n\}_{n \in \mathbb{N}} \subseteq E$ converging to x in \mathcal{X} and for any $T > 0$ it holds

$$\lim_{n \rightarrow +\infty} \sup_{t \in [0, T]} \|X(t, x) - X(t, x_n)\| = 0,$$

where $\{X(t, x_n)\}_{t \geq 0}$ is the unique mild solution of (2.3) with initial datum x_n (see (i)).

Moreover for any $p \geq 1$, there exist two positive constants C_p, κ_p such that

$$\|X(t, x)\|_{\mathcal{O}}^p \leq C_p \left(e^{-\kappa_p t} \|x\|_{\mathcal{O}}^p + \|W_A(t)\|_{\mathcal{O}}^p + \int_0^t e^{-\kappa_p(t-s)} \|F(W_A(s))\|_{\mathcal{O}}^p ds \right),$$

for any $t > 0$ and $x \in \mathcal{O}$, being either $\mathcal{O} = E$ or $\mathcal{O} = \mathcal{X}$. Finally, there exists $\eta > 0$ such that

$$\|X(t, x_1) - X(t, x_2)\| \leq e^{-\eta t} \|x_1 - x_2\|, \quad t > 0, \quad x_1, x_2 \in \mathcal{X}. \quad (2.8)$$

We point out that all the (in)equalities in the above theorem and in what follows, if not otherwise specified, have to be understood to hold for \mathbb{P} -a.e. $\omega \in \Omega$.

3. LOGARITHMIC HARNACK INEQUALITY: THE LIPSCHITZ CONTINUOUS CASE

(LHI) are a classical tools to prove regularity results for transition semigroups associated to stochastic differential equations. They have already been proved in several settings and more is known in finite dimension. In this case, (LHI) are satisfied by solutions of the Cauchy problems associated to the equation

$$\frac{d}{dt} u(t, \xi) = \mathcal{A}(t) u(t, \xi), \quad t > 0, \quad \xi \in \mathbb{R}^n;$$

where $\mathcal{A}(t)\psi := \text{Tr}(Q(t)D^2\psi) + \langle b(t, x), D\psi \rangle$ for any smooth function ψ assuming suitable conditions on Q and b (see, for example, [9, 17, 19, 44]). In infinite dimension such kind of estimates have been already proved in [6] and in [38] for the transition semigroups associated to (2.3). In [6], (LHI) are proved assuming that $C^{-1} \in \mathcal{L}(\mathcal{X})$ and the dissipativity of F whereas in [38] they are proved assuming that C depends on the spatial variable and F is Lipschitz continuous and dissipative along both \mathcal{X} and H_C . Here we generalize the results in [38] when $C \in \mathcal{L}(X)$ assuming either F Lipschitz continuous or dissipative along H_C and in this last case without requiring that $C^{-1} \in \mathcal{L}(\mathcal{X})$.

We start by assuming the following set of hypotheses.

Hypotheses 3.1. *Hypotheses 2.6 are satisfied with $E = \mathcal{X}$. Furthermore we assume that $F : \mathcal{X} \rightarrow \mathcal{X}$ is a Fréchet differentiable and Lipschitz continuous function with Lipschitz constant L_F , and there exists $\zeta_{\mathcal{X}} > 0$ such that*

$$\langle (A + \mathcal{D}F(x))h, h \rangle \leq -\zeta_{\mathcal{X}} \|h\|^2, \quad x, h \in \mathcal{X}. \quad (3.1)$$

Remark 3.2. We point out that the Lipschitz continuity of F together with (3.1) yields the existence of $\bar{\zeta} \in \mathbb{R}$ such that $A - \bar{\zeta} \text{Id}_{\mathcal{X}}$ and $F - \bar{\zeta} \text{Id}_{\mathcal{X}}$ are both dissipative. Moreover, we remark that the following weaker form of (3.1) can be assumed in Hypotheses 3.1 instead of (3.1)

$$\langle A(h_1 - h_2) + F(x + h_1) - F(x + h_2), h_1 - h_2 \rangle \leq -\zeta_{\mathcal{X}} \|h_1 - h_2\|^2, \quad h_1, h_2 \in \mathcal{X}.$$

For the sake of clarity in the proofs of this section we decided to assume (3.1).

Under Hypotheses 3.1 the stochastic problem (2.3) admits a unique mild solution $\{X(t, x)\}_{t \geq 0}$ satisfying (2.8) and allows us to define the family of operators $\{P(t)\}_{t \geq 0}$ as

$$P(t)\varphi(x) := \mathbb{E}[\varphi(X(t, x))], \quad x \in \mathcal{X}, \quad t \geq 0, \quad \varphi \in B_b(\mathcal{X}). \quad (3.2)$$

Further regularity properties of the mild solution $\{X(t, x)\}_{t \geq 0}$ of (2.3) will be needed in the sequel.

Proposition 3.3. *Assume Hypotheses 3.1 hold true and let $T > 0$. The map $x \mapsto \{X(t, x)\}_{t \in [0, T]}$ from \mathcal{X} into $\mathcal{X}^2([0, T])$ (see Definition 2.1) is Gateaux differentiable and for any $x, y \in \mathcal{X}$, its Gateaux derivative is the unique mild solution of*

$$\begin{cases} dY_x(t) = [A + \mathcal{D}F(X(t, x))]Y_x(t)dt, & t \in (0, T]; \\ Y_x(0) = y, \end{cases} \quad (3.3)$$

namely, for every $x, y \in \mathcal{X}$ the process $\{\mathcal{D}_G X(t, x)y\}_{t \in [0, T]}$ satisfies

$$\mathcal{D}_G X(t, x)y = e^{tA}y + \int_0^t e^{(t-s)A} \mathcal{D}F(X(s, x)) \mathcal{D}_G X(s, x)y ds, \quad t \in [0, T]. \quad (3.4)$$

In addition it holds that

$$\|\mathcal{D}_G X(t, x)y\| \leq e^{-\zeta_X t} \|y\|, \quad t \in [0, T], \quad x, y \in \mathcal{X}. \quad (3.5)$$

Proof. The first part of the statement is classical and can be found in [8, Theorem 9.8] or [28]. To prove estimate (3.5) we assume that $\{Y_x(t, y)\}_{t \geq 0}$ is a strict solution of (3.3), otherwise we can proceed as in [2, Proposition 3.6] or [4, Proposition 6.2.2] approximating $\{Y_x(t, y)\}_{t \geq 0}$ by means of a sequence of more regular processes. In this case, scalarly multiplying the stochastic differential equation in (3.3) by $Y_x(t, y)$ and using (3.1) we obtain

$$\frac{1}{2} \frac{d}{dt} \|Y_x(t, y)\|^2 = \langle [A + \mathcal{D}F(X(t, x))]Y_x(t, y), Y_x(t, y) \rangle \leq -\zeta_X \|Y_x(t, y)\|^2,$$

whence we infer (3.5). \square

Our proof of (LHI) relies on a gradient estimate for the transition semigroup $\{P(t)\}_{t \geq 0}$ (see (3.12)). To obtain such result we need some additional hypotheses on A, C and F .

Hypotheses 3.4. *Assume that the part of A in H_C , denoted by A_C , generates a contractive strongly continuous semigroup e^{tA_C} in H_C and that one of the following two conditions is satisfied:*

(i) *there exists $\gamma \in (0, 1)$ such that*

$$e^{tA}(\mathcal{X}) \subseteq C^{1/2}(\mathcal{X}), \quad \|C^{-1/2}e^{tA}\|_{\mathcal{L}(\mathcal{X})} \leq Kt^{-\gamma} \quad (3.6)$$

for any $t > 0$ and some positive constant K independent of t ;

(ii) *$F = C^{1/2}G$ for some Fréchet differentiable and Lipschitz continuous function $G : \mathcal{X} \rightarrow \mathcal{X}$ with Lipschitz constant L_G .*

We refer to Section 5 for some examples of operators A and C verifying Hypotheses 3.4.

The following proposition gives us information about the continuity of $h \mapsto \mathcal{D}_G X(t, x)h$ as a linear map in H_C .

Proposition 3.5. *Assume Hypotheses 3.1 and 3.4 hold true. For any $t > 0$ and $x \in \mathcal{X}$, the operator $\mathcal{D}_G X(t, x)$ belongs to $\mathcal{L}(H_C)$. More precisely*

$$\|\mathcal{D}_G X(t, x)h\|_C \leq K_1 \|h\|_C, \quad t > 0, \quad x \in \mathcal{X}, \quad h \in H_C, \quad (3.7)$$

for some positive constant K_1 independent of t, x and h .

Proof. We start by assuming that Hypotheses 3.4(i) hold true. For any $t > 0$, $x \in \mathcal{X}$ and $h \in H_C$, by (3.4) and the definition of $\|\cdot\|_C$,

$$\|\mathcal{D}_G X(t, x)h\|_C \leq \|e^{tA_C}h\|_C + \int_0^t \|C^{-1/2}e^{(t-s)A} \mathcal{D}F(X(s, x)) \mathcal{D}_G X(s, x)h\|_C ds$$

By the Lipschitz continuity of F , the contractivity of $\{e^{tA_C}\}_{t \geq 0}$ in H_C , Hypotheses 3.4(i) and (3.5) we have

$$\begin{aligned} \|\mathcal{D}_G X(t, x)h\|_C &\leq \|h\|_C + KL_F \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})} \|h\|_C \int_0^t \frac{e^{-\zeta_X s}}{(t-s)^\gamma} ds \\ &\leq \|h\|_C + KL_F \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})} \|h\|_C \left(\int_0^1 \frac{1}{s^\gamma} ds + \int_1^t e^{-\zeta_X(t-s)} ds \right) \end{aligned}$$

$$\leq \left[1 + KL_F \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})} \left(\frac{1}{1-\gamma} + \frac{1}{\zeta_X} \right) \right] \|h\|_C$$

whence the claim, with $K_1 := 1 + KL_F \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})} ((1-\gamma)^{-1} + \zeta_X^{-1})$.

Now we assume that Hypotheses 3.4(ii) are satisfied. In this case, using again (3.4) we have

$$\|\mathcal{D}_G X(t, x)h\|_C \leq \|e^{tA_C} h\|_C + \int_0^t \|e^{(t-s)A_C} C^{1/2} \mathcal{D}G(X(s, x)) \mathcal{D}_G X(s, x)h\|_C ds$$

for any $t > 0, x \in \mathcal{X}$ and $h \in H_C$. Thus, the Lipschitz continuity of G , the contractivity of $\{e^{tA_C}\}_{t \geq 0}$ in H_C and (3.5) yield that

$$\begin{aligned} \|\mathcal{D}_G X(t, x)h\|_C &\leq \|e^{tA_C} h\|_C + \int_0^t \|e^{(t-s)A_C} C^{1/2} \mathcal{D}G(X(s, x)) \mathcal{D}_G X(s, x)h\|_C ds \\ &\leq \|h\|_C + \int_0^t \|C^{1/2} \mathcal{D}G(X(s, x)) \mathcal{D}_G X(s, x)h\|_C ds \\ &= \|h\|_C + \int_0^t \|\mathcal{D}G(X(s, x)) \mathcal{D}_G X(s, x)h\| ds \\ &\leq \|h\|_C + L_G \int_0^t \|\mathcal{D}_G X(s, x)h\| ds \\ &\leq \|h\|_C + L_G \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})} \|h\|_C \int_0^t e^{-\zeta_X s} ds \\ &= \|h\|_C + L_G \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})} \|h\|_C \left(\frac{1}{\zeta_X} - \frac{e^{-\zeta_X t}}{\zeta_X} \right) \\ &\leq \left(1 + \frac{L_G \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})}}{\zeta_X} \right) \|h\|_C, \end{aligned}$$

whence the claim follows, with $K_1 := 1 + L_G \zeta_X^{-1} \|C^{1/2}\|_{\mathcal{L}(\mathcal{X})}$. \square

Remark 3.6. We could remove the requirement $\zeta_X > 0$ in Hypotheses 3.1 and replace the condition (3.6) with the following: there exist $c > 0$ and $\gamma \in (0, 1)$ such that

$$e^{tA}(\mathcal{X}) \subseteq C^{1/2}(\mathcal{X}), \quad \|C^{-1/2} e^{tA}\|_{\mathcal{L}(\mathcal{X})} \leq K e^{ct} t^{-\gamma}.$$

In this case (3.7) would be changed as follows: for any $T > 0$ there exist $K_1(T)$, such that for any $t > 0, x \in \mathcal{X}$ and $h \in H_C$ we have

$$\|\mathcal{D}_G X(t, x)h\|_C \leq K_1(T) \|h\|_C.$$

Now, we prove that for any fixed $t > 0$, the map $\mathcal{X} \ni x \mapsto X(t, x) \in \mathcal{X}$ is H_C -Gateaux differentiable.

Proposition 3.7. Assume that Hypotheses 3.1 and 3.4 hold true. For any $t > 0$, the map $x \mapsto X(t, x)$, from \mathcal{X} to \mathcal{X} , is H_C -Gateaux differentiable \mathbb{P} -a.e. and its H_C -Gateaux derivative along $h \in H_C$ is given by $\mathcal{D}_G X(t, x)h$.

Proof. We restrict ourselves to prove the claim when Hypotheses 3.4(i) hold true since the other case can be obtain similarly arguing as in Proposition 3.5. First of all we prove that for any fixed $t > 0, x \in \mathcal{X}$ and $h \in H_C$, the function $\varphi_{x,h}^t : \mathbb{R} \rightarrow \mathcal{X}$ defined as

$$\varphi_{x,h}^t(r) := X(t, x + rh) - X(t, x),$$

is H_C -valued. Using (3.3) we obtain

$$\varphi_{x,h}^t(r) = r e^{tA} h + \int_0^t e^{(t-s)A} [F(X(s, x + rh)) - F(X(s, x))] ds, \quad r > 0.$$

Thus, Hypotheses 3.4(i), the definition of $\|\cdot\|_C$, the Lipschitz continuity of F and estimate (2.8) infer

$$\begin{aligned}
 \|\varphi_{x,h}^t(r)\|_C &\leq \|re^{tA}h\|_C + \int_0^t \|e^{(t-s)A}[F(X(s, x+rh)) - F(X(s, x))]\|_C ds \\
 &\leq \|rC^{-1/2}e^{tA}h\| + \int_0^t \|C^{-1/2}e^{(t-s)A}[F(X(s, x+rh)) - F(X(s, x))]\| ds \\
 &\leq K|r|t^{-\gamma}\|h\| + \int_0^t \|C^{-1/2}e^{(t-s)A}\|_{\mathcal{L}(X)}\|F(X(s, x+rh)) - F(X(s, x))\| ds \\
 &\leq K|r|t^{-\gamma}\|h\| + KL_F \int_0^t (t-s)^{-\gamma}\|X(s, x+rh) - X(s, x)\| ds \\
 &\leq K|r|t^{-\gamma}\|h\| + KL_F|r\|\|h\| \int_0^t (t-s)^{-\gamma}e^{-\eta s} ds \\
 &\leq K|r\|\|h\| \left(t^{-\gamma} + L_F \frac{t^{1-\gamma}}{1-\gamma} \right) < +\infty.
 \end{aligned}$$

So $\varphi_{x,h}^t$ is H_C -valued.

Thanks to Proposition 3.5, to conclude we just need to prove (2.2) with $\varphi_{x,h}$ being replaced by $\varphi_{x,h}^t$. For any $x \in \mathcal{X}$, $h \in H_C$ and $t, r > 0$, by (3.4) and (3.6) we have

$$\begin{aligned}
 &\mathbb{E} \left[\left\| \frac{1}{r} \varphi_{x,h}^t(r) - \mathcal{D}_G X(t, x)h \right\|_C \right] \\
 &= \mathbb{E} \left[\left\| \int_0^t C^{-1/2}e^{(t-s)A} \left(\frac{F(X(s, x+rh)) - F(X(s, x))}{r} - \mathcal{D}F(X(s, x))\mathcal{D}_G X(s, x)h \right) ds \right\| \right] \\
 &\leq K \int_0^t \frac{1}{(t-s)^\gamma} \mathbb{E} \left[\left\| \frac{F(X(s, x+rh)) - F(X(s, x))}{r} - \mathcal{D}F(X(s, x))\mathcal{D}_G X(s, x)h \right\| \right] ds.
 \end{aligned}$$

By (2.8), Proposition 3.3, the fact that $\gamma \in (0, 1)$ and the fact that F is Fréchet differentiable and Lipschitz continuous we can apply the dominated convergence theorem to deduce that

$$\lim_{r \rightarrow 0} \mathbb{E} \left[\left\| \frac{1}{r} \varphi_{x,h}^t(r) - \mathcal{D}_G X(t, x)h \right\|_C \right] = 0,$$

which concludes the proof. \square

The following proposition is a chain rule for H_C -differentiable functions.

Proposition 3.8. *Assume Hypotheses 3.1 and 3.4 hold true. If $G : \mathcal{X} \rightarrow \mathbb{R}$ is a Fréchet differentiable function, then G is H_C -Fréchet differentiable and for any $t \geq 0$, $x \in \mathcal{X}$ and $h \in H_C$*

$$[\mathcal{D}_C (G \circ X(t, \cdot))(x), h]_C = [\mathcal{D}_C G(X(t, x)), \mathcal{D}_G X(t, x)h]_C.$$

Proof. The proof follows the same ideas of [3, Corollary 21] with some minor changes. \square

A key tool to prove the (LHI) stated in Theorem 3.12 is a finite-dimensional approximation procedure which allows us to approximate the transition semigroup $\{P(t)\}_{t \geq 0}$ by means of a sequence of transition semigroups $\{P^n(t)\}_{t \geq 0}$ associated to suitable finite-dimensional stochastic differential equations. The idea of such approximation comes from [6]. Here, for the sake of completeness and to point out the minimal assumptions needed for such kind of procedure, we recall it and we provide a proof of the main approximation result (Proposition 3.10).

We need one more condition.

Hypotheses 3.9. *Assume that H_C is dense in \mathcal{X} and that there exists a sequence of A -invariant and C -invariant finite dimensional subspaces $\mathcal{X}_n \subseteq \text{Dom}(A_C)$ such that $\bigcup_{n=1}^\infty \mathcal{X}_n$ is dense in H_C .*

Hypotheses 3.9 hold true for instance, if A is a self-adjoint positive operator and C admits a continuous inverse $C^{-1} \in \mathcal{L}(X)$ or if A and C are simultaneously diagonalizable.

In view of Hypothesis 3.9 we can consider $\{e_k\}_{k \in \mathbb{N}} \in \text{Dom}(A_C)$ such that for any $n \in \mathbb{N}$

$$\mathcal{X}_n = \text{span}\{e_1, \dots, e_n\},$$

and the family $\{e_k \mid k \in \mathbb{N}\}$ is an orthonormal basis of H_C . Further, let $\pi_n : \mathcal{X} \rightarrow \mathcal{X}_n$ be the orthogonal projection with respect to $(\mathcal{X}, \langle \cdot, \cdot \rangle)$, for any $n \in \mathbb{N}$ we can define $A_n : \mathcal{X} \rightarrow \mathcal{X}_n$, $C_n : \mathcal{X} \rightarrow \mathcal{X}_n$ and $F_n : \mathcal{X} \rightarrow \mathcal{X}_n$ as

$$A_n := \pi_n A \pi_n (= A \pi_n), \quad C_n := \pi_n C \pi_n (= C \pi_n) \quad \text{and} \quad F_n := \pi_n F \pi_n$$

Now, fix $n \in \mathbb{N}$ and consider

$$\begin{cases} dX_n(t) = [A_n X_n(t) + F_n(X_n(t))]dt + C_n^{1/2} dW_n(t), & t > 0, \\ X_n(0) = x \in \mathcal{X}_n. \end{cases} \quad (3.8)$$

Here $W_n(t) := \pi_n W(t) = \sum_{k=1}^n \langle W(t), e_k \rangle e_k$.

It is straightforward to see that A_n, C_n and F_n satisfy Hypotheses 3.1. Moreover, being C an injective operator it follows that C_n is bijective hence $C_n^{1/2} \mathcal{X}_n = \mathcal{X}_n$ for any $n \in \mathbb{N}$. Therefore, fixed $x \in \mathcal{X}_n$, by Theorem 2.8 we can deduce existence and uniqueness of a mild solution $\{X_n(t, x)\}_{t \geq 0}$ of (3.8) and consequently well-posedness for the associated transition semigroup defined for $f \in B_b(\mathcal{X}_n)$ as

$$P_n(t)f(x) := \mathbb{E}[f(X_n(t, x))], \quad t > 0, x \in \mathcal{X}_n. \quad (3.9)$$

we recall that the map $(t, x) \mapsto (P_n(t)f)(x)$ solves the Cauchy problem associated to the second order elliptic operator \mathcal{N}_n acting on cylindrical smooth functions as follows

$$\mathcal{N}_n \varphi(x) = \frac{1}{2} \text{Tr}[C_n D^2 \varphi(x)] + \langle A_n x + F_n(x), D\varphi(x) \rangle, \quad x \in \mathcal{X}_n, n \in \mathbb{N}.$$

Now we are able to state the main finite-dimensional approximation result. The proof of this result is the same of [6, Proposition 3.1(i)], with some minor changes. We decided to give it here for the sake of completeness.

Proposition 3.10. *Assume that Hypotheses 3.1 and 3.9 hold true. For any $f \in C_b(\mathcal{X})$, $t \geq 0$ and $x \in \mathcal{X}_{n_0}$, for some $n_0 \in \mathbb{N}$, it holds*

$$\lim_{n \rightarrow +\infty} P_n(t)f(x) = P(t)f(x).$$

Proof. Let $\{X(t, x)\}_{t \geq 0}$ be the unique mild solution of (2.3). For any $t \geq 0$, we set $Z(t) := X(t) - W_A(t)$, where $W_A(t)$ is defined in (2.4). For any fixed $x \in \mathcal{X}_{n_0}$ the process $\{Z(t)\}_{t \geq 0}$ is the unique mild solution of the problem

$$\begin{cases} dZ(t) = (AZ(t) + F(Z(t) + W_A(t)))dt, & t > 0; \\ Z(0) = x \in \mathcal{X}_{n_0}, \end{cases}$$

which satisfies

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|Z(t)\|^2 \right] < +\infty$$

Since

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|W_A(t)\|^2 \right] < +\infty,$$

by the dominated convergence theorem, it is easy to see that $\pi_n W_A(t)$ converges to $W_A(t)$ in $L^2(\Omega, \mathbb{P})$, as n tends to infinity.

Setting $W_n(t) := \pi_n W(t)$ and

$$W_{A_n}(t) := \int_0^t e^{(t-s)A_n} C_n^{1/2} dW_n(s), \quad t \geq 0,$$

then the process $Z_n(t) := X_n(t, x) - W_{A_n}(t)$, ($n \geq n_0$) satisfies

$$\begin{cases} dZ_n(t) = (A_n Z_n(t) + F_n(Z_n(t) + W_{A_n}(t)))dt, & t > 0; \\ Z_n(0) = x \in \mathcal{X}_{n_0}. \end{cases}$$

Now we split the proof in two steps. In the first one we show that $W_{A_n}(t) - W_A(t)$ converges to 0 in $L^2(\omega, \mathbb{P})$ and that, consequently $\|Z_n(t)\|$ is uniformly bounded with respect to n . In the second one we complete the proof.

Step 1. Using that $e_{|x_n}^{tA} = e^{tA_n}$ for every $t \geq 0$ we can write

$$W_A(s) - W_{A_n}(s) = \int_0^s e^{(s-r)A} (C^{1/2} - \pi_n C^{1/2} \pi_n) dW(r), \quad s \in [0, T]$$

By the Itô formula we deduce

$$\mathbb{E}(\|W_A(s) - W_{A_n}(s)\|^2) = \int_0^s \left\| e^{(s-r)A} (C^{1/2} - \pi_n C^{1/2} \pi_n) \right\|_{\mathcal{L}(X)}^2 dr$$

and, since the integrand converges to zero as $n \rightarrow \infty$ uniformly with respect to $r \in (0, s)$, by the dominated convergence theorem we get the claim. Now, scalarly multiplying $dZ_n(t) = (A_n Z_n(t) + F_n(Z_n(t) + W_{A_n}(t)))dt$ by $Z_n(t)$ and using the Hypotheses 3.1 with A and F being replaced by A_n and F_n we deduce (see (2.6))

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|Z_n(t)\|^2 &= \langle A_n Z_n(t) + F_n(Z_n(t) + W_{A_n}(t)), Z_n(t) \rangle \\ &= \langle A_n Z_n(t) + F_n(Z_n(t) + W_{A_n}(t)) \pm F_n(W_{A_n}(t)), Z_n(t) \rangle \\ &\leq \zeta x \|Z_n(t)\|^2 + \langle F_n(W_{A_n}(t)), Z_n(t) \rangle \\ &\leq \left(\zeta x + \frac{1}{2} \right) \|Z_n(t)\|^2 + \frac{1}{2} \|F_n(W_{A_n}(t))\|^2 \\ &\leq \left(\zeta x + \frac{1}{2} \right) \|Z_n(t)\|^2 + M 2^{-1} (1 + \|W_{A_n}(t)\|^2), \quad t > 0. \end{aligned}$$

The Gronwall lemma and the uniform boundedness of $\|W_{A_n}(t)\|$ with respect to n allows to deduce that $\|Z_n(t)\|$ is uniformly bounded with respect to n .

Step 2. To conclude the proof we show that $Z_n(t) - Z(t)$ converges to 0 in $L^2(\Omega, \mathbb{P})$ as $n \rightarrow \infty$. This fact will imply that $X_n(t, x)$ converges to $X(t, x)$ in $L^2(\Omega, \mathbb{P})$ as $n \rightarrow \infty$ and by (3.2) and (3.9) we conclude. By Hypothesis 3.9 we have

$$\begin{aligned} d(Z(t) - Z_n(t)) &= (AZ(t) - A_n Z_n(t) + F(X(t, x)) - F_n(X_n(t, x)))dt \\ &= (A(Z(t) - Z_n(t)) + F(X(t, x)) - F_n(X_n(t, x)))dt, \quad t > 0; \end{aligned}$$

so, scalarly multiplying by $Z(t) - Z_n(t)$ we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|Z(t) - Z_n(t)\|^2 &= \langle (A(Z(t) - Z_n(t)) + F(X(t, x)) - F_n(X_n(t, x))), Z(t) - Z_n(t) \rangle \\ &= \langle (A(Z(t) - Z_n(t)) + F(Z(t) + W_A(t)) \\ &\quad - F_n(Z_n(t) + W_{A_n}(t))), Z(t) - Z_n(t) \rangle. \end{aligned}$$

Adding and subtracting the terms $F(Z_n(t) + W_A(t))$ and $F(Z_n(t) + W_{A_n}(t))$ and using Hypotheses 3.1 we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|Z(t) - Z_n(t)\|^2 &\leq \zeta x \|Z(t) - Z_n(t)\|^2 \\ &\quad + \langle F(Z_n(t) + W_A(t)) - F(Z_n(t) + W_{A_n}(t)), Z(t) - Z_n(t) \rangle \\ &\quad + \langle F(Z_n(t) + W_{A_n}(t)) - \pi_n F(Z_n(t) + W_{A_n}(t)), Z(t) - Z_n(t) \rangle \\ &\leq \zeta x \|Z(t) - Z_n(t)\|^2 \\ &\quad + \|F(Z_n(t) + W_A(t)) - F(Z_n(t) + W_{A_n}(t))\| \|Z(t) - Z_n(t)\| \\ &\quad + \|(\text{Id}_X - \pi_n)F(Z_n(t) + W_{A_n}(t))\| \|Z(t) - Z_n(t)\| \end{aligned}$$

$$\begin{aligned}
&\leq \zeta_X \|Z(t) - Z_n(t)\|^2 + L_F \|W_A(t) - W_{A_n}(t)\| \|Z(t) - Z_n(t)\| \\
&\quad + \frac{1}{2} \|(\text{Id}_X - \pi_n)F(Z_n(t) + W_{A_n}(t))\|^2 + \frac{1}{2} \|Z(t) - Z_n(t)\|^2 \\
&\leq (\zeta_X + 1) \|Z(t) - Z_n(t)\|^2 + \frac{1}{2} L_F^2 \|W_A(t) - W_{A_n}(t)\|^2 \\
&\quad + \frac{1}{2} \|(\text{Id}_X - \pi_n)F(Z_n(t) + W_{A_n}(t))\|^2.
\end{aligned}$$

where in the last two lines we have used the Young inequality.

$$\begin{aligned}
&\frac{1}{2} \|Z(t) - Z_n(t)\|^2 \\
&\leq \int_0^t \left(\zeta_X + \frac{1}{2} \right) \|Z(s) - Z_n(s)\|^2 + \|F(X(s)) - F(X_n(s))\|^2 + \|(1 - \pi_n)F(X_n(s))\|^2 ds \\
&\leq \int_0^t \left(\zeta_X + \frac{1}{2} + 2L_F^2 \right) \|Z(s) - Z_n(s)\|^2 + 2L_F^2 \|W_A(s) - W_n(s)\|^2 + \|(1 - \pi_n)F(X_n(s))\|^2 ds.
\end{aligned}$$

Applying again the Gronwall lemma we obtain

$$\begin{aligned}
\|Z(t) - Z_n(t)\|^2 &\leq L_F^2 e^{2(\zeta_X + 1)t} \int_0^t \|W_A(s) - W_{A_n}(s)\|^2 ds \\
&\quad + e^{2(\zeta_X + 1)t} \int_0^t \|(\text{Id}_X - \pi_n)F(Z_n(s) + W_{A_n}(s))\|^2 ds \quad (3.10)
\end{aligned}$$

and using the results in Step 1 we infer that the right hand side of (3.10) vanishes as $n \rightarrow \infty$, concluding the proof. \square

We point out that the assumption of C -invariance of \mathcal{X}_n can be dropped in order to prove Proposition 3.10. However it is essential to apply the results in the previous sections to the finite-dimensional approximating operators. Indeed, assuming Hypotheses 3.9, it is easy to see that if A, C and F satisfy Hypothesis 3.4(i) then A_n, C_n and F_n satisfy them as well with the same constants. So all the results of the previous sections hold true even for the mild solution of (3.8) and for the semigroup in (3.9). In particular for every $h \in H_{C_n}$

$$\|\mathcal{D}_G X_n(t, x)h\|_{C_n} \leq K_1 \|h\|_{C_n}, \quad \text{for } x \in \mathcal{X}_n \text{ and } \mathbb{P}\text{-a.e.}, \quad (3.11)$$

where K_1 is the same constant appearing in Proposition 3.5.

The next result is a gradient estimate which is interesting in its own right. For our aims it is fundamental to prove Theorem 3.12.

Theorem 3.11. *Assume Hypotheses 3.1, 3.4 and 3.9 hold true. For any $\varphi \in C_b^1(\mathcal{X})$, $P(t)\varphi$ is Fréchet differentiable and for any $t > 0$*

$$\|\mathcal{D}_C P(t)\varphi(x)\|_C \leq K_1 P(t) \|\mathcal{D}_C \varphi(x)\|_C, \quad x \in \mathcal{X}; \quad (3.12)$$

and

$$\|\mathcal{D}_{C_n} P_n(t)\varphi(x)\|_{C_n} \leq K_1 P(t) \|\mathcal{D}_{C_n} \varphi(x)\|_{C_n}, \quad x \in \mathcal{X}_n, \quad n \in \mathbb{N}. \quad (3.13)$$

Proof. We restrict ourselves to prove estimate (3.12) since (3.13) can be obtained similarly using (3.11) in place of (3.7). By (3.5), Proposition 3.8 and [34, Fact 1.13(b), p. 8] it holds that $P(t)\varphi$ is both Fréchet differentiable and H_C -Fréchet differentiable (see Remark 2.4). Moreover, thanks to Propositions 3.5 and 3.8 we obtain

$$\begin{aligned}
[\mathcal{D}_C P(t)\varphi(x), h]_C &= [\mathcal{D}_C \mathbb{E}[\varphi(X(t, x))], h]_C \\
&= \mathbb{E}[[\mathcal{D}_C \varphi(X(t, x)), \mathcal{D}_G X(t, x)h]_C] \\
&\leq \mathbb{E}[\|\mathcal{D}_C \varphi(X(t, x))\|_C \|\mathcal{D}_G X(t, x)h\|_C] \\
&\leq K_1 \|h\|_C P(t) \|\mathcal{D}_C \varphi(x)\|_C,
\end{aligned}$$

for any $t \geq 0$, $x \in \mathcal{X}$ and $h \in H_C$. Now (3.12) follows by a standard argument. \square

We are ready to prove the main theorem of this section.

Theorem 3.12. *Assume Hypotheses 3.1, 3.4 and 3.9 hold true. For any $\varphi \in B_b(\mathcal{X})$ and $p > 1$ it holds*

$$|P(t)\varphi(x+h)|^p \leq P(t)|\varphi(x)|^p \exp\left(\frac{pK_1^2}{t(p-1)}\|h\|_C^2\right), \quad t > 0, \quad x \in \mathcal{X}, \quad h \in H_C. \quad (3.14)$$

Here K_1 is the constant appearing in Proposition 3.5.

Proof. Fix $n, n_0 \in \mathbb{N}$ with $n > n_0$ and $x, h \in \mathcal{X}_{n_0}$. We claim that estimate (3.14) holds true with $P(t)$, C , \mathcal{X} and φ replaced respectively by $P_n(t)$, C_n , \mathcal{X}_n and f , being $f \in \mathcal{F}_b^2(\mathcal{X})$ with positive infimum. To this aim, fix $\varepsilon > 0$ and $f \in \mathcal{F}_b^2(\mathcal{X})$ be such that $f(x) \geq \varepsilon$ for any $x \in \mathcal{X}$. We set

$$g_{n,n_0}(r, x) := P_n(r)f(x), \quad x \in \mathcal{X}_{n_0}, \quad r \geq 0,$$

and we note that for any $r > 0$, the function $g_{n,n_0} : [0, T] \times \mathcal{X}_{n_0} \rightarrow \mathbb{R}$ belongs to $C^{1,2}([0, T] \times \mathcal{X}_{n_0})$ solves

$$\begin{cases} D_r g_{n,n_0}(r, x) = \mathcal{N}_n g_{n,n_0}(r, x), & r > 0, \quad x \in \mathcal{X}_{n_0}; \\ g_{n,n_0}(0, x) = f(x), & x \in \mathcal{X}_{n_0}. \end{cases}$$

and satisfies $g_{n,n_0}(r, x) \geq \varepsilon$ for any $r \geq 0$ and $x \in \mathcal{X}_{n_0}$ (see [25, Theorem 1.2.5]). Now fix $t > 0$ and consider the function

$$G_n(r) := (P_n(t-r)g_{n,n_0}^p(r, \cdot))(x + rt^{-1}h), \quad r \in [0, t], \quad x, h \in \mathcal{X}_{n_0}.$$

For the sake of simplicity we let $\psi_h(r) := x + rt^{-1}h$. We differentiate the map $r \mapsto \ln(G(r))$.

$$\begin{aligned} \frac{d}{dr} \ln G_n(r) &= (G_n(r))^{-1} P_n(t-r) \left(-\mathcal{N}_n g_{n,n_0}^p(r, \cdot) + D_r g_{n,n_0}^p(r, \cdot) \right) (\psi_h(r)) \\ &\quad + (tG_n(r))^{-1} \langle \mathcal{D}P_n(t-r)g_{n,n_0}^p(\psi_h(r)), h \rangle. \end{aligned} \quad (3.15)$$

where we used that the semigroup and its generator commute on smooth functions. A straightforward computation yields that

$$-\mathcal{N}_n g_{n,n_0}^p + D_r g_{n,n_0}^p = -p(p-1)g_{n,n_0}^{p-2} \|C_n^{1/2} \mathcal{D}g_{n,n_0}\|_{\mathcal{X}_n}^2 \quad (3.16)$$

whereas by estimate (3.13) we infer

$$\begin{aligned} &\langle \mathcal{D}P_n(t-r)g_{n,n_0}^p(\psi_h(r)), h \rangle \\ &= \langle C_n^{1/2} \mathcal{D}P_n(t-r)g_{n,n_0}^p(\psi_h(r)), C_n^{-1/2}h \rangle \\ &\leq K_1 \|C_n^{-1/2}h\|_{\mathcal{X}_n} P_n(t-r) \left[pg_{n,n_0}^{p-1}(r, \cdot) \|C_n^{1/2} \mathcal{D}g_{n,n_0}(r, \cdot)\|_{\mathcal{X}_n} \right] (\psi_h(r)). \end{aligned} \quad (3.17)$$

Hence, using (3.16) and (3.17) in (3.15) we obtain

$$\frac{d}{dr} \ln G_n(r) \leq (G_n(r))^{-1} P_n(t-r) [-p(p-1)g_{n,n_0}^p(r, \cdot)(-v^2 + \beta v)](\psi_h(r))$$

where

$$v := g_{n,n_0}^{-1}(r, \psi_h(r)) \|C_n^{1/2} \mathcal{D}g_{n,n_0}(r, \psi_h(r))\|_{\mathcal{X}_n}, \quad \beta := \frac{K_1}{t(p-1)} \|C_n^{-1/2}h\|_{\mathcal{X}_n}.$$

The elementary inequality $a^2 - ab + 4^{-1}b^2 \geq 0$ which holds true for any $a, b \in \mathbb{R}$ allows us to estimate $-v^2 + \beta v \leq \beta^2$ and, consequently, by the positivity of $P_n(t)$ (see again [25, Theorem 1.2.5]), to deduce that

$$\frac{d}{dr} \ln G_n(r) \leq \frac{pK_1^2}{t^2(p-1)} \|C_n^{-1/2}h\|_{\mathcal{X}_n}^2.$$

whence, integrating from 0 to t with respect to r we obtain the claim.

Now, by a standard approximation argument, the Jensen inequality, Proposition 3.10 and the fact that $\|h\|_{C_n}$ converges to $\|h\|_C$ for any $h \in H_C$ as $n \rightarrow \infty$, we get

$$|P(t)f(x+h)|^p \leq P(t)|f(x)|^p \exp\left(\frac{pK_1^2}{t(p-1)}\|h\|_{H_C}^2\right), \quad t > 0, \quad (3.18)$$

for any $f \in \mathcal{FC}_b^2(\mathcal{X})$ and $x, h \in \bigcup_{i \in \mathbb{N}} \mathcal{X}_i$ (recall that $C_n^{1/2} \mathcal{X}_n = \mathcal{X}_n$ for any $n \in \mathbb{N}$).

A double approximation argument allows to extend (3.18) first to bounded and continuous functions and then to Borel bounded functions (see [18, Lemma 2.6 and Theorem 4.1(b)] and [23, Theorem 6.3]).

Finally, let us fix $x \in \mathcal{X}$ and $h \in H_C$, by Hypothesis 3.9 we can find two sequences $(x_n)_{n \in \mathbb{N}}$ and $(h_m)_{m \in \mathbb{N}}$ belonging to $\bigcup_{i \in \mathbb{N}} \mathcal{X}_i$ converging respectively to x in \mathcal{X} and h in H_C as $n, m \rightarrow \infty$ (recall that, by Hypotheses 3.9, H_C is dense in \mathcal{X}). Writing (3.18) with x and h being replaced by x_n and h_m we deduce

$$|P(t)f(x_n + h_m)|^p \leq P(t)|f(x_n)|^p \exp\left(\frac{pK_1^2}{t(p-1)}\|h_m\|_C^2\right), \quad t > 0, f \in B_b(\mathcal{X}). \quad (3.19)$$

By the continuity of the map $x \mapsto P(t)f(x)$, we complete the proof, letting n, m tend to ∞ in (3.19). \square

4. LOGARITHMIC HARNACK INEQUALITY: THE DISSIPATIVE CASE

The aim of this section is proving some (LHI) when no hypotheses of global Lipschitzianity for F is done but only some m -dissipativity along H_C . The main tool is again an approximation procedure this time using the Yosida approximants,

The standing assumptions we consider here are Hypotheses 2.6 which allows us to deal with the mild solution to (2.3) with initial datum $x \in E$ and also guarantees the existence of two positive constants C_p and κ_p such that for every $x \in E$ and $t \geq 0$

$$\|X(t, x)\|_E^p \leq C_p \left(e^{-\kappa_p t} \|x\|_E^p + \|W_A(t)\|_E^p + \int_0^t e^{-\kappa_p(t-s)} \|F(W_A(s))\|_E^p ds \right), \quad \mathbb{P}\text{-a.e.} \quad (4.1)$$

Consequently, the map $(t, x) \mapsto \|X(t, x)\|_E$ is bounded as a function from $[0, +\infty) \times E$ to \mathbb{R} .

We start by recalling the definition of the Yosida approximants for the function F . For any $\delta > 0$ and $x \in \mathcal{X}$ we let $J_\delta(x) \in \text{Dom}(F)$ be the unique solution of

$$y - \delta(F(y) - \zeta_F y) = x.$$

The existence of $J_\delta(x)$, for every $x \in \mathcal{X}$ and $\delta > 0$, is guaranteed by [7, Proposition 5.3.3]. We define $F_\delta : \mathcal{X} \rightarrow \mathcal{X}$ as

$$F_\delta(x) := F(J_\delta(x)), \quad x \in \mathcal{X}, \delta > 0.$$

Remark 4.1. We point out that $\text{Dom}(F_\delta) = \mathcal{X}$ and that Hypotheses 2.6 guarantee that, for any $\delta > 0$ and $x \in E$, $J_\delta(x)$ belongs to E , too. In particular, since E is preserved by the action of F , it turns out that $F_\delta(E) \subseteq E$.

Now, we set

$$\bar{\zeta}_F := \begin{cases} |\zeta_F|^{-1} & \text{if } \zeta_F \neq 0 \\ +\infty & \text{if } \zeta_F = 0 \end{cases}$$

and we list some useful properties of the Yosida approximants.

Lemma 4.2. *If Hypothesis 2.6(iii) holds true, then*

$$\lim_{\delta \rightarrow 0} \|J_\delta(y) - y\|_E = 0, \quad \lim_{\delta \rightarrow 0} \|(F_\delta)_{|_E}(y) - F_{|_E}(y)\|_E = 0, \quad y \in E. \quad (4.2)$$

For any $\delta \in (0, \bar{\zeta}_F)$, the functions $F_\delta - \zeta_F \text{Id}_{\mathcal{X}}$ and $(F_\delta)_{|_E} - \zeta_F \text{Id}_E$ are m -dissipative on \mathcal{X} and E , respectively. Moreover for any $\delta > 0$ it hold

$$\|J_\delta(y) - y\|_E \leq \delta(M + M\|y\|_E^m + \zeta_F\|y\|_E), \quad y \in E; \quad (4.3)$$

$$\|(F_\delta)_{|_E}(y)\|_E \leq (3 + \delta\zeta_F)\|F_{|_E}(y)\| + (2\zeta_F + \delta\zeta_F^2)\|y\|_E. \quad y \in E; \quad (4.4)$$

and

$$\|F_\delta(x_1) - F_\delta(x_2)\| \leq \left(\frac{2}{\delta} + \zeta_F\right) \|x_1 - x_2\|, \quad x_1, x_2 \in \mathcal{X}; \quad (4.5)$$

$$\|(F_\delta)_{|_E}(y_1) - (F_\delta)_{|_E}(y_2)\|_E \leq \left(\frac{2}{\delta} + \zeta_F\right) \|y_1 - y_2\|_E, \quad y_1, y_2 \in E. \quad (4.6)$$

Proof. By applying [7, Proposition 5.5.3] with $f = F - \zeta_F \text{Id}_\mathcal{X}$ and $f = F_{|_E} - \zeta_F \text{Id}_E$ we get immediately (4.2), the m -dissipativity of $F_\delta - \zeta_F \text{Id}_\mathcal{X}$ and $(F_\delta)_{|_E} - \zeta_F \text{Id}_E$ and estimates (4.5) and (4.6). Moreover [4, Proposition A.2.2(4)] yields that

$$\|J_\delta(y) - y\|_E \leq \delta \|F_{|_E}(y) - \zeta_F y\|_E, \quad y \in E, \delta > 0. \quad (4.7)$$

Estimate (4.3) thus follows by (2.6) and (4.7). Hence it remains to prove (4.4). By [7, Proposition 5.5.3(ii)] we can estimate $\|(F_\delta)_{|_E}(y) - \zeta_F J_\delta(y)\|_E \leq \|F_{|_E}(y) - \zeta_F y\|_E$ for any $y \in E$ and $\delta > 0$. Using this fact together with (4.7) we get

$$\begin{aligned} \|(F_\delta)_{|_E}(y)\|_E &\leq \|F_{|_E}(y)\|_E + \|(F_\delta)_{|_E}(y) - F_{|_E}(y)\|_E \\ &\leq \|F_{|_E}(y)\|_E + \|(F_\delta)_{|_E}(y) - \zeta_F J_\delta(y)\|_E + \|F_{|_E}(y) - \zeta_F y\|_E + \zeta_F \|J_\delta(y) - y\|_E \\ &\leq \|F_{|_E}(y)\|_E + 2\|F_{|_E}(y) - \zeta_F y\|_E + \delta \zeta_F \|F_{|_E}(y) - \zeta_F y\|_E \\ &\leq (3 + \delta \zeta_F) \|F_{|_E}(y)\|_E + (2\zeta_F + \delta \zeta_F^2) \|y\|_E \end{aligned}$$

for any $\delta > 0$ and $y \in E$. Hence (4.4) is proved. \square

Theorem 2.8, Remark 4.1 and Lemma 4.2 imply that for every $\delta \in (0, 1)$ and $x \in \mathcal{X}$ the problem

$$\begin{cases} dX_\delta(t) = [AX_\delta(t) + F_\delta(X_\delta(t))]dt + C^{1/2}dW(t), & t > 0; \\ X_\delta(0) = x \in \mathcal{X} \end{cases}$$

has a unique mild solution $\{X_\delta(t, x)\}_{t \geq 0}$ satisfying (see formula (4.4))

$$\|X_\delta(t, x)\|_E^p \leq C_p \left(e^{-\kappa_p t} \|x\|_E^p + \|W_A(t)\|_E^p + \int_0^t e^{-\kappa_p(t-s)} (\|F(W_A(s))\|_E^p + \|W_A(s)\|_E^p) ds \right), \quad (4.8)$$

for any $p \geq 1$, $t > 0$, $x \in E$ and some positive constants κ_p and $C_p = C_p(\zeta_F)$. Consequently the semigroup

$$P_\delta(t)\varphi(x) := \mathbb{E}[\varphi(X_\delta(t, x))]$$

is well defined for any $\varphi \in B_b(\mathcal{X})$. Moreover, (4.3) and (4.4) are crucial in order to prove, as the next proposition shows, that $P_\delta(t)$ approximates the semigroup $P(t)$ (see Theorem 2.8) as $\delta \rightarrow 0$.

Proposition 4.3. *If Hypotheses 2.6 hold true, then for any $T > 0$, $\varphi \in C_b(\mathcal{X})$ and $x \in E$*

$$\lim_{\delta \rightarrow 0} \sup_{t \in [0, T]} \|X_\delta(t, x) - X(t, x)\|_E = 0, \quad \mathbb{P}\text{-a.e.}; \quad (4.9)$$

$$\lim_{\delta \rightarrow 0} |P_\delta(t)\varphi(x) - P(t)\varphi(x)| = 0, \quad t > 0. \quad (4.10)$$

Proof. First of all let us observe that (4.10) immediately follows by (4.9). Therefore we just prove (4.9). To this aim we start pointing out that (4.8) implies that the function $(t, x) \mapsto \|X_\delta(t, x)\|_E$, as a map from $[0, T] \times E$ into \mathbb{R} , is bounded by a positive constant $C = C(T, x)$ independent of δ . This fact, together with estimates (4.1) and (4.3) implies that for any $\delta > 0$, $x \in E$ and $T > 0$

$$K(T, x) := \sup_{t \in [0, T]} (\|J_\delta(X_\delta(t, x))\|_E + \|X_\delta(t, x)\|_E + \|X(t, x)\|_E) < +\infty \quad (4.11)$$

and $K(T, x)$ is independent of δ . So by the local Lipschitzianity of F (Hypotheses 2.6(iii)) there exists $L := L(x, T) > 0$ such that

$$\begin{aligned} \|F_\delta(X_\delta(t, x)) - F(X(t, x))\|_E &= \|F(J_\delta(X_\delta(t, x))) - F(X(t, x))\|_E \\ &\leq L \|J_\delta(X_\delta(t, x)) - X(t, x)\|_E \\ &\leq L \|J_\delta(X_\delta(t, x)) - X_\delta(t, x)\|_E + L \|X_\delta(t, x) - X(t, x)\|_E. \end{aligned} \quad (4.12)$$

By (4.3), (4.11) and (4.12) we can conclude that

$$\|F_\delta(X_\delta(t, x)) - F(X(t, x))\|_E \leq \delta M' + L \|X_\delta(t, x) - X(t, x)\|_E. \quad (4.13)$$

for some positive $M' = M'(K, M, L, m, \zeta_F)$. Thus, by using the definition of mild solution and estimate (4.13) we obtain

$$\|X_\delta(t, x) - X(t, x)\|_E \leq \delta M_0 M' \int_0^t e^{(t-s)\eta_0} ds + M_0 L \int_0^t e^{(t-s)\eta_0} \|X_\delta(t, x) - X(t, x)\|_E ds.$$

Applying the Gronwall lemma we complete the proof. \square

As announced we need an additional assumption of dissipativity on F .

Hypotheses 4.4. *It holds that:*

- (i) $F(\text{Dom}(F) \cap H_C) \subseteq H_C$ and $F|_{H_C} - \zeta_F \text{Id}_{H_C}$ is m -dissipative (ζ_F is the constant appearing in Hypothesis 2.6(iii)).
- (ii) A and C are simultaneously diagonalizable and exists a basis $\{e_k\}_{k \in \mathbb{N}}$ of H_C consisting of eigenvectors of A and C .

Remark 4.5. We need to point out two observations that we will use in this section.

- (i) Arguing as in Lemma 4.2 it is easy to prove that under Hypotheses 2.6(iii) and 4.4(i), the function $(F|_{H_C})_\delta - \zeta_F \text{Id}_{H_C}$ is dissipative as a function from H_C to H_C for any $\delta \in (0, \bar{\zeta}_F)$.
- (ii) If Hypothesis 4.4(ii) holds true, then H_C is dense in \mathcal{X} and Hypotheses 3.9 are immediately satisfied with $\mathcal{X}_n := \text{span}\{e_1, \dots, e_n\}$.

Under Hypotheses 2.6, 4.4, for any $\delta \in (0, \bar{\zeta}_F)$ and $n \in \mathbb{N}$ we let $\{P_{\delta,n}(t)\}_{t \geq 0}$ be the transition semigroup associated to

$$\begin{cases} dX_{\delta,n}(t) = [A_n X_{\delta,n}(t) + (F_\delta)_n(X_{\delta,n}(t))]dt + C_n^{1/2} dW_n(t), & t > 0; \\ X_{\delta,n}(0) = z \in \mathcal{X}_n. \end{cases} \quad (4.14)$$

where A_n , C_n and $(F_\delta)_n$ are defined in Section 3.

Proposition 4.6. *Under Hypotheses 2.6 and 4.4 the following estimate*

$$\|\mathcal{D}_{C_n} P_{\delta,n}(t)f(x)\|_{C_n} \leq e^{t\zeta_X} P_{\delta,n}(t) \|\mathcal{D}_{C_n} f(x)\|_{C_n}, \quad x \in \mathcal{X}_n$$

holds true for any $\delta \in (0, \bar{\zeta}_F)$, $n \in \mathbb{N}$, $t > 0$, $f \in \mathcal{FC}_b^{1,n}(\mathcal{X})$ and $\zeta_X := \zeta_A + \zeta_F$.

Proof. We fix $n \in \mathbb{N}$, $\delta \in (0, \bar{\zeta}_F)$ and $x, y \in \mathcal{X}_n$ and let us consider $\{X_{n,\delta}(t, x)\}_{t \geq 0}$ and $\{X_{n,\delta}(t, y)\}_{t \geq 0}$ be the mild solutions of (4.14) with initial datum x and y respectively. We assume that $\{X_{n,\delta}(t, x)\}_{t \geq 0}$ and $\{X_{n,\delta}(t, y)\}_{t \geq 0}$ are strict solutions of (4.14), otherwise we proceed as in [2, Proposition 3.6] or [4, Proposition 6.2.2] approximating them by means of a sequence of more regular processes. For any $t \geq 0$ we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|X_{n,\delta}(t, x) - X_{n,\delta}(t, y)\|_{C_n}^2 &= \langle A(X_{n,\delta}(t, x) - X_{n,\delta}(t, y)), X_{n,\delta}(t, x) - X_{n,\delta}(t, y) \rangle_{C_n} \\ &\quad + \langle (F_\delta)_n(X_{n,\delta}(t, x)) - (F_\delta)_n(X_{n,\delta}(t, y)), X_{n,\delta}(t, x) - X_{n,\delta}(t, y) \rangle_{C_n}. \end{aligned}$$

Since \mathcal{X}_n is a finite dimensional space and the operators A_n and C_n are simultaneously diagonalizable, they commute. Thus, thanks to Hypotheses 2.6 and 4.4(i) (see also Remark 4.5(i)), for any $t \geq 0$ we obtain

$$\frac{d}{dt} \|X_{n,\delta}(t, x) - X_{n,\delta}(t, y)\|_{C_n}^2 \leq -2\zeta_X \|X_{n,\delta}(t, x) - X_{n,\delta}(t, y)\|_{C_n}^2,$$

where $\zeta_X = \zeta_A + \zeta_F$. By the Gronwall inequality, for any $t \geq 0$ and $x, y \in \mathcal{X}_n$ we obtain

$$\|X_{n,\delta}(t, x) - X_{n,\delta}(t, y)\|_{C_n} \leq e^{-t\zeta_X} \|x - y\|_{C_n}.$$

Therefore, for any $f \in \mathcal{FC}_b^{1,n}(\mathcal{X})$, $t \geq 0$ and $x \in \mathcal{X}_n$ we have

$$\begin{aligned} \|\mathcal{D}_{C_n} P_{\delta,n}(t)f(x)\|_{C_n} &= \lim_{y \rightarrow x} \frac{|P_{\delta,n}(t)f(x) - P_{\delta,n}(t)f(y)|}{\|x - y\|_{C_n}} \\ &= \lim_{y \rightarrow x} \left(\frac{|P_{\delta,n}(t)f(x) - P_{\delta,n}(t)f(y)|}{\|X(t, x) - Y(t, y)\|_{C_n}} \right) \left(\frac{\|X(t, x) - Y(t, y)\|_{C_n}}{\|x - y\|_{C_n}} \right) \end{aligned}$$

$$\begin{aligned}
&= \lim_{y \rightarrow x} \left(\frac{|\mathbb{E}(f(X_{\delta,n}(t, x))) - \mathbb{E}(f(X_{\delta,n}(t, y)))|}{\|X(t, x) - Y(t, y)\|_{C_n}} \right) \left(\frac{\|X(t, x) - Y(t, y)\|_{C_n}}{\|x - y\|_{C_n}} \right) \\
&\leq e^{-t\zeta_x} P_{\delta,n}(t) \|\mathcal{D}_{C_n} f(x)\|_{C_n}
\end{aligned}$$

whence the claim. \square

Theorem 4.7. *Assume that Hypotheses 2.6 and 4.4 hold true. For any $\varphi \in B_b(\mathcal{X})$, $t > 0$, $x \in \mathcal{X}$ and $p > 1$ it holds*

$$|P(t)\varphi(x+h)|^p \leq P(t)|\varphi(x)|^p \exp\left(\frac{pe^{2t\zeta_x}}{t(p-1)}\|h\|_C^2\right), \quad h \in H_C \cap E. \quad (4.15)$$

Proof. Using Proposition 4.6 and arguing exactly as in the proof of Theorem 3.12 we obtain

$$|P_\delta(t)\varphi(x+h)|^p \leq P_\delta(t)|\varphi(x)|^p \exp\left(\frac{pe^{2t\zeta_x}}{t(p-1)}\|h\|_C^2\right). \quad (4.16)$$

for any $\delta \in (0, \bar{\zeta}_F)$, $p > 1$, $t > 0$, $\varphi \in C_b(\mathcal{X})$, $x \in X$ and $h \in H_C$. Now, using (4.10) and letting $\delta \rightarrow 0$ in (4.16) we get that

$$|P(t)\varphi(x+h)|^p \leq P(t)|\varphi(x)|^p \exp\left(\frac{pe^{2t\zeta_x}}{t(p-1)}\|h\|_C^2\right). \quad (4.17)$$

for any $p > 1$, $t > 0$, $\varphi \in C_b(\mathcal{X})$, $x \in E$ and $h \in H_C \cap E$. Using the fact that E is densely embedded in \mathcal{X} and the continuity of $P(t)\varphi$ we can extend estimate (4.17) for any $x \in \mathcal{X}$. Finally, using the monotone class theorem as in the proof of Theorem 3.12 we complete the proof. \square

Remark 4.8. We point out that if $H_C \cap E$ is dense in H_C , then (4.15) will hold true for any $h \in H_C$.

5. SOME CONSEQUENCES AND EXAMPLES

In the first part of this section we collect some corollaries of Theorems 3.12 and 4.7. Finally, we exhibit some classes of examples to which all of our results can be applied.

We start by stating and proving some classical consequences of the (LHI) for which we refer to [38, Corollary 1.2] and [43, Section 1.3.1]. In the sequel μ will denote an invariant measure associated with $P(t)$, that is a Borel probability measure on \mathcal{X} such that

$$\int_{\mathcal{X}} P(t)f d\mu = \int_{\mathcal{X}} f d\mu, \quad t > 0, f \in B_b(\mathcal{X}).$$

Sufficient conditions that guarantee the existence of such a measure can be found in [4, Chapter 8], [8, Chapter 6] and [8, Chapter 11]. In this case $P(t)$ extends to a positive contractive semigroup in $L^p(\mathcal{X}, \mu) =: L^p_\mu$, the L^p spaces related to the measure μ , (see [33, Theorem 2.14]). For simplicity we write $\mu(f)$ to denote $\int_{\mathcal{X}} f d\mu$.

Corollary 5.1. *Assume that Hypotheses 2.6 and 4.4 hold true.*

(i) *For any positive $f \in B_b(\mathcal{X})$, $t > 0$, $x \in \mathcal{X}$ and $h \in H_C \cap E$*

$$[P(t)(\ln f)](x+h) \leq \ln P(t)f(x) + \frac{e^{2t\zeta_x}}{t}\|h\|_C^2. \quad (5.1)$$

(ii) *For every $f \in B_b(\mathcal{X})$ and $x \in \mathcal{X}$ it holds*

$$\lim_{\substack{\|h\|_C \rightarrow 0 \\ h \in H_C \cap E}} P(t)f(x+h) = P(t)f(x). \quad (5.2)$$

(iii) *The following entropy-cost inequality holds true*

$$\mu((P^*(t)f) \ln(P^*(t)f)) \leq \frac{e^{2t\zeta_x}}{t} W(f\mu, \mu)^2,$$

for any positive function $f \in L^2_\mu$ such that $\mu(f) = 1$. Here $\{P^*(t)\}_{t \geq 0}$ is the adjoint semi-group of $\{P(t)\}_{t \geq 0}$ in L^2_μ and W denotes the L^2 -Wasserstein distance with respect to the cost function $(x, y) \mapsto \|x - y\|_C$, namely for any two probability measure μ_1, μ_2 on \mathcal{X}

$$W(\mu_1, \mu_2)^2 := \inf \left\{ \int_{\mathcal{X} \times \mathcal{X}} \|x - y\|_C^2 \pi(dx, dy) \mid \pi \in \mathcal{C}(\mu_1, \mu_2) \right\},$$

where $\mathcal{C}(\mu_1, \mu_2)$ is the set of all the couplings of μ_1 and μ_2 and we let $\|x - y\|_C = +\infty$, if $x - y$ does not belong to $H_C \cap E$.

Proof. A proof of (i) can be found in [43, Section 1.3.1]. We now prove (ii). It suffices to prove (5.2) for a non-negative function $f \in B_b(\mathcal{X})$. Indeed the general case can be obtained writing $f = f^+ - f^-$, being f^+ and f^- the positive and the negative part of f . So, let us fix a non-negative function f and for any $\varepsilon > 0$ we set $f_\varepsilon := 1 + \varepsilon f$. Recalling that $r \leq \ln(1 + r) + r^2$ for any $r \geq 0$ we get for every $x \in \mathcal{X}$

$$\ln f_\varepsilon(x) = \ln(1 + \varepsilon f(x)) \geq \varepsilon f(x) - \varepsilon^2 f^2(x) \geq \varepsilon f(x) - \varepsilon^2 \|f\|_\infty^2. \quad (5.3)$$

Now applying (5.1) to f_ε , using (5.3) and dividing by ε we get for every $x \in \mathcal{X}$ and $h \in H_C \cap E$

$$P(t)f(x + h) - \varepsilon \|f\|_\infty^2 \leq \frac{1}{\varepsilon} \ln P(t)(1 + \varepsilon f(x)) + \frac{e^{2t\zeta x}}{\varepsilon t} \|h\|_C^2. \quad (5.4)$$

Taking the supremum limit as $\|h\|_C \rightarrow 0$ with $h \in H_C \cap E$ and then letting $\varepsilon \rightarrow 0$ we get

$$\limsup_{\substack{\|h\|_C \rightarrow 0 \\ h \in H_C \cap E}} P(t)f(x + h) \leq P(t)f(x).$$

Recalling that $\ln(1 + r) \leq r$ for any $r > -1$ and arguing as above we get that for any $\varepsilon > 0$, $x \in \mathcal{X}$ and $h \in H_C \cap E$

$$P(t) \left(\frac{1 + \varepsilon f(x)}{\varepsilon} \right) - \frac{e^{2t\zeta x}}{\varepsilon t} \|h\|_C^2 \leq \frac{1}{\varepsilon} \ln P(t)(1 + \varepsilon f(x - h)) \leq P(t)f(x - h).$$

Taking the infimum limit as $\|h\|_C \rightarrow 0$ with $h \in H_C \cap E$ and then letting $\varepsilon \rightarrow 0$ we get

$$P(t)f(x) \leq \liminf_{\substack{\|h\|_C \rightarrow 0 \\ h \in H_C \cap E}} P(t)f(x - h). \quad (5.5)$$

Since $H_C \cap E$ is a linear space then applying (5.5) to $-h$ we get

$$P(t)f(x) \leq \liminf_{\substack{\|h\|_C \rightarrow 0 \\ h \in H_C \cap E}} P(t)f(x + h). \quad (5.6)$$

By (5.4) and (5.6) we get (5.2).

Now a standard argument allows us to prove (iii) for a bounded Borel and positive function f with $\mu(f) = 1$. Writing (5.1) with $P^*(t)f$ in place of f , we get

$$[(P(t)f)(\ln P^*(t)f)](x) \leq \ln(P(t)P^*(t)f(y)) + \frac{e^{2t\zeta x}}{t} \|x - y\|_C^2 \quad (5.7)$$

for any $t > 0$, $x, y \in \mathcal{X}$ such that $x - y \in H_C \cap E$. Integrating both sides of (5.7) with respect to $\pi \in \mathcal{C}(f\mu, \mu)$ we get

$$\mu((P^*(t)f)(\ln P^*(t)f)) \leq \mu(\ln P(t)P^*(t)f) + \frac{e^{2t\zeta x}}{t} \int_{\mathcal{X} \times \mathcal{X}} \|x - y\|_C^2 \pi(dx, dy).$$

To conclude it is sufficient to observe that the Jensen inequality yields that

$$\mu(\ln P(t)P^*(t)f) \leq \ln \mu(P(t)P^*(t)f) = \ln \mu(f) = 0,$$

whence the claim. \square

Remark 5.2. We stress that Corollary 5.1 remains true (with the constant $e^{2t\zeta x}$ replaced by K_1^2) if we assume that the hypotheses of Theorem 3.12 hold true.

Another classical consequence of (LHI) is a hypercontractivity type estimate for the semigroup $P(t)$ in L_μ^p . Such estimate relies on the Hölder inequality and some integrability conditions with respect to μ of some exponential functions.

Corollary 5.3. *Assume that the hypotheses of Theorem 3.12 hold true and that there exists an invariant measure μ for the semigroup $P(t)$. If, in addition there exists $\varepsilon > 0$ such that*

$$\int_{\mathcal{X}} \int_{\mathcal{X}} e^{\varepsilon \|x-y\|_C^2} \mu(dx) \mu(dy) < +\infty, \quad (5.8)$$

then, for any $p \geq 2$, there exists $t_0 > 0$ and a positive constant C such that

$$\|P(t)f\|_{L_\mu^p} \leq C \|f\|_{L_\mu^2} \quad (5.9)$$

for any $t \geq t_0$ and any $f \in L_\mu^2$.

Proof. Let us consider $f \in L_\mu^2$ and $\vartheta \in (1, 2)$. By (3.14) we deduce that for any $t > 0$

$$\begin{aligned} \int_{\mathcal{X}} |P(t)f|^{2\vartheta}(x) \mu(dx) &= \int_{\mathcal{X}} \int_{\mathcal{X}} |P(t)f|^{2\vartheta}(x) \mu(dx) \mu(dy) \\ &= \int_{\mathcal{X}} \int_{\mathcal{X}} |P(t)f(x)|^\vartheta (|P(t)f(x)|^2)^{\vartheta/2} \mu(dx) \mu(dy) \\ &\leq \int_{\mathcal{X}} \int_{\mathcal{X}} |P(t)f(x)|^\vartheta (P(t)f^2(y))^{\vartheta/2} e^{\frac{\vartheta K^2}{t} \|x-y\|_C^2} \mu(dx) \mu(dy) \\ &=: \int_{\mathcal{X}} \int_{\mathcal{X}} h(x, y) g(x, y) \mu(dx) \mu(dy), \end{aligned}$$

where $h(x, y) := |P(t)f(x)|^\vartheta (P(t)f^2(y))^{\vartheta/2}$ and $g(x, y) := e^{\frac{\vartheta K^2}{t} \|x-y\|_C^2}$. Applying the Hölder inequality with respect to the measure $\mu \otimes \mu$ we get

$$\int_{\mathcal{X}} \int_{\mathcal{X}} h(x, y) g(x, y) \mu(dx) \mu(dy) \leq \|h\|_{L_{\mu \otimes \mu}^{2/\vartheta}} \|g\|_{L_{\mu \otimes \mu}^{2/(2-\vartheta)}}.$$

Now, the invariance of μ and the contractivity of $P(t)$ in L_μ^2 allow us to estimate

$$\|h\|_{L_{\mu \otimes \mu}^{2/\vartheta}} = \|P(t)f\|_{L_\mu^2}^\vartheta \|f\|_{L_\mu^2}^\vartheta \leq \|f\|_{L_\mu^2}^{2\vartheta}.$$

Moreover, being

$$\|g\|_{L_{\mu \otimes \mu}^{2/(2-\vartheta)}} = \int_{\mathcal{X}} \int_{\mathcal{X}} e^{\frac{2\vartheta}{2-\vartheta} \frac{K^2}{t} \|x-y\|_C^2} \mu(dx) \mu(dy) =: C(\vartheta, t)$$

condition (5.8) ensures that there exists $\bar{t} > 0$ such that $C(\vartheta, t) < +\infty$ for any $t \geq \bar{t}$ and any $\vartheta \in (1, 2)$. Consequently

$$\|P(t)f\|_{L_\mu^{2\vartheta}} \leq (C(\vartheta, t))^{\frac{2-\vartheta}{4\vartheta}} \|f\|_{L_\mu^2}.$$

i.e., $P(t)$ maps L_μ^2 into $L_\mu^{2\vartheta}$ for $t \geq \bar{t}$. Since $\vartheta > 1$, $P(t)$ actually improves summability of the initial datum when $t \geq \bar{t}$. To go further we use the semigroup law. Indeed, if $f \in L_\mu^2$, then $P(\bar{t})f \in L_\mu^{2\vartheta}$, i.e. $|P(\bar{t})f|^\vartheta \in L_\mu^2$. Using again the first part of the proof, we deduce that $P(t)|P(\bar{t})f|^\vartheta \in L_\mu^{2\vartheta}$ for $t \geq \bar{t}$. Since, by the Jensen inequality and the positivity of $P(t)$ we can estimate

$$\begin{aligned} +\infty &> \|P(t)|P(\bar{t})f|^\vartheta\|_{L_\mu^{2\vartheta}}^{2\vartheta} = \int_{\mathcal{X}} |P(t)|P(\bar{t})f|^{2\vartheta} d\mu \\ &\geq \int_{\mathcal{X}} |P(t)|P(\bar{t})f|^{2\vartheta^2} d\mu \\ &= \int_{\mathcal{X}} (P(t)|P(\bar{t})f|)^{2\vartheta^2} d\mu \\ &\geq \int_{\mathcal{X}} |P(t)P(\bar{t})f|^{2\vartheta^2} d\mu \end{aligned}$$

$$= \int_{\mathcal{X}} |P(t + \bar{t})f|^{2\vartheta^2} d\mu, \quad t \geq \bar{t}$$

we infer that $P(t)$ maps L_μ^2 into $L_\mu^{2\vartheta^2}$ for any $t \geq 2\bar{t}$. Iterating this procedure we can prove that for any $p > 2$ there exists $t_0 = t_0(p) > 0$ such that $P(t)$ maps L_μ^2 into L_μ^p for any $t \geq t_0$ and estimate (5.9) holds true. \square

Remark 5.4. Note that the result in Corollary 5.3 and (5.9) continue to hold true if we assume the hypotheses of Theorem 4.7, that $\zeta_{\mathcal{X}} < 0$ and that (5.8) is satisfied with $\varepsilon = 1$.

Now we collect some examples of operators to which the results of Sections 3 and 4 and of Corollaries 5.1 and 5.3 can be applied.

Example 5.5. A suitable choice for A and C .

Let \mathcal{X} be a Hilbert space. For a positive and compact operator $Q \in \mathcal{L}(\mathcal{X})$, we set

$$A := -(1/2)Q^{-\beta} : Q^\beta(\mathcal{X}) \subseteq \mathcal{X} \rightarrow \mathcal{X}, \quad C := Q^{2\alpha},$$

with $\alpha, \beta \geq 0$ such that (2.7) is verified.

Let $\{e_k\}_{k \in \mathbb{N}}$ be an orthonormal basis of \mathcal{X} consisting of eigenvectors of Q , and let $\{\lambda_k\}_{k \in \mathbb{N}}$ be the eigenvalues associated with $\{e_k\}_{k \in \mathbb{N}}$. Since Q is a compact and positive operator, there exists $k_0 \in \mathbb{N}$ such that $0 < \lambda_k \leq \lambda_{k_0}$, for any $k \in \mathbb{N}$. Without loss of generality we can assume that $k_0 = 1$. Hence, for any $x \in Q^\beta(\mathcal{X})$

$$\langle Ax, x \rangle = -\frac{1}{2} \sum_{k=1}^{+\infty} \lambda_k^{-\beta} \langle x, e_k \rangle^2 \leq -\frac{1}{2} \lambda_1^{-\beta} \|x\|^2. \quad (5.10)$$

Moreover, by the properties of Q , $\text{Dom}(A) = Q^\beta(\mathcal{X})$ is dense in \mathcal{X} , so A generates a strongly continuous and contraction semigroup in \mathcal{X} and, so Hypotheses 2.6 are satisfied. Further, let us consider A_C , the part of A in $H_C = H_{Q^\alpha}$ and recall that

$$\text{Dom}(A_C) := \{x \in Q^\alpha(\mathcal{X}) \cap Q^\beta(\mathcal{X}) \mid Ax \in Q^\alpha(\mathcal{X})\}.$$

By (5.10), for any $x \in \text{Dom}(A_C)$, we have

$$[Ax, x]_C = \langle Q^{-\alpha} Ax, Q^{-\alpha} x \rangle = \langle AQ^{-\alpha} x, Q^{-\alpha} x \rangle \leq -\frac{\|Q^{-\alpha} x\|^2}{2\lambda_1^\beta} = -\frac{\|x\|_C^2}{2\lambda_1^\beta}. \quad (5.11)$$

Since $Q^{\alpha+\beta}(\mathcal{X})$ is dense in \mathcal{X} and $Q^{-\alpha}$ is a close operator in \mathcal{X} , then $Q^{\alpha+\beta}(\mathcal{X})$ is dense in H_C , moreover $Q^{\alpha+\beta}(\mathcal{X}) \subseteq \text{Dom}(A_C)$. Hence A generates a strongly continuous and contraction semigroup in H_C . By the compactness and positivity of Q , we get that H_C is dense in \mathcal{X} . Moreover Hypotheses 3.9 is verified with $\mathcal{X}_n := \text{span}\{e_1, \dots, e_n\}$. Let G be a Lipschitz continuous function, it easy to see that $F = Q^\alpha G$ is Lipschitz continuous too with Lipschitz constant $\|Q^\alpha\|_{\mathcal{L}(\mathcal{X})} L_G$. Moreover due to the choice of Q we have $\|Q^\alpha\|_{L_G} \leq \lambda_1^\alpha L_G$ and so

$$\langle Q^\alpha \mathcal{D}G(x)h, h \rangle \leq \lambda_1^\alpha L_G \|h\|^2, \quad x, h \in \mathcal{X}.$$

If we assume $L_G < (2\lambda_1^{\alpha+\beta})^{-1}$ then condition (3.1) is verified. So Hypotheses 3.4(ii) and Hypotheses 3.9 are satisfied too. If, in addition Hypotheses 3.4(i) are satisfied, then Theorem 3.12 can be applied. This is the case when, for instance, $(A, \text{Dom}(A))$ is a sectorial operator in \mathcal{X} as the next example shows.

Example 5.6. An example in $L^2([0, 1], \lambda)$.

In the previous example one can take as \mathcal{X} the Hilbert space $L^2([0, 1], \lambda)$ where λ is the Lebesgue measure and by $-Q^{-1}$ the realization in $L^2([0, 1], \lambda)$ of the second order derivative with Dirichlet boundary conditions. With these choices Q turns out to be a positive and trace class operator in \mathcal{X} . Thus, if A and C are defined as above, then estimate (5.10) and (5.11) hold true with $\lambda_1 = \pi^{-2}$ and A generates a strongly continuous and analytic semigroup e^{tA} satisfying

$$\|e^{tA}\|_{\mathcal{L}(\mathcal{X})} \leq e^{-(1/2)\pi^2 t}, \quad t \geq 0$$

see [8, Chapter 4]. Moreover by [26, Proposition 2.1.1], if $\alpha < \beta$ then (3.6) is verified with $\gamma = \alpha/\beta$. If, in addition F is a Lipschitz continuous function with Lipschitz constant $L_F < (1/2)\pi^{2\beta}$, then (3.1) is verified. In particular Hypotheses 3.4(i) and Hypotheses 3.9 are verified. Choosing $\alpha = 1/2 < \beta$, then H_C is the space $W_0^{1,2}([0, 1], \lambda)$ and (2.7) is satisfied too and Theorem 3.12 can be applied.

Example 5.7. Infinite dimensional polynomial.

In this example we exhibit a class of functions satisfying Hypotheses 4.4. To this aim we recall the notion of infinite dimensional polynomial (see [5, 13, 31]). For every $n \in \mathbb{N}$, we say that a map $V : \mathcal{X}^n \rightarrow \mathcal{X}$ is n -multilinear if it is linear in each variable separately. A n -multilinear map V is said to be symmetric if

$$V(x_1, \dots, x_n) = V(x_{\sigma(1)}, \dots, x_{\sigma(n)}), \quad (5.12)$$

for any permutation σ of the index set $\{1, \dots, n\}$. We say that a function $P_n : \mathcal{X} \rightarrow \mathcal{X}$ is a homogeneous polynomial of degree $n \in \mathbb{N}$ if there exists a n -multilinear symmetric map V such that for every $x \in \mathcal{X}$

$$P_n(x) = V(x, \dots, x).$$

We consider the function $F : \mathcal{X} \rightarrow \mathcal{X}$ defined by

$$F(x) := P_n(x) + \zeta_F x,$$

where $x \in \mathcal{X}$, $\zeta_F \in \mathbb{R}$ and P_n is a homogeneous polynomial of degree n such that,

$$\langle V(h, x, \dots, x), h \rangle \leq 0, \quad (5.13)$$

for any $h, x \in \mathcal{X}$. By [5, Theorem 3.4], there exists $d > 0$ such that

$$\|F(x)\| \leq d(1 + \|x\|^n), \quad x \in \mathcal{X}. \quad (5.14)$$

Moreover, for any $x, h \in \mathcal{X}$, we have

$$\mathcal{D}P_n(x)h = nV(h, x, \dots, x),$$

and so, by (5.13), for any $x, y \in \mathcal{X}$, we obtain

$$\langle F(x) - F(y), x - y \rangle \leq \zeta_F \|x - y\|^2. \quad (5.15)$$

In order to give a more concrete application we place ourself in the same scenario as in Example 5.6 with $\alpha = 1/2 < \beta$. Consider a version of $K \in L^2([0, 1]^4, \lambda)$ which is symmetric (see (5.12)) and set

$$[P_3(f)](\xi) := \int_0^1 \int_0^1 \int_0^1 K(\xi_1, \xi_2, \xi_3, \xi) f(\xi_1) f(\xi_2) f(\xi_3) d\xi_1 d\xi_2 d\xi_3 \quad (5.16)$$

with $f \in L^2([0, 1])$. P_3 is a homogeneous polynomial of degree three on $L^2([0, 1], \lambda)$ (see [13, Exercise 1.73]). Moreover, condition (5.13) holds true whenever K has negative values. Indeed observe that, for $f_1, f_2, f_3 \in L^2([0, 1], \lambda)$,

$$V(f_1, f_2, f_3) = \int_0^1 \int_0^1 \int_0^1 K(\xi_1, \xi_2, \xi_3, \xi) f_1(\xi_1) f_2(\xi_2) f_3(\xi_3) d\xi_1 d\xi_2 d\xi_3,$$

and for $f, h \in L^2([0, 1], \lambda)$

$$\langle V(h, f, f), h \rangle = \int_0^1 \int_0^1 \int_0^1 \int_0^1 K(\xi_1, \xi_2, \xi_3, \xi) f(\xi_1) f(\xi_2) h(\xi_3) h(\xi) d\xi_1 d\xi_2 d\xi_3 d\xi.$$

A standard argument allows to deduce that $\langle V(h, f, f), h \rangle = 0$ if, and only if, $f = 0$ λ -a.e. or $h = 0$ λ -a.e. So by the continuity of $\langle V(h, f, f), h \rangle$ with respect to h (for a fixed f) and the fact that

$$\langle V(-h, f, f), -h \rangle = \langle V(h, f, f), h \rangle,$$

the claim follows. In addition, if we assume that K has weak derivative with respect to the fourth variable belonging to $L^2([0, 1]^4, \lambda)$, then for any $f \in W^{1,2}([0, 1], \lambda)$ it holds that $F(f) = P_3(f) + \zeta_F f$ belongs to $W^{1,2}([0, 1], \lambda)$ (see (5.16)) and its weak derivative is

$$[F(f)]'(\xi) = \int_0^1 \int_0^1 \int_0^1 \frac{\partial K}{\partial \xi}(\xi_1, \xi_2, \xi_3, \xi) f(\xi_1) f(\xi_2) f(\xi_3) d\xi_1 d\xi_2 d\xi_3 + \zeta_F f'(\xi). \quad (5.17)$$

If we assume that $(\partial K / \partial \xi) \in L^2([0, 1]^4, \lambda)$ has a symmetric version (see (5.12)) with negative value, then by (5.17), estimate (5.14) and (5.15) are verified in $W^{1,2}([0, 1], \lambda)$. Hence Hypotheses 4.4(i) is verified and with the choices of A and C in the Example 5.6, all the results in Section 4 can be applied.

Example 5.8. A reaction-diffusion system.

Assume that $\mathcal{X} = L^2([0, 1], \lambda)$ (where λ is the Lebesgue measure), $E = C([0, 1])$, A is the realization in $L^2([0, 1], \lambda)$ of the second order derivative operator with Dirichlet boundary condition and $C = \text{Id}_{\mathcal{X}}$. In order to define the function F we consider a decreasing function $\varphi \in C^1(\mathbb{R})$ such that

$$|\varphi'(\xi)| \leq d_1(1 + |\xi|^m), \quad \xi \in \mathbb{R},$$

for some constants $d_1 > 0$ and $m \in \mathbb{N}$. Let $\zeta_F > 0$. We set

$$[F(f)](\xi) = \begin{cases} \varphi(f(\xi)) - \frac{\zeta_F}{2} f(\xi)^2, & f \in C([0, 1]), \xi \in [0, 1]; \\ 0, & \text{otherwise.} \end{cases}$$

By [4, Section 6.1, Lemma 6.1.2 and Lemma 8.2.1] and [8, Example D.7] it follows that Hypotheses 4.4 are verified, in particular (3.1) is satisfied. Finally, taking into account the results in Examples 5.5 and 5.6 (when $\alpha = 0$ and $\beta = 1$), we can conclude that Hypotheses 3.4(i) and Hypotheses 3.9 are verified too and so Theorem 4.7 can be applied.

REFERENCES

- [1] Aronson, D. G., Serrin, J., Local behaviour of solutions of quasi-linear parabolic equations, *Arch. Rat. Mech. Anal.* **25** (1967), 81–123.
- [2] Bignamini, D.A., L^2 -theory for transition semigroups associated to dissipative systems., *Arxiv e-prints*, 2021. **arXiv**: 2110.05271.
- [3] Bignamini, D.A., Ferrari, S., Regularizing Properties of (Non-Gaussian) Transition Semigroups in Hilbert Spaces, *Potential Anal.* (2021), **DOI**:10.1007/s11118-021-09931-2.
- [4] Cerrai, S., *Second order PDE's in finite and infinite dimension*, Lecture Notes in Mathematics, vol. 1762, Springer-Verlag, Berlin, 2001.
- [5] Chae, S.B. *Holomorphy and Calculus in Normed Spaces*, Chapman & Hall/CRC Pure and Applied Mathematics, vol. 92, Taylor & Francis Group, New York and Basel, 1985.
- [6] Da Prato, G., and Röckner, M., Wang, F.-Y., Singular stochastic equations on Hilbert spaces: Harnack inequalities for their transition semigroups, *J. Funct. Anal.* **257** (2009), 992–1017.
- [7] Da Prato, G., Zabczyk, J., *Ergodicity for Infinite-Dimensional Systems*, London Mathematical Society Lecture Note Series, vol. 229, Cambridge University Press, Cambridge, 1996.
- [8] Da Prato, G., Zabczyk, J., *Stochastic equations in infinite dimensions*, Encyclopedia of Mathematics and its Applications, vol. 152, Second edition, Cambridge University Press, Cambridge, 2014.
- [9] Deng, C.S., Zhang, S.Q., Log-Harnack inequalities for Markov semigroups generated by non-local Grushin type operators, *Math. Nachr.* **291** (2018), 1055–1074.
- [10] De Giorgi, E., Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari, *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat.* **3** (1957), 25–43.
- [11] DiBenedetto, E., Gianazza, U., Vespri, V., Forward, backward and elliptic harnack inequalities for non-negative solutions to certain singular parabolic partial differential equations, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **9** (2010) 385–422.
- [12] DiBenedetto, E., Gianazza, U., Vespri, V., Harnack estimates for quasi-linear degenerate parabolic differential equations, *Acta Math.* **200** (2008), 181–209.
- [13] Dineen, S., *Complex Analysis on Infinite Dimensional Spaces*, Springer Monographs in Mathematics, Springer-Verlag, London, 1999.
- [14] Dunford, N., Schwartz, J.T., *Linear operators. Part I*, Wiley Classics Library, John Wiley & Sons, Inc., New York, 1988.
- [15] Dunford, N., Schwartz, J.T., *Linear operators. Part II*, Wiley Classics Library, John Wiley & Sons, Inc., New York, 1988.

- [16] Engel, K.-J., Nagel, R., *A short course on operator semigroups*, Universitext, Springer, New York, 2006.
- [17] Es-Sarhir, A., von Renesse, M.-K., Scheutzow, M., Harnack inequality for functional SDEs with bounded memory, *Electron. Commun. Probab.* **14** (2009), 560–565.
- [18] Goldys, B., Kocan, M., Diffusion semigroups in spaces of continuous functions with mixed topology, *J. Differential Equations* **173** (2001), 17–39.
- [19] Gordina, M., Röckner, M., Wang, F.-Y., Dimension-independent Harnack inequalities for subordinated semigroups, *Potential Anal.* **34** (2011), 293–307.
- [20] Hadamard, J., Extension à l'équation de la chaleur d'un théorème de A. Harnack, *Rend. Circ. Mat. Palermo* **2** (1954), 337–346.
- [21] Harnack, A., *Die Grundlagen der Theorie des logarithmischen Potentials und der eindeutigen Potentialfunktion in der Ebene*, V. G. Teubner, Leipzig, 1887.
- [22] Huang, X., Zhang, S.Q., Mild solutions and Harnack inequality for functional stochastic partial differential equations with Dini drift, *J. Theoret. Probab.* **32** (2019), 303–329.
- [23] Jacod, J., Protter, P., *Probability essentials*, Universitext, Springer-Verlag, Berlin, 2003.
- [24] Kassmann, M., *Harnack inequalities: an introduction*, Bound. Value Probl. (2007), Art. ID 81415, 21.
- [25] Lorenzi, L., *Analytical Methods for Markov Semigroups*, CRC Press, second ed., 2017.
- [26] Lunardi, A., *Analytic semigroups and optimal regularity in parabolic problems*, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 1995.
- [27] Lv, W., Huang, X., Harnack and shift Harnack inequalities for degenerate (functional) stochastic partial differential equations with singular drifts, *J. Theoret. Probab.* **34** (2021), 827–851.
- [28] Masiero, F., Stochastic optimal control problems and parabolic equations in Banach spaces, *SIAM J. Control Optim.* **47** (2008), 251–300.
- [29] Moser, J., On Harnack's theorem for elliptic differential equations, *Comm. Pure Appl. Math.* **14** (1961), 577–591.
- [30] Moser, J., A Harnack inequality for parabolic differential equations, *Comm. Pure Appl. Math.* **17** (1964), 101–134.
- [31] Mujica, J., *Complex Analysis in Banach Spaces*, North-Holland Mathematics Studies, vol. 120, North-Holland, Amsterdam, 1985.
- [32] Nash, J., Parabolic equations, *Proc. Nat. Acad. Sci. U.S.A.* **43** (1957), 754–758.
- [33] Ouhabaz, E.M., *Analysis of heat equations on domains*, London Mathematical Society Monographs Series, vol. 31, Princeton University Press, Princeton, NJ, 2005.
- [34] Phelps, R.R., *Convex Functions, Monotone Operators and Differentiability*, Lecture Notes in Mathematics, vol. 1364, Springer-Verlag, Berlin Heidelberg, 1993.
- [35] Pini, B., Sulla soluzione generalizzata di Wiener per il primo problema di valori al contorno nel caso parabolico, *Rend. Sem. Mat. Univ. Padova* **23** (1954), 422–434.
- [36] Reed, M., Simon, B., *Methods of modern mathematical physics. I. Functional analysis*, Academic Press, New York-London, 1972.
- [37] Röckner, M., Wang, F.-Y., Harnack and functional inequalities for generalized Mehler semigroups, *J. Funct. Anal.* **203** (2003), 237–261.
- [38] Röckner, M., Wang, F.-Y., Log-Harnack inequality for stochastic differential equations in Hilbert spaces and its consequences, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **13** (2010), 27–37.
- [39] Serrin, J., On the Harnack inequality for linear elliptic equations, *J. Analyse Math.* **4** (1955/56), 292–308.
- [40] Trudinger, N.S., On Harnack type inequalities and their application to quasilinear elliptic equations, *Commun. Pure Appl. Math.* **20** (1967), 721–747.
- [41] Wang, F.-Y., Logarithmic Sobolev inequalities on noncompact Riemannian manifolds, *Probab. Theory Related Fields* **109** (1997), 417–424.
- [42] Wang, F.-Y., Equivalence of dimension-free Harnack inequality and curvature condition, *Integral Equations Operator Theory* **48** (2004), 547–552.
- [43] Wang, F.-Y., *Harnack inequalities for stochastic partial differential equations*, SpringerBriefs in Mathematics, Springer, New York, 2013.
- [44] Wang, F.-Y., Yuan, C., Harnack inequalities for functional SDEs with multiplicative noise and applications, *Stochastic Process. Appl.* **121** (2011), 2692–2710.

L.A. & S.F.: DIPARTIMENTO DI MATEMATICA E FISICA “ENNIO DE GIORGI”, UNIVERSITÀ DEL SALENTO, VIA PER ARNESANO, I-73100 LECCE, ITALY

D.A.B.: DIPARTIMENTO DI SCIENZE MATEMATICHE, FISICHE E INFORMATICHE, PLESSO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI PARMA, PARCO AREA DELLE SCIENZE 53/A, I-43124 PARMA, ITALY

Email address: luciana.angiuli@unisalento.it

Email address: davideaugusto.bignamini@unimore.it

Email address: simone.ferrari@unisalento.it