arXiv:2111.13407v2 [math.AP] 29 Nov 2021

Preprint to...

On solvability of the non-local problem for the fractional mixed-type equation with Bessel operator

Bakhodirjon Toshtemirov
V.I.Romanovskiy Institute of Mathematics, Uzbekistan and
Ghent University, Belgium.
bakhodirjon.toshtemirov@ugent.be

Abstract. *The non-local problem is considered for the partial differential equation of mixed-type with Bessel operator and fractional order. Explicit solution is represented by Fourier-Bessel series in the given domain. It is established the connection between the given data and the uniquely solvability of the problem.*

Keywords: Mixed-type equation; the bi-ordinal Hilfer derivative; hyper-Bessel fractional differential opertor

1 Introduction

Fractional differential equations plays a significant role, because of its multiple applications in engineering, chemistry, biology and other parts of science and modeling the real-life problems [1]-[3]. Studying boundary value problems for linear and non-linear fractional differential equations with Riemann-Liouville and Caputo fractional derivative have becoming interesting targets simultaneously [4]-[7].

The quality and the types of articles have been changed when the generalized Riemann-Liouville differential operators (later called Hilfer derivative) used in the scientific field with its interesting application [8], [9]. To tell the truth, the generalization of the Riemann-Lioville fractional differential operators was already announced by M. M. Dzherbashian and A. B. Nersesian [10] in 1968, but because of some reasons it was not familiar with many mathematicians around the globe till the translation of this work published in the journal of Fract. Calc. Appl. An. [11]. We also refer some papers [12]-[13] devoted studying some problems with the Dzherbashian-Nersesian differential operator which has the following form

$$D_{0x}^{\sigma_n} g(x) = I_{0x}^{1-\gamma_n} D_{0x}^{\gamma_{n-1}} ... D_{0x}^{\gamma_1} D_{0x}^{\gamma_0}, \quad n \in \mathbb{N}, \ x > 0 \tag{1}$$

where I_{0x}^{α} and D_{0x}^{α} are the Riemann-Liouville fractional integral and the Riemann-Liouville fractional derivative of order α respectively, $\sigma_n \in (0, n]$ which is defined by

$$\sigma_n = \sum_{j=0}^{n} \gamma_j - 1 > 0, \ \gamma_j \in (0, 1].$$

V.I.Bulavatsky considered specific generalization of Hilfer fractional derivative [14], the particular case of the Dzherbashian-Nersesian differential operator which forms as follows [15]:

$$D_{0\pm}^{(\alpha,\beta)\mu} g(t) = I_{0\pm}^{\mu(n-\alpha)} \left(\pm \frac{d}{dt} \right)^n I_{0\pm}^{(1-\mu)(n-\beta)} g(t) \tag{2}$$

This operator is similar to Hilfer derivative in terms of its interpolation concept between the Riemann-Liouville and Caputo fractional derivatives, specifically, i.e.

$$D_{a+}^{(\alpha,\beta)\mu} g(t) = \begin{cases} D_{a+}^{\beta} g(t), & \mu = 0, \\ {}^{C}D_{a+}^{\alpha} g(t), & \mu = 1. \end{cases}$$

In [15] the Cauchy problem is investigated for the ordinary differential equation involving the right-sided bi-ordinal Hilfer fractional derivative:

$$\begin{cases} D_{0-}^{(\alpha,\beta)\mu} u(t) = \lambda u(t) + g(t), \\ \lim\limits_{t \to 0-} I_{0-}^{2-\gamma} u(t) = \xi_0, \\ \lim\limits_{t \to 0-} \frac{d}{dt} I_{0-}^{2-\gamma} u(t) = \xi_1, \end{cases} \tag{3}$$

where $1 < \alpha, \beta \leq 2$, $\gamma = \beta + \mu(2 - \beta)$, $\xi_0, \xi_1 \in \mathbb{R}$, $g(t)$ is the given function.

Lemma 1. Let $g(t) \in C[-T,0],\ g'(t) \in L_1(-T,0)$. Then the solution of the problem (3) as follows:

$$u(t) = \xi_0(-t)^{\gamma-2}E_{\delta,\gamma-1}[\lambda(-t)^\delta] - \xi_1(-t)^{\gamma-1}E_{\delta,\gamma}[\lambda(-t)^\delta] +$$

$$+\int_t^0 (z-t)^{\delta-1}E_{\delta,\delta}\left[\lambda(z-t)^\delta\right] g(z)dz, \tag{4}$$

where $\delta = \beta + \mu(\alpha-\beta)$, $\gamma = \beta + \mu(2-\beta)$.

We notice that while have been investigated the initial-boundary value or non-local problems involving popular class of differential operators like Riemann-Liouville, Caputo [16], [17], Hilfer fractional derivatives, Hadamard [18], Hilfer-Hadamard, Prabhakar, Atangana-Baleanu [19] the interest to another type of the differential operators is increased by many scientists, for instance, the hyper-Bessel differential operator [20], [21] is becoming main target of research. The importance of the hyper-Bessel differential operator is increasing since introduced by Dimovski [22] because of its applications in science. For example, in [23] authors used hyper-Bessel differential operator to investigate heat diffusion equation for discribing Brownian motion. In [24] it is investigated fractional relaxation equation the regularized Caputo-like counterpart of the hyper-Bessel operator which has the following form

$${}^C\left(t^\theta \frac{d}{dt}\right)^\alpha f(t) = (1-\theta)^\alpha t^{-\alpha(1-\theta)} D_{1-\theta}^{-\alpha,\alpha}\left(f(t) - f(0)\right) \tag{5}$$

where

$$D_\beta^{\gamma,\delta} f(t) \prod_{j=1}^{n}\left(\gamma + j + \frac{t}{\beta}\frac{d}{dt}\right)\left(I_\beta^{\gamma+\delta,n-\delta} f(t)\right),$$

is a Erdelyi-Kober fractional derivative and $I_\beta^{\gamma,\delta}$ is a Erdelyi-Kober (E-K) fractional integral defined as [25]

$$I_\beta^{\gamma,\delta} f(t) = \frac{t^{-\beta(\gamma+\delta)}}{\Gamma(\delta)}\int_0^t (t^\beta - \tau^\beta)^{\delta-1}\tau^{\beta\gamma} f(\tau) d(\tau^\beta),$$

which can be reduced up with weight to $I_{0+}^q f(t)$ Riemann-Liouville fractional integral at $\gamma = 0$ and $\beta = 1$. In [26] Fatma Al-Musalhi, et al. considered direct and inverse problems for the fractional diffusion equation with the regularized Caputo-like counterpart of the hyper-Bessel operator and proved the theorem of existence and uniqueness of the solution.

Several local and nonlocal boundary value problems for mixed-type equations, ie, elliptic-hyperbolic and hyperbolic type equations were published [27]-[29]. The interesting point is that the conjugation conditions are taken according to the considered mixed-type equations and domains [30], [31].

2 Formulation of the problem and its investigation

In the present work, we investigate the following fractional differential equation of mixed type involving the regularized Caputo-like counterpart of the hyper-Bessel operator and the bi-ordinal Hilfer derivative in $\Omega = \Omega_1 \cup \Omega_2 \cup Q$ domain:

$$f(x,t) = \begin{cases} {}^C\left(t^\theta \dfrac{\partial}{\partial t}\right)^{\alpha_1} u(x,t) - \dfrac{1}{x}u_x(x,t) - u_{xx}(x,t), & (x,t) \in \Omega_1, \\ D_{0-}^{(\alpha_2,\beta_2)\mu} u(x,t) - \dfrac{1}{x}u_x(x,t) - u_{xx}(x,t), & (x,t) \in \Omega_2, \end{cases} \tag{6}$$

where $\Omega_1 = \{(x,t) : 0 < x < 1,\ 0 < t < T\}$, $\Omega_2 = \{(x,t) : 0 < x < 1,\ -T < t < 0\}$, $Q = \{(x,t) : 0 < x < 1,\ t = 0\}$, $0 < \alpha_1 \le 1$, $\theta < 1$, $1 < \alpha_2, \beta_2 \le 2$, $0 \le \mu \le 1$,
$D_{0-}^{(\alpha_2,\beta_2)\mu}$ is the right-sided bi-ordinal Hilfer fractional derivative in the form (2) and
${}^C\left(t^\theta \frac{d}{dt}\right)^{\alpha_1}$ is the regularized Caputo-like counterpart of the hyper-Bessel fractional differential operator defined as (5).

Problem. Find a solution of eq.(6) in Ω, which satisfies the following regularity conditions

$$u(x,t) \in C(\Omega), \quad u(\cdot,t) \in C^1 \cap C^2(\Omega_1 \cup \Omega_2),$$
$${}^C\left(t^\theta \frac{\partial}{\partial t}\right)^\alpha u(x,t) \in C(\Omega_1), \qquad D_{0-}^{(\alpha_2,\beta_2)\mu} u(x,t) \in C(\Omega_2)$$

along with the boundary conditions

$$\lim_{t\to 0+} x u_x(x,t) = 0, \;\; u(1,t) = 0, \quad -T \le t \le T, \tag{7}$$

non-local condition

$$\sum_{i=1}^{m} p_i I_{0-}^{(1-\mu)(2-\beta_2)} u(x,\xi_i) = u(x,T), \;\; 0 \le x \le 1, \tag{8}$$

and the gluing conditions

$$\lim_{t\to 0-} I_{0-}^{(1-\mu)(2-\beta_2)} u(x,t) = \lim_{t\to 0+} u(x,t), \quad 0 \le x \le 1, \tag{9}$$

$$\lim_{t\to 0-} \frac{d}{dt} I_{0-}^{(1-\mu)(2-\beta_2)} u(x,t) = \lim_{t\to 0+} t^{1-(1-\theta)\alpha_1} u_t(x,t), \quad 0 < x < 1, \tag{10}$$

where $-T \le \xi_1 < \xi_2 < .... < \xi_m \le 0, \;\; f(x,t)$ is a given function.

Using the separation variables method we obtain the following spectral problem

$$X''(x) - \frac{1}{x} X'(x) + \lambda^2 X(x) = 0, \tag{11}$$

$$\lim_{x\to 0+} x X(x) = 0, X(1) = 0, \tag{12}$$

(9) is a Bessel equation of order zero; furthermore, (9), (10) is a self-adjoint problem and its eigenfunctions are the Bessel functions given as follows

$$X_k(x) = J_0(\lambda_k x), \;\; k = 1, 2, ..., \tag{13}$$

and the eigenvalues λ_k, are the positive zeros of $J_0(x)$, i.e.

$$\lambda_k = \pi k - \frac{\pi}{4}.$$

The system of eigenfunctions $\{X_k\}$ forms an orthogonal basis in $L^2(0,1)$, hence we can write sought function and given function in the form of series expansions as follows:

$$u(x,t) = \sum_{k=1}^{\infty} u_k(t) J_0(x), \tag{14}$$

$$f(x,t) = \sum_{k=1}^{\infty} f_k(t) J_0(x), \tag{15}$$

where $u_k(t)$ is not knownt yet, and $f_k(t)$ is the coefficient of Fourier-Bessel series, i.e.

$$f_k(t) = \frac{2}{J_1^2(\lambda_k)} \int_0^1 x f(x,t) J_0(\lambda_k x) dx.$$

Let us introduce new notations:

$$\lim_{t\to 0-} I_{0-}^{(1-\mu)(2-\beta_2)} u(x,t) = \varphi(x), \;\; 0 \le x \le 1, \tag{16}$$

$$\lim_{t\to 0-} \frac{d}{dt} I_{0-}^{(1-\mu)(2-\beta_2)} u(x,t) = \psi(x), \;\; 0 < x < 1, \tag{17}$$

$$\lim_{t\to 0+} u(x,t) = \tau(x), \quad 0 \le x \le 1, \tag{18}$$

here $\varphi(x)$, $\tau(x)$ and $\psi(x)$ are unknown functions to be found later.

Further, after substituting (14) and (15) into the eq.(6) and initial conditions (16), (17), (18), we obtain the following prdoblems

$$\begin{cases} {}^{C}\Big(t^{\theta}\dfrac{d}{dt}\Big)^{\alpha_1}u_k(t)+\lambda_k^2u_k(t)=f_k(t), \\ \qquad\qquad u_k(0+)=\tau_k, \end{cases} \tag{19}$$

and

$$\begin{cases} D_{0-}^{(\alpha_2,\beta_2)\mu}u_k(t)+\lambda_k^2u_k(t)=f_k(t), \\ I_{0-}^{(1-\mu)(2-\beta_2)}u_k(0-)=\varphi_k, \\ \quad \dfrac{d}{dt}I_{0-}^{(1-\mu)(2-\beta_2)}u_k(0-)=\psi_k, \end{cases} \tag{20}$$

in Ω_1 and Ω_2 respectively.

The problem (19) was studied in [26] and we can write the solution of the considered problem in Ω_1:

$$u(x,t)=\sum_{k=1}^{\infty}\Big[\tau_kE_{\alpha_1,1}\Big(-\frac{{\lambda_k}^2}{p^{\alpha_1}}t^{p\alpha_1}\Big)+G_k(t)\Big]J_0(\lambda_kx), \tag{21}$$

here $p=1-\theta$ and

$$G_k(t)=\frac{1}{p^{\alpha_1}\Gamma(\alpha_1)}\int_0^t\big(t^p-\tau^p\big)^{\alpha_1-1}f_k(\tau)d(\tau^p)-$$

$$-\frac{{\lambda_k}^2}{p^{2\alpha_1}}\int_0^t\big(t^p-\tau^p\big)^{2\alpha_1-1}E_{\alpha_1,2\alpha_1}\Big[-\frac{{\lambda_k}^2}{p^{\alpha_1}}(t^p-\tau^p)^{\alpha_1}\Big]f_k(\tau)d(\tau^p),$$

where τ_k is not known yet.

As we mentioned above the problem (20) studied in [15] and using the Lemma 1 we can write the solution of (20) and then, considering (14) the solution of considered problem in Ω_2 domain can be represented in the following form

$$u(x,t)=\sum_{k=1}^{+\infty}\varphi_k(-t)^{\gamma_2-2}E_{\delta_2,\gamma_2-1}[-\lambda_k^2(-t)^{\delta_2}]J_0(\lambda_kx)-$$

$$-\sum_{k=1}^{+\infty}\psi_k(-t)^{\gamma_2-1}E_{\delta_2,\gamma_2}[-\lambda_k^2(-t)^{\delta_2}]J_0(\lambda_kx)+$$

$$+\sum_{k=1}^{+\infty}\int_t^0(z-t)^{\delta_2-1}E_{\delta_2,\delta_2}[-\lambda_k^2(z-t)^{\delta_2}]f_k(z)dzJ_0(\lambda_kx), \tag{22}$$

where $\gamma_2=\beta_2+\mu(2-\beta_2)$, $\delta_2=\beta_2+\mu(\alpha_2-\beta_2)$ and φ_k, ψ_k are not known yet.

After substituting (21) and (22) into gluing conditions with considering (14), (15) we obtain the following system of equations with respect to τ_k, φ_k and ψ_k:

$$\begin{cases} \psi_k=-\frac{\lambda_k^2}{\Gamma(\alpha_1)}\tau_k, \\ \qquad \tau_k=\varphi_k \end{cases} \tag{23}$$

Considering non-local condition (8) and from (23) we find unknowns as follows

$$\tau_k=\varphi_k=\frac{F_k}{\Delta_k}, \tag{24}$$

$$\psi_k=\frac{-\lambda_k^2}{p^{\alpha_1}\Gamma(\alpha_1)}\frac{F_k}{\Delta_k}, \tag{25}$$

where

$$\Delta_k=\sum_{i=1}^{m}p_i\Big[E_{\delta_2,1}(-\lambda_k^2(-\xi_i))+\frac{\lambda_k^2\xi_i}{p^{\alpha_1}\Gamma(\alpha_1)}E_{\delta_2,2}(-\lambda_k^2(-\xi_i))\Big]-E_{\alpha_1,1}\Big(-\frac{\lambda_k^2}{p^{\alpha_1}}T^{\alpha_1p}\Big)$$

$$F_k = G_k(T) - \sum_{i=1}^{m} p_i \int_{\xi_i}^{0} (s-\xi_i)^{\delta_2-\gamma_2+1} E_{\delta_2,\delta_2-\gamma_2+2}(-\lambda_k^2 (s-\xi_i)^{\delta_2}) f_k(s) ds,$$

$$G_k(T) = \frac{1}{p^{\alpha_1}\Gamma(\alpha_1)} \int_0^T \left(T^p - \tau^p\right)^{\alpha_1-1} f_k(\tau) d(\tau^p) -$$

$$-\frac{{\lambda_k}^2}{p^{2\alpha_1}} \int_0^T \left(T^p - \tau^p\right)^{2\alpha_1-1} E_{\alpha_1,2\alpha_1}\Big[-\frac{{\lambda_k}^2}{p^{\alpha_1}}(T^p-\tau^p)^{\alpha_1}\Big] f_k(\tau) d(\tau^p).$$

If $\Delta_k \neq 0$, then we can find τ_k, φ_k, ψ_k unknowns uniquely.

First we show that $\Delta_k \neq 0$ for sufficiently large k. For this intention we use the following lemma obtained from the properties of Wright-type function studied by A. Pskhu in [32].

Lemma 2. [32] If $\pi \geq |arg z| > \frac{\pi\alpha}{2} + \varepsilon,\ \varepsilon > 0$, then the following relations are valid for $z \to \infty$:

$$\lim_{|z|\to\infty} E_{\alpha,\beta}(z) = 0,$$

$$\lim_{|z|\to\infty} z E_{\alpha,\beta}(z) = -\frac{1}{\Gamma(\beta-\alpha)}.$$

By using the lemma 2, we can calculate the behaviour of Δ_k at $k \to \infty$

$$\lim_{k\to+\infty} \Delta_k = \lim_{|z_1|\to+\infty} \sum_{i=1}^{m} p_i \big[E_{\delta_2,1}(z_1) + \frac{1}{\Gamma(\alpha_1)p^{\alpha_1}} z_1 E_{\delta_2,2}(z_1)\big] -$$

$$-\lim_{|z_2|\to+\infty} E_{\alpha_1,1}(z_2) = \sum_{i=1}^{m} \frac{p_i}{\Gamma(\alpha_1)p^{\alpha_1}\Gamma(2-\delta_2)}$$

where $z_1 = -\lambda_k^2(-\xi_i)$, $z_2 = -\frac{\lambda_k^2}{p^{\alpha_1}} T^{\alpha_1 p}$, $\lambda_k = \pi k - \frac{\pi}{4}$.

From the last equality it is seen that $\Delta_k \neq 0$ for sufficiently large k.

2.1 Unequeness of the solution

To show the uniqueness of the solution, it is enough to prove that homogeneous problem has a trivial solution.

Let us first consider the following integral

$$u_k(t) = \frac{2}{J_1^2(\lambda_k)} \int_0^1 x u(x,t) J_0(\lambda_k x) dx,\ k = 1,2,3,...., \tag{26}$$

Then we introduce another function based on (26)

$$v_\varepsilon(t) = \frac{2}{J_1^2(\lambda_k)} \int_\varepsilon^{1-\varepsilon} x u(x,t) J_0(\lambda_k x) dx,\ k = 1,2,3,...., \tag{27}$$

Applying ${}^C\left(t^\theta \frac{\partial}{\partial t}\right)^{\alpha_1}$ and $D_{0-}^{(\alpha_2,\beta_2)\mu}$ to (27) and using the equation (6) in homogeneous case with respect to t

$${}^C\left(t^\theta \frac{\partial}{\partial t}\right)^{\alpha_1} v_\varepsilon(t) = \frac{2}{J_1^2(\lambda_k)} \int_\varepsilon^{1-\varepsilon} {}^C\left(t^\theta \frac{\partial}{\partial t}\right)^{\alpha_1} u(x,t) x J_0(\lambda_k x) dx =$$

$$= \frac{2}{J_1^2(\lambda_k)} \int_\varepsilon^{1-\varepsilon} \Big[u_{xx}(x,t) + \frac{1}{x} u_x(x,t)\Big] x J_0(\lambda_k x) dx = \frac{-2\lambda_k^2}{J_1^2(\lambda_k)} \int_\epsilon^{1-\epsilon} u(x,t) x J_0(\lambda_k x) dx$$

$$D_{0-}^{(\alpha_2,\beta_2)\mu} v_\varepsilon(t) = \frac{2}{J_1^2(\lambda_k)} \int_\varepsilon^{1-\varepsilon} D_{0-}^{(\alpha_2,\beta_2)\mu} u(x,t) x J_0(\lambda_k x) dx =$$

$$= \frac{2}{J_1^2(\lambda_k)} \int\limits_{\varepsilon}^{1-\varepsilon} \Big[u_{xx}(x,t) + \frac{1}{x} u_x(x,t)\Big] x J_0(\lambda_k x) dx = \frac{-2\lambda_k^2}{J_1^2(\lambda_k)} \int\limits_{\epsilon}^{1-\epsilon} u(x,t) x J_0(\lambda_k x) dx$$

and integrating by parts twice the right sides of the equalities on $t \in (0,T)$ and $t \in (-T,0)$, respectively, and passing to the limit on $\varepsilon \to +0$ yield

$$\begin{cases} {}^C\Big(t^\theta \frac{d}{dt}\Big)^{\alpha_1} u_k(t) + \lambda^2 u_k(t) = 0, \ \ t > 0, \\ D_{0-}^{(\alpha_2,\beta_2)\mu} u_k(t) + \lambda^2 u_k(t) = 0, \ \ t < 0. \end{cases} \tag{28}$$

Considering conditions in (19), (20) in homogeneous case, (28) has a solution $u_k(t) = 0$ if $\Delta_k \neq 0$. Then from (26) and the completeness of the system $X_k(x)$ in the space $L_2[0,l]$, $u(x,t) \equiv 0$ in $\overline{\Omega}$. This completes the prove of uniqueness of the solution of the main problem.

2.2 Existence of the solution

In the below, we present the well-known lemma about the upper bound of the Mittag-Leffler function and the theorem related to Fourier-Bessel series for showing the existence of the solution.

Lemma 3. Let $\alpha < 2$, $\beta \in R$ and $\frac{\pi\alpha}{2} < \mu < min\{\pi, \pi\alpha\}$, $M^* > 0$. Then,the following estimate hold

$$|E_{\alpha,\beta}(z)| \leq \frac{M^*}{1+|z|}, \ \ \mu \leq |arg z| \leq \pi, \ \ |z| \geq 0.$$

Theorem 1. Let $f(x)$ be a function defined on the interval [0, 1] such that $f(x)$ is differentiable $2s$ times ($s \geq 1$) and

- $f(0) = f'(0) = ... = f^{(2s-1)}(0) = 0$
- $f^{(2s)}(x)$ is bounded (this derivative may not exist at certain points)
- $f(1) = f'(1) = ... = f^{(2s-2)}(1) = 0$

then the following inequalities satisfied by the Fourier-Bessel coefficients of $f(x)$:

$$|f_k| \leq \frac{M}{\lambda_k^{2s-\frac{1}{2}}}.$$

Considering above lemma 3 and theorem 1, we show the upper bound of $G_k(t)$:

$$| \ G_k(t) \ | \leq \int\limits_0^t |t^p - \tau^p|^{\alpha_1 - 1} |f_k(\tau)| d(\tau^p) +$$

$$+ \frac{\lambda_k^2}{p^{2\alpha_1}} \int\limits_0^t |t^p - \tau^p|^{2\alpha_1 - 1} |E_{\alpha_1, 2\alpha_1}\big[-\frac{\lambda_k^2}{p^{\alpha_1}} (t^p - \tau^p)^{\alpha_1}\big]| |f_k(\tau)| d(\tau^p) \leq$$

$$\leq \frac{1}{p^{\alpha_1}\Gamma(\alpha_1)} \int\limits_0^t |t^p - \tau^p|^{\alpha_1 - 1} \frac{M}{{\lambda_k}^{7/2}} d(\tau^p) + \frac{\lambda_k^2}{p^{2\alpha_1}} \int\limits_0^t \frac{p^{\alpha_1} |t^p - \tau^p|^{2\alpha_1 - 1} M^*}{p^{\alpha_1} + \lambda_k^2 |t^p - \tau^p|^{\alpha_1}} \frac{M}{{\lambda_k}^{7/2}} d(\tau^p) \leq$$

$$\leq \Big[\frac{M}{{\lambda_k}^{7/2} p^{\alpha_1}\Gamma(\alpha_1)} + \frac{M^* M}{p^{\alpha_1} {\lambda_k}^{7/2}}\Big] \int\limits_0^t |t^p - \tau^p|^{\alpha_1 - 1} d(\tau^p) \leq \frac{1}{{\lambda_k}^{7/2}} \Big[\frac{M}{p^{\alpha_1}\Gamma(\alpha_1)} + \frac{M^* M}{p^{\alpha_1}}\Big] \frac{|t|^{\alpha_1 p}}{\alpha_1} \leq$$

$$\leq \frac{M_1}{{\lambda_k}^{7/2}} |t|^{\alpha_1 p} \leq \frac{M_1}{{\lambda_k}^{7/2}} |T|^{\alpha_1 p}, \qquad M_1 = \frac{M}{p^{\alpha_1}\Gamma(\alpha_1 + 1)} + \frac{M^* M}{\alpha_1 p^{\alpha_1}},$$

By using the last inequality, lemma 3 and theorem 1, we can write the upper bound of F_k:

$$|F_k| \leq |G_k(T)| + \sum_{i=1}^m p_i \int\limits_{\xi_i}^0 |s - \xi_i|^{\delta_2 - \gamma_2 + 1} |E_{\delta_2, \delta_2 - \gamma_2 + 2}(-{\lambda_k}^2 (s - \xi_i)^{\delta_2})| |f_k(s)| ds \leq$$

$$\leq \frac{M_1 T^{\alpha_1 p}}{{\lambda_k}^{7/2}} + \sum_{i=1}^{m} p_i \int_{\xi_i}^{0} |s-\xi_i|^{\delta_2-\gamma_2+1} \frac{M^*}{1+\lambda_k^2|s-\xi_i|^{\delta_2}} \frac{M}{\lambda_k^{7/2}} ds \leq$$

$$\leq \frac{M_1 T^{\alpha_1 p}}{{\lambda_k}^{7/2}} + \sum_{i=1}^{m} p_i \int_{\xi_i}^{0} |s-\xi_i|^{1-\gamma_2} \frac{MM^*}{\lambda_k^{11/2}} \leq \frac{M_1 T^{\alpha_1 p}}{{\lambda_k}^{7/2}} + \sum_{i=1}^{m} p_i \frac{(-\xi_i)^{2-\gamma_2}}{(2-\gamma_2)} \frac{MM^*}{\lambda_k^{11/2}} \leq,$$

$$\leq \frac{M_2}{\lambda_k^{7/2}}, \quad M_2 = M_1 T^{\alpha_1 p} + \sum_{i=1}^{m} \frac{p_i MM^* (-\xi_i)^{2-\gamma_2}}{(2-\gamma_2){\lambda_k}^2},$$

or

$$|F_k| \leq \frac{M_2}{\lambda_k^{7/2}}. \tag{29}$$

Now considering (29) and $\Delta_k \neq 0$ then, we write upper bounds of τ_k, φ_k, ψ_k in (24) and (25).

$$|\tau_k| = |\varphi_k| \leq |\frac{1}{\Delta_k}||F_k| \leq \frac{M_2}{|\Delta_k|\lambda_k^{7/2}}, \tag{30}$$

$$|\psi_k| = \Big|\frac{-\lambda_k^2}{p^{\alpha_1}\Gamma(\alpha_1)}\Big|\Big|\frac{F_k}{\Delta_k}\Big| \leq \frac{M_2}{p^{\alpha_1}\Gamma(\alpha_1)|\Delta_k|\lambda_k^{3/2}}. \tag{31}$$

For proving the existence of the solution, we need to show uniform convergence of series representations of $u(x,t)$, $u_x(x,t)$, $u_{xx}(x,t)$, ${}^C\left(t^\theta \frac{\partial}{\partial t}\right)^\alpha u(x,t)$ and $D_{0-}^{(\alpha_2,\beta_2)\mu} u(x,t)$ by using the solution (21) and (22) in Ω_1 and Ω_2 respectively.

According to the last inequality and theorem 1, then we can present the existence of the solutions in both domains.

$$|u(x,t)| = \sum_{k=1}^{\infty} |u_k(t)||J_0(\lambda_k x)| \leq \sum_{k=1}^{\infty} |u_k(t)| \leq \sum_{k=1}^{\infty} \Big[|\tau_k||E_{\alpha_1,1}(-\frac{\lambda_k^2}{p^{\alpha_1}} t^{\alpha_1 p})| + |G_k(t)|\Big] \leq$$

$$\leq \sum_{k=1}^{\infty} \Big(\frac{p^{\alpha_1}}{p^{\alpha_1} + {\lambda_k}^2 |t^{p\alpha_1}|} \frac{M_2}{|\Delta_k|\lambda_k^{7/2}} + \frac{M_1 T^{\alpha_1 p}}{{\lambda_k}^{7/2}}\Big)$$

One can shows that the series representation of $u(x,t)$ is bounded by convergent numerical series and by Weierstrass M-test, the series of $u(x,t)$ converges uniformly in Ω_1.

Now we remind some properties of Bessel functions [33]: $J_0'(x) = -J_1(x)$;
$2J_1'(x) = J_0(x) - J_2(x)$ and asymptotic formula $|J_\nu(\lambda)kx)| \leq \frac{2A}{\sqrt{\lambda_k x)}}, \quad \nu > -1/2$,
$A = const$.

It is not difficult to see that the series representation of $u_{xx}(x,t)$ is bigger than $u_x(x,t)$ hence it is enough to show the uniform convergency of $u_{xx}(x,t)$. By using these properties we have

$$|u_{xx}(x,t)| \leq \sum_{k=1}^{\infty} |u_k(t)|\Big|\frac{d^2}{dx^2} J_0(\lambda_k x)\Big| = \sum_{k=1}^{\infty} |u_k(t)| \frac{\lambda_k^2}{2}\Big|J_2(\lambda_k x) - J_0(\lambda_k x)\Big|$$

And as a similar way of $u(x,t)$ we can show that

$$|u_{xx}(x,t)| \leq \sum_{k=1}^{\infty} \Big(\frac{p^{\alpha_1}}{p^{\alpha_1} + {\lambda_k}^2 |t^{p\alpha_1}|} \frac{M_2}{|\Delta_k|\lambda_k^{7/2}} + \frac{M_1 T^{\alpha_1 p}}{{\lambda_k}^{7/2}}\Big) \frac{2A}{\sqrt{\lambda_k x)}}.$$

From the last inequality we can see that the series representation of $u_{xx}(x,t)$ is bounded by convergent series. According to Weierstrass M-test, the series of $u_{xx}(x,t)$ converges uniformly in Ω_1.

The uniform convergency of ${}^C\left(t^\theta \frac{\partial}{\partial t}\right)^\alpha u(x,t)$ which is defined as

$${}^C\left(t^\theta \frac{\partial}{\partial t}\right)^\alpha u(x,t) = u_{xx}(x,t) - \frac{1}{x} u_x(x,t) + f(x,t)$$

is similar to the way of showing convergency of the series representation of $u_{xx}(x,t)$.

In Ω_2 domain it is enough to show the uniform convergency of $u_{xx}(x,t)$ which is bigger than other series. Hence the convergency of the series of $u(x,t)$, $u_x(x,t)$, $D_{0-}^{(\alpha_2,\beta_2)\mu}u(x,t)$ can be derived from the uniform convergency of $u_{xx}(x,t)$.

From theorem 1 and lemma 3, in Ω_2 we can have

$$|u_{xx}(x,t)| \leq \sum_{k=1}^{\infty} |u_k(t)| \frac{\lambda_k^2}{2} \Big| J_2(\lambda_k x) - J_0(\lambda_k x) \Big| \leq \sum_{k=1}^{\infty} \lambda_k^2 |\varphi_k| |(-t)^{\gamma_2 - 2}| |E_{\delta_2,\gamma_2-1}(-\lambda_k^2(-t)^{\delta_2})| +$$

$$+ \sum_{k=1}^{\infty} \lambda_k^2 |\psi_k| |(-t)^{\gamma_2-1}| |E_{\delta_2,\gamma_2}(-\lambda_k^2(-t)^{\delta_2})| + \sum_{k=1}^{+\infty} \lambda_k^2 \int_t^0 |z-t|^{\delta_2-1} |E_{\delta_2,\delta_2}[-\lambda_k^2(z-t)^{\delta_2}]| |f_k(z)| dz \leq$$

$$\leq \sum_{k=1}^{\infty} \Big[\frac{\lambda_k^2 M_2}{|\Delta_k| \lambda_k^{7/2}} \frac{|(-t)^{\gamma_2-2}| M^*}{1+\lambda_k^2 |(-t)^{\delta_2}|} + \frac{\lambda_k^2 M_2}{|\Delta_k| \lambda_k^{3/2} p^{\alpha_1} \Gamma(\alpha_1)} \frac{|(-t)^{\gamma_2-1}| M^*}{1+\lambda_k^2 |(-t)^{\delta_2}|} \Big] +$$

$$+ \sum_{k=1}^{+\infty} \lambda_k^2 \int_t^0 |z-t|^{\delta_2-1} \frac{M^*}{1+\lambda_k^2 |(z-t)^{\delta_2}|} \frac{M}{\lambda_k^{7/2}} dz \leq$$

$$\leq \sum_{k=1}^{\infty} \frac{1}{\lambda_k^{3/2}} \Big[\frac{M_2 M^* T^{\gamma_2-2}}{|\Delta_k| (1+{\lambda_k}^2 T^{\delta_2})} + \frac{M_2 M^* T^{\gamma_2-\delta_2-1}}{|\Delta_k|} + \frac{M M^* \ln(1+\lambda_k^2 T^{\delta_2})}{\delta_2 \lambda_k^2} \Big] \leq \sum_{k=1}^{\infty} \frac{M_3}{\lambda_k^{3/2}}$$

where $\lim\limits_{\lambda_k \to \infty} \frac{\ln(1+\lambda_k^2 T^{\delta_2})}{\delta_2 \lambda_k^2} < \infty$ according to l'Hopital's rule. It can be seen that the series representation of $u_{xx}(x,t)$ is bounded by convergent numerical series and due to Weierstrass M-test, the series of $u_{xx}(x,t)$ converges uniformly in Ω_2

Using (30), (31) and theorem 1, lemma 3, we can show the uniform convergence of $u(x,t)$, $u_x(x,t)$, and $D_{0-}^{(\alpha_2,\beta_2)\mu}u(x,t)$ as a similar method used for $u_{xx}(x,t)$.

Finally we have proved the uniqueness and existence of the solution to the considered problem as stated in the following theorem.

Theorem 2. Let $\Delta_k \neq 0$ and $f(x,t)$ satisfies the following conditions

- $f(0,t) = f_x'(0,t) = ... = f_x'''(0,t) = 0$;
- $f(1,t) = f_x'(1,t) = f_x''(1,t) = 0$;
- $\frac{\partial^4}{\partial x^4} f(x,t)$ is bounded,

then the considered problem's unique solution exists.

3 Acknowledgments

The author is very grateful to his supervisor prof. Erkinjon Karimov for his valuable suggestions and to the reviewers in terms of improving the quality of this research.

References

[1] F. Mainardi, *Fractional calculus and waves in linear Viscoelasticity*, Imperial College Press, 2010.

[2] Kilbas A. A., Srivastava H. M., Trujillo J. J. *Theory and Applications of Fractional Differential Equations*, volume 204. North-Holland Mathematics Studies. Amsterdam: Elsevier, 2006.

[3] O. S. Iyiola, F. D. Zaman, *A fractional diffusion equation model for cancer tumor*, AIP Adv.4 (2014) 107-121

[4] Bashir Ahmad, Soha Hamdan, Ahmed Alsaedi, Sotiris K. Ntouyas. *study of a nonlinear coupled system of three fractional differential equations with nonlocal coupled boundary conditions* Advances in Difference Equations (2021) 2021:278 https://doi.org/10.1186/s13662-021-03440-7

[5] Cuesta, Eduardo, Kirane, Mokhtar, Alsaedi, Ahmed and Ahmad, Bashir. "On the sub-diffusion fractional initial value problem with time variable order" Advances in Nonlinear Analysis, vol. 10, no. 1, 2021, pp. 1301-1315. https://doi.org/10.1515/anona-2020-0182

[6] Kerbal, S., Kadirkulov, B. J., Kirane, M. *Direct and inverse problems for a Samarskii-Ionkin type problem for a two dimensional fractional parabolic equation.* Progress in Fractional Differentiation and Applications, 4(**3**), 147-160. https://doi.org/10.18576/pfda/040301

[7] Ahmad, B., Nieto, J.J. *Riemann-Liouville fractional integro-differential equations with fractional nonlocal integral boundary conditions.* Bound Value Probl 2011, 36 (2011). https://doi.org/10.1186/1687-2770-2011-36

[8] Hilfer R. *Applications of Fractional Calculus in Physics.* Singapore: World Scientific, 2000.

[9] Hilfer R., Luchko Y., Tomovski ˇZ. *Operational method for the solution of fractional differential equations with generalized Riemann-Liouville fractional derivatives.* Fract. Calc. Appl. Anal. 12(3), 2009, pp.299-318

[10] M. M. Dzhrbashyan, A. B. Nersesyan, *Fractional Derivatives and the Cauchy Problem for Fractional Differential Equations*, Izv. Akad. Nauk Armyan. SSR. 3, No 1 (1968), 3-29.

[11] M. M. Dzherbashian, A. B. Nersesian, *Fractional derivatives and Cauchy problem for differential equations of fractional order.* Fract. Calc. Appl. Anal. 23, No 6 (2020), 1810-1836. https://doi.org/10.1515/fca-2020-0090.

[12] Anwar Ahmad, Muhammad Ali, Salman A. Malik. *Inverse problems for diffusion equation with fractional Dzherbashian-Nersesian operator.* arXiv:2105.05040v1

[13] Bogatyreva F.T. *Initial value problem for fractional order equation with constant coefficients,* Vestnik KRAUNC. Fiz.-mat. nauki. 2016, 16: 4-1, 21- 26. DOI: 10.18454/2079- 6641-2016-16-4-1-21-26

[14] V. M. Bulavatsky. *Closed form of the solutions of some boundary problems for anomalous diffusion equation with Hilfer's generalized derivative.* Cybernetics and Systems Analysis. 30(4), 2014, pp. 570-577

[15] Karimov E. T., Toshtemirov B. H. Non-local boundary value problem for a mixed-type equation involving the bi-ordinal Hilfer fractional differential operators. Uzbek Mathematical Journal, 2021, Volume 65, Issue 2, pp.61-77 DOI: 10.29229/uzmj.2021-2-5

[16] Ahmad, B., Hamdan, S., Alsaedi, A. et al. *A study of a nonlinear coupled system of three fractional differential equations with nonlocal coupled boundary conditions.* Adv Differ Equ 2021, 278 (2021). https://doi.org/10.1186/s13662-021-03440-7

[17] Bashir Ahmad, Ahmed Alsaedi, Mokhtar Kirane and Ramiz G. Tapdigoglu (2017) *An inverse problem for space and time fractional evolution equations with an involution perturbation*, Quaestiones Mathematicae, 40:2, 151-160, DOI: 10.2989/16073606.2017.1283370

[18] Ahmad, B., Ntouyas, S.K., Alsaedi, A. et al. *A study of a coupled system of Hadamard fractional differential equations with nonlocal coupled initial-multipoint conditions.* Adv Differ Equ 2021, 33 (2021). https://doi.org/10.1186/s13662-020-03198-4

[19] Meiirkhan Borikhanov, Mokhtar Kirane, Berikbol T. Torebek, *Maximum principle and its application for the nonlinear time-fractional diffusion equations with Cauchy-Dirichlet conditions,* Applied Mathematics Letters, Volume 81, 2018, Pages 14-20, ISSN 0893-9659, https://doi.org/10.1016/j.aml.2018.01.012.

[20] V. Kiryakova, *Transmutation method for solving hyper-Bessel differential equations based on the Poisson-Dimovski transformation.* Fract. Calc. Appl. Anal. 11, No 3 (2008), 299–316; at http://www.math.bas.bg/complan/fcaa.

[21] B. Toshtemirov. Frankl-type problem for a mixed type equation associated hyper-Bessel differential operator. Montes Taurus J. Pure Appl. Math. 3(3), 2021, pp. 327-333.

[22] I. Dimovski, *Operational calculus of a class of differential operators*. C. R. Acad. Bulg. Sci. 19, No 12 (1966), 1111–1114.

[23] R. Garra, E.Orsingher, F. Polito, *Fractional diffusion with time-varying coefficients*. J. of Math. Phys. 56, No 9 (2015), 1–19.

[24] R. Garra, A. Giusti, F. Mainardi, G. Pagnini, *Fractional relaxation with time-varying coefficient.* Fract. Calc. Appl. Anal., Vol. 17, No 2 (2014), pp. 424-439; DOI: 10.2478/s13540- 014-0178-0

[25] V. Kiryakova, *Generalized Fractional Calculus and Applications.* Longman-J. Wiley, Harlow-N.York (1994).

[26] Fatma Al-Musalhi, Nasser Al-Salti and Erkinjon Karimov. *Initial boundary value problems for fractional differential equation with hyper-Bessel operator*. Fract. Calc. Appl. Anal., Vol. 21, No 1 (2018), pp. 200-219, DOI: 10.1515/fca-2018-0013

[27] P. Agarwal, A. Berdyshev and E. Karimov. *Solvability of a non-local problem with integral form transmitting condition for mixed type equation with Caputo fractional derivative*. Results in Mathematics. 71(3), 2017, pp.1235-1257

[28] E.T.Karimov, B.H.Toshtemirov. Tricomi type problem with integral conjugation condition for a mixed type equation with the hyper-Bessel fractional differential operator, Bulleten of the Institute of Mathematics, 4(1) 9-14, (2019)

[29] E.T.Karimov. *Boundary value problems for parabolic-hyperbolic type equations with spectral parameter.* PhD Thesis, Tashkent, 2006.

[30] S. Kerbal, E. Karimov, N. Rakhmatullayeva. *Non-local boundary problem with integral form transmitting condition for fractional mixed type equation in a composite domain*. Mat. Model. Nat. Phenom. 12(3), 2017, pp. 95-104.

[31] Berdyshev A.S., Eshmatov B. E., Kadirkulov B.J. *Boundary value problems for fourth-order mixed type equation with fractional derivative.* Electronic Journal of Differential Equations, 2016(36), pp. 1-11.

[32] A. V. Pskhu, *Partial Differential Equations of Fractional Order*, Moscow, Nauka. 2005. [in Russian].

[33] G.P. Tolstov, Fourier series (translated by R.A. Silverman), Prentice Hall, Inc., Englewood Cliffs, N. J., (1962)

This figure "test.png" is available in "png" format from:

http://arxiv.org/ps/2111.13407v2