

Robust Permutation Tests in Linear Instrumental Variables Regression*

Purevdorj Tuvaandorj[†]

York University

Abstract

This paper develops permutation versions of identification-robust tests in linear instrumental variables regression. Unlike the existing randomization and rank-based tests in which independence between the instruments and the error terms is assumed, the permutation Anderson-Rubin (AR), Lagrange Multiplier (LM) and Conditional Likelihood Ratio (CLR) tests are asymptotically similar and robust to conditional heteroskedasticity under standard exclusion restriction i.e. the orthogonality between the instruments and the error terms. Moreover, when the instruments are independent of the structural error term, the permutation AR tests are exact, hence robust to heavy tails. As such, these tests share the strengths of the rank-based tests and the wild bootstrap AR tests. Numerical illustrations corroborate the theoretical results.

Keywords: Anderson-Rubin statistic, Asymptotic and exact tests, Conditional likelihood ratio statistic, Heteroskedasticity, Identification, Lagrange Multiplier statistic

*I thank the editor, two anonymous referees, Dmitry Arkhangelsky, Xavier D'Haultfœuille, Antoine Djogbenou, Hiroyuki Kasahara, Uros Petronijevic, and the participants of the Econometric Society European Winter Meeting 2019 and the 37th Canadian Econometrics Study Group (CESG) Meeting 2021 for helpful comments. I gratefully acknowledge financial support from the LA&PS Minor Research Grant, York University. All errors are my own.

[†]tpujee@yorku.ca

1 Introduction

The use of instrumental variables (IVs) is widespread across many disciplines. One of the main inferential issues that require taking careful account in the IV regression is to protect against a possible weak correlation between instruments and endogenous regressors e.g. treatment variable. To this end, it is desirable to use *identification-robust* tests that are *similar*, at least asymptotically, when the instruments have an arbitrary degree of explanatory power (identification strength) for the endogenous regressor, and have good power when the instruments are informative.¹

This paper proposes permutation (randomization) test versions of heteroskedasticity-robust [Anderson and Rubin \(1949\)](#)’s AR, [Kleibergen \(2002\)](#)’s LM and [Andrews and Guggenberger \(2019a\)](#)’s CLR tests in IV regression.² These tests are asymptotically similar (in a uniform sense) and heteroskedasticity-robust under the standard exclusion restriction condition i.e. the orthogonality (or uncorrelatedness) between the error terms and the instruments. When the latter assumption is replaced by independence of the instruments and the error term in the structural equation, the permutation AR tests become exact hence robust to heavy tails.

Permutation inference is attractive because of its exactness when relevant assumptions hold. It also bears a natural connection to IV methods since IVs provide an exogenous variation independent of the unobserved confounders for the endogenous regressor in the IV regression and the permutation inference typically seeks to exploit independence of two sets of variables by permuting the elements of one variable holding the other fixed to replicate the distribution of a test statistic at hand.

Despite the link, the literature on randomization inference in IV models is scarce. [Im-](#)

¹For a comprehensive treatment of similar tests, see Chapter 4 of [Lehmann and Romano \(2005\)](#) and [Linnik \(2008\)](#).

²The LM test is a score test that uses an outer-product-of-the-gradient information matrix estimator, and the CLR test is a heteroskedasticity-robust version of the CLR test proposed by [Moreira \(2003\)](#) which is a LR test with *Neyman structure*, see Chapter 4.3 of [Lehmann and Romano \(2005\)](#) for the latter.

[bens and Rosenbaum \(2005\)](#) develop exact permutation tests in the IV model where the instruments are assigned randomly.³ Their test statistic takes the form of an inner product between the instruments (or transformations thereof e.g. ranks) and the structural error terms evaluated under the null hypothesis. Exact permutation tests are obtained by permuting the former while holding the latter fixed. Given the advantage of the permutation method in an IV setting as exemplified by [Imbens and Rosenbaum \(2005\)](#), we broaden its scope by proposing the permutation versions of the trinity of aforementioned identification-robust test statistics commonly used in practice. Numerical results corroborate the good properties of the proposed permutation tests.

Recently, [DiCiccio and Romano \(2017\)](#) propose permutation tests for correlation between a pair of random vectors and regression coefficients. [DiCiccio and Romano \(2017\)](#) show that using an appropriate studentization argument, one can obtain tests that are asymptotically pivotal under the assumption of zero correlation between the two random variables instead of the usual independence assumption. The results are then generalized to heteroskedastic linear regression where the error term is conditionally mean-independent of the regressors.

We extend the result of [DiCiccio and Romano \(2017\)](#) to the IV setting and propose two permutation AR (PAR) statistics, denoted as PAR_1 and PAR_2 respectively. The PAR_1 statistic is based on permuting the rows of instrument matrix that shifts endogenous regressors, and the PAR_2 statistic is based on permuting the null-restricted residuals in the structural equation. We establish the finite sample validity of the permutation AR tests under an independence assumption akin to the one used in [Andrews and Marmer \(2008\)](#). We then show their asymptotic validity under the usual exogeneity condition allowing for conditional heteroskedasticity. Thus, our result differs from that of [Imbens and Rosenbaum \(2005\)](#) who consider a finite-population framework where the IVs are randomly assigned

³This leads to what is referred to as the design-based approach by [Abadie et al. \(2020\)](#) who make an explicit distinction between design-based and sampling-based uncertainties and propose robust standard errors to account for each uncertainty in a regression model.

and the quantities that do not depend on the IVs are held fixed. In addition, we consider permutation LM (PLM) and CLR (PCLR) statistics. Following [Freedman and Lane \(1983\)](#), we permute the residuals from the first-stage estimation of the reduced-form equation and generate a permutation right-hand side endogenous variables. Based on the latter and the permuted null-restricted residuals in the structural equation, the permutation LM statistic is then constructed. In the PCLR tests, we condition on the nuisance parameter estimates and permute the null-restricted residuals of the structural equation only as in the PAR_2 test to generate the conditional permutation distribution.

We compare the robust permutation tests to both the identification-robust rank-based tests of [Andrews and Marmer \(2008\)](#) and [Andrews and Soares \(2007\)](#), and the wild bootstrap tests of [Davidson and MacKinnon \(2012\)](#) through simulations. In terms of the control of type I error, we find that the robust permutation tests outperform the rank-based tests under the standard exclusion restriction, and perform on par with or have an edge over the wild bootstrap tests in the heteroskedastic designs considered.

As a main technical contribution, we show the uniform asymptotic similarity of the proposed permutation tests and derive their asymptotic power under local alternatives and strong identification which, to our best knowledge, have not been shown before in the context of bootstrap and randomization inference in the IV model. In doing so, we adapt the results of [Andrews and Guggenberger \(2017a,b, 2019a,b\)](#) and [Andrews et al. \(2020\)](#) in a nontrivial way.

Finally, we provide an analysis of the PAR tests with a diverging number of IVs and characterize conditions under which the tests remain asymptotically valid.

Discussion of the related literature

The literature on linear IV model is vast. Earlier surveys on this topic are provided by [Stock et al. \(2002\)](#), [Dufour \(2003\)](#), and [Andrews and Stock \(2007\)](#), and recent contributions include [Moreira and Moreira \(2019\)](#), [Young \(2022\)](#) and the survey of [Andrews et al. \(2019\)](#).

[Andrews and Marmer \(2008\)](#) develop a rank-based AR statistic under the indepen-

dence assumption between the instruments and the structural error term. Under their assumption, the PAR_1 test proposed here is exact while the PAR_2 test is so when there are no included non-constant exogenous variables. [Andrews and Soares \(2007\)](#) consider a rank-version of the CLR statistic of [Moreira \(2003\)](#). The latter test employs asymptotic critical value after conditioning on a rank-based nuisance parameter estimate and is robust to heavy-tailed errors but not to heteroskedasticity.

A strand of literature that focuses on bootstrap inference in the linear IV model include, among others, [Davidson and MacKinnon \(2008\)](#), [Moreira et al. \(2009\)](#), and [Davidson and MacKinnon \(2012\)](#) where they document an improved performance of bootstrap inference over asymptotic inference. The results of this paper complement these studies.

The key for our results is the fact that the identification-robust PAR, PLM and PCLR statistics are studentized. The use of studentization to obtain an improved test is not new; a well-known example is the higher-order accuracy of the bootstrap- t confidence interval, as opposed to the percentile bootstrap, in the one-sample problem, see, for instance, Chapter 15.5 of [Lehmann and Romano \(2005\)](#). [So and Shin \(1999\)](#) develop a persistence-robust test for autoregressive models based on a Cauchy estimator and a studentized statistic. In the context of permutation tests, [Neuhaus \(1993\)](#) proposes a studentized permutation statistic in a two-sample problem with randomly censored data. For further extensions and applications of permutation tests based on studentized statistic, see [Janssen \(1997\)](#), [Neubert and Brunner \(2007\)](#), [Pauly \(2011\)](#) and [Chung and Romano \(2013, 2016\)](#).

Several recent papers that consider randomization inference in different contexts include [Chernozhukov et al. \(2009\)](#), [Canay et al. \(2017\)](#), [Bugni et al. \(2018\)](#), [Canay and Kamat \(2018\)](#), [Ganong and Jäger \(2018\)](#) and [Dufour et al. \(2019\)](#).

The paper is organized as follows. Section 2 introduces the model and the identification-robust test statistics and develops the permutation tests. Section 3 provides simulation results comparing the performance of alternative test procedures. Section 4 presents an empirical application. We briefly conclude in Section 5. Supplemental Appendix contains

all the proofs of our theoretical results and additional simulation evidence. We also provide the replication R codes of the empirical applications.

2 Robust permutation tests

2.1 Model setup

We develop permutation-based tests for a restriction on the $d \times 1$ vector of structural coefficients

$$H_0 : \theta = \theta_0, \quad (2.1)$$

in the linear IV regression model:

$$y = Y\theta + X\gamma + u, \quad (2.2)$$

$$Y = W\Gamma + X\Psi + V, \quad (2.3)$$

where $y = [y_1, \dots, y_n]'$ is $n \times 1$ vector of endogenous variables, $Y = [Y_1, \dots, Y_n]'$ is $n \times d$ matrix of right-hand side endogenous variables, $X = [X_1, \dots, X_n]'$ is $n \times p$ matrix of right-hand side exogenous variables whose first column is the $n \times 1$ vector of ones $\iota = [1, \dots, 1]'$, $W = [W_1, \dots, W_n]'$ is $n \times k$ ($k \geq d$) matrix of instrumental variables that does not include ι and the exogenous variables in X , $u = [u_1, \dots, u_n]'$ is $n \times 1$ vector of structural error terms, $V = [V_1, \dots, V_n]'$ with $V_i = [V_{i1}, \dots, V_{id}]'$, is $n \times d$ matrix of reduced-form error terms; γ is $p \times 1$ parameter vector, and Γ and Ψ are $k \times d$ and $p \times d$ matrices of reduced-form coefficients, respectively.

Let $X_i = [X_{i1}, \dots, X_{ip}]'$, $W_i = [W_{i1}, \dots, W_{ik}]'$, $Y_i = [Y_{i1}, \dots, Y_{id}]'$ and $Z = [Z_1, \dots, Z_n]' \equiv M_X W$, where $M_A \equiv I_n - P_A \equiv I_n - A(A'A)^{-1}A'$ for a matrix A of full column rank. Rewrite the equations (2.2)-(2.3) as

$$y = Y\theta + X\gamma + u, \quad (2.4)$$

$$Y = Z\Gamma + X\xi + V, \quad (2.5)$$

where $\xi \equiv (X'X)^{-1}X'W\Gamma + \Psi$. Let $Y = [Y_1, \dots, Y_d]$, where $Y_s = [Y_{1s}, \dots, Y_{ns}]'$ denotes the s -th column of Y , $s = 1, \dots, d$. Define $\tilde{u}_i(\theta) \equiv y_i - Y_i'\theta - X_i'(X'X)^{-1}X'(y - Y\theta)$, $\tilde{u}(\theta) = M_X(y - Y\theta) = [\tilde{u}_1(\theta), \dots, \tilde{u}_n(\theta)]'$, $\tilde{Y} = [\tilde{Y}_1, \dots, \tilde{Y}_n]'$ $\equiv M_X Y$, and

$$\hat{m}(\theta) \equiv n^{-1}Z'\tilde{u}(\theta) = n^{-1}Z'(y - Y\theta), \quad (2.6)$$

$$\hat{G}_s(\theta) \equiv n^{-1}Z'Y_s, \quad \hat{C}_s(\theta) \equiv n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{Y}_{is} \tilde{u}_i(\theta), \quad (2.7)$$

$$\hat{\Sigma}(\theta) \equiv n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{u}_i(\theta)^2, \quad \hat{J}(\theta) = [\hat{J}_1(\theta), \dots, \hat{J}_d(\theta)], \quad (2.8)$$

$$\hat{J}_s(\theta) \equiv \hat{G}_s(\theta) - \hat{C}_s(\theta) \hat{\Sigma}(\theta)^{-1} \hat{m}(\theta), \quad s = 1, \dots, d, \quad (2.9)$$

where \tilde{Y}_{is} is the s -th element of \tilde{Y}_i . The heteroskedasticity-robust AR test statistic is defined as

$$\text{AR} = \text{AR}(W, X, \tilde{u}(\theta_0)) \equiv n \hat{m}(\theta_0)' \hat{\Sigma}(\theta_0)^{-1} \hat{m}(\theta_0).$$

The null asymptotic distribution of the AR statistic is chi-square regardless of the values of Γ because $n^{-1/2}Z'(y - Y\theta_0)$ is asymptotically normal under (2.1). Another asymptotically pivotal test statistic is the following heteroskedasticity-robust version of Kleibergen (2002)'s LM statistic

$$\text{LM} = \text{LM}(Z, \tilde{u}(\theta_0), Y) \equiv n \hat{m}(\theta_0)' \hat{\Sigma}(\theta_0)^{-1} \hat{J}(\theta_0) \left(\hat{J}(\theta_0)' \hat{\Sigma}(\theta_0)^{-1} \hat{J}(\theta_0) \right)^{-1} \hat{J}(\theta_0)' \hat{\Sigma}(\theta_0)^{-1} \hat{m}(\theta_0). \quad (2.10)$$

As defined, $\hat{J}_s(\theta_0)$ in (2.9) is a vector of residuals from the regression of the sample Jacobian in (2.7) on the sample moment function in (2.6). When Γ has full rank i.e. the identifying power of W is strong, $\hat{J}(\theta_0)$ converges to a matrix of full rank under mild conditions, and the LM statistic is asymptotically chi-squared under (2.1). However, even when Γ is “small” so that the noise and signal parts of Y are of a similar magnitude or the former dominates the latter, $\hat{J}(\theta_0)$ and $\hat{m}(\theta_0)$ are asymptotically independent after suitable normalizations, and consequently, the LM statistic is still chi-squared in the limit.

The CLR-type test statistics considered next have an approximate Neyman structure such that their conditional asymptotic distributions given sufficient statistics for the matrix

of nuisance parameters Γ are independent of Γ . Define

$$\hat{\mathcal{S}} \equiv \hat{\Sigma}(\theta_0)^{-1/2} n^{1/2} \hat{m}(\theta_0) \in \mathbb{R}^k, \quad (2.11)$$

$$T(\hat{\Sigma}(\theta_0), \hat{J}(\theta_0), \tilde{\mathcal{V}}) \equiv \hat{\Sigma}(\theta_0)^{-1/2} n^{1/2} \hat{J}(\theta_0) \left((\theta_0, I_d) \Omega^\epsilon(\hat{\Sigma}(\theta_0), \tilde{\mathcal{V}})^{-1} (\theta_0, I_d)' \right)^{1/2} \in \mathbb{R}^{k \times d}, \quad (2.12)$$

where $\Omega^\epsilon(\cdot, \cdot) \in \mathbb{R}^{(d+1) \times (d+1)}$ is an eigenvalue-adjusted version of the matrix defined as

$$\Omega(\hat{\Sigma}(\theta), \tilde{\mathcal{V}}) = \{\Omega_{ij}\}_{1 \leq i, j \leq d+1} \in \mathbb{R}^{(d+1) \times (d+1)}, \quad \Omega_{ij} \equiv \text{tr}(K_{ij}(\tilde{\mathcal{V}})' \hat{\Sigma}(\theta)^{-1}) / k \in \mathbb{R},^4 \quad (2.13)$$

and the $k \times k$ matrix $K_{ij}(\tilde{\mathcal{V}})$ is the (i, j) submatrix of $K(\tilde{\mathcal{V}})$ given by

$$K(\tilde{\mathcal{V}}) \equiv (B(\theta_0)' \otimes I_k) \tilde{\mathcal{V}} (B(\theta_0) \otimes I_k) \in \mathbb{R}^{k(d+1) \times k(d+1)}, \quad B(\theta) \equiv \begin{bmatrix} 1 & 0_{1 \times d} \\ -\theta & -I_d \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}. \quad (2.14)$$

With a suitable choice of $\tilde{\mathcal{V}}$, the transformation in (2.12) yields an asymptotically sufficient statistic for Γ that is independent of the statistic $\hat{\mathcal{S}}$ in (2.11). The CLR-type statistics proposed by [Andrews and Guggenberger \(2019a,b\)](#) are

$$\text{CLR}_a = \text{CLR}(\hat{\mathcal{S}}, \hat{\mathcal{T}}_a) \equiv \hat{\mathcal{S}}' \hat{\mathcal{S}} - \lambda_{\min} \left[(\hat{\mathcal{S}}, \hat{\mathcal{T}}_a)' (\hat{\mathcal{S}}, \hat{\mathcal{T}}_a) \right], \quad (2.15)$$

$$\text{CLR}_b = \text{CLR}(\hat{\mathcal{S}}, \hat{\mathcal{T}}_b) \equiv \hat{\mathcal{S}}' \hat{\mathcal{S}} - \lambda_{\min} \left[(\hat{\mathcal{S}}, \hat{\mathcal{T}}_b)' (\hat{\mathcal{S}}, \hat{\mathcal{T}}_b) \right], \quad (2.16)$$

where $\lambda_{\min}(\cdot)$ denotes the smallest eigenvalue of a matrix, and

$$\hat{\mathcal{T}}_a = T(\hat{\Sigma}(\theta_0), \hat{J}(\theta_0), \hat{\mathcal{V}}_a(\theta_0)), \quad \hat{\mathcal{V}}_a(\theta) \equiv \begin{bmatrix} \hat{\Sigma}(\theta) & \hat{C}(\theta) \\ \hat{C}(\theta)' & \hat{\Sigma}^G \end{bmatrix} \in \mathbb{R}^{k(d+1) \times k(d+1)}, \quad (2.17)$$

$$\hat{C}(\theta) \equiv [\hat{C}_1(\theta), \dots, \hat{C}_d(\theta)] \in \mathbb{R}^{k \times kd},$$

$$\hat{\Sigma}^G = \{\hat{\Sigma}_{st}^G\}_{1 \leq s, t \leq d} \in \mathbb{R}^{kd \times kd}, \quad \hat{\Sigma}_{st}^G \equiv n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{Y}_{is} \tilde{Y}_{it} - \left(n^{-1} \sum_{i=1}^n Z_i \tilde{Y}_{is} \right) \left(n^{-1} \sum_{i=1}^n Z_i \tilde{Y}_{it} \right)' \in \mathbb{R}^{k \times k}, \quad (2.18)$$

$$\begin{aligned}
\hat{\mathcal{T}}_b &= T(\hat{\Sigma}(\theta_0), \hat{J}(\theta_0), \hat{\mathcal{V}}_b(\theta_0)), \\
\hat{\mathcal{V}}_b(\theta) &\equiv n^{-1} \sum_{i=1}^n [(\varepsilon_i(\theta) - \hat{\varepsilon}_{in}(\theta))(\varepsilon_i(\theta) - \hat{\varepsilon}_{in}(\theta))'] \otimes (Z_i Z_i') \in \mathbb{R}^{k(d+1) \times k(d+1)}, \quad (2.19) \\
\varepsilon_i(\theta) &\equiv [\tilde{u}_i(\theta), -\tilde{Y}_i']' \in \mathbb{R}^{d+1}, \quad \tilde{Y} = [\tilde{Y}_1, \dots, \tilde{Y}_n]' = M_X Y \in \mathbb{R}^{n \times d}, \\
\hat{\varepsilon}_{in}(\theta) &\equiv \varepsilon(\theta)' Z (Z' Z)^{-1} Z_i \in \mathbb{R}^{d+1}, \quad \varepsilon(\theta) \equiv [\varepsilon_1(\theta), \dots, \varepsilon_n(\theta)]' \in \mathbb{R}^{n \times (d+1)}. \quad (2.20)
\end{aligned}$$

The CLR_a and CLR_b statistics are denoted as QLR and QLR_P , respectively, in [Andrews and Guggenberger \(2019b\)](#), and differ only in their definitions of $\hat{\mathcal{V}}_a(\theta_0)$ and $\hat{\mathcal{V}}_b(\theta_0)$. [Andrews and Guggenberger \(2019b\)](#) show that the CLR_b statistic is asymptotically equivalent to [Moreira \(2003\)](#)'s CLR statistic in the homoskedastic linear IV regression with fixed instruments and multiple endogenous variables under all strengths of identification while the CLR_a statistic has the same property under only certain strengths of identification but its general version is more broadly applicable.

2.2 Main results

We begin by recalling the basic notion of randomization tests from Chapter 15.2 of [Lehmann and Romano \(2005\)](#). Let \mathbb{G}_n be the set of all permutations $\pi = [\pi(1), \dots, \pi(n)]$ of $\{1, \dots, n\}$, and denote a permutation version of a generic test statistic R by $\text{PR} = R^\pi$. The distribution function corresponding to permutations in \mathbb{G}_n is

$$\hat{F}_n^{\text{PR}}(x) \equiv \frac{1}{n!} \sum_{\pi \in \mathbb{G}_n} 1(R^\pi \leq x), \quad (2.21)$$

⁴The eigenvalue-adjustment of [Andrews and Guggenberger \(2019b\)](#) is as follows. Let A be a nonzero positive semi-definite matrix of dimension $p \times p$ that has a spectral decomposition $A = N \Delta N'$, where $\Delta = \text{diag}(\lambda_1, \dots, \lambda_p)$, $\lambda_1 \geq \dots \geq \lambda_p \geq 0$, is the diagonal matrix that consists of the eigenvalues of A , and N is an orthogonal matrix of the corresponding eigenvectors. Given a constant $\epsilon > 0$, the eigenvalue adjusted matrix is defined as $A^\epsilon \equiv N \Delta^\epsilon N'$, where $\Delta^\epsilon \equiv \text{diag}(\max\{\lambda_1, \lambda_1 \epsilon\}, \dots, \max\{\lambda_p, \lambda_p \epsilon\})$. [Andrews and Guggenberger \(2019a\)](#) recommend $\epsilon = 0.01$. We refer to [Andrews and Guggenberger \(2019a\)](#) for further properties of the eigenvalue-adjustment procedure.

where $1(\cdot)$ is the indicator function. The computation of (2.21) and related quantities (at a particular value of x) for permutation tests entails computing $n!$ test statistics which is impractical even for n relatively small. So instead, a stochastic approximation based on uniform random draws from \mathbb{G}_n is often used. To this end, let us fix $N \in \{1, \dots, |n!|\}$ and consider a subset of permutations $\mathbb{G}'_n = \{\pi_1, \dots, \pi_N\} \subseteq \mathbb{G}_n$, where π_1 is the identity permutation, and π_2, \dots, π_N are i.i.d. uniformly distributed on \mathbb{G}_n . The permutation distribution function corresponding to \mathbb{G}'_n is

$$\tilde{F}_N^{\text{PR}}(x) \equiv \frac{1}{N} \sum_{\pi \in \mathbb{G}'_n} 1(R^\pi \leq x). \quad (2.22)$$

Let $R_{(1)}^\pi \leq \dots \leq R_{(N)}^\pi$ be the order statistics of $\{R^\pi : \pi \in \mathbb{G}'_n\}$, and for a nominal significance level $\alpha \in (0, 1)$, define

$$r \equiv N - I(N\alpha), \quad (2.23)$$

$$\begin{aligned} N^+ &\equiv |\{j = 1, \dots, N : R_{(j)}^\pi > R_{(r)}^\pi\}|, \quad N^0 \equiv |\{j = 1, \dots, N : R_{(j)}^\pi = R_{(r)}^\pi\}|, \\ a^{\text{PR}} &\equiv (N\alpha - N^+)/N^0, \end{aligned} \quad (2.24)$$

where $I(\cdot)$ denotes the integer part of a number. Clearly, N^+ and N^0 are the number of values $R_{(j)}^\pi = \text{PR}_{(j)}, j = 1, \dots, N$, that are greater than $R_{(r)}^\pi$ and equal to $R_{(r)}^\pi$, respectively.

A randomization test is then defined as

$$\phi_n^{\text{PR}} \equiv \begin{cases} 1 & \text{if } R > R_{(r)}^\pi, \\ a^{\text{PR}} & \text{if } R = R_{(r)}^\pi, \\ 0 & \text{if } R < R_{(r)}^\pi. \end{cases} \quad (2.25)$$

The finite sample results presented below hold for any N but we will assume that $N \rightarrow \infty$ as $n \rightarrow \infty$ for the asymptotic results. We next describe the robust permutation test statistics.

Let $W_\pi = [W_{\pi(1)}, \dots, W_{\pi(n)}]'$ and $\tilde{u}_\pi(\theta) = [\tilde{u}_{\pi(1)}(\theta), \dots, \tilde{u}_{\pi(n)}(\theta)]'$ be an instrument matrix and a residual vector obtained by permuting the rows of W and the elements of $\tilde{u}(\theta)$, respectively, for a permutation $\pi \in \mathbb{G}'_n$. Set

$$\tilde{\Sigma}_u(\theta) \equiv \text{diag}(\tilde{u}_1(\theta)^2, \dots, \tilde{u}_n(\theta)^2), \quad \hat{m}^\pi(\theta) \equiv n^{-1} Z' \tilde{u}_\pi(\theta), \quad \hat{\Sigma}^\pi(\theta) \equiv n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{u}_{\pi(i)}(\theta)^2.$$

We consider two heteroskedasticity-robust permutation AR statistics defined as:

$$\text{PAR}_1 \equiv \text{AR}(W_\pi, X, \tilde{u}(\theta_0)) = \tilde{u}(\theta_0)' M_X W_\pi (W_\pi' M_X \tilde{\Sigma}_u(\theta_0) M_X W_\pi)^{-1} W_\pi' M_X \tilde{u}(\theta_0), \quad (2.26)$$

$$\text{PAR}_2 \equiv \text{AR}(W, X, \tilde{u}_\pi(\theta_0)) = n \hat{m}^\pi(\theta_0)' \hat{\Sigma}^\pi(\theta_0)^{-1} \hat{m}^\pi(\theta_0). \quad (2.27)$$

The PAR_1 statistic is based on the permutation of the rows of the instrument matrix W , and the PAR_2 statistic uses the permutation of the null-restricted residuals $\tilde{u}(\theta_0)$. The finite sample validity of these tests are shown under the following condition.

Assumption 1. $\{(W'_i, X'_i, u_i)'\}_{i=1}^n$ are i.i.d., and W_i and $[X'_i, u_i]'$ are independently distributed.

Andrews and Marmer (2008) develop exact rank-based AR tests under assumptions similar to Assumption 1. When W is independent of X and u , $W' M_X \tilde{u}(\theta_0)$ and $W'_\pi M_X \tilde{u}(\theta_0)$ have the same distribution under $H_0 : \theta = \theta_0$, so the PAR_1 test is exact. On the other hand, the PAR_2 test is not, in general, exact because $W' M_X \tilde{u}(\theta_0)$ and $W'_\pi M_X \tilde{u}_\pi(\theta_0)$ may not have identical distributions. However, when $X = \iota$ in Assumption 1, Z and u_π are independent, $Z' \tilde{u}(\theta_0)$ and $Z' \tilde{u}_\pi(\theta_0)$ are identically distributed, and consequently, the PAR_2 test is exact. The following result summarizes the finite sample validity of the PAR tests.

Proposition 2.1 (Finite sample validity). *Under Assumption 1 and $H_0 : \theta = \theta_0$, $E[\phi_n^{\text{PAR}_1}] = \alpha$ for $\alpha \in (0, 1)$. If $X = \iota$ in Assumption 1, then $E[\phi_n^{\text{PAR}_2}] = \alpha$.*

Even if $[X'_i, u_i]'$ and W_i are not distributed independently but satisfy the orthogonality condition $E[(W'_i, X'_i)' u_i] = 0$, we show that the permutation AR tests are asymptotically similar and heteroskedasticity-robust.

The argument for permutation LM statistic is more involved because it is difficult to construct an estimator of the reduced-form coefficients Γ whose (asymptotic) distribution remains invariant to a permutation of the data. The OLS estimators of the reduced-form coefficients and the corresponding residuals are

$$\hat{\Gamma} = (Z'Z)^{-1} Z'Y, \quad \hat{\xi} = (X'X)^{-1} X'Y, \quad \hat{V} = Y - Z\hat{\Gamma} - X\hat{\xi} = [\hat{V}_1, \dots, \hat{V}_n]' = [\hat{V}_{1,1}, \dots, \hat{V}_{1,d}],$$

where $\hat{V}_{\cdot s} = [\hat{V}_{1s}, \dots, \hat{V}_{ns}]'$ is the s -th column of \hat{V} , $s = 1, \dots, d$. Now permute the residuals $\hat{V}_\pi = [\hat{V}'_{\pi(1)}, \dots, \hat{V}'_{\pi(n)}]'$, and let

$$Y^\pi = [Y_{\cdot 1}^\pi, \dots, Y_{\cdot d}^\pi] \equiv Z\hat{\Gamma} + X\hat{\xi} + \hat{V}_\pi,$$

where $Y_{\cdot s}^\pi = [Y_{1s}^\pi, \dots, Y_{ns}^\pi]'$ denotes the s -th column of Y^π . The idea of permuting the residuals to obtain asymptotically valid randomization tests appears in [Freedman and Lane \(1983\)](#) and [DiCiccio and Romano \(2017\)](#).

The Jacobian estimator used in the permutation LM statistic is

$$\hat{J}^\pi(\theta) = [\hat{J}_1^\pi(\theta), \dots, \hat{J}_d^\pi(\theta)], \quad (2.28)$$

$$\hat{J}_s^\pi(\theta) \equiv n^{-1} Z' Y_{\cdot s}^\pi - \hat{C}_s^\pi(\theta) \hat{\Sigma}^\pi(\theta)^{-1} \hat{m}^\pi(\theta), \quad (2.29)$$

$$\hat{C}_s^\pi(\theta) \equiv n^{-1} \sum_{i=1}^n Z_i Z_i' \hat{V}_{\pi(i)s} \tilde{u}_{\pi(i)}(\theta), \quad s = 1, \dots, d. \quad (2.30)$$

It is not difficult to see that in strongly identified models where Γ is a fixed matrix of full rank, $\hat{J}^\pi(\theta_0) - n^{-1} Z' Z \Gamma \xrightarrow{p^\pi} 0$ in probability, where $\xrightarrow{p^\pi}$ denotes the convergence in probability with respect to the probability measure of π uniformly distributed over \mathbb{G}_n conditional on the data. The permutation LM statistic is then defined as

$$\text{PLM} \equiv \text{LM}(Z, \tilde{u}_\pi(\theta_0), Y^\pi) = n \hat{m}^\pi(\theta_0)' \hat{\Sigma}^\pi(\theta_0)^{-1/2} P_{\hat{\Sigma}^\pi(\theta_0)^{-1/2} \hat{J}^\pi(\theta_0)} \hat{\Sigma}^\pi(\theta_0)^{-1/2} \hat{m}^\pi(\theta_0). \quad (2.31)$$

Finally, we define the permutation CLR statistics as follows:

$$\text{PCLR}_a \equiv \text{CLR}(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}_a) = \hat{\mathcal{S}}^{\pi'} \hat{\mathcal{S}}^\pi - \lambda_{\min} \left[(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}_a)' (\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}_a) \right], \quad (2.32)$$

$$\text{PCLR}_b \equiv \text{CLR}(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}_b) = \hat{\mathcal{S}}^{\pi'} \hat{\mathcal{S}}^\pi - \lambda_{\min} \left[(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}_b)' (\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}_b) \right], \quad (2.33)$$

where $\hat{\mathcal{T}}_a$ and $\hat{\mathcal{T}}_b$ are as defined in (2.17) and (2.19), and

$$\hat{\mathcal{S}}^\pi \equiv n^{1/2} \hat{\Sigma}^\pi(\theta_0)^{-1/2} \hat{m}^\pi(\theta_0) = \left(\sum_{i=1}^n Z_i Z_i' \tilde{u}_{\pi(i)}(\theta_0)^2 \right)^{-1/2} Z' \tilde{u}_\pi(\theta_0). \quad (2.34)$$

Analogously to (2.21) and (2.22), define the permutation distributions of the statistic $\text{PCLR} = \text{CLR}(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}) \in \{\text{PCLR}_a, \text{PCLR}_b\}$ given $\hat{\mathcal{T}} \in \{\hat{\mathcal{T}}_a, \hat{\mathcal{T}}_b\}$, respectively, as $\hat{F}_n^{\text{PCLR}}(x, \hat{\mathcal{T}}) \equiv$

$(n!)^{-1} \sum_{\pi \in \mathbb{G}_n} 1(\text{CLR}(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}) \leq x)$ and

$$\tilde{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) \equiv N^{-1} \sum_{\pi \in \mathbb{G}'_n} 1(\text{CLR}(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}) \leq x). \quad (2.35)$$

Let $\text{PCLR}_{(r)}(\hat{\mathcal{T}})$ be the r -th order statistic of $\{\text{CLR}(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}}) : \pi \in \mathbb{G}'_n\}$ for r defined in (2.23), and a^{PCLR} be as in (2.24). The nominal level α PCLR test ϕ_n^{PCLR} is 1 if $\text{CLR}(\hat{\mathcal{S}}, \hat{\mathcal{T}}) > \text{PCLR}_{(r)}(\hat{\mathcal{T}})$, a^{PCLR} if $\text{CLR}(\hat{\mathcal{S}}, \hat{\mathcal{T}}) = \text{PCLR}_{(r)}(\hat{\mathcal{T}})$ and 0 otherwise.

Let P denote the distribution of the vector $(W'_i, X'_i, u_i, V'_i)'$ and we shall index the relevant quantities by P in the sequel. Define

$$e_i \equiv [u_i, V'_i]', \quad \Sigma_P^e \equiv \text{Var}_P[e_i] = \begin{bmatrix} \sigma_P^2 & \Sigma_P^{uV} \\ \Sigma_P^{Vu} & \Sigma_P^V \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}, \quad (2.36)$$

$$Z_i^* \equiv W_i - \text{E}_P[W_i X'_i] (\text{E}_P[X_i X'_i])^{-1} X_i \in \mathbb{R}^k,$$

$$\mathcal{V}_{P,a} \equiv \text{Var}_P \begin{bmatrix} Z_i^* u_i \\ \text{vec}(Z_i^* (Z_i^{*'} \Gamma + V'_i)) \end{bmatrix} = \begin{bmatrix} \Sigma_P & C_P \\ C'_P & \Sigma_P^G \end{bmatrix} \in \mathbb{R}^{k(d+1) \times k(d+1)}, \quad \Sigma_P \equiv \text{E}_P[Z_i^* Z_i^{*'} u_i^2] \in \mathbb{R}^{k \times k},$$

$$C_P = [C_{P1}, \dots, C_{Pd}] \in \mathbb{R}^{k \times kd}, \quad C_{Ps} \equiv \text{E}_P[Z_i^* Z_i^{*'} u_i (Z_i^{*'} \Gamma_s + V_{is})] \in \mathbb{R}^{k \times k}, \quad s = 1, \dots, d,$$

$$\Sigma_P^G = \{\Sigma_{P,st}^G\}, \quad \Sigma_{P,st}^G \equiv \text{E}_P[Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma_s + V_{is})(Z_i^{*'} \Gamma_t + V_{it})] - \text{E}_P[Z_i^* Z_i^{*'} \Gamma_s] \text{E}_P[Z_i Z_i^{*'} \Gamma_t'],$$

$$s, t = 1, \dots, d, \quad (2.37)$$

with Γ_s denoting the s -th column of Γ , and

$$\mathcal{V}_{P,b} \equiv \text{E}_P \left[\begin{pmatrix} u_i^2 & -u_i V'_i \\ -u_i V_i & V_i V'_i \end{pmatrix} \otimes (Z_i^* Z_i^{*'}) \right] \in \mathbb{R}^{k(d+1) \times k(d+1)}. \quad (2.38)$$

We maintain the following assumptions to develop the asymptotic results.

Assumption 2. For some $\delta, \delta_1 > 0$ and $M_0 < \infty$,

(a) $\{(W'_i, X'_i, u_i, V'_i)'\}_{i=1}^n$ are i.i.d. with distribution P ,

(b) $\text{E}_P[u_i(W'_i, X'_i)] = 0$,

(c) $\text{E}_P[\| (W'_i, X'_i, u_i)' \|^{4+\delta}] < M_0$,

$$(d) \lambda_{\min}(A) \geq \delta_1 \text{ for } A \in \{E_P[(W'_i, X'_i)'(W'_i, X'_i)], E_P[Z_i^* Z_i^{*'}], \Sigma_P, \sigma_P^2\},$$

$$(e) E_P[V_i(W'_i, X'_i)] = 0,$$

$$(f) E_P[\|V_i\|^{4+\delta}] < M_0,$$

$$(g) \lambda_{\min}(A) \geq \delta_1 \text{ for } A \in \{\Sigma_P^V - \Sigma_P^V (\sigma_P^2)^{-1} \Sigma_P^V, \mathcal{V}_{P,a}\}.$$

Next we define the parameter space for the permutation tests.

$$\mathcal{P}_0^{\text{PAR}_1} = \mathcal{P}_0^{\text{PAR}_2} \equiv \{P : \text{Assumption 2(a)-(d) hold}\}, \quad (2.39)$$

$$\mathcal{P}_0^{\text{PLM}} \equiv \{P : \text{Assumption 2 holds}\}, \quad (2.40)$$

$$\mathcal{P}_0^{\text{PCLR}} \equiv \{P : \text{Assumption 2(a)-(f) hold}\}, \text{ PCLR} \in \{\text{PCLR}_a, \text{PCLR}_b\}. \quad (2.41)$$

Assumption 2 does not impose virtually any restriction on the degree of association between Y and W (after controlling for the effect of the exogenous covariates), thus allowing for an arbitrary identification strength on the part of the instruments.

The distribution P in Assumption 2(a) is allowed to vary with the sample size i.e. $P = P_n$. For simplicity, we suppress the dependence on n . Assumption 2(c) and (f) impose finite $4 + \delta$ moment on the error terms and the exogenous variables, and are slightly stronger than the cross moment restrictions used in Andrews and Guggenberger (2019a). For pointwise asymptotic results or when P does not vary with n , weaker restrictions would be sufficient.

The independence assumption between the instruments and the error terms, which is typically made in the randomization test literature (see, for example, Imbens and Rosenbaum (2005)), is not maintained here; the instruments are only assumed to satisfy the standard exogeneity conditions in Assumption 2(b) and (e). If independence assumptions are maintained, as shown in Proposition 2.1, the PAR tests are exact and do not require the finite $4 + \delta$ moment assumptions.

Assumption 2(d) and (g) require that the second moment matrices of the instruments and exogenous covariates, and covariance matrices are (uniformly) nonsingular, and are

similar to the assumptions employed by [Andrews and Guggenberger \(2017a, 2019a\)](#) and [Andrews et al. \(2020\)](#) when developing identification-robust (but not singularity-robust) AR, LM and CLR tests. These conditions can be restrictive but can potentially be relaxed as in [Andrews and Guggenberger \(2019a\)](#) (see also [Dufour and Taamouti \(2007\)](#)) using generalized inverses of the relevant matrices. However, doing so would entail a considerable complication in the proof, and so, for simplicity, we maintain Assumption 2(d) and (g).

The condition $\lambda_{\min}(\mathcal{V}_{P,a}) \geq \delta_1$ in Assumption 2(g) plays an important role for showing the asymptotic similarity of the LM test, and it can be substituted by the condition $\lambda_{\min}(E_P[(e_i e_i') \otimes (Z_i^* Z_i^{*'})]) \geq \delta_1$, see Section 3.2 of [Andrews and Guggenberger \(2017a\)](#). The condition $\lambda_{\min}(\Sigma_P^V - \Sigma_P^{Vu}(\sigma_P^2)^{-1}\Sigma_P^{uV}) \geq \delta_1$ is used in the asymptotic distribution of the PLM statistic.

The main result of this paper is given in the following theorem which establishes the uniform asymptotic validity of the robust permutation tests.

Theorem 2.2 (Asymptotic similarity). *Suppose that $H_0 : \theta = \theta_0$ holds and $N \rightarrow \infty$ as $\rightarrow \infty$. Then, for $\alpha \in (0, 1)$ and the permutation statistics*

$$\text{PR} \in \{\text{PAR}_1, \text{PAR}_2, \text{PLM}, \text{PCLR}_a, \text{PCLR}_b\}$$

with the corresponding parameter spaces $\mathcal{P}_0 \in \{\mathcal{P}_0^{\text{PAR}_1}, \mathcal{P}_0^{\text{PAR}_2}, \mathcal{P}_0^{\text{PLM}}, \mathcal{P}_0^{\text{PCLR}_a}, \mathcal{P}_0^{\text{PCLR}_b}\}$, respectively,

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_0} E_P[\phi_n^{\text{PR}}] = \liminf_{n \rightarrow \infty} \inf_{P \in \mathcal{P}_0} E_P[\phi_n^{\text{PR}}] = \alpha.$$

Theorem 2.2 shows that the permutation tests are asymptotically similar (hence asymptotically of correct size) i.e. the asymptotic null rejection probability is equal to the nominal significance over a large class of data generating processes. Thus, the advantage of the PAR_1 and PAR_2 tests relative to the existing tests is that they inherit the desirable properties of both the exact and asymptotic AR tests (including other simulation-based tests that are asymptotically valid); the PAR tests are exact under the same assumptions as the rank-based tests are, and asymptotically pivotal and heteroskedasticity-robust under the assumptions used to show the validity of the asymptotic tests.

The main technical tools for the asymptotics of the permutation statistics are the combinatorial central limit theorem based on a Lindeberg type condition (Motoo (1956)) and a variance formula (the equation (10) of Hoeffding (1951) and Lemma S.3.4 of DiCiccio and Romano (2017)) for the sums

$$n^{-1/2} \sum_{i=1}^n Z_i \tilde{u}_{\pi(i)}(\theta_0), \quad n^{-1/2} \sum_{i=1}^n Z_i \hat{V}_{\pi(i)s}, \quad s = 1, \dots, d. \quad (2.42)$$

The fact that the intercept term ι is included in X in (2.2)-(2.3) implies that the two quantities in (2.42) have mean zero with respect to the distribution of π , which, in turn, plays an important role for the asymptotic validity of the permutation tests (see also Remark 3.1 of DiCiccio and Romano (2017)).

The proof of asymptotic similarity of the permutation tests uses the generic method of Andrews et al. (2020) for establishing the asymptotic size of tests; in particular, the derivation of the asymptotic distribution of the PCLR statistics relies to a great extent on Theorems 9.1 and 16.6 of Andrews and Guggenberger (2019a,b).

Next, we shall derive the asymptotic distribution of the permutation statistics under the local alternatives $\theta_n = \theta_0 + h_\theta n^{-1/2}$ with $h_\theta \in \mathbb{R}^d$ fixed, assuming strong identification. Let $\chi_l^2(\eta^2)$ denote noncentral chi-square random variable with degrees of freedom l and noncentrality parameter η^2 .

Proposition 2.3 (Asymptotic local power under strong identification). *Let Assumption 2 hold, and assume that the model is strongly or semi-strongly identified in the sense that the smallest singular value τ_{dn} of $\Sigma_P^{-1/2} G_P ((\theta_0, I_d) \Omega^\epsilon (\Sigma_P, \mathcal{V}_P)^{-1} (\theta_0, I_d)')^{1/2}$, where $G_P \equiv E_P[Z_i^* Z_i^{*'}] \Gamma$ and $\mathcal{V}_P \in \{\mathcal{V}_{P,a}, \mathcal{V}_{P,b}\}$, satisfies $n^{1/2} \tau_{dn} \rightarrow \infty$ as $n \rightarrow \infty$. In addition, assume that $G = \lim_{n \rightarrow \infty} G_P$ and $\Sigma = \lim_{n \rightarrow \infty} \Sigma_P$ exist, and $N \rightarrow \infty$ as $n \rightarrow \infty$. Then, under $H_1 : \theta_n = \theta_0 + h_\theta n^{-1/2}$, $AR \xrightarrow{d} \chi_k^2(\eta^2)$, $LM \xrightarrow{d} \chi_d^2(\eta^2)$, $CLR_a \xrightarrow{d} \chi_d^2(\eta^2)$ and $CLR_b \xrightarrow{d}$*

$\chi_d^2(\eta^2)$, where $\eta^2 \equiv h'_\theta G' \Sigma^{-1} G h_\theta$, and

$$\sup_{x \in \mathbb{R}} |\tilde{F}_N^{\text{PAR}_1}(x) - P[\chi_k^2 \leq x]| \xrightarrow{p} 0, \quad (2.43)$$

$$\sup_{x \in \mathbb{R}} |\tilde{F}_N^{\text{PAR}_2}(x) - P[\chi_k^2 \leq x]| \xrightarrow{p} 0, \quad (2.44)$$

$$\sup_{x \in \mathbb{R}} |\tilde{F}_N^{\text{PLM}}(x) - P[\chi_d^2 \leq x]| \xrightarrow{p} 0, \quad (2.45)$$

$$\sup_{x \in \mathbb{R}} |\tilde{F}_N^{\text{PCLR}_a}(x, \hat{\mathcal{T}}_a) - P[\chi_d^2 \leq x]| \xrightarrow{p} 0, \quad (2.46)$$

$$\sup_{x \in \mathbb{R}} |\tilde{F}_N^{\text{PCLR}_b}(x, \hat{\mathcal{T}}_b) - P[\chi_d^2 \leq x]| \xrightarrow{p} 0. \quad (2.47)$$

It follows that the asymptotic local power of the PAR tests is equal to that of the asymptotic AR test given by $P[\chi_k^2(\eta^2) > r_{1-\alpha}(\chi_k^2)]$, where $r_{1-\alpha}(\chi_k^2)$ is the $1 - \alpha$ quantile of χ_k^2 random variable. Moreover, under strong identification, the PLM and PCLR tests, and the asymptotic LM and CLR tests have the same asymptotic local power equal to $P[\chi_d^2(\eta^2) > r_{1-\alpha}(\chi_d^2)]$.

2.3 PAR tests with a diverging number of IVs

This section considers an extension of the PAR tests to cases where the number of IVs, k , may grow with the sample size n . This extension comprises two main parts. First, the derivation of the asymptotic distribution of the PAR statistics requires a proof strategy different from the fixed- k case. This is accomplished by interpreting a dominant term in the PAR statistic (after centering and scaling) as a double-indexed permutation statistic. Second, we derive the asymptotic distribution of the AR statistic under the condition that $k^3/n \rightarrow 0$ as $n \rightarrow \infty$, which is the same rate used by [Andrews and Stock \(2007\)](#) who establish the asymptotic validity of the AR, LM and CLR tests in homoskedastic linear IV models.

For simplicity, we assume that $X = \iota$, although the proof can be extended to incorporate a fixed number exogenous covariates in a straightforward manner. In the present case, the PAR_1 and PAR_2 tests are (distributionally) equivalent because $W'_\pi M_\iota u(\theta_0) = \sum_{i=1}^n (W_{\pi(i)} -$

$\bar{W})u_i(\theta_0) = \sum_{i=1}^n (W_i - \bar{W})u_{\pi^{-1}(i)}(\theta_0) = Z'u_{\pi^{-1}}(\theta_0)$ and $W'_\pi M_\iota \tilde{\Sigma}_u(\theta_0) M_\iota W_\pi = \sum_{i=1}^n (W_{\pi(i)} - \bar{W})(W_{\pi(i)} - \bar{W})' \tilde{u}_i(\theta_0)^2 = \sum_{i=1}^n (W_i - \bar{W})(W_i - \bar{W})' \tilde{u}_{\pi^{-1}(i)}(\theta_0)^2$, where $\bar{W} \equiv n^{-1}W'_\iota$ and π^{-1} is the inverse of π i.e. $\pi \circ \pi^{-1}$ is identity permutation, and is uniformly distributed over \mathbb{G}_n . When Z and $u(\theta_0)$ are independent under $H_0 : \theta = \theta_0$, the PAR tests are still exact since Proposition 2.1 holds for any k provided the statistics are well-defined.

Let $P_{ij} \equiv Z'_i(Z'Z)^{-1}Z_j$ denote the (i, j) element of P_Z , $\sum_{i \neq j}$ denote the summation over distinct values of the indices $i, j = 1, \dots, n$, and $O_{a.s.}(\cdot)$ abbreviate $O(\cdot)$ almost surely. The asymptotic validity of the PAR tests are established under the following conditions in addition to Assumption 2(a),(b) and (d).

Assumption 3. As $k, n \rightarrow \infty$,

- (a) $\max_{1 \leq i \leq n} \left(\sum_{j=1, j \neq i}^n |P_{ij}| \right) = O_{a.s.} \left(\frac{\sum_{i \neq j} P_{ij}^2}{n \max_{i, j, i \neq j} |P_{ij}|} \right),$
- (b) $\frac{\sum_{i \neq j} P_{ij}^2}{n^2 \max_{i, j, i \neq j} P_{ij}^2} \xrightarrow{a.s.} 0,$
- (c) $k^{-1} \sum_{i=1}^n P_{ii}^2 \xrightarrow{a.s.} 0,$
- (d) $\sup_{j \leq k} (E[W_{1j}^4 u_1^4] + E[W_{1j}^4] + E[u_1^4]) < M_0 < \infty.$

Assumptions 3 is key for deriving the asymptotic distribution of the PAR statistics under the diverging- k asymptotics. Assumptions 3(a) states that the row sums of the off-diagonal elements of P_Z are roughly equal, Assumptions 3(b) requires that the off-diagonal elements of P_Z are small relative to its largest element. By standard arguments, $\sum_{i=1}^n P_{ii}^2 \leq \sum_{i=1}^n P_{ii} = k$, hence $k^{-1} \sum_{i=1}^n P_{ii}^2 \leq 1$. Assumption 3(c) makes a stronger requirement that the left-hand side of the latter inequality actually converges to 0.

To gain insight into Assumption 3(a)-(c), let W be a matrix of group dummy variables i.e. the j -th column of W has entries $W_{ij} = 1, i = n_{j-1} + 1, \dots, n_{j-1} + n_j, j = 1, \dots, k$, with $n_0 = 0, n = \sum_{j=1}^k n_j$, and 0 otherwise. Let $n_{\min} \equiv \min_{1 \leq j \leq k} n_j$ and $n_{\max} \equiv \max_{1 \leq j \leq k} n_j$. By direct calculations, one can see that Assumption 3(a) is equivalent to $(1 - n_{\max}^{-1})nn_{\min}^{-1}/(k - \sum_{j=1}^k n_j^{-1}) = O_{a.s.}(1)$, which holds if n_{\max}/n_{\min} is bounded.

Assumption 3(b) is equivalent to $(k - \sum_{j=1}^k n_j^{-1})/(n^2/n_{\min})$ being bounded. The latter holds because $(k - \sum_{j=1}^k n_j^{-1})/(n^2/n_{\min}) \leq k^{-2}(k - \sum_{j=1}^k n_j^{-1}) \leq k^{-2}(k - n^{-1}k^2) \leq 1$ using $(n/n_{\min}) \leq k$ and $\sum_{j=1}^k n_j^{-1} \geq k^2/(\sum_{j=1}^k n_j)$ which follows from the convexity of $x \mapsto x^{-1}$. Finally, 3(c) holds if $n_{\min} \xrightarrow{a.s.} \infty$, since $k^{-1} \sum_{i=1}^n P_{ii}^2 = k^{-1} \sum_{j=1}^k n_j^{-2} \leq n_{\min}^{-2} \xrightarrow{a.s.} 0$.

Assumption 3(d) replaces Assumption 2(c). The boundedness of the cross moment $E[W_{1j}^4 u_1^4]$ is needed because we allow for a growing k which, in turn, imposes a restriction on the consistency of the covariance matrix estimator. Furthermore, the marginal moment conditions in Assumption 3(d) are slightly weaker than Assumption 2(c) because we only show the tests have asymptotically correct level, as opposed to size (uniform validity). The main results of this section is as follows.

Proposition 2.4. *Let Assumptions 2(a),(b), (d), and 3 hold with P independent of n . Assume that $k, N \rightarrow \infty$ and $k^3/n \rightarrow 0$ as $n \rightarrow \infty$. If $H_0 : \theta = \theta_0$ is true, then, for $\alpha \in (0, 1)$ and the permutation statistics $\text{PAR} \in \{\text{PAR}_1, \text{PAR}_2\}$, it holds that*

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PAR}}] = \alpha.$$

The asymptotic distribution of the PAR statistics is obtained as follows. Letting $\tilde{\sigma}^2 \equiv n^{-1} \tilde{u}(\theta_0)' \tilde{u}(\theta_0)$, we can rewrite

$$\begin{aligned} \text{PAR}_2 = & \tilde{u}_\pi(\theta_0)' Z \left(\left(\sum_{i=1}^n Z_i Z_i' \tilde{u}_{\pi(i)}(\theta_0)^2 \right)^{-1} - (Z' Z)^{-1} \tilde{\sigma}^{-2} \right) Z' \tilde{u}_\pi(\theta_0) \\ & + \sum_{i=1}^n P_{ii} \tilde{u}_{\pi(i)}(\theta_0)^2 \tilde{\sigma}^{-2} + \sum_{i \neq j} P_{ij} \tilde{u}_{\pi(i)}(\theta_0) \tilde{u}_{\pi(j)}(\theta_0) \tilde{\sigma}^{-2}. \end{aligned} \quad (2.48)$$

It can be shown that the first summand in the above expression is negligible while the second summand converges to k . The key insight for establishing the asymptotic distribution of the third summand is to view it as a double-indexed permutation statistic of the form $\sum_{i \neq j} a_{ij} b_{\pi(i)\pi(j)}$ to which one can apply a CLT established by [Pham et al. \(1989\)](#).

Recently, [Mikusheva and Sun \(2022\)](#) propose a heteroskedasticity-robust Jackknife AR (JAR) test that is based on $\sum_{i \neq j} P_{ij} \tilde{u}_i(\theta_0) \tilde{u}_j(\theta_0)$ and is asymptotically valid under the condition $k/n \rightarrow \tau \in (0, 1)$ among others. Our proof strategy is likely to be useful for

deriving the asymptotic distribution of a permutation version of their statistic. We leave this extension as well as those of the PLM and PCLR statistics under conditions less stringent than those in Assumption 3 to future research.

3 Simulations

This section presents a simulation evidence on the performance of the proposed permutation tests. The data were generated according to

$$\begin{aligned} y_i &= Y_i\theta + \gamma_1 + X'_{2i}\gamma_2 + u_i, \\ Y_i &= W'_i\Gamma + \psi_1 + X'_{2i}\Psi_2 + V_i, \quad i = 1, \dots, n, \end{aligned}$$

where $\theta = 0$, $d = 1$, $\Gamma = (1, \dots, 1)'\sqrt{\lambda/(nk)}(k \times 1)$ and $\lambda \in \{0.1, 4, 20\}$, X_{2i} is $(p - 1) \times 1$ vector of non-constant included exogenous variables with $p \in \{1, 5\}$, and $V_i = \rho u_i + \sqrt{1 - \rho^2}\epsilon_i$ with $\rho = 0.5$. The parameters $\gamma_1, \gamma_2, \psi_1$, and Ψ_2 are set equal to 0. As specified below, W_i has zero mean and unit covariance matrix except for the first part of simulations in Section 3.1, hence $\lambda = n\Gamma' E[W_i W'_i] \Gamma \approx \Gamma' W' W \Gamma$. We consider the values $\lambda \in \{0.1, 4, 20\}$ which correspond to very weak, weak and strong identification. The tested restriction is $H_0 : \theta = \theta_0 = 0$.

We implement the heteroskedasticity-robust asymptotic tests denoted as AR, LM, CLR_a, CLR_b, their permutation versions PAR₁, PAR₂, PLM, PCLR_a, PCLR_b, the normal score and Wilcoxon score rank-based AR tests of Andrews and Marmer (2008) denoted as RARn and RARw, respectively, the normal score and Wilcoxon score rank-based CLR statistics of Andrews and Soares (2007) denoted as RCLRn and RCLRw, respectively, and the wild bootstrap AR and LM tests of Davidson and MacKinnon (2012) denoted as WAR and WLM, respectively. For both the asymptotic and permutation CLR-type tests, we do not make the eigenvalue-adjustment (i.e. $\epsilon = 0$ in footnote 4).

In all simulations below, the null rejection probabilities of the tests (including CLR, PCLR, the rank-based and the wild bootstrap tests considered below) are computed using

2000 replications with 999 simulated samples for each replication while we display some power curves associated with the tests in Supplemental Appendix. We consider the following three cases in turn.

3.1 Heavy tails

To examine the finite sample validity, we consider heavy-tailed observations where each element of the random vector $(W'_i, X'_{2i}, u_i, \epsilon_i)'$ is drawn independently from standard Cauchy distribution, and set $k \in \{5, 10\}$ $n \in \{50, 100\}$, and $\lambda = 4$.

Table 1 presents the null rejection probabilities. The asymptotic tests all underreject. The PAR_1 test has nearly correct levels as predicted by Theorem 2.1 in all cases, and the PAR_2 test, which is exact only when $p = 1$, has more accurate rejection rates in the case $p = 1$ than in the case $p = 5$. The RAR_n , RAR_w , RCLR_n and RCLR_w tests are all robust against heavy-tailed errors when the IVs, the exogenous covariates and the error terms are independent as stated in Assumption 2.1. The RAR_n and RAR_w tests are exact and this is borne out in the simulation results. The RCLR_n and RCLR_w tests slightly underreject which may be attributed to their asymptotic nature.

Note, however, that, under the independence assumption and heavy-tailed errors, the rank-based tests have better power properties than the asymptotic tests, see Andrews and Soares (2007) and Andrews and Marmer (2008). In such cases, the permutation tests are likely to be dominated by the rank-based tests in terms of power given the local asymptotic equivalence between the permutation tests and the asymptotic tests under strong identification established in Proposition 2.3.

3.2 Independence vs. standard exclusion restriction

The next set of simulations examines the role of studentization for obtaining valid permutation tests and highlights the difference between the independence and the standard exclusion restriction assumption in homoskedastic setting. Two cases are considered:

1. $(W'_i, X'_{2i}, u_i, \epsilon_i)' \sim t_5[0, I_{k+p+1}]$, where $t_5[0, I_{k+p+1}]$ stands for $(k + p + 1)$ -variate t -distribution with degrees of freedom 5 and covariance matrix I_{k+p+1} . Under this setting, these random variables have finite fourth moments, and $(W'_i, X'_{2i})'$ satisfies the standard exclusion restriction in Assumption 2: $E[(u_i, V'_i)'(W'_i, X'_i)] = 0$ but $(u_i, \epsilon_i)'$ and $(W'_i, X'_i)'$ are dependent;⁵
2. $(W'_i, X'_{2i}, u_i, \epsilon_i)' \sim N[0, I_{k+p+1}]$. Clearly, in this case the instruments and the error terms are independent.

The result for the overidentified models with $n = 100$ and $k = 5$ is reported in Table 2. The permutation tests perform better than the asymptotic tests regardless of whether the instruments and the error terms are independent or not. The rank-based tests, displayed in the bottom part of Table 2, have nearly correct level in the independent case but overreject in the dependent case.

The results for the just-identified design with $k = 1$ is similar to the over-identified case, thus is relegated to Supplemental Appendix.

3.3 Conditional heteroskedasticity

The next part of the simulations considers designs with conditional heteroskedasticity. The data are generated according to (2.2) and (2.3) with

$$u_i = W_{i1}v_i,$$

$$V_i = \rho u_i + \sqrt{1 - \rho^2}\epsilon_i,$$

where W_{i1} is the first element of W_i , and similarly to the previous case

1. $(W'_i, X'_{2i}, v_i, \epsilon_i)' \sim t_5[0, I_{k+p+1}]$;
2. $(W'_i, X'_{2i}, v_i, \epsilon_i)' \sim N[0, I_{k+p+1}]$.

⁵Because the distribution is fixed i.e. does not vary with the sample size, the permutation tests remain asymptotically valid under the finite fourth moment assumption.

The sample size and the number of instruments are $n = 100$ and $k \in \{2, 5, 10\}$. The results for $\lambda = 4$ are displayed in Tables 3. The asymptotic AR test tends to underreject and performs poorly compared to the LM statistic. The permutation tests perform better than their asymptotic counterparts in most cases, and on par with the wild bootstrap in the independent case. However, when the instruments satisfy the standard exogeneity condition, and there are included exogenous variables, the permutation tests appear to have an edge over the wild bootstrap AR test which overrejects. Qualitatively, the same observations made in the homoskedastic case for the rank-based tests are also observed in the heteroskedastic case as the rank-based tests reject by a substantial margin in the dependent case. See Supplemental Appendix for additional simulation evidence for the case $\lambda \in \{0.1, 20\}$.

4 Empirical application

Bazzi and Clemens (2013) re-examine cross-sectional IV regression results about the effect of foreign aid on economic growth from several published studies. The variables in the “aid-growth” IV regressions estimated by Rajan and Subramanian (2008) and Bazzi and Clemens (2013) are

- y_i : the average annual growth of per capita GDP;
- Y_i : the foreign-aid receipts to GDP ratio (Aid/GDP);
- W_{i1} : a variable constructed from aid-recipient population size, aid-donor population size, colonial relationship, and language traits (see Appendix A of Bazzi and Clemens (2013));
- X_i : constant, initial per capita GDP, initial level of policy, initial level of life expectancy, geography, institutional quality, initial inflation, initial M2/GDP, initial budget balance/GDP, Revolutions, and Ethnic fractionalization.

Table 1: Null rejection probabilities at 5% level. Cauchy distributed IVs, controls and error terms.

$\lambda = 4$	$n = 50$				$n = 100$			
	$p = 1$		$p = 5$		$p = 1$		$p = 5$	
	$k = 5$	$k = 10$	$k = 5$	$k = 10$	$k = 5$	$k = 10$	$k = 5$	$k = 10$
	Tests							
AR	0.95	0.55	0.85	0.25	0.50	1.05	0.25	0.40
LM	2.35	4.25	2.05	3.10	2.10	4.45	1.85	3.45
CLR _a	1.45	1.70	1.15	0.75	0.75	1.95	0.55	1.30
CLR _b	2.10	3.60	1.60	2.60	1.75	3.85	1.25	3.25
PAR ₁	4.40	5.30	4.70	5.60	4.05	4.95	4.45	4.70
PAR ₂	4.40	5.45	3.40	3.50	3.95	4.75	3.25	3.05
RAR _n	4.25	4.55	4.10	4.15	5.10	4.05	4.90	3.75
RAR _w	4.10	4.85	4.30	4.55	5.15	3.95	4.55	3.60
RCLR _n	3.15	3.75	4.20	5.10	4.55	3.50	3.85	4.00
RCLR _w	2.35	3.25	3.35	4.25	3.50	2.80	3.25	3.45

Note: AR, LM, and CLR are the heteroskedasticity-robust asymptotic tests, PAR₁ and PAR₂ denote the robust permutation AR tests, RAR_n and RAR_w denote the normal score and Wilcoxon score rank-based AR tests of [Andrews and Marmer \(2008\)](#), and RCLR_n and RCLR_w denote the normal score and Wilcoxon score rank-based CLR tests of [Andrews and Soares \(2007\)](#) respectively. The number of replications is 2000, and the number of permutation samples for each replication is 999 ($N = 1000$).

Table 2: Null rejection probabilities at 5% level. Over-identified homoskedastic model with $k = 5$, $n = 100$.

$n = 100$	$\lambda = 0.1$				$\lambda = 4$				$\lambda = 20$			
$k = 5$	$t_5[0, I_{k+p+1}]$		$N[0, I_{k+p+1}]$		$t_5[0, I_{k+p+1}]$		$N[0, I_{k+p+1}]$		$t_5[0, I_{k+p+1}]$		$N[0, I_{k+p+1}]$	
Tests	$p = 1$	$p = 5$	$p = 1$	$p = 5$	$p = 1$	$p = 5$	$p = 1$	$p = 5$	$p = 1$	$p = 5$	$p = 1$	$p = 5$
AR	3.15	4.50	4.50	4.25	3.15	4.50	4.50	4.25	3.15	4.50	4.50	4.25
LM	2.90	4.30	4.15	4.25	3.20	4.05	3.70	3.70	3.00	4.10	3.25	3.85
CLR _a	2.90	3.95	4.00	4.05	3.20	4.15	3.30	3.55	2.95	4.25	3.45	3.75
CLR _b	2.80	3.95	4.05	4.05	3.20	4.10	3.35	3.45	2.55	4.00	3.30	3.95
PAR ₁	5.00	5.55	5.45	4.75	5.00	5.55	5.45	4.75	5.00	5.55	5.45	4.75
PAR ₂	5.05	6.85	5.60	5.45	5.05	6.85	5.60	5.45	5.05	6.85	5.60	5.45
PLM	5.10	5.60	5.40	5.15	5.00	6.05	5.05	5.05	4.70	6.15	4.95	5.45
PCLR _a	4.80	6.25	5.20	4.85	4.30	5.50	4.20	4.30	4.25	5.20	3.70	4.15
PCLR _b	6.20	7.25	5.35	5.10	6.30	6.70	4.65	4.75	5.80	7.00	4.30	5.00
RAR _n	18.65	15.70	5.25	4.45	18.65	15.70	5.25	4.45	18.65	15.70	5.25	4.45
RAR _w	12.40	10.20	5.35	4.10	12.40	10.20	5.35	4.10	12.40	10.20	5.35	4.10
RCLR _n	13.85	14.55	3.35	3.70	13.15	14.50	3.15	3.75	11.10	12.65	4.00	3.50
RCLR _w	10.40	12.90	4.45	4.55	10.20	11.90	4.20	4.65	9.15	10.90	4.55	4.35
WAR	5.30	7.45	5.35	4.10	5.30	7.45	5.35	4.10	5.30	7.45	5.35	4.10
WLM	4.65	5.85	4.95	4.80	4.20	6.40	4.40	4.05	4.50	5.95	4.30	4.35

Note: $t_5[0, I_{k+p+1}]$ and $N[0, I_{k+p+1}]$ correspond to $(W'_i, X'_{2i}, u_i, \epsilon_i)' \sim t_5[0, I_{k+p+1}]$ (dependent but uncorrelated) and $(W'_i, X'_{2i}, u_i, \epsilon_i)' \sim N[0, I_{k+p+1}]$ (independent), respectively. PLM and PCLR denote the robust permutation LM and CLR tests, WAR and WLM are the wild bootstrap AR and LM tests of [Davidson and MacKinnon \(2012\)](#) and the remaining tests are as in Table 1. The number of replications is 2000, and the number of permutation samples for each replication is 999 ($N = 1000$).

Table 3: Null rejection probabilities at 5% level. Conditionally heteroskedastic design with weak identification ($\lambda = 4$).

$n = 100$	$p = 1$						$p = 5$					
	$t_5[0, I_{k+p+1}]$			$N[0, I_{k+p+1}]$			$t_5[0, I_{k+p+1}]$			$N[0, I_{k+p+1}]$		
	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$
Tests												
AR	2.35	1.75	0.60	4.05	2.90	1.40	4.30	2.40	1.15	5.25	3.25	2.60
LM	3.10	3.95	3.70	4.55	3.85	3.80	4.50	2.80	2.50	6.00	3.60	4.00
CLR _a	2.45	2.10	1.00	4.15	3.20	1.55	4.25	2.55	1.25	5.70	2.85	3.15
CLR _b	2.40	2.25	1.50	4.10	3.15	1.85	4.15	2.45	1.30	5.65	3.00	3.45
PAR ₁	3.60	5.15	4.35	4.85	5.60	5.20	7.40	7.55	8.85	5.95	4.70	4.70
PAR ₂	3.35	4.85	4.35	4.90	5.30	5.00	6.35	5.50	4.95	6.40	5.30	5.35
PLM	4.20	5.35	5.80	5.15	4.90	5.05	6.20	3.90	4.60	6.30	4.45	6.10
PCLR _a	3.70	5.55	4.15	4.90	4.70	4.05	6.05	4.65	3.90	6.45	4.55	5.65
PCLR _b	3.80	6.20	8.75	5.05	5.20	5.40	6.20	5.80	7.85	6.85	5.05	7.10
RAR _n	22.30	27.30	32.55	13.45	10.20	8.05	23.55	24.80	29.85	13.75	10.10	8.70
RAR _w	14.35	17.90	20.10	10.40	8.85	6.65	15.95	16.00	17.55	10.40	8.25	7.35
RCLR _n	19.45	22.40	29.95	11.05	7.50	6.05	29.25	31.25	41.50	11.85	8.50	7.15
RCLR _w	13.70	17.50	19.55	9.40	7.45	6.35	25.00	26.20	36.00	10.10	8.25	7.55
WAR	4.50	5.80	5.55	4.75	5.20	4.90	8.65	10.80	12.60	5.75	4.40	4.95
WLM	3.20	3.50	4.15	4.00	4.05	4.90	6.60	5.85	7.00	5.55	3.90	5.30

Note: $t_5[0, I_{k+p+1}]$ and $N[0, I_{k+p+1}]$ correspond to $(W'_i, X'_{2i}, v_i, \epsilon_i)' \sim t_5[0, I_{k+p+1}]$ (dependent but uncorrelated) and $(W'_i, X'_{2i}, v_i, \epsilon_i)' \sim N[0, I_{k+p+1}]$ (independent), respectively. The error terms are generated as $u_i = W_{i1}v_i, V_i = \rho u_i + \sqrt{1 - \rho^2}\epsilon_i$, where W_{i1} is the first element of W_i and $\rho = 0.5$. The tests are in Table 2. The number of replications is 2000, and the number of permutation samples for each replication is $N = 999$.

The data cover the period 1970-2000 and the sample size is $n = 78$. [Bazzi and Clemens \(2013\)](#) consider the following four specifications (Table 1, Columns 1-4 therein):

- Specification 1: The baseline specification of [Rajan and Subramanian \(2008\)](#) (Table 4, Column 2) with the variables as above;
- Specification 2: log population is included in the second stage of Specification 1;
- Specification 3: $W_{i2} = \log \text{population}$ replaces the instrument W_{i1} in Specification 1;
- Specification 4: An instrument, W_{i3} , constructed from the colonial ties indicators only replaces W_{i1} in Specification 1.

Specifications 1-4 are just-identified as there is a single instrument for the single endogenous regressor, hence we only consider the AR and PAR test results. In addition, we consider Specifications 5-7 with IVs $(W_{i1}, W_{i2})'$, $(W_{i2}, W_{i3})'$ and $(W_{i1}, W_{i2}, W_{i3})'$, respectively, to examine whether the over-identified specifications could have any instrumentation power. W_{i2} is included in the latter three cases because it is the strongest instrument as reported in [Bazzi and Clemens \(2013\)](#). We also include the confidence interval based on the two-stage least squares t -test, denoted as t_{2sls} .

Tables 4 and 5 report the 95% confidence intervals for the coefficient on Aid/GDP. In all specifications, the Breusch-Pagan test p -values from the two separate reduced-form regressions of the endogenous variables on the exogenous variables show an evidence of heteroskedasticity.

The results for Specifications 1-4 in Table 4 qualitatively agree with the homoskedastic CLR confidence intervals reported in [Bazzi and Clemens \(2013\)](#). The permutation confidence intervals are nearly identical but slightly shorter than the asymptotic and wild bootstrap confidence intervals in Specifications 1 and 3. The infinite confidence intervals in Specifications 2 and 4 that do not use log population or the instruments based on it indicate that the population size instrument used in the other specifications has indeed the most identifying power.

The above results also extend to Table 5. Some noteworthy findings are as follows. The WAR confidence intervals are wider than the AR and PAR confidence intervals. The PLM confidence intervals are shorter than the LM and WLM confidence intervals, but still wider than the other confidence intervals. Somewhat surprisingly, in Specification 6, the AR, CLR_a , PAR and PCLR confidence intervals exclude 0 thereby pointing towards borderline significant effect of the foreign aid on growth despite the rather small value of the first-stage F -statistic 17.90. Also, in Specification 7, the PCLR confidence intervals exclude 0 while the opposite holds for the asymptotic CLR confidence intervals which underscore the utility of the proposed tests.

However, these results should be interpreted with caution because the population size might affect the growth through multiple channels as argued by [Bazzi and Clemens \(2013\)](#). Overall, the uncertainty regarding the effect of foreign aid in the sample data may be expressed more accurately by the permutation and the wild bootstrap confidence intervals as the unobserved confounders in the “aid-growth” regression are more likely to be orthogonal to, rather than independent of the population size.

5 Conclusion

The PAR_1 test shares the strengths of the rank/permutation tests and the wild bootstrap tests because it is exact under independence and heteroskedasticity-robust under standard assumptions. This test is recommended when the IVs are randomly assigned, hence independent of the exogenous covariates and the error term. The PAR_2 test also shows a satisfactory performance in the simulations. Because the PAR tests do not require the correct specification of the first-stage equation as they directly uses the IVs in the equation $\hat{m}(\theta_0) = n^{-1}Z'\tilde{u}(\theta_0)$ ([Dufour and Taamouti, 2007](#)), they have robustness advantages relative to the other permutation tests.

Moreover, the PCLR tests are found to perform reasonably well in the simulations. Assuming that (2.2)-(2.3) are correctly specified, as we have shown, the PCLR tests are

Table 4: Confidence intervals for the coefficient on Aid/GDP in [Rajan and Subramanian \(2008\)](#) and [Bazzi and Clemens \(2013\)](#) data.

	Spec. 1		Spec. 2		Spec. 3		Spec. 4
Test statistics	95% CI	Length	95% CI	95% CI	Length	95% CI	
t_{2sls}	$[-0.05, 0.24]$	0.29	$[-5.14, 6.96]$	$[-0.06, 0.21]$	0.27	$[-973.38, 941.49]$	
Hom. AR	$[-0.02, 0.28]$	0.30	$(-\infty, \infty)$	$[-0.03, 0.25]$	0.28	$(-\infty, \infty)$	
RARn	$[-0.05, 0.29]$	0.34	$(-\infty, \infty)$	$[-0.06, 0.28]$	0.34	$(-\infty, \infty)$	
RARw	$[-0.06, 0.20]$	0.26	$(-\infty, \infty)$	$[-0.07, 0.16]$	0.23	$(-\infty, \infty)$	
RCLRn	$[-0.03, 0.26]$	0.29	$(-\infty, \infty)$	$[-0.04, 0.22]$	0.26	$(-\infty, \infty)$	
RCLRw	$[-0.04, 0.18]$	0.22	$(-\infty, \infty)$	$[-0.05, 0.16]$	0.21	$(-\infty, \infty)$	
AR	$[-0.01, 0.29]$	0.30	$(-\infty, \infty)$	$[-0.02, 0.25]$	0.27	$(-\infty, \infty)$	
PAR ₁	$[-0.01, 0.30]$	0.31	$(-\infty, \infty)$	$[-0.03, 0.27]$	0.30	$(-\infty, \infty)$	
PAR ₂	$[-0.01, 0.28]$	0.29	$(-\infty, \infty)$	$[-0.02, 0.25]$	0.27	$(-\infty, \infty)$	
PNS	$[-0.04, 0.37]$	0.41	$(-\infty, \infty)$	$[-0.05, 0.30]$	0.35	$(-\infty, \infty)$	
WAR	$[-0.02, 0.32]$	0.34	$(-\infty, \infty)$	$[-0.03, 0.28]$	0.31	$(-\infty, \infty)$	
F -statistic	31.63		0.13		36.30		0.00
Breusch-Pagan p -values	{0.03, 0.00}		{0.03, 0.00}		{0.02, 0.00}		{0.01, 0.00}

Note: The number of permutation samples is 1999 ($N = 2000$). The sample size is 78. Specifications 1-4 are just-identified. t_{2sls} and Hom. AR are the homoskedastic two-stage least squares t and AR tests, PNS denotes the non-studentized permutation test, and the remaining tests are as in Tables 1 and 2. The Breusch-Pagan test p -values are computed using the fitted values in the reduced-form regressions of y_i and Y_i on the instrument and exogenous covariates respectively.

Table 5: Confidence intervals for the coefficient on Aid/GDP in [Rajan and Subramanian \(2008\)](#) and [Bazzi and Clemens \(2013\)](#) data.

Test statistics	Spec. 5		Spec. 6		Spec. 7	
	95% CI	Length	95% CI	Length	95% CI	Length
t_{2sls}	[-0.06, 0.22]	0.28	[-0.06, 0.21]	0.27	[-0.06, 0.21]	0.27
Hom. AR	[-0.05, 0.31]	0.36	[-0.02, 0.26]	0.28	[-0.04, 0.32]	0.36
Hom. LM	[-0.63, 0.26]	0.89	[-0.63, 0.28]	0.91	[-0.63, 0.28]	0.91
Hom. CLR	[-0.03, 0.26]	0.29	[-0.03, 0.28]	0.31	[-0.03, 0.28]	0.31
RARn	[-0.08, 0.41]	0.49	[-0.06, 0.32]	0.38	[-0.09, 0.40]	0.49
RARw	[-0.08, 0.29]	0.37	[-0.06, 0.23]	0.29	[-0.09, 0.27]	0.36
RCLRn	[-0.03, 0.24]	0.27	[-0.03, 0.25]	0.28	[-0.03, 0.25]	0.28
RCLRw	[-0.05, 0.17]	0.22	[-0.03, 0.17]	0.20	[-0.04, 0.17]	0.21
AR	[-0.04, 0.34]	0.38	[0.04, 0.32]	0.28	[0.01, 0.42]	0.41
LM	[-0.43, 0.27]	0.70	[-0.61, 0.36]	0.97	[-0.69, 0.35]	1.04
CLR _a	[-0.03, 0.28]	0.31	[0.02, 0.36]	0.34	[0.00, 0.35]	0.35
CLR _b	[-0.02, 0.28]	0.30	[0.00, 0.36]	0.36	[-0.03, 0.34]	0.37
PAR ₁	[-0.04, 0.35]	0.39	[0.04, 0.34]	0.30	[0.01, 0.43]	0.42
PAR ₂	[-0.04, 0.34]	0.38	[0.04, 0.31]	0.27	[0.02, 0.38]	0.36
PLM	[-0.43, 0.25]	0.68	[-0.60, 0.33]	0.93	[-0.65, 0.33]	0.98
PCLR _a	[-0.03, 0.27]	0.30	[0.02, 0.34]	0.32	[0.01, 0.35]	0.34
PCLR _b	[-0.02, 0.27]	0.29	[0.02, 0.34]	0.32	[0.01, 0.35]	0.34
PNS	[-0.04, 0.39]	0.43	[-0.06, 0.32]	0.38	[-0.04, 0.35]	0.39
WAR	[-0.05, 0.44]	0.49	[0.00, 0.33]	0.33	[-0.04, 0.48]	0.52
WLM	[-0.64, 0.30]	0.94	[-0.66, 0.30]	0.96	[-0.66, 0.33]	0.99
F -statistic	17.98		17.90		11.86	
Sargan p -value	0.42		0.08		0.21	
Breusch-Pagan p -values	{0.03, 0.00}		{0.02, 0.00}		{0.02, 0.00}	

Note: The number of permutation samples is 1999 ($N = 2000$). The sample size is 78. Specifications 5-7 are over-identified. Hom. LM and Hom. CLR are the homoskedastic LM and CLR tests, Sargan p -value is the p -value associated with the Sargan test of overidentifying restrictions, and the remaining tests are as in Tables 4.

similar and equivalent to the CLR tests of [Andrews and Guggenberger \(2019a\)](#) asymptotically. In homoskedastic IV models, the latter tests are asymptotically equivalent or reduce to the CLR test of [Moreira \(2003\)](#) which is known to be optimal when there is single endogenous regressor ([Andrews et al., 2006](#)). However, despite being equivalent to the PCLR tests under strong identification, the PLM test shows an inferior performance in terms of power. Based on the properties mentioned and the simulation evidence, we recommend reporting both the PAR and PCLR test results in practice.

Looking ahead, we plan to address the important issue of cluster and identification-robust inference in IV models using the current approach in another work.

Supplementary material

Supplemental Appendix: This file contains all the proofs of the results in the paper and additional numerical evidence. (pdf)

R codes: This file contains the replication R codes of the empirical analyses as well as the companion datasets described in the paper. (zip)

References

- ABADIE, A., ATHEY, S., IMBENS, G. W. and WOOLDRIDGE, J. M. (2020). Sampling-based vs. Design-based Uncertainty in Regression Analysis. *Econometrica*, **88** 265–296.
- ANDERSON, T. W. and RUBIN, H. (1949). Estimation of the Parameters of a Single Equation in a Complete System of Stochastic Equations. *Annals of Mathematical Statistics*, **20** 46–63.
- ANDREWS, D. W. and GUGGENBERGER, P. (2019a). Identification-and Singularity-Robust Inference for Moment Condition Models. *Quantitative Economics*, **10** 1703–1746.

- ANDREWS, D. W. K., CHENG, X. and GUGGENBERGER, P. (2020). Generic Results for Establishing the Asymptotic Size of Confidence Sets and Tests. *Journal of Econometrics*, **218** 496–531.
- ANDREWS, D. W. K. and GUGGENBERGER, P. (2017a). Asymptotic Size of Kleibergen’s LM and Conditional LR Tests for Moment Condition Models. *Econometric Theory*, **33** 1046–1080.
- ANDREWS, D. W. K. and GUGGENBERGER, P. (2017b). Supplemental material to ”Asymptotic Size of Kleibergen’s LM and Conditional LR Tests for Moment Condition Models”. *Econometric Theory*, **33**.
- ANDREWS, D. W. K. and GUGGENBERGER, P. (2019b). Supplemental material to “Identification-and Singularity-Robust Inference for Moment Condition Models”. *Quantitative Economics*, **10**.
- ANDREWS, D. W. K. and MARMER, V. (2008). Exactly Distribution-Free Inference in Instrumental Variables Regression with Possibly Weak Instruments. *Journal of Econometrics*, **142** 183–200.
- ANDREWS, D. W. K., MOREIRA, M. J. and STOCK, J. H. (2006). Optimal Two-Sided Invariant Similar Tests for Instrumental Variables Regression. *Econometrica*, **74** 715–752.
- ANDREWS, D. W. K. and SOARES, G. (2007). Rank Tests for Instrumental Variables Regression with Weak Instruments. *Econometric Theory*, **23** 1033–1082.
- ANDREWS, D. W. K. and STOCK, J. H. (2007). Inference with Weak Instruments. In *Advances in Econometrics: Proceedings of the Ninth World Congress of the Econometric Society*, vol. 3.
- ANDREWS, I., STOCK, J. H. and SUN, L. (2019). Weak Instruments in Instrumental Variables Regression: Theory and Practice. *Annual Review of Economics*, **11** 727–753.
- BAZZI, S. and CLEMENS, M. A. (2013). Blunt Instruments: Avoiding Common Pitfalls in Identifying the Causes of Economic Growth. *American Economic Journal: Macroeconomics*, **5** 152–86.

- BUGNI, F. A., CANAY, I. A. and SHAIKH, A. M. (2018). Inference under covariate-adaptive randomization. *Journal of the American Statistical Association*, **113** 1784–1796.
- CANAY, I. A. and KAMAT, V. (2018). Approximate Permutation Tests and Induced Order Statistics in the Regression Discontinuity Design. *The Review of Economic Studies*, **85** 1577–1608.
- CANAY, I. A., ROMANO, J. P. and SHAIKH, A. M. (2017). Randomization Tests under an Approximate Symmetry Assumption. *Econometrica*, **85** 1013–1030.
- CHERNOZHUKOV, V., HANSEN, C. and JANSSEN, M. (2009). Finite Sample Inference for Quantile Regression Models. *Journal of Econometrics*, **152** 93–103.
- CHUNG, E. and ROMANO, J. P. (2013). Exact and Asymptotically Robust Permutation Tests. *Annals of Statistics*, **41** 484–507.
- CHUNG, E. and ROMANO, J. P. (2016). Multivariate and Multiple Permutation Tests. *Journal of Econometrics*, **193** 76–91.
- DAVIDSON, R. and MACKINNON, J. G. (2008). Bootstrap Inference in a Linear Equation Estimated by Instrumental Variables. *The Econometrics Journal*, **11** 443–477.
- DAVIDSON, R. and MACKINNON, J. G. (2012). Wild Bootstrap Tests for IV Regression. *Journal of Business & Economic Statistics*, **28** 128–144.
- DICICCIO, C. J. and ROMANO, J. P. (2017). Robust Permutation Tests for Correlation and Regression Coefficients. *Journal of the American Statistical Association*, **112** 1211–1220.
- DUFOUR, J.-M. (2003). Identification, Weak Instruments and Statistical Inference in Econometrics. *Canadian Journal of Economics*, **36** 767–808.
- DUFOUR, J.-M., FLACHAIRE, E. and KHALAF, L. (2019). Permutation Tests for Comparing Inequality Measures. *Journal of Business & Economic Statistics*, **37** 457–470.

- DUFOUR, J.-M. and TAAMOUTI, M. (2007). Further Results on Projection-Based Inference in IV Regressions with Weak, Collinear or Missing Instruments. *Journal of Econometrics*, **139** 133–153.
- FREEDMAN, D. and LANE, D. (1983). A Nonstochastic Interpretation of Reported Significance Levels. *Journal of Business & Economic Statistics*, **1** 292–298.
- GANONG, P. and JÄGER, S. (2018). A Permutation Test for the Regression Kink Design. *Journal of the American Statistical Association*, **113** 494–504.
- HOEFFDING, W. (1951). A Combinatorial Central Limit Theorem. *Annals of Mathematical Statistics*, **22** 558–566.
- IMBENS, G. W. and ROSENBAUM, P. R. (2005). Robust, Accurate Confidence Intervals with a Weak Instrument: Quarter of Birth and Education. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, **168** 109–126.
- JANSSEN, A. (1997). Studentized Permutation Tests for Non-IID Hypotheses and the Generalized Behrens-Fisher Problem. *Statistics & Probability letters*, **36** 9–21.
- KLEIBERGEN, F. (2002). Pivotal Statistics for Testing Structural Parameters in Instrumental Variables Regression. *Econometrica*, **70** 1781–1803.
- LEHMANN, E. L. and ROMANO, J. P. (2005). *Testing Statistical Hypotheses*. 3rd ed. Springer.
- LINNIK, J. V. (2008). *Statistical Problems with Nuisance Parameters*, vol. 20. American Mathematical Soc.
- MIKUSHEVA, A. and SUN, L. (2022). Inference with Many Weak Instruments. *The Review of Economic Studies*, **89** 2663–2686.
- MOREIRA, H. and MOREIRA, M. J. (2019). Optimal Two-Sided Tests for Instrumental Variables Regression with Heteroskedastic and Autocorrelated Errors. *Journal of Econometrics*, **213** 398–433.

- MOREIRA, M. J. (2003). A Conditional Likelihood Ratio Test for Structural Models. *Econometrica*, **71** 1027–1048.
- MOREIRA, M. J., PORTER, J. R. and SUAREZ, G. A. (2009). Bootstrap Validity for the Score Test When Instruments May be Weak. *Journal of Econometrics*, **149** 52–64.
- MOTOO, M. (1956). On the Hoeffding’s Combinatorial Central Limit Theorem. *Annals of the Institute of Statistical Mathematics*, **8** 145–154.
- NEUBERT, K. and BRUNNER, E. (2007). A Studentized Permutation Test for the Non-Parametric Behrens–Fisher Problem. *Computational Statistics & Data Analysis*, **51** 5192–5204.
- NEUHAUS, G. (1993). Conditional Rank Tests for the Two-Sample Problem under Random Censorship. *Annals of Statistics*, **21** 1760–1779.
- PAULY, M. (2011). Discussion about the Quality of F-Ratio Resampling Tests for Comparing Variances. *Test*, **20** 163–179.
- PHAM, D. T., MÖCKS, J. and SROKA, L. (1989). Asymptotic Normality of Double-Indexed Linear Permutation Statistics. *Annals of the Institute of Statistical Mathematics*, **41** 415–427.
- RAJAN, R. G. and SUBRAMANIAN, A. (2008). Aid and Growth: What Does the Cross-Country Evidence Really Show? *The Review of Economics and Statistics*, **90** 643–665.
- SO, B. S. and SHIN, D. W. (1999). Cauchy Estimators for Autoregressive Processes with Applications to Unit Root Tests and Confidence Intervals. *Econometric Theory*, **15** 165–176.
- STOCK, J. H., WRIGHT, J. H. and YOGO, M. (2002). A Survey of Weak Instruments and Weak Identification in Generalized Method of Moments. *Journal of Business and Economic Statistics*, **20** 518–529.
- YOUNG, A. (2022). Consistency without Inference: Instrumental Variables in Practical Application. *European Economic Review*, **147** 104112.

Robust Permutation Tests in Linear Instrumental Variables Regression Supplemental Appendix

Purevdorj Tuvaandorj*

York University

Abstract

This supplemental appendix contains the proofs of the results in the main paper and additional numerical evidence.

In Section [SA-2.1](#), we recall asymptotic results for triangular arrays of random variables and some results of [Andrews and Guggenberger \(2019b\)](#) which will be used later in the proof of asymptotic similarity of the proposed tests .

In Section [SA-3](#), we provide the proof of the main results, Theorem [2.2](#) and Proposition [2.3](#), for the case of just-identified model with no non-constant exogenous controls, single endogenous regressor and single IV.

The finite sample validity of the PAR tests (Proposition [2.1](#) in the main paper) is proved in Section [SA-4](#).

Section [SA-5](#) establishes the uniform asymptotic similarity (Theorem [2.2](#) in the main paper). In Section [SA-5.1](#), we first define the drifting sequences of parameters needed for the uniform validity of the permutation tests. Section [SA-5.2](#) provides some asymptotic results for quantities that depend on the sample data only. Section [SA-5.3](#) presents the main results about the asymptotic distribution of key permutation quantities (Lemma [SA-5.1](#)) under the drifting parameter sequences.

Sections [SA-5.4](#), [SA-5.5](#) and [SA-5.6](#) prove the results for the PAR, PLM and PCLR tests respectively.

The asymptotic distributions of the test statistics under strong or semi-strong identification, and local alternatives (Proposition [2.3](#) in the main paper) are derived in Sections [SA-6](#).

Section [SA-7](#) provides the proof of Proposition [2.4](#) in the main paper about the asymptotic validity of the PAR tests under a diverging number of IVs.

*tpujee@yorku.ca

Section SA-8 presents the proofs of some of the results of Section SA-2 and others.

Additional simulation results on the test size and power, and empirical application are provided in Section SA-9 and SA-10.

Contents

SA-1	Notation and abbreviations	5
SA-2	Initial supplementary results	5
SA-2.1	Technical lemmas	5
SA-2.2	Some results of Andrews and Guggenberger (2019a,b)	8
SA-3	Proofs of Theorem 2.2 and Proposition 2.3: Just-identified model with single endogenous regressor and IV	9
SA-3.1	Drifting sequences of parameters	10
SA-3.2	Asymptotic size	11
SA-3.3	Asymptotic power	15
SA-4	Proof of Proposition 2.1 (Finite sample validity)	22
SA-5	Proof of Theorem 2.2 (Asymptotic similarity)	23
SA-5.1	Drifting sequences of parameters	23
SA-5.2	Probability limits of some key sample quantities	26
SA-5.2.1	Probability limits of $\hat{\mathcal{V}}_a(\theta_0)$, $\hat{\mathcal{V}}_b(\theta_0)$ and \hat{J}	27
SA-5.3	Asymptotic distributions of key permutation quantities	33
SA-5.4	PAR tests	35
SA-5.5	PLM test	40
SA-5.6	PCLR tests	42
SA-5.6.1	Asymptotic distribution of the PCLR statistics	44
SA-5.6.2	Completing the proof	45
SA-6	Proof of Proposition 2.3 (Asymptotic local power under strong identification)	47
SA-6.1	Probability limits of $\hat{\mathcal{V}}_a(\theta_0)$ and $\hat{\mathcal{V}}_b(\theta_0)$ under local alternatives	47
SA-6.2	AR and PAR statistics	49
SA-6.3	LM and PLM statistics	51
SA-6.4	CLR and PCLR statistics	53

SA-7	Proof of Proposition 2.4 (Many IVs)	54
SA-8	Proofs of supplementary results	65
SA-8.1	Lemma SA-2.2	65
SA-8.2	Lemma SA-2.3	65
SA-8.3	Lemma SA-2.5	65
SA-8.4	Lemma SA-5.1	65
SA-8.5	Theorem SA-5.3	74
SA-8.6	Lemma SA-5.2	78
SA-9	Additional simulation evidence	88
SA-9.1	Role of studentization	88
SA-9.2	Conditional heteroskedasticity with strong and very weak identification	89
SA-9.3	Power	89
SA-10	Additional empirical application	93

SA-1 Notation and abbreviations

Hereafter, $\stackrel{d}{=}$ denotes distributional equality, $1(\cdot)$ denotes the indicator function, ι is the $n \times 1$ vector of ones, and $\|\cdot\|$ denotes the Frobenius norm of a matrix. We use the abbreviations SLLN for the strong law of large of numbers, WLLN for the weak law of large numbers for independent $L^{1+\delta}$ -bounded random variables, CLT for the central limit theorem, CMT for the continuous mapping theorem, and SV and SVD for “singular value” and “singular value decomposition”, respectively.

Let $(\Omega_0, \mathcal{F}, P)$ be the probability triplet underlying the observations (y, Y, W, X) , and P^π denote the probability measure of a random permutation π uniformly distributed over \mathbb{G}_n . $\xrightarrow{p^\pi}$ and $\xrightarrow{d^\pi}$ denote the convergence in probability and convergence in distribution with respect to P^π .

$U_n = o_p(R_n)$ means that $U_n = R_n V_n$ and $V_n \xrightarrow{p} 0$ and $U_n = O_p(R_n)$ means that $U_n = R_n V_n$ with $V_n = O_p(1)$ i.e. for any $\epsilon > 0$, there exists a constant $M_\epsilon > 0$ such that $\limsup_{n \rightarrow \infty} P[|V_n| > M_\epsilon] < \epsilon$. Moreover, $o_{a.s.}(\cdot)$ and $O_{a.s.}(\cdot)$ stand for $o(\cdot)$ P -almost surely and $O(\cdot)$ P -almost surely, respectively. Whenever a stochastic convergence along a subsequence $\{m\}$ of $\{n\}$ is considered, the stochastic order notations are used for random variables indexed by $\{m\}$ with the corresponding properties.

We write “ $T_n^\pi \xrightarrow{d^\pi} T$ in P -probability” and “ $T_n^\pi \xrightarrow{d^\pi} T$ P -almost surely” if a permutation statistic $T_n^\pi \in \mathbb{R}^m$ converges in distribution to a random vector T on a set with probability approaching to 1 and with probability 1, respectively, i.e. $P^\pi[T_n^\pi \leq x] \xrightarrow{p} P[T \leq x]$ and $P^\pi[T_n^\pi \leq x] \xrightarrow{a.s.} P[T \leq x]$ as $n \rightarrow \infty$ for every x at which $x \mapsto P[T \leq x]$ is continuous. We write “ $U_n - V_n \xrightarrow{p^\pi} 0$ in P -probability”, if two random sequences U_n and V_n satisfy $P^\pi[\|U_n - V_n\| > \epsilon] \xrightarrow{p} 0$ as $n \rightarrow \infty$ for any $\epsilon > 0$.

SA-2 Initial supplementary results

SA-2.1 Technical lemmas

We use the following auxiliary results in the proof of asymptotic results. The first result is a version of the [Hoeffding \(1951\)](#)’s combinatorial CLT established by [Motoo \(1956\)](#) (see also [Schneller \(1988\)](#)) which is based on a Lindeberg-type condition.

Lemma SA-2.1 (Combinatorial CLT). *For n^2 real scalars $c_n(i, j)$, $i, j = 1, \dots, n$ with $n > 1$, and*

a random permutation π uniformly distributed over \mathbb{G}_n , consider the sum

$$S_n \equiv \sum_{i=1}^n c_n(i, \pi(i))$$

with $E_{P^\pi}[S_n] = n^{-1} \sum_{i=1}^n \sum_{j=1}^n c_n(i, j)$ and $\text{Var}_{P^\pi}[S_n] = \frac{1}{n-1} \sum_{i=1}^n \sum_{j=1}^n g_n^2(i, j)$, where

$$g_n(i, j) = c_n(i, j) - n^{-1} \sum_{i=1}^n c_n(i, j) - n^{-1} \sum_{j=1}^n c_n(i, j) + n^{-2} \sum_{i=1}^n \sum_{j=1}^n c_n(i, j).$$

If $g_n(i, j) \neq 0$ for some (i, j) , so that $\text{Var}_{P^\pi}[S_n] > 0$, and for any $\epsilon > 0$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n \frac{g_n^2(i, j)}{\frac{1}{n-1} \sum_{i=1}^n \sum_{j=1}^n g_n^2(i, j)} 1 \left(\frac{|g_n(i, j)|}{\left(\frac{1}{n-1} \sum_{i=1}^n \sum_{j=1}^n g_n^2(i, j) \right)^{1/2}} > \epsilon \right) = 0, \quad (\text{SA-2.1})$$

then as $n \rightarrow \infty$

$$\frac{S_n - E_{P^\pi}[S_n]}{\sqrt{\text{Var}_{P^\pi}[S_n]}} \xrightarrow{d^\pi} N[0, 1].$$

The next result is a triangular array SLLN.

Lemma SA-2.2 (Corollary to Theorem 1 of [Hu et al. \(1989\)](#)). *Let $\{Z_{nk} : k = 1, \dots, n; n = 1, 2, \dots\}$ be an array of i.i.d. random variables such that $E[|Z_{nk}|^{2p}] < \infty$ for some $1 \leq p < 2$. Then,*

$$n^{-1/p} \sum_{k=1}^n (Z_{nk} - E[Z_{nk}]) \xrightarrow{a.s.} 0. \quad (\text{SA-2.2})$$

We will also use the following fact.

Lemma SA-2.3. *If $\{X_i\}_{i=1}^n$ and X are defined on the probability space $(\Omega_0, \mathcal{F}, P)$ and $X_n \xrightarrow{p} X$, then $X_n \xrightarrow{p^\pi} X$ in P -probability.*

We will use the following result of [Pham et al. \(1989\)](#) to establish the limiting distributions of the permutation statistics under growing k . See [Barbour and Eagleson \(1986\)](#), [Zhao et al. \(1997\)](#) and [Barbour and Chen \(2005\)](#) for related results. Let $\sum_{i,j}$ denote $\sum_{i=1}^n \sum_{j=1}^n$.

Lemma SA-2.4 (Theorem 3.1 of [Pham et al. \(1989\)](#)). *Let $\{a_{ij}\}_{i,j=1}^n$ and $\{b_{ij}\}_{i,j=1}^n$ be arrays of*

scalars such that $a_{ii} = b_{ii} = 0$, $a_{ij} = a_{ji}$ for $i, j = 1, \dots, n$. Suppose that

$$\max \left(\sum_{j=1}^n |a_{ij}| \right) = O \left(\frac{\sum_{i,j} a_{ij}^2}{n \max_{i,j} |a_{ij}|} \right), \quad (\text{SA-2.3})$$

$$\frac{\sum_{i,j} a_{ij}^2}{n^2 \max_{i,j} a_{ij}^2} \rightarrow 0 \text{ as } n \rightarrow \infty, \quad (\text{SA-2.4})$$

$$\sum_{i,j} b_{ij} = 0, \quad (\text{SA-2.5})$$

$$\sum_{i=1}^n b_{i+}^2 = O \left(\sum_{i,j} b_{ij}^2 \right), \quad b_{i+} \equiv \sum_{j=1}^n b_{ij}. \quad (\text{SA-2.6})$$

Then,

$$\frac{\sum_{i \neq j} a_{ij} b_{\pi(i)\pi(j)}}{\left[2n^{-2} \left(\sum_{i,j} a_{ij}^2 \right) \left(\sum_{i,j} b_{ij}^2 \right) \right]^{1/2}} \xrightarrow{d\pi} N[0, 1].$$

The following lemma is standard and will be used in the proof of Proposition 2.4.

Lemma SA-2.5. *For real-valued vectors $a_1, \dots, a_n \in \mathbb{R}^k$ and scalars b_1, \dots, b_n such that $\sum_{i=1}^n b_i = 0$, and π uniformly distributed over \mathbb{G}_n , it holds that*

$$\mathbb{E}_{P^\pi} \left[\left\| \sum_{i=1}^n a_i b_{\pi(i)} \right\|^2 \right] = \frac{1}{n-1} \left(\sum_{i=1}^n \|a_i\|^2 - n \left\| \sum_{i=1}^n a_i \right\|^2 \right) \left(\sum_{i=1}^n b_i^2 \right). \quad (\text{SA-2.7})$$

To derive the asymptotic distribution of the AR statistic under a diverging number of IVs, we use Lemma 7 of [Andrews and Stock \(2007b\)](#) stated below, which, in turn, is established using Theorem 1 of [Hall \(1984\)](#) among other arguments.

Lemma SA-2.6 (Corollary to Lemma 7 of [Andrews and Stock \(2007a\)](#)). *Suppose that*

(a) $\{\xi_{ni}, \hat{\Sigma}_{ni}, i \leq n, n \geq 1\}$ is a triangular array of random vectors and matrices that satisfies

(i) $\xi_{ni} \in \mathbb{R}^k$ for all $i \leq n$, where $k = k_n$;

(ii) for each $n \geq 1$, $\{\xi_{ni}\}$ are i.i.d. across $i \leq n$;

(iii) $\mathbb{E}[\xi_{ni}] = \mu_{n\xi}$, $\lambda_{n\xi}^* = n\mu_{n\xi}' \Sigma_{n\xi}^{-1} \mu_{n\xi}$;

(iv) $V[\xi_{ni}] = \Sigma_{n\xi} \in \mathbb{R}^{k \times k}$, and $\hat{\Sigma}_{n\xi}$ is a random $k \times k$ matrix that satisfies $\|\hat{\Sigma}_{n\xi} - \Sigma_{n\xi}\| = o_p(k^{-1/2})$;

(v) $\sup_{\ell \leq k, n \geq 1} \mathbb{E}[\xi_{ni\ell}^4] < \infty$, where $\xi_{ni} = [\xi_{ni1}, \dots, \xi_{nik}]'$;

(vi) $k \rightarrow \infty$ as $n \rightarrow \infty$;

(vii) $\frac{k^3}{n} \rightarrow 0$;

(viii) $\inf_{n \geq 1} \lambda_{\min}(\Sigma_{n\xi}) > 0$;

(b) $\xi_{ni} = \xi_{n1i} + D_{n\xi}\xi_{n2i}$, where $D_{n\xi} \in \mathbb{R}^{k \times k}$ is a non-random matrix;

(c) $\{\xi_{n2i} : i \leq n\}$ are i.i.d. across $i \leq n$ with $\|E[\xi_{n2i}]\|^2 = O(k^2/n)$;

(d) $\hat{D}_{n\xi} \in \mathbb{R}^{k \times k}$ is a random matrix that satisfies $\|\hat{D}_{n\xi} - D_{n\xi}\| = o_p(k^{-1})$;

(e) $\hat{\xi}_{ni} = \xi_{n1i} + \hat{D}_{n\xi}\xi_{n2i}$.

Then,

$$(2k)^{-1/2} \left(n^{-1/2} \sum_{i=1}^n [\hat{\Sigma}_{n\xi}^{-1/2} \hat{\xi}_{ni}]' n^{-1/2} \sum_{j=1}^n [\hat{\Sigma}_{n\xi}^{-1/2} \hat{\xi}_{nj}] - k \right) - (2k)^{-1/2} \lambda_{n\xi}^* \xrightarrow{d} N[0, 1]. \quad (\text{SA-2.8})$$

SA-2.2 Some results of Andrews and Guggenberger (2019a,b)

For a non-random matrix $\mathcal{J} \in \mathbb{R}^{(k-q) \times (d-q)}$, let $r_{k,d,q}(\mathcal{J}, 1 - \alpha)$ denote the $1 - \alpha$ quantile of

$$\mathcal{Z}'\mathcal{Z}^* - \lambda_{\min}[(\mathcal{J}, \mathcal{Z}_2^*)'(\mathcal{J}, \mathcal{Z}_2^*)], \quad \mathcal{Z}^* = [\mathcal{Z}_1^*, \mathcal{Z}_2^{*'}]' \sim N[0_{k \times 1}, I_k], \quad \mathcal{Z}_1^* \in \mathbb{R}^q, \quad \mathcal{Z}_2^* \in \mathbb{R}^{k-q}. \quad (\text{SA-2.9})$$

The following results due to Andrews and Guggenberger (2019b) will be used in the proof of the asymptotic similarity of the PCLR tests.

Lemma SA-2.7 (Lemma 27.3 of Andrews and Guggenberger (2019b)). *Let $\mathcal{Z} = [\mathcal{Z}_1', \mathcal{Z}_2']' \sim N[0_{k \times 1}, I_k]$, $\mathcal{Z}_1 \in \mathbb{R}^q$, $\mathcal{Z}_2 \in \mathbb{R}^{k-q}$, and $\Upsilon(\tau^c) \equiv [\text{diag}(\tau^c), 0_{(d-q) \times (k-d)}] \in \mathbb{R}^{(k-q) \times (d-q)}$, where $\tau^c \equiv (\tau_{(q+1)}^c, \dots, \tau_d^c)' \in \mathbb{R}^{d-q}$ with $\tau_{q+1}^c \geq \dots, \tau_d^c \geq 0$, and $k \geq d \geq 1$ and $0 \leq q \leq d$. Then, the distribution function of*

$$\text{CLR}_\infty(\tau^c) \equiv \mathcal{Z}'\mathcal{Z} - \lambda_{\min}[(\Upsilon(\tau^c), \mathcal{Z}_2)'(\Upsilon(\tau^c), \mathcal{Z}_2)] \quad (\text{SA-2.10})$$

is continuous and strictly increasing at its $1 - \alpha$ quantile $r_{k,d,q}(\Upsilon(\tau^c), 1 - \alpha)$ for all $\alpha \in (0, 1)$.

Lemma SA-2.8 (Lemma 16.2 of Andrews and Guggenberger (2019b)). *Let \mathcal{J}^c be a $(k-q) \times (d-q)$ matrix with the singular value decomposition $A^c \Upsilon^c B^{c'}$, where A^c is a $(k-q) \times (k-q)$ orthogonal matrix of eigenvectors of $\mathcal{J}^c \mathcal{J}^{c'}$, B^c is a $(d-q) \times (d-q)$ orthogonal matrix of eigenvectors of*

$\mathcal{J}'\mathcal{J}^c$, and Υ^c is the $(k-q) \times (d-q)$ matrix with the $d-q$ singular values $\tau_{q+1}^c \geq \dots \geq \tau_d^c$ of \mathcal{J}^c as its first $d-q$ diagonal elements and zeros elsewhere. Then, for $r_{k,d,q}(\cdot, 1-\alpha)$ defined in (SA-2.9)

$$r_{k,d,q}(\mathcal{J}^c, 1-\alpha) = r_{k,d,q}(\Upsilon^c, 1-\alpha),$$

and the distribution of the random variable in (SA-2.10) is identical to that of (SA-2.9) with \mathcal{J}^c in place of \mathcal{J} .

Using the notations introduced in Section SA-5.1 and onwards, a result in Lemma 16.4 of Andrews and Guggenberger (2019b) reads as follows:

$$n^{1/2}[\hat{m}, \hat{J} - G_n, H_n \hat{J} U_n T_n] \xrightarrow{d} [m_h, J_h, \Delta_h], \quad (\text{SA-2.11})$$

where (J_h, Δ_h) and m_h are independent,

$$\begin{bmatrix} m_h \\ \text{vec}(J_h) \end{bmatrix} \sim N \left[0_{k \times (d+1)}, \begin{pmatrix} h_{5,m} & 0_{k \times dk} \\ 0_{k \times dk} & \Phi_h \end{pmatrix} \right], \quad \Phi_h \equiv \lim_{n \rightarrow \infty} (\Sigma_n^G - C_n' \Sigma_n^{-1} C_n)$$

whenever the limit exists, and

$$\begin{aligned} \Delta_h &\equiv [\Delta_{h,q}, \Delta_{h,d-q}] \in \mathbb{R}^{k \times d}, \quad \Delta_{h,q} \equiv h_{3,q} \in \mathbb{R}^{k \times q}, \quad \Delta_{h,d-q} \equiv h_3 \tilde{h}_{1,d-q} + h_{5,m}^{-1/2} J_h h_{91} h_{2,d-q} \in \mathbb{R}^{k \times (d-q)}, \\ \tilde{h}_{1,d-q} &\equiv \begin{bmatrix} 0_{q \times (d-q)} \\ \text{diag}(h_{1,q+1}, \dots, h_{1,d}) \\ 0_{(k-d) \times (d-q)} \end{bmatrix}, \quad h_{91} \equiv ((\theta_0, I_d) \Omega^\epsilon(h_9) (\theta_0, I_d)')^{1/2}, \quad \Omega = \Omega(\Sigma, K(\mathcal{V})). \end{aligned} \quad (\text{SA-2.12})$$

SA-3 Proofs of Theorem 2.2 and Proposition 2.3: Just-identified model with single endogenous regressor and IV

This section presents the proof of the asymptotic validity of the permutation tests in the just-identified case with single endogenous regressor, Y , and single IV, W . This case is encountered often in practice. In the just-identified IV regression, the AR, LM, CLR_a and CLR_b statistics are identical, and so are the PAR, PLM, PCLR_a and PCLR_b statistics.

Andrews et al. (2020) show the uniform asymptotic similarity of the asymptotic AR, LM and CLR statistics in the IV model with one endogenous regressor. By contrast, we derive the asymptotic distribution of the AR statistic under local $\theta = \theta_0 + n^{-1/2} h_\theta$, with h_θ fixed, and fixed $\theta \neq \theta_0$ alternatives, and consider permutation tests.

Compared to the case of multiple endogenous variables, the proof of the asymptotic validity results and the description of the drifting parameters are considerably simpler as there are only two possible identification categories. For simplicity, we assume that there are no exogenous controls except for a constant regressor. The proof can be extended directly to incorporate exogenous covariates by replacing $Z_i^* \equiv W_i - E_P[W_i]$ by $Z_i^* \equiv W_i - E_P[W_i X_i'] (E_P[X_i X_i'])^{-1} X_i$ below, and modifying the corresponding drifting parameters and their limits accordingly. We will use Lemma SA-5.1 for deriving the asymptotic distribution of the PAR₂ statistic.

SA-3.1 Drifting sequences of parameters

Define

$$\lambda_{1,P}^* \equiv |\Gamma| \in \mathbb{R}, \quad (\text{SA-3.1})$$

$$\lambda_{3,P}^* \equiv \Gamma/|\Gamma| \in \mathbb{R}, \quad (\text{SA-3.2})$$

$$\lambda_{5,P}^* \equiv E_P \left[\begin{pmatrix} Z_i^* u_i \\ Z_i^* V_i \end{pmatrix} \begin{pmatrix} Z_i^* u_i \\ Z_i^* V_i \end{pmatrix}' \right] \in \mathbb{R}^{2 \times 2}, \quad (\text{SA-3.3})$$

$$\lambda_{7,P}^* \equiv \text{Var}_P[Z_i^*] = \text{Var}_P[W_i] \in \mathbb{R}, \quad (\text{SA-3.4})$$

$$\lambda_{8,P}^* \equiv \Sigma_P^e = \text{Var}_P[(u_i, V_i)'] \in \mathbb{R}^{2 \times 2}, \quad (\text{SA-3.5})$$

$$\lambda_{10,P}^* \equiv P, \quad (\text{SA-3.6})$$

$$\lambda_P^* \equiv (\lambda_{1,P}^*, \lambda_{3,P}^*, \lambda_{5,P}^*, \lambda_{7,P}^*, \lambda_{8,P}^*, \gamma, \Psi, \lambda_{10,P}^*). \quad (\text{SA-3.7})$$

¹ Let $\lambda_{1,P}^* \equiv 1$ if $\Gamma = 0$. Given the model, under $H_0 : \theta = \theta_0$, the distribution of the data is determined by $(\lambda_{1,P}^*, \lambda_{3,P}^*, \lambda_{5,P}^*, \lambda_{7,P}^*, \gamma, \Psi, \lambda_{10,P}^*)'$. The parameter $\lambda_{8,P}^*$ is needed for the asymptotic properties of the permutation statistics. We consider the limits of the drifting parameters introduced above:

$$n^{1/2} \lambda_{1,P_n}^* \rightarrow h_1^* \geq 0, \quad (\text{SA-3.8})$$

$$\lambda_{3,P_n}^* \rightarrow h_3^*, \quad (\text{SA-3.9})$$

$$\lambda_{5,P_n}^* \rightarrow h_5^* = \begin{bmatrix} h_{5,m}^* & h_{5,2}^* \\ h_{5,2}^* & h_{5,3}^* \end{bmatrix}, \quad (\text{SA-3.10})$$

¹These parameters are indexed by their correspondences with the drifting parameters introduced in Section SA-5.1.

$$\lambda_{7,P_n}^* \rightarrow h_7^*, \quad (\text{SA-3.11})$$

$$\lambda_{8,P_n}^* \rightarrow h_8^* \equiv \Sigma^e = \begin{bmatrix} \sigma^2 & \Sigma^{uV} \\ \Sigma^{Vu} & \Sigma^V \end{bmatrix}, \quad (\text{SA-3.12})$$

where, for instance, the upper-left element of h_5^* , $h_{5,m}^* = \Sigma \in \mathbb{R}$, is the limit of Σ_P . Define

$$\begin{aligned} \Lambda^* &\equiv \{\lambda_P^* : P \in \mathcal{P}_0^{\text{PAR}_2}\}, \\ h_n^*(\lambda_P^*) &\equiv [n^{1/2}\lambda_{1,P}^*, \lambda_{3,P}^*, \text{vech}(\lambda_{5,P}^*)', \lambda_{7,P}^*, \text{vech}(\lambda_{8,P}^*)']', \end{aligned} \quad (\text{SA-3.13})$$

and

$$\mathcal{H}^* \equiv \{h^* \in (\mathbb{R} \cup \{\pm\infty\})^9 : h_{w_n}^*(\lambda_{P_{w_n}}^*) \rightarrow h^* \text{ for some subsequence } \{w_n\} \text{ of } \{n\}\} \quad (\text{SA-3.14})$$

$$\text{and some sequence } \{\lambda_{P_{w_n}}^* \in \Lambda^* : n \geq 1\}. \quad (\text{SA-3.15})$$

The following assumption corresponds to Assumption B* of [Andrews and Guggenberger \(2019b\)](#) and [Andrews et al. \(2020\)](#), and is used for verifying the asymptotic similarity of the tests (see Corollary 2.1 of [Andrews et al. \(2020\)](#) and also Assumption 2).

Assumption 1. For any subsequence $\{w_n\}$ of $\{n\}$ and any sequence $\{\lambda_{P_{w_n}}^* \in \Lambda : n \geq 1\}$ for which $h_{w_n}^*(\lambda_{P_{w_n}}^*) \rightarrow h^* \in \mathcal{H}^*$, $E_{P_{w_n}}[\phi_{w_n}^{\text{PAR}_2}] \rightarrow \alpha$ for some $\alpha \in (0, 1)$.

In the next section, we will verify this assumption. Also, for notational convenience, we will write $\hat{m} = \hat{m}(\theta_0)$, $\hat{\Sigma} = \hat{\Sigma}(\theta_0)$ and $\hat{\Sigma}^\pi = \hat{\Sigma}^\pi(\theta_0)$.

SA-3.2 Asymptotic size

In the just-identified model with $d = k = 1$ and $X = \iota$, the Jacobian $\hat{J}(\theta_0)$ in the PLM statistic is scalar and hence drops out of the statistic by the scale invariance of the projection matrix. This makes PAR₂ and PLM statistics identical. Furthermore, both $\hat{\mathcal{S}}^\pi$ and $\hat{\mathcal{T}}$ are scalars, so

$$\lambda_{\min}[(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}})'(\hat{\mathcal{S}}^\pi, \hat{\mathcal{T}})] = \frac{1}{2} \left(\hat{\mathcal{S}}^{\pi 2} + \hat{\mathcal{T}}^2 - \sqrt{(\hat{\mathcal{S}}^{\pi 2} + \hat{\mathcal{T}}^2)^2 + 4(\hat{\mathcal{S}}^{\pi 2} \hat{\mathcal{T}}^2 - (\hat{\mathcal{S}}^\pi \hat{\mathcal{T}})^2)} \right) = 0. \quad (\text{SA-3.16})$$

As a result, the PCLR statistics simplify to $\hat{\mathcal{S}}^{\pi 2}$, hence is identical to the PAR₂ statistic. Also, the PAR₁ and PAR₂ test statistics are identically distributed because, letting $\bar{W} \equiv n^{-1} \sum_{i=1}^n W_i$, $W_\pi' M_\iota u(\theta_0) = \sum_{i=1}^n (W_{\pi(i)} - \bar{W}) u_i(\theta_0) = \sum_{i=1}^n (W_i - \bar{W}) u_{\pi^{-1}(i)}(\theta_0) = Z' u_{\pi^{-1}}(\theta_0)$ and

$$W_\pi' M_\iota \tilde{\Sigma}_u(\theta_0) M_\iota W_\pi = \sum_{i=1}^n (W_{\pi(i)} - \bar{W})(W_{\pi(i)} - \bar{W})' \tilde{u}_i(\theta_0)^2 = \sum_{i=1}^n (W_i - \bar{W})(W_i - \bar{W})' \tilde{u}_{\pi^{-1}(i)}(\theta_0)^2,$$

where π^{-1} is the inverse of π i.e. $\pi \circ \pi^{-1}$ is identity permutation, and is uniformly distributed over \mathbb{G}_n . Therefore, it suffices to establish the asymptotic size of the PAR_2 statistic in this just-identified case.

[Andrews et al. \(2020\)](#) show that

$$n^{1/2}\hat{m} = n^{1/2}\hat{m}(\theta_0) = n^{-1/2}Z'u(\theta_0) = n^{-1/2}\sum_{i=1}^n Z_i^* u_i + Dn^{-1/2}\sum_{i=1}^n u_i \xrightarrow{d} m_h \sim N[0, h_{5,m}^*], \quad (\text{SA-3.17})$$

and $\text{AR} = n\hat{m}'\hat{\Sigma}^{-1}\hat{m} \xrightarrow{d} \text{AR}_\infty \equiv (\Sigma^{-1/2}m_h)^2 \sim \chi_1^2$ with $\Sigma = h_{5,m}^*$.

Next we establish the asymptotic distribution of the PAR_2 . To this end, we first determine the limit of the covariance matrix estimator.

The limit of $\hat{\Sigma}^\pi$ $= n^{-1}\sum_{i=1}^n Z_i^2 \tilde{u}_{\pi(i)}(\theta_0)^2$. Note that by Lemma S.3.4 of [DiCiccio and Romano \(2017\)](#) (see also [Hoeffding \(1951\)](#)),

$$\mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W] = (n^{-1}Z'Z) \left(n^{-1}\sum_{i=1}^n \tilde{u}_i(\theta_0)^2 \right) \xrightarrow{a.s.} \sigma^2 h_7^*, \quad (\text{SA-3.18})$$

where the convergence follows from the CMT, and

$$n^{-1}Z'Z = n^{-1}W'W - n^{-1}W'\iota n^{-1}\iota'W \xrightarrow{a.s.} h_7^* \equiv \lim_{n \rightarrow \infty} \text{Var}[W_i], \quad (\text{SA-3.19})$$

$$n^{-1}\sum_{i=1}^n \tilde{u}_i(\theta_0)^2 = n^{-1}u'u - (n^{-1}u'\iota)^2 \xrightarrow{a.s.} \sigma^2 \equiv \lim_{n \rightarrow \infty} \text{Var}[u_i], \quad (\text{SA-3.20})$$

which, in turn, hold by the SLLN (Lemma [SA-2.2](#)), CMT, ([SA-3.11](#)) and ([SA-3.12](#)). Since $n^{-1}\iota'u \xrightarrow{p} 0$ and $n^{-1}\sum_{i=1}^n u_i^4 - \mathbb{E}[u_i^4] \xrightarrow{p} 0$, by Minkowski's inequality, the triangular array WLLN and the CMT

$$n^{-1}\sum_{i=1}^n Z_i^4 = n^{-1}\sum_{i=1}^n (W_i - \bar{W})^4 \leq 8n^{-1}\sum_{i=1}^n W_i^4 + 8\bar{W}^4 = O_p(1), \quad (\text{SA-3.21})$$

$$n^{-1}\sum_{i=1}^n \tilde{u}_i(\theta_0)^4 \leq 8n^{-1}\sum_{i=1}^n u_i^4 + 8\bar{u}^4 = O_p(1). \quad (\text{SA-3.22})$$

Therefore, using (SA-3.21) and (SA-3.22), we have

$$\begin{aligned}
\text{Var}_{P^\pi}[\hat{\Sigma}^\pi|y, Y, W] &= \text{Var}_{P^\pi} \left[n^{-1} \sum_{i=1}^n Z_i^2 \tilde{u}_{\pi(i)}(\theta_0)^2 | y, Y, W \right] \\
&= \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n Z_i^4 - \left(n^{-1} \sum_{i=1}^n Z_i^2 \right)^2 \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 - \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^2 \right)^2 \right) \\
&\leq \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n Z_i^4 \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 \right) \\
&\xrightarrow{p} 0,
\end{aligned} \tag{SA-3.23}$$

where the second equality follows from Lemma S.3.4 of DiCiccio and Romano (2017). From (SA-3.23) and Chebyshev's inequality, for any $\epsilon > 0$

$$P^\pi[|\hat{\Sigma}^\pi - \mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi|y, Y, W]| > \epsilon] \leq \text{Var}_{P^\pi}[\hat{\Sigma}^\pi|y, Y, W]/\epsilon^2 \xrightarrow{p} 0. \tag{SA-3.24}$$

Thus, $\hat{\Sigma}^\pi - \mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi|y, Y, W] \xrightarrow{p^\pi} 0$ in P -probability. The latter combined with (SA-3.18) and the triangle inequality yields $|\hat{\Sigma}^\pi - \sigma^2 h_7^*| \leq |\mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi|y, Y, W] - \sigma^2 h_7^*| + |\hat{\Sigma}^\pi - \mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi|y, Y, W]| \xrightarrow{p^\pi} 0$ in P -probability, hence

$$n^{-1} \sum_{i=1}^n Z_i^2 \tilde{u}_{\pi(i)}(\theta_0)^2 \xrightarrow{p^\pi} \sigma^2 h_7^* \quad \text{in } P\text{-probability.} \tag{SA-3.25}$$

We next determine the asymptotic distribution of $n^{1/2} \hat{m}^\pi = n^{-1/2} Z' u_\pi(\theta_0)$.

The asymptotic distribution of $n^{1/2} \hat{m}^\pi = n^{-1/2} W' \tilde{u}_\pi(\theta_0)$. By Lemma SA-5.1 (and Lemma SA-2.1),

$$\frac{n^{-1/2} W' \tilde{u}_\pi(\theta_0)}{((n-1)^{-1} \tilde{u}(\theta_0)' \tilde{u}(\theta_0) n^{-1} W' M_\ell W)^{1/2}} \xrightarrow{d^\pi} N[0, 1] \quad \text{in } P\text{-probability.} \tag{SA-3.26}$$

The Lyapunov-type sufficient conditions for (SA-3.30) are given in (SA-5.68) and (SA-5.69) which can be restated as follows: for some $\delta > 0$

$$n^{-1} \sum_{i=1}^n |\tilde{u}_i(\theta_0)|^{2+\delta/2} = O_{a.s.}(1), \tag{SA-3.27}$$

$$n^{-1} \sum_{i=1}^n |W_i|^{2+\delta/2} = O_{a.s.}(1). \tag{SA-3.28}$$

(SA-3.27) is verified by an argument similar to (SA-3.22) and the SLLN (Lemma SA-2.2). (SA-3.28) holds by the SLLN. From (SA-3.19) and (SA-3.20),

$$(n-1)^{-1} \tilde{u}(\theta_0)' \tilde{u}(\theta_0) n^{-1} W' M_\ell W \xrightarrow{a.s.} \sigma^2 h_7^*. \tag{SA-3.29}$$

From (SA-3.26), (SA-3.29) and Slutsky's lemma,

$$n^{-1/2}W'\tilde{u}_\pi(\theta_0) \xrightarrow{d^\pi} m_h^\pi \sim N[0, \sigma^2 h_7^*] \quad \text{in } P\text{-probability.} \quad (\text{SA-3.30})$$

Then, (SA-3.25), (SA-3.30) and the CMT together imply

$$\text{PAR}_2 \xrightarrow{d^\pi} m_h^{\pi'} (\sigma^2 h_7^*)^{-1} m_h^\pi \sim \chi_1^2 \quad \text{in } P\text{-probability.} \quad (\text{SA-3.31})$$

Let $r_{1-\alpha}(\chi_1^2)$ denote the $1 - \alpha$ quantile of χ_1^2 distribution. Next we will show that $\text{PAR}_{2(r)} \xrightarrow{p} r_{1-\alpha}(\chi_1^2)$. For π_2, \dots, π_N i.i.d. uniformly distributed over \mathbb{G}_n , let

$$\bar{F}_N^{\text{PAR}_2}(x) \equiv \frac{1}{N-1} \sum_{i=2}^N 1(\text{AR}^{\pi_i} \leq x). \quad (\text{SA-3.32})$$

By Theorem 12.2.18 of [Lehmann and Romano \(2005\)](#), for some universal constant C_0 and any positive integer $N \geq 2$

$$P^\pi \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - \hat{F}_n^{\text{PAR}_1}(x) \right| > \epsilon \right] \leq C_0 \exp(-2(N-1)\epsilon^2). \quad (\text{SA-3.33})$$

Using the law of iterated expectations and letting $N \rightarrow \infty$ in (SA-3.33),

$$P \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_2}(x) - \hat{F}_n^{\text{PAR}_2}(x) \right| > \epsilon \right] = E_P \left[P^\pi \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_2}(x) - \hat{F}_n^{\text{PAR}_2}(x) \right| > \epsilon \right] \right] \rightarrow 0. \quad (\text{SA-3.34})$$

By the triangle inequality, (SA-3.31) and (SA-3.34)

$$\begin{aligned} \sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_2}(x) - P[\chi_1^2 \leq x] \right| &\leq \sup_{x \in \mathbb{R}} \left| \hat{F}_n^{\text{PAR}_2}(x) - P[\chi_1^2 \leq x] \right| + \sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_2}(x) - \hat{F}_n^{\text{PAR}_2}(x) \right| \\ &\xrightarrow{p} 0. \end{aligned} \quad (\text{SA-3.35})$$

Therefore, the permutation distribution $\hat{F}_N^{\text{PAR}_2}(x)$ of PAR_2 on \mathbb{G}'_n satisfies

$$\tilde{F}_N^{\text{PAR}_2}(x) = N^{-1} 1(\text{AR}^{\pi_1} \leq x) + \frac{N-1}{N} \bar{F}_N^{\text{PAR}_2}(x) \xrightarrow{p} P[\chi_1^2 \leq x]. \quad (\text{SA-3.36})$$

Since the cdf of χ_1^2 distribution is continuous and strictly increasing at its $1 - \alpha$ quantile, we have $\text{PAR}_{2(r)} \xrightarrow{p} r_{1-\alpha}(\chi_1^2)$ by Lemma 11.2.1 of [Lehmann and Romano \(2005\)](#). By definition, $E_P[\phi_n^{\text{PAR}_2}] = P[\text{AR} > \text{PAR}_{2(r)}] + a^{\text{PAR}_2} P[\text{AR} = \text{PAR}_{2(r)}]$, hence

$$P[\text{AR} > \text{PAR}_{2(r)}] \leq E_P[\phi_n^{\text{PAR}_2}] \leq P[\text{AR} \geq \text{PAR}_{2(r)}].$$

By Slutsky's lemma (Corollary 11.2.3 of [Lehmann and Romano \(2005\)](#)), we have $\lim_{n \rightarrow \infty} P[\text{AR} > \text{PAR}_{2(r)}] = \lim_{n \rightarrow \infty} P[\text{AR} \geq \text{PAR}_{2(r)}] = P[\text{AR}_\infty > r_{1-\alpha}(\chi_1^2)]$ because of the continuity of the χ_1^2 distribution function at its $1 - \alpha$ quantile. Therefore,

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PAR}_2}] = P[\text{AR}_\infty > r_{1-\alpha}(\chi_1^2)] = \alpha.$$

The results above hold for any subsequence $\{w_n\}$ of $\{n\}$. Thus, Assumption 1 is verified for the PAR_2 statistic. By Proposition 16.3 of [Andrews and Guggenberger \(2019b\)](#), we obtain the desired result.

SA-3.3 Asymptotic power

We derive the asymptotic power functions in the two exhaustive cases $h_1^* < \infty$ (Case 1) and $h_1^* = \infty$ (Case 2) below. Let $D \equiv E_P[W_i] - \bar{W}$. On noting that for $2 \leq q \leq 4$ and $0 \leq q + r + s \leq 4$, $E_P[(|Z_i^*|^q |u_i|^r |V_i|^s)^{1+\delta/4}] < \infty$ by Cauchy-Schwarz inequality (the case of $q = 4$ is trivial because $r = s = 0$), by the triangular array WLLN

$$D \xrightarrow{p} 0, \tag{SA-3.37}$$

$$\bar{u} \equiv n^{-1} \sum_{i=1}^n u_i \xrightarrow{p} 0, \tag{SA-3.38}$$

$$\bar{V} \equiv n^{-1} \sum_{i=1}^n V_i \xrightarrow{p} 0, \tag{SA-3.39}$$

$$n^{-1} \sum_{i=1}^n |Z_i^*|^q |u_i|^r |V_i|^s - E_P[|Z_i^*|^q |u_i|^r |V_i|^s] \xrightarrow{p} 0, \quad 2 \leq q \leq 4, \quad 0 \leq q + r + s \leq 4, \tag{SA-3.40}$$

$$n^{-1} \sum_{i=1}^n Z_i^{*2} u_i^2 - E_P[Z_i^{*2} u_i^2] \xrightarrow{p} 0, \tag{SA-3.41}$$

$$n^{-1} \sum_{i=1}^n Z_i^{*2} V_i^2 - E_P[Z_i^{*2} V_i^2] \xrightarrow{p} 0, \tag{SA-3.42}$$

$$n^{-1} \sum_{i=1}^n Z_i^{*2} u_i V_i - E_P[Z_i^{*2} V_i u_i] \xrightarrow{p} 0, \tag{SA-3.43}$$

$$n^{-1} \sum_{i=1}^n Z_i' Z_i - E_P[Z_i^{*2}] = n^{-1} \sum_{i=1}^n (Z_i^* + D)^2 - E_P[Z_i^{*2}] \xrightarrow{p} 0. \tag{SA-3.44}$$

Since $\tilde{u}(\theta_0) = M_\iota(y - Y\theta_0) = M_\iota(W\Gamma + V)(\theta - \theta_0) + M_\iota u$, we have

$$\tilde{u}_i(\theta_0) = (Z_i^* + D)\Gamma(\theta - \theta_0) + (V_i - \bar{V})(\theta - \theta_0) + u_i - \bar{u}, \quad (\text{SA-3.45})$$

$$\begin{aligned} \hat{\Sigma} &= \hat{\Sigma}(\theta_0) = n^{-1} \sum_{i=1}^n Z_i^2 \tilde{u}_i(\theta_0)^2 \\ &= n^{-1} \sum_{i=1}^n (Z_i^* + D)^2 ((Z_i^* + D)\Gamma(\theta - \theta_0) + (V_i - \bar{V})(\theta - \theta_0) + u_i - \bar{u})^2. \end{aligned} \quad (\text{SA-3.46})$$

Case 1: $h_1^* < \infty$. In this case, using (SA-3.1) and (SA-3.2), we have

$$\Gamma = |\Gamma|\Gamma/|\Gamma| = \lambda_{1,P}^* \lambda_{3,P}^* = n^{-1/2} (n^{1/2} \lambda_{1,P}^*) \lambda_{3,P}^* \rightarrow 0 \times h_1^* h_3^* = 0. \quad (\text{SA-3.47})$$

AR statistic. Let θ be fixed. Then, using the triangle inequality, (SA-3.38)-(SA-3.43) and the CMT, we obtain after some algebra that

$$\begin{aligned} &\hat{\Sigma} - \mathbb{E}_P[Z_i^{*2}(V_i(\theta - \theta_0) + u_i)^2] \\ &= n^{-1} \sum_{i=1}^n (Z_i^* + D)^2 ((Z_i^* + D)\Gamma(\theta - \theta_0) + (V_i - \bar{V})(\theta - \theta_0) + u_i - \bar{u})^2 \\ &\quad - \mathbb{E}_P[Z_i^{*2}(V_i(\theta - \theta_0) + u_i)^2] \\ &\xrightarrow{p} 0. \end{aligned} \quad (\text{SA-3.48})$$

Since $\mathbb{E}_P[Z_i^{*2}(V_i(\theta - \theta_0) + u_i)^2] - (h_{5,m}^* + 2h_{5,2}^*(\theta - \theta_0) + h_{5,3}^*(\theta - \theta_0)^2) \rightarrow 0$ using (SA-3.10), we have

$$\hat{\Sigma} - (h_{5,m}^* + 2h_{5,2}^*(\theta - \theta_0) + h_{5,3}^*(\theta - \theta_0)^2) \xrightarrow{p} 0. \quad (\text{SA-3.49})$$

By the bivariate Lyapunov's CLT followed by the CMT, and (SA-3.49)

$$n^{-1/2} Z'(V(\theta - \theta_0) + u) \xrightarrow{d} N[0, h_{5,m}^* + 2h_{5,2}^*(\theta - \theta_0) + h_{5,3}^*(\theta - \theta_0)^2]. \quad (\text{SA-3.50})$$

Moreover, by (SA-3.8), (SA-3.9), (SA-3.11), (SA-3.44), (SA-3.50) and Slutsky's lemma, we obtain

$$\begin{aligned} n^{1/2} \hat{m} &= n^{-1/2} Z' \tilde{u}(\theta_0) \\ &= n^{-1/2} Z'(Z\Gamma(\theta - \theta_0) + V(\theta - \theta_0) + u) \\ &= n^{-1/2} Z' Z\Gamma(\theta - \theta_0) + n^{-1/2} Z'(V(\theta - \theta_0) + u) \\ &= n^{-1} Z' Z n^{1/2} |\Gamma|(\Gamma/|\Gamma|)(\theta - \theta_0) + n^{-1/2} Z'(V(\theta - \theta_0) + u) \\ &\xrightarrow{d} N[h_7^* h_1^* h_3^*(\theta - \theta_0), (h_{5,m}^* + 2h_{5,2}^*(\theta - \theta_0) + h_{5,3}^*(\theta - \theta_0)^2)]. \end{aligned} \quad (\text{SA-3.51})$$

By the CMT, (SA-3.49) and (SA-3.51)

$$\text{AR} \xrightarrow{d} \text{AR}_\infty \sim \chi_1^2(\eta_1^2), \quad (\text{SA-3.52})$$

where $\eta_1^2 \equiv (h_7^* h_1^* h_3^* (\theta - \theta_0))^2 (h_{5,m}^* + 2h_{5,2}^* (\theta - \theta_0) + h_{5,3}^* (\theta - \theta_0)^2)^{-1}$. As a result, under weak identification, the AR statistic does not diverge even under fixed alternatives. Under $\theta = \theta_0 + h_\theta n^{-1/2}$, arguments as in (SA-3.49) and (SA-3.51) yield

$$\hat{\Sigma} - h_{5,m}^* \xrightarrow{p} 0, \quad (\text{SA-3.53})$$

$$n^{1/2} \hat{m} \xrightarrow{d} N[0, h_{5,m}^*]. \quad (\text{SA-3.54})$$

(SA-3.53), (SA-3.54) and the CMT together yield

$$\text{AR} \xrightarrow{d} \text{AR}_\infty \sim \chi_1^2. \quad (\text{SA-3.55})$$

So in this case, the asymptotic distribution is the same as that under the null hypothesis.

PAR statistic. Next we derive the asymptotic distribution of the permutation statistics. Note that $n^{-1} Z' Z \xrightarrow{a.s.} h_7^*$ ((SA-3.19)), $n^{-1} V' Z \xrightarrow{a.s.} 0$, $n^{-1} Z' u \xrightarrow{a.s.} 0$, $n^{-1} V' M_l V \xrightarrow{a.s.} \Sigma^V$, $n^{-1} u' M_l V \xrightarrow{a.s.} \Sigma^{uV}$ and $n^{-1} u' M_l u \xrightarrow{a.s.} \sigma^2$ ((SA-3.20)), by the SLLN, CMT, (SA-3.11) and (SA-3.12).

Let $\theta \neq \theta_0$ be fixed. Then, using the convergences above and (SA-3.47)

$$\begin{aligned} & n^{-1} \tilde{u}(\theta_0)' \tilde{u}(\theta_0) - \mathbb{E}_P[u_i(u_i + V_i(\theta - \theta_0))] \\ &= n^{-1} (Z\Gamma(\theta - \theta_0) + M_l V(\theta - \theta_0) + M_l u)' (Z\Gamma(\theta - \theta_0) + M_l V(\theta - \theta_0) + M_l u) \\ &\quad - \mathbb{E}_P[u_i(u_i + V_i(\theta - \theta_0))] \\ &= (\theta - \theta_0)^2 \Gamma^2 n^{-1} Z' Z + (\theta - \theta_0)^2 n^{-1} V' M_l V + u' M_l u + 2(\theta - \theta_0)^2 \Gamma n^{-1} Z' M_l V(\theta - \theta_0) \\ &\quad + 2(\theta - \theta_0)^2 \Gamma n^{-1} Z' M_l u + (\theta - \theta_0) n^{-1} V' M_l u - \mathbb{E}_P[u_i(u_i + V_i(\theta - \theta_0))] \\ &\xrightarrow{a.s.} 0. \end{aligned} \quad (\text{SA-3.56})$$

Hence,

$$n^{-1} \tilde{u}(\theta_0)' \tilde{u}(\theta_0) \xrightarrow{a.s.} (\sigma^2 + \Sigma^{Vu}(\theta - \theta_0)). \quad (\text{SA-3.57})$$

Under $\theta = \theta_0 + h n^{-1/2}$,

$$n^{-1} \tilde{u}(\theta_0)' \tilde{u}(\theta_0) \xrightarrow{a.s.} \sigma^2. \quad (\text{SA-3.58})$$

By Jensen's inequality and (SA-3.45),

$$\begin{aligned}
& n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 \\
&= n^{-1} \sum_{i=1}^n (Z_i \Gamma(\theta - \theta_0) + (V_i - \bar{V})(\theta - \theta_0) + u_i - \bar{u})^4 \\
&\leq 27n^{-1} \sum_{i=1}^n Z_i^4 \Gamma^4(\theta - \theta_0)^4 + 27n^{-1} \sum_{i=1}^n (V_i - \bar{V})^4 (\theta - \theta_0)^4 + 27n^{-1} \sum_{i=1}^n (u_i - \bar{u})^4 \\
&= O_p(1),
\end{aligned} \tag{SA-3.59}$$

where the last equality holds by (SA-3.21), (SA-3.22), and the fact that $n^{-1} \sum_{i=1}^n (V_i - \bar{V})^4 = O_p(1)$ which follows from arguments similar to (SA-3.22) and the WLLN. Therefore, as in (SA-3.23)

$$\begin{aligned}
\text{Var}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W] &= \text{Var}_{P^\pi} \left[n^{-1} \sum_{i=1}^n Z_i^2 \tilde{u}_{\pi(i)}(\theta_0)^2 | y, Y, W \right] \\
&\leq \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n Z_i^4 \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 \right) \\
&\xrightarrow{P} 0.
\end{aligned} \tag{SA-3.60}$$

By Chebyshev's inequality, for any $\epsilon > 0$

$$P^\pi[|\hat{\Sigma}^\pi - \mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W]| > \epsilon] \leq \text{Var}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W] / \epsilon^2 \xrightarrow{P} 0. \tag{SA-3.61}$$

When θ is fixed, using (SA-3.19) (SA-3.57), $\mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W] \xrightarrow{a.s.} (\sigma^2 + \Sigma^{Vu}(\theta - \theta_0))h_7^*$, hence by the triangle inequality and (SA-3.61)

$$\hat{\Sigma}^\pi \xrightarrow{P^\pi} (\sigma^2 + \Sigma^{Vu}(\theta - \theta_0))h_7^* \quad \text{in } P\text{-probability.} \tag{SA-3.62}$$

When $\theta = \theta_0 + h_\theta n^{-1/2}$, $\mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W] \xrightarrow{a.s.} \sigma^2 h_7^*$, hence

$$\hat{\Sigma}^\pi \xrightarrow{P^\pi} \sigma^2 h_7^* \quad \text{in } P\text{-probability.} \tag{SA-3.63}$$

Next we verify the conditions for the combinatorial CLT

$$(n^{-1} \tilde{u}(\theta_0)' \tilde{u}(\theta_0))^{-1/2} (n^{-1} Z' Z)^{-1/2} n^{-1/2} Z' \tilde{u}(\theta_0) \xrightarrow{d^\pi} N[0, 1] \quad P\text{-almost surely.} \tag{SA-3.64}$$

The Lyapunov-type sufficient conditions for (SA-3.64) (see (SA-5.68) and (SA-5.69)) are: for some

$\delta > 0$

$$n^{-1} \sum_{i=1}^n |\tilde{u}_i(\theta_0)|^{2+\delta/2} = O_{a.s.}(1), \quad (\text{SA-3.65})$$

$$n^{-1} \sum_{i=1}^n |W_i|^{2+\delta/2} = O_{a.s.}(1). \quad (\text{SA-3.66})$$

(SA-3.66) is already verified in (SA-3.28). (SA-3.65) holds because using the convexity of $x \mapsto |x|^{2+\delta/2}$

$$\begin{aligned} & n^{-1} \sum_{i=1}^n |\tilde{u}_i(\theta_0)|^{2+\delta/2} \\ &= n^{-1} \sum_{i=1}^n |Z_i \Gamma(\theta - \theta_0) + (V_i - \bar{V})(\theta - \theta_0) + u_i - \bar{u}|^{2+\delta/2} \\ &\leq 3^{1+\delta/2} n^{-1} \left(\sum_{i=1}^n |Z_i|^{2+\delta/2} |\Gamma(\theta - \theta_0)|^{2+\delta/2} + \sum_{i=1}^n |V_i - \bar{V}|^{2+\delta/2} |\theta - \theta_0|^{2+\delta/2} + \sum_{i=1}^n |u_i - \bar{u}|^{2+\delta/2} \right) \\ &= O_{a.s.}(1), \end{aligned} \quad (\text{SA-3.67})$$

where the last equality holds because by the SLLN

$$n^{-1} \sum_{i=1}^n |Z_i|^{2+\delta/2} \leq 2^{1+\delta/2} n^{-1} \sum_{i=1}^n |W_i|^{2+\delta/2} + 2^{1+\delta/2} |\bar{W}|^{2+\delta/2} = O_{a.s.}(1),$$

$n^{-1} \sum_{i=1}^n |V_i - \bar{V}|^{2+\delta/2} = O_{a.s.}(1)$ and $n^{-1} \sum_{i=1}^n |u_i - \bar{u}|^{2+\delta/2} = O_{a.s.}(1)$ as in (SA-3.27). If θ is fixed, (SA-3.44), (SA-3.57) and (SA-3.64) give

$$n^{1/2} \hat{m}^\pi = n^{-1/2} Z' \tilde{u}_\pi(\theta_0) \xrightarrow{d^\pi} N[0, (\sigma^2 + \Sigma^{Vu}(\theta - \theta_0)) h_7^*] \text{ in } P\text{-probability.} \quad (\text{SA-3.68})$$

If $\theta = \theta_0 + h_\theta n^{-1/2}$, (SA-3.44), (SA-3.58) and (SA-3.64) yield

$$n^{1/2} \hat{m}^\pi = n^{-1/2} Z' \tilde{u}_\pi(\theta_0) \xrightarrow{d^\pi} N[0, h_7^* \sigma^2] \text{ in } P\text{-probability.} \quad (\text{SA-3.69})$$

Finally, by (SA-3.62), (SA-3.63), (SA-3.68), (SA-3.69) and the CMT, we obtain under both θ fixed and $\theta = \theta_0 + h_\theta n^{-1/2}$ that

$$\text{PAR}_2 \xrightarrow{d^\pi} \chi_1^2 \text{ in } P\text{-probability.} \quad (\text{SA-3.70})$$

Test power. Using Lemma 11.2.1 of Lehmann and Romano (2005), it follows from (SA-3.70) that $\text{PAR}_{2(r)} \xrightarrow{p} r_{1-\alpha}(\chi_1^2)$. From (SA-3.52), (SA-3.55) and Slutsky's lemma, $\text{AR} - \text{PAR}_{2(r)} \xrightarrow{d} 0$

$\text{AR}_\infty - r_{1-\alpha}(\chi_1^2)$. Then, by continuity of the cdf of the (non)-central χ_1^2 distribution at all positive points,

$$\lim_{n \rightarrow \infty} P[\text{AR} > \text{PAR}_{2(r)}] = \lim_{n \rightarrow \infty} P[\text{AR} \geq \text{PAR}_{2(r)}] = P[\text{AR}_\infty > r_{1-\alpha}(\chi_1^2)].$$

By the sandwich theorem,

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PAR}_2}] = P[\text{AR}_\infty > r_{1-\alpha}(\chi_1^2)].$$

If θ is fixed, from (SA-3.52)

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PAR}_2}] = P[\chi_1^2(\eta_1^2) > r_{1-\alpha}(\chi_1^2)],$$

where $\eta_1^2 = (h_7^* h_1^* h_3^* (\theta - \theta_0))^2 (h_{5,m}^* + 2h_{5,2}^* (\theta - \theta_0) + h_{5,3}^* (\theta - \theta_0)^2)^{-1}$.

If $\theta = \theta_0 + h_\theta n^{-1/2}$, from (SA-3.55)

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PAR}_2}] = P[\chi_1^2 > r_{1-\alpha}(\chi_1^2)] = \alpha.$$

Thus, in both cases, the test is not consistent.

Case $h_1^* = \infty$. In this case, using (SA-3.8) and (SA-3.9)

$$|\Gamma| > 0 \quad \text{for all } n \text{ large, and } n^{1/2}\Gamma = n^{1/2}|\Gamma|(\Gamma/|\Gamma|) \rightarrow h_3^* h_1^* = \infty. \quad (\text{SA-3.71})$$

AR statistic. If $\theta \neq \theta_0$ is fixed, using (SA-3.44), (SA-3.71) and the fact that $n^{-1/2}Z'(V(\theta - \theta_0) + u) = O_p(1)$ in (SA-3.50), we obtain

$$\begin{aligned} n^{1/2}\hat{m} &= n^{-1/2}Z'\tilde{u}(\theta_0) \\ &= n^{-1/2}Z'Z\Gamma(\theta - \theta_0) + n^{-1/2}Z'(V(\theta - \theta_0) + u) \\ &= n^{-1}Z'Zn^{1/2}|\Gamma|(\Gamma/|\Gamma|)(\theta - \theta_0) + n^{-1/2}Z'(V(\theta - \theta_0) + u) \\ &\xrightarrow{p} \infty. \end{aligned} \quad (\text{SA-3.72})$$

Using the inequality $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$, we have

$$\begin{aligned} \hat{\Sigma} &\leq n^{-1} \sum_{i=1}^n Z_i^2 (Z_i \Gamma(\theta - \theta_0) + (V_i - \bar{V})(\theta - \theta_0) + u_i - \bar{u})^2 \\ &\leq 3n^{-1} \sum_{i=1}^n Z_i^2 [Z_i^2 \Gamma^2(\theta - \theta_0)^2 + (V_i - \bar{V})^2(\theta - \theta_0)^2 + (u_i - \bar{u})^2] \\ &= 3n^{-1} \sum_{i=1}^n [Z_i^4 \Gamma^2(\theta - \theta_0)^2 + Z_i^2 (V_i - \bar{V})^2(\theta - \theta_0)^2 + Z_i^2 (u_i - \bar{u})^2]. \end{aligned}$$

Then,

$$\begin{aligned} \text{AR} &= n \hat{m}^2 \hat{\Sigma}^{-1} \\ &\geq n \hat{m}^2 \left(3n^{-1} \sum_{i=1}^n [Z_i^4 \Gamma^2(\theta - \theta_0)^2 + Z_i^2(V_i - \bar{V})^2(\theta - \theta_0)^2 + Z_i^2(u_i - \bar{u})^2] \right)^{-1}. \end{aligned} \quad (\text{SA-3.73})$$

From (SA-3.37)-(SA-3.44), $n^{-1} \sum_{i=1}^n Z_i^4 \Gamma^2(\theta - \theta_0)^2 = O_p(1)$, $n^{-1} \sum_{i=1}^n Z_i^2(V_i - \bar{V})^2(\theta - \theta_0)^2 = O_p(1)$ and $n^{-1} \sum_{i=1}^n Z_i^2(u_i - \bar{u})^2 = n^{-1} \sum_{i=1}^n (Z_i^* + D)^2(u_i - \bar{u})^2 \xrightarrow{p} h_{5,m}^* > 0$. Using the latter three and (SA-3.72) in (SA-3.73), we then obtain

$$\text{AR} \xrightarrow{p} \infty. \quad (\text{SA-3.74})$$

If $\theta = \theta_0 + h_\theta n^{-1/2}$, using (SA-3.9), (SA-3.37)-(SA-3.44), (SA-3.71) and some calculations

$$\begin{aligned} |\Gamma|^{-2} \hat{\Sigma} &= n^{-1} \sum_{i=1}^n Z_i^2 \left(Z_i(\Gamma|\Gamma|^{-1})h_\theta n^{-1/2} + (V_i - \bar{V})h_\theta(|\Gamma|^{-1}n^{-1/2}) + |\Gamma|^{-1}(u_i - \bar{u}) \right)^2 \\ &= |\Gamma|^{-2} n^{-1} \sum_{i=1}^n Z_i^{*2} u_i^2 + o_p(1) \\ &= |\Gamma|^{-2} h_{5,m}^* + o_p(1). \end{aligned} \quad (\text{SA-3.75})$$

Moreover, by (SA-3.9), (SA-3.11), (SA-3.44), the fact that $n^{-1} Z'V \xrightarrow{p} 0$ and some calculations

$$\begin{aligned} n^{1/2} |\Gamma|^{-1} \hat{m} &= n^{-1/2} |\Gamma|^{-1} Z' \tilde{u}(\theta_0) \\ &= n^{-1/2} |\Gamma|^{-1} Z' (Z\Gamma(\theta - \theta_0) + V(\theta - \theta_0) + u) \\ &= n^{-1} Z' Z(\Gamma/|\Gamma|)h_\theta + |\Gamma|^{-1} n^{-1/2} Z' (Vh_\theta n^{-1/2} + u) \\ &= h_7^* h_3^* h_\theta + |\Gamma|^{-1} n^{-1/2} Z' u + o_p(1). \end{aligned} \quad (\text{SA-3.76})$$

Thus, (SA-3.75), (SA-3.76), the fact that $n^{-1/2} Z' u \xrightarrow{d} N[0, h_{5,m}^*]$ and the CMT yield

$$\begin{aligned} \text{AR} &= (n^{1/2} |\Gamma|^{-1} \hat{m})^2 (|\Gamma|^{-2} \hat{\Sigma})^{-1} \\ &= (h_7^* h_3^* h_\theta + |\Gamma|^{-1} n^{-1/2} Z' u)^2 \Gamma^2 (h_{5,m}^*)^{-1} + o_p(1) \\ &= (|\Gamma| h_7^* h_3^* h_\theta (h_{5,m}^*)^{-1/2} + (h_{5,m}^*)^{-1/2} n^{-1/2} Z' u)^2 + o_p(1) \\ &\xrightarrow{d} \chi_1^2(\eta_2^2), \end{aligned} \quad (\text{SA-3.77})$$

where $\eta_2^2 \equiv \lim_{n \rightarrow \infty} \Gamma^2 h_7^{*2} h_3^{*2} h_\theta^2 (h_{5,m}^*)^{-1}$ is assumed to exist.

PAR statistic. The asymptotic distribution of the PAR_2 statistic is still χ_1^2 by arguments analogous to Case 1 after noting that (i) $n^{-1}\tilde{u}(\theta_0)'\tilde{u}(\theta_0) - \text{E}_P[(Z_i^*\Gamma + V_i)(\theta - \theta_0) + u_i]^2] \xrightarrow{p} 0$ if θ is fixed, and $n^{-1}\tilde{u}(\theta_0)'\tilde{u}(\theta_0) \xrightarrow{p} \sigma^2$ if $\theta = \theta_0 + n^{-1/2}h_\theta$; (ii) $n^{-1}\sum_{i=1}^n \tilde{u}_i(\theta_0)^4 = O_p(1)$, which is required for the consistency of $\hat{\Sigma}^\pi$, and $n^{-1}\sum_{i=1}^n \tilde{u}_i(\theta_0)^{2+\delta/2} = O_{a.s.}(1)$, which is required for the combinatorial CLT, in both $\theta = \theta_0 + n^{-1/2}h_\theta$ and fixed θ cases. Thus, the details are omitted.

Test power. Since, as shown in Case 1,

$$\lim_{n \rightarrow \infty} \text{E}_P[\phi_n^{\text{PAR}_2}] = P[\text{AR}_\infty > r_{1-\alpha}(\chi_1^2)],$$

if $\theta \neq \theta_0$ is fixed, from (SA-3.74)

$$\lim_{n \rightarrow \infty} \text{E}_P[\phi_n^{\text{PAR}_2}] = P[\infty > r_{1-\alpha}(\chi_1^2)] = 1,$$

so the test is consistent. If $\theta = \theta_0 + h_\theta n^{-1/2}$, from (SA-3.77)

$$\lim_{n \rightarrow \infty} \text{E}_P[\phi_n^{\text{PAR}_2}] = P[\chi_1^2(\eta_2^2) > r_{1-\alpha}(\chi_1^2)],$$

with $\eta_2^2 = \lim_{n \rightarrow \infty} \Gamma^2 h_7^{*2} h_3^{*2} h_\theta^2 (h_{5,m}^*)^{-1}$.

SA-4 Proof of Proposition 2.1 (Finite sample validity)

Since $\tilde{\Sigma}_u(\theta_0) = \text{diag}(\tilde{u}_1(\theta_0)^2, \dots, \tilde{u}_n(\theta_0)^2)$ and $\tilde{u}_i(\theta_0) = u_i - X_i'(X'X)^{-1}X'u$ under $H_0 : \theta = \theta_0$, we may write $\tilde{\Sigma}_u(\theta_0) = \tilde{\Sigma}_u(u, X)$. Given $\mathbb{G}'_n = \{\pi_1, \dots, \pi_N\}$ with π_1 being the identity permutation, we have $\text{AR}^{\pi_j} = \text{AR}(W_{\pi_j}, X, M_X u)$, $1 \leq j \leq N$, with $\text{AR}_{\pi_1} = \text{AR}$. For any $\pi \in \mathbb{G}'_n$,

$$\begin{aligned} \text{PAR}_1 &= \text{AR}(W_\pi, X, M_X u) \\ &= u' M_X W_\pi (W_\pi' M_X \tilde{\Sigma}_u(u, X) M_X W_\pi)^{-1} W_\pi' M_X u \\ &\stackrel{d}{=} u' M_X W (W' M_X \tilde{\Sigma}_u(u, X) M_X W)^{-1} W' M_X u \\ &= \text{AR}, \end{aligned} \tag{SA-4.1}$$

where the equality in distribution follows from Assumption 1 (in the main paper). For the order statistics $\text{PAR}_{1(1)} \leq \dots \leq \text{PAR}_{1(N)}$ of $\text{AR}^{\pi_1}, \dots, \text{AR}^{\pi_N}$ and $\pi \in \mathbb{G}'_n$, define $\tilde{\phi}_n(W_\pi, X, u)$ to be equal to 1 if $\text{AR}(W_\pi, X, M_X u) > \text{PAR}_{(r)}$, $(N\alpha - N^+)/N^0$ if $\text{AR}(W_\pi, X, M_X u) = \text{PAR}_{(r)}$, and 0

otherwise, where N^+ and N^0 are given in the text. Clearly, $\phi_n^{\text{PAR}_1} = \tilde{\phi}_n(W_{\pi_1}, X, u)$. It follows from (SA-4.1) that for all $1 \leq j \leq N$,

$$\phi_n^{\text{PAR}_1} \stackrel{d}{=} \tilde{\phi}_n(W_{\pi_j}, X, u). \quad (\text{SA-4.2})$$

Thus,

$$\begin{aligned} \mathbb{E}[\phi_n^{\text{PAR}_1}] &= \frac{1}{N} \mathbb{E} \left[\sum_{j=1}^N \tilde{\phi}_n(W_{\pi_j}, X, u) \right] \\ &= \frac{1}{N} \mathbb{E} \left[N^+ + \frac{N\alpha - N^+}{N^0} N^0 \right] \\ &= \alpha, \end{aligned}$$

where the first equality uses (SA-4.2), and the second equality follows from the definitions of $\tilde{\phi}_n(W_{\pi_j}, X, u)$, N^+ and N^0 . The proof of the PAR_2 test is similar, thus is omitted.

SA-5 Proof of Theorem 2.2 (Asymptotic similarity)

SA-5.1 Drifting sequences of parameters

Following Andrews and Guggenberger (2019b), we first introduce some notations, and define the DGPs. Henceforth, we shall suppress the argument θ_0 in some quantities, and write, for example, $\hat{m}(\theta_0) = \hat{m}$, $\hat{\Sigma}(\theta_0) = \hat{\Sigma}$, $\hat{J}(\theta_0) = \hat{J}$, $\hat{m}^\pi(\theta_0) = \hat{m}^\pi$, $\hat{\Sigma}^\pi(\theta_0) = \hat{\Sigma}^\pi$, $\hat{J}^\pi(\theta_0) = \hat{J}^\pi$.

Define $G_P \equiv \mathbb{E}_P[Z_i^* Z_i^{*'}] \Gamma$, $\hat{H} \equiv \hat{\Sigma}^{-1/2}$, $H_P \equiv \Sigma_P^{-1/2}$, $\hat{U} \equiv \left((\theta_0, I_d) \Omega^\epsilon(\hat{\Sigma}, \tilde{\mathcal{V}})^{-1}(\theta_0, I_d)' \right)^{1/2}$ and $U_P \equiv \left((\theta_0, I_d) \Omega^\epsilon(\Sigma_P, \mathcal{V}_P)^{-1}(\theta_0, I_d)' \right)^{1/2}$, where $\tilde{\mathcal{V}} = \hat{\mathcal{V}}_a(\theta_0)$, $\mathcal{V}_P = \mathcal{V}_{P,a}$ (defined in (2.17)-(2.18) and (2.35) of the main paper) and $\tilde{\mathcal{V}} = \hat{\mathcal{V}}_b(\theta_0)$, $\mathcal{V}_P = \mathcal{V}_{P,b}$ (defined in (2.19)-(2.20) and (2.36) of the main paper) for the CLR_a , PCLR_a and CLR_b , PCLR_b statistics, respectively.

The SVD of $H_P G_P U_P$ is given as follows. Let $\kappa_{1P} \geq \dots \geq \kappa_{dP}$ and B_P denote the ordered eigenvalues, and a $d \times d$ orthogonal matrix of corresponding eigenvectors, respectively, of

$$U_P' G_P' H_P H_P' G_P U_P, \quad (\text{SA-5.1})$$

and $\kappa_{1P} \geq \dots \geq \kappa_{kP}$ and A_P denote the ordered eigenvalues, and a $k \times k$ orthogonal matrix of corresponding eigenvectors, respectively, of

$$H_P G_P U_P U_P' G_P' H_P'. \quad (\text{SA-5.2})$$

Let $\tau_{1P} \geq \dots \geq \tau_{dP} \geq 0$ denote the singular values of

$$H_P G_P U_P. \quad (\text{SA-5.3})$$

Define

$$\lambda_{1,P} \equiv (\tau_{1P}, \dots, \tau_{dP})' \in \mathbb{R}^d, \quad (\text{SA-5.4})$$

$$\lambda_{2,P} \equiv B_P \in \mathbb{R}^{d \times d}, \quad (\text{SA-5.5})$$

$$\lambda_{3,P} \equiv A_P \in \mathbb{R}^{k \times k}, \quad (\text{SA-5.6})$$

$$\lambda_{4,P} \equiv G_P = E_P[Z_i^* Z_i^{*'}] \Gamma \in \mathbb{R}^{k \times d}, \quad (\text{SA-5.7})$$

$$\lambda_{5,P} \equiv \mathcal{V}_{P,a} = E_P \left[\begin{pmatrix} Z_i^* u_i \\ \text{vec}(Z_i^* (Z_i^{*'} \Gamma + V_i')) \end{pmatrix} \begin{pmatrix} Z_i^* u_i \\ \text{vec}(Z_i^* (Z_i^{*'} \Gamma + V_i')) \end{pmatrix}' \right] \in \mathbb{R}^{(d+1)k \times (d+1)k}, \quad (\text{SA-5.8})$$

$$\lambda_{6,P} \equiv [\lambda_{6,1P}, \dots, \lambda_{6,(d-1)P}]' = \left[\frac{\tau_{2P}}{\tau_{1P}}, \dots, \frac{\tau_{dP}}{\tau_{(d-1)P}} \right]' \in [0, 1]^{d-1}, \quad (\text{SA-5.9})$$

$$\lambda_{7,P} \equiv E_P[(W_i', X_i')'(W_i', X_i')] \in \mathbb{R}^{(k+p) \times (k+p)}, \quad (\text{SA-5.10})$$

$$\lambda_{8,P} \equiv \Sigma_P^e = \text{Var}_P[(u_i, V_i')'] \in \mathbb{R}^{(d+1) \times (d+1)}, \quad (\text{SA-5.11})$$

$$\lambda_{9,P} \equiv (\Sigma_P, K(\mathcal{V}_P)) \in (\mathbb{R}^{k \times k}, \mathbb{R}^{(d+1)k \times (d+1)k}), \quad \mathcal{V}_P \in \{\mathcal{V}_{P,a}, \mathcal{V}_{P,b}\}, \quad (\text{SA-5.12})$$

$$\lambda_{10,P} \equiv P, \quad (\text{SA-5.13})$$

$$\lambda_P \equiv (\lambda_{1,P}, \dots, \lambda_{10,P})'. \quad (\text{SA-5.14})$$

As in [Andrews and Guggenberger \(2019b\)](#), the drifting parameters $\lambda_{1,P}, \dots, \lambda_{6,P}, \lambda_{9,P}, \lambda_{10,P}$ are used in the asymptotic results for the AR, LM, CLR statistics. The parameter vector $\lambda_{1,P}$ characterizes the identification strength. The sequences $\lambda_{7,P}$ and $\lambda_{8,P}$ are needed additionally for the asymptotic properties of the permutation statistics.

Define

$$\Lambda \equiv \{\lambda_P : P \in \mathcal{P}_0^{\text{PCLR}}\},$$

$$h_n(\lambda_P) \equiv [n^{1/2} \lambda_{1,P}, \lambda_{2,P}, \lambda_{3,P}, \lambda_{4,P}, \lambda_{5,P}, \lambda_{6,P}, \lambda_{7,P}, \lambda_{8,P}, \lambda_{9,P}], \quad (\text{SA-5.15})$$

and

$$\mathcal{H} \equiv \{h \in (\mathbb{R} \cup \{\pm\infty\})^{\dim(h_n(\lambda_{P_n}))} : h_{w_n}(\lambda_{P_{w_n}}) \rightarrow h \text{ for some subsequence } \{w_n\} \text{ of } \{n\}\} \quad (\text{SA-5.16})$$

$$\text{and some sequence } \{\lambda_{P_{w_n}} \in \Lambda : n \geq 1\}. \quad (\text{SA-5.17})$$

The following assumption corresponds to Assumption B* of [Andrews and Guggenberger \(2019b\)](#) which is key for establishing the asymptotic similarity of the tests.

Assumption 2. For any subsequence $\{w_n\}$ of $\{n\}$ and any sequence $\{\lambda_{P_{w_n}} \in \Lambda : n \geq 1\}$ for which $h_{w_n}(\lambda_{P_{w_n}}) \rightarrow h \in \mathcal{H}$, $E_{P_{w_n}}[\phi_{w_n}^{\text{PR}}] \rightarrow \alpha$ for some $\alpha \in (0, 1)$.

Let $\{\lambda_{P_n} \in \Lambda : n \geq 1\}$ be a sequence such that $h_n(\lambda_{P_n}) \rightarrow h \in \mathcal{H}$, and

$$n^{1/2}\tau_{jP_n} \rightarrow h_{1,j} \geq 0, \quad j = 1, \dots, d, \quad (\text{SA-5.18})$$

$$\lambda_{j,P_n} \rightarrow h_j, \quad j = 2, \dots, 8, \quad (\text{SA-5.19})$$

$$\lambda_{5,mP_n} = E_{P_n}[Z_i^* Z_i^{*'} u_i^2] \rightarrow h_{5,m} = \Sigma, \quad (\text{SA-5.20})$$

$$\lambda_{6,jP_n} \rightarrow h_{6,j}, \quad j = 1, \dots, d-1, \quad (\text{SA-5.21})$$

$$\lambda_{7,P_n} \rightarrow h_7 \equiv \begin{bmatrix} Q_{ww} & Q_{wx} \\ Q_{xw} & Q_{xx} \end{bmatrix}, \quad (\text{SA-5.22})$$

$$\lambda_{8,P_n} \rightarrow h_8 \equiv \Sigma^e = \begin{bmatrix} \sigma^2 & \Sigma^{uV} \\ \Sigma^{Vu} & \Sigma^V \end{bmatrix}, \quad (\text{SA-5.23})$$

$$\lambda_{9,P_n} \rightarrow h_9 \equiv (\Sigma, K(\mathcal{V})), \quad (\text{SA-5.24})$$

where $\lambda_{5,mP_n} \in \mathbb{R}^{k \times k}$ is the upper-left sub-matrix of λ_{5,P_n} with the corresponding limit $h_{5,m} = \Sigma$, h_7 and h_8 are partitioned conformably with λ_{7,P_n} and λ_{8,P_n} , respectively, and for both $(\Sigma_P, K(\mathcal{V}_{P,a}))$ and $(\Sigma_P, K(\mathcal{V}_{P,b}))$, the limit is denoted by the same variable h_9 with \mathcal{V} in (SA-5.24) denoting the limit of $\mathcal{V}_P \in \{\mathcal{V}_{P,a}, \mathcal{V}_{P,b}\}$ defined in (SA-5.12).

Consider $q = q_h \in \{0, \dots, d\}$ such that

$$h_{1,j} = \infty \quad \text{for } 1 \leq j \leq q,$$

$$h_{1,j} < \infty \quad \text{for } q+1 \leq j \leq d.$$

The normalization matrices used to derive the limiting distribution of the test statistics are defined as follows:

$$S_{P_n} \equiv \text{diag} \left((n^{1/2}\tau_{1P_n})^{-1}, \dots, (n^{1/2}\tau_{qP_n})^{-1}, 1, \dots, 1 \right) \in \mathbb{R}^{d \times d}, \quad (\text{SA-5.25})$$

$$T_{P_n} \equiv B_{P_n} S_{P_n} \in \mathbb{R}^{d \times d}. \quad (\text{SA-5.26})$$

In the sequel, we index the quantities indexed by $P = P_n$ by n ; for example, write $\Sigma_P = \mathbb{E}_P[Z_i^* Z_i^{*'} u_i^2] = \Sigma_n$, $G_P = \mathbb{E}_P[Z_i^* Z_i^{*'}] \Gamma = G_n$, $C_{Ps} = \mathbb{E}_P[Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma + V_{is}) u_i] = C_{ns}$, $\lambda_{w_n} = \lambda_{P_{w_n}}$ and $\lambda_n = \lambda_{P_n}$.

Let $\hat{\kappa}_1 \geq \dots \geq \hat{\kappa}_d \geq 0$ denote the eigenvalues of

$$n \hat{U}' \hat{J}' \hat{H} \hat{H}' \hat{J} \hat{U}.$$

Partition

$$A_n = [A_{n,q}, A_{n,k-q}], \quad A_{n,q} \in \mathbb{R}^{k \times q}, \quad A_{n,k-q} \in \mathbb{R}^{k \times (k-q)}, \quad (\text{SA-5.27})$$

$$B_n = [B_{n,q}, B_{n,d-q}], \quad B_{n,q} \in \mathbb{R}^{d \times q}, \quad B_{n,d-q} \in \mathbb{R}^{d \times (d-q)}, \quad (\text{SA-5.28})$$

$$h_2 = [h_{2,q}, h_{2,d-q}], \quad h_{2,q} \in \mathbb{R}^{d \times q}, \quad h_{2,d-q} \in \mathbb{R}^{d \times (d-q)}, \quad (\text{SA-5.29})$$

$$h_3 = [h_{3,q}, h_{3,k-q}], \quad h_{3,q} \in \mathbb{R}^{k \times q}, \quad h_{3,k-q} \in \mathbb{R}^{k \times (k-q)}. \quad (\text{SA-5.30})$$

Define

$$\Upsilon_n \equiv \begin{bmatrix} \Upsilon_{n,q} & 0_{q \times (d-q)} \\ 0_{(d-q) \times q} & \Upsilon_{n,d-q} \\ 0_{(k-d) \times q} & 0_{(k-d) \times (d-q)} \end{bmatrix} \in \mathbb{R}^{k \times d}, \quad \Upsilon_{n,q} \equiv \text{diag}(\tau_{1n}, \dots, \tau_{qn}) \in \mathbb{R}^{q \times q},$$

$$\Upsilon_{n,d-q} \equiv \text{diag}(\tau_{(q+1)n}, \dots, \tau_{dn}) \in \mathbb{R}^{(d-q) \times (d-q)}. \quad (\text{SA-5.31})$$

SA-5.2 Probability limits of some key sample quantities

The proof of the asymptotic similarity of the permutation tests involves the derivation of the asymptotic distributions of the AR, LM, CLR_a and CLR_b statistics under the drifting sequences of parameters given above which can be derived following [Andrews and Guggenberger \(2017a,b, 2019a,b\)](#) based on the moment function $m_i^*(\theta) = Z_i^* u_i^*(\theta)$, where $u_i^*(\theta) \equiv Z_i^{*'} \Gamma(\theta - \theta_0) + u_i + V_i'(\theta - \theta_0)$, and the corresponding sample moment function $\hat{m}^*(\theta) \equiv n^{-1} \sum_{i=1}^n Z_i^* u_i^*(\theta)$. Note here that the test statistics are based on $\hat{m}(\theta) \equiv n^{-1} Z'(y - Y\theta) = n^{-1} Z' Z \Gamma(\theta - \theta_0) + n^{-1} Z' u + n^{-1} Z' V(\theta - \theta_0)$ rather than $\hat{m}^*(\theta)$. Therefore, since

$$\hat{m}^*(\theta) = n^{-1} Z^{*'} Z^* \Gamma(\theta - \theta_0) + n^{-1} Z^{*'} u + n^{-1} Z^{*'} V(\theta - \theta_0),$$

where $Z^* \equiv W - X(\mathbb{E}_P[X_i X_i'])^{-1} \mathbb{E}_P[X_i W_i']$, one needs to account for the randomness that results when Z^* is replaced by Z in order to derive the asymptotic distributions. Since the latter does not

entail any substantial change in the proof of [Andrews and Guggenberger \(2017a,b, 2019a,b\)](#), we do not provide them here to avoid overlap.

However, following [Andrews and Guggenberger \(2019b\)](#), we shall determine the probability limits of certain sample random quantities that appear in the permutation statistics but do not depend on the random permutation π , under the drifting parameter sequences.

SA-5.2.1 Probability limits of $\hat{\mathcal{V}}_a(\theta_0)$, $\hat{\mathcal{V}}_b(\theta_0)$ and \hat{J} .

By the triangular array WLLN and the CMT, for $s = 1, \dots, d$, $n^{-1} \sum_{i=1}^n Z_i \tilde{Y}_{is} = n^{-1} Z' \tilde{Y}_{\cdot s} = n^{-1} Z' Z \Gamma_s + n^{-1} Z' V_{\cdot s} = E_P[Z_i^* Z_i^{*'}] \Gamma_s + o_p(1)$,

$$D \equiv E_P[W_i X_i'] (E_P[X_i X_i'])^{-1} - W' X (X' X)^{-1} \xrightarrow{p} 0, \quad (\text{SA-5.32})$$

$$(X' X)^{-1} X' V_{\cdot s} \xrightarrow{p} 0, \quad (\text{SA-5.33})$$

$$n^{-1} X' u \xrightarrow{p} 0. \quad (\text{SA-5.34})$$

We may rewrite

$$Z_i = W_i - W' X (X' X)^{-1} X_i = W_i - E_P[W_i X_i'] (E_P[X_i X_i'])^{-1} X_i + D X_i = Z_i^* + D X_i,$$

$$\tilde{Y}_{is} = Z_i' \Gamma_s + V_{is} - X_i' (X' X)^{-1} X' V_{\cdot s} = Z_i^{*'} \Gamma_s + V_{is} + X_i' (D' \Gamma_s - (X' X)^{-1} X' V_{\cdot s}).$$

Note that $\|Z_i^*\|^4, \|Z_i^*\|^3 \|X_i\|, \|Z_i^*\|^3 \|V_i\|, \|Z_i^*\|^2 \|X_i\| \|V_i\|, \|Z_i^*\|^2 \|X_i\|^2, \|Z_i^*\|^2 \|V_i\|^2, \|Z_i^*\| \|X_i\|^3, \|Z_i^*\| \|X_i\|^2 \|V_i\|, \|Z_i^*\| \|X_i\| \|V_i\|^2, \|X_i\|^3 \|V_i\|, \|X_i\|^2 \|V_i\|^2$, have finite $1+\delta/4$ moments by the Cauchy-Schwarz inequality. By [\(SA-5.32\)](#), [\(SA-5.33\)](#), [\(SA-5.34\)](#), the triangular array WLLN and the CMT, for $s, t = 1, \dots, d$

$$\begin{aligned} n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{Y}_{is} \tilde{Y}_{it} &= n^{-1} \sum_{i=1}^n (Z_i^* + D X_i) (Z_i^* + D X_i)' (Z_i^{*'} \Gamma_s + V_{is} + X_i' (D' \Gamma_s - (X' X)^{-1} X' V_{\cdot s})) \\ &\quad (Z_i^{*'} \Gamma_t + V_{it} + X_i' (D' \Gamma_t - (X' X)^{-1} X' V_{\cdot t})) \\ &= n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma_s + V_{is}) (Z_i^{*'} \Gamma_t + V_{it}) + o_p(1) \\ &= E_P[Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma_s + V_{is}) (Z_i^{*'} \Gamma_t + V_{it})] + o_p(1). \end{aligned} \quad (\text{SA-5.35})$$

Similarly,

$$\begin{aligned}
\hat{C}_s &= n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{Y}_{is} \tilde{u}_i(\theta_0) \\
&= n^{-1} \sum_{i=1}^n (Z_i^* + DX_i)(Z_i^* + DX_i)' (Z_i^{*'} \Gamma_s + V_{is} + X_i'(D' \Gamma_s - (X'X)^{-1} X' V_{,s})) \\
&\quad (u_i - X_i'(X'X)^{-1} X' u) \\
&= n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma_s + V_{is}) u_i + o_p(1) = E_P[Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma_s + V_{is}) u_i] + o_p(1) = C_{ns} + o_p(1),
\end{aligned} \tag{SA-5.36}$$

and

$$\begin{aligned}
\hat{\Sigma} &= n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{u}_i(\theta_0)^2 = n^{-1} \sum_{i=1}^n (Z_i^* + DX_i)(Z_i^* + DX_i)' (u_i - X_i'(X'X)^{-1} X' u)^2 \\
&= E_P[Z_i^* Z_i^{*'} u_i^2] + o_p(1) = \Sigma_n + o_p(1) = h_{5,m} + o_p(1).
\end{aligned} \tag{SA-5.37}$$

By (SA-5.35), (SA-5.36), (SA-5.37) and the CMT (see also the equations (27.73) and (27.74) of Andrews and Guggenberger (2019b))

$$\begin{aligned}
\hat{\mathcal{V}}_a(\theta_0) &= \mathcal{V}_{n,a} + o_p(1) = h_5 + o_p(1), \\
K(\hat{\mathcal{V}}_a(\theta_0)) &= K(\mathcal{V}_{n,a}) + o_p(1) = K(h_5) + o_p(1), \\
(\hat{\Sigma}, K(\hat{\mathcal{V}}_a(\theta_0))) &= (h_{5,m}, K(h_5)) + o_p(1), \\
\Omega^\epsilon(\hat{\Sigma}, K(\hat{\mathcal{V}}_a(\theta_0))) &= \Omega^\epsilon(h_{5,m}, K(h_5)) + o_p(1) = \Omega^\epsilon(h_9) + o_p(1),
\end{aligned} \tag{SA-5.38}$$

where the last equality holds by the continuity of the eigenvalue-adjusted matrix, see Andrews and Guggenberger (2019a), p.1717. To determine the limit of $\hat{\mathcal{V}}_b(\theta_0)$, note that $\varepsilon_i(\theta_0) = [\tilde{u}_i(\theta_0), -Y_i' + X_i'(X'X)^{-1} X' Y]'$ and

$$\varepsilon_i(\theta_0) - \begin{bmatrix} 0_{1 \times k} \\ -\Gamma' \end{bmatrix} Z_i = \begin{bmatrix} u_i - u' X (X'X)^{-1} X_i \\ -V_i + V' X (X'X)^{-1} X_i \end{bmatrix} = v_i - U' X (X'X)^{-1} X_i, \tag{SA-5.39}$$

where $v_i \equiv [u_i, -V_i']'$ and $U \equiv [u, -V]$. Letting $\tilde{Q} \equiv (Z'Z)^{-1} Z'U$, we have

$$\hat{\varepsilon}_{in}(\theta_0) - [0_{k \times 1}, -\Gamma']' Z_i = \varepsilon(\theta_0)' Z (Z'Z)^{-1} Z_i - [0_{k \times 1}, -\Gamma']' Z_i = \tilde{Q}' Z_i. \tag{SA-5.40}$$

Note here that $\tilde{Q} = O_p(n^{-1/2})$ because $n^{-1} Z'Z = n^{-1} W'W - n^{-1} W'X (X'X)^{-1} X'W \xrightarrow{a.s.} Q_{zz}$

which follows from $E_P[Z_i^* Z_i^{*'}] \rightarrow Q_{ww} - Q_{wx} Q_{xx}^{-1} Q_{xw} \equiv Q_{zz}$, the WLLN and the CMT, and

$$n^{-1/2} Z'U = n^{-1/2} W'U - n^{-1} W'X (n^{-1} X'X)^{-1} n^{-1/2} X'U = O_p(1)$$

which holds due to the triangular array CLT, the WLLN and the CMT. Rewrite

$$\begin{aligned}\hat{\mathcal{V}}_b(\theta_0) &= n^{-1} \sum_{i=1}^n \left\{ (\varepsilon_i(\theta_0) - \hat{\varepsilon}_{in}(\theta_0))(\varepsilon_i(\theta_0) - \hat{\varepsilon}_{in}(\theta_0))' \right\} \otimes (Z_i Z_i') \\ &= n^{-1} \sum_{i=1}^n \left\{ (\varepsilon_i(\theta_0) - [0_{k \times 1}, -\Gamma]' Z_i)(\varepsilon_i(\theta_0) - [0_{k \times 1}, -\Gamma]' Z_i)' \right\} \otimes (Z_i Z_i')\end{aligned}\quad (\text{SA-5.41})$$

$$+ n^{-1} \sum_{i=1}^n \left\{ (\varepsilon_i(\theta_0) - [0_{k \times 1}, -\Gamma]' Z_i)([0_{k \times 1}, -\Gamma]' Z_i - \hat{\varepsilon}_{in}(\theta_0))' \right\} \otimes (Z_i Z_i') \quad (\text{SA-5.42})$$

$$+ n^{-1} \sum_{i=1}^n \left\{ ([0_{k \times 1}, -\Gamma]' Z_i - \hat{\varepsilon}_{in}(\theta_0))(\varepsilon_i(\theta_0) - [0_{k \times 1}, -\Gamma]' Z_i)' \right\} \otimes (Z_i Z_i') \quad (\text{SA-5.43})$$

$$+ n^{-1} \sum_{i=1}^n \left\{ ([0_{k \times 1}, -\Gamma]' Z_i - \hat{\varepsilon}_{in}(\theta_0))([0_{k \times 1}, -\Gamma]' Z_i - \hat{\varepsilon}_{in}(\theta_0))' \right\} \otimes (Z_i Z_i'). \quad (\text{SA-5.44})$$

Using (SA-5.39), the term (SA-5.41) can be rewritten as

$$\begin{aligned}& n^{-1} \sum_{i=1}^n \left\{ (\varepsilon_i(\theta_0) - [0_{k \times 1}, -\Gamma]' Z_i)(\varepsilon_i(\theta_0) - [0_{k \times 1}, -\Gamma]' Z_i)' \right\} \otimes (Z_i Z_i') \\ &= n^{-1} \sum_{i=1}^n (v_i - U' X (X' X)^{-1} X_i)(v_i - U' X (X' X)^{-1} X_i)' \otimes (Z_i Z_i') \\ &= n^{-1} \sum_{i=1}^n \left(v_i v_i' - U' X (X' X)^{-1} X_i v_i' - v_i X_i' (X' X)^{-1} X' U \right. \\ &\quad \left. + U' X (X' X)^{-1} X_i X_i' (X' X)^{-1} X' U \right) \otimes (Z_i Z_i').\end{aligned}$$

Let v_{ij} and v_{il} denote the j and l -th elements of v_i , $j, l = 1, \dots, d+1$. By the Cauchy-Schwarz inequality $\mathbb{E}_P[\|v_{ij} v_{il} Z_i^* Z_i^{*'}\|^{1+\delta/4}] \leq (\mathbb{E}_P[\|v_i\|^{4+\delta}])^{1/2} (\mathbb{E}_P[\|Z_i^*\|^{4+\delta}])^{1/2} < \infty$. Similarly,

$$\mathbb{E}_P[\|v_{ij} v_{il} X_i X_i'\|^{1+\delta/4}] < \infty, \quad \mathbb{E}_P[\|v_{ij} v_{il} Z_i^* X_i'\|^{1+\delta/4}] < \infty.$$

Then, the WLLN and (SA-5.32) yield

$$\begin{aligned}& n^{-1} \sum_{i=1}^n v_i v_i' \otimes (Z_i Z_i') - \mathbb{E}_P[v_i v_i' \otimes (Z_i^* Z_i^{*'})] \\ &= n^{-1} \sum_{i=1}^n v_i v_i' \otimes ((Z_i^* + D X_i)(Z_i^* + D X_i)') - \mathbb{E}_P[v_i v_i' \otimes (Z_i^* Z_i^{*'})] \xrightarrow{p} 0.\end{aligned}\quad (\text{SA-5.45})$$

Using the Cauchy-Schwarz inequality again,

$$\mathbb{E}_P[\|X_{ij} v_{il} Z_i^* Z_i^{*'}\|^{1+\delta/4}] \leq (\mathbb{E}_P[\|Z_i^*\|^{4+\delta}])^{1/2} (\mathbb{E}_P[|X_{ij}|^{4+\delta}])^{1/4} (\mathbb{E}_P[\|v_i\|^{4+\delta}])^{1/4} < \infty,$$

for $j = 1, \dots, p$ and $l = 1, \dots, d+1$. Similarly,

$$\mathbb{E}_P[\|X_{ij} v_{il} X_i Z_i^{*'}\|^{1+\delta/4}] < \infty, \quad \mathbb{E}_P[\|X_{ij} v_{il} X_i X_i'\|^{1+\delta/4}] < \infty.$$

Therefore, by the WLLN and (SA-5.32)

$$\begin{aligned} & n^{-1} \sum_{i=1}^n X_{ij} v_{il} Z_i Z_i' - \mathbb{E}_P[X_{ij} v_{il} Z_i^* Z_i^{*'}] \\ &= n^{-1} \sum_{i=1}^n X_{ij} v_{il} (Z_i^* + DX_i)(Z_i^* + DX_i)' - \mathbb{E}_P[X_{ij} v_{il} Z_i^* Z_i^{*'}] \xrightarrow{p} 0, \end{aligned}$$

and

$$n^{-1} \sum_{i=1}^n v_i X_i' (X'X)^{-1} X'U \xrightarrow{p} 0. \quad (\text{SA-5.46})$$

By the Cauchy-Schwarz inequality $\mathbb{E}_P[\|X_{ij} X_{il} Z_i^* Z_i^{*'}\|^{1+\delta/4}] \leq \mathbb{E}_P[\|X_i\|^{4+\delta}]^{1/2} (\mathbb{E}_P[\|Z_i^*\|^{4+\delta}])^{1/2} < \infty$ and similarly

$$\mathbb{E}_P[\|X_{ij} X_{il} X_i Z_i^{*'}\|^{1+\delta/4}] < \infty \quad \text{and} \quad \mathbb{E}_P[\|X_{ij} X_{il} X_i X_i'\|^{1+\delta/4}] < \infty.$$

Then, by the WLLN and (SA-5.32)

$$\begin{aligned} & n^{-1} \sum_{i=1}^n U' X (X'X)^{-1} X_i X_i' (X'X)^{-1} X'U \otimes (Z_i Z_i') \\ &= n^{-1} \sum_{i=1}^n U' X (X'X)^{-1} X_i X_i' (X'X)^{-1} X'U \otimes ((Z_i^* + DX_i)(Z_i^* + DX_i)') \xrightarrow{p} 0. \end{aligned} \quad (\text{SA-5.47})$$

Combining (SA-5.45), (SA-5.46) and (SA-5.47), we obtain

$$n^{-1} \sum_{i=1}^n \left\{ (\varepsilon_i - [0_{k \times 1}, -\Gamma]' Z_i)(\varepsilon_i - [0_{k \times 1}, -\Gamma]' Z_i)' \right\} \otimes (Z_i Z_i') - \mathbb{E}_P[(v_i v_i') \otimes (Z_i^* Z_i^{*'})] \xrightarrow{p} 0. \quad (\text{SA-5.48})$$

We consider the term in (SA-5.42) next. Since by the Cauchy-Schwarz inequality

$$\begin{aligned} \mathbb{E}_P[\|v_{ij} Z_{il}^* Z_i^* Z_i^{*'}\|^{1+\delta/4}] &\leq (\mathbb{E}_P[\|v_{ij}\|^{4+\delta}])^{1/4} (\mathbb{E}_P[\|Z_{il}^*\|^{4+\delta}])^{1/4} (\mathbb{E}_P[\|Z_i^*\|^{4+\delta}])^{1/2} < \infty, \\ \mathbb{E}_P[\|X_{ij} Z_{il}^* Z_i^* Z_i^{*'}\|^{1+\delta/4}] &\leq (\mathbb{E}_P[\|X_{ij}\|^{4+\delta}])^{1/4} (\mathbb{E}_P[\|Z_{il}^*\|^{4+\delta}])^{1/4} (\mathbb{E}_P[\|Z_i^*\|^{4+\delta}])^{1/2} < \infty \end{aligned}$$

and the above moment bounds hold when $Z_i^* Z_i^{*'}$ is replaced by $X_i Z_i^{*'}$ and $X_i X_i'$. Using $\tilde{Q} = O_p(n^{-1/2})$, (SA-5.32), the WLLN and the CMT, the terms in (SA-5.42) and (SA-5.43) are $o_p(1)$:

$$\begin{aligned} & n^{-1} \sum_{i=1}^n \left\{ (\varepsilon_i - [0_{k \times 1}, -\Gamma]' Z_i)([0_{k \times 1}, -\Gamma]' Z_i - \hat{\varepsilon}_{in})' \right\} \otimes (Z_i Z_i') \\ &= -n^{-1} \sum_{i=1}^n (v_i (Z_i^* + DX_i)' \tilde{Q} - U' X (X'X)^{-1} X_i (Z_i^* + DX_i)' \tilde{Q}) \otimes ((Z_i^* + DX_i)(Z_i^* + DX_i)') \\ &\xrightarrow{p} 0. \end{aligned} \quad (\text{SA-5.49})$$

Finally, consider the last term in (SA-5.44). Since $E[\|Z_{ij}^* Z_{il}^* Z_i^* Z_i^{*'}\|^{1+\delta/4}] \leq E[\|Z_i^*\|^{4+\delta}] < \infty$, $E[\|Z_{ij}^* Z_{il}^* X_i Z_i^{*'}\|^{1+\delta/4}] < \infty$, and $E[\|Z_{ij}^* Z_{il}^* X_i X_i'\|^{1+\delta/4}] < \infty$, by (SA-5.32) and the WLLN

$$\begin{aligned}
& n^{-1} \sum_{i=1}^n \left\{ ([0_{k \times 1}, -\Gamma]' Z_i - \hat{\varepsilon}_{in})([0_{k \times 1}, -\Gamma]' Z_i - \hat{\varepsilon}_{in})' \right\} \otimes (Z_i Z_i') \\
&= n^{-1} \sum_{i=1}^n \tilde{Q}' Z_i Z_i' \tilde{Q} \otimes (Z_i Z_i') \\
&= n^{-1} \sum_{i=1}^n \tilde{Q}' ((Z_i^* + DX_i)(Z_i^* + DX_i)') \tilde{Q} \otimes ((Z_i^* + DX_i)(Z_i^* + DX_i)') \\
&\xrightarrow{p} 0.
\end{aligned} \tag{SA-5.50}$$

Therefore, (SA-5.48), (SA-5.49), (SA-5.50) and the CMT together with the continuity of the eigenvalue-adjusted matrix give

$$\begin{aligned}
\hat{\mathcal{V}}_b(\theta_0) &= \mathcal{V}_{n,b} + o_p(1), \\
K(\hat{\mathcal{V}}_b(\theta_0)) &= K(\mathcal{V}_{n,b}) + o_p(1), \\
(\hat{\Sigma}, K(\hat{\mathcal{V}}_b(\theta_0))) &= (h_{5,m}, K(\mathcal{V})) + o_p(1) \equiv h_9 + o_p(1), \\
\Omega^\epsilon(\hat{\Sigma}, K(\hat{\mathcal{V}}_b(\theta_0))) &= \Omega^\epsilon(h_9) + o_p(1).
\end{aligned} \tag{SA-5.51}$$

Next we determine the limit of $n^{1/2}(\hat{J} - G_n)$. Rewrite

$$\begin{aligned}
n^{1/2}(\hat{G}_n - G_n) &= n^{-1/2}(n^{-1} Z' Y - E_P[Z_i^* Z_i^{*'}] \Gamma) \\
&= n^{-1/2} Z' Z \Gamma - n^{1/2} E_P[Z_i^* Z_i^{*'}] \Gamma + n^{-1/2} Z' V \\
&= n^{-1/2} Z' V + (n^{-1/2} W' W - n^{1/2} E_P[W_i W_i']) \Gamma \\
&\quad - \left(n^{-1/2} W' X - n^{1/2} E_P[W_i X_i'] \right) (X' X)^{-1} X' W \Gamma \\
&\quad - E_P[W_i X_i'] (n^{-1} X' X)^{-1} \left(n^{-1/2} X' W - n^{1/2} E_P[W_i X_i'] \right) \Gamma \\
&\quad - E_P[W_i X_i'] \left(n^{1/2} (n^{-1} X' X)^{-1} - n^{1/2} (E_P[X_i X_i'])^{-1} \right) E_P[X_i W_i'] \Gamma.
\end{aligned} \tag{SA-5.52}$$

On noting that $n^{-1/2} Z' V = n^{-1/2} W' V - n^{-1} W' X (n^{-1} X' X)^{-1} X' V = n^{-1/2} Z^{*'} V + o_p(1)$, $n^{-1/2} Z' u = n^{-1/2} W' u - n^{-1} W' X (n^{-1} X' X)^{-1} X' u = n^{-1/2} Z^{*'} u + o_p(1)$, by the multivariate triangular array

Lyapunov CLT

$$\begin{bmatrix} n^{1/2}\text{vec}(n^{-1}W'W - E_P[W_iW_i']) \\ n^{1/2}\text{vec}(n^{-1}W'X - E_P[W_iX_i']) \\ n^{1/2}\text{vec}(n^{-1}X'X - E_P[X_iX_i']) \\ n^{-1/2}\text{vec}(Z'V) \\ n^{-1/2}Z'u \end{bmatrix} = \begin{bmatrix} n^{1/2}\text{vec}(n^{-1}W'W - E_P[W_iW_i']) \\ n^{1/2}\text{vec}(n^{-1}W'X - E_P[W_iX_i']) \\ n^{1/2}\text{vec}(n^{-1}X'X - E_P[X_iX_i']) \\ n^{-1/2}\text{vec}(Z^*V) \\ n^{-1/2}Z^*u \end{bmatrix} + o_p(1) \xrightarrow{d} \begin{bmatrix} \text{vec}(\mathcal{Z}_{ww}) \\ \text{vec}(\mathcal{Z}_{wx}) \\ \text{vec}(\mathcal{Z}_{xx}) \\ \text{vec}(\mathcal{Z}_{zv}) \\ \mathcal{Z}_{zu} \end{bmatrix}, \quad (\text{SA-5.53})$$

where $Z^* = W - X(E_P[X_iX_i'])^{-1}E_P[X_iW_i']$ and

$$\begin{aligned} \mathcal{Z}_{ww} &\equiv [\mathcal{Z}_{ww,1}, \dots, \mathcal{Z}_{ww,k}], \quad \mathcal{Z}_{ww,j} \sim N[0, \lim_{n \rightarrow \infty} \text{Var}_P[W_iW_{ij}']], \quad j = 1, \dots, k, \\ \mathcal{Z}_{wx} &\equiv [\mathcal{Z}_{wx,1}, \dots, \mathcal{Z}_{wx,p}], \quad \mathcal{Z}_{wx,l} \sim N[0, \lim_{n \rightarrow \infty} \text{Var}_P[W_iX_{il}']], \quad l = 1, \dots, p, \\ \mathcal{Z}_{xx} &\equiv [\mathcal{Z}_{xx,1}, \dots, \mathcal{Z}_{xx,p}], \quad \mathcal{Z}_{xx,s} \sim N[0, \lim_{n \rightarrow \infty} \text{Var}_P[X_iX_{is}']], \quad s = 1, \dots, p, \\ \mathcal{Z}_{zv} &\equiv [\mathcal{Z}_{zv,1}, \dots, \mathcal{Z}_{zv,d}], \quad \mathcal{Z}_{zv,r} \sim N[0, \lim_{n \rightarrow \infty} E_P[Z_i^*Z_i^{*'}V_{ir}^2]], \quad r = 1, \dots, d, \\ \mathcal{Z}_{zu} &\sim N[0, \lim_{n \rightarrow \infty} E_P[Z_i^*Z_i^{*'}u_i^2]]. \end{aligned}$$

By the triangular array WLLN,

$$n^{-1}W'W \xrightarrow{p} Q_{ww}, \quad n^{-1}W'X \xrightarrow{p} Q_{wx}, \quad n^{-1}X'X \xrightarrow{p} Q_{xx}. \quad (\text{SA-5.54})$$

Next we derive the asymptotic distribution of $n^{1/2}((n^{-1}X'X)^{-1} - (E_P[X_iX_i'])^{-1})$. Let $f : \text{vec}(A) \rightarrow \text{vec}(A^{-1})$, where $A \in \mathbb{R}^{p \times p}$ is positive definite. The Jacobian of the mapping evaluated at $\text{vec}(A) = \text{vec}(Q_{xx})$ is $\dot{f} = -(A^{-1})' \otimes A^{-1}|_{A=Q_{xx}} = -Q_{xx}^{-1} \otimes Q_{xx}^{-1}$. Since $n^{1/2}(\text{vec}(n^{-1}X'X) - \text{vec}(E_P[X_iX_i'])) \xrightarrow{d} \text{vec}(\mathcal{Z}_{xx})$, by the delta method

$$\begin{aligned} n^{1/2}((n^{-1}X'X)^{-1} - (E_P[X_iX_i'])^{-1}) &\xrightarrow{d} \text{vec}_{p,p}^{-1}(\dot{f} \text{vec}(\mathcal{Z}_{xx})) \\ &= -\text{vec}_{p,p}^{-1}((Q_{xx}^{-1} \otimes Q_{xx}^{-1})\text{vec}(\mathcal{Z}_{xx})) \\ &= -Q_{xx}^{-1}\mathcal{Z}_{xx}Q_{xx}^{-1}, \end{aligned} \quad (\text{SA-5.55})$$

where $\text{vec}_{p,p}^{-1}$ denotes the inverse vec operator with the property that $\text{vec}_{p,p}^{-1}(\text{vec}(A)) = A$. Since $n^{1/2}((n^{-1}X'X)^{-1} - (E_P[X_iX_i'])^{-1}) = -Q_{xx}^{-1}n^{1/2}(n^{-1}X'X - E_P[X_iX_i'])Q_{xx}^{-1} + o_p(1)$ by (SA-5.53) and (SA-5.55), the convergence in (SA-5.55) holds jointly with (SA-5.53). Using (SA-5.53)-(SA-5.55)

in (SA-5.52) and invoking the Slutsky's lemma,

$$n^{1/2}(\hat{G} - G_n) \xrightarrow{d} G_h \equiv (\mathcal{Z}_{ww} - \mathcal{Z}_{wx}Q_{xx}^{-1}Q_{xw} - Q_{wx}Q_{xx}^{-1}\mathcal{Z}_{xw} + Q_{wx}Q_{xx}^{-1}\mathcal{Z}_{xx}Q_{xx}^{-1}Q_{xw})\Gamma + \mathcal{Z}_{zv}. \quad (\text{SA-5.56})$$

Then, from (SA-5.36), (SA-5.37), (SA-5.53), (SA-5.56) and the CMT

$$\begin{aligned} n^{1/2}\text{vec}(\hat{J} - G_n) &= n^{-1/2} \sum_{i=1}^n \text{vec}(Z_i Y_i' - \mathbb{E}_P[Z_i^* Z_i^{*'}] \Gamma) - [\hat{C}'_1, \dots, \hat{C}'_d]' \hat{\Sigma}^{-1} n^{1/2} \hat{m} \\ &= \text{vec}((\mathcal{Z}_{ww} - \mathcal{Z}_{wx}Q_{xx}^{-1}Q_{xw} - Q_{wx}Q_{xx}^{-1}\mathcal{Z}_{xw} + Q_{wx}Q_{xx}^{-1}\mathcal{Z}_{xx}Q_{xx}^{-1}Q_{xw})\Gamma + \mathcal{Z}_{zv}) \\ &\quad - [C'_{P1}, \dots, C'_{Pd}]' \Sigma_P^{-1} \mathcal{Z}_{zu} + o_p(1), \\ &= \text{vec}(G_h) - [C'_{P1}, \dots, C'_{Pd}]' \Sigma_P^{-1} \mathcal{Z}_{zu} + o_p(1). \end{aligned} \quad (\text{SA-5.57})$$

SA-5.3 Asymptotic distributions of key permutation quantities

In this section, we derive the asymptotic distributions of key quantities that underlie the permutation test statistics under the drifting sequence of parameters.

Conditioning events. We first describe the events on which we condition when deriving the asymptotic results for the permutation tests. From (SA-5.57), for any $\epsilon > 0$ we can find a constant U_0 and $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$.

$$\Omega_1 \equiv \{\omega \in \Omega_0 : \|n^{1/2}(\hat{J} - G_n)\| < U_0\}, \quad P[\Omega_1^c] < \epsilon/2. \quad (\text{SA-5.58})$$

We shall establish that for $\text{PR} \in \{n^{1/2}(\hat{J}^\pi - \hat{G}), n^{1/2}H_n\hat{J}^\pi U_n T_n, \text{PLM}, \text{PCLR}_a, \text{PCLR}_b\}$ and all $\omega \in \Omega_1$

$$\sup_{x \in \mathbb{R}} |P^\pi[\text{PR} \leq x] - P[\text{PR}_\infty \leq x]| \xrightarrow{p} 0, \quad (\text{SA-5.59})$$

where PR_∞ is the limiting random variable. When (SA-5.59) holds, for any $\epsilon, \epsilon_1 > 0$, there exists $n_1 \in \mathbb{N}$ such that $P[\sup_{x \in \mathbb{R}} |P^\pi[\text{PR} \leq x] - P[\text{PR}_\infty \leq x]| > \epsilon_1] \leq \epsilon/2$ for all $\omega \in \Omega_1$ and $n \geq n_1$. Then, by the Boole-Bonferroni inequality, for all $n \geq \max\{n_0, n_1\}$

$$\begin{aligned} &P[\omega \in \Omega_0 : |P^\pi[\text{PR} \leq x] - P[\text{PR}_\infty \leq x]| > \epsilon_1] \\ &\leq P[\omega \in \Omega_1 : |P^\pi[\text{PR} \leq x] - P[\text{PR}_\infty \leq x]| > \epsilon_1] + P[\Omega_1^c] \\ &\leq \epsilon. \end{aligned} \quad (\text{SA-5.60})$$

Since $\epsilon > 0$ is arbitrary, it will follow that

$$\sup_{x \in \mathbb{R}} |P^\pi[\text{PR} \leq x] - P[\text{PR}_\infty \leq x]| \xrightarrow{p} 0. \quad (\text{SA-5.61})$$

This means that we can use the distributional results such as that in (SA-5.57) in the conditioning events to obtain the convergence in P^π -distribution in P -probability, see [Mason and Newton \(1992\)](#) for a related approach.

The following lemma is a permutation analog of Lemma 16.4 (see also (SA-2.11)) of [Andrews and Guggenberger \(2019b\)](#).

Lemma SA-5.1. *Under all sequences $\{\lambda_n \in \Lambda : n \geq 1\}$,*

$$n^{1/2}[\hat{m}^\pi, \hat{J}^\pi - \hat{G}, H_n \hat{J}^\pi U_n T_n] \xrightarrow{d^\pi} [m_h^\pi, J_h^\pi, \Delta_h^\pi] \quad \text{in } P\text{-probability,}$$

where

(a)

$$\begin{bmatrix} m_h^\pi \\ \text{vec}(J_h^\pi) \end{bmatrix} \sim N \left[0_{k \times (d+1)}, \begin{pmatrix} \sigma^2 Q_{zz} & 0_{k \times dk} \\ 0_{k \times dk} & \Phi_h^\pi \end{pmatrix} \right], \quad \Phi_h^\pi \equiv (\Sigma^V - \Sigma^{Vu}(\sigma^2)^{-1}\Sigma^{uV}) \otimes Q_{zz},$$

(b)

$$\begin{aligned} \Delta_h^\pi &\equiv [\Delta_{h,q}^\pi, \Delta_{h,d-q}^\pi] \in \mathbb{R}^{k \times d}, \quad \Delta_{h,q}^\pi \equiv h_{3,q} \in \mathbb{R}^{k \times q}, \\ \Delta_{h,d-q}^\pi &\equiv h_3 \tilde{h}_{1,d-q} + h_{5,m}^{-1/2} (G_h + J_h^\pi) h_{91} h_{2,d-q} \in \mathbb{R}^{k \times (d-q)}, \end{aligned}$$

where G_h is defined in (SA-5.56), and $\tilde{h}_{1,d-q}$ and h_{91} are defined in (SA-2.12),

(c) (J_h^π, Δ_h^π) and m_h^π are independent,


(d) When $U_n \equiv I_d$ (as opposed to $U_P \equiv ((\theta_0, I_d)\Omega^\epsilon(\Sigma_P, \mathcal{V}_P)^{-1}(\theta_0, I_d)')^{1/2}$ defined in Section SA-5.1) with the eigenvalues, the orthogonal eigenvectors and the SVs of the matrices (SA-5.1)-(SA-5.3) defined accordingly as in Section SA-5.1, the above results also hold with $\Delta_{h,d-q}^\pi \equiv h_3 \tilde{h}_{1,d-q} + h_{5,m}^{-1/2} (G_h + J_h^\pi) h_{2,d-q}$ where G_h is as defined in (SA-5.56), and $\tilde{h}_{1,d-q}$ is as defined in (SA-2.12),

(e) Under all subsequences $\{w_n\}$ and all sequences $\{\lambda_{w_n,n} : n \geq 1\} \in \Lambda$, the convergence results above hold with n replaced with w_n .

SA-5.4 PAR tests

The diagram in Figure 1 gives a roadmap to the proof of the PAR test. DP abbreviates the drifting sequences of parameters and AG abbreviates [Andrews and Guggenberger \(2019b\)](#). The (uniform) asymptotic similarity of the permutation tests are established using Proposition 16.3 of [Andrews and Guggenberger \(2019b\)](#). The null asymptotic distribution of the AR statistic follows from Theorem 16.6 of [Andrews and Guggenberger \(2019b\)](#) which establishes the asymptotic distribution of the CLR statistics (thus yields that of the AR statistic).

Figure 1: Proof of the PAR tests



Notes: This figure depicts the main results needed for the asymptotic validity of the PAR tests and their dependencies. A solid arrow indicates that a result is a prerequisite of a next result. DP abbreviates the drifting sequences of parameters and AG stands for [Andrews and Guggenberger \(2019b\)](#).

Let $\tilde{W} = [\tilde{W}_1, \dots, \tilde{W}_n]' \equiv M_l W$. Since ι is a column of X , $M_l M_X = M_X$. Write

$$\begin{aligned}
 W'_\pi M_X \tilde{\Sigma}_u M_X W_\pi &= W'_\pi M_l \tilde{\Sigma}_u M_l W_\pi - W'_\pi M_l X (X' X)^{-1} X' \tilde{\Sigma}_u M_l W_\pi - W'_\pi M_l \tilde{\Sigma}_u X (X' X)^{-1} X' M_l W_\pi \\
 &\quad + W'_\pi M_l X (X' X)^{-1} X' \tilde{\Sigma}_u X (X' X)^{-1} X' M_l W_\pi.
 \end{aligned} \tag{SA-5.62}$$

Consider the first term $n^{-1} W'_\pi M_l \tilde{\Sigma}_u M_l W_\pi = n^{-1} \tilde{W}'_\pi \tilde{\Sigma}_u \tilde{W}_\pi = n^{-1} \sum_{i=1}^n \tilde{W}_{\pi(i)} \tilde{W}'_{\pi(i)} \tilde{u}_i(\theta_0)^2$ on the right-hand side of the expression above. Proceeding similarly to (SA-8.25) and noting that $\text{Var}_P[W_i] - (Q_{ww} - Q_w Q'_w) \rightarrow 0$, where Q_w denotes the column of Q_{wx} corresponding to the element 1 of X_i for Q_{ww} and Q_{wx} defined in (SA-5.22), we obtain

$$n^{-1} \sum_{i=1}^n \tilde{W}_{\pi(i)} \tilde{W}'_{\pi(i)} \tilde{u}_i(\theta_0)^2 - (Q_{ww} - Q_w Q'_w) \sigma^2 \xrightarrow{P^\pi} 0 \quad \text{in } P\text{-probability.} \tag{SA-5.63}$$

Next we determine the limit of $n^{-1}W'_\pi M_\iota X$. Note that letting $[\tilde{X}_1, \dots, \tilde{X}_n]' \equiv M_\iota X$, $\tilde{W}_i = [\tilde{W}_{i1}, \dots, \tilde{W}_{ik}]'$, $\tilde{W}_{ij} = W_{ij} - \bar{W}_j$, $\bar{W}_j \equiv n^{-1} \sum_{i=1}^n W_{ij}$, and $\tilde{X}_i = [\tilde{X}_{i1}, \dots, \tilde{X}_{ip}]'$,

$$\begin{aligned} \mathbb{E}_{P^\pi}[n^{-1}W'_\pi M_\iota X|y, Y, W, X] &= \left(n^{-1} \sum_{i=1}^n \tilde{W}_i\right) \left(n^{-1} \sum_{i=1}^n \tilde{X}_i\right) = 0, \\ \text{Var}_{P^\pi} \left[n^{-1} \sum_{i=1}^n \tilde{W}_{\pi(i)j} \tilde{X}_{il} | y, Y, W, X \right] &= \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n \tilde{W}_{ij}^2 \right) \left(n^{-1} \sum_{i=1}^n \tilde{X}_{il}^2 \right) \rightarrow 0, \\ j &= 1, \dots, k, \quad l = 1, \dots, p, \end{aligned}$$

where the convergence uses $n^{-1} \sum_{i=1}^n \tilde{W}_{ij}^2 - (\mathbb{E}_P[W_{ij}^2] - (\mathbb{E}_P[W_{ij}])^2) \xrightarrow{a.s.} 0$ and $n^{-1} \sum_{i=1}^n \tilde{X}_{il}^2 - (\mathbb{E}_P[X_{il}^2] - (\mathbb{E}_P[X_{il}])^2) \xrightarrow{a.s.} 0$, which, in turn, hold by the triangular array SLLN and the CMT. By Chebyshev's inequality, we obtain

$$n^{-1}W'_\pi M_\iota X \xrightarrow{p^\pi} 0 \quad \text{a.s.} \quad (\text{SA-5.64})$$

Consider the part $n^{-1}X'\tilde{\Sigma}_u M_\iota W_\pi = n^{-1} \sum_{i=1}^n X_i \tilde{W}'_{\pi(i)} \tilde{u}_i(\theta_0)^2$ in the second term of (SA-5.62).

Using the Cauchy-Schwarz inequality twice,

$$\|n^{-1}X'\tilde{\Sigma}_u M_\iota W_\pi\| \leq \left(n^{-1} \sum_{i=1}^n \|X_i\|^4\right)^{1/4} \left(n^{-1} \sum_{i=1}^n \|\tilde{W}_i\|^4\right)^{1/4} \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4\right)^{1/2} = O_p(1), \quad (\text{SA-5.65})$$

where we used $n^{-1} \sum_{i=1}^n \|X_i\|^4 = O_p(1)$ and $n^{-1} \sum_{i=1}^n \|\tilde{W}_i\|^4 \leq n^{-1} \sum_{i=1}^n (8\|W_i\|^4 + 8\|\tilde{W}\|^4) = O_p(1)$ which hold by the triangular array WLLN, and (SA-8.22). Furthermore, note that

$$\begin{aligned} n^{-1} \sum_{i=1}^n X_{ij} X_{il} u_i X'_i - \mathbb{E}_P[X_{ij} X_{il} u_i X'_i] &\xrightarrow{p} 0, \\ n^{-1} \sum_{i=1}^n X_{ij} X_{il} X_i X'_i - \mathbb{E}_P[X_{ij} X_{il} X_i X'_i] &\xrightarrow{p} 0, \quad j, l = 1, \dots, p, \end{aligned}$$

by the WLLN, $E_P[\|X_{ij}X_{il}u_iX_i'\|^{1+\delta/4}] \leq (E_P[\|X_i\|^{4+\delta}])^{1/2}(E_P[|u_i|^{4+\delta}])^{1/4}(E_P[\|X_i\|^{4+\delta}])^{1/4} < \infty$ by Cauchy-Schwarz inequality, and $E[\|X_i\|^{4+\delta}] < \infty$. Therefore, we have

$$\begin{aligned}
& n^{-1}X'\tilde{\Sigma}_uX - E_P[X_iX_i'u_i^2] \\
&= n^{-1}\sum_{i=1}^n X_iX_i'\tilde{u}_i(\theta_0)^2 - E_P[X_iX_i'u_i^2] \\
&= n^{-1}\sum_{i=1}^n X_iX_i'(u_i(\theta_0))^2 - 2u_iX_i'(X'X)^{-1}X'u + u'X(X'X)^{-1}X_iX_i'(X'X)^{-1}X'u \\
&\quad - E_P[X_iX_i'u_i^2] \\
&\xrightarrow{p} 0,
\end{aligned} \tag{SA-5.66}$$

where the last line uses the WLLN, CMT, $E_P[\|X_iX_i'u_i^2\|^{1+\delta/4}] \leq (E_P[\|X_i\|^{4+\delta}])^{1/2}(E_P[|u_i|^{4+\delta}])^{1/2}$, $n^{-1}X'u \xrightarrow{p} 0$ and $(X'X)^{-1} - (E_P[X_iX_i'])^{-1} \xrightarrow{p} 0$ which follow from the WLLN and the CMT. Using (SA-5.63), (SA-5.64), (SA-5.65), (SA-5.66), and the CMT in (SA-5.62), we obtain

$$n^{-1}W'_\pi M_X \tilde{\Sigma}_u M_X W_\pi - (Q_{ww} - Q_w Q'_w)\sigma^2 \xrightarrow{p} 0 \quad \text{in } P\text{-probability.} \tag{SA-5.67}$$

Next, we determine the limit of $n^{-1/2}\tilde{W}'_\pi \tilde{u}(\theta_0)$. By the c_r inequality and the triangular array SLLN,

$$\begin{aligned}
& n^{-1}\sum_{i=1}^n |\tilde{u}_i(\theta_0)|^{2+\delta} - 2^{1+\delta} E_P[|u_i|^{2+\delta}] \\
&\leq 2^{1+\delta}(n^{-1}\sum_{i=1}^n |u_i|^{2+\delta} - E_P[|u_i|^{2+\delta}]) + n^{-1}\sum_{i=1}^n \|X_i\|^{2+\delta}\|(X'X)^{-1}X'u\|^{2+\delta} \\
&\xrightarrow{a.s.} 0,
\end{aligned} \tag{SA-5.68}$$

and for any $t \in \mathbb{R}^k$

$$\begin{aligned}
n^{-1}\sum_{i=1}^n \|t'\tilde{W}_i\|^{2+\delta} &\leq \|t\|^{2+\delta}n^{-1}\sum_{i=1}^n \|\tilde{W}_i\|^{2+\delta} \\
&\leq \|t\|^{2+\delta}2^{1+\delta}\left(n^{-1}\sum_{i=1}^n \|W_i\|^{2+\delta} + \|n^{-1}\sum_{i=1}^n W_i\|^{2+\delta}\right) \\
&= O_{a.s.}(1),
\end{aligned} \tag{SA-5.69}$$

$$\begin{aligned}
\tilde{D} &\equiv \frac{1}{n-1}\sum_{i=1}^n \sum_{j=1}^n g_n^2(i, j) = \frac{1}{n(n-1)}\sum_{i=1}^n \sum_{j=1}^n (t'\tilde{W}_i \tilde{u}_j(\theta_0))^2 = \frac{1}{n-1}\sum_{i=1}^n \tilde{u}_i(\theta_0)^2 n^{-1}\sum_{i=1}^n (t'\tilde{W}_i)^2 \\
&= \sigma^2 t' \text{Var}_P[W_i]t + o_{a.s.}(1).
\end{aligned} \tag{SA-5.70}$$

Note that by (SA-5.67), (SA-5.69) and (SA-5.70), for any $\epsilon > 0$

$$\begin{aligned}
& n^{-2} \sum_{i=1}^n \sum_{j=1}^n \frac{(t' \tilde{W}_i \tilde{u}_j(\theta_0))^2}{\tilde{D}} 1 \left(\frac{n^{-1} (t' \tilde{W}_i \tilde{u}_j(\theta_0))^2}{\tilde{D}} > \epsilon^2 \right) \\
&= n^{-2} \tilde{D}^{-1} \sum_{i=1}^n \sum_{j=1}^n \frac{|t' \tilde{W}_i \tilde{u}_j(\theta_0)|^{2+\delta}}{|t' \tilde{W}_i \tilde{u}_j(\theta_0)|^\delta} 1 \left(|t' \tilde{W}_i \tilde{u}_j(\theta_0)|^\delta > \epsilon^\delta \tilde{D}^{\delta/2} n^{\delta/2} \right) \\
&\leq n^{-\delta/2} \epsilon^{-\delta} \tilde{D}^{-1-\delta/2} n^{-2} \sum_{i=1}^n \sum_{j=1}^n |t' \tilde{W}_i \tilde{u}_j(\theta_0)|^{2+\delta} \\
&\leq n^{-\delta/2} \epsilon^{-\delta} \tilde{D}^{-1-\delta/2} n^{-1} \sum_{i=1}^n \|t' \tilde{W}_i\|^{2+\delta} n^{-1} \sum_{j=1}^n |\tilde{u}_j(\theta_0)|^{2+\delta} \\
&\xrightarrow{a.s.} 0.
\end{aligned} \tag{SA-5.71}$$

Thus,

$$\lim_{n \rightarrow \infty} n^{-2} \sum_{i=1}^n \sum_{j=1}^n \frac{(t' \tilde{W}_i \tilde{u}_j(\theta_0))^2}{\tilde{D}} 1 \left(\frac{n^{-1} (t' \tilde{W}_i \tilde{u}_j(\theta_0))^2}{\tilde{D}} > \epsilon^2 \right) = 0. \tag{SA-5.72}$$

On noting that we may write $n^{-1/2} W'_\pi M_X \tilde{u}(\theta_0) = n^{-1/2} \tilde{W}'_\pi \tilde{u}(\theta_0) = n^{-1/2} \tilde{W}'_\pi \tilde{u}_{\tilde{\pi}}(\theta_0)$ for a permutation $\tilde{\pi}$ uniformly distributed over \mathbb{G}_n , Lemma SA-2.1 gives

$$n^{-1/2} t' \tilde{W}'_\pi \tilde{u}(\theta_0) \xrightarrow{d^\pi} N[0, t'(Q_{ww} - Q_w Q'_w) t \sigma^2] \quad P\text{-almost surely.} \tag{SA-5.73}$$

Using the Cramér-Wold device

$$n^{-1/2} t' \tilde{W}'_\pi \tilde{u}(\theta_0) \xrightarrow{d^\pi} N[0, (Q_{ww} - Q_w Q'_w) \sigma^2] \quad P\text{-almost surely.} \tag{SA-5.74}$$

Finally, from the Slutsky's lemma, (SA-5.67), (SA-5.74) followed by the Polya's theorem (Theorem 11.2.9 of Lehmann and Romano (2005)), we obtain $\text{PAR}_1 \xrightarrow{d^\pi} \text{PAR}_{1,\infty} \sim \chi_k^2$ in P -probability, and

$$\sup_{x \in \mathbb{R}} |\hat{F}_n^{\text{PAR}_1}(x) - P[\chi_k^2 \leq x]| \xrightarrow{p} 0. \tag{SA-5.75}$$

The above result holds for any subsequence $\{w_n\}$ of $\{n\}$.

Next we will show that $\text{PAR}_{1(r)} \xrightarrow{p} r_{1-\alpha}(\chi_k^2)$, where $r_{1-\alpha}(\chi_k^2)$ denotes the $1 - \alpha$ quantile of χ_k^2 distribution. First, recall that $\mathbb{G}'_n = \{\pi_1, \pi_2, \dots, \pi_N\}$, where π_1 is the identity permutation and π_2, \dots, π_N are i.i.d. uniformly distributed over \mathbb{G}_n . Let

$$\bar{F}_N^{\text{PAR}_1}(x) \equiv \frac{1}{N-1} \sum_{i=2}^N 1(\text{AR}^{\pi_i} \leq x). \tag{SA-5.76}$$

By Theorem 12.2.18 of [Lehmann and Romano \(2005\)](#), there exists a universal constant C_0 such that for any positive integer $N \geq 2$

$$P^\pi \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - \hat{F}_n^{\text{PAR}_1}(x) \right| > \epsilon \right] \leq C_0 \exp(-2(N-1)\epsilon^2). \quad (\text{SA-5.77})$$

By the iterated expectation

$$\begin{aligned} & P \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - \hat{F}_n^{\text{PAR}_1}(x) \right| > \epsilon \right] \\ &= E_P \left[1 \left(\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - \hat{F}_n^{\text{PAR}_1}(x) \right| > \epsilon \right) \right] \\ &= E_P \left[E_{P^\pi} \left[1 \left(\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - \hat{F}_n^{\text{PAR}_1}(x) \right| > \epsilon \right) \middle| y, Y, W, X \right] \right] \\ &= E_P \left[P^\pi \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - \hat{F}_n^{\text{PAR}_1}(x) \right| > \epsilon \right] \right] \\ &\rightarrow 0, \end{aligned} \quad (\text{SA-5.78})$$

where the convergence holds by letting $N \rightarrow \infty$ in [\(SA-5.77\)](#). Then, by the triangle inequality, [\(SA-5.75\)](#) and [\(SA-5.78\)](#)

$$\begin{aligned} \sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - P[\chi_k^2 \leq x] \right| &\leq \sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PAR}_1}(x) - \hat{F}_n^{\text{PAR}_1}(x) \right| + \sup_{x \in \mathbb{R}} \left| \hat{F}_n^{\text{PAR}_1}(x) - P[\chi_k^2 \leq x] \right| \\ &\xrightarrow{p} 0. \end{aligned} \quad (\text{SA-5.79})$$

Therefore, the empirical cdf $\tilde{F}_N^{\text{PAR}_1}(x)$ of PAR_1 on \mathbb{G}'_n satisfies

$$\tilde{F}_N^{\text{PAR}_1}(x) = N^{-1} 1(\text{AR}^{\pi_1} \leq x) + \frac{N-1}{N} \bar{F}_N^{\text{PAR}_1}(x) \xrightarrow{p} P[\chi_k^2 \leq x]. \quad (\text{SA-5.80})$$

Since χ_k^2 distribution function is continuous and strictly increasing at its $1 - \alpha$ quantile, we have $\text{PAR}_{1(r)} \xrightarrow{p} r_{1-\alpha}(\chi_k^2)$ by Lemma 11.2.1 of [Lehmann and Romano \(2005\)](#) and [\(SA-5.80\)](#). By definition, $E_P[\phi_n^{\text{PAR}_1}] = P[\text{AR} > \text{PAR}_{1(r)}] + a^{\text{PAR}_1} P[\text{AR} = \text{PAR}_{1(r)}]$, hence

$$P[\text{AR} > \text{PAR}_{1(r)}] \leq E_P[\phi_n^{\text{PAR}_1}] \leq P[\text{AR} \geq \text{PAR}_{1(r)}].$$

By Theorem 16.6 of [Andrews and Guggenberger \(2019b\)](#), $\text{AR} \xrightarrow{d} \text{AR}_\infty \sim \chi_k^2$. By Slutsky's lemma (Corollary 11.2.3 of [Lehmann and Romano \(2005\)](#)), we have $\lim_{n \rightarrow \infty} P[\text{AR} > \text{PAR}_{1(r)}] = \lim_{n \rightarrow \infty} P[\text{AR} \geq \text{PAR}_{1(r)}] = P[\text{AR}_\infty > r_{1-\alpha}(\chi_k^2)]$ because of the continuity of the χ_k^2 distribution function at its $1 - \alpha$ quantile. Therefore,

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PAR}_1}] = P[\text{AR}_\infty > r_{1-\alpha}(\chi_k^2)] = \alpha.$$

Thus, Assumption 2 is verified for the PAR_1 statistic. By Proposition 16.3 of Andrews and Guggenberger (2019b), we obtain the desired result.

PAR₂ test: By (SA-8.44), we have $\text{PAR}_2 \xrightarrow{d^\pi} \chi_k^2$ in P -probability, and by the Polya's theorem


$$\sup_{x \in \mathbb{R}} |\hat{F}_n^{\text{PAR}_2}(x) - P[\chi_k^2 \leq x]| \xrightarrow{p} 0.$$

The remaining argument is analogous to the PAR_1 test.

SA-5.5 PLM test

The diagram in Figure 2 gives a roadmap to the proof of the PLM test. DP abbreviates the drifting sequences of parameters and AG abbreviates Andrews and Guggenberger (2019b).

Figure 2: Proof of the PLM test



Notes: This figure depicts the main results needed for the asymptotic validity of the PLM test and their dependencies. A solid arrow indicates that a result is a prerequisite of a next result. DP abbreviates the drifting sequences of parameters, AG stands for Andrews and Guggenberger (2019b), and AG'17a, b abbreviate Andrews and Guggenberger (2017a,b)

Note first that because Σ_n and T_n are nonsingular

$$\begin{aligned} \text{PLM} &= n \hat{m}'^{\pi'} \hat{\Sigma}^{\pi-1/2} P_{\hat{\Sigma}^{\pi-1/2} \hat{J}^{\pi}} \hat{\Sigma}^{\pi-1/2} \hat{m}^{\pi} \\ &= n \hat{m}'^{\pi'} \hat{\Sigma}^{\pi-1/2} P_{(\hat{\Sigma}^{\pi-1/2} \Sigma_n^{1/2}) n^{1/2} \Sigma_n^{-1/2} \hat{J}^{\pi} T_n} \hat{\Sigma}^{\pi-1/2} \hat{m}^{\pi}. \end{aligned}$$

By Lemma SA-5.1(d), $n^{1/2}\Sigma_n^{-1/2}\hat{J}^\pi T_n \xrightarrow{d^\pi} \Delta_h^\pi = [\Delta_{h,q}^\pi, \Delta_{h,d-q}^\pi] = [h_{3,q}, h_3\tilde{h}_{1,d-q} + h_{5,m}^{-1/2}(G_h + J_h^\pi)h_{2,d-q}]$, where $h_{5,m} = \Sigma$. The only random term conditional on the data is $h_{5,m}^{-1/2}J_h^\pi h_{2,d-q}$. We apply Corollary 16.2 of Andrews and Guggenberger (2017b) with $p = d$, $q_* = q$, $\Delta_{q_*} = h_{3,q} \in \mathbb{R}^{k \times q}$, $\Delta_{p-q_*} = \Delta_{h,d-q}^\pi \in \mathbb{R}^{k \times (d-q)}$, $M = h'_3 \in \mathbb{R}^{k \times k}$, $M_1 = h'_{3,q} \in \mathbb{R}^{q \times k}$, $M_2 = h'_{3,k-q} \in \mathbb{R}^{(k-q) \times k}$, $\xi_2 \in \mathbb{R}^{d-q}$ and $\Delta = \Delta_h^\pi \in \mathbb{R}^{k \times d}$. Note that for $\xi_2 \in \mathbb{R}^{d-q}$ with $\|\xi_2\| = 1$,

$$\begin{aligned}
& \text{Var}_{P^\pi}(h'_{3,k-q}\Sigma^{-1/2}J_h^\pi h_{2,d-q}\xi_2) \\
&= \text{Var}_{P^\pi}(\text{vec}(h'_{3,k-q}\Sigma^{-1/2}J_h^\pi h_{2,d-q}\xi_2)) \\
&= \text{Var}_{P^\pi}\left(\left((h_{2,d-q}\xi_2)' \otimes (h'_{3,k-q}\Sigma^{-1/2})\right)(\text{vec}(J_h^\pi))\right) \\
&= \left((h_{2,d-q}\xi_2)' \otimes (h'_{3,k-q}\Sigma^{-1/2})\right) \text{Var}_{P^\pi}(\text{vec}(J_h^\pi)) \left((h_{2,d-q}\xi_2) \otimes (\Sigma^{-1/2}h_{3,k-q})\right) \\
&= \left((h_{2,d-q}\xi_2)' \otimes (h'_{3,k-q}\Sigma^{-1/2})\right) \left((\Sigma^V - \Sigma^{Vu}(\sigma^2)^{-1}\Sigma^{uV}) \otimes Q_{zz}\right) \left((h_{2,d-q}\xi_2) \otimes (\Sigma^{-1/2}h_{3,k-q})\right), \\
&= (\xi_2' h'_{2,d-q}(\Sigma^V - \Sigma^{Vu}(\sigma^2)^{-1}\Sigma^{uV})h_{2,d-q}\xi_2)(h'_{3,k-q}\Sigma^{-1/2}Q_{zz}\Sigma^{-1/2}h_{3,k-q}),
\end{aligned}$$

where the second equality uses the formula $\text{vec}(ABC) = (C' \otimes A)\text{vec}(B)$ and the fourth equality uses Lemma SA-5.1(a). Observe that $\xi_2' h'_{2,d-q}(\Sigma^V - \Sigma^{Vu}(\sigma^2)^{-1}\Sigma^{uV})h_{2,d-q}\xi_2$ is scalar, and

$$\frac{\xi_2' h'_{2,d-q}(\Sigma^V - \Sigma^{Vu}(\sigma^2)^{-1}\Sigma^{uV})h_{2,d-q}\xi_2}{\xi_2' h'_{2,d-q}h_{2,d-q}\xi_2} \geq \lambda_{\min}(\Sigma^V - \Sigma^{Vu}(\sigma^2)^{-1}\Sigma^{uV}) > 0.$$

Moreover, using $h'_{3,k-q}h_{3,k-q} = I_{k-q}$

$$\begin{aligned}
\lambda_{\min}(h'_{3,k-q}\Sigma^{-1/2}Q_{zz}\Sigma^{-1/2}h_{3,k-q}) &= \min_{\eta \in \mathbb{R}^{k-q} \setminus \{0\}} \frac{\eta' h'_{3,k-q}\Sigma^{-1/2}Q_{zz}\Sigma^{-1/2}h_{3,k-q}\eta}{\eta'\eta} \\
&= \min_{\eta \in \mathbb{R}^{k-q} \setminus \{0\}} \frac{\eta' h'_{3,k-q}\Sigma^{-1/2}Q_{zz}\Sigma^{-1/2}h_{3,k-q}\eta}{\eta' h'_{3,k-q}h_{3,k-q}\eta} \\
&\geq \min_{a \in \mathbb{R}^k \setminus \{0\}} \frac{a'\Sigma^{-1/2}Q_{zz}\Sigma^{-1/2}a}{a'a} \\
&\geq \lambda_{\min}(\Sigma^{-1})\lambda_{\min}(Q_{zz}) \\
&> 0.
\end{aligned}$$

Therefore, $\text{rank}(\text{Var}_{P^\pi}(h'_{3,k-q}\Sigma^{-1/2}J_h^\pi h_{2,d-q}\xi_2)) = k - q \geq d - q$, and by Corollary 16.2 of Andrews and Guggenberger (2017b), $P^\pi[\text{rank}(\Delta_h^\pi) = d] = 1$ in P -probability. Since $(\Sigma^\pi)^{-1/2}\Sigma^{1/2} = (\sigma^2 Q_{zz})^{-1/2}\Sigma^{1/2}$ is nonsingular, $P^\pi[\text{rank}((\Sigma^\pi)^{-1/2}\Sigma^{1/2}\Delta_h^\pi) = d] = 1$ in P -probability. As shown in the proof of Theorem 11.1 of Andrews and Guggenberger (2017b), the mapping defined as

$f : \mathbb{R}^{k \times d} \rightarrow \mathbb{R}^{k \times k}$, $f(D) = D(D'D)^{-1}D'$ is continuous at $D \in \mathbb{R}^{k \times d}$ that has full column rank with probability one. Then,


$$\begin{aligned} \text{PLM} &= n \hat{m}^{\pi'} \hat{\Sigma}^{\pi-1/2} P_{(\hat{\Sigma}^{\pi-1/2} \Sigma_n^{1/2}) n^{1/2} \Sigma_n^{-1/2} \hat{J}^{\pi} T_n} \hat{\Sigma}^{\pi-1/2} \hat{m}^{\pi} \\ &\xrightarrow{d^{\pi}} m_h^{\pi'} \Sigma^{\pi-1/2} P_{(\Sigma^{\pi})^{-1/2} \Sigma^{1/2} \Delta_h^{\pi}} \Sigma^{\pi-1/2} m_h^{\pi} \\ &\sim \chi_d^2 \quad \text{in } P\text{-probability,} \end{aligned}$$

where the convergence follows from the CMT, Lemma SA-5.1 and the full column rank property established above, and the last line holds because using the independence between Δ_h^{π} and m_h^{π} , the full column rank of Δ_h^{π} and $\Sigma^{\pi-1/2} m_h^{\pi} \sim N[0, I_k]$, the equality in distribution holds conditional on Δ_h^{π} , hence also holds unconditionally. The null asymptotic distribution of the sample LM statistic follows from Theorem 4.1 of Andrews and Guggenberger (2017a) because the parameter space assumptions in the equations (3.3), (3.9) and (3.10) of Andrews and Guggenberger (2017a) are implied by Assumption 2. The asymptotic similarity of the PLM test is similar to the PAR tests, thus is omitted.

SA-5.6 PCLR tests

The diagram in Figure 3 gives a roadmap to the proof of the PCLR test. Lemmas 16.2, 27.3, 16.4 and 26.1 of Andrews and Guggenberger (2019b) are the key results for their Theorem 16.6 that establishes the asymptotic distribution of the CLR statistics under the drifting sequences of parameters. We will use their Lemma 16.2 and Lemma 27.3 (restated in Section SA-2.2) to establish the asymptotic validity of the PCLR tests. Lemmas SA-5.1 and SA-5.2 of this paper are the permutation analogs of their Lemma 16.4 and Lemma 26.1, and are the key results for the asymptotic distribution of the PCLR statistics (Theorem SA-2.2). Combining the results of the CLR and PCLR statistics, we obtain the asymptotic validity of the PCLR tests.

Figure 3: Proof of the PCLR test



Notes: This figure depicts the main results needed for the asymptotic validity of the PCLR tests and their dependencies. A solid arrow indicates that a result is a prerequisite of a next result. A dotted line means analogous results. A solid line means related results. DP abbreviates the drifting sequences of parameters and AG abbreviates [Andrews and Guggenberger \(2019b\)](#).

Define

$$\hat{J}^+ \equiv [\hat{J}, \hat{H}^{-1} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi] \in \mathbb{R}^{k \times (d+1)}, \quad (\text{SA-5.81})$$

$$\hat{U}^+ \equiv \begin{bmatrix} \hat{U} & 0_{d \times 1} \\ 0_{1 \times d} & 1 \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}, \quad U_n^+ \equiv \begin{bmatrix} U_n & 0_{d \times 1} \\ 0_{1 \times d} & 1 \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}, \quad (\text{SA-5.82})$$

$$h_{91}^+ \equiv \begin{bmatrix} h_{91} & 0_{d \times 1} \\ 0_{1 \times d} & 1 \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}, \quad B_n^+ \equiv \begin{bmatrix} B_n & 0_{d \times 1} \\ 0_{1 \times d} & 1 \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}, \quad (\text{SA-5.83})$$

where h_{91} is as defined in (SA-2.12). Also, let

$$\begin{aligned} B_n^+ &= [B_{n,q}^+, B_{n,d+1-q}^+], \quad B_{n,q}^+ \in \mathbb{R}^{(d+1) \times q}, \quad B_{n,d+1-q}^+ \in \mathbb{R}^{(d+1) \times (d+1-q)}, \\ J_n^+ &\equiv [G_n, 0_{k \times 1}] \in \mathbb{R}^{k \times (d+1)}, \quad \Upsilon_n^+ \equiv [\Upsilon_n, 0_{k \times 1}] \in \mathbb{R}^{k \times (d+1)}, \\ S_n^+ &\equiv \text{diag}((n^{1/2}\tau_{1n})^{-1}, \dots, (n^{1/2}\tau_{qn})^{-1}, 1, \dots, 1) = \begin{bmatrix} S_n & 0_{d \times 1} \\ 0_{1 \times d} & 1 \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}. \end{aligned}$$

Let $\hat{\kappa}_1^+ \geq \dots \geq \hat{\kappa}_{d+1}^+ \geq 0$ denote the eigenvalues of

$$n \hat{U}^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ \hat{U}^+. \quad (\text{SA-5.84})$$

SA-5.6.1 Asymptotic distribution of the PCLR statistics

The following lemma is a permutation analog of Lemma 26.1 of Andrews and Guggenberger (2019b) and will be used for deriving the asymptotic distribution of the PCLR statistics.

Lemma SA-5.2 (Rates of convergence of sample eigenvalues). *Under all sequences $\{\lambda_{n,h} : n \geq 1\}$ with $\lambda_{n,h} \in \Lambda$ and $q \geq 1$,*

$$(a) \quad \hat{\kappa}_j^+ \xrightarrow{p^\pi} \infty \text{ in } P\text{-probability for all } j \leq q,$$

$$(b) \quad \hat{\kappa}_j^+ = o_{p^\pi}((n^{1/2}\tau_{ln})^2) \text{ in } P\text{-probability for all } l \leq q \text{ and } j = q+1, \dots, d+1,$$

Under all subsequences $\{w_n\}$ and all sequences $\{\lambda_{w_n,h} : n \geq 1\}$ with $\lambda_{w_n,h} \in \Lambda$, the same results hold with n replaced with w_n .

The proof of Lemma SA-5.2 is essentially the same as the proofs of Lemma 26.1 of Andrews and Guggenberger (2019b) and Lemma 17.1 of Andrews and Guggenberger (2017b) with only minor (mostly notational) modifications. For completeness, we reproduce the arguments in Section SA-8.6 after the proof of Theorem SA-5.3. Next we state a result analogous to Theorem 16.6 of Andrews and Guggenberger (2019b).

Theorem SA-5.3 (Asymptotic distribution of the PCLR statistic). *Under all sequences $\{\lambda_{n,h} : n \geq 1\}$ with $\lambda_{n,h} \in \Lambda$,*

$$\text{PCLR} \xrightarrow{d^\pi} \text{PCLR}_\infty \equiv \mathcal{Z}^{\pi'} \mathcal{Z}^\pi - \lambda_{\min} [(\Delta_{h,d-q}, \mathcal{Z}^\pi)' h_{3,k-q} h'_{3,k-q} (\Delta_{h,d-q}, \mathcal{Z}^\pi)] \quad \text{in } P\text{-probability},$$

where $\mathcal{Z}^\pi \equiv (\sigma^2 Q_{zz})^{-1/2} m_h^\pi \sim N[0, I_k]$, and the convergence holds jointly with the convergence in Lemma SA-5.1. If $q = d$, $\text{PCLR}_\infty = \mathcal{Z}^{\pi'} h_{3,d} h'_{3,d} \mathcal{Z}^\pi \sim \chi_d^2$. Under all subsequences $\{w_n\}$ and all sequences $\{\lambda_{w_n, h} : n \geq 1\}$ with $\lambda_{w_n, h} \in \Lambda$, the above results hold with n replaced with w_n .

SA-5.6.2 Completing the proof

By Theorem SA-5.3,

$$\text{PCLR} \xrightarrow{d^\pi} \text{PCLR}_\infty = \mathcal{Z}^{\pi'} \mathcal{Z}^\pi - \lambda_{\min} [(\Delta_{h,d-q}, \mathcal{Z}^\pi)' h_{3,k-q} h'_{3,k-q} (\Delta_{h,d-q}, \mathcal{Z}^\pi)] \quad \text{in } P\text{-probability,}$$

hence at any continuity point x of $P[\text{PCLR}_\infty \leq x]$

$$|\hat{F}_n^{\text{PCLR}}(x, \hat{\mathcal{T}}) - P[\text{PCLR}_\infty \leq x]| \xrightarrow{p} 0. \quad (\text{SA-5.85})$$

For π_2, \dots, π_N i.i.d. uniformly distributed over \mathbb{G}_n , let

$$\bar{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) \equiv \frac{1}{N-1} \sum_{i=2}^N 1(\text{CLR}(\hat{\mathcal{S}}^{\pi_i}, \hat{\mathcal{T}}) \leq x). \quad (\text{SA-5.86})$$

By Theorem 12.2.18 of Lehmann and Romano (2005), for some universal constant C_0 and any positive integer $N \geq 2$

$$P^\pi \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) - \hat{F}_n^{\text{PCLR}}(x, \hat{\mathcal{T}}) \right| > \epsilon \right] \leq C_0 \exp(-2(N-1)\epsilon^2). \quad (\text{SA-5.87})$$

Using the law of iterated expectations and letting $N \rightarrow \infty$ in (SA-5.87),

$$\begin{aligned} P \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) - \hat{F}_n^{\text{PCLR}}(x, \hat{\mathcal{T}}) \right| > \epsilon \right] &= E_P \left[P^\pi \left[\sup_{x \in \mathbb{R}} \left| \bar{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) - \hat{F}_n^{\text{PCLR}}(x, \hat{\mathcal{T}}) \right| > \epsilon \right] \right] \\ &\rightarrow 0. \end{aligned} \quad (\text{SA-5.88})$$

By the triangle inequality, (SA-5.85) and (SA-5.89)

$$\begin{aligned} \left| \bar{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) - P[\text{PCLR}_\infty \leq x] \right| &\leq \left| \hat{F}_n^{\text{PCLR}}(x, \hat{\mathcal{T}}) - P[\text{PCLR}_\infty \leq x] \right| \\ &\quad + \left| \bar{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) - \hat{F}_n^{\text{PCLR}}(x, \hat{\mathcal{T}}) \right| \\ &\xrightarrow{p} 0. \end{aligned} \quad (\text{SA-5.89})$$

From (SA-5.89), the permutation distribution $\tilde{F}_N^{\text{PCLR}}(x)$ of PCLR on \mathbb{G}'_n satisfies

$$\tilde{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) = N^{-1} (\text{CLR}(\hat{\mathcal{S}}^{\pi_1}, \hat{\mathcal{T}}) \leq x) + \frac{N-1}{N} \bar{F}_N^{\text{PCLR}}(x, \hat{\mathcal{T}}) \xrightarrow{p} P[\text{PCLR}_\infty \leq x]. \quad (\text{SA-5.90})$$

Denote the SVs of $h'_{3,q}\Delta_{h,d-q}$ by $\bar{\tau}_h^{(2)} = [\bar{\tau}_{(q+1)h}, \dots, \bar{\tau}_{dh}]'$. Then, $r_{k,d,q}(h'_{3,k-q}\Delta_{h,d-q}, 1 - \alpha) = r_{k,d,q}(\bar{\tau}_h^{(2)}, 1 - \alpha)$ and the distribution of PCLR_∞ is continuous and strictly increasing at its $1 - \alpha$ quantile by Lemma SA-2.7 and SA-2.8 because conditional on the data (e.g. $\omega \in \Omega_1$ with Ω_1 defined in (SA-5.58)), $h'_{3,q}\Delta_{h,d-q}$ is nonrandom. Let $\text{PCLR}_{(r)} = \text{PCLR}_{(r)}(\hat{\mathcal{T}})$. It follows from (SA-5.89) and Lemma 11.2.1 of Lehmann and Romano (2005) (see also Theorem 15.2.3 therein) that

$$\text{PCLR}_{(r)} \xrightarrow{p} r_{k,d,q}(h'_{3,k-q}\Delta_{h,d-q}, 1 - \alpha). \quad (\text{SA-5.91})$$

By Lemma 16.6 of Andrews and Guggenberger (2019b),

$$\text{CLR} \xrightarrow{d} \text{CLR}_\infty \equiv \mathcal{Z}'\mathcal{Z} - \lambda_{\min} [(\Delta_{h,d-q}, \mathcal{Z})' h_{3,k-q} h'_{3,k-q} (\Delta_{h,d-q}, \mathcal{Z})], \quad \mathcal{Z} \sim N[0_{k \times 1}, I_k]. \quad (\text{SA-5.92})$$

By the CMT, (SA-5.91) and (SA-5.92)

$$\text{CLR} - \text{PCLR}_{(r)} \xrightarrow{d} \text{CLR}_\infty - r_{k,d,q}(h'_{3,k-q}\Delta_{h,d-q}, 1 - \alpha). \quad (\text{SA-5.93})$$

As shown in Andrews and Guggenberger (2019b) p.72,

$$P[\text{CLR}_\infty = r_{k,d,q}(h'_{3,k-q}\Delta_{h,d-q}, 1 - \alpha)] = 0. \quad (\text{SA-5.94})$$

Then,

$$\begin{aligned} P[\text{CLR} > \text{PCLR}_{(r)}] &\rightarrow P[\text{CLR}_\infty > r_{k,d,q}(h'_{3,k-q}\Delta_{h,d-q}, 1 - \alpha)] \\ &= E[P[\text{CLR}_\infty > r_{k,d,q}(h'_{3,k-q}\Delta_{h,d-q}, 1 - \alpha) | \Delta_{h,d-q}]] \\ &= \alpha, \end{aligned}$$

where the convergence follows from (SA-5.93) and (SA-5.94), the first equality holds by the law of total expectations, and the last equality holds because the conditional rejection probability is α . (SA-5.94) also implies that $\lim_{n \rightarrow \infty} P[\text{CLR} \geq \text{PCLR}_{(r)}] = \alpha$. Since $E_P[\phi_n^{\text{PCLR}}] = P[\text{CLR} > \text{PCLR}_{(r)}] + a^{\text{PCLR}} P[\text{CLR} = \text{PCLR}_{(r)}]$,

$$P[\text{CLR} > \text{PCLR}_{(r)}] \leq E_P[\phi_n^{\text{PCLR}}] \leq P[\text{PCLR} \geq \text{PCLR}_{(r)}].$$

By the sandwich rule,

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PCLR}}] = \alpha.$$

The above result holds for any subsequence $\{w_n\}$ of $\{n\}$. Thus, Assumption 2 holds for the PCLR statistic. Using Proposition 16.3 of Andrews and Guggenberger (2019b), we obtain the desired result.

SA-6 Proof of Proposition 2.3 (Asymptotic local power under strong identification)

SA-6.1 Probability limits of $\hat{\mathcal{V}}_a(\theta_0)$ and $\hat{\mathcal{V}}_b(\theta_0)$ under local alternatives

Under $H_1 : \theta_n = \theta_0 + h_\theta n^{-1/2}$,

$$\tilde{u}_i(\theta_0) = (W_i' \Gamma + V_i') h_\theta n^{-1/2} - X_i'(X'X)^{-1} X'(W\Gamma + V) h_\theta n^{-1/2} + u_i - X_i'(X'X)^{-1} X'u.$$

Note that $\|Z_i^*\|^2 \|W_i\|^2$, $\|Z_i^*\|^2 \|X_i\|^2$, $\|Z_i^*\|^2 \|V_i\|^2$, $\|Z_i^*\|^2 u_i^2$, $\|Z_i^*\|^2 \|W_i\| \|X_i\|$, $\|Z_i^*\|^2 \|W_i\| \|V_i\|$, $\|Z_i^*\|^2 \|W_i\| |u_i|$, $\|Z_i^*\|^2 \|X_i\| \|V_i\|$, $\|Z_i^*\|^2 \|X_i\| |u_i|$, $\|Z_i^*\|^2 \|V_i\| |u_i|$ have finite $1 + \delta/4$ moments by the Cauchy-Schwarz inequality. The same moment bounds are also obtained when $\|Z_i^*\|^2$ in the previous quantities is replaced by $\|X_i\|^2$ and $\|Z_i^*\| \|X_i\|$. By the triangular array WLLN, CMT and the fact that $D \xrightarrow{p} 0$ in (SA-5.32),

$$\begin{aligned} \hat{\Sigma} &= n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{u}_i(\theta_0)^2 = n^{-1} \sum_{i=1}^n (Z_i^* + DX_i)(Z_i^* + DX_i)' \\ &\quad ((W_i' \Gamma + V_i') h_\theta n^{-1/2} - X_i'(X'X)^{-1} X'(W\Gamma + V) h_\theta n^{-1/2} + u_i - X_i'(X'X)^{-1} X'u)^2 \\ &= E_P[Z_i^* Z_i^{*'} u_i^2] + o_p(1) = \Sigma_n + o_p(1) = h_{5,m} + o_p(1). \end{aligned} \tag{SA-6.1}$$

Proceeding similarly to the argument in (SA-6.1),

$$\begin{aligned} \hat{C}_s &= n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{Y}_{is} \tilde{u}_i(\theta_0) \\ &= n^{-1} \sum_{i=1}^n (Z_i^* + DX_i)(Z_i^* + DX_i)' (Z_i^{*'} \Gamma_s + V_{is} + X_i'(D' \Gamma_s - (X'X)^{-1} X' V_s)) \\ &\quad ((W_i' \Gamma + V_i') h_\theta n^{-1/2} - X_i'(X'X)^{-1} X'(W\Gamma + V) h_\theta n^{-1/2} + u_i - X_i'(X'X)^{-1} X'u) \\ &= n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma_s + V_{is}) u_i + o_p(1) = E_P[Z_i^* Z_i^{*'} (Z_i^{*'} \Gamma_s + V_{is}) u_i] + o_p(1) = C_{ns} + o_p(1). \end{aligned} \tag{SA-6.2}$$

The limit in (SA-5.35) remains unchanged under the local alternatives. Thus, from (SA-6.1) and (SA-6.2), we have

$$\hat{\mathcal{V}}_a(\theta_0) = \mathcal{V}_{n,a} + o_p(1). \tag{SA-6.3}$$

Next to determine limit of $\hat{\mathcal{V}}_b(\theta_0)$, note first that

$$\varepsilon_i(\theta_0) - \begin{bmatrix} 0_{1 \times k} \\ -\Gamma' \end{bmatrix} Z_i = \begin{bmatrix} (W_i' \Gamma + V_i') h_\theta n^{-1/2} - X_i'(X'X)^{-1} X'(W\Gamma + V) h_\theta n^{-1/2} + u_i - X_i'(X'X)^{-1} X'u \\ -V_i + V'X(X'X)^{-1} X_i \end{bmatrix}$$

and

$$\hat{\varepsilon}_{in}(\theta_0) - \begin{bmatrix} 0_{1 \times k} \\ -\Gamma' \end{bmatrix} Z_i = \begin{bmatrix} n^{-1/2} h'_\theta \Gamma' Z_i + n^{-1/2} h'_\theta V' Z (Z' Z)^{-1} Z_i + u' Z (Z' Z)^{-1} Z_i \\ -V' Z (Z' Z)^{-1} Z_i \end{bmatrix}.$$

Hence,

$$\varepsilon_i(\theta_0) - \hat{\varepsilon}_{in}(\theta_0) = \begin{bmatrix} a_i \\ 0_{d \times 1} \end{bmatrix} + \begin{bmatrix} u_i - X'_i (X' X)^{-1} X' u - u' Z (Z' Z)^{-1} Z_i \\ -V_i + V' X (X' X)^{-1} X_i + V' Z (Z' Z)^{-1} Z_i \end{bmatrix}, \quad (\text{SA-6.4})$$

where

$$a_i \equiv (W'_i \Gamma + V'_i) h_\theta n^{-1/2} - X'_i (X' X)^{-1} X' (W \Gamma + V) h_\theta n^{-1/2} + n^{-1/2} h'_\theta \Gamma' Z_i + n^{-1/2} h'_\theta V' Z (Z' Z)^{-1} Z_i. \quad (\text{SA-6.5})$$

Next we show that $n^{-1} \sum_{i=1}^n a_i^2 Z_i Z'_i \xrightarrow{p} 0$. For $D_i \in \{W_i, X_i, V_i\}$ and $j + l \leq 4, j \geq 0, l \geq 2$, by the c_r inequality, and the WLLN,

$$n^{-1} \sum_{i=1}^n \|D_i\|^j \|Z_i\|^l \leq n^{-1} \sum_{i=1}^n \|D_i\|^j 2^{l-1} (\|W_i\|^l + \|W' X (X' X)^{-1}\|^l \|X_i\|^l) = O_p(1). \quad (\text{SA-6.6})$$

Therefore, using (SA-6.6) and $n^{-1} X' X = O_p(1), n^{-1} X' W = O_p(1), n^{-1} Z' Z = O_p(1), n^{-1} Z' V = O_p(1), n^{-1} X' V = O_p(1)$ which hold by the WLLN and the CMT,

$$\begin{aligned} & \|n^{-1} \sum_{i=1}^n a_i^2 Z_i Z'_i\| \\ & \leq \|h_\theta\|^2 n^{-1} \left(n^{-1} \sum_{i=1}^n \|W'_i \Gamma + V'_i\|^2 \|Z_i\|^2 + \|(X' X)^{-1} X' (W \Gamma + V)\|^2 n^{-1} \sum_{i=1}^n \|X_i\|^2 \|Z_i\|^2 \right. \\ & \quad + \|\Gamma\|^2 n^{-1} \sum_{i=1}^n \|Z_i\|^3 + \|V' Z (Z' Z)^{-1}\|^2 n^{-1} \sum_{i=1}^n \|Z_i\|^3 \\ & \quad + 2\|(X' X)^{-1} X' (W \Gamma + V)\| n^{-1} \sum_{i=1}^n \|X_i\| \|Z_i\|^2 + 2\|\Gamma\| n^{-1} \sum_{i=1}^n \|W'_i \Gamma + V'_i\| \|Z_i\|^3 \\ & \quad + 2\|V' Z (Z' Z)^{-1}\| n^{-1} \sum_{i=1}^n \|W'_i \Gamma + V'_i\| \|Z_i\|^3 \\ & \quad + 2\|(X' X)^{-1} X' (W \Gamma + V)\| \|\Gamma\| n^{-1} \sum_{i=1}^n \|X_i\| \|Z_i\|^3 \\ & \quad + 2\|V' Z (Z' Z)^{-1}\| \|(X' X)^{-1} X' (W \Gamma + V)\| n^{-1} \sum_{i=1}^n \|X_i\| \|Z_i\|^3 \\ & \quad \left. + 2\|V' Z (Z' Z)^{-1}\| \|\Gamma\| n^{-1} \sum_{i=1}^n \|Z_i\|^4 \right) \\ & \xrightarrow{p} 0. \end{aligned} \quad (\text{SA-6.7})$$

Similarly, we have

$$n^{-1} \sum_{i=1}^n a_i (u_i - X_i'(X'X)^{-1}X'u - u'Z(Z'Z)^{-1}Z_i) \otimes (Z_iZ_i') \xrightarrow{p} 0, \quad (\text{SA-6.8})$$

$$n^{-1} \sum_{i=1}^n a_i (-V_i + V'X(X'X)^{-1}X_i + V'Z(Z'Z)^{-1}Z_i) \otimes (Z_iZ_i') \xrightarrow{p} 0. \quad (\text{SA-6.9})$$

As shown in (SA-5.48)

$$\begin{aligned} & n^{-1} \sum_{i=1}^n \begin{pmatrix} u_i - X_i'(X'X)^{-1}X'u - u'Z(Z'Z)^{-1}Z_i \\ -V_i + V'X(X'X)^{-1}X_i + V'Z(Z'Z)^{-1}Z_i \end{pmatrix} \begin{pmatrix} u_i - X_i'(X'X)^{-1}X'u - u'Z(Z'Z)^{-1}Z_i \\ -V_i + V'X(X'X)^{-1}X_i + V'Z(Z'Z)^{-1}Z_i \end{pmatrix}' \\ & \otimes (Z_iZ_i') \\ & - \mathbb{E}_P \left[\begin{pmatrix} u_i^2 & -u_iV_i' \\ -u_iV_i & V_iV_i' \end{pmatrix} \otimes (Z_i^*Z_i^{*'}) \right] \xrightarrow{p} 0. \end{aligned} \quad (\text{SA-6.10})$$

By (SA-6.4), (SA-6.7), (SA-6.8), (SA-6.9), (SA-6.10) and the CMT, we have

$$\hat{\mathcal{V}}_b(\theta_0) - \mathcal{V}_{n,b} \xrightarrow{p} 0. \quad (\text{SA-6.11})$$

SA-6.2 AR and PAR statistics

Since, under $H_1 : \theta_n = \theta_0 + h_\theta n^{-1/2}$, $y - Y\theta_0 = Yh_\theta n^{-1/2} + X\gamma + u$, using $n^{-1}W'M_XW \xrightarrow{a.s.} Q_{zz}$ and $n^{-1}W'M_XV \xrightarrow{a.s.} 0$, the convergence in Lemma 16.4 of Andrews and Guggenberger (2019b) and Slutsky's lemma we obtain

$$\begin{aligned} n^{1/2}\hat{m}(\theta_0) &= n^{-1/2}W'M_X(y - Y\theta_0) \\ &= n^{-1/2}W'M_X(Yh_\theta n^{-1/2} + X\gamma + u) \\ &= n^{-1}W'M_XW\Gamma h_\theta + n^{-1}W'M_XVh_\theta + n^{-1/2}W'M_Xu \\ &\xrightarrow{d} N[Gh_\theta, \Sigma], \end{aligned} \quad (\text{SA-6.12})$$

where $\Sigma = \lim_{n \rightarrow \infty} \mathbb{E}_P[Z_i^*Z_i^{*'}u_i^2]$ and $G = \lim_{n \rightarrow \infty} \mathbb{E}_P[Z_i^*Z_i^{*'}]\Gamma = h_4$. (SA-6.1), (SA-6.12) and the CMT together yield $\text{AR} \xrightarrow{d} \chi_k^2(\eta^2)$.

Next we derive the asymptotic distribution of the permutation statistics. Since $n^{-1}W'M_XW \xrightarrow{a.s.} Q_{zz}$, $n^{-1}V'M_XW \xrightarrow{a.s.} 0$, $n^{-1}W'M_Xu \xrightarrow{a.s.} 0$, $n^{-1}V'M_XV \xrightarrow{a.s.} \Sigma^V$, $n^{-1}u'M_XV \xrightarrow{a.s.} \Sigma^{uV}$ and

$n^{-1}u'M_Xu \xrightarrow{a.s.} \sigma^2$, by the SLLN and CMT

$$\begin{aligned}
& n^{-1}\tilde{u}(\theta_0)'\tilde{u}(\theta_0) \\
&= n^{-1} \left(M_X W \Gamma h_\theta n^{-1/2} + M_X V h_\theta n^{-1/2} + M_X u \right)' \left(M_X W \Gamma h_\theta n^{-1/2} + M_X V h_\theta n^{-1/2} + M_X u \right) \\
&\xrightarrow{a.s.} \sigma^2.
\end{aligned} \tag{SA-6.13}$$

By Jensen's inequality, the WLLN, CMT, and (SA-8.22)

$$\begin{aligned}
& n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 - E_P[u_i^4] \\
&\leq 27n^{-1} \sum_{i=1}^n \|W'_i \Gamma + V'_i\|^4 \|h_\theta\|^4 n^{-2} + 27n^{-1} \sum_{i=1}^n \|X_i\|^4 \|(X'X)^{-1}X'(W\Gamma + V)\|^4 \|h_\theta\|^4 n^{-2} \\
&\quad + 27n^{-1} \sum_{i=1}^n (u_i - X'_i(X'X)^{-1}X'u)^4 - E_P[u_i^4] \\
&\xrightarrow{p} 0,
\end{aligned} \tag{SA-6.14}$$

where for the first term we used $n^{-1} \sum_{i=1}^n \|W'_i \Gamma + V'_i\|^4 \leq 8n^{-1} \sum_{i=1}^n (\|W_i\|^4 \|\Gamma\|^4 + \|V_i\|^4) = O_p(1)$ which holds by the c_r inequality and the WLLN. (SA-5.63) holds by (SA-6.13), (SA-6.14) and an argument analogous to the consistency of $\hat{\Sigma}^\pi$ in (SA-8.25). (SA-5.65) holds by (SA-6.14). (SA-5.66) holds by an argument similar to (SA-6.1). Combining these and following an argument analogous to that in (SA-5.67) then give

$$n^{-1}W'_\pi M_X \tilde{\Sigma}_u M_X W_\pi - (Q_{ww} - Q_w Q'_w) \sigma^2 \xrightarrow{p} 0 \quad \text{in } P\text{-probability.} \tag{SA-6.15}$$

Furthermore, the convergence in (SA-8.25) holds here using (SA-6.13) and (SA-6.14) in (SA-8.20) and (SA-8.24), respectively:

$$\hat{\Sigma}^\pi \xrightarrow{p^\pi} \sigma^2 Q_{zz} \quad \text{in } P\text{-probability.} \tag{SA-6.16}$$

(SA-5.70) holds by (SA-6.13). A bound analogous to (SA-5.68) is obtained by an argument similar to (SA-6.14). Thus, (SA-5.71) and (SA-5.72) hold. Therefore, by Theorem SA-2.1

$$\begin{aligned}
n^{-1/2}W'_\pi M_X(y - Y\theta_0) &= n^{-1/2}W'_\pi \tilde{u}_\pi(\theta_0) \xrightarrow{d^\pi} N[0, (Q_{ww} - Q_w Q'_w) \sigma^2] \quad P\text{-almost surely,} \\
n^{1/2}\hat{m}^\pi(\theta_0) &= n^{-1/2}Z' \tilde{u}_\pi(\theta_0) \xrightarrow{d^\pi} N[0, Q_{zz} \sigma^2] \quad P\text{-almost surely.}
\end{aligned} \tag{SA-6.17}$$

Finally, by (SA-6.15), (SA-6.16), (SA-6.17) and Slutsky's lemma, we obtain that $\text{PAR}_1 \xrightarrow{d^\pi} \chi_k^2$ and $\text{PAR}_2 \xrightarrow{d^\pi} \chi_k^2$ in P -probability.

SA-6.3 LM and PLM statistics

By definition, $q = d$ if and only if $n^{1/2}\tau_{dn} \rightarrow \infty$. Decompose $n^{1/2}H_n\hat{J}U_nT_n = n^{1/2}H_nG_nU_nT_n + n^{1/2}H_n(\hat{J} - G_n)U_nT_n$. By an analogous argument to that in (SA-8.33) (see also Lemma 16.4 and the equations (25.5)-(25.6) of Andrews and Guggenberger (2019b)),

$$n^{1/2}H_nG_nU_nT_n \xrightarrow{d} h_{3,q} \quad (\text{SA-6.18})$$

because the second term involving $B_{n,d-q}$ in (SA-8.33) drops out when $q = d$. The asymptotic distribution of $n^{1/2}(\hat{G} - G_n)$ remains invariant under $H_1 : \theta_n = \theta_0 + h_\theta n^{-1/2}$. This combined with (SA-6.1), (SA-6.2) and (SA-6.12) yields

$$n^{1/2}(\hat{J} - G_n) = O_p(1). \quad (\text{SA-6.19})$$

Therefore, $n^{1/2}H_n(\hat{J} - G_n)U_nT_n = H_n n^{1/2}(\hat{J} - G_n)U_n B_{n,q} n^{-1/2} \Upsilon_{n,q}^{-1} = o_p(1)$ because $n^{1/2}\tau_{dn} \rightarrow \infty$, and

$$n^{1/2}H_n\hat{J}U_nT_n \xrightarrow{d} h_{3,q}. \quad (\text{SA-6.20})$$

Using $\hat{\Sigma}^{-1/2}H_n^{-1} \xrightarrow{p} I_k$ which holds by (SA-6.1) and the CMT, (SA-6.1), (SA-6.20) and the CMT

$$\begin{aligned} \text{LM} &= n \hat{m}' \hat{\Sigma}^{-1/2} P_{\hat{\Sigma}^{-1/2} \hat{J}} \hat{\Sigma}^{-1/2} \hat{m} = n \hat{m}' \hat{\Sigma}^{-1/2} P_{\hat{\Sigma}^{-1/2} H_n^{-1} H_n \hat{J} U_n T_n} \hat{\Sigma}^{-1/2} \hat{m} \\ &\xrightarrow{d} \chi_d^2(\eta^2), \end{aligned} \quad (\text{SA-6.21})$$

where we used $P_{\hat{\Sigma}^{-1/2} H_n^{-1} H_n \hat{J} U_n T_n} \xrightarrow{p} P_{h_{3,q}}$, and the fact that U_n and T_n are nonsingular, and $h_{3,q} = h_{3,d}$ has full column rank d .

Next consider the PLM statistic. To see that (SA-8.15) holds under the local alternatives, note that by Jensen's inequality, the SLLN and (SA-8.15),

$$\begin{aligned} n^{-1} \sum_{i=1}^n |\tilde{u}_i(\theta_0)|^{2+\delta/2} &\leq 3^{1+\delta/2} n^{-1} \sum_{i=1}^n \|W_i' \Gamma + V_i'\|^{2+\delta/2} \|h_\theta n^{-1/2}\|^{2+\delta/2} \\ &\quad + 3^{1+\delta} n^{-1} \sum_{i=1}^n \|X_i\|^{2+\delta/2} \|(X'X)^{-1} X'(W\Gamma + V)\|^{2+\delta/2} \|h_\theta n^{-1/2}\|^{2+\delta/2} \\ &\quad + 3^{1+\delta/2} n^{-1} \sum_{i=1}^n |u_i - X_i'(X'X)^{-1} X'u|^{2+\delta/2} \\ &= O_{a.s.}(1), \end{aligned} \quad (\text{SA-6.22})$$

where $n^{-1} \sum_{i=1}^n \|W_i' \Gamma + V_i'\|^{2+\delta/2} \leq 2^{1+\delta/2} n^{-1} \sum_{i=1}^n (\|W_i\|^{2+\delta/2} \|\Gamma\|^{2+\delta/2} + \|V_i\|^{2+\delta/2}) = O_{a.s.}(1)$ by the c_r inequality and the triangular array SLLN. (SA-8.17) holds by (SA-6.22), hence (SA-8.18) holds. Also, by the SLLN and the CMT

$$n^{-1} \tilde{u}(\theta_0)' \hat{V} \tag{SA-6.23}$$

$$\begin{aligned} &= n^{-1} (M_X W \Gamma h_\theta n^{-1/2} + M_X V h_\theta n^{-1/2} + M_X u)' (I_n - P_Z - P_X) V, \\ &= n^{-1/2} h_\theta' n^{-1} V' (M_X - M_X W (W' M_X W)^{-1} W' M_X) V + n^{-1} u' (M_X - M_X W (W' M_X W)^{-1} W' M_X) V \\ &\xrightarrow{a.s.} \Sigma u V. \end{aligned} \tag{SA-6.24}$$

Hence, (SA-8.12) holds. By (SA-8.18) and Lemma SA-2.1, we then obtain (SA-8.6). Note that (SA-8.28) holds because (SA-8.29) holds by (SA-6.14). Also, (SA-8.26) holds by (SA-6.24). This verifies (SA-8.30). Combining the latter with (SA-8.6) and (SA-6.16), we obtain

$$n^{1/2}(\hat{J}^\pi - \hat{G}) = O_{p^\pi}(1) \quad \text{in } P\text{-probability.} \tag{SA-6.25}$$

Thus, following the proof of Lemma SA-5.1, noting that the term involving $B_{n,d-q}$ in (SA-8.33) evacuates when $q = d$, and using (SA-6.19) and (SA-6.25), we obtain

$$\begin{aligned} n^{1/2} H_n \hat{J}^\pi U_n T_n &= n^{1/2} H_n G_n U_n T_n + n^{1/2} H_n (\hat{G} - G_n) U_n T_n + n^{1/2} H_n (\hat{J}^\pi - \hat{G}) U_n T_n \\ &\xrightarrow{p^\pi} \Delta_h^\pi = h_{3,d} \quad \text{in } P\text{-probability,} \end{aligned}$$

where $\Delta_h^\pi = h_{3,q} = h_{3,d} = \Delta_h$ has full column rank. Since $(\Sigma^\pi)^{-1/2} \Sigma^{1/2} = (\sigma^2 Q_{zz})^{-1/2} \Sigma^{1/2}$ is nonsingular, $P^\pi[\text{rank}((\Sigma^\pi)^{-1/2} \Sigma^{1/2} \Delta_h^\pi) = d] = 1$ with P -probability one. Then,

$$\begin{aligned} \text{PLM} &= n \hat{m}^{\pi'} \hat{\Sigma}^{\pi-1/2} P_{\hat{\Sigma}^{\pi-1/2} \hat{J}^\pi} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi \\ &= n \hat{m}^{\pi'} \hat{\Sigma}^{\pi-1/2} P_{(\hat{\Sigma}^{\pi-1/2} \Sigma_n^{1/2}) n^{1/2} \Sigma_n^{-1/2} \hat{J}^\pi U_n T_n} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi \\ &\xrightarrow{d^\pi} m_h^{\pi'} \Sigma^{\pi-1/2} P_{(\Sigma^\pi)^{-1/2} \Sigma^{1/2} \Delta_h^\pi} \Sigma^{\pi-1/2} m_h^\pi \\ &\sim \chi_d^2 \quad \text{in } P\text{-probability,} \end{aligned} \tag{SA-6.26}$$

where the second equality holds because Σ_n and T_n are nonsingular, the convergence follows from the CMT, the full column rank property verified above, and the last line uses (SA-6.17) and the fact that $P_{(\Sigma^\pi)^{-1/2} \Sigma^{1/2} \Delta_h^\pi}$ is idempotent with rank d .

SA-6.4 CLR and PCLR statistics

The asymptotic distributions of the CLR and PCLR statistics under the sequence of local alternatives and strong or semi-strong identification with $n^{1/2}\tau_{dn} \rightarrow \infty$, are derived proceeding similarly to Proposition SA-5.3 and Lemma 28.1 of Andrews and Guggenberger (2019b). By definition, $q = d$ if and only if $n^{1/2}\tau_{dn} \rightarrow \infty$ in which case $Q_2^+(\kappa)$ defined in (SA-8.52) is scalar. From (SA-8.60) which also holds under the local alternatives because $\hat{J} - G_n = O_p(n^{-1/2})$ as verified in (SA-6.19), all the remaining quantities $n^{1/2}\hat{\Sigma}^{\pi-1/2}\hat{m}^\pi, \hat{\Sigma}, \hat{\mathcal{V}} \in \{\hat{\mathcal{V}}_a(\theta_0), \hat{\mathcal{V}}_b(\theta_0)\}$ have the same limits as under the null as shown in (SA-6.1), (SA-6.3), (SA-6.11), (SA-6.16), (SA-6.17), and

$$Q_2^+(\hat{\kappa}_j^+) = M_{d+1-q}^+ - \hat{\kappa}_j^+(I_{d+1-q} + o_p(1)), \quad (\text{SA-6.27})$$

where

$$M_{d+1-q}^+ \equiv nB_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}H_n'h_{3,k-q}h'_{3,k-q}H_n\hat{J}^+U_n^+B_{n,d+1-q}^+ + o_p(1). \quad (\text{SA-6.28})$$

Since $|Q_2^+(\hat{\kappa}_{d+1}^+)| = |M_{d+1-q}^+ - \hat{\kappa}_{d+1}^+(I_{d+1-q} + o_p(1))| = 0$ with P^π -probability approaching 1, in P -probability, we obtain

$$\begin{aligned} \hat{\kappa}_{d+1}^+ &= nB_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}H_n'h_{3,k-q}h'_{3,k-q}H_n\hat{J}^+U_n^+B_{n,d+1-q}^+(1 + o_p(1)) + o_p(1), \\ &= n\hat{m}^{\pi'}\hat{\Sigma}^{\pi-1/2}h_{3,k-q}h'_{3,k-q}\hat{\Sigma}^{\pi-1/2}\hat{m}^\pi + o_p(1) \quad \text{in } P\text{-probability.} \end{aligned} \quad (\text{SA-6.29})$$

When $q = d$ and under the local alternatives $H_1 : \theta_n = \theta_0 + h_\theta n^{-1/2}$, as shown in (SA-6.20), $n^{1/2}\hat{H}\hat{J}^\pi U_n B_n S_n \xrightarrow{d^\pi} \Delta_h^\pi = h_{3,q}$ as the term $\Delta_{h,d-q}^\pi$ evacuates. Then,

$$\begin{aligned} \text{PCLR} &= n\hat{m}^{\pi'}\hat{\Sigma}^{\pi-1}\hat{m}^\pi - \hat{\kappa}_{d+1}^+ \quad (\text{SA-6.30}) \\ &= n\hat{m}^{\pi'}\hat{\Sigma}^{\pi-1/2}(I_k - h_{3,k-q}h'_{3,k-q})\hat{\Sigma}^{\pi-1/2}\hat{m}^\pi + o_p(1) \\ &= n\hat{m}^{\pi'}\hat{\Sigma}^{\pi-1/2}h_{3,q}h'_{3,q}\hat{\Sigma}^{\pi-1/2}\hat{m}^\pi + o_p(1) \\ &= n\hat{m}^{\pi'}\hat{\Sigma}^{\pi-1/2}P_{h_{3,q}}\hat{\Sigma}^{\pi-1/2}\hat{m}^\pi + o_p(1) \\ &= n\hat{m}^{\pi'}\hat{\Sigma}^{\pi-1/2}P_{(\hat{\Sigma}^{\pi-1/2}\Sigma_n^{1/2})n^{1/2}\Sigma_n^{-1/2}\hat{J}^\pi U_n T_n}\hat{\Sigma}^{\pi-1/2}\hat{m}^\pi + o_p(1) \\ &\xrightarrow{d^\pi} \chi_d^2 \quad \text{in } P\text{-probability,} \end{aligned}$$

where the second equality uses (SA-6.29), the third and the fourth equalities use $P_{n^{1/2}\hat{H}\hat{J}^\pi U_n B_n S_n} = P_{h_{3,q}} + o_p(1) = h_{3,q}(h'_{3,q}h_{3,q})^{-1}h'_{3,q} + o_p(1) = h_{3,q}h'_{3,q} + o_p(1)$ and (SA-6.12) since $h'_{3,q}h_{3,q} = I_d$, and the convergence follows from (SA-6.26).

Recall that by (SA-6.18) (see also Lemma SA-5.1, Lemma 10.3 of Andrews and Guggenberger (2017b) and Lemma 16.4 of Andrews and Guggenberger (2019b)), $n^{1/2}\hat{H}\hat{J}U_nB_nS_n \xrightarrow{d} h_{3,q}$. By replacing $\hat{\Sigma}^{\pi-1/2}\hat{m}^\pi$ in (SA-6.29) and (SA-6.30) with $\hat{\Sigma}^{-1/2}\hat{m}$ and using (SA-6.12), the asymptotic distribution of the CLR statistic is obtained similarly:

$$\begin{aligned}\text{CLR} &= n\hat{m}'\hat{\Sigma}^{-1/2}h_{3,q}h'_{3,q}\hat{\Sigma}^{-1/2}\hat{m} + o_p(1) \\ &= n\hat{m}'\hat{\Sigma}^{-1/2}P_{\hat{\Sigma}^{-1/2}H_n^{-1}H_n\hat{J}U_nT_n}\hat{\Sigma}^{-1/2}\hat{m} \\ &= n\hat{m}'\hat{\Sigma}^{-1/2}P_{\hat{\Sigma}^{-1/2}\hat{J}}\hat{\Sigma}^{-1/2}\hat{m} + o_p(1) \\ &\xrightarrow{d} \chi_d^2(\eta^2),\end{aligned}$$

where the second and the third equalities use

$$P_{\hat{\Sigma}^{-1/2}\hat{J}} = P_{n^{1/2}\hat{H}\hat{J}U_nB_nS_n} = P_{h_{3,q}} + o_p(1) = h_{3,q}(h'_{3,q}h_{3,q})^{-1}h'_{3,q} + o_p(1) = h_{3,q}h'_{3,q} + o_p(1),$$

and the convergence uses (SA-6.21).

SA-7 Proof of Proposition 2.4 (Many IVs)

As before, we use the shorthand notations $u_i = u_i(\theta_0)$, $\tilde{u}_i = \tilde{u}_i(\theta_0)$, $\tilde{u} = \tilde{u}(\theta_0)$, $\hat{\Sigma} = \hat{\Sigma}(\theta_0)$ and $\hat{\Sigma}^\pi = \hat{\Sigma}^\pi(\theta_0) = n^{-1}\sum_{i=1}^n Z_i Z_i' \tilde{u}_{\pi(i)}(\theta_0)^2$. In addition, we let $\tilde{\Sigma} \equiv n^{-1}Z'Z\tilde{\sigma}^2$, where $\tilde{\sigma}^2 \equiv n^{-1}\tilde{u}'\tilde{u}$. The proof consists of three steps. In Step 1 and 2, we derive the asymptotic distribution of the PAR and AR statistics, respectively. Step 3 completes the proof. Since Step 3 is analogous to the proofs of the PAR tests with k fixed, it will be brief.

Step 1: Asymptotic distribution of the PAR statistic via double-index permutation CLT.

As in (2.48), we may rewrite

$$\begin{aligned}\frac{\text{PAR} - k}{(2k)^{1/2}} &= \frac{n^{-1/2}\tilde{u}'_\pi Z\tilde{\Sigma}^{-1}n^{-1/2}Z'\tilde{u}_\pi - k}{(2k)^{1/2}} + \frac{n^{-1/2}\tilde{u}'_\pi Z(\hat{\Sigma}^{\pi-1} - \tilde{\Sigma}^{-1})n^{-1/2}Z'\tilde{u}_\pi}{(2k)^{1/2}} \\ &= \frac{\sum_{i=1}^n P_{ii}\tilde{u}_{\pi(i)}^2\tilde{\sigma}^{-2} - k}{(2k)^{1/2}} + \frac{\sum_{i \neq j} P_{ij}\tilde{u}_{\pi(i)}\tilde{u}_{\pi(j)}\tilde{\sigma}^{-2}}{(2k)^{1/2}} + \frac{n^{-1/2}\tilde{u}'_\pi Z(\hat{\Sigma}^{\pi-1} - \tilde{\Sigma}^{-1})n^{-1/2}Z'\tilde{u}_\pi}{(2k)^{1/2}}.\end{aligned}\tag{SA-7.1}$$

Below, the asymptotic distribution of the PAR statistic in (SA-7.1) is derived in three substeps. The first substep shows that the first summand of (SA-7.1) is negligible asymptotically. The second

substep establishes the asymptotic normality of the second summand using Lemma SA-2.4. The last substep shows that the third term also vanishes asymptotically.

Substep 1: $(2k)^{-1/2} \left(\sum_{i=1}^n P_{ii} \tilde{u}_{\pi(i)}^2 \tilde{\sigma}^{-2} - k \right) \xrightarrow{p^\pi} 0$ P -almost surely.

From Lemma S.3.4 of DiCiccio and Romano (2017), the fact that $\sum_{i=1}^n P_{ii} = \text{tr} \left(\sum_{i=1}^n Z'_i (Z'Z)^{-1} Z_i \right) = \text{tr} \left((Z'Z)^{-1} Z'Z \right) = k$ and the definition of $\tilde{\sigma}^2$

$$\mathbb{E}_{P^\pi} \left[\sum_{i=1}^n P_{ii} \tilde{u}_{\pi(i)}^2 \tilde{\sigma}^{-2} | y, Y, X, W \right] = \left(\sum_{i=1}^n P_{ii} \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i^2 \tilde{\sigma}^{-2} \right) = k, \quad (\text{SA-7.2})$$

and

$$\begin{aligned} \text{Var}_{P^\pi} \left[(2k)^{-1/2} \sum_{i=1}^n P_{ii} \tilde{u}_{\pi(i)}^2 \tilde{\sigma}^{-2} | y, Y, X, W \right] &= (2k)^{-1} \frac{1}{n-1} \left(\sum_{i=1}^n P_{ii}^2 - n \left(n^{-1} \sum_{i=1}^n P_{ii} \right)^2 \right) \\ &\quad \left(\sum_{i=1}^n \tilde{u}_i^4 \tilde{\sigma}^{-4} - n \left(n^{-1} \sum_{i=1}^n \tilde{u}_i^2 \tilde{\sigma}^{-2} \right)^2 \right) \\ &\leq (2k)^{-1} \left(\sum_{i=1}^n P_{ii}^2 \right) \frac{1}{n-1} \left(\sum_{i=1}^n \tilde{u}_i^4 \tilde{\sigma}^{-4} \right) \\ &\xrightarrow{a.s.} 0. \end{aligned} \quad (\text{SA-7.3})$$

where we used Assumption 3(c), $(n-1)^{-1} \sum_{i=1}^n \tilde{u}_i^4 = O_{a.s.}(1)$ and $\tilde{\sigma}^{-4} = O_{a.s.}(1)$ which follow from arguments similar to (SA-8.22) and the SLLN. Therefore, by Chebyshev's inequality, (SA-7.2) and (SA-7.3)

$$(2k)^{-1/2} \left(\sum_{i=1}^n P_{ii} \tilde{u}_{\pi(i)}^2 \tilde{\sigma}^{-2} - k \right) \xrightarrow{p^\pi} 0 \quad P\text{-almost surely.} \quad (\text{SA-7.4})$$

Substep 2: Asymptotic distribution of $(2k)^{-1/2} \sum_{i \neq j} P_{ij} \tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} \tilde{\sigma}^{-2}$.

We will apply Lemma SA-2.4 to determine the asymptotic distribution of

$$(2k)^{-1/2} \sum_{i \neq j} P_{ij} \tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} \tilde{\sigma}^{-2}. \quad (\text{SA-7.5})$$

To this end, we first center the summands of (SA-7.5). Since $\tilde{u}'\iota = 0$,

$$\begin{aligned} \mathbb{E}_{P^\pi} \left[\sum_{i \neq j} P_{ij} \tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} \tilde{\sigma}^{-2} \middle| y, Y, W, X \right] &= \left(\sum_{i \neq j} P_{ij} \right) \frac{1}{n(n-1)} \sum_{i \neq j} \tilde{u}_i \tilde{u}_j \tilde{\sigma}^{-2} \\ &= \left(\sum_{i \neq j} P_{ij} \right) \frac{1}{n(n-1)} \sum_{i=1}^n \left(\tilde{u}_i \left(\sum_{j=1}^n \tilde{u}_j - \tilde{u}_i \right) \right) \tilde{\sigma}^{-2} \\ &= -\frac{1}{n-1} \sum_{i \neq j} P_{ij}. \end{aligned} \quad (\text{SA-7.6})$$

After centering by (SA-7.6),

$$\sum_{i \neq j} P_{ij} \tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} \tilde{\sigma}^{-2} + \frac{1}{n-1} \sum_{i \neq j} P_{ij} = \sum_{i \neq j} P_{ij} \tilde{\sigma}^{-2} \left(\tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} + \frac{\tilde{\sigma}^2}{n-1} \right). \quad (\text{SA-7.7})$$

Put $a_{ij} = P_{ij}$ and $b_{ij} = \tilde{u}_i \tilde{u}_j + \frac{\tilde{\sigma}^2}{n-1}$ if $i \neq j$, and $a_{ij} = b_{ij} = 0$ if $i = j$ in Lemma SA-2.4. Conditions (SA-2.3) and (SA-2.4) hold by Assumptions 3(a) and (b). Condition (SA-2.5) holds because of the centering done above. Next, we verify Condition (SA-2.6). By some algebra,

$$b_{i+} = \sum_{j=1}^n b_{ij} = \sum_{j=1, j \neq i}^n \left(\tilde{u}_i \tilde{u}_j + \frac{\tilde{\sigma}^2}{n-1} \right) = \tilde{u}_i \left(\sum_{j=1}^n \tilde{u}_j - \tilde{u}_i \right) + \tilde{\sigma}^2 = -\tilde{u}_i^2 + \tilde{\sigma}^2, \quad (\text{SA-7.8})$$

hence

$$\sum_{i=1}^n b_{i+}^2 = \sum_{i=1}^n \tilde{u}_i^4 - n\tilde{\sigma}^4. \quad (\text{SA-7.9})$$

On noting that $\sum_{i \neq j} \tilde{u}_i^2 \tilde{u}_j^2 = (\sum_{i=1}^n \tilde{u}_i^2)^2 - \sum_{i=1}^n \tilde{u}_i^4 = n^2 \tilde{\sigma}^4 - \sum_{i=1}^n \tilde{u}_i^4$ and $\sum_{i \neq j} \tilde{u}_i \tilde{u}_j = -\sum_{i=1}^n \tilde{u}_i^2 = -n\tilde{\sigma}^2$, we have

$$\begin{aligned} \sum_{i \neq j} b_{ij}^2 &= \sum_{i \neq j} \left(\tilde{u}_i \tilde{u}_j + \frac{\tilde{\sigma}^2}{n-1} \right)^2 \\ &= \sum_{i \neq j} \left(\tilde{u}_i^2 \tilde{u}_j^2 + 2 \frac{\tilde{\sigma}^2}{n-1} \tilde{u}_i \tilde{u}_j + \frac{\tilde{\sigma}^4}{(n-1)^2} \right) \\ &= n^2 \tilde{\sigma}^4 - \sum_{i=1}^n \tilde{u}_i^4 - \frac{n}{n-1} \tilde{\sigma}^4 \end{aligned} \quad (\text{SA-7.10})$$

Since $n^{-1} \sum_{i=1}^n \tilde{u}_i^4 = O_{a.s.}(1)$ and $\tilde{\sigma}^2 = O_{a.s.}(1)$ in (SA-7.9) and (SA-7.10),

$$\frac{\sum_{i=1}^n b_{i+}^2}{\sum_{i,j} b_{ij}^2} \xrightarrow{a.s.} 0. \quad (\text{SA-7.11})$$

Condition (SA-2.6) follows. Moreover,

$$\begin{aligned}\sum_{i,j} a_{ij}^2 &= \sum_{i \neq j} P_{ij}^2 = \sum_{i,j} Z'_i (Z'Z)^{-1} Z_j Z'_j (Z'Z)^{-1} Z_i - \sum_i P_{ii}^2 \\ &= \text{tr} \left(\sum_i (Z'Z)^{-1} Z_i Z'_i \right) - \sum_i P_{ii}^2 = k - \sum_i P_{ii}^2.\end{aligned}\tag{SA-7.12}$$

From (SA-7.7), (SA-7.10), (SA-7.12) and Lemma SA-2.4, we then obtain

$$T^\pi \equiv \frac{\sum_{i \neq j} P_{ij} \left(\tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} + \frac{\tilde{\sigma}^2}{n-1} \right)}{\left[2(k - \sum_i P_{ii}^2)(\tilde{\sigma}^4 - n^{-2} \sum_{i=1}^n \tilde{u}_i^4 - n^{-2} \frac{n}{n-1} \tilde{\sigma}^4) \right]^{1/2}} \xrightarrow{d^\pi} N[0, 1] \quad P\text{-almost surely.}\tag{SA-7.13}$$

Since $k^{-1} \sum_i P_{ii}^2 \xrightarrow{a.s.} 0$ by Assumption 3(c), and $\tilde{\sigma}^4 - \sigma^4 - n^{-2} \sum_{i=1}^n \tilde{u}_i^4 - n^{-2} \frac{n}{n-1} \tilde{\sigma}^4 \xrightarrow{a.s.} 0$ which follows from $\tilde{\sigma}^2 \xrightarrow{a.s.} \sigma^2 = \mathbb{E}_P[u_1^2]$ and the CMT, for the denominator of T^π in (SA-7.13) we have

$$(2k)^{-1/2} \left[2k(1 - k^{-1} \sum_i P_{ii}^2) \left(\tilde{\sigma}^4 - n^{-2} \sum_{i=1}^n \tilde{u}_i^4 - n^{-2} \frac{n}{n-1} \tilde{\sigma}^4 \right) \right]^{1/2} \xrightarrow{a.s.} \sigma^2,\tag{SA-7.14}$$

Thus,

$$\begin{aligned}& \frac{\sum_{i \neq j} P_{ij} \tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} \tilde{\sigma}^{-2}}{(2k)^{1/2}} \\ &= (2k)^{-1/2} \sum_{i \neq j} P_{ij} \left(\tilde{u}_{\pi(i)} \tilde{u}_{\pi(j)} + \frac{\tilde{\sigma}^2}{n-1} \right) \tilde{\sigma}^{-2} - (2k)^{-1/2} \frac{1}{n-1} \sum_{i \neq j} P_{ij} \\ &= T^\pi \frac{\left[2k(1 - k^{-1} \sum_i P_{ii}^2)(\tilde{\sigma}^4 - n^{-2} \sum_{i=1}^n \tilde{u}_i^4 - n^{-2} \frac{n}{n-1} \tilde{\sigma}^4) \right]^{1/2} \tilde{\sigma}^{-2}}{(2k)^{1/2}} + (2k)^{-1/2} \frac{k}{n-1} \\ &\xrightarrow{d^\pi} N[0, 1] \quad P\text{-almost surely,}\end{aligned}\tag{SA-7.15}$$

where the second equality uses $\sum_{i \neq j} P_{ij} = \sum_{i=1}^n (Z'_i (Z'Z)^{-1} Z'_i - P_{ii}) = -k$ which follows from $\sum_{i=1}^n P_{ii} = \text{tr}((Z'Z)^{-1} Z'Z) = k$ and $Z'_i = 0$, and the convergence uses (SA-7.13), (SA-7.14), $k^{1/2}/(n-1) \rightarrow 0$ and Slutsky's lemma.

Substep 3: $(2k)^{-1/2} n^{-1/2} \tilde{u}'_\pi Z (\hat{\Sigma}^{\pi-1} - \tilde{\Sigma}^{-1}) n^{-1/2} Z' \tilde{u}_\pi \xrightarrow{p^\pi} 0$ in P -probability.

First, we will establish that

$$\|\hat{\Sigma}^\pi - \tilde{\Sigma}\| = o_{p^\pi}(k^{-1/2}) \text{ in } P\text{-probability.}\tag{SA-7.16}$$

Let $a_i \equiv Z_i Z_{ij}$, $j = 1, \dots, k$ and $b_i = \tilde{u}_i^2 - \tilde{\sigma}^2$ in Lemma SA-2.5. The j -th column of $\hat{\Sigma}^\pi - \tilde{\Sigma}$ can be written as $n^{-1} \sum_{i=1}^n a_i b_{\pi(i)}$. Since $\sum_{i=1}^n b_i = 0$, Lemma SA-2.5 gives

$$\mathbb{E}_{P^\pi} \left[\left\| n^{-1} \sum_{i=1}^n Z_i Z_{ij} (\tilde{u}_{\pi(i)}^2 - \tilde{\sigma}^2) \right\|^2 \middle| y, Y, X, W \right] \leq \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n Z'_i Z_i Z_{ij}^2 \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i^4 \right). \quad (\text{SA-7.17})$$

Using $\|\hat{\Sigma}^\pi - \tilde{\Sigma}\|^2 = \sum_{j=1}^k \left\| n^{-1} \sum_{i=1}^n Z_i Z_{ij} (\tilde{u}_{\pi(i)}^2 - \tilde{\sigma}^2) \right\|^2$ and (SA-7.17),

$$\mathbb{E}_{P^\pi} [\|\hat{\Sigma}^\pi - \tilde{\Sigma}\|^2 | y, Y, X, W] \leq \frac{1}{n-1} \left(n^{-1} \sum_{j=1}^k \sum_{i=1}^n Z'_i Z_i Z_{ij}^2 \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i^4 \right). \quad (\text{SA-7.18})$$

Since, $\mathbb{E}_P[Z_{ij}^4] = \mathbb{E}_P[(W_{ij} - n^{-1} \sum_{i=1}^n W_{ij})^4] \leq 8\mathbb{E}_P[W_{ij}^4] + 8n^{-1} \sum_{i=1}^n \mathbb{E}_P[W_{ij}^4] = O(1)$, we have $\mathbb{E}_P[\sum_{j=1}^k Z'_i Z_i Z_{ij}^2] = \mathbb{E}_P[Z'_i Z_i (\sum_{j=1}^k Z_{ij}^2)] = \mathbb{E}_P[(\sum_{j=1}^k Z_{ij}^2)^2] \leq k^2 \mathbb{E}_P[k^{-1} \sum_{j=1}^k Z_{ij}^4] = O(k^2)$ and $n^{-1} \sum_{j=1}^k \sum_{i=1}^n \mathbb{E}_P[Z'_i Z_i Z_{ij}^2] = O(n^{-1} k^2)$. By Markov's inequality and $k^3/n \rightarrow 0$, we have

$$kn^{-1} \sum_{j=1}^k \sum_{i=1}^n Z'_i Z_i Z_{ij}^2 = O_p(n^{-1} k^3) = o_p(1). \quad (\text{SA-7.19})$$

Using Markov's inequality once again along with (SA-7.18), (SA-7.19) and the fact that $n^{-1} \sum_{i=1}^n \tilde{u}_i^4 = O_p(1)$, we obtain

$$\begin{aligned} P^\pi \left[k \|\hat{\Sigma}^\pi - \tilde{\Sigma}\|^2 > \epsilon \right] &\leq \epsilon^{-1} k \mathbb{E}_{P^\pi} [\|\hat{\Sigma}^\pi - \tilde{\Sigma}\|^2 | y, Y, X, W] \\ &= o_p(1). \end{aligned} \quad (\text{SA-7.20})$$

Therefore, (SA-7.16) holds.

From Substeps 1-2, we have

$$\|\tilde{\Sigma}^{-1/2} n^{-1/2} Z' \tilde{u}_\pi\|^2 = O_{p^\pi}(k) \text{ in } P\text{-probability.} \quad (\text{SA-7.21})$$

By Lemma SA-7.25 below, we have $\|\tilde{\Sigma} - \sigma^2 \mathbb{E}_P[Z_i^* Z_i^{*'}]\| = o_p(k^{-1/2})$ where $Z_i^* = W_i - \mathbb{E}_P[W_i]$, and by Weyl's inequality (see e.g. Vershynin (2018), Theorem 4.5.3), $|\lambda_{\min}(\tilde{\Sigma}) - \lambda_{\min}(\sigma^2 \mathbb{E}_P[Z_i^* Z_i^{*'}])| \leq \|\tilde{\Sigma} - \sigma^2 \mathbb{E}_P[Z_i^* Z_i^{*'}]\|$, hence $\lambda_{\min}(\tilde{\Sigma}) = O_p(1)$. Similarly, since $|\lambda_{\min}(\hat{\Sigma}^\pi) - \lambda_{\min}(\tilde{\Sigma})| \leq \|\hat{\Sigma}^\pi - \tilde{\Sigma}\| = o_{p^\pi}(k^{1/2})$ in P -probability by virtue of (SA-7.16), we obtain $\lambda_{\min}(\hat{\Sigma}^\pi) = O_{p^\pi}(1)$ in P -probability. Andrews and Stock (2007b) (the equation (38) therein) establish the following inequality: for a $k \times 1$ vector A_n and $k \times k$ symmetric positive definite matrices C_n and D_n :

$$|A_n'(C_n^{-2} - D_n^{-2})A_n| \leq \|D_n^2 - C_n^2\| \lambda_{\min}^{-1}(C_n^2) \|C_n^{-1} A_n\|^2 + \|D_n^2 - C_n^2\|^2 \|C_n^{-1} A_n\|^2 \lambda_{\max}^2(C_n^{-1}) \lambda_{\min}^{-2}(D_n). \quad (\text{SA-7.22})$$

Let $A_n = n^{-1/2}Z'\tilde{u}_\pi$, $C_n = \tilde{\Sigma}^{1/2}$ and $D_n = \hat{\Sigma}^{\pi 1/2}$ in the inequality above. Then, using (SA-7.16), (SA-7.21), and the facts that $\lambda_{\min}(\tilde{\Sigma}) = O_p(1)$ and $\lambda_{\min}(\hat{\Sigma}^\pi) = O_{p^\pi}(1)$ in P -probability in (SA-7.22)

$$\begin{aligned}
|n^{-1/2}\tilde{u}_\pi'Z(\hat{\Sigma}^{\pi-1} - \tilde{\Sigma}^{-1})n^{-1/2}Z'\tilde{u}_\pi| &\leq \|\hat{\Sigma}^\pi - \tilde{\Sigma}\|\lambda_{\min}^{-1}(\tilde{\Sigma})\|\tilde{\Sigma}^{-1/2}n^{-1/2}Z'\tilde{u}_\pi\|^2 \\
&+ \|\hat{\Sigma}^\pi - \tilde{\Sigma}\|^2\|\tilde{\Sigma}^{-1/2}n^{-1/2}Z'\tilde{u}_\pi\|^2(1/\lambda_{\min}^2(\tilde{\Sigma}^{1/2}))\lambda_{\min}^{-2}(\hat{\Sigma}^{\pi 1/2}) \\
&= o_{p^\pi}(k^{-1/2})O_p(1)O_{p^\pi}(k) + o_{p^\pi}(k^{-1})O_{p^\pi}(k)O_p(1)O_{p^\pi}(1) \\
&= o_{p^\pi}(k^{1/2}) \text{ in } P\text{-probability,}
\end{aligned} \tag{SA-7.23}$$

as required. Finally, collecting (SA-7.4), (SA-7.15) and (SA-7.23) in (SA-7.1), and invoking Slutsky's lemma, we obtain

$$\frac{\text{PAR} - k}{(2k)^{1/2}} \xrightarrow{d^\pi} N[0, 1] \quad \text{in } P\text{-probability.} \tag{SA-7.24}$$

Step 2: Asymptotic distribution of the AR statistic.

The derivation of the asymptotic distribution of the AR statistic consists of two substeps. In the first substep, we show the consistency of the covariance matrix estimator, and in the second substep, we derive the limiting distribution of the AR statistic using Lemma SA-2.6.

Substep 1: Consistency of the covariance matrix estimator.

We will show under the conditions of Proposition 2.4 that

$$\|\hat{\Sigma} - \Sigma\| = o_p(k^{-1/2}). \tag{SA-7.25}$$

Using $\bar{W} = n^{-1}\sum_{i=1}^n W_i$, we may rewrite

$$\begin{aligned}
\hat{\Sigma} &= n^{-1}\sum_{i=1}^n Z_i Z_i' \tilde{u}_i^2 = n^{-1}\sum_{i=1}^n (Z_i^* - (\bar{W} - E_P[W_i]))(Z_i^* - (\bar{W} - E_P[W_i]))' \tilde{u}_i^2 \\
&= n^{-1}\sum_{i=1}^n Z_i^* Z_i^{*'} \tilde{u}_i^2 - n^{-1}\sum_{i=1}^n (\bar{W} - E_P[W_i]) Z_i^{*'} \tilde{u}_i^2 - n^{-1}\sum_{i=1}^n Z_i^* (\bar{W} - E_P[W_i])' \tilde{u}_i^2 \\
&\quad + n^{-1}\sum_{i=1}^n (\bar{W} - E_P[W_i])(\bar{W} - E_P[W_i])' \tilde{u}_i^2.
\end{aligned} \tag{SA-7.26}$$

In what follows, we derive the limits of each summands on the RHS of (SA-7.26).

The limit of $n^{-1}\sum_{i=1}^n Z_i^* Z_i^{*'} \tilde{u}_i^2$.

By expansion,

$$n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} \tilde{u}_i^2 = n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} u_i^2 - 2n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} u_i \bar{u} + 2n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} \bar{u}^2, \quad (\text{SA-7.27})$$

where $\bar{u} = n^{-1} \sum_{i=1}^n u_i$. For the first summand of (SA-7.27), by Markov's inequality and the i.i.d. assumption, for any $\epsilon > 0$

$$\begin{aligned} P \left[k \left\| n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} u_i^2 - \mathbb{E}_P[Z_i^* Z_i^{*'} u_i^2]) \right\|^2 > \epsilon \right] \\ \leq \frac{k}{\epsilon} \mathbb{E}_P \left[\text{tr} \left\{ \left(n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} u_i^2 - \mathbb{E}_P[Z_i^* Z_i^{*'} u_i^2]) \right) \left(n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} u_i^2 - \mathbb{E}_P[Z_i^* Z_i^{*'} u_i^2]) \right)' \right\} \right] \\ = \frac{k}{n\epsilon} \text{tr} \left\{ \mathbb{E}_P \left[(Z_i^* Z_i^{*'} u_i^2 - \mathbb{E}_P[Z_i^* Z_i^{*'} u_i^2]) (Z_i^* Z_i^{*'} u_i^2 - \mathbb{E}_P[Z_i^* Z_i^{*'} u_i^2])' \right] \right\} \\ = \frac{k}{n\epsilon} (\text{tr} \{ \mathbb{E}_P [Z_i^* Z_i^{*'} u_i^2 Z_i^* Z_i^{*'} u_i^2] \} - \text{tr} \{ \mathbb{E}_P [Z_i^* Z_i^{*'} u_i^2] \mathbb{E}_P [Z_i^* Z_i^{*'} u_i^2] \}). \end{aligned} \quad (\text{SA-7.28})$$

For the first term of (SA-7.28), by the convexity of $x \mapsto x^2$ and Assumption 3(d)

$$\begin{aligned} n^{-1} k \text{tr} \{ \mathbb{E}_P [Z_i^* Z_i^{*'} u_i^2 Z_i^* Z_i^{*'} u_i^2] \} &= n^{-1} k \mathbb{E}_P [(Z_i^{*'} Z_i^*)^2 u_i^4] \\ &= n^{-1} k^3 \mathbb{E}_P \left[\left(k^{-1} \sum_{j=1}^k Z_{ij}^{*2} \right)^2 u_i^4 \right] \\ &\leq n^{-1} k^3 \left(k^{-1} \sum_{j=1}^k \mathbb{E}_P [Z_{ij}^{*4} u_i^4] \right) \\ &= o(1). \end{aligned} \quad (\text{SA-7.29})$$

For the second term of (SA-7.28), we have

$$\begin{aligned} n^{-1} k \text{tr} \{ \mathbb{E}_P [Z_i^* Z_i^{*'} u_i^2] \mathbb{E}_P [Z_i^* Z_i^{*'} u_i^2] \} &= n^{-1} k \left\| \mathbb{E}_P [Z_i^* Z_i^{*'} u_i^2] \right\|^2 \\ &= n^{-1} k \sum_{j=1}^k \mathbb{E}_P [Z_{ij}^{*'} Z_{ij}^* u_i^2] \mathbb{E}_P [Z_{ij}^* Z_{ij}^{*'} u_i^2] \\ &= n^{-1} k \sum_{j=1}^k \sum_{\ell=1}^k (\mathbb{E}_P [Z_{i\ell}^* Z_{ij}^* u_i^2])^2 \\ &= O(n^{-1} k^3) \\ &= o(1), \end{aligned} \quad (\text{SA-7.30})$$

where we used $\mathbb{E}_P [Z_{i\ell}^* Z_{ij}^* u_i^2] \leq (\mathbb{E}_P [Z_{i\ell}^{*4}])^{1/4} (\mathbb{E}_P [Z_{ij}^{*4}])^{1/4} (\mathbb{E}_P [u_i^4])^{1/2} = O(1)$ which holds by Cauchy-Schwarz inequality and Assumption 3(d). Therefore, it follows from (SA-7.28), (SA-7.29)

and (SA-7.30) that

$$\left\| n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} u_i^2 - \mathbb{E}_P[Z_i^* Z_i^{*'} u_i^2]) \right\| = o_p(k^{-1/2}). \quad (\text{SA-7.31})$$

Next we verify for the second summand of (SA-7.27) that

$$\left\| n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} u_i \bar{u} \right\| = O_p(kn^{-1/2}) = o_p(k^{-1/2}). \quad (\text{SA-7.32})$$

Once again using Markov's inequality and arguments analogous to (SA-7.28), for any $\epsilon > 0$

$$\begin{aligned} P \left[k \left\| n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} u_i - \mathbb{E}[Z_i^* Z_i^{*'} u_i]) \right\|^2 > \epsilon \right] &\leq kn^{-1} \mathbb{E}_P[(Z_i^{*'} Z_i^*)^2 u_i^2] \\ &= O(n^{-1} k^3) \\ &= o(1). \end{aligned} \quad (\text{SA-7.33})$$

Furthermore, since $\|\mathbb{E}_P[Z_i^* Z_i^{*'} u_i]\| \leq \mathbb{E}_P[\|Z_i^* Z_i^{*'} u_i\|] \leq \mathbb{E}_P[\|Z_i^*\|^2 |u_i|] = k \mathbb{E}_P[k^{-1} \sum_{j=1}^k Z_{ij}^{*2} |u_i|] = O(k)$, by the triangle inequality, (SA-7.33) and the fact that $\bar{u} = O_p(n^{-1/2})$

$$\begin{aligned} \left\| n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} u_i \bar{u} \right\| &\leq \left\| n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} u_i - \mathbb{E}_P[Z_i^* Z_i^{*'} u_i]) \right\| |\bar{u}| + \|\mathbb{E}_P[Z_i^* Z_i^{*'} u_i]\| |\bar{u}| \\ &= o_p(k^{-1/2} n^{-1/2}) + O_p(kn^{-1/2}) = O_p(kn^{-1/2}). \end{aligned} \quad (\text{SA-7.34})$$

(SA-7.32) follows. Similarly, we have

$$\begin{aligned} &P \left[k \left\| n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} - \mathbb{E}_P[Z_i^* Z_i^{*'}]) \right\|^2 > \epsilon \right] \\ &\leq \frac{k}{\epsilon} \mathbb{E}_P \left[\text{tr} \left\{ \left(n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} - \mathbb{E}_P[Z_i^* Z_i^{*'}]) \right) \left(n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} - \mathbb{E}_P[Z_i^* Z_i^{*'}]) \right) \right\} \right] \\ &= \frac{k}{n\epsilon} \text{tr} \{ \mathbb{E}_P [(Z_i^* Z_i^{*'} - \mathbb{E}_P[Z_i^* Z_i^{*'}]) (Z_i^* Z_i^{*'} - \mathbb{E}_P[Z_i^* Z_i^{*'}])] \} \\ &= \frac{k}{n\epsilon} (\mathbb{E}_P[(Z_i^{*'} Z_i^*)^2] - \|\mathbb{E}_P[Z_i^* Z_i^{*'}]\|^2) \\ &\leq \frac{k^3}{n} \mathbb{E}_P[(k^{-1} Z_i^{*'} Z_i^*)^2] \\ &= o(1). \end{aligned} \quad (\text{SA-7.35})$$

Furthermore, since $\|\mathbb{E}_P[Z_i^* Z_i^{*'}]\| = \left(\sum_{j=1}^k \sum_{\ell=1}^\ell (\mathbb{E}_P[Z_{ij}^* Z_{i\ell}^{*'}])^2 \right)^{1/2} = O(k)$, by the triangle inequality and (SA-7.35)

$$\left\| n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} \right\| \leq \left\| n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} - \mathbb{E}_P[Z_i^* Z_i^{*'}]) \right\| + \|\mathbb{E}_P[Z_i^* Z_i^{*'}]\| = o_p(k^{-1/2}) + O_p(k) = O_p(k). \quad (\text{SA-7.36})$$

Since $\bar{u}^2 = O_p(n^{-1})$, it follows from (SA-7.36) that for the last summand on the RHS of (SA-7.27)

$$\left\| n^{-1} \sum_{i=1}^n Z_i^* Z_i^{*'} \right\| \bar{u}^2 = O_p(n^{-1}k). \quad (\text{SA-7.37})$$

Combining (SA-7.31), (SA-7.32) and (SA-7.37) in (SA-7.27), we obtain

$$\left\| n^{-1} \sum_{i=1}^n (Z_i^* Z_i^{*'} \tilde{u}_i^2 - E_P[Z_i^* Z_i^{*'} u_i^2]) \right\| = o_p(k^{-1/2}). \quad (\text{SA-7.38})$$

The limit of $n^{-1} \sum_{i=1}^n (\bar{W} - E_P[W_i]) Z_i^{*'} \tilde{u}_i^2$. We will next show that

$$\begin{aligned} \left\| (\bar{W} - E_P[W_i]) n^{-1} \sum_{i=1}^n Z_i^* \tilde{u}_i^2 \right\| &= \|\bar{W} - E_P[W_i]\| \left\| n^{-1} \sum_{i=1}^n Z_i^* u_i^2 - 2\bar{u} n^{-1} \sum_{i=1}^n Z_i^* u_i + n^{-1} \sum_{i=1}^n Z_i^* \bar{u}^2 \right\| \\ &= o_p(k^{-2}). \end{aligned} \quad (\text{SA-7.39})$$

By Markov's inequality for any $\epsilon > 0$

$$\begin{aligned} P \left[k^2 \left\| n^{-1} \sum_{i=1}^n (W_i - E_P[W_i]) \right\|^2 > \epsilon \right] &\leq n^{-1} k^2 \text{tr} \{ E[W_i W_i'] - E[W_i] E[W_i]' \} \\ &\leq n^{-1} k^3 E[k^{-1} W_i' W_i] = o(1), \end{aligned}$$

hence (see also Lemma 7(d) of Andrews and Stock (2007b))

$$\left\| n^{-1} \sum_{i=1}^n (W_i - E_P[W_i]) \right\| = o_p(k^{-1}). \quad (\text{SA-7.40})$$

By Markov's and Cauchy-Schwarz inequalities, for any $\epsilon > 0$

$$\begin{aligned} P \left[k^2 \left\| n^{-1} \sum_{i=1}^n (Z_i^* u_i^2 - E_P[Z_i^* u_i^2]) \right\|^2 > \epsilon \right] &\leq \frac{k^2}{n\epsilon} (E_P[Z_i^{*'} Z_i^* u_i^4] - \|E_P[Z_i^* u_i^2]\|^2) \\ &\leq \frac{k^2}{n\epsilon} E_P[Z_i^{*'} Z_i^* u_i^4] \\ &= \frac{k^2}{n\epsilon} \sum_{j=1}^k E_P[Z_{ij}^{*2} u_i^4] \\ &\leq \frac{k^2}{n\epsilon} \sum_{j=1}^k (E_P[Z_{ij}^{*4} u_i^4])^{1/2} (E_P[u_i^4])^{1/2} \\ &= O(k^3 n^{-1}) \\ &= o(1), \end{aligned}$$

so we have

$$n^{-1} \sum_{i=1}^n Z_i^* u_i^2 = o_p(k^{-1}) \quad (\text{SA-7.41})$$

Proceeding similarly to above and using $\bar{u} = O_p(n^{-1/2})$

$$\bar{u} n^{-1} \sum_{i=1}^n Z_i^* u_i = O_p(n^{-1/2}) o_p(k^{-1}) = o_p(n^{-1/2} k^{-1}). \quad (\text{SA-7.42})$$

From (SA-7.40) and $\bar{u}^2 = O_p(n^{-1})$, we have

$$n^{-1} \sum_{i=1}^n Z_i^* \bar{u}^2 = o_p(k^{-1} n^{-1}). \quad (\text{SA-7.43})$$

Combining (SA-7.40), (SA-7.41), (SA-7.42) and (SA-7.43), we have

$$\begin{aligned} \left\| (\bar{W} - E_P[W_i]) n^{-1} \sum_{i=1}^n Z_i^* \tilde{u}_i^2 \right\| &= \|\bar{W} - E_P[W_i]\| \left\| n^{-1} \sum_{i=1}^n Z_i^* u_i^2 - 2\bar{u} n^{-1} \sum_{i=1}^n Z_i^* u_i + n^{-1} \sum_{i=1}^n Z_i^* \bar{u}^2 \right\| \\ &= o_p(k^{-1})(o_p(k^{-1}) + O_p(n^{-1/2})o_p(k^{-1}) + o_p(k^{-1} n^{-1})) \end{aligned} \quad (\text{SA-7.44})$$

give (SA-7.39).

The limit of $n^{-1} \sum_{i=1}^n (\bar{W} - E_P[W_i])(\bar{W} - E_P[W_i])' \tilde{u}_i^2$.

Using the triangle inequality, (SA-7.40) and the fact that $n^{-1} \sum_{i=1}^n \tilde{u}_i^2 = O_p(1)$

$$\begin{aligned} \left\| n^{-1} \sum_{i=1}^n (\bar{W} - E_P[W_i])(\bar{W} - E_P[W_i])' \tilde{u}_i^2 \right\| &\leq \|\bar{W} - E_P[W_i]\|^2 n^{-1} \sum_{i=1}^n \tilde{u}_i^2 \\ &= o_p(k^{-2}). \end{aligned} \quad (\text{SA-7.45})$$

Finally, using (SA-7.31), (SA-7.39) and (SA-7.45) in (SA-7.26) and the triangle inequality, we obtain (SA-7.25).

Substep 2: Asymptotic distribution of AR statistic.

Set in Lemma SA-2.6, $\hat{\xi}_{ni} = Z_i u_i$, $\hat{\xi}_{ni}^* = Z_i^* u_i$, $\hat{\Sigma}_{n\xi} = \hat{\Sigma}$, $\Sigma_{n\xi} = \Sigma$, $\hat{D}_{n\xi} = [\bar{W}, 0_{k \times (k-1)}]$, $D_{n\xi} = [E_P[W_i], 0_{k \times (k-1)}]$, $\xi_{n1i} = W_i u_i$ and $\xi_{n2\xi} = [u_i, 0'_{k-1}]'$. Then, Conditions (a)(i)-(iii) and (v)-(viii) are satisfied with $\mu_{n\xi} = 0$ and $\lambda_{n\xi}^* = 0$ by the conditions of Proposition 2.4. Condition (a)(iv) follows from (SA-7.25). Condition (c) is trivially satisfied since $E_P[u_i] = 0$. Conditions (b) and (e) hold by definitions. Finally, Condition (d) holds by (SA-7.40). Therefore, it follows from (SA-2.8) that

$$(2k)^{-1/2}(\text{AR} - k) = (2k)^{-1/2} \left(n^{-1/2} \sum_{i=1}^n [\hat{\Sigma}_{n\xi}^{-1/2} \hat{\xi}_{ni}]' n^{-1/2} \sum_{j=1}^n [\hat{\Sigma}_{n\xi}^{-1/2} \hat{\xi}_{nj}] - k \right) \xrightarrow{d} N[0, 1].$$

Step 3: Completing the proof.

Since, from (SA-7.24), $(2k)^{-1/2}(\text{PAR} - k) \xrightarrow{d^\pi} N[0, 1]$ in P -probability, and $N[0, 1]$ is increasing at its $1 - \alpha$ quantile $z_{1-\alpha}$, $(2k)^{-1/2}(\text{PAR}_{(r)} - k) \xrightarrow{p} z_{1-\alpha}$. Then, since $(2k)^{-1/2}(\text{AR} - k) \xrightarrow{d} N[0, 1]$, proceeding similarly to the proof of the PAR tests for k fixed, we obtain

$$\lim_{n \rightarrow \infty} E_P[\phi_n^{\text{PAR}}] = \alpha.$$

SA-8 Proofs of supplementary results

SA-8.1 Lemma SA-2.2

By the triangle, c_r and Jensen's inequalities, $E[|Z_{nk} - E[Z_{nk}]|^{2p}] \leq 2^{2p-1}(E[|Z_{nk}|^{2p}] + |E[Z_{nk}]|^{2p}) < \infty$. Applying Theorem 1 of [Hu et al. \(1989\)](#) with $X_{nk} = Z_{nk} - E[Z_{nk}]$ yields that $n^{-1/p} \sum_{k=1}^n X_{nk}$ converges completely to 0 i.e. for any $\varepsilon > 0$

$$\sum_{n=1}^{\infty} P \left[\left| n^{-1/p} \sum_{k=1}^n X_{nk} \right| > \varepsilon \right] < \infty. \quad (\text{SA-8.1})$$

From Borel-Cantelli lemma, it follows that $n^{-1/p} \sum_{k=1}^n X_{nk} \xrightarrow{a.s.} 0$.

SA-8.2 Lemma SA-2.3

The proof is analogous to the bootstrap version of the result, see [Hansen \(2022\)](#), Theorem 10.1.

SA-8.3 Lemma SA-2.5

By direct calculations,

$$\begin{aligned} E_{P^\pi} \left[\left\| \sum_{i=1}^n a_i b_{\pi(i)} \right\|^2 \right] &= E_{P^\pi} \left[\left(\sum_{i=1}^n a_i b_{\pi(i)} \right)' \left(\sum_{j=1}^n a_j b_{\pi(j)} \right) \right] \\ &= E_{P^\pi} \left[\sum_{i=1}^n a_i' a_i b_{\pi(i)}^2 \right] + E_{P^\pi} \left[\sum_{i \neq j}^n a_i' a_j b_{\pi(i)} b_{\pi(j)} \right] \\ &= \left(\sum_{i=1}^n a_i' a_i \right) \left(n^{-1} \sum_{i=1}^n b_i^2 \right) + \left(\sum_{i \neq j}^n a_i' a_j \right) \left(\frac{1}{n(n-1)} \sum_{i \neq j} b_i b_j \right). \quad (\text{SA-8.2}) \end{aligned}$$

On noting that $\sum_{i \neq j} a_i' a_j = \sum_{i=1}^n a_i' \left(\sum_{j=1}^n a_j - a_i \right) = \left\| \sum_{i=1}^n a_i \right\|^2 - \sum_{i=1}^n \|a_i\|^2$, and similarly, $\sum_{i \neq j} b_i b_j = -\sum_{i=1}^n b_i^2$ in (SA-8.2), we obtain (SA-2.7).

SA-8.4 Lemma SA-5.1

The proof is divided into three main steps. In the first step, we prove the multivariate normality of some key permutation quantities. In the second step, we prove the consistency of the covariance matrix estimates. The final step shows the desired result of the lemma.

Step 1: Multivariate normality.

By the triangular array SLLN (Lemma SA-2.2), $n^{-1}W'W - E_P[W_i W_i'] \xrightarrow{a.s.} 0$, $n^{-1}X'W - E_P[X_i W_i'] \xrightarrow{a.s.} 0$ and $n^{-1}X'X - E_P[X_i X_i'] \xrightarrow{a.s.} 0$, and by (SA-5.22) $E_P[W_i W_i'] \rightarrow Q_{ww}$, $E_P[X_i W_i'] \rightarrow Q_{xw}$, and $E_P[X_i X_i'] \rightarrow Q_{xx}$. Thus,

$$E_P[Z_i^* Z_i^{*'}] \rightarrow Q_{ww} - Q_{wx} Q_{xx}^{-1} Q_{xw} \equiv Q_{zz}, \quad (\text{SA-8.3})$$

and by the CMT

$$n^{-1}Z'Z = n^{-1}W'W - n^{-1}W'X(X'X)^{-1}X'W \xrightarrow{a.s.} Q_{zz}. \quad (\text{SA-8.4})$$

Note that $n^{1/2}(\hat{J}_s^\pi - n^{-1}Z'Z\hat{\Gamma}_s) = n^{1/2}Z'\hat{V}_{\pi,s} - \hat{C}_s^\pi \hat{\Sigma}^\pi n^{1/2}\hat{m}^\pi$, $s = 1, \dots, d$, where $\hat{\Gamma}_s = (Z'Z)^{-1}Z'Y_s$ and $\hat{V}_{\pi,s}$ is the s -th column of \hat{V}_π . Next we will show that

$$\begin{bmatrix} n^{-1/2}Z'M_l \tilde{u}_\pi(\theta_0) \\ n^{-1/2}\text{vec}(Z'M_l \hat{V}_\pi) \end{bmatrix} \xrightarrow{d^\pi} \begin{bmatrix} m_h^\pi \\ G_h^\pi \end{bmatrix} \sim N[0, \Sigma^e \otimes Q_{zz}] \quad P\text{-almost surely.} \quad (\text{SA-8.5})$$

Using the Cramér-Wold device, it suffices to prove that for any $[t'_0, t'_1, \dots, t'_d]'$ with $t_l \in \mathbb{R}^k$, $l = 0, \dots, d$

$$t' \begin{bmatrix} n^{-1/2}Z'M_l \tilde{u}_\pi(\theta_0) \\ n^{-1/2}\text{vec}(Z'M_l \hat{V}_\pi) \end{bmatrix} \xrightarrow{d^\pi} N[0, t'(\Sigma^e \otimes Q_{zz})t] \quad P\text{-almost surely.} \quad (\text{SA-8.6})$$

Rewrite

$$\begin{aligned} t' \begin{bmatrix} n^{-1/2}Z'M_l \tilde{u}_\pi(\theta_0) \\ n^{-1/2}\text{vec}(Z'M_l \hat{V}_\pi) \end{bmatrix} &= n^{-1/2} \sum_{i=1}^n \left(t'_0 Z_i \tilde{u}_{\pi(i)}(\theta_0) + t'_1 Z_i \hat{V}_{\pi(i)1} + \dots + t'_d Z_i \hat{V}_{\pi(i)d} \right) \\ &= n^{-1/2} \sum_{i=1}^n Z_i' [t_0, t_1, \dots, t_d] [\tilde{u}_{\pi(i)}(\theta_0), \hat{V}_{\pi(i)1}, \dots, \hat{V}_{\pi(i)d}]' \\ &= n^{-1/2} \sum_{i=1}^n Z_i' \zeta_{\pi(i)}, \end{aligned}$$

where $\zeta_i \equiv [t_0, t_1, \dots, t_d] [\tilde{u}_i(\theta_0), \hat{V}_{i1}, \dots, \hat{V}_{id}]'$, $i = 1, \dots, n$. We verify the condition (SA-2.1) of Lemma SA-2.1 with $c_n(i, j) \equiv n^{-1/2}Z_i' \zeta_j$. Since

$$\begin{aligned} E_{P^\pi} \left[n^{-1/2} \sum_{i=1}^n t'_0 Z_i \tilde{u}_{\pi(i)}(\theta_0) | y, Y, W, X \right] &= n^{1/2} \left(n^{-1} \sum_{i=1}^n t'_0 Z_i \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0) \right) = 0, \\ E_{P^\pi} \left[n^{-1/2} \sum_{i=1}^n t'_l Z_i \hat{V}_{\pi(i)l} | y, Y, W, X \right] &= n^{1/2} \left(n^{-1} \sum_{i=1}^n t'_l Z_i \right) \left(n^{-1} \sum_{i=1}^n \hat{V}_{il} \right) = 0, \quad l = 1, \dots, d, \end{aligned}$$

we have

$$\begin{aligned} \mathbb{E}_{P^\pi} \left[n^{-1/2} \sum_{i=1}^n t'_0 Z_i \tilde{u}_{\pi(i)}(\theta_0) | y, Y, W, X \right] &= n^{1/2} \left(n^{-1} \sum_{i=1}^n t'_0 Z_i \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0) \right) = 0, \\ \mathbb{E}_{P^\pi} \left[n^{-1/2} \sum_{i=1}^n t'_0 Z_i \hat{V}_{\pi(i)l} | y, Y, W, X \right] &= n^{1/2} \left(n^{-1} \sum_{i=1}^n t'_0 Z_i \right) \left(n^{-1} \sum_{i=1}^n \hat{V}_{il} \right) = 0, \quad l = 1, \dots, d, \end{aligned}$$

and $g_n(i, j) = c_n(i, j)$ in Lemma SA-2.1. Next we will show that

$$n^{-1} \tilde{e}' \tilde{e} = n^{-1} \sum_{i=1}^n \tilde{e}_i \tilde{e}'_i = n^{-1} \begin{bmatrix} \tilde{u}(\theta_0)' \tilde{u}(\theta_0) & \tilde{u}(\theta_0)' \hat{V} \\ \hat{V}' \tilde{u}(\theta_0) & \hat{V}' \hat{V} \end{bmatrix} \xrightarrow{a.s.} \Sigma^e, \quad (\text{SA-8.7})$$

or equivalently

$$n^{-1} \hat{V}' \hat{V} \xrightarrow{a.s.} \Sigma^V, \quad (\text{SA-8.8})$$

$$n^{-1} \hat{V}' \tilde{u}(\theta_0) \xrightarrow{a.s.} \Sigma^{Vu}, \quad (\text{SA-8.9})$$

$$n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^2 \xrightarrow{a.s.} \sigma^2. \quad (\text{SA-8.10})$$

By the SLLN (Lemma SA-2.2), $n^{-1} W' V - \mathbb{E}_P[W_i V'_i] \xrightarrow{a.s.} 0$, hence $n^{-1} W' V \xrightarrow{a.s.} 0$ and similarly $n^{-1} X' V \xrightarrow{a.s.} 0$. Combining these with $n^{-1} W' X \xrightarrow{a.s.} Q_{wx}$ and $n^{-1} X' X \xrightarrow{a.s.} Q_{xx}$, and using the CMT,

$$n^{-1} Z' V = n^{-1} W' V - n^{-1} W' X (n^{-1} X' X)^{-1} n^{-1} X' V \xrightarrow{a.s.} 0. \quad (\text{SA-8.11})$$

Thus, on noting that $\hat{V} = Y - Z\hat{F} - X\hat{\xi} = (I_n - P_Z - P_X)V$, by the SLLN and CMT

$$n^{-1} \hat{V}' \hat{V} - \mathbb{E}_P[V_i V'_i] = n^{-1} V' V - n^{-1} V' P_Z V - n^{-1} V' P_X V - \mathbb{E}_P[V_i V'_i] \xrightarrow{a.s.} 0$$

and $\mathbb{E}_P[V_i V'_i] \rightarrow \Sigma^V$, we have $n^{-1} \hat{V}' \hat{V} \xrightarrow{a.s.} \Sigma^V$ which verifies (SA-8.8). Analogously to (SA-8.11), $n^{-1} Z' u \xrightarrow{a.s.} 0$ and $n^{-1} X' u \xrightarrow{a.s.} 0$. Then, (SA-8.9) follows using (SA-8.4) and on noting that

$$\begin{aligned} n^{-1} \hat{V}' \tilde{u}(\theta_0) - \mathbb{E}_P[V_i u_i] &= n^{-1} V' \tilde{u}(\theta_0) - n^{-1} V' P_Z \tilde{u}(\theta_0) - n^{-1} V' P_X \tilde{u}(\theta_0) - \mathbb{E}_P[V_i u_i] \\ &= n^{-1} V' u - n^{-1} V' X (X' X)^{-1} X' u - \mathbb{E}_P[V_i u_i] - n^{-1} V' Z (Z' Z)^{-1} Z' u \\ &\xrightarrow{a.s.} 0, \end{aligned}$$

and $\mathbb{E}_P[V_i u_i] \rightarrow \Sigma^{Vu}$. (SA-8.10) holds because $n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^2 - \mathbb{E}_P[u_i^2] = n^{-1} u' u - \mathbb{E}_P[u_i^2] - n^{-1} u' X (X' X)^{-1} X' u \xrightarrow{a.s.} 0$ by the SLLN and CMT, and $\sigma_n^2 \equiv \mathbb{E}_P[u_i^2] \rightarrow \sigma^2$. Furthermore, note

that

$$\begin{aligned}
\hat{D} &\equiv \frac{1}{n-1} \sum_{i=1}^n \sum_{j=1}^n g_n^2(i, j) \\
&= \frac{1}{n(n-1)} \sum_{i=1}^n \sum_{j=1}^n \zeta_i' Z_j Z_j' \zeta_i \\
&= \frac{1}{n-1} \sum_{i=1}^n \zeta_i' \left(n^{-1} \sum_{j=1}^n Z_j Z_j' \right) \zeta_i \\
&= \frac{1}{n-1} \sum_{i=1}^n \zeta_i' (Q_{zz} + o_{a.s.}(1)) \zeta_i \\
&= \frac{1}{n-1} \sum_{i=1}^n \zeta_i' Q_{zz} \zeta_i + \frac{1}{n-1} \sum_{i=1}^n \zeta_i' \zeta_i o_{a.s.}(1) \\
&= \frac{1}{n-1} \sum_{i=1}^n t'(\tilde{e}_i \otimes I_k) Q_{zz} (\tilde{e}_i' \otimes I_k) t + \frac{1}{n-1} \sum_{i=1}^n \zeta_i' \zeta_i o_{a.s.}(1) \\
&= \frac{1}{n-1} \sum_{i=1}^n t'(\tilde{e}_i \tilde{e}_i' \otimes Q_{zz}) t + o_{a.s.}(1) \\
&= t'(\Sigma^e \otimes Q_{zz}) t + o_{a.s.}(1),
\end{aligned} \tag{SA-8.12}$$

where the third equality follows from (SA-8.4), the second to the last equality uses (SA-8.7) and the fact that $n^{-1} \sum_{i=1}^n \zeta_i' \zeta_i = n^{-1} \sum_{i=1}^n t'(\tilde{e}_i \tilde{e}_i' \otimes I_k) t \xrightarrow{a.s.} t'(\Sigma^e \otimes I_k) t$ which follows from (SA-8.7) and the Slutsky's lemma. By the triangular array SLLN,

$$n^{-1} \sum_{i=1}^n \|V_i\|^{2+\delta} = O_{a.s.}(1), \quad n^{-1} \sum_{i=1}^n \|X_i\|^{2+\delta} = O_{a.s.}(1), \quad n^{-1} \sum_{i=1}^n \|W_i\|^{2+\delta} = O_{a.s.}(1), \tag{SA-8.13}$$

$$n^{-1} \sum_{i=1}^n \|Z_i\|^{2+\delta} \leq 2^{1+\delta} \left(n^{-1} \sum_{i=1}^n \|W_i\|^{2+\delta} + \|X_i\|^{1+\delta} \|(X'X)^{-1} X'W\|^{2+\delta} \right) = O_{a.s.}(1). \tag{SA-8.14}$$

By the c_r inequality and the triangular array SLLN,

$$\begin{aligned}
n^{-1} \sum_{i=1}^n |\tilde{u}_i(\theta_0)|^{2+\delta} &\leq 2^{1+\delta} n^{-1} \sum_{i=1}^n |u_i|^{2+\delta} + 2^{1+\delta} n^{-1} \sum_{i=1}^n \|X_i\|^{2+\delta} \|(X'X)^{-1} X'u\|^{2+\delta} \\
&= O_{a.s.}(1).
\end{aligned} \tag{SA-8.15}$$

Since $\hat{V} = (I_n - P_Z - P_X)V$, $\hat{V}_i = V_i - Z_i'(Z'Z)^{-1}Z'V - X_i'(X'X)^{-1}X'V$, and by Jensen's inequality

(applied with the function $f(x) = x^{2+\delta}, x > 0$), (SA-8.13) and (SA-8.14)

$$\begin{aligned} n^{-1} \sum_{i=1}^n \|\hat{V}_i\|^{2+\delta} &\leq 3^{1+\delta} \left(n^{-1} \sum_{i=1}^n \|V_i\|^{2+\delta} + n^{-1} \sum_{i=1}^n \|Z_i\|^{2+\delta} \|(n^{-1} Z' Z)^{-1} n^{-1} Z' V\|^{2+\delta} \right. \\ &\quad \left. + n^{-1} \sum_{i=1}^n \|X_i\|^{2+\delta} \|(n^{-1} X' X)^{-1} n^{-1} X' V\|^{2+\delta} \right) = O_{a.s.}(1). \end{aligned} \quad (\text{SA-8.16})$$

By the c_r inequality, (SA-8.15) and (SA-8.16)

$$\begin{aligned} n^{-1} \sum_{i=1}^n \|\zeta_i\|^{2+\delta} &\leq 2^{1+\delta} \|t_0\|^{2+\delta} n^{-1} \sum_{i=1}^n |\tilde{u}_i(\theta_0)|^{2+\delta} + 2^{1+\delta} \|[t_1, \dots, t_d]\|^{2+\delta} n^{-1} \sum_{i=1}^n \|\hat{V}_i\|^{2+\delta} \\ &= O_{a.s.}(1). \end{aligned} \quad (\text{SA-8.17})$$

We show that for \hat{D} defined in (SA-8.12) and any $\epsilon > 0$

$$\lim_{n \rightarrow \infty} n^{-2} \sum_{i=1}^n \sum_{j=1}^n \frac{|Z'_i \zeta_j|^2}{\hat{D}} 1 \left(\frac{n^{-1} |Z'_i \zeta_j|^2}{\hat{D}} > \epsilon^2 \right) = 0. \quad (\text{SA-8.18})$$

Note that

$$\begin{aligned} n^{-2} \sum_{i=1}^n \sum_{j=1}^n \frac{|Z'_i \zeta_j|^2}{\hat{D}} 1 \left(\frac{n^{-1} |Z'_i \zeta_j|^2}{\hat{D}} > \epsilon^2 \right) &= n^{-2} \hat{D}^{-1} \sum_{i=1}^n \sum_{j=1}^n \frac{|Z'_i \zeta_j|^{2+\delta}}{|Z'_i \zeta_j|^\delta} 1 \left(|Z'_i \zeta_j|^\delta > \epsilon^\delta \hat{D}^{\delta/2} n^{\delta/2} \right) \\ &\leq n^{-\delta/2} \epsilon^{-\delta} \hat{D}^{-1-\delta/2} n^{-2} \sum_{i=1}^n \sum_{j=1}^n |Z'_i \zeta_j|^{2+\delta} \\ &\leq n^{-\delta/2} \epsilon^{-\delta} \hat{D}^{-1-\delta/2} n^{-1} \sum_{i=1}^n \|Z_i\|^{2+\delta} n^{-1} \sum_{i=1}^n \|\zeta_i\|^{2+\delta} \\ &\xrightarrow{a.s.} 0, \end{aligned} \quad (\text{SA-8.19})$$

where the convergence uses (SA-8.14) and (SA-8.17). Thus, the Lindeberg-type condition in (SA-2.1) is satisfied. Lemma SA-2.1 and (SA-8.12) yield (SA-8.6).

Step 2: Consistency of the covariance matrix estimates.

Next we determine the limits of $\hat{C}_s^\pi, s = 1, \dots, d$, and $\hat{\Sigma}^\pi$. Note first that by Lemma S.3.4 of DiCiccio and Romano (2017) (see also Hoeffding (1951)),

$$\begin{aligned} E_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W, X] &= E_{P^\pi} \left[n^{-1} \sum_{i=1}^n Z_i Z'_i \tilde{u}_{\pi(i)}(\theta_0)^2 | y, Y, W, X \right] \\ &= (n^{-1} Z' Z) (n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^2) \xrightarrow{a.s.} \sigma^2 Q_{zz}, \end{aligned} \quad (\text{SA-8.20})$$

where the convergence follows from the CMT, (SA-8.4) and (SA-8.10). On noting that $n^{-1}X'u \xrightarrow{p} 0$ and $n^{-1} \sum_{i=1}^n u_i^4 - \mathbb{E}_P[u_i^4] \xrightarrow{p} 0$, by Minkowski's inequality, the triangular array WLLN and the CMT,

$$\begin{aligned} n^{-1} \sum_{i=1}^n \|Z_i\|^4 &= n^{-1} \sum_{i=1}^n \|W_i - W'X(X'X)^{-1}X_i\|^4 \\ &\leq 8n^{-1} \sum_{i=1}^n \|W_i\|^4 + \|W'X(X'X)^{-1}\|^4 8n^{-1} \sum_{i=1}^n \|X_i\|^4 \\ &= O_p(1), \end{aligned} \tag{SA-8.21}$$

$$n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 \leq 8n^{-1} \sum_{i=1}^n u_i^4 + 8n^{-1} \sum_{i=1}^n \|X_i\|^4 \|(X'X)^{-1}X'u\|^4 = O_p(1), \tag{SA-8.22}$$

hence

$$n^{-1} \sum_{i=1}^n Z_{ij}^2 Z_{il}^2 \leq \left(n^{-1} \sum_{i=1}^n Z_{ij}^4 \right)^{1/2} \left(n^{-1} \sum_{i=1}^n Z_{il}^4 \right)^{1/2} = O_p(1) \quad j, l = 1, \dots, k. \tag{SA-8.23}$$

Therefore, letting $\hat{\Sigma}_{jl}^\pi$ denote the $(j, l), j, l = 1, \dots, k$, element of $\hat{\Sigma}^\pi$, (SA-8.22) and (SA-8.23) give

$$\begin{aligned} &\text{Var}_{P^\pi}[\hat{\Sigma}_{jl}^\pi | y, Y, W, X] \\ &= \text{Var}_{P^\pi} \left[n^{-1} \sum_{i=1}^n Z_{ij} Z_{il} \tilde{u}_{\pi(i)}(\theta_0)^2 | y, Y, W, X \right] \\ &= \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n Z_{ij}^2 Z_{il}^2 - (n^{-1} \sum_{i=1}^n Z_{ij} Z_{il})^2 \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 - (n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^2)^2 \right) \\ &\leq \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n Z_{ij}^2 Z_{il}^2 \right) \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 \right) \\ &\xrightarrow{p} 0, \end{aligned} \tag{SA-8.24}$$

where the second equality follows from Lemma S.3.4 of DiCiccio and Romano (2017). Combining (SA-8.20) and (SA-8.24), and using Chebyshev's inequality, $P^\pi[|\hat{\Sigma}_{jl}^\pi - \mathbb{E}_{P^\pi}[\hat{\Sigma}_{jl}^\pi | y, Y, W, X]| > \epsilon] \leq \text{Var}_{P^\pi}[\hat{\Sigma}_{jl}^\pi | y, Y, W, X] / \epsilon^2 \xrightarrow{p} 0$. Thus $\hat{\Sigma}^\pi - \mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W, X] \xrightarrow{p^\pi} 0$ in P -probability. Combining this with (SA-8.20) and using the triangle inequality, $\|\hat{\Sigma}^\pi - \sigma^2 Q_{zz}\| \leq \|\mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W, X] - \sigma^2 Q_{zz}\| + \|\hat{\Sigma}^\pi - \mathbb{E}_{P^\pi}[\hat{\Sigma}^\pi | y, Y, W, X]\| \xrightarrow{p^\pi} 0$ in P -probability, hence

$$n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{u}_{\pi(i)}(\theta_0)^2 \xrightarrow{p^\pi} \sigma^2 Q_{zz} \quad \text{in } P\text{-probability.} \tag{SA-8.25}$$

Furthermore, by (SA-8.4), (SA-8.9) and the CMT, for $s = 1, \dots, d$

$$\begin{aligned}
\mathbb{E}_{P^\pi}[\hat{C}_s^\pi | y, Y, W, X] &= \mathbb{E}_{P^\pi} \left[n^{-1} \sum_{i=1}^n Z_i Z_i' \hat{V}_{\pi(i)s} \tilde{u}_{\pi(i)}(\theta_0) | y, Y, W, X \right] \\
&= (n^{-1} Z' Z) (n^{-1} \sum_{i=1}^n \hat{V}_{is} \tilde{u}_i(\theta_0)) \\
&\xrightarrow{a.s.} Q_{zz} \Sigma_s^{Vu},
\end{aligned} \tag{SA-8.26}$$

where Σ_s^{Vu} denotes the s -th element of Σ^{Vu} . Since $\hat{V}_i = V_i - Z_i'(Z'Z)^{-1}Z'V - X_i'(X'X)^{-1}X'V$, applying Minkowski's inequality twice and using $n^{-1} \sum_{i=1}^n \|V_i\|^4 = O_p(1)$ and $n^{-1} \sum_{i=1}^n \|X_i\|^4 = O_p(1)$ which hold by the triangular array WLLN,

$$\begin{aligned}
n^{-1} \sum_{i=1}^n \|\hat{V}_i\|^4 &\leq 8n^{-1} \sum_{i=1}^n \|V_i\|^4 + 64n^{-1} \sum_{i=1}^n \|Z_i\|^4 \|(n^{-1} Z' Z)^{-1} n^{-1} Z' V\|^4 \\
&\quad + 64n^{-1} \sum_{i=1}^n \|X_i\|^4 \|(n^{-1} X' X)^{-1} n^{-1} X' V\|^4 = O_p(1).
\end{aligned} \tag{SA-8.27}$$

Then, letting $\hat{C}_{s,jl}^\pi$ denote the (j, l) , $j, l = 1, \dots, k$, element of \hat{C}_s^π , we have

$$\begin{aligned}
&\text{Var}_{P^\pi}[\hat{C}_{s,jl}^\pi | y, Y, W, X] \\
&= \text{Var}_{P^\pi} \left[n^{-1} \sum_{i=1}^n Z_{ij} Z_{il} \hat{V}_{\pi(i)s} \tilde{u}_{\pi(i)}(\theta_0) | y, Y, W, X \right] \\
&= \frac{1}{n-1} \left(n^{-1} \sum_{i=1}^n Z_{ij}^2 Z_{il}^2 - \left(n^{-1} \sum_{i=1}^n Z_{ij} Z_{il} \right)^2 \right) \left(n^{-1} \sum_{i=1}^n \hat{V}_{is}^2 \tilde{u}_i(\theta_0)^2 - \left(n^{-1} \sum_{i=1}^n \hat{V}_{is} \tilde{u}_i(\theta_0) \right)^2 \right) \\
&\xrightarrow{p} 0,
\end{aligned} \tag{SA-8.28}$$

where the convergence follows from (SA-8.4), (SA-8.9), (SA-8.23), and

$$n^{-1} \sum_{i=1}^n \hat{V}_{is}^2 \tilde{u}_i(\theta_0)^2 \leq \left(n^{-1} \sum_{i=1}^n \hat{V}_{is}^4 \right)^{1/2} \left(n^{-1} \sum_{i=1}^n \tilde{u}_i(\theta_0)^4 \right)^{1/2} = O_p(1), \tag{SA-8.29}$$

which in turn holds by (SA-8.22) and (SA-8.27). Therefore, from (SA-8.26), (SA-8.28) and Chebyshev's inequality

$$\hat{C}_s^\pi = n^{-1} \sum_{i=1}^n Z_i Z_i' \hat{V}_{\pi(i)s} \tilde{u}_{\pi(i)}(\theta_0) \xrightarrow{p^\pi} Q_{zz} \Sigma_s^{Vu} \quad \text{in } P\text{-probability.} \tag{SA-8.30}$$

Step 3: Completing the proof.

Let $G_h^\pi = [G_{h,1}^\pi, \dots, G_{h,d}^\pi]$. Using (SA-8.5), (SA-8.25) and (SA-8.30) along with the CMT, for $s = 1, \dots, d$

$$\begin{aligned}
& n^{1/2} \hat{J}_s^\pi - n^{-1/2} Z' Z \hat{F}_s \\
&= n^{-1/2} Z' M_\iota \hat{V}_{\pi,s} - \left(n^{-1} \sum_{i=1}^n Z_i Z_i' \hat{V}_{\pi(i)s} \tilde{u}_{\pi(i)}(\theta_0) \right) \left(n^{-1} \sum_{i=1}^n Z_i Z_i' \tilde{u}_{\pi(i)}(\theta_0)^2 \right)^{-1} n^{-1/2} Z' \tilde{u}_\pi(\theta_0) \\
&\xrightarrow{d^\pi} G_{h,s}^\pi - \Sigma_s^{Vu}(\sigma^2)^{-1} m_h^\pi \quad \text{in } P\text{-probability.}
\end{aligned} \tag{SA-8.31}$$

This proves the part (a). Next we determine the limit of $H_n \hat{J}^\pi U_n T_n$ following Andrews and Guggenberger (2019b). Write

$$n^{1/2} H_n \hat{J}^\pi U_n T_n = n^{1/2} H_n G_n U_n T_n + n^{1/2} H_n (\hat{G} - G_n) U_n T_n + n^{1/2} H_n (\hat{J}^\pi - \hat{G}) U_n T_n. \tag{SA-8.32}$$

Using (SA-5.25), (SA-5.26), (SA-5.27), (SA-5.28), (SA-5.31) and the SVD $H_n G_n U_n = A_n \Upsilon_n B_n'$, the first term on the right-hand side of the equation (SA-8.32) can be expressed as

$$\begin{aligned}
n^{1/2} H_n G_n U_n T_n &= n^{1/2} H_n G_n U_n B_n S_n \\
&= n^{1/2} H_n G_n U_n (B_{n,q}, B_{n,d-q}) \begin{bmatrix} \Upsilon_{n,q}^{-1} n^{-1/2} & 0_{q \times (d-q)} \\ 0_{(d-q) \times q} & I_{d-q} \end{bmatrix} \\
&= [H_n G_n U_n B_{n,q} \Upsilon_{n,q}^{-1}, n^{1/2} H_n G_n U_n B_{n,d-q}] \\
&= [A_n \Upsilon_n B_n' B_{n,q} \Upsilon_{n,q}^{-1}, n^{1/2} H_n G_n U_n B_{n,d-q}] \\
&= [A_{n,q}, n^{1/2} H_n G_n U_n B_{n,d-q}],
\end{aligned} \tag{SA-8.33}$$

where we used $\Upsilon_n B_n' B_{n,q} \Upsilon_{n,q}^{-1} = \Upsilon_n B_n' B_n [I_q, 0_{q \times (d-q)}]' \Upsilon_{n,q}^{-1} = [I_q, 0_{q \times (d-q)}]'$. For the second submatrix in (SA-8.33),

$$\begin{aligned}
n^{1/2} H_n G_n U_n B_{n,d-q} &= n^{1/2} A_n \Upsilon_n B_n' B_{n,d-q} \\
&= n^{1/2} A_n \Upsilon_n [0_{(d-q) \times q}, I_{d-q}]' \\
&= A_n [0_{(d-q) \times q}, n^{1/2} \Upsilon_{n,d-q}, 0_{(d-q) \times (k-d)}]' \\
&\rightarrow h_3 [0_{(d-q) \times q}, \text{diag}(h_{1,q+1}, \dots, h_{1,d}), 0_{(d-q) \times (k-d)}]' \\
&= h_3 \tilde{h}_{1,d-q},
\end{aligned} \tag{SA-8.34}$$

where $\tilde{h}_{1,d-q}$ is defined in (SA-2.12). Furthermore,

$$\begin{aligned}
n^{1/2}H_n(\hat{G} - G_n)U_nT_n &= n^{1/2}H_n(\hat{G} - G_n)U_nB_nS_n \\
&= n^{1/2}H_n(\hat{G} - G_n)U_n[B_{n,q}, B_{n,d-q}] \begin{bmatrix} \Upsilon_{n,q}^{-1}n^{-1/2} & 0_{q \times (d-q)} \\ 0_{(d-q) \times q} & I_{d-q} \end{bmatrix} \\
&= [H_n(\hat{G} - G_n)U_nB_{n,q}\Upsilon_{n,q}^{-1}, n^{1/2}H_n(\hat{G} - G_n)U_nB_{n,d-q}]. \tag{SA-8.35}
\end{aligned}$$

Using $H_n \rightarrow \Sigma^{-1/2}$, (SA-5.56), $U_n \rightarrow h_{91}$ where h_{91} is defined in (SA-2.12), $B_{n,q} \rightarrow h_{3,q}$, and $n^{1/2}\tau_{jn} \rightarrow \infty$ for all $j \leq q$,

$$H_n(\hat{G} - G_n)U_nB_{n,q}\Upsilon_{n,q}^{-1} = H_n n^{1/2}(\hat{G} - G_n)U_nB_{n,q}(n^{1/2}\Upsilon_{n,q})^{-1} = o_p(1). \tag{SA-8.36}$$

Since $B_{n,d-q} \rightarrow h_{2,d-q}$,

$$n^{1/2}H_n(\hat{G} - G_n)U_nB_{n,d-q} \xrightarrow{d} \Sigma^{-1/2}G_h h_{91} h_{2,d-q}. \tag{SA-8.37}$$

Using (SA-8.36) and (SA-8.37) in (SA-8.35),

$$n^{1/2}H_n(\hat{G} - G_n)U_nT_n \xrightarrow{d} [0_{k \times q}, \Sigma^{-1/2}G_h h_{91} h_{2,d-q}]. \tag{SA-8.38}$$

Moreover, rewrite the the third term on the right-hand side of the equation (SA-8.32) as

$$n^{1/2}H_n(\hat{J}^\pi - \hat{G})U_nT_n = n^{1/2}H_n(\hat{J}^\pi - \hat{G})U_nB_nS_n \tag{SA-8.39}$$

$$\begin{aligned}
&= n^{1/2}H_n(\hat{J}^\pi - \hat{G})U_n[B_{n,q}, B_{n,d-q}] \begin{bmatrix} \Upsilon_{n,q}^{-1}n^{-1/2} & 0_{q \times (d-q)} \\ 0_{(d-q) \times q} & I_{d-q} \end{bmatrix} \\
&= [H_n(\hat{J}^\pi - \hat{G})U_nB_{n,q}\Upsilon_{n,q}^{-1}, n^{1/2}H_n(\hat{J}^\pi - \hat{G})U_nB_{n,d-q}]. \tag{SA-8.40}
\end{aligned}$$

In addition, by the part (a) of Lemma SA-5.1, $n^{1/2}(\hat{J}^\pi - \hat{G}) \xrightarrow{d^\pi} J_h^\pi$ in P -probability. Using an argument analogous to those in (SA-8.36) and (SA-8.37) and Lemma SA-2.3,

$$H_n(\hat{J}^\pi - \hat{G})U_nB_{n,q}\Upsilon_{n,q}^{-1} = o_{p^\pi}(1) \quad \text{in } P\text{-probability}, \tag{SA-8.41}$$

$$n^{1/2}H_n(\hat{J}^\pi - \hat{G})U_nB_{n,d-q} \xrightarrow{d^\pi} \Sigma^{-1/2}J_h^\pi h_{91} h_{2,d-q} \quad \text{in } P\text{-probability}. \tag{SA-8.42}$$

Using (SA-8.41) and (SA-8.42) in (SA-8.40),

$$n^{1/2}H_n(\hat{J}^\pi - \hat{G})U_nT_n \xrightarrow{d^\pi} [0_{k \times q}, \Sigma^{-1/2}J_h^\pi h_{91} h_{2,d-q}] \quad \text{in } P\text{-probability}. \tag{SA-8.43}$$

Combining (SA-8.33), (SA-8.34), (SA-8.38), (SA-8.43), and using $A_{n,q} \rightarrow h_{3,q} = \Delta_{h,q}^\pi$ and an argument similar to that in (SA-5.61) in (SA-8.32), we obtain

$$n^{1/2} H_n \hat{J}^\pi U_n T_n \xrightarrow{d^\pi} (\Delta_{h,q}, h_3 \tilde{h}_{1,d-q} + \Sigma^{-1/2} (G_h + J_h^\pi) h_{91} h_{2,d-q}) \quad \text{in } P\text{-probability.}$$

This proves the part (b). Since $\Delta_{h,d-q}^\pi$ is a deterministic function of J_h^π , the part (c) follows. The part (d) follows similarly to (c). The part (e) is also analogous to the parts above.

SA-8.5 Theorem SA-5.3

By Lemma SA-5.1, $n^{1/2} \hat{m}^\pi \xrightarrow{d^\pi} m_h^\pi \sim N[0_{k \times 1}, \sigma^2 Q_{zz}]$ in P -probability and by (SA-8.25) $\hat{\Sigma}^\pi \xrightarrow{p^\pi} \sigma^2 Q_{zz}$ in P -probability. Thus, by Slutsky's lemma

$$\text{PAR}_2 \xrightarrow{d^\pi} m_h^{\pi'} (\sigma^2 Q_{zz})^{-1} m_h^\pi = \mathcal{Z}^{\pi'} \mathcal{Z}^\pi \sim \chi_k^2 \quad \text{in } P\text{-probability.} \quad (\text{SA-8.44})$$

By definition, $\hat{H} \hat{J}^+ \hat{U}^+ = [\hat{H} \hat{J} \hat{U}, \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi]$. As defined in (SA-5.84), the eigenvalues of $n \hat{U}^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ \hat{U}^+$ are $\hat{\kappa}_1^+, \dots, \hat{\kappa}_{d+1}^+$. Then,

$$\hat{\kappa}_{d+1}^+ = \lambda_{\min}[n(\hat{H} \hat{J} \hat{U}, \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi)' (\hat{H} \hat{J} \hat{U}, \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi)] = \lambda_{\min}[n \hat{U}^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ \hat{U}^+].$$

Now premultiply the determinantal equation

$$|n \hat{U}^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ \hat{U}^+ - \kappa I_{d+1}| = 0$$

by $|S_n^{+'} B_n^{+'} U_n^{+'} (\hat{U}^+)^{-1}|$ and postmultiply by $|(\hat{U}^+)^{-1} U_n^+ B_n^+ S_n^+|$ to obtain

$$|Q^+(\kappa)| = 0, \quad Q^+(\kappa) \equiv n S_n^{+'} B_n^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_n^+ S_n^+ - \kappa S_n^{+'} B_n^{+'} U_n^{+'} (\hat{U}^+)^{-1} (\hat{U}^+)^{-1} U_n^+ B_n^+ S_n^+. \quad (\text{SA-8.45})$$

Since $|S_n^+|, |B_n^+|, |U_n^+| > 0$ and $|\hat{U}^+| > 0$ with P^π -probability approaching 1 (wppa 1), the eigenvalues $\{\hat{\kappa}_j^+, 1 \leq j \leq d+1\}$ above also solve $|Q^+(\kappa)| = 0$ wppa 1.

Letting $S_{n,q} \equiv \text{diag}((n^{1/2} \tau_{1n})^{-1}, \dots, (n^{1/2} \tau_{qn})^{-1}) = (n^{1/2} \Upsilon_{n,q})^{-1} \in \mathbb{R}^{q \times q}$,

$$B_n^+ S_n^+ = [B_{n,q}^+ S_{n,q}, B_{n,d+1-q}^+]. \quad (\text{SA-8.46})$$

Thus, $\hat{H} \hat{J}^+ U_n^+ B_n^+ S_n^+ = [\hat{H} \hat{J}^+ U_n^+ B_{n,q}^+ S_{n,q}, \hat{H} \hat{J}^+ U_n^+ B_{n,d+1-q}^+]$, $n^{1/2} H_n \hat{J}^+ U_n^+ B_{n,q}^+ S_{n,q} = n^{1/2} H_n \hat{J} U_n B_{n,q} S_{n,q}$, and by the SVD, $H_n J_n U_n = A_n \Upsilon_n B_n'$, where $\Upsilon_n = [\text{diag}(\tau_{1n}, \dots, \tau_{dn}), 0_{d \times (k-d)}]'$. Since $A_n =$

$$[A_{n,q}, A_{n,k-q}], A_{n,q} \in \mathbb{R}^{k \times q}, A_{n,k-q} \in \mathbb{R}^{k \times (k-q)},$$

$$\begin{aligned}
n^{1/2} \hat{H} \hat{J}^+ U_n^+ B_{n,q}^+ S_{n,q} &= n^{1/2} \hat{H} H_n^{-1} H_n \hat{J} U_n B_{n,q} S_{n,q} \\
&= n^{1/2} \hat{H} H_n^{-1} H_n G_n U_n B_{n,q} S_{n,q} + n^{1/2} \hat{H} H_n^{-1} H_n (\hat{J} - G_n) U_n B_{n,q} S_{n,q} \\
&= \hat{H} H_n^{-1} A_n \Upsilon_n B_n' B_n [I_q, 0_{q \times (d-q)}]' \Upsilon_{n,q}^{-1} + n^{1/2} \hat{H} H_n^{-1} H_n (\hat{J} - G_n) U_n B_{n,q} (n^{1/2} \Upsilon_{n,q})^{-1} \\
&= A_n [I_q, 0_{q \times (d-q)}]' + o_{p^\pi}(1) \\
&= A_{n,q} + o_{p^\pi}(1) \\
&\xrightarrow{p^\pi} h_{3,q} = \Delta_{h,q} \quad \text{in } P\text{-probability,}
\end{aligned} \tag{SA-8.47}$$

where the third equality holds by the definitions of $S_{n,q}$, Υ_n and $\Upsilon_{n,q}$, the fourth equality holds by $B_n' B_n = I_d$, $\hat{J} - G_n = O_p(n^{-1/2})$ (shown in (SA-5.57)), $\hat{H} H_n^{-1} \xrightarrow{p} I_k$ which in turn holds by (SA-5.37), $H_n \rightarrow \Sigma^{-1/2}$ and the CMT, $\|H_n\| < \infty$, $U_n = O(1)$ by definition, $B_n = O(1)$ by the orthogonality of B_n , and $n^{1/2} \tau_{jn} \rightarrow \infty$ for all $j \leq q$, and the convergence uses the definition of $\Delta_{h,q}$ and Lemma SA-2.3. Using (SA-5.57), (SA-8.34), $B_{n,d-q} \rightarrow h_{2,d-q}$, $H_n \rightarrow h_{5,m}^{-1/2}$ and $U_n \rightarrow h_{91}$,

$$\begin{aligned}
n^{1/2} H_n \hat{J} U_n B_{n,d-q} &= n^{1/2} H_n G_n U_n B_{n,d-q} + n^{1/2} H_n (\hat{J} - G_n) U_n B_{n,d-q} \\
&\xrightarrow{d} h_3 \tilde{h}_{1,d-q} + h_{5,m}^{-1/2} J_h h_{91} h_{2,d-q} = \Delta_{h,d-q}.
\end{aligned} \tag{SA-8.48}$$

Then, under $H_0 : \theta = \theta_0$, using (SA-8.48) and an argument analogous to that in (SA-5.60)

$$\begin{aligned}
n^{1/2} H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ &= [n^{1/2} H_n \hat{J} U_n, H_n \hat{H}^{-1} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi] B_{n,d+1-q}^+ \\
&= [n^{1/2} H_n \hat{J} U_n B_{n,d-q}, n^{1/2} H_n \hat{H}^{-1} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi] \\
&\xrightarrow{d^\pi} [\Delta_{h,d-q}, \mathcal{Z}^\pi] \quad \text{in } P\text{-probability,}
\end{aligned} \tag{SA-8.49}$$

where the convergence holds by Slutsky's lemma, $\hat{\Sigma}^{\pi-1/2} \hat{m}^\pi \xrightarrow{d^\pi} \mathcal{Z}^\pi$ in P -probability, $\hat{H} H_n^{-1} \xrightarrow{p^\pi} I_k$ in P -probability and (SA-8.48). Define

$$\hat{E}^+ = \begin{bmatrix} E_1^+ & E_2^+ \\ E_2^{+'} & E_3^+ \end{bmatrix} \equiv B_n^{+'} U_n^{+'} (\hat{U}^+)^{-1'} (\hat{U}^+)^{-1} U_n^+ B_n^+ - I_{d+1}, \tag{SA-8.50}$$

where $E_1^+ \in \mathbb{R}^{q \times q}$, $E_2^+ \in \mathbb{R}^{q \times (d+1-q)}$ and $E_3^+ \in \mathbb{R}^{(d+1-q) \times (d+1-q)}$. By the CMT, (SA-5.38) and (SA-5.51), $\hat{U} - U_n \xrightarrow{p} 0$ with U_n positive definite, hence $(\hat{U}^+)^{-1} U_n^+ \xrightarrow{p} I_{d+1}$. Using the CMT again in conjunction with $B_n^{+'} B_n^+ = I_{d+1}$, we have $\hat{E}^+ \xrightarrow{p} 0_{(d+1) \times (d+1)}$ and by Lemma SA-2.3

$$\hat{E}^+ \xrightarrow{p^\pi} 0_{(d+1) \times (d+1)} \quad \text{in } P\text{-probability.} \tag{SA-8.51}$$

Using (SA-8.47), (SA-8.49), (SA-8.50) and the fact that $\Delta'_{h,q}\Delta_{h,q} = h'_{3,q}h_{3,q} = \lim_{n \rightarrow \infty} A'_{n,q}A_{n,q} = I_q$ in (SA-8.45) yields

$$\begin{aligned} Q^+(\kappa) &= \begin{bmatrix} nS'_{n,q}B_{n,q}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,q}^+S_{n,q} & nS'_{n,q}B_{n,q}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,d+1-q}^+ \\ nB_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,q}^+S_{n,q} & nB_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,d+1-q}^+ \end{bmatrix} - \kappa S_n^{+'}(\hat{E}^+ + I_{d+1})S_n^+ \\ &= \begin{bmatrix} I_q + o_{p^\pi}(1) - \kappa S_{n,q}E_1^+S_{n,q} - \kappa S_{n,q}^2 & n^{1/2}A'_{n,q}H_n\hat{J}^+U_n^+B_{n,d+1-q}^+ + o_{p^\pi}(1) - \kappa S_{n,q}E_2^+ \\ n^{1/2}B_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}H_nA_{n,q} + o_{p^\pi}(1) - \kappa E_2^{+'}S_{n,q} & nB_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}H_nH_n\hat{J}^+U_n^+B_{n,d+1-q}^+ - \kappa E_3^+ - \kappa \end{bmatrix}. \end{aligned}$$

If $q = 0$, $B_n^+ = B_{n,d+1-q}^+$ and

$$\begin{aligned} nB_n^{+'}\hat{U}^{+'}\hat{J}^+\hat{H}'\hat{H}\hat{J}_n^+\hat{U}^+B_n^+ &= nB_n^{+'}((U_n^+)^{-1}\hat{U}^+)'(B_n^+)^{-1'}B_n^{+'}U_n^{+'}\hat{J}^{+'}H'_n(\hat{H}H_n^{-1})' \\ &\quad (\hat{H}H_n^{-1})(H_n\hat{J}^+U_n^+B_n^+)(B_n^+)^{-1}((U_n^+)^{-1}\hat{U}^+)B_n^+ \\ &\xrightarrow{d^\pi} (\Delta_{h,d-q}, (\sigma^2 Q_{zz})^{-1/2}m_h^\pi)'(\Delta_{h,d-q}, (\sigma^2 Q_{zz})^{-1/2}m_h^\pi), \end{aligned}$$

where the convergence uses (SA-8.49), $(U_n^+)^{-1}\hat{U}^+ \xrightarrow{p^\pi} I_{d+1}$ and $\hat{H}H_n^{-1} \xrightarrow{p^\pi} I_k$ in P -probability.

Using the CMT in conjunction with the fact that the smallest eigenvalue of a matrix is a continuous function of the matrix, and $h_{3,k-q}h'_{3,k-q} = h'_3h_3 = I_k$, we obtain the result stated in the theorem.

Next, we consider the case $q \geq 1$. Using the formula of partitioned matrix determinant and provided $Q_1^+(\kappa)$ is nonsingular, $|Q^+(\kappa)| = |Q_1^+(\kappa)||Q_2^+(\kappa)|$, where

$$\begin{aligned} Q_1^+(\kappa) &\equiv I_q + o_{p^\pi}(1) - \kappa S_{n,q}E_1^+S_{n,q} - \kappa S_{n,q}^2, \\ Q_2^+(\kappa) &\equiv nB_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}H'_nH_n\hat{J}^+U_n^+B_{n,d+1-q}^+ - \kappa E_3^+ - \kappa I_{d+1-q} + o_{p^\pi}(1) \\ &\quad - [n^{1/2}B_{n,d+1-q}^{+'}U_n^{+'}\hat{J}^{+'}H_nA_{n,q} + o_{p^\pi}(1) - \kappa E_2^{+'}S_{n,q}](I_q + o_{p^\pi}(1) - \kappa S_{n,q}E_1^+S_{n,q} - \kappa S_{n,q}^2)^{-1} \\ &\quad [n^{1/2}A'_{n,q}H_n\hat{J}^+U_n^+B_{n,d+1-q}^+ + o_{p^\pi}(1) - \kappa S_{n,q}E_2^+]. \end{aligned} \tag{SA-8.52}$$

Moreover, by Lemma SA-5.2 and $E_1^+ = o_{p^\pi}(1)$,

$$\hat{\kappa}_j^+ S_{n,q}^2 = o_{p^\pi}(1), \quad \hat{\kappa}_j^+ S_{n,q}E_1^+S_{n,q} = o_{p^\pi}(1), \quad j = q+1, \dots, d+1, \tag{SA-8.53}$$

in P -probability. Thus,

$$Q_1^+(\hat{\kappa}_j) = I_q + o_{p^\pi}(1) - \hat{\kappa}_j^+ S_{n,q}E_1^+S_{n,q} - \hat{\kappa}_j^+ S_{n,q}^2 = I_q + o_{p^\pi}(1),$$

in P -probability, and so

$$|Q_1^+(\hat{\kappa}_j)| \neq 0, \quad j = q+1, \dots, d+1, \text{ wppa } 1 \text{ in } P\text{-probability.} \tag{SA-8.54}$$

Furthermore,

$$E_3^+ = o_{p^\pi}(1), \quad (\text{SA-8.55})$$

$$n^{1/2} H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ = O_{p^\pi}(1), \quad (\text{SA-8.56})$$

$$S_{n,q} = o(1), \quad (\text{SA-8.57})$$

$$E_2^+ = o_{p^\pi}(1), \quad (\text{SA-8.58})$$

$$\hat{\kappa}_j^+ E_2^{+'} S_{n,q} (I_q + o_{p^\pi}(1)) S_{n,q} E_2^+ = \hat{\kappa}_j^+ E_2^{+'} S_{n,q}^2 E_2^+ + \hat{\kappa}_{d+1}^+ E_2^{+'} S_{n,q}^2 E_2^+ o_{p^\pi}(1) = o_{p^\pi}(1), \quad (\text{SA-8.59})$$

where (SA-8.55) and (SA-8.58) follow from (SA-8.51), (SA-8.56) holds by (SA-8.49), and (SA-8.59) uses (SA-8.53) and (SA-8.58). Therefore, using the results in the previous display

$$\begin{aligned} Q_2^+(\hat{\kappa}_j^+) &= n B_{n,d+1-q}^{+'} U_n^{+'} \hat{J}^{+'} H_n' H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ + o_{p^\pi}(1) \\ &\quad - [n^{1/2} B_{n,d+1-q}^{+'} U_n^{+'} \hat{J}^{+'} H_n A_{n,q} + o_{p^\pi}(1)] (I_q + o_{p^\pi}(1)) [n^{1/2} A_{n,q}' H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ + o_{p^\pi}(1)] \\ &\quad - \hat{\kappa}_j^+ \left[I_{d+1-q} + E_3^+ - (n^{1/2} B_{n,d+1-q}^{+'} U_n^{+'} \hat{J}^{+'} H_n A_{n,q} + o_{p^\pi}(1)) (I_q + o_{p^\pi}(1)) S_{n,q} E_2^+ \right. \\ &\quad \left. - E_2^{+'} S_{n,q}' (I_q + o_{p^\pi}(1)) (n^{1/2} A_{n,q}' H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ + o_{p^\pi}(1)) + \hat{\kappa}_j^+ E_2^{+'} S_{n,q} (I_q + o_{p^\pi}(1)) S_{n,q} E_2^+ \right] \\ &= n B_{n,d+1-q}^{+'} U_n^{+'} \hat{J}^{+'} H_n' H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ + o_{p^\pi}(1) \\ &\quad - [n^{1/2} B_{n,d+1-q}^{+'} U_n^{+'} \hat{J}^{+'} H_n A_{n,q} + o_{p^\pi}(1)] (I_q + o_{p^\pi}(1)) [n^{1/2} A_{n,q}' H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ + o_{p^\pi}(1)] \\ &\quad - \hat{\kappa}_j^+ (I_{d+1-q} + o_{p^\pi}(1)) \\ &= M_{d+1-q}^+ - \hat{\kappa}_j^+ (I_{d+1-q} + o_{p^\pi}(1)), \end{aligned} \quad (\text{SA-8.60})$$

where

$$M_{d+1-q}^+ \equiv n B_{n,d+1-q}^{+'} U_n^{+'} \hat{J}^{+'} H_n' h_{3,k-q} h_{3,k-q}' H_n \hat{J}^+ U_n^+ B_{n,d+1-q}^+ + o_{p^\pi}(1). \quad (\text{SA-8.61})$$

From (SA-8.45), (SA-8.54) and $|Q^+(\kappa)| = |Q_1^+(\kappa)| |Q_2^+(\kappa)|$, $|Q_2^+(\hat{\kappa}_j^+)| = |M_{d+1-q}^+ - \hat{\kappa}_j^+ (I_{d+1-q} + o_{p^\pi}(1))| = 0, j = q+1, \dots, d+1$, wppa 1 in P -probability. Therefore, $\{\hat{\kappa}_j : j = q+1, \dots, d+1\}$ are the $d+1-q$ eigenvalues of $(I_{d+1-q} + o_{p^\pi}(1))^{-1/2} M_{d+1-q}^+ (I_{d+1-q} + o_{p^\pi}(1))^{-1/2}$ wppa 1 in P -probability. Using (SA-8.49) in (SA-8.61),

$$(I_{d+1-q} + o_{p^\pi}(1))^{-1/2} M_{d+1-q}^+ (I_{d+1-q} + o_{p^\pi}(1))^{-1/2} \xrightarrow{d^\pi} (\Delta_{h,d-q}, \mathcal{Z}^\pi)' (\Delta_{h,d-q}, \mathcal{Z}^\pi) \quad \text{in } P\text{-probability.} \quad (\text{SA-8.62})$$

On noting that the vector of eigenvalues of a matrix is continuous function of the matrix elements, invoking the CMT yields the convergence in P^π -distribution of $\{\hat{\kappa}_j : j = q + 1, \dots, d + 1\}$ to the vector of eigenvalues of $(\Delta_{h,d-q}, \mathcal{Z}^\pi)'(\Delta_{h,d-q}, \mathcal{Z}^\pi)$ in P -probability, and so

$$\lambda_{\min}[n\hat{U}^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^{+}\hat{U}^{+}] \xrightarrow{d^\pi} \lambda_{\min}[(\Delta_{h,d-q}, \mathcal{Z}^\pi)'(\Delta_{h,d-q}, \mathcal{Z}^\pi)] \quad \text{in } P\text{-probability.} \quad (\text{SA-8.63})$$

Combining (SA-8.63) with (SA-8.44) completes the proof of the asymptotic distribution of the PCLR statistic. When $q = d$, $U_n^+ B_{n,d+1-q}^+ = [0_{1 \times d}, 1]'$ and

$$n^{1/2} H_n \hat{J}^{+} U_n^+ B_{n,d+1-q}^+ = n^{1/2} H_n [\hat{J}, \hat{H}^{-1} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi] [0_{1 \times d}, 1]' = n^{1/2} H_n \hat{H}^{-1} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi \xrightarrow{d^\pi} \mathcal{Z}^\pi \quad (\text{SA-8.64})$$

in P -probability. Therefore, using (SA-8.64) in (SA-8.61), (SA-8.62) and (SA-8.63) gives

$$\lambda_{\min}[n\hat{U}^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^{+}\hat{U}^{+}] \xrightarrow{d^\pi} \mathcal{Z}^{\pi'} h_{3,k-q} h'_{3,k-q} \mathcal{Z}^\pi \sim \chi_{k-d}^2.$$

Since $h_3 h'_3 = I_k$, $\text{PCLR}_\infty = \mathcal{Z}^{\pi'} \mathcal{Z}^\pi - \mathcal{Z}^{\pi'} h_{3,k-q} h'_{3,k-q} \mathcal{Z}^\pi = \mathcal{Z}^{\pi'} h_{3,d} h'_{3,d} \mathcal{Z}^\pi \sim \chi_d^2$. The proof for subsequences follows similarly. Also, the convergence holds jointly with Lemma SA-5.1 because we only used the convergence of the permutation quantity $\hat{\mathcal{S}}^\pi \equiv n^{1/2} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi$ conditional on the convergence of other quantities computed from the observed data e.g. $n^{1/2}(\hat{J} - G_n)$ as shown in Lemma SA-5.1.

SA-8.6 Lemma SA-5.2

In view of Lemma SA-2.3, $o_p(1)$ terms are $o_{P^\pi}(1)$ in P -probability. For simplicity, we suppress the qualifier in P -probability. Also, whenever we use the convergence result $n^{1/2}(\hat{J} - G_n) = O_p(1)$ given in (SA-5.57), we are conditioning on the event $\omega \in \Omega_1$ as discussed in Section SA-5.2. The result for subsequences follow similarly, thus we provide only the full sequence result.

Recall that by definition, $h_{6,j} \equiv \lim_{n \rightarrow \infty} \tau_{(j+1)n} / \tau_{jn}$, $j = 1, \dots, d$, and $h_{6,q} = 0$ if $q < d + 1$. Define $h_{6,q} = 0$ if $q = d + 1$. Because $\tau_{1n} \geq \dots \geq \tau_{(d+1)n} \geq 0$, $h_{6,j} \in [0, 1]$. When $h_{6,j} = \lim_{n \rightarrow \infty} \tau_{(j+1)n} / \tau_{jn} > 0$, $\{\tau_{jn} : n \geq 1\}$ and $\{\tau_{(j+1)n} : n \geq 1\}$ have the same orders of magnitude. Next divide the first q SVs $\tau_{1n} \geq \dots \geq \tau_{qn}$ into groups such that the SVs within each group have the same orders of magnitude. Denote the number of groups by G_h with $q \geq G_h \geq 1$, and the first and last indices of the SVs in the g -th group by r_g and r_g^* respectively. Thus, $r_1 = 1$, $r_g^* = r_{g+1} - 1$ and $r_{G_h}^* = q$ with the definition $r_{G_h+1} \equiv q + 1$. By construction, $h_{6,j} > 0$ for all $j = r_g, \dots, r_g^* - 1$,

$g = 1, \dots, G_h$. Also, $\lim_{n \rightarrow \infty} \tau_{j'n}/\tau_{jn} = 0$ for any $j \in g$ and $j' \in g'$ with $g \neq g'$. When $d = 1$, $G_h = 1$ and $r_1 = r_1^* = 1$.

Step 1: The first group of eigenvalues.

Recall that the eigenvalues of $n\hat{U}^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+\hat{U}^+$ are denoted as $\hat{\kappa}_1^+, \dots, \hat{\kappa}_{d+1}^+$. Then,

$$\hat{\kappa}_{d+1}^+ = \lambda_{\min}[n(\hat{H}\hat{J}\hat{U}, \hat{\Sigma}^{\pi-1/2}\hat{m}^\pi)'(\hat{H}\hat{J}\hat{U}, \hat{\Sigma}^{\pi-1/2}\hat{m}^\pi)] = \lambda_{\min}[n\hat{U}^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+\hat{U}^+].$$

Now premultiply the determinantal equation

$$|n\hat{U}^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+\hat{U}^+ - \kappa I_{d+1}| = 0$$

by $|\tau_{r_1 n}^{-2} n^{-1} B_n^{+'} U_n^{+'} (\hat{U}^+)^{-1'}|$ and postmultiply by $|(\hat{U}^+)^{-1} U_n^+ B_n^+|$ to obtain

$$|\tau_{r_1 n}^{-2} B_n^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_n^+ - (n^{1/2} \tau_{r_1 n})^{-2} \kappa B_n^{+'} U_n^{+'} (\hat{U}^+)^{-1'} (\hat{U}^+)^{-1} U_n^+ B_n^+| = 0. \quad (\text{SA-8.65})$$

Since $|B_n^+|, |U_n^+| > 0$ and $|\hat{U}^+| > 0$ with P^π -probability approaching 1 (wppa 1), and $\tau_{r_1 n} > 0$ for n large given $n^{1/2} \tau_{r_1 n} \rightarrow \infty$ for $r_1 \leq q$, the eigenvalues $\{\hat{\kappa}_j^+, 1 \leq j \leq d+1\}$ above also solve (SA-8.65) wppa 1. Thus, $\{(n^{1/2} \tau_{r_1 n})^{-2} \hat{\kappa}_j^+ : 1 \leq j \leq d+1\}$ solve

$$|\tau_{r_1 n}^{-2} B_n^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_n^+ - \kappa (I_{d+1} + \hat{E}^+)| = 0, \quad (\text{SA-8.66})$$

where

$$\hat{E}^+ = \begin{bmatrix} \hat{E}_1^+ & \hat{E}_2^+ \\ \hat{E}_2^{+'} & \hat{E}_3^+ \end{bmatrix} = B_n^{+'} U_n^{+'} (\hat{U}^+)^{-1'} (\hat{U}^+)^{-1} U_n^+ B_n^+ - I_{d+1}, \quad (\text{SA-8.67})$$

where $\hat{E}_1^+ \in \mathbb{R}^{r_1^* \times r_1^*}$, $\hat{E}_2^+ \in \mathbb{R}^{r_1^* \times (d+1-r_1^*)}$ and $\hat{E}_3^+ \in \mathbb{R}^{(d+1-r_1^*) \times (d+1-r_1^*)}$.² Note that $\hat{H}\hat{J}^+ U_n^+ B_n^+ = [\hat{H}\hat{J}U_n B_n, \hat{\Sigma}^{\pi-1/2}\hat{m}^\pi]$ and $\hat{H}J_n^+ U_n^+ B_n^+ = [\hat{H}G_n U_n B_n, 0_{k \times 1}]$, and by the SVD, $H_n G_n U_n = A_n \Upsilon_n B_n'$. Let $O(\tau_{r_2 n}/\tau_{r_1 n})^{d_1 \times d_2}$ denote a $d_1 \times d_2$ matrix with $O(\tau_{r_2 n}/\tau_{r_1 n})$ elements on the main diagonal

² $\hat{E}_j^+, j = 1, 2, 3$, and $E_j^+, j = 1, 2, 3$, in (SA-8.50) are different matrices because they have different dimensions.

and zeros elsewhere. Thus,

$$\begin{aligned}
& \tau_{r_1 n}^{-1} \hat{H} \hat{J}^+ U_n^+ B_n^+ \\
&= \tau_{r_1 n}^{-1} \hat{H} J_n^+ U_n^+ B_n^+ + \tau_{r_1 n}^{-1} \hat{H} (\hat{J}^+ - J_n^+) U_n^+ B_n^+, \\
&= \tau_{r_1 n}^{-1} [\hat{H} G_n U_n B_n, 0_{k \times 1}] + (n^{1/2} \tau_{r_1 n})^{-1} [\hat{H} n^{1/2} (\hat{J} - G_n) U_n B_n, n^{1/2} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi] \\
&= \tau_{r_1 n}^{-1} [\hat{H} H_n^{-1} A_n \Upsilon_n B_n', 0_{k \times 1}] + O_{p^\pi}((n^{1/2} \tau_{r_1 n})^{-1}), \\
&= \tau_{r_1 n}^{-1} (I_k + o_{p^\pi}(1)) A_n \Upsilon_n^+ + O_{p^\pi}((n^{1/2} \tau_{r_1 n})^{-1}), \\
&= (I_k + o_{p^\pi}(1)) A_n \begin{bmatrix} h_{6, r_1^*}^* + o(1) & 0_{r_1^* \times (d-r_1^*)} & 0_{r_1^* \times 1} \\ 0_{(d-r_1^*) \times r_1^*} & O(\tau_{r_2 n} / \tau_{r_1 n})^{(d-r_1^*) \times (d-r_1^*)} & 0_{(d-r_1^*) \times 1} \\ 0_{(k-d) \times r_1^*} & 0_{(k-d) \times (d-r_1^*)} & 0_{(k-d) \times 1} \end{bmatrix} + O_{p^\pi}((n^{1/2} \tau_{r_1 n})^{-1}),
\end{aligned} \tag{SA-8.68}$$

$$\stackrel{p^\pi}{\longrightarrow} h_3 \begin{bmatrix} h_{6, r_1^*}^* & 0_{r_1^* \times (d+1-r_1^*)} \\ 0_{(k-r_1^*) \times r_1^*} & 0_{(k-r_1^*) \times (d+1-r_1^*)} \end{bmatrix}, \quad h_{6, r_1^*}^* \equiv \text{diag}(1, h_{6,1}, h_{6,1} h_{6,2}, \dots, \prod_{l=1}^{r_1^*-1} h_{6,l}),$$

where the third equality holds by $B_n' B_n = I_d$, $\hat{J} - G_n = O_p(n^{-1/2})$ as shown in (SA-5.57), $n^{1/2} \hat{\Sigma}^{\pi-1/2} \hat{m}^\pi = O_{p^\pi}(1)$ as shown in Lemma SA-5.1, $\hat{H} H_n^{-1} \stackrel{p}{\longrightarrow} I_k$, $\|H_n\| < \infty$, $U_n = O(1)$ by definition, and $B_n = O(1)$ by the orthogonality of B_n , the fifth equality uses $\tau_{jn} / \tau_{r_1 n} = \prod_{l=1}^{j-1} (\tau_{(l+1)n} / \tau_{ln}) = \prod_{l=1}^{j-1} h_{6,l} + o(1)$, $j = 2, \dots, r_1^*$, and $\tau_{jn} / \tau_{r_1 n} = O(\tau_{jn} / \tau_{r_1 n})$, $j = r_2, \dots, d+1$, which holds by $\tau_{1n} \geq \dots \tau_{(d+1)n}$, and the convergence uses $A_n \rightarrow h_3$, $\tau_{r_2 n} / \tau_{r_1 n} \rightarrow 0$ and $n^{1/2} \tau_{r_1 n} \rightarrow \infty$ since $r_1 \leq q$. Therefore, using $h_3' h_3 = I_k$

$$\begin{aligned}
\tau_{r_1 n}^{-2} B_n^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_n^+ & \stackrel{p^\pi}{\longrightarrow} \begin{bmatrix} h_{6, r_1^*}^* & 0_{r_1^* \times (d+1-r_1^*)} \\ 0_{(k-r_1^*) \times r_1^*} & 0_{(k-r_1^*) \times (d+1-r_1^*)} \end{bmatrix}' h_3' h_3 \begin{bmatrix} h_{6, r_1^*}^* & 0_{r_1^* \times (d+1-r_1^*)} \\ 0_{(k-r_1^*) \times r_1^*} & 0_{(k-r_1^*) \times (d+1-r_1^*)} \end{bmatrix} \\
&= \begin{bmatrix} h_{6, r_1^*}^{*2} & 0_{r_1^* \times (d+1-r_1^*)} \\ 0_{(d+1-r_1^*) \times r_1^*} & 0_{(d+1-r_1^*) \times (d+1-r_1^*)} \end{bmatrix}.
\end{aligned}$$

Since $\hat{U} - U_n \stackrel{p}{\longrightarrow} 0$ and $U_n \rightarrow h_{g1}$ positive definite, $\hat{U}^+ - U_n^+ \stackrel{p}{\longrightarrow} 0$ and $U_n^+ \rightarrow h_{g1}^+$. Thus, $(\hat{U}^+)^{-1} U_n^+ \stackrel{p}{\longrightarrow} I_{d+1}$ and using the CMT in conjunction with $B_n^{+'} B_n^+ = I_{d+1}$, we have

$$\hat{E}^+ = B_n^{+'} U_n^{+'} \hat{U}^{+'-1} \hat{U}^{+-1} U_n^+ B_n^+ - I_{d+1} \stackrel{p^\pi}{\longrightarrow} 0_{(d+1) \times (d+1)}. \tag{SA-8.69}$$

Thus, $\{(n^{1/2} \tau_{r_1 n})^{-2} \hat{\kappa}_j^+ : j = 1, \dots, d+1\}$ solve $|\tau_{r_1 n}^{-2} B_n^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_n^+ - \kappa I_{d+1}| = 0$ with wppa 1. Then, since the ordered vector of eigenvalues of a matrix is a continuous function of the

matrix, using Slutsky's lemma

$$(n^{-1}\hat{\kappa}_1^+, \dots, n^{-1}\hat{\kappa}_{r_1^*}^+) \xrightarrow{p^\pi} (1, h_{6,1}^2, h_{6,1}^2 h_{6,2}^2, \dots, \prod_{l=1}^{r_1^*-1} h_{6,l}^2), \quad (\text{SA-8.70})$$

$$n^{-1}\hat{\kappa}_j^+ \xrightarrow{p^\pi} \infty, \quad j = 1, \dots, r_1^*, \quad (\text{SA-8.71})$$

because $n^{1/2}\tau_{r_1 n} \rightarrow \infty$ given $r_1 \leq q$ and $h_{6,l} > 0$ for all $l = 1, \dots, r_1^* - 1$. The same argument also yields that the remaining $d + 1 - r_1^*$ eigenvalues $\{(n^{1/2}\tau_{r_1 n})^{-2}\hat{\kappa}_j^+ : j = r_1^* + 1, \dots, d + 1\}$ of $|\tau_{r_1 n}^{-2} B_n^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_n^+ - \kappa I_{d+1}| = 0$ satisfy

$$(n^{1/2}\tau_{r_1 n})^{-2}\hat{\kappa}_j^+ \xrightarrow{p^\pi} 0, \quad j = r_1^* + 1, \dots, d + 1. \quad (\text{SA-8.72})$$

Let B_{n,j_1,j_2}^+ denote the $(d + 1) \times (j_2 - j_1)$ matrix that consists of $j_1 + 1, \dots, j_2$ columns of B_n^+ for $0 \leq j_1 < j_2 \leq d + 1$. Thus, we may write $B_n^+ = [B_{n,0,r_1^*}^+, B_{n,r_1^*,d+1}^+]$. Proceeding similarly to [\(SA-8.68\)](#),

$$\begin{aligned} \tau_{r_1 n}^{-1} \hat{H}^{-1} \hat{J}^+ U_n^+ B_{n,0,r_1^*}^+ &= (I_k + o_p(1)) A_n \begin{bmatrix} h_{6,r_1^*} + o(1) \\ 0_{(k-r_1^*) \times r_1^*} \end{bmatrix} + O_{p^\pi}((n^{1/2}\tau_{r_1 n})^{-1}), \\ \tau_{r_1 n}^{-1} \hat{H}^{-1} \hat{J}^+ U_n^+ B_{n,r_1^*,d+1}^+ &= (I_k + o_p(1)) A_n \begin{bmatrix} 0_{r_1^* \times (d+1-r_1^*)} \\ O\left(\frac{\tau_{r_2 n}}{\tau_{r_1 n}}\right)^{(k-r_1^*) \times (d+1-r_1^*)} \end{bmatrix} + O_{p^\pi}((n^{1/2}\tau_{r_1 n})^{-1}). \end{aligned}$$

From the last two expressions,

$$\begin{aligned} \varrho_n &\equiv \tau_{r_1 n}^{-2} B_{n,0,r_1^*}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n,r_1^*,d+1}^+ \\ &= \begin{bmatrix} h_{6,r_1^*} + o(1) \\ 0_{(k-r_1^*) \times r_1^*} \end{bmatrix}' A_n' (I_k + o_p(1)) A_n \begin{bmatrix} 0_{r_1^* \times (d+1-r_1^*)} \\ O\left(\frac{\tau_{r_2 n}}{\tau_{r_1 n}}\right)^{(k-r_1^*) \times (d+1-r_1^*)} \end{bmatrix} + O_{p^\pi}((n^{1/2}\tau_{r_1 n})^{-1}) \\ &= \begin{bmatrix} h_{6,r_1^*} + o(1) \\ 0_{(k-r_1^*) \times r_1^*} \end{bmatrix}' \begin{bmatrix} I_{r_1^*} + o_{p^\pi}(1) & o_{p^\pi}(1)^{r_1^* \times (k-r_1^*)} \\ o_{p^\pi}(1)^{(k-r_1^*) \times r_1^*} & I_{k-r_1^*} + o_{p^\pi}(1) \end{bmatrix} \begin{bmatrix} 0_{r_1^* \times (d+1-r_1^*)} \\ O\left(\frac{\tau_{r_2 n}}{\tau_{r_1 n}}\right)^{(k-r_1^*) \times (d+1-r_1^*)} \end{bmatrix} + O_{p^\pi}((n^{1/2}\tau_{r_1 n})^{-1}) \\ &= o_{p^\pi}(\tau_{r_2 n}/\tau_{r_1 n}) + O_{p^\pi}((n^{1/2}\tau_{r_1 n})^{-1}). \end{aligned} \quad (\text{SA-8.73})$$

Define

$$\begin{aligned} \hat{\xi}_1(\kappa) &\equiv \tau_{r_1 n}^{-2} B_{n,0,r_1^*}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n,0,r_1^*}^+ - \kappa(I_{r_1^*} + \hat{E}_1^+) \in \mathbb{R}^{r_1^* \times r_1^*}, \\ \hat{\xi}_2(\kappa) &\equiv \varrho_n - \kappa \hat{E}_2^+ \in \mathbb{R}^{r_1^* \times (d+1-r_1^*)}, \\ \hat{\xi}_3(\kappa) &\equiv \tau_{r_1 n}^{-2} B_{n,r_1^*,d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n,r_1^*,d+1}^+ - \kappa(I_{d+1-r_1^*} + \hat{E}_1^+) \in \mathbb{R}^{d+1-r_1^* \times (d+1-r_1^*)}. \end{aligned}$$

Next we verify that $(n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+, j = r_1^* + 1, \dots, d+1$, cannot solve the determinantal equation $|\hat{\xi}_1(\kappa)| = 0$ with wppa 1. First, note that

$$\tau_{r_1n}^{-2}B_{n,0,r_1^*}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,0,r_1^*}^+ \xrightarrow{p^\pi} h_{6,r_1^*}^2. \quad (\text{SA-8.74})$$

Then, using (SA-8.69), (SA-8.72) and (SA-8.74), for $j = r_1^* + 1, \dots, d+1$,

$$\begin{aligned} \tilde{\xi}_{j1} &\equiv \hat{\xi}_1((n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+) \\ &= \tau_{r_1n}^{-2}B_{n,0,r_1^*}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,0,r_1^*}^+ - (n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+(I_{r_1^*} + \hat{E}_1^+) \\ &= h_{6,r_1^*}^{*2} + o_{p^\pi}(1) - o_{p^\pi}(1)(I_{r_1^*} + o_{p^\pi}(1)) \\ &= h_{6,r_1^*}^{*2} + o_{p^\pi}(1). \end{aligned} \quad (\text{SA-8.75})$$

Note that $\lambda_{\min}(h_{6,r_1^*}^{*2}) > 0$ because $h_{6,l} > 0$ for all $l = 1, \dots, r_1^* - 1$. Hence,

$$|\hat{\xi}_1((n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+)| \neq 0 \quad (\text{SA-8.76})$$

wppa 1. Using the determinant formula for partitioned matrices,

$$0 = |\tau_{r_1n}^{-2}B_{n,0,r_1^*}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,0,r_1^*}^+ - \kappa(I_{d+1} + \hat{E}^+)| \quad (\text{SA-8.77})$$

$$\begin{aligned} &= \left| \begin{bmatrix} \hat{\xi}_1(\kappa) & \hat{\xi}_2(\kappa) \\ \hat{\xi}_2(\kappa)' & \hat{\xi}_3(\kappa) \end{bmatrix} \right| \\ &= |\hat{\xi}_1(\kappa)| |\hat{\xi}_{3n}(\kappa) - \hat{\xi}_2(\kappa)' \hat{\xi}_1(\kappa)^{-1} \hat{\xi}_2(\kappa)| \\ &= |\hat{\xi}_1(\kappa)| |\tau_{r_1n}^{-2}B_{n,r_1^*,d+1}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,r_1^*,d+1}^+ - \varrho_n' \hat{\xi}_1(\kappa)^{-1} \varrho_n - \kappa(I_{d+1-r_1^*} + \hat{E}_3^+ - \hat{E}_2^{+'} \hat{\xi}_1(\kappa)^{-1} \varrho_n \\ &\quad - \varrho_n' \hat{\xi}_1(\kappa)^{-1} \hat{E}_2^+ + \kappa \hat{E}_2^{+'} \hat{\xi}_1(\kappa)^{-1} \hat{E}_2^+)|, \end{aligned} \quad (\text{SA-8.78})$$

for κ equal to any solution $(n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+$ to the equation (SA-8.77). From (SA-8.76), $(n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+, j = r_1^* + 1, \dots, d+1$, solve $|\hat{\xi}_3(\kappa) - \hat{\xi}_2(\kappa)' \hat{\xi}_1(\kappa)^{-1} \hat{\xi}_2(\kappa)| = 0$. Thus,

$$\begin{aligned} &|\tau_{r_1n}^{-2}B_{n,r_1^*,d+1}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,r_1^*,d+1}^+ - o((\tau_{r_2n}/\tau_{r_1n})^2) + O_{p^\pi}((n^{1/2}\tau_{r_1n})^{-2}) \\ &\quad - (n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+(I_{d+1-r_1^*} + \hat{E}_{j2})| = 0, \end{aligned}$$

where $\hat{E}_{j2} \equiv \hat{E}_3 - \hat{E}_2' \tilde{\xi}_{j1}^{-1} \varrho_n - \varrho_n' \tilde{\xi}_{j1} \hat{E}_2 + (n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+ \hat{E}_2' \tilde{\xi}_{j1}^{-1} \hat{E}_2 \in \mathbb{R}^{(d+1-r_1^*) \times (d+1-r_1^*)}$. Multiplying the equation in the previous display by $\tau_{r_2n}^{-2}/\tau_{r_1n}^{-2}$ and using $O_{p^\pi}((n^{1/2}\tau_{r_1n})^{-2}) = o_{p^\pi}(1)$ which follows from $r_2 \leq q$ and $n^{1/2}\tau_{jn} \rightarrow \infty$ for all $j \leq q$,

$$|\tau_{r_2n}^{-2}B_{n,r_1^*,d+1}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,r_1^*,d+1}^+ + o_{p^\pi}(1) - (n^{1/2}\tau_{r_2n})^{-2}\hat{\kappa}_j^+(I_{d+1-r_1^*} + \hat{E}_{j2})| = 0, \quad (\text{SA-8.79})$$

Thus, $\{(n^{1/2}\tau_{r_2n})^{-2}\hat{\kappa}_j^+ : j = r_1^* + 1, \dots, d+1\}$ solve

$$|\tau_{r_2n}^{-2}B_{n,r_1^*,d+1}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,r_1^*,d+1}^+ + o_{p^\pi}(1) - \kappa(I_{d+1-r_1^*} + \hat{E}_{j2})| = 0. \quad (\text{SA-8.80})$$

On noting that $\hat{E}_2 = o_{p^\pi}(1)$ and $\hat{E}_3 = o_{p^\pi}(1)$ by (SA-8.69), $\tilde{\xi}_{j1}^{-1} = O_{p^\pi}(1)$ by (SA-8.75), and $\varrho_n = o_{p^\pi}(1)$ using (SA-8.73), $\tau_{r_2n} \leq \tau_{r_1n}, n^{1/2}\tau_{r_1n} \rightarrow 0$, and $(n^{1/2}\tau_{r_1n})^{-2}\hat{\kappa}_j^+ = o_{p^\pi}(1)$ for $j = r_1^* + 1, \dots, d+1$ by (SA-8.72),

$$\hat{E}_{j2} = o_{p^\pi}(1). \quad (\text{SA-8.81})$$

Now the same argument is repeated for the subsequent groups of indices. Specifically, replace (SA-8.66) and (SA-8.69) by (SA-8.80) and (SA-8.81), and replace $j = r_1^* + 1, \dots, d+1$, \hat{E} , B_n^+ , τ_{r_1n} , τ_{r_2n} , r_1^* , $d+1-r_1^*$, $h_{6,r_1^*}^+$, $B_{n,0,r_1^*}^+$, and $B_{n,r_1^*,d+1}^+$ by $j = r_2^* + 1, \dots, d+1$, \hat{E}_{j2} , $B_{n,d+1-r_1^*}^+$, τ_{r_2n} , τ_{r_3n} , $r_2^* - r_1^*$, $d+1-r_2^*$, $h_{6,r_2^*}^+ \equiv \text{diag}(1, h_{6,r_1^*+1}, h_{6,r_1^*+1}h_{6,r_1^*+2}, \dots, \prod_{l=r_1^*+1}^{r_2^*-1} h_{6,l}) \in \mathbb{R}^{(r_2^*-r_1^*) \times (r_2^*-r_1^*)}$, $B_{n,r_1^*,r_2^*}^+$, and $B_{n,r_2^*,d+1}^+$, respectively. Moreover, \hat{E}_{j2} is partitioned further as

$$\hat{E}_{j2} = \begin{bmatrix} \hat{E}_{1j2} & \hat{E}_{2j2} \\ \hat{E}'_{2j2} & \hat{E}_{3j2} \end{bmatrix}, \quad \hat{E}_{1j2} \in \mathbb{R}^{r_2^* \times r_2^*}, \hat{E}_{2j2} \in \mathbb{R}^{r_2^* \times (d+1-r_1^*-r_2^*)}, \hat{E}_{3j2} \in \mathbb{R}^{(d+1-r_1^*-r_2^*) \times (d+1-r_1^*-r_2^*)}.$$

The analog of (SA-8.80) is that $\{(n^{1/2}\tau_{r_3n})^{-2}\hat{\kappa}_j^+ : j = r_2^* + 1, \dots, d+1\}$ solve

$$|\tau_{r_3n}^{-2}B_{n,r_2^*,d+1}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,r_2^*,d+1}^+ + o_{p^\pi}(1) - \kappa(I_{d+1-r_2^*} + \hat{E}_{j3})| = 0, \quad (\text{SA-8.82})$$

where

$$\begin{aligned} \hat{E}_{j3} &\equiv \hat{E}_{3j2} - \hat{E}'_{2j2}\tilde{\xi}_{1j2}^{-1}\varrho_{2n} - \varrho'_{2n}\tilde{\xi}_{1j2}^{-1}\hat{E}_{2j2} + (n^{1/2}\tau_{r_2n})^{-2}\hat{\kappa}_j^+\hat{E}'_{2j2}\tilde{\xi}_{1j2}^{-1}\hat{E}_{2j2}, \\ \tilde{\xi}_{1j2} &\equiv \hat{\xi}_{1j2}((n^{1/2}\tau_{r_2n})^{-2}\hat{\kappa}_j^+) \\ &= \tau_{r_2n}^{-2}B_{n,r_1^*,r_2^*}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,r_1^*,r_2^*}^+ - (n^{1/2}\tau_{r_2n})^{-2}\hat{\kappa}_j^+(I_{r_2^*-r_1^*} + \hat{E}_{1j2}) \\ &= h_{6,r_2^*}^{*2} + o_{p^\pi}(1) - o_{p^\pi}(1)(I_{r_2^*-r_1^*} + o_{p^\pi}(1)) \\ &= h_{6,r_2^*}^{*2} + o_{p^\pi}(1), \\ \varrho_{2n} &\equiv \tau_{r_2n}^{-2}B_{n,r_1^*,r_2^*}^{+'}U_n^{+'}\hat{J}^{+'}\hat{H}'\hat{H}\hat{J}^+U_n^+B_{n,r_2^*,d+1}^+. \end{aligned}$$

Analogously to (SA-8.70), (SA-8.71) and (SA-8.72), it then follows that

$$\hat{\kappa}_j^+ \xrightarrow{p^\pi} \infty, \quad j = r_2, \dots, r_2^*, \quad (\text{SA-8.83})$$

$$(n^{1/2}\tau_{r_2n})^{-2}\hat{\kappa}_j^+ = o_{p^\pi}(1), \quad j = r_2^* + 1, \dots, d+1. \quad (\text{SA-8.84})$$

Then, repeating the previous steps $G_h - 2$ more times gives (as verified below)

$$\hat{\kappa}_j^+ \xrightarrow{p^\pi} \infty, \quad j = 1, \dots, r_{G_h}^*, \quad (\text{SA-8.85})$$

$$(n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j^+ = o_{p^\pi}(1), \quad j = r_g^* + 1, \dots, d+1, \quad g = 1, \dots, G_h. \quad (\text{SA-8.86})$$

Since $r_{G_h}^* = q$, the part (a) of the lemma follows from (SA-8.84). Setting $g = G_h$ in (SA-8.86) and noting that $r_{G_h}^* = q$ yield

$$(n^{1/2} \tau_{r_{G_h} n})^{-2} \hat{\kappa}_j^+ = o_{p^\pi}(1), \quad j = q+1, \dots, d+1. \quad (\text{SA-8.87})$$

If $r_{G_h} = r_{G_h}^* = q$, from (SA-8.87) $(n^{1/2} \tau_{qn})^{-2} \hat{\kappa}_j^+ = o_{p^\pi}(1)$ for $j = q+1, \dots, d+1$. If $r_{G_h} < r_{G_h}^* = q$, using $h_{6,l} > 0$ for all $l = r_{G_h}, \dots, r_{G_h}^* - 1$,

$$\lim_{n \rightarrow \infty} \frac{\tau_{qn}}{\tau_{r_{G_h} n}} = \lim_{n \rightarrow \infty} \frac{\tau_{r_{G_h}^* n}}{\tau_{r_{G_h} n}} = \prod_{j=r_{G_h}}^{r_{G_h}^*-1} h_{6,j} > 0. \quad (\text{SA-8.88})$$

Combining (SA-8.87) and (SA-8.88), $(n^{1/2} \tau_{qn})^{-2} \hat{\kappa}_j^+ = o_{p^\pi}(1)$ for $j = q+1, \dots, d+1$. On noting that $\tau_{ln} \geq \tau_{qn}$ for $l \leq q$, the part (b) of the lemma follows. The result for all subsequences $\{w_n\}$ and $\{\lambda_{w_n, h} \in \Lambda : n \geq 1\}$ holds analogously. The formal induction proof of (SA-8.85) and (SA-8.86) are provided next.

Step 2: Using induction to complete the proof.

Denote o_{gp} denote a symmetric $(d+1 - r_{g-1}^*) \times (d+1 - r_{g-1}^*)$ matrix whose $(l, m), l, m = 1, \dots, d+1 - r_{g-1}^*$, element is $o_{p^\pi}(\tau_{(r_g^*+l)n} \tau_{(r_g^*+l)n} / \tau_{r_g n}^2) + O_{p^\pi}((n^{1/2} \tau_{r_g n})^{-1})$. $o_{gp} = o_{p^\pi}(1)$ because $r_{g-1}^* + l \geq r_g$ for $l \geq 1$ as τ_{jn} 's are nonincreasing in j , and $n^{1/2} \tau_{r_g n} \rightarrow \infty$ for $g = 1, \dots, G_h$. The goal is to show that $\{(n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j^+ : j = r_{g-1}^* + 1, \dots, d+1\}$ solve

$$|\tau_{r_g n}^{-2} B_{n, r_{g-1}^*, d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n, r_{g-1}^*, d+1}^+ + o_{gp} - \kappa(I_{d+1-r_{g-1}^*} + \hat{E}_{jg})| = 0, \quad (\text{SA-8.89})$$

for some $(d+1 - r_{g-1}^*) \times (d+1 - r_{g-1}^*)$ symmetric matrices $\hat{E}_{jg} = o_{p^\pi}(1)$ and o_{gp} . This is shown using induction over $g = 1, \dots, G_h$. For $g = 1$, the above holds upon noting that $\hat{E}_{jg} = \hat{E}$, $o_{gp} = 0$, $r_{g-1}^* = r_0^* \equiv 0$ and $B_{n, r_{g-1}^*, d+1}^+ = B_{n, 0, d+1}^+ = B_n^+$. Now suppose that (SA-8.82) holds for g with

$\hat{E}_{jg} = o_{p^\pi}(1)$ and o_{gp} . Proceeding similarly to the argument in (SA-8.68)

$$\begin{aligned} & \tau_{r_g n}^{-1} \hat{H} \hat{J}^+ U_n^+ B_{n, r_{g-1}^*, d+1}^+ \\ &= \tau_{r_g n}^{-1} (I_k + o_{p^\pi}(1)) A_n \begin{bmatrix} 0_{r_{g-1}^* \times (d+1-r_g^*)} \\ \text{diag}(\tau_{r_g n}, \dots, \tau_{(d+1)n}) \\ 0_{(k-d-1) \times (d+1-r_{g-1}^*)} \end{bmatrix} + O_{p^\pi}((n^{1/2} \tau_{r_g n})^{-1}), \quad (\text{SA-8.90}) \\ & \xrightarrow{p^\pi} h_3 \begin{bmatrix} 0_{r_{g-1}^* \times (r_g^* - r_{g-1}^*)} & 0_{r_{g-1}^* \times (d+1-r_g^*)} \\ h_{6, r_g^*}^* & 0_{(r_g^* - r_{g-1}^*) \times (d+1-r_g^*)} \\ 0_{(k-r_g^*) \times (r_g^* - r_{g-1}^*)} & 0_{(k-r_g^*) \times (d+1-r_g^*)} \end{bmatrix}, \quad h_{6, r_g^*}^* \equiv \text{diag}(1, h_{6, r_g^*}, \dots, \prod_{l=1}^{r_g^*-1} h_{6, l}) \in \mathbb{R}^{(r_g^* - r_{g-1}^*) \times (r_g^* - r_{g-1}^*)}, \end{aligned}$$

and $h_{r_g^*}^* \equiv 1$ for $r_g^* = 1$. From (SA-8.90) and $h_3' h_3 = I_k = \lim_{n \rightarrow \infty} A_n' A_n$,

$$\tau_{r_g n}^{-2} B_{n, r_{g-1}^*, d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n, r_{g-1}^*, d+1}^+ \xrightarrow{p^\pi} \begin{bmatrix} h_{6, r_g^*}^{*2} & 0_{(r_g^* - r_{g-1}^*) \times (d+1-r_g^*)} \\ 0_{(d+1-r_g^*) \times (r_g^* - r_{g-1}^*)} & 0_{(d+1-r_g^*) \times (d+1-r_g^*)} \end{bmatrix}. \quad (\text{SA-8.91})$$

Using (SA-8.82) which is assumed to hold in the induction step and $o_{gp} = o_{p^\pi}(1)$, wppa 1, $\{(n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j : j = r_{g-1}^* + 1, \dots, d+1\}$ solve

$$|(I_{d+1-r_{g-1}^*} + \hat{E}_{jg})^{-1} \tau_{r_g n}^{-2} B_{n, r_{g-1}^*, d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n, r_{g-1}^*, d+1}^+ + o_{p^\pi}(1) - \kappa I_{d+1-r_{g-1}^*}| = 0. \quad (\text{SA-8.92})$$

By the same arguments that led to (SA-8.70)-(SA-8.72), (SA-8.91), (SA-8.92) and the induction assumption that $\hat{E}_{jg} = o_{p^\pi}(1)$,

$$\hat{\kappa}_j^+ \xrightarrow{p^\pi} \infty, \quad j = r_{g-1}^*, \dots, r_g^*, \quad (\text{SA-8.93})$$

$$(n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j^+ = o_{p^\pi}(1), \quad j = r_g^* + 1, \dots, d+1. \quad (\text{SA-8.94})$$

Let o_{gp}^* denote an $(r_g^* - r_{g-1}^*) \times (d+1-r_g^*)$ matrix whose elements in column $j, j = 1, \dots, d+1-r_g^*$, are $o_{p^\pi}(\tau_{(r_g^*+j)n}/\tau_{r_g n}) + O_{p^\pi}((n^{1/2} \tau_{r_g n})^{-1})$. Then $o_{gp}^* = o_{p^\pi}(1)$. Next replacing $B_{n, r_{g-1}^*, d+1}^+$ in (SA-8.68)

with $B_{n,r_g^*,d+1}$ and B_{n,r_{g-1}^*,r_g^*} , and multiplying the resulting matrices give

$$\begin{aligned}
\varrho_{gn} &\equiv \tau_{r_g n}^{-2} B_{n,r_{g-1}^*,r_g^*}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n,r_g^*,d+1}^+ \\
&= \begin{bmatrix} 0_{r_{g-1}^* \times (r_g^* - r_{g-1}^*)} \\ \text{diag}(\tau_{(r_{g-1}+1)n}, \dots, \tau_{r_g^* n}) / \tau_{r_g n} \\ 0_{(k-r_g^*) \times (r_g^* - r_{g-1}^*)} \end{bmatrix}' A'_n (I_k + o_{p^\pi}(1)) A_n \begin{bmatrix} 0_{r_{g-1}^* \times (r_g^* - r_{g-1}^*)} \\ \text{diag}(\tau_{(r_{g-1}+1)n}, \dots, \tau_{r_g^* n}) / \tau_{r_g n} \\ 0_{(k-r_g^*) \times (r_g^* - r_{g-1}^*)} \end{bmatrix} \\
&\quad + O_{p^\pi}((n^{1/2} \tau_{r_g n})^{-1}) \\
&= o_{gp}^* \in \mathbb{R}^{(r_g^* - r_{g-1}^*) \times (d+1-r_g^*)}, \tag{SA-8.95}
\end{aligned}$$

where $\text{diag}(\tau_{(r_{g-1}+1)n}, \dots, \tau_{r_g^* n}) / \tau_{r_g n} = h_{6,r_g^*}^* + o(1)$, and the last equality uses $A'_n (I_k + o_{p^\pi}(1)) A_n = I_k + o_{p^\pi}(1)$. Partition the $(d+1-r_{g-1}^*) \times (d+1-r_g^*)$ matrices o_{gp} and \hat{E}_{jg} as:

$$o_{gp} = \begin{bmatrix} o_{1gp} & o_{2gp} \\ o'_{2gp} & o_{3gp} \end{bmatrix}, \quad \hat{E}_{jg} = \begin{bmatrix} \hat{E}_{1jg} & \hat{E}_{2jg} \\ \hat{E}'_{2jg} & \hat{E}_{3jg} \end{bmatrix}, \quad o_{1gp}, \hat{E}_{1jg} \in \mathbb{R}^{(r_g^* - r_{g-1}^*) \times (r_g^* - r_{g-1}^*)},$$

$o_{2gp}, \hat{E}_{2jg} \in \mathbb{R}^{(r_g^* - r_{g-1}^*) \times (d+1-r_g^*)}$, $o_{3gp}, \hat{E}_{3jg} \in \mathbb{R}^{(d+1-r_g^*) \times (d+1-r_g^*)}$, $j = r_{g-1}^* + 1, \dots, d+1$ and $g = 1, \dots, G_h$.

Define

$$\begin{aligned}
\hat{\xi}_{1jg}(\kappa) &\equiv \tau_{r_g n}^{-2} B_{n,r_{g-1}^*,r_g^*}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n,r_{g-1}^*,r_g^*}^+ + o_{1gp} - \kappa(I_{r_g^* - r_{g-1}^*} + \hat{E}_{1jg}) \in \mathbb{R}^{(r_g^* - r_{g-1}^*) \times (r_g^* - r_{g-1}^*)}, \\
\hat{\xi}_{2jg}(\kappa) &\equiv \varrho_{gn} + o_{2gp} - \kappa \hat{E}_{2jg} \in \mathbb{R}^{(r_g^* - r_{g-1}^*) \times (d+1-r_g^*)}, \\
\hat{\xi}_{3jg}(\kappa) &\equiv \tau_{r_g n}^{-2} B_{n,r_g^*,d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n,r_g^*,d+1}^+ + o_{3gp} - \kappa(I_{d+1-r_g^*} + \hat{E}_{3jg}) \in \mathbb{R}^{d+1-r_g^* \times (d+1-r_g^*)}.
\end{aligned}$$

From (SA-8.82), $\{(n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j : j = r_{g-1}^* + 1, \dots, d+1\}$ solve

$$0 = |\tau_{r_g n}^{-2} B_{n,r_{g-1}^*,d+1}^{+'} U_n^{+'} \hat{J}_n^{+'} \hat{H}' \hat{H} \hat{J}_n^+ U_n^+ B_{n,r_{g-1}^*,d+1}^+ + o_{gp} - \kappa(I_{d+1-r_{g-1}^*} + \hat{E}_{jgn})| \tag{SA-8.96}$$

$$\begin{aligned}
&= \left| \begin{bmatrix} \hat{\xi}_{1jg}(\kappa) & \hat{\xi}_{2jg}(\kappa) \\ \hat{\xi}_{2jg}(\kappa)' & \hat{\xi}_{3jg}(\kappa) \end{bmatrix} \right| \\
&= |\hat{\xi}_{1jg}(\kappa)| |\hat{\xi}_{3jg}(\kappa) - \hat{\xi}_{2jg}(\kappa)' \hat{\xi}_{1jg}(\kappa)^{-1} \hat{\xi}_{2jg}(\kappa)| \\
&= |\hat{\xi}_{1jg}(\kappa)| |\tau_{r_g n}^{-2} B_{n,r_g^*,d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^+ U_n^+ B_{n,r_g^*,d+1}^+ + o_{3gp} - (\varrho_{gn} + o_{2gp})' \hat{\xi}_{1jgn}(\kappa)^{-1} (\varrho_{gn} + o_{2gp}) \\
&\quad - \kappa(I_{d+1-r_g^*} + \hat{E}_{3jg} - \hat{E}'_{2jg} \hat{\xi}_{1jg}(\kappa)^{-1} (\varrho_{gn} + o_{2gp})) \\
&\quad - (\varrho_{gn} + o_{2gp})' \hat{\xi}_{1jg}(\kappa)^{-1} \hat{E}_{2jg} + \kappa \hat{E}'_{2jg} \hat{\xi}_{1jg}(\kappa)^{-1} \hat{E}_{2jg}|, \tag{SA-8.97}
\end{aligned}$$

where the second equality holds by the formula of partitioned matrix determinant provided

$$\hat{\xi}_{1jg} \left((n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j \right), j = r_{g-1}^* + 1, \dots, d+1 \}$$

is nonsingular wppa 1, a fact that is verified below. By the same argument as in (SA-8.75), $\hat{E}_{1jg} = o_{p^\pi}(1)$ which holds by definition, and $o_{gp} = o_{p^\pi}(1)$,

$$\tilde{\xi}_{1jg} \equiv \hat{\xi}_{1jg}((n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j) = h_{6,r_g^*}^{*2} + o_{p^\pi}(1). \quad (\text{SA-8.98})$$

Since $\lambda_{\min}(h_{6,r_g^*}^{*2}) > 0$, $\hat{\xi}_{1jg n}((n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j), j = r_{g-1}^* + 1, \dots, d+1$, is nonsingular wppa 1. Therefore,

$$\begin{aligned} 0 = & |\tau_{r_g n}^{-2} B_{n,r_g^*,d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^{+'} U_n^{+'} B_{n,r_g^*,d+1}^{+} + o_{3gp} - (\varrho_{gn} + o_{2gp})' \tilde{\xi}_{1jg}^{-1} (\varrho_{gn} + o_{2gp}) \\ & + (n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j^{+} (I_{d+1-r_g^*} + \hat{E}_{j(g+1)n})|, \end{aligned} \quad (\text{SA-8.99})$$

where

$$\begin{aligned} \hat{E}_{j(g+1)} \equiv & \hat{E}_{3jg} - \hat{E}_{2jg}' \tilde{\xi}_{j1g} (\varrho_{gn} + o_{2gp}) - (\varrho_{gn} + o_{2gp})' \tilde{\xi}_{j1g} \hat{E}_{2jg} \\ & + (n^{1/2} \tau_{r_g n})^{-2} \hat{\kappa}_j^{+} \hat{E}_{2jg}' \tilde{\xi}_{1jg}^{-1} \hat{E}_{2jg} \in \mathbb{R}^{(d+1-r_g^*) \times (d+1-r_g^*)}. \end{aligned} \quad (\text{SA-8.100})$$

Furthermore,

$$\begin{aligned} o_{3gp} - (\varrho_{gn} + o_{2gp})' \tilde{\xi}_{1jg}^{-1} (\varrho_{gn} + o_{2gp}) &= o_{3gp} - (o_{gp}^* + o_{2gp})' (h_{6,r_g^*}^{*-2} + o_{p^\pi}(1)) (o_{gp}^* + o_{2gp}) \\ &= o_{3gp} - o_{gp}^{*'} o_{gp}^* \\ &= (\tau_{r_{g+1}n}^2 / \tau_{r_g n}^2) o_{(g+1)p}, \end{aligned} \quad (\text{SA-8.101})$$

where

1. the first equality follows from (SA-8.95) and (SA-8.98);
2. the second equality uses $o_{2gp} = o_{gp}^*$. The latter holds because the $(j, m), j = 1, \dots, r_g^* - r_{g-1}^*, m = 1, \dots, d+1 - r_g^*$ element is

$$o_{p^\pi}(\tau_{(r_{g-1}^*+j)n} \tau_{(r_g^*+m)n} / \tau_{r_g n}^2) + O_{p^\pi}((n^{1/2} \tau_{r_g n})^{-1}) = o_{p^\pi}(\tau_{(r_g^*+m)n} / \tau_{r_g n}) + O_{p^\pi}((n^{1/2} \tau_{r_g n})^{-1}),$$

since $r_{g-1}^* + j \geq r_g$ and $(h_{6,r_g^*}^{*-2} + o_{p^\pi}(1)) o_{gp}^* = o_{gp}^*$ which in turn holds because $h_{6,r_g^*}^{*2}$ is diagonal and $\lambda_{\min}(h_{6,r_g^*}^{*2}) > 0$;

3. the third equality uses the fact that the $(j, m), j, m = 1, \dots, d+1-r_g^*$, element of $(\tau_{r_g n}^2/\tau_{r_{g+1} n}^2)o_{3gp}$ is of order

$$\begin{aligned} & o_{p^\pi}(\tau_{(r_g^*+j)n}\tau_{(r_g^*+m)n}/\tau_{r_g n}^2)(\tau_{r_g n}^2/\tau_{r_{g+1} n}^2) + O_{p^\pi}((n^{1/2}\tau_{r_g n})^{-1})(\tau_{r_g n}^2/\tau_{r_{g+1} n}^2) \\ &= o_{p^\pi}(\tau_{(r_g^*+j)n}\tau_{(r_g^*+m)n}/\tau_{r_{g+1} n}^2) + O_{p^\pi}((n^{1/2}\tau_{r_{g+1} n})^{-1})(\tau_{r_g n}/\tau_{r_{g+1} n}) \end{aligned}$$

which is the same order as the (j, m) element of $o_{(g+1)p}$ using $\tau_{r_g n}/\tau_{r_{g+1} n} \leq 1$;

4. the third equality also uses the fact that the $(j, m), j, m = 1, \dots, d+1-r_g^*$, element of $(\tau_{r_g n}^2/\tau_{r_{g+1} n}^2)o_{gp}^*o_{gp}^*$ is the sum of two terms that are of orders

$$\begin{aligned} & o_{p^\pi}(\tau_{(r_g^*+j)n}\tau_{(r_g^*+m)n}/\tau_{r_g n}^2)(\tau_{r_g n}^2/\tau_{r_{g+1} n}^2) = o_{p^\pi}(\tau_{(r_g^*+j)n}\tau_{(r_g^*+m)n}/\tau_{r_{g+1} n}^2), \\ & O_{p^\pi}((n^{1/2}\tau_{r_g n})^{-2})(\tau_{r_g n}^2/\tau_{r_{g+1} n}^2) = O_{p^\pi}((n^{1/2}\tau_{r_{g+1} n})^{-2}), \end{aligned}$$

respectively. Thus, $(\tau_{r_g n}^2/\tau_{r_{g+1} n}^2)o_{gp}^*o_{gp}^*$ is $o_{(g+1)p}$.

For $j = r_g^* + 1, \dots, d+1$, since $\hat{E}_{2jg} = o_{p^\pi}(1)$ and $\hat{E}_{3jg} = o_{p^\pi}(1)$ by (SA-8.82), $\tilde{\xi}_{1jg} = O_{p^\pi}(1)$ by (SA-8.98), $\varrho_{gn} + o_{2gp} = o_{p^\pi}(1)$ by (SA-8.95) and since $o_{gp}^* = o_{p^\pi}(1)$ and $(n^{1/2}\tau_{r_g n})^{-2}\hat{\kappa}_j^+ = o_{p^\pi}(1)$ by (SA-8.84), it follows that

$$\hat{E}_{j(g+1)} = o_{p^\pi}(1). \quad (\text{SA-8.102})$$

Using (SA-8.101) and (SA-8.102) in (SA-8.99),

$$|\tau_{r_{g+1} n}^{-2} B_{n, r_g^*, d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^{+'} U_n^{+} B_{n, r_g^*, d+1}^{+} + o_{(g+1)p} - (n^{1/2}\tau_{r_{g+1} n})^{-2} \hat{\kappa}_j^+ (I_{d+1-r_g^*} + \hat{E}_{j(g+1)})| = 0, \quad (\text{SA-8.103})$$

Thus, wppa 1 $\{(n^{1/2}\tau_{r_{g+1} n})^{-2} \hat{\kappa}_j^+ : j = r_{g+1}, \dots, d+1\}$ solve

$$|\tau_{r_{g+1} n}^{-2} B_{n, r_g^*, d+1}^{+'} U_n^{+'} \hat{J}^{+'} \hat{H}' \hat{H} \hat{J}^{+'} U_n^{+} B_{n, r_g^*, d+1}^{+} + o_{(g+1)p} - \kappa(I_{d+1-r_g^*} + \hat{E}_{j(g+1)})| = 0. \quad (\text{SA-8.104})$$

This completes the induction step and (SA-8.82) holds for all $g = 1, \dots, G_h$.

SA-9 Additional simulation evidence

SA-9.1 Role of studentization

Section 3.2 of the main paper highlights the difference between the independence and the standard exclusion restriction assumption in homoskedastic setting. The next simulation examines the role of

studentization for obtaining valid permutation tests. The data generating process is the same as in the main paper. In addition to the tests, for the just-identified design with $k = 1$, we consider a non-studentized randomization test, labelled as PNS, that rejects when $W'\tilde{u}(\theta_0)$, $\tilde{u}(\theta_0) = M_l(y - Y\theta_0)$, is below 0.25 or above 0.975 quantiles of the permutation distribution of $W'\tilde{u}_\pi(\theta_0)$.

Table 1 reports the empirical level of the tests in the case with $k = 1$ across different sample sizes. When the error terms and the instruments satisfy the exogeneity condition but are dependent, the PNS test overrejects; somewhat unexpectedly, the overrejection becomes more severe as the sample size grows. The rank-type tests also show some size distortions and it does not improve as the sample size increases. The right panel of Table 1 shows that when the instruments and the error terms are independent, the empirical level of the asymptotic, permutation and the rank-based CLR-type tests are close to the nominal significance level.

SA-9.2 Conditional heteroskedasticity with strong and very weak identification

In addition to the two identification scenarios $\lambda = 4$ considered in Section 3.3 of the main paper, we take up the case of very weak identification $\lambda = 0.1$ and strong identification $\lambda = 20$. The remaining part of the simulation design is the same as in the paper. The results are displayed in Table 2 and remain invariant to the case $\lambda = 4$. To reiterate, the rank-based tests reject by a substantial margin in the dependent case. The asymptotic AR test tends to underreject. The null rejection rates of the permutation tests are better than their asymptotic counterparts in most cases. Moreover, they perform on par with the wild bootstrap in the independent case and have an edge over the wild bootstrap AR test when the instruments satisfy the standard exogeneity condition, and there are included exogenous variables.

SA-9.3 Power

We use the following four DGPs: $(W'_i, X'_{2i}, v_i, \epsilon_i)' \sim t_5[0, I_{k+p+1}]$, $n = 100$, $\rho = 0.5$, $p = 1$, $k \in \{1, 5\}$, $\lambda \in \{4, 20\}$ considered in Section 3.3 of the main paper. The four DGPs combine two possible identification scenarios, weak vs. strong, with two different numbers of IVs (just vs. over-identification). The true parameter values are set as in the main paper. The power plots are obtained by testing the grid points $\{-2, -1.9, \dots, 1.9, 2\}$ with 1000 replications and $N = 200$ permutations for each replication.

Table 1: Null rejection probabilities at 5% level. Just-identified model with $k = p = 1$.

$\lambda = 4$	$t_5[0, I_{k+p+1}]$				$N[0, I_{k+p+1}]$			
n	50	100	200	400	50	100	200	400
AR	3.75	5.00	4.55	4.25	4.60	5.50	4.85	5.80
LM	3.75	5.00	4.55	4.25	4.60	5.50	4.85	5.80
CLR _a	3.55	4.75	4.75	4.30	4.55	5.50	4.70	5.65
CLR _b	3.55	4.75	4.75	4.30	4.55	5.50	4.70	5.65
PAR ₁	4.50	5.05	4.75	4.40	4.75	5.70	5.05	5.85
PAR ₂	4.25	5.40	4.90	4.30	4.75	5.65	4.90	5.90
PLM	4.25	5.40	4.90	4.30	4.75	5.65	4.90	5.90
PCLR _a	4.25	5.40	4.90	4.30	4.75	5.65	4.90	5.90
PCLR _b	4.25	5.40	4.90	4.30	4.75	5.65	4.90	5.90
RARn	10.35	11.15	12.35	12.00	4.90	5.45	4.90	6.05
RARw	7.65	9.30	8.70	8.35	4.60	5.35	4.85	5.65
RCLRn	8.10	9.65	11.40	11.65	3.45	4.45	4.25	5.50
RCLRw	7.10	8.95	8.65	8.10	4.55	5.20	5.05	5.75
WAR	4.65	6.00	5.35	4.80	4.40	5.45	4.85	5.60
WLM	4.65	6.00	5.35	4.80	4.40	5.45	4.85	5.60
PNS	15.30	16.90	17.80	19.40	5.20	6.00	5.55	6.20

Note: In the left panel, $(W'_i, u_i, \epsilon_i)' \sim t_5[0, I_{k+p+1}]$ and the single instrument W_i and the structural error term u_i are orthogonal but dependent. In the right panel, $(W'_i, u_i, \epsilon_i)' \sim N[0, I_{k+p+1}]$ and W_i and u_i are independent. AR, LM, and CLR are the heteroskedasticity-robust asymptotic tests, PAR₁ and PAR₂ denote the robust permutation AR tests, PLM and PCLR are the robust permutation LM and CLR tests, RARn and RARw denote the normal score and Wilcoxon score rank-based AR tests of [Andrews and Marmer \(2008\)](#), RCLRn and RCLRw are the normal score and Wilcoxon score rank-based CLR tests of [Andrews and Soares \(2007\)](#), WAR and WLM are the wild bootstrap AR and LM tests of [Davidson and MacKinnon \(2012\)](#) and PNS denotes the non-studentized permutation test. The number of replications is 2000, and the number of permutation samples for each replication is $N = 999$.

Table 2: Null rejection probabilities at 5% level. Conditionally heteroskedastic design with weak identification ($\lambda = 0.1$).

$n = 100$	$p = 1$						$p = 5$					
	$t_5[0, I_{k+p+1}]$			$N[0, I_{k+p+1}]$			$t_5[0, I_{k+p+1}]$			$N[0, I_{k+p+1}]$		
	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$
Tests												
AR	2.35	1.75	0.60	4.05	2.90	1.40	4.30	2.40	1.15	5.25	3.25	2.60
LM	2.95	4.35	3.55	4.90	4.30	4.30	5.25	3.35	2.85	6.00	4.30	3.85
CLR _a	2.40	2.05	0.80	4.05	3.65	2.00	4.10	2.40	1.35	5.30	3.30	3.00
CLR _b	2.45	2.25	1.25	4.10	3.60	2.20	4.25	2.60	1.25	5.25	3.45	3.00
PAR ₁	3.60	5.15	4.35	4.85	5.60	5.20	7.40	7.55	8.85	5.95	4.70	4.70
PAR ₂	3.35	4.85	4.35	4.90	5.30	5.00	6.35	5.50	4.95	6.40	5.30	5.35
PLM	4.45	5.95	6.20	5.45	5.80	6.35	6.80	4.65	5.05	6.50	5.60	5.95
PCLR _a	3.60	5.20	4.80	5.50	5.45	4.35	6.65	4.70	4.10	6.35	5.25	5.70
PCLR _b	3.65	6.40	8.45	5.50	5.65	5.15	6.80	5.20	7.55	6.35	5.50	6.85
RAR _n	22.30	27.30	32.55	13.45	10.20	8.05	23.55	24.80	29.85	13.75	10.10	8.70
RAR _w	14.35	17.90	20.10	10.40	8.85	6.65	15.95	16.00	17.55	10.40	8.25	7.35
RCLR _n	19.75	25.70	32.30	11.40	8.30	6.40	32.25	33.75	44.75	13.20	8.75	7.85
RCLR _w	14.35	18.05	21.95	9.75	8.20	6.60	27.00	27.55	38.65	11.15	8.55	8.60
WAR	4.50	5.80	5.55	4.75	5.20	4.90	8.65	10.80	12.60	5.75	4.40	4.95
WLM	4.65	3.90	3.95	5.65	4.20	6.30	7.00	6.80	7.00	5.25	4.30	5.45

Note: $t_5[0, I_{k+p+1}]$ and $N[0, I_{k+p+1}]$ correspond to $(W'_i, X'_{2i}, v_i, \epsilon_i)' \sim t_5[0, I_{k+p+1}]$ (dependent but uncorrelated) and $(W'_i, X'_{2i}, v_i, \epsilon_i)' \sim N[0, I_{k+p+1}]$ (independent), respectively. The error terms are generated as $u_i = W_{i1}v_i, V_i = \rho u_i + \sqrt{1 - \rho^2}\epsilon_i$, where W_{i1} is the first element of W_i and $\rho = 0.5$. The tests are in Table 1. The number of replications is 2000, and the number of permutation samples for each replication is $N = 999$.

Table 3: Null rejection probabilities at 5% level. Conditionally heteroskedastic design with strong identification ($\lambda = 20$).

$n = 100$	$p = 1$						$p = 5$					
	$t_5[0, I_{k+p+1}]$			$N[0, I_{k+p+1}]$			$t_5[0, I_{k+p+1}]$			$N[0, I_{k+p+1}]$		
	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$	$k = 2$	$k = 5$	$k = 10$
Tests												
AR	2.35	1.75	0.60	4.05	2.90	1.40	4.30	2.40	1.15	5.25	3.25	2.60
LM	3.05	3.20	2.40	4.90	3.00	3.15	4.55	3.10	2.10	5.65	3.95	3.90
CLR _a	2.25	2.00	0.95	4.60	2.80	1.85	4.55	2.70	1.45	5.70	3.50	3.20
CLR _b	2.45	2.40	1.35	4.65	2.95	2.20	4.70	2.80	1.75	5.80	3.55	3.25
PAR ₁	3.60	5.15	4.35	4.85	5.60	5.20	7.40	7.55	8.85	5.95	4.70	4.70
PAR ₂	3.35	4.85	4.35	4.90	5.30	5.00	6.35	5.50	4.95	6.40	5.30	5.35
PLM	4.35	4.35	4.50	5.45	4.20	4.75	5.95	4.40	4.20	6.45	4.95	5.90
PCLR _a	4.35	4.55	3.55	4.95	3.40	3.20	5.85	4.10	3.30	6.35	4.10	4.60
PCLR _b	4.50	6.00	8.25	5.00	3.90	4.65	6.00	5.35	7.25	6.50	4.90	6.65
RAR _n	22.30	27.30	32.55	13.45	10.20	8.05	23.55	24.80	29.85	13.75	10.10	8.70
RAR _w	14.35	17.90	20.10	10.40	8.85	6.65	15.95	16.00	17.55	10.40	8.25	7.35
RCLR _n	16.75	17.90	23.45	9.80	7.05	5.85	25.35	24.15	33.65	11.60	6.25	6.95
RCLR _w	12.80	13.80	16.75	8.85	7.05	5.95	22.05	21.15	28.75	9.85	6.25	6.90
WAR	4.50	5.80	5.55	4.75	5.20	4.90	8.65	10.80	12.60	5.75	4.40	4.95
WLM	3.60	3.85	3.70	4.90	4.75	4.55	6.15	5.25	7.00	5.80	3.65	4.95

Note: As in Table 3.

Except for the RCLR tests which are not heteroskedasticity-robust, we plot only the AR-type tests in Figure 4 and Figure 5 as they are identical to their LM and CLR counterparts. Relative to the other tests, the permutation tests show more accurate control of the Type I error and this is translated into the power. The wild bootstrap tests slightly overreject while the asymptotic tests underreject under the null. The slight power advantages of the wild bootstrap and disadvantages of the asymptotic tests relative to the permutation tests can be attributed this.

We can see that the rank-based tests are not robust to heteroskedasticity. Similar findings carry over to the over-identified case in Figures 6 and 7 though with more pronounced differences. All tests that control the size have lower power when identification is weak and the LM-type tests exhibit non-monotonic power curves. The performances of the PAR and PCLR tests appear to be promising.


SA-10 Additional empirical application

Becker and Woessmann (2010) study the effect of Protestantism on education before the Industrialization using counties and towns-level data from 1816 Prussia. To unravel the direction of causation between a better education of Protestants and industrialization in 1870s, the authors estimate IV regressions where the dependent variable is a number of primary schools per 1000 inhabitants, and the endogenous regressor is a fraction of Protestants instrumented by *the distance to Wittenberg*, the city from which Protestantism were spread in a roughly concentric manner in Martin Luther's times.

The authors consider several different specifications which differ in terms of the exogenous controls used: for the counties-level data,

- Specification 1: constant;
- Specification 2: latitude, longitude, a product of latitude and longitude, percentage of population living in towns;
- Specification 3: latitude, longitude, a product of latitude and longitude, percentage of population living in towns, percentage of population younger than 15 years of age, percentage of population older than 60 years of age, number of horses per capita, number of bulls per


Figure 4: Power curves: Just-identified model with $k = 1$ and $\lambda = 4$



Notes: the horizontal line corresponds to the nominal level of the tests (5%). The tests are as in the main paper.


$N = 200$ and 1000 replications.

Figure 5: Power curves: Just-identified model with $k = 1$ and $\lambda = 20$




Notes: same as in Figure 4.

Figure 6: Power curves: Over-identified model with $k = 5$ and $\lambda = 4$



Notes: same as in Figure 4.

Figure 7: Power curves: Over-identified model with $k = 5$ and $\lambda = 20$



Notes: same as in Figure 4.

capita, looms per capita, share of farm laborers in total population and tonnage of transport ships;

and for the towns-level data,

- Specification 4: latitude, longitude, a product of latitude and longitude, percentage of population younger than 15 years of age, percentage of population older than 60 years of age, looms per capita, buildings w/massive walls, businesses per capita and retailers per capita.

For further details of the study and the arguments underlie the use of the distance instrument, see [Becker and Woessmann \(2010\)](#).

Since there is a single instrument, all specifications are just-identified. The sample sizes are 293 for the counties-level data and 156 for the towns-level data in the sample for 1816 Prussia. We construct confidence intervals for the coefficient on the share of protestants using 1999 simulated samples for the asymptotic CLR, wild bootstrap, permutation and rank-based confidence intervals. We include confidence intervals from the homoskedastic two-stage least squares t test, t_{2sls} , and a non-studentized permutation test, PNS, based on $W'\tilde{u}(\theta_0)$, $\tilde{u}(\theta_0) = M_X(y - Y\theta_0)$, that rejects when it is in the lower or upper tails of the permutation distribution of $W'\tilde{u}_\pi(\theta_0)$ in all of the specifications.

The 95% confidence intervals and their lengths based on inverting different tests are reported in Table 4. In all of the specifications, the heteroskedasticity-robust asymptotic, permutation and wild bootstrap confidence intervals are nearly identical. Interestingly, they are shorter than the homoskedastic AR, RARn and RARw confidence intervals. The RCLRn and RCLRw confidence intervals are comparable to the permutation confidence intervals for the counties-level data, and shorter for the towns-level data, however these tests do not guard against heteroskedasticity. As the Breusch-Pagan test for heteroskedasticity in the reduced-form regression of the dependent variable and the endogenous regressor on the instrument and the exogenous covariates have p -values 0.01366 and 2.22×10^{-16} for Specification 1, 1.0916×10^{-6} and 0.0033 for Specification 2, 5.3147×10^{-7} and 6.8536×10^{-6} for Specification 3, and 0.14162 and 7.0197×10^{-5} for Specification 4, there is a strong evidence of heteroskedasticity and so the heteroskedasticity-robust methods should be more reliable. The non-studentized permutation test which is exact under independence (Assumption 1) and does not account for conditional heteroskedasticity yields a confidence interval considerably

wider than the one based on the robust permutation tests.

Using the heteroskedasticity and identification-robust confidence intervals, we confirm the findings of [Becker and Woessmann \(2010\)](#) that there is a significant effect of the share of Protestants on the school supply in the counties-level data.

Table 4: Confidence intervals for the coefficient on the share of Protestants in [Becker and Woessmann \(2010\)](#) data.

	Counties, $n = 293$						Towns, $n = 156$	
	Spec. 1		Spec. 2		Spec. 3		Spec. 4	
	95% CI	Length	95% CI	Length	95% CI	Length	95% CI	Length
t_{2sls}	[0.24, 1.27]	1.03	[0.94, 1.80]	0.86	[0.98, 1.94]	0.96	[−0.48, 1.03]	1.51
Hom. AR	[0.21, 1.25]	1.04	[0.95, 1.80]	0.85	[0.98, 1.95]	0.97	[−0.48, 1.05]	1.53
RAR _n	[0.35, 1.28]	0.93	[0.83, 2.04]	1.21	[0.86, 2.14]	1.28	[−0.49, 1.06]	1.55
RAR _w	[0.46, 1.31]	0.85	[0.79, 1.81]	1.02	[0.81, 1.97]	1.16	[−0.51, 0.98]	1.49
RCLR _n	[0.32, 1.28]	0.96	[0.99, 1.78]	0.79	[1.07, 1.89]	0.82	[−0.29, 0.73]	0.93
RCLR _w	[0.45, 1.32]	0.87	[1.03, 1.68]	0.65	[1.11, 1.83]	0.72	[−0.26, 0.68]	0.94
AR	[0.27, 1.22]	0.95	[1.07, 1.69]	0.62	[1.10, 1.86]	0.76	[−0.51, 0.97]	1.48
PAR ₁	[0.28, 1.21]	0.93	[1.07, 1.69]	0.62	[1.07, 1.87]	0.80	[−0.53, 0.98]	1.51
PAR ₂	[0.27, 1.23]	0.96	[1.07, 1.70]	0.63	[1.10, 1.86]	0.76	[−0.51, 0.95]	1.46
PNS	[0.22, 1.25]	1.03	[0.77, 2.03]	1.26	[0.72, 2.22]	1.50	[−0.89, 1.59]	2.48
WAR	[0.25, 1.23]	0.98	[1.07, 1.69]	0.62	[1.10, 1.88]	0.78	[−0.56, 0.98]	1.54
F -statistic	80.78		174.65		143.78		79.50	
Breusch-Pagan p -values	{0.01, 0.00}		{0.00, 0.00}		{0.00, 0.00}		{0.14, 0.00}	

Note: The number of permutation samples is 1999 ($N = 2000$). t_{2sls} and Hom. AR are the homoskedastic two-stage least squares t and AR tests, PNS denotes the non-studentized permutation test, and the remaining tests are as in Tables 1 and 2. The Breusch-Pagan test p -values are computed using the fitted values in the reduced-form regressions of y_i and Y_i on the instrument and exogenous covariates respectively.

References

- ANDREWS, D. W. and GUGGENBERGER, P. (2019a). Identification-and Singularity-Robust Inference for Moment Condition Models. *Quantitative Economics*, **10** 1703–1746.
- ANDREWS, D. W. K., CHENG, X. and GUGGENBERGER, P. (2020). Generic Results for Establishing the Asymptotic Size of Confidence Sets and Tests. *Journal of Econometrics*, **218** 496–531.
- ANDREWS, D. W. K. and GUGGENBERGER, P. (2017a). Asymptotic Size of Kleibergen’s LM and Conditional LR Tests for Moment Condition Models. *Econometric Theory*, **33** 1046–1080.
- ANDREWS, D. W. K. and GUGGENBERGER, P. (2017b). Supplemental material to ”Asymptotic Size of Kleibergen’s LM and Conditional LR Tests for Moment Condition Models”. *Econometric Theory*, **33**.
- ANDREWS, D. W. K. and GUGGENBERGER, P. (2019b). Supplemental material to “Identification-and Singularity-Robust Inference for Moment Condition Models”. *Quantitative Economics*, **10**.
- ANDREWS, D. W. K. and MARMER, V. (2008). Exactly Distribution-Free Inference in Instrumental Variables Regression with Possibly Weak Instruments. *Journal of Econometrics*, **142** 183–200.
- ANDREWS, D. W. K. and SOARES, G. (2007). Rank Tests for Instrumental Variables Regression with Weak Instruments. *Econometric Theory*, **23** 1033–1082.
- ANDREWS, D. W. K. and STOCK, J. H. (2007a). Inference with Weak Instruments. In *Advances in Econometrics: Proceedings of the Ninth World Congress of the Econometric Society*, vol. 3.
- ANDREWS, D. W. K. and STOCK, J. H. (2007b). Testing with Many Weak Instruments. *Journal of Econometrics*, **138** 24–46.
- BARBOUR, A. D. and CHEN, L. H. Y. (2005). The Permutation Distribution of Matrix Correlation Statistics. In *Stein’s Method and Applications* (A. D. Barbour and L. H. Y. Chen, eds.). Singapore University Press, 223–246.

- BARBOUR, A. D. and EAGLESON, G. K. (1986). Random Association of Symmetric Arrays. *Stochastic Analysis and Applications*, **4** 239–281.
- BECKER, S. O. and WOESSMANN, L. (2010). The Effect of Protestantism on Education before the Industrialization: Evidence from 1816 Prussia. *Economics Letters*, **107** 224–228.
- DAVIDSON, R. and MACKINNON, J. G. (2012). Wild Bootstrap Tests for IV Regression. *Journal of Business & Economic Statistics*, **28** 128–144.
- DICICCIO, C. J. and ROMANO, J. P. (2017). Robust Permutation Tests for Correlation and Regression Coefficients. *Journal of the American Statistical Association*, **112** 1211–1220.
- HALL, P. (1984). Central Limit Theorem for Integrated Square Error of Multivariate Nonparametric Density Estimators. *Journal of Multivariate Analysis*, **14** 1–16.
- HANSEN, B. E. (2022). *Econometrics*. Princeton University Press.
- HOEFFDING, W. (1951). A Combinatorial Central Limit Theorem. *Annals of Mathematical Statistics*, **22** 558–566.
- HU, T.-C., MORICZ, F. and TAYLOR, R. L. (1989). Strong Laws of Large Numbers for Arrays of Rowwise Independent Random Variables. *Acta Mathematica Hungarica*, **54** 153–162.
- LEHMANN, E. L. and ROMANO, J. P. (2005). *Testing Statistical Hypotheses*. 3rd ed. Springer.
- MASON, D. M. and NEWTON, M. A. (1992). A Rank Statistics Approach to the Consistency of a General Bootstrap. *The Annals of Statistics*, **20** 1611 – 1624.
- MOTOO, M. (1956). On the Hoeffding’s Combinatorial Central Limit Theorem. *Annals of the Institute of Statistical Mathematics*, **8** 145–154.
- PHAM, D. T., MÖCKS, J. and SROKA, L. (1989). Asymptotic Normality of Double-Indexed Linear Permutation Statistics. *Annals of the Institute of Statistical Mathematics*, **41** 415–427.
- SCHNELLER, W. (1988). A Short Proof of Motoo’s Combinatorial Central Limit Theorem using Stein’s Method. *Probability Theory and Related Fields*, **78** 249–252.

- VERSHYNIN, R. (2018). *High-Dimensional Probability: An Introduction with Applications in Data Science*, vol. 47. Cambridge University Press.
- ZHAO, L., BAI, Z., CHAO, C.-C. and LIANG, W.-Q. (1997). Error Bound in a Central Limit Theorem of Double-Indexed Permutation Statistics. *Annals of Statistics*, **25** 2210–2227.