

Nonlocal extension of causal thermodynamics of the isotropic cosmic fluid

Alexander B. Balakin^{1,*} and Alexei S. Ilin^{1,†}

¹*Department of General Relativity and Gravitation, Institute of Physics,
Kazan Federal University, Kremlevskaya str. 16a, Kazan 420008, Russia*

(Dated: November 30, 2021)

We establish the nonlocal generalization of the Israel-Stewart model for the relativistic causal thermodynamics of the cosmic fluid, which evolves in the homogeneous isotropic Universe. Based on the second law of thermodynamics we derive the integro-differential master equation for the nonequilibrium pressure scalar and reduce it to the differential equation of the second order in time derivatives. We show that this master equation can be considered as the relativistic analog of the Burgers equation, which is known in the classical theory of viscoelasticity. We obtain the nonlinear key equation, which contains the energy density scalar only, and analyze two exact solutions of the model. The effective temperature is considered to be associated with the barotropic equation of state of the cosmic fluid.

PACS numbers: 04.40.-b, 05.70.Ln

Keywords: causal thermodynamics, nonlocal interactions

I. INTRODUCTION

A. Prologue

The relativistic causal thermodynamics elaborated by Israel and Stewart [1] presents the remarkable page of the story of the development of the irreversible thermodynamics. The second law of the phenomenological thermodynamics, which declares that the entropy production σ of a closed physical system should be non-negative $\sigma \geq 0$, is the basic element of all versions of the relativistic phenomenological thermodynamics. The entropy production scalar σ is defined as the covariant divergence of the entropy flux four-vector S^k , i.e., $\sigma = \nabla_k S^k$. The difference between the versions of the irreversible thermodynamics, which have been formulated till now, is encoded in the structure of the four-vector S^k . The Eckart's version of this quantity [2] is known to have the form

$$S^k_{(\text{Eckart})} = s_0 n U^k + \frac{1}{T} q^k, \quad (1)$$

where n is the scalar of particle number density, T is the temperature, s_0 is the scalar of entropy per one particle, U^k is the timelike medium velocity four-vector, and q^k is the spacelike heat-flux four-vector. The Israel-Stewart version of the entropy flux four-vector is

$$S^k_{(IS)} = S^k_{(\text{Eckart})} + \frac{1}{T} q^l \left[\delta^k_l \alpha_0 \Pi + \alpha_1 \Pi^k_{l(0)} \right] - \frac{1}{2T} U^k \left[\beta_0 \Pi^2 - \beta_1 q^m q_m + \beta_2 \Pi^{mn}_{(0)} \Pi_{mn(0)} \right]. \quad (2)$$

In fact, Israel and Stewart have added all the possible terms of the second order with respect to the non-equilibrium quantities: the heat-flux four-vector q^k , the

scalar part Π of the non-equilibrium pressure Π_{ik} , and its traceless shear part $\Pi_{mn(0)}$. We use standardly the decomposition $\Pi_{ik} = \Pi_{ik(0)} + \frac{1}{3} \Delta_{ik} \Pi$, where $\Delta_{ik} \equiv g_{ik} - U_i U_k$ is the projector. New phenomenological parameters α_0 , α_1 , β_0 , β_1 and β_2 were considered as functions of the temperature T .

The standard Gibbs equation includes the scalar s_0

$$De + PD \left(\frac{1}{n} \right) = TDs_0, \quad (3)$$

where e describes the energy density per one particle $e = \frac{W}{n}$; W is the energy density scalar; P is the isotropic equilibrium Pascal pressure; D is the convective derivative defined as $D = U^k \nabla_k$ with the covariant derivative ∇_k . The listed quantities form the stress-energy tensor of the medium

$$T^{ik} = W U^i U^k + U^i q^k + U^k q^i - \Delta^{ik} P + \Pi^{ik}. \quad (4)$$

The heat flux four-vector is orthogonal to the velocity four-vector, $q^k U_k = 0$, and the symmetric non-equilibrium pressure tensors satisfies the condition $\Pi^{ik} U_k = 0$.

B. Relativistic causal thermodynamics of the isotropic homogeneous Universe

Numerous applications of the causal thermodynamics to the cosmological models of the Friedmann type (see, e.g., [3] - [15]) were based on the evident fact that due to the claimed spacetime symmetry one has to put the spacelike heat-flux four-vector to zero, $q^k \equiv 0$, and to consider the tensor $\Pi_{mn(0)}$ to be vanishing. Also, all the thermodynamic quantities are considered to be the functions of the cosmological time only, and the velocity four-vector is chosen to be of the form $U^i = \delta_0^i$. These requirements are connected with the idea of inheritance of the symmetry of the spacetime metric

$$ds^2 = dt^2 - a^2(t) [dx^2 + dy^2 + dz^2], \quad c = 1, \quad (5)$$

*Electronic address: Alexander.Balakin@kpfu.ru

†Electronic address: alexeyilinjukeu@gmail.com

by all the physical quantities: the energy density scalar, Pascal pressure, etc. With these assumptions, the entropy production scalar is calculated to have the form

$$\sigma = \frac{\Pi}{T} \left[\frac{1}{3} \Theta - \beta_0 D\Pi - \frac{1}{2} T \Pi \nabla_l \left(\frac{\beta_0 U^l}{T} \right) \right] \geq 0, \quad (6)$$

and the non-equilibrium pressure Π appeared as the solution to the equation

$$\beta_0 D\Pi + \Pi \left[\frac{1}{9\zeta} + \frac{T}{2} (\Theta + D) \left(\frac{\beta_0}{T} \right) \right] = \frac{1}{3} \Theta. \quad (7)$$

The phenomenologically introduced function ζ plays the role of the bulk viscosity coefficient; the scalar $\Theta \equiv \nabla_k U^k$ is the expansion scalar of the medium flow.

C. Motivation and structure of the work

According to the standard terminology accepted in the classical theory of viscoelasticity (see, e.g., [16]), and in the theory of irreversible thermodynamics of fluid (see, e.g., [17]), when $\beta_0=0$, the equation (7) describes the analog of the model of the Newton fluid, for which the stress tensor is proportional to the time derivative of the strain tensor. In the classical linear theory of viscoelasticity the strain tensor $\varepsilon_{\mu\nu} = \frac{1}{2} \left(\frac{\partial \mathcal{U}_\nu}{\partial x^\mu} + \frac{\partial \mathcal{U}_\mu}{\partial x^\nu} \right)$ contains the displacement three-vector \mathcal{U}^ν (Greek indices take values 1, 2, 3). For illustration only, we use for this object the term ϵ without indices. Since in the classical viscoelasticity the three velocity of the medium V^ν is equal to the time derivative of the displacement vector, i.e., $V^\nu = \dot{\mathcal{U}}^\nu$, we see that $\dot{\varepsilon}_{\mu\nu} = \frac{1}{2} \left(\frac{\partial V_\nu}{\partial x^\mu} + \frac{\partial V_\mu}{\partial x^\nu} \right)$, thus, for the Newton fluid we can use the compact formula $\pi = \eta \dot{\varepsilon}$, where symbol π indicates the stress tensor, and the η is the viscosity coefficient. When $\beta_0 \neq 0$, the equation (7) describes the analog of the Maxwell model of viscoelasticity, for which $\tau \dot{\pi} + \pi = \nu \dot{\varepsilon}$. The analog of the Hooke's term, which in classical viscoelasticity is of the form $\pi = E\epsilon$, does not appear in the equation (7), however, it would be interesting to study the analog of the classical model, associated with the constitutive equation of the type $\tau \dot{\pi} + \mu \pi = \nu \dot{\varepsilon} + E\epsilon$, which is extensively used in the transient and steady-state rheologies, as well as in the theory of viscoelastic media and materials (see, e.g., [18]). This equation is associated with the Burgers model (see, e.g., [19, 20]), for which the constitutive equation is of the second order in time derivatives for the stress π

$$\tau \ddot{\pi} + \mu \dot{\pi} + \omega \pi = \nu \dot{\varepsilon} + E\epsilon, \quad (8)$$

and contains the strain ϵ with the first and second time derivatives. When one establishes the relativistic theory of elasticity and viscoelasticity, the problem appears how to define the covariant version of the displacement four-vector, or of the covariant strain tensor (see, e.g., [21–23]), however, there are no problems with the formulation of the covariant analog of the tensor $\dot{\varepsilon}_{\mu\nu}$. In this

sense, when we address to the covariant generalization of the Burgers equation, we avoid the mentioned problem. In this work we show that the covariant analog of the Burgers equation for the isotropic system can be formulated using the nonlocal extension of the Israel-Stewart theory.

The paper is organized as follows. In Section II we describe the formalism of the nonlocal extension of the Israel-Stewart theory based on the Friedmann spacetime platform. In Section III we derive the key equations of the model and analyze them using two examples. Section IV contains conclusions.

II. NONLOCAL EXTENSION OF THE ISRAEL-STEWART THEORY

A. The formalism

1. Evolutionary equation for the non-equilibrium pressure Π

We consider now the decomposition of the entropy flux four-vector for the spatially homogeneous isotropic cosmological model in the following form:

$$S^k = s_0 n U^k - \frac{1}{2T} U^k \beta_0 \Pi^2 + \frac{1}{2} \gamma_0 n U^k (D^{-1}\Pi)^2, \quad (9)$$

where γ_0 is a constant. The term $D^{-1}\Pi$ is defined as follows:

$$D(D^{-1}\Pi) = \Pi, \quad D^{-1}\Pi = \int d\tau \Pi, \quad d\tau = U_k dx^k. \quad (10)$$

The coefficient $\frac{1}{2}\gamma_0 n U^k$ is chosen so that its divergence is equal to zero due to the conservation law for the particle number

$$\nabla_k (n U^k) = 0, \quad \Rightarrow Dn + n\Theta = 0. \quad (11)$$

Now the entropy production scalar takes the form

$$\sigma = \frac{\Pi}{T} \left[\frac{\Theta}{3} - \beta_0 D\Pi - \frac{T}{2} \Pi \nabla_l \left(\frac{\beta_0 U^l}{T} \right) + \gamma_0 n T (D^{-1}\Pi) \right]. \quad (12)$$

This quantity is non-negative, when

$$\frac{\Pi}{9\zeta} = \frac{\Theta}{3} - \beta_0 D\Pi - \frac{T}{2} \Pi \nabla_l \left(\frac{\beta_0 U^l}{T} \right) + \gamma_0 n T (D^{-1}\Pi). \quad (13)$$

Clearly, the equation (13) recovers the definition of the non-equilibrium pressure appeared in the Israel-Stewart theory, if $\gamma_0 = 0$.

The integro-differential equation (13) can be rewritten in the form of differential equation of the second order

$$D \left[\frac{\Pi}{9n\zeta T} - \frac{\Theta}{3nT} + \frac{\beta_0}{nT} D\Pi + \frac{\Pi}{2n} (\Theta + D) \left(\frac{\beta_0}{T} \right) \right] = \gamma_0 \Pi. \quad (14)$$

We can rewrite this equation in the form

$$\begin{aligned} & \frac{\beta_0}{nT} D^2 \Pi + D \Pi \left[D \left(\frac{\beta_0}{nT} \right) + \frac{1}{9nT\zeta} + \frac{1}{2n} (\Theta + D) \left(\frac{\beta_0}{T} \right) \right] + \\ & + \Pi \left\{ -\gamma_0 + D \left(\frac{1}{9nT\zeta} \right) + D \left[\frac{1}{2n} (\Theta + D) \left(\frac{\beta_0}{T} \right) \right] \right\} = \\ & = D \left(\frac{1}{3nT} \Theta \right), \end{aligned} \quad (15)$$

analogous to the Burgers equation.

2. Evolution of the temperature T

Since in the isotropic Universe, which we consider in our model, the heat-flux four-vector vanishes, $q^i = 0$, and since the spatial gradient of the temperature is equal to zero, we cannot obtain the standard equation for the temperature evolution by the Cattaneo scheme [24]. We need now some additional equation for the temperature depending on the cosmological time only. The corresponding procedure was described, e.g., in the works [6, 9, 25–27]. Let us recall the main idea of this procedure. When we consider the energy density $W(n, T)$ and the Pascal pressure $P(n, T)$ to be functions of the particle number density n and temperature T , we have to require that

$$\frac{\partial^2 s_0}{\partial n \partial T} = \frac{\partial^2 s_0}{\partial T \partial n}. \quad (16)$$

This means that the entropy is considered as the state function with continuous derivatives up to the second order. Using the Gibbs equation (3) and the condition (16), we obtain immediately the compatibility equation

$$n \frac{\partial W}{\partial n} + T \frac{\partial P}{\partial T} = W + P, \quad (17)$$

which has to link the functions $W(n, T)$ and $P(n, T)$.

When the equation of state for the fluid is linear barotropic, $P = \gamma W$, i.e., the state functions are linked directly and thus, the functions T and n do not take part in the formulation of the equation of state, the functions T and n become the latent ones. Evolution of the function $n(t)$ is predicted by the law (11); we need also the equation for the function $T(t)$. For the linear barotropic equation of state the energy density scalar has to satisfy the linear differential equation in partial derivatives

$$n \frac{\partial W}{\partial n} + \gamma T \frac{\partial W}{\partial T} = (1 + \gamma) W. \quad (18)$$

Solving the characteristic equations

$$\frac{dn}{n} = \frac{dT}{\gamma T} = \frac{dW}{(1 + \gamma)W}, \quad (19)$$

we obtain the energy density scalar in the following form

$$W = n^{1+\gamma} \mathcal{F} \left(\frac{T}{n^\gamma} \right), \quad (20)$$

where \mathcal{F} is arbitrary function of its argument. Equivalently, one can express the temperature as follows:

$$T = n^\gamma \mathcal{G} \left(\frac{W^{1+\gamma}}{n^\gamma} \right), \quad (21)$$

where \mathcal{G} is arbitrary function of its argument. In this formula for the particle number scalar $n(t)$ we can use the result of integration of the equation (11), namely,

$$n(t) = n_0 \left(\frac{a(t_0)}{a(t)} \right)^3, \quad n_0 = n(t_0). \quad (22)$$

It is interesting to attract attention to two special cases.

(i) When the function \mathcal{G} in (21) is linear; then the scalar n disappears, and we obtain

$$T = \left(\frac{W}{\omega_0} \right)^{\frac{\gamma}{1+\gamma}} \Rightarrow W = \omega_0 T^{\frac{1+\gamma}{\gamma}}. \quad (23)$$

When $\gamma = \frac{1}{3}$, we deal with the Stefan-Boltzmann law $W = \varkappa T^4$, thus one can indicate (23) as a generalization of this law.

(ii) When $\gamma = -1$, we deal with the formulas $W = \mathcal{F}(Tn)$. This model can be associated with the dark energy, since now $W + P = 0$.

Below we will indicate the temperature found using this auxiliary procedure as the effective temperature.

3. Evolution of the energy density scalar

The conservation law $\nabla_k T^{ik} = 0$ can be reduced to

$$\dot{W} + \left[W + P - \frac{1}{3} \Pi \right] \Theta = 0, \quad (24)$$

where Θ can be expressed in terms of the Hubble function $H(t) = \frac{\dot{a}}{a}$, i.e., $\Theta = 3H$. Here and below the dot denotes the derivative with respect to cosmological time. Then we add the Einstein equation

$$3H^2 = \kappa W, \quad (25)$$

and obtain the set of four equations (15), (24), (25), (21) for four unknown functions $\Pi(t)$, $T(t)$, $W(t)$, $H(t)$.

B. Truncated model and the key equation

Our goal is to derive and analyze the so-called key equation, which includes only one unknown function, say,

the energy density scalar W . In order to reach some analytic progress in calculations, we can use the ansatz about the structure of the phenomenologically introduced coefficients ζ and β_0 . First of all, we assume that the bulk viscosity coefficient depends on the temperature T and on the expansion scalar Θ as follows: $\zeta(T, \Theta) = \frac{\zeta_0}{T\Theta}$, where ζ_0 is some constant. This assumption is motivated by the results of numerous investigations of the bulk viscosity coefficient for various media and materials: generally, this coefficient decreases when the temperature grows. Also we assume that $\beta_0(T) = h_0 \cdot T$, where h_0 is some constant. The second assumption relates to the findings that the relaxation time associated with the coefficient β_0 grows when the temperature increases. Of course, these assumptions should be considered as an ansatz, which has to be verified later. Now we use the relationship $\Theta = 3H$ and simplify the equation (15) distinguishing two cases: $h_0 \neq 0$ and $h_0 = 0$.

1. *The key equation for the model with $h_0 \neq 0$*

When $h_0 \neq 0$ we obtain from (15)

$$\begin{aligned} & \ddot{\Pi} + H\dot{\Pi} \left(\frac{1}{3\zeta_0 h_0} + \frac{9}{2} \right) + \\ & + \Pi \left[\left(\dot{H} + 3H^2 \right) \left(\frac{1}{3\zeta_0 h_0} + \frac{3}{2} \right) - \frac{\gamma_0}{h_0} n \right] = \\ & = \frac{H}{h_0 T} \left(\frac{\dot{H}}{H} + 3H - \frac{\dot{T}}{T} \right). \end{aligned} \quad (26)$$

In order to simplify this equation, we introduce the new dimensionless variable $x = \frac{a(t)}{a(t_0)}$ and replace the derivative with respect to time using the relationship $\frac{d}{dt} = xH \frac{d}{dx}$. We obtain from (26) the equation

$$\begin{aligned} & x^2 \Pi''(x) + x \Pi'(x) \left[\frac{11}{2} + \frac{1}{3\zeta_0 h_0} + x \frac{H'}{H} \right] + \\ & + \Pi \left[\left(x \frac{H'}{H} + 3 \right) \left(\frac{1}{3\zeta_0 h_0} + \frac{3}{2} \right) - \frac{\gamma_0 n_0}{h_0 x^3 H^2} \right] = \\ & = \frac{1}{h_0 T} \left(x \frac{H'}{H} + 3 - x \frac{T'}{T} \right). \end{aligned} \quad (27)$$

Here and below the prime denotes the derivative with respect to the variable x . Using the Einstein equation (25) we make the replacements $H^2 \rightarrow \frac{\kappa}{3} W$, and $\frac{H'}{H} = \frac{W'}{2W}$. Using the conservation law (24) we can write now

$$\Pi = xW'(x) + 3(1 + \gamma)W. \quad (28)$$

Finally, we express the temperature and its derivative via the W and n , using (21) and keeping in mind that $n = \frac{n_0}{x^3}$

from (22). Thus, we obtain the nonlinear equation of the third order for the function $W(x)$ only; this is the key equation, it has the form

$$\begin{aligned} & x^3 W''' + x^2 W'' \left[3\gamma + \frac{21}{2} + \frac{1}{3\zeta_0 h_0} + x \frac{W'}{2W} \right] + \\ & + xW' \left[x \frac{W'}{2W} \left(\frac{1}{3\zeta_0 h_0} + \frac{11}{2} + 3\gamma \right) + \frac{17+9\gamma}{6\zeta_0 h_0} + \frac{5}{4}(23+15\gamma) \right] + \\ & + 3(1 + \gamma)W \left(\frac{1}{\zeta_0 h_0} + \frac{9}{2} \right) = \\ & = \frac{1}{h_0 T} \left(x \frac{W'}{2W} + 3 - x \frac{T'}{T} \right) + \frac{3\gamma_0 n_0}{h_0 \kappa x^3} \left[x \frac{W'}{W} + 3(1 + \gamma) \right]. \end{aligned} \quad (29)$$

2. *The key equation for the model with $h_0 = 0$*

When $h_0 = 0$ and thus $\beta_0 = 0$, the equation for Π obtained in the framework of the Israel-Stewart theory, is of the type typical for the Newtonian fluid, $\Pi = \frac{1}{3}\Theta$. However, the nonlocal extension of the Israel-Stewart theory, presented in our paper, gives the equation of the first order in time derivative:

$$D\Pi + \Pi n T \zeta \left[D \left(\frac{1}{n T \zeta} \right) - 9\gamma_0 \right] = 3n T \zeta D \left(\frac{\Theta}{n T} \right), \quad (30)$$

The procedure similar to the one used for the case $h_0 \neq 0$ gives the nonlinear equation of the second order for the energy density scalar

$$\begin{aligned} & x^2 W'' + xW' \left[x \frac{W'}{2W} + \frac{17+9\gamma}{2} \right] + 9(1 + \gamma)W = \\ & = \frac{3\zeta_0}{T} \left(x \frac{W'}{2W} + 3 - x \frac{T'}{T} \right) + \frac{9\zeta_0 \gamma_0 n_0}{\kappa x^3} \left[x \frac{W'}{W} + 3(1 + \gamma) \right]. \end{aligned} \quad (31)$$

With key equations (29) or (31) we are ready to analyze some exact solutions for the nonlocal model.

III. EXACT SOLUTIONS TO THE KEY EQUATIONS

A. Solution of the de Sitter type

If we consider $W(x) = W_* = const$, we obtain that $H = H_0 = \sqrt{\frac{\kappa W_*}{3}}$, and $a(t) = a(t_0) e^{H_0(t-t_0)}$, i.e., we deal with the solution of the de Sitter type, and the quantity κW_* plays the role of effective cosmological constant. If

to put $W(x) = W_*$ to the key equation (29), we conclude that the temperature $T(x)$ has to satisfy the equation

$$x \left(\frac{1}{T} \right)' + \frac{3}{T} = 3(1+\gamma) \left[W_* \left(\frac{1}{\zeta_0} + \frac{9}{2} h_0 \right) - \frac{3\gamma_0 n_0}{\kappa x^3} \right]. \quad (32)$$

The solution to this equation is

$$\frac{1}{T(x)} = \frac{1}{x^3 T(1)} + (1+\gamma) W_* \left(\frac{1}{\zeta_0} + \frac{9}{2} h_0 \right) \left(1 - \frac{1}{x^3} \right) - 9(1+\gamma) \frac{\gamma_0 n_0}{\kappa x^3} \log x. \quad (33)$$

Asymptotically, when $x \rightarrow \infty$, the inverse temperature tends to the constant

$$\frac{1}{T(\infty)} = (1+\gamma) W_* \left(\frac{1}{\zeta_0} + \frac{9}{2} h_0 \right). \quad (34)$$

Taking into account that according to (21) we have for this case that

$$\frac{1}{T} = x^{3\gamma} \mathcal{H}(x^{-3\gamma}), \quad (35)$$

where $\mathcal{H}(z)$ is arbitrary function of its argument, one can represent the function $\mathcal{H}(z)$ as follows:

$$\begin{aligned} \mathcal{H}(z) &= (1+\gamma) W_* \left(\frac{1}{\zeta_0} + \frac{9}{2} h_0 \right) z + \\ &+ \left[\frac{1}{T(1)} - (1+\gamma) W_* \left(\frac{1}{\zeta_0} + \frac{9}{2} h_0 \right) \right] z^{\frac{1+\gamma}{\gamma}} + \\ &+ 3 \frac{(1+\gamma) \gamma_0 n_0}{\gamma \kappa} z^{\frac{1+\gamma}{\gamma}} \log z, \end{aligned} \quad (36)$$

In other words, when the temperature is presented by the admissible function (35), (36), we obtain the exact solution of the non-locally extended Israel-Stewart model, which can be characterized as the de Sitter type solution.

B. Power-law type solutions

1. The key equation for the energy density scalar

The key equation is nonlinear in the function W and in its first derivative $W'(x)$. In order to find its exact particular solutions we assume the energy density scalar to have the power - law form $W = W_* x^\lambda$. Now the key equation (29) takes the form

$$\begin{aligned} &\frac{3}{2} x^\lambda h_0 W_* \left\{ \lambda^3 + \lambda^2 \left[\frac{1}{2} (13 + 6\gamma) + \frac{1}{3\zeta_0 h_0} \right] + \right. \\ &\left. + \lambda \left[\frac{1}{2} (27 + 21\gamma) + \frac{5 + 3\gamma}{3\zeta_0 h_0} \right] + 2(1+\gamma) \left(\frac{9}{2} + \frac{1}{\zeta_0 h_0} \right) \right\} = \end{aligned}$$

$$= x \left(\frac{1}{T} \right)' + \left(3 + \frac{\lambda}{2} \right) \left(\frac{1}{T} \right) + \frac{3\gamma_0 n_0}{\kappa x^3} [\lambda + 3(1+\gamma)]. \quad (37)$$

When $h_0 = 0$, this equation can be rewritten as

$$\begin{aligned} &\frac{1}{2\zeta_0} x^\lambda W_* [\lambda^2 + \lambda(5 + 3\gamma) + 6(1+\gamma)] = \\ &= x \left(\frac{1}{T} \right)' + \left(3 + \frac{\lambda}{2} \right) \left(\frac{1}{T} \right) + \frac{3\gamma_0 n_0}{\kappa x^3} [\lambda + 3(1+\gamma)]. \end{aligned} \quad (38)$$

The non-equilibrium pressure Π can be written now as

$$\Pi(x) = P(x) \frac{1}{\gamma} [\lambda + 3(1+\gamma)], \quad (39)$$

i.e., the ratio $\frac{\Pi(x)}{P(x)}$ is constant and is predetermined by the parameters λ and γ .

2. Evolution of the effective temperature

Our ansatz is that the effective temperature satisfies the equation

$$x \left(\frac{1}{T} \right)' + \left(3 + \frac{\lambda}{2} \right) \left(\frac{1}{T} \right) + \frac{3\gamma_0 n_0}{\kappa x^3} [\lambda + 3(1+\gamma)] = 0. \quad (40)$$

The solution to this equation is

$$\frac{1}{T(x)} = x^{-3} \left\{ \frac{x^{-\frac{\lambda}{2}}}{T(1)} + \frac{6\gamma_0 n_0}{\kappa \lambda} [\lambda + 3(1+\gamma)] [x^{-\frac{\lambda}{2}} - 1] \right\}. \quad (41)$$

As for the admissible function $\tilde{\mathcal{H}}(z)$ associated with this solution, we have to use the formula

$$\frac{1}{T(x)} = x^{3\gamma} \tilde{\mathcal{H}} \left(x^{\frac{\gamma}{1+\gamma}} [\lambda + 3(1+\gamma)] \right), \quad (42)$$

where $\tilde{\mathcal{H}}(z)$ is again an arbitrary function of its argument. We can find now that

$$\begin{aligned} \tilde{\mathcal{H}}(z) &= \left\{ \frac{1}{T(1)} + \frac{6\gamma_0 n_0}{\kappa \lambda} [\lambda + 3(1+\gamma)] \right\} z^{-\frac{1+\gamma}{2\gamma} \left[\frac{\lambda + 6(1+\gamma)}{\lambda + 3(1+\gamma)} \right]} - \\ &- \frac{6\gamma_0 n_0}{\kappa \lambda} [\lambda + 3(1+\gamma)] z^{-\frac{3(1+\gamma)^2}{\gamma[\lambda + 3(1+\gamma)]}}. \end{aligned} \quad (43)$$

Asymptotically, at $x \rightarrow \infty$ we obtain that $T \rightarrow 0$, if $\lambda < -6$. When $\lambda = -6$, we law for the inverse temperature is

$$\frac{1}{T(x)} = \frac{1}{T(1)} + \frac{3\gamma_0 n_0}{\kappa} (1-\gamma) (1-x^{-3}), \quad (44)$$

and thus the asymptotic value $T(\infty)$ is constant.

3. Geometric properties of the model

The Hubble function $H(x)$ found in terms of the reduced scale factor x

$$H(x) = \sqrt{\frac{\kappa W_*}{3}} x^{\frac{\lambda}{2}}, \quad (45)$$

allows us to recover the dependence $a(t)$ by the standard formula

$$t - t_0 = \int_1^{\frac{a(t)}{a(t_0)}} \frac{dx}{xH(x)}. \quad (46)$$

We obtain immediately that

$$\frac{a(t)}{a(t_0)} = \left[1 - \frac{\lambda}{2}(t - t_0) \sqrt{\frac{\kappa W_*}{3}} \right]^{-\frac{2}{\lambda}}. \quad (47)$$

In terms of the cosmological time the Hubble function is

$$H(t) = \sqrt{\frac{\kappa W_*}{3}} \left[1 - \frac{\lambda}{2}(t - t_0) \sqrt{\frac{\kappa W_*}{3}} \right]^{-1}. \quad (48)$$

The acceleration parameter has now the form

$$-q \equiv \frac{\ddot{a}}{aH^2} = 1 + \frac{\lambda}{2}. \quad (49)$$

Universe expands with acceleration, if $\lambda > -2$.

4. Characteristic equation and its solutions

When the temperature satisfies the equation (40), the parameter λ has to satisfy the cubic characteristic equation

$$\lambda^3 + \lambda^2 \left[\frac{1}{2}(13 + 6\gamma) + \frac{1}{3\zeta_0 h_0} \right] + \lambda \left[\frac{1}{2}(27 + 21\gamma) + \frac{5 + 3\gamma}{3\zeta_0 h_0} \right] + 2(1 + \gamma) \left(\frac{9}{2} + \frac{1}{\zeta_0 h_0} \right) = 0. \quad (50)$$

Depending on values of two parameters γ and $\xi \equiv \frac{1}{3\zeta_0 h_0}$ this equation can have one real solution plus two complex conjugated, or three real solutions $\lambda_1, \lambda_2, \lambda_3$. These real roots can be different, say, $\lambda_1 > \lambda_2 > \lambda_3$; two of them can coincide, $\lambda_1 = \lambda_2 \neq \lambda_3$, or three roots can coincide $\lambda_1 = \lambda_2 = \lambda_3$. These solutions can be standardly presented by the Cardano formulas and describe various regimes of the cosmological expansion.

As an illustration, we consider only one special case $\gamma = -1$, for which the characteristic equation

$$\lambda^3 + \lambda^2 \left(\frac{7}{2} + \xi \right) + \lambda(3 + 2\xi) = 0 \quad (51)$$

has the following roots:

$$\lambda_1 = 0, \quad \lambda_2 = -2, \quad \lambda_3 = -\frac{1}{2}(3 + 2\xi). \quad (52)$$

Clearly, the root $\lambda_1 = 0$ converts the power-law solution to the de Sitter one. Finally, when $h_0 = 0$, the characteristic equation (50) converts into the quadratic equation

$$\lambda^2 + \lambda(5 + 3\gamma) + 6(1 + \gamma) = 0, \quad (53)$$

the solutions to which are $\lambda_1 = -2, \lambda_2 = -3(1 + \gamma)$.

IV. CONCLUSIONS

We presented the nonlocal extension of the Israel-Stewart causal thermodynamics of the cosmic fluid, which inherits the isotropy and homogeneity of the expanding Universe. We would like to formulate briefly the following results of the analysis of the model as a whole, and of the exact solutions, which we obtained for two cases: the de Sitter type and the power-law solutions.

1. The established nonlocal extension of the Israel-Stewart causal thermodynamics is the relativistic analog of the Burgers model known in classical viscoelasticity (please, compare (8) with (26)).

2. For the linear barotropic equation of state of the cosmic fluid we reconstructed the properties of the effective temperature (21).

3. The de Sitter type solution is provided by the specific choice of the effective temperature (33), which tends asymptotically to the constant; the effective cosmological constant is presented by the value κW_* , where W_* is a stationary value of the energy density of the cosmic fluid; the de Sitter type solution coincides with power-law solution, when the value $\lambda = 0$ happens to be the root of the characteristic equation (see, e.g., (51)).

4. The parameter λ , which predetermines the properties of the power-law solution of the model, satisfies the cubic characteristic equation (50), and thus can describe at most three stages in the Universe history, which correspond to three real roots of the characteristic equation; the value of this parameter depends on two constants: the effective bulk viscosity ζ_0 and the effective relaxation parameter h_0 .

5. For the case $\lambda < 0$ the scale factor of the expanding Universe is monotonic and is presented by the power-law function (47).

6. The effective temperature vanishes asymptotically, when the parameter λ satisfies the inequality $\lambda < -6$ (see, e.g., (41)).

Acknowledgments

The work was supported by the Russian Science Foundation (Grant No 21-12-00130), and partially by the Kazan Federal University Strategic Academic Leadership Program.

References

- [1] W. Israel, J.M. Stewart, Transient relativistic thermodynamics and kinetic theory, *Ann. Phys.* 118 (1979) 341.
- [2] C. Eckart, The thermodynamics of irreversible processes. III. Relativistic theory of the simple fluid, *Phys. Rev.* 58 (1940) 919.
- [3] R. Maartens, J. Triginer, Density perturbations with relativistic thermodynamics, *Phys. Rev. D* 56 (1997) 4640.
- [4] W. Zimdahl, Cosmological particle production, causal thermodynamics, and inflationary expansion, *Phys. Rev. D* 61 (2000) 083511.
- [5] W. Zimdahl, D. Pavon, R. Maartens, Reheating and causal thermodynamics, *Phys. Rev. D* 55 (1997) 4681.
- [6] W. Zimdahl, Bulk Viscous Cosmology, *Phys. Rev. D* 53 (1996) 5483.
- [7] L. Herrera, D. Pavon, Hyperbolic theories of dissipation: Why and when do we need them? *Physica A* 307 (2002) 121.
- [8] E. Bittencourt, L.G. Gomes, R. Klippert, Bianchi-I cosmology from causal thermodynamics, *Class. Quantum Grav.* 34 (2017) 045010.
- [9] R. Maartens, Causal Thermodynamics in Relativity, arXiv:astro-ph/9609119.
- [10] N. Cruz, A. Hernandez-Almada, O. Cornejo-Perez, Constraining a causal dissipative cosmological model, *Phys. Rev. D* 100 (2019) 083524.
- [11] M. Cruz, S. Lepe, S.D. Odintsov, Thermodynamically allowed phantom cosmology with viscous fluid, *Phys. Rev. D* 98 (2018) 083515.
- [12] M. Cruz, N. Cruz, S. Lepe, Accelerated and decelerated expansion in a causal dissipative cosmology, *Phys. Rev. D* 96 (2017) 124020.
- [13] M. Cataldo, N. Cruz, S. Lepe, Viscous dark energy and phantom evolution, *Phys. Lett. B* 619 (2005)5.
- [14] M.K. Mak, T. Harko, Full causal dissipative cosmologies with stiff matter, *Int. J. Mod. Phys. D* 13 (2004) 273.
- [15] R. Maartens, J. Triginer, Acoustic oscillations and viscosity, *Phys. Rev. D* 58 (1998) 123507.
- [16] R.M. Christensen, Theory of viscoelasticity, Dover Publications Inc. Mineola, New York, 2003.
- [17] D. Jou, J. Casas - Vázquez, G. Lebon, Extended Irreversible Thermodynamics, Berlin, Springer Verlag, 1996.
- [18] G. Rumpker, D. Wolf, Viscoelastic relaxation of a Burgers half-space: implications for the interpretation of the Fennoscandian uplift, *Geophys. J. Int.* 124 (1996) 541.
- [19] J.M. Burgers, Mechanical considerations, model systems, phenomenological theories of relaxation and of viscosity, *Verh. K. Akad. Wet. Amsterdam* 1 15(3) (1935) 5.
- [20] J. Malek, K.R. Rajagopal, K. Tuma, Derivation of the Variants of the Burgers Model Using a Thermodynamic Approach and Appealing to the Concept of Evolving Natural Configurations, *Fluids* 3 (2018) 69.
- [21] B. Mashhoon, Nonlocal gravity, Oxford University Press, Oxford, 2017.
- [22] D. Puetzfeld, Yu. N. Obukhov, F.W. Hehl, Constitutive law of nonlocal gravity, *Phys. Rev. D* 99 (2019) 104013.
- [23] J.D. Brown, Elasticity Theory in General Relativity, *Class. Quantum Grav.* 38 (2021) 085017.
- [24] C. Cattaneo, Sulla Condizione Del Calore, *Atti Del Semin. Matem. E Fis. Della Univ. Modena* 3 (1948) 83.
- [25] W. Zimdahl, A.B. Balakin, Inflation in a self-interacting gas universe, *Phys. Rev. D* 58 (1998) 063503.
- [26] W. Zimdahl, J. Gariel, G. Le Denmat, Generalised equilibrium of cosmological fluids in second-order thermodynamics, *Class. Quant. Grav.* 16 (1999) 3207.
- [27] W. Zimdahl, A.B. Balakin, Cosmological thermodynamics and deflationary gas universe, *Phys. Rev. D* 63 (2001) 023507.