

# ON THE SET OF POINTS AT WHICH AN INCREASING CONTINUOUS SINGULAR FUNCTION HAS A NONZERO FINITE DERIVATIVE

MARTA KOSSACZKÁ AND LUDEK ZAJÍČEK

ABSTRACT. Sánchez, Viader, Paradís and Carrillo (2016) proved that there exists an increasing continuous singular function  $f$  on  $[0, 1]$  such that the set  $A_f$  of points where  $f$  has a nonzero finite derivative has Hausdorff dimension 1 in each subinterval of  $[0, 1]$ . We prove a stronger (and optimal) result showing that a set  $A_f$  as above can contain any prescribed  $F_\sigma$  null subset of  $[0, 1]$ .

## 1. INTRODUCTION

By a singular function we mean (following [5]) a continuous nonconstant function with zero derivative almost everywhere. Motivated by the observation that well-known singular functions have at no point a nonzero finite derivative, the authors of [5] constructed an increasing singular function  $f$  on  $[0, 1]$  such that the set  $A_f$  of points where  $f$  has a nonzero finite derivative is uncountable. Later it was shown in [6] that such  $A_f$  can be dense in  $[0, 1]$  and in [7] that  $A_f$  can have Hausdorff dimension 1 in each subinterval of  $[0, 1]$ .

We prove in the present note a stronger result showing that a set  $A_f$  as above can contain any prescribed  $F_\sigma$  null subset of  $[0, 1]$ , and that this result on the size of sets  $A_f$  is optimal. More precisely, we prove the following result.

**Theorem 1.1.** *Let  $A \subset [0, 1]$ . Then the following conditions are equivalent.*

- (i) *There exists a strictly increasing continuous singular function  $f$  on  $[0, 1]$  such that  $f$  has a nonzero finite derivative at each point of  $A$ .*
- (ii)  *$A$  is a subset of an  $F_\sigma$  Lebesgue null set.*

## 2. PROOF

Before the proof of Theorem 1.1, we will introduce some notation and prove two lemmas.

*Notation 2.1.* The symbol  $\lambda$  stands for the Lebesgue measure on  $\mathbb{R}$ . Recall that, by our definition, each singular function is continuous and nonconstant. For a function defined on an interval  $I$ , its limits and derivatives are computed with respect to  $I$ .

**Lemma 2.2.** *Let  $J = [a_1, a_2]$  be a closed interval. Let  $H \subset J$  be an  $F_\sigma$  set with  $\lambda(H) = 0$ . Then there exists a nondecreasing singular function  $\varphi$  on  $J$  such that*

- (i)  $\varphi(a_i) = a_i$ ,  $i = 1, 2$ ;
- (ii)  $\varphi'(x) = 0$ ,  $x \in H$ .

*Proof.* Since  $J \setminus H$  is  $G_\delta$  and dense in  $[0, 1]$ , its intersection with a null dense  $G_\delta$  set is a null  $G_\delta$  set  $S \subset J \setminus H$  which is uncountable by Baire category theorem. Since  $S$  is uncountable and  $G_\delta$ , [4, Theorem 13.6] implies that  $S$  contains a homeomorphic copy  $C$

---

2010 *Mathematics Subject Classification.* 26A30 .

*Key words and phrases.* increasing singular function, nonzero finite derivative,  $F_\sigma$  null set .

of the Cantor set. We can choose (e.g. by [2, Corollary 2.8.]) a nonatomic Borel finite measure  $\nu$  on  $C$  whose support is  $C$ . Now it is easy to see that the function  $\varphi$  defined by

$$\varphi(x) = \frac{(a_2 - a_1)}{\nu(C)} \nu(C \cap [a_1, x]) + a_1, \quad x \in J,$$

is nondecreasing singular and satisfies conditions (i) and (ii).  $\square$

We will say that  $P \subset (a, b)$  is an “infinite partition of  $(a, b)$ ” if  $a$  and  $b$  are the only accumulation points of  $P$ . Clearly,  $P$  has this property if and only if there exists a sequence  $(p_z)_{z \in \mathbb{Z}}$  such that  $P = \{p_z : z \in \mathbb{Z}\}$ ,  $p_z < p_{z+1}$ ,  $z \in \mathbb{Z}$ ,  $\inf\{p_z : z \in \mathbb{Z}\} = a$  and  $\sup\{p_z : z \in \mathbb{Z}\} = b$ . Any such sequence  $(p_z)_{z \in \mathbb{Z}}$  will be called an “ordering of  $P$ ”.

We will show that, for each  $(a, b)$ , there exists an infinite partition  $P$  of  $(a, b)$  such that, for each ordering  $(p_z)_{z \in \mathbb{Z}}$  of  $P$  and each  $z \in \mathbb{Z}$ ,

$$(1) \quad p_{z+1} - p_z < \min\{|p_z - a|^2, |p_{z+1} - b|^2\}.$$

To this end, choose an arbitrary infinite partition  $P^*$  of  $(0, 1)$ , its ordering  $(p_z^*)_{z \in \mathbb{Z}}$  and, for each  $z \in \mathbb{Z}$ , choose a finite partition  $p_z^* = t_{z,0} < t_{z,1} < \dots < t_{z,k_z} = p_{z+1}^*$  of  $[p_z^*, p_{z+1}^*]$  such that  $\max\{t_{z,i} - t_{z,i-1} : 1 \leq i \leq k_z\} < \min((p_z^* - a)^2, (b - p_{z+1}^*)^2)$ . Now it is easy to see that  $P := \{t_{z,i} : z \in \mathbb{Z}, 0 \leq i \leq k_z\}$  has the desired property.

**Lemma 2.3.** *Let  $\{0, 1\} \subset F \subset [0, 1]$  be a closed set with  $\lambda(F) = 0$ . Let  $M \subset (0, 1) \setminus F$  be an  $F_\sigma$  set with  $\lambda(M) = 0$ . Then there exists a nondecreasing singular function  $g$  on  $[0, 1]$  such that*

- (i)  $g(z) = z$ ,  $z \in F$ ;
- (ii)  $g'(z) = 0$ ,  $z \in M$ ;
- (iii)  $|g(x) - x| \leq \text{dist}^2(x, F)$ ,  $x \in [0, 1]$ .

*Proof.* Set  $G = (0, 1) \setminus F$ . Then  $G = \bigcup_{\alpha \in A} I_\alpha$ , where  $(I_\alpha)_{\alpha \in A}$  is a (nonempty countable) disjoint system of open intervals. For each  $\alpha \in A$ , choose an infinite partition  $P_\alpha$  of  $I_\alpha = (a_\alpha, b_\alpha)$  and its ordering  $(p_z^\alpha)_{z \in \mathbb{Z}}$  such that (1) holds for  $[a, b] := [a_\alpha, b_\alpha]$  and  $p_z := p_z^\alpha$ . For any  $\alpha \in A$  and  $z \in \mathbb{Z}$  we apply Lemma 2.2 to  $[a_1, a_2] := [p_z^\alpha, p_{z+1}^\alpha]$  and  $H := (M \cap [p_z^\alpha, p_{z+1}^\alpha]) \cup \{p_z^\alpha, p_{z+1}^\alpha\}$  and obtain a nondecreasing singular function  $\varphi_z^\alpha$  on  $[p_z^\alpha, p_{z+1}^\alpha]$  such that

$$(2) \quad \varphi_z^\alpha(p_z^\alpha) = p_z^\alpha, \quad \varphi_z^\alpha(p_{z+1}^\alpha) = p_{z+1}^\alpha,$$

$$(3) \quad (\varphi_z^\alpha)'_+(p_z^\alpha) = (\varphi_z^\alpha)'_-(p_{z+1}^\alpha) = 0 \quad \text{and} \quad (\varphi_z^\alpha)'(x) = 0, \quad x \in M \cap (p_z^\alpha, p_{z+1}^\alpha).$$

For each  $\alpha \in A$ , put  $g_\alpha(x) := \varphi_z^\alpha(x)$ ,  $x \in [p_z^\alpha, p_{z+1}^\alpha]$ . By (2), the definition is correct and  $g_\alpha$  is a continuous nondecreasing function on  $I_\alpha$  which has a.e. zero derivative. Moreover, (3) implies that  $g'_\alpha(x) = 0$  for each  $x \in M \cap I_\alpha$ .

Finally put  $g(x) := x$  if  $x \in F$  and  $g(x) := g_\alpha(x)$  if  $x \in I_\alpha$ . Then  $g$  is clearly nondecreasing on all  $[0, 1]$ , has a.e. zero derivative, is continuous on each  $I_\alpha$  and properties (i) and (ii) hold.

The inequality of (iii) is trivial for  $x \in F$ . If  $x \in [0, 1] \setminus F$ , then  $x \in I_\alpha = (a_\alpha, b_\alpha)$  for some  $\alpha \in A$  and we can choose  $z \in \mathbb{Z}$  such that  $x \in [p_z^\alpha, p_{z+1}^\alpha]$ . Then we have  $g(x) = g_\alpha(x) = \varphi_z^\alpha(x)$  and, since  $\varphi_z^\alpha(x) \in [p_z^\alpha, p_{z+1}^\alpha]$ , we obtain

$$|g(x) - x| = |\varphi_z^\alpha(x) - x| \leq p_{z+1}^\alpha - p_z^\alpha \leq \min\{|p_z^\alpha - a_\alpha|^2, |p_{z+1}^\alpha - b_\alpha|^2\} \leq \text{dist}^2(x, F)$$

and so (iii) is proved.

It remains to prove that  $g$  is continuous on all  $[0, 1]$ . We know that  $g$  is continuous on  $G$  and (iii) implies that, if  $a \in F$  is fixed, we have for each  $x \in [0, 1]$

$$|g(x) - g(a)| = |g(x) - a| \leq |g(x) - x| + |x - a| \leq |x - a|^2 + |x - a| \rightarrow 0, \quad x \rightarrow a.$$

□

**Proof of Theorem.**

To prove the implication “(i)  $\Rightarrow$  (ii)”, suppose that  $f$  is a singular increasing function on  $[0, 1]$  and  $0 < f'(x) < \infty$  for each  $x \in A$ . Let  $f^*$  be a continuous extension of  $f$  to  $\mathbb{R}$  and denote  $E_f := \{x \in (0, 1) : -\infty < f'(x) < \infty\}$ . Then  $f'(x) = \lim_{n \rightarrow \infty} n(f^*(x + 1/n) - f^*(x))$ ,  $x \in E_f$ , and so  $f'|_{E_f}$  is a first Baire class function on  $E_f$ . Consequently  $P := \{x \in E_f : f'(x) > 0\}$  is an  $F_\sigma$  subset in the space  $E_f$  (see e.g. [1, Theorem 10.12]) and consequently there exists an  $F_\sigma$  set  $H \subset [0, 1]$  such that  $P = E_f \cap H$ . Since  $f$  is singular,  $P$  is a null set,  $[0, 1] \setminus E_f$  is a null set and consequently also  $H \subset P \cup ([0, 1] \setminus E_f)$  is null. Therefore  $A \subset P \cup \{0, 1\}$  is contained in an  $F_\sigma$  null set  $H \cup \{0, 1\}$ .

To prove the implication “(ii)  $\Rightarrow$  (i)”, let  $A \subset M$ , where  $M \subset [0, 1]$  is an  $F_\sigma$  null set. Write  $M = \bigcup_{n=1}^\infty F_n$ , where  $F_n$  are closed subsets of  $[0, 1]$ . Without any loss of generality we can assume that  $M$  is dense in  $[0, 1]$ ,  $F_1 \subset F_2 \subset F_3 \subset \dots$  and  $\{0, 1\} \subset F_1$ .

Set  $M_n := M \setminus F_n$ ,  $n \in \mathbb{N}$ . Then  $M_n$  is an  $F_\sigma$  set with  $\lambda(M_n) = 0$  and  $M_n \subset (0, 1) \setminus F_n$ . Applying Lemma 2.3 to the sets  $F_n$  and  $M_n$  we obtain nondecreasing singular functions  $g_n$  on  $[0, 1]$  such that, for each  $n \in \mathbb{N}$ ,

(4) the conditions (i) - (iii) of Lemma 2.3 hold for  $g := g_n$ ,  $F := F_n$ ,  $M := M_n$ .

Now set

$$f(x) = \sum_{n=1}^\infty \frac{1}{2^n} g_n(x), \quad x \in [0, 1].$$

Clearly  $f$  is continuous and nondecreasing on  $[0, 1]$ . According to the Fubini theorem on derivative of a sum of monotone functions [3, Theorem 17.18] the function  $f$  has zero derivative almost everywhere.

Now we will prove that

(5) at each point  $a \in M$ , there exists a nonzero finite  $f'(a)$ .

So fix an arbitrary  $a \in M$ . Let  $n \in \mathbb{N}$  be the natural number such that  $a \in F_n \setminus F_{n-1}$ , (where  $F_0 := \emptyset$ ). If  $1 \leq k < n$ , then  $a \in M_k$  and so (4) implies

$$\left(\frac{1}{2^k} g_k\right)'(a) = 0.$$

Consequently, to prove  $0 < f'(a) < \infty$ , it is sufficient to show that

$$r'_n(a) = \sum_{k=n}^\infty \frac{1}{2^k}, \quad \text{where } r_n = \sum_{k=n}^\infty \frac{1}{2^k} g_k.$$

And this equality is true since, using (4) and  $a \in F_k$  for  $k \geq n$ , we obtain

$$\begin{aligned} |r_n(x) - r_n(a) - \sum_{k=n}^\infty \frac{1}{2^k} (x - a)| &\leq \sum_{k=n}^\infty \frac{1}{2^k} |g_k(x) - g_k(a) - (x - a)| \\ &= \sum_{k=n}^\infty \frac{1}{2^k} |g_k(x) - x| \leq \sum_{k=n}^\infty \frac{1}{2^k} \text{dist}^2(x, F_k) \leq |x - a|^2 = o(|x - a|), \quad x \rightarrow a. \end{aligned}$$

Observe that  $f$  is strictly increasing since it is nondecreasing,  $M$  is dense and (5) holds. Using (5) and  $A \subset M$ , we conclude that (i) holds, since  $f$  has all desired properties.

#### REFERENCES

- [1] A.M. Bruckner, J.B.Bruckner, B.S.Thomson, *Real Analysis*, Upper Saddle River, NJ, Prentice Hall, (1997).
- [2] D. Hebert, H. Lacey, *On supports of regular Borel measures*, Pac. J. Math., **27** (1968), 101–118.
- [3] E. Hewitt, K. R. Stromberg, *Real and Abstract Analysis. A Modern Treatment of the Theory of a Real Variable*, New York, Springer-Verlag, (1969).
- [4] A. Kechris, *Classical Descriptive Set Theory*, New York, Springer-Verlag, (1995).
- [5] J. F. Sánchez, P. Viader, J. Paradís, M. D. Carrillo, *A singular function with a non-zero finite derivative*, Nonlinear Anal., **75** (2012), 5010–5014.
- [6] J. F. Sánchez, P. Viader, J. Paradís, M. D. Carrillo, *A singular function with a non-zero finite derivative on a dense set*, Nonlinear Anal., **95** (2014), 703–713.
- [7] J. F. Sánchez, P. Viader, J. Paradís, M. D. Carrillo, *A singular function with a non-zero finite derivative on a dense set with Hausdorff dimension one*, J. Math. Anal. Appl., **434** (2016), 713–728.

*Email address:* kossaczka@karlin.mff.cuni.cz

*Email address:* zajicek@karlin.mff.cuni.cz

CHARLES UNIVERSITY, FACULTY OF MATHEMATICS AND PHYSICS, SOKOLOVSKÁ 83, 186 75 PRAHA 8-KARLÍN, CZECH REPUBLIC