

# The Fixed- $b$ Limiting Distribution and the ERP of HAR Tests Under Nonstationarity\*

ALESSANDRO CASINI<sup>†</sup>

University of Rome Tor Vergata

30th November 2021

## Abstract

We show that the limiting distribution of HAR test statistics under fixed- $b$  asymptotics is not pivotal when the data are nonstationary. It takes the form of a complicated function of Gaussian processes and depends on the second moments of the relevant series (e.g., of the regressors and errors for the case of the linear regression model). Hence, existing fixed- $b$  inference methods based on stationarity are not theoretically valid in general. The nuisance parameters entering the fixed- $b$  limiting distribution can be consistently estimated under small- $b$  asymptotics but only with nonparametric rate of convergence. We show that the error in rejection probability (ERP) is an order of magnitude larger than that under stationarity and is also larger than that of HAR tests based on HAC estimators under conventional asymptotics. These theoretical results reconcile with recent finite-sample evidence showing that fixed- $b$  HAR tests can perform poorly when the data are nonstationary. They can be conservative under the null hypothesis and have non-monotonic power under the alternative hypothesis irrespective of how large the sample size is.

**JEL Classification:** C12, C13, C18, C22, C32, C51

**Keywords:** Asymptotic expansion, Fixed- $b$ , HAC standard errors, HAR inference, Long-run variance, Nonstationarity.

---

\*I am grateful to Pierre Perron for his support and advice. I thank Andrew Chesher, Jungbin Hwang, Oliver Linton, Julius Vainora, Daniel Whitem and seminar participants at University College London, University of Cambridge and University of Connecticut for comments.

<sup>†</sup>Department of Economics and Finance, University of Rome Tor Vergata, Via Columbia 2, Rome 00133, IT. Email: [alessandro.casini@uniroma2.it](mailto:alessandro.casini@uniroma2.it).

# 1 Introduction

The construction of standard errors robust to autocorrelation and heteroskedasticity is important for empirical work because economic and financial time series exhibit temporal dependence. The early literature focused on heteroskedasticity and autocorrelation consistent (HAC) estimators of the asymptotic variance of test statistics (or simply the long-run variance (LRV) of the relevant series) [see, e.g., [Newey and West \(1987; 1994\)](#), [Andrews \(1991\)](#), [Andrews and Monahan \(1992\)](#), [Hansen \(1992\)](#), [de Jong and Davidson \(2000\)](#)]. This approach aims at devising a good estimate of the LRV. Over the last twenty years, the literature has focused on methods based on fixed- $b$  asymptotics. These involve an inconsistent estimate of the LRV that keeps the bandwidth at a fixed fraction of the sample size. This approach was initiated by [Kiefer, Vogelsang, and Bunzel \(2000\)](#) and [Kiefer and Vogelsang \(2002a; 2002b\)](#). They developed the analysis assuming stationarity and showed that valid heteroskedasticity and autocorrelation robust (HAR) inference is feasible even without a consistent estimator of the LRV. Inconsistency results in a pivotal nonstandard limiting distribution whose critical value can be obtained by simulations (e.g., a  $t$ -statistic on a coefficient in the linear regression model will not follow asymptotically a standard normal distribution but a distribution involving a ratio of Gaussian processes). Theoretical results based on asymptotic expansions suggested that fixed- $b$  HAR test statistics exhibit an error in rejection probability (ERP) that is smaller than that associated to test statistics based on HAC estimators [see [Jansson \(2004\)](#) and [Sun, Phillips, and Jin \(2008\)](#)]. This supported extensive finite-sample evidence in the literature documenting that the fixed- $b$  approach leads to HAR test statistics with more accurate null rejection rates when the data are stationary with strong temporal dependence than those associated to test statistics based on HAC estimators. Since then the literature has mostly concentrated on various refinements of fixed- $b$  HAR inference while maintaining the stationarity assumption, mostly to have tests having null rejection rates closer to the nominal level.

Although stationarity rarely holds in economic and financial time series, the literature has surprisingly ignored investigating the theoretical and empirical properties of existing fixed- $b$  HAR inference when stationarity does not hold.<sup>1</sup> Nonstationarity can occur for several reasons: changes in the moments of the relevant time series induced by changes in the model parameters that govern the data (think about the Great Moderation with the decline in variance for many macroeconomic variables, the effects of the financial crisis of 2007–2008 or of the COVID-19 pandemic); smooth changes in the distributions governing the data that arise from transitory dynamics from one

---

<sup>1</sup>By nonstationary we mean non-constant moments. As in the literature, we consider processes whose sum of absolute autocovariances is finite. This rules out processes with unbounded second moments (e.g., unit root).

regime to another. Unfortunately, the theoretical properties of fixed- $b$  HAR inference change substantially when stationarity does not hold. The contribution of the paper is to establish such theoretical results and discuss their relevance for inference in empirical work.

We show that the limiting distribution of HAR test statistics under fixed- $b$  asymptotics is not pivotal when the data are nonstationary. It takes the form of a complicated function of Gaussian processes and depends on the second moments of the relevant series. For example, in the case of the linear regression model, it depends on the second moments of the regressors and errors. Hence, existing fixed- $b$  inference methods based on stationarity are not theoretically valid in general. The nuisance parameters entering the fixed- $b$  limiting distribution can be consistently estimated under small- $b$  asymptotics but only with nonparametric rate of convergence. We develop asymptotic expansions under nonstationarity and we show that the ERP is an order of magnitude larger than that obtained under stationarity by Jansson (2004) and Sun, Phillips, and Jin (2008) [cf.  $O(T^{-\gamma})$  with  $\gamma < 1/2$  versus  $O(T^{-1})$  where  $T$  is the sample size]. Further, we show that the ERP of fixed- $b$  HAR tests is also larger than that of HAR tests based on HAC estimators. It follows from our results that if one uses existing fixed- $b$  methods based on the pivotal fixed- $b$  limiting distribution obtained under stationarity but the data are nonstationary, then the ERP does not even converge to zero as the sample size increases because that is not the correct limiting distribution.

The pivotal property breaks down because under nonstationarity the LRV estimator that uses a fixed-bandwidth is not asymptotically proportional to the LRV. A non-pivotal limiting distribution results in a much more complex type of inference in practice. It is beyond the scope of this paper to investigate how best to use the non-pivotal limiting distribution for inference. The increase in the ERP from the stationary case arises from fact that the nuisance parameters have to be estimated. It is the discrepancy between these estimates and their probability limits that is reflected in the leading term of the asymptotic expansion.

Our theoretical results reconcile with recent finite-sample evidence that showed that fixed- $b$  HAR tests can perform poorly when the data are nonstationary. These issues have been documented extensively by Belotti et al. (2021), Casini (2021b), Casini and Perron (2021) and Casini, Deng, and Perron (2021) who considered  $t$ -tests in the linear regression models as well as HAR tests outside the linear regression model, and a variety of data-generating processes. They provided evidence that fixed- $b$  HAR tests can be severely undersized and can exhibit non-monotonic power. The more nonstationary the data are, the stronger the distortions. This is especially visible in HAR inference contexts characterized by a stationary null hypothesis and a nonstationary alternative hypothesis [e.g., tests for structural breaks, tests for regime-switching, tests for time-

varying parameters and threshold effects, and tests for forecast evaluation]. In such cases, the power of fixed- $b$  HAR tests can be zero irrespective of how large the sample size is and how far the alternative is from the null value.

Although we discuss the original fixed- $b$  approach as proposed by [Kiefer, Vogelsang, and Bunzel \(2000\)](#) and [Kiefer and Vogelsang \(2002a; 2002b\)](#), the same issues apply to the refinements considered more recently by several authors; for references see, e.g., [Casini \(2021b\)](#) and [Lazarus, Lewis, and Stock \(2020\)](#). For example, [Sun, Phillips, and Jin \(2008\)](#) proposed to use the second-order corrected critical value based on the expansion of the pivotal fixed- $b$  limiting distribution which, however, is not valid under nonstationarity. A similar feature applies to the more recent frequency-domain methods proposed by [Lazarus, Lewis, and Stock \(2020\)](#). On the other hand, using consistent LRV estimators is valid also under nonstationarity and results in an invariant standard asymptotic distribution. The theoretical high-order refinement provided by fixed- $b$  HAR inference relative to HAC-based HAR inference which was justified by a smaller ERP under stationarity, does not hold under nonstationarity in which case the opposite is true.

The remainder of the paper is organized as follows. Section 2 introduces the statistical problem in the well-known setting of the linear regression model. In Section 3 we study the limiting distribution of  $t$ - and  $F$ -type test statistics. Section 4 develops the asymptotic expansions and presents the results on the ERP. Section 5 concludes the paper. The supplemental material [cf. [Casini \(2021a\)](#)] contains all mathematical proofs.

## 2 HAR Testing in the Linear Regression Model

We consider the linear regression model

$$y_t = x_t' \beta_0 + e_t, \quad t = 1, 2, \dots, T, \quad (2.1)$$

where  $\beta_0 \in \Theta \subset \mathbb{R}^p$ ,  $y_t$  is an observation on the dependent variable,  $x_t$  is a  $p$ -vector of regressors and  $e_t$  is an unobserved disturbance that is autocorrelated and possibly conditionally heteroskedastic, and  $\mathbb{E}(e_t | x_t) = 0$ . The problem addressed is testing linear hypotheses about  $\beta_0$ . We consider the ordinary least squares (OLS) estimator  $\hat{\beta} = (\sum_{t=1}^T x_t x_t')^{-1} \sum_{t=1}^T x_t y_t$ . Let  $V_t = x_t e_t$ . Define  $S_{[Tr]} = \sum_{t=1}^{[Tr]} V_t$  where  $[Tr]$  denotes the integer part of  $Tr$ . Using ordinary manipulations,

$$\sqrt{T}(\hat{\beta} - \beta_0) = \left( T^{-1} \sum_{t=1}^T x_t x_t' \right)^{-1} T^{-1/2} S_T.$$

The variance of  $T^{-1/2}S_T$  plays an important role for constructing tests about  $\beta_0$ . Its exact formula depends on the assumptions about  $\{V_t\}$ . We begin with the following notational conventions.

A function  $g(\cdot) : [0, 1] \mapsto \mathbb{R}$  is said to be piecewise (Lipschitz) continuous if it is (Lipschitz) continuous except on a set of discontinuity points that has zero Lebesgue measure. A matrix is said to be piecewise (Lipschitz) continuous if each of its element is piecewise (Lipschitz) continuous. Let  $W_p(r)$  denote a  $p$ -vector of independent standard Wiener processes where  $r \in [0, 1]$ . We use  $\xrightarrow{\mathbb{P}}$ ,  $\Rightarrow$  and  $\xrightarrow{d}$  to denote convergence in probability, weak convergence and convergence in distribution, respectively. The following assumptions are sufficient to establish the asymptotic distribution of the test statistics. Let  $\Omega(u)$  denote some  $p \times p$  positive semidefinite matrix.

**Assumption 2.1.**  $T^{-1/2}S_{[Tr]} \Rightarrow \int_0^r \Sigma(u) dW_p(u)$  where  $\Sigma(u)$  is given by the Cholesky decomposition  $\Omega(u) = \Sigma(u) \Sigma(u)'$  and is piecewise continuous with  $\sup_{u \in [0, 1]} \|\Sigma(u)\| < \infty$ .

**Assumption 2.2.**  $T^{-1} \sum_{t=1}^{[Tr]} x_t x_t' \xrightarrow{\mathbb{P}} \int_0^r Q(u) du$  uniformly in  $r$  where  $Q(u)$  is piecewise continuous with  $\sup_{u \in [0, 1]} \|Q(u)\| < \infty$ .

Assumption 2.1 states a functional law for nonstationary processes [see, e.g., Aldous (1978) and Merlevède, Peligrad, and Utev (2019)]. If  $\{V_t\}$  is second-order stationary, then  $\Sigma(u) = \Sigma$  for all  $u$  and Assumption 2.1 reduces to  $T^{-1/2}S_{[Tr]} \Rightarrow \Sigma W_p(r)$ . The fixed- $b$  literature has routinely used the assumption of second-order stationarity [see, e.g., Kiefer, Vogelsang, and Bunzel (2000), Jansson (2004), Sun, Phillips, and Jin (2008) and Lazarus, Lewis, and Stock (2020)]. We relax this assumption substantially as we allow for general time-variation in the second moments of the regressors and errors which encompasses most of the nonstationary processes used in econometrics and statistics. For example, it allows for structural breaks, regime-switching, time-varying parameters and segmented local stationarity in the second moments of  $\{V_t\}$ . With regards to the temporal dependence, Assumption 2.1 holds under a variety of regularity conditions. For example, standard mixing conditions and (time-varying) invertible ARMA processes are allowed.

Assumption 2.2 allows for structural breaks as well as smooth variation in the second moments of the regressors.<sup>2</sup> The fixed- $b$  literature required  $Q(u) = Q$  for all  $u$  in which case Assumption 2.2 reduces to  $T^{-1} \sum_{t=1}^{[Tr]} x_t x_t' \xrightarrow{\mathbb{P}} rQ$ . The latter is quite restrictive in practice. The uniform convergence, boundness and positive definiteness of  $Q(\cdot)$  are satisfied for a fairly general class of

---

<sup>2</sup>Assumption 2.2 also allows for polynomial trending regressors as long as they are written in the form  $(t/T)^l$  ( $l \geq 0$ ), or more generally, written as a piecewise continuous function of the time trend, say  $g(t/T)$ .

processes. As in previous works, Assumption 2.1-2.2 rule out unit roots and long memory. Let

$$\text{Var} \left( T^{-1/2} S_T \right) = \sum_{k=-T+1}^{T-1} \Gamma_{T,k}, \quad \Gamma_{T,k} = \begin{cases} T^{-1} \sum_{t=k+1}^T \mathbb{E}(V_t V_{t-k}') & \text{for } k \geq 0 \\ T^{-1} \sum_{t=-k+1}^T \mathbb{E}(V_{t+k} V_t') & \text{for } k < 0 \end{cases}. \quad (2.2)$$

Under Assumption 2.1-2.2, the limit of  $\text{Var}(T^{-1/2} S_T)$  is given by [cf. Casini (2021b)]

$$\lim_{T \rightarrow \infty} \text{Var} \left( T^{-1/2} S_T \right) \triangleq \Omega = \int_0^1 c(u, 0) du + \sum_{k=1}^{\infty} \int_0^1 \left( c(u, k) + c(u, k)' \right) du,$$

where  $c(u, k) = \mathbb{E}(V_{[Tu]} V_{[Tu]-k}) + O(T^{-1})$ . By the Cholesky decomposition  $\Omega(u) = \Sigma(u) \Sigma(u)'$  and so  $\Omega = \int_0^1 \Omega(u) du$ . Note that  $\Omega = 2\pi \int_0^1 f(u, 0) du$  where  $f(u, 0)$  is the local spectral density matrix of  $\{V_t\}$  at rescaled time  $u$  and frequency 0. For  $u$  a continuity point,  $f(u, \omega)$  is defined implicitly by the relation  $\mathbb{E}(V_{[Tu]} V_{[Tu]-k}) = \int_{-\pi}^{\pi} e^{i\omega k} f(u, \omega) d\omega$ ; see Casini (2021b) for more details. If  $\{V_t\}$  is second-order stationary, then  $\Omega = \Sigma \Sigma' = 2\pi f(0)$  since  $f(u, 0) = f(0)$ .

Under Assumption 2.1-2.2, it directly follows, using standard arguments, that

$$\sqrt{T} \left( \hat{\beta} - \beta_0 \right) \xrightarrow{d} \overline{Q}^{-1} \Omega^{1/2} W_p(1) \sim \mathcal{N} \left( \mathbf{0}, \overline{Q}^{-1} \Omega \overline{Q}^{-1} \right), \quad (2.3)$$

where  $\Omega^{1/2}$  is the matrix square-root of  $\Omega$  and  $\overline{Q} \triangleq \int_0^1 Q(u) du$ . Under second-order stationarity  $Q(u) = Q$ ,  $\Sigma(u) = \Sigma$  and (2.3) reduces to

$$\sqrt{T} \left( \hat{\beta} - \beta_0 \right) \xrightarrow{d} Q^{-1} \Sigma W_p(1) \sim \mathcal{N} \left( \mathbf{0}, Q^{-1} \Omega Q^{-1} \right). \quad (2.4)$$

The classical approach to testing hypotheses about  $\beta_0$  is based on studentization. Provided that a consistent estimator of  $\overline{Q}^{-1} \Omega \overline{Q}^{-1}$  can be constructed, it is possible to construct a test statistic whose asymptotic distribution is free of nuisance parameters. The term  $\overline{Q}$  can be consistently estimated straightforwardly using  $\hat{Q} = T^{-1} \sum_{t=1}^T x_t x_t'$ . Consistent estimators of  $\Omega$  are known as HAC estimators [see, e.g., Newey and West (1987), Andrews (1991), de Jong and Davidson (2000) and Casini (2021b)]. HAC estimators take the following general form,

$$\hat{\Omega}_{\text{HAC}} \triangleq \sum_{k=-T+1}^{T-1} K(b_T k) \hat{\Gamma}(k),$$

where

$$\hat{\Gamma}(k) = \begin{cases} T^{-1} \sum_{t=k+1}^T \hat{V}_t \hat{V}_{t-k}' & \text{for } k \geq 0 \\ T^{-1} \sum_{t=-k+1}^T \hat{V}_{t+k} \hat{V}_t' & \text{for } k < 0 \end{cases}, \quad (2.5)$$

$\hat{V}_t = x_t \hat{e}_t$  and  $\{\hat{e}_t\}$  are the OLS residuals,  $K(\cdot)$  is a kernel and  $b_T$  is a bandwidth sequence. Under  $b_T \rightarrow 0$  at an appropriate rate, we have  $\hat{\Omega}_{\text{HAC}} \xrightarrow{\mathbb{P}} \Omega$ . An alternative to  $\hat{\Omega}_{\text{HAC}}$  is the double-kernel HAC (DK-HAC) estimator, say  $\hat{\Omega}_{\text{DK-HAC}}$ , proposed by [Casini \(2021b\)](#) to flexibly account for nonstationarity.  $\hat{\Omega}_{\text{DK-HAC}}$  uses an additional kernel for smoothing over time; see [Casini \(2021b\)](#) for details. Under appropriate conditions on the bandwidths, we have  $\hat{\Omega}_{\text{DK-HAC}} \xrightarrow{\mathbb{P}} \Omega$ . Hence, equipped with either  $\hat{\Omega}_{\text{HAC}}$  or  $\hat{\Omega}_{\text{DK-HAC}}$ , HAR inference is standard because test statistics follow asymptotically standard distributions.

An alternative approach to HAR inference relies on inconsistent estimation of  $\Omega$ . [Kiefer and Vogelsang \(2002a, 2002b\)](#) proposed to use the following estimator,

$$\hat{\Omega}_{\text{fixed-}b} \triangleq \sum_{k=-T+1}^{T-1} K(k/T) \hat{\Gamma}(k),$$

which is equivalent to  $\hat{\Omega}_{\text{HAC}}$  with  $b_T = T^{-1}$ .<sup>3</sup>  $\hat{\Omega}_{\text{fixed-}b}$  is inconsistent for  $\Omega$ . [Kiefer, Vogelsang, and Bunzel \(2000\)](#) showed that an asymptotic distribution theory for HAR tests is possible even with an inconsistent estimate of  $\Omega$ . One first has to derive the limiting distribution of  $\hat{\Omega}_{\text{fixed-}b}$  under the null hypothesis. Then, one can use it to obtain the limiting distribution of the test statistic of interest which typically involves a ratio of Gaussian processes. Thus, from the inconsistency of  $\hat{\Omega}_{\text{fixed-}b}$ , HAR test statistics will not follow asymptotically standard distributions.

### 3 Fixed- $b$ Limiting Distribution of HAR Tests

In this section we study the limiting distribution of the HAR tests for linear hypothesis in the linear regression model under fixed- $b$  asymptotics when the data are nonstationary. Consider testing the null hypothesis  $H_0 : R\beta_0 = r$  against the alternative hypothesis  $H_1 : R\beta_0 \neq r$  where  $R$  is a  $q \times p$

---

<sup>3</sup>The label fixed- $b$  refers to that fact that one can define more generally  $\hat{\Omega}_{\text{fixed-}b} = \sum_{k=-T+1}^{T-1} K(k/(bT)) \hat{\Gamma}(k)$  with  $b \in (0, 1]$  fixed.

matrix with rank  $q$  and  $r$  is a  $q \times 1$  vector. Using  $\hat{\Omega}_{\text{fixed}-b}$  an  $F$ -test can be constructed as follows:

$$F_{\text{fixed}-b} = T \left( R\hat{\beta} - r \right)' \left[ R\hat{Q}^{-1} \hat{\Omega}_{\text{fixed}-b} \hat{Q}^{-1} R' \right]^{-1} \left( R\hat{\beta} - r \right) / q.$$

For testing one restriction,  $q = 1$ , one can use the following  $t$ -statistic:

$$t_{\text{fixed}-b} = \frac{T^{1/2} \left( R\hat{\beta} - r \right)}{\sqrt{R\hat{Q}^{-1} \hat{\Omega}_{\text{fixed}-b} \hat{Q}^{-1} R'}}.$$

Let  $B_p(r) = W_p(r) - rW_p(1)$  denote the  $p \times 1$  vector of Brownian bridges. Consider the following class of kernels,

$$\begin{aligned} \mathbf{K} = \{ K(\cdot) : \mathbb{R} \rightarrow [-1, 1] : K(0) = 1, K(x) = K(-x), \forall x \in \mathbb{R} \\ \int_{-\infty}^{\infty} K^2(x) dx < \infty, K(\cdot) \text{ is continuous at } 0 \}. \end{aligned} \quad (3.1)$$

Examples of kernels in  $\mathbf{K}$  include the Truncated, Bartlett, Parzen, Quadratic Spectral (QS) and Tukey-Hanning kernels. [Kiefer, Vogelsang, and Bunzel \(2000\)](#) and [Kiefer and Vogelsang \(2002b\)](#) showed under stationarity that

$$\hat{\Omega}_{\text{fixed}-b} \Rightarrow \Sigma \left( \int_0^1 \int_0^1 -K''(r-s) B_p(r) B_p(s)' dr ds \right) \Sigma', \quad (3.2)$$

for  $K \in \mathbf{K}$  with  $K''(x)$  assumed to exist for  $x \in [-1, 1]$  and to be continuous.<sup>4</sup> A key feature of the result in (3.2) is that  $\hat{\Omega}_{\text{fixed}-b}$  is asymptotically proportional to  $\Omega$  through  $\Sigma\Sigma'$ . The null asymptotic distributions of  $F_{\text{fixed}-b}$  and  $t_{\text{fixed}-b}$  under stationarity are given, respectively, by

$$F_{\text{fixed}-b} \Rightarrow W_q(1)' \left[ \int_0^1 \int_0^1 -K''(r-s) B_q(r) B_q(s)' dr ds \right]^{-1} W_q(1) / q,$$

---

<sup>4</sup>Note that  $K''(x)$  does not exist for some popular kernels. This is the case for the Bartlett kernel for which  $K''(0)$  does not exist. However, [Kiefer and Vogelsang \(2002a\)](#) showed that for the Bartlett kernels it holds that

$$\hat{\Omega}_{\text{fixed}-b} \Rightarrow 2\Sigma \left( \int_0^1 B_p(r) B_p(r)' dr \right) \Sigma'.$$

Recall that the Bartlett kernel is defined as  $K_{\text{BT}}(x) = 1 - |x|$  for  $|x| \leq 1$  and  $K_{\text{BT}}(x) = 0$  otherwise.



and

$$t_{\text{fixed}-b} \Rightarrow \frac{W_1(1)}{\sqrt{\int_0^1 \int_0^1 -K''(r-s) B_q(r) B_q(s) dr ds}}.$$

Both null distributions are pivotal. Thus, valid testing is possible without consistent estimation of  $\Omega$ . This result crucially hinges on stationarity. To see this, consider  $t_{\text{fixed}-b}$  for the single-regressor case ( $p = 1$ ) and for the null hypothesis  $H_0 : \beta_0 = 0$ . Its numerator and denominator are asymptotically equivalent to, respectively,  $Q^{-1}\Sigma W_1(1)$  and

$$Q^{-1}\Sigma \left( \int_0^1 \int_0^1 -K''(r-s) B_1(r) B_1(s) dr ds \right)^{1/2}.$$

Since  $W_1(1)$  and  $B_1(r)$  are independent,  $t_{\text{fixed}-b}$  is a ratio of two independent random variables. The factor  $Q^{-1}\Sigma$  cancels because it appears in both numerator and denominator. It follows that the asymptotic null distribution is pivotal. We show that this argument break downs when stationarity does not hold. Under nonstationarity the factor in the denominator corresponding to  $Q^{-1}\Sigma$  will depend on the rescaled time  $s$  and  $r$ , and enter the integrand. Thus, it will not cancel out.

We now present the results about the fixed- $b$  limiting distribution of the HAR tests. We begin with the following theorem which provides the limiting distribution of  $\hat{\Omega}_{\text{fixed}-b}$ .

**Theorem 3.1.** *Let Assumption 2.1-2.2 hold and  $K \in \mathbf{K}$ . Then, we have: (i) If  $K''(x)$  exists for  $x \in [-1, 1]$  and is continuous, then*

$$\begin{aligned} \hat{\Omega}_{\text{fixed}-b} &\Rightarrow - \int_0^1 \int_0^1 K''(r-s) \left( \int_0^r \Sigma(u) dW_p(u) - \left( \int_0^r Q(u) du \right) \bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right) \\ &\quad \times \left( \int_0^s \Sigma(u) dW_p(u) - \left( \int_0^s Q(u) du \right) \bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right)' dr ds \\ &\triangleq \mathcal{G}. \end{aligned} \tag{3.3}$$

(ii) If  $K(x) = K_{\text{BT}}(x)$ , then

$$\begin{aligned} \hat{\Omega}_{\text{fixed}-b} &\Rightarrow 2 \int_0^1 \left( \int_0^r \Sigma(u) dW_p(u) - \left( \int_0^r Q(u) du \right) \bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right) \\ &\quad \times \left( \int_0^r \Sigma(u) dW_p(u) - \left( \int_0^r Q(u) du \right) \bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right)' dr \\ &\triangleq \mathcal{G}_{\text{BT}}. \end{aligned} \quad (3.4)$$

Theorem 3.1 show that, unlike in the stationary case,  $\hat{\Omega}_{\text{fixed}-b}$  is not asymptotically proportional to  $\Omega$ . This anticipates that asymptotically pivotal tests for null hypotheses involving  $\beta_0$  cannot be constructed. The limiting distribution depends on  $K''(\cdot)$ , and most importantly on  $\Sigma(\cdot)$  and  $Q(\cdot)$  so that it is not free of nuisance parameters.

We now present the limiting distribution of  $F_{\text{fixed}-b}$  and  $t_{\text{fixed}-b}$  under  $H_0$ .

**Theorem 3.2.** *Let Assumption 2.1-2.2 hold and  $K \in \mathbf{K}$ . Then, we have: (i) If  $K''(x)$  exists for  $x \in [-1, 1]$  and is continuous, then*

$$F_{\text{fixed}-b} \Rightarrow \left( R\bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right)' \left( R\bar{Q}^{-1} \mathcal{G} \bar{Q}^{-1} R' \right)^{-1} R\bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) / q,$$

where  $\mathcal{G}$  is defined in (3.3). If  $q = 1$ , then

$$t_{\text{fixed}-b} \Rightarrow \frac{R\bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u)}{\sqrt{R\bar{Q}^{-1} \mathcal{G} \bar{Q}^{-1} R'}}.$$

(ii) If the Bartlett kernel is used,  $K(x) = K_{\text{BT}}(x)$ , then

$$F_{\text{fixed}-b} \Rightarrow \left( R\bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right)' \left( R\bar{Q}^{-1} \mathcal{G}_{\text{BT}} \bar{Q}^{-1} R' \right)^{-1} R\bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) / q,$$

and for  $q = 1$ ,

$$t_{\text{fixed}-b} \Rightarrow \frac{R\bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u)}{\sqrt{R\bar{Q}^{-1} \mathcal{G}_{\text{BT}} \bar{Q}^{-1} R'}}.$$

Theorem 3.2 shows that the asymptotic distribution of the  $F$  and  $t$  test statistics under fixed- $b$  asymptotics under nonstationarity are not pivotal. This contrasts with the stationary case. Consequently, fixed- $b$  inference based on stationarity is not theoretically valid under nonstationarity.

The limiting distributions of  $F_{\text{fixed}-b}$  and  $t_{\text{fixed}-b}$  depend on nuisance parameters such as the time-varying autocovariance function of  $\{V_t\}$  through  $\Sigma(\cdot)$  and the second moments of the regressors through  $Q(\cdot)$ . An inspection of the proof shows that it is practically impossible to make  $F_{\text{fixed}-b}$  and  $t_{\text{fixed}-b}$  pivotal by studentization based on any sequence of inconsistent covariance matrix estimates. This follows because the long-run variance is time-varying and, as noted above, this breaks down the property that both numerator and denominator are asymptotically proportional to  $\Omega$  so that the nuisance parameters cancel out.

Theorem 3.2 suggests that valid inference under fixed- $b$  asymptotics is going to be more complex in terms of practical implementation relative to when the data are stationary. In the literature, complexity in the implementation has been recognized as a strong disadvantage for the success of a given method in empirical work [see, e.g., Lazarus, Lewis, and Stock (2020)]. The simplest way to use Theorem 3.2 for conducting inference is to replace the nuisance parameters by consistent estimates. This means constructing estimates of  $\Sigma(u)$ ,  $Q(u)$  and  $\overline{Q}$ . For  $\overline{Q}$  the argument is straightforward. As under stationarity, one can use  $\hat{Q} = T^{-1} \sum_{t=1}^T x_t x_t'$  since  $\hat{Q} - \overline{Q} \xrightarrow{\mathbb{P}} 0$  also under nonstationarity. More complex is the case for  $\Sigma(u)$  and  $Q(u)$ . Nonparametric estimators for  $\Sigma(u)$  and  $Q(u)$  can be constructed. This requires introducing bandwidths and kernels as well as a criterion for their choice. Then, one plugs-in these estimates into the limit distribution and the critical value can be obtained by simulations. However, since  $\Sigma(u)$ ,  $Q(u)$  and  $\overline{Q}$  depend on the data, the critical values need to be obtained on a case-by-case basis.

Before concluding this section, we discuss the properties of fixed- $b$  HAR inference when  $\{V_t\}$  follows more general forms of nonstationary, i.e.,  $\{V_t\}$  does not satisfy Assumption 2.1-2.2. Assumption 2.1-2.2 are satisfied if  $\{V_t\}$  is, e.g., segmented locally stationary, locally stationary and, of course, stationary. However, if  $\{V_t\}$  is a sequence of unconditionally heteroskedastic random variables such that  $Q(s)$  and  $\Sigma(s)$  do not satisfy the smoothness restrictions in Assumption 2.1-2.2 then Theorem 3.1-3.2 do not hold. For example, consider  $V_t = \rho_t V_{t-1} + u_t$  where  $u_t \sim i.i.d. \mathcal{N}(0, 1)$  and  $\rho_t \in (-1, 1)$  for all  $t$ . Segmented local stationarity corresponds to  $\rho_t$  being piecewise continuous, local stationarity corresponds to  $\rho_t$  being continuous and stationarity corresponds to  $\rho_t$  being constant. If  $\rho_t$  does not satisfy any of these restrictions, Assumption 2.1-2.2 do not hold. For unconditionally heteroskedastic random variables the asymptotic distributions of  $F_{\text{fixed}-b}$  and  $t_{\text{fixed}-b}$  remain unknown since they cannot be characterized. Thus, for general nonstationary random variables fixed- $b$  inference based on the asymptotic distribution is infeasible. This highlights one major difference from HAR inference based on consistent estimation of  $\Omega$ . HAC and DK-HAC estimators are consistent for  $\Omega$  also for general nonstationary random variables so that HAR test statistics

follow asymptotically the usual standard distributions. For example, a  $t$ -statistic studentized by a HAC or DK-HAC estimator will follow asymptotically a standard normal.

## 4 Error in Rejection Probability in a Gaussian Location Model

We develop high-order asymptotic expansions and obtain the ERP of fixed- $b$  HAR tests. The results in [Velasco and Robinson \(2001\)](#), [Jansson \(2004\)](#) and [Sun, Phillips, and Jin \(2008\)](#) suggest that under stationarity the ERP of  $F_{\text{fixed-}b}$  and of  $t_{\text{fixed-}b}$  are smaller than those of the conventional HAC-based HAR tests. We show that the opposite is true when stationarity does not hold.

Consider the location model  $y_t = \beta_0 + e_t$  ( $t = 1, \dots, T$ ). We have  $V_t = e_t$ . Under the assumption that  $V_t$  is stationary and Gaussian, [Velasco and Robinson \(2001\)](#) developed second-order Edgeworth expansions and showed that

$$\mathbb{P}(t_{\text{HAC}} \leq z) - \Phi(z) = d(z) (Tb_T)^{-1/2} + o\left((Tb_T)^{-1/2}\right), \quad (4.1)$$

for any  $z \in \mathbb{R}$  where

$$t_{\text{HAC}} = \frac{\sqrt{T}(\hat{\beta} - \beta_0)}{\sqrt{\hat{\Omega}_{\text{HAC}}}},$$

$b_T \rightarrow 0$ ,  $\Phi(\cdot)$  is the distribution function of the standard normal and  $d(\cdot)$  is an odd function. The ERP is the leading term of the right-hand side of (4.1). Since  $b_T = O(T^{-\eta})$  with  $0 < \eta < 1$ , the ERP of  $t_{\text{HAC}}$  is  $O(T^{-\gamma})$  with  $\gamma < 1/2$ . It follows that the leading term of  $\mathbb{P}(F_{\text{HAC}} \leq c)$  where  $F_{\text{HAC}} = T(\hat{\beta} - \beta_0)^2 / \hat{\Omega}_{\text{HAC}}$  is of the form  $2d(\sqrt{c})T^{-\gamma} = O(T^{-\gamma})$  for any  $c > 0$ .

[Jansson \(2004\)](#) and [Sun, Phillips, and Jin \(2008\)](#) showed that<sup>5</sup>

$$\mathbb{P}(F_{\text{fixed-}b} \leq c) - \mathbb{P}\left(\frac{W_1(1)^2}{\int_0^1 \int_0^1 -K''(r-s) B_1(r) B_1(s)' dr ds} \leq c\right) = O(T^{-1}). \quad (4.2)$$

Thus,  $F_{\text{fixed-}b}$  has a smaller ERP than  $F_{\text{HAC}}$  [cf.  $O(T^{-1})$  versus  $O(T^{-\gamma})$ ]. This implies that the rate of convergence of  $F_{\text{fixed-}b}$  to its (nonstandard) limiting null distribution is faster than the rate of convergence of  $F_{\text{HAC}}$  to a  $\chi_1^2$ . These results reconciled with finite-sample evidence in the literature showing that the null rejection rates of  $F_{\text{fixed-}b}$  and  $t_{\text{fixed-}b}$  are more accurate than those of  $F_{\text{HAC}}$

---

<sup>5</sup>Actually [Jansson \(2004\)](#) showed that the bound was  $O(T^{-1} \log T)$ . Using a different proof strategy, [Sun, Phillips, and Jin \(2008\)](#) showed that the  $\log T$  term can be dropped.

and  $t_{\text{HAC}}$ , respectively, when that data are stationary.

We now address the question of whether these results extend to nonstationarity. It turns out that the answer is negative. This provides an analytical explanation for the Monte Carlo experiments that have appeared recently in [Casini \(2021b\)](#), [Casini, Deng, and Perron \(2021\)](#) and [Casini and Perron \(2021\)](#) who found serious distortions in the rejection rates of fixed- $b$  HAR tests under the null and alternative hypotheses when the data are nonstationary. These distortions being often much larger than those corresponding to the conventional HAC-based HAR tests.

Theorem 3.2 showed that the fixed- $b$  HAR tests are not pivotal. Thus, to conduct inference based on the fixed- $b$  asymptotic distribution one has to construct consistent estimates of its nuisance parameters. We introduce a general nonparametric estimator of  $\Sigma(u)$ . Let

$$\begin{aligned} \mathbf{K}_2 = \{K_2(\cdot) : \mathbb{R} \rightarrow [0, \infty] : K_2(x) = K_2(1-x), \int K_2(x) dx = 1, \\ K_2(x) = 0, \text{ for } x \notin [0, 1], K_2(\cdot) \text{ is continuous}\}, \end{aligned}$$

and

$$\hat{\Omega}(u) = \hat{\Sigma}^2(u) = \sum_{k=-T+1}^{T-1} K_{h_1}(h_1 k) \hat{c}_{T,h_2}(u, k), \quad (4.3)$$

where  $K_{h_1}(\cdot) \in \mathbf{K}$ ,  $h_1$  is a bandwidth sequence satisfying  $h_1 \rightarrow 0$ ,

$$\hat{c}_{T,h_2}(u, k) = (Th_2)^{-1} \sum_{s=|k|+1}^T K_{h_2} \left( \frac{(\lfloor Tu \rfloor - (s - |k|/2))/T}{h_2} \right) \hat{V}_s \hat{V}_{s-|k|},$$

with  $K_{h_2}(\cdot) \in \mathbf{K}_2$  and  $h_2$  is a bandwidth sequence satisfying  $h_2 \rightarrow 0$ . Since  $x_t = 1$  for all  $t$ , we have  $Q(u) = 1$  for all  $u \in [0, 1]$  in (3.3)-(3.4). Thus, we set  $\hat{Q}(u) = 1$  for all  $u \in [0, 1]$ . For arbitrary  $x_t$ , one can take

$$\hat{Q}(u) = (Th_2)^{-1} \sum_{s=1}^T K_{h_2} \left( \frac{(\lfloor Tu \rfloor - s)/T}{h_2} \right) x_s^2.$$

As in the literature, we focus on the simple location model with Gaussian errors. The Gaussianity assumption can be relaxed by considering distributions with, for example, Gram-Charlier representations at the expenses of more complex derivations [see, e.g., [Phillips \(1980\)](#)]. The following assumption on  $V_t$  facilitates the development of the higher order expansions and is weaker than the one used by [Sun, Phillips, and Jin \(2008\)](#) since they also imposed second-order stationarity.

**Assumption 4.1.**  $\{V_t\}$  is a mean-zero Gaussian process with  $\sup_{1 \leq t \leq T} \sum_{k=-\infty}^{\infty} k^2 |\mathbb{E}(V_t V_{t-k})| < \infty$ .

In order to develop the asymptotic expansions we use the following conditions on the kernel which were also used by [Andrews \(1991\)](#) and [Sun, Phillips, and Jin \(2008\)](#).

**Assumption 4.2.** (i)  $K(x) : \mathbb{R} \rightarrow [0, 1]$  is symmetric and satisfies  $K(0) = 1$ ,  $\int_0^\infty x K(x) dx < \infty$  and  $|K(x) - K(y)| \leq C_1 |x - y|$  for all  $x, y \in \mathbb{R}$  and some  $C_1 < \infty$ .

(ii)  $q_0 \geq 1$  where  $q_0$  is the Parzen characteristic exponent defined by

$$q_0 = \max \left\{ \tilde{q} : \tilde{q} \in \mathbb{Z}_+, \overline{K}_{\tilde{q}} = \lim_{x \rightarrow 0} \frac{1 - K(x)}{|x|^{\tilde{q}}} < \infty \right\}.$$

(iii)  $K(x)$  is positive semidefinite, i.e., for any square integrable function  $g(x)$ ,  $\int_0^\infty \int_0^\infty K(s - t) g(s) g(t) ds dt \geq 0$ .

All of the commonly used kernels with the exception of the truncated kernel satisfy Assumption 4.2-(i, ii). [Sun, Phillips, and Jin \(2008\)](#) required piecewise smoothness on  $K(\cdot)$  instead of the Lipschitz condition. Part (iii) ensures that the associated LRV estimator is positive semidefinite. The commonly used kernels that satisfy part (i, iii) are the Bartlett, Parzen and quadratic spectral (QS) kernels. For the Bartlett kernel,  $q_0 = 1$ , while for the Parzen and QS kernels,  $q_0 = 2$ .

As in [Sun, Phillips, and Jin \(2008\)](#), we present the asymptotic expansion for the test statistic studentized by the more general estimator  $\hat{\Omega}_{\text{fixed}-b}$  that keeps the bandwidth  $b \in (0, 1]$  at a fixed fraction of the sample size,

$$\hat{\Omega}_{\text{fixed}-b} = \sum_{k=-T+1}^{T-1} K\left(\frac{k}{Tb}\right) \hat{\Gamma}(k).$$

In the previous section,  $\hat{\Omega}_{\text{fixed}-b}$  was defined with  $b = 1$ . Let  $K_b = K(\cdot/b)$ . Lemma [S.A.1](#) in the supplement extends Theorem [3.1](#) to  $\hat{\Omega}_{\text{fixed}-b}$  using  $b \in (0, 1]$  and the kernels that satisfy Assumption [4.2](#). Under Assumption [2.1](#) and [4.1-4.2](#), Lemma [S.A.1](#) shows that  $\hat{\Omega}_{\text{fixed}-b} \Rightarrow \mathcal{G}_b$  where

$$\mathcal{G}_b \triangleq \int_0^1 \int_0^1 K_b(r - s) d\tilde{B}_1(r) d\tilde{B}_1(s),$$

and

$$\tilde{B}_1(r) = \int_0^r \left( \Sigma(u) dW_1(u) - r \left( \int_0^1 \Sigma(u) dW_1(u) \right) \right).$$

Let  $C_\Omega = \sup_{s \in [0,1]} \Omega(s)$ ,  $C_{2,\Omega} = \max\{C_\Omega, 1\}$ ,

$$t_{\text{fixed}-b} \triangleq \frac{\sqrt{T}(\hat{\beta} - \beta_0)}{\sqrt{\hat{\Omega}_{\text{fixed}-b}}},$$

and

$$\begin{aligned} \hat{\mathcal{G}}_b &= \int_0^1 \int_0^1 K_b(r-s) \left( \int_0^r \hat{\Sigma}(u) dW_1(u) - r \int_0^1 \hat{\Sigma}(u) dW_1(u) \right) \\ &\quad \times \left( \int_0^s \hat{\Sigma}(u) dW_1(u) - s \int_0^1 \hat{\Sigma}(u) dW_1(u) \right) dr ds. \end{aligned}$$

**Theorem 4.1.** *Let Assumption 2.1, 4.1-4.2,  $h_1 \rightarrow 0$ ,  $h_2 \rightarrow 0$  and  $Th_1h_2 \rightarrow \infty$  hold. Provided that  $b$  is fixed such that  $b < 1/(16C_{2,\Omega} \int_{-\infty}^{\infty} |K(x)| dx)$ , we have*

$$\sup_{z \in \mathbb{R}_+} \left| \mathbb{P}(|t_{\text{fixed}-b}| \leq z) - \mathbb{P}\left(\left| \frac{\int_0^1 \hat{\Sigma}(u) dW_1(u)}{\sqrt{\hat{\mathcal{G}}_b}} \right| \leq z \right) \right| = O((Th_1h_2)^{-1/2}).$$

Comments: 1. Theorem 4.1 shows that the ERP associated to  $t_{\text{fixed}-b}$  is  $O((Th_1h_2)^{-1/2})$ . This is an order of magnitude larger than the ERP associated to  $t_{\text{fixed}-b}$  under stationarity,  $O(T^{-1})$ , where the latter was established by Jansson (2004) and Sun, Phillips, and Jin (2008). The increase in the ERP of  $t_{\text{fixed}-b}$  is the price one has to pay for not having a pivotal distribution under nonstationarity. This is intuitive. Without a pivotal distribution, one has to obtain estimates of the nuisance parameters. However, the nuisance parameters can be consistently estimated only under small- $b$  asymptotics. The latter estimates enjoy a nonparametric rate of convergence which then results in a larger ERP since it is the discrepancy between these estimates and their probability limits that is reflected in the leading term of the asymptotic expansion.

2. Existing fixed- $b$  methods use the critical value from the pivotal fixed- $b$  limiting distribution obtained under the assumption of stationarity. Our results suggest that the ERP associated to such fixed- $b$  HAR tests is  $O(1)$ . This follows because that critical value is not theoretically valid, i.e., it is from the pivotal fixed- $b$  limiting distribution which, however, is different from the non-pivotal fixed- $b$  limiting distribution under nonstationarity. Thus, as  $T \rightarrow \infty$  the ERP does not converge to zero, implying large distortions in the null rejection rates even for unbounded sample sizes.

3. It is useful to compare Theorem 4.1 with the results for the ERP associated to  $t_{\text{HAC}}$ . Under stationarity Velasco and Robinson (2001) showed that the ERP associated to  $t_{\text{HAC}}$  is  $O((Tb_T)^{-1/2})$

where  $b_T \rightarrow 0$ . [Casini, Deng, and Perron \(2021\)](#) showed that under nonstationarity the ERP associated to  $t_{\text{HAC}}$  has the same order as under stationarity, i.e.,  $O((Tb_T)^{-1/2})$ . Thus, it is sufficient to compare  $O((Th_1h_2)^{-1/2})$  and  $O((Tb_T)^{-1/2})$ . Since  $h_1$  and  $b_T$  are the bandwidths used for smoothing over lagged autocovariances, they have a similar order. It follows that the ERP associated to  $t_{\text{fixed}-b}$  is larger than that associated to  $t_{\text{HAC}}$ . In addition, the ERP associated to  $t_{\text{fixed}-b}$  based on existing fixed- $b$  methods that rely on stationarity is much larger than the ERP associated to  $t_{\text{HAC}}$  since the former is  $O(1)$ .

4. Theorem [4.1](#) implies that the theoretical properties of fixed- $b$  inference changes substantially depending on whether the data are stationarity or not. In particular, it suggests that the approximations based on fixed- $b$  asymptotics obtained under stationarity in the literature are not valid and do not provide a good approximation when stationarity does not hold. This contrasts to HAR inference tests based on consistent long-run variance estimators which are valid also under nonstationarity and have the same asymptotic distribution regardless of whether the data are stationary or not.

5. Overall, the theoretical results contrast with the mainstream conclusion in the literature that fixed- $b$  HAR inference is theoretically superior to HAR inference based on consistent long-run variance estimators. Similar comments apply to more recent inference methods [see, e.g., [Lazarus, Lewis, and Stock \(2020\)](#) and [Lazarus, Lewis, Stock, and Watson \(2018\)](#)] since they are based on fixed- $b$  asymptotics as well as stationarity.

6. Our theoretical results complement the recent finite-sample evidence in [Belotti et al. \(2021\)](#), [Casini \(2021b\)](#), [Casini and Perron \(2021\)](#) and [Casini, Deng, and Perron \(2021\)](#). Their simulation results showed that existing fixed- $b$  HAR inference tests perform poorly in terms of the accuracy of the null rejection rates and of power when stationarity does not hold. They considered  $t$ -tests in the linear regression models and HAR tests outside the linear regression model, and a variety of data-generating processes. They provided evidence that fixed- $b$  HAR tests can be severely undersized and can exhibit non-monotonic power. Some of these issues are generated by the low frequency contamination induced by nonstationarity which biases upward each sample autocovariance  $\hat{\Gamma}(k)$ . Since  $\hat{\Omega}_{\text{fixed}-b}$  uses many lagged autocovariances as  $b$  is fixed, it is inflated which then results in size distortions and lower power. To avoid redundancy, in this paper we do not repeat the above-mentioned simulation analysis but refer to those works for more details.

7. The fixed- $b$  limiting distribution under nonstationarity is complex to use in practice as it depends in a complicated way on nuisance parameters. It is beyond the scope of the paper to investigate how best to use the non-pivotal fixed- $b$  limiting distribution. If one wants to use fixed- $b$



inference without relying on the limiting distribution and consider other procedures such as e.g., bootstrap-based autocorrelation robust tests, modification of the test statistic<sup>6</sup>, etc., then one has to face the challenge that these methods are not as simple as HAC-based inference and therefore would find little traction in empirical work as argued by [Lazarus, Lewis, and Stock \(2020\)](#).

8. The requirement  $Th_1h_2 \rightarrow \infty$  is a standard condition for consistency of nonparametric estimators such as  $\hat{\Omega}(u)$ . The requirement  $b < 1/(16C_{2,\Omega} \int_{-\infty}^{\infty} |K(x)|dx)$  is similar to the one used by [Sun, Phillips, and Jin \(2008\)](#). It can be relaxed at the expenses of more complex derivations.

9. As remarked in the last paragraph of Section 3, for unconditionally heteroskedastic random variables, fixed- $b$  HAR inference is infeasible. Thus, the associated ERP does not convergence to zero. In contrast, HAR inference based on consistent long-run variance estimator is valid and the associated ERP is again  $O((Tb_T)^{-1/2})$  with  $b_T \rightarrow 0$ .

## 5 Conclusions

This paper has shown that the theoretical properties of fixed- $b$  HAR inference change depending on whether the data are stationary or not. Under nonstationarity we established that fixed- $b$  HAR test statistics have a limiting distribution that is not pivotal and that their ERP is an order of magnitude larger than that under stationarity and is larger than that of HAR tests based on traditional HAC estimators. These theoretical results reconcile with recent finite-sample evidence showing that fixed- $b$  HAR test statistics can perform poorly when the data are nonstationary, both in terms of distortions in the null rejection rates and of non-monotonic power. Our theoretical results contrast with the mainstream conclusion in the literature that fixed- $b$  HAR inference is theoretically superior to HAR inference based on consistent long-run variance estimators.

## Supplemental Materials

The supplement for online publication [cf. [Casini \(2021a\)](#)] contains the proofs of the results.

---

<sup>6</sup>[Hwang and Sun \(2017\)](#) proposed a modification to the trinity of test statistics in the two-step GMM setting and showed that the modified test statistics are asymptotically  $F$  distributed under fixed- $b$  asymptotics.

## References

- ALDOUS, D. (1978): “Stopping Times and Tightness,” *Annals of Probability*, 6(2), 335–340.
- ANDREWS, D. W. K. (1991): “Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimation,” *Econometrica*, 59(3), 817–858.
- ANDREWS, D. W. K., AND J. C. MONAHAN (1992): “An Improved Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimator,” *Econometrica*, 60(4), 953–966.
- BELOTTI, F., A. CASINI, L. CATANIA, S. GRASSI, AND P. PERRON (2021): “Simultaneous Bandwidths Determination for Double-Kernel HAC Estimators and Long-Run Variance Estimation in Nonparametric Settings,” *arXiv preprint arXiv:2103.00060*.
- CASINI, A. (2021a): “Supplement to ”The Fixed-b Limiting Distribution and the ERP of HAR Tests Under Nonstationarity”,” *Unpublished Manuscript, Department of Economics and Finance, University of Rome Tor Vergata*.
- (2021b): “Theory of Evolutionary Spectra for Heteroskedasticity and Autocorrelation Robust Inference in Possibly Misspecified and Nonstationary Models,” *arXiv preprint arXiv:2103.02981*.
- CASINI, A., T. DENG, AND P. PERRON (2021): “Theory of Low Frequency Contamination from Nonstationarity and Misspecification: Consequences for HAR Inference,” *arXiv preprint arXiv:2103.01604*.
- CASINI, A., AND P. PERRON (2021): “Minimax MSE Bounds and Nonlinear VAR Prewhitening for Long-Run Variance Estimation Under Nonstationarity,” *arXiv preprint arXiv:2103.02235*.
- DE JONG, R., AND J. DAVIDSON (2000): “Consistency of Kernel Estimators of Heteroskedastic and Autocorrelated Covariance Matrices,” *Econometrica*, 68(2), 407–423.
- HANSEN, B. (1992): “Consistent Covariance Matrix Estimation for Dependent Heterogeneous Processes,” *Econometrica*, 60(4), 967–972.
- HWANG, J., AND Y. SUN (2017): “Asymptotic F and t Tests in an Efficient GMM Setting,” *Journal of Econometrics*, 198(2), 277–295.
- JANSSON, M. (2004): “The Error in Rejection Probability of Simple Autocorrelation Robust Tests,” *Econometrica*, 72(3), 937–946.
- KIEFER, N. M., AND T. J. VOGELSANG (2002a): “Heteroskedasticity-Autocorrelation Robust Standard Errors using the Bartlett Kernel without Truncation,” *Econometrica*, 70(5), 2093–2095.
- (2002b): “Heteroskedasticity-Autocorrelation Robust Testing Using Bandwidth Equal to Sample Size,” *Econometric Theory*, 18(6), 1350–1366.

- KIEFER, N. M., T. J. VOGELSANG, AND H. BUNZEL (2000): “Simple Robust Testing of Regression Hypotheses,” *Econometrica*, 69(3), 695–714.
- LAZARUS, E., D. J. LEWIS, AND J. H. STOCK (2020): “The Size-Power Tradeoff in HAR Inference,” *Econometrica*, 89(5), 2497–2516.
- LAZARUS, E., D. J. LEWIS, J. H. STOCK, AND M. W. WATSON (2018): “HAR Inference: Recommendations for Practice,” *Journal of Business and Economic Statistics*, 36(4), 541–559.
- MERLEVÈDE, F., M. PELIGRAD, AND S. UTEV (2019): “Functional CLT for Martingale-like Nonstationary Dependent Structures,” *Bernoulli*, 25(4B), 3203–3233.
- NEWKEY, W. K., AND K. D. WEST (1987): “A Simple Positive Semidefinite, Heteroskedastic and Autocorrelation Consistent Covariance Matrix,” *Econometrica*, 55(3), 703–708.
- (1994): “Automatic Lag Selection in Covariance Matrix Estimation,” *Review of Economic Studies*, 61(4), 631–653.
- PHILLIPS, P. C. B. (1980): “Finite Sample Theory and the Distributions of Alternative Estimators of the Marginal Propensity to Consume,” *Review of Economic Studies*, 47(1), 183–224.
- SUN, Y., P. C. B. PHILLIPS, AND S. JIN (2008): “Optimal Bandwidth Selection in Heteroskedasticity-Autocorrelation Robust Testing,” *Econometrica*, 76(1), 175–194.
- VELASCO, C., AND P. M. ROBINSON (2001): “Edgeworth Expansions for Spectral Density Estimates and Studentized Sample Mean,” *Econometric Theory*, 17(3), 497–539.

# Supplement to “The Fixed- $b$ Limiting Distribution and the ERP of HAR Tests Under Nonstationarity”

ALESSANDRO CASINI

Department of Economics and Finance  
University of Rome Tor Vergata

30th November 2021

## Abstract

This supplemental material is for online publication. It contains the proofs of the results.

## S.A Mathematical Proofs

### S.A.1 Proof of Theorem 3.1

Define

$$Q_T(r) = T^{-1} \sum_{t=1}^{\lfloor Tr \rfloor} x_t x'_t, \quad X_T(r) = T^{-1/2} S_{\lfloor Tr \rfloor}.$$

We begin with part (i). Let

$$D_T(r) = T^2 \left[ \left( K\left(\frac{\lfloor Tr \rfloor + 1}{T}\right) - K\left(\frac{\lfloor Tr \rfloor}{T}\right) \right) - \left( K\left(\frac{\lfloor Tr \rfloor}{T}\right) - K\left(\frac{\lfloor Tr \rfloor - 1}{T}\right) \right) \right].$$

By symmetry of  $K(\cdot)$ , it follows the symmetry of  $D_T(\cdot)$ . Since  $K''(r)$  is assumed to exist, we have  $\lim_{T \rightarrow \infty} D_T(r) = K''(r)$ . The convergence is uniform in  $r$  since  $K''(r)$  is continuous. From Assumption 2.1-2.2 it follows that  $(Q_T(r), X_T(r)', D_T(r)) \Rightarrow (\int_0^r Q(u) du, (\int_0^r \Sigma(u) dW_p(u))', K''(r))$  jointly.

Define  $K_{i,j} = ((i-j)/T)$ . We have

$$\hat{\Omega}_{\text{fixed}-b} = T^{-1} \sum_{i=1}^T \sum_{j=1}^T K_{i,j} \hat{V}_i \hat{V}_j' = T^{-1} \sum_{i=1}^T \hat{V}_i \left( \sum_{j=1}^T K_{i,j} \hat{V}_j' \right).$$

Note that

$$\begin{aligned} & T^2 ((K_{i,j} - K_{i,j+1}) - (K_{i+1,j} - K_{i+1,j+1})) \\ &= -T^2 \left[ \left( K\left(\frac{i-j+1}{T}\right) - K\left(\frac{i-j}{T}\right) \right) - \left( K\left(\frac{i-j}{T}\right) - K\left(\frac{i-j-1}{T}\right) \right) \right] \\ &= D_T((i-j)/T) \end{aligned} \tag{S.1}$$

Define  $\hat{S}_t = \sum_{j=1}^t \hat{V}_j$ . Note that  $\hat{S}_T = \mathbf{0}$  by the normal equations for OLS. We have

$$\begin{aligned} T^{-1/2} \hat{S}_{\lfloor Tr \rfloor} &= T^{-1/2} S_{\lfloor Tr \rfloor} - T^{-1} \sum_{t=1}^{\lfloor Tr \rfloor} x_t x'_t \left( T^{-1} \sum_{t=1}^T x_t x'_t \right)^{-1} T^{-1/2} S_T \\ &= X_T(r) - Q_T(r) Q_T(1)^{-1} X_T(1). \end{aligned} \tag{S.2}$$

Using the identity

$$\sum_{l=1}^T a_l b_l = \sum_{l=1}^{T-1} \left( (a_l - a_{l+1}) \sum_{j=1}^l b_j \right) + a_T \sum_{j=1}^T b_j, \tag{S.3}$$

first applied to  $\sum_{j=1}^T K_{i,j} \hat{V}_j'$  and then again to the sum over  $i$ , [Kiefer and Vogelsang \(2002b\)](#) showed that

$$\hat{\Omega}_{\text{fixed}-b} = T^{-1} \sum_{i=1}^{T-1} T^{-1} \sum_{j=1}^{T-1} T^2 ((K_{i,j} - K_{i,j+1}) - (K_{i+1,j} - K_{i+1,j+1})) T^{-1/2} \hat{S}_i T^{-1/2} \hat{S}_j'. \tag{S.4}$$

Using (S.1)-(S.2) in (S.4) we have

$$\begin{aligned}\widehat{\Omega}_{\text{fixed}-b} &= \int_0^1 \int_0^1 -D_T(r-s) \left[ X_T(r) - Q_T(r) Q_T(1)^{-1} X_T(1) \right] \left[ X_T(s) - Q_T(s) Q_T(1)^{-1} X_T(1) \right]' dr ds \\ &\Rightarrow - \int_0^1 \int_0^1 K''(r-s) \left( \int_0^r \Sigma(u) dW_p(u) - \left( \int_0^r Q(u) du \right) \overline{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right) \\ &\quad \times \left( \int_0^s \Sigma(u) dW_p(u) - \left( \int_0^s Q(u) du \right) \overline{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right)' dr ds,\end{aligned}$$

where we have used Assumption 2.1-2.2 and the continuous mapping theorem since  $\widehat{\Omega}_{\text{fixed}-b}$  is a continuous functions of  $(Q_T(r), X_T(r)', D_T(r))$ .

We now move to part (ii). Suppose that the Bartlett kernel  $K_{\text{BT}}$  is used. We have

$$\widehat{\Omega}_{\text{fixed}-b} = T^{-1} \sum_{i=1}^T \sum_{j=1}^T \widehat{V}_i \left( 1 - \frac{|i-j|}{T} \right) \widehat{V}_j'.$$

Using the identity (S.3) and  $\widehat{S}_T = \mathbf{0}$ , Kiefer and Vogelsang (2002a) showed that

$$\widehat{\Omega}_{\text{fixed}-b} = 2T^{-2} \sum_{i=1}^T \widehat{S}_i \widehat{S}_i'.$$

Using (S.2) and Assumption 2.1-2.2 we yield,

$$\begin{aligned}\widehat{\Omega}_{\text{fixed}-b} &\Rightarrow 2 \int_0^1 \left( \int_0^r \Sigma(u) dW_p(u) - \left( \int_0^r Q(u) du \right) \overline{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right) \\ &\quad \times \left( \int_0^r \Sigma(u) dW_p(u) - \left( \int_0^r Q(u) du \right) \overline{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right)' dr,\end{aligned}$$

which concludes the proof.  $\square$

## S.A.2 Proof of Theorem 3.2

We begin with part (i). Using Theorem 3.1 we have

$$\begin{aligned}F_{\text{fixed}-b} &= \left( RQ_T(1)^{-1} X_T(1) \right)' \left( RQ_T(1)^{-1} \widehat{\Omega}_{\text{fixed}-b} Q_T(1)^{-1} R' \right)^{-1} RQ_T(1)^{-1} X_T(1) / q \\ &\Rightarrow \left( R\overline{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) \right)' \left( R\overline{Q}^{-1} \mathcal{G} \overline{Q}^{-1} R' \right)^{-1} R\overline{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u) / q\end{aligned}$$

where  $\mathcal{G}$  is defined in (3.3). If  $q = 1$ , then

$$t_{\text{fixed}-b} = \frac{RQ_T(1)^{-1} X_T(1)}{\sqrt{RQ_T(1)^{-1} \widehat{\Omega}_{\text{fixed}-b} Q_T(1)^{-1} R'}}$$

$$\Rightarrow \frac{R\bar{Q}^{-1} \int_0^1 \Sigma(u) dW_p(u)}{\sqrt{R\bar{Q}^{-1} \mathcal{G} \bar{Q}^{-1} R'}}.$$

Part (ii) can be proved in a similar manner.  $\square$

### S.A.3 Proof of Theorem 4.1

Throughout the proof, let  $\hat{\Omega}_b = \hat{\Omega}_{\text{fixed}-b}$  and

$$Z_{T,0}(z) \triangleq \mathbb{P} \left( \left| \frac{\sqrt{T}(\hat{\beta} - \beta_0)}{\sqrt{\hat{\Omega}_b}} \right| \leq z \right), \quad Z_0(z) \triangleq \mathbb{P} \left( \left| \frac{\int_0^1 \Sigma(u) dW_1(u)}{\sqrt{\mathcal{G}}} \right| \leq z \right),$$

$$\hat{Z}_0(z) \triangleq \mathbb{P} \left( \left| \frac{\int_0^1 \hat{\Sigma}(u) dW_1(u)}{\sqrt{\hat{\mathcal{G}}_b}} \right| \leq z \right).$$

We have

$$\sup_{z \in \mathbb{R}_+} |Z_{T,0}(z) - \hat{Z}_0(z)| \leq \sup_{z \in \mathbb{R}_+} |Z_{T,0}(z) - Z_0(z)| + \sup_{z \in \mathbb{R}_+} |Z_0(z) - \hat{Z}_0(z)|$$

$$\triangleq D_1 + D_2.$$

We show that  $D_1 = O(T^{-1})$  and  $D_2 = O((Th_1h_2)^{-1/2})$ . We begin with some preliminary lemmas. The first lemma below generalizes Theorem 3.1 to allow for general kernels satisfying Assumption 4.2 and for a  $p$ -vector  $\hat{V}_t$ . Let

$$\tilde{B}_p(r) = \int_0^r \Sigma(u) dW_p(u) - r \left( \int_0^1 \Sigma(u) dW_p(u) \right),$$

and

$$K_b^*(r, s) \triangleq K_b(r - s) - \int_0^1 K_b(r - t) dt - \int_0^1 K_b(\tau - s) d\tau + \int_0^1 \int_0^1 K_b(t - \tau) dt d\tau.$$

**Lemma S.A.1.** *Let Assumption 2.1 and 4.2 hold. We have:*

- (i)  $\hat{\Omega}_b \Rightarrow \mathcal{G}_b$  where  $\mathcal{G}_b = \int_0^1 \int_0^1 K_b(r - s) d\tilde{B}_p(r) d\tilde{B}_p(s)'$ ;
- (ii)  $\mu_b = \mathbb{E}(\mathcal{G}_b) = \int_0^1 K_b^*(s, s) \Omega(s) ds$ .

*Proof of Lemma S.A.1.* We begin with part (i). Since  $K(\cdot)$  is positive semidefinite, Mercer's Theorem implies that

$$K(r - s) = \sum_{n=1}^{\infty} \frac{1}{\lambda_n} g_n(r) g_n(s), \quad (\text{S.5})$$

where  $\lambda_n^{-1} > 0$  are the eigenvalues of  $K(\cdot)$  and  $g_n(\cdot)$  are the corresponding eigenfunctions, i.e.,  $g_n(s) = \lambda_n \int_0^1 K(r - s) g_n(r) dr$ . The convergence of the right-hand side over  $(r, s) \in [0, 1] \times [0, 1]$  is uniform.

Using Assumption 2.1 and (S.5) we have

$$\begin{aligned}
\hat{\Omega}_b &= \frac{1}{T} \sum_{t=1}^T \sum_{s=1}^T K_b \left( \frac{t-s}{T} \right) \hat{V}_t \hat{V}_s' \\
&= \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \frac{1}{T} \sum_{t=1}^T \sum_{s=1}^T g_n(t/(bT)) g_n(s/(bT)) \hat{V}_t \hat{V}_s' \\
&= \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \left( \frac{1}{\sqrt{T}} \sum_{t=1}^T \hat{V}_t g_n(t/(bT)) \right) \left( \frac{1}{\sqrt{T}} \sum_{s=1}^T g_n(s/(bT)) \hat{V}_s' \right) \\
&\Rightarrow \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \left( \int_0^1 g_n(r/b) d\tilde{B}_p(r) \right) \left( \int_0^1 g_n(s/b) d\tilde{B}_p(s) \right)' \\
&= \int_0^1 \int_0^1 \sum_{n=1}^{\infty} \frac{1}{\lambda_n} g_n(r/b) g_n(s/b) d\tilde{B}_p(r) d\tilde{B}_p(s)' \\
&= \int_0^1 \int_0^1 K_b(r-s) d\tilde{B}_p(r) d\tilde{B}_p(s)' \\
&= \mathcal{G}_b
\end{aligned}$$

For part (ii), after some algebraic manipulations we can write

$$\mathcal{G}_b = \int_0^1 \int_0^1 K_b^*(r, s) \Sigma(r) dW_p(r) (\Sigma(s) dW_p(s))'.$$

It follows that

$$\begin{aligned}
\mathbb{E}(\mathcal{G}_b) &= \mathbb{E} \left( \int_0^1 \int_0^1 K_b^*(r, s) \Sigma(r) dW_p(r) (\Sigma(s) dW_p(s))' \right) \\
&= \int_0^1 K_b^*(s, s) \Sigma(s) \Sigma(s)' ds \\
&= \int_0^1 K_b^*(s, s) \Omega(s) ds,
\end{aligned}$$

which concludes the proof.  $\square$

Let  $p = 1$ . Sun, Phillips, and Jin (2008) showed that  $K_b^*(r, s)$  is positive semidefinite. Thus, using Mercer's theorem, we have

$$K_b^*(r, s) = \sum_{n=1}^{\infty} \lambda_n^* g_n^*(r) g_n^*(s), \quad (\text{S.6})$$

where  $\lambda_n^* > 0$  are the eigenvalues of  $K_b^*(\cdot, \cdot)$  and  $g_n^*(r)$  are the corresponding eigenfunctions, i.e.,  $\lambda_n^* g_n^*(s) = \int_0^{\infty} K_b^*(r, s) g_n^*(r) dr$ . Since  $\Sigma(s) > 0$ , we can write

$$K_b^*(r, s) \Sigma(r) \Sigma(s) = \sum_{n=1}^{\infty} \lambda_n^* g_n^{**}(r) g_n^{**}(s),$$



where  $g_n^{**}(r) = \Sigma(r) g_n^*(r)$ . Then,  $\lambda_n^* g_n^{**}(s) = \int_0^\infty K_b^*(r, s) \Sigma(r) \Sigma(s) g_n^{**}(r) dr$ . It follows that  $\mathcal{G}_b = \sum_{n=1}^\infty \lambda_n^* Z_n^2$  where  $Z_n \sim i.i.d. \mathcal{N}(0, 1)$ . Thus, the characteristic function of  $\Omega^{-1}(\mathcal{G}_b - \mu_b)$  is given by

$$\psi(t) = \mathbb{E} \left( e^{it\Omega^{-1}(\mathcal{G}_b - \mu_b)} \right) = \prod_{n=1}^\infty \left( 1 - 2i\lambda_n^* \Omega^{-1} t \right)^{-1/2} \left( e^{-it\Omega^{-1}\mu_b} \right), \quad (\text{S.7})$$

and the cumulant generating function is

$$\log \psi(t) = \sum_{m=2}^\infty \left( 2^{m-1} (m-1)! \sum_{n=1}^\infty \left( \lambda_n^* \Omega^{-1} \right)^m \right) \frac{(it)^m}{m!}. \quad (\text{S.8})$$

Let  $\kappa_j$  ( $j = 1, \dots$ ) be the  $j$ th cumulant of  $\Omega^{-1}(\mathcal{G}_b - \mu_b)$ . Then

$$\kappa_1 = 0 \quad \text{and} \quad \kappa_m = 2^{m-1} (m-1)! \sum_{n=1}^\infty \left( \lambda_n^* \Omega^{-1} \right)^m \quad \text{for } m \geq 2. \quad (\text{S.9})$$

Let  $\tau_{m+1} = \tau_1$ . For  $m \geq 2$ , some algebraic manipulations show that

$$\begin{aligned} \kappa_m &= 2^{m-1} (m-1)! \Omega^{-m} \sum_{n=1}^\infty \left( (g_n^{**}(s))^{-1} \int_0^\infty K_b^*(r, s) \Sigma(r) \Sigma(s) g_n^{**}(r) dr \right)^m \\ &= 2^{m-1} (m-1)! \Omega^{-m} \int_0^1 \cdots \int_0^1 \left( \prod_{j=1}^m \Omega(\tau_j) K_b^*(\tau_j, \tau_{j+1}) \right) d\tau_1 \cdots d\tau_m. \end{aligned} \quad (\text{S.10})$$

Let  $\Xi_m = \Omega^{-m} \mathbb{E}((\mathcal{G}_b - \mu_b)^m)$  for  $m \geq 1$ .

**Lemma S.A.2.** *Let  $\overline{C}_1 = 4 \int_{-\infty}^\infty |K(v)| dv$ ,  $C_\Omega = \sup_{s \in [0, 1]} \Omega(s)$ ,  $D_m > 0$  be a constant depending on  $m$  and Assumption 4.2 hold. Then, for  $m \geq 1$ , we have*

$$|\kappa_m| \leq 2^m (m-1)! \Omega^{-m} C_\Omega^m \left( \overline{C}_1 b \right)^{m-1}, \quad (\text{S.11})$$

and

$$|\Xi_m| \leq D_m 2^{2m} m! \Omega^{-m} C_\Omega^m \left( \overline{C}_1 b \right)^{m-1}. \quad (\text{S.12})$$

*Proof of Lemma S.A.2.* From eq. (A.3) in Sun, Phillips, and Jin (2008),<sup>7</sup>

$$\left| \int_0^1 \cdots \int_0^1 \left( \prod_{j=1}^m K_b^*(\tau_j, \tau_{j+1}) \right) d\tau_1 \cdots d\tau_m \right| \leq 2 \left( \sup_s \int_0^1 |K_b^*(r, s)| dr \right)^{m-1}. \quad (\text{S.13})$$

We have

$$\sup_s \int_0^1 |K_b^*(r, s)| dr \leq b \overline{C}_1, \quad (\text{S.14})$$

---

<sup>7</sup>Any reference to equations in Sun, Phillips, and Jin (2008) corresponds to the long version of the working paper available in Sun's webpage.

from which it follows that

$$\left| \int_0^1 \cdots \int_0^1 \left( \prod_{j=1}^m K_b^*(\tau_j, \tau_{j+1}) \right) d\tau_1 \cdots d\tau_m \right| \leq 2 (b\overline{C}_1)^{m-1}. \quad (\text{S.15})$$

Using (S.15) and some algebraic manipulations we have for  $m \geq 2$ ,

$$\begin{aligned} \kappa_m &= 2^{m-1} (m-1)! \Omega^{-m} \sum_{n=1}^{\infty} (\lambda_n^*)^m \\ &= 2^{m-1} (m-1)! \Omega^{-m} \int_0^1 \cdots \int_0^1 \left( \prod_{j=1}^m \Omega(\tau_j) K_b^*(\tau_j, \tau_{j+1}) \right) d\tau_1 \cdots d\tau_m \\ &\leq 2^{m-1} (m-1)! \Omega^{-m} C_{\Omega}^m \int_0^1 \cdots \int_0^1 \left| \prod_{j=1}^m K_b^*(\tau_j, \tau_{j+1}) \right| d\tau_1 \cdots d\tau_m \\ &\leq 2^m (m-1)! \Omega^{-m} C_{\Omega}^m (b\overline{C}_1)^{m-1}, \end{aligned} \quad (\text{S.16})$$

where  $C_{\Omega} = \sup_{s \in [0,1]} \Omega(s)$ . The moments  $\{\Xi_j\}$  and cumulants  $\{\kappa_j\}$  are related by the following

$$\Xi_m = \sum_{\pi_p} \frac{m!}{(j_1!)^{m_1} (j_2!)^{m_2} \cdots (j_l!)^{m_l}} \frac{1}{m_1! m_2! \cdots m_l!} \prod_{j=\pi_p} \kappa_j, \quad (\text{S.17})$$

where the sum is taken over all partitions  $\pi_p \in \Pi$  such that

$$\pi_p = \left[ \underbrace{j_1, \dots, j_1}_{m_1 \text{ times}}, \underbrace{j_2, \dots, j_2}_{m_2 \text{ times}}, \dots, \underbrace{j_l, \dots, j_l}_{m_l \text{ times}} \right], \quad (\text{S.18})$$

for some integer  $l$ , sequence  $\{j_i\}_{i=1}^l$  such that  $j_1 > j_2 > \cdots > j_l$  and  $m = \sum_{i=1}^l m_i j_i$ .

Using (S.16)-(S.18) yield

$$\begin{aligned} |\Xi_m| &< 2^m m! \Omega^{-m} C_{\Omega}^m (\overline{C}_1 b)^{m-1} \sum_{\pi} \frac{(j_1)^{-m_1} (j_2)^{-m_2} \cdots (j_l)^{-m_l}}{m_1! m_2! \cdots m_l!} j_1^m \\ &\leq D_m 2^{2m} m! \Omega^{-m} C_{\Omega}^m (\overline{C}_1 b)^{m-1}, \end{aligned} \quad (\text{S.19})$$

where the last inequality follows from

$$\sum_{\pi_p} \frac{(j_1)^{-m_1} (j_2)^{-m_2} \cdots (j_l)^{-m_l}}{m_1! m_2! \cdots m_l!} \leq \sum_{\pi_p} \frac{1}{m_1! m_2! \cdots m_l!} < 2^m, \quad (\text{S.20})$$

and  $D_m = \sup_{\pi_p \in \Pi} (j_1 \in \pi_p)$ .  $\square$

We now develop an asymptotic expansion of  $Z_{T,0} = \mathbb{P}(\sqrt{T}(\hat{\beta} - \beta_0)/\sqrt{\hat{\Omega}_b} \leq z)$  for  $\beta = \beta_0 + d/\sqrt{T}$ . When  $d = 0$  (resp.,  $d \neq 0$ ) the expansion can be used to approximate the size (resp., power) of the

$t$ -statistic. Since  $V_t$  is autocorrelated,  $\hat{\beta}$  and  $\hat{\Omega}_b$  are statistically dependent. Thus, we decompose  $\hat{\beta}$  and  $\hat{\Omega}_b$  into statistically independent components. Let  $V = (V_1, \dots, V_T)'$ ,  $y = (y_1, \dots, y_T)'$ ,  $l_T = (1, \dots, 1)'$  and  $\Upsilon_T = \text{Var}(V)$ . The GLS estimator of  $\beta$  is  $\tilde{\beta} = (l_T' \Upsilon_T^{-1} l_T)^{-1} l_T' \Upsilon_T^{-1} y$ . Then,

$$\hat{\beta} - \beta = \tilde{\beta} - \beta + (l_T' l_T)^{-1} l_T' \tilde{V}, \quad (\text{S.21})$$

where  $\tilde{V} = (I - l_T(l_T' \Upsilon_T^{-1} l_T)^{-1} l_T' \Upsilon_T^{-1})V$ , which is statistically independent of  $\tilde{\beta} - \beta$ . Since  $\hat{\Omega}_b$  can be written as a quadratic form in  $\tilde{V}$ , it is also statistically independent of  $\tilde{\beta} - \beta$ . From [Casini \(2021b\)](#)

$$\Omega_T \triangleq \text{Var}(\sqrt{T}(\hat{\beta} - \beta)) = T^{-1} l_T' \Upsilon_T l_T = \Omega + O(T^{-1}), \quad (\text{S.22})$$

where  $\Omega = 2\pi \int_0^1 f(u, 0) du$ . Similarly, one can show that

$$\tilde{\Omega}_T \triangleq \text{Var}(\sqrt{T}(\tilde{\beta} - \beta)) = T(l_T' \Upsilon_T^{-1} l_T)^{-1} = \Omega + O(T^{-1}). \quad (\text{S.23})$$

Therefore  $T^{-1/2} l_T' \tilde{V} = \mathcal{N}(0, O(T^{-1}))$ . As in eq. (45) in [Sun, Phillips, and Jin \(2008\)](#), using the independence of  $\tilde{\beta}$  and  $(\tilde{V}, \hat{\Omega}_b)$ , we have

$$\begin{aligned} \mathbb{P}\left(\sqrt{T}(\hat{\beta} - \beta_0)/\sqrt{\hat{\Omega}_b} \leq z\right) \\ = \mathbb{P}\left(\sqrt{T}(\tilde{\beta} - \beta)/\sqrt{\tilde{\Omega}_T} + d/\sqrt{\tilde{\Omega}_T} \leq z\sqrt{\hat{\Omega}_b/\tilde{\Omega}_T}\right) + O(T^{-1}), \end{aligned} \quad (\text{S.24})$$

uniformly over  $z \in \mathbb{R}$  where  $\Phi$  and  $\varphi$  are the cdf and pdf of the standard normal distribution, respectively.

Similarly, uniformly over  $z \in \mathbb{R}$ , we have

$$\mathbb{P}\left(\sqrt{T}(\hat{\beta} - \beta_0)/\sqrt{\hat{\Omega}_b} \leq -z\right) = \mathbb{P}\left(\sqrt{T}(\tilde{\beta} - \beta)/\sqrt{\tilde{\Omega}_T} + c/\sqrt{\tilde{\Omega}_T} \leq -z\sqrt{\hat{\Omega}_b/\tilde{\Omega}_T}\right) + O(T^{-1}).$$

It follows that

$$\begin{aligned} Z_{T,d}(z) &= \mathbb{P}\left(\left|\sqrt{T}(\hat{\beta} - \beta_0)/\sqrt{\hat{\Omega}_b}\right| \leq z\right) \\ &= \mathbb{P}\left(\left(\sqrt{T}(\tilde{\beta} - \beta)/\sqrt{\tilde{\Omega}_T} + d/\sqrt{\tilde{\Omega}_T}\right)^2 \leq z^2 \hat{\Omega}_b/\tilde{\Omega}_T\right) + O(T^{-1}) \\ &= \mathbb{E}\left(G_{\tilde{d}}(z^2 \hat{\Omega}_b/\tilde{\Omega}_T)\right) = \mathbb{E}\left(G_{\tilde{d}}(z^2 \zeta_{b,T})\right) + O(T^{-1}), \end{aligned} \quad (\text{S.25})$$

uniformly over  $z \in \mathbb{R}_+$ , where  $G_{\tilde{d}}(z) = G(z; \tilde{d})$  is the cdf of a non-central chi-squared  $\chi_1(\tilde{d}^2)$  with noncentrality parameter  $\tilde{d}^2 = d^2/\Omega_T$  and  $\zeta_{b,T} = \hat{\Omega}_b/\Omega_T$ . Note that  $\zeta_{b,T} \Rightarrow \mathcal{G}_b/\Omega$ . Let  $\mu_{b,T} = \mathbb{E}(\zeta_{b,T})$  and consider a fourth-order Taylor expansion of  $\zeta_{b,T}$  around its mean,

$$\begin{aligned} Z_{T,d}(z) &= G_{\tilde{d}}(\mu_{b,T} z^2) + \frac{1}{2} G_{\tilde{d}}''(\mu_{b,T} z^2) \mathbb{E}(\zeta_{b,T} - \mu_{b,T})^2 z^4 \\ &\quad + \frac{1}{6} G_{\tilde{d}}'''(\mu_{b,T} z^2) \mathbb{E}(\zeta_{b,T} - \mu_{b,T})^3 z^6 + O(\mathbb{E}(\zeta_{b,T} - \mu_{b,T})^4) + O(T^{-1}), \end{aligned} \quad (\text{S.26})$$

where the  $O(\cdot)$  term holds uniformly over  $z \in \mathbb{R}_+$ .

Using (S.25) we have

$$\begin{aligned} Z_{T,0}(z) - Z_0(z) &= \mathbb{P}\left(\left|\frac{\sqrt{T}(\hat{\beta} - \beta_0)}{\sqrt{\hat{\Omega}_b}}\right| \leq z\right) - \mathbb{P}\left(\left|\frac{\int_0^1 \Sigma(u) dW_1(u)}{\sqrt{\mathcal{G}_b}}\right| \leq z\right) \\ &= \mathbb{E}\left(F_\chi\left(z^2 \zeta_{b,T}\right)\right) - \mathbb{E}\left(F_\chi\left(z^2 \mathcal{G}_b / \Omega\right)\right) + O\left(T^{-1}\right), \end{aligned} \quad (\text{S.27})$$

where  $F_\chi(\cdot) = G_0(\cdot)$  is the cdf of the  $\chi_1^2$  distribution. Next, we compute the cumulants of both  $\zeta_{b,T} - \mu_{b,T}$  and  $\Omega^{-1}(\mathcal{G}_b - \mu_b)$  where  $\mu_b = \mathbb{E}(\mathcal{G}_b)$ . Note that  $\zeta_{b,T}$  is a quadratic form in a Gaussian vector since  $\hat{\Omega}_b = T^{-1} \hat{V}' W_b \hat{V} = T^{-1} V' A_T W_b A_T V$ , where  $W_b$  is  $T \times T$  with  $(j, s)$ -th element  $K_b((j-s)/T)$  and  $A_T = I_T - l_T l_T' / T$ . The characteristic function of  $\zeta_{b,T} - \mu_{b,T}$  is given by

$$\psi_{b,T}(t) = \left| I - 2it \frac{\Upsilon_T A_T W_b A_T}{T \Omega_T} \right|^{-1/2} \exp(-it \mu_{b,T}), \quad (\text{S.28})$$

where  $\Upsilon_T = \mathbb{E}(uu')$  and the cumulant generating function is

$$\log(\psi_{b,T}(t)) = -\frac{1}{2} \log \left| I - 2it \frac{\Upsilon_T A_T W_b A_T}{T \Omega_T} \right| - it \mu_{b,T} = \sum_{m=1}^{\infty} \kappa_{m,T} \frac{(it)^m}{m!}, \quad (\text{S.29})$$

where  $\kappa_{m,T}$  is the  $m$ th cumulant of  $\zeta_{b,T} - \mu_{b,T}$ . Note that  $\kappa_{1,T} = 0$  and

$$\kappa_{m,T} = 2^{m-1} (m-1)! T^{-m} (\Omega_T)^{-m} \text{Tr}((\Upsilon_T A_T W_b A_T)^m), \quad m \geq 2. \quad (\text{S.30})$$

**Lemma S.A.3.** *Let Assumption 4.1-4.2 hold. We have: (i)  $\mu_{b,T} = \Omega^{-1} \mu_b + O(T^{-1})$ ; (ii)  $\kappa_{m,T} = \kappa_m + O(m! 2^m T^{-2} (\overline{C}_1 b)^{m-2})$  uniformly over  $m \geq 1$ ; (iii)  $\Xi_{m,T} = \mathbb{E}(\zeta_{b,T} - \mu_{b,T})^m = \Xi_m + O(m! 2^m T^{-2} (\overline{C}_1 b)^{m-2})$ .*

*Proof of Lemma S.A.3.* Note that  $\mu_{b,T} = (T \Omega_T)^{-1} \text{Tr}(\Upsilon_T A_T W_b A_T)$ . Let  $\widetilde{W}_b = A_T W_b A_T$ , where its  $(j, s)$ -th element is given by

$$\begin{aligned} \widetilde{K}_b\left(\frac{j}{T}, \frac{s}{T}\right) &= K_b\left(\frac{j-s}{T}\right) - \frac{1}{T} \sum_{p=1}^T K_b\left(\frac{j-p}{T}\right) \\ &\quad - \frac{1}{T} \sum_{q=1}^T K_b\left(\frac{q-s}{T}\right) + \frac{1}{T^2} \sum_{p=1}^T \sum_{q=1}^T K_b\left(\frac{p-q}{T}\right). \end{aligned} \quad (\text{S.31})$$

Let  $\Gamma_{r_1/T}(r_1 - r_2) = \mathbb{E}(V_{r_1} V_{r_2})$ . We have

$$\begin{aligned} &\text{Tr}(\Upsilon_T \widetilde{W}_b) \\ &= \sum_{1 \leq r_1, r_2 \leq T} \mathbb{E}(V_{r_1} V_{r_2}) \widetilde{K}_b\left(\frac{r_1}{T}, \frac{r_2}{T}\right) \\ &= \sum_{r_2=1}^T \sum_{h=1-r_2}^{T-r_2} \Gamma_{r_2/T}(-h) \widetilde{K}_b\left(\frac{r_2+h}{T}, \frac{r_2}{T}\right) \end{aligned} \quad (\text{S.32})$$

$$= \left( \sum_{h=1}^{T-1} \sum_{r_2=1}^{T-h} + \sum_{h=1-T}^0 \sum_{r_2=1-h}^T \right) \Gamma_{r_2/T}(-h) \widetilde{K}_b \left( \frac{r_2+h}{T}, \frac{r_2}{T} \right).$$

Using the Lipschitz property of  $K(\cdot)$  and the fact that  $\sup_{r_2, h} |\Gamma_{r_2/T}(-h)| < \infty$  which follows from Assumption 4.1, some algebra shows that

$$\begin{aligned} & \sum_{r_2=1}^{T-h} \Gamma_{r_2/T}(-h) \widetilde{K}_b \left( \frac{r_2+h}{T}, \frac{r_2}{T} \right) \\ &= \sum_{r_2=1}^{T-h} \Gamma_{r_2/T}(-h) K_b \left( \frac{h}{T} \right) - \frac{1}{T} \sum_{r_2=1}^{T-h} \sum_{p=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{r_2+h-p}{T} \right) \\ & \quad - \frac{1}{T} \sum_{r_2=1}^{T-h_1} \sum_{q=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{q-r_2}{T} \right) + \sum_{r_2=1}^{T-h_1} \frac{1}{T^2} \sum_{p=1}^T \sum_{q=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{p-q}{T} \right) \\ &= -\frac{1}{T} \sum_{r_2=1}^T \sum_{p=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{r_2-p}{T} \right) - \frac{1}{T} \sum_{r_2=1}^T \sum_{q=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{q-r_2}{T} \right) \\ & \quad + \sum_{r_2=1}^T \frac{1}{T^2} \sum_{p=1}^T \sum_{q=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{p-q}{T} \right) + \sum_{r_2=1}^{T-h} \Gamma_{r_2/T}(-h) K_b \left( \frac{h}{T} \right) + O(|h|) \\ &= -\frac{1}{T} \sum_{r_2=1}^T \sum_{p=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{r_2-p}{T} \right) + \sum_{r_2=1}^{T-h} \Gamma_{r_2/T}(-h) K_b \left( \frac{h}{T} \right) + O(|h|) + o(1) \\ &= \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) K_b(0) - \frac{1}{T} \sum_{r_2=1}^T \sum_{p=1}^T \Gamma_{r_2/T}(-h) K_b \left( \frac{r_2-p}{T} \right) \\ & \quad + \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \left( K_b \left( \frac{h}{T} \right) - K_b(0) \right) + O(|h|) + o(1) \\ &= \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \widetilde{K}_b \left( \frac{r_2}{T}, \frac{r_2}{T} \right) + \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \left( K_b \left( \frac{h}{T} \right) - K_b(0) \right) + O(|h|) + o(1). \end{aligned} \tag{S.33}$$

The same arguments yield

$$\begin{aligned} & \sum_{r_2=1-h}^T \Gamma_{r_2/T}(-h) \widetilde{K}_b \left( \frac{r_2+h}{T}, \frac{r_2}{T} \right) \\ &= \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \widetilde{K}_b \left( \frac{r_2}{T}, \frac{r_2}{T} \right) + \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \left( K_b \left( \frac{h}{T} \right) - K_b(0) \right) + O(|h|) + o(1). \end{aligned} \tag{S.34}$$

Using (S.33)-(S.34) into (S.32), we have

$$\begin{aligned} & \text{Tr} \left( \Upsilon_T \widetilde{W}_b \right) \\ &= \sum_{h=-\infty}^{\infty} \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \widetilde{K}_b \left( \frac{r_2}{T}, \frac{r_2}{T} \right) + \sum_{h=-\infty}^{\infty} \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \left( K_b \left( \frac{h}{T} \right) - K_b(0) \right) + O(1) \end{aligned} \tag{S.35}$$

$$\begin{aligned}
&= \sum_{h=-\infty}^{\infty} \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \widetilde{K}_b\left(\frac{r_2}{T}, \frac{r_2}{T}\right) + (Tb)^{-q} \sum_{r_2=1}^T \sum_{h=-\infty}^{\infty} |h|^q \Gamma_{r_2/T}(-h) \left( \frac{K(h/Tb) - K(0)}{|h/(Tb)|^q} \right) + O(1) \\
&= \sum_{h=-\infty}^{\infty} \sum_{r_2=1}^T \Gamma_{r_2/T}(-h) \widetilde{K}_b\left(\frac{r_2}{T}, \frac{r_2}{T}\right) + (Tb)^{-q} q_0 \sum_{r_2=1}^T \sum_{h=-\infty}^{\infty} |h|^q \Gamma_{r_2/T}(-h) (1 + o(1)) + O(1).
\end{aligned}$$

By Theorem 2.1 in [Casini \(2021b\)](#) and Lemma 4.1 in [Casini, Deng, and Perron \(2021\)](#),

$$\sum_{h=-\infty}^{\infty} \Gamma_{s/T}(-h) = 2\pi f(s/T, 0) \left(1 + O\left(\frac{1}{T}\right)\right), \quad (\text{S.36})$$

and

$$\frac{1}{T} \sum_{h=-\infty}^{\infty} \sum_{s=1}^T \Gamma_{s/T}(-h) \widetilde{K}_b\left(\frac{s}{T}, \frac{s}{T}\right) = \int_0^1 \Omega(u) K_b^*(u, u) du + O\left(\frac{1}{T}\right), \quad (\text{S.37})$$

where we have used  $2\pi f(u, 0) = \Omega(u)$ . Using Assumption 4.1 and (S.37), we yield

$$\begin{aligned}
\mu_{b,T} &= \Omega^{-1} \int_0^1 \Omega(s) K_b^*(s, s) ds \\
&\quad + (Tb)^{-q} q_0 \left( \Omega_T^{-1} T^{-1} \sum_{r_2=1}^T \sum_{h=-\infty}^{\infty} |h|^q \Gamma_{r_2/T}(-h) \right) (1 + o(1)) + O\left(\frac{1}{T}\right).
\end{aligned} \quad (\text{S.38})$$

Since  $\mu_b = \Omega^{-1} \mathbb{E}(\mathcal{G}_b) = \Omega^{-1} \int_0^1 K_b^*(s, s) \Omega(s) ds$ ,  $b$  is fixed and  $q \geq 1$ , we have  $\mu_{b,T} = \mu_b + O(T^{-1})$ .

Next, we consider part (ii). For  $m > 1$ , let  $r_{2m+1} = r_1$ ,  $r_{2m+2} = r_2$  and  $h_{m+1} = h_1$ . Using the same argument used for the case  $m = 1$  in (S.33) and using eq. (A.26) in [Sun, Phillips, and Jin \(2008\)](#), we have

$$\begin{aligned}
\text{Tr}\left((\Upsilon_T \widetilde{W}_b)^m\right) &= \sum_{r_1, r_2, \dots, r_{2m+1}=1}^T \prod_{j=1}^m \mathbb{E}\left(V_{r_{2j-1}} V_{r_{2j}}\right) \widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+1}}{T}\right) \\
&= \sum_{r_1, r_2, \dots, r_{2m+1}=1}^T \sum_{h_1=1-r_2}^{T-r_2} \sum_{h_2=1-r_4}^{T-r_4} \cdots \sum_{h_m=1-r_{2m}}^{T-r_{2m}} \prod_{j=1}^m \Gamma_{r_{2j}/T}(-h_j) \widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+1} + h_{j+1}}{T}\right) \\
&= \left( \sum_{h_1=1}^{T-1} \sum_{r_2=1}^{T-h_1} + \sum_{h_1=1-T}^0 \sum_{r_2=1-h_1}^T \right) \cdots \left( \sum_{h_m=1}^{T-1} \sum_{r_{2m}=1}^{T-h_m} + \sum_{h_m=1-T}^0 \sum_{r_{2m}=1-h_m}^T \right) \\
&\quad \times \prod_{j=1}^m \Gamma_{r_{2j}/T}(-h_j) \widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+1} + h_{j+1}}{T}\right) \\
&= L_1 + L_2,
\end{aligned} \quad (\text{S.39})$$

where

$$L_1 = \left( \sum_{h_1=1}^{T-1} \sum_{r_2=1}^{T-h_1} + \sum_{h_1=1-T}^0 \sum_{r_2=1-h_1}^T \right) \cdots \left( \sum_{h_m=1}^{T-1} \sum_{r_{2m}=1}^{T-h_m} + \sum_{h_m=1-T}^0 \sum_{r_{2m}=1-h_m}^T \right) \quad (\text{S.40})$$

$$\prod_{j=1}^m \Gamma_{r_{2j}/T}(-h_j) \widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+1} + h_{j+1}}{T}\right),$$

and

$$L_2 = O\left(\left(\sum_{h_1=1}^{T-1} \sum_{r_2=1}^{T-h_1} + \sum_{h_1=1-T}^0 \sum_{r_2=1-h_1}^T\right) \cdots \left(\sum_{h_m=1}^{T-1} \sum_{r_{2m}=1}^{T-h_m} + \sum_{h_m=1-T}^0 \sum_{r_{2m}=1-h_m}^T\right)\right. \\ \left.\prod_{j=1}^m \Gamma_{r_{2j}/T}(-h_j) \left(\frac{|h_{j+1}|}{Tb}\right)\right). \quad (\text{S.41})$$

Using the same arguments as in eq. (A.30)-(A.31) in [Sun, Phillips, and Jin \(2008\)](#), we can show that

$$L_1 = \sum_{h=-\infty}^{\infty} \sum_r \prod_{j=1}^m \Gamma_{r_{2j}/T}(-h_j) \left(\widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+2}}{T}\right)\right) + O\left(2mT^{m-2} (\overline{C}_1 b)^{m-2}\right),$$

where  $O(2mT^{m-2}((\overline{C}_1 b))^{m-2})$  follows from

$$\sum_{h_1=-\infty}^{\infty} \sum_{r_2=1}^T \cdots \sum_{h_a=-\infty}^{\infty} \sum_{r_{2a}=1}^{-h_a} \cdots \sum_{h_m=-\infty}^{\infty} \sum_{r_{2m}=1}^T \prod_{j=1}^m \left(\sup_{r_{2j}} |\Gamma_{r_{2j}/T}(-h_j)|\right) |h_a| \prod_{j \neq a} \left|\widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+2}}{T}\right)\right| \\ \leq \left[\sup_t \sum_{s=1}^T \widetilde{K}_b\left(\frac{s}{T}, \frac{t}{T}\right)\right]^{m-2} \left(\sum_{h=-\infty}^{\infty} \sup_s |\Gamma_{s/T}(-h)|\right)^{m-1} \left(\sum_{h_a=-\infty}^{\infty} \sup_s |\Gamma_{s/T}(-h_a)| |h_a|\right) \\ \leq O\left(2mT^{m-2} (\overline{C}_1 b)^{m-2}\right),$$

uniformly over  $m$  for some integer  $a$  such that  $1 \leq a \leq m$ . A similar argument yields that  $L_2 = o(2mT^{m-2}(\overline{C}_1 b)^{m-2})$  uniformly over  $m$ . Thus,

$$\text{Tr}\left(\left(\Upsilon_T \widetilde{W}_b\right)^m\right) = \sum_{h=-\infty}^{\infty} \sum_r \prod_{j=1}^m \Gamma_{r_{2j}/T}(-h_j) \left(\widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+2}}{T}\right)\right) + O\left(2mT^{m-2} (\overline{C}_1 b)^{m-2}\right),$$

and using  $\tau_1 = \tau_{m+1}$  we yield

$$\kappa_{m,T} = 2^{m-1} (m-1)! T^{-m} \Omega_T^{-m} \text{Tr}\left((\Upsilon_T A_T W_b A_T)^m\right) \quad (\text{S.42}) \\ = 2^{m-1} (m-1)! \Omega_T^{-m} \left(T^{-m} \sum_r \Omega(r_{2j}/T) \widetilde{K}_b\left(\frac{r_{2j}}{T}, \frac{r_{2j+2}}{T}\right) + O\left(2mT^{-2} (\overline{C}_1 b)^{m-2}\right)\right) \\ = 2^{m-1} (m-1)! \Omega^{-m} \left(\int_0^1 \cdots \int_0^1 \left(\prod_{j=1}^m \Omega(\tau_j) K_b^*(\tau_j, \tau_{j+1})\right) d\tau_1 \cdots d\tau_m + O\left(2mT^{-2} (\overline{C}_1 b)^{m-2}\right)\right) \\ = \kappa_m + O\left(2mT^{-2} (\overline{C}_1 b)^{m-2}\right),$$

uniformly over  $m$ .

Next, we consider part (iii). From (S.17) and part (ii), we have uniformly over  $m$ ,

$$\begin{aligned}\Xi_{m,T} &= \Xi_m + O\left(\frac{m!2^m}{T^2} (\overline{C}_1 b)^{m-2} \sum_{\pi} \frac{m!}{m_1! m_2! \cdots m_k!}\right) \\ &= \Xi_m + O\left(\frac{m!2^{2m}}{T^2} (\overline{C}_1 b)^{m-2}\right),\end{aligned}\tag{S.43}$$

where we have used  $\sum_{\pi} \frac{m!}{m_1! m_2! \cdots m_k!} < 2^m$ .  $\square$

*Proof of Theorem 4.1.* Let  $F_{\chi}^{(m)}(\cdot)$  denote the  $m$ th derivative of  $F_{\chi}(\cdot)$ . Since  $F_{\chi}(\cdot)$  is a bounded function, we can write

$$\begin{aligned}\mathbb{P}\left(\left|\frac{\int_0^1 \Sigma(u) dW_1(u)}{\sqrt{\mathcal{G}_b}}\right| \leq z\right) &= \lim_{C \rightarrow \infty} \mathbb{E}\left(F_{\chi}\left(z^2 \mathcal{G}_b / \Omega\right) \mathbf{1}(|\mathcal{G}_b - \mu_b| \leq \Omega C)\right) \\ &= \lim_{C \rightarrow \infty} \mathbb{E} \sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)}\left(\mu_b z^2 / \Omega\right) \Omega^{-m} (\mathcal{G}_b - \mu_b)^m z^{2m} \mathbf{1}\{|\mathcal{G}_b - \mu_b| \leq \Omega C\} \\ &= \lim_{C \rightarrow \infty} \sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)}\left(\mu_b z^2 / \Omega\right) \Xi_m z^{2m} \mathbf{1}\{|\mathcal{G}_b - \mu_b| \leq \Omega C\},\end{aligned}\tag{S.44}$$

where  $\Xi_m = \Omega^{-m} \mathbb{E}((\mathcal{G}_b - \mu_b)^m)$ . Since  $F_{\chi}(z^2)$  decays exponentially as  $z \rightarrow \infty$ , there exists a constant  $C_2 > 0$  such that  $|F_{\chi}^{(m)}(\mu_b z^2 / \Omega) z^{2m}| < C_2$  for all  $m$ . Using Lemma S.A.2, we yield

$$\begin{aligned}\left|\sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)}\left(\mu_b z^2 / \Omega\right) \Xi_m z^{2m}\right| &\leq C_2 \sum_{m=1}^{\infty} \frac{1}{m!} \Xi_m \leq C_2 D \sum_{m=1}^{\infty} \frac{1}{m!} 2^{2m} m! C_{\Omega}^m (\overline{C}_1 b)^{m-1} \\ &= C_2 D (\overline{C}_1 b)^{-1} \sum_{m=1}^{\infty} (4C_{\Omega} \overline{C}_1 b)^m,\end{aligned}\tag{S.45}$$

where  $D_m \leq D$  for some  $D < \infty$ . The right-hand side of (S.45) is bounded in view of  $b < 1/(4C_{\Omega} \overline{C}_1)$ . This implies that

$$\mathbb{P}\left(\left|\frac{\int_0^1 \Sigma(u) dW_1(u)}{\sqrt{\mathcal{G}_b}}\right| \leq z\right) = \sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)}\left(\mu_b z^2 / \Omega\right) \Xi_m z^{2m},\tag{S.46}$$

provided that  $b < 1/(4C_{\Omega} \overline{C}_1)$ .

From (S.25) we have

$$Z_{T,0}(z) = \mathbb{P}\left(\left|\frac{\sqrt{T}(\hat{\beta} - \beta_0)}{\sqrt{\hat{\Omega}_b}}\right| \leq z\right) = \mathbb{E}\left(F_{\chi}\left(z^2 \zeta_{b,T}\right)\right) + O\left(T^{-1}\right).\tag{S.47}$$

Using a similar argument as in (S.44),

$$\mathbb{E}\left(F_{\chi}\left(z^2 \zeta_{b,T}\right)\right) - \sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)}\left(\mu_{b,T} z^2\right) \Xi_{m,T} z^{2m} \rightarrow 0,\tag{S.48}$$



uniformly over  $T$  since by Lemma S.A.3-(iii) we have

$$\Xi_{m,T} = \Xi_m + O\left(\frac{2^{2m}m!}{T^2}(\bar{C}_1 b)^{m-2}\right),$$

uniformly in  $m$  and  $|F_\chi^{(m)}(\mu_{b,T} z^2) z^{2m}| < C_2$  for some constant  $C_2 > 0$  for all  $m$  so that

$$\left| \sum_{m=1}^{\infty} \frac{1}{m!} F_\chi^{(m)}(\mu_{b,T} z^2) \Xi_{m,T} z^{2m} \right| \leq C_2 \sum_{m=1}^{\infty} \frac{1}{m!} |\Xi_m| + \frac{C_2}{T^2} \sum_{m=1}^{\infty} 2^{2m} (\bar{C}_1 b)^{m-2} < \infty,$$

provided that  $b < 1/(4\bar{C}_1)$ . Note that  $b < 1/(16C_{2,\Omega} \int_{-\infty}^{\infty} |k(x)| dx) < 1/(4\bar{C}_1)$  by assumption. It follows that

$$Z_{T,0}(z) = \sum_{m=1}^{\infty} \frac{1}{m!} F_\chi^{(m)}(\mu_{b,T} z^2) \Xi_{m,T} z^{2m} + O(T^{-1}), \quad (\text{S.49})$$

uniformly over  $z \in \mathbb{R}_+$ .

By Lemma S.A.3-(i), we have

$$F_\chi^{(m)}(\mu_{b,T} z^2) = F_\chi^{(m)}(\mu_b z^2 / \Omega) + O(F_\chi^{(m+1)}(\mu_b z^2 / \Omega) z^2 T^{-1}). \quad (\text{S.50})$$

Combining (S.46) and (S.49)-(S.50) leads to

$$\begin{aligned} & |Z_{T,0}(z) - Z_0(z)| \\ &= \left| \sum_{m=1}^{\infty} \frac{1}{m!} F_\chi^{(m)}(\mu_{b,T} z^2) \Xi_{m,T} z^{2m} - \sum_{m=1}^{\infty} \frac{1}{m!} F_\chi^{(m)}(\mu_b z^2) \Xi_m z^{2m} \right| + O\left(\frac{1}{T}\right) \\ &= \left| \sum_{m=1}^{\infty} \frac{1}{m!} F_\chi^{(m)}(\mu_b z^2) z^{2m} (\Xi_{m,T} - \Xi_m) \right| + O\left(\frac{1}{T}\right) \\ &= \left| \sum_{m=1}^{\infty} \frac{1}{m!} F_\chi^{(m)}(\mu_b z^2) z^{2m} O\left(\frac{m! 2^{2m} (\bar{C}_1 b)^{m-2}}{T^2}\right) \right| + O\left(\frac{1}{T}\right) \\ &= O\left(\frac{1}{T^2} \sum_{m=1}^{\infty} 2^{2m} (\bar{C}_1 b)^{m-2}\right) + O\left(\frac{1}{T}\right) \\ &= O\left(\frac{1}{T}\right). \end{aligned} \quad (\text{S.51})$$

uniformly over  $z \in \mathbb{R}$  where we have used Lemma S.A.3-(iii). Hence,  $D_1 = O(T^{-1})$ .

Let  $\hat{\Omega} = \int_0^1 \hat{\Omega}(u) du$  where  $\hat{\Omega}(u)$  was defined in (4.3). Note that  $\hat{\mathcal{G}}_b = \mathcal{G}_b + O((Th_1 h_2)^{-1/2})$  by definition of  $\hat{\Omega}(u)$ . Using this and proceeding as in (S.44), we yield

$$\begin{aligned} \hat{Z}_0(z) &= \mathbb{P}\left(\left|\frac{\int_0^1 \hat{\Sigma}(u) dW_p(u)}{\sqrt{\hat{\mathcal{G}}_b}}\right| \leq z\right) \\ &= \lim_{C \rightarrow \infty} \mathbb{E}\left(F_\chi\left(z^2 \hat{\mathcal{G}}_b / \hat{\Omega}\right) \mathbf{1}\left(|\hat{\mathcal{G}}_b - \hat{\mu}_b| \leq \hat{\Omega} C\right)\right) \end{aligned} \quad (\text{S.52})$$

$$\begin{aligned}
&= \lim_{C \rightarrow \infty} \mathbb{E} \sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)} \left( \hat{\mu}_b z^2 / \hat{\Omega} \right) \hat{\Omega}^{-m} \left( \hat{\mathcal{G}}_b - \hat{\mu}_b \right)^m z^{2m} \mathbf{1} \left\{ \left| \hat{\mathcal{G}}_b - \hat{\mu}_b \right| \leq \Omega C \right\} \\
&= \lim_{C \rightarrow \infty} \mathbb{E} \sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)} \left( \mu_b z^2 / \Omega \right) \Omega^{-m} \left( \mathcal{G}_b - \mu_b \right)^m z^{2m} \mathbf{1} \left\{ \left| \mathcal{G}_b - \mu_b \right| \leq \Omega C \right\} + O((Th_1 h_2)^{-1/2}) \\
&= \lim_{C \rightarrow \infty} \sum_{m=1}^{\infty} \frac{1}{m!} F_{\chi}^{(m)} \left( \mu_b z^2 / \Omega \right) \Xi_m z^{2m} \mathbf{1} \left\{ \left| \mathcal{G}_b - \mu_b \right| \leq \Omega C \right\} + O \left( (Th_1 h_2)^{-1/2} \right),
\end{aligned}$$

uniformly in  $z \in \mathbb{R}_+$ . This implies  $D_2 = O((Th_1 h_2)^{-1/2})$ .  $\square$

## References

- CASINI, A. (2021): “Theory of Evolutionary Spectra for Heteroskedasticity and Autocorrelation Robust Inference in Possibly Misspecified and Nonstationary Models,” *arXiv preprint arXiv:2103.02981*.
- CASINI, A., T. DENG, AND P. PERRON (2021): “Theory of Low Frequency Contamination from Nonstationarity and Misspecification: Consequences for HAR Inference,” *arXiv preprint arXiv:2103.01604*.
- KIEFER, N. M., AND T. J. VOGELSANG (2002a): “Heteroskedasticity-Autocorrelation Robust Standard Errors using the Bartlett Kernel without Truncation,” *Econometrica*, 70(5), 2093–2095.
- (2002b): “Heteroskedasticity-Autocorrelation Robust Testing Using Bandwidth Equal to Sample Size,” *Econometric Theory*, 18(6), 1350–1366.
- SUN, Y., P. C. B. PHILLIPS, AND S. JIN (2008): “Optimal Bandwidth Selection in Heteroskedasticity-Autocorrelation Robust Testing,” *Econometrica*, 76(1), 175–194.