

# Improvements on some partial trace inequalities for positive semidefinite block matrices\*

Yongtao Li\*

School of Mathematics, Hunan University  
Changsha, Hunan, 410082, P.R. China

January 24, 2022

Dedicated to Prof. Weijun Liu on his 60th birthday

## Abstract

We study matrix inequalities involving partial traces for positive semidefinite block matrices. First of all, we present a new method to prove a celebrated result of Choi [Linear Algebra Appl. 516 (2017)]. Our method also allows us to prove a generalization of another result of Choi [Linear Multilinear Algebra 66 (2018)]. Furthermore, we shall give an improvement on a recent result of Li, Liu and Huang [Operators and Matrices 15 (2021)]. In addition, we include with some majorization inequalities involving partial traces for two by two block matrices, and also provide inequalities related to the unitarily invariant norms as well as the singular values, which can be viewed as slight extensions of two results of Lin [Linear Algebra Appl. 459 (2014)] and [Electronic J. Linear Algebra 31 (2016)].

**Key words:** Partial transpose; Partial traces; Cauchy and Khinchin; Positive semidefinite.

**2010 Mathematics Subject Classification.** 15A45, 15A60, 47B65.

## 1 Introduction

The space of  $m \times n$  complex matrices is denoted by  $\mathbb{M}_{m \times n}$ ; if  $m = n$ , we write  $\mathbb{M}_n$  for  $\mathbb{M}_{n \times n}$  and if  $n = 1$ , we use  $\mathbb{C}^m$  for  $\mathbb{M}_{m \times 1}$ . By convention, if  $A \in \mathbb{M}_n$  is positive semidefinite, we write  $A \geq 0$ . For Hermitian matrices  $A$  and  $B$  with the same size,  $A \geq B$  means that  $A - B$  is positive semidefinite, i.e.,  $A - B \geq 0$ . We denote by  $\mathbb{M}_m(\mathbb{M}_n)$  the set of  $m \times m$  block matrices with each block in  $\mathbb{M}_n$ . Each element of  $\mathbb{M}_m(\mathbb{M}_n)$  is also viewed as an  $mn \times mn$  matrix with numerical entries and usually written as  $A = [A_{i,j}]_{i,j=1}^m$ , where  $A_{ij} \in \mathbb{M}_n$ . We

---

\*E-mail addresses: ytli0921@hnu.edu.cn (Y. Li).

denote by  $A \otimes B$  the Kronecker product of  $A$  with  $B$ , that is, if  $A = [a_{i,j}] \in \mathbb{M}_m$  and  $B \in \mathbb{M}_n$ , then  $A \otimes B \in \mathbb{M}_m(\mathbb{M}_n)$  whose  $(i,j)$  block is  $a_{i,j}B$ . We write  $A \circ B$  for the Hadamard product if  $A$  and  $B$  have the same size.

In the present paper, we are mainly concentrated on the block positive semidefinite matrices. Given  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$ , the partial transpose of  $A$  is defined as

$$A^\tau = [A_{j,i}]_{i,j=1}^m = \begin{bmatrix} A_{1,1} & \cdots & A_{m,1} \\ \vdots & \ddots & \vdots \\ A_{1,m} & \cdots & A_{m,m} \end{bmatrix}.$$

This is different from the usual transpose, which is defined as

$$A^T = [A_{j,i}^T]_{i,j=1}^m = \begin{bmatrix} A_{1,1}^T & \cdots & A_{m,1}^T \\ \vdots & \ddots & \vdots \\ A_{1,m}^T & \cdots & A_{m,m}^T \end{bmatrix}.$$

The primary application of partial transpose is materialized in quantum information theory [17, 15, 34]. It is easy to see that  $A \geq 0$  does not necessarily imply  $A^\tau \geq 0$ . If both  $A$  and  $A^\tau$  are positive semidefinite, then  $A$  is said to be positive partial transpose (or PPT for short). For more explanations and applications of the PPT matrices, we recommend a comprehensive monograph [6], and see, e.g., [27, 20, 41, 14] for recent results. Next we introduce the definition of two partial traces  $\text{tr}_1 A$  and  $\text{tr}_2 A$  of  $A$ .

$$\text{tr}_1 A = \sum_{i=1}^m A_{i,i} \quad \text{and} \quad \text{tr}_2 A = [\text{tr} A_{i,j}]_{i,j=1}^m,$$

where  $\text{tr}(\cdot)$  stands for the usual trace. Clearly, we have  $\text{tr}_1 A \in \mathbb{M}_n$  and  $\text{tr}_2 A \in \mathbb{M}_m$ .

It is believed that there are many elegant matrix inequalities that are arose from the probability theory and quantum information theory in the literature. As we all know, these two partial trace maps are linear and trace-preserving. Furthermore, if  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$  is positive semidefinite, it is easy to see that both  $\text{tr}_1 A$  and  $\text{tr}_2 A$  are also positive semidefinite; see, e.g., [38, p. 237] or [39, Theorem 2.1]. Over the years, various results involving partial transpose and partial traces have been obtained in the literature, e.g., [3, 9, 11, 13, 24]. We introduce the background and recent progress briefly. In the following results, we always assume that  $A \in \mathbb{M}_m(\mathbb{M}_n)$  is a positive semidefinite block matrix.

(R1) Choi [9] presented by using induction on  $m$  that  $I_m \otimes \text{tr}_1 A^\tau \geq A^\tau$ .

(R2) Zhang [41] revisited Choi's result and proved  $(\text{tr}_2 A^\tau) \otimes I_n \geq A^\tau$ .

(R3) Choi [11] further extended the result of Zhang and proved

$$I_m \otimes \text{tr}_1 A^\tau \geq \pm A^\tau, \tag{1}$$

and

$$(\operatorname{tr}_2 A^\tau) \otimes I_n \geq \pm A^\tau. \quad (2)$$

(R4) Ando [3] (or see [30] for an alternative proof) revealed a well connection between the first partial trace and the second partial trace, and established

$$(\operatorname{tr} A) I_{mn} - (\operatorname{tr}_2 A) \otimes I_n \geq I_m \otimes (\operatorname{tr}_1 A) - A. \quad (3)$$

(R5) Motivated by Choi's result (1) and Ando's result (3), Li, Liu and Huang [24] (or see [25] for a unified treatment) proved recently an analogous complement, which states that

$$(\operatorname{tr} A) I_{mn} - (\operatorname{tr}_2 A) \otimes I_n \geq \pm (I_m \otimes (\operatorname{tr}_1 A) - A), \quad (4)$$

and they also obtained

$$(\operatorname{tr} A) I_{mn} \pm (\operatorname{tr}_2 A) \otimes I_n \geq A \pm I_m \otimes (\operatorname{tr}_1 A). \quad (5)$$

We will present some partial trace inequalities for positive semidefinite block matrices which improve the abovementioned results. The paper is organized as follows. In Section 2, we shall give a new method to prove Choi's result (R1). This new method can allow us to show an improvement on inequality (1); see Theorems 2.1 and 2.2. In Section 3, we present inequalities for the second partial trace and prove an improvement on inequality (2); see Theorem 3.1. In Section 4, we shall prove some inequalities involving both the first and second partial trace. Our results can also be viewed as the well improvements on inequalities (4) and (5); see Theorems 4.2, 4.3 and 4.4. In Section 5, we give an application on Cauchy–Khinchin's inequality by using Theorems 4.2 and 4.3; see Corollary 5.2. In Section 6, we study the majorization inequalities related to partial traces for two by two block matrices; see Theorem 6.2. We also prove some inequalities about the unitarily invariant norms as well as the singular values, which extend slightly two recent results of Lin [27] and [31]; see Theorems 6.4 and 6.7.

## 2 Inequalities about the first partial trace

In this section, we first give a short and succinct proof of Choi's result (R1). For convenience, we write it as the following theorem. The original proof stated in [9, Theorem 2] uses a standard decomposition of positive semidefinite matrices and then applies the inductive technique. Our proof is quite different and more transparent than the original proof. As an application of our method, we then present an improvement on inequality (1) of Choi's result (R3).

**Theorem 2.1** [9] Let  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then

$$I_m \otimes \text{tr}_1 A^\tau \geq A^\tau.$$

**Proof.** Denote  $D_A := A_{1,1} \oplus A_{2,2} \oplus \cdots \oplus A_{m,m}$ . We are going to prove

$$I_m \otimes \text{tr}_1 A^\tau + (m-2)D_A \geq A^\tau + (m-2)D_A,$$

which is the same as

$$\begin{aligned} & \begin{bmatrix} (m-1)A_{1,1} + \sum_{i \neq 1} A_{i,i} & 0 & \cdots & 0 \\ 0 & (m-1)A_{2,2} + \sum_{i \neq 2} A_{i,i} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & (m-1)A_{m,m} + \sum_{i \neq m} A_{i,i} \end{bmatrix} \\ \geq & \begin{bmatrix} (m-1)A_{1,1} & A_{2,1} & \cdots & A_{m,1} \\ A_{1,2} & (m-1)A_{2,2} & \cdots & A_{m,2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1,m} & A_{2,m} & \cdots & (m-1)A_{m,m} \end{bmatrix}. \end{aligned} \quad (6)$$

It suffices to show that for every pair  $(i, j)$  with  $1 \leq i < j \leq m$ ,

$$\begin{bmatrix} & i\text{-th} & & j\text{-th} & \\ & \vdots & & \vdots & \\ \cdots & A_{i,i} + A_{j,j} & \cdots & 0 & \cdots \\ & \vdots & & \vdots & \\ \cdots & 0 & \cdots & A_{i,i} + A_{j,j} & \cdots \\ & \vdots & & \vdots & \end{bmatrix} \geq \begin{bmatrix} & \vdots & & \vdots & \\ \cdots & A_{i,i} & \cdots & A_{j,i} & \cdots \\ & \vdots & & \vdots & \\ \cdots & A_{i,j} & \cdots & A_{j,j} & \cdots \\ & \vdots & & \vdots & \end{bmatrix}. \quad (7)$$

Indeed, we can sum inequalities (7) over all  $1 \leq i < j \leq m$ , which leads to the desired (6). Note that the omitted blocks in (7) are zero matrices, so we need to prove

$$\begin{bmatrix} A_{i,i} + A_{j,j} & 0 \\ 0 & A_{i,i} + A_{j,j} \end{bmatrix} \geq \begin{bmatrix} A_{i,i} & A_{j,i} \\ A_{i,j} & A_{j,j} \end{bmatrix}.$$

This inequality immediately holds by observing that

$$\begin{bmatrix} A_{j,j} & -A_{j,i} \\ -A_{i,j} & A_{i,i} \end{bmatrix} = \begin{bmatrix} 0 & -I_n \\ I_n & 0 \end{bmatrix} \begin{bmatrix} A_{i,i} & A_{i,j} \\ A_{j,i} & A_{j,j} \end{bmatrix} \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}.$$

Hence, we complete the proof. ■

As promised, we shall provide an improvement on inequality (1).

**Theorem 2.2** Let  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then

$$I_m \otimes \text{tr}_1 A^\tau \geq -A^\tau + 2D_A,$$

where  $D_A = A_{1,1} \oplus A_{2,2} \oplus \cdots \oplus A_{m,m}$ .

**Proof.** Our treatment is similar with the previous proof. We intend to prove

$$I_m \otimes \text{tr}_1 A^\tau - mD_A \geq -A^\tau - (m-2)D_A.$$

This inequality can be written as

$$\begin{aligned} & \begin{bmatrix} -(m-1)A_{1,1} + \sum_{i \neq 1} A_{i,i} & 0 & \cdots & 0 \\ 0 & -(m-1)A_{2,2} + \sum_{i \neq 2} A_{i,i} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -(m-1)A_{m,m} + \sum_{i \neq m} A_{i,i} \end{bmatrix} \\ & \geq \begin{bmatrix} -(m-1)A_{1,1} & -A_{2,1} & \cdots & -A_{m,1} \\ -A_{1,2} & -(m-1)A_{2,2} & \cdots & -A_{m,2} \\ \vdots & \vdots & \ddots & \vdots \\ -A_{1,m} & -A_{2,m} & \cdots & -(m-1)A_{m,m} \end{bmatrix}. \end{aligned}$$

Using a similar treatment as to the proof of Theorem 2.1, it is sufficient to prove

$$\begin{bmatrix} -A_{i,i} + A_{j,j} & 0 \\ 0 & -A_{j,j} + A_{i,i} \end{bmatrix} \geq \begin{bmatrix} -A_{i,i} & -A_{j,i} \\ -A_{i,j} & -A_{j,j} \end{bmatrix}$$

for every pair  $(i, j)$  with  $1 \leq i < j \leq m$ , which follows by noting that

$$\begin{bmatrix} A_{j,j} & A_{j,i} \\ A_{i,j} & A_{i,i} \end{bmatrix} = \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix} \begin{bmatrix} A_{i,i} & A_{i,j} \\ A_{j,i} & A_{j,j} \end{bmatrix} \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix}.$$

Thus, this completes the proof. ■

**Corollary 2.3** Let  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$  be Hermitian. Then

$$(m-1)\lambda_{\max}(A)I_{mn} + 2D_A \geq I_m \otimes \text{tr}_1 A^\tau + A^\tau \geq (m-1)\lambda_{\min}(A)I_{mn} + 2D_A,$$

where  $D_A = A_{1,1} \oplus A_{2,2} \oplus \cdots \oplus A_{m,m}$ .

**Proof.** The required result holds from Theorem 2.2 by replacing  $A$  with  $A - \lambda_{\min}(A)I_{mn}$  and  $\lambda_{\max}(A)I_{mn} - A$ . We leave the detailed proof to the readers. ■

It is easy to see that  $D_A$  is positive semidefinite whenever  $A$  is positive semidefinite. So Theorem 2.2 can yield the following corollary, which is partially an improvement on Choi's results (R3).

**Corollary 2.4** Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then

$$I_m \otimes \text{tr}_1 A^\tau \geq -A^\tau + (m-1)\lambda_{\min}(A)I_{mn}.$$

### 3 Inequalities about the second partial trace

We present some inequalities of the second partial trace in this section. First of all, we shall give an improvement on inequality (2). As a byproduct of Theorem 2.1 and the forthcoming Theorem 3.1, we can get Corollary 3.4 and Corollary 3.5, which were early established by Choi [11, Theorem 2] and Horodecki et al. [17] respectively.

To illustrate the relations between the first and second partial traces, we shall apply a useful technique, which was recently introduced by Choi in [11]. Assume that  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$ , where  $A_{i,j} = [a_{r,s}^{i,j}]_{r,s=1}^n$ . We define  $B_{r,s} := [a_{r,s}^{i,j}]_{i,j=1}^m \in \mathbb{M}_m$  and

$$\tilde{A} := [B_{r,s}]_{r,s=1}^n \in \mathbb{M}_n(\mathbb{M}_m). \quad (8)$$

Next, we make a brief review of some useful properties. Clearly, we have

$$\tilde{A}^\tau = (\tilde{A}^\tau)^T = \widetilde{(A^\tau)^T}. \quad (9)$$

Indeed, we can observe that

$$\tilde{A}^\tau = [B_{s,r}]_{r,s=1}^n = \left[ [a_{s,r}^{i,j}]_{i,j=1}^m \right]_{r,s=1}^n.$$

Since  $A^\tau = [A_{j,i}]_{i,j=1}^m = \left[ [a_{r,s}^{j,i}]_{r,s=1}^n \right]_{i,j=1}^m$ , we then get

$$(\tilde{A}^\tau)^T = \left( \left[ [a_{r,s}^{j,i}]_{r,s=1}^n \right]_{r,s=1}^m \right)^T = \left[ [a_{s,r}^{i,j}]_{i,j=1}^m \right]_{r,s=1}^n$$

and

$$\widetilde{(A^\tau)^T} = \left[ [a_{s,r}^{i,j}]_{r,s=1}^n \right]_{i,j=1}^m = \left[ [a_{s,r}^{i,j}]_{i,j=1}^m \right]_{r,s=1}^n.$$

In addition, it is easy to check that

$$\mathrm{tr}_2 A = [\mathrm{tr} A_{i,j}]_{i,j=1}^m = \left[ \sum_{r=1}^n a_{r,r}^{i,j} \right]_{i,j=1}^m = \sum_{r=1}^n [a_{r,r}^{i,j}]_{i,j=1}^m = \sum_{r=1}^n B_{r,r} = \mathrm{tr}_1 \tilde{A}. \quad (10)$$

For any  $X = [x_{i,j}]_{i,j=1}^m \in \mathbb{M}_m$  and  $Y = [y_{r,s}]_{r,s=1}^n \in \mathbb{M}_n$ , we know that

$$X \otimes Y = [x_{i,j} Y]_{i,j=1}^m = \left[ [x_{i,j} y_{r,s}]_{r,s=1}^n \right]_{i,j=1}^m.$$

Then it follows that

$$\widetilde{X \otimes Y} = \left[ [x_{i,j} y_{r,s}]_{i,j=1}^m \right]_{r,s=1}^n = [y_{r,s} X]_{r,s=1}^n = Y \otimes X. \quad (11)$$

Clearly, both  $A$  and  $\tilde{A}$  are matrices of order  $mn \times mn$ . Furthermore,  $A$  and  $\tilde{A}$  are unitarily similar; see, e.g., [10] or [22]. The similarity implies that  $\tilde{A}$  is positive semidefinite whenever  $A$  is positive semidefinite.

**Theorem 3.1** *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then*

$$(\mathrm{tr}_2 A^\tau) \otimes I_n \geq -A^\tau + 2A^\tau \circ J,$$

where  $J$  is the  $m \times m$  block matrix with each block  $I_n$ .

**Proof.** Replacing  $A$  with  $\tilde{A}$  in Theorem 2.2, we have

$$I_n \otimes \mathrm{tr}_1 \tilde{A}^\tau \geq -\tilde{A}^\tau + 2D_{\tilde{A}}.$$

Applying the relation (9), we can get

$$I_n \otimes \mathrm{tr}_1(\widetilde{A^\tau})^T \geq -(\widetilde{A^\tau})^T + 2D_{\tilde{A}}.$$

Observe that  $X \geq Y$  implies  $\tilde{X} \geq \tilde{Y}$ . Invoking the facts (10) and (11), we obtain

$$(\mathrm{tr}_2(A^\tau)^T) \otimes I_n = \mathrm{tr}_1(\widetilde{A^\tau})^T \otimes I_n \geq -(A^\tau)^T + 2\widetilde{D_{\tilde{A}}}. \quad (12)$$

Recall in (8) that  $B_{r,s} = [a_{r,s}^{i,j}]_{i,j=1}^m \in \mathbb{M}_m$ . A direct computation reveals that

$$\widetilde{D_{\tilde{A}}} = \widetilde{\mathrm{diag}(B_{1,1}, B_{2,2}, \dots, B_{n,n})} = [A_{i,j} \circ I_n]_{i,j=1}^m.$$

Note that  $(X \otimes Y)^T = X^T \otimes Y^T$  and

$$(\widetilde{D_{\tilde{A}}})^T = [(A_{j,i} \circ I_n)^T]_{i,j=1}^m = [A_{j,i} \circ I_n]_{i,j=1}^m = A^\tau \circ J.$$

Taking transpose in both sides of (12) yields the required result. ■

We replace  $A$  with  $A - \lambda_{\min}(A)I_{mn}$  and  $\lambda_{\max}(A)I_{mn} - A$  in Theorem 3.1, which yields the following corollary.

**Corollary 3.2** *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be Hermitian. Then*

$$(n-1)\lambda_{\max}(A)I_{mn} + 2A^\tau \circ J \geq (\mathrm{tr}_2 A^\tau) \otimes I_n + A^\tau \geq (n-1)\lambda_{\min}(A)I_{mn} + 2A^\tau \circ J,$$

where  $J$  is the  $m \times m$  block matrix with each block  $I_n$ .

**Corollary 3.3** *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then*

$$(\mathrm{tr}_2 A^\tau) \otimes I_n \geq -A^\tau + (n-1)\lambda_{\min}(A)I_{mn}.$$

From Theorem 2.1 and Theorem 3.1, we get the following corollary.

**Corollary 3.4** [11, Theorem 2] *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then*

$$(\mathrm{tr}_2 A^\tau) \otimes I_n \geq \pm A^\tau.$$

**Proof.** We recall that  $J$  denotes the  $m \times m$  block matrix with each block being  $I_n$ . Invoking the positivity of  $J$  and  $A^T$ , we know that the Hadamard product  $A^T \circ J = A^T \circ J$  is positive semidefinite. Thus Theorem 3.1 yields the required inequality

$$(\mathrm{tr}_2 A^\tau) \otimes I_n \geq -A^\tau.$$

On the other hand, replacing  $A$  with  $\tilde{A}$  in Theorem 2.1 leads to

$$I_n \otimes \mathrm{tr}_1 \tilde{A}^\tau \geq \tilde{A}^\tau.$$

Applying the identity (9), we get

$$(\mathrm{tr}_2 (A^\tau)^T) \otimes I_n = \left( \mathrm{tr}_1 \widetilde{(A^\tau)^T} \right) \otimes I_n = (\mathrm{tr}_1 \tilde{A}^\tau) \otimes I_n = I_n \otimes \mathrm{tr}_1 \tilde{A}^\tau \geq \tilde{A}^\tau = (A^\tau)^T.$$

This completes the proof by taking the transpose in both sides. ■

Recall that a positive semidefinite matrix  $A$  is said to be PPT if it has positive partial transpose, i.e., both  $A$  and  $A^\tau$  are positive semidefinite. The following corollary is a direct consequence of Theorem 2.1 and Corollary 3.4. It was first proved by Horodecki et al. [17] for the separability of mixed states in quantum information theory; see [15] for references to the physics literature.

**Corollary 3.5** [17] *If  $A \in \mathbb{M}_m(\mathbb{M}_n)$  is PPT, then*

$$I_m \otimes (\mathrm{tr}_1 A) \geq A \quad \text{and} \quad (\mathrm{tr}_2 A) \otimes I_n \geq A.$$

## 4 Inequalities about two partial traces

In this section, we shall present inequalities involving both the first and second partial traces, which are improvements on Li–Liu–Huang’s result (R5). Recall that a map  $\Phi : \mathbb{M}_n \rightarrow \mathbb{M}_k$  is called positive if it maps positive semidefinite matrices to positive semidefinite matrices. A map  $\Phi : \mathbb{M}_n \rightarrow \mathbb{M}_k$  is said to be  $m$ -positive if for  $[A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$ ,

$$[A_{i,j}]_{i,j=1}^m \geq 0 \quad \Rightarrow \quad [\Phi(A_{i,j})]_{i,j=1}^m \geq 0.$$

We say that  $\Phi$  is *completely positive* if it is  $m$ -positive for every integer  $m \geq 1$ . On the other hand, a map  $\Phi : \mathbb{M}_n \rightarrow \mathbb{M}_k$  is called  $m$ -copositive if

$$[A_{i,j}]_{i,j=1}^m \geq 0 \quad \Rightarrow \quad [\Phi(A_{j,i})]_{i,j=1}^m \geq 0.$$

Similarly, the map  $\Phi$  is *completely copositive* if it is  $m$ -copositive for all  $m \geq 1$ . Furthermore,  $\Phi$  is called completely PPT if the block matrix  $[\Phi(A_{i,j})]_{i,j=1}^m$  is PPT for all  $m \geq 1$  whenever  $[A_{i,j}]_{i,j=1}^m \geq 0$ , i.e.,  $\Phi$  is both completely positive and completely copositive. It is well known



that both the determinant map and the trace map are completely PPT; see [38, p. 221, 237] and see [6, Chapter 3] for more standard results of completely positive maps.

In 2014, Lin [27] proved that  $\Phi(X) = (\text{tr}X)I + X$  is a completely PPT map. Two years later, Lin [30] obtained that  $\Psi(X) = (\text{tr}X)I - X$  is a completely copositive map. In the present section, we first provide an alternative proof of Lin's results, our proof is substantially based on Corollary 3.4, it is of course quite different and concise from [27] and [30]. Moreover, we obtain some new matrix inequalities involving both the first and the second partial trace. Our results are improvements on inequalities (4) and (5).

**Theorem 4.1** [27, 30] *The map  $\Phi(X) = (\text{tr}X)I + X$  is completely PPT and  $\Psi(X) = (\text{tr}X)I - X$  is completely copositive.*

The first part of Theorem 4.1 was proved in [27, Theorem 1.1] by combining a trick of block partition with an inequality of Hadamard product. The second part in Theorem 4.1 was shown in [30, Proposition 2.1] by applying a standard criterion. We shall show that Theorem 4.1 is essentially an equivalent version of Corollary 3.4.

**Proof.** Let  $A = [A_{i,j}]_{i,j=1}^m$  be positive semidefinite. Since the trace map is completely positive, i.e.,  $\text{tr}_2 A = [\text{tr}A_{i,j}]_{i,j=1}^m$  is positive semidefinite (see [38, p. 237]), so is  $[\Phi(A_{i,j})]_{i,j=1}^m = [\text{tr}A_{i,j}]_{i,j=1}^m \otimes I_n + A$ . In other words,  $\Phi(X) = (\text{tr}X)I + X$  is a completely positive map. The remaining proof is to show that  $\Phi$  and  $\Psi$  are completely copositive. We can observe that

$$[\Phi(A_{j,i})]_{i,j=1}^m = [\text{tr}(A_{j,i})I_n + A_{j,i}]_{i,j=1}^m = (\text{tr}_2 A^\tau) \otimes I_n + A^\tau.$$

Correspondingly, we have

$$[\Psi(A_{j,i})]_{i,j=1}^m = [\text{tr}(A_{j,i})I_n - A_{j,i}]_{i,j=1}^m = (\text{tr}_2 A^\tau) \otimes I_n - A^\tau.$$

Corollary 3.4 guarantees that both  $[\Phi(A_{j,i})]_{i,j=1}^m$  and  $[\Psi(A_{j,i})]_{i,j=1}^m$  are positive semidefinite. Thus, the maps  $\Phi$  and  $\Psi$  are completely copositive. ■

**Remark.** We remark here that the map  $\Psi(X) = (\text{tr}X)I - X$  is not completely positive since it is even not 2-positive. For example, we set  $E = [E_{i,j}]_{i,j=1}^2 \in \mathbb{M}_2(\mathbb{M}_n)$ , where  $E_{i,j} \in \mathbb{M}_n$  is the matrix unit, i.e., an  $n$ -square matrix with 1 in the  $(i,j)$ -th entry and 0 elsewhere. Clearly,  $E = [E_{i,j}]_{i,j=1}^2$  is positive semidefinite. However,

$$[\Psi(E_{i,j})]_{i,j=1}^2 = \begin{bmatrix} I_n - E_{1,1} & -E_{1,2} \\ -E_{2,1} & I_n - E_{2,2} \end{bmatrix} \not\geq 0.$$

Therefore,  $\Psi$  is not a completely PPT map.

We now conclude briefly with some interesting applications of Theorem 4.1; see [23] for more details. The first application is to establish trace inequalities for  $2 \times 2$  block matrices.

In 2014, Lin [27, Proposition 2.5] proved the 2-copositivity of  $\Phi(X) = (\text{tr}X)I + X$  and then obtained the following trace inequality (13). More precisely, if  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$  is positive semidefinite, then

$$\text{tr}A\text{tr}C - |\text{tr}B|^2 \geq \text{tr}(AC) - \text{tr}(B^*B). \quad (13)$$

We remark that (13) is a byproduct formulated first in [4, Example 1] in connection with the subadditivity of  $q$ -entropy (the Tsallis entropy) and then also reproved in [40]. Note that the positivity of the block matrix entails  $\text{tr}A\text{tr}C - |\text{tr}B|^2 \geq 0$ ; see [38, p. 237] or [5, Theorem IX.5.10]. However, the right hand side of (13) might be negative. It has been shown that the PPT condition on the block matrix can ensure  $\text{tr}AC \geq \text{tr}B^*B$ ; see [28, Theorem 2.1]. Motivated by this observation, Kittaneh and Lin [19] further generalized (13) to the following (14) by an elegant self-improved technique.

$$\text{tr}A\text{tr}C - |\text{tr}B|^2 \geq \text{tr}(B^*B) - \text{tr}(AC). \quad (14)$$

Moreover they proved by applying the 2-copositivity of  $\Psi(X) = (\text{tr}X)I - X$  that

$$\text{tr}A\text{tr}C + |\text{tr}B|^2 \geq \text{tr}(AC) + \text{tr}(B^*B). \quad (15)$$

It is worth noting that the 2-copositivity of  $\Psi$  could also lead to the previous (14); see [23] for more details.

Another attractive application of Theorem 4.1 states that if  $A \in \mathbb{M}_m(\mathbb{M}_n)$  is positive semidefinite, by using the 2-copositivity of  $\Psi(X) = (\text{tr}X)I - X$ , one can obtain Ando's result (3). On the other hand, by employing the 2-copositivity of  $\Phi(X) = (\text{tr}X)I + X$ , Li, Liu and Huang established inequalities (4) and (5). Last but not least, one may observe an interesting phenomenon that (4) is similar to (13) and (14) in mathematical writing form. Correspondingly, (5) is also similar to (14) and (15).

In what follows, we will illustrate a new application of Theorem 4.1, which together with Theorem 2.2 yields an improvement on inequality (5).

**Theorem 4.2** *Let  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then*

$$(\text{tr}A)I_{mn} + (\text{tr}_2A) \otimes I_n \geq A + I_m \otimes (\text{tr}_1A) + 2(\text{tr}_2D_A) \otimes I_n - 2D_A,$$

where  $D_A = A_{1,1} \oplus A_{2,2} \oplus \cdots \oplus A_{m,m}$ .

**Proof.** The desired inequality can be written as

$$\begin{bmatrix} \sum_{i \neq 1} (\text{tr}A_{i,i})I_n & (\text{tr}A_{1,2})I_n & \cdots & (\text{tr}A_{1,m})I_n \\ (\text{tr}A_{2,1})I_n & \sum_{i \neq 2} (\text{tr}A_{i,i})I_n & \cdots & (\text{tr}A_{2,m})I_n \\ \vdots & \vdots & \ddots & \vdots \\ (\text{tr}A_{m,1})I_n & (\text{tr}A_{m,2})I_n & \cdots & \sum_{i \neq m} (\text{tr}A_{i,i})I_n \end{bmatrix}$$

$$\geq \begin{bmatrix} \sum_{i \neq 1} A_{i,i} & A_{1,2} & \cdots & A_{1,m} \\ A_{2,1} & \sum_{i \neq 2} A_{i,i} & \cdots & A_{2,m} \\ \vdots & \vdots & & \vdots \\ A_{m,1} & A_{m,2} & \cdots & \sum_{i \neq m} A_{i,i} \end{bmatrix}.$$

It is easy to see from Theorem 2.2 that

$$\begin{bmatrix} \sum_{i \neq 1} A_{i,i} & A_{2,1} & \cdots & A_{m,1} \\ A_{1,2} & \sum_{i \neq 2} A_{i,i} & \cdots & A_{m,2} \\ \vdots & \vdots & & \vdots \\ A_{1,m} & A_{2,m} & \cdots & \sum_{i \neq m} A_{i,i} \end{bmatrix} \geq 0.$$

By Theorem 4.1,  $\Psi(X) = (\text{tr}X)I - X$  is completely copositive. Then applying  $\Psi$  to the above positive semidefinite block matrix yields the required inequality. ■

Note that  $(\text{tr}A_{ii})I_n \geq A_{ii}$  for each integer  $i$ , then  $(\text{tr}_2 D_A) \otimes I_n \geq D_A$ , i.e.,

$$2(\text{tr}_2 D_A) \otimes I_n - 2D_A \geq 0.$$

So Theorem 4.2 is indeed a generalization of (5). In view of symmetry of definitions of  $\text{tr}_1$  and  $\text{tr}_2$ , one can easily obtain the following equivalent theorem.

**Theorem 4.3** *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then*

$$(\text{tr}A)I_{mn} - (\text{tr}_2 A) \otimes I_n \geq A - I_m \otimes (\text{tr}_1 A) + 2(I_m \otimes \text{tr}_1 A - A) \circ J,$$

where  $J$  is the  $m \times m$  block matrix with each block  $I_n$ .

The next result is an analogue of Theorem 4.2.

**Theorem 4.4** *Let  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite. Then*

$$(\text{tr}A)I_{mn} + (\text{tr}_2 A) \otimes I_n + I_m \otimes (\text{tr}_1 A) + A \geq 2(\text{tr}_2 D_A) \otimes I_n + 2D_A,$$

where  $D_A = A_{1,1} \oplus A_{2,2} \oplus \cdots \oplus A_{m,m}$ .

**Proof.** The idea of the proof is similar to that in Theorem 4.2. It is equivalent to show

$$\begin{bmatrix} \sum_{i \neq 1} (\text{tr}A_{ii})I_n & (\text{tr}A_{12})I_n & \cdots & (\text{tr}A_{1m})I_n \\ (\text{tr}A_{21})I_n & \sum_{i \neq 2} (\text{tr}A_{ii})I_n & \cdots & (\text{tr}A_{2m})I_n \\ \vdots & \vdots & & \vdots \\ (\text{tr}A_{m1})I_n & (\text{tr}A_{m2})I_n & \cdots & \sum_{i \neq m} (\text{tr}A_{ii})I_n \end{bmatrix}$$

$$\geq - \begin{bmatrix} \sum_{i \neq 1} A_{ii} & A_{12} & \cdots & A_{1m} \\ A_{21} & \sum_{i \neq 2} A_{ii} & \cdots & A_{2m} \\ \vdots & \vdots & & \vdots \\ A_{m1} & A_{m2} & \cdots & \sum_{i \neq m} A_{ii} \end{bmatrix}.$$

This inequality follows from Theorem 2.2 and the completely copositivity of  $\Phi(X) = (\text{tr}X)I + X$  in Theorem 4.1. ■

Recently, Choi [9] proved by replacing  $A$  by  $A - \lambda_{\min}(A)I_{mn}$  and  $\lambda_{\max}(A)I_{mn} - A$  in Ando's result (3) that for any Hermitian  $A \in \mathbb{M}_m(\mathbb{M}_n)$ , we have

$$I_n \otimes (\text{tr}_1 A) + (\text{tr}_2 A) \otimes I_n \geq A + (\text{tr}A)I_{mn} - (m-1)(n-1)\lambda_{\max}(A)I_{mn},$$

and

$$I_n \otimes (\text{tr}_1 A) + (\text{tr}_2 A) \otimes I_n \leq A + (\text{tr}A)I_{mn} - (m-1)(n-1)\lambda_{\min}(A)I_{mn}.$$

Correspondingly, we can get from Li-Liu-Huang's result (4) that

**Proposition 4.5** *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be Hermitian. Then*

$$(\text{tr}A)I_{mn} - (\text{tr}_2 A) \otimes I_n \geq \pm(I_m \otimes (\text{tr}_1 A) - A) + (m-1)(n \mp 1)\lambda_{\min}(A)I_{mn},$$

and

$$(\text{tr}A)I_{mn} - (\text{tr}_2 A) \otimes I_n \leq \pm(I_m \otimes (\text{tr}_1 A) - A) + (m-1)(n \mp 1)\lambda_{\max}(A)I_{mn}.$$

On the other hand, one could obtain from (5) that

**Proposition 4.6** *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be Hermitian. Then*

$$(\text{tr}A)I_{mn} + (\text{tr}_2 A) \otimes I_n \geq I_m \otimes (\text{tr}_1 A) + A + (m+1)(n-1)\lambda_{\min}(A)I_{mn},$$

and

$$(\text{tr}A)I_{mn} + (\text{tr}_2 A) \otimes I_n \leq I_m \otimes (\text{tr}_1 A) + A + (m+1)(n-1)\lambda_{\max}(A)I_{mn}.$$

Motivated by Theorem 4.2 and Theorem 4.3, we believe highly that it is possible to improve Ando's result (3). In other words, for every positive semidefinite matrix  $A \in \mathbb{M}_m(\mathbb{M}_n)$ , is there a positive semidefinite matrix  $T$  satisfying the following inequality?

$$(\text{tr}A)I_{mn} + A \geq I_m \otimes (\text{tr}_1 A) + (\text{tr}_2 A) \otimes I_n + T.$$

## 5 Extensions on Cauchy–Khinchin’s inequality

In this section, we shall provide some applications of Theorem 4.2 and Theorem 4.3 in the field of numerical inequalities. The celebrated Cauchy–Khinchin inequality (see [36, Theorem 1]) states that if  $X = [x_{ij}]$  is an  $m \times n$  real matrix, then

$$\left( \sum_{i=1}^m \sum_{j=1}^n x_{ij} \right)^2 + mn \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 \geq m \sum_{i=1}^m \left( \sum_{j=1}^n x_{ij} \right)^2 + n \sum_{j=1}^n \left( \sum_{i=1}^m x_{ij} \right)^2. \quad (16)$$

It was proved that this inequality has many applications on problems related to the directed graph in combinatorics; see, e.g., [1, 12, 21] for more details. In 2016, Lin [30] provided a simple proof using Ando’s inequality (3). Under the similar lines of the proof, Li, Liu and Huang [24] proved the following generalizations by applying (4) and (5).

**Corollary 5.1** [24] *Let  $X = [x_{ij}]$  be an  $m \times n$  real matrix. Then*

$$mn \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 - n \sum_{j=1}^n \left( \sum_{i=1}^m x_{ij} \right)^2 \geq \left| m \sum_{i=1}^m \left( \sum_{j=1}^n x_{ij} \right)^2 - \left( \sum_{i=1}^m \sum_{j=1}^n x_{ij} \right)^2 \right|,$$

and

$$mn \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 \pm n \sum_{j=1}^n \left( \sum_{i=1}^m x_{ij} \right)^2 \geq \left( \sum_{i=1}^m \sum_{j=1}^n x_{ij} \right)^2 \pm m \sum_{i=1}^m \left( \sum_{j=1}^n x_{ij} \right)^2. \quad (17)$$

The following corollary gives a further improvement on inequalities (17).

**Corollary 5.2** *Let  $X = [x_{ij}]$  be an  $m \times n$  real matrix. Then*

$$(m-2)n \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 + n \sum_{j=1}^n \left( \sum_{i=1}^m x_{ij} \right)^2 \geq \left( \sum_{i=1}^m \sum_{j=1}^n x_{ij} \right)^2 + (m-2) \sum_{i=1}^m \left( \sum_{j=1}^n x_{ij} \right)^2,$$

and

$$m(n-2) \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 + m \sum_{i=1}^m \left( \sum_{j=1}^n x_{ij} \right)^2 \geq \left( \sum_{i=1}^m \sum_{j=1}^n x_{ij} \right)^2 + (n-2) \sum_{j=1}^n \left( \sum_{i=1}^m x_{ij} \right)^2.$$

**Proof.** We only prove the first inequality since the second is similar. Let  $J_n$  be an  $n$ -square matrix with all entries 1. Setting  $A = J_m \otimes J_n$  in Theorem 4.2. Clearly,  $\text{tr}_1 A = mJ_n$  and  $\text{tr}_2 A = nJ_m$ . Additionally, we have  $D_A = J_n \oplus \cdots \oplus J_n$ , which leads to  $(\text{tr}_2 D_A) \otimes I_n = (nI_m) \otimes I_n$ . Therefore,

$$(m-2)nI_{mn} + nJ_m \otimes I_n \geq J_m \otimes J_n + (m-2)I_m \otimes J_n. \quad (18)$$

Let  $\text{vec } X = [x_{11}, \dots, x_{1n}, x_{21}, \dots, x_{2n}, \dots, x_{m1}, \dots, x_{mn}]^T \in \mathbb{R}^{mn}$  be the column vector determined by matrix  $X$ . Then some calculations give the following equalities:

$$\begin{aligned} (\text{vec } X)^T I_{mn} (\text{vec } X) &= \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2, \\ (\text{vec } X)^T (J_m \otimes I_n) (\text{vec } X) &= \sum_{j=1}^n \left( \sum_{i=1}^m x_{ij} \right)^2, \\ (\text{vec } X)^T (J_m \otimes J_n) (\text{vec } X) &= \left( \sum_{i=1}^m \sum_{j=1}^n x_{ij} \right)^2, \\ (\text{vec } X)^T (I_m \otimes J_n) (\text{vec } X) &= \sum_{i=1}^m \left( \sum_{j=1}^n x_{ij} \right)^2. \end{aligned}$$

Thus the desired inequality is equivalent to

$$\begin{aligned} &(\text{vec } X)^T ((m-2)nI_{mn} + nJ_m \otimes I_n) (\text{vec } X) \\ &\geq (\text{vec } X)^T (J_m \otimes J_n + (m-2)I_m \otimes J_n) (\text{vec } X). \end{aligned}$$

This inequality follows immediately from the matrix inequality (18). ■

**Remark.** We remark that inequality (18) can also be proved by a standard argument on eigenvalues by noting that  $J_{mn}, I_{mn}, I_m \otimes J_n$  and  $J_m \otimes I_n$  mutually commute. Moreover, Corollary 5.2 is indeed an improvement on (17) since the Cauchy–Schwarz inequality implies  $2n \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 \geq 2 \sum_{i=1}^m \left( \sum_{j=1}^n x_{ij} \right)^2$ . In addition, setting  $A = J_m \otimes J_n$ , we can see that  $A = A^T$  and so  $A$  is PPT. Unfortunately, our results stated in Sections 2 and 3 (For instance, Theorems 2.1, 2.2 and 3.1) will lead to some trivial numerical inequalities.

## 6 More inequalities for two by two block matrices

In this section, we will investigate the  $2 \times 2$  block positive semidefinite matrices. First of all, we introduce the notion of majorization. For a vector  $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$ , we arrange the coordinates of  $\mathbf{x}$  in non-increasing order  $x_1^\downarrow \geq \dots \geq x_n^\downarrow$  and denote  $\mathbf{x}^\downarrow = (x_1^\downarrow, \dots, x_n^\downarrow)$ . Given  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ , we say that  $\mathbf{x}$  is weakly majorized by  $\mathbf{y}$ , written as  $\mathbf{x} \prec_w \mathbf{y}$ , if

$$\sum_{i=1}^k x_i^\downarrow \leq \sum_{i=1}^k y_i^\downarrow \quad \text{for } k = 1, 2, \dots, n.$$

We say that  $\mathbf{x}$  is majorized by  $\mathbf{y}$ , denoted by  $\mathbf{x} \prec \mathbf{y}$ , if  $\mathbf{x} \prec_w \mathbf{y}$  and the sum of all entries of  $\mathbf{x}$  equal to the sum of all entries of  $\mathbf{y}$ . The majorization theory has become a rich research field with far-reaching applications to a wide number of areas, we refer to the recent monograph [33] or [38, Chapter 10] for comprehensive surveys on this subject.

Historically, the first example of majorization arising in matrix theory is usually attributed to Issai Schur, which states that the diagonal elements of a Hermitian matrix  $H$  are majorized by its eigenvalues, i.e.,  $\mathbf{d}(H) \prec \boldsymbol{\lambda}(H)$ . This majorization provided a new and profound understanding on Hadamard's determinant inequality; see, e.g., [16, p. 514]. Due to Schur's discovery, a large number of other majorization inequalities have been found in the context of matrix analysis. For instance, if  $A = [A_{i,j}]_{i,j=1}^m \in \mathbb{M}_m(\mathbb{M}_n)$  is Hermitian, then Schur's inequality implies  $\mathbf{d}(H) \prec \boldsymbol{\lambda}(A_{1,1} \oplus \cdots \oplus A_{m,m})$ . Furthermore, if  $A$  is positive semidefinite, it is well-known (see [16, p. 259] and [33, p. 308]) that

$$\boldsymbol{\lambda}(A_{1,1} \oplus \cdots \oplus A_{m,m}) \prec \boldsymbol{\lambda}(A) \prec \boldsymbol{\lambda}(A_{1,1}) + \cdots + \boldsymbol{\lambda}(A_{m,m}). \quad (19)$$

Moreover, Rotfeld and Thompson [33, p. 330] obtained an analogous complement,

$$\boldsymbol{\lambda}(A_{1,1} \oplus \cdots \oplus A_{m,m}) \prec \boldsymbol{\lambda}(\text{tr}_1 A) \prec \boldsymbol{\lambda}(A_{1,1}) + \cdots + \boldsymbol{\lambda}(A_{m,m}). \quad (20)$$

In the sequel, we shall provide a comparison between (19) and (20) under the PPT condition. First of all, we recall the following lemma, which was proved by Hiroshima [15] in the language of quantum information theory; see [29] for an alternative proof.

**Lemma 6.1** [15, 29] *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be positive semidefinite.*

- (1) *If  $I_m \otimes \text{tr}_1 A \geq A$ , then  $\boldsymbol{\lambda}(A) \prec \boldsymbol{\lambda}(\text{tr}_1 A)$ .*
- (2) *If  $(\text{tr}_2 A) \otimes I_n \geq A$ , then  $\boldsymbol{\lambda}(A) \prec \boldsymbol{\lambda}(\text{tr}_2 A)$ .*

The first theorem in this section is the following majorization inequalities.

**Theorem 6.2** *Let  $A \in \mathbb{M}_m(\mathbb{M}_n)$  be PPT. Then*

$$\boldsymbol{\lambda}(A) \prec \boldsymbol{\lambda}(\text{tr}_1 A) \quad \text{and} \quad \boldsymbol{\lambda}(A) \prec \boldsymbol{\lambda}(\text{tr}_2 A).$$

*Similarly, we have*

$$\boldsymbol{\lambda}(A^T) \prec \boldsymbol{\lambda}(\text{tr}_1 A) \quad \text{and} \quad \boldsymbol{\lambda}(A^T) \prec \boldsymbol{\lambda}(\text{tr}_2 A).$$

**Proof.** This is a direct consequence of Corollary 3.5 and Lemma 6.1. ■

In particular, for the case of  $2 \times 2$  block matrices, we remark here that inequalities involving the first partial trace in Theorem 6.2 was partially proved by Bourin, Lee and Lin [7, 8] by making use of a simple but useful decomposition lemma for positive semidefinite matrices. Moreover, other special cases can be found in [26, 35].

**Corollary 6.3** *Suppose that  $H \in \mathbb{M}_2(\mathbb{M}_n)$  is a block matrix partitioned as*

$$H = \begin{bmatrix} M & K \\ K^* & N \end{bmatrix}.$$

(1) [7, 26] If  $K$  is Hermitian, then  $\lambda(H) \prec \lambda(M + N)$ .

(2) [35, Corollary 5] If  $K$  is skew-Hermitian, then  $\lambda(H) \prec \lambda(M + N)$ .

Over the past few years,  $2 \times 2$  block positive partial transpose matrices play an important role in matrix analysis and quantum information, such as the separability of mixed states and the subadditivity of  $q$ -entropies; see [4, 28] for related topics and references to the physics literature. It is extremely meaningful and significant to finding  $2 \times 2$  block PPT matrices. To the author's best knowledge, the most famous example is commonly regarded as the Hua matrix; see [18, 2, 37, 28, 32] for more details.

In 2014, Lin [27] proved that if  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{M}_2(\mathbb{M}_n)$  is positive semidefinite, then

$$\begin{bmatrix} (\operatorname{tr}A)I + A & (\operatorname{tr}B)I + B \\ (\operatorname{tr}B^*)I + B^* & (\operatorname{tr}C)I + C \end{bmatrix}$$

is PPT. In 2017, Choi [9] also showed an extremely similar result, which states that

$$\begin{bmatrix} (\operatorname{tr}A)I + C & (\operatorname{tr}B)I - B \\ (\operatorname{tr}B^*)I - B^* & (\operatorname{tr}C)I + A \end{bmatrix} \quad (21)$$

is positive semidefinite. In the next theorem, we shall prove that the  $2 \times 2$  block matrix in (21) is further PPT.

**Theorem 6.4** *Let  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{M}_2(\mathbb{M}_n)$  be positive semidefinite. Then*

$$\begin{bmatrix} (\operatorname{tr}A)I + C & (\operatorname{tr}B)I - B \\ (\operatorname{tr}B^*)I - B^* & (\operatorname{tr}C)I + A \end{bmatrix}$$

*is PPT.*

**Proof.** In view of (21), we only need to prove

$$\begin{bmatrix} (\operatorname{tr}A)I + C & (\operatorname{tr}B^*)I - B^* \\ (\operatorname{tr}B)I - B & (\operatorname{tr}C)I + A \end{bmatrix} \geq 0.$$

Note that

$$\begin{bmatrix} C & -B^* \\ -B & C \end{bmatrix} = \begin{bmatrix} 0 & -I \\ I & 0 \end{bmatrix} \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} \geq 0.$$

It suffices to show

$$\begin{bmatrix} (\operatorname{tr}A)I & (\operatorname{tr}B^*)I \\ (\operatorname{tr}B)I & (\operatorname{tr}C)I \end{bmatrix} \geq 0,$$



which follows from the complete positivity of trace map. ■

A norm  $\|\cdot\|$  on  $\mathbb{M}_n$  is called unitarily invariant if  $\|UAV\| = \|A\|$  for any  $A \in \mathbb{M}_n$  and any unitary matrices  $U, V \in \mathbb{M}_n$ . The unitarily invariant norm of a matrix is closely related to its singular values; see, e.g., [38, pp. 372–376] and [5, pp. 91–98].

**Corollary 6.5** *Let  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{M}_2(\mathbb{M}_n)$  be positive semidefinite. Then*

$$2\|(\operatorname{tr}B)I \pm B\| \leq \|(\operatorname{tr}(A+C))I + A + C\|$$

for any unitarily invariant norm.

**Proof.** Invoking a theorem of von Neumann (see [38, p. 375]), it is sufficient to prove

$$2\mathbf{s}((\operatorname{tr}B)I \pm B) \prec_w \mathbf{s}((\operatorname{tr}(A+C))I + A + C),$$

where  $\mathbf{s}(X)$  is the vector consisting of singular values of  $X$ . There is a well-known fact that if  $X = \begin{bmatrix} Y & Z \\ Z^* & W \end{bmatrix}$  is positive semidefinite, then  $2s_i(Z) \leq s_i(X)$  for every  $i = 1, 2, \dots, n$ . With the help of this fact, it is straightforward to get the desired inequality by combining Theorem 6.4 and Theorem 6.2. ■

The second partial trace inequality in Theorem 6.2 yields an elegant corollary.

**Corollary 6.6** *Let  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{M}_2(\mathbb{M}_n)$  be positive semidefinite. Then*

$$\|(\operatorname{tr}B)I + B\| \leq \frac{n+1}{2} \left\| \begin{bmatrix} \operatorname{tr}A & \operatorname{tr}B \\ \operatorname{tr}B^* & \operatorname{tr}C \end{bmatrix} \right\|$$

for any unitarily invariant norm.

At the end of this paper, we conclude with the following inequality involving singular values, which is a stronger inequality than Corollary 6.5. Theorem 6.7 was partially proved in [31]. To some extent, our proof grows out of that of Lin with some simplifications.

**Theorem 6.7** *Let  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$  be positive semidefinite. Then*

$$2s_j((\operatorname{tr}B)I \pm B) \leq s_j((\operatorname{tr}(A+C))I + A + C).$$

To proceed the proof, we need to present the following two lemmas.

**Lemma 6.8** [5, p. 262] *For any  $M, N \in \mathbb{M}_{n \times m}$  and  $j = 1, 2, \dots, n$ , we have*

$$2s_j(MN^*) \leq s_j(M^*M + N^*N).$$

**Lemma 6.9** For any  $M, N \in \mathbb{M}_{n \times m}$  and  $j = 1, 2, \dots, n$ , we have

$$\begin{aligned} \lambda_j(M^*M + N^*N) &\leq \lambda_j(MM^* + NN^*) \\ &\quad + \frac{1}{2}\text{tr}(M^*M + N^*N - M^*N - N^*M). \end{aligned}$$

**Proof.** By a direct computation, we can get

$$\begin{aligned} 2\lambda_j(MM^* + NN^*) &= \lambda_j((M + N)(M + N)^* + (M - N)(M - N)^*) \\ &\geq \lambda_j((M + N)(M + N)^*) + \lambda_n((M - N)(M - N)^*) \\ &\geq \lambda_j((M + N)^*(M + N)) \\ &= \lambda_j(2(M^*M + N^*N) - (M - N)^*(M - N)) \\ &\geq 2\lambda_j((M^*M + N^*N)) - \lambda_1((M - N)^*(M - N)), \end{aligned}$$

where the first and last inequality hold by Weyl's eigenvalue inequality (see [5, p. 63]). Moreover, the positivity of  $(M - N)^*(M - N)$  leads to

$$\begin{aligned} \lambda_1((M - N)^*(M - N)) &\leq \text{tr}((M - N)^*(M - N)) \\ &= \text{tr}(M^*M + N^*N - M^*N - N^*M). \end{aligned}$$

This completes the proof. ■

**Proof of Theorem 6.7** Since  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$  is positive semidefinite, we may write

$$\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} = \begin{bmatrix} MM^* & MN^* \\ NM^* & NN^* \end{bmatrix}$$

for some  $M, N \in \mathbb{M}_{n \times 2n}$ . Then the desired inequality is the same as

$$2s_j(\text{tr}(MN^*)I \pm MN^*) \leq s_j(\text{tr}(MM^* + NN^*)I + MM^* + NN^*).$$

By the Weyl's inequality of singular value and Lemma 6.8, we have

$$\begin{aligned} 2s_j(\text{tr}(MN^*)I \pm MN^*) &\leq 2s_j(\pm MN^*) + 2s_1((\text{tr}MN^*)I) \\ &= 2s_j(MN^*) + 2|\text{tr}(MN^*)| \\ &\leq s_j(M^*M + N^*N) + 2|\text{tr}(MN^*)|, \end{aligned}$$

On the other hand, we observe that

$$s_j(\text{tr}(MM^* + NN^*)I + MM^* + NN^*) = \lambda_j(MM^* + NN^*) + \text{tr}(MM^* + NN^*).$$

It is sufficient to prove that for every  $M, N \in \mathbb{M}_{n \times m}$ ,

$$s_j(M^*M + N^*N) + 2|\text{tr}(MN^*)| \leq \lambda_j(MM^* + NN^*) + \text{tr}(MM^* + NN^*). \quad (22)$$

We may assume without loss of generality that  $\operatorname{tr}(MN^*) \geq 0$  in (22), since it is clear that (22) is invariant when we replace  $M$  with  $e^{i\theta}M$  for every  $\theta \in [0, 2\pi]$ . Therefore,

$$\begin{aligned} & s_j(M^*M + N^*N) + |\operatorname{tr}(MN^*)| \\ = & \lambda_j(M^*M + N^*N) + \frac{1}{2}\operatorname{tr}(M^*N + N^*M) \\ \leq & \lambda_j(MM^* + NN^*) + \frac{1}{2}\operatorname{tr}(M^*M + N^*N), \end{aligned} \quad (23)$$

where the last inequality holds by Lemma 6.9. Note that

$$\begin{bmatrix} \operatorname{tr}MM^* & \operatorname{tr}MN^* \\ \operatorname{tr}NM^* & \operatorname{tr}NN^* \end{bmatrix}$$

is a positive semidefinite matrix, we have

$$|\operatorname{tr}(MN^*)| = \frac{1}{2}\operatorname{tr}(MN^* + NM^*) \leq \frac{1}{2}\operatorname{tr}(MM^* + NN^*). \quad (24)$$

The desired inequality (22) now follows by combining (23) and (24). ■

**Corollary 6.10** *For any  $X \in \mathbb{M}_n$ , we have*

$$2s_j((\operatorname{tr}X)I \pm X) \leq s_j((\operatorname{tr}(|X| + |X^*|))I + |X| + |X^*|),$$

where  $|X| = (X^*X)^{1/2}$ .

**Proof.** Observe that for any square matrix  $X$ , the following block matrix

$$\begin{bmatrix} |X| & X^* \\ X & |X^*| \end{bmatrix}$$

is positive semidefinite by applying the singular value decomposition or polar decomposition; see, e.g., [38, pp. 82–83]. Hence, the required inequality holds by Theorem 6.7. ■

## Acknowledgments

This paper is dedicated to Prof. Weijun Liu (Central South University) on his 60th birthday, October 22 of the lunar calendar in 2021. The author would like to thank Prof. Minghua Lin for bringing the topic on partial traces to his attention. Thanks also go to Prof. Fuzhen Zhang and Prof. Yuejian Peng for valuable comments which improve the presentation. The author is also grateful to Prof. Tsuyoshi Ando for the correspondence [3] before its publication, which initiates our research. This work was supported by NSFC (Grant No. 11931002).

## References

- [1] R. Ahlswede, G.O.H. Katona, Graphs with maximal number of adjacent pairs of edges, *Acta Math. Acad. Sci. Hungar.* 32(1978) 97–120.
- [2] T. Ando, Positivity of operator-matrices of Hua-type, *Banach J. Math. Anal.* 2 (2008) 1–8.
- [3] T. Ando, Matrix inequalities involving partial traces, ILAS Conference, 2014.
- [4] A. Desenyei, D. Petz, Partial subadditivity of entropies, *Linear Algebra Appl.* 439 (2013) 3297–3305.
- [5] R. Bhatia, *Matrix Analysis*, GTM, vol. 169, Springer-Verlag, New York, 1997.
- [6] R. Bhatia, *Positive Definite Matrices*, Princeton University Press, Princeton, 2007.
- [7] J.-C. Bourin, E.-Y. Lee, M. Lin, On a decomposition lemma for positive semi-definite block-matrices, *Linear Algebra Appl.* 437 (2012) 1906–1912.
- [8] J.-C. Bourin, E.-Y. Lee, M. Lin, Positive matrices partitioned into a small number of Hermitian blocks, *Linear Algebra Appl.* 438 (2013) 2591–2598.
- [9] D. Choi, Inequalities related to partial transpose and partial trace, *Linear Algebra Appl.* 516 (2017) 1–7.
- [10] D. Choi, Inequalities related to trace and determinant of positive semidefinite block matrices, *Linear Algebra Appl.* 532 (2017) 1–7.
- [11] D. Choi, Inequalities about partial transpose and partial traces, *Linear Multilinear Algebra* 66 (2018) 1619–1625.
- [12] D. de Caen, An upper bound on the sum of squares of degrees in a graph, *Discrete Math.* 185 (1998) 245–248.
- [13] X. Fu, P.-S. Lau, T.-Y. Tam, Linear maps of positive partial transpose matrices and singular value inequalities, *Math. Inequal. Appl.* 23 (4) (2020) 1459–1468.
- [14] X. Fu, P.-S. Lau, T.-Y. Tam, Inequalities on  $2 \times 2$  block positive semidefinite matrices, *Linear Multilinear Algebra* (2021), <https://doi.org/10.1080/03081087.2021.1969327>.
- [15] T. Hiroshima, Majorization criterion for distillability of a bipartite quantum state, *Phys. Rev. Lett.* 91 (2003), 057902.
- [16] R.A. Horn, C.R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.

- [17] M. Horodecki, P. Horodecki and R. Horodecki, Separability of mixed states: Necessary and sufficient conditions, *Phys. Lett. A* 223 (1996) 1–8.
- [18] L.-K. Hua, Inequalities involving determinants (in Chinese), *Acta Math. Sin.* 5 (1955) 463–470. See also *Trans. Am. Math. Soc. Ser. II* 32 (1963) 265–272.
- [19] F. Kittaneh, M. Lin, Trace inequalities for positive semidefinite block matrices, *Linear Algebra Appl.* 524 (2017) 153–158.
- [20] E.-Y. Lee, The off-diagonal block of a PPT matrix, *Linear Algebra Appl.* 486 (2015) 449–453.
- [21] J.-S. Li, Y.-L. Pan, de Caen’s inequality and bounds on the largest Laplacian eigenvalue of a graph, *Linear Algebra Appl.* 328 (2001) 153–160.
- [22] Y. Li, L. Feng, Z. Huang, W. Liu, Inequalities regarding partial trace and partial determinant, *Math. Inequal. Appl.* 23 (2020) 477–485.
- [23] Y. Li, Y. Huang, L. Feng, W. Liu, Some applications of two completely copositive maps, *Linear Algebra Appl.* 590 (2020) 124–132.
- [24] Y. Li, W. Liu, Y. Huang, A new matrix inequality involving partial traces, *Operators and Matrices* 15 (2021), no. 3, 1189–1199.
- [25] Y. Li, Extensions of some matrix inequalities related to trace and partial traces, 19 pages, arXiv:2003.04520v2.
- [26] M. Lin, H. Wolkowicz, An eigenvalue majorization inequality for positive semidefinite block matrices, *Linear Multilinear Algebra* 60 (2012) 1365–1368.
- [27] M. Lin, A completely PPT map, *Linear Algebra Appl.* 459 (2014) 404–410.
- [28] M. Lin, Inequalities related to  $2 \times 2$  block PPT matrices, *Operators and Matrices* 9 (2015), no. 4, 917–924.
- [29] M. Lin, Some applications of a majorization inequality due to to Bapat and Sunder, *Linear Algebra Appl.* 469 (2015) 510–517.
- [30] M. Lin, A determinantal inequality involving partial traces, *Canad. Math. Bull.* 59 (2016) 585–591.
- [31] M. Lin, A singular value inequality related to a linear map, *Electronic J. Linear Algebra* 31 (2016) 120–124.
- [32] M. Lin, The Hua matrix and inequalities related to contractive matrices, *Linear Algebra Appl.* 511 (2016) 22–30.

- [33] A.W. Marshall, I. Olkin, B.C. Arnold, *Inequalities: Theory of Majorization and Its Applications*, 2nd ed., Springer, New York, 2011.
- [34] D. Petz, *Quantum Information Theory and Quantum Statistics. Theoretical and Mathematical Physics*, Springer, Berlin, 2008.
- [35] R. Turkmen, V.E. Paksoy, F. Zhang, Some inequalities of majorization type, *Linear Algebra Appl.* 437 (2012) 1305–1316.
- [36] E. R. van Dam, A Cauchy–Khinchin matrix inequality, *Linear Algebra Appl.* 280 (1998) 163–172.
- [37] G. Xu, C. Xu, F. Zhang, Contractive matrices of Hua type, *Linear Multilinear Algebra* 59 (2011) 159–172.
- [38] F. Zhang, *Matrix Theory: Basic Results and Techniques*, 2nd edition, Springer, New York, 2011.
- [39] F. Zhang, Positivity of matrices with generalized matrix functions. *Acta Math. Sin. (Engl. Ser.)* 28 (2012) 1779–1786.
- [40] F. Zhang, Trace inequality for positive block matrices, *IMAGE Solution* 50-3.1, *Bull. Internat. Linear Algebra Soc.* (Fall 2013).
- [41] P. Zhang, On some inequalities related to positive block matrices, *Linear Algebra Appl.* 576 (2019) 258–267.