

# EXPONENTIAL INTEGRATORS FOR SECOND-ORDER IN TIME PARTIAL DIFFERENTIAL EQUATIONS

ALEXANDER OSTERMANN AND DUY PHAN

ABSTRACT. Two types of second-order in time partial differential equations (PDEs), namely semilinear wave equations and semilinear beam equations are considered. To solve these equations with exponential integrators, we present an approach to compute efficiently the action of the matrix exponential as well as those of related matrix functions. Various numerical simulations are presented that illustrate this approach.

## 1. INTRODUCTION

We consider semilinear damped wave equations with damping term, structural (visco-elastic) damping term, and mass term

$$\partial_{tt}^2 u - \alpha \Delta u - \beta \Delta(\partial_t u) + \gamma \partial_t u + \delta u = g(u) + h(\partial_t u), \quad (x, t) \in \Omega \times (0, +\infty), \quad (1a)$$

$$u|_{\partial\Omega} = 0, \quad (x, t) \in \partial\Omega \times (0, +\infty), \quad (1b)$$

$$u(0, x) = p(x), \quad \partial_t u(0, x) = q(x), \quad x \in \Omega \quad (1c)$$

on a bounded and open domain  $\Omega \subset \mathbb{R}^d$  with smooth compact boundary  $\partial\Omega$ . The term  $\Delta(\partial_t u)$  is the structural (visco-elastic) damping while the term  $\partial_t u$  is the damping term. We assume that  $\beta, \gamma$ , and  $\delta$  are three non-negative coefficients. Moreover, the diffusion coefficient  $\alpha$  must be positive. The initial data  $p$  and  $q$  are chosen from the usual energy space  $(p, q) \in H_0^1(\Omega) \times L^2(\Omega)$ . Concerning the nonlinear term, we recall some particular equations from the literature:

- (i) the perturbed sine-Gordon equation (see [PZ06, GM91, CCD08])

$$\partial_{tt}^2 u - \alpha \Delta u - \beta \Delta(\partial_t u) + \gamma \partial_t u = \sin(u) + k(x); \quad (2)$$

- (ii) the perturbed wave equation of quantum mechanics (see [PZ06, GM91, CCD08])

$$\partial_{tt}^2 u - \alpha \Delta u - \beta \Delta(\partial_t u) = |u|^q u + |\partial_t u|^p (\partial_t u), \quad p, q \geq 0. \quad (3)$$

Another type of second-order in time PDE is the Euler–Bernoulli beam equation with Kelvin–Voigt damping

$$\partial_{tt}^2 u + \partial_{xx}^2 (\alpha \partial_{xx}^2 u + \beta \partial_{xx}^2 (\partial_t u)) + \gamma \partial_t u + \delta u = g(u), \quad (x, t) \in (0, L) \times (0, +\infty), \quad (4a)$$

$$u(0, x) = p(x), \quad \partial_t u(0, x) = q(x), \quad x \in (0, L), \quad (4b)$$

*Date:* Tuesday 30<sup>th</sup> November, 2021.

*Key words and phrases.* semilinear wave equations, semilinear beam equations, exponential integrators, computation of matrix functions.

where  $u(t, x)$  denotes the deflection of the beam of its rigid body motion at time  $t$  and position  $x$ . For positive physical parameter  $\alpha > 0$  and  $\beta \geq 0$ , the moment function is

$$m(t, x) = \alpha \partial_{xx}^2 u(t, x) + \beta \partial_{xx}^2 (\partial_t u)(t, x).$$

The first derivative of the moment  $m(t, x)$  with respect to the variable  $x$  represents the shear force. The following boundary conditions will be considered where  $\xi \in \{0, L\}$ :

- (a) Hinged end:  $u(t, \xi) = 0, m(t, \xi) = 0$ .
- (b) Clamped end:  $u(t, \xi) = 0, \partial_x u(t, \xi) = 0$ .
- (c) Free end:  $m(t, \xi) = 0, \partial_x m(t, \xi) = 0$ .
- (d) Sliding end:  $\partial_x u(t, \xi) = 0, m(t, \xi) = 0$ .

Depending on the set up of the beam model, various combinations of boundary conditions are of interest, for example: hinged-hinged boundary conditions

$$u(t, 0) = 0, m(t, 0) = 0, u(t, L) = 0, m(t, L) = 0.$$

Concerning semilinear beam equations, in [EM19a, EM19b, AEY11], a nonlinear term  $g(u) = -ku^3$ ,  $k \in \mathbb{R}$  was used when the authors considered a railway track model.

Both problems (1) and (4) can be rewritten as abstract ordinary differential equations in a product space  $X = H \times L^2(\Omega)$  by denoting a new variable  $y(t, x) = (u(t, x), \partial_t u(t, x))'$  as follows

$$\dot{y}(t) = \mathcal{A}y(t) + \mathcal{F}(y(t)),$$

where

$$\mathcal{A} = \begin{bmatrix} 0 & I \\ -\alpha(-\Delta) - \delta I & -\beta(-\Delta) - \gamma I \end{bmatrix} \quad \text{for (1),}$$

$$\mathcal{A} = \begin{bmatrix} 0 & I \\ -\alpha \partial_{xxxx}^4 - \delta I & -\beta \partial_{xxxx}^4 - \gamma I \end{bmatrix} \quad \text{for (4),}$$

and

$$\mathcal{F}(w_1, w_2) = \begin{bmatrix} 0 \\ g(w_1) + h(w_2) \end{bmatrix}.$$

The space  $H$  and the domain of the operator  $\mathcal{A}$  will be chosen to be consistent with the boundary conditions. Here and henceforth, the transpose of a matrix  $E$  is denoted by  $E'$ .

These types of equations have been studied extensively in many fields of mathematics. For damped wave equations, see [PZ06, GM91, CCD08, CF21, CCD08, PZ06, Pon85, GM91, ITY13]; for Euler–Bernoulli beam equations, see [MS15, Mor20, BI97, EM19a, EM19b, PP20, PP21, IM98, LL98, AEY11] and references therein. The time discretization of these equations, to the best of our knowledge, is usually carried out by standard integration schemes such as Runge–Kutta methods or multistep methods. In this article, we will consider exponential integrators to solve this class of PDEs. By spatial

discretization of (1) or of (4), we get a semi-discretization of the equation in matrix form

$$\dot{y}(t) = Ay(t) + F(y(t)), \quad y(0) = y_0 = (p, q)', \quad (5)$$

where

$$A = \begin{bmatrix} 0 & I \\ -\alpha S - \delta I & -\beta S - \gamma I \end{bmatrix}. \quad (6)$$

and  $S \in \mathbb{R}^{k \times k}$  is the discretized version of the operator  $(-\Delta)$  or  $\partial_{xxxx}^4$ . The linear part of (17)

$$\dot{y}(t) = Ay(t), \quad y(0) = y_0 = (p, q)', \quad (7)$$

can be solved exactly, e.g.,

$$y(t) = e^{tA}y_0, \quad t > 0 \quad (8)$$

For the undamped wave equations (i.e.  $\beta = \gamma = 0$  in (6)), an explicit form of the matrix exponential  $e^{tA}$  by using the matrix sine and matrix cosine functions is well-known (see [HO10, Section 3.2]). Based on this formula, Gautschi (in [Gau61]) and Deuffhard (in [Deu79]) developed a number of schemes to tackle semilinear second-order differential equations. Nevertheless, when damping terms appear in (6), a direct approach to compute the matrix exponential  $e^{tA}$  is more involved and not yet discussed in the literature. Therefore, in this paper, we firstly present an approach to exactly evaluate the matrix exponential of (6).

Let us briefly explain our procedure to compute the matrix exponential. We start by employing two linear transformations to represent the matrix  $A$  as  $A = \tilde{Q}PCP'\tilde{Q}'$  where the new matrix  $C$  is a block diagonal matrix, i.e.  $C = \text{diag}(G_1, \dots, G_n)$ . Each block  $G_i$  is a  $2 \times 2$  matrix. The exponential of this matrix  $G_i$  will be computed explicitly. Regarding its eigenvalues, a suitable formula will be constructed. In this way the matrix exponential can be computed cheaply even for large time. We also want to discuss the cases  $\beta = \gamma = \delta = 0$  and  $\beta, \gamma \ll \alpha$  (see [IM98, Section 3] for typical physical parameters). In the second case, solving the equation using finite differences in time (for example: the Crank–Nicolson scheme) is more expensive. In both cases, the matrix  $G_i$  has usually two conjugate complex eigenvalues. To reduce the computation cost, we avoid complex arithmetic. The exact matrix exponential will not only give us a huge advantage to solve the class of linear damped wave equations or linear beam equations but also be valuable in computing solutions of semilinear problems. The numerical schemes for the full equation (17) were constructed by incorporating the exact solution of (7) in an appropriate way. In the literature, these methods were investigated by many authors (see, e.g., [DOT17, HO10, Kro05, SW92, CM02, LO14, HO05a, HO05b]). To employ these known exponential integrators, the core point is the computation of related matrix functions  $\varphi_k(tA)$ . Analogously, we will use two linear transformations and compute the action of the matrix functions  $\varphi_k(tG_i)$ . Explicit formulas will be established in the same way as for computing the matrix exponential  $e^{tG_i}$ . Concerning the computation of matrix functions, we refer to the review work by Higham and Al-Mohy [HAM10] as well as the monograph by Higham [Hig08].

The outline of the paper is as follows. We start with the discussion of computing the matrix exponential  $e^{tA}$  in section 2. Two linear transformations  $P$  and  $Q$  will be presented. The computations of the matrix exponential  $e^{tG_i}$  will be discussed in three different cases. In simulations, instead of computing the matrix exponential, we will rather compute its action to an arbitrary vector. A detailed instruction will be presented in remark 2.7. In section 3 we recall some exponential integrators and discuss an approach to compute the related matrix functions  $\varphi_k(tA)$ . The procedure will be summarized in section 3.3. In section 4, we will present some numerical examples of semi-linear equations. The operators  $(-\Delta)$  and  $\partial_{xxxx}^4$  will be discretized by finite differences. We will use exponential integrators for the time integration of these examples.

## 2. EXACT MATRIX EXPONENTIAL

In this section, we propose an approach to compute efficiently the matrix exponential  $e^{tA}$  for a matrix  $A$  of the form (6). With this at hand, the solution of linear system (7) can be evaluated for arbitrary time  $t > 0$  in a fast and reliable way.

**2.1. Two linear transformations.** The key idea is to transform  $A$  to a simple block-diagonal matrix for which the exponential can be computed cheaply.

**Lemma 2.1.** *Assume that there exist an orthogonal matrix  $Q$  and a diagonal matrix  $D = \text{diag}\{\lambda_1, \dots, \lambda_n\}$  such that  $S = QDQ'$ , then the matrix  $A$  can be transformed to the block form*

$$B = \begin{bmatrix} 0 & I \\ D_1 & D_2 \end{bmatrix},$$

where  $D_1$  and  $D_2$  are two diagonal matrices.

*Proof.* By substituting  $S = QDQ'$  and  $QQ' = I$  into (6), we get that

$$\begin{aligned} A &= \begin{bmatrix} 0 & QQ' \\ -\alpha QDQ' - \delta QQ' & -\beta QDQ' - \gamma QQ' \end{bmatrix} \\ &= \begin{bmatrix} Q & 0 \\ 0 & Q \end{bmatrix} \begin{bmatrix} 0 & I \\ -\alpha D - \delta I & -\beta D - \gamma I \end{bmatrix} \begin{bmatrix} Q' & 0 \\ 0 & Q' \end{bmatrix}. \end{aligned}$$

The proof is complete by identifying two diagonal matrices  $D_1 = -\alpha D - \delta I$  and  $D_2 = -\beta D - \gamma I$ .  $\square$

**Lemma 2.2.** *Consider  $P \in \mathbb{R}^{2n \times 2n}$  the permutation matrix satisfying*

$$P_{i,2i-1} = 1, \quad P_{i+n,2i} = 1 \quad \text{for } 1 \leq i \leq n, \quad P_{k,l} = 0 \text{ else.} \quad (9)$$

*The matrix  $B$  can be transformed under the permutation  $P$  to a block diagonal matrix  $C$ , i.e.  $B = PCP'$ , where*

$$C = \begin{bmatrix} G_1 & 0 & \cdots & 0 \\ 0 & G_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \cdots & G_n \end{bmatrix} \quad \text{with} \quad G_i = \begin{bmatrix} 0 & 1 \\ -\alpha\lambda_i - \delta & -\beta\lambda_i - \gamma \end{bmatrix}.$$

*Proof.* Following the definitions of the matrices  $B$  and  $C$ , for  $1 \leq i \leq n$  we have

$$\begin{aligned} B_{i,n+i} &= 1, & B_{n+i,i} &= -\alpha\lambda_i, & B_{n+i,n+i} &= -\beta\lambda_i - \gamma, \\ C_{2i-1,2i} &= 1, & C_{2i,2i-1} &= -\alpha\lambda_i, & C_{2i,2i} &= -\beta\lambda_i - \gamma. \end{aligned}$$

We will prove that  $B = PCP'$ . Indeed, for  $1 \leq i \leq n$ , we have

$$\begin{aligned} (PCP')_{n+i,n+i} &= P_{n+i,2i}(CP')_{2i,n+i} = (CP')_{2i,n+i} = C_{2i,2i}P'_{2i,n+i} \\ &= C_{2i,2i}P_{i+n,2i} = C_{2i,2i} = -\beta\lambda_i - \gamma = B_{n+i,n+i}, \\ (PCP')_{i,n+i} &= P_{i,2i-1}(CP')_{2i-1,n+i} = (CP')_{2i-1,n+i} = C_{2i-1,2i}P'_{2i,n+i} \\ &= C_{2i-1,2i} = 1 = B_{i,n+i}, \\ (PCP')_{n+i,i} &= P_{n+i,2i}(CP')_{2i,i} = (CP')_{2i,i} = C_{2i,2i-1}P'_{2i-1,i} \\ &= C_{2i,2i-1} = -\alpha\lambda_i - \delta = B_{n+i,i}. \end{aligned}$$

We will not be concerned with the remaining elements of  $B$  and  $C$  since they are all zero. Thus, the proof is complete.  $\square$

**Example 2.3.** For  $n = 2$  and  $n = 3$  the permutation matrices  $P$  have the following form

$$P_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad P_3 = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Next, we recall some important properties of matrix functions (see [Hig08, Theorem 1.13] or [HAM10, Theorem 2.3]).

**Theorem 2.4.** Let  $A \in \mathbb{C}^{k \times k}$  and  $f$  be defined on the spectrum of  $A$ . Then

- (a)  $f(A)$  commutes with  $A$ ;
- (b)  $f(A') = f(A)'$ ;
- (c)  $f(XAX^{-1}) = Xf(A)X^{-1}$ .
- (d) The eigenvalues of  $f(A)$  are  $f(\lambda_i)$ , where  $\lambda_i$  are the eigenvalues of  $A$ .
- (e) If  $A = (A_{ij})$  is block triangular then  $F = f(A)$  is block triangular with the same block structure as  $A$ , and  $F_{ii} = f(A_{ii})$ .
- (f) If  $A = \text{diag}(A_{11}, A_{22}, \dots, A_{mm})$  is block diagonal then

$$f(A) = \text{diag}(f(A_{11}), f(A_{22}), \dots, f(A_{mm})).$$

A direct consequence of this theorem is the following result.

**Theorem 2.5.** Assume that there exist an orthogonal matrix  $Q$  and a diagonal matrix  $D = \text{diag}\{\lambda_1, \dots, \lambda_n\}$  such that  $S = QDQ'$ . Then, for  $t > 0$ , the exponential of the matrix  $tA$  is computed as follows

$$e^{tA} = \begin{bmatrix} Q & 0 \\ 0 & Q \end{bmatrix} P \begin{bmatrix} e^{tG_1} & 0 & \dots & 0 \\ 0 & e^{tG_2} & \dots & 0 \\ \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \dots & e^{tG_n} \end{bmatrix} P' \begin{bmatrix} Q' & 0 \\ 0 & Q' \end{bmatrix}, \quad (10)$$

where  $P \in \mathbb{R}^{2n \times 2n}$  is defined by (9) and  $G_i = \begin{bmatrix} 0 & 1 \\ -\alpha\lambda_i - \delta & -\beta\lambda_i - \gamma \end{bmatrix}$ .

*Proof.* The two lemmas 2.1 and 2.2 imply that

$$A = \begin{bmatrix} Q & 0 \\ 0 & Q \end{bmatrix} B \begin{bmatrix} Q' & 0 \\ 0 & Q' \end{bmatrix} = \begin{bmatrix} Q & 0 \\ 0 & Q \end{bmatrix} P \begin{bmatrix} G_1 & 0 & \cdots & 0 \\ 0 & G_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \cdots & e^{G_n} \end{bmatrix} P' \begin{bmatrix} Q' & 0 \\ 0 & Q' \end{bmatrix}.$$

Formula (10) is proved by using the properties (c) and (f) in Theorem 2.4.  $\square$

**Remark 2.6.** We need to compute the matrix exponential of the small matrices  $tG_i$ . This will be presented in the next section. The exponential matrix  $e^{tG_i}$  can be computed explicitly by using formula (13), (14), or (15) depending on the sign of  $(\beta\lambda_i + \gamma)^2 - 4(\alpha\lambda_i + \delta)$ .

**Remark 2.7.** In practical situations, to minimize the computational cost, we will compute the action of the matrix exponential to a vector instead of computing it explicitly. In (10),  $P$  and  $Q$  are two square matrices of orders  $2n$  and  $n$ , respectively. Since  $P$  is a permutation matrix, it can be however stored as a column matrix with  $n$  entries by indicating the positions of the non-zero elements, for example:

$$P_3 = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 \\ 3 \\ 5 \\ 2 \\ 4 \\ 6 \end{bmatrix}, \quad P'_3 = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 \\ 4 \\ 2 \\ 5 \\ 3 \\ 6 \end{bmatrix}$$

The block matrix  $[e^{tG_i}]$  can be stored as a  $2 \times 2n$  matrix. Given a compound vector  $v_0 = [a, b]'$ , where  $a$  and  $b$  are two column vectors with  $n$  entries, we start by evaluating a new vector  $v_1 = [Q'a, Q'b]'$ . Next, the action of the permutation matrix  $P'$  to the vector  $v_1$  is the reorder of its entries to get a new vector  $v_2$ . Then we compute the multiplication of the block exponential matrix with the vector  $v_3$  by cheaply multiplying each block  $2 \times 2$  matrix  $e^{tG_i}$  with two corresponding elements of  $v_2$ . Analogously for last two steps, we get an exact valuation of the action of matrix exponential to an arbitrary vector.

**2.2. The matrix exponential  $G_i$ .** From (10), instead of evaluating the matrix exponential of  $A \in \mathbb{R}^{2n \times 2n}$ , we need to compute the matrix exponential of each  $G_i \in \mathbb{R}^{2 \times 2}$ . In this section, we give some explicit formulas.

**Theorem 2.8.** *Assume that  $f$  is an analytic function. For a  $2 \times 2$  matrix  $G$ , the matrix function  $f(G)$  can be computed explicitly as*

$$f(G) = \frac{f(z_1) - f(z_2)}{z_1 - z_2} G + \frac{z_1 f(z_2) - z_2 f(z_1)}{z_1 - z_2} I, \quad (11)$$

where  $z_1$  and  $z_2$  are the two distinct eigenvalues of the matrix  $G$ . In case the matrix  $G$  has a double eigenvalue  $z_1$ , we get

$$f(G) = f'(z_1)G + (f(z_1) - f'(z_1)z_1) I. \quad (12)$$

*Proof.* Let  $p(z)$  be the characteristic polynomial of the matrix  $G$  and assume for a moment that the equation  $p(z) = 0$  has two distinct roots  $z_1$  and  $z_2$ . The Cayley–Hamilton theorem states that  $p(G) = 0$ . Since  $f$  is an analytic function, we know that

$$f(z) = \sum_{k=0}^{\infty} a_k z^k.$$

The function  $f$  can be rewritten in the form  $f(z) = q(z)p(z) + r(z)$  where  $q(z)$  is some quotient and  $r(z)$  is a remainder polynomial with  $0 \leq \deg r(z) < 2$ . From  $p(G) = 0$ , we obtain

$$f(G) = r(G) = d_1 G + d_0 I.$$

To complete the proof, we determine the coefficients  $d_1$  and  $d_0$ . From  $f(z_1) = r(z_1)$  and  $f(z_2) = r(z_2)$ , we obtain that

$$d_1 = \frac{f(z_1) - f(z_2)}{z_1 - z_2}, \quad d_2 = \frac{z_1 f(z_2) - z_2 f(z_1)}{z_1 - z_2}.$$

In case of a double eigenvalue  $z_1$ , we use the conditions  $f(z_1) = r(z_1)$  and  $f'(z_1) = r'(z_1)$ . As a consequence, we obtain that  $d_1 = f'(z_1)$  and  $d_0 = f(z_1) - f'(z_1)z_1$ .  $\square$

To reduce the computational cost, we try to avoid complex arithmetic.

**Lemma 2.9.** *Assume that the matrix  $G$  is of the form*

$$G = \begin{bmatrix} 0 & 1 \\ -\alpha\lambda - \delta & -\beta\lambda - \gamma \end{bmatrix}$$

and denote  $m = -\frac{1}{2}(\beta\lambda + \gamma)$ .

(i) *If  $(\beta\lambda + \gamma)^2 > 4(\alpha\lambda + \delta)$ , denoting  $n = \frac{1}{2}\sqrt{(\beta\lambda + \gamma)^2 - 4(\alpha\lambda + \delta)}$ , the exponential matrix  $e^{tG}$  can be computed explicitly as follows*

$$e^{tG} = \frac{e^{t(m+n)} - e^{t(m-n)}}{2n} \begin{bmatrix} -m - n & 1 \\ n^2 - m^2 & m - n \end{bmatrix} + e^{t(m+n)} I. \quad (13)$$

(ii) *If  $(\beta\lambda + \gamma)^2 = 4(\alpha\lambda + \delta)$ , we obtain that*

$$e^{tG} = e^{tm} \begin{bmatrix} 1 - tm & t \\ -tm^2 & tm + 1 \end{bmatrix}. \quad (14)$$

(iii) *If  $(\beta\lambda + \gamma)^2 < 4(\alpha\lambda + \delta)$ , denoting  $n = \frac{1}{2}\sqrt{4(\alpha\lambda + \delta) - (\beta\lambda + \gamma)^2}$ , we get that*

$$e^{tG} = \frac{e^{tm} \sin(tn)}{n} \begin{bmatrix} -m & 1 \\ -n^2 - m^2 & m \end{bmatrix} + e^{tm} \cos(tn) I. \quad (15)$$

*Proof.* Let  $z_1$  and  $z_2$  be the two eigenvalues of the matrix  $tG$ . Thus  $z_1$  and  $z_2$  satisfy the characteristic equation  $z^2 + (\beta\lambda + \gamma)tz + (\alpha\lambda + \delta)t^2 = 0$ . By using formula (11), we obtain that

$$e^{tG} = \frac{e^{z_1} - e^{z_2}}{z_1 - z_2} tG + \frac{z_1 e^{z_2} - z_2 e^{z_1}}{z_1 - z_2} I. \quad (16)$$

The discriminant of the characteristic equation is

$$D = ((\beta\lambda + \gamma)^2 - 4(\alpha\lambda + \delta)) t^2.$$

We consider three cases:

- (i) If  $D > 0$  or  $(\beta\lambda + \gamma)^2 > 4(\alpha\lambda + \delta)$ , the two real roots of the characteristic equation are  $z_1 = tm + tn$  and  $z_2 = tm - tn$ , where  $m = -\frac{1}{2}(\beta\lambda + \gamma)$  and  $n = \frac{1}{2}\sqrt{(\beta\lambda + \gamma)^2 - 4(\alpha\lambda + \delta)}$ . From the definitions of the parameters  $m$  and  $n$ , we get that  $-\alpha\lambda - \delta = n^2 - m^2$  and  $-\beta\lambda - \gamma = 2m$ . We will simplify the two coefficients in formula (16):

$$\begin{aligned} \frac{e^{z_1} - e^{z_2}}{z_1 - z_2} &= \frac{e^{t(m+n)} - e^{t(m-n)}}{2tn}, \\ \frac{z_1 e^{z_2} - z_2 e^{z_1}}{z_1 - z_2} &= \frac{(m+n)e^{t(m-n)} - (m-n)e^{t(m+n)}}{2n} \\ &= e^{t(m+n)} - (m+n) \frac{e^{t(m+n)} - e^{t(m-n)}}{2n}. \end{aligned}$$

By substituting the two simplified coefficients into (16), we get that

$$\begin{aligned} e^{tG} &= \frac{e^{t(m+n)} - e^{t(m-n)}}{2n} (G - (m+n)I) + e^{t(m+n)} I \\ &= \frac{e^{t(m+n)} - e^{t(m-n)}}{2n} \left( \begin{bmatrix} 0 & 1 \\ n^2 - m^2 & 2m \end{bmatrix} - \begin{bmatrix} m+n & 0 \\ 0 & m+n \end{bmatrix} \right) + e^{t(m+n)} I \\ &= \frac{e^{t(m+n)} - e^{t(m-n)}}{2n} \begin{bmatrix} -m-n & 1 \\ n^2 - m^2 & m-n \end{bmatrix} + e^{t(m+n)} I. \end{aligned}$$

- (ii) If  $D = 0$  or  $(\beta\lambda + \gamma)^2 = 4(\alpha\lambda + \delta)$ , the characteristic equation has only one root  $z_1 = tm$ , where  $m = -\frac{1}{2}(\beta\lambda + \gamma)$ . In this case, we have

$$\begin{aligned} e^{tG} &= e^{z_1}(tG) + e^{z_1}(1 - z_1)I \\ &= e^{tm} t \begin{bmatrix} 0 & 1 \\ -m^2 & 2m \end{bmatrix} + e^{tm}(1 - tm) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = e^{tm} \begin{bmatrix} 1 - tm & t \\ -tm^2 & tm + 1 \end{bmatrix}. \end{aligned}$$

- (iii) If  $D < 0$  or  $(\beta\lambda + \gamma)^2 < 4(\alpha\lambda + \delta)$ , the characteristic equation has two conjugate complex roots  $z_1 = tm + itn$  and  $z_2 = tm - itn$ , where

$$m = -\frac{1}{2}(\beta\lambda + \gamma), \quad n = \frac{1}{2}\sqrt{4(\alpha\lambda + \delta) - (\beta\lambda + \gamma)^2}.$$

Since  $z_2 = \overline{z_1}$ , we infer that  $e^{z_2} = \overline{e^{z_1}}$  and  $z_2 e^{z_1} = \overline{z_1 e^{z_2}} = \overline{z_1 e^{z_2}}$ . We analogously simplify the two coefficients in (16) as follows

$$\begin{aligned} \frac{e^{z_1} - e^{z_2}}{z_1 - z_2} &= \frac{e^{z_1} - \overline{e^{z_1}}}{z_1 - \overline{z_1}} = \frac{2\operatorname{Im}(e^{z_1})}{2\operatorname{Im}(z_1)} = \frac{e^{tm} \sin(tn)}{tn}, \\ \frac{z_1 e^{z_2} - z_2 e^{z_1}}{z_1 - z_2} &= \frac{\operatorname{Im}(z_1 e^{\overline{z_1}})}{\operatorname{Im}(z_1)} = e^{tm} \cos(tn) - m \frac{e^{tm} \sin(tn)}{n}. \end{aligned}$$

By substituting the two simplified coefficients into (16) and noting that  $-\alpha\lambda - \delta = -n^2 - m^2$  and  $-\beta\lambda - \gamma = 2m$ , we get that

$$\begin{aligned} e^{tG} &= \frac{e^{tm} \sin(tn)}{n} (G - mI) + e^{tm} \cos(tn) I \\ &= \frac{e^{tm} \sin(tn)}{n} \begin{bmatrix} -m & 1 \\ -n^2 - m^2 & m \end{bmatrix} + e^{tm} \cos(tn) I. \end{aligned}$$

This concludes the proof.  $\square$



**Remark 2.10.** Formula (15) is useful in computations. In practical situations, when  $\beta, \gamma$ , and  $\delta$  are often much smaller than  $\alpha$ . For example, for beam equations with typical physical parameters proposed by Ito and Morris in [IM98, Section 3], the matrix  $tG$  has usually two complex eigenvalues.

### 3. EXPONENTIAL INTEGRATORS

**3.1. Exponential integrators for semilinear problems.** We consider semi-linear differential equations of the form

$$\dot{y}(t) = Ay(t) + F(y(t)). \tag{17}$$

The solution of this equation at time  $t_{n+1} = t_n + \tau_n$ ,  $t_0 = 0, n \in \mathbb{N}$  is given by the variation-of-constants formula

$$y(t_{n+1}) = e^{\tau_n A} y(t_n) + \int_0^{\tau_n} e^{(\tau_n - \tau)A} F(y(t_n + \tau)) d\tau.$$

For the numerical solution of (17), we recall a general class of one-step exponential integrators from [HO05a, HO05b, HO10]

$$y_{n+1} = e^{\tau_n A} y_n + \tau_n \sum_{i=1}^s b_i(\tau_n A) F_{ni},$$

$$Y_{ni} = e^{c_j \tau_n A} y_n + \tau_n \sum_{j=1}^{i-1} a_{ij}(\tau_n A) F_{nj},$$

$$F_{nj} = F(Y_{nj}).$$

The coefficients are as usually collected in a Butcher tableau

$c_1$				
$c_2$	$a_{21}(\tau_n A)$			
$\vdots$	$\vdots$	$\ddots$		
$c_s$	$a_{s1}(\tau_n A)$	$\dots$	$a_{s,s-1}(\tau_n A)$	
	$b_1(\tau_n A)$	$\dots$	$b_{s-1}(\tau_n A)$	$b_s(\tau_n A)$

The method coefficients  $a_{ij}$  and  $b_i$  are constructed from a family of functions  $\varphi_k$  evaluated at the matrix  $(\tau_n A)$ . We next recall this family  $\varphi_k$ , which was introduced before in [HO05a, HO05b, HO10].

**Corollary 3.1.** *Consider the entire functions*

$$\varphi_k(z) = \int_0^1 e^{(1-\theta)z} \frac{\theta^{k-1}}{(k-1)!} d\theta, \quad k \geq 1, \quad \varphi_0(z) = e^z.$$

*These functions satisfy the following properties*

- (i)  $\varphi_k(0) = \frac{1}{k!}$ ;
- (ii) they satisfy the recurrence relation

$$\varphi_{k+1}(z) = \frac{\varphi_k(z) - \varphi(0)}{z};$$

- (iii) the Taylor expansion of the function  $\varphi_k$  is

$$\varphi_k(z) = \sum_{n=0}^{\infty} \frac{z^n}{(n+k)!}.$$

To simplify notation, we denote

$$\varphi_{k,j} = \varphi_k(c_j \tau_n A), \quad \varphi_k = \varphi_k(\tau_n A).$$

Next, we recall five exponential integrators that will be used in our numerical examples.

**Example 3.2.** For  $s = 1$ , the exponential Euler method has the form

$$y_{n+1} = e^{\tau_n A} y_n + \tau_n \varphi_1(\tau_n A) F(y_n). \quad (18)$$

$$(EI-E1) \quad \begin{array}{c|c} 0 & \\ \hline c_1 & \varphi_1 \end{array}$$

We denote this method by EI-E1.

**Example 3.3.** For  $s = 2$ , we recall a second-order method proposed by Strehmel and Weiner in [SW92, Section 4.5.3]:

$$(EI-SW21) \quad \begin{array}{c|cc} 0 & & \\ c_2 & c_2 \varphi_{1,2} & \\ \hline & \varphi_1 - \frac{1}{c_2} \varphi_2 & \frac{1}{c_2} \varphi_2 \end{array}$$

A simplified version, where only  $\varphi_1$  is used, is also proposed by Strehmel and Weiner

$$(EI-SW22) \quad \begin{array}{c|cc} 0 & & \\ c_2 & c_2 \varphi_{1,2} & \\ \hline & \left(1 - \frac{1}{2c_2}\right) \varphi_1 & \frac{1}{c_2} \varphi_1 \end{array}$$

**Example 3.4.** For  $s = 4$ , we recall two schemes. The first one, proposed by Krogstad in [Kro05], is given by

$$(EI-K4) \quad \begin{array}{c|ccc} 0 & & & \\ \frac{1}{2} & \frac{1}{2} \varphi_{1,2} & & \\ \frac{1}{2} & \frac{1}{2} \varphi_{1,3} - \varphi_{2,3} & \varphi_{2,3} & \\ 1 & \varphi_{1,4} - 2\varphi_{2,4} & 0 & 2\varphi_{2,4} \\ \hline & \varphi_1 - 3\varphi_2 + 4\varphi_3 & 2\varphi_2 - 4\varphi_3 & 2\varphi_2 - 4\varphi_3 \quad -\varphi_2 + 4\varphi_3 \end{array}$$

The second method is suggested by Strehmel and Weiner (see [SW92, Example 4.5.5])

$$(EI-SW4) \quad \begin{array}{c|ccc} 0 & & & \\ \frac{1}{2} & \frac{1}{2} \varphi_{1,2} & & \\ \frac{1}{2} & \frac{1}{2} \varphi_{1,3} - \frac{1}{2} \varphi_{2,3} & \frac{1}{2} \varphi_{2,3} & \\ 1 & \varphi_{1,4} - 2\varphi_{2,4} & -2\varphi_{2,4} & 4\varphi_{2,4} \\ \hline & \varphi_1 - 3\varphi_2 + 4\varphi_3 & 0 & 4\varphi_2 - 8\varphi_3 \quad -\varphi_2 + 4\varphi_3 \end{array}$$

**3.2. Computing matrix functions of  $tA$ .** To apply these exponential integrators to semilinear problems, we next introduce an approach to explicitly compute the matrix functions  $\varphi_k(tA)$ . We first present an analogous version of Theorem 2.5.

**Theorem 3.5.** *Assume that there exist an orthogonal matrix  $Q$  and a diagonal matrix  $D = \text{diag}\{\lambda_1, \dots, \lambda_n\}$  such that  $S = QDQ'$ . Then, for  $t > 0$  and  $k \geq 1$ , the functions  $\varphi_k(tA)$  are computed as follows*

$$\varphi_k(tA) = \begin{bmatrix} Q & 0 \\ 0 & Q \end{bmatrix} P \begin{bmatrix} \varphi_k(tG_1) & 0 & \cdots & 0 \\ 0 & \varphi_k(tG_2) & \cdots & 0 \\ \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \cdots & \varphi_k(tG_n) \end{bmatrix} P' \begin{bmatrix} Q' & 0 \\ 0 & Q' \end{bmatrix}, \quad (19)$$

where  $P \in \mathbb{R}^{2n \times 2n}$  is given in (9) and  $G_i = \begin{bmatrix} 0 & 1 \\ -\alpha\lambda_i - \delta & -\beta\lambda_i - \gamma \end{bmatrix}$ .

The matrix functions  $\varphi_k(tG_i)$  are computed explicitly. The actual formula depends on the sign of  $(\beta\lambda_i + \gamma)^2 - 4(\alpha\lambda_i + \delta)$ . Next, we will present two lemmas concerning these functions.

**Lemma 3.6.** *Assume that the matrix  $G$  is of the form*

$$G = \begin{bmatrix} 0 & 1 \\ -\alpha\lambda - \delta & -\beta\lambda - \gamma \end{bmatrix}$$

and denote  $m = -\frac{1}{2}(\beta\lambda + \gamma)$ .

(i) *If  $(\beta\lambda + \gamma)^2 > 4(\alpha\lambda + \delta)$ , denoting  $n = \frac{1}{2}\sqrt{(\beta\lambda + \gamma)^2 - 4(\alpha\lambda + \delta)}$ , the matrix functions  $\varphi_k(tG)$  can be computed explicitly as follows*

$$\varphi_k(tG) = \frac{\varphi_k^+ - \varphi_k^-}{2n} \begin{bmatrix} -m - n & 1 \\ n^2 - m^2 & m - n \end{bmatrix} + \varphi_k^+ I, \quad (20)$$

where  $\varphi_k^+ = \varphi_k(t(m+n))$  and  $\varphi_k^- = \varphi_k(t(m-n))$ .

(ii) *If  $(\beta\lambda + \gamma)^2 = 4(\alpha\lambda + \delta)$ , we obtain that*

$$\varphi_k(tG) = \varphi_k'(tm) \begin{bmatrix} -tm & t \\ -tm^2 & tm \end{bmatrix} + \varphi_k(tm) I, \quad (21)$$

where the derivative  $\varphi_k'(z)$  can be computed recursively

$$\begin{aligned} \varphi_0(z) &= \varphi_0'(z) = e^z, \\ \varphi_{k+1}'(z) &= \frac{\varphi_k'(z) - \varphi_{k+1}(z)}{z}, \quad \varphi_{k+1}(z) = \frac{\varphi_k(z) - \varphi_k(0)}{z}. \end{aligned}$$

*Proof.* By applying Theorem 2.8, the proof follows the lines of the first two points in the proof of Lemma 2.9.  $\square$

The last lemma concentrates on the case of two complex eigenvalues of  $G_i$ . Again the idea is to compute the matrix functions without explicitly using complex numbers. It is inspired by formula (15) above.

**Lemma 3.7.** *In the case  $(\beta\lambda + \gamma)^2 < 4(\alpha\lambda + \delta)$ , the matrix  $G$  has two conjugate complex eigenvalues  $z_1$  and  $z_2$  with  $z_1 = tm + itn$ , where*

$$m = -\frac{1}{2}(\beta\lambda + \gamma), \quad n = \frac{1}{2}\sqrt{4(\alpha\lambda + \delta) - (\beta\lambda + \gamma)^2}.$$

The matrix  $\varphi_k(tG)$  can be explicitly computed as follows

$$\varphi_k(tG) = \frac{i_k}{n} \begin{bmatrix} -m & 1 \\ -n^2 - m^2 & m \end{bmatrix} + r_k I. \quad (22)$$

Here, the two coefficients  $i_k$  and  $r_k$  depend on  $(t, m, n)$  and can be computed recursively as follows

$$i_0 = e^{tm} \sin(tn), \quad r_0 = e^{tm} \cos(tn), \quad (23a)$$

$$i_k = \frac{1}{t(m^2 + n^2)} \left( mi_{k-1} - n \left( r_{k-1} - \frac{1}{(k-1)!} \right) \right), \quad (23b)$$

$$r_k = \frac{1}{t(m^2 + n^2)} \left( ni_{k-1} + m \left( r_{k-1} - \frac{1}{(k-1)!} \right) \right). \quad (23c)$$

*Proof.* By using formula (11), we obtain that

$$\varphi_k(tG) = \frac{\varphi_k(z_1) - \varphi_k(z_2)}{z_1 - z_2} tG + \frac{z_1 \varphi_k(z_2) - z_2 \varphi_k(z_1)}{z_1 - z_2} I.$$

First, we note that  $\varphi_k(z_2) = \overline{\varphi_k(z_1)}$  because  $\varphi_k$  has real coefficients. Thus we can simplify as follows

$$\frac{\varphi_k(z_1) - \varphi_k(z_2)}{z_1 - z_2} = \frac{\operatorname{Im}(\varphi_k(z_1))}{\operatorname{Im} z_1}, \quad \frac{z_1 \varphi_k(z_2) - z_2 \varphi_k(z_1)}{z_1 - z_2} = \frac{\operatorname{Im}(z_1 \overline{\varphi_k(z_1)})}{\operatorname{Im} z_1}.$$

Next, we rewrite the recursion as follows

$$\varphi_{k+1}(z_1) = \frac{\varphi_k(z_1) - \varphi_k(0)}{z_1} = \frac{(\varphi_k(z_1) - \varphi_k(0)) \overline{z_1}}{|z_1|^2}.$$

To simplify notation, we denote  $i_k = \operatorname{Im}(\varphi_k(z_1))$  and  $r_k = \operatorname{Re}(\varphi_k(z_1))$ . Thus we obtain that

$$\begin{aligned} i_{k+1} &= \operatorname{Im}(\varphi_{k+1}(z_1)) = \frac{1}{|z_1|^2} \operatorname{Im} \left( (\varphi_k(z_1) - \varphi_k(0)) \overline{z_1} \right) \\ &= \frac{1}{t^2(m^2 + n^2)} \left( \operatorname{Im}(\varphi_k(z_1) - \varphi_k(0)) \operatorname{Re}(\overline{z_1}) + \operatorname{Re}(\varphi_k(z_1) - \varphi_k(0)) \operatorname{Im}(\overline{z_1}) \right) \\ &= \frac{1}{t(m^2 + n^2)} \left( mi_k - n \left( r_k - \frac{1}{k!} \right) \right), \\ r_{k+1} &= \operatorname{Re}(\varphi_{k+1}(z_1)) = \frac{1}{|z_1|^2} \operatorname{Re} \left( (\varphi_k(z_1) - \varphi_k(0)) \overline{z_1} \right) \\ &= \frac{1}{t^2(m^2 + n^2)} \left( \operatorname{Re}(\varphi_k(z_1) - \varphi_k(0)) \operatorname{Re}(\overline{z_1}) - \operatorname{Im}(\varphi_k(z_1) - \varphi_k(0)) \operatorname{Im}(\overline{z_1}) \right) \\ &= \frac{1}{t(m^2 + n^2)} \left( ni_k + m \left( r_k - \frac{1}{k!} \right) \right). \end{aligned}$$

Besides, we also get that  $\operatorname{Im}(z_1 \overline{\varphi_k(z_1)}) = tnr_k - tmi_k$ . This finally yields that

$$\begin{aligned} \varphi_k(tG) &= \frac{i_k}{n} G + \frac{nr_k - mi_k}{n} I = \frac{i_k}{n} (G - mI) + r_k I \\ &= \frac{i_k}{n} \begin{bmatrix} -m & 1 \\ -n^2 - m^2 & m \end{bmatrix} + r_k I, \end{aligned}$$

which completes the proof.  $\square$

**3.3. Summary of the integration procedure.** The above described procedure can be summarized in two main parts. The **P**reparation part, which is done once at the beginning, consists of three steps:

- P1:** Discretize the operator  $(-\Delta)$  or  $\partial_{xxxx}^4$  as a square symmetric matrix  $S$  (e.g., by finite differences, see also section 4).
- P2:** Find an orthogonal matrix  $Q$  and a diagonal matrix  $D = \text{diag}\{\lambda_1, \dots, \lambda_n\}$  such that  $S = QDQ'$ . The matrix  $D$  is stored as a vector.
- P3:** Create a column vector which stores the positions of all non-zero entries of the permutation matrix  $P$  by using formula (9).

The **M**ain part is used to compute the action of a matrix functions. This is required in the time stepping. Computing this action consists of two steps:

- M1:** Compute the matrix functions  $\varphi_k(tG_i)$  by using formulas (13), (14), or (15) for  $\varphi_0(tG_i) = e^{tG_i}$ ; formulas (20), (21), or (22) for  $\varphi_k(tG_i)$  with  $k \geq 1$ .
- M2:** Compute the action of the matrix functions  $\varphi_k(tA)$  using formula (10) for  $k = 0$  and formula (19) for  $k \geq 1$ .

#### 4. NUMERICAL EXAMPLES

**4.1. Semilinear wave equations.** We consider a 1D semilinear wave equation on  $\Omega = (0, \ell)$

$$\partial_{tt}^2 u - \alpha \partial_{xx}^2 u - \beta \partial_{xxt}^3 u + \delta u + \gamma \partial_t u = g(u), \quad 0 < x < \ell, \quad t > 0, \quad (24a)$$

$$u(t, 0) = 0, \quad u(t, \ell) = 0, \quad (24b)$$

$$u(0, x) = p(x), \quad u_t(0, x) = q(x). \quad (24c)$$

We consider the product space  $X = H_0^1(\Omega) \times L^2(\Omega)$  and rewrite (24) in abstract form

$$\dot{y}(t) = \mathcal{A}y(t) + \mathcal{F}(y(t)), \quad y(0) = (p, q)',$$

where  $\mathcal{A} : D(\mathcal{A}) \rightarrow X$  is the operator defined by

$$\begin{aligned} \mathcal{A} \begin{bmatrix} u \\ w \end{bmatrix} &= \begin{bmatrix} 0 & I \\ -\alpha(-\partial_{xx}^2) - \delta I & -\beta(-\partial_{xx}^2) - \gamma I \end{bmatrix} \begin{bmatrix} u \\ w \end{bmatrix} \\ &= \begin{bmatrix} w \\ -\alpha(-\partial_{xx}^2 u) - \beta(-\partial_{xx}^2 w) - \delta u - \gamma w \end{bmatrix}, \\ D(\mathcal{A}) &= (H^2(\Omega) \cap H_0^1(\Omega))^2, \\ \mathcal{F} \begin{pmatrix} u \\ w \end{pmatrix} &= \begin{bmatrix} 0 \\ f(u) \end{bmatrix}. \end{aligned}$$

Define the closed self-adjoint positive operator  $\mathcal{A}_0$  on  $L^2(0, \ell)$  by

$$\begin{aligned} \mathcal{A}_0 \phi &= -\partial_{xx}^2 \phi, \\ D(\mathcal{A}_0) &= \{\phi \in H^2(0, \ell) \mid \phi(0) = \phi(1) = 0\}. \end{aligned}$$


We use symmetric finite differences to discretize the operator  $\mathcal{A}_0$ . For this, the space interval  $(0, \ell)$  is divided equidistantly by the nodes  $x_i = i\Delta x$ ,  $i \in$

$\{0, \dots, N + 1\}$ , where  $N$  is a given natural number and  $\Delta x = \frac{\ell}{N+1}$ . Then, the discrete operator  $\mathcal{A}_0$  is given by the matrix  $S_w \in \mathbb{R}^{N \times N}$  defined by

$$S_w = \frac{1}{\Delta x^2} \begin{bmatrix} 2 & -1 & 0 & \dots & 0 \\ -1 & 2 & -1 & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & -1 & 2 & 1 \\ 0 & \dots & 0 & -1 & 2 \end{bmatrix}.$$

In the four examples below, we consider the space interval  $\Omega = (0, 1)$  with  $N = 200$ .

**Example 4.1.** Consider equation (24) with  $\alpha = \pi^2$ ,  $\beta = 0.01$ ,  $\delta = 0$ ,  $\gamma = 0.01$ . The nonlinear source term is  $g(u) = \sin(u)$ . This is a perturbed sine-Gordon equation of the form (2). The initial conditions are  $p(x) = 5 \sin(2\pi x)$  and  $q(x) = 0$ . We use three different schemes namely EI-E1, EI-SW21, and EI-K4 to compute the solution at time  $T = 6$  with  $M \in \{5, 10, 20, \dots, 5 \cdot 2^{11}\}$  time steps. The reference solution plotted in Figure 1b is computed by using EI-SW4 with  $M = 200000$  time steps. We compute the discrete  $L_2$ -errors between these approximate solutions obtained with the three mentioned schemes and the reference solution. The errors are plotted in figure 1a. The expected convergence rate is observed for each schemes. Even though we use an coarse time mesh with  $M = 5$  and  $\Delta t = 1.2$ , the error here is quite small (approximate  $10^{-2}$ ).



a) Convergence rates.

b) Reference solution at  $T = 6$ .

FIGURE 1. Example 4.1

**Example 4.2.** Consider equation (24) with  $\alpha = 100$ ,  $\beta = 0.01$ ,  $\delta = 0$ ,  $\gamma = 0.001$ . The nonlinear source term is  $g(u) = u|u|$ . The initial conditions are

$$p(x) = \begin{cases} 2x & \text{if } x \leq \frac{1}{2}, \\ -2x + 2 & \text{if } x > \frac{1}{2}, \end{cases} \quad q(x) = \pi^2 \sin(\pi x).$$

We use three different schemes, namely EI-E1, EI-SW22, and EI-SW4 to compute the solution at time  $T = 15$  with  $M \in \{20, 40, 80, \dots, 20 \cdot 2^{12}\}$  time steps. The reference solution, plotted in figure 2b, is computed by using EI-K4 with  $M = 200000$ . The  $L_2$ -errors are plotted in Figure 2a. The  $L_2$ -error

for all schemes is quite small, approximately  $10^{-2}$ , although we use a coarse mesh (for example  $M = 20$ ,  $\Delta t = 0.75$ ).




FIGURE 2. Example 4.2

**Example 4.3.** Consider equation (24) with  $\alpha = 15$ ,  $\beta = 10^{-3}$ ,  $\delta = 1$ ,  $\gamma = 10^{-6}$ . The nonlinear source term is  $g(u) = u^3$ . The initial conditions are  $p(x) = 10 \sin(3\pi x)$ ,  $q(x) = -10 \cos(3\pi x)$ . We use all five schemes, namely EI-E1, EI-SW21, EI-SW4, and EI-K4 to compute the solution at time  $T = 30$  with  $M \in \{20, 40, 80, \dots, 20 \cdot 2^{13}\}$  time steps. We plot the reference solution computed with EI-K4 and  $M = 300000$  in figure 3b. The  $L_2$ -errors are plotted in Figure 3a. Again the expected convergence rates are observed for each schemes. Comparing the examples above, the exponential Euler method (EI-E1) works extremely bad here. The other methods still preserve the convergence rate.




FIGURE 3. Example 4.3

**Example 4.4.** The last example for wave equation (24) concerns discontinuous initial conditions

$$p(x) = \begin{cases} -1 & \text{if } x \leq \frac{1}{2}, \\ 5 & \text{if } x > \frac{1}{2}, \end{cases} \quad q(x) = 0.$$

The other parameters are  $\alpha = 5$ ,  $\beta = 10^{-3}$ ,  $\delta = 1$ ,  $\gamma = 10^{-4}$ . The nonlinear term is  $g(u) = |u|$ . The approximate solutions at  $T = 3$  are computed by using five exponential integrators, namely EI-E1, EI-SW21, EI-SW22, EI-SW4, and EI-K4 with  $M \in \{20, 40, \dots, 20 \cdot 2^{13}\}$  time steps. The reference solution computed with  $M = 300000$  time steps by using EI-SW4 is plotted in figure 4b. The  $L_2$ -errors are plotted in Figure 4a. We observe an order reduction to order 2 for the two fourth-order exponential integrators EI-SW4 and EI-K4 while the other integrators preserve their convergence rate.




FIGURE 4. Example 4.4

**Remark 4.5.** Another common approach for solving (24) consists in merging these damping terms with the nonlinear terms and then employing various exponential integrators to solve the resulting semilinear equation. However this approach would be actually more expensive and it also lacks the accuracy.

**4.2. Railway track model.** Assume that a track beam is made of Kelvin-Voigt material. The resulting railway track model is a semilinear PDE on  $\Omega = (0, \ell)$

$$\partial_{tt}^2 u + \partial_{xx}^2 (\alpha \partial_{xx}^2 u + \beta \partial_{xx}^2 (\partial_t u)) + \gamma \partial_t u + \delta u = -mu^3, \quad x \in \Omega, t > 0, \quad (25a)$$

$$u(t, 0) = u(t, \ell) = 0, \quad (25b)$$

$$\alpha \partial_{xx}^2 u(t, 0) + \beta \partial_{xx}^2 (\partial_t u)(t, 0) = \alpha \partial_{xx}^2 u(t, \ell) + \beta \partial_{xx}^2 (\partial_t u)(t, \ell) = 0, \quad (25c)$$

$$u(0, x) = p(x), \quad \partial_t u(0, x) = q(x). \quad (25d)$$

Denote the closed self-adjoint positive operator  $\mathcal{A}_0$  on  $L^2(0, \ell)$  as

$$\mathcal{A}_0 \phi := \partial_{xxxx}^4 \phi,$$

$$D(\mathcal{A}_0) := \{\phi \in H^4(\Omega) \mid \phi(0) = \phi(\ell) = 0, \phi''(0) = \phi''(\ell) = 0\}.$$

Concerning the analysis of the linear operators, we refer to the literature [Mor20, BI97]. We also use finite differences to discretize the operator  $\mathcal{A}_0$  with an equidistant space mesh  $x_i = i\Delta x$ ,  $i \in \{0, \dots, N+1\}$  where  $N$  is a given integer number and  $\Delta x = \frac{\ell}{N+1}$ . Then the discrete operator  $\mathcal{A}_0$  is




given by the matrix  $S_b \in \mathbb{R}^{N \times N}$ , defined as

$$S_b = \frac{1}{\Delta x^4} \begin{bmatrix} 5 & -4 & 1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & 0 \\ -4 & 6 & -4 & 1 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & 0 \\ 1 & -4 & 6 & -4 & 1 & 0 & \dots & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -4 & 6 & -4 & 1 & \dots & 0 & 0 & 0 & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & \dots & \dots & \dots & \ddots & 1 & -4 & 6 & -4 & 1 \\ 0 & \dots & \dots & \dots & \dots & \dots & \ddots & 0 & 1 & -4 & 6 & -4 \\ 0 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 1 & -4 & 5 \end{bmatrix}.$$


**Example 4.6.** Consider equation (25) with  $\alpha = 15$ ,  $\beta = 3\text{E-}6$ ,  $\delta = 10$ ,  $\gamma = 3\text{E} - 4$ . The nonlinear term is  $g(u) = -5u^3$ . The initial conditions are

$$p(x) = 5e^{-100(x-\frac{2}{3})^2}, \quad q(x) = 0.$$

For our numerical solution, the space interval  $\Omega = (0, 1)$  is divided into 300 equidistant subintervals. We compute approximate solutions at  $T = 5$  with the exponential integrators EI-SW22 and EI-SW4 with  $M \in \{20, 40, \dots, 20 \cdot 2^{13}\}$  time steps. We compare these numerical results with the reference solution evaluated by EI-K4 with  $M = 200000$  time steps. The errors are plotted in figure 5a.



a) Convergence Rate.



b) Reference solution at  $T = 5$ .

FIGURE 5. Example 4.5

## 5. CONCLUSION

We presented an approach to cheaply compute the action of the matrix exponential  $e^{tA}$  as well as the action of the matrix functions  $\varphi_k(tA)$  by employing two linear transformations. Thus, the solution of certain linear

differential equations can be computed in a fast and efficient way. By applying the exponential integrators in the literature, we can solve the semilinear wave equations and semilinear beam equations.

Note that the described procedure can be extended to the case

$$A = \begin{bmatrix} A_1 & A_2 \\ A_3 & A_4 \end{bmatrix}, \quad \text{where } [A_i, A_j] = 0 \quad \text{for } 1 \leq i, j \leq 4, \quad i \neq j.$$

Indeed, under the above assumption, the four matrices  $A_i$  share the same eigenvalues and eigenvectors. Thus, there exist a matrix  $Q$  and four corresponding diagonal matrices  $D_i$  such that  $A_i = QD_iQ'$  for  $1 \leq i \leq 4$ . This implies that

$$A = \begin{bmatrix} Q & 0 \\ 0 & Q \end{bmatrix} \begin{bmatrix} D_1 & D_2 \\ D_3 & D_4 \end{bmatrix} \begin{bmatrix} Q' & 0 \\ 0 & Q' \end{bmatrix}$$

Thus, the scheme can be analogously applied by evaluating the exponential of each  $2 \times 2$  block matrix  $\begin{bmatrix} \lambda_{1,i} & \lambda_{2,i} \\ \lambda_{3,i} & \lambda_{4,i} \end{bmatrix}$  where  $\lambda_{k,i}$  is an entry of the diagonal matrix  $D_k$  with  $1 \leq k \leq 4$ .

## REFERENCES

- [AEY11] M. Ansari, E. Esmailzadeh, and D. Younesian. Frequency analysis of finite beams on nonlinear Kelvin–Voigt foundation under moving loads. *Journal of Sound and Vibration*, 330:1455–1471, 2011.
- [BI97] H. T. Banks and K. Ito. Approximation in LQR problems for infinite dimensional systems with unbounded input operators. *Journal of Mathematical Systems, Estimation and Control*, 7:1–34, 1997.
- [CCD08] A.N. Carvalho, J.W. Cholewa, and T. Dlotko. Strongly damped wave problems: bootstrapping and regularity of solutions. *Journal of Differential Equations*, 244:2310–2333, 2008.
- [CF21] W. Chen and A. Z. Fino. Blow-up of solutions to semilinear strongly damped wave equations with different nonlinear terms in an exterior domain. *Mathematical Methods in the Applied Sciences*, 44:6787–6807, 2021.
- [CM02] S.M. Cox and P.C. Matthews. Exponential time differencing for stiff systems. *Journal of Computational Physics*, 176:430–455, 2002.
- [Deu79] P. Deuffhard. A study of extrapolation methods based on multistep schemes without parasitic solutions. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 30:177–189, 1979.
- [DOT17] V. N. V. Do, T. H. Ong, and C. H. Thai. Dynamic responses of Euler–Bernoulli beam subjected to moving vehicles using isogeometric approach. *Applied Mathematical Modelling*, 51, 2017.
- [EM19a] M. S. Edalatzadeh and K. A. Morris. Optimal actuator design for semilinear systems. *SIAM J. Control Optim.*, 57:2992–3020, 2019.
- [EM19b] M. S. Edalatzadeh and K. A. Morris. Stability and well-posedness of a nonlinear railway track model. *IEEE Control Systems Letters*, 3:162–167, 2019.
- [Gau61] W. Gautschi. Numerical integration of ordinary differential equations based on trigonometric polynomials. *Numerische Mathematik*, 3:381–397, 1961.
- [GM91] J. M. Ghidaglia and A. Marzocchi. Longtime behaviour of strongly damped wave equations, global attractors and their dimension. *SIAM Journal on Mathematical Analysis*, 22:879–895, 1991.
- [HAM10] N. J. Higham and A. H. Al-Mohy. Computing matrix functions. *Acta Numerica*, pages 159–208, 2010.
- [Hig08] N. J. Higham. *Functions of Matrices*. Society for Industrial and Applied Mathematics, 2008.

- [HO05a] M. Hochbruck and A. Ostermann. Explicit exponential Runge–Kutta methods for semilinear parabolic problems. *SIAM Journal on Numerical Analysis*, 43:1069–1090, 2005.
- [HO05b] M. Hochbruck and A. Ostermann. Exponential Runge–Kutta methods for parabolic problems. *Applied Numerical Mathematics*, 53:323–339, 2005.
- [HO10] M. Hochbruck and A. Ostermann. Exponential integrators. *Acta Numerica*, 19:209–286, 2010.
- [IM98] K. Ito and K. Morris. An approximation theory of solutions to operator Riccati equations for  $H^\infty$  control. *SIAM J. Control Optim.*, 36:82–99, 1998.
- [ITY13] R. Ikehata, G. Todorova, and B. Yordanov. Wave equations with strong damping in Hilbert spaces. *Journal of Differential Equations*, 254:3352–3368, 2013.
- [Kro05] S. Krogstad. Generalized integrating factor methods for stiff PDEs. *Journal of Computational Physics*, 203:72–88, 2005.
- [LL98] K. Liu and Z. Liu. Exponential decay of energy of the Euler–Bernoulli beam with locally distributed Kelvin–Voigt damping. *SIAM J. Control Optim.*, 36:1086–1098, 1998.
- [LO14] V. T. Luan and A. Ostermann. Explicit exponential Runge–Kutta methods of high order for parabolic problems. *Journal of Computational and Applied Mathematics*, 256:168–179, 2014.
- [Mor20] K. A. Morris. *Controller Design for Distributed Parameter Systems*. Springer International Publishing, 2020.
- [MS15] K. Mattsson and V. Stiernström. High-fidelity numerical simulation of the dynamic beam equation. *Journal of Computational Physics*, 286:194–213, 2015.
- [Pon85] G. Ponce. Global existence of small solutions to a class of nonlinear evolution equations. *Nonlinear Analysis: Theory, Methods & Applications*, 9:399–418, 1985.
- [PP20] L. Paunonen and D. Phan. Reduced order controller design for robust output regulation. *IEEE Transactions on Automatic Control*, 65:2480–2493, 2020.
- [PP21] D. Phan and L. Paunonen. Finite-dimensional controllers for robust regulation of boundary control systems. *Mathematical Control & Related Fields*, 11:95–117, 2021.
- [PZ06] V. Pata and S. Zelik. Smooth attractors for strongly damped wave equations. *Nonlinearity*, 19:1495–1506, 2006.
- [SW92] K. Strehmel and R. Weiner. *Linear-implizite Runge-Kutta-Methoden und ihre Anwendung*. Vieweg+Teubner Verlag, 1992.

*Email address:* `duy.phan-duc@uibk.ac.at`

INSTITUT FÜR MATHEMATIK, LEOPOLD-FRANZENS-UNIVERSITÄT INNSBRUCK, TECHNIKERSTRASSE 13, A-6020 INNSBRUCK, AUSTRIA.