Equiseparability on Terminal Wiener Index and Distances

Sulphikar A

Department of Computer Science and Engineering National Institute of Technology , Trichy Tamilnadu INDIA

Abstract

Teminal Wiener index is one of the commonly used topological index in mathematical chemistry. If two or more chemical compounds have the same terminal Wiener index then they will have similar physico-chemical properties. In this work we propose a new method for constructing equiseparable trees w.r.t terminal Wiener index. The existing method is based on the number of pendent vertices but the proposed method is based on distance parameters.

keywords

Terminal Wiener index, Wiener index, Distance in graphs, Equiseparability.

1 Introduction

Given a graph G = (V, E), the Wiener index W(G) of G is defined as

$$W(G) = \sum_{(u,v)\in V} d(u,v).$$

$$\tag{1}$$

where d(u, v) is the distance between vertices u and v of G.[2, 3]

Let T be an n-vertex tree and e = uv an edge of T. Denote by $n_u(e|T)$ (resp. $n_v(e|T)$) the number of vertices of T lying on one side of the edge e, closer to vertex u (resp. v). Then W(T) also satisfies the following relation:

$$W(T) = \sum_{e=uv} n_u(e|T)n_v(e|T)$$
⁽²⁾

Let T be an n-vertex tree with k pendent vertices. Then the terminal Wiener index TW(T) [1, 6, 7] of T is defined as the sum of the distances between all pairs of pendent vertices of T. That is

$$TW(T) = \sum_{1 \le i < j \le k} d(v_i, v_j) \tag{3}$$

where v_i, v_j are pendent vertices in T. Let P(T) be the set of pendent vertices of T. Terminal Wiener index can also be calculated as

$$TW(T) = \frac{1}{2} \sum_{u \in P(T)} d^+(u)$$
(4)

where $d^+(u)$ is the sum of the distances between u and pendent vertices of T. For an edge e = uv, $p_u(e)(\text{resp. } p_v(e))$ denotes the number of vertices closer to u(resp. v) than v(resp. u). The numbers $p_u(e)$ and $p_v(e)$ satisfy the relation

$$d^{+}(u) - d^{+}(v) = p_{v}(e) - p_{u}(e)$$
(5)

Let P_n be a path on n vertices. It is easy to see that $d^+(v) = n - 1$ for any $v \in V(P_n)$. Let uv be a pendent edge of a tree T with deg(u) = 1. Then $d^+(u) - d^+(v) = p - 2$. For a star S_n on n vertices, $d^+(v) = n - 1$ if v is the nonpendent vertex and $d^+(v) = 2(n - 2)$ otherwise.

Definition 1.1. [12] Assuming that $p_u(e) \leq p_v(e)$, two trees T' and T" of order n with the same number of pendent vertices are said to be equiseparable w.r.t terminal Wiener index if their edges $e'_1, e'_2, \dots e'_{n-1}$ and $e''_1, e''_2, \dots e''_{n-1}$ can be labelled such that the equality $p_u(e'_i|T') = p_u(e''_i|T'')$ holds for all $i = 1, 2, \dots n - 1$. $p_u(e'_i|T')$ denotes $p_u(e'_i)$ in T'.

The concept of Wiener index has been extensively used in chemistry. If two or more chemical compounds have equiseparable trees, then those compounds will have similar physico-chemical properties which can not be distinguished by means of indices like Wiener index and terminal Wiener index[12]. Terminal distance matrices were used in the mathematical modelling of proteins and genetic codes[10]. One way of generating equiseparable graphs with respect to Wiener index can be found in [4]. Construction of a class of equiseparable trees with respect to terminal Wiener index can be found in [12]. A detailed study of equiseparable molecules with respect to Wiener index can be found in [5, 7, 8]. It is well known that almost all trees have a terminal equiseparable mate.[12]

Our contribution:

In this work, we propose a new method to generate trees having the same terminal Wiener index. This method is based on distances between vertices. Using this method it is possible to construct many tree pairs with same terminal Wiener index.

This paper is organized as follows: In Section 2 we explain some methods for construction of equiseparable tree pairs with respect to terminal Wiener index only. In Section 3 we explain some methods for constructing large class of trees that are equiseparable with respect to terminal Wiener index.

2 Construction of equiseparable trees with respect to terminal Wiener Index

We use the notation $d^+(u)$ to denote the sum of distances between u and the pendent vertices of Twhere $u \in V(T)$. We start with a lemma given below.

Lemma 2.1. Let T_a and T_b be two trees of orders n_1 and n_2 respectively. Let $u \in V(T_a)$, $v \in V(T_b)$ and $T_a.T_b(u, v)$ denotes the new tree obtained from T_a and T_b by identifying u and v. Let the number of pendent vertices in T_a and T_b be l_1 and l_2 respectively.

If both u and v are nonpendent vertices, then

$$TW(T_a, T_b(u, v)) = TW(T_a) + TW(T_b) + l_2d^+(u) + l_1d^+(v).$$
(6)

The above lemma can be extended to three trees as follows.

Lemma 2.2. Let T_a, T_b and T_c be three trees of orders n_1, n_2 and n_3 respectively. Let $u \in V(T_a)$, $v \in V(T_b), w \in V(T_c)$ and $T_a.T_b.T_c(u, v, w)$ denotes the new tree obtained from T_a , T_b and T_c by identifying u, v and w. Let the number of pendent vertices in T_a , T_b and T_c be l_1, l_2 and l_3 respectively. If u, v and w are nonpendent vertices, then

$$TW(T_a, T_b, T_c(u, v, w)) = TW(T_a) + TW(T_b) + TW(T_c) + (l_2 + l_3)d^+(u) + (l_1 + l_3)d^+(v) + (l_1 + l_2)d^+(w).$$
(7)

Lemma 2.3. Let T be a tree composed of two disjoint trees T_a and T_b of orders n_1 and n_2 respectively. Let $u \in V(T_a)$, $v \in V(T_b)$ and uv be a cut-edge in T. Let the number of leaves in T_a and T_b be l_1 and l_2 respectively. In T, for a vertex $x \in V(T)$ let $d^+(x)$ denote the sum of the distances of the form d(x, l) where l is a leaf in T. Then

$$TW(T) = TW(T_a) + TW(T_b) + l_2 d^+(u) + l_1 d^+(v) + l_1 l_2.$$
(8)

Equiseparability w.r.t the Wiener index and equiseparability w.r.t the terminal Wiener index are entirely different. So, it is useful to find some rules for constructing equiseparable trees w.r.t the terminal Wiener index only. The following theorem can be used to construct equiseparable trees with respect to terminal Wiener index.

Theorem 2.1. ([12]) Let T, X and Y be arbitrary trees, each with more than two vertices. Let T_1 be obtained from T by identifying the vertices u and s and by identifying the vertices v and r. Let T_2 be obtained from T by identifying the vertices u and r, and by identifying the vertices v and s. If $p_x - p_s = p_y - p_r$, then T_1 and T_2 are equiseparable w.r.t terminal Wiener index. p_x and p_y denote the number of pendent vertices of fragments X and Y, respectively. $p_s = 1$ if s is a pendent vertex of X;otherwise it is equal to 0. p_r is defined similar to p_s .




Figure 1: Two trees T_1 and T_2 constructed by method in theorem 2.1

Fig.1 shows the construction of equiseparable trees w.r.t terminal Wiener idex. We will prove that the condition in theorem 2.1 is sufficient but not necessary to generate equiseparable trees w.r.t terminal Wiener index. We extend theorem 2.1 as

Theorem 2.2. Let T, X and Y be arbitrary trees, each with at least three vertices. Let T_3 be obtained from T by identifying the vertices u and s and by identifying the vertices v and r. Let T_4 be obtained from T by identifying the vertices u and r, and by identifying the vertices v and s. If **either** $p_x - p_s =$ $p_y - p_r$ or $d^+(u) = d^+(v)$, then T_3 and T_4 are equiseparable w.r.t terminal Wiener index. p_x, p_y, p_t, p_s and p_r are same as in theorem 2.1.

Proof. By lemma 2.2, we get

$$TW(T_1) = TW(T) + TW(X) + TW(Y) + d^+(s)(p_t + p_y - p_r) + d^+(r)(p_t + p_x - p_s) + (p_x - p_s)(p_y - p_r)d(u, v) + d^+(u)(p_x - p_s) + d^+(v)(p_y - p_r) - p_s d^+(s) - p_r d^+(r)$$
(9)

and

$$TW(T_2) = TW(T) + TW(X) + TW(Y) + d^+(s)(p_t + p_y - p_r) + d^+(r)(p_t + p_x - p_s) + (p_x - p_s)(p_y - p_r)d(u, v) + d^+(u)(p_y - p_r) + d^+(v)(p_x - p_s) - p_s d^+(s) - p_r d^+(r).$$
(10)

where the last two terms in (9) and (10) respectively are correction factors for TW(X) and TW(Y)in case s and r are pendent. From (9) and (10), for $TW(T_1) = TW(T_2)$

 $d^{+}(u)(p_{x} - p_{s}) + d^{+}(v)(p_{y} - p_{r}) = d^{+}(u)(p_{y} - p_{r}) + d^{+}(v)(p_{x} - p_{s}).$ This will be true if either (a) $p_{x} - p_{s} = p_{y} - p_{r} \text{ or (b) } d^{+}(u) = d^{+}(v) \text{ holds.}$

The above theorem shows that it is possible to construct equiseparable trees w.r.t terminal Wiener index based on distance parameters also. Consider the following trees T_3 and T_4 .




Figure 2: Equiseparable trees w.r.t terminal Wiener index constructed by method in theorem 2.2.

In Fig.2 we can see that $p_x - p_s \neq p_y - p_r$ but $TW(T_3) = TW(T_4)$. For vertices u and v, $d^+(u) = d^+(v)$. The following theorem is based on partitioning of integers.

Theorem 2.3. Let T be a tree of order n. Let n_1, n_2 and n_3, n_4 be two different partitions of n. Let T' be tree obtained by connecting some nonpendent vertex of P_{n_1} and P_{n_2} by an edge. Let T'' be tree obtained by connecting some nonpendent vertex of P_{n_3} and P_{n_4} by an edge. Then T' and T'' are equiseparable w.r.t terminal Wiener index.

Proof. Since $TW(P_{n_1}) + TW(P_{n_2}) = TW(P_{n_3}) + TW(P_{n_4}) = n-2$ and $l_1 = l_2 = 2$ and $d^+(u) + d^+(v) = n-2$, by lemma 2.3 both T' and T'' are equiseparable w.r.t terminal Wiener index.

Consider two trees T' and T'' constructed using theorem 2.3.




Figure 3: Two trees equiseparable w.r.t terminal Wiener index.

In fig.3 $n_1 = 4, n_2 = 4, n_3 = 5$ and $n_4 = 3$. It is easy to see that TW(T') = TW(T'') = 22. In the rest of the paper, we use p_t to denote the number of pendent vertices in a tree T.

Theorem 2.4. Let Z be an arbitrary tree; $u \in V(Z)$, tree T' is obtained from T by identifying the vertices u and i and T'' be obtained from T by identifying the vertices u and j. If $d^+(i) = d^+(j)$, then T' and T'' are equiseparable w.r.t terminal Wiener index. See Fig. 4




Figure 4: Two trees T' and T'' equiseparable w.r.t terminal Wiener index.

Proof. By lemma 2.1, we get

$$TW(T') = TW(T) + TW(Z) + d^{+}(i)p_{z} + d^{+}(u)p_{t}$$
(11)

and

$$TW(T'') = TW(T) + TW(Z) + d^{+}(j)p_{z} + d^{+}(u)p_{t}.$$
(12)

where p_t and p_z denote the number of pendent vertices in T and Z respectively. From (11) and (12), for TW(T') = TW(T'') the equality $d^+(i) = d^+(j)$ must be true.

Theorem 2.5. Let T be a tree with vertices $i_1, i_2, ..., i_n$. Let Z_a 's be arbitrary trees for $1 \le a \le t$. Let T_1 be the tree obtained by identifying i_a and u_a in Z_a . T_2 is obtained by identifying $i_a + j$ and u_a in Z_a . If $d^+(i_a) = d^+(i_a + j)$ for every $1 \le a \le t$, then the resulting two trees are equiseparable w.r.t terminal Wiener index. See Fig. 5




Figure 5: Two trees equiseparable w.r.t terminal Wiener index.

Proof. By lemma 2.1, we get

$$TW(T_1) = TW(T) + \sum_{i=1}^{t} TW(Z_i) + \sum_{1 \le a \le t} (p_{z_a}d^+(i_a) + p_td^+(u_a))$$
(13)

and

$$TW(T_2) = TW(T) + \sum_{i=1}^{t} TW(Z_i) + \sum_{1 \le a \le t} (p_{z_a}d^+(i_a + j) + p_td^+(u_a)).$$
(14)

for some j. For $TW(T_1) = TW(T_2)$ it must be true that $d^+(i_a) = d^+(i_a + j)$ for every $1 \le a \le t$ and for some j.

3 Large families of trees with same terminal Wiener index

Let P_{2k} be a path on 2k vertices labelled as $v_1, v_2...v_{2k}$. Let X and Y be two different trees with same number of vertices. Let $s \in V(X)$ and $t \in V(Y)$ such that $p_x - p_s = p_y - p_t$ where p_x and p_y denote the number of pendent vertices in X and Y respectively. $p_s = 1$ if s is a pendent vertex of X; otherwise it is equal to 0. p_t is defined similar to p_s . We can construct a set STW(T,X,Y) of trees equiseparable with respect to terminal Wiener index as follows.

Each element of the set is obtained from a copy of T, k copies of X and k copies of Y. Fragments of X are attached(via their vertices s) to k among the vertices $v_1, v_2...v_{2k}$ of T. Fragments of Y are attached(via their vertices t) to the remaining k vertices among $v_1, v_2...v_{2k}$ of T.

Theorem 3.1. The trees in the set STW(T, X, Y) have the same terminal Wiener index.

Proof. Let T' and T'' be two trees belonging to the set STW(T, X, Y). It is easy to see that $p_u(e_i|T') = p_u(e_i|T'')$ for corresponding edges of X and Y in T' and T'' respectively. Considering the corresponding edges of P_{2k} in T' and T'', we can see that

$$p_u(v_i v_{i+1} | T') = p_u(v_i v_{i+1} | T'') = ik^2(2k - 1)$$

Thus, for every pair of corresponding edges $p_u(e_i|T') = p_u(e_i|T'')$ is true.

Fig. 6 shows two trees T_5 and T_6 constructed from trees T, X and Y using the above method. Here $k = 2, p_s = 1$ and $p_t = 0$.




Figure 6: Two trees equiseparable with respect to terminal Wiener index.

We use the notation T.T'(u, v) to denote the tree obtained from T and T' by identifying the vertices u and v.

Theorem 3.2. Let T be a tree with two vertices u and v such that $d^+(u) = d^+(v)$. Let T' be another tree with a vertex x. Then T.T'(u, x) and T.T'(v, x) are equiseparable w.r.t terminal Wiener index.

Proof. Substituting $T_a = T$, $T_b = T'$ and $d^+(u) = d^+(v)$ in lemma 2.1 we get the result.




Figure 7: Two nonisomorphic trees with the same terminal Wiener index.

In T_8 of Fig. 7, we can note that $d^+(w) = d^+(v) = 13$. Therefore by theorem 3.2 we can construct an infinite number of non-isomorphic tree pairs with the same terminal Wiener index.

Theorem 3.3. Let T_1 and T_2 be two nonisomorphic trees with same terminal Wiener index. Let $u \in V(T_1)$ and $v \in V(T_2)$ be two nonpendent vertices such that $d^+(u) = d^+(v)$. Let T' be another tree with a vertex x. Then $T_1 \cdot T'(u, x)$ and $T_2 \cdot T'(v, x)$ are nonisomorphic trees with same terminal Wiener index.

Proof. By lemma 2.1

$$TW(T_1.T'(u,x)) = TW(T_1) + TW(T') + l_2d^+(u) + l_1d^+(x)$$
 and
$$TW(T_2.T'(v,x)) = TW(T_2) + TW(T') + l_2d^+(v) + l_1d^+(x)$$

Since $TW(T_1) = TW(T_2)$ and $d^+(u) = d^+(v)$ the result follows.

For both the non-isomorphic trees T_7 and T_8 in Fig. 7, we have both $TW(T_7) = TW(T_8) = 57$. We note that in both these trees, $d^+(u) = d^+(w) = 13$. Therefore attaching a path at u in T_7 and w in T_8 will give rise to another tree pair with the same terminal Wiener index. By using different paths we can construct an infinite number of non-isomorphic tree pairs with the same terminal Wiener index.

4 Conclusion

In this work we proposed some new methods to generate equiseparable trees w.r.t terminal Wiener index. Equiseparable trees are mainly used in chemistry to identify molecules with similar properties. Our future work include the study of the following two classes of trees:Trees equiseparable w.r.t Wiener index but not equiseparable w.r.t terminal Wiener index and trees equiseparable w.r.t terminal Wiener index but not equiseparable w.r.t Wiener index.

References

- Meryam Zeryouh, Mohamed El Marraki, Mohamed Essalih Terminal Wiener Index of Star-Tree and Path-Tree, Applied Mathematical Sciences, Vol. 9, 39(2015), 1919 - 1929.
- [2] Y. H. Chen, X. D. Zhang, On Wiener and terminal Wiener indices of trees, MATCH Commun. Math. Comput. Chem. 70 (2013) 591-602.
- [3] A.A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: theory and applications, Acta Appl. Math. 66 (2001) 211-249.
- [4] A. A. Dobrynin, I. Gutman, Boris Furtula, Equiseparable chemical trees, Journal of serbian chemical society 68(2003) 549-555.
- [5] I. Gutman, Boris Furtula, D. Vukicević, Biljana Arsic, Equiseparable molecules and molecular graphs, Indian Journal of Chemistry 43 (2004) 7-10.
- [6] I. Gutman, B. Furtula and M. Petrović, Terminal Wiener index, J. Math. Chem. 46 (2009) 522-531.
- [7] I. Gutman and B. Furtula(2010) A survey on terminal Wiener index, in I. Gutman and B. Furtula(Eds.) Novel Molecular Structure Descriptors - Theory and Applications, Kragujevac, 173-190.
- [8] I. Gutman, B. Furtula and Arsic, Equiseparable chemical trees, J. Serbian . Chem. 68 (2003) 549-555.
- [9] I. Gutman, T. Kortvelyesi, Wiener indices and molecular surfaces, Z. Naturforsch.50a(1995) 669-671.
- [10] Horvat B., Pisanski T., Randić M, Terminal polynomials and star-like graphs, MATCH Commun. Math. Comput. Chem. 60 (2008) 493-512.
- [11] Sujuan Wang, Xiaofeng Guo, Trees with extremal Wiener indices, MATCH Commun. Math. Comput. Chem. 60 (2008) 609-622.
- [12] Xiaotie Deng, Jie Zhang, Equiseparability on terminal Wiener Index, Applied Mathematics Letters, 25 (2012) 580-85.
- [13] Xiaotie Deng, Jie Zhang, Equiseparability on terminal Wiener Index, Algorithmic aspects in information and management, Springer-Verlag, Berlin, (2009) 166-74.