

EXISTENCE OF GINZBURG-LANDAU MINIMIZERS WITH OPTIMAL BOUNDARY DATA AND APPLICATIONS

PAUL LAURAIN, ROMAIN PETRIDES

ABSTRACT. We perform the classical Ginzburg-Landau analysis originated from the celebrated paper by Bethuel, Brezis, Hélein for optimal boundary data. More precisely, we give optimal regularity assumptions on the boundary curve of planar domains and Dirichlet boundary data on them. When the Dirichlet boundary data is the tangent vector field of the boundary curve, our framework allows us to define a natural energy minimizing frame for simply connected domains enclosing Weil-Petersson curves.

The so-called simplified Ginzburg-Landau functional on a domain Ω

$$E_\varepsilon(u) = \frac{1}{2} \int_\Omega |\nabla u|^2 + \frac{1}{4\varepsilon^2} \int_\Omega (1 - |u|^2)^2$$

and all its variations were intensively studied after the celebrated paper by Bethuel, Brezis, Hélein [1], since energies of this type are useful to study phase transition problems. From this seminal work, a lot of attention has been given to the behavior of the critical points: see e.g. Pacard and Rivière [29], Sandier and Serfaty [33], Farina and Mironescu [14], Millot and Pisante [27], Ignat, Nguyen, Slastikov and Zarnescu [21] and the references therein.

One other well known usefulness of this energy is to build kinds of “Dirichlet energy minimizing” maps with singularities adapted to the strong topological constraints given by some geometrical problems. This is our central motivation. For instance, we cannot find any smooth nor H^1 unit vector field extensions with finite energy of maps $\partial\Omega \rightarrow \mathbb{S}^1$ of any topological degree. Therefore, we have to relax the constraint on the set of admissible vector fields in the variational problem. Vector fields do not necessarily have unit norm anymore, but as a compensation, we add a term to the energy which drastically penalizes vector fields that do not have unit norm as $\varepsilon \rightarrow 0$. Classical works by [1], [36], prove the convergence as $\varepsilon \rightarrow 0$ of minimizers of E_ε to harmonic maps (with singularities), that are minimizers of the so-called renormalized energy. In the same spirit, studying the Allen-Cahn equation is a valuable technique to build new minimal surfaces, see [17] and [12].

In the current paper, we aim at weakening the known regularity assumptions on the boundary data for the asymptotic analysis as $\varepsilon \rightarrow 0$ for minimizers of the Ginzburg-Landau energy E_ε on simply connected domains Ω and $u : \Omega \rightarrow \mathbb{R}^2$ satisfying a Dirichlet boundary condition $u = g$ on $\Gamma := \partial\Omega$ for some functions $g : \Gamma \rightarrow \mathbb{S}^1$ with a prescribed degree. Not only the weakenings hold for the regularity of the boundary data g but also for the regularity of the domain Ω . A complete study was given in [14], for data $g \in H^{\frac{1}{2}}(\Gamma)$ such that $\deg(g) = 0$ and \mathcal{C}^1 domains, but to our knowledge, the classical Ginzburg-Landau analysis coming from the originated paper [1] has never been completely given for $\deg(g) \neq 0$ with $g \in H^{\frac{1}{2}}(\Gamma)$, nor with domains Ω with lower regularity than \mathcal{C}^1 .

Before stating our result, we would like to give some motivations for these weakenings of assumptions. One by-product of Ginzburg-Landau theory is to produce $u_* : \Omega \rightarrow S^1$ which is harmonic with $u_* = g$ on the boundary (of course u_* is singular if $\deg(g) \neq 0$). In the special case $g = \tau$ being the tangent unit vector field of the boundary, one can easily produce a family of such harmonic maps by considering a uniformization of the domain. Indeed, see proposition 4.3, if $f_a : \mathbb{D} \rightarrow \Omega$ is a biholomorphic map such that $f(0) = a \in \Omega$, then $u_a = \frac{\partial_\theta f}{|\partial_\theta f|}$ is a desired harmonic map. As we show in proposition 0.3, using the renormalized energy, u_* is the one that maximizes $|f'_a(0)|$. More precisely, the expression of the renormalized energy associated to a uniformization f has the form

$$\int_{\mathbb{D}} \left| \frac{f''}{f'} \right|^2 + 2\pi \ln |f'(0)|,$$

which is closely related to the Loewner energy of the boundary curve, see section 1 for more details. Hence our goal is to make the asymptotic analysis of Ginzburg-Landau work for domains whose boundary curves have the minimal regularity that makes this energy finite. Those curves are known as Weil-Petersson curves are chord-arc curves whose unit tangent vector field lies in $H^{\frac{1}{2}}$ (see theorem-definition due to Bishop). More details, on the geometric motivations of our result are given in section 1.

Coming back to the analysis, we first state an optimal result on the weakest regularity of Ω such that the classical definition of $H^{\frac{1}{2}}(\Gamma)$ makes sense:

Theorem 0.1. *Let Ω be a domain such that $\Gamma := \partial\Omega$ is a chord-arc curve and $g \in H^{\frac{1}{2}}(\Gamma, S^1)$ with $\deg(g) = 1$, then there is $u_\varepsilon \in H^1(\Omega)$ which minimizes, E_ε on the admissible set*

$$\mathcal{A} = \{u \in H^1(\Omega, \mathbb{R}^2); u = g \text{ a.e on } \Gamma\}.$$

Moreover, for any sequence $\varepsilon_k \rightarrow 0$, there are a subsequence $\varepsilon_{k_l} \rightarrow 0$, $a \in \Omega$ and $u_* \in C^\infty(\Omega \setminus \{a\}, S^1) \cap H_{loc}^1(\overline{\Omega} \setminus \{a\})$ such that

- (i) u_* is a harmonic map,
- (ii) $u_* = g$ on $\partial\Omega$,
- (iii) $\deg\left((u_*)|_{\partial B(a, \rho)}\right) = 1$ for $\rho > 0$ small enough,
- (iv) $u_{\varepsilon_{k_l}} \rightarrow u_*$ in $C_{loc}^\infty(\Omega \setminus \{a\})$ as $l \rightarrow +\infty$,
- (v) $u_{\varepsilon_{k_l}} \rightarrow u_*$ in $H_{loc}^1(\overline{\Omega} \setminus \{a\})$ as $l \rightarrow +\infty$,
- (vi) $|u_{\varepsilon_{k_l}}| \rightarrow 1$ in $L_{loc}^\infty(\overline{\Omega} \setminus \{a\})$ as $l \rightarrow +\infty$.

This theorem can be easily generalized to the case $\deg(g) \in \mathbb{Z}^*$, since once the difficulty due to the weak regularity of the data at the boundary is overcome for one bad disk, then it is overcome for all. Hence, Theorems VI.1 and VI.2 of [2] hold true in our setting. In the case $\deg(g) = 0$, it is even easier, since we do not need to remove any singularity and section 3.4 is not necessary. Notice that for \mathcal{C}^1 domains with $H^{\frac{1}{2}}$ data at the boundary, the case $\deg(g) = 0$ was solved in [14].

The assumptions of this theorem are optimal in the sense that the presence of the Dirichlet energy requires the $H^{\frac{1}{2}}$ condition on g , and $H^{\frac{1}{2}}(\Gamma)$ only makes sense if Γ is a chord-arc curve. However, in the proof we only use that the pullback of g by a uniformization of the domain belongs to $H^{\frac{1}{2}}(S^1, S^1)$. Hence, taking this as a definition of $H^{\frac{1}{2}}(\Gamma, S^1)$ and defining $\mathcal{A} = \{v \circ f^{-1}\} + H_0^1(\Omega, \mathbb{R}^2)$, where v is the

harmonic extension of $g \circ f$ for some uniformization $f : \mathbb{D} \rightarrow \Omega$, we can even weaken the regularity of the boundary.

Theorem 0.2. *Let Ω be a quasidisk and $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ in sense of definition 2.3, then the conclusion of Theorem 0.1 holds.*

Indeed, one main idea to handle the low regularity of Ω is to pull back all the problem on the unit disk \mathbb{D} thanks to one uniformization map $f : \mathbb{D} \rightarrow \Omega$. The unit disk is then a pleasant smooth domain, but we have to pay one price : the pullback functions $\tilde{u} = u \circ f$ for $u \in \mathcal{A}$ now belong to the H^1 space associated to the Riemannian metric $|f'(z)|^2 (dx^2 + dy^2)$ which is the weighted space of functions having the following norm finite:

$$\|u\|_{H^1(\Omega)}^2 = \|\tilde{u}\|_{H^1(\mathbb{D}, |f'|^2)}^2 := \int_{\mathbb{D}} \tilde{u}^2 |f'(z)|^2 dz + \int_{\mathbb{D}} |\nabla \tilde{u}|^2 dz .$$

If Ω is a smooth domain, the weight $|f'(z)|^2$ satisfies a Harnack inequality *until the boundary*, but if Ω has low regularity, $|f'(z)|^2$ may go to $+\infty$ or 0 as $|z| \rightarrow 1$ so that the geometry might be wildly different to the Euclidean one close to the boundary. In our setting, we assume that Ω is a quasidisk. It is a sufficient regularity for the domain in order to make the behavior of $|f'(z)|^2$ close to the boundary good enough to perform a Ginzburg-Landau analysis on sequences of minimizers u_ε . One reason is because the shape of disks $\mathbb{D}_r(x)$ centered in $x \in \Omega$ is not too deformed after the pullback by f , even when x is close to Γ . This is because uniformization maps of quasidisks can be extended to quasiconformal maps on \mathbb{C} .

For instance, one striking and fundamental lemma in the classical Ginzburg-Landau analysis is the uniform bound of $\varepsilon |\nabla u_\varepsilon|$ (see Step B.1 [1]) on Ω . It is primordial to control the behavior of bad disks (small disks centered on points x such that $|u_\varepsilon(x)| \leq \frac{1}{2}$). Indeed, this estimate leads to the η -compactness (there is a quantum of energy around points such that $|u_\varepsilon| \leq \frac{1}{2}$), see Lemma 24 of [31]. Classically, this estimate is a consequence of the elliptic equation satisfied by u_ε as a critical point of E_ε :

$$-\Delta u_\varepsilon = \frac{1}{\varepsilon^2} (1 - |u_\varepsilon|^2) u_\varepsilon \text{ in } \Omega .$$

The argument relies on the bound $\|u_\varepsilon\|_\infty \leq 1$ coming from a maximum principle argument and then on a rescaling argument on the equation. However, with low regularity on Ω , we cannot perform any maximum principle argument nor any rescaling argument close to the boundary. We first need to assume that u_ε is a minimizer of the energy in order to have the uniform bound $\|u_\varepsilon\|_\infty \leq 1$, and then, a rescaling argument only allows that $\varepsilon |\nabla u_\varepsilon(x)|$ is uniformly bounded on a set of points $x \in \Omega$ such that $\frac{d(x, \partial\Omega)}{\varepsilon}$ is uniformly lower bounded. The main novelty of our paper (see Proposition 3.3) is that there is $\eta_0 > 0$ such that for the points $x_\varepsilon \in \Omega$ satisfying $d(x_\varepsilon, \partial\Omega) \leq \eta_0 \varepsilon$, we have $|u_\varepsilon(x_\varepsilon)| \geq \frac{1}{2}$. We crucially use the quasidisk assumption here. Thanks to this proposition, the classical ‘‘bad disks’’ cannot intersect the boundary of Ω and the classical steps of asymptotic analysis for Ginzburg-Landau can be managed.

Notice that Theorem 0.1 is only stated for simply connected domains and in Proposition 0.3 below, we give a link between the Ginzburg-Landau problem and the uniformization of domains. However, our analysis provides all the tools for

domains Ω such that $\partial\Omega$ is a disjoint union of chord-arc closed curves, and $H^{\frac{1}{2}}$ data of any degree on these closed curves. Then it would be interesting to give new constructions of uniformizations in the spirit of [41], [22], see also [28].

As already remarked, a uniformization $f : \mathbb{D} \rightarrow \Omega$ defined a harmonic map. Here we give a more precise description in terms of Coulomb frame. We set $p = f(0)$ and we let

$$u = \frac{f_\theta}{|f_\theta|} \circ f^{-1} \text{ and } v = \frac{f_r}{|f_r|} \circ f^{-1}$$

the pushforward in Ω of the normalized angular and radial derivatives of f . Let $\omega = \langle du, v \rangle$ the Cartan form associated to (v, u) . We prove in Proposition 4.3 that this frame is Coulomb and that

$$\star\omega = d(\mu + G_p) \text{ in } \Omega ,$$

where G_p is the Dirichlet Green function with respect to p and that

$$\left| \frac{f''}{f'} \right|^2 = |\nabla(\mu \circ f)|^2 \text{ in } \mathbb{D} \text{ and then } \int_{\mathbb{D}} \left| \frac{f''}{f'} \right|^2 dz = \int_{\Omega} |\nabla\mu|^2 dz .$$

By definition, the last integral is finite if the boundary curve is Weil-Petersson. As briefly explained above, looking at the case $g = \tau$, where τ is the tangent vector field on Γ such that the frame (ν, τ) is a direct frame, where ν is the out-pointing normal of the bounded simply connected domain Ω , the minimizer in Theorem 0.1 satisfies the following result, proved in section 4.3:

Theorem 0.3. *We assume that $\Gamma := \partial\Omega$ is a Weil-Petersson curve. Let $u : \Omega \rightarrow \mathbb{S}^1$ be given by Theorem 0.1 with Dirichlet condition $u = \tau$ on Γ . Then $u = u_a$ is the pushforward on Ω of normalized angular derivative of uniformization maps f_a such that $f_a(0) = a$, where a is the singularity the Ginzburg-Landau problem. Moreover, the uniformization maps f_a such that $f_a(0) = a$ maximize $|f'(0)|$ among all the uniformization maps f .*

From this proposition, we figured out that the renormalized energy of frames built with Ginzburg-Landau minimization is deeply related to the Loewner energy of the domain, see [39] for a detailed presentation about the Loewner Energy. However, in some sense, only a half part of the Loewner energy appears in our construction since the frame is only defined on Ω . Therefore, in [26], Michelat and Wang give some reinterpretation of the Loewner energy of a domain in terms of this new frame energy setting. Their work is complementary to ours.

We also believe that our techniques perfectly adapt to the Ginzburg-Landau problem on surfaces. Some very interesting works are already given on closed immersed surfaces (see [20]). In fact, the present article, with the one of [26], are the first stones to define a renormalized frame energy in the case of surfaces with boundary but also with low regularity, ideally some weak immersion with boundary as the one defined by Rivière for closed surface, see [32].

Acknowledgements : This paper is part of a common project between Michelat and Wang and the two authors of the current paper, on the links between the Loewner energy and the Willmore energy. We would like to thank both of them for all the fruitful discussions on the topic - more precisely, Yilin for her explanation of the Loewner Energy and Bishop's work, and Alexis for our conversations about Ginzburg-Landau - especially, Section 3.4 owes him a lot.

1. MOTIVATIONS

By an idea coming from Chern [11] (see also section 5.4 of [18]), using moving frames in the Cartan formalism is more flexible in order to build conformal coordinates. This point of view is particularly interesting in higher dimension, where we can replace the absence of conformal coordinates by the existence of Coulomb frames. We briefly remind why searching an optimal frame gives rise to conformal coordinates in dimension 2. The main idea consists in building a Coulomb frame in a neighborhood U of a point on a surface Σ , i.e. an orthonormal family $\vec{e}_1, \vec{e}_2 \in \Gamma(TU)$ such that

$$(1.1) \quad d * (\langle \vec{e}_1, d\vec{e}_2 \rangle) = 0.$$

This could be done by minimizing the Dirichlet energy of the frame, assuming that locally the total curvature is small enough. Then we deduce from (1.1) that there is $\lambda \in C^\infty(U)$ such that

$$(1.2) \quad [e^\lambda \vec{e}_1, e^\lambda \vec{e}_2] = 0.$$

We obtain the desired conformal coordinates after integration.

It is important to note that *a priori*, this construction is purely local for two reasons. The first one is that if Σ has some topology then the space of frames can be empty. The second is that to get some control on the frame we need that the total curvature does not exceed a certain level (see [18] and [34] for more details). Of course, on planar domains, there are no such problem. In this case, the minimizing Coulomb frame is given by the trivial frame but it doesn't see any geometry of the domain. However, since the boundary has a natural frame given by a tangent and a normal unit vector field, it seems interesting to consider it as Dirichlet boundary data for the Coulomb frame.

Let us consider a simply connected domain Ω . A tangent unit vector field of $\Gamma := \partial\Omega$ can be seen as a map from Γ to \mathbb{S}^1 of degree one. Then, it does not have any regular extension in Ω which still takes values into \mathbb{S}^1 . Anyway, if we are able to construct a (singular) Coulomb frame with the tangent and exterior normal unit vector fields as Dirichlet boundary data, then by integrating it, we find a conformal map from the disk to our domain, which is nothing but a uniformization of the domain.

From this remark, we observe in proposition 4.3 that one natural *global* moving frame (v, u) is defined by the pushforward of the orthonormal polar coordinates on the disk by a uniformization map $f : \mathbb{D} \rightarrow \Omega$ defined by

$$(1.3) \quad u = \frac{f_\theta}{|f_\theta|} \circ f^{-1} \text{ and } v = \frac{f_r}{|f_r|} \circ f^{-1},$$

where f_θ and f_r denotes respectively the angular and radial derivatives of f . As expected, this frame has a singularity, located at $f(0)$. Note that there is as much frame as uniformization. The advantage of the frame approach is that since the frame is built by minimization then it should give a “best” Coulomb frame and hence a “best” uniformization.

Nevertheless, as already pointed above, the space of smooth (even H^1) frames is empty by some degree obstruction. The problem of finding harmonic extensions of vector fields despite some topological obstruction is exactly the goal of the Ginzburg-Landau Theory. Thanks to the seminal work by Bethuel, Brezis and Hélein, we know that on a smooth simply connected domain Ω there exists $u \in C^\infty(\Omega \setminus \{a_0\})$




FIGURE 1. A generic uniformization




FIGURE 2. The uniformization given by the 'minimizing' frame

such that u is tangent to the boundary and $(-iu, u)$ is a coulomb frame. Moreover a_0 minimizes the renormalized energy

$$W(a) = \lim_{\delta \rightarrow 0} \left(\int_{\Omega \setminus \mathbb{D}_\delta(a)} |\nabla w|^2 - 2\pi \ln \frac{1}{\delta} \right),$$

where w is the canonical harmonic vector field matching with u on the boundary and having its singularity at a . Hence among all the uniformizations, the “best” one is the one such that $f(0) = a_0$. In fact, a_0 is not necessary unique, but it is unique if the domain is not far from a disk, see [9] for the convex case. Since we can conversely associate a harmonic vector field to some uniformization, thanks to (1.3), we can associate the following energy to each uniformization, see proposition 4.20,

$$(1.4) \quad W(f) = \int_{\mathbb{D}} \left| \frac{f''}{f'} \right|^2 + 2\pi \ln |f'(0)|.$$

Then remarking that

$$\int_{\mathbb{D}} \left| \frac{f''}{f'} \right|^2 + 4\pi \ln |f'(0)|$$

is independent of the uniformization, the “best” one can be defined as the one that maximizes $|f'(0)|$. Of course, our method is not the first to determine the “best” conformal representation of a simply connected domain. For instance, in section 4.2 of [15], Flucher looks for an optimal conformal rearrangement by studying the Robin function of the domain. His motivation was the study of extremal functions for the Moser-Trudinger inequality. To our knowledge, it is the first time that the point of view of optimal frames is adopted, especially for domains with very low regularity.

In our construction of a “best” uniformization presented above, we used the classical Ginzburg-Landau theory which is only valid for smooth domains. Having a careful look to the classical proofs, it seems that the regularity of the boundary can be reduced to C^2 but not much less since those proofs make use of the Pohozaev identity which requires a control on the derivative of the boundary data, that is to say here the derivative of the tangent unit vector field.

Hence, our main motivation is to be able to have the existence of a minimizer for the Ginzburg-Landau functional as soon as (1.4) makes sense. In fact (1.4) is very similar to the Loewner energy of a Jordan curve Γ , given by

$$E_L(\Gamma) = \frac{1}{\pi} \left(\int_{\mathbb{D}} \left| \frac{f''}{f'} \right|^2 + \int_{\mathbb{D}} \left| \frac{g''}{g'} \right|^2 + 4\pi \ln \left| \frac{f'(0)}{g'(\infty)} \right| \right),$$

where $f : \mathbb{D} \rightarrow \Omega$, $g : \mathbb{D}^c \rightarrow \Omega^c$ are two uniformization with $\partial\Omega = \Gamma$. It is well-known that the class of domains to consider is the one enclosed by a Weil-Petersson curve (see next section for definition). Those curves can be roughly thought as curves whose tangent unit vector field belongs to $H^{\frac{1}{2}}$.

Finally, to get the most general result for minimizers of the Ginzburg-Landau energy, we decouple the regularity of the boundary $\partial\Omega$ and the regularity of the boundary data g and we are able to prove the existence of a minimizer as soon as the boundary is chord-arc (not necessarily Weil-Petersson) and the boundary data $H^{\frac{1}{2}}$ in a classical sense. Moreover, giving a natural sense to the notion of $H^{\frac{1}{2}}$ maps for quasicircles (see next section), we are able to assume only that the boundary is a quasicircle. It is clear that this result is optimal, since the $H^{\frac{1}{2}}$ assumption cannot

be weakened due to the presence of the Dirichlet energy, and Weil-Petersson curves are quasicircles. We then give the existence of a best uniformization in this weak setting.

To conclude, we want to notice that the uniformization was not exactly our first motivation. As explained in the introduction, this work has started by some discussions with Michelat and Wang on the way to rely quantitatively the Willmore energy to the Loewner Energy. To that purpose, we realized that we need the existence of a minimizer of the Ginzburg-Landau energy on a surface with a Weil-Petersson boundary. The present paper is then devoted to the case of a planar domains. It is interesting in itself since we have some purely Ginzburg-Landau result with optimal regularity and a nice application to the problem of finding a best uniformization. From their side, Michelat and Wang have established some clear link of the frame approach and the Loewner-Energy. Indeed, they prove the existence of a consistent generalization of the frame energy (1.4) on Ω to a new frame energy defined on frames (v_1, u_1) and (v_2, u_2) defined on both sides of the curve Γ viewed on the sphere by stereographic projection, denoted here by $W_{v,u}(\Gamma)$ (see [26] for a precise definition of this frame energy):

Theorem 1.1 (Michelat, Wang [26], Theorem A). *Let $\Gamma \subset \mathbb{S}^2$ be a Weil-Petersson curve. Let Ω_1, Ω_2 be the two connected components of $\Omega \setminus \Gamma$. For $p_1 \in \Omega_1$ and $p_2 \in \Omega_2$, there are harmonic frames $(v_1, u_1) : \Omega_1 \setminus \{p_1\} \rightarrow U\Omega_1$ and $(v_2, u_2) : \Omega_2 \setminus \{p_2\} \rightarrow U\Omega_2$ such that for conformal maps $f_1 : \mathbb{D} \rightarrow \Omega_1$ and $f_2 : \mathbb{D} \rightarrow \Omega_2$ such that $f_1(0) = p_1$ and $f_2(0) = p_2$, then*

$$E_L(\Gamma) = \frac{1}{\pi} W_{v,u}(\Gamma) + 2 \ln |\nabla f_1(0)| + 2 \ln |\nabla f_2(0)| - 12 \ln(2),$$

Where $W_{v,u}(\Gamma)$ is the renormalized energy of the frames (v_1, u_1) and (v_2, u_2) .

2. NOTATIONS AND DEFINITIONS

- \mathbb{D} denotes the open unit disk of the complex plane \mathbb{C} identified with \mathbb{R}^2 .
- $\mathbb{D}_r = r\mathbb{D}$ and $\mathbb{D}_r(x)$ the open disk centered at x with radius r .
- $S_r = \partial\mathbb{D}_r$
- $A_{r_1, r_2} = \mathbb{D}_{r_2} \setminus \mathbb{D}_{r_1}$
- A^c is the complement of a given set A .

We remind some definition about quasiconformal maps, see [19] for more details.

Theorem-Definition 2.1 (quasiconformal maps). *Let U, V two open sets of \mathbb{C} and $K \geq 1$. A mapping $f : U \rightarrow V$ is K -quasiconformal if it is a homeomorphism and*

- i) $f \in H_{loc}^1(U)$,
- ii) $\left| \frac{\partial f}{\partial \bar{z}} \right| \leq \frac{K-1}{K+1} \left| \frac{\partial f}{\partial z} \right|$ almost everywhere.

It is equivalent to be L -quasisymmetric of modulus η that is to say, there exists $\eta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a homeomorphism (depending only on K) such that for any three distinct points $x, y, z \in U$ we have

$$\left| \frac{f(x) - f(y)}{f(x) - f(z)} \right| \leq \eta \left(\left| \frac{x - y}{x - z} \right| \right).$$

Definition 2.1 (quasidisk, quasicircle). *A quasidisk is the image of \mathbb{D} by a quasiconformal map from \mathbb{C} into itself. A quasicircle is the boundary of a quasidisk.*

Quasicircle can be very rough since it can even be a Julia set. Nevertheless, it can't be any Jordan curves since the cross-ratio must be bounded as shown by the following proposition

Proposition 2.1. *A Jordan curve $\Gamma \subset \mathbb{C}$ containing 1 is a quasicircle if and only if there exists a constant C such that for any $z_1, z_2, z_3 \in \Gamma$ that appears in that order¹ on Γ , we have*

$$(2.1) \quad |z_1 - z_2| \leq C|z_1 - z_3|.$$

We will make a central use of the following theorem which permits to extend uniformization of quasidisks to quasiconformal maps from \mathbb{C} into itself.

Theorem 2.1. *Any uniformization map from \mathbb{D} to a quasidisk admits a K -quasiconformal extension from \mathbb{C} to itself.*

Now we are restricting our attention to a special class of quasicircles, the class of Weil-Petersson curves. Recently Bishop, see [5], gives no less than 26 equivalent definitions for Weil-Petersson curves. The most adapted one to our setting is the following, which makes him said that 'roughly speaking, Weil-Petersson curves are to L^2 as quasicircles are to L^∞ '.

Theorem-Definition 2.2 (Definition 1 of [5]). *A Jordan curve Γ enclosing a domain Ω is Weil-Petersson if it is a quasicircle and it admits a uniformization map $f : \mathbb{D} \rightarrow \Omega$ such that*

$$\int_{\mathbb{D}} \left| \frac{f''}{f'} \right|^2 dz < \infty.$$

In particular it is rectifiable and a chord-arc curve, i.e. there exists C such that for any subarc $\gamma \subset \Gamma$, we have

$$l(\gamma) \leq C|x - y|,$$

where x, y are the end points of γ .

Definition 2.2 ($H^{\frac{1}{2}}$ maps for chord-arc curves). *Let $n \geq 1$ and Γ be a chord-arc curves, in particular rectifiable. If $g : \Gamma \rightarrow \mathbb{R}^n$ satisfies*

$$\int_{\Gamma} \int_{\Gamma} \left| \frac{g(x) - g(y)}{|x - y|} \right|^2 d\mu(x) d\mu(y) < \infty,$$

where μ is the measure of Γ , we say that $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{R}^n)$ and we denote $\|g\|_{H^{\frac{1}{2}}(\Gamma)}$ the left-hand side of the previous inequality. In particular, $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ if $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{R}^2)$ and $g(x) \in \mathbb{S}^1$ for a.e. $x \in \Gamma$.

In the previous definition we could have only assumed that Γ is rectifiable, but we will need the following important proposition. It is due to Bishop, see page 16 of [5], and it relies on the fact that quasymmetric maps preserve the $H^{\frac{1}{2}}$ -norm by Ahlfors and Beurling, see page 129 of [3].

Theorem 2.2. *Let Γ be a chord-arc curve enclosing a domain Ω and $f : \mathbb{D} \rightarrow \Omega$ be a uniformization, then there exists $C > 0$ depending only on Γ , such that for every $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{R}^n)$ we have*

$$\frac{1}{C} \|g\|_{H^{\frac{1}{2}}(\Gamma)} \leq \|g \circ f\|_{H^{\frac{1}{2}}(\mathbb{S}^1)} \leq C \|g\|_{H^{\frac{1}{2}}(\Gamma)}.$$

¹Which makes sense since $1 \in \Gamma$.

This theorem leads us to the following definition of $H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ maps

Definition 2.3 ($H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ maps). *Let Γ be a quasicircle enclosing a domain Ω , and $f : \mathbb{D} \rightarrow \Omega$ a uniformization. We say that $g : \Gamma \rightarrow \mathbb{S}^1$ is in $H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ if $g \circ f \in H^{\frac{1}{2}}(\mathbb{S}^1, \mathbb{S}^1)$ and we set $\|g\|_{H^{\frac{1}{2}}(\Gamma)} = \|g \circ f\|_{H^{\frac{1}{2}}(\mathbb{S}^1)}$.*

Notice that there are subtleties with this definition. Indeed, let v be the harmonic extension of $g \circ f \in H^{\frac{1}{2}}(\mathbb{S}^1, \mathbb{S}^1)$. Then, knowing both $v \in H^1(\mathbb{D}, \mathbb{R}^2)$ and $v \circ f^{-1} \in H^1(\Omega, \mathbb{R}^2)$ makes our definition consistent. We explain why in the appendix B.

Now we define the degree of a weak maps, see [7] and the appendix of [6].

Theorem 2.3. *Let $g \in H^{\frac{1}{2}}(\mathbb{S}^1, \mathbb{S}^1)$, then there exists an integer n and $\phi \in H^{\frac{1}{2}}(\mathbb{S}^1, \mathbb{R})$, unique up to a multiple of 2π , such that $g = z^n e^{i\phi}$. The integer n is called the degree of g , and is denoted $\deg(g)$, moreover we have²*

$$(2.2) \quad \deg(g) = \frac{1}{2\pi} \int_{\mathbb{S}^1} g \wedge g_\theta d\theta.$$

Thanks to Theorem 2.2, we have

Theorem-Definition 2.3. *For Γ a chord-arc curve and $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$, we define $\deg(g, \Gamma) = \deg(g \circ f)$, where f is any uniformization map of the enclosing domain of Γ .*

In particular, see property 5 and corollary 8 of [7], we have

Theorem 2.4. *Let Ω a domain whose boundary consists of a finite number of quasicircle $(\Gamma_i)_{i=1}^k$, then if $u \in H^1(\Omega, \mathbb{S}^1)$ then*

$$\sum_{i=1}^k \deg(u|_{\Gamma_i}) = 0.$$

In particular, for $a \in \Omega$ and $u \in H^1(\Omega \setminus \{a\}, \mathbb{S}^1)$, we set $\deg(u, a) = \deg(u|_{\partial B(a,r)})$ for $r > 0$ small enough.

3. PROOF OF THEOREM 0.1

3.1. Upper and lower log estimates for the energy. Let $u_\varepsilon \in \mathcal{A}$ be such that $E_\varepsilon(u_\varepsilon) = \inf_{u \in \mathcal{A}} E_\varepsilon(u)$. Such a minimizer exists because the functional E_ε is coercive. Since u_ε comes from a minimization problem, by a test function argument, we have the following upper bound on the energy, see Theorem III.1 of [2]. There exists $D_0 > 0$ such that, for any $\varepsilon > 0$,

$$(3.1) \quad E_\varepsilon(u_\varepsilon) \leq \pi |\ln \varepsilon| + D_0.$$

Then we want to prove the following key estimate,

$$(3.2) \quad \frac{1}{4\varepsilon^2} \int_{\Omega} (1 - |u_\varepsilon|^2)^2 dx \leq C$$

for some constant C independent from ε , depending only on Ω and g .

In their seminal work Bethuel, Brezis and Hélein proved this estimate using a Pohožaev identity, but in our case since the boundary data is only $H^{\frac{1}{2}}$ we can't apply this idea.

²This formula is well defined since $g \in H^{\frac{1}{2}}$ and $g_\theta \in H^{-\frac{1}{2}}$.

Our argument, inspired from [13], is based on an extension of the sequence of minimizers $u_\varepsilon : \Omega \rightarrow \mathbb{R}^2$ to some bigger smooth domain $\Omega' \supset \Omega$ whose boundary data is smooth on $\partial\Omega'$.

Lemma 3.1 (extension lemma). *Let Ω a domain with a quasicircle as boundary and $g \in H^{\frac{1}{2}}(\partial\Omega, \mathbb{S}^1)$ there is a smooth simply connected domain Ω' such that $\overline{\Omega} \subset \Omega'$ and a function $\hat{u} : \Omega' \setminus \Omega \rightarrow \mathbb{R}^2$ such that*

- $\hat{u} = g$ on $\partial\Omega$,
- $\hat{u} \in C^\infty(\overline{\Omega'} \setminus \overline{\Omega})$,
- $\hat{u} \in H^1(\Omega' \setminus \Omega)$ and whose energy depends only on Ω and $\|g\|_{H^{\frac{1}{2}}}$,
- $|\hat{u}| = 1$ on $\overline{\Omega'} \setminus \Omega$,
- $\deg(\hat{u}|_{\partial\Omega'}) = \deg(g)$.

Proof. Let $h : \mathbb{D}^c \rightarrow \Omega^c$ be a uniformization map, since Ω is a quasidisk, thanks to Theorem 2.1, h admits a quasiconformal extension from \mathbb{C} to itself, we still denote h . We set $\tilde{g} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ by $\tilde{g} = g \circ h$, by conformal invariance, see Theorem 2.2, \tilde{g} is in $H^{\frac{1}{2}}(\mathbb{S}^1, \mathbb{S}^1)$. Let $\tilde{u} \in H^1(\mathbb{D}^c) \cap C^\infty(\mathbb{D}^c, \mathbb{D})$ be the harmonic extension of \tilde{g} , finally we set $v = \tilde{u} \circ h^{-1}$. Since h and \tilde{u} are smooth outside $\overline{\mathbb{D}^c}$ then v is smooth on $\overline{\Omega}^c$. Since h is conformal on $\overline{\mathbb{D}^c}$, the energy of v is bounded and depends only on $\|\tilde{g}\|_{H^{\frac{1}{2}}}$, i.e on Ω and g by Theorem 2.2. Finally, thanks to Lemma A.2, there exists $\delta > 0$ such that $\frac{3}{4} \leq |\tilde{u}| \leq \frac{5}{4}$ on $\mathbb{D}_{1+\delta} \setminus \mathbb{D}$. Then $\hat{u} = \frac{v}{|v|}$ satisfies the desired properties on $\Omega' = g(\mathbb{D}_{1+\delta})$ since it has the same property than v and the last property is given by Theorem 2.4. \square

Then, thanks to Lemma 3.1, the function $\hat{u}_\varepsilon : \Omega' \rightarrow \mathbb{R}^2$ defined as an extension of u_ε such that

$$\hat{u}_\varepsilon = u_\varepsilon \text{ in } \Omega \text{ and } \hat{u}_\varepsilon = \hat{u} \text{ in } \Omega' \setminus \Omega$$

can be used as a test function for the classical Ginzburg-Landau problem on the smooth domain Ω' , minimizing the energy

$$E_\varepsilon^{\Omega'}(v) = \frac{1}{2} \int_{\Omega'} |\nabla v|^2 + \frac{1}{4\varepsilon^2} \int_{\Omega'} (1 - |v|^2)^2$$

on the set of admissible functions $\mathcal{A}^{\Omega'} = \{v \in W^{1,2}(\Omega'); v = \hat{u} \text{ on } \partial\Omega'\}$. We recall that \hat{u} is a smooth function on $\partial\Omega'$. Since

$$E_{2\varepsilon}(u_\varepsilon) = E_{2\varepsilon}^{\Omega'}(\hat{u}_\varepsilon) - \frac{1}{2} \int_{\Omega' \setminus \Omega} |\nabla \hat{u}|^2 \geq \inf_{v \in \mathcal{A}^{\Omega'}} E_{2\varepsilon}^{\Omega'}(v) - \frac{1}{2} \int_{\Omega' \setminus \Omega} |\nabla \hat{u}|^2$$

and by the classical lower bound on the energy, see estimate (1.9) in [13], we finally get

$$(3.3) \quad E_{2\varepsilon}(u_\varepsilon) \geq \pi |\ln \varepsilon| - D_1$$

for some constant D_1 depending only on Ω and τ .

Now, using (3.1) and (3.3) we obtain

$$\frac{1}{4\varepsilon^2} \int_{\Omega} (1 - |u_\varepsilon|^2)^2 = \frac{4}{3} (E_\varepsilon(u_\varepsilon) - E_{2\varepsilon}(u_\varepsilon)) \leq \frac{4}{3} (D_0 + D_1).$$

Setting $C = \frac{4}{3} (D_0 + D_1)$ ends the proof of (3.2).

3.2. A priori estimates for sequences of minimizers. In this section we prove two fundamental estimates, the first one, proposition 3.3, asserts that u_ε can't be too small near the boundary, and the second one, proposition , is known as η -compactness. Their proof does not require that u_ε is a minimizer but they only need as assumption the uniform boundness of u_ε . This boundness is an easy consequence of the maximum principle in the smooth case, but in our weak setting we are not able to prove it without assuming that the u_ε is a minimizer. Which leads us to ask the following question:

Open question : are there sequences of critical point of E_ε with fixed boundary data in $H^{\frac{1}{2}}(\partial\Omega, \mathbb{S}^1)$ which are not uniformly bounded ?

Any critical point u_ε of E_ε satisfies the Euler-Lagrange equation

$$(3.4) \quad \begin{cases} -\Delta u_\varepsilon = \frac{1}{\varepsilon^2} (1 - |u_\varepsilon|^2) u_\varepsilon & \text{in } \Omega \\ u_\varepsilon =_{a.e} g & \text{on } \partial\Omega \end{cases}$$

and we have those very first estimates on u_ε , which are true as soon as $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$. In fact in [14] they assume that boundary is Lipschitz but the argument works as soon as we have a minimizer.

Proposition 3.1 (Lemma 3.2 of [14]). *If u_ε is a minimizer of E_ε , then*

$$(3.5) \quad \|u_\varepsilon\|_{L^\infty(\Omega)} \leq 1.$$

Notice that (3.5) is true for any critical solution u_ε to E_ε if Ω is smooth enough to apply the maximum principle argument, see proposition 2 of [1]. The next proposition is a classical elliptic estimate true independently of the regularity of Ω .

Proposition 3.2 (Lemma A.1 [1]). *There is a constant $C_1 > 0$ such that for any $\varepsilon > 0$, and for any $x \in \Omega$,*

$$(3.6) \quad |\nabla u_\varepsilon|(x) \leq C_1 \|u_\varepsilon\|_\infty \left(\frac{1}{\varepsilon} + \frac{1}{\text{dist}(x, \partial\Omega)} \right).$$

In particular, if (3.5) holds, $\|u_\varepsilon\|_\infty$ can be removed.

In the following proposition, we prove that if u_ε is uniformly bounded then $|u_\varepsilon(x)|$ has to be far from 0 as soon as x is close to $\partial\Omega$ at the scale ε . This hypothesis of uniform boundness of u_ε is satisfied in the case of minimizers thanks to proposition 3.1.

Proposition 3.3. *Let u_ε be solutions of (3.4) which are uniformly bounded with respect to ε , then there are $\varepsilon_0, \eta_0 > 0$ depending only on $\sup_\varepsilon \|u_\varepsilon\|_\infty$, Ω and g such that for any $0 < \varepsilon \leq \varepsilon_0$, for any $x \in \Omega$*

$$(3.7) \quad \text{dist}(x, \partial\Omega) \leq \eta_0 \varepsilon \Rightarrow |u_\varepsilon(x)| \geq \frac{1}{2}.$$

Proof. First of all, let us remind that thanks to our assumption and (3.2), there exists $C > 0$ such that

$$\frac{1}{4\varepsilon^2} \int_\Omega (1 - |u_\varepsilon|^2)^2 dx \leq C \text{ and } \|u_\varepsilon\|_\infty \leq C.$$

We aim at finding η_0 and ε_0 such that for any $0 < \varepsilon \leq \varepsilon_0$ such that for any $y_\varepsilon \in \partial\Omega$,

$$(3.8) \quad \inf_{x \in D(y_\varepsilon, \eta_0 \varepsilon) \cap \Omega} |u_\varepsilon(x)| \geq \frac{1}{2}.$$

Let $f : \mathbb{D} \rightarrow \Omega$ be a uniformization of Ω . Since $\Gamma = \partial\Omega$ is a quasicircle, there is a K -quasiconformal extension $f : \mathbb{C} \rightarrow \mathbb{C}$ of f , still denoted f . We let $v_\varepsilon = u_\varepsilon \circ f : \mathbb{D} \rightarrow \mathbb{R}^2$ be the pullback of u_ε by f . We have by conformal invariance, and because $u_\varepsilon \in H^{\frac{1}{2}}(\Gamma)$ that

$$\begin{cases} -\Delta v_\varepsilon = |f'|^2 \frac{1-|v_\varepsilon|^2}{\varepsilon^2} v_\varepsilon & \text{in } \mathbb{D} \\ v_\varepsilon = g \circ f \in H^{\frac{1}{2}}(\mathbb{S}^1) & \text{on } \mathbb{S}^1 \end{cases}$$

Let $\lambda_0 > 0$ we shall fix later, we set $D_\varepsilon = f^{-1}(\mathbb{D}_{\lambda_0 \varepsilon}(y_\varepsilon) \cap \Omega)$. We split $v_\varepsilon = a_\varepsilon + b_\varepsilon + h$ so that

$$\begin{cases} -\Delta a_\varepsilon = \mathbf{1}_{D_\varepsilon} |f'|^2 \frac{1-|v_\varepsilon|^2}{\varepsilon^2} v_\varepsilon & \text{in } \mathbb{D} \\ a_\varepsilon = 0 & \text{on } \mathbb{S}^1 \\ -\Delta b_\varepsilon = (1 - \mathbf{1}_{D_\varepsilon}) |f'|^2 \frac{1-|v_\varepsilon|^2}{\varepsilon^2} v_\varepsilon & \text{in } \mathbb{D} \\ b_\varepsilon = 0 & \text{on } \mathbb{S}^1 \\ -\Delta h = 0 & \text{in } \mathbb{D} \\ h = v_\varepsilon = g \circ f & \text{on } \mathbb{S}^1 \end{cases}$$

Claim 1: For λ_0 small enough $\|a_\varepsilon\|_\infty \leq \frac{1}{8}$.

By Lemma C.1, we have

$$(3.9) \quad \|a_\varepsilon\|_{L^\infty} + \|\nabla a_\varepsilon\|_{L^2} \leq C_0 \left(\left\| (1 - |z|^2)^2 \Delta a_\varepsilon \right\|_{L^\infty} + \|\Delta a_\varepsilon\|_{L^1} \right).$$

We aim at estimating the right-hand terms. By Koebe distortion theorem, see Corollary 1.4 [30], we have that

$$(3.10) \quad \frac{\text{dist}(f(z), \partial\Omega)}{1 - |z|^2} \leq |f'(z)| \leq 4 \frac{\text{dist}(f(z), \partial\Omega)}{1 - |z|^2}.$$

In particular, if $z \in D_\varepsilon$,

$$(3.11) \quad |f'(z)|^2 \frac{1 - |v_\varepsilon(z)|^2}{\varepsilon^2} |v_\varepsilon(z)| \leq 4C \frac{\lambda_0^2}{(1 - |z|^2)^2}.$$

We also have that, by change of variables,

$$(3.12) \quad \begin{aligned} \int_{D_\varepsilon} |f'|^2 \frac{1 - |v_\varepsilon|^2}{\varepsilon^2} |v_\varepsilon| dz &= \int_{f(D_\varepsilon)} \frac{1 - |u_\varepsilon|^2}{\varepsilon^2} |u_\varepsilon| dz \\ &\leq \left(\int_{\Omega} \frac{(1 - |u_\varepsilon|^2)^2}{\varepsilon^2} dz \right)^{\frac{1}{2}} \left(C \frac{|f(D_\varepsilon)|}{\varepsilon^2} \right)^{\frac{1}{2}} \\ &\leq 2C \sqrt{2\pi} \lambda_0. \end{aligned}$$

Combining (3.9), (3.11) and (3.12), we obtain

$$(3.13) \quad \|a_\varepsilon\|_{L^\infty} + \|\nabla a_\varepsilon\|_{L^2} \leq C \left(4\lambda_0^2 + 2\sqrt{2\pi} \lambda_0 \right).$$

Choosing λ_0 small enough such that $C(4\lambda_0^2 + 2\sqrt{2\pi}\lambda_0) \leq \frac{1}{8}$, we obtain that

$$(3.14) \quad \|a_\varepsilon\|_{L^\infty} \leq \frac{1}{8}.$$

Claim 2: There exists $r_0 > 0$ such that $\|b_\varepsilon\|_{L^\infty(f^{-1}(\mathbb{D}_{r_0\lambda_0\varepsilon}(y_\varepsilon)\cap\Omega))} \leq \frac{1}{8}$.

Let

$$\delta_\varepsilon = \frac{\text{dist}(f^{-1}(y_\varepsilon), \partial(f^{-1}(\mathbb{D}_{\lambda_0\varepsilon}(y_\varepsilon))))}{2},$$

then

$$(3.15) \quad \mathbb{D}_{\delta_\varepsilon}(f^{-1}(y_\varepsilon)) \subset f^{-1}(\mathbb{D}_{\lambda_0\varepsilon}(y_\varepsilon))$$

We set $\tilde{b}_\varepsilon(z) = b_\varepsilon(f^{-1}(y_\varepsilon) + \delta_\varepsilon z)$. Then, let $B_\varepsilon = \frac{\mathbb{D}_{\delta_\varepsilon}(f^{-1}(y_\varepsilon))}{\delta_\varepsilon} \cap \mathbb{D}$, we have that

$$\begin{cases} \Delta \tilde{b}_\varepsilon = 0 & \text{in } B_\varepsilon \\ \tilde{b}_\varepsilon = 0 & \text{on } \partial B_\varepsilon \cap \mathbb{D}. \end{cases}$$

By standard elliptic estimates, there is a constant $D > 0$ independent from $\varepsilon > 0$ such that

$$(3.16) \quad \|\nabla \tilde{b}_\varepsilon\|_{L^\infty(B_\varepsilon \cap \mathbb{D}_{\frac{1}{2}})} \leq D \|\tilde{b}_\varepsilon\|_{L^\infty(B_\varepsilon)}.$$

We also know by (3.14), the fact that v_ε is bounded and that $h \in \mathbb{D}$ as harmonic extension of a \mathbb{S}^1 valued function, that

$$(3.17) \quad \|\tilde{b}_\varepsilon\|_{L^\infty(B_\varepsilon)} \leq \|b_\varepsilon\|_{L^\infty} \leq \|v_\varepsilon\|_{L^\infty} + \|a_\varepsilon\|_{L^\infty} + \|h\|_{L^\infty} \leq C + \frac{1}{8} + 1 \leq C + 2.$$

In particular, since we have that $\tilde{b}_\varepsilon = 0$ on $\partial B_\varepsilon \cap \mathbb{D}$, thanks to (3.16) and (3.17), there is a radius $0 < \rho_0 < \frac{1}{2}$ such that

$$(3.18) \quad \|b_\varepsilon\|_{L^\infty(\mathbb{D} \cap \mathbb{D}_{\rho_0\delta_\varepsilon}(f^{-1}(y_\varepsilon)))} = \|\tilde{b}_\varepsilon\|_{L^\infty(B_\varepsilon \cap \mathbb{D}_{\rho_0})} \leq \frac{1}{8}.$$

Finally since f^{-1} is also quasiconformal, see Corollary 4.5.10 of [19], thanks to Theorem 2.1, for every $x \in \partial\mathbb{D}_{\lambda_0\varepsilon}(y_\varepsilon)$ and $y \in \mathbb{D}_{r_0\lambda_0\varepsilon}(y_\varepsilon)$ we have

$$\frac{|f^{-1}(y) - f^{-1}(y_\varepsilon)|}{|f^{-1}(x) - f^{-1}(y_\varepsilon)|} \leq \eta(r_0)$$

which gives

$$|f^{-1}(y) - f^{-1}(y_\varepsilon)| \leq \eta(r_0)(2\eta(1)\delta_\varepsilon)$$

hence there is a constant $r_0 > 1$, smaller than 1, depending only on the quasiconformal constant K of f^{-1} , that is to say only on Ω , such that

$$(3.19) \quad f^{-1}(\mathbb{D}_{r_0\lambda_0\varepsilon}(y_\varepsilon) \cap \Omega) \subset \mathbb{D}_{\rho_0\delta_\varepsilon}(f^{-1}(y_\varepsilon))$$

hence (3.18) and (3.19) give that

$$(3.20) \quad \|\tilde{b}_\varepsilon\|_{L^\infty(f^{-1}(\mathbb{D}_{r_0\lambda_0\varepsilon}(y_\varepsilon)\cap\Omega))} \leq \frac{1}{8},$$

which achieves the proof of claim 2.

Now, we set $\eta_0 = r_0\lambda_0$. To complete the proof it suffices to prove that $\|h\|_{L^\infty(f^{-1}(\mathbb{D}_{\eta_0\varepsilon}(y_\varepsilon)\cap\Omega))} \geq \frac{3}{4}$. Notice that the $H^{\frac{1}{2}}(\mathbb{S}^1)$ norm of $h = v_\varepsilon = u_\varepsilon \circ f = g \circ f$

is independent of ε , hence the result follows because h is uniformly proper by Lemma A.2. \square

The following proposition is an easy consequence of Proposition 3.2 and Proposition 3.3.

Proposition 3.4. *Let u_ε be solutions of (3.4) which are uniformly bounded with respect to ε , then there are $\varepsilon_0, \eta_0, \delta_0 > 0$ depending only on $\sup_\varepsilon \|u_\varepsilon\|_\infty$, Ω and g such that for any $0 < \varepsilon \leq \varepsilon_0$, for any $x \in \Omega$*

$$(3.21) \quad |u_\varepsilon(x)| \leq \frac{1}{2} \Rightarrow \frac{1}{4\varepsilon^2} \int_{\mathbb{D}_{\eta_0\varepsilon}(x)} (1 - |u_\varepsilon|^2)^2 dx \geq \delta_0.$$

3.3. Weak convergence to u_* . From now to the end of the subsection, we follow the lines of part 3 in [36]. First, the following claim is a very simple consequence of (3.2), Lemma 3.1 and Proposition 3.4.

Claim 3.1. *There are $\eta_0 > 0$ and $J_0 \in \mathbb{N}$ depending only on Ω, g such that for any disjoint collection of balls $(\mathbb{D}_{\eta_0\varepsilon}(x_j^\varepsilon))_{1 \leq j \leq J}$ such that $|u_\varepsilon(x_j^\varepsilon)| \leq \frac{1}{2}$, then $J \leq J_0$.*

By a classical Vitali's covering lemma, we can find a family of disjoint balls $(\mathbb{D}_{\eta_0\varepsilon}(x_j^\varepsilon))_{1 \leq j \leq J_\varepsilon}$ such that $|u_\varepsilon(x_j^\varepsilon)| \leq \frac{1}{2}$ for any $1 \leq j \leq J_\varepsilon$ and

$$\left\{ x \in \Omega; |u_\varepsilon(x)| \leq \frac{1}{2} \right\} \subset \bigcup_{1 \leq j \leq J_\varepsilon} \mathbb{D}_{5\eta_0\varepsilon}(x_j^\varepsilon).$$

By Claim 3.1, $J_\varepsilon \leq J_0$ and for some radius $r > 0$, we denote by

$$\Omega'_{\varepsilon,r} = \Omega' \setminus \bigcup_{1 \leq j \leq J_\varepsilon} \mathbb{D}_r(x_j^\varepsilon) \text{ and } \Omega_{\varepsilon,r} = \Omega'_{\varepsilon,r} \cap \Omega$$

where we recall that Ω' is the extension of Ω given by Lemma 3.1. $\Omega'_{\varepsilon,r}$ is defined for technical reasons since we have the property $\bigcup_{1 \leq j \leq J_\varepsilon} \mathbb{D}_r(x_j^\varepsilon) \subset \Omega'$ for r small enough.

Notice that by definition, $|u_\varepsilon(x)| \geq \frac{1}{2}$ for $x \in \Omega_{\varepsilon,r}$ if $r \geq 5\nu_0\varepsilon$ and that it is also true for the extension \hat{u}_ε given by Lemma 3.1 then:

$$|\hat{u}_\varepsilon(x)| \geq \frac{1}{2} \text{ for } x \in \Omega'_{\varepsilon,r} \text{ if } r \geq 5\nu_0\varepsilon$$

Let us remind the following general proposition originally due to Brezis, Merle and Rivière [8].

Proposition 3.5 ([36] Proposition 3.4 and [37]). *Let $0 < r_0 \leq r_1 \leq r_2$ and $C > 0$. Then for any $v \in H^1(A_{r_0,r_2})$ such that $\frac{1}{2} \leq |v| \leq 1$ a.e on A_{r_0,r_1} and such that v has degree d on every circle S_r for $r_1 \leq r \leq r_2$, and such that*

$$\frac{1}{r_1^2} \int_{A_{r_0,r_2}} (1 - |v|^2)^2 \leq C \text{ and } \int_{A_{r_0,r_2}} |\nabla v|^2 \leq C |\ln r_0|,$$

then for any $r_1 \leq r \leq r_2$,

$$\int_{A_{r_1,r}} |\nabla v|^2 \geq 2\pi d^2 \left(\ln \left(\frac{r}{r_1} \right) - D \right) - D$$

where D only depends on C .

The next proposition is a consequence of (3.1), Proposition 3.5 applied on the extended function \hat{u}_ε on an accurate disjoint union of annuli centered at points in the set $(x_j^\varepsilon)_{1 \leq j \leq J_\varepsilon}$ almost filling $(\Omega')_{\varepsilon, r}$ and the fact that the degree of \hat{u}_ε is 1 on $\partial\Omega'$.

Proposition 3.6 ([36] Proposition 3.3). *There are constants $\kappa > 0$, C depending only on Ω and g such that for any $\varepsilon < r < \delta$,*

$$(3.22) \quad \int_{\Omega'_{\varepsilon, r}} |\nabla u_\varepsilon|^2 \leq 2\pi |\ln r| + C$$

Claim 3.2 (Remark that follows Proposition 3.3 of [36]). *Let $\varepsilon_k \rightarrow 0$ be a sequence, Then there is a subsequence ε_{k_l} such that $(J_{\varepsilon_{k_l}})_{l \in \mathbb{N}}$ is constant and such that for any $1 \leq j \leq J := J_{\varepsilon_{k_l}}$ there is $a_j \in \bar{\Omega}$ such that $x_j^{\varepsilon_{k_l}} \rightarrow a_j$ as $l \rightarrow +\infty$. We denote*

$$S = \{a_j; 1 \leq j \leq J\}$$

Moreover there is $a \in S$ and $u_\star \in H_{loc}^1(\bar{\Omega} \setminus \{a\}) \cap C^\infty(\Omega \setminus \{a\})$ such that $u_{\varepsilon_{k_l}} \rightarrow u_\star$ in $C_{loc}^\infty(\Omega \setminus S)$ as $l \rightarrow +\infty$. Moreover, we have that

$$(3.23) \quad \int_{\Omega \setminus \mathbb{D}_r(a)} |\nabla u_\star|^2 \leq 2\pi \ln\left(\frac{1}{r}\right) + \kappa.$$

$$(3.24) \quad \int_{\mathbb{D}_{r_2}(a) \setminus \mathbb{D}_{r_1}(a)} |\nabla u_\star|^2 \leq 2\pi \ln\left(\frac{r_2}{r_1}\right) + \kappa'.$$

and that u_\star is a harmonic map into \mathbb{S}^1 .

Notice that combining proposition 3.5 and Claim 3.2, a is a degree one singularity for u_\star and the other points of $S \setminus \{a\}$ are degree zero singularities. In the next subsections, we aim at proving that $a \in \Omega$ is an interior point, that $S \setminus \{a\} = \emptyset$ and that $u_{\varepsilon_{k_l}} \rightarrow u_\star$ in $H_{loc}^1(\bar{\Omega} \setminus \{a\})$ as $l \rightarrow +\infty$. These properties will complete the proof of (i)-(v) of Theorem 0.1. The last point of Theorem 0.1 is a direct consequence of Proposition 7.1 [14].

3.4. The degree one singularity is an interior point of the domain. We aim at proving that $a \in \Omega$. Let us assume by contradiction that $a \in \partial\Omega$ and without loss of generality that $a = 0$.


As in the proof of the convergence of u_ε , u_\star is extended to a smooth domain Ω' thanks to Lemma 3.1. In order to light the notation, we will denote its extension u_\star . Moreover we can also apply Lemma 3.1 to Ω^c in order to get the following claim

Claim 3.3. *Let Γ a quasicircle and $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ then there exists a smooth neighborhood Ω'' of Γ diffeomorphic to an annulus and $u_0 \in H^1(\Omega'', \mathbb{S}^1)$ such that $u = g$ on Γ .*

We can assume that $\Omega'' \setminus \Omega = \Omega' \setminus \Omega$. Let $t > 0$ small enough such that $\mathbb{D}_t \subset \Omega''$.

We first proof that on $S_t \cap \Omega'$ there is always an arc with angle uniformly bounded from below.

Claim 3.4. *there exists $\delta > 0$ such that $\mathbb{D}_t \cap \Omega'$ contains a connected arc of arc length greater than δt , we denote it A'_t .*

FIGURE 3. Uniform control on $S_t \cap \Omega'$

Proof. We choose some orientation on Γ , let a_t^- the last point of $S_t \cap \Gamma$ before 0 and a_t^+ the first point of $S_t \cap \Gamma$ after 0. We denote by Γ_- the open arc of Γ from a_t^+ to 0. Thanks to (2.1), there is no point of Γ_- in $B(a_t^+, \frac{t}{C}) \cap \Gamma$. Let A_t^1 and A_t^2 the two open connected component of $S_t \setminus \{a_t^-, a_t^+\}$, and let a_t^1 (resp. a_t^2) the closest point to a_t^+ of $\Gamma_- \cap A_t^1$ (resp. $\Gamma_- \cap A_t^2$). By the previous remark, the distance of those points to a_t^+ is bigger than $\frac{t}{2C}$. Hence there is the two open arc starting at the a_t^i of arc length $\frac{t}{2C}$ which contains no point of Γ , one of this two is included in Ω' . \square

By (2.2) and the fact 0 is a degree one singularity, we have

$$\begin{aligned} 1 = \deg(u_*, 0) &= \frac{1}{2\pi} \int_{S_t} u_* \wedge u_{*\theta} d\theta \\ &= \frac{1}{2\pi} \int_{A_t^1} u_* \wedge u_{*\theta} d\theta + \frac{1}{2\pi} \int_{\Gamma \setminus A_t^1} u_* \wedge u_{*\theta} d\theta. \end{aligned}$$

Then, we have

$$\frac{1}{2\pi} \left| \int_{A_t^1} u_* \wedge u_{*\theta} d\theta \right| \leq \frac{1}{2\pi} \|u_0\|_{H^{\frac{1}{2}}(S_t)}^2 \leq C \|u_0\|_{H^1(A_t)}^2,$$

where $A_t = \mathbb{D}_{2t} \setminus \mathbb{D}_{\frac{t}{2}}$, and

$$\frac{1}{2\pi} \left| \int_{\Gamma \setminus A_t^1} u_* \wedge u_{*\theta} d\theta \right| \leq \frac{1}{2\pi} \int_{\Gamma \setminus A_t^1} |u_{*\theta}| d\theta \leq \frac{1}{2\pi} \left(\int_{\Gamma \setminus A_t^1} |u_{*\theta}|^2 d\theta \right)^{\frac{1}{2}} \sqrt{(2\pi - \delta)t}$$

Here we use the fact that the length of A_t^1 is at least δt . Then, we have

$$1 - C \|\nabla u_0\|_{L^2(A_t)}^2 \leq \frac{\sqrt{(2\pi - \delta)t}}{2\pi} \left(\int_{S_t} |u_{*\theta}|^2 d\theta \right)^{\frac{1}{2}},$$

hence by squaring this identity

$$\frac{4\pi^2}{(2\pi - \delta)t} (1 - 2C \|\nabla u_0\|_{L^2(A_t)}^2) \leq \int_{S_t} |u_{*\theta}|^2 d\theta,$$

But, thanks to Fubini Theorem, we have

$$\begin{aligned}
\int_{r_1}^{r_2} \frac{\|\nabla u_0\|_{L^2(A_t)}^2}{t} dt &= \int_{r_1}^{r_2} \frac{1}{t} \int_{A_{\frac{r_1}{2}, 2r_2}} |\nabla u_0|^2 \mathbf{1}_{\frac{t}{2} \leq |x| \leq 2t} dx dt \\
&= \int_{A_{\frac{r_1}{2}, 2r_2}} |\nabla u_0|^2 \int_{r_1}^{r_2} \frac{1}{t} dt dx \\
&= \ln(4) \int_{A_{\frac{r_1}{2}, 2r_2}} |\nabla u_0|^2 dx.
\end{aligned}$$

Finally we get

$$\frac{4\pi^2}{(2\pi - \delta)t} \ln\left(\frac{r_2}{r_1}\right) - C \|\nabla u_0\|_{L^2(A_{\frac{r_1}{2}, 2r_2})} \leq \int_{A_{r_1, r_2}} |\nabla u_*|^2 dx.$$

But thanks to (3.24), we know that

$$\int_{A_{r_1, r_2}} |\nabla u_*|^2 dx \leq 2\pi \ln\left(\frac{r_2}{r_1}\right) + C,$$

which leads to a contradiction taking $\frac{r_2}{r_1}$ small enough.

3.5. Final convergence arguments.

Proposition 3.7. *With the notations of Claim 3.2, we have that $u_{\varepsilon_{k_l}} \rightarrow u_*$ in $H_{loc}^1(\bar{\Omega} \setminus \{a\})$ and for any compact set $K \Subset \bar{\Omega} \setminus \{a\}$,*

$$(3.25) \quad \frac{1}{4\varepsilon_{k_l}^2} \int_K (1 - |u_{\varepsilon_{k_l}}|^2)^2 \rightarrow 0 \text{ as } l \rightarrow +\infty$$

Proof. Let $r_0 > 0$ small enough be such that $\mathbb{D}_{r_0}(a) \cap (\partial\Omega \cup S) = \emptyset$. Since $u_{\varepsilon_{k_l}}$ is bounded in $H^1(\Omega \setminus \mathbb{D}_{r_0}(a))$, $u_{\varepsilon_{k_l}}$ weakly converges to some function u_* in $H^1(\Omega \setminus \mathbb{D}_{r_0}(a))$. Moreover, u_* is a harmonic map into \mathbb{S}^1 and u_* satisfies

$$(3.26) \quad \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_*|^2 \leq \liminf_{l \rightarrow +\infty} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_{\varepsilon_{k_l}}|^2.$$

From now to the end of the proof of the proposition we denote $\varepsilon := \varepsilon_{k_l}$.

Let w_ε be the solution the following minimization problem

$$\frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla w_\varepsilon|^2 = \inf_{u \in \mathcal{A}_\varepsilon} \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u|^2$$

on the admissible set

$$\mathcal{A}_\varepsilon = \left\{ u \in H^1(\bar{\Omega} \setminus \mathbb{D}_{r_0}(a), \mathbb{S}^1); u = \frac{u_\varepsilon}{|u_\varepsilon|} \text{ on } \partial(\Omega \setminus \mathbb{D}_{r_0}(a)) \right\}.$$

As explained in Claim 3.3, we can find u_ε^0 is defined a tubular neighborhood Ω'' of $\partial\Omega$ which is in $C^\infty(\Omega'' \setminus \mathbb{S}^1) \cap H^1(\Omega'')$ and such that $u_\varepsilon^0 = u_\varepsilon$ on $\partial\Omega$. Up to reduce r_0 , we can assume that $\mathbb{D}(a, r_0) \cap \Omega'' = \emptyset$. Then we set $\omega = \Omega \setminus \Omega''$. Let $\Psi : [a, b] \times \mathbb{S}^1 \rightarrow \bar{\omega} \setminus \mathbb{D}_{r_0}(a)$ be a smooth diffeomorphism such that $\Psi(\{a\} \times \mathbb{S}^1) = \partial\omega$ and $\Psi(\{b\} \times \mathbb{S}^1) = S(a, r_0)$. For instance, Ψ can be build with a uniformization map from an annulus to $\bar{\omega} \setminus \mathbb{D}_{r_0}(a)$. We set $a_\varepsilon = u_\varepsilon^0 \circ \Psi|_{\{a\} \times \mathbb{S}^1}$ and $b_\varepsilon = u_\varepsilon \circ \Psi|_{\{b\} \times \mathbb{S}^1}$.

Since $\deg(a_\varepsilon) = \deg(b_\varepsilon)$, there is a smooth homotopy $H_\varepsilon : [a, b] \times \mathbb{S}^1 \rightarrow \mathbb{S}^1$ such that $H_\varepsilon(a, \cdot) = a_\varepsilon$ and $H_\varepsilon(b, \cdot) = b_\varepsilon$ and the map

$$v_\varepsilon := \begin{cases} H_\varepsilon \circ \Psi^{-1} & \text{in } \omega \setminus \mathbb{D}_{r_0}(a) \\ u_\varepsilon^0 & \text{in } \Omega \setminus \omega \end{cases}$$

satisfies $v_\varepsilon \in \mathcal{A}_\varepsilon$.

It is clear that since $w_\varepsilon \rightarrow u_\star$ in $H^{\frac{1}{2}}(\partial(\Omega \setminus \mathbb{D}_{r_0}(a)))$ as $l \rightarrow +\infty$, $w_{\varepsilon_{k_l}}$ converges in $H^1(\Omega \setminus \mathbb{D}_{r_0}(a))$ to some function w_\star such that $w_\star = u_\star$ on $\partial(\Omega \setminus \mathbb{D}_{r_0}(a))$ and solution of

$$\frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla w_\star|^2 = \inf_{u \in \mathcal{A}_\star} \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u|^2$$

on the admissible set

$$\mathcal{A}_\star = \{u \in H^1(\overline{\Omega} \setminus \mathbb{D}_{r_0}(a), \mathbb{S}^1); u = u_\star \text{ on } \partial(\Omega \setminus \mathbb{D}_{r_0}(a))\}.$$

In particular $u_\star \in \mathcal{A}_\star$ and

$$(3.27) \quad \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla w_\star|^2 \leq \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_\star|^2$$

We let f_ε be the solution of

$$\begin{cases} \Delta f_\varepsilon = 0 & \text{in } \Omega \setminus \mathbb{D}_{r_0}(a) \\ f_\varepsilon = 1 & \text{on } \partial\Omega \\ f_\varepsilon = |u_\varepsilon| & \text{on } S(a, r_0) \end{cases}$$

Testing the function

$$h_\varepsilon := \begin{cases} f_\varepsilon w_\varepsilon & \text{in } \Omega \setminus \mathbb{D}_{r_0}(a) \\ u_\varepsilon & \text{in } \mathbb{D}_{r_0}(a) \end{cases}$$

for the Ginzburg-Landau energy, we have that

$$E_\varepsilon(u_\varepsilon) \leq E_\varepsilon(h_\varepsilon).$$

so that since $h_\varepsilon = u_\varepsilon$ in $\mathbb{D}_{r_0}(a)$ and $|h_\varepsilon| = f_\varepsilon$ in $\Omega \setminus \mathbb{D}_{r_0}(a)$,

$$(3.28) \quad \begin{aligned} & \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_\varepsilon|^2 + \frac{1}{4\varepsilon^2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} (1 - |u_\varepsilon|^2)^2 \\ & \leq \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} f_\varepsilon^2 |\nabla w_\varepsilon|^2 + \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla f_\varepsilon|^2 + \frac{1}{4\varepsilon^2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} (1 - f_\varepsilon^2)^2 \end{aligned}$$

We set $\varphi_\varepsilon = \frac{1-f_\varepsilon}{\varepsilon^2}$. Knowing that $f_\varepsilon \leq 1$ by the maximum principle, we then easily obtain that

$$(3.29) \quad \begin{aligned} & \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} f_\varepsilon^2 |\nabla w_\varepsilon|^2 + \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla f_\varepsilon|^2 + \frac{1}{4\varepsilon^2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} (1 - f_\varepsilon^2)^2 \\ & \leq \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla w_\varepsilon|^2 + \varepsilon^2 \left(\frac{\varepsilon^2}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla \varphi_\varepsilon|^2 + \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} \varphi_\varepsilon^2 \right) \end{aligned}$$

Now, we aim at proving that φ_ε is a non negative uniformly bounded function in $H^1 \cap L^\infty(\Omega \setminus \mathbb{D}_{r_0}(a))$. Indeed, φ_ε is a harmonic function, uniformly bounded in

$\mathcal{C}^\infty(S(a, r_0))$ since we know by Claim 3.2 that

$$\frac{1 - |u_\varepsilon|^2}{\varepsilon^2} \rightarrow |\nabla u_\star|^2 \text{ on } \mathcal{C}^\infty(S(a, r_0)).$$

Setting $\tilde{\varphi}_\varepsilon = \varphi_\varepsilon \circ f$, where $f : \mathbb{D} \rightarrow \Omega$ is a uniformization map, we have that

$$\begin{cases} \Delta \tilde{\varphi}_\varepsilon = 0 & \text{in } \mathbb{D} \setminus f^{-1}(\mathbb{D}_{r_0}(a)) \\ \tilde{\varphi}_\varepsilon = 0 & \text{on } \mathbb{S}^1 \\ \tilde{\varphi}_\varepsilon = \frac{1 - |u_\varepsilon \circ f|^2}{\varepsilon^2} & \text{on } f^{-1}(S(a, r_0)) \end{cases}$$

and by standard elliptic regularity, $\tilde{\varphi}_\varepsilon$ is uniformly bounded. Coming back to $\varphi_\varepsilon = \tilde{\varphi}_\varepsilon \circ f^{-1}$, using the conformal invariance of the energy, we obtain the expected result.

Now, using (3.28), (3.29) and the uniform boundedness of φ_ε in $H^1 \cap L^\infty$, we obtain

$$(3.30) \quad \begin{aligned} \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_\varepsilon|^2 + \frac{1}{4\varepsilon^2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} (1 - |u_\varepsilon|^2)^2 \\ \leq \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla w_\varepsilon|^2 + O(\varepsilon^2) \end{aligned}$$

Passing to the limit on (3.30) and using (3.27) and (3.26), we obtain

$$\begin{aligned} \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_\star|^2 &\leq \liminf_{\varepsilon \rightarrow 0} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_\varepsilon|^2 \\ &\leq \limsup_{\varepsilon \rightarrow 0} \left(\int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_\varepsilon|^2 + \frac{1}{4\varepsilon^2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} (1 - |u_\varepsilon|^2)^2 \right) \\ &\leq \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla w_\star|^2 \leq \frac{1}{2} \int_{\Omega \setminus \mathbb{D}_{r_0}(a)} |\nabla u_\star|^2 \end{aligned}$$

so that choosing r_0 as small as necessary, we obtain strong $H_{loc}^1(\Omega \setminus \{a\})$ convergence of u_ε and (3.25). \square

We obtain from Proposition 3.7 and proposition 3.4 that $S \setminus \{a\} = \emptyset$, so that the proof of Theorem 0.1 is complete.

4. MINIMIZATION OF THE RENORMALIZED ENERGY AND THE EXAMPLE OF TANGENTIAL BOUNDARY DATA

4.1. Renormalized energy of canonical harmonic maps. Let Ω be a quasidisk and Γ its boundary. In this section, we give an H^1 version of the definition of canonical harmonic maps with a prescribed degree 1 singularity $a \in \Omega$ and of its renormalized energy. The following definition is an H^1 version of section I.3 of [2].

Theorem-Definition 4.1 (Canonical harmonic map). *Let $a \in \Omega$. Let $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ and $\deg(g, \Gamma) = 1$. There is a unique map $u : \Omega \setminus \{a\} \rightarrow \mathbb{S}^1$ such that*

- $u \in H_{loc}^1(\overline{\Omega} \setminus \{a\})$,
- $\deg(u, a) = 1$,
- $u = g$ on Γ ,

- $\star\omega = d\Phi$, where $\omega = \langle du, iu \rangle$ and $\Phi = \tilde{\Phi} \circ f^{-1}$, where $f : \mathbb{D} \rightarrow \Omega$ is a uniformization map such that $f(0) = a$ and $\tilde{\Phi}$ is the unique solution (up to a constant) of

$$(4.1) \quad \begin{cases} \Delta \tilde{\Phi} = 2\pi\delta_0 & \text{in } \mathbb{D}, \\ \partial_r \tilde{\Phi} = \langle \partial_\theta (g \circ f), i(g \circ f) \rangle & \text{on } \mathbb{S}^1. \end{cases}$$

Moreover $u : \Omega \setminus \{a\} \rightarrow \mathbb{S}^1$ is a harmonic map. We say that u is the canonical harmonic map of degree 1 associated to the data (g, a) .

Proof. STEP 1 : Uniqueness of u . Let u_1 and u_2 satisfying all the properties. Let U be a simply connected domain such that $U \Subset \Omega \setminus \{a\}$. Then, we can write $u_1 = e^{i\psi_1}$ and $u_2 = e^{i\psi_2}$ for $\psi_1, \psi_2 \in H^1(U)$. Then we have that

$$d(\psi_1 - \psi_2) = \langle du_1, iu_1 \rangle - \langle du_2, iu_2 \rangle = \star d\Phi - \star d\Phi = 0$$

Therefore there is a constant α_U such that $|\alpha_U| = 1$ and $u_2 = \alpha_U u_1$ in U . Since Ω is connected, we get that $\alpha := \alpha_U$ does not depend on U and $u_2 = \alpha u_1$ in Ω . We finally get that $\alpha = 1$ since on Γ we have $u_1 = g = u_2$.

STEP 2 : $\tilde{\Phi}$ is well defined by the Green representation formula. Let $\mathcal{G}_x(y)$ the Green function such that

$$(4.2) \quad \begin{cases} \Delta \mathcal{G}_x = 2\pi\delta_x & \text{in } \mathbb{D} \\ \partial_r \mathcal{G}_x = 0 & \text{on } \mathbb{S}^1 \\ \int_{\mathbb{S}^1} \mathcal{G}_x = 0 & \text{on } \mathbb{S}^1 \end{cases}$$

Then for $x \in \mathbb{D}$, since $g \circ f \in H^{\frac{1}{2}}(\mathbb{S}^1)$, $\partial_\theta (g \circ f) \in H^{-\frac{1}{2}}$ and $\mathcal{G}_x \in C^\infty$

$$\tilde{\Phi}(x) = \mathcal{G}_0(x) + \int_{\mathbb{S}^1} \mathcal{G}_x \langle \partial_\theta (g \circ f), i(g \circ f) \rangle$$

is a solution of (4.1).

STEP 3 : Construction of u . We pullback all the expected properties of u by $f : \mathbb{D} \rightarrow \Omega$, a uniformization map such that $f(0) = a$. We set

$$\tilde{g} = g \circ f \quad \tilde{u} = u \circ f \quad \tilde{\omega} = \langle d\tilde{u}, i\tilde{u} \rangle$$

Then, u is nothing but $u = \tilde{u} \circ f^{-1}$, where \tilde{u} is the canonical harmonic map of degree 1 on the disk \mathbb{D} associated to the data $(g \circ f, 0)$ which satisfies the desired property thanks to section 1.3 of [2]. □

Definition 4.1 (Renormalized energy of canonical harmonic maps). *Let $u : \Omega \rightarrow \mathbb{S}^1$ be the canonical harmonic map with data $a \in \Omega$, $g \in H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$: the renormalized energy is well defined by*

$$W(a) = \lim_{\delta \rightarrow 0} \left(\int_{\Omega \setminus \mathbb{D}_\delta(a)} |\nabla u|^2 - 2\pi \ln \frac{1}{\delta} \right)$$

4.2. Energy minimizing Ginzburg-Landau solutions minimize the renormalized energy. Let $u := u_*$ a minimizing solution of the Ginzburg-Landau problem given by Theorem 0.1. We set $v = -iu$, so that $v : \Omega \setminus \{a\} \rightarrow \mathbb{S}^1$ is a unit vector field such that, for any point $x \in \Omega \setminus \{a\}$, $(v(x), u(x))$ is an direct orthonormal basis. We set

$$\omega = \langle du, v \rangle = -\langle dv, u \rangle$$

the Cartan form in Ω .

Let G be the Dirichlet Green function with respect to a in Ω , i.e satisfying

$$(4.3) \quad \begin{cases} \Delta G = 2\pi\delta_a & \text{in } \Omega \\ G = 0 & \text{on } \Gamma . \end{cases}$$

Proposition 4.1. *There is a harmonic function μ in Ω , such that*

$$(4.4) \quad \star\omega = d(\mu + G) .$$

Then u is the canonical harmonic map of degree 1 with respect to the data (g, a) .

Proof. We know that $du \in L^p(\Omega)$ for any $1 < p < 2$ (see e.g. [36], page 1620), so that $\omega \in L^p(\Omega)$. We have that

$$\star d\omega = \star(\langle du \wedge dv \rangle) = \langle \nabla^\perp u, \nabla v \rangle = 0$$

in $\Omega \setminus \{a\}$ since $(v, u) = i(-u, v)$. Then the support of the distribution $\star d\omega$ is $\{a\}$. Since $\omega \in L^p$, we can write

$$(4.5) \quad d\omega = \alpha\delta_a .$$

Let's compute α .

$$(4.6) \quad \alpha = \int_\Omega d\omega = \int_{\partial\Omega} \omega = \int_{\mathbb{S}^1} \tilde{\omega} = \int_{\partial\Omega} \langle \tilde{u}_\theta, \tilde{v} \rangle = 2\pi \deg(g) = 2\pi .$$

We then have that

$$(4.7) \quad d(\omega - \star dG) = 0$$

Moreover, we prove that there is $\beta \in \mathbb{R}$ such that

$$(4.8) \quad d(\star\omega) = \beta\delta_a$$

in the sense of distributions in Ω . Indeed, again, since $\omega \in L^p(\Omega)$, we just have to prove that the support of the distribution $d(\star\omega)$ is $\{a\}$. Indeed, we first have that

$$\star d(\star\omega) = \langle \Delta u, v \rangle + \star(\langle \star du \wedge dv \rangle) = \langle \Delta u, v \rangle + \langle \nabla u, \nabla v \rangle$$

in $\Omega \setminus \{a\}$. Since $u : \Omega \setminus \{a\} \rightarrow \mathbb{S}^1$ is a harmonic map we have that

$$\langle \Delta u, v \rangle = 0$$

in $\Omega \setminus \{a\}$. Moreover, we have that

$$(4.9) \quad 0 = \Delta(\langle u, v \rangle) = \langle \Delta u, v \rangle + \langle u, \Delta v \rangle + 2\langle \nabla u, \nabla v \rangle = 2\langle \nabla u, \nabla v \rangle \text{ in } \Omega \setminus \{a\}$$

since $\langle u, v \rangle = 0$ and $(v, u) = i(-u, v)$. We obtain (4.8). Then we have that

$$(4.10) \quad d(\star\omega - dG) = \beta\delta_a$$

which gives that $h = \star\omega - dG - \beta\star dG$ satisfies

$$(4.11) \quad dh = 0 \text{ and } d(\star h) = 0 .$$

Then h is a smooth harmonic form in Ω . There is a smooth function $\tilde{\mu}$ such that $h = d\tilde{\mu}$. To complete the proof of Claim 4.1, we finally prove that $\beta = 0$. From 3.23 we know that there is a constant $C > 0$ such that for $\delta < \frac{\text{dist}(a, \Gamma)}{2}$,

$$\int_{\Omega \setminus B_a(\delta)} |\nabla u|^2 \leq 2\pi \ln \left(\frac{1}{\delta} \right) + C.$$

Knowing that $|\nabla u|^2 = |\langle \nabla u, v \rangle|^2 = |\omega|^2$, we have

$$(4.12) \quad 2\pi \ln \left(\frac{1}{\delta} \right) + C \geq \int_{\Omega \setminus B_a(\delta)} |\omega|^2 = \int_{\Omega \setminus B_a(\delta)} |\nabla G + \beta \nabla^\perp G + \nabla \mu|^2.$$

Since μ is smooth on Ω and ∇G is bounded in L^p for any $p < 2$, we have a constant C' independent from $\delta < \frac{d(a, \Gamma)}{2}$ such that

$$\left| \int_{\Omega \setminus B_a(\delta)} 2 \langle \nabla G + \beta \nabla^\perp G, \nabla \mu \rangle + |\nabla \mu|^2 \right| \leq C'$$

Then

$$\int_{\Omega \setminus B_a(\delta)} |\nabla G + \beta \nabla^\perp G|^2 = (1 + \beta^2) \int_{\Omega \setminus B_a(\delta)} |\nabla G|^2 + 2\beta \int_{\Omega \setminus B_a(\delta)} \langle \nabla G, \nabla^\perp G \rangle$$

Since $\langle \nabla G, \nabla^\perp G \rangle = 0$, we then can write (4.12) as

$$(4.13) \quad 2\pi \ln \left(\frac{1}{\delta} \right) + C \geq (1 + \beta^2) \int_{\Omega \setminus B_a(\delta)} |\nabla G|^2 - C' \geq 2\pi (1 + \beta^2) \ln \left(\frac{1}{\delta} \right) - C'',$$

for a constant C'' independent from δ , since $G(x) = \ln|x - a| + w(x)$ on Ω , where w is a harmonic function. Dividing by $\ln \left(\frac{1}{\delta} \right)$ and letting $\delta \rightarrow 0$ gives $\beta = 0$.

Now it is easy to check that u is the canonical harmonic map of degree 1 with respect to the data (g, a) and the proof of Claim 4.1 is complete. \square

Finally, we also have, in our setting, the fact that the singularity of the minimizer minimizes the renormalized energy, the proof is exactly the same as the one of Theorem VIII.1 [2].

Proposition 4.2. *The singularity a of the minimizing Ginzburg-Landau solution u is a minimizer of the renormalized energy $W(a)$.*

4.3. The special case of tangential boundary data. In this section we prove the Proposition 0.3, when the boundary data $g = \tau$, where τ is the tangent vector field on Γ such that the frame (ν, τ) is a direct frame, where ν is the out-pointing normal of Ω . We have $\tau = -i\nu$.

In the following proposition 4.3 we aim at proving that canonical harmonic maps with respect to tangential and normal data are exactly push forwards in Ω of the polar orthonormal frame of the disk (with singularity at 0) by uniformization maps.

Let $f : \mathbb{D} \rightarrow \Omega$ be a uniformization map of Ω . We set $u : \Omega \rightarrow \mathbb{S}^1$

$$u = \frac{f_\theta}{|f_\theta|} \circ f^{-1}$$

and $v : \Omega \rightarrow \mathbb{S}^1$

$$v = \frac{f_r}{|f_r|} \circ f^{-1}$$

where f_r and f_θ are the radial and angular derivatives of f defined by

$$f_r := \partial_r f = \frac{xf_x + yf_y}{\sqrt{x^2 + y^2}} = \frac{xf_x + yf_y}{r}$$

where $r = \sqrt{x^2 + y^2}$ is the radial coordinate and

$$f_\theta := \partial_\theta f = -yf_x + xf_y$$

where $f_x := \partial_x f$ and $f_y := \partial_y f$ are the partial derivatives with respect to the coordinates of $z = x + iy$.

Proposition 4.3. *Let $a \in \Omega$. For any uniformization map $f : \mathbb{D} \rightarrow \Omega$ such that $f(0) = a$, $u = \frac{f_\theta}{|f_\theta|} \circ f^{-1}$ and $v = \frac{f_r}{|f_r|} \circ f^{-1}$ are the canonical harmonic maps of degree 1 associated to the data (τ, a) and (ν, a) .*

Proof. Let $f : \mathbb{D} \rightarrow \Omega$ be a uniformization map such that $f(0) = a$. Notice that since f is a conformal map, (v, u) is an orthonormal frame in Ω and notice also that on Γ , $u = \tau$ and $v = \nu$. We now set

$$\tilde{\omega} = \langle d\tilde{u}, \tilde{v} \rangle = -\langle d\tilde{v}, \tilde{u} \rangle$$

the Cartan form of (\tilde{v}, \tilde{u}) where

$$\tilde{v} = \frac{f_r}{|f_r|} \text{ and } \tilde{u} = \frac{f_\theta}{|f_\theta|}$$

and

$$\tilde{\mu} = \ln |f_x| = \ln |f_y| = \ln |f_r| = \ln \left(\frac{|f_\theta|}{r} \right)$$

so that

$$\begin{cases} f_r = e^{\tilde{\mu}} \tilde{v} \\ f_\theta = e^{\tilde{\mu} + \tilde{G}} \tilde{u} \end{cases}$$

and

$$\begin{cases} \tilde{v} = (xf_x + yf_y) e^{-\tilde{\mu} - \tilde{G}} \\ \tilde{u} = (-yf_x + xf_y) e^{-\tilde{\mu} - \tilde{G}} \end{cases}$$

where $\tilde{G}(x, y) = \ln r$ is the Green function of \mathbb{D} with Dirichlet boundary condition with respect to the singularity 0.

Let's prove the following equation:

$$(4.14) \quad \star \tilde{\omega} = d(\tilde{\mu} + \tilde{G})$$

in \mathbb{D} . We have that

$$\begin{aligned} \langle \tilde{u}_x, \tilde{v} \rangle &= \langle -yf_{xx} + f_y + xf_{xy}, xf_x + yf_y \rangle e^{-2\tilde{\mu} - 2\tilde{G}} \\ &= \left(-xyf_{xx} \cdot f_x - y^2 f_{xx} \cdot f_y + xf_y \cdot f_x + y|f_y|^2 + x^2 f_{xy} \cdot f_x + xyf_{xy} \cdot f_y \right) e^{-2\tilde{\mu} - 2\tilde{G}} \\ &= \left(-\frac{xy}{2} \left(|f_x|^2 \right)_x + \frac{y^2}{2} \left(|f_y|^2 \right)_y + y|f_y|^2 + \frac{x^2}{2} \left(|f_x|^2 \right)_y + \frac{xy}{2} \left(|f_y|^2 \right)_x \right) e^{-2\tilde{\mu} - 2\tilde{G}} \\ &= (-xy\tilde{\mu}_x + y^2\tilde{\mu}_y + y + x^2\tilde{\mu}_y + xy\tilde{\mu}_x) e^{-2\tilde{G}} \\ &= \tilde{\mu}_y + \tilde{G}_y \end{aligned}$$

where we used that $f_{xx} = -f_{yy}$ since f is harmonic $\Delta f = 0$. We also have that

$$\begin{aligned}
\langle \tilde{u}_y, \tilde{v} \rangle &= \langle -f_x - yf_{xy} + xf_{yy}, xf_x + yf_y \rangle e^{-2\tilde{\mu}-2\tilde{G}} \\
&= \left(-x|f_x|^2 - yf_x \cdot f_y - xyf_{xy} \cdot f_x - y^2 f_{xy} \cdot f_y + x^2 f_{yy} \cdot f_x + xyf_{yy} \cdot f_y \right) e^{-2\tilde{\mu}-2\tilde{G}} \\
&= \left(-x|f_x|^2 - \frac{xy}{2} \left(|f_x|^2 \right)_y - \frac{y^2}{2} \left(|f_y|^2 \right)_x - \frac{x^2}{2} \left(|f_x|^2 \right)_x + \frac{xy}{2} \left(|f_y|^2 \right)_y \right) e^{-2\tilde{\mu}-2\tilde{G}} \\
&= (-x - xy\tilde{\mu}_y - y^2\tilde{\mu}_x - x^2\tilde{\mu}_x + xy\tilde{\mu}_y) e^{-2\tilde{G}} \\
&= - \left(\tilde{\mu}_x + \tilde{G}_x \right)
\end{aligned}$$

We proved (4.14). Moreover, since (\tilde{u}, \tilde{v}) is an orthonormal frame, we have

$$\star d\tilde{\omega} = \star (\langle d\tilde{u} \wedge d\tilde{v} \rangle) = \langle \nabla^\perp \tilde{u}, \nabla \tilde{v} \rangle = 0 \text{ in } \mathbb{D} \setminus \{0\}$$

since $(\tilde{u}, \tilde{v}) = i(\tilde{v}, -\tilde{u})$. Then the support of the distribution $\star d\tilde{\omega}$ is $\{0\}$. Since by (4.14) $\tilde{\omega} \in L^p_{loc}(\mathbb{D})$ for $1 \leq p < 2$, we can write

$$(4.15) \quad d\tilde{\omega} = \alpha \delta_0,$$

where δ_0 is the Dirac mass supported in $\{0\}$. Let's compute α .

$$(4.16) \quad \alpha = \int_{\mathbb{D}} d\tilde{\omega} = \int_{\mathbb{S}^1} \tilde{\omega} = \int_{\mathbb{S}^1} \langle \tilde{u}_\theta, \tilde{v} \rangle = 2\pi \deg(\tilde{u}, \mathbb{S}^1) = 2\pi.$$

This degree is well defined since $\tilde{u} \in H^{\frac{1}{2}}(\mathbb{S}^1)$ because Γ is Weil-Petersson. We then have that

$$(4.17) \quad \Delta \tilde{\mu} = d(\tilde{\omega} - \star d\tilde{G}) = 0.$$

In Ω , we have $u = \tilde{u} \circ f^{-1}$ and $v = \tilde{v} \circ f^{-1}$. We set

$$\omega = \langle du, v \rangle = (f^{-1})' \tilde{\omega} \circ f^{-1}$$

the Cartan form on Ω and $G = \tilde{G} \circ f^{-1}$ the Dirichlet-Green function with respect to $a := f(0)$. We obtain from (4.14)

$$(4.18) \quad \star \omega = d(\mu + G)$$

denoting $\mu = \tilde{\mu} \circ f^{-1}$, where we have that $\tilde{\Phi} = (\mu + G) \circ f$ is a solution of (4.1).

Notice that Γ is a Weil-Petersson curve if and only if

$$|\nabla \tilde{\mu}|^2 = \left| \frac{f''}{f'} \right|^2 \in L^1(\mathbb{D})$$

if and only if $|\nabla \mu|^2 \in L^1(\Omega)$ where $\mu = \tilde{\mu} \circ f^{-1}$ by conformal invariance of the Dirichlet energy. In particular by (4.18), $u \in H^1_{loc}(\overline{\Omega} \setminus \{a\})$ and $\omega \in L^2_{loc}(\overline{\Omega} \setminus \{a\})$. Therefore, we have all the necessary regularity to have $u = \tau \in H^{\frac{1}{2}}(\Gamma)$ and the proof of the proposition is complete. \square

Remark 4.1. Notice that by Proposition 4.1, $\tilde{\mu} = \mu \circ f$ satisfies the following equation

$$(4.19) \quad \begin{cases} \Delta \tilde{\mu} = 0 & \text{in } \mathbb{D} \\ \partial_r \tilde{\mu} = \langle \partial_\theta \tilde{u}, \tilde{v} \rangle - 1 & \text{on } \partial \mathbb{D}, \end{cases}$$

where ∂_r and ∂_θ denote the radial and angular derivatives on the disk, and $\tilde{u} = u \circ f$, $\tilde{v} = v \circ f$. Equation (4.19) is nothing but the Liouville equation up to the boundary, since $\langle u_\tau, v \rangle$ is the curvature of Γ , see [10] for more details about prescribing geodesic curvature.

Now, we can reformulate the renormalized energy in terms of the uniformization

Proposition 4.4. *Let $f : \mathbb{D} \rightarrow \Omega$ be a uniformization map such that $f(0) = a$. Then*

$$(4.20) \quad W(a) = W_0 - 2\pi \ln |f'(0)|$$

where

$$W_0 = \int_{\mathbb{D}} \left| \frac{f''}{f'} \right|^2 + 4\pi \ln |f'(0)|$$

does not depend on the choice of the uniformization $f : \mathbb{D} \rightarrow \Omega$ nor in a .

Proof. The fact that W_0 only depends on Ω is well known, see lemma 3.4 page 86 of [38]. As in the proof of Proposition 4.3, we let $\tilde{\mu} = \ln |f'|$. Since Γ is a Weil-Petersson curve,

$$|\nabla \tilde{\mu}|^2 = \left| \frac{f''}{f'} \right|^2 \in L^1(\mathbb{D})$$

so that $|\nabla \mu|^2 \in L^1(\Omega)$ where $\mu = \tilde{\mu} \circ f^{-1}$ by conformal invariance of the Dirichlet energy. In particular by (4.18), $\omega \in L^2_{loc}(\bar{\Omega} \setminus \{a\})$. Then by a simple computation and by (4.18),

$$\begin{aligned} \int_{\Omega \setminus B_\rho(a)} |\omega|^2 &= \int_{\Omega \setminus B_\rho(a)} |\nabla G|^2 + 2 \int_{\Omega \setminus B_\rho(a)} \langle \nabla G, \nabla \mu \rangle + \int_{\Omega \setminus B_\rho(a)} |\nabla \mu|^2 \\ &= - \int_{\partial B_\rho(a)} G \partial_\nu G - 2 \int_{\partial B_\rho(a)} G \partial_\nu \mu + \int_{\Omega \setminus B_\rho(a)} |\nabla \mu|^2 \\ &= - \int_{\partial B_\rho(a)} (\ln |x - a| + h) \partial_\nu (\ln |x - a|) + O(\rho \ln \rho) \\ &\quad + \int_{\Omega \setminus B_\rho(a)} |\nabla \mu|^2 \\ &= 2\pi \ln \left(\frac{1}{\rho} \right) + \int_{\Omega \setminus B_\rho(a)} |\nabla \mu|^2 - 2\pi h(a) + o(1) \end{aligned}$$

as $\rho \rightarrow 0$, where $h(x) = G(x, a) - \ln |x - a|$ is a smooth harmonic function in Ω . Since $\ln |x| = G(f(x), a)$, we have that

$$(4.21) \quad \begin{aligned} h(a) &= h(f(0)) \\ &= \lim_{z \rightarrow 0} (G(f(z)) - \ln |f(z) - a|) \\ &= \lim_{z \rightarrow 0} \left(- \ln \left| \frac{f(z) - a}{z} \right| \right) \\ &= - \ln |f'(0)| \end{aligned}$$

so that

$$W(a) = \int_{\mathbb{D}} |\nabla \tilde{\mu}|^2 + 2\pi \ln |f'(0)| .$$

We then have the expected identity (4.20). \square

Remark 4.2. A minimizer $a \in \Omega$ of W is $a = f(0)$ where f is a maximizer of the following maximization problem

$$\max_{f: \mathbb{D} \rightarrow \Omega} |f'(0)|$$

where the maximum stands on the set of uniformizing maps of Ω . Or equivalently, fixing one uniformization map $f_0: \mathbb{D} \rightarrow \Omega$, a minimizer is $a = f_0(\omega_0)$ where $\omega_0 \in \mathbb{D}$ is chosen as a maximizer for

$$(4.22) \quad |f'_0(\omega_0)| \left(1 - |\omega_0|^2\right) = \max_{\omega \in \mathbb{D}} |f'_0(\omega)| \left(1 - |\omega|^2\right) .$$

Indeed, we can write any uniformization map as $f = f_0 \circ \psi^{-1}$, where

$$\psi(z) = e^{i\theta} \frac{z - \omega}{\bar{\omega}z - 1}$$

for $\omega \in \mathbb{D}$ and $\theta \in \mathbb{R}$ so that since $\psi(\omega) = 0$

$$|f'(0)| = \left| f'_0(\omega) (\psi^{-1})'(0) \right| = \left| \frac{f'_0(\omega)}{\psi'(\omega)} \right| = |f'_0(\omega)| \left(1 - |\omega|^2\right) .$$

As we can see in (4.21), $-\ln |f'(0)|$ is nothing but the mass of the Dirichlet-Green function of Ω at the point a where f is defined as a uniformization map such that $f(0) = a$.

Remark 4.3. We know by section 3.4 that a minimizer of the renormalized energy cannot be realized on the boundary of Ω . In the case of tangential data, it is a simple consequence of the previous remark and Koebe's distortion theorem. Indeed, a point ω_0 satisfying (4.22) must belong to \mathbb{D} since

$$|f'_0(\omega)| \left(1 - |\omega|^2\right) \leq \text{dist}(f_0(\omega), \partial\Omega) \rightarrow 0 \text{ as } |\omega| \rightarrow 1$$

so that $a = f_0(\omega_0) \in \Omega$.

5. A NEW POINT OF VIEW ON THE RIEMANN MAPPING THEOREM

We assume in this section that Γ is a smooth³ Jordan curve. As a side remark, we aim at proving the Riemann mapping theorem for the smooth bounded domain Ω enclosing Γ .

Let (v, u) be the frame defined in subsection 4.2 with tangential and normal boundary data, and $\omega = \langle du, v \rangle$ be the associated Cartan form. We let $\Phi = \mu + G$ where G is defined by (4.3) and μ is given by Proposition (4.1). Notice that since we assumed that Ω is smooth, we never needed any uniformization map of Ω to build the moving frame (v, u) , since the map u is given from the classical Ginzburg-Landau setting. We compute the following Lie bracket in $\Omega \setminus \{a\}$:

$$\begin{aligned} [e^{\Phi} v, e^{\Phi} u] &= e^{2\Phi} \{[v, u] + d\Phi(v)u - d\Phi(u)v\} \\ &= e^{2\Phi} \{\nabla_v u - \nabla_u v + d\Phi(v)u - d\Phi(u)v\} \\ &= e^{2\Phi} \{(\omega(u) + d\Phi(v))u + (-d\Phi(u) + \omega(v))v\} . \end{aligned}$$

Let (α, β) the dual basis of (v, u) in $T^*\Omega$, we have that

$$\star\omega = \star(\omega(v)\alpha + \omega(u)\beta) = \omega(v)\beta - \omega(u)\alpha ,$$

³In fact we can assume C^2 but not much less as explained in remark 5.1 below.

where $\star\alpha = \beta$ and $\star\beta = -\alpha$. Therefore, we have that

$$[e^\Phi v, e^\Phi u] = e^{2\Phi} \{(-\star\omega(v) + d\Phi(v))u + (-d\Phi(u) + \star\omega(u))v\} = 0,$$

since $\star\omega = d\Phi$. We have the following result, which is a global version of the Frobenius theorem in our context

Proposition 5.1. *There is a homeomorphism $\psi :]-\infty, 0] \times \mathbb{R}/\rho\mathbb{Z} \rightarrow \bar{\Omega} \setminus \{a\}$ such that*

$$(5.1) \quad \begin{cases} \partial_s \psi = e^{\Phi \circ \psi} v \circ \psi \\ \partial_\theta \psi = e^{\Phi \circ \psi} u \circ \psi. \end{cases}$$

In particular, $\psi :]-\infty, 0[\times \mathbb{R}/\rho\mathbb{Z} \rightarrow \Omega \setminus \{a\}$ is a conformal diffeomorphism.

Proof. We know by the Frobenius theorem (see for instance [40]), that for any $x \in \Omega \setminus \{a\}$, there is a map $\psi_U : U \rightarrow \Omega \setminus \{a\}$, defined on an open set U such that $0 \in U$, $\psi_U(0) = x$ and $\psi(s, \theta)$ satisfies

$$(5.2) \quad \begin{cases} \partial_s \psi_U = e^{\Phi \circ \psi_U} v \circ \psi_U \\ \partial_\theta \psi_U = e^{\Phi \circ \psi_U} u \circ \psi_U. \end{cases}$$

Moreover, ψ_U is unique in the sense that two such maps are equal on the intersection of their open sets U of definition. Notice also that such an application ψ_U is a local diffeomorphism. Let (U_x, ψ_x) be the maximal solution: U_x is defined as the union of open sets U such that there is a map ψ_U satisfying $\psi_U(0) = x$ and (5.2). By uniqueness in the Frobenius theorem, $\psi_x(y) = \psi_U(y)$ is perfectly defined independently of U and satisfies $\psi_x(0) = x$ and (5.2). We aim at proving that up to factorisation and translation on the domain of definition, ψ_x is the map we are looking for.

All along the proof we denote φ_{X_1} and φ_{X_2} the flows along the vector fields $X_1 := e^\Phi v$ and $X_2 := e^\Phi u$. Notice that the partial maps of ψ_x are paths along the flow: if $x_0 = \psi_x(s_0, \theta_0)$,

$$\psi_x(s_0 + s, \theta_0) = \varphi_{X_1}^s(x_0) \text{ and } \psi_x(s_0, \theta_0 + \theta) = \varphi_{X_2}^\theta(x_0).$$

Notice also that we can extend X_1 and X_2 by smooth vector-fields on an open set $\tilde{\Omega}$ such that $\bar{\Omega} \subset \tilde{\Omega}$, and that there is no stationary point on $\tilde{\Omega} \setminus \{a\}$ for φ_{X_1} and φ_{X_2} . In particular, the flows are perfectly defined up to the boundary in $\bar{\Omega}$. We recall that $\partial\Omega$ is a cycle of φ_{X_2} by assumption. Moreover, the only stationary point of φ_{X_1} and φ_{X_2} is a since by definition of Φ as a Green function on a , the only vanishing point of e^Φ is a .

STEP 1 : *We prove that for any x , ψ_x is surjective*

Notice that if $x, y \in \Omega \setminus \{a\}$, we have that

$$\psi_x(U_x) \cap \psi_y(U_y) \neq \emptyset \Rightarrow \psi_x(U_x) = \psi_y(U_y).$$

Indeed, assume that $z = \psi_x(u) = \psi_y(v)$ for $u \in U_x$ and $v \in U_y$. By uniqueness in the Frobenius theorem, and by maximality of (U_x, ψ_x) and (U_y, ψ_y) , one has $U_x = U_y + \{b - a\}$ and $\psi_x(z) = \psi_y(z + v - u)$ for any $z \in U_x$. Since for any $y \in \Omega$, ψ_y is a local diffeomorphism, then if $y \in \psi_x(U_x)$, there is an open set U such that $\psi_y(U)$ is an open neighbourhood of y in $\psi_x(U_x)$. Therefore $\psi_x(U_x)$ is open. Denoting $C = \{C(x); x \in \Omega \setminus \{a\}\}$, where

$$C(x) = \{y \in \Omega \setminus \{a\}; \psi_x(U_x) = \psi_y(U_y)\},$$

we can write

$$\Omega \setminus \{a\} = \bigcup_{C(x) \in C} \psi_x(U_x)$$

as a disjoint union of open sets. By connectedness, the cardinal of C is 1 and $\psi_x(U_x) = \Omega \setminus \{a\}$ for any $x \in \Omega \setminus \{a\}$.

From now on, we drop the index x in the definition of $U := U_x$ and $\psi := \psi_x$

STEP 2 : *We prove the existence of β such that $U = (-\infty, \beta) \times \mathbb{R}$ and such that we can extend ψ on $(-\infty, \beta] \times \mathbb{R}$ so that $\psi(\{\beta\} \times \mathbb{R}) = \partial\Omega$ and ψ is a local homeomorphism up to the boundary.*

We first prove that for any $(s_0, \theta_0) \in U$, we have $\{s_0\} \times \mathbb{R} \subset U$. Indeed, let $(s_0, \theta_0) \in U$, $x_0 = \psi(s_0, \theta_0)$, and let (γ, δ) be the maximal interval such that $\gamma < \theta_0 < \delta$ and $\{s_0\} \times (\gamma, \delta) \subset U$. Let's prove that $\delta = +\infty$ (proving that $\gamma = -\infty$ is analogous).

Since $\bar{\Omega}$ is compact, up to a subsequence of $\theta_n \rightarrow \delta$ one has $\lim_{\theta_n \rightarrow \delta} \psi(s, \theta_n) = l \in \bar{\Omega}$. If $l \in \Omega \setminus \{a\}$ and $\delta < +\infty$, by the Frobenius theorem, one can extend ψ by a map $\hat{\psi}$ satisfying $\hat{\psi}(s, \delta) = l$ and (5.2), defined on an open neighbourhood of (s, δ) and we contradict the maximality of δ . Therefore, $l \in \partial\Omega \cup \{a\}$ or $\delta = +\infty$. Now, we recall that $\psi(s_0, \theta) = \varphi_{X_2}^{\theta - \theta_0}(x_0)$ is the flow associated to X_2 . If $l = a$, then $\lim_{\theta \rightarrow \delta} \psi(s, \theta) = a$ and since a is a stationary point, we have that $\delta = +\infty$. Similarly, if $l \in \partial\Omega$, since $\partial\Omega$ is a cycle of φ_{X_2} , we must have $\delta = +\infty$. In all the cases, $\delta = +\infty$.

Now, let $\beta = \sup\{s \in \mathbb{R}; \{s\} \times \mathbb{R} \subset \Omega\}$.

We prove that $\beta < +\infty$ and that for any $\theta \in \mathbb{R}$, $\lim_{z \rightarrow (\beta, \theta)} \psi(y)$ exists and belongs to $\partial\Omega$.

The vector field $X_1 = e^\Phi v$ is transversal to $\partial\Omega$ (and outward pointing on $\partial\Omega$). Let $y \in \partial\Omega$. Let $F :]-\varepsilon, \varepsilon[\times]-\varepsilon, \varepsilon[\rightarrow V$ be a straightening of X_1 at the neighbourhood V of $y = F(0, 0)$: ie F is a diffeomorphism and $F^*(X_1) = \partial_{x_1}$. We have that $F(]-\varepsilon, 0[\times]-\varepsilon, \varepsilon[) \subset \Omega$. Let $(s_0, \theta_0) \in U$ be such that $x_0 = \psi(s_0, \theta_0) = F(-\frac{\varepsilon}{2}, 0)$. Recalling that $\psi(s, \theta_0) = \varphi_{X_1}^{s - s_0}(x_0)$, we deduce the expected result.

Now, we let $\alpha = \inf\{\tilde{\alpha} \in (-\infty, \beta); (\tilde{\alpha}, \beta) \times \mathbb{R} \subset U\}$. With the same methods as previously thanks to the minimality of α and using the flow associated to X_1 by $\psi(s, \theta) = \varphi_{X_1}^{s - s_0}(x_0)$ one can prove that $\alpha = -\infty$. This ends the proof of STEP 2.

STEP 3 : *We prove the existence of $\rho > 0$ such that $\psi(s, \theta) = \psi(s, \theta + \rho)$ for any $(s, \theta) \in U$.*

It is equivalent to prove that for any $x \in \psi(U)$, we have that $\varphi_{X_2}(x)$ is a cycle.

Let $y \in \partial\Omega$. Since $\varphi_{X_2}(y)$ is a cycle, there is a minimal $\rho > 0$ such that $\varphi_{X_2}^\rho(y) = y$. We aim at proving that $\varphi_{X_2}^\rho(x) = x$ for any $x \in \psi(U)$.

Let $x_0 \in \psi(U)$ and $(s_0, \theta_0) \in U$ be such that $x_0 = \psi(s_0, \theta_0)$. From the computations we made before Proposition 5.1, we have that $[X_1, X_2] = 0$ in $\Omega \setminus \{a\}$. Therefore, the flows φ_{X_1} and φ_{X_2} commute. Then by STEP 2,

$$\varphi_{X_2}^\rho(x_0) = \varphi_{X_1}^{s_0 - s} \circ \varphi_{X_2}^\rho \circ \varphi_{X_1}^{s - s_0}(x_0),$$

for any $s_0 \leq s < \beta$. By STEP 2 again, we can pass to the limit as $s \rightarrow \beta$ and we get

$$(5.3) \quad \varphi_{X_2}^\rho(x_0) = \varphi_{X_1}^{s_0 - \beta} \circ \varphi_{X_2}^\rho \circ \varphi_{X_1}^{\beta - s_0}(x_0).$$

Since $\varphi_{X_1}^{\beta-s_0}(x_0) \in \partial\Omega$, we have that $\varphi_{X_2}^\rho(\varphi_{X_1}^{\beta-s_0}(x_0)) = \varphi_{X_1}^{\beta-s_0}(x_0)$. We then get STEP 3 from (5.3).

STEP 4 : *We prove the Proposition*

We assume that $\rho > 0$ given by STEP 3 is minimal. Now we can factorize ψ into a surjective map we still denote $\psi : (-\infty, \beta] \times \mathbb{R}/\rho\mathbb{Z} \rightarrow \bar{\Omega} \setminus \{a\}$.

Let's prove that this map is now injective. Let $z_1 = (s_1, \theta_1)$ and $z_2 = (s_2, \theta_2)$ such that $\psi(z_2) = \psi(z_1)$. Then $\psi(z) = \psi(z + z_2 - z_1)$. Assume that $s_1 \leq s_2$. We then have that $\psi(\{\beta - s_2 + s_1\} \times \mathbb{R}) = \partial\Omega$. Then we must have $s_1 = s_2$. By minimality of ρ , we get that $\theta_1 = \theta_2$. Then ψ is injective. Since ψ is bijective and a local homeomorphism, it is a global homeomorphism.

Moreover, since $\psi : (-\infty, \beta) \times \mathbb{R}/\rho\mathbb{Z} \rightarrow \Omega \setminus \{a\}$ is bijective smooth and open, it is a diffeomorphism. By equation (5.2), it is also a conformal diffeomorphism. Finally, we replace ψ by $\psi(z - (0, \beta))$ and we complete the proof of Proposition 5.1. \square

We set a holomorphic function $f : \mathbb{D} \setminus \{0\} \rightarrow \Omega \setminus \{a\}$, defined by

$$(5.4) \quad f(re^{i\theta}) = \psi\left(\frac{\rho}{2\pi}(\ln r, \theta)\right),$$

where ψ is the map given by claim (5.1). The implicit formula $f(e^{\frac{2\pi w}{\rho}}) = \psi(w)$ holds and one can directly extend f to a bi-holomorphic map $f : \mathbb{D} \rightarrow \Omega$ by $f(0) = a$. This ends the proof of the Riemann mapping theorem.

Remark 5.1. *We cannot ask for low regularity of Ω to prove the result by this method for two reasons : C^1 regularity of Ω is needed to use classical construction of Ginzburg-Landau minimizers (without using the uniformization map as in the current paper for Weil-Petersson curves). Moreover, we use that the unit tangent vector field on Γ is Lipschitz in order to apply the Cauchy-Lipschitz theorem and define the flow of vector fields. Using some methods from dynamical systems, it may be possible to prove the existence of a closed orbit in the interior, which would play the same role as the boundary orbit and would permit to conclude.*

APPENDIX A. TECHNICAL LEMMAS

The result of this section are well known, we prove it in a way more adapted to our paper for the sake of completeness. First we need a basic estimate on the Poisson Kernel

Lemma A.1. *Let $M > 0$, then*

$$(A.1) \quad \lim_{r \rightarrow 1^-} (1-r) \int_{\delta}^{2\pi - M(1-r)} \frac{1}{1+r^2-2r\cos(\theta)} d\theta = 8 \arctan\left(\frac{1}{M}\right).$$

Proof. We fix $M > 0$ and we set $\delta = M(1 - r)$, then we have

$$\begin{aligned}
(1-r) \int_{\delta}^{2\pi-\delta} \frac{1}{1+r^2-2r\cos(\theta)} d\theta &= (1-r) \int_{-\pi+\delta}^{\pi-\delta} \frac{1}{1+r^2+2r\cos(\theta)} d\theta \\
&= 2(1-r) \int_0^{\pi-\delta} \frac{1}{1+r^2+2r\cos(\theta)} d\theta \\
&= 2(1-r) \int_0^{\tan(\frac{\pi-\delta}{2})} \frac{2}{1+r^2+2r\frac{1-t^2}{1+t^2}} \frac{2}{1+t^2} dt \\
&= \frac{4(1-r)}{(1+r)^2} \int_0^{\tan(\frac{\pi-\delta}{2})} \frac{1}{1+(\frac{1-r}{1+r})^2 t^2} dt \\
&= 4(1+r) \int_0^{\frac{1-r}{1+r} \tan(\frac{\pi-\delta}{2})} \frac{1}{1+u^2} du \\
&= 4(1+r) \arctan\left(\frac{1-r}{1+r} \tan\left(\frac{\pi-\delta}{2}\right)\right),
\end{aligned}$$

finally the result follows from $\tan\left(\frac{\pi}{2} - x\right) \sim_0 \frac{1}{x}$. \square

Then we are in position to prove that the harmonic extension of a map in $H^{\frac{1}{2}}(\mathbb{S}^1, \mathbb{S}^1)$ is proper, more precisely

Lemma A.2. *Let $v \in H^{1/2}(\mathbb{S}^1, \mathbb{S}^1)$ and \tilde{v} its harmonic extension on \mathbb{D} . We denote, for $z \in \mathbb{D}$, I_z the subarc of \mathbb{S}^1 centred at $\frac{z}{|z|}$ of length $2(1 - |z|)$, and for subarc $I \subset \mathbb{S}^1$ we set*

$$v_I = \frac{1}{|I|} \int_I v d\theta.$$

Then

$$(A.2) \quad \lim_{M \rightarrow +\infty} \limsup_{r \rightarrow 1} \sup_{|z| \geq r} |\tilde{v}(z) - v_{MI_z}| = 0.$$

Moreover for every $\epsilon > 0$ there exists $M > 0$ and $r < 1$ such that if $|z| \geq r$ there exists $S_z \subset MI_z$ such that

$$|S_z| \geq \frac{|MI_z|}{2}$$

and

$$|\tilde{v}(z) - v(x)| \leq \epsilon \text{ for all } x \in S_z,$$

in particular \tilde{v} is proper.

The result and proof are sketch at beginning of section 8 of [4], we give details for the sake of completeness.

Proof. Let us remind that, $H^{1/2} \subset \text{VMO} \subset \text{BMO}$, see §1.2 [7]⁴. So we have, by Lemma 3 of [7], that, for every $M > 0$,

$$(A.3) \quad \lim_{|z| \rightarrow 1} \frac{1}{|MI_z|} \int_{MI_z} |v - v_{MI_z}| d\theta = 0 \text{ uniformly in } z.$$

⁴In this reference it is proved on \mathbb{R} but it is also true on \mathbb{S}^1 by conformal invariance of all those spaces, see corollary VI.1.3 [16]

Moreover, \tilde{v} is given by the Poisson formula,

$$\tilde{v}(z) = \int_0^{2\pi} P_z(\theta) v(e^{i\theta}) d\theta,$$

where $P_z(\theta) = \frac{1}{2\pi} \frac{1-|z|^2}{|z-e^{i\theta}|^2}$. Hence, let $M > 0$, we have, using the fact that $\int_{\mathbb{S}^1} P_z d\theta = 1$ and $P_z \geq 0$, that

$$|\tilde{v}(z) - v_{MI_z}| \leq \int_{MI_z} P_z |v - v_{MI_z}| d\theta + \int_{(MI_z)^c} P_z |v - v_{MI_z}| d\theta$$

Let $\epsilon > 0$, thanks to (A.1) and the fact that $|v| = 1$ and $|v_{MI_z}| \leq 1$, we have, for M large enough, that

$$\lim_{|z| \rightarrow 1} \int_{(MI_z)^c} P_z |v - v_{MI_z}| d\theta \leq \frac{\epsilon}{4}$$

Once M is fixed, since v is in VMO, we have, thanks to (A.3)), that

$$\int_{MI_z} P_z |v - v_{MI_z}| d\theta \leq \frac{2M}{\pi |MI_z|} \int_{MI_z} |v - v_{MI_z}| d\theta \leq \frac{\epsilon}{4},$$

which proves the first part of the lemma.

Moreover, v satisfies the John-Nirenberg theorem, see (3') of [23], there exist $\beta, C > 0$ independent of v , such for any arc $I \subset \mathbb{S}^1$ we have

$$\frac{1}{|I|} \int_I e^{\frac{\beta|v-v_I|}{\kappa}} d\theta \leq C,$$

where

$$\kappa = \sup_{I' \subset I} \frac{1}{|I'|} \int_{I'} |v - v_{I'}| d\theta,$$

In particular,

$$|\{x \in MI_z \mid |v(x) - v_{MI_z}| \geq \frac{\epsilon}{2}\}| \leq C |MI_z| e^{-\frac{\beta\epsilon}{2\kappa}}.$$

Hence by (A.3), for $1 - |z|$ small enough, there exists $x \in MI_z$ such that

$$(A.4) \quad |g_{MI_z} - g(x)| \leq \frac{\delta}{2}$$

which achieves the proof of the lemma. \square

APPENDIX B. NOTE ON THE EXTENDED DEFINITION OF $H^{\frac{1}{2}}(\Gamma)$

In this section, we comment and justify the general definition of $H^{\frac{1}{2}}(\Gamma, \mathbb{S}^1)$ (see definition 2.3).

First, let $\Omega \subset \mathbb{R}^2$ be a bounded open set and $f : \mathbb{D} \rightarrow \Omega$ be a uniformization. Then for any $u : \Omega \rightarrow \mathbb{R}^n$, we have

$$(B.1) \quad u \circ f \in H_0^1(\mathbb{D}, \mathbb{R}^n) \Leftrightarrow u \in H_0^1(\Omega, \mathbb{R}^n).$$

Notice that it is equivalent to $H_0^1(\mathbb{D}, \mathbb{R}^n) = H_0^1(\mathbb{D}, |f'|^2, \mathbb{R}^n)$ by pullback of functions with f . In fact (B.1) is just an application of the conformal invariance of the Dirichlet energy and the classical Poincaré inequality on \mathbb{D} and on Ω :

$$\int_{\mathbb{D}} |u \circ f|^2 \leq C_{\mathbb{D}} \int_{\mathbb{D}} |\nabla(u \circ f)|^2 = C_{\mathbb{D}} \int_{\Omega} |\nabla u|^2 \leq C_{\mathbb{D}} \|u\|_{H^1(\Omega)}$$

$$\int_{\Omega} |u|^2 \leq C_{\Omega} \int_{\Omega} |\nabla u|^2 = C_{\Omega} \int_{\mathbb{D}} |\nabla (u \circ f)|^2 \leq C_{\Omega} \|u \circ f\|_{H^1(\mathbb{D})}^2$$

for any $u \in C_c^{\infty}(\Omega)$. Notice that the Poincaré inequality in Ω can also be seen as the classical Hardy inequality since $\sup_{z \in \mathbb{D}} |f'(z)|^2 (1 - |z|)^2 < +\infty$ by Koebe's theorem.

Now, we would ideally like to have $H^1(\mathbb{D}, |f'(z)|^2) = H^1(\mathbb{D})$ too in order to give a sense to definition 2.3. Indeed, we would like the function $g : \Gamma \rightarrow \mathbb{R}^n$ to be some trace of a $H^1(\Omega)$ function u . Pulling back to \mathbb{D} by f , this means that $g \circ f : \Gamma \rightarrow \mathbb{R}^n$ is some trace of the function $u \circ f \in H^1(\mathbb{D}, |f'(z)|^2)$. However, $g \circ f : \Gamma \rightarrow \mathbb{R}^n$ is well-defined as a $H^{\frac{1}{2}}(\mathbb{S}^1)$ trace only if it is a trace of a $H^1(\mathbb{D})$ function.

Then, we assume that $g \circ f$ admits some extension in $H^1(\mathbb{D}, |f'(z)|^2) \cap H^1(\mathbb{D})$ since those sets are not a priori equal. We then a priori ask more restrictions for being in $H^{\frac{1}{2}}(\Gamma, \mathbb{R}^n)$ than in definition 2.3:

Definition B.1 ($H^{\frac{1}{2}}(\Gamma, \mathbb{R}^n)$ maps). *Let $n \geq 1$, Γ be a quasicircle enclosing a domain Ω , and $f : \mathbb{D} \rightarrow \Omega$ a uniformization. We say that $g : \Gamma \rightarrow \mathbb{R}^n$ is in $H^{\frac{1}{2}}(\Gamma, \mathbb{R}^n)$ if $g \circ f \in H^{\frac{1}{2}}(\mathbb{S}^1, \mathbb{R}^n)$ and if $g \circ f$ admits an extension $v \in H^1(\mathbb{D}, \mathbb{R}^n)$ such that*

$$(B.2) \quad \int_{\mathbb{D}} |v|^2(z) |f'(z)|^2(z) dz < +\infty$$

that is to say $v \circ f^{-1} \in H^1(\Omega, \mathbb{S}^1)$.

Once we know that such an extension v exists, then by (B.1), the set of $H^1(\mathbb{D})$ extensions of $g \circ f$: $\{v\} + H_0^1(\mathbb{D})$ is equal to the pullback of the natural set of $H^1(\Omega)$ extensions of g : $\{v \circ f^{-1}\} + H_0^1(\Omega)$. This makes the definition consistent. In particular, the harmonic extension v of $g \circ f$ satisfies $v \circ f^{-1} \in H^1(\Omega)$.

In the special case where g is bounded, then (B.2) is automatically satisfied since $H^1 \cap L^{\infty}(\mathbb{D}, |f'(z)|^2) = H^1 \cap L^{\infty}(\mathbb{D})$ and the harmonic extension v of $g \circ f$ is also bounded. This shows that definition 2.3 makes sense.

APPENDIX C. WENTE TYPE ESTIMATE

We give a proof of a Wente type lemma, which is inspired of Theorem A.4 of [24] (see also Theorem 1.100 [35] and proof of Theorem 1.2 [25]).

Lemma C.1. *There exists $C > 0$ such that if $f \in L^1(\mathbb{D})$ such that*

$$\sup_{\mathbb{D}} |f(x)(1 - |x|)^2| < +\infty,$$

then, there exists $\phi \in W_0^{1,2}(\mathbb{D}) \cap L^{\infty}(\mathbb{D})$ such that

$$\Delta \phi = f,$$

and

$$\|\phi\|_{\infty} + \|\nabla \phi\|_2 \leq C(\|f\|_1 + \|f(1 - |x|)^2\|_{\infty}).$$

Proof. It is classical that, by integration by part, the L^{∞} -estimate implies the L^2 -estimate. Hence we set

$$\phi(x) = \int_{\mathbb{D}} G(x, y) f(y) dy,$$

where G is the Green function of the Laplacian. Then, we remark that the singularity of the Green function can be decomposed as a sum of bump functions with dyadic support, see Theorem A.4 of [24]: there exists $C > 0$ such that

$$-C \leq G(x, y) - l_x(y) \leq C,$$

with

$$l_x(y) = - \sum_{j=1}^{\infty} \Theta \left(\frac{2^j(x-y)}{1-|x|} \right),$$

where $\Theta \in C_c^\infty(\mathbb{R}, [0, 1])$ is a bump function, such as $\text{supp}(\Theta) \subset [-\frac{1}{8}, \frac{1}{8}]$ and $\Theta \equiv 1$ on $[-\frac{1}{16}, \frac{1}{16}]$. Then

$$|\phi(x)| \leq C \|f\|_1 + \sum_{j=1}^{\infty} \int_{\mathbb{D}} \Theta \left(\frac{2^j(x-y)}{1-|x|} \right) f(y) dy.$$

Then we estimate each integral of the right hand

$$\begin{aligned} \int_{\mathbb{D}} \Theta \left(\frac{2^j(x-y)}{1-|x|} \right) f(y) dy &\leq \int_{B(x, 2^{-j-3}(1-|x|))} f(y) dy \\ \text{(C.1)} \quad &\leq \|f(1-|x|)^2\|_\infty \int_{B(x, 2^{-j-3}(1-|x|))} \frac{1}{(1-|y|)^2} dy \\ &\leq \frac{\pi}{(2^{j+3}-1)^2} \|f(1-|x|)^2\|_\infty \end{aligned}$$

Finally we just have to sum this inequality to get the result. \square

REFERENCES

- [1] Fabrice Bethuel, Haïm Brezis, and Frédéric Hélein. Asymptotics for the minimization of a Ginzburg-Landau functional. *Calc. Var. Partial Differential Equations*, 1(2):123–148, 1993.
- [2] Fabrice Bethuel, Haïm Brezis, and Frédéric Hélein. *Ginzburg-Landau vortices*. Modern Birkhäuser Classics. Birkhäuser/Springer, Cham, 2017. Reprint of the 1994 edition [MR1269538].
- [3] A. Beurling and L. Ahlfors. The boundary correspondence under quasiconformal mappings. *Acta Math.*, 96:125–142, 1956.
- [4] Christopher J. Bishop. Function theoretic characterizations of weil-petersson curves.
- [5] Christopher J. Bishop. Weil-petersson curves, beta-numbers, and minimal surfaces.
- [6] Anne Boutet de Monvel-Berthier, Vladimir Georgescu, and Radu Purice. A boundary value problem related to the Ginzburg-Landau model. *Comm. Math. Phys.*, 142(1):1–23, 1991.
- [7] H. Brezis and L. Nirenberg. Degree theory and BMO. I. Compact manifolds without boundaries. *Selecta Math. (N.S.)*, 1(2):197–263, 1995.
- [8] Haïm Brezis, Frank Merle, and Tristan Rivière. Quantization effects for $-\Delta u = u(1-|u|^2)$ in \mathbf{R}^2 . *Arch. Rational Mech. Anal.*, 126(1):35–58, 1994.
- [9] Luis A. Caffarelli and Avner Friedman. Convexity of solutions of semilinear elliptic equations. *Duke Math. J.*, 52(2):431–456, 1985.
- [10] K. C. Chang and J. Q. Liu. A prescribing geodesic curvature problem. *Math. Z.*, 223(2):343–365, 1996.
- [11] Shiing-shen Chern. An elementary proof of the existence of isothermal parameters on a surface. *Proc. Amer. Math. Soc.*, 6:771–782, 1955.
- [12] Otis Chodosh and Christos Mantoulidis. Minimal surfaces and the Allen-Cahn equation on 3-manifolds: index, multiplicity, and curvature estimates. *Ann. of Math. (2)*, 191(1):213–328, 2020.
- [13] Manuel del Pino and Patricio L. Felmer. On the basic concentration estimate for the Ginzburg-Landau equation. *Differential Integral Equations*, 11(5):771–779, 1998.
- [14] Alberto Farina and Petru Mironescu. Uniqueness of vortexless Ginzburg-Landau type minimizers in two dimensions. *Calc. Var. Partial Differential Equations*, 46(3-4):523–554, 2013.

- [15] Martin Flucher. Extremal functions for the Trudinger-Moser inequality in 2 dimensions. *Comment. Math. Helv.*, 67(3):471–497, 1992.
- [16] John B. Garnett. *Bounded analytic functions*, volume 236 of *Graduate Texts in Mathematics*. Springer, New York, first edition, 2007.
- [17] Marco A. M. Guaraco. Min-max for phase transitions and the existence of embedded minimal hypersurfaces. *J. Differential Geom.*, 108(1):91–133, 2018.
- [18] F. Hélein. *Harmonic maps, conservation laws and moving frames*, volume 150 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, second edition, 2002. Translated from the 1996 French original, With a foreword by James Eells.
- [19] John Hamal Hubbard. *Teichmüller theory and applications to geometry, topology, and dynamics. Vol. 1*. Matrix Editions, Ithaca, NY, 2006. Teichmüller theory, With contributions by Adrien Douady, William Dunbar, Roland Roeder, Sylvain Bonnot, David Brown, Allen Hatcher, Chris Hruska and Sudeb Mitra, With forewords by William Thurston and Clifford Earle.
- [20] R. Ignat and R. L. Jerrard. Renormalized energy between vortices in some Ginzburg-Landau models on 2-dimensional Riemannian manifolds. *Arch. Ration. Mech. Anal.*, 239(3):1577–1666, 2021.
- [21] Radu Ignat, Luc Nguyen, Valeriy Slastikov, and Arghir Zarnescu. On the uniqueness of minimisers of Ginzburg-Landau functionals. *Ann. Sci. Éc. Norm. Supér. (4)*, 53(3):589–613, 2020.
- [22] Miao Jin, Xianfeng Gu, Ying He, and Yalin Wang. *Conformal geometry*. Springer, Cham, 2018. Computational algorithms and engineering applications.
- [23] F. John and L. Nirenberg. On functions of bounded mean oscillation. *Comm. Pure Appl. Math.*, 14:415–426, 1961.
- [24] Tobias Lamm and Longzhi Lin. Estimates for the energy density of critical points of a class of conformally invariant variational problems. *Adv. Calc. Var.*, 6(4):391–413, 2013.
- [25] Paul Laurain and Romain Petrides. Existence of min-max free boundary disks realizing the width of a manifold. *Adv. Math.*, 352:326–371, 2019.
- [26] Alexis Michelat and Yilin Wang. Loewner energy as the renormalised energy of moving frames and surfaces of finite willmore energy bounding weil-petersson curves. *preprint*, 2021.
- [27] Vincent Millot and Adriano Pisante. Symmetry of local minimizers for the three-dimensional Ginzburg-Landau functional. *J. Eur. Math. Soc. (JEMS)*, 12(5):1069–1096, 2010.
- [28] Édouard Oudet, Chiu-Yen Kao, and Braxton Osting. Computation of free boundary minimal surfaces via extremal Steklov eigenvalue problems. *ESAIM Control Optim. Calc. Var.*, 27:Paper No. 34, 30, 2021.
- [29] Frank Pacard and Tristan Rivière. *Linear and nonlinear aspects of vortices*, volume 39 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 2000. The Ginzburg-Landau model.
- [30] Ch. Pommerenke. *Boundary behaviour of conformal maps*, volume 299 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992.
- [31] Tristan Rivière. Asymptotic analysis for the Ginzburg-Landau equations. *Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8)*, 2(3):537–575, 1999.
- [32] Tristan Rivière. Weak immersions of surfaces with L^2 -bounded second fundamental form. In *Geometric analysis*, volume 22 of *IAS/Park City Math. Ser.*, pages 303–384. Amer. Math. Soc., Providence, RI, 2016.
- [33] Etienne Sandier and Sylvia Serfaty. *Vortices in the magnetic Ginzburg-Landau model*, volume 70 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 2007.
- [34] Reiner Michael Schätzle. Estimation of the conformal factor under bounded Willmore energy. *Math. Z.*, 274(3-4):1341–1383, 2013.
- [35] Stephen Semmes. A primer on Hardy spaces, and some remarks on a theorem of Evans and Müller. *Comm. Partial Differential Equations*, 19(1-2):277–319, 1994.
- [36] M. Struwe. On the asymptotic behavior of minimizers of the Ginzburg-Landau model in 2 dimensions. *Differential Integral Equations*, 7(5-6):1613–1624, 1994.
- [37] Michael Struwe. Erratum: “On the asymptotic behavior of minimizers of the Ginzburg-Landau model in 2 dimensions”. *Differential Integral Equations*, 8(1):224, 1995.

- [38] Leon A. Takhtajan and Lee-Peng Teo. Weil-Petersson metric on the universal Teichmüller space. *Mem. Amer. Math. Soc.*, 183(861):viii+119, 2006.
- [39] Yilin Wang. Equivalent descriptions of the Loewner energy. *Invent. Math.*, 218(2):573–621, 2019.
- [40] Frank W. Warner. *Foundations of differentiable manifolds and Lie groups*, volume 94 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1983. Corrected reprint of the 1971 edition.
- [41] Xiaotian Yin, Junfei Dai, Shing-Tung Yau, and Xianfeng Gu. Slit map: conformal parameterization for multiply connected surfaces. In *Advances in geometric modeling and processing*, volume 4975 of *Lecture Notes in Comput. Sci.*, pages 410–422. Springer, Berlin, 2008.

(Paul Laurain) INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ DE PARIS, BÂTIMENT SOPHIE GERMAIN, CASE 7052, 75205 PARIS CEDEX 13, FRANCE & DMA, ECOLE NORMALE SUPÉRIEURE, CNRS, PSL RESEARCH UNIVERSITY, 75005 PARIS.

Email address: paul.laurain@imj-prg.fr

(Romain Petrides) INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ DE PARIS, BÂTIMENT SOPHIE GERMAIN, 75205 PARIS CEDEX 13, FRANCE

Email address: romain.petrides@imj-prg.fr