

Clausius Equation for Horizons in $F(R)$ Gravity

Bohuslav Matouš

Department of Theoretical Physics and Astrophysics

Faculty of Science, Masaryk University

Kotlářská 2, 611 37, Brno

Czech Republic

E-mail: bmatous@mail.muni.cz

North-Bohemian Observatory and Planetarium in Teplice

Koperníkova 3062, 415 01, Teplice

Czech Republic

December 20, 2021

Abstract

Covariant Hamiltonian formulation of $F(R)$ gravity is used to find variation of surface Hamiltonian which leads to formulation of temperature and entropy of a horizon. These thermodynamic quantities are then used to establish an analogy between Einstein equation and Clausius equation. The obtained results are compared to those from classical General Relativity.

1 Introduction

The idea of interpreting Einstein equations as a thermodynamic equation originated in work of Jacobson [1] where he found analogy between Einstein equation and Clausius equation using Rindler horizons and Raychadhauri equation. His method was generalised for $F(R)$ gravitational theory in later works [2][3][22] where this gravitational theory was seen as non-equilibrium thermodynamics and the contribution from new gravity term was interpreted as an addition to entropy which caused a dissipative nature of gravity.

Other approach to study thermodynamic aspects of gravity was introduced in works of Padmanabhan and his colleagues [4][5][6]. Their approach focuses strongly on Lagrangian and variation principle. It is known that Lagrangian for general relativity can be separated into two parts, the quadratic bulk term and a surface term. When deriving the equations of motion with fixed variation of metric tensor and its conjugated momentum, the surface term vanishes, but in general it can be shown that there is a relationship between the two parts of Lagrangian [7][8] and that both of them contain information about space-time configuration. This brought interest to the surface Lagrangian which was until then eclipsed by the bulk Lagrangian. It was found in [9] that action of the surface Lagrangian on a horizon leads directly to the temperature and the entropy of such horizon. This has put the surface Lagrangian into spotlight of thermodynamic research of gravity and this surface Lagrangian has been since then thoroughly examined.

The description of gravity is mostly done in Lagrangian formalism that is caused by the principle of covariance. If one wants to switch from the Lagrangian formalism

to Hamiltonian formalism using classical Hamiltonian theory the covariance gets broken because the momentum is defined by partial derivative of a Lagrangian with respect to time derivative of metric. To preserve the covariance, a covariant Hamiltonian formulation of field theory has to be used. Weyl-De Donder theory [10][11][12] is such theory. The Weyl-De Donder theory defines the momentum as a partial derivative of a Lagrangian with respect to each partial derivative of coordinate. That basically means that the momenta have dimensions one higher than the coordinates.

Covariant Hamiltonian formulation of gravity was presented by Hořava [13] and by Parattu [14]. In the latter, it was shown that the situation simplifies significantly when the role of coordinate is not acted by simple metric but a slightly different coordinate $f^{ik} = \sqrt{-g}g^{ik}$. This coordinate is not a new one but it has historical significance [15].

As mentioned in the first paragraph, the original Jacobson's theory was generalised for $F(R)$ gravity. This article works also with the $F(R)$ theory and aims to find Clausius equations on horizons but it is using the Padmanabhan's approach and examines the covariant Hamiltonians of horizons using method presented in [14]. To do so, covariant formulation of $F(R)$ gravity presented in [16] is used.

This paper uses the East Coast convention with metric signature $(-, +, +, +)$ and Latin indices running over 0..3 interval while the Greek ones over 1..3. The fundamental constants c, G, \hbar, k_B are treated as equal to one. The second section of this article reviews the covariant formulation of $F(R)$ gravity and introduces the method used to examine the horizons. Third and fourth sections deals with the example metrics and provides their thermodynamic results. Fifth section is conclusion which summarises the results.

2 Review of Covariant Formulation of $F(R)$ Gravity

The main objective of this article is to find thermodynamic equations on a horizon. To do so, the method from [14] is used but it is generalized to the case of $F(R)$ gravity. The covariant formulation of $F(R)$ gravity was done in [16]. Let's summarise it quickly to provide overview of the situation. In the given theory, it is an advantage to introduce new two scalar fields A and B and write the Lagrangian density as

$$\mathcal{L} = \frac{1}{16\pi} \sqrt{-g} [F(B) + A(R - B)] , \quad (1)$$

the equation of motion for B leads to relation $A = F'(B)$ where prime denotes derivative with respect to B . This equation of motion can be used to simplify the Lagrangian density by removing the influence of scalar field A completely, leaving us with the Lagrangian density

$$\mathcal{L} = \frac{1}{16\pi} \sqrt{-g} [F(B) + F'(R - B)] . \quad (2)$$

Solving the equation of motion for scalar field B for this Lagrangian leads to $R = B$ with condition $F''(B) \neq 0$. Breaking on this condition leads to standard General Relativity. The next step is to split the Lagrangian density into two parts: a bulk term and a surface term. The bulk term is

$$16\pi \mathcal{L}_{bulk} = \sqrt{-g} F'(B) (g^{ba} \Gamma_{ad}^c \Gamma_{bc}^d - g^{bc} \Gamma_{ad}^a \Gamma_{bc}^d) - \sqrt{-g} F''(B) (\partial_c B) (g^{bd} \Gamma_{bd}^c - g^{bc} \Gamma_{bd}^d) + \sqrt{-g} [F(B) - F'(B)B] . \quad (3)$$

The density of surface Lagrangian, which is the more important one for the analysis of thermodynamics of horizon, has the form of

$$\mathcal{L}_{sur} = \frac{1}{16\pi} \partial_c [\sqrt{-g} F'(B) (g^{bd} \Gamma_{bd}^c - g^{bc} \Gamma_{bd}^d)] . \quad (4)$$

The bulk term might be then used for formulation of covariant Hamiltonian. To do so, the coordinate is changed and instead of metric tensor, tensor f^{ik} , where $f^{ik} = \sqrt{-g}g^{ik}$, is used. Momentum conjugate to f^{ik} is

$$N_{ik}^c = -\frac{F'(B)}{16\pi}\Gamma_{ik}^c + \frac{F'(B)}{32\pi}(\Gamma_{id}^d\delta_k^c + \Gamma_{kd}^d\delta_i^c) + \frac{F''(B)}{32\pi}(\partial_i B\delta_k^c + \partial_k B\delta_i^c) + \frac{F''(B)}{32\pi}\partial_d B f^{dc} f_{ik}, \quad (5)$$

and momentum conjugate to B is

$$p^n = \frac{F''}{16\pi}(\Gamma_{bd}^b f^{dn} - \Gamma_{bd}^n f^{bd}), \quad (6)$$

which can be used to express surface Lagrangian density (4) as

$$\mathcal{L}_{sur} = -\partial_c \left(\frac{F'}{F''} p^c \right). \quad (7)$$

This is the starting point of our analysis. The surface Lagrangian density is used to find thermodynamic variables for a horizon. This can be done, since the action of surface Lagrangian on the horizon is proportional to TS [17]. Thus, when working with static metric, the following definition [14][17] can be used

$$H_{sur} = -\frac{\partial A_{sur}}{\partial \tau} = TS. \quad (8)$$

Now let's find the formula for variation of surface Hamiltonian. Using formula from [14] stating that if $\mathcal{L}_{sur} = \partial_c(\sqrt{-g}V^c)$ then $\delta H_{sur} = -\int \delta(\sqrt{-g}V^n)$ where n is a coordinate normal to the surface. Using this for our Lagrangian (7) yields

$$\delta H_{sur} = \int d^2 x_{\parallel} F' \delta \frac{p^n}{F''} + \int d^2 x_{\parallel} \frac{p^n}{F''} \delta F', \quad (9)$$

where the integral is over the variables of the horizon only, these coordinates are denoted as x_{\parallel} . In the GR case [14], the variation of surface Hamiltonian took form $\delta H \sim \delta(\text{coordinate} \cdot \text{momentum})$. In $F(R)$ gravity case, the form is similar $\delta H \sim \delta(\text{function of coordinate} \cdot \text{momentum})$. To get the GR scheme from the $F(R)$ one, $\frac{F'}{F''} = B$ condition has to be fulfilled. Which is so only in case that $F = \alpha B^2 + \beta$ for any α, β .

The problematic term of (9) is F'' which appears in denominators. It is non-zero from definition but it makes the form of Hamiltonian's variation complicated. If one recalls the definition (6) a substitution is clearly visible. Let's define vector q^n , metric part of momentum conjugate to B as

$$q^n = 16\pi \frac{p^n}{F''} = \Gamma_{bd}^b f^{dn} - \Gamma_{bd}^n f^{bd}, \quad (10)$$

this allows us to formulate variation of surface Hamiltonian in a more convenient form as

$$\delta H_{sur} = \frac{1}{16\pi} \int d^2 x_{\parallel} F' \delta q^n + \frac{1}{16\pi} \int d^2 x_{\parallel} q^n \delta F'. \quad (11)$$

From this term, temperature and entropy shall be obtained. Vector q^n brings another advantage when compared to results using GR [14] because $q^n = (-\sqrt{-g}V^c)^{GR}$. Defining Hamiltonian density \mathcal{H}_{sur} as $\delta H_{sur} = \int d^2 x_{\parallel} \delta \mathcal{H}_{sur}$ leads to

$$\mathcal{H}_{sur}^{F(R)} = \mathcal{H}_{sur}^{GR} F' \sim T^{GR} (S^{GR} F'), \quad (12)$$

from this it is visible that expected result is that the definition of temperature remains the same and the result for entropy shall be extended with a factor F' .¹

Having found variation of Hamiltonian as $T\delta S + S\delta T$, the next step is to look for an analogy between Clausius and Einstein equation. To do so, the Einstein equations are evaluated on the horizon and modified to a form containing $T\delta S$ member. In the $F(R)$ gravity case, we have these following equations of motion

$$G_{ik} = \frac{1}{2} \frac{F}{F'} g_{ik} - \frac{1}{2} R g_{ik} - \frac{1}{F'} \Delta F' g_{ik} + \frac{1}{F'} \nabla_k \nabla_i F' + \frac{8\pi}{F'} T_{ik}, \quad (13)$$

$$B = R, \quad (14)$$

where ∇_i denotes covariant derivation and Δ is the Laplacian $\Delta = \nabla^i \nabla_i$. The second equation can be applied and the first equation can be rewritten into the following form

$$G_{ik} = \frac{8\pi}{F'} T_{ik} + \frac{1}{2} \mathcal{U} g_{ik} + \frac{1}{F'} \mathcal{V}_{ik}, \quad (15)$$

where $\mathcal{U} = \frac{F}{F'} - R$ is a scalar function of R and its form depends solely on $F(R)$, and $\mathcal{V}_{ik} = \Delta F' g_{ik} - \nabla_k \nabla_i F'$ is tensor part and depends on the metric as well as function $F(R)$.

Once the Einstein equations are evaluated on the horizon the knowledge of $T\delta S$ resulting from (11) can be applied and the equations are expressed in the form of $T\delta S = \dots$. Then the thermodynamic interpretation of the members on the right hand side shall be performed to reveal the Clausius equation.

The next two sections deal with spherical symmetric metric and general static spacetime metric and provide resulting Clausius equations.

3 Spherical case

The first case which is analysed is the spherically symmetric case. The metric is taken in the following form

$$ds^2 = -a dt^2 + \frac{1}{a} dr^2 + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\phi^2, \quad (16)$$

where a is a function of r which satisfies conditions $a(r_h) = 0$ and $a'(r_h) = 2\kappa \neq 0$ for a parameter r_h and surface gravity κ . The normal coordinate in this case is radius r . Vector q^i , the metric part of the B -momentum, can be easily calculated using (10) as

$$q^r = \sin \vartheta (4ar + a'r^2). \quad (17)$$

Substituting this vector q^n to equation (11) leads to the variation of surface Hamiltonian in form

$$\delta H_{sur} = \frac{1}{16\pi} \int d\vartheta \int d\phi \sin \vartheta [F' (4\kappa r_h \delta r_h + 2r_h^2 \delta \kappa) + \delta F' (2\kappa r_h^2)]. \quad (18)$$

After the integration, it can be modified to the simplified form using the formula for surface area $A = 4\pi r_h^2$

$$\delta H_{sur} = \frac{\kappa}{2\pi} \delta \left(F' \frac{A}{4} \right) + \delta \left(\frac{\kappa}{2\pi} \right) F' \frac{A}{4}, \quad (19)$$

¹This addition to entropy is also expected. When using the entropy tensor $P_{cd}^{ab} = \frac{\partial \mathcal{L}}{\partial R_{ab}^{cd}}$ [17] it is easily visible that for $F(R)$ gravity it takes form of $P_{cd}^{ab} = F' P_{cd}^{ab}|_{GR}$ so the entropy tensor has the same factor F' comparing to General Relativity and that is exactly what was found in (12).

which is equivalent to the relationship $\delta E = T\delta S + S\delta T$. It is clearly visible that the GR definition of temperature $T = \frac{\kappa}{2\pi}$ holds for the $F(R)$ gravity case. While the definition of entropy $S = F' \frac{A}{4}$ differs only in factor F' . This factor goes to one in the GR limit. This result concurs with our expectations from (12) as well as with the GR result from [14].

To move further, Einstein equations on the horizon need to be evaluated. The Einstein tensor G_k^i takes the GR form

$$G_t^t = G_r^r = \frac{a'r + a - 1}{r^2}, \quad G_\vartheta^\vartheta = G_\phi^\phi = \frac{2a' + a''r}{2r}. \quad (20)$$

For simplicity let's assume that tensor T_k^i takes the form of perfect fluid. The extra members of Einstein equations for $F(R)$ gravity are \mathcal{U} which cannot be further simplified without knowing $F(R)$ and the tensor \mathcal{V}_k^i which has to be evaluated as a function of $F(R)$ and its derivatives. In the process of calculating \mathcal{V}_k^i , we shall make use of the second equation of motion $B = R$ and take F as a function of scalar curvature R directly. We can also make use of spherical symmetry where the scalar curvature is a function of the radius only. Being interested only in the equations of motion on horizon allows us to neglect terms containing solely a since it vanishes on the surface. Having taken all this into consideration, the tensor \mathcal{V} can be written as

$$\mathcal{V}_i^k = -F'' \partial_{\underline{r}} R (g^{kj} \Gamma_{ji}^{\underline{r}} - \delta_i^k g^{mj} \Gamma_{mj}^{\underline{r}}), \quad (21)$$

where \underline{r} denotes that r is a coordinate and shall not be summed over. Using the Christoffel symbols and evaluating them on the horizon allows us to formulate Einstein equations on the horizon

$$\frac{2\kappa r_h - 1}{r_h^2} = \frac{8\pi}{F'} T_t^t + \frac{1}{2} \mathcal{U} - \frac{F''}{F'} \kappa \partial_r R, \quad (22)$$

$$\frac{2\kappa}{r_h} = \frac{8\pi}{F'} T_t^t + \frac{1}{2} \mathcal{U} - 2 \frac{F''}{F'} \kappa \partial_r R. \quad (23)$$

In the upcoming part, only the first one is used. At first, κ is expressed using the first equation

$$\kappa = \frac{1}{2r_h} + \frac{4\pi}{F'} T_t^t r_h + \frac{1}{4} \mathcal{U} r_h - \frac{1}{2} \frac{F''}{F'} \kappa r_h \partial_r R. \quad (24)$$

Now we take $T\delta S$ part of the equation (19) and substitute κ in it yielding

$$T\delta S = \kappa \frac{r_h^2}{2} \delta F' + \frac{F' \delta r_h}{2} + 4\pi r_h^2 \delta r_h T_t^t + \frac{1}{4} \mathcal{U} F' r_h^2 \delta r_h - \frac{1}{2} F'' \kappa r_h^2 (\partial_r R) \delta r_h. \quad (25)$$

Using the fact that $\delta F' = F'' \partial_r R \delta r_h$ the first member and the last member cancel each other. Leaving the equation in the form of

$$T\delta S = \frac{F' \delta r_h}{2} + 4\pi r_h^2 \delta r_h T_t^t + \frac{1}{4} \mathcal{U} F' r_h^2 \delta r_h. \quad (26)$$

This is the point where the members on the right hand side shall be interpreted as thermodynamic quantities. The Misner-Sharp energy for the horizon can be found in the first order as $E = \frac{r_h F'}{2}$ [18]. Thus, the first term on the right hand side can be interpreted as variation of energy. Comparing it to the GR result, it is visible that it differs only in the factor F' which goes to one in the GR limit. The second member on the right hand side contains a variation of volume and it can be easily interpreted as $p_M \delta V$ member, where $T_t^t = p_M$, which is exactly the same as in the GR case. The only additional member to the GR case is the last one on the right hand side. There are two possible ways of its interpretation. The first one is to interpret it as a pressure using

$$p_\Lambda \delta V = \frac{1}{4} \mathcal{U} F' r_h^2 \delta r_h = \frac{1}{16\pi} \mathcal{U} F' \delta V, \quad (27)$$

with a new pressure

$$p_\Lambda = \frac{F - F'R}{16\pi}, \quad (28)$$

if the function $F(R)$ is taken as $F(R) = R + \Lambda$, this pressure would take form of $p_\Lambda = \frac{\Lambda}{16\pi}$. This allows us to see it as a kind of generalization of dark energy pressure.

The second interpretation was done in [19] where the spherical symmetric metric in $F(R)$ gravity was also analysed using a different approach. They interpreted the last member as an additional internal entropy

$$T\delta S^i = \frac{1}{4}\mathcal{U}F'r_h^2\delta r_h = T\frac{A}{4}\frac{F'R - F}{a'}\delta r_h. \quad (29)$$

Having analysed spherically symmetric case, we have found horizon temperature and entropy. The temperature was the same as in the GR case and the entropy differs only in factor of F' . Then the thermodynamic equation was looked for yielding equation with one new extra member compared to the GR case. Two possible interpretations were formulated for this equation

$$T\delta S = \delta E + p_M\delta V + p_\Lambda\delta V, \quad T\delta S = \delta E + p_M\delta V - T\delta S^i, \quad (30)$$

where in the first one, the extra member was interpreted as a pressure arising from the function F . The second possible interpretation is as a new entropy which depends not only on horizon surface but also on the function $F(R)$ and the surface gravity.

4 General static spacetime

The second examined case is the general static spacetime. As in the previous case, the metric is static, however there is general two dimensional metric on horizon. Explicitly, we have

$$ds^2 = -N^2 dt^2 + dn^2 + \sigma_{AB} dx^A dx^B, \quad (31)$$

where major Latin indices A, B, \dots denote the indices of the horizon and take value 2 or 3. The coordinate n corresponds to the normal direction to the horizon. The horizon is expected at the point $n = 0$. Near the horizon, we expect the following expansions in n [20]

$$N = \kappa n + \mathcal{O}(n^3), \quad \sigma_{AB} = \bar{\sigma}_{AB} + \frac{1}{2}\hat{\sigma}_{AB}n^2 + \mathcal{O}(n^3), \quad (32)$$

where κ is the surface gravity of the horizon. To formulate the variation of surface Hamiltonian, the first step is to use the metric to find vector q^n

$$q^n = 2\dot{N}\sqrt{\sigma} + N\sqrt{\sigma}\sigma^{AB}\dot{\sigma}_{AB}, \quad (33)$$

where dot indicates derivative with respect to the normal coordinate n . Using the limit on horizon simplifies the form of the vector q^n into

$$q^n|_{n \rightarrow 0} = 2\kappa\sqrt{\bar{\sigma}}, \quad (34)$$

where $\bar{\sigma}$ is the limit of σ on horizon. The surface Hamiltonian's variation follows from (11) in the form of

$$\delta H_{sur} = \frac{1}{16\pi} \int d^2x_{\parallel} F' \left(2\sqrt{\bar{\sigma}}\delta\kappa + 2\kappa\delta\sqrt{\bar{\sigma}} \right) + \frac{1}{16\pi} \int d^2x_{\parallel} 2\kappa\sqrt{\bar{\sigma}}\delta F'. \quad (35)$$

Having performed integration using formula for surface area $A = \int \sqrt{\sigma} d^2x_{\parallel}$ yields

$$\delta H_{sur} = \frac{\kappa}{2\pi} \delta \left(\frac{F'A}{4} \right) + \frac{F'A}{4} \delta \left(\frac{\kappa}{2\pi} \right). \quad (36)$$

When interpreting the variation as $\delta H = T\delta S + S\delta T$, the result is identical to the one derived in the spherical case. Again as in the spherical symmetric case, the temperature remains same as in GR while the entropy takes a factor of F' .

To formulate the thermodynamic equation on the horizon, Einstein equations have to be examined. Focusing on coordinates n and t , it is easy to find members of Einstein tensor. Non-zero are only the diagonal terms G_t^t and G_r^r . Albeit they differ in general, when doing limit on the horizon both take the same form [20]

$$G_t^t|_H = \frac{1}{2}\sigma_2 - \frac{1}{2}\bar{R}_\parallel = G_n^n|_H, \quad (37)$$

where $\sigma_2 = \sigma^{AB}\hat{\sigma}_{AB}$ and R_\parallel is a Ricci scalar calculated using the two-dimensional metric σ . Since the method of its calculation preserves behaviour with respect to coordinate n , it is reasonable to expect that near the horizon R_\parallel behaves as $R_\parallel = \bar{R}_\parallel + \frac{1}{2}\hat{R}_\parallel n^2$.

To obtain the right hand side of Einstein equations, tensor \mathcal{V} has to be calculated. The calculation is more complicated than in the spherical case since the scalar curvature now depends on normal coordinate as well as both horizontal coordinates. Important is also the form of Ricci scalar curvature

$$R = -2\frac{\ddot{N}}{N} - \frac{\dot{N}}{N}\sigma^{AB}\dot{\sigma}_{AB} - \sigma^{AB}\ddot{\sigma}_{AB} + \frac{3}{4}\sigma^{AC}\sigma^{BD}\dot{\sigma}_{AB}\dot{\sigma}_{CD} - \frac{1}{4}\sigma^{AC}\sigma^{BD}\dot{\sigma}_{AC}\dot{\sigma}_{BD} + R_\parallel, \quad (38)$$

with the following expansion in n

$$R \approx -2\sigma_2 + \bar{R}_\parallel - \frac{1}{4}\sigma_2^2 n^2 + \frac{3}{4}\sigma^{AB}\sigma^{CD}\hat{\sigma}_{AD}\hat{\sigma}_{BC}n^2 + \frac{1}{2}\hat{R}_\parallel n^2. \quad (39)$$

This expansion provides us with important result, that $\partial_n R$ is vanishing on the horizon. Then the general form of both diagonal terms can be calculated. Similarly to the Einstein tensor, the both members are different, however they limit to the same value on the horizon

$$\mathcal{V}_t^t|_H = \mathcal{V}_n^n|_H = -D^A [F'' D_A (\bar{R}_\parallel - 2\sigma_2)] + F'' \left(\frac{1}{2}\sigma_2^2 - \frac{3}{2}\sigma_3 - \hat{R}_\parallel \right), \quad (40)$$

where D denotes covariant derivative with respect to the two-dimensional metric σ and $\sigma_3 = \sigma^{AB}\sigma^{CD}\hat{\sigma}_{AD}\hat{\sigma}_{BC}$. On the horizon, the Einstein equation for nn and tt components are

$$\frac{1}{2}\sigma_2 - \frac{1}{2}\bar{R}_\parallel = \frac{8\pi}{F'} T_t^t + \frac{1}{2}\mathcal{U} - \frac{1}{F'} D^A [F'' D_A (\bar{R}_\parallel - 2\sigma_2)] + \frac{F''}{F'} \left(\frac{1}{2}\sigma_2^2 - \frac{3}{2}\sigma_3 - \hat{R}_\parallel \right). \quad (41)$$

To be able to analyse $T\delta S$, variation of $\sqrt{\sigma}$ is needed. To find it, the method from [14] is used. A new parameter Λ is introduced. It is the affine parameter corresponding to the tangent vectors of the outgoing null geodesics. Near the horizon, we find that it behaves as

$$\Lambda = \bar{\Lambda} + \frac{1}{2}\kappa n^2 + \mathcal{O}(n^3), \quad (42)$$

where $\bar{\Lambda}$ is value of Λ on the horizon and it is constant with respect to n . If we take the $T\delta S$ part from (35) we are able to write it as

$$T\delta S = \frac{1}{16\pi} \int d^2 x_\parallel 2\kappa \left(\sqrt{\sigma} \delta F' + F' \delta \sqrt{\sigma} \right). \quad (43)$$

It is clearly visible that our intention is to replace both variations in this formula with variation with respect to the affine parameter Λ . The relationships between variations follow as

$$\delta \sqrt{\sigma} = \frac{1}{2\kappa} \sqrt{\sigma} \sigma_2 \delta \Lambda, \quad \delta F' = \frac{\partial F'}{\partial R} \frac{\partial R}{\partial n} \frac{\partial n}{\partial \Lambda} \delta \Lambda = \frac{F''}{\kappa} \left(\frac{3}{2}\sigma_3 + \hat{R}_\parallel - \frac{1}{2}\sigma_2^2 \right) \delta \Lambda. \quad (44)$$

Using these formulae in (43) together with σ_2 expressed from the left hand side of Einstein equation (41) results in an integral

$$T\delta S = \frac{1}{16\pi} \int d^2x_{\parallel} \sqrt{\bar{\sigma}} \{F'R_{\parallel} + 16\pi T_t^t + F'\mathcal{U} - 2D^A [F''D_A(\bar{R}_{\parallel} - 2\sigma_2)]\} \delta\Lambda. \quad (45)$$

Stokes theorem [21] can be applied on the last term and since the horizon is a closed surface, this term will vanish. Analogically to the spherical symmetric case, the term originated from tensor \mathcal{V}_k^i vanishes. The first term can be interpreted as variation of energy, analogically to the spherical case, it consists of the classical energy term multiplied by a factor of F' . The second member of the right hand side is a classical pressure term. The third term offers two interpretations as in the previous case. The first and straightforward interpretation is a generalized dark energy pressure

$$T\delta S = \delta E + p\delta V + p_{\lambda}\delta V, \quad p_{\lambda} = \frac{F - F'R}{16\pi}, \quad (46)$$

where the definition of the dark energy pressure is identical to the one in the spherical symmetric case. With a little care we can take the second path and interpret the last term as an internal entropy term

$$T\delta S = \delta E + p\delta V - T\delta S^i, \quad \delta S^i = \frac{A}{4} \frac{F'R - F}{2\dot{N}} \delta\Lambda. \quad (47)$$

Again, it is visible that the definition of the internal entropy reminds the one from the spherical case.

5 Conclusions

Spherically symmetric metric and general static spacetime metric were analysed. For both metrics, variation of surface Hamiltonian was found. In both cases, it was interpreted as $T\delta S + S\delta T$. Temperature definition is in both cases identical to the GR temperature while the entropy takes an extra factor of F' .

Einstein equations were used to find Clausius equation on the horizon. The part of the Einstein equations specific for $F(R)$ gravity was split into a scalar \mathcal{U} defined only by the function F and a tensor \mathcal{V}_k^i which mixes function F with metric. In both cases, the terms coming from \mathcal{V}_k^i vanish on the horizon. The only additional term comes from the \mathcal{U} part. In both cases, there are two possible interpretations of such term. The first one is to see it as an extra pressure, a generalized dark energy pressure. The second one is to see that term as an additional entropy.

Vanishing of the term from \mathcal{V}_k^i is an interesting behaviour but having examined only two example metrics does not provide enough insight to determine whether it is a general behaviour or a coincidence, since there might exist cases when this member does not vanish but creates a new addition to the Clausius equation instead.

Both of the presented physical interpretations of the new member of Clausius equation shall be also examined further to see if there is a method to find which is more suitable.

References

- [1] T. Jacobson, *Thermodynamics of space-time: The Einstein equation of state*, Phys. Rev. Lett. **75** (1995), 1260-1263 doi:10.1103/PhysRevLett.75.1260 [arXiv:gr-qc/9504004 [gr-qc]].

- [2] G. Chirco and S. Liberati, *Non-equilibrium Thermodynamics of Spacetime: The Role of Gravitational Dissipation*, Phys. Rev. D **81** (2010), 024016 doi:10.1103/PhysRevD.81.024016 [arXiv:0909.4194 [gr-qc]].
- [3] C. Eling, R. Guedens and T. Jacobson, *Non-equilibrium thermodynamics of space-time*, Phys. Rev. Lett. **96** (2006), 121301 doi:10.1103/PhysRevLett.96.121301 [arXiv:gr-qc/0602001 [gr-qc]].
- [4] T. Padmanabhan, *Classical and quantum thermodynamics of horizons in spherically symmetric space-times*, Class. Quant. Grav. **19** (2002), 5387-5408 doi:10.1088/0264-9381/19/21/306 [arXiv:gr-qc/0204019 [gr-qc]].
- [5] T. Padmanabhan, *Gravity from space-time thermodynamics*, Astrophys. Space Sci. **285** (2003), 407 doi:10.1023/A:1025448712533 [arXiv:gr-qc/0209088 [gr-qc]].
- [6] T. Padmanabhan, *A Physical Interpretation of Gravitational Field Equations*, AIP Conf. Proc. **1241** (2010) no.1, 93-108 doi:10.1063/1.3462738 [arXiv:0911.1403 [gr-qc]].
- [7] A. Mukhopadhyay and T. Padmanabhan, *Holography of gravitational action functionals*, Phys. Rev. D **74** (2006), 124023 doi:10.1103/PhysRevD.74.124023 [arXiv:hep-th/0608120 [hep-th]].
- [8] S. Kolekar and T. Padmanabhan, *Holography in Action*, Phys. Rev. D **82** (2010), 024036 doi:10.1103/PhysRevD.82.024036 [arXiv:1005.0619 [gr-qc]].
- [9] T. Padmanabhan, *Holographic gravity and the surface term in the Einstein-Hilbert action*, Braz. J. Phys. **35** (2005), 362-372 doi:10.1590/S0103-97332005000200023 [arXiv:gr-qc/0412068 [gr-qc]].
- [10] Th. De Donder, *Théorie Invariantive Du Calcul des Variations*, (Gauthier-Villars and Cie.,Paris, 1930).
- [11] H. Weyl, *Geodesic Fields in the Calculus of Variation for Multiple Integrals*, Annals of Mathematics, **36**, p.607.
- [12] J. Struckmeier and A. Redelbach, *Covariant Hamiltonian field theory*, Int. J. Mod. Phys. **E17** (2008), 435-491 doi:10.1142/S0218301308009458 [arXiv:0811.0508 [math-ph]].
- [13] P. Hořava, *On a covariant Hamilton-Jacobi framework for the Einstein-Maxwell theory*, Class. Quant. Grav. **8** (1991), 2069-2084 doi:10.1088/0264-9381/8/11/01.
- [14] K. Parattu, B. R. Majhi and T. Padmanabhan, *Structure of the gravitational action and its relation with horizon thermodynamics and emergent gravity paradigm*, Phys. Rev. D **87** (2013) no.12, 124011 doi:10.1103/PhysRevD.87.124011 [arXiv:1303.1535 [gr-qc]].
- [15] A. Eddington, *The Mathematical Theory of Relativity*. Cambridge University Press, Cambridge, UK, 2 ed., 1924.
- [16] J. Klusoň and B. Matouš, *Covariant Hamiltonian Formalism for $F(R)$ -Gravity*, Gen. Rel. Grav. **53** (2021) no.11, 100 doi:10.1007/s10714-021-02868-2 [arXiv:2008.00659 [gr-qc]].
- [17] T. Padmanabhan, *Structural Aspects Of Gravitational Dynamics And The Emergent Perspective Of Gravity*, AIP Conf. Proc. **1483** (2012) no.1, 212-238 doi:10.1063/1.4756971 [arXiv:1208.1375 [hep-th]].
- [18] R. G. Cai, L. M. Cao, Y. P. Hu and N. Ohta, *Generalized Misner-Sharp Energy in $f(R)$ Gravity*, Phys. Rev. D **80** (2009), 104016 doi:10.1103/PhysRevD.80.104016 [arXiv:0910.2387 [hep-th]].
- [19] M. Akbar and R. G. Cai, *Thermodynamic Behavior of Field Equations for $f(R)$ Gravity*, Phys. Lett. B **648** (2007), 243-248 doi:10.1016/j.physletb.2007.03.005 [arXiv:gr-qc/0612089 [gr-qc]].

- [20] A. J. M. Medved, D. Martin and M. Visser, *Dirty black holes: Space-time geometry and near horizon symmetries*, *Class. Quant. Grav.* **21** (2004), 3111-3126 doi:10.1088/0264-9381/21/13/003 [arXiv:gr-qc/0402069 [gr-qc]].
- [21] E. Poisson, *A Relativist's Toolkit: The Mathematics of Black-Hole Mechanics*, doi:10.1017/CBO9780511606601
- [22] K. Bamba, C. Q. Geng, S. Nojiri and S. D. Odintsov, *Equivalence of modified gravity equation to the Clausius relation*, *EPL* **89** (2010) no.5, 50003 doi:10.1209/0295-5075/89/50003 [arXiv:0909.4397 [hep-th]].