

Thermal Dark Matter from Freezeout of Inverse Decays

Ronny Frumkin^{1,*}, Yonit Hochberg^{1,†}, Eric Kuflik^{1,‡} and Hitoshi Murayama^{2,3,4,5§}

¹*Racah Institute of Physics, Hebrew University of Jerusalem, Jerusalem 91904, Israel*

²*Ernest Orlando Lawrence Berkeley National Laboratory,
University of California, Berkeley, CA 94720, USA*

³*Department of Physics, University of California, Berkeley, CA 94720, USA*

⁴*Kavli Institute for the Physics and Mathematics of the Universe (WPI),
University of Tokyo, Kashiwa 277-8583, Japan and*

⁵*The Institute for AI and Beyond, The University of Tokyo, Tokyo 113-8655, Japan*

We propose a new thermal dark matter candidate whose abundance is determined by the freezeout of inverse decays. The relic abundance depends parametrically only on a decay width, while matching the observed value requires that the coupling determining the width—and the width itself—should be exponentially small. The dark matter is therefore very weakly coupled to the Standard Model, evading conventional searches. This INverse DecaY (‘INDY’) dark matter can be discovered by searching for the long-lived particle that decays into the dark matter at future planned experiments.

INTRODUCTION

The identity of dark matter (DM) is one of the most pressing open questions in modern day physics. While the weakly interacting massive particle (WIMP) paradigm has long guided the particle physics community, the absence of experimental evidence for the WIMP at colliders, direct-detection and indirect-detection experiments stresses the importance of considering DM beyond the WIMP. Indeed, recent years have seen a surge of new DM ideas (see *e.g.* Refs. [1–15]) which utilize various processes in the early universe.

One such process is inverse decay, where a DM particle is produced through the inverse decay of a heavier particle in the dark sector. Thus far, decays have been considered in the literature in the context of freeze-in DM [16], where a slow inverse decay of a bath particle slowly freezes in the DM abundance; as a process maintaining chemical equilibrium within the dark sector or with the SM [7, 8, 13, 17–22]; and in other dark matter frameworks [16–18, 23–26]. The effects of inverse decays on dark matter depletion have been considered as a contributing reaction [27, 28], but never as the main process for setting the dark matter abundance. In this work, and in a companion paper [29], we study the freezeout of inverse decays as the mechanism to set the relic abundance of DM.

This *Letter* is organized as follows. We begin by outlining the basic idea for freezeout of inverse decays and derive an analytical understanding of the mechanism. We then solve the Boltzmann equations of the system and obtain the DM parameter space. Finally we present a model that realizes the mechanism.

BASIC IDEA

Here we show that the freezeout of inverse decays can be responsible for the relic abundance of DM. Consider

a dark matter particle χ and an unstable dark sector particle ψ that has a decay that contains some number of χ particles in the final state. For simplicity we will consider a simple decay and inverse decay

$$\psi \longleftrightarrow \chi + \phi \quad (1)$$

motivated by a Z_2 symmetry in the dark sector. (Other inverse decay topologies can be considered as well.) Here ϕ can be a dark sector or visible particle that is assumed to be in equilibrium with the bath. (Later we will take a concrete model in which ϕ is a dark photon that kinetically mixes with hypercharge.)

The Boltzmann equation for the abundance of χ , assuming that ϕ is in equilibrium, is:

$$\dot{n}_\chi + 3Hn_\chi = \Gamma \left(n_\psi - n_\chi \frac{n_\psi^{\text{eq}}}{n_\chi^{\text{eq}}} \right), \quad (2)$$

with Γ the decay rate of $\psi \rightarrow \chi\phi$. We assume that the DM is cold, namely that it freezes out when non-relativistic; our numerics presented later on confirm this. We can thus ignore the thermally averaged time dilation that would normally appear in the collision term.

Approximate analytic solutions to the Boltzmann equations can be obtained in the instantaneous freezeout approximation, but will not always suffice. The inverse decay rate is falling off exponentially as $e^{-(m_\psi - m_\chi)/T}$ (rather than $e^{-m_\chi/T}$ for the well-studied WIMP), which is not necessarily fast enough to assume that instantaneous freezeout occurs. Further consideration must also be taken into account because the decays and inverse decays may not actually be in equilibrium before they completely decouple.

We begin by calculating the relic abundance when decays are in equilibrium. This will give us an approximate range of parameter space—couplings and masses—necessary to reproduce the observed abundance. We first assume that ψ is always in chemical equilibrium with the Standard Model (SM) bath. This can be achieved

through rapid annihilations of ψ into bath particles (either SM bath particles, or particles that are in equilibrium with the SM). The Boltzmann Eq. (2) can be rewritten as

$$\frac{\partial Y_\chi}{\partial x} = -\frac{\Gamma}{Hx} \left(Y_\chi \frac{n_\psi^{\text{eq}}}{n_\chi^{\text{eq}}} - Y_\psi^{\text{eq}} \right), \quad (3)$$

where $Y = n/s$ is the yield and $x = m_\chi/T$. In the instantaneous freezeout approximation, the χ particle departs the chemical equilibrium when the coefficient of the collision term becomes of order unity,

$$\Gamma \frac{n_\psi^{\text{eq}}}{n_\chi^{\text{eq}}} \simeq xH \quad \text{for } x = x_f. \quad (4)$$

The freezeout abundance of χ can be determined by solving Eq. (4) for x_f and using that at the time of freezeout, χ is approximately in equilibrium:

$$n_\chi(x_f) \simeq n_\chi^{\text{eq}}(x_f) = g_\chi \left(\frac{m^2}{2\pi x_f} \right)^{3/2} e^{-x_f}. \quad (5)$$

We parameterize the decay rate as

$$\Gamma \equiv \alpha_{\text{decay}} m_\psi \sqrt{1 - 2 \frac{(m_\phi^2 + m_\chi^2)}{m_\psi^2} + \frac{(m_\phi^2 - m_\chi^2)^2}{m_\psi^4}}. \quad (6)$$

Then requiring that we match the observed DM abundance, given by $n_\chi(x_f) \sim m^2 T_{\text{eq}}$, we arrive at the requisite relationship between the DM mass and coupling,

$$m_\chi \simeq \alpha_{\text{decay}}^{\frac{1}{1+\Delta}} (m_{\text{pl}} T_{\text{eq}}^\Delta)^{\frac{1}{1+\Delta}}. \quad (7)$$

Here $\Delta \equiv (m_\psi - m_\chi)/m_\chi$, $T_{\text{eq}} = 0.8$ eV is the temperature at matter radiation equality, and $m_{\text{pl}} = 2.4 \times 10^{18}$ GeV is the reduced Planck mass. This formula may remind the reader of other dark matter freezeout mechanisms, and would produce similar parametric dependence on the temperature of equality and Planck scale in the WIMP and SIMP [1] cases for $\Delta = 1$ and $\Delta = 2$, respectively. We therefore see that for small mass splittings $\Delta < 1$, much heavier mass DM for the same size coupling (or much smaller coupling for same mass) is needed to produce the relic abundance, when compared to the WIMP. For this reason, inverse decays can achieve super heavy dark matter with masses beyond the WIMP unitarity bound, in the spirit of Ref. [13], where an inverse chain of dark sector particles set the abundance [30].

The above also indicates when this analysis is valid. Provided that the rate over Hubble becomes of order unity before x_f , the system will come close to equilibrium and there will be no sensitivity to the initial conditions.

Looking at Eq. (4), the rate over Hubble is not monotonically decreasing, but rather reaches a maximum at $x = \Delta^{-1}$. Equilibrium is achieved if

$$\Gamma \gtrsim \Delta H_m, \quad (8)$$

where $H_m \equiv H(x = 1)$. If Eq. (8) is not satisfied, then the decays and inverse decays are not sufficient to thermalize the dark matter to chemical equilibrium before freezeout. In this case the initial χ abundance may be much higher or much lower than the equilibrium abundance when it becomes non-relativistic. However, we will see in the next section that the final abundance can still remain very insensitive to initial conditions, where the observed relic abundance is obtained for almost the same parameters.

PHASES OF INVERSE DECAY

We can now setup the system of interest in full. We consider the decay and inverse decay processes of $\psi \leftrightarrow \chi\phi$ along with an annihilation process of $\psi\psi \rightarrow \tilde{\phi}\tilde{\phi}$, with ϕ and $\tilde{\phi}$ indicating a particle in equilibrium with the SM bath—the latter process maintaining chemical equilibrium between ψ and the SM. The Boltzmann equations for the system are given by:

$$\dot{n}_\chi + 3Hn_\chi = -\langle\Gamma\rangle \left(\frac{n_\psi^{\text{eq}}}{n_\chi^{\text{eq}}} n_\chi - n_\psi \right), \quad (9)$$

$$\dot{n}_\psi + 3Hn_\psi = \langle\Gamma\rangle \left(\frac{n_\psi^{\text{eq}}}{n_\chi^{\text{eq}}} n_\chi - n_\psi \right) - \langle\sigma v\rangle (n_\psi^2 - n_\psi^{\text{eq}^2}).$$

Here $\langle\Gamma\rangle$ represents the thermally averaged (time dilation included) decay rate of $\psi \rightarrow \chi\phi$, and $\langle\sigma v\rangle$ represents the thermally averaged cross section for the annihilation $\psi\psi \rightarrow \tilde{\phi}\tilde{\phi}$. We parameterize the cross section as

$$\langle\sigma v\rangle \equiv \frac{\alpha_{\text{ann}}^2}{m_\psi^2}, \quad (10)$$

and the decay rate as in Eq (6).

Fig. 1 shows our solutions to the Boltzmann equations, α_{decay} versus α_{ann} , that reproduces the observed relic abundance of DM, for various mass splitting and DM masses. There are several distinct phases, each indicating a different mechanism for setting the relic abundance of DM, with some dependence on the initial conditions. Here we focus on two of the phases, assuming that the dark matter is in equilibrium with the bath prior to becoming non-relativistic.

Along the horizontal branch, the relic abundance of χ is being set by the annihilations of ψ into other particles. This is possible because the rapid inverse decays between χ and ψ keep the two particles in chemical equilibrium. Once α_{decay} is large enough such that χ and ψ are in




FIG. 1. **Left:** Annihilations versus decays phase diagram, for various mass splittings and DM masses (solid colored curves). The two new phases are evident: The first is the horizontal branch, where the relic abundance of the DM is set through the annihilations of another dark particle, with which it is in chemical equilibrium via decays and inverse decays. The second is the vertical branch, where the relic abundance of the DM is set by the *freezeout of the inverse decay* of the DM particle; we dub this ‘INDY’ DM. **Right:** The decay coupling versus DM mass that reproduces the observed relic abundance along the vertical branch, for various mass splittings. We plot the numerical solution to the Boltzmann equations, with the dashed region of the curves indicating where the effective coupling α_{ann} that we use to parameterize the cross section exceeds 100.

chemical equilibrium, its precise value does not matter, and it is α_{ann} that controls the abundance. This branch is similar in spirit to coannihilations [3, 11], however here the chemical contact between the DM and the particle whose number changing process is determining the relic abundance is set by fast decays and inverse decays, rather than by $2 \rightarrow 2$ processes such as annihilations.

The value of α_{ann} that reproduces the correct abundance along this branch can be found in the instantaneous freezeout approximation using the same steps as from Eq. (4) to Eq. (7), but here x_f is determined by $n_\psi \langle \sigma v \rangle \simeq H$. Doing so, one finds the relationship

$$m_\chi \simeq \alpha_{\text{ann}}^{\frac{2}{2+\Delta}} (m_{\text{pl}} T_{\text{eq}}^{1+\Delta})^{\frac{1}{2+\Delta}}. \quad (11)$$

In contrast, along the vertical branch, the relic abundance of χ is being set by the freezeout of inverse decays of the DM. As χ inverse decays into ψ , ψ rapidly annihilated away into other light particles. For large enough α_{ann} , its precise value does not play a role, and the relic abundance of χ is set when its inverse decay shuts off and it freezes out. This corresponds to the case studied analytically in the previous section, which leads to the DM mass-coupling relationship of Eq. (7).

While both branches present a new mechanism for establishing the DM abundance in the early universe, we focus here on the more novel of the two — the vertical branch, where freezeout of inverse decays of DM set its abundance. We dub this ‘INverse DecaY’ (INDY) dark matter. The parameter space has another two phases, if we assume that initial abundance of χ is very small: freeze-in, and freeze-in and freezeout (of inverse decays),

which is very similar to the vertical branch. We leave a detailed investigation of the horizontal branch, the other additional phases, and many further details to our companion paper [29], and focus here on INDY dark matter.

In Fig. 1 we show α_{decay} as a function of DM mass along the vertical branch, for various mass splittings, where INDY DM is achieved. We show the exact solutions to the Boltzmann equations, in qualitative agreement with the analytic approximations of Eq. (7). Note that the minimum value of α_{ann} needed to be in the INDY phase is given by Eq. (11). Therefore the mass of the DM is bounded above by the unitarity of the $\langle \sigma v \rangle$ cross section. Moreover, since $\Delta > 0$, the INDY mechanism predicts lighter DM than the standard WIMP case (which would correspond to $\Delta = 0$ in Eq. (11)).

Fig. 2 depicts the evolution of the densities of χ along the vertical branch of INDYs for some benchmark values of mass, splitting and couplings. The two panels illustrate that the DM density can be set by the freezeout of the inverse decay where χ either maintains chemical equilibrium with ψ until freezeout occurs (*left panel*), or χ can decouple at early times and still freezeout in a similar manner (*right panel*). In both cases, DM is indeed freezing out non-relativistically, justifying our back-of-the-envelope computation.

A comment is in order regarding initial conditions. In the above we have assumed an equilibrium distribution for the DM χ at very early times. However in some regions along the vertical branch, where the coupling α_{decay} is very small, decays and inverse decays may not be strong enough to bring χ into equilibrium in the




FIG. 2. Example solutions to the Boltzmann equations along the vertical branch, where the relic abundance of DM is set by the freezeout of the inverse decays of the DM. For both panels, we show the curves for initial conditions $Y_\chi = Y_\chi^{\text{eq}}$ (solid) and $Y_\chi = 0$ (dashed), and the corresponding couplings α_{decay} for INDY DM, freeze-in and freezeout (FIFO), and freeze-in (FI). **Left:** χ is thermalized via the inverse decays, and maintains an equilibrium distribution until freezeout of its inverse decays. **Right:** the DM χ departs from chemical equilibrium early on (due to the smaller α_{decay}), and later freezes out.

early universe. To understand the impact of the initial conditions, we assess the extent to which varying them changes the relic abundance. We find our results to be insensitive to the initial conditions over a broad range of DM masses and mass splittings, allowing many orders of magnitude change in the initial condition without significantly modifying the relic abundance. If the dark matter inverse decays thermalize before they become non-relativistic, then the final relic abundance is essentially insensitive to the initial conditions. Additionally, one can consider the case in which the inverse decays are never in equilibrium. In this case, starting with a near equilibrium abundance or even no abundance of dark matter (in which case dark matter freezes in and then freezes out) produces almost the identical parameter space to match the DM relic abundance. This is evident in the right panel of Fig. 2, where relic abundance for starting with an equilibrium value (labeled INDY) and starting with a small abundance (labeled FIFO) produces nearly identical α_{decay} values. We relegate a detailed discussion of initial conditions and the relation to other phases of the parameter space to our companion paper [29].

MODEL

As proof of concept, we now present a model that realizes INDY dark matter. Consider a $U(1)_d$ gauge theory with gauge coupling e_d , two Dirac fermions χ and ψ , and a complex scalar ϕ , with charges 0, 1, and 1 respectively. The general renormalizable Lagrangian contains masses for the fermion, a spontaneous symmetry breaking po-

tential for the scalar, and the Yukawa interaction

$$\mathcal{L}_{\text{Yukawa}} = -y\phi^*\bar{\chi}\psi + h.c. \quad (12)$$

The field ϕ acquires a VEV through spontaneous symmetry breaking, which gives a mass m_{A_d} to the dark photon and sources a mixing term between the fermions. This leads to interactions between ψ , χ and the dark photon that yield decays and inverse decays. Mapping to the parametrization of Eqs. (10) and (6), we have [31–33]:

$$\alpha_{\text{decay}} = \frac{y^2 \left((m_\chi - m_\psi)^2 - m_{A_d}^2 \right) \left((m_\chi + m_\psi)^2 + 2m_{A_d}^2 \right)}{32\pi m_\psi^2 (m_\chi - m_\psi)^2} \quad (13)$$

and

$$\alpha_{\text{ann}}^2 = \frac{e_d^4 \sqrt{1 - \frac{m_{A_d}^2}{m_\psi^2}} \left(m_\psi^2 - m_{A_d}^2 \right)}{32\pi \left(m_\psi^2 - \frac{1}{2}m_{A_d}^2 \right)^2} m_\psi^2. \quad (14)$$

Large values of y and low values of e_d correspond to large α_{decay} and represents solutions on the horizontal branch of Fig. 1, while small values of y and large values of e_d correspond to large values of α_{ann} and represents solutions along the vertical branch of INDY dark matter.

For dark matter masses at the MeV scale and above, other processes such as co-scattering [10] with the dark photon can make small corrections to the curves in Fig. 1. On the horizontal branch, co-scattering plays no role as it does not change the number of $n_\chi + n_\psi$ particles. INDY dark matter requires large values of α_{ann} , which correspond to large values of dark coupling $e_d \sim \mathcal{O}(10^{-2} - 1)$, resulting in the dark photon mass being similar to the




FIG. 3. Allowed dark photon parameter space. Solid colored curves indicate the minimal kinetic mixing as a function of dark photon mass within the model, for various values of mass splitting Δ and with $m_{A_d} = 0.9\Delta m_\chi$. We show existing limits [34, 43, 45–52] in shaded gray, with future projections [43, 52–62] indicated by the colored dashed curves.

mass splitting between χ and ψ . Since the ratio between the rates of the co-scattering process and the inverse decay is proportional to the dark photon number density, the rate of the co-scattering processes decreases rapidly when $x \sim \Delta^{-1} \sim m_\chi/m_{A_d}$, leaving the inverse decay as the dominant process. Co-scattering with the scalar and co-annihilation processes are small for similar reasons, and the annihilation process of $\chi\chi \rightarrow A_d A_d$ is suppressed by y^4 . The model can thus accommodate the INDY mechanism. As a benchmark, for $m_\chi = 200$ MeV, $m_\psi = 230$ MeV, $m_{A_d} = 27$ MeV, $m_\phi = 60$ MeV, $e_d = 0.32$ and $y = 7.7 \times 10^{-9}$ and $\epsilon = 10^{-4}$ we obtain the observed relic abundance of DM particles along the vertical INDY branch, evading current constraints from BBN and cosmology [34–44]. Equilibrium with the bath is maintained via dark photon decay and inverse decays, $A_d \leftrightarrow e^+e^-$.

PHENOMENOLOGY

At the mechanism level, χ does not couple directly to the SM. In the model above, all interactions of the χ particle are suppressed by $y \sim 10^{-8}$, and the couplings to the SM are further suppressed by $\epsilon \ll 1$. As a result, direct detection of DM in the laboratory and indirect detection of DM annihilations in the galaxy will be highly suppressed, and thus INDY DM can be expected to evade these conventional searches. For other examples of exponentially small couplings to the SM from coannihilations, see Ref. [63].

The dark photon can be searched for directly. Fig. 3

shows the relevant parameter space for a visible decaying dark photon. Existing limits are indicated in shaded gray [34, 43, 45–52]. Solid curves indicate the lower limit on the kinetic mixing parameter ϵ for given mass splitting Δ , where we have fixed $m_{A_d} = 0.9\Delta m_\chi$. Below the solid curves, the SM and dark sector are not thermalized, while to the right of the curve, one does not reproduce the DM relic abundance since the annihilation rate is too small. As is evident, the allowed parameter space is expected to be probed by future experiments, indicated by the dashed curves [43, 52–62] (bounds and projections are taken from the recent compilation in Ref. [40]).

Discovery of the INDY dark matter candidate can hopefully be made via the direct production of the ψ particle, which decays into χ and either additional invisible products or SM products. The ψ particle is typically long lived; for sub-GeV INDY DM, the lifetime can vary from $\mathcal{O}(\text{mm})$ to $\mathcal{O}(10^4 \text{ km})$. In the model presented above, we have the decay chain $\psi \rightarrow \chi A_d$, $A_d \rightarrow e^+e^-$. This decay chain can be searched for in beam dump experiments where off-shell dark photons can be produced. A similar scenario was recently considered in Ref. [52], which showed that such decay chains can be probed in current and future beam dump experiments such as NA62 [64–68], and SeaQuest/DarkQuest/LongQuest [59, 69, 70].

Acknowledgments. The work of YH is supported by the Israel Science Foundation (grant No. 1112/17), by the Binational Science Foundation (grant No. 2018140), by the I-CORE Program of the Planning Budgeting Committee (grant No. 1937/12), and by the Azrieli Foundation. EK is supported by the Israel Science Foundation (grant No. 1111/17), by the Binational Science Foundation (grants No. 2016153 and 2020220) and by the I-CORE Program of the Planning Budgeting Committee (grant No. 1937/12). The work of HM was supported by the Director, Office of Science, Office of High Energy Physics of the U.S. Department of Energy under the Contract No. DE-AC02-05CH11231, by the NSF grant PHY-1915314, by the Binational Science Foundation (grant No. 2018140), by the JSPS Grant-in-Aid for Scientific Research JP20K03942, MEXT Grant-in-Aid for Transformative Research Areas (A) JP20H05850, JP20A203, by WPI, MEXT, Japan, the Institute for AI and Beyond of the University of Tokyo, and Hamamatsu Photonics, K.K.

* ronny.frumkin@mail.huji.ac.il

† yonit.hochberg@mail.huji.ac.il

‡ eric.kuflik@mail.huji.ac.il

§ hitoshi@berkeley.edu, hitoshi.murayama@ipmu.jp

- [1] Y. Hochberg, E. Kuflik, T. Volansky, and J. G. Wacker, Phys. Rev. Lett. **113**, 171301 (2014), arXiv:1402.5143 [hep-ph].
- [2] Y. Hochberg, E. Kuflik, H. Murayama, T. Volansky,

- and J. G. Wacker, Phys. Rev. Lett. **115**, 021301 (2015), arXiv:1411.3727 [hep-ph].
- [3] K. Griest and D. Seckel, Phys. Rev. **D43**, 3191 (1991).
- [4] R. T. D’Agnolo and J. T. Ruderman, Phys. Rev. Lett. **115**, 061301 (2015), arXiv:1505.07107 [hep-ph].
- [5] E. Kuflik, M. Perelstein, N. R.-L. Lorier, and Y.-D. Tsai, Phys. Rev. Lett. **116**, 221302 (2016), arXiv:1512.04545 [hep-ph].
- [6] E. Kuflik, M. Perelstein, N. R.-L. Lorier, and Y.-D. Tsai, JHEP **08**, 078 (2017), arXiv:1706.05381 [hep-ph].
- [7] J. A. Dror, E. Kuflik, and W. H. Ng, Phys. Rev. Lett. **117**, 211801 (2016), arXiv:1607.03110 [hep-ph].
- [8] J. A. Dror, E. Kuflik, B. Melcher, and S. Watson, Phys. Rev. **D97**, 063524 (2018), arXiv:1711.04773 [hep-ph].
- [9] J. Kopp, J. Liu, T. R. Slatyer, X.-P. Wang, and W. Xue, JHEP **12**, 033 (2016), arXiv:1609.02147 [hep-ph].
- [10] R. T. D’Agnolo, D. Pappadopulo, and J. T. Ruderman, Phys. Rev. Lett. **119**, 061102 (2017), arXiv:1705.08450 [hep-ph].
- [11] R. T. D’Agnolo, C. Mondino, J. T. Ruderman, and P.-J. Wang, JHEP **08**, 079 (2018), arXiv:1803.02901 [hep-ph].
- [12] P. J. Fitzpatrick, H. Liu, T. R. Slatyer, and Y.-D. Tsai, (2020), arXiv:2011.01240 [hep-ph].
- [13] H. Kim and E. Kuflik, Phys. Rev. Lett. **123**, 191801 (2019), arXiv:1906.00981 [hep-ph].
- [14] P. Asadi, E. D. Kramer, E. Kuflik, G. W. Ridgway, T. R. Slatyer, and J. Smirnov, (2021), arXiv:2103.09822 [hep-ph].
- [15] P. Asadi, E. D. Kramer, E. Kuflik, G. W. Ridgway, T. R. Slatyer, and J. Smirnov, (2021), arXiv:2103.09827 [hep-ph].
- [16] L. J. Hall, K. Jedamzik, J. March-Russell, and S. M. West, JHEP **03**, 080 (2010), arXiv:0911.1120 [hep-ph].
- [17] A. Berlin, D. Hooper, and G. Krnjaic, Phys. Lett. B **760**, 106 (2016), arXiv:1602.08490 [hep-ph].
- [18] D. E. Morrissey, D. Poland, and K. M. Zurek, JHEP **07**, 050 (2009), arXiv:0904.2567 [hep-ph].
- [19] T. Cohen, D. J. Phalen, A. Pierce, and K. M. Zurek, Phys. Rev. D **82**, 056001 (2010), arXiv:1005.1655 [hep-ph].
- [20] P. Bandyopadhyay, E. J. Chun, and J.-C. Park, JHEP **06**, 129 (2011), arXiv:1105.1652 [hep-ph].
- [21] M. Farina, D. Pappadopulo, J. T. Ruderman, and G. Trevisan, JHEP **12**, 039 (2016), arXiv:1607.03108 [hep-ph].
- [22] Y. Hochberg, E. Kuflik, and H. Murayama, Phys. Rev. **D99**, 015005 (2019), arXiv:1805.09345 [hep-ph].
- [23] J. L. Feng, A. Rajaraman, and F. Takayama, Phys. Rev. Lett. **91**, 011302 (2003), arXiv:hep-ph/0302215.
- [24] M. Kaplinghat, Phys. Rev. D **72**, 063510 (2005), arXiv:astro-ph/0507300.
- [25] T. Moroi and L. Randall, Nucl. Phys. B **570**, 455 (2000), arXiv:hep-ph/9906527.
- [26] B. S. Acharya, G. Kane, S. Watson, and P. Kumar, Phys. Rev. D **80**, 083529 (2009), arXiv:0908.2430 [astro-ph.CO].
- [27] M. Garny, J. Heisig, B. Lülzf, and S. Vogl, Phys. Rev. D **96**, 103521 (2017), arXiv:1705.09292 [hep-ph].
- [28] C. E. Yaguna, Phys. Lett. B **669**, 139 (2008), arXiv:0811.0485 [hep-ph].
- [29] R. Frumkin, Y. Hochberg, E. Kuflik, and H. Murayama, work in progress.
- [30] Y. Hochberg, E. Kuflik, and T. Silverwater, work in progress.
- [31] V. Shtabovenko, R. Mertig, and F. Orellana, Comput. Phys. Commun. **256**, 107478 (2020), arXiv:2001.04407 [hep-ph].
- [32] V. Shtabovenko, R. Mertig, and F. Orellana, Comput. Phys. Commun. **207**, 432 (2016), arXiv:1601.01167 [hep-ph].
- [33] R. Mertig, M. Bohm, and A. Denner, Comput. Phys. Commun. **64**, 345 (1991).
- [34] R. H. Parker, C. Yu, W. Zhong, B. Estey, and H. Müller, Science **360**, 191 (2018), arXiv:1812.04130 [physics.atom-ph].
- [35] D. Banerjee *et al.*, Phys. Rev. Lett. **123**, 121801 (2019), arXiv:1906.00176 [hep-ex].
- [36] J. P. Lees *et al.* (BaBar), Phys. Rev. Lett. **119**, 131804 (2017), arXiv:1702.03327 [hep-ex].
- [37] S. Andreas, C. Niebuhr, and A. Ringwald, Phys. Rev. D **86**, 095019 (2012), arXiv:1209.6083 [hep-ph].
- [38] J. Blümlein and J. Brunner, Physics Letters B **731**, 320 (2014).
- [39] N. Collaboration, Physics Letters B **746**, 178 (2015).
- [40] Y.-D. Tsai, R. McGehee, and H. Murayama, (2020), arXiv:2008.08608 [hep-ph].
- [41] M. Ibe, S. Kobayashi, Y. Nakayama, and S. Shirai, Journal of High Energy Physics **2020** (2020), 10.1007/jhep04(2020)009.
- [42] N. Sabti, J. Alvey, M. Escudero, M. Fairbairn, and D. Blas, Journal of Cosmology and Astroparticle Physics **2020**, 004–004 (2020).
- [43] J. Alexander *et al.* (2016) arXiv:1608.08632 [hep-ph].
- [44] M. Bauer, P. Foldenauer, and J. Jaeckel, JHEP **07**, 094 (2018), arXiv:1803.05466 [hep-ph].
- [45] A. Fradette, M. Pospelov, J. Pradler, and A. Ritz, Phys. Rev. D **90**, 035022 (2014), arXiv:1407.0993 [hep-ph].
- [46] J. H. Chang, R. Essig, and S. D. McDermott, JHEP **01**, 107 (2017), arXiv:1611.03864 [hep-ph].
- [47] E. Hardy and R. Lasenby, JHEP **02**, 033 (2017), arXiv:1611.05852 [hep-ph].
- [48] M. Pospelov and Y.-D. Tsai, Phys. Lett. B **785**, 288 (2018), arXiv:1706.00424 [hep-ph].
- [49] D. Banerjee *et al.* (NA64), Phys. Rev. Lett. **120**, 231802 (2018), arXiv:1803.07748 [hep-ex].
- [50] R. Aaij *et al.* (LHCb), Phys. Rev. Lett. **120**, 061801 (2018), arXiv:1710.02867 [hep-ex].
- [51] R. Aaij *et al.* (LHCb), Phys. Rev. Lett. **124**, 041801 (2020), arXiv:1910.06926 [hep-ex].
- [52] Y.-D. Tsai, P. deNiverville, and M. X. Liu, Phys. Rev. Lett. **126**, 181801 (2021), arXiv:1908.07525 [hep-ph].
- [53] A. Celentano (HPS), J. Phys. Conf. Ser. **556**, 012064 (2014), arXiv:1505.02025 [physics.ins-det].
- [54] I. Jaegle (Belle), Phys. Rev. Lett. **114**, 211801 (2015), arXiv:1502.00084 [hep-ex].
- [55] P. Ilten, J. Thaler, M. Williams, and W. Xue, Phys. Rev. D **92**, 115017 (2015), arXiv:1509.06765 [hep-ph].
- [56] S. Alekhin *et al.*, Rept. Prog. Phys. **79**, 124201 (2016), arXiv:1504.04855 [hep-ph].
- [57] P. Ilten, Y. Soreq, J. Thaler, M. Williams, and W. Xue, Phys. Rev. Lett. **116**, 251803 (2016), arXiv:1603.08926 [hep-ph].
- [58] A. Caldwell *et al.*, (2018), arXiv:1812.11164 [physics.acc-ph].
- [59] A. Berlin, S. Gori, P. Schuster, and N. Toro, Phys. Rev. D **98**, 035011 (2018), arXiv:1804.00661 [hep-ph].
- [60] A. Berlin, N. Blinov, G. Krnjaic, P. Schuster, and N. Toro, Phys. Rev. D **99**, 075001 (2019),

- arXiv:1807.01730 [hep-ph].
- [61] A. Ariga *et al.* (FASER), *Phys. Rev. D* **99**, 095011 (2019), arXiv:1811.12522 [hep-ph].
- [62] C. NA62 (NA62 Collaboration), *2018 NA62 Status Report to the CERN SPSC*, Tech. Rep. (CERN, Geneva, 2018).
- [63] R. T. D’Agnolo, D. Pappadopulo, J. T. Ruderman, and P.-J. Wang, *Phys. Rev. Lett.* **124**, 151801 (2020), arXiv:1906.09269 [hep-ph].
- [64] E. Cortina Gil *et al.* (NA62), *JINST* **12**, P05025 (2017), arXiv:1703.08501 [physics.ins-det].
- [65] B. Döbrich (NA62), *Frascati Phys. Ser.* **66**, 312 (2018), arXiv:1807.10170 [hep-ex].
- [66] B. Döbrich, J. Jaeckel, and T. Spadaro, *JHEP* **05**, 213 (2019), [Erratum: *JHEP* **10**, 046 (2020)], arXiv:1904.02091 [hep-ph].
- [67] C.-W. Chiang and P.-Y. Tseng, *Phys. Lett. B* **767**, 289 (2017), arXiv:1612.06985 [hep-ph].
- [68] E. Cortina Gil *et al.* (NA62), *JHEP* **05**, 182 (2019), arXiv:1903.08767 [hep-ex].
- [69] S. Gardner, R. J. Holt, and A. S. Tadepalli, *Phys. Rev. D* **93**, 115015 (2016), arXiv:1509.00050 [hep-ph].
- [70] M. X. Liu, *Mod. Phys. Lett. A* **32**, 1730008 (2017).