

CLEAR: Boosted Ly α Transmission of the Intergalactic Medium in UV bright Galaxies

INTAE JUNG,^{1,2,3} CASEY PAPOVICH,^{4,5} STEVEN L. FINKELSTEIN,⁶ RAYMOND C. SIMONS,⁷ VICENTE ESTRADA-CARPENTER,^{4,5}
BREN E. BACKHAUS,⁸ NIKKO J. CLERI,^{4,5} KRISTIAN FINLATOR,^{9,10} MAURO GIAVALISCO,¹¹ ZHIYUAN JI,¹¹ JASLEEN MATHARU,^{4,5}
IVELINA MOMCHEVA,⁷ AMBER N. STRAUGHN,¹ AND JONATHAN R. TRUMP⁸

¹*Astrophysics Science Division, Goddard Space Flight Center, Greenbelt, MD 20771, USA*

²*Department of Physics, The Catholic University of America, Washington, DC 20064, USA*

³*Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, MD 20771*

⁴*Department of Physics and Astronomy, Texas A&M University, College Station, TX, 77843-4242 USA*

⁵*George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, TX, 77843-4242 USA*

⁶*Department of Astronomy, The University of Texas at Austin, Austin, TX, 78759 USA*

⁷*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD, 21218 US*

⁸*Department of Physics, University of Connecticut, Storrs, CT 06269, USA*

⁹*Department of Astronomy, New Mexico State University, Las Cruces, New Mexico, USA*

¹⁰*Cosmic Dawn Center (DAWN) at the Niels Bohr Institute, University of Copenhagen and National Space Institute, Technical University of Denmark*

¹¹*Astronomy Department, University of Massachusetts, Amherst, MA, 01003 USA*

ABSTRACT

Reionization is an inhomogeneous process, thought to begin in small ionized bubbles of the intergalactic medium (IGM) around overdense regions of galaxies. Recent Lyman-alpha (Ly α) studies during the epoch of reionization show growing evidence that ionized bubbles formed earlier around brighter galaxies, suggesting higher IGM transmission of Ly α from these galaxies. We investigate this problem using IR slitless spectroscopy from the Hubble Space Telescope (HST) Wide-Field Camera 3 (WFC3) G102 grism observations of 148 galaxies selected via photometric redshifts at $6.0 < z < 8.2$. These galaxies have spectra extracted from the CANDELS Ly α Emission at Reionization (CLEAR) survey. We combine the CLEAR data for 275 galaxies with the Keck/DEIMOS+MOSFIRE dataset from the Texas Spectroscopic Search for Ly α Emission at the End of Reionization Survey. We then constrain the Ly α equivalent-width (EW) distribution at $6.0 < z < 8.2$, which is described by an exponential form, $dN/dEW \propto \exp(-EW)/W_0$, with the characteristic e -folding scale width (W_0). We confirm a significant drop of the Ly α strength (or W_0) at $z > 6$. Furthermore, we compare the redshift evolution of W_0 between galaxies at different UV luminosities. The UV-bright ($M_{UV} < -21$, or $L_{UV} > L^*$) galaxies show weaker evolution with a decrease of $0.4 (\pm 0.2)$ dex in W_0 at $z > 6$ while UV-faint ($M_{UV} > -21$, or $L_{UV} < L^*$) galaxies exhibit a significant drop by a factor of $0.7-0.8 (\pm 0.2)$ dex in W_0 from $z < 6$ to $z > 6$. Our results add to the accumulating evidence that UV-bright galaxies exhibit boosted Ly α transmission in the IGM, suggesting that reionization completes sooner in regions proximate to galaxies of higher UV luminosity.

1. Introduction

Reionization marks the last major phase transition of the universe, when the first stars and galaxies ionized the intergalactic medium (IGM) neutral hydrogen (H I) gas. Galaxies in the early universe are inherently coupled with the process of reionization, as the galaxies were likely the primary sources of ionizing photons (e.g., Robertson et al. 2015; McQuinn 2016; Finkelstein et al. 2019a; Dayal & Ferrara 2018) while the contribution from active galactic nucleus (AGN) activity could also play an important (though subdominant)

role at $z \gtrsim 6$ (e.g., Matsuoka et al. 2018; Kulkarni et al. 2019; Robertson 2021). Thus, observations of this epoch also provide key information for studying the dominant source of the ionizing photons: galaxies in the early universe. However, constraining the ionizing photon budget during the epoch of reionization (EoR) is still extremely difficult due to current observational limits as well as a poorly constrained ionizing photon escape fraction which depends on galaxy physical conditions (e.g., Finkelstein et al. 2012a, 2015, 2019b; Robertson et al. 2013, 2015; Bouwens et al. 2015, 2016; Kimm et al. 2019; Yoo et al. 2020; Ocvirk et al. 2021). Understanding the temporal and spatial evolution of reionization by tracing the IGM H I fraction provides a key constraint on the ionizing emissivity required from galaxies as a function of redshift.

As Lyman-alpha ($\text{Ly}\alpha$) photons are resonantly scattered by H I in the IGM, an analysis of $\text{Ly}\alpha$ can be used to trace the existence of H I gas in the IGM at different points in the history of the universe (e.g., Miralda-Escudé & Rees 1998; Rhoads & Malhotra 2001; Malhotra & Rhoads 2004; Dijkstra et al. 2014). This technique uses follow-up spectroscopic observations, targeting high- z candidate galaxies, to measure the strength of $\text{Ly}\alpha$ emission from galaxies in the reionization era. Initial studies using $\text{Ly}\alpha$ spectroscopy have found an apparent deficit of $\text{Ly}\alpha$ emission at $z > 6.5$ (e.g., Fontana et al. 2010; Pentericci et al. 2011; Finkelstein et al. 2013; Pentericci et al. 2014; Caruana et al. 2012, 2014; Curtis-Lake et al. 2012; Mallery et al. 2012; Ono et al. 2012; Schenker et al. 2012, 2014; Treu et al. 2012, 2013; Tilvi et al. 2014; Vanzella et al. 2014; Schmidt et al. 2016; De Barros et al. 2017; Fuller et al. 2020), implying an increasing H I fraction in the IGM from $z \sim 6 \rightarrow 7$, although other $\text{Ly}\alpha$ systematics with galaxy evolutionary features are a factor in these measurements (e.g., Finkelstein et al. 2012b; Yang et al. 2017; Tang et al. 2019; Trainor et al. 2019; Du et al. 2020; Hassan & Gronke 2021; Weiss et al. 2021).

Recent $\text{Ly}\alpha$ studies suggest a more complicated picture of reionization. For instance, Pentericci et al. (2018) provide $\text{Ly}\alpha$ fraction ($f_{\text{Ly}\alpha}$) measurements at $z \sim 6$ and $z \sim 7$ ($f_{\text{Ly}\alpha} = N_{\text{LAE}}/N_{\text{LBG}}$, where N_{LAE} is the number of spectroscopically-confirmed $\text{Ly}\alpha$ -emitting galaxies, and N_{LBG} is the number of high- z candidate galaxies that were targeted in spectroscopic observations). Their results show a possible flattening or a steady increase in the redshift evolution the $\text{Ly}\alpha$ fraction from $z \sim 5 \rightarrow 6$ and a relatively smoother evolution from $z \sim 6 \rightarrow 7$, compared to previous studies (e.g., Stark et al. 2011; Tilvi et al. 2014), implying a more extended ending to reionization (this finding is also supported by Kulkarni et al. 2019; Fuller et al. 2020). Furthermore, while Zheng et al. (2017), Castellano et al. (2018), and Tilvi et al. (2020) report observations of an ionized bubble via $\text{Ly}\alpha$ observations at $z \gtrsim 7$, non/rare detections of $\text{Ly}\alpha$ in Hoag et al. (2019) and Mason et al. (2019) suggest a significantly neutral fraction in the IGM at $z \sim 7.5$. Specifically, Hoag et al. (2019) report a high neutral fraction of $88^{+5}_{-10}\%$ at $z \sim 7.6$. More recently, Jung et al. (2020) analyzed deep NIR observations in GOODS-N, suggesting a modestly-ionized universe with the inferred IGM neutral fraction of $49^{+19}_{-19}\%$ at $z \sim 7.6$, lower than other $\text{Ly}\alpha$ studies at the same redshifts. Additionally, Hu et al. (2021) disclose a protocluster structure at $z \sim 6.9$ which consists of 16 spectroscopically-confirmed $\text{Ly}\alpha$ emitting galaxies. One way to reconcile these apparently contrasting recent findings is if reionization is complex and inhomogeneous, and/or if there are large spatial and temporal variations in the history of reionization.

Taken together, the evidence from recent studies suggests that $\text{Ly}\alpha$ visibility during the EoR may evolve differently in UV bright and faint galaxies (e.g., Oesch et al. 2015; Zitrin et al. 2015; Roberts-Borsani et al. 2016; Stark et al. 2017; Zheng et al. 2017; Mason et al. 2018b). This is explained as UV bright galaxies are likely located in highly ionized bubbles which were created by a wealth of ionizing photons produced by those galaxies as well as a potentially larger number of nearby fainter galaxies (Finkelstein et al. 2019b, Larson et al. 2021 in preparation). Endsley et al. (2021a) provide additional evidence for accelerated reionization around massive galaxies. Particularly, they find a higher $\text{Ly}\alpha$ detection rate at $z \simeq 7$ from massive galaxies with strong [O III]+H β emission, which reflects enhanced ionizing photo production rates (e.g., Roberts-Borsani et al. 2016; Tang et al. 2019, 2021a; Endsley et al. 2021b), arguing for higher $\text{Ly}\alpha$ equivalent-widths (EWs) from the strong [O III]+H β emitting population as similar as shown at lower redshifts of $z \simeq 2 - 3$ (Tang et al. 2021b). This may be related to the galaxies' specific star-formation rates (sSFRs) and ionization (Backhaus et al. 2021, Papovich et al., in prep), although this remains tenuous as there is not yet any conclusive evidence for a significant enhancement of Lyman continuum escape fraction found for galaxies with higher [O III]+H β galaxies at $z \sim 3$ (Saxena et al. 2021). Therefore, it is prudent to look for indications that evolution of $\text{Ly}\alpha$ emission in galaxies depends on UV luminosity in the epoch of reionization.

In this study, we present an analysis of the CANDELS $\text{Ly}\alpha$ Emission At Reionization (CLEAR) observations in the CANDELS GOODS fields (Grogin et al. 2011; Koekoemoer et al. 2011). As discussed below, CLEAR uses slitless IR spectroscopy from the HST WFC3. Therefore, the CLEAR data provide (unbiased) constraints on $\text{Ly}\alpha$ emission in galaxies at $6.0 < z < 8.2$. This allows us to constrain the evolution of the $\text{Ly}\alpha$ EW in these galaxies. We use these data to study the evolution as a function of both redshift and galaxy UV absolute magnitude. Section 2 describes the CLEAR high- z galaxy dataset (data reduction, sample selection, and emission-line and continuum-detection search) and the additional Keck DEIMOS and MOSFIRE observations from the Texas Spectroscopic Search for $\text{Ly}\alpha$ Emission at the End of Reionization Survey (Jung et al. 2018, 2020), finalizing the $6.0 < z < 8.2$ galaxy dataset for the $\text{Ly}\alpha$ analysis. In Section 3, we present our results, which includes the measurements of the $\text{Ly}\alpha$ EW distribution and the IGM transmission to $\text{Ly}\alpha$. We summarize and discuss our findings in Section 4. In this work, we assume the Planck cosmology (Planck Collaboration et al. 2016) with $H_0 = 67.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.308$, and $\Omega_\Lambda = 0.692$. The Hubble Space Telescope (HST) F435W, F606W, F775W, F814W, F850LP, F105W, F125W, F140W, and F160W bands are referred to as B_{435} , V_{606} , i_{775} , I_{814} , z_{850} , Y_{105} , J_{125} , JH_{140}

and H_{160} , respectively. All magnitudes are given in the AB system (Oke & Gunn 1983), and all errors presented in this paper represent 1σ uncertainties (or central 68% confidence ranges), unless stated otherwise. All EWs discussed in this paper represent the rest-frame values, unless defined otherwise.

2. Data: $6.0 < z < 8.2$ Galaxies

2.1. CLEAR HST/grism Survey

The CLEAR Experiment is a cycle 23 HST observing program (Program GO-14227, PI: C. Papovich), which observed 12 fields in the CANDELS GOODS fields to 10 to 12-orbit depth with the G102 grism in the HST/WFC3 camera. Each field was observed at 3 position angles (separated by >10 degrees) to properly correct galaxy spectra from contamination. The goal of the CLEAR survey is to measure the distribution of $\text{Ly}\alpha$ emission in galaxies during the epoch of reionization at $6.0 < z < 8.2$. The feasibility of detecting $\text{Ly}\alpha$ emission from HST/WFC3 grism observations has been proved in previous studies, such as Schmidt et al. (2016) from the GLASS survey (Treu et al. 2015), and Tilvi et al. (2016) and Larson et al. (2018) from the FIGS survey (Pirzkal et al. 2017). The dataset also provides constraints on the stellar populations of $1 < z < 2$ galaxies. Previous work on the metallicities, ages, and formation histories of massive galaxies at $1 < z < 2$ has been published in Estrada-Carpenter et al. (2019, 2020), and the gas-phase metallicity gradients of star-forming galaxies is investigated in Simons et al. (2020). Also, Cleri et al. (2020) studied Paschen- β as a star-formation rate indicator in low redshift galaxies, using the CLEAR dataset.

The dataset has been extended to include all publicly-available HST/WFC3 G102 and G141 grism observations in the CLEAR fields. For processing the grism observations, the grism redshift and line analysis software GRIZLI (Brammer 2019) has been utilized, which retrieves the raw observations and performs astrometric alignment, contamination modeling, extracting spectra, and fitting continuum and emission line models. The full details of the grism data reduction and spectral extractions are described in Estrada-Carpenter et al. (2019, 2020) as well as Simons et al. (2020).

While the primary CLEAR spectral extractions were made for galaxies based on the 3D-HST GOODS catalog (Skelton et al. 2014), we extracted the CLEAR spectra of high- z galaxies based on the updated HST CANDELS photometry and its segmentation maps from Finkelstein et al. (2021). The photometric selection of high- z candidate galaxies was done following the criteria described in Section 3.2 in Finkelstein et al. (2015). In brief, the selection is based on the full photometric redshift probability distribution functions of $P(z)$ calculated by EAZY (Brammer et al. 2008) rather than simply using the best-fit redshifts. This includes, for example, the integral of $P(z)$ under the primary redshift peak must be

$>70\%$ of the total $P(z)$ (for the full detail, refer to Finkelstein et al. 2015). Additionally, visual inspection was performed for removing any artifacts (e.g., diffraction spikes, nearby bright sources) and checking the quality of photometry. This resulted in 180, $6.0 < z_{\text{phot}} < 8.2$ galaxies in the CLEAR fields. However, checking the quality of the extracted spectra, in total 148 CLEAR spectra were collected after removing 32 spectra due to technical issues in treating the removal and corrections from contaminating spectrum from nearby galaxies using GRIZLI (e.g., we removed objects where residuals from contaminating spectra from nearby bright galaxies were too severe compared to the expected continuum or line flux for our objects).

2.2. Emission Line and Continuum Detection in CLEAR

In order to search $\text{Ly}\alpha$ emission lines or continuum objects detected from these high- z targets, we utilized the grism spectroscopy from the full sample of the CLEAR spectra of 148, $6.0 < z_{\text{phot}} < 8.2$ galaxies. In this subsection, we describe the process we used to determine if any emission lines or continuum spectra are present in the spectra of each galaxy.

2.2.1. Emission-Line Search

We first found 24 galaxies from the GRIZLI run with candidate emission lines detected at significance $>3\sigma$ from their co-added 1-dimensional (1D) spectra. For these galaxies, we inspected their 2-dimensional (2D) spectra and those of nearby objects to see if the emission lines are a result of contamination. Additionally, we attempted to secure detections from multiple PAs, requiring $>2\sigma$ detections in the data taken at different orientations. This assists in our effort to rule out artefacts in the data including contamination from nearby objects (as these will be different in each PA).

Our emission-line search identified only one tentative $\text{Ly}\alpha$ detection at high confidence. This object (GN4.5461276) shows an emission line that would correspond to $z_{\text{grism}} = 6.513 \pm 0.005$ with a detection significance of $\simeq 4\sigma$. All other candidates showing potential $\text{Ly}\alpha$ emission either correspond to residuals of nearby contaminating sources (16 candidates) or possible spurious sources that show the emission in only a single PA (7 candidates). In the case of the emission line detected in GN4.5461276, the HST ACS z_{850} -band data challenges this measurement: if the emission line (interpreted as $\text{Ly}\alpha$) is real, then it should contribute 37 ± 8 nJy to the z_{850} -band flux. However, the measured z_{850} -band flux of this galaxy in the HST/CANDELS observations is only $f_{\nu}(z_{850}) = 6.3 \pm 8.3$ nJy. Thus, it may require further observational evidence for validating this tentative detection as $\text{Ly}\alpha$, and we do not include it in the remainder of our analysis for the $\text{Ly}\alpha$ EW distribution and the IGM transmission. Instead, we discuss the details of this tentative detection in the Appendix. Additionally, we suggest that it would be

crucial to include an automated and improved removal process of contaminating emission-line sources in slitless spectroscopic surveys. This will be very important in the study of emission lines in data from the next generation space telescopes such as the James Webb Space Telescope (JWST), Euclid, or Nancy Grace Roman Space Telescope (NGRST).

2.2.2. Continuum-Detection Search

We also inspected 42 sources with potential continuum detections identified from our visual inspection of the coadded 1D and 2D grism spectra. We vetted these further to determine if they are bona fide continuum sources. First, we estimated Y_{105} , J_{125} , or H_{160} magnitudes from the CLEAR spectra of the 42 candidates by applying the HST filter transmission curves to the reduced co-added 1D CLEAR spectra and compared them to the known HST magnitudes of the sources from the HST/CANDELS photometry from [Finkelstein et al. \(2015\)](#). We required agreement within the 2σ uncertainties in this comparison. The continuum flux is estimated by summing all fluxes after the continuum break, and the continuum break is located where the estimated signal-to-noise (S/N) value of the continuum flux is maximized. Thus, we required a secure (i.e., $>3\sigma$) continuum detection redward the Lyman- α continuum break measured from the GRIZLI-extracted 1D spectrum.

From the selection above, we obtained 15 candidates which satisfy the continuum selection criteria. On this subset, we then performed additional visual inspections of the galaxies' 2D and 1D spectra taken at the different PAs to check for possible contamination from nearby sources. From these inspections, we identified contaminating sources in 6 of these 15 candidates, and no other candidates display clear continuum detection in multiple PAs due to the limited observing depth (as the depth in each individual PA is $\sim 1/3$ rd the exposure time of the total). In general, therefore, the depth of the data is insufficient for reliable detection of the continuum in our target galaxies of $J_{125} \gtrsim 26$ at individual PA-depth (~ 3 orbits). However, we identify possible continuum in one source (GN2_23734), which we discuss in the Appendix. This galaxy shows possibly faint continua in both G102 and G141 spectra. However, confirmation of this source would require deeper observations to locate the continuum break confidently.

While the current analysis of the slitless grism observations has achieved a great success in removing contamination from moderate brightness objects, this remains a factor when studying emission in faint galaxies. For example, even 1% residuals from the spectra of contaminating sources leave a signal as bright as that from objects of interest for magnitude differences of $\Delta m = 5$ mag (as will be the case for 27th-magnitude galaxies near galaxies brighter than 22nd magnitude). Mitigating the effects of contamination will be impor-

tant as detecting galaxy continuum breaks provides a means to confirm high- z galaxies without Ly α : this will be particularly useful into the epoch of reionization where observable Ly α emission may be less frequent (e.g., [Rhoads et al. 2013](#); [Watson et al. 2015](#); [Oesch et al. 2016](#)).

In summary, we do not identify any convincing Ly α emission or continuum detected galaxies from the CLEAR spectra. However, this in itself is an important finding as the CLEAR data for $148 \ 6.0 < z_{\text{phot}} < 8.2$ candidate galaxies still provides constraints on the evolution of the Ly α EW distribution. This is because some bright Ly α emission lines could (even *should*) have been detected with the CLEAR observing depth if the Ly α EW distribution does not evolve from $z \lesssim 6$ to $z \gtrsim 6$. The non-detections in CLEAR rule out the existence of large EW Ly α sources in the CLEAR sample. We utilize the CLEAR spectra of these $148 \ 6.0 < z_{\text{phot}} < 8.2$ candidate galaxies for the remainder of this study to set constraints on the Ly α EW distribution.

2.3. Supplemental data from Keck DEIMOS and MOSFIRE Observations

The Texas Spectroscopic Search for Ly α Emission at the End of Reionization Survey ([Jung et al. 2018, 2019, 2020](#)) obtained deep spectroscopic data of over 120 $z > 6$ galaxies in the GOODS-N fields (65 galaxies from DEIMOS and 62 galaxies from MOSFIRE observations). These data provided the largest number of confirmed Ly α emitters at $z > 7$ with which to constrain the IGM H I fraction at $z \sim 7.6$ ([Jung et al. 2020](#)). Using these data, [Jung et al. \(2020\)](#) noted tentative evidence that the Ly α EW distribution at $z > 7$ depends on the galaxies' UV luminosities. Here we use the CLEAR sample to explore this finding, as the CLEAR data provides constraints on Ly α emission for a larger sample of faint galaxies.

CLEAR aims to measure the Ly α EW distribution during the reionization era at $z > 6$. Also, CLEAR is a blind survey with slitless spectroscopy, providing galaxy spectra for a wide dynamic range of UV magnitudes. Even non-detections in CLEAR are still highly constraining, placing upper limits of Ly α visibility ([Treu et al. 2012](#); [Jung et al. 2018](#); [Mason et al. 2019](#)). The combination of the CLEAR data and Keck data then provided important, complementary data. This allows us to explore the evolution of the Ly α EW distribution and its dependence on UV absolute magnitude (M_{UV}) at higher significance than possible with either dataset separately.

2.4. Physical Properties of Combined Sample

We derived the properties of our 148 sources in CLEAR by performing spectral energy distribution (SED) fitting with the [Bruzual & Charlot \(2003\)](#) stellar population model, following [Jung et al. \(2019, 2020\)](#). We utilized the HST ACS and




Figure 1. The left panel shows the M_{UV} distribution of target galaxies in the CLEAR dataset as a function of redshift, and the $M_{star}-M_{UV}$ relation in the right panel. All 148 galaxies in the CLEAR sample are shown as open circles while the targets from the Keck observations are denoted as filled blue (DEIMOS: 65 galaxies) and red (MOSFIRE: 62 galaxies) circles. A wide dynamic range of UV magnitudes in the CLEAR sample apparently show an unbiased nature of the blind survey, compared to the targeted Keck observations. The blue solid line in the right panel shows the fiducial $z \sim 7$ relation of Song et al. (2016a) with the shaded area of its uncertainty.

WFC3 broadband photometry (B_{435} , V_{606} , i_{775} , I_{814} , z_{850} , Y_{105} , J_{125} , JH_{140} and H_{160}) in addition to Spitzer/IRAC 3.6 μ m and 4.5 μ m band fluxes. Here, we use photometric redshift measurements that have been updated in Finkelstein et al. (2021) based on updated CANDELS photometry, including deeper I_{814} and Spitzer/IRAC data (where those authors utilized a deblending technique to measure IRAC photometry more accurately using the HST images as priors). A Salpeter (1955) initial mass function was assumed with a stellar mass range of 0.1-100 $M_{\odot}$, and we allowed a metallicity range of 0.005-1.0 $Z_{\odot}$. We explored a range of exponential models of star formation histories with exponentially varying timescales ($\tau = 10$ Myr, 100 Myr, 1 Gyr, 10 Gyr, 100 Gyr, -300 Myr, -1 Gyr, -10 Gyr). We use the Calzetti (2001) dust attenuation curve with $E(B-V)$ values ranging from 0 to 0.8 for modeling galaxy spectra and add nebular emission lines, adopting the Inoue (2011) emission-line ratio, as done in Salmon et al. (2015). The IGM attenuation was applied to the SED models based on Madau (1995).

Physical properties were obtained from the best-fit models, minimizing χ^2 . The uncertainties of physical parameters were estimated from SED fitting with 1000 Monte Carlo (MC) realizations of the perturbed photometric fluxes, according to their photometric errors for individual galaxies. UV magnitudes (M_{UV}) of galaxies were calculated from the averaged fluxes over a 100Å-bandpass (at 1450-1550Å) from the best-fit models, which were not dust-corrected. In our SED fitting, we fixed redshifts with the best-fit photometric redshifts calculated by EAZY.

In the left panel of Figure 1 we present the M_{UV} distribution of our CLEAR targets as a function of redshift with the existing Keck DEIMOS and MOSFIRE targets from Jung

et al. (2018, 2020). The black open circles show the CLEAR sample, and the blue and red filled circles denote targets from DEIMOS and MOSFIRE observations, respectively. The CLEAR galaxy sample does not show any noticeable bias, randomly distributed over a range of $-18 \gtrsim M_{UV} \gtrsim -22$, giving more UV faint ($M_{UV} > -19$) targets than the Keck observations. However, the Keck observations are much deeper than CLEAR in general, providing significantly lower EW limits for those targeted galaxies. Figure 1 also shows the stellar mass (M_{star}) - M_{UV} distribution of our CLEAR galaxies in the right panel, compared to the fiducial relation derived at $z \sim 7$ by Song et al. (2016b). Similar to the Keck sample, most of the CLEAR targets are placed on the fiducial $M_{star}-M_{UV}$ relation at $z \sim 7$ of Song et al. (2016b) with no significant selection bias from the typical high- z galaxy population, although there are a few UV faint and massive galaxies located above the fiducial relation.

3. Ly α EW Distribution and IGM Transmission

3.1. Measuring the Ly α EW Distribution at $6.0 < z < 8.2$

The Ly α EW distribution is commonly described by a declining exponential form,

$$P(EW) \propto \exp^{-EW/W_0}, \quad (1)$$

where EW is the Ly α rest-frame EW, and W_0 is a e -folding scale width (e.g., Cowie et al. 2010). Jung et al. (2018) introduced an improved methodology of measuring the Ly α EW distribution, motivated by earlier studies (e.g., Treu et al. 2012, 2013), which utilizes galaxies without detected emission lines as well as Ly α detected objects. Following Jung et al., we constructed a template of an expected number of Ly α emitters as a function of detection signifi-

cance, by accounting for all types of data incompleteness including instrumental wavelength coverage, the wavelength-dependent $\text{Ly}\alpha$ detection limit, the UV-continuum flux, and the photometric redshift probability distribution function, $P(z)$, based on the combined dataset of the CLEAR spectra. We then applied this to the CLEAR dataset, and combined these results with those from our previous work using the Keck/DEIMOS+MOSFIRE observations in Jung et al. (2018, 2019, 2020) to measure the $\text{Ly}\alpha$ EW distribution.

To calculate the expected number of $\text{Ly}\alpha$ emitters for a given model we did the following. We first estimated the detection sensitivity for an unresolved emission line in the grism spectrum for each object. The detection limit was inferred through Monte Carlo simulations with the 1D spectra by assigning a mock $\text{Ly}\alpha$ emission line to the reduced 1D spectra for all objects. We recovered the line flux of this mock emission line with its error from the Gaussian line fitting with the IDL MPFIT package (Markwardt 2009). This mock emission line was created to have an intrinsic line profile equal to the best-fit line profile from one of the highest-S/N $\text{Ly}\alpha$ emission detected in z7_GND_42912 from the MOSFIRE observations of Jung et al. (2019, 2020), and we smoothed the line profile to match the G102 resolution ($\sim 45\text{\AA}$). We note that the choice of the emission line profile has a negligible effect given the relatively low resolution of the G102 grism data ($R \sim 200$). We consider our targets to be point-like sources in our 1D mock emission simulation. This assumption is appropriate as our target galaxies at this redshift are small, with a typical effective radius (R_{eff}) of $R_{\text{eff}} < 1$ kpc in general (or $\lesssim 3$ pixels at the resolution of *HST*/WFC3) and there is evidence that these kinds of galaxies are even smaller at higher redshifts (Shibuya et al. 2019). Additionally, in GRIZLI, our 1D spectra are extracted using the source segmentation maps from the photometric catalog, thus the extracted 1D spectra include the contribution estimated from the (rest-frame UV) morphology as measured by the Y_{105} band. We may not capture the contribution of $\text{Ly}\alpha$ from extended halos in 1D spectra (e.g., Leclercq et al. 2017; Wisotzki et al. 2018). However, an appropriate modeling of the extended component of $\text{Ly}\alpha$ at this redshift is beyond the scope of this study. In future, for example, JWST NIRSpec IFU observations will resolve the extended component of $\text{Ly}\alpha$.

From these simulation emission lines we can derive emission-line flux limits for the CLEAR G102 data. The top panel of Figure 2 shows the 3σ detection limits over the wavelength range of the G102 grism, which corresponds to a redshift range of for $\text{Ly}\alpha$ emission at $6.0 \lesssim z \lesssim 8.2$. In the figure, the colored dots show the detection limits for all the 148 individual galaxies in our sample; the red curve shows the median detection limit for all spectra. The bottom panel of Figure 2 shows the corresponding (median) 3σ rest-frame




Figure 2. (Top) 3σ emission-line detection limits across the wavelengths corresponding to the WFC3 G102 redshift coverage of $\text{Ly}\alpha$ at $6.0 < z < 8.2$. The red curve represents the median detection limit, and the detection limits of the 148 individual galaxies analyzed here are shown as colored dots. The 3σ detection limit reaches down to below $\sim 1 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ around $1.0 - 1.1 \mu\text{m}$ where the G102 grism transmission is high. (Bottom) The median 3σ rest-frame EW limits as a function of wavelength for varying apparent UV magnitudes (as labeled).

EW limits for galaxies as a function of apparent UV magnitude. The EW limits were estimated by dividing the median 3σ emission-line flux limits ($f_{3\sigma\text{-limit}}$) by the continuum level (f_{cont}) and $(1 + z_{\text{Ly}\alpha})$ at each corresponding wavelength as below.

$$\text{EW}_{3\sigma\text{-limit}} [\text{\AA}] = \frac{f_{3\sigma\text{-limit}}}{f_{\text{cont}}(1 + z_{\text{Ly}\alpha})} \quad (2)$$

After detection limits estimated, we constructed a template of an expected number of detected $\text{Ly}\alpha$ emission lines as a function of detection significance. The template construction was done in a Monte Carlo (MC) fashion, by simulating mock $\text{Ly}\alpha$ emission lines for entire target galaxies, following Jung et al. (2018, 2020). The MC simulation of individual mock emission lines has three main steps: (i) we allocate the wavelength of the simulated $\text{Ly}\alpha$ emission, randomly taken based on an object's $P(z)$; (ii) at each redshift, we simulate a $\text{Ly}\alpha$ line flux given the galaxy's magnitude and EW, where the latter is drawn from the distribution in Equation 1 over a range for the assumed $\text{Ly}\alpha$ EW scale width, $W_0 = 5\text{--}200 \text{\AA}$; (iii) finally, we measure a detection significance for this simulated emission line based on the precomputed emission-line detection limits taken from above. For each choice of W_0 , we carried out 1000 sets of MC simulations of the mock emission lines for the entire target galaxies, which produces a distribution of the expected number of $\text{Ly}\alpha$ detections as a function of detection significance.




Figure 3. Similar to Figure 11 in Jung et al. (2020): the redshift dependence of the Ly α EW e -folding scale width (W_0) up to $z \sim 8$. The figure includes a compilation of the previous W_0 measurements (Gronwall et al. 2007; Ouchi et al. 2008; Nilsson et al. 2009; Guaita et al. 2010; Hu et al. 2010; Kashikawa et al. 2011; Ciardullo et al. 2012; Zheng et al. 2014; Wold et al. 2014, 2017; Hashimoto et al. 2017; Jung et al. 2018; Pentericci et al. 2018; Santos et al. 2020; Jung et al. 2020). Our updated W_0 measurements with CLEAR are denoted by red, large symbols: horizontal bars at $z \sim 6.5$ and a filled circle at $z \sim 7.6$. The $z \sim 7.6$ measurement is displayed at $z = 7.5$ to avoid an overlap with the Jung et al. (2020) data point. Our study confirms a decreased W_0 at $z > 6$, whereas there is little/no redshift evolution of W_0 reported in the literature at $z < 6$.

Lastly, we fit the data for each galaxy to the suite of simulated templates over the range of redshift and emission line EW. We then combined these results to construct a probability distribution function (PDF) for the e -folding scale width (W_0) of the Ly α EW distribution. We utilized a Markov Chain Monte Carlo (MCMC) algorithm for this fitting. As counting the number of Ly α emission line detections is a general Poisson problem, we used a Poisson likelihood. We used the Cash (1979) statistic to describe the Poisson likelihood. We then used a Metropolis-Hastings MCMC sampler (Metropolis et al. 1953; Hastings 1970) to construct the PDF of W_0 , generating 10^5 MCMC chains. For more information on this methodology, see Jung et al. (2019, 2020).

3.2. The Evolution of Ly α EW at $z > 6$

Jung et al. (2018) constrained the characteristic e -folding scale width W_0 of the Ly α EW distribution at $6 < z < 7$ using data from Keck/DEIMOS for a sample of galaxies. They measured 1σ and 2σ upper limits of $W_0 < 36\text{\AA}$ and $< 125\text{\AA}$, respectively. Similarly, Jung et al. (2020) measured $W_0 = 32^{+14}_{-9}\text{\AA}$ for galaxies at $7.0 < z < 8.2$ using data from Keck/MOSFIRE observations. These values are shown in Figure 3.

Here, we report new measurements on the evolution of the Ly α EW from the combined analysis of the CLEAR and Keck datasets. The CLEAR data provide important constraints on the evolution of Ly α as the lack of detections implies strong evolution in the number of sources with high Ly α EWs at these redshifts. The updated constraints improve the constraint on the EW e -folding scale width, W_0 , to be $< 17\text{\AA}$ (1σ) and $< 28\text{\AA}$ (2σ) at $6.0 < z < 7.0$ and $W_0 = 27^{+10}_{-7}\text{\AA}$ at $7.0 < z < 8.2$ (these are shown as red, large symbols in Figure 3). These measurements confirm the decline of W_0 already seen at $6.0 < z < 7.0$ compared to $z < 6$.

3.3. The Dependence of the Ly α EW on UV Magnitude

Observations at lower redshifts ($2 \lesssim z \lesssim 6$) show the the Ly α EW e -folding scale width (W_0) decreases with increasing UV luminosity (e.g., Ando et al. 2006; Stark et al. 2010; Schaerer et al. 2011; Cassata et al. 2015; Furusawa et al. 2016; Wold et al. 2017; Hashimoto et al. 2017; Oyarzún et al. 2017; Santos et al. 2020). In contrast, Jung et al. (2020) reported tentative evidence that W_0 increases with increasing UV luminosity at $z > 6$ from the brightest galaxies ($M_{\text{UV}} < -21$). Although the initial evidence was weak due to the large measurement errors, this finding is consistent with a picture of reionization where the IGM around brighter




Figure 4. (Left) Measurements of the e -folding scale width (W_0) of the Ly α EW distribution at different M_{UV} . Our W_0 measurements from this study (combining the CLEAR and Keck samples) at three M_{UV} bins are shown as red filled circles, compared with those from lower- z observations in Santos et al. (2020). Our W_0 measurement from the brightest UV magnitude galaxies show weaker evolution from $z < 6$ to $z > 6$ while those from fainter UV magnitude galaxies present a significant drop from $z < 6$ to $z > 6$. (Right) Ly α Transmission in the IGM ($T_{IGM}^{Ly\alpha}$) at different M_{UV} . Our result indicates that the IGM transmission to Ly α during the reionization epoch is high from UV bright galaxies, compared to those from UV faint ones. The gray lines in the backgrounds in both panels represent 1000 random draws from the 10^5 MCMC chains that were generated for constructing the PDF of W_0 . We find that 76% of the MCMC chains show an upturn in W_0 (or a greater W_0) in the UV brightest bin as shown in the left panel. In addition, we find that 91% of the chains suggest an increasing trend between the Ly α IGM transmission and the (increasing) UV luminosity in the right panel.

Table 1. Ly α EW e -folding Scale Width (W_0) and the Ly α IGM Transmission ($T_{IGM}^{Ly\alpha}$) at Different M_{UV}

M_{UV} (L_{UV}/L^* ^a)	N_{galaxy}	W_0		$T_{IGM}^{Ly\alpha}$
		$z \sim 2-6$	$6.0 < z < 8.2$	$6.0 < z < 8.2$
		Santos et al. (2020)	This study	
$-20 < M_{UV} < -19$ ($0.15 < L_{UV}/L^* < 0.4$) ^b	77	$178^{+13}_{-13} \text{ \AA}$	$29^{+16}_{-13} \text{ \AA}$	$0.16^{+0.11}_{-0.08}$
$-21 < M_{UV} < -20$ ($0.4 < L_{UV}/L^* < 1.0$)	92	$73^{+10}_{-10} \text{ \AA}$	$14^{+5}_{-4} \text{ \AA}$	$0.18^{+0.11}_{-0.07}$
$-22 < M_{UV} < -21$ ($1.0 < L_{UV}/L^* < 2.5$)	30	$54^{+11}_{-11} \text{ \AA}$	$20^{+10}_{-6} \text{ \AA}$	$0.37^{+0.31}_{-0.17}$

^a $L_{UV}/L^* = 10^{0.4(M_{UV}^* - M_{UV})}$, where M_{UV}^* is the characteristic magnitude of a typical galaxy with an approximate value of $M_{UV}^* \simeq -21$ at $z \sim 7$ (Finkelstein et al. 2015).

^b Our sample in this faintest UV magnitude group could be incomplete in the CANDELS GOODS Wide fields due to the ~ 1 magnitude shallower detection limits in the Wide fields than those in the CANDELS GOODS Deep fields as shown in Figure 1 (left). Thus, we also calculated W_0 in $-20 < M_{UV} < -19$ excluding objects in the Wide fields, which provides $W_0 = 30^{+20}_{-15} \text{ \AA}$. Although it is slightly increased from our fiducial value listed in the table, the change is insignificant and does not alter our conclusion.

galaxies becomes ionized earlier than that of fainter objects, allowing for more efficient Ly α -photon escape (e.g., Zheng et al. 2017; Mason et al. 2018a; Endsley et al. 2021a).

CLEAR provides a means to improve the constraint on the evolution of W_0 as a function of absolute UV magnitude as the CLEAR data cover many more lower-luminosity objects. We therefore recalculated W_0 from the combined dataset of CLEAR and Keck/DEIMOS+MOSFIRE observations in dif-

ferent bins of UV absolute magnitude (M_{UV}). These W_0 measurements are listed in the third column in Table 1 and shown in the left panel of Figure 4 as filled red dots. From this analysis, the CLEAR data improve the evidence that the e -folding scale width (W_0) of the Ly α EW distribution at $6.0 < z < 8.2$ shows a possible upturn of W_0 in the brightest UV magnitude bin, $-22 < M_{UV} < -21$, although it is consistent with no upturn at 1σ level. The PDF of W_0

is constructed from 10^5 MCMC chains as described in Section 3.1. We then inspect what fraction of the MCMC chains have an upturn of W_0 in the brightest UV magnitude bin. We find that 76% of the MCMC chains show such an upturn (or a greater W_0). In addition, we find that 91% of the chains show an increasing trend between the Ly α IGM transmission and the (increasing) UV luminosity as shown in Figure 4. We discuss this further in the next subsection. This is in strong contrast with the measurements at $2 < z < 6$ that show W_0 decreases with increasing UV luminosity (Santos et al. 2020, shown as black open circles), indicating that at $z > 6$ the Ly α emission for UV-brighter galaxies is less affected by the IGM compared to UV-fainter galaxies.

We note that we cannot completely rule out the possibility of the redshift evolution of galaxy physical properties (e.g., ISM condition), which alters the Ly α properties significantly even without the IGM effect (Hassan & Gronke 2021). However, such decreasing trend of W_0 has been reported from most lower redshifts studies at $z < 6$ (see references earlier in this section), and evolutionary features of Ly α -emitting galaxies are not expected to be extreme particularly between $z < 6$ and $z > 6$.

3.4. Ly α Transmission in the IGM

Our W_0 measurements indicate different evolution of the Ly α EW distribution between bright and faint galaxies into the epoch of reionization, $z \gtrsim 6$. As shown in Figure 4, the W_0 values obtained for the UV-fainter objects ($M_{\text{UV}} > -21$) at $z > 6$ are substantially lower than those at lower redshift ($2 < z < 6$), whereas there is milder evolution in W_0 for the UV-brighter objects ($M_{\text{UV}} < -21$) from $z < 6$ to $z > 6$.

The evolution of W_0 leads to constraints on the relative Ly α transmission, $T_{\text{IGM}}^{\text{Ly}\alpha}$ which we define as the ratio of observed-to-expected (intrinsic) rest-frame EW, $\text{EW}_{\text{obs}}/\text{EW}_{\text{int}}$, therefore,

$$T_{\text{IGM}, z > 6}^{\text{Ly}\alpha} = \frac{\text{EW}_{z > 6}}{\text{EW}_{z < 6}} \quad (3)$$

where we take $\text{EW}_{z < 6}$ as the intrinsic EW after the IGM became completely ionized. Thus, we adopted the measurements of the EW distribution at $2 < z < 6$ from Santos et al. (2020) as the intrinsic values and calculated the IGM transmission of Ly α at $z > 6$ with our W_0 measurements.

The estimated $T_{\text{IGM}}^{\text{Ly}\alpha}$ values are listed in the last column in Table 1. Tentatively, the fact that the IGM transmission for Ly α photons appears to be higher for the UV-brightest galaxies can be interpreted as evidence that the IGM is highly ionized and more transparent around the brightest galaxies at $z > 6$, albeit this result will require further confirmation with larger and more sensitive datasets from future telescopes, such as JWST and NGRST. In contrast, the decrease in IGM

transmission for fainter galaxies ($M_{\text{UV}} > -21$) can be interpreted as evidence that the IGM is more neutral where Ly α photons suffer significantly more attenuation. This is consistent with a picture of the spatially inhomogeneous process of reionization where large ionized bubbles were often created by bright galaxies (e.g., Zheng et al. 2017; Castellano et al. 2018; Jung et al. 2019, 2020; Hu et al. 2021; Endsley et al. 2021b; Park et al. 2021; Qin et al. 2021), and we expand on this interpretation in the next section.

We caution that our definition of the Ly α IGM transmission with the ratio of observed-to-expected (intrinsic) EW could be overly simplistic as it cannot describe variations of the Ly α IGM transmission depending the reionization history and ionized bubble sizes as well as Ly α emission properties of individual galaxies. In principle, the Ly α transmission can be modeled using resionization simulations (and this has been the focus of several previous studies, e.g., Mesinger et al. 2015; Mason et al. 2018a; Weinberger et al. 2018). However, such approaches still depend on other assumptions (e.g., the reionization history, the evolution of ionized bubble sizes, and M_{UV} – halo mass relation). Here, our intent is to provide results that are model-independent from the latest sets of Ly α observations combined with our empirical approach. In future work, it will be very useful to extend these investigations to study how the Ly α IGM transmission varies depending on the reionization history and the characteristic ionized bubble sizes around galaxies (e.g., Garel et al. 2021; Park et al. 2021; Smith et al. 2021) as well as the intrinsic Ly α emission properties of galaxies during the epoch of reionization such as Ly α velocity offset (e.g., Stark et al. 2017; Hashimoto et al. 2018; Hutchison et al. 2019).

4. Conclusions and Discussion

We extracted HST slitless grism spectra of 148 high- z galaxies in the GOODS fields from the CLEAR survey. Combining our high- z dataset from CLEAR to the existing Keck Ly α observations from the Texas Spectroscopic Search for Ly α Emission at the End of Reionization Survey (Jung et al. 2020), we provide an improved measurement of the e -folding scale width (W_0) of the Ly α EW distribution at $z > 6$ and examine the dependence of W_0 on the UV luminosity (M_{UV}). Our findings are summarized as follows.

1. We inspected the reduced CLEAR spectra of high- z candidate galaxies in order to search for Ly α emission-lines or continuum-break detections. The search finds only a tentative ($\sim 4\sigma$) Ly α detection from a $z_{\text{grism}} = 6.513 \pm 0.005$ galaxy and a continuum detection from a $z_{\text{grism}} = 6.2_{-0.8}^{+2.5}$ galaxy. We discuss these more in the Appendix, although we cannot confirm their nature as high- z sources from the current data due to the incomplete contamination removal and the shallow observing depth.




Figure 5. Cartoon summarizing the inhomogeneous process of reionization at $z > 6$ universe consistent with the data and interpretation here. Galaxies with higher UV luminosity have larger ionized bubbles around them. This allows a higher $\text{Ly}\alpha$ transmission to the IGM. These UV bright galaxies are likely to be centered in overdense regions where they cluster with fainter galaxies (and these may be beyond current detection limits). The combination of ionizing photons from the galaxies in the overdense regions may provide significant ionizing photon budget around UV bright galaxies (Finkelstein et al. 2019b; Endsley et al. 2021a, Larson et al. 2021, in prepration). In contrast, UV-fainter galaxies have more neutral gas around them (i.e., they sit in smaller ionized bubbles), thus the $\text{Ly}\alpha$ transmission to the IGM is suppressed. At lower redshift, where the IGM is completely ionized the $\text{Ly}\alpha$ emission has overall high transmission for all galaxies. In the figure, the darker (black) color shows a more ionized medium while a neutral gas is represented as bright (white) color. UV-brighter galaxies are shown with larger symbols.

2. With the combined dataset of CLEAR and the Keck observations, we compute the 1σ (2σ) upper limit of W_0 at $<17\text{\AA}$ ($<28\text{\AA}$) at $6.0 < z < 7.0$ and $27^{+10}_{-7}\text{\AA}$ at $7.0 < z < 8.2$. This is illustrated in Figure 3. This confirms a significant drop of W_0 previously suggested at $6.0 < z < 7.0$. The CLEAR data improved these constraints on the evolution of W_0 compared to the previous analysis using only the Keck data. This is primarily a consequence of the spectral measurements for an increased number of galaxies, pushing to lower UV luminosities ($L_{\text{UV}} \approx 0.025L^*$) in the CLEAR sample.
3. We constrained the W_0 values and estimated the IGM transmission of $\text{Ly}\alpha$ at $z > 6$ as a function of UV absolute magnitude. This is shown in Figure 4. The $\text{Ly}\alpha$ EW distribution at $6.0 < z < 8.2$ show a possible upturn of W_0 in the UV brightest galaxies ($-22 < M_{\text{UV}} < -21$) at a 76% confidence level, in contrast with the strong decline in W_0 with increasing

UV brightness for galaxies at $z < 6$. Consequently, the IGM transmission to $\text{Ly}\alpha$ at $z > 6$ appears to be higher for the UV brightest galaxies with 91% confidence.

4. We interpret the evolution in W_0 as a function of UV luminosity as a change in the transmission of the IGM to $\text{Ly}\alpha$ at $z \gtrsim 6$ that depends on UV luminosity. This is consistent with a picture where the IGM around the brightest galaxies ($-22 < M_{\text{UV}} < -21$) is largely ionized, allowing high $\text{Ly}\alpha$ transmission. In this picture, the IGM is spatially inhomogeneous where large ionized bubbles are created by UV bright galaxies, while the IGM around fainter galaxies is more neutral, leading to strong attenuation of $\text{Ly}\alpha$ photons and lower transmission.

This picture builds on other recent observations that support an inhomogeneous nature of the IGM during reionization. We illustrate this picture in Figure 5. Studies of the IGM neutral fraction from $\text{Ly}\alpha$ emission show disparity in

their measurements at $z \sim 7-8$ (Mason et al. 2018a, 2019; Hoag et al. 2019; Jung et al. 2020). Particularly, a higher Ly α detection rate reported in Jung et al. (2020) suggests an overdense and highly ionized region in the GOODS-N field. Recent studies of the Lyman-alpha damping wing in QSO spectra also provide a large scatter of the IGM neutral fraction at the same redshift (McGreer et al. 2015; Davies et al. 2018; Wang et al. 2020; Yang et al. 2020). One explanation for such disagreement between the IGM neutral fraction measurements is that reionization did not occur spatially-uniformly, but in an inhomogeneous way (e.g., Finlator et al. 2009; Pentericci et al. 2014; Katz et al. 2019).

In the same context, recent findings of clustered LAEs from bright galaxies in the middle phase of reionization (Zheng et al. 2017; Castellano et al. 2018; Tilvi et al. 2020; Jung et al. 2020; Hu et al. 2021; Endsley et al. 2021b) suggest high Ly α visibility from these bright galaxies, implying inhomogeneous reionization caused by individual/groups of bright galaxies. This picture is further supported by Ly α detections from two of the brightest $z > 8.5$ galaxies (Zitrin et al. 2015; Naidu et al. 2020, Larson et al. 2021 in preparation), which implies that reionization proceeds more rapidly and/or earlier around brighter galaxies.

Our analysis of the Ly α transmission at $z > 6$ as a function UV magnitude shows a high Ly α transmission of the IGM in UV bright galaxies ($M_{UV} < -21$), while Ly α visibility from fainter galaxies ($-21 < M_{UV} < -19$) is rapidly suppressed at $z > 6$. This is in general consistent with a picture of the spatially inhomogeneous process of reionization where bright galaxies often created large ionized bubbles which allow enhanced Ly α transmission to the IGM nearby these bright sources, which is illustrated in Figure 5.

This picture is consistent with theoretical predictions from reionization simulations where reionization begins in overdense regions and progresses into low-density regions (e.g., Finlator et al. 2009; Katz et al. 2019). In particular, Park et al. (2021) calculate Ly α transmissivity of the IGM from the Cosmic Dawn II simulation (Ocvirk et al. 2020), showing that UV-bright galaxies tend to reside in “bubbles” as they ionize large volumes and are located in overdense regions where nearby fainter galaxies also can contribute to the ionizing emissivity (Finkelstein et al. 2019b; Endsley et al. 2021a, Larson et al. 2021 in preparation). These fainter galaxies sit in large ionized bubbles are expected to have higher Ly α transmission than isolated faint objects (Qin et al. 2021). Most of these faint objects are likely under current detection limit, but could be probed in larger samples (which

will be available from Nancy Grace Roman Space Telescope, NGRST) or tested in future observations sensitive to fainter emission from James Webb Space Telescope (JWST) and the next generation of 20-30 m-class telescopes. This model also naturally predicts that Ly α emission shows a larger velocity offset from systemic for brighter galaxies, which further facilitates the escape of Ly α photons through the IGM (Mason et al. 2018b; Garel et al. 2021). The existing data from Ly α supports this picture.

Our results suggest significantly different evolution of the Ly α transmission in the IGM in the middle phase of reionization in varying UV-magnitude galaxies. As the Ly α transmission to the IGM is the key quantity in constraining the neutral fraction of the IGM with Ly α observations (Mason et al. 2018a; Hoag et al. 2019; Jung et al. 2020; Morales et al. 2021; Wold et al. 2021), it is critical to understand the dependence of IGM transmission of Ly α on galaxy UV luminosity in the future. This requires deeper spectroscopic observations and larger sample sizes, both of which will need to cover a wide dynamic range in galaxy UV luminosity. This will be possible by grism observations with NGRST, which will be ideal for such studies of Ly α emission from galaxies during the epoch of reionization. Furthermore, the pending launch of JWST will provide data we can use to study Ly α systematics (e.g., velocity offset) with other emission line observations for galaxies in the reionization era.

We thank our colleagues on the CLEAR team for their valuable conversations and contributions. This work is based on data obtained from the Hubble Space Telescope through program number GO-14227. Support for Program number GO-14227 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555. I.J. acknowledges support from NASA under award number 80GSFC21M0002. SLF acknowledges support from NASA through ADAP award 80NSSC18K0954. VEC acknowledges support from the NASA Headquarters under the Future Investigators in NASA Earth and Space Science and Technology (FINESST) award 19-ASTRO19-0122, as well as support from the Hagler Institute for Advanced Study at Texas A&M University. This work benefited from generous support from the George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy at Texas A&M University. This work was supported in part by NASA contract NNG16PJ33C, the Studying Cosmic Dawn with WFIRST Science Investigation Team.

Appendix

A. Ly α -Emitter Candidate at $z = 6.513$

We performed emission-line search as described in Section 2.2.1, which results in no convincing Ly α detections from the entire sample except for one tentative detection from a $z_{\text{grism}} = 6.513 \pm 0.005$ galaxy (GN4_5461276). Although the detection of Ly α emission is possibly spurious from the galaxy with given data, it is worth presenting the promising features of the detected emission line here as the physical properties of the emission line would be rather extreme if confirmed. Further observations are required for validating the nature of the detected emission line.

We display the individual 2D spectra obtained from 4 PAs with the HST H -band stamp of GN4_5461276 in Figure 6 where we detect significant emission-line features from 2 (PA=93 and PA=173) of the 4 PAs. The all PA combined 1D and 2D spectra are shown in Figure 7, which shows a formal $>4\sigma$ detection of the emission line.




Figure 6. (Left) Four individual PA 2D spectra of the emission-line candidate galaxy (GN4_5461276). White (black) shows regions with positive (negative) flux in the spectra. The topmost and bottom-most panels (PA = 93 and 173) show positive emission features. These are marked with yellow arrows. The middle two panels (PA=133 and 127) show no such emission features. (Right) HST H -band image centered at GN4_5461276 showing the spectral dispersion directions of the four PAs (as labeled). The two PAs (93 and 173) showing positive emission do not share any potential contaminating sources. The two intermediate PAs (127 and 133) have possible contamination from the large galaxy in the bottom right of the image.

To rule out the possibility of being contaminated emission from nearby sources, we checked the known redshifts of the nearby bright sources and concluded that the emission line is not the zeroth-order reflection of an emission line associated with the nearby sources. We fit the Gaussian function to the emission line to estimate the physical properties of the detected emission line. The S/N value was estimated from Gaussian fitting on 1000 simulated spectra by fluctuating the 1D fluxes with the 1D noise level. The emission line flux was calculated from the best-fit Gaussian (red in Figure 7), and the redshift was derived from the peak of the best-fit Gaussian curve. The derived physical quantities are listed in the bottom panel of the plot, which shows its extreme values of a rest-frame EW ($>1000\text{\AA}$) and a line width (FWHM $>100\text{\AA}$).

To check the possibility of being a low- z interloper, we compare the best-fit SED models between high- z (Ly α) and low- z ([O II]) interpretations with the HST stamp images in Figure 8. For the high- z interpretation, the Ly α contribution to HST fluxes were subtracted for the SED fitting. From the best-fit models, the high- z solution is preferred than the low- z solution, but both interpretations remain viable with the given data. Thus, further observational evidence is required to confirm this tentative Ly α detection.

B. Continuum Detection from a $z = 6.2$ Galaxy

We inspected CLEAR spectra of high- z candidates to see if they present any continuum features as described in Section 2.2.2. It results in one continuum candidate from a $z_{\text{grism}} = 6.2^{+2.5}_{-0.8}$ galaxy. The continuum is detected at $>4\sigma$ at wavelengths redward of a possible continuum break. Figure 9 shows the grism spectra of GN2_23734, combining both G102 and G140 spectra. The extracted 1D spectrum is shown as a black histogram with the noise level shown as dashed-line curves. The continuum fit given




Figure 7. All PA combined 1D (bottom) and 2D (top) spectrum of the emission line detected from GN4_5461276. In the bottom panel, the measured spectrum is shown as the black histogram with the noise level showed as the dashed curves. The red curve represents the a model Gaussian curve fit to the the emission line.





Figure 8. Images and SED models for GN4_5461276. (Top) The HST ACS/WFC3 “postage-stamp images” for this galaxy. The yellow arrows indicate the locations of the object in individual stamps. (Bottom) The best-fit SED models between high- z (red: Ly α) and low- z (blue: [O II]). The faint blue and red curves represent the SED fitting for each 100 MC realization set of the high- z and low- z cases. For the high- z solution, Ly α contribution is subtracted from HST photometry. The inset figure shows the probability distribution function of the photometric redshift (solid curve) with the grism redshift (vertical dashed line). Although the high- z solution ($\chi^2 = 1.7$) is slightly preferred compared to the low- z ($\chi^2 = 3.1$), it is difficult to draw conclusion with given data.

by GRIZLI is presented as a red curve, and we display the best-fit SED (blue), which is obtained from the SED fitting based on the object’s HST+Spitzer/IRAC photometry (see Section 2.4). In the SED fitting, we used the best-fit grism redshift ($z_{\text{grism}} = 6.2$) derived by GRIZLI. Although the grism continuum-fit spectrum seems comparable to the best-fit SED, we consider it tentative

given the current noise level in the grism spectrum. Nevertheless, we report this tentative detection of the continuum break as it provides a possible path to confirm other high- z galaxies without $\text{Ly}\alpha$. This is particularly useful for galaxies in the epoch of reionization where the $\text{Ly}\alpha$ transmission is lower (see Section 2.2.2 and discussion in, e.g., Rhoads et al. 2013; Watson et al. 2015; Oesch et al. 2016). It will be valuable to target this galaxy (and others) with spectroscopy from future space-based telescopes (e.g., JWST, NGRST).




Figure 9. The 1D grism spectrum of a continuum-detection candidate galaxy (GN2_23734) at $z_{\text{grism}} = 6.2^{+2.5}_{-0.8}$. The continuum is formally detected at wavelengths redward of the $\text{Ly}\alpha$ -break ($S/N_{\text{continuum}} > 4$). The black histogram shows the 1D spectrum, which combines both of the G102 and G140 spectra of GN2_23734. The noise level is shown as dashed curves. The red curve represents the continuum fit derived by GRIZLI, and the best-fit SED is displayed as the blue curve, which is obtained from SED fitting based on the object’s HST+Spitzer/IRAC photometry.

C. Summary of $6.0 < z < 8.2$ Candidate Galaxies in CLEAR

As discussed in Section 2, we use the CLEAR spectra of 148 $6.0 < z < 8.2$ candidate galaxies for exploring the galaxy UV magnitude dependency of the IGM transmission to $\text{Ly}\alpha$ during the epoch of reionization. Table 2 summarizes the entire 148 $6.0 < z < 8.2$ CLEAR galaxies in order of increasing photometric redshift, which includes the 3σ rest-EW upper limits on $\text{Ly}\alpha$ in the last column.

Table 2. Summary of CLEAR $6.0 < z < 8.2$ Targets

ID ^a	R.A. (J2000.0)	Decl. (J2000.0)	J_{125}	M_{UV}^b	z_{phot}^c	$\text{EW}_{\text{Ly}\alpha}^d$ (Å)
GS4_28545	53.142085	-27.779851	$26.12^{+0.01}_{-0.01}$	$-20.63^{+0.02}_{-0.03}$	$6.00^{+0.02}_{-0.02}$	<82.2
GS4_5428651	53.162267	-27.796050	$29.90^{+0.17}_{-0.15}$	$-16.03^{+0.21}_{-0.13}$	$6.00^{+5.33}_{-0.85}$	<1332.7
GS4_30518	53.159459	-27.771855	$28.59^{+0.06}_{-0.06}$	$-18.29^{+0.05}_{-0.02}$	$6.01^{+0.08}_{-0.08}$	<142.8
GN5_5453006	189.147675	62.310692	$26.18^{+0.10}_{-0.09}$	$-21.41^{+0.03}_{-0.09}$	$6.02^{+0.09}_{-0.08}$	<14.0
GS4_25097	53.143880	-27.793083	$27.85^{+0.02}_{-0.02}$	$-18.97^{+0.01}_{-0.02}$	$6.03^{+0.07}_{-0.07}$	<123.9
GN1_38168	189.320618	62.377956	$25.29^{+0.29}_{-0.23}$	$-22.97^{+0.08}_{-0.03}$	$6.03^{+0.04}_{-0.05}$	<3.3
GS2_49117	53.128395	-27.679525	$26.10^{+0.10}_{-0.09}$	$-20.41^{+0.08}_{-0.13}$	$6.05^{+0.31}_{-0.40}$	<55.2

Table 2 continued

Table 2 (continued)

ID ^a	R.A. (J2000.0)	Decl. (J2000.0)	J_{125}	M_{UV} ^b	z_{phot} ^c	EW _{Lyα} ^d (Å)
GN2_22377	189.231018	62.252335	26.71 ^{+0.07} _{-0.06}	-20.03 ^{+0.07} _{-0.07}	6.06 ^{+0.10} _{-0.09}	<49.0
GS4_24418	53.139299	-27.795872	26.85 ^{+0.15} _{-0.13}	-19.88 ^{+0.05} _{-0.05}	6.07 ^{+0.09} _{-0.09}	<53.9
ERSPRIME_43483	53.070813	-27.706732	26.07 ^{+0.06} _{-0.06}	-20.66 ^{+0.05} _{-0.05}	6.07 ^{+0.09} _{-0.09}	<36.0
GS4_29439	53.165805	-27.776102	28.55 ^{+0.08} _{-0.08}	-18.23 ^{+0.05} _{-0.07}	6.09 ^{+0.08} _{-0.08}	<155.4
GS4_29085	53.151986	-27.778248	25.44 ^{+0.00} _{-0.00}	-21.36 ^{+0.00} _{-0.01}	6.10 ^{+0.04} _{-0.05}	<8.7
GN1_37875	189.292984	62.366493	25.96 ^{+0.26} _{-0.21}	-20.30 ^{+0.15} _{-0.17}	6.12 ^{+0.24} _{-0.26}	<54.7
GN4_5463603	189.346069	62.255791	27.21 ^{+0.09} _{-0.08}	-19.64 ^{+0.08} _{-0.07}	6.13 ^{+0.25} _{-0.82}	<279.9
GS2_49517	53.144795	-27.676513	26.54 ^{+0.17} _{-0.15}	-20.24 ^{+0.14} _{-0.18}	6.14 ^{+0.27} _{-0.29}	<45.9
GS5_40046	53.105382	-27.723476	25.56 ^{+0.07} _{-0.06}	-21.10 ^{+0.07} _{-0.09}	6.15 ^{+0.19} _{-0.27}	<33.0
GN5_5453623	189.159882	62.307541	25.76 ^{+0.05} _{-0.04}	-20.43 ^{+0.09} _{-0.06}	6.15 ^{+0.27} _{-0.30}	<75.6
GS4_24191	53.152647	-27.797311	26.81 ^{+0.00} _{-0.00}	-20.01 ^{+0.05} _{-0.00}	6.15 ^{+0.04} _{-0.03}	<48.8
GS4_24855	53.146101	-27.794571	26.77 ^{+0.00} _{-0.00}	-19.87 ^{+0.01} _{-0.02}	6.17 ^{+0.06} _{-0.07}	<72.0
GS3_5443046	53.155751	-27.746295	27.31 ^{+0.14} _{-0.13}	-19.39 ^{+0.09} _{-0.09}	6.17 ^{+0.18} _{-0.19}	<103.7
GS2_48937	53.123742	-27.680620	27.48 ^{+0.22} _{-0.18}	-19.29 ^{+0.10} _{-0.20}	6.17 ^{+0.29} _{-0.40}	<89.3
GS3_35639	53.155782	-27.746150	26.61 ^{+0.14} _{-0.13}	-20.28 ^{+0.09} _{-0.08}	6.17 ^{+0.11} _{-0.11}	<39.6
GS5_44454	53.120956	-27.702285	26.19 ^{+0.13} _{-0.12}	-20.61 ^{+0.14} _{-0.11}	6.18 ^{+0.17} _{-0.22}	<26.4
ERSPRIME_43514	53.071191	-27.706498	26.30 ^{+0.09} _{-0.08}	-20.40 ^{+0.07} _{-0.07}	6.18 ^{+0.13} _{-0.13}	<45.3
GN3_30188	189.256210	62.291283	27.02 ^{+0.05} _{-0.05}	-19.67 ^{+0.06} _{-0.04}	6.20 ^{+0.09} _{-0.10}	<242.8
GN7_5429834	189.095474	62.227612	27.77 ^{+0.23} _{-0.19}	-19.14 ^{+0.17} _{-0.13}	6.20 ^{+0.26} _{-0.63}	<112.1
GN3_5450671	189.263977	62.318367	27.64 ^{+0.17} _{-0.15}	-19.49 ^{+0.08} _{-0.07}	6.21 ^{+0.10} _{-0.09}	<94.7
GN4_5461825	189.323929	62.271130	28.16 ^{+0.12} _{-0.11}	-18.76 ^{+0.09} _{-0.08}	6.22 ^{+0.20} _{-0.22}	<174.3
GN4_5446203	189.305573	62.271095	27.96 ^{+0.22} _{-0.18}	-17.87 ^{+0.88} _{-0.49}	6.22 ^{+2.25} _{-4.08}	<14672.0
GN4_23534	189.296127	62.257969	26.88 ^{+0.13} _{-0.12}	-19.98 ^{+0.14} _{-0.10}	6.23 ^{+0.21} _{-0.33}	<47.7
GN2_5432549	189.252869	62.235577	27.63 ^{+0.09} _{-0.08}	-19.33 ^{+0.06} _{-0.05}	6.24 ^{+0.11} _{-0.12}	<128.9
ERSPRIME_5455619	53.069652	-27.697069	25.63 ^{+0.02} _{-0.02}	-21.06 ^{+0.02} _{-0.02}	6.24 ^{+0.13} _{-0.13}	<67.6
GN5_5453421	189.157303	62.309128	27.85 ^{+0.17} _{-0.15}	-19.07 ^{+0.16} _{-0.11}	6.24 ^{+0.33} _{-0.88}	<146.2
GN4_21677	189.318161	62.249264	26.91 ^{+0.10} _{-0.09}	-19.95 ^{+0.10} _{-0.08}	6.28 ^{+0.15} _{-0.15}	<61.5
GN7_16571	189.114838	62.224430	25.73 ^{+0.04} _{-0.04}	-21.08 ^{+0.05} _{-0.05}	6.30 ^{+0.06} _{-0.07}	<18.0
GS5_5445164	53.114526	-27.737678	26.95 ^{+0.17} _{-0.15}	-19.81 ^{+0.18} _{-0.16}	6.30 ^{+0.32} _{-4.74}	<87.2
GN2_23734	189.211990	62.258846	26.24 ^{+0.06} _{-0.06}	-20.43 ^{+0.04} _{-0.04}	6.31 ^{+0.18} _{-0.21}	<67.4
GS4_29685	53.156131	-27.775849	26.42 ^{+0.00} _{-0.00}	-20.22 ^{+0.01} _{-0.01}	6.32 ^{+0.06} _{-0.07}	<38.0
GN2_17132	189.197571	62.226990	26.30 ^{+0.04} _{-0.04}	-20.48 ^{+0.03} _{-0.03}	6.32 ^{+0.08} _{-0.08}	<34.7
GN2_19472	189.219406	62.238247	27.70 ^{+0.08} _{-0.08}	-18.97 ^{+0.08} _{-0.07}	6.32 ^{+0.20} _{-0.21}	<174.7
GS2_47924	53.146726	-27.686787	26.96 ^{+0.08} _{-0.07}	-19.86 ^{+0.10} _{-0.12}	6.33 ^{+0.30} _{-0.29}	<68.9
GS5_40767	53.118464	-27.719486	26.36 ^{+0.11} _{-0.10}	-20.43 ^{+0.10} _{-0.12}	6.34 ^{+0.24} _{-0.23}	<35.4
GS4_28201	53.156040	-27.780970	27.86 ^{+0.02} _{-0.02}	-19.04 ^{+0.01} _{-0.01}	6.34 ^{+0.04} _{-0.56}	<71.6
GN3_35300	189.238388	62.327126	25.51 ^{+0.08} _{-0.08}	-21.26 ^{+0.12} _{-0.09}	6.35 ^{+0.19} _{-0.19}	<22.4
GS4_23893	53.153257	-27.798288	27.69 ^{+0.01} _{-0.01}	-18.87 ^{+0.02} _{-0.01}	6.38 ^{+0.07} _{-0.06}	<180.2
GN1_37866	189.323151	62.366215	25.99 ^{+0.16} _{-0.14}	-20.11 ^{+0.10} _{-0.13}	6.39 ^{+0.57} _{-1.08}	<76.4
GS3_34282	53.149862	-27.752839	27.25 ^{+0.29} _{-0.23}	-19.82 ^{+0.15} _{-0.12}	6.42 ^{+0.24} _{-0.64}	<224.2
GN1_37461	189.318008	62.353180	26.50 ^{+0.12} _{-0.11}	-20.22 ^{+0.09} _{-0.08}	6.43 ^{+0.14} _{-0.30}	<65.5
GS4_27209	53.173343	-27.784645	28.05 ^{+0.03} _{-0.03}	-18.90 ^{+0.01} _{-0.02}	6.44 ^{+0.06} _{-0.04}	<88.9

Table 2 continued

Table 2 (*continued*)

ID ^a	R.A. (J2000.0)	Decl. (J2000.0)	J_{125}	M_{UV} ^b	z_{phot} ^c	$EW_{\text{Ly}\alpha}$ ^d (Å)
ERSPRIME_44510	53.049653	-27.701990	$26.85^{+0.05}_{-0.04}$	$-19.86^{+0.04}_{-0.07}$	$6.45^{+0.07}_{-0.08}$	<98.9
GN4_27400	189.277420	62.276405	$25.92^{+0.04}_{-0.04}$	$-20.95^{+0.04}_{-0.04}$	$6.45^{+0.09}_{-0.10}$	<26.0
GS2_5457820	53.114407	-27.685098	$26.91^{+0.16}_{-0.14}$	$-19.31^{+0.09}_{-0.13}$	$6.47^{+0.47}_{-0.49}$	<360.3
GS4_5430868	53.155602	-27.788733	$29.47^{+0.05}_{-0.05}$	$-17.48^{+0.08}_{-0.03}$	$6.48^{+0.13}_{-0.12}$	<355.1
GN2_20362	189.240814	62.242630	$26.91^{+0.06}_{-0.05}$	$-20.04^{+0.05}_{-0.04}$	$6.52^{+0.08}_{-0.08}$	<53.5
GS3_40377	53.172480	-27.721434	$26.42^{+0.08}_{-0.07}$	$-20.52^{+0.05}_{-0.07}$	$6.54^{+0.13}_{-0.14}$	<38.0
ERSPRIME_39697	53.064237	-27.724698	$27.11^{+0.10}_{-0.09}$	$-19.77^{+0.08}_{-0.10}$	$6.55^{+0.14}_{-0.15}$	<83.6
GN5_33256	189.178146	62.310635	$27.41^{+0.06}_{-0.06}$	$-19.36^{+0.08}_{-0.06}$	$6.57^{+0.31}_{-0.33}$	<128.3
GN7_16422	189.177979	62.223713	$26.32^{+0.05}_{-0.05}$	$-20.49^{+0.07}_{-0.05}$	$6.58^{+0.11}_{-0.12}$	<71.4
GN7_14851	189.111633	62.215374	$26.54^{+0.03}_{-0.03}$	$-20.33^{+0.03}_{-0.04}$	$6.58^{+0.06}_{-0.07}$	<43.4
GN4_5438687	189.295410	62.252560	$28.66^{+0.28}_{-0.22}$	$-18.25^{+0.24}_{-0.13}$	$6.59^{+0.52}_{-5.20}$	<330.2
GS4_28784	53.169042	-27.778832	$28.06^{+0.03}_{-0.03}$	$-18.79^{+0.02}_{-0.03}$	$6.61^{+0.06}_{-0.06}$	<129.3
GN5_34340	189.187347	62.318821	$27.27^{+0.08}_{-0.07}$	$-19.71^{+0.06}_{-0.07}$	$6.63^{+0.18}_{-0.17}$	<66.0
GN2_5465437	189.239700	62.248108	$26.35^{+0.01}_{-0.01}$	$-20.59^{+0.01}_{-0.01}$	$6.65^{+0.04}_{-0.05}$	<39.9
GS4_5431395	53.161653	-27.787043	$27.97^{+0.05}_{-0.05}$	$-18.88^{+0.07}_{-0.03}$	$6.67^{+0.13}_{-0.14}$	<128.6
ERSPRIME_43078	53.055896	-27.708708	$26.74^{+0.09}_{-0.08}$	$-20.27^{+0.06}_{-0.06}$	$6.68^{+0.12}_{-0.13}$	<65.6
GS3_5447463	53.167998	-27.728187	$28.31^{+0.24}_{-0.20}$	$-18.34^{+0.17}_{-0.14}$	$6.70^{+0.63}_{-0.79}$	<689.3
GN4_5462173	189.308792	62.267353	$26.66^{+0.05}_{-0.05}$	$-20.12^{+0.04}_{-0.06}$	$6.71^{+0.25}_{-0.34}$	<125.9
GN7_5430295	189.088989	62.229202	$28.07^{+0.15}_{-0.13}$	$-18.96^{+0.11}_{-0.09}$	$6.72^{+0.25}_{-0.26}$	<278.5
GN5_31683	189.165039	62.300194	$26.35^{+0.01}_{-0.01}$	$-20.59^{+0.04}_{-0.02}$	$6.73^{+0.05}_{-0.05}$	<35.5
GN5_32053	189.140656	62.302368	$27.40^{+0.15}_{-0.13}$	$-19.47^{+0.09}_{-0.14}$	$6.73^{+0.46}_{-0.57}$	<157.3
GN5_33584	189.156357	62.313087	$27.11^{+0.08}_{-0.07}$	$-19.67^{+0.08}_{-0.08}$	$6.75^{+0.50}_{-0.45}$	<112.4
GN7_5428106	189.104630	62.222271	$27.49^{+0.14}_{-0.12}$	$-19.59^{+0.10}_{-0.08}$	$6.75^{+0.18}_{-0.20}$	<69.1
GS4_26311	53.151601	-27.787910	$28.07^{+0.04}_{-0.03}$	$-18.91^{+0.02}_{-0.03}$	$6.76^{+0.07}_{-0.07}$	<123.5
GN5_5454319	189.140610	62.305511	$26.14^{+0.17}_{-0.14}$	$-20.73^{+0.10}_{-0.13}$	$6.77^{+0.32}_{-0.40}$	<46.7
GS4_20530	53.154950	-27.815805	$25.89^{+0.08}_{-0.07}$	$-20.86^{+0.04}_{-0.11}$	$6.79^{+0.07}_{-0.07}$	<35.6
GN4_5461276 ^e	189.281143	62.274857	$28.10^{+0.17}_{-0.15}$	$-18.01^{+0.24}_{-0.26}$	$6.80^{+0.56}_{-0.60}$	-
ERSPRIME_45005	53.066739	-27.699829	$26.09^{+0.18}_{-0.16}$	$-20.86^{+0.13}_{-0.17}$	$6.81^{+0.30}_{-0.49}$	<29.6
GN4_26634	189.282623	62.272507	$27.01^{+0.08}_{-0.07}$	$-19.90^{+0.05}_{-0.09}$	$6.82^{+0.24}_{-0.26}$	<89.3
GS5_41330	53.094447	-27.716946	$25.53^{+0.10}_{-0.09}$	$-21.37^{+0.09}_{-0.10}$	$6.82^{+0.16}_{-0.16}$	<16.8
ERSPRIME_39962	53.041104	-27.723434	$26.60^{+0.14}_{-0.12}$	$-20.43^{+0.10}_{-0.08}$	$6.83^{+0.12}_{-0.16}$	<55.2
GN3_5452837	189.220932	62.311447	$27.84^{+0.12}_{-0.11}$	$-19.16^{+0.06}_{-0.11}$	$6.83^{+0.25}_{-0.25}$	<154.1
GN5_34059	189.166245	62.316494	$26.36^{+0.05}_{-0.05}$	$-20.57^{+0.04}_{-0.05}$	$6.87^{+0.15}_{-0.14}$	<40.7
GS3_35821	53.150039	-27.745016	$26.18^{+0.10}_{-0.09}$	$-20.77^{+0.07}_{-0.08}$	$6.88^{+0.18}_{-0.21}$	<27.4
GS4_20622	53.155370	-27.815248	$25.01^{+0.03}_{-0.03}$	$-21.88^{+0.01}_{-0.03}$	$6.88^{+0.04}_{-0.03}$	<14.0
GN2_5430417	189.223526	62.229626	$27.77^{+0.14}_{-0.12}$	$-19.25^{+0.09}_{-0.10}$	$6.89^{+0.20}_{-0.22}$	<126.7
GN5_32855	189.134445	62.307865	$26.36^{+0.06}_{-0.06}$	$-20.52^{+0.05}_{-0.07}$	$6.90^{+0.21}_{-0.19}$	<57.7
GS4_25335	53.177375	-27.792132	$27.17^{+0.03}_{-0.03}$	$-19.70^{+0.02}_{-0.02}$	$6.90^{+0.09}_{-0.09}$	<77.8
GN7_15746	189.095871	62.220078	$26.14^{+0.03}_{-0.03}$	$-20.72^{+0.03}_{-0.03}$	$6.91^{+0.16}_{-0.15}$	<42.0
GS2_5452228	53.136772	-27.710775	$26.41^{+0.08}_{-0.08}$	$-20.48^{+0.07}_{-0.08}$	$6.91^{+0.15}_{-0.15}$	<48.0
GS4_22623	53.158407	-27.804372	$28.69^{+0.08}_{-0.07}$	$-18.04^{+0.06}_{-0.03}$	$6.91^{+0.21}_{-0.20}$	<330.6
GN4_24681	189.353012	62.263409	$25.66^{+0.02}_{-0.02}$	$-21.40^{+0.02}_{-0.02}$	$6.94^{+0.05}_{-0.06}$	<43.0

Table 2 continued

Table 2 (continued)

ID ^a	R.A. (J2000.0)	Decl. (J2000.0)	J_{125}	M_{UV}^b	z_{phot}^c	$EW_{Ly\alpha}^d$ (Å)
GN5_35003	189.178085	62.324314	$26.84^{+0.06}_{-0.05}$	$-20.15^{+0.06}_{-0.05}$	$6.94^{+0.16}_{-0.16}$	<68.1
GN3_5455960	189.279175	62.299282	$27.40^{+0.10}_{-0.09}$	$-19.64^{+0.10}_{-0.09}$	$6.95^{+0.26}_{-0.30}$	<173.9
GS4_23998	53.160569	-27.797819	$28.99^{+0.04}_{-0.04}$	$-17.73^{+0.02}_{-0.01}$	$6.97^{+0.07}_{-0.08}$	<306.7
GN3_5455082	189.229721	62.302711	$28.16^{+0.11}_{-0.10}$	$-18.91^{+0.07}_{-0.08}$	$6.97^{+0.31}_{-0.37}$	<291.4
GN7_11318	189.116867	62.198677	$26.66^{+0.25}_{-0.20}$	$-20.23^{+0.21}_{-0.15}$	$6.98^{+0.36}_{-0.37}$	<175.6
ERSPRIME_38389	53.066731	-27.731170	$27.51^{+0.08}_{-0.07}$	$-19.48^{+0.06}_{-0.05}$	$6.98^{+0.21}_{-0.28}$	<343.1
GN5_33361	189.177856	62.311707	$26.40^{+0.06}_{-0.06}$	$-20.59^{+0.04}_{-0.05}$	$6.99^{+0.20}_{-0.21}$	<35.5
GN4_26575	189.283737	62.272240	$26.90^{+0.04}_{-0.04}$	$-19.89^{+0.04}_{-0.09}$	$6.99^{+0.31}_{-0.31}$	<166.5
GN2_24019	189.275925	62.260296	$27.14^{+0.05}_{-0.05}$	$-19.85^{+0.04}_{-0.04}$	$7.01^{+0.18}_{-0.18}$	<262.4
GN2_17220	189.201050	62.227440	$27.08^{+0.07}_{-0.07}$	$-19.87^{+0.06}_{-0.04}$	$7.02^{+0.21}_{-0.21}$	<77.4
GN7_5428506	189.150253	62.223709	$28.01^{+0.22}_{-0.19}$	$-19.26^{+0.15}_{-0.07}$	$7.03^{+0.27}_{-0.43}$	<133.2
GN5_34042	189.166092	62.316406	$27.75^{+0.17}_{-0.15}$	$-19.13^{+0.14}_{-0.20}$	$7.03^{+0.60}_{-5.54}$	<232.7
GN4_5441249	189.310867	62.260452	$27.20^{+0.03}_{-0.03}$	$-19.76^{+0.03}_{-0.02}$	$7.04^{+0.14}_{-0.13}$	<98.6
GN1_37724	189.273300	62.360783	$25.80^{+0.18}_{-0.15}$	$-21.18^{+0.12}_{-0.08}$	$7.05^{+0.21}_{-0.21}$	<25.4
GN5_31436	189.166931	62.298553	$28.24^{+0.17}_{-0.15}$	$-19.03^{+0.10}_{-0.10}$	$7.06^{+0.25}_{-5.53}$	<261.7
GN3_34711	189.254974	62.321877	$27.16^{+0.20}_{-0.17}$	$-19.82^{+0.14}_{-0.17}$	$7.06^{+0.34}_{-0.34}$	<113.6
GN2_5464240	189.232895	62.247395	$28.68^{+0.20}_{-0.17}$	$-18.45^{+0.08}_{-0.12}$	$7.06^{+0.36}_{-5.52}$	<399.1
GN2_20956	189.205231	62.245541	$27.64^{+0.08}_{-0.07}$	$-19.20^{+0.04}_{-0.08}$	$7.07^{+0.31}_{-0.32}$	<213.9
GN7_12243	189.114151	62.203175	$25.47^{+0.05}_{-0.05}$	$-21.49^{+0.08}_{-0.07}$	$7.08^{+0.17}_{-0.20}$	<26.2
GS4_5435288	53.153420	-27.774471	$27.30^{+0.22}_{-0.18}$	$-19.62^{+0.06}_{-0.18}$	$7.11^{+0.33}_{-5.53}$	<111.9
GN7_16103	189.091812	62.221901	$27.32^{+0.06}_{-0.06}$	$-19.73^{+0.05}_{-0.06}$	$7.14^{+0.26}_{-0.25}$	<103.2
GN4_5463508	189.307037	62.255745	$28.26^{+0.09}_{-0.09}$	$-18.82^{+0.06}_{-0.08}$	$7.18^{+0.29}_{-0.27}$	<282.2
GN3_5449006	189.286926	62.277760	$27.42^{+0.06}_{-0.05}$	$-19.59^{+0.04}_{-0.05}$	$7.19^{+0.17}_{-0.18}$	<299.1
GN3_5454301	189.225098	62.306057	$28.58^{+0.31}_{-0.24}$	$-18.51^{+0.17}_{-0.21}$	$7.19^{+0.53}_{-5.35}$	<442.2
GN2_21995	189.205307	62.250763	$26.44^{+0.06}_{-0.06}$	$-20.59^{+0.05}_{-0.04}$	$7.22^{+0.18}_{-0.19}$	<39.5
GN4_25192	189.264938	62.265793	$26.06^{+0.02}_{-0.02}$	$-20.95^{+0.02}_{-0.02}$	$7.22^{+0.14}_{-0.14}$	<41.5
GN1_37619	189.278732	62.357456	$26.21^{+0.16}_{-0.14}$	$-20.99^{+0.08}_{-0.06}$	$7.23^{+0.13}_{-0.13}$	<37.7
GN7_5429974	189.091492	62.228096	$27.87^{+0.12}_{-0.11}$	$-19.13^{+0.10}_{-0.09}$	$7.28^{+0.61}_{-0.98}$	<305.1
GS4_5427001	53.152865	-27.801940	$29.31^{+0.16}_{-0.14}$	$-17.40^{+0.09}_{-0.09}$	$7.30^{+0.52}_{-0.57}$	<848.5
GS3_37079	53.169588	-27.738066	$26.76^{+0.16}_{-0.14}$	$-20.30^{+0.11}_{-0.11}$	$7.32^{+0.32}_{-0.35}$	<55.0
GN3_35055	189.273392	62.324780	$26.15^{+0.31}_{-0.24}$	$-20.90^{+0.17}_{-0.13}$	$7.35^{+0.29}_{-0.30}$	<36.0
GN5_32031	189.157898	62.302376	$25.54^{+0.02}_{-0.02}$	$-21.52^{+0.02}_{-0.02}$	$7.37^{+0.15}_{-0.15}$	<27.8
GN2_21790	189.199692	62.249802	$27.55^{+0.06}_{-0.06}$	$-19.48^{+0.05}_{-0.04}$	$7.37^{+0.26}_{-0.25}$	<113.0
GN3_33949	189.222000	62.315758	$26.98^{+0.07}_{-0.06}$	$-19.99^{+0.06}_{-0.05}$	$7.39^{+0.19}_{-0.18}$	<79.0
GN5_33655	189.133728	62.313564	$26.59^{+0.10}_{-0.09}$	$-20.52^{+0.07}_{-0.07}$	$7.43^{+0.21}_{-0.22}$	<64.3
GS4_23143	53.155096	-27.801771	$27.83^{+0.01}_{-0.01}$	$-19.22^{+0.00}_{-0.00}$	$7.44^{+0.14}_{-0.14}$	<154.4
GS4_27055	53.161714	-27.785390	$26.80^{+0.03}_{-0.03}$	$-20.34^{+0.01}_{-0.02}$	$7.44^{+0.10}_{-0.09}$	<38.4
GS4_27958	53.138064	-27.781866	$29.04^{+0.05}_{-0.05}$	$-18.27^{+0.02}_{-0.02}$	$7.48^{+0.23}_{-0.24}$	<846.2
GN4_23416	189.333069	62.257233	$25.91^{+0.02}_{-0.02}$	$-21.19^{+0.01}_{-0.02}$	$7.54^{+0.24}_{-0.26}$	<66.7
GS3_36060	53.172602	-27.743931	$25.96^{+0.07}_{-0.06}$	$-21.11^{+0.04}_{-0.05}$	$7.54^{+0.39}_{-0.41}$	<43.6
GN2_20084	189.249817	62.241226	$26.32^{+0.03}_{-0.03}$	$-20.82^{+0.07}_{-0.03}$	$7.58^{+0.15}_{-0.15}$	<39.7
GS4_5432952	53.170708	-27.782452	$29.12^{+0.09}_{-0.08}$	$-17.68^{+0.05}_{-0.10}$	$7.63^{+0.23}_{-5.84}$	<916.7

Table 2 continued

Table 2 (*continued*)

ID ^a	R.A. (J2000.0)	Decl. (J2000.0)	J_{125}	M_{UV} ^b	z_{phot} ^c	EW _{Lyα} ^d (Å)
GN7_11906	189.117416	62.201412	26.47 ^{+0.27} _{-0.21}	-20.61 ^{+0.23} _{-0.13}	7.65 ^{+0.65} _{-0.70}	<42.2
GN7_16124	189.084152	62.222023	25.79 ^{+0.03} _{-0.03}	-21.37 ^{+0.04} _{-0.03}	7.68 ^{+0.37} _{-0.34}	<34.2
GN2_5436513	189.245880	62.244953	28.04 ^{+0.17} _{-0.14}	-19.07 ^{+0.12} _{-0.11}	7.69 ^{+0.36} _{-0.64}	<230.4
GN2_23331	189.197769	62.256966	27.67 ^{+0.12} _{-0.10}	-19.52 ^{+0.08} _{-0.08}	7.71 ^{+0.32} _{-0.45}	<274.8
GN2_20338	189.203125	62.242481	26.48 ^{+0.07} _{-0.06}	-20.66 ^{+0.04} _{-0.06}	7.72 ^{+0.20} _{-0.21}	<42.2
GN5_36116	189.161255	62.334435	25.73 ^{+0.05} _{-0.05}	-21.47 ^{+0.04} _{-0.04}	7.75 ^{+0.54} _{-0.52}	<44.5
GN4_28143	189.300125	62.280354	27.23 ^{+0.21} _{-0.18}	-19.90 ^{+0.14} _{-0.13}	7.75 ^{+0.48} _{-0.78}	<99.6
GN4_22848	189.336029	62.254650	27.26 ^{+0.08} _{-0.07}	-19.71 ^{+0.07} _{-0.09}	7.89 ^{+0.18} _{-0.45}	<165.6
GN7_14602	189.168625	62.214306	26.95 ^{+0.25} _{-0.20}	-20.05 ^{+0.52} _{-0.27}	7.91 ^{+0.55} _{-0.52}	<186.5
ERSPRIME_5454792	53.072163	-27.700729	26.64 ^{+0.14} _{-0.12}	-20.70 ^{+0.09} _{-0.09}	7.92 ^{+0.66} _{-0.69}	<69.5
GN3_5454841	189.214523	62.305443	27.62 ^{+0.12} _{-0.11}	-19.50 ^{+0.08} _{-0.11}	7.95 ^{+0.35} _{-0.38}	<240.2
GS4_26328	53.164701	-27.788230	27.43 ^{+0.04} _{-0.04}	-19.80 ^{+0.04} _{-0.01}	7.95 ^{+0.17} _{-0.16}	<73.6
GN3_34236	189.287003	62.318020	26.49 ^{+0.19} _{-0.16}	-20.71 ^{+0.12} _{-0.13}	8.00 ^{+0.38} _{-0.41}	<62.4
GN4_5439303	189.333542	62.254501	28.57 ^{+0.20} _{-0.17}	-18.75 ^{+0.16} _{-0.18}	8.15 ^{+0.61} _{-0.97}	<516.4
GN3_33315	189.224442	62.311329	25.94 ^{+0.02} _{-0.02}	-21.39 ^{+0.02} _{-0.02}	8.18 ^{+0.10} _{-0.09}	<38.7

^aThe listed IDs are encoded with their observed CLEAR fields and matched 3D-HST IDs. We use 5400000 + IDs from Finkelstein et al. (2015) for 3D-HST unmatched objects.

^b M_{UV} is estimated from the averaged flux over a 1450 – 1550 Å bandpass from the best-fit galaxy SED model.

^cWe present the 1σ range of z_{phot} .

^d 3σ upper limits, measured from the median flux limits from individual spectra.

^eThis object is not included in the analysis for the Ly α EW distribution and the IGM transmission due to the tentative emission-line detection.

References

- Ando, M., Ohta, K., Iwata, I., et al. 2006, ApJL, 645, L9, doi: [10.1086/505652](https://doi.org/10.1086/505652)
- Backhaus, B. E., Trump, J. R., Cleri, N. J., et al. 2021, arXiv e-prints, arXiv:2109.08147. <https://arxiv.org/abs/2109.08147>
- Bouwens, R. J., Smit, R., Labbé, I., et al. 2016, ApJ, 831, 176, doi: [10.3847/0004-637X/831/2/176](https://doi.org/10.3847/0004-637X/831/2/176)
- Bouwens, R. J., Illingworth, G. D., Oesch, P. A., et al. 2015, ApJ, 803, 34, doi: [10.1088/0004-637X/803/1/34](https://doi.org/10.1088/0004-637X/803/1/34)
- Brammer, G. 2019, Grizli: Grism redshift and line analysis software. <http://ascl.net/1905.001>
- Brammer, G. B., van Dokkum, P. G., & Coppi, P. 2008, ApJ, 686, 1503, doi: [10.1086/591786](https://doi.org/10.1086/591786)
- Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000, doi: [10.1046/j.1365-8711.2003.06897.x](https://doi.org/10.1046/j.1365-8711.2003.06897.x)
- Calzetti, D. 2001, New Astron., 45, 601, doi: [10.1016/S1387-6473\(01\)00144-0](https://doi.org/10.1016/S1387-6473(01)00144-0)
- Caruana, J., Bunker, A. J., Wilkins, S. M., et al. 2012, MNRAS, 427, 3055, doi: [10.1111/j.1365-2966.2012.21996.x](https://doi.org/10.1111/j.1365-2966.2012.21996.x)
- . 2014, MNRAS, 443, 2831, doi: [10.1093/mnras/stu1341](https://doi.org/10.1093/mnras/stu1341)
- Cash, W. 1979, ApJ, 228, 939, doi: [10.1086/156922](https://doi.org/10.1086/156922)
- Cassata, P., Tasca, L. A. M., Le Fèvre, O., et al. 2015, A&A, 573, A24, doi: [10.1051/0004-6361/201423824](https://doi.org/10.1051/0004-6361/201423824)
- Castellano, M., Pentericci, L., Vanzella, E., et al. 2018, ApJL, 863, L3, doi: [10.3847/2041-8213/aad59b](https://doi.org/10.3847/2041-8213/aad59b)
- Ciardullo, R., Gronwall, C., Wolf, C., et al. 2012, ApJ, 744, 110, doi: [10.1088/0004-637X/744/2/110](https://doi.org/10.1088/0004-637X/744/2/110)
- Cleri, N. J., Trump, J. R., Backhaus, B. E., et al. 2020, arXiv e-prints, arXiv:2009.00617. <https://arxiv.org/abs/2009.00617>
- Cowie, L. L., Barger, A. J., & Hu, E. M. 2010, ApJ, 711, 928, doi: [10.1088/0004-637X/711/2/928](https://doi.org/10.1088/0004-637X/711/2/928)
- Curtis-Lake, E., McLure, R. J., Pearce, H. J., et al. 2012, MNRAS, 422, 1425, doi: [10.1111/j.1365-2966.2012.20720.x](https://doi.org/10.1111/j.1365-2966.2012.20720.x)
- Davies, F. B., Hennawi, J. F., Bañados, E., et al. 2018, ApJ, 864, 142, doi: [10.3847/1538-4357/aad6dc](https://doi.org/10.3847/1538-4357/aad6dc)
- Dayal, P., & Ferrara, A. 2018, PhR, 780, 1, doi: [10.1016/j.physrep.2018.10.002](https://doi.org/10.1016/j.physrep.2018.10.002)
- De Barros, S., Pentericci, L., Vanzella, E., et al. 2017, A&A, 608, A123, doi: [10.1051/0004-6361/201731476](https://doi.org/10.1051/0004-6361/201731476)
- Dijkstra, M., Wyithe, S., Haiman, Z., Mesinger, A., & Pentericci, L. 2014, MNRAS, 440, 3309, doi: [10.1093/mnras/stu531](https://doi.org/10.1093/mnras/stu531)

- Du, X., Shapley, A. E., Tang, M., et al. 2020, *ApJ*, 890, 65, doi: [10.3847/1538-4357/ab67b8](https://doi.org/10.3847/1538-4357/ab67b8)
- Endsley, R., Stark, D. P., Charlot, S., et al. 2021a, *MNRAS*, 502, 6044, doi: [10.1093/mnras/stab432](https://doi.org/10.1093/mnras/stab432)
- Endsley, R., Stark, D. P., Chevallard, J., & Charlot, S. 2021b, *MNRAS*, 500, 5229, doi: [10.1093/mnras/staa3370](https://doi.org/10.1093/mnras/staa3370)
- Estrada-Carpenter, V., Papovich, C., Momcheva, I., et al. 2019, *ApJ*, 870, 133, doi: [10.3847/1538-4357/aaf22e](https://doi.org/10.3847/1538-4357/aaf22e)
- , 2020, *ApJ*, 898, 171, doi: [10.3847/1538-4357/aba004](https://doi.org/10.3847/1538-4357/aba004)
- Finkelstein, S., Bradac, M., Casey, C., et al. 2019a, *BAAS*, 51, 221. <https://arxiv.org/abs/1903.04518>
- Finkelstein, S. L., Papovich, C., Salmon, B., et al. 2012a, *ApJ*, 756, 164, doi: [10.1088/0004-637X/756/2/164](https://doi.org/10.1088/0004-637X/756/2/164)
- Finkelstein, S. L., Papovich, C., Ryan, R. E., et al. 2012b, *ApJ*, 758, 93, doi: [10.1088/0004-637X/758/2/93](https://doi.org/10.1088/0004-637X/758/2/93)
- Finkelstein, S. L., Papovich, C., Dickinson, M., et al. 2013, *Nature*, 502, 524, doi: [10.1038/nature12657](https://doi.org/10.1038/nature12657)
- Finkelstein, S. L., Ryan, Jr., R. E., Papovich, C., et al. 2015, *ApJ*, 810, 71, doi: [10.1088/0004-637X/810/1/71](https://doi.org/10.1088/0004-637X/810/1/71)
- Finkelstein, S. L., D'Aloisio, A., Paardekooper, J.-P., et al. 2019b, *ApJ*, 879, 36, doi: [10.3847/1538-4357/ab1ea8](https://doi.org/10.3847/1538-4357/ab1ea8)
- Finkelstein, S. L., Bagley, M., Song, M., et al. 2021, arXiv e-prints, arXiv:2106.13813. <https://arxiv.org/abs/2106.13813>
- Finlator, K., Özel, F., Davé, R., & Oppenheimer, B. D. 2009, *MNRAS*, 400, 1049, doi: [10.1111/j.1365-2966.2009.15521.x](https://doi.org/10.1111/j.1365-2966.2009.15521.x)
- Fontana, A., Vanzella, E., Pentericci, L., et al. 2010, *ApJL*, 725, L205, doi: [10.1088/2041-8205/725/2/L205](https://doi.org/10.1088/2041-8205/725/2/L205)
- Fuller, S., Lemaux, B. C., Bradač, M., et al. 2020, *ApJ*, 896, 156, doi: [10.3847/1538-4357/ab959f](https://doi.org/10.3847/1538-4357/ab959f)
- Furusawa, H., Kashikawa, N., Kobayashi, M. A. R., et al. 2016, *ApJ*, 822, 46, doi: [10.3847/0004-637X/822/1/46](https://doi.org/10.3847/0004-637X/822/1/46)
- Garel, T., Blaizot, J., Rosdahl, J., et al. 2021, *MNRAS*, 504, 1902, doi: [10.1093/mnras/stab990](https://doi.org/10.1093/mnras/stab990)
- Grogin, N. A., Kocevski, D. D., Faber, S. M., et al. 2011, *ApJS*, 197, 35, doi: [10.1088/0067-0049/197/2/35](https://doi.org/10.1088/0067-0049/197/2/35)
- Gronwall, C., Ciardullo, R., Hickey, T., et al. 2007, *ApJ*, 667, 79, doi: [10.1086/520324](https://doi.org/10.1086/520324)
- Guaita, L., Gawiser, E., Padilla, N., et al. 2010, *ApJ*, 714, 255, doi: [10.1088/0004-637X/714/1/255](https://doi.org/10.1088/0004-637X/714/1/255)
- Hashimoto, T., Garel, T., Guiderdoni, B., et al. 2017, *A&A*, 608, A10, doi: [10.1051/0004-6361/201731579](https://doi.org/10.1051/0004-6361/201731579)
- Hashimoto, T., Laporte, N., Mawatari, K., et al. 2018, *Nature*, 557, 392, doi: [10.1038/s41586-018-0117-z](https://doi.org/10.1038/s41586-018-0117-z)
- Hassan, S., & Gronke, M. 2021, *ApJ*, 908, 219, doi: [10.3847/1538-4357/abd554](https://doi.org/10.3847/1538-4357/abd554)
- Hastings, W. K. 1970, *Biometrika*, 57, 97, doi: [10.1093/biomet/57.1.97](https://doi.org/10.1093/biomet/57.1.97)
- Hoag, A., Bradač, M., Huang, K., et al. 2019, *ApJ*, 878, 12, doi: [10.3847/1538-4357/ab1de7](https://doi.org/10.3847/1538-4357/ab1de7)
- Hu, E. M., Cowie, L. L., Barger, A. J., et al. 2010, *ApJ*, 725, 394, doi: [10.1088/0004-637X/725/1/394](https://doi.org/10.1088/0004-637X/725/1/394)
- Hu, W., Wang, J., Infante, L., et al. 2021, *Nature Astronomy*, doi: [10.1038/s41550-021-01322-2](https://doi.org/10.1038/s41550-021-01322-2)
- Hutchison, T. A., Papovich, C., Finkelstein, S. L., et al. 2019, *ApJ*, 879, 70, doi: [10.3847/1538-4357/ab22a2](https://doi.org/10.3847/1538-4357/ab22a2)
- Inoue, A. K. 2011, *MNRAS*, 415, 2920, doi: [10.1111/j.1365-2966.2011.18906.x](https://doi.org/10.1111/j.1365-2966.2011.18906.x)
- Jung, I., Finkelstein, S. L., Livermore, R. C., et al. 2018, *ApJ*, 864, 103, doi: [10.3847/1538-4357/aad686](https://doi.org/10.3847/1538-4357/aad686)
- Jung, I., Finkelstein, S. L., Dickinson, M., et al. 2019, *ApJ*, 877, 146, doi: [10.3847/1538-4357/ab1bde](https://doi.org/10.3847/1538-4357/ab1bde)
- , 2020, *ApJ*, 904, 144, doi: [10.3847/1538-4357/abbd44](https://doi.org/10.3847/1538-4357/abbd44)
- Kashikawa, N., Shimasaku, K., Matsuda, Y., et al. 2011, *ApJ*, 734, 119, doi: [10.1088/0004-637X/734/2/119](https://doi.org/10.1088/0004-637X/734/2/119)
- Katz, H., Kimm, T., Haehnelt, M. G., et al. 2019, *MNRAS*, 483, 1029, doi: [10.1093/mnras/sty3154](https://doi.org/10.1093/mnras/sty3154)
- Kimm, T., Blaizot, J., Garel, T., et al. 2019, *MNRAS*, 486, 2215, doi: [10.1093/mnras/stz989](https://doi.org/10.1093/mnras/stz989)
- Koekemoer, A. M., Faber, S. M., Ferguson, H. C., et al. 2011, *ApJS*, 197, 36, doi: [10.1088/0067-0049/197/2/36](https://doi.org/10.1088/0067-0049/197/2/36)
- Kulkarni, G., Keating, L. C., Haehnelt, M. G., et al. 2019, *MNRAS*, 485, L24, doi: [10.1093/mnras/slz025](https://doi.org/10.1093/mnras/slz025)
- Larson, R. L., Finkelstein, S. L., Pirzkal, N., et al. 2018, *ApJ*, 858, 94, doi: [10.3847/1538-4357/aab893](https://doi.org/10.3847/1538-4357/aab893)
- Leclercq, F., Bacon, R., Wisotzki, L., et al. 2017, *A&A*, 608, A8, doi: [10.1051/0004-6361/201731480](https://doi.org/10.1051/0004-6361/201731480)
- Madau, P. 1995, *ApJ*, 441, 18, doi: [10.1086/175332](https://doi.org/10.1086/175332)
- Malhotra, S., & Rhoads, J. E. 2004, *ApJL*, 617, L5, doi: [10.1086/427182](https://doi.org/10.1086/427182)
- Mallery, R. P., Mobasher, B., Capak, P., et al. 2012, *ApJ*, 760, 128, doi: [10.1088/0004-637X/760/2/128](https://doi.org/10.1088/0004-637X/760/2/128)
- Markwardt, C. B. 2009, in *Astronomical Society of the Pacific Conference Series*, Vol. 411, *Astronomical Data Analysis Software and Systems XVIII*, ed. D. A. Bohlender, D. Durand, & P. Dowler, 251. <https://arxiv.org/abs/0902.2850>
- Mason, C. A., Treu, T., Dijkstra, M., et al. 2018a, *ApJ*, 856, 2, doi: [10.3847/1538-4357/aab0a7](https://doi.org/10.3847/1538-4357/aab0a7)
- Mason, C. A., Treu, T., de Barros, S., et al. 2018b, *ApJL*, 857, L11, doi: [10.3847/2041-8213/aabbab](https://doi.org/10.3847/2041-8213/aabbab)
- Mason, C. A., Fontana, A., Treu, T., et al. 2019, *MNRAS*, 485, 3947, doi: [10.1093/mnras/stz632](https://doi.org/10.1093/mnras/stz632)
- Matsuoka, Y., Strauss, M. A., Kashikawa, N., et al. 2018, *ApJ*, 869, 150, doi: [10.3847/1538-4357/aaec7a](https://doi.org/10.3847/1538-4357/aaec7a)
- McGreer, I. D., Mesinger, A., & D'Odorico, V. 2015, *MNRAS*, 447, 499, doi: [10.1093/mnras/stu2449](https://doi.org/10.1093/mnras/stu2449)
- McQuinn, M. 2016, *ARA&A*, 54, 313, doi: [10.1146/annurev-astro-082214-122355](https://doi.org/10.1146/annurev-astro-082214-122355)
- Mesinger, A., Aykotalp, A., Vanzella, E., et al. 2015, *MNRAS*, 446, 566, doi: [10.1093/mnras/stu2089](https://doi.org/10.1093/mnras/stu2089)

- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., & Teller, E. 1953, *JChPh*, 21, 1087, doi: [10.1063/1.1699114](https://doi.org/10.1063/1.1699114)
- Miralda-Escudé, J., & Rees, M. J. 1998, *ApJ*, 497, 21, doi: [10.1086/305458](https://doi.org/10.1086/305458)
- Morales, A., Mason, C., Bruton, S., et al. 2021, arXiv e-prints, arXiv:2101.01205. <https://arxiv.org/abs/2101.01205>
- Naidu, R. P., Tacchella, S., Mason, C. A., et al. 2020, *ApJ*, 892, 109, doi: [10.3847/1538-4357/ab7cc9](https://doi.org/10.3847/1538-4357/ab7cc9)
- Nilsson, K. K., Tapken, C., Møller, P., et al. 2009, *A&A*, 498, 13, doi: [10.1051/0004-6361/200810881](https://doi.org/10.1051/0004-6361/200810881)
- Ocvirk, P., Lewis, J. S. W., Gillet, N., et al. 2021, arXiv e-prints, arXiv:2105.01663. <https://arxiv.org/abs/2105.01663>
- Ocvirk, P., Aubert, D., Sorce, J. G., et al. 2020, *MNRAS*, 496, 4087, doi: [10.1093/mnras/staa1266](https://doi.org/10.1093/mnras/staa1266)
- Oesch, P. A., van Dokkum, P. G., Illingworth, G. D., et al. 2015, *ApJL*, 804, L30, doi: [10.1088/2041-8205/804/2/L30](https://doi.org/10.1088/2041-8205/804/2/L30)
- Oesch, P. A., Brammer, G., van Dokkum, P. G., et al. 2016, *ApJ*, 819, 129, doi: [10.3847/0004-637X/819/2/129](https://doi.org/10.3847/0004-637X/819/2/129)
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713, doi: [10.1086/160817](https://doi.org/10.1086/160817)
- Ono, Y., Ouchi, M., Mobasher, B., et al. 2012, *ApJ*, 744, 83, doi: [10.1088/0004-637X/744/2/83](https://doi.org/10.1088/0004-637X/744/2/83)
- Ouchi, M., Shimasaku, K., Akiyama, M., et al. 2008, *ApJS*, 176, 301, doi: [10.1086/527673](https://doi.org/10.1086/527673)
- Oyarzún, G. A., Blanc, G. A., González, V., Mateo, M., & Bailey, John I., I. 2017, *ApJ*, 843, 133, doi: [10.3847/1538-4357/aa7552](https://doi.org/10.3847/1538-4357/aa7552)
- Park, H., Jung, I., Song, H., et al. 2021, arXiv e-prints, arXiv:2105.10770. <https://arxiv.org/abs/2105.10770>
- Pentericci, L., Fontana, A., Vanzella, E., et al. 2011, *ApJ*, 743, 132, doi: [10.1088/0004-637X/743/2/132](https://doi.org/10.1088/0004-637X/743/2/132)
- Pentericci, L., Vanzella, E., Fontana, A., et al. 2014, *ApJ*, 793, 113, doi: [10.1088/0004-637X/793/2/113](https://doi.org/10.1088/0004-637X/793/2/113)
- Pentericci, L., Vanzella, E., Castellano, M., et al. 2018, *A&A*, 619, A147, doi: [10.1051/0004-6361/201732465](https://doi.org/10.1051/0004-6361/201732465)
- Pirzkal, N., Malhotra, S., Ryan, R. E., et al. 2017, *ApJ*, 846, 84, doi: [10.3847/1538-4357/aa81cc](https://doi.org/10.3847/1538-4357/aa81cc)
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13, doi: [10.1051/0004-6361/201525830](https://doi.org/10.1051/0004-6361/201525830)
- Qin, Y., Wyithe, J. S. B., Oesch, P. A., et al. 2021, arXiv e-prints, arXiv:2108.03675. <https://arxiv.org/abs/2108.03675>
- Rhoads, J. E., & Malhotra, S. 2001, *ApJL*, 563, L5, doi: [10.1086/338477](https://doi.org/10.1086/338477)
- Rhoads, J. E., Malhotra, S., Stern, D., et al. 2013, *ApJ*, 773, 32, doi: [10.1088/0004-637X/773/1/32](https://doi.org/10.1088/0004-637X/773/1/32)
- Roberts-Borsani, G. W., Bouwens, R. J., Oesch, P. A., et al. 2016, *ApJ*, 823, 143, doi: [10.3847/0004-637X/823/2/143](https://doi.org/10.3847/0004-637X/823/2/143)
- Robertson, B. E. 2021, arXiv e-prints, arXiv:2110.13160. <https://arxiv.org/abs/2110.13160>
- Robertson, B. E., Ellis, R. S., Furlanetto, S. R., & Dunlop, J. S. 2015, *ApJL*, 802, L19, doi: [10.1088/2041-8205/802/2/L19](https://doi.org/10.1088/2041-8205/802/2/L19)
- Robertson, B. E., Furlanetto, S. R., Schneider, E., et al. 2013, *ApJ*, 768, 71, doi: [10.1088/0004-637X/768/1/71](https://doi.org/10.1088/0004-637X/768/1/71)
- Salmon, B., Papovich, C., Finkelstein, S. L., et al. 2015, *ApJ*, 799, 183, doi: [10.1088/0004-637X/799/2/183](https://doi.org/10.1088/0004-637X/799/2/183)
- Salpeter, E. E. 1955, *ApJ*, 121, 161, doi: [10.1086/145971](https://doi.org/10.1086/145971)
- Santos, S., Sobral, D., Matthee, J., et al. 2020, *MNRAS*, 493, 141, doi: [10.1093/mnras/staa093](https://doi.org/10.1093/mnras/staa093)
- Saxena, A., Pentericci, L., Ellis, R. S., et al. 2021, arXiv e-prints, arXiv:2109.03662. <https://arxiv.org/abs/2109.03662>
- Schaerer, D., de Barros, S., & Stark, D. P. 2011, *A&A*, 536, A72, doi: [10.1051/0004-6361/201117685](https://doi.org/10.1051/0004-6361/201117685)
- Schenker, M. A., Ellis, R. S., Konidaris, N. P., & Stark, D. P. 2014, *ApJ*, 795, 20, doi: [10.1088/0004-637X/795/1/20](https://doi.org/10.1088/0004-637X/795/1/20)
- Schenker, M. A., Stark, D. P., Ellis, R. S., et al. 2012, *ApJ*, 744, 179, doi: [10.1088/0004-637X/744/2/179](https://doi.org/10.1088/0004-637X/744/2/179)
- Schmidt, K. B., Treu, T., Bradač, M., et al. 2016, *ApJ*, 818, 38, doi: [10.3847/0004-637X/818/1/38](https://doi.org/10.3847/0004-637X/818/1/38)
- Shibuya, T., Ouchi, M., Harikane, Y., & Nakajima, K. 2019, *ApJ*, 871, 164, doi: [10.3847/1538-4357/aaf64b](https://doi.org/10.3847/1538-4357/aaf64b)
- Simons, R. C., Papovich, C., Momcheva, I., et al. 2020, arXiv e-prints, arXiv:2011.03553. <https://arxiv.org/abs/2011.03553>
- Skelton, R. E., Whitaker, K. E., Momcheva, I. G., et al. 2014, *ApJS*, 214, 24, doi: [10.1088/0067-0049/214/2/24](https://doi.org/10.1088/0067-0049/214/2/24)
- Smith, A., Kannan, R., Garaldi, E., et al. 2021, arXiv e-prints, arXiv:2110.02966. <https://arxiv.org/abs/2110.02966>
- Song, M., Finkelstein, S. L., Livermore, R. C., et al. 2016a, *ApJ*, 826, 113, doi: [10.3847/0004-637X/826/2/113](https://doi.org/10.3847/0004-637X/826/2/113)
- Song, M., Finkelstein, S. L., Ashby, M. L. N., et al. 2016b, *ApJ*, 825, 5, doi: [10.3847/0004-637X/825/1/5](https://doi.org/10.3847/0004-637X/825/1/5)
- Stark, D. P., Ellis, R. S., Chiu, K., Ouchi, M., & Bunker, A. 2010, *MNRAS*, 408, 1628, doi: [10.1111/j.1365-2966.2010.17227.x](https://doi.org/10.1111/j.1365-2966.2010.17227.x)
- Stark, D. P., Ellis, R. S., & Ouchi, M. 2011, *ApJL*, 728, L2, doi: [10.1088/2041-8205/728/1/L2](https://doi.org/10.1088/2041-8205/728/1/L2)
- Stark, D. P., Ellis, R. S., Charlot, S., et al. 2017, *MNRAS*, 464, 469, doi: [10.1093/mnras/stw2233](https://doi.org/10.1093/mnras/stw2233)
- Tang, M., Stark, D. P., Chevallard, J., & Charlot, S. 2019, *MNRAS*, 489, 2572, doi: [10.1093/mnras/stz2236](https://doi.org/10.1093/mnras/stz2236)
- Tang, M., Stark, D. P., Chevallard, J., et al. 2021a, *MNRAS*, 501, 3238, doi: [10.1093/mnras/staa3454](https://doi.org/10.1093/mnras/staa3454)
- , 2021b, *MNRAS*, doi: [10.1093/mnras/stab705](https://doi.org/10.1093/mnras/stab705)
- Tilvi, V., Papovich, C., Finkelstein, S. L., et al. 2014, *ApJ*, 794, 5, doi: [10.1088/0004-637X/794/1/5](https://doi.org/10.1088/0004-637X/794/1/5)
- Tilvi, V., Pirzkal, N., Malhotra, S., et al. 2016, *ApJL*, 827, L14, doi: [10.3847/2041-8205/827/1/L14](https://doi.org/10.3847/2041-8205/827/1/L14)
- Tilvi, V., Malhotra, S., Rhoads, J. E., et al. 2020, *ApJL*, 891, L10, doi: [10.3847/2041-8213/ab75ec](https://doi.org/10.3847/2041-8213/ab75ec)
- Trainor, R. F., Strom, A. L., Steidel, C. C., et al. 2019, *ApJ*, 887, 85, doi: [10.3847/1538-4357/ab4993](https://doi.org/10.3847/1538-4357/ab4993)
- Treu, T., Schmidt, K. B., Trenti, M., Bradley, L. D., & Stiavelli, M. 2013, *ApJL*, 775, L29, doi: [10.1088/2041-8205/775/1/L29](https://doi.org/10.1088/2041-8205/775/1/L29)

- Treu, T., Trenti, M., Stiavelli, M., Auger, M. W., & Bradley, L. D. 2012, *ApJ*, 747, 27, doi: [10.1088/0004-637X/747/1/27](https://doi.org/10.1088/0004-637X/747/1/27)
- Treu, T., Schmidt, K. B., Brammer, G. B., et al. 2015, *ApJ*, 812, 114, doi: [10.1088/0004-637X/812/2/114](https://doi.org/10.1088/0004-637X/812/2/114)
- Vanzella, E., Fontana, A., Pentericci, L., et al. 2014, *A&A*, 569, A78, doi: [10.1051/0004-6361/201424285](https://doi.org/10.1051/0004-6361/201424285)
- Wang, F., Davies, F. B., Yang, J., et al. 2020, *ApJ*, 896, 23, doi: [10.3847/1538-4357/ab8c45](https://doi.org/10.3847/1538-4357/ab8c45)
- Watson, D., Christensen, L., Knudsen, K. K., et al. 2015, *Nature*, 519, 327, doi: [10.1038/nature14164](https://doi.org/10.1038/nature14164)
- Weinberger, L. H., Kulkarni, G., Haehnelt, M. G., Choudhury, T. R., & Puchwein, E. 2018, *MNRAS*, 479, 2564, doi: [10.1093/mnras/sty1563](https://doi.org/10.1093/mnras/sty1563)
- Weiss, L. H., Bowman, W. P., Ciardullo, R., et al. 2021, arXiv e-prints, arXiv:2103.12126. <https://arxiv.org/abs/2103.12126>
- Wisotzki, L., Bacon, R., Brinchmann, J., et al. 2018, *Nature*, 562, 229, doi: [10.1038/s41586-018-0564-6](https://doi.org/10.1038/s41586-018-0564-6)
- Wold, I. G. B., Barger, A. J., & Cowie, L. L. 2014, *ApJ*, 783, 119, doi: [10.1088/0004-637X/783/2/119](https://doi.org/10.1088/0004-637X/783/2/119)
- Wold, I. G. B., Finkelstein, S. L., Barger, A. J., Cowie, L. L., & Rosenwasser, B. 2017, *ApJ*, 848, 108, doi: [10.3847/1538-4357/aa8d6b](https://doi.org/10.3847/1538-4357/aa8d6b)
- Wold, I. G. B., Malhotra, S., Rhoads, J., et al. 2021, arXiv e-prints, arXiv:2105.12191. <https://arxiv.org/abs/2105.12191>
- Yang, H., Malhotra, S., Rhoads, J. E., et al. 2017, *ApJ*, 838, 4, doi: [10.3847/1538-4357/aa6337](https://doi.org/10.3847/1538-4357/aa6337)
- Yang, J., Wang, F., Fan, X., et al. 2020, *ApJL*, 897, L14, doi: [10.3847/2041-8213/ab9c26](https://doi.org/10.3847/2041-8213/ab9c26)
- Yoo, T., Kimm, T., & Rosdahl, J. 2020, *MNRAS*, 499, 5175, doi: [10.1093/mnras/staa3187](https://doi.org/10.1093/mnras/staa3187)
- Zheng, Z.-Y., Wang, J.-X., Malhotra, S., et al. 2014, *MNRAS*, 439, 1101, doi: [10.1093/mnras/stu054](https://doi.org/10.1093/mnras/stu054)
- Zheng, Z.-Y., Wang, J., Rhoads, J., et al. 2017, *ApJL*, 842, L22, doi: [10.3847/2041-8213/aa794f](https://doi.org/10.3847/2041-8213/aa794f)
- Zitrin, A., Labbé, I., Belli, S., et al. 2015, *ApJL*, 810, L12, doi: [10.1088/2041-8205/810/1/L12](https://doi.org/10.1088/2041-8205/810/1/L12)