

Quantum Regression theorem for multi-time correlators : A detailed analysis in the Heisenberg Picture

Sakil Khan,^{1,*} Bijay Kumar Agarwalla,^{1,†} and Sachin Jain^{1,‡}

¹*Department of Physics, Indian Institute of Science Education and Research, Pune 411008, India*

(Dated: December 1, 2021)

Quantum regression theorem is a very useful result in open quantum system and extensively used for computing multi-point correlation functions. Traditionally it is derived for two-time correlators in the Markovian limit employing the Schrödinger picture. In this paper we make use of the Heisenberg picture to derive quantum regression theorems for multi-time correlation functions which in the special limit reduce to the well known two-time regression theorem. For multi-time correlation function we find that the regression theorem takes the same form as it takes for two-time correlation function with a mild restriction that one of the times should be greater than all the other time variables. Interestingly, the Heisenberg picture also allows us to derive analogue of regression theorem for out-of-time-ordered correlators (OTOCs). We further extend our study for the case of non-Markovian dynamics and report the modifications to the standard quantum regression theorem. We illustrate all of the above results using the paradigmatic dissipative spin-boson model.

I. INTRODUCTION

Correlation functions are important dynamical quantities which are often related to experimentally measurable quantities. In the context of open quantum system [1–6] with the system of interest following a Markovian dynamics, Quantum Regression Theorem (QRT)[4–7] turns out to be one of the most useful and practical tools to compute the correlation functions [4]. However, it is limited to certain special kind of time arrangements and thus allowing to compute only special types of correlation functions. The QRT states that the knowledge of time evolution of a single-point function is sufficient to determine the time evolution of two-point or multi-point correlation functions. More explicitly, the validity of QRT requires that there exists a complete set of system operators $A_\mu, \mu = 1, 2, \dots$ such that

$$\frac{d}{dt}\langle A_\mu(t) \rangle = \sum_\lambda M_{\mu\lambda}\langle A_\lambda(t) \rangle. \quad (1)$$

Then the QRT for two and three-point function reads as,

$$\begin{aligned} \frac{d}{d\tau}\langle O(t)A_\mu(t+\tau) \rangle &= \sum_\lambda M_{\mu\lambda}\langle O(t)A_\lambda(t+\tau) \rangle, \\ \frac{d}{d\tau}\langle O_1(t)A_\mu(t+\tau)O_2(t) \rangle &= \sum_\lambda M_{\mu\lambda}\langle O_1(t)A_\lambda(t+\tau)O_2(t) \rangle, \\ \frac{d}{d\tau}\langle A_\mu(t+\tau)O_1(t)O_3(t) \rangle &= \sum_\lambda M_{\mu\lambda}\langle A_\lambda(t+\tau)O_1(t)O_3(t) \rangle. \end{aligned} \quad (2)$$

Interestingly, it is easy to generalize the QRT in Eq. (2) for arbitrary N -point correlation functions of the form

$$\langle A_1(t)A_2(t+\tau)A_3(t) \cdots A_n(t) \rangle. \quad (3)$$

where the position of the operator with argument $t + \tau$ can be arbitrary. Note that the above QRTs are given for multi-point correlation functions which are dependent on two-times t and τ . Recently there has been a lot of research activity to understand systems that follow non-Markovian dynamics [8–12] and attempt has been made to show violation of regression theorems for such systems [13–19]. In spite of its great importance and interest, there has been a lack of systematic derivation of Regression type theorem for cases beyond two-time correlation function and its extension for non-Markovian systems. In general, regression theorem may not hold for general time configurations such as

$$\langle A(t_1)B(t_2)C(t_3) \rangle \quad (4)$$

or more generally

$$\langle A_1(t_1)A_2(t_2) \cdots A_n(t_n) \rangle. \quad (5)$$

Here we would like to understand if there exist QRT type relations for such class of correlation functions including the out-of-time-ordered correlators (OTOCs) [20] which are a special class of correlation functions [21].

As in the Schrödinger picture it is difficult to define multi-time correlators of the above form, it is therefore more appropriate and suitable to consider such correlators in the Heisenberg picture [22]. We therefore, in this paper, derive regression theorem for multi-time correlators, in the Markovian limit, using the Heisenberg picture, as discussed in Ref. [22]. We also extend our study for the non-Markovian systems as well.

The paper is organised as follows: In section II, we first start with deriving the QRT for two-point functions and then extend our analysis to three, four and further generalize to N -point functions with general time arrangements. We also point out for what special time arrangements the QRT may not hold. In section III, we derive a regression like expression for the OTOC. In section IV, we illustrate all these above findings for a paradigmatic dissipative spin-boson model. In section V, we give a

* sakil.khan@students.iiserpune.ac.in

† bijay@iiserpune.ac.in

‡ sachin.jain@iiserpune.ac.in

systematic derivation of the equation similar to QRT for non-Markovian systems. We skip most of the lengthy derivations to the appendix to keep the discussion in the main text transparent.

II. QRT USING THE HEISENBERG PICTURE

In this section we derive QRT for multi-time correlation functions. We make extensive use of the Heisenberg picture formulation for open quantum system, recently described in [22] and for completeness we also review this formalism in Appendix A and further discuss the Markovian limit in Appendix A 2. Our first aim here is to derive the well-known forms of QRTs for two-point and special three-point functions, as discussed in the introduction, using the Heisenberg picture. We then aim for generalizing the QRTs for more generic N -point correlation functions defined with multi-time arguments. We also discuss the limitations of the QRT in the Markovian limit and also generalize our study to non-Markovian systems.

Let the start by writing the Hamiltonian of the total sy $H = H_S + H_R + \lambda H_{SR}$, where H_S is the Hamiltonian of the system of interest, H_R represents the Hamiltonian for the bath (reservoir), and H_{SR} is the coupling Hamiltonian between the system and the bath. We keep the parameter λ to keep track of the order of the perturbation with respect to system-bath interaction. We also make the standard choice for the initial condition of the total density operator at $t = 0$ by considering a product initial state between the system and the bath and write $\rho_{SR}(t = 0) = \rho_S \otimes \rho_R$. The interaction between the system and the bath is turned on at $t = 0^+$. Since in the Heisenberg picture, the operators evolve in time, it is important to define the reduced system operators. The one point reduced operator is defined as $O_S(t) = \text{Tr}_R[O(t)\rho_R]$ where the operator $O(t)$ evolves unitarily with respect to the full Hamiltonian H . The expectation value of the operator O at time t then can be written as,

$$\langle O(t) \rangle = \text{Tr}_S[\text{Tr}_R[O(t)\rho_R]\rho_S] = \text{Tr}_S[O_S(t)\rho_S]. \quad (6)$$

In a similar manner, one can define arbitrary N -point reduced operator as

$$(O_1(t_1)O_2(t_2)\dots O_N(t_N))_S = \text{Tr}_R[O_1(t_1)O_2(t_2)\dots O_N(t_N)\rho_R]. \quad (7)$$

This definition of reduced operator has a property that $(O_1(t_1)O_2(t_2)\dots O_N(t_N))_S \neq O_{1S}(t_1)O_{2S}(t_2)\dots O_N(t_N)$ as a result of finite system-bath coupling.

To derive the QRT in the Heisenberg picture we write an analogous equation like equation (1) in the Heisenberg picture by assuming that there exists a complete set of reduced system operators $A_{\mu S}(t)$ that satisfies [23] the relation

$$\frac{d}{dt}A_{\mu S}(t) = \sum_{\lambda} M_{\mu\lambda}A_{\lambda S}(t). \quad (8)$$

The above equation implies that the operators form a closed set between themselves.

A. QRT for two-point correlation functions

To keep our discussion simple, we first focus on deriving the QRT for two-point correlation functions. Following the definition in (7) for two-point reduced operators, one can write [22] (please see Appendix A for details of the derivation)

$$(O_1(t_1)O_2(t_2))_S = O_{1S}(t_1)O_{2S}(t_2) + I[O_{1S}(t_1), O_{2S}(t_2)] \quad (9)$$

where $O_{1S}(t_1), O_{2S}(t_2)$ are reduced one-point operators and $I[O_{1S}(t_1), O_{2S}(t_2)]$ is called the irreducible part capturing the information about coupled system-bath dynamics. [24]

One can explicitly calculate the quantity I upto the second-order of the system-bath coupling (λ^2) in both Markovian and non-Markovian limits. Given the above expression, the two-point correlation functions can be easily computed by performing an additional trace over the initial system density operator i.e.,

$$\langle O_1(t_1)O_2(t_2) \rangle = \text{Tr}_S[(O_1(t_1)O_2(t_2))_S\rho_S(0)] \quad (10)$$

To derive the QRT, we set, $O_2 = A_{\mu}, O_1 = O$, and consider $t_2 > t_1$. Now taking derivative with respect to t_2 in equation (9), the first term of the of the right hand side immediately gives a QRT like expression

$$\frac{d}{dt_2}(O_S(t_1)A_{\mu S}(t_2)) = \sum_{\lambda} M_{\mu\lambda}(O_S(t_1)A_{\lambda S}(t_2)). \quad (11)$$

thanks to Eq. (8). Now interestingly, one can show that for $t_2 > t_1$ and in the Markovian limit, the irreducible part $I[O_S(t_1), A_{\mu S}(t_2)]$ also satisfies a regression like expression, given as,

$$\frac{d}{dt_2}I[O_S(t_1), A_{\mu S}(t_2)] = \sum_{\lambda} M_{\mu\lambda}I[O_S(t_1), A_{\lambda S}(t_2)]. \quad (12)$$

(please see equation (B1)) As a result of the above two equations, we receive the QRT for arbitrary two-point system operators as

$$\frac{d}{dt_2}(O(t_1)A_{\mu}(t_2))_S = \sum_{\lambda} M_{\mu\lambda}(O(t_1)A_{\lambda}(t_2))_S, \quad (13)$$

from which we trivially receive the standard QRT in terms of correlation functions,

$$\frac{d}{dt_2}\langle O(t_1)A_{\mu}(t_2) \rangle = \sum_{\lambda} M_{\mu\lambda}\langle O(t_1)A_{\lambda}(t_2) \rangle, \quad (14)$$

which matches with (2). Furthermore, in the expression of $I[O_S(t_1), A_{\mu S}(t_2)]$, it can be shown that (see equations

(A16),(A25)-(A27)) if we swap the position of $O_S(t_1)$ and $A_{\mu S}(t_2)$, equation (12) still gets respected, i.e.,

$$\frac{d}{dt_2} I[A_{\mu S}(t_2), O_S(t_1)] = \sum_{\lambda} M_{\mu\lambda} I[A_{\lambda S}(t_2), O_S(t_1)] \quad (15)$$

and as a result we receive another form of QRT as,

$$\frac{d}{dt_2} \langle A_{\mu}(t_2) O(t_1) \rangle = \sum_{\lambda} M_{\mu\lambda} \langle A_{\lambda}(t_2) O(t_1) \rangle. \quad (16)$$

It is important to note that, if we consider the other time sequence i.e., if $t_2 < t_1$ and take derivative with respect to t_2 , what we receive does not obey the standard QRT.

$$\begin{aligned} (O_1(t_1)O_2(t_2)O_3(t_3))_S &= O_{1S}(t_1)O_{2S}(t_2)O_{3S}(t_3) + W_{1,2,3} \left\{ O_{1S}(t_1) I[O_{2S}(t_2), O_{3S}(t_3)] \right\} \\ &+ W_{1,2,3} \left\{ I[O_{1S}(t_1), O_{2S}(t_2)] O_{3S}(t_3) \right\} + W_{1,2,3} \left\{ I[O_{1S}(t_1), O_{3S}(t_3)] O_{2S}(t_2) \right\}, \end{aligned} \quad (17)$$

where the operator $W_{1,2,3}$ ensures that the operator product is ordered such that O_{1S} comes before O_{2S} , and O_{2S} comes before O_{3S} . (please see A1b more details). The three-point multi-time correlation functions can be computed as,

$$\langle O_1(t_1)O_2(t_2)O_3(t_3) \rangle = \text{Tr}_S[(O_1(t_1)O_2(t_2)O_3(t_3))_S \rho_S(0)] \quad (18)$$

Now once again, to derive the QRT we first set, $O_3 = A_{\mu}$ and assume $t_i < t_3$ with $i = 1, 2$. Taking derivative of the equation (17) with respect to t_3 , 1st term of the right hand side gives

$$\begin{aligned} \frac{d}{dt_3} (O_{1S}(t_1)O_{2S}(t_2)A_{\mu S}(t_3)) \\ = \sum_{\lambda} M_{\mu\lambda} (O_{1S}(t_1)O_{2S}(t_2)A_{\lambda S}(t_3)) \end{aligned} \quad (19)$$

where we have used Eq.(8). Now once again one can show that (please see (B8) of appendix B2) for the time sequence $t_i < t_3$ with $i = 1, 2$ and invoking the Markovian limit, the second term of the right hand side of equation (17) satisfies the equation

$$\begin{aligned} \frac{d}{dt_3} W_{1,2,3} \{ O_{1S}(t_1) I[O_{2S}(t_2), A_{\mu S}(t_3)] \} \\ = \sum_{\lambda} M_{\mu\lambda} W_{1,2,3} \{ O_{1S}(t_1) I[O_{2S}(t_2), A_{\lambda S}(t_3)] \} \end{aligned} \quad (20)$$

Interestingly, the third and the fourth term also follow identical equations like above. Finally, summing up all these contribution, we receive a multi-time QRT like form

These results is what one also receives by working in the Schrödinger picture. We next discuss the generalization of our analysis for higher-point and multi-time correlators.

B. QRT for three-point, multi-time correlation functions

Following similar steps as before, we can define the three-point multi-time reduced operators which upto the second-order of system-bath coupling (λ^2 order) is given as [22],

involving the reduced system operators,

$$\begin{aligned} \frac{d}{dt_3} (O_1(t_1)O_2(t_2)A_{\mu}(t_3))_S \\ = \sum_{\lambda} M_{\mu\lambda} (O_1(t_1)O_2(t_2)A_{\lambda}(t_3))_S \end{aligned} \quad (21)$$

from which we trivially receive the QRT for the three-point correlation functions as,

$$\frac{d}{dt_3} \langle O_1(t_1)O_2(t_2)A_{\mu}(t_3) \rangle = \sum_{\lambda} M_{\mu\lambda} \langle O_1(t_1)O_2(t_2)A_{\lambda}(t_3) \rangle. \quad (22)$$

Interestingly if we swap the position of $O_2(t_2)$ and $A_{\mu}(t_3)$, one can obtain a similar QRT. The equations (22) is the regression theorem for three-point correlation functions that involve three different times t_1, t_2 , and t_3 . This above regression type relation is of much more general form than the standard QRT derived using the Schrödinger picture. In fact, in special limits $t_1 = t_2$ the above QRT in (22) reduces to the standard result

$$\frac{d}{dt_3} \langle O_1(t_1)A_{\mu}(t_3)O_2(t_1) \rangle = \sum_{\lambda} M_{\mu\lambda} \langle O_1(t_1)A_{\lambda}(t_3)O_2(t_1) \rangle \quad (23)$$

Note that the QRT in our case holds irrespective of the position of $A_{\mu}(t_3)$. However in equation (22), if we take derivative with respect to t_1 or t_2 instead of t_3 (maximum time), interestingly, we do not receive a QRT like relation.

C. QRT for four-point and general N-point, multi-time correlation functions

Following almost similar steps as before, one can work out the regression theorem for four-point functions and in

fact, it is possible to generalize this analysis for N -point correlation functions, in the Markovian limit. Here we present the central results (see appendix B3 for more details). For four-point, multi-time correlation functions we receive a QRT,

$$\begin{aligned} & \frac{d}{dt_4} \langle O_1(t_1) O_2(t_2) O_3(t_3) A_\mu(t_4) \rangle \\ &= \sum_{\lambda} M_{\mu\lambda} \langle O_1(t_1) O_2(t_2) O_3(t_3) A_\lambda(t_4) \rangle. \end{aligned} \quad (24)$$

Let us emphasize that (24) holds as long as $t_4 > t_i$ with $i = 1, 2, 3$. Also note that (24) holds irrespective of the position of $A_\mu(t_4)$ as was observed for the QRT for three-point functions. This entire analysis can be generalised to N -point multi-time correlation functions as,

$$\begin{aligned} & \frac{d}{dt_N} \langle O_1(t_1) O_2(t_2) \dots O_{N-1}(t_{N-1}) A_\mu(t_N) \rangle \\ &= \sum_{\lambda} M_{\mu\lambda} \langle O_1(t_1) O_2(t_2) \dots O_{N-1}(t_{N-1}) A_\lambda(t_N) \rangle \end{aligned} \quad (25)$$

where, $t_i < t_N$ and $i = 1, 2, \dots, N-1$. Once again, the operator $A_\mu(t_N)$ can take any place and if we take derivative with respect to t_k instead of the highest time t_N , then we do not receive the regression type formula.

III. QRT FOR OUT-OF-TIME-ORDERED CORRELATORS

As an application of the developed formalism, we now extend our analysis to compute the out of time ordered correlator (OTOC)[25] which is an excellent measure of many body localization, quantum chaos and scrambling [26–32]. OTOC has received significant attention in recent times with its applicability ranging from quantum information theory to condensed matter physics to quantum gravity. Very recently, OTOC has found its application in the context of open quantum systems as the coupling of the system with a dissipative or dephasing bath naturally leads to information scrambling [33, 34]. Motivated by this, in this section we derive QRT like formula for OTOC correlators. Details of the derivation are provided in appendix C. Here we present the main result. Let us first define four-point reduced operator of the form

$$(O_1(t_1) A_\mu(t_2) O_3(t_1) A_\nu(t_2))_S \quad (26)$$

where the system reduced operator $A_{\mu S}(t_2)$ satisfies Eq. (8). One can then receive the following regression

type formula for the OTOC,

$$\begin{aligned} & \frac{d}{dt_2} \langle O_1(t_1) A_\mu(t_2) O_3(t_1) A_\nu(t_2) \rangle \\ &= \sum_{\lambda} M_{\mu\lambda} \langle O_1(t_1) A_\lambda(t_2) O_3(t_1) A_\nu(t_2) \rangle \\ &+ \sum_{\lambda'} M_{\nu\lambda'} \langle O_1(t_1) A_\mu(t_2) O_3(t_1) A_{\lambda'}(t_2) \rangle \\ &+ \langle W_{1,2,3,4} \{ O_{1S}(t_1) F[A_{\mu S}(t_2), A_{\nu S}(t_2)] O_{3S}(t_1) \} \rangle \end{aligned} \quad (27)$$

where we assume $t_2 > t_1$. The above equation is almost identical to the regression theorem except for the last term. The explicit form of the operator $F[A_{\mu S}(t_2), A_{\nu S}(t_2)]$ is given in the appendix (Eq. (C5)). Once again, it is important to realize that this particular structure appears as a result of taking derivative with respect to t_2 . If we take derivative with respect to t_1 instead of t_2 , such regression type form does not appear. Eq.(25) and Eq.(27) are the central results of this paper.

Interestingly, in Ref. [21], a regression type theorem for OTOC was recently derived using a different approach. The central result obtained there is very similar to what we have derived in Eq.(27).

IV. EXAMPLE : DISSIPATIVE SPIN-BOSON MODEL

In this section we illustrate the above derived results for the paradigmatic dissipative spin-boson model by calculating various correlation functions in the Heisenberg picture. In particular, we verify the QRTs for two, three and four-point correlators. In order to perform this calculations, we first workout the master equation for reduced one point operator and then proceed to calculate the multi-time and multi-point correlation functions. The details of the derivations are provided in appendix D.

The Hamiltonian for a dissipative spin-1/2 system, coupled to a bath consisting of an infinite collection of harmonic oscillators with different normal mode frequencies, can be written as

$$\begin{aligned} H &= H_S + H_R + H_{SR} \\ &= \frac{\omega_0}{2} \sigma_z + \sum_k \omega_k b_k^\dagger b_k + \sum_k \alpha_k (b_k^\dagger \sigma_- + b_k \sigma_+) \end{aligned} \quad (28)$$

where ω_0 is the frequency of the qubit. The bath is characterized by the eigen-mode frequency ω_k referring to the k -th oscillator with $b_k (b_k^\dagger)$ is the corresponding annihilation (creation) operator. The last term in the above Hamiltonian represents the standard dissipative coupling term between the spin 1/2 system and the harmonic bath with α_k being the coupling strength between the k -th mode and the qubit. In what follows, for simplicity, we work in the zero-temperature limit ($T = 0$).

We first derive the master equation correct upto the second-order of the system-bath coupling and further

make secular approximation [6]. One can show that the reduced one point operator obeys the following equation at zero temperature $T = 0$

$$\frac{d}{dt}O_S(t) = i\frac{\omega'_0}{2}[\sigma_z, O_S(t)] + \frac{\gamma}{2}[2\sigma_+O_S(t)\sigma_- - \sigma_+\sigma_-O_S(t) - O_S(t)\sigma_+\sigma_-] \quad (29)$$

Where $\omega'_0 = \omega_0 + \Delta$ is the renormalized frequency of the qubit due to the coupling with the environment with

$$\Delta = \text{P.V.} \int_0^\infty \frac{g(\omega')|\alpha(\omega')|^2 d\omega'}{\omega_0 - \omega'}. \quad (30)$$

Here $g(\omega)$ represents the density of states of the bath oscillators and $\gamma = 2\pi g(\omega_0)|\alpha(\omega_0)|^2$ represents the net decay rate. Using the above master equation for the operator, it is easy to show that,

$$\frac{d}{dt} \begin{pmatrix} \sigma_{xS} \\ \sigma_{yS} \\ \sigma_{zS} \end{pmatrix} = \begin{pmatrix} -\frac{\gamma}{2} & -\omega'_0 & 0 \\ \omega'_0 & -\frac{\gamma}{2} & 0 \\ 0 & 0 & -\gamma \end{pmatrix} \begin{pmatrix} \sigma_{xS} \\ \sigma_{yS} \\ \sigma_{zS} \end{pmatrix} - \gamma \begin{pmatrix} 0_{2 \times 2} \\ 0_{2 \times 2} \\ I_{2 \times 2} \end{pmatrix} \quad (31)$$

where $0_{2 \times 2}$ and $I_{2 \times 2}$ are the 2×2 null matrix and the identity matrix, respectively. It is therefore easy to see that for this model there exists two different set of closed

operators that follow equation (8). These are given as,

$$\begin{aligned} A_\mu &= \{\sigma_z, I_{2 \times 2}\}, & M_{\mu\lambda} &= \begin{bmatrix} -\gamma & -\gamma \\ 0 & 0 \end{bmatrix} \\ A_\nu &= \{\sigma_x, \sigma_y\}, & M'_{\nu\lambda'} &= \begin{bmatrix} -\frac{\gamma}{2} & -\omega'_0 \\ \omega'_0 & -\frac{\gamma}{2} \end{bmatrix} \end{aligned} \quad (32)$$

With this in hand, we are now ready to asses the validity of QRT results that we derived in the previous section.

A. Two-point correlation function

Let us first verify the regression theorem for two-point function (13) for this model. To do that let us first set, $O_1 = \sigma_x$, $A_\mu = \{\sigma_x, \sigma_y\}$ in equation (13) and assume $t_1 < t_2$. Following the calculation in the Heisenberg picture for the two-point reduced operator, we receive

$$\left(\sigma_x(t_1)\sigma_x(t_2) \right)_S = \begin{pmatrix} (1 - \frac{\gamma}{2}(t_2 - t_1))e^{-i\omega'_0(t_2-t_1)} + 2i\gamma t_1 \sin \omega'_0(t_2 - t_1) & 0 \\ 0 & (1 - \frac{\gamma}{2}(t_2 - t_1))e^{i\omega'_0(t_2-t_1)} \end{pmatrix}, \quad (33)$$

$$\left(\sigma_x(t_1)\sigma_y(t_2) \right)_S = \begin{pmatrix} i(1 - \frac{\gamma}{2}(t_2 - t_1))e^{-i\omega'_0(t_2-t_1)} - 2i\gamma t_1 \cos \omega'_0(t_2 - t_1) & 0 \\ 0 & -i(1 - \frac{\gamma}{2}(t_2 - t_1))e^{i\omega'_0(t_2-t_1)} \end{pmatrix}, \quad (34)$$

and as a result the left hand side of the equation (13) gives

$$\begin{aligned} \frac{d}{dt_2} \left(\sigma_x(t_1)\sigma_x(t_2) \right)_S &= -\frac{\gamma}{2} \begin{pmatrix} e^{-i\omega'_0(t_2-t_1)} & 0 \\ 0 & e^{i\omega'_0(t_2-t_1)} \end{pmatrix} \\ &- \omega'_0 \begin{pmatrix} i(1 - \frac{\gamma}{2}(t_2 - t_1))e^{-i\omega'_0(t_2-t_1)} - 2i\gamma t_1 \cos \omega'_0(t_2 - t_1) & 0 \\ 0 & -i(1 - \frac{\gamma}{2}(t_2 - t_1))e^{i\omega'_0(t_2-t_1)} \end{pmatrix} \end{aligned} \quad (35)$$

Similarly, the right hand side of the equation (13) can be computed and we receive,

$$\begin{aligned} &-\frac{\gamma}{2} \left(\sigma_x(t_1)\sigma_x(t_2) \right)_S - \omega'_0 \left(\sigma_x(t_1)\sigma_y(t_2) \right)_S \\ &= -\frac{\gamma}{2} \begin{pmatrix} e^{-i\omega'_0(t_2-t_1)} & 0 \\ 0 & e^{i\omega'_0(t_2-t_1)} \end{pmatrix} \end{aligned} \quad (36)$$

$$- \omega'_0 \begin{pmatrix} i(1 - \frac{\gamma}{2}(t_2 - t_1))e^{-i\omega'_0(t_2-t_1)} - 2i\gamma t_1 \cos \omega'_0(t_2 - t_1) & 0 \\ 0 & -i(1 - \frac{\gamma}{2}(t_2 - t_1))e^{i\omega'_0(t_2-t_1)} \end{pmatrix} \quad (37)$$

These above two equations therefore shows the validity

of QRT at the level of two-point reduced operators. As

an immediate consequence, we conclude that the QRT follows for two-point correlation functions Eq. (14) for arbitrary initial density matrix for the system.

B. Three-point, multi-time correlation function

We next move to three-point correlation function as given in Eq. (21). We first set, $O_1 = O_2 = \sigma_x$, $A_\mu = \{\sigma_x, \sigma_y\}$ and assume $t_1 < t_2 < t_3$. The left hand side of the equation (21) can be computed following the Heisenberg picture (please see the details of the calculation in appendix D),

$$(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3))_S = \begin{pmatrix} 0 & (1 - \frac{1}{2}\gamma(t_1 + t_3 - t_2))e^{i\omega'_0(t_1+t_3-t_2)} \\ (1 - \frac{1}{2}\gamma(-t_1 + t_2 + t_3))e^{-i\omega'_0(t_1+t_3-t_2)} & 0 \\ +\gamma(t_2 - t_1)e^{i\omega'_0(-t_1-t_2+t_3)} & 0 \end{pmatrix} \quad (38)$$

and

$$(\sigma_x(t_1)\sigma_x(t_2)\sigma_y(t_3))_S = \begin{pmatrix} 0 & -i(1 - \frac{1}{2}\gamma(t_1 + t_3 - t_2))e^{i\omega'_0(t_1+t_3-t_2)} \\ i(1 - \frac{1}{2}\gamma(-t_1 + t_2 + t_3))e^{-i\omega'_0(t_1+t_3-t_2)} & 0 \\ -i\gamma(t_2 - t_1)e^{i\omega'_0(-t_1-t_2+t_3)} & 0 \end{pmatrix} \quad (39)$$

Now taking derivative of the first equation with respect to the maximum time t_3 , we get,

$$\begin{aligned} \frac{d}{dt_3}(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3))_S &= -\frac{1}{2}\gamma \begin{pmatrix} 0 & e^{i\omega'_0(t_1+t_3-t_2)} \\ e^{-i\omega'_0(t_1+t_3-t_2)} & 0 \end{pmatrix} \\ &- \omega'_0 \begin{pmatrix} 0 & -i(1 - \frac{1}{2}\gamma(t_1 + t_3 - t_2))e^{i\omega'_0(t_1+t_3-t_2)} \\ i(1 - \frac{1}{2}\gamma(-t_1 + t_2 + t_3))e^{-i\omega'_0(t_1+t_3-t_2)} & 0 \\ -i\gamma(t_2 - t_1)e^{i\omega'_0(-t_1-t_2+t_3)} & 0 \end{pmatrix} \end{aligned} \quad (40)$$

Now, one can calculate the RHS of (21) and obtain,

$$\begin{aligned} &-\frac{1}{2}\gamma(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3))_S - \omega'_0(\sigma_x(t_1)\sigma_x(t_2)\sigma_y(t_3))_S \\ &= -\frac{1}{2}\gamma \begin{pmatrix} 0 & e^{i\omega'_0(t_1+t_3-t_2)} \\ e^{-i\omega'_0(t_1+t_3-t_2)} & 0 \end{pmatrix} \\ &- \omega'_0 \begin{pmatrix} 0 & -i(1 - \frac{1}{2}\gamma(t_1 + t_3 - t_2))e^{i\omega'_0(t_1+t_3-t_2)} \\ i(1 - \frac{1}{2}\gamma(-t_1 + t_2 + t_3))e^{-i\omega'_0(t_1+t_3-t_2)} & 0 \\ -i\gamma(t_2 - t_1)e^{i\omega'_0(-t_1-t_2+t_3)} & 0 \end{pmatrix} \end{aligned} \quad (41)$$

Equations (40) and (41) gives the identical result, this verifies the regression theorem (21). Note that, we choose a particular order of time but note that we can show the equation (21) holds as long as $t_i < t_3$ with $i = 1, 2$. There are no constrains on the order of (t_1, t_2) .

C. Four-point, multi-time correlation function

Now we want to verify the regression theorem for four-point function (24) in this example. To do that let us

first set, $O_1 = O_2 = O_3 = \sigma_x$, $A_\mu = \{\sigma_x, \sigma_y\}$ in equation (24) and assume $t_1 < t_2 < t_3 < t_4$. We then compute the following reduced operators which turns out to be diagonal. More explicitly,

$$(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3)\sigma_x(t_4))_S = \begin{pmatrix} (1 - \frac{1}{2}\gamma(t_1 + t_3 - t_2 - t_4))e^{i\omega'_0(t_1+t_3-t_2-t_4)} \\ + \gamma t_1 e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \\ + \gamma(t_3 - t_2)e^{i\omega'_0(t_1-t_2-t_3+t_4)} \\ 0 \\ 0 \\ (1 - \frac{1}{2}\gamma(t_1 + t_2 + t_3 + t_4))e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \\ + \gamma t_1 e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \\ + \gamma t_3 e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \end{pmatrix} \quad (42)$$

and,

$$(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3)\sigma_y(t_4))_S = \begin{pmatrix} i(1 - \frac{1}{2}\gamma(t_1 + t_3 - t_2 - t_4))e^{i\omega'_0(t_1+t_3-t_2-t_4)} \\ -i\gamma t_1 e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \\ -i\gamma(t_3 - t_2)e^{i\omega'_0(t_1-t_2-t_3+t_4)} \\ 0 \\ 0 \\ -i(1 - \frac{1}{2}\gamma(t_1 + t_2 + t_3 + t_4))e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \\ -i\gamma t_1 e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \\ -i + \gamma t_3 e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \end{pmatrix} \quad (43)$$

Taking derivative of the equation (42) with respect to t_4 , we receive the QRT as,

$$\begin{aligned} \frac{d}{dt_4}(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3)\sigma_x(t_4))_S &= -\frac{1}{2}\gamma(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3))\sigma_x(t_4)_S \\ &\quad - \omega'_0(\sigma_x(t_1)\sigma_x(t_2)\sigma_x(t_3)\sigma_y(t_4))_S \end{aligned} \quad (44)$$

This immediately verifies equation (24). Note that, we choose here a particular order of time but note that one can show the validity of the QRT as long as $t_i < t_4$ with $i = 1, 2, 3$. There are no constrains on the order of (t_1, t_2, t_3) .

D. Verification of OTOC

We next provide one example to asses the validity of our expression for OTOC. For the spin-boson model, it is easy to compute the following four-point reduced operator and one receives,

$$\frac{d}{dt_2}(\sigma_x(t_1)\sigma_z(t_2)\sigma_x(t_1)\sigma_z(t_2))_S = 2\gamma \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (45)$$

It is easy to check that the corresponding RHS of the OTOC also gives the same result. In a similar way, OTOC can be checked for

$$\frac{d}{dt_2}(\sigma_z(t_1)\sigma_z(t_2)\sigma_z(t_1)\sigma_z(t_2))_S = -8\gamma \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \quad (46)$$

V. GENERALIZATION OF QRT FOR NON-MARKOVIAN CASE

The results for QRT presented in the previous sections can be extended for the non-Markovian case. For simplicity, We here focus on systems with bosonic bath and

linear system-bath interaction but one can generalize this study for a more generic type of system-bath interaction as well. We derive a Lindblad type equation upto order λ^2 for this setup which takes into account the non-Markovian evolution. We then derive the correction to the QRT for the non-Markovian case by focusing only on the two-point correlation functions [14]. Extension to higher point multi-time correlators can be similarly obtained even for the non-Markovian case.

A. Lindblad type non-Markovian equation for one-point reduced operator

Let the Hamiltonian of the composite system+ bath is,

$$\begin{aligned} H &= H_S + H_R + H_{SR} \\ &= H_S + \sum \omega_k b_k^\dagger b_k + \sum g_k (L b_k^\dagger + L^\dagger b_k). \end{aligned} \quad (47)$$

Here, the system is coupled with the bath through a generic system operator L . Let us assume that the initial density operator of the total system can be written as, $\rho_{SR}(0) = \rho_S \otimes \rho_R$. Now following the master equation in the Heisenberg picture, (please see equation (A10)), one can show that the reduced density operator for the system obeys the following non-Markovian master equation at zero temperature ($T = 0$) and is correct up to the

second order of system-bath coupling,

$$\begin{aligned} \frac{d}{dt}O_S(t) &= i[H_S, O_S(t)] + \int_0^t d\tau \alpha(\tau) [L^\dagger(0), O_S(t)] \tilde{L}(-\tau) \\ &+ \int_0^t d\tau \alpha^*(\tau) \tilde{L}^\dagger(-\tau) [O_S(t), L(0)], \end{aligned} \quad (48)$$

where, $\tilde{L}(t) = U_0(t)LU_0^\dagger(t)$ is the coupled system operator in the interaction picture with $U_0(t)$ represents the free evolution due to the Hamiltonian H_S . $\alpha(\tau)$ denotes the bath correlation function which is given as,

$$\alpha(\tau) = \sum_k |g_k|^2 \text{Tr}_R [\tilde{b}_k(0)\tilde{b}_k^\dagger(-\tau)\rho_R] = \sum_k |g_k|^2 e^{-i\omega_k\tau}. \quad (49)$$

B. Extension of QRT to non-Markovian case

Having obtained the non-Markovian master equation in the schrödinger picture, we now extend the QRT for the non-Markovian dynamics. To achieve that let us first assume that there exists a complete set of system operators $A_{\mu S}(t)$ such that

$$\frac{d}{dt}A_{\mu S}(t) = \sum_\lambda M_{\mu\lambda}(t) A_{\lambda S}(t) \quad (50)$$

Note the crucial explicit time dependence in $M_{\mu\lambda}(t)$ for the non-Markovian case which was independent of time for the Markovian dynamics. Using the above results, it is easy to derive QRT for the non-Markovian case and is given by (see appendix E for derivation)

$$\begin{aligned} \frac{d}{dt_1} (A_\mu(t_1)O(t_2))_S &= \sum_\lambda M_{\mu\lambda}(t_1) (A_\lambda(t_1)O(t_2))_S \\ &- \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) A_{\mu S}(t_1) \tilde{L}^\dagger(-t_1) O_S(t_2) \tilde{L}(-\tau_2) \\ &- \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) \tilde{L}^\dagger(-t_1) A_{\mu S}(t_1) \tilde{L}(-\tau_2) O_S(t_2) \\ &+ \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) \tilde{L}^\dagger(-t_1) A_{\mu S}(t_1) O_S(t_2) \tilde{L}(-\tau_2) \\ &+ \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) A_{\mu S}(t_1) \tilde{L}^\dagger(-t_1) \tilde{L}(-\tau_2) O_S(t_2) \end{aligned} \quad (51)$$

where we have assumed $t_1 > t_2$ and that the operator A_μ satisfies Eq. (50). The above equation is the extension of QRT to the non-Markovian dynamics. In the Markovian

limit Eq. (51) correctly reproduce the standard QRT, as given in Eq. (11). Note that, In the Markovian limit, the bath correlation function $\alpha(\tau_2 - t_1)$ is non-vanishing only if τ_2 lies in the range from $t_1 - \tau_B$ to $t_1 + \tau_B$ where τ_B corresponds to a characteristic time scale of the bath dynamics. However, the upper limit of the integration in equation (51) is t_2 which is always less than t_1 . This implies the bath correlation function, $\alpha(\tau_2 - t_1) \rightarrow 0$ in the given range of τ_2 . Eq. (51) thus reduces to the QRT for two-point function with the last four terms disappear in the Markovian limit.

VI. DISCUSSION

While defining the multi-time correlation function, it is natural to work in the Heisenberg picture. Even though, multi-time correlation function is of great physical importance, Heisenberg picture has not got that much attention in the context of quantum open system. In this paper we make use of recently developed Heisenberg Picture technique [22] to calculate correlation functions in the Markovian limit and derive quantum regression theorem. In particular, we generalize the regression theorem for multi-time correlation functions with general time arguments. What we observe is that that the form of regression theorem remains the same for two or multi-time correlation functions as long as some mild restriction on the time arrangements is met. We also derived regression theorem for OTOC in the Markovian limit and find that regression theorem gets modification from the known two-time regression theorem. We further extend our study to the non-Markovian dynamics. However in this case the QRT receives a complicated correction term with two-point correlation function requiring information about four-point function.

As a possible future direction, one of the interesting problem would be to use the Heisenberg picture [22] and to go beyond the standard second-order perturbation scheme. For example, one can consider exactly solvable systems such as the spin-boson dephasing model, or the well-known Caldeira-Leggett model and investigate the possibility to sum up the perturbation series exactly in the Heisenberg picture.

Acknowledgements: Work of SJ is supported by Ramanujan Fellowship. Work of SK is supported by CSIR fellowship with Grant Number 09/0936(11643)/2021-EMR-I. BKA acknowledges the MATRICS grant MTR/2020/000472 from SERB, Government of India. BKA also thanks the Shastri Indo-Canadian Institute for providing financial support for this research work in the form of a Shastri Institutional Collaborative Research Grant (SICRG). The authors would also like to thank the people of India for their steady support in basic research.

[1] A. Smirne, N. Megier, and B. Vacchini, *Quantum* **5**, 439 (2021).

[2] E. B. Davies, *Quantum Theory of Open Systems*. (Aca-

- demec Press, London, 1976).
- [3] R. Alicki and K. Lendi, *Quantum Dynamical Semigroups and Applications*. (Springer, Berlin, 1987).
- [4] H. J. Carmichael, *An Open System Approach to Quantum Optics, Lecture Notes in Physics* (Springer, 1993).
- [5] C. W. Gardiner and P. Zoller, *Quantum Noise* (Springer, 2000).
- [6] H. Breuer and F. Petruccione., *The Theory of Open Quantum Systems*. (Oxford University Press, 2002).
- [7] M. Cosacchi, T. Seidelmann, M. Cygorek, A. Vagov, D. E. Reiter, and V. M. Axt, *Phys. Rev. Lett.* **127**, 100402 (2021).
- [8] M. Ban, *Phys. Rev. A* **96**, 042111 (2017).
- [9] B. Bellomo, R. Lo Franco, and G. Compagno, *Phys. Rev. Lett.* **99**, 160502 (2007).
- [10] K.-L. Liu and H.-S. Goan, *Phys. Rev. A* **76**, 022312 (2007).
- [11] B. Wang, Z.-Y. Xu, Z.-Q. Chen, and M. Feng, *Phys. Rev. A* **81**, 014101 (2010).
- [12] J.-H. An and W.-M. Zhang, *Phys. Rev. A* **76**, 042127 (2007).
- [13] G. Guarnieri, A. Smirne, and B. Vacchini, *Phys. Rev. A* **90**, 022110 (2014).
- [14] D. Alonso and I. de Vega, *Phys. Rev. A* **75**, 052108 (2007).
- [15] M. Ban, *Phys. Rev. A* **96**, 042111 (2017).
- [16] H.-S. Goan, C.-C. Jian, and P.-W. Chen, *Phys. Rev. A* **82**, 012111 (2010).
- [17] H.-P. Breuer, E.-M. Laine, J. Piilo, and B. Vacchini, *Rev. Mod. Phys.* **88**, 021002 (2016).
- [18] H.-P. Breuer, **45**, 154001 (2012).
- [19] Á. Rivas, S. F. Huelga, and M. B. Plenio, **77**, 094001 (2014).
- [20] OTOC is defined as $\langle O_1(t_1)O_2(t_2)O_3(t_1)O_4(t_2) \rangle$. Even though it's a two-time correlator it's form significantly differ from equation (3).
- [21] P. D. Blocher and K. Mølmer, *Phys. Rev. A* **99**, 033816 (2019).
- [22] N. Karve and R. Loganayagam, (2020), [arXiv:2011.15118 \[quant-ph\]](https://arxiv.org/abs/2011.15118).
- [23] In one of the examples discussed later, we explicitly construct operator A_μ .
- [24] See equation (A16) and (A25)-(A27) in the appendix A 2 for details.
- [25] A. I. Larkin and Y. N. Ovchinnikov, *Soviet Journal of Experimental and Theoretical Physics* **28**, 1200 (1969).
- [26] S. V. Syzranov, A. V. Gorshkov, and V. Galitski, *Phys. Rev. B* **97**, 161114 (2018).
- [27] B. Swingle and D. Chowdhury, *Phys. Rev. B* **95**, 060201 (2017).
- [28] I. L. Aleiner, L. Faoro, and L. B. Ioffe, *Annals of Physics* **375**, 378 (2016).
- [29] E. B. Rozenbaum, S. Ganeshan, and V. Galitski, *Phys. Rev. Lett.* **118**, 086801 (2017).
- [30] Y. Chen, "Universal logarithmic scrambling in many body localization," (2016), [arXiv:1608.02765 \[cond-mat.dis-nn\]](https://arxiv.org/abs/1608.02765).
- [31] Y. Huang, Y.-L. Zhang, and X. Chen, *Annalen der Physik* **529**, 1600318, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/andp.201600318>.
- [32] A. A. Patel, D. Chowdhury, S. Sachdev, and B. Swingle, *Phys. Rev. X* **7**, 031047 (2017).
- [33] G. Styliaris, N. Anand, and P. Zanardi, *Phys. Rev. Lett.* **126**, 030601 (2021).
- [34] S. V. Syzranov, A. V. Gorshkov, and V. Galitski, *Phys. Rev. B* **97**, 161114 (2018).
- [35] We can explicitly express the one-point image operator in terms of one point reduced operator[22].
- [36] Throughout the paper we are going to use the Einstein summation convention that repeated indices are summed.
- [37] It can't be expressed simply as the multiplication of two one pint reduced operator but it's a function of one point reduced operator and it starts from λ^2 order.
- [38] With out loss of generality, we use $t_2 > t_1$ for rest of the discussion.
- [39] We shall see that this observation is true for general multi time correlation function. In Schrodinger picture as well, we have seen that regression theorem holds only for derivatives with respect to highest time.

Appendix A: Review of Heisenberg picture dynamics

In this Appendix for completeness, we briefly review the Heisenberg picture results as obtained by [22]. We follow the notation and convention introduced in this reference. Let the Hamiltonian of the total system is $H = H_S + H_R + \lambda H_{SR}$, where we introduce an extra parameter λ to keep track of the order of perturbation in terms of system-bath coupling. Let us assume that the interaction between system and bath is turned on at $t = 0$. Before turning the interaction, system and bath were decoupled and their total density matrix can be written as, $\rho_{SR} = \rho_S \otimes \rho_R$. In the Heisenberg picture the density matrix is time independent. The expectation value of any operator can be written as,

$$\langle O(t) \rangle = \text{Tr}_S[\text{Tr}_R[O(t)\rho_R]\rho_S] = \text{Tr}_S[O_S(t)\rho_S] \quad (\text{A1})$$

where the reduced one point operator define as, $O_S(t) = \text{Tr}_R[O(t)\rho_R]$. Similarly, we can define N -point reduced operator as

$$(O_1(t_1)O_2(t_2)\dots O_N(t_N))_S = \text{Tr}_R[O_1(t_1)O_2(t_2)\dots O_N(t_N)\rho_R]. \quad (\text{A2})$$

This definition has a property that $(O_1(t_1)O_2(t_2))_S \neq O_{1S}(t_1)O_{2S}(t_2)$ However, we can express the reduced N -point operator in terms of one-point reduced operators using what are called image operators [22]. The image operator of any operator $O(t)$ is defined as,

$$O_{\alpha\beta}(t) = T_\alpha^\dagger O(t) T_\beta \quad (\text{A3})$$

where, $T_\alpha = |i\alpha\rangle \langle i|$ where $\{|i\rangle\}$ are orthonormal basis of H_S and $\{|\alpha\rangle\}$ are orthonormal basis of H_R . It is easy to see that

$$T_\alpha T_\alpha^\dagger = 1. \quad (\text{A4})$$

One can show that the N -point image operators can be written in terms of one-point image operators as

$$(O_1(t_1)O_2(t_2)\dots O_N(t_N))_{\alpha\beta} = O_{1\alpha\gamma_1}(t_1)O_{2\gamma_1\gamma_2}(t_2)\dots O_{N\gamma_{N-1}\beta}(t_N). \quad (\text{A5})$$

The N -point reduced operators defined in (A2) can also be expressed in terms of one point image operators by inserting (A4) and using (A5) as follows[35]

$$(O_1(t_1)O_2(t_2)\dots O_N(t_N))_S = O_{1\alpha\gamma_1}(t_1)O_{2\gamma_1\gamma_2}(t_2)\dots O_{N\gamma_{N-1}\beta}(t_N)\rho_{R\beta\alpha} \quad (\text{A6})$$

We next consider a general form of the interaction Hamiltonian between system and bath and write

$$H_{SR} = \sum_i S^i \otimes R^i \quad (\text{A7})$$

where, S^i is a Hermitian operator acting on the system's Hilbert space, and R^i is a hermitian operator in the bath's Hilbert space. The corresponding image operators of H_{SR} are (using equation (A3)):

$$H_{SR\alpha\gamma} = \sum_i S^i R_{\alpha\gamma}^i, \quad (\text{A8})$$

Here, $R_{\alpha\gamma}^i$ is the α, γ th element of R^i ($|\alpha\rangle, |\gamma\rangle$ are the eigenstates of bath Hamiltonian H_R). We define the interaction picture operators $\tilde{H}_{SR\alpha\gamma}(t)$ as,

$$\tilde{H}_{SR\alpha\gamma}(t) = \sum_i \tilde{S}^i(t) \tilde{R}_{\alpha\gamma}^i(t), \quad (\text{A9})$$

where, $\tilde{S}^i(t) = U_0(t)S^iU_0^\dagger(t) = \sum_\omega S_\omega^i e^{i\omega t}$ and $\tilde{R}_{\alpha\gamma}^i(t) = R_{\alpha\gamma}^i e^{-i(E_\alpha - E_\gamma)t}$; here $U_0(t) = e^{-iH_S t}$ and E_α, E_γ are the eigenvalues of bath Hamiltonian H_R . Then the exact equation (written to all orders in λ) that satisfies the reduced one point system operator $O_S(t)$ is[22],

$$\frac{d}{dt}O_S(t) = i[H_S, O_S(t)] + \sum_{n=1}^{\infty} \sum_{k=0}^{\infty} \sum_{n_1, n_2, \dots, n_k=1}^{\infty} (-1)^k \lambda^{n+n_1+\dots+n_k} D_t P_s^n P_s^{n_1} \dots P_s^{n_k} O_S(t) \quad (\text{A10})$$

where the super-operator $D_t P_s^n$ is defined as [36]

$$\begin{aligned}
D_t P_s^n A(t) &= \sum_{r=0}^n i^{n-2r} U_0^\dagger(t) \frac{d}{dt} [U_0(t) K_{\gamma\alpha}^{(n-r)\dagger}(t) U_0^\dagger(t)] U_0(t) A(t) K_{\gamma\beta}^r(t) \rho_{B\beta\alpha} \\
&+ \sum_{r=0}^n i^{n-2r} K_{\gamma\alpha}^{(n-r)\dagger}(t) A(t) U_0^\dagger(t) \frac{d}{dt} [U_0(t) K_{\gamma\beta}^r(t) U_0^\dagger(t)] U_0(t) \rho_{B\beta\alpha}
\end{aligned} \tag{A11}$$

here,

$$\begin{aligned}
K_{\alpha\beta}^r(t) &= e^{i(E_\alpha - E_\beta)t} U_0^\dagger(t) \tilde{K}_{\alpha\beta}^r(t) U_0(t), \\
\tilde{K}_{\alpha\beta}^n(t) &= \int_0^t dt_1 \dots \int_0^{t_{n-1}} dt_n \tilde{H}_{SR\alpha\gamma_1}(t_1) \dots \tilde{H}_{SR\gamma_{n-1}\beta}(t_n).
\end{aligned} \tag{A12}$$

with

$$\tilde{K}_{\alpha\beta}^0(t) = \delta_{\alpha\beta}. \tag{A13}$$

1. Correlation function in the Heisenberg picture

a. Expression for two-point reduced operators

We now compute the two-point reduced operator in the Heisenberg picture which can be written as [22]:

$$(O_1(t_1) O_2(t_2))_S = O_{1S}(t_1) O_{2S}(t_2) + I[O_{1S}(t_1), O_{2S}(t_2)] \tag{A14}$$

$$\begin{aligned}
I[O_{1S}(t_1), O_{2S}(t_2)] &= \sum_{n_0^1, l_0^1, n_0^2, l_0^2} \left((-i\lambda)^{n_0^1} K_{\gamma_0\alpha}^{n_0^1}(t_1) \right)^\dagger O_{1S}(t_1) \left((-i\lambda)^{l_0^1} K_{\gamma_0\gamma}^{l_0^1}(t_1) \right) \\
&\left((-i\lambda)^{n_0^2} K_{\gamma_0\gamma}^{n_0^2}(t_2) \right)^\dagger O_{2S}(t_2) \left((-i\lambda)^{l_0^2} K_{\gamma_0\beta}^{l_0^2}(t_2) \right) \rho_{R\beta\alpha}
\end{aligned} \tag{A15}$$

such that $n_0^1 + l_0^1 = 1$ and $n_0^2 + l_0^2 = 1$, so these are the following four possible combinations,

1. $n_0^1 = 0, l_0^1 = 1$ and $n_0^2 = 0, l_0^2 = 1$
2. $n_0^1 = 1, l_0^1 = 0$ and $n_0^2 = 0, l_0^2 = 1$
3. $n_0^1 = 0, l_0^1 = 1$ and $n_0^2 = 1, l_0^2 = 0$
4. $n_0^1 = 1, l_0^1 = 0$ and $n_0^2 = 1, l_0^2 = 0$

Where $I[O_{1S}(t_1), O_{2S}(t_2)]$ is the irreducible part[37]. This irreducible part can be expressed upto λ^2 order, following Eq.(A6) as,

We can then write I as the sum over these four combinations i.e.,

$$I = I_1 + I_2 + I_3 + I_4. \tag{A16}$$

One can write down the expressions for each I_i . For example, using (A12), (A13), (A15), we obtain

$$I_1 = -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} O_{1S}(t_1) S_\omega^i O_{2S}(t_2) S_{\omega'}^j \int_0^{t_1} d\tau_1 e^{-i\omega\tau_1} \int_0^{t_2} d\tau_2 e^{-i\omega'\tau_2} Tr_R[\tilde{R}^i(-\tau_1) \tilde{R}^j(-\tau_2) \rho_R] \tag{A17}$$

where, $\tau_1 = t_1 - t'_1$ and $\tau_2 = t_2 - t'_2$.

Similarly, we can show that the other contributions

gives,

$$\begin{aligned}
I_2 &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} S_{\omega}^{i\dagger} O_{1S}(t_1) O_{2S}(t_2) S_{\omega'}^{j\dagger} \int_0^{t_1} d\tau_1 e^{i\omega\tau_1} \int_0^{t_2} d\tau_2 e^{-i\omega'\tau_2} \text{Tr}_R[\tilde{R}^i(-\tau_1) \tilde{R}^j(-\tau_2) \rho_R] \\
I_3 &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} O_{1S}(t_1) S_{\omega}^i O_{2S}(t_2) S_{\omega'}^{j\dagger} \int_0^{t_1} d\tau_1 e^{-i\omega\tau_1} \int_0^{t_2} d\tau_2 e^{i\omega'\tau_2} \text{Tr}_R[\tilde{R}^i(-\tau_1) \tilde{R}^j(-\tau_2) \rho_R] \\
I_4 &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} S_{\omega}^{i\dagger} O_{1S}(t_1) S_{\omega'}^{j\dagger} O_{2S}(t_2) \int_0^{t_1} d\tau_1 e^{i\omega\tau_1} \int_0^{t_2} d\tau_2 e^{i\omega'\tau_2} \text{Tr}_R[\tilde{R}^i(-\tau_1) \tilde{R}^j(-\tau_2) \rho_R]
\end{aligned} \tag{A18}$$

Note that, all these expressions are correct upto order λ^2 and is valid for general dynamics.

b. Expression for three-point reduced operator

As before, we can work out expressions for three point reduced operator upto λ^2 order. We receive [22]

$$\begin{aligned}
&(O_1(t_1)O_2(t_2)O_3(t_3))_S \\
&= O_{1S}(t_1)O_{2S}(t_2)O_{3S}(t_3) \\
&+ W_{1,2,3} \left\{ I[O_{1S}(t_1), O_{2S}(t_2)]O_{3S}(t_3) \right\} \\
&+ W_{1,2,3} \left\{ O_{1S}(t_1)I[O_{2S}(t_2), O_{3S}(t_3)] \right\} \\
&+ W_{1,2,3} \left\{ I[O_{1S}(t_1), O_{3S}(t_3)]O_{2S}(t_2) \right\}
\end{aligned} \tag{A19}$$

The operator $W_{1,2,3}$ makes sure that the operator product is ordered such that O_{1S} comes before O_{2S} , and O_{2S}

comes before O_{3S} . Let us illustrate this by one example. Considering the last term of the above equation (A19) we get,

$$\begin{aligned}
&W_{1,2,3} \left\{ I[O_{1S}(t_1), O_{3S}(t_3)]O_{2S}(t_2) \right\} \\
&= \lambda^2 \sum_{n_0^1, l_0^1, n_0^2, l_0^2} i^{n_0^1+n_0^2-l_0^1-l_0^2} \left(K_{\gamma_0\alpha}^{n_0^1}(t_1) \right)^\dagger O_{1S}(t_1) K_{\gamma_0\gamma}^{l_0^1}(t_1) O_{2S}(t_2) \\
&\quad * \left(K_{\gamma_0\gamma}^{n_0^2}(t_3) \right)^\dagger O_{3S}(t_3) K_{\gamma_0\beta}^{l_0^2}(t_3) \rho_{R\beta\alpha}
\end{aligned} \tag{A20}$$

such that $n_0^1 + l_0^1 = 1$ and $n_0^2 + l_0^2 = 1$.

c. Expression for four-point reduced operators

We can now calculate the four point function as well. The four-point reduced operator (upto λ^2 order) is [22],

$$\begin{aligned}
&(O_1(t_1)O_2(t_2)O_3(t_3)O_4(t_4))_S \\
&= O_{1S}(t_1)O_{2S}(t_2)O_{3S}(t_3)O_{4S}(t_4) + W_{1,2,3,4} \left\{ I[O_{1S}(t_1), O_{2S}(t_2)]O_{3S}(t_3)O_{4S}(t_4) \right\} \\
&+ W_{1,2,3,4} \left\{ O_{1S}(t_1)I[O_{2S}(t_2), O_{3S}(t_3)]O_{4S}(t_4) \right\} + W_{1,2,3,4} \left\{ I[O_{1S}(t_1), O_{3S}(t_3)]O_{2S}(t_2)O_{4S}(t_4) \right\} \\
&+ W_{1,2,3,4} \left\{ I[O_{4S}(t_4), O_{3S}(t_3)]O_{2S}(t_2)O_{1S}(t_1) \right\} + W_{1,2,3,4} \left\{ I[O_{1S}(t_1), O_{4S}(t_4)]O_{3S}(t_3)O_{2S}(t_2) \right\} \\
&+ W_{1,2,3,4} \left\{ O_{1S}(t_1)I[O_{2S}(t_2), O_{4S}(t_4)]O_{3S}(t_3) \right\}
\end{aligned} \tag{A21}$$

where functions W and I has the same property as is already discussed previously. For our purpose we need to calculate the explicit form of the function I . Below we give an explicit example.

2. Results in the Markovian limit

In the Markovian limit [22] the bath correlation functions i.e., $\text{Tr}_R[\tilde{R}(t)\tilde{R}(t-\tau)\rho_R]$ is a rapidly decaying function of τ only, using this property we can show that the general equation (A10) reduces to the well-known master

equation for reduced operator [22]

$$\begin{aligned} \frac{d}{dt}O_S(t) &= iH_S O_S(t) \\ &+ (i\lambda)^2 \sum_{\omega, \omega'} \sum_{i, j} J^{ij}(\omega) [S_\omega^{i\dagger} S_{\omega'}^j O_S(t) - S_\omega^{i\dagger} O_S(t) S_{\omega'}^j] + h.c. \end{aligned} \quad (\text{A22})$$

where $J^{ij}(\omega)$ is the Fourier transformation of the bath correlation functions and S_ω^i is the Fourier decomposition $\tilde{S}^i(t)$ and are given as,

$$\begin{aligned} J^{ij}(\omega) &= \int_0^\infty d\tau e^{-i\omega\tau} \text{Tr}_R[\tilde{R}^i(0)\tilde{R}^j(-\tau)\rho_R] \\ \tilde{S}^i(t) &= \sum_\omega S_\omega^i e^{i\omega t} \end{aligned} \quad (\text{A23})$$

One can simplify further the expressions for the I in the Markovian limit. Let us first analyse Eq. (A17) in this limit. As in the Markovian limit the bath correlation functions i.e., $\text{Tr}_R[\tilde{R}(-\tau_1)\tilde{R}(-\tau_2)\rho_R]$ is a rapidly decaying function of $\tau_2 - \tau_1$ only, this implies, we can ignore the bath correlation after a characteristic time scale τ_B , determined by bath dynamics. Using this fact, it becomes easy to analyse equation (A17). Let us first calculate the τ_2 integration by assuming $t_2 > t_1$. [38] Significant contribution to the integral will come with in the range $|\tau_2 - \tau_1| \leq \tau_B$ which gives $\tau_1 - \tau_B \leq \tau_2 \leq \tau_B + \tau_1$. Using this fact, we can write equation (A17) as

$$\begin{aligned} I_1 &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} O_{1S}(t_1) S_\omega^i O_{2S}(t_2) S_{\omega'}^j \int_0^{t_1} d\tau_1 e^{-i\omega\tau_1} \int_{\tau_1 - \tau_B}^{\tau_1 + \tau_B} d\tau_2 e^{-i\omega'\tau_2} \text{Tr}_R[\tilde{R}^i(-\tau_1)\tilde{R}^j(-\tau_2)\rho_R] \\ &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} O_{1S}(t_1) S_\omega^i O_{2S}(t_2) S_{\omega'}^j \int_0^{t_1} d\tau_1 e^{-i\omega\tau_1} \int_{-\tau_B}^{\tau_B} d\tau e^{-i\omega'(\tau_1 - \tau)} \text{Tr}_R[\tilde{R}^i(0)\tilde{R}^j(\tau)\rho_R] \end{aligned} \quad (\text{A24})$$

where in the last line we have used the variable $\tau = \tau_1 - \tau_2$ to rewrite the integral. In the Markovian limit,

the correlation function decays very fast beyond τ_B , this implies that the integration limit in the last line of (A24) can be extended to infinity. This gives

$$\begin{aligned} I_1 &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} O_{1S}(t_1) S_\omega^i O_{2S}(t_2) S_{\omega'}^j \int_0^{t_1} d\tau_1 e^{-i(\omega + \omega')\tau_1} \int_{-\infty}^{\infty} d\tau e^{i\omega'\tau} \text{Tr}_R[\tilde{R}^i(0)\tilde{R}^j(\tau)\rho_R] \\ &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} O_{1S}(t_1) S_\omega^i O_{2S}(t_2) S_{\omega'}^j \alpha_1(\omega, \omega', t_1) \beta_1^{ij}(\omega') \end{aligned} \quad (\text{A25})$$

where in the last line we have defined α_1 and β_1 for simplicity and their explicit expression is given below (A27). By following the identical steps we can find I_2, I_3 and I_4

where,

$$\begin{aligned} I_2 &= \lambda^2 \sum_{\omega, \omega'} \sum_{i, j} S_\omega^{i\dagger} O_{1S}(t_1) O_{2S}(t_2) S_{\omega'}^j \alpha_2(\omega, \omega', t_1) \beta_2^{ij}(\omega'), \\ I_3 &= \lambda^2 \sum_{\omega, \omega'} \sum_{i, j} O_{1S}(t_1) S_\omega^i S_{\omega'}^{j\dagger} O_{2S}(t_2) \alpha_3(\omega, \omega', t_1) \beta_3^{ij}(\omega'), \\ I_4 &= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} S_\omega^{i\dagger} O_{1S}(t_1) S_{\omega'}^{j\dagger} O_{2S}(t_2) \alpha_4(\omega, \omega', t_1) \beta_4^{ij}(\omega'). \end{aligned} \quad (\text{A26})$$

$$\begin{aligned}
\alpha_1(\omega, \omega', t_1) &= \int_0^{t_1} d\tau_1 e^{-i(\omega+\omega')\tau_1} & \beta_1^{ij}(\omega') &= \int_{-\infty}^{\infty} d\tau e^{i\omega'\tau} \text{Tr}_R[\tilde{R}^i(0)\tilde{R}^j(\tau)\rho_R] \\
\alpha_2(\omega, \omega', t_1) &= \int_0^{t_1} d\tau_1 e^{i(\omega-\omega')\tau_1} & \beta_3^{ij}(\omega') &= \int_{-\infty}^{\infty} d\tau e^{-i\omega'\tau} \text{Tr}_R[\tilde{R}^i(0)\tilde{R}^j(\tau)\rho_R] \\
\alpha_3(\omega, \omega', t_1) &= \int_0^{t_1} d\tau_1 e^{-i(\omega-\omega')\tau_1} & \beta_2^{ij}(\omega') &= \beta_1^{ij}(\omega') \\
\alpha_4(\omega, \omega', t_1) &= \int_0^{t_1} d\tau_1 e^{i(\omega+\omega')\tau_1} & \beta_4^{ij}(\omega') &= \beta_3^{ij}(\omega')
\end{aligned} \tag{A27}$$

Using equation (A25), (A26) we get the explicit form of the irreducible part I ((A14)) in the Markovian limit.

We can further simplify the expression of α 's defined in equation (A27) using the secular approximation. Let us first substitute, $s_1 = \lambda^2\tau_1$ and $\sigma = \lambda^2t_1$ in the expression of α_1 define in (A27),

$$\alpha_1(\omega, \omega', t_1) = \lambda^2 \int_0^{t_1} d\tau_1 e^{i(\omega+\omega')\tau_1} = \lim_{\lambda \rightarrow 0} \int_0^{\sigma} ds_1 e^{i\frac{(\omega+\omega')}{\lambda^2}s_1} \tag{A28}$$

Now the Riemann-Lebesgue lemma states that, if $f(t)$ is an integrable function in $[a, b]$ then

$$\lim_{x \rightarrow \infty} \int_a^b dt e^{ixt} f(t) = 0. \tag{A29}$$

By comparing the equation (A28) with the Riemann-Lebesgue lemma (A29), we can conclude that α_1 is non-zero only when $(\omega + \omega') = 0$ i.e. $\alpha_1(\omega, \omega', t_1) = \lambda^2 t_1 \delta_{\omega, -\omega'}$. Similarly, we can show that, $\alpha_4(\omega, \omega', t_1) = \alpha_1(\omega, \omega', t_1) = \lambda^2 t_1 \delta_{\omega, -\omega'}$ and $\alpha_2(\omega, \omega', t_1) = \alpha_3(\omega, \omega', t_1) = \lambda^2 t_1 \delta_{\omega, \omega'}$.

In the Markovian-secular limit, the master equation for the one-point reduced operator Eq. (A22) further simplifies to,

$$\begin{aligned}
\frac{d}{dt} O_S(t) &= iH_S O_S(t) \\
&+ (i\lambda)^2 \sum_{\omega} \sum_{i,j} J^{ij}(\omega) [S_{\omega}^{i\dagger} S_{\omega}^j O_S(t) - S_{\omega}^{i\dagger} O_S(t) S_{\omega}^j] + h.c.
\end{aligned} \tag{A30}$$

This master equation is used to derive Eq. (29) for the dissipative spin-boson model in the main text.

Appendix B: Some further details on QRT in the Markovian limit

In this Appendix, we discuss some further details on QRT in the Markovian limit and in particular some useful properties of the function I as also discussed in the main text. One of the interesting properties of the function I is to note that its dependence on t_2 is quite simple. This in turn helps us to establish Regression theorem quite easily.

1. Irreducible part for two-point function

Let us first set, $O_2 = A_{\mu}$ and assume $t_1 < t_2$ in equation (A14). Taking derivative of the equation (A14) with respect to t_2 , 1st term of the right hand side of the equation (A14) trivially gives the regression type form (using equation (8)). Equations (A16), (A25)-(A27) gives us the explicit form of the 2nd term i.e. irreducible part of the equation (A14). Notice that, the t_2 dependency in the irreducible part $I[O_S(t_1), A_{\mu S}(t_2)]$ comes from $A_{\mu S}(t_2)$ only. So if we take derivative of I with respect to t_2 , it simply gives

$$\frac{d}{dt_2} I[O_S(t_1), A_{\mu S}(t_2)] = \sum_{\lambda} M_{\mu\lambda} I[O_S(t_1), A_{\lambda S}(t_2)] \tag{B1}$$

In addition, in the expression of $I[O_S(t_1), A_{\mu S}(t_2)]$ (see equation (A25)-(A27)), it is easy to notice that if we swap the position of $O_S(t_1)$ and $A_{\mu S}(t_2)$, the equation (B1) still holds i.e.

$$\frac{d}{dt_2} I[A_{\mu S}(t_2), O_S(t_1)] = \sum_{\lambda} M_{\mu\lambda} I[A_{\lambda S}(t_2), O_S(t_1)] \tag{B2}$$

Interestingly, If we consider $t_2 < t_1$ and take derivative of the equation (A14) with respect to t_2 , then we will not get the regression type form. The simple reason behind this in Heisenberg picture is that in the expression of I , t_2 dependency comes from both $A_{\mu S}$ and α 's defined in (A27). Now taking derivative with respect to t_2 will give rise to complicated terms[39] i.e.

$$\frac{d}{dt_2} \langle O(t_1) A_{\mu}(t_2) \rangle \neq \sum_{\lambda} M_{\mu\lambda} \langle O(t_1) A_{\lambda}(t_2) \rangle \tag{B3}$$

2. Irreducible part for three-point function

Now let us first set, $O_3 = A_{\mu}$ and assume $t_i < t_3$ with $i = 1, 2$ in equation (A19). Taking derivative of the equation (A19) with respect to t_3 , 1st term of the right

hand side gives (using equation (8)),

$$\begin{aligned} & \frac{d}{dt_3}(O_{1S}(t_1)O_{2S}(t_2)A_{\mu S}(t_3)) \\ &= \sum_{\lambda} M_{\mu\lambda}(O_{1S}(t_1)O_{2S}(t_2)A_{\lambda S}(t_3)) \end{aligned} \quad (\text{B4})$$

Now the 3rd term of the right hand side of the equation (A19) is (using equation (A16)),

$$\begin{aligned} & W_{1,2,3}\{O_{1S}(t_1)I[O_{2S}(t_2), A_{\mu S}(t_3)]\} \\ &= W_{1,2,3}\{O_{1S}(t_1)(I_1 + I_2 + I_3 + I_4)\} \end{aligned} \quad (\text{B5})$$

$$W_{1,2,3}\{O_{1S}(t_1)I_1[O_{2S}(t_2), A_{\mu S}(t_3)]\} = -\lambda^2 \sum_{\omega, \omega'} \sum_{i,j} O_{1S}(t_1)O_{2S}(t_2)S_{\omega}^i A_{\mu S}(t_3)S_{\omega'}^j \alpha_1(\omega, \omega', t_2) \beta_1^{ij}(\omega') \quad (\text{B6})$$

If we differentiate the equation (B6) with respect to t_3 , we get (using equation (8))

$$\begin{aligned} & \frac{d}{dt_3}W_{1,2,3}\{O_{1S}(t_1)I_1[O_{2S}(t_2), A_{\mu S}(t_3)]\} \\ &= \sum_{\lambda} M_{\mu\lambda}W_{1,2,3}\{O_{1S}(t_1)I_1[O_{2S}(t_2), A_{\lambda S}(t_3)]\} \end{aligned} \quad (\text{B7})$$

since in the expression of $W_{1,2,3}\{O_{1S}(t_1)I_1[O_{2S}(t_2), A_{\mu S}(t_3)]\}$, t_3 dependency comes from $A_{\mu S}(t_3)$ only. Similarly, we can show that all the other terms of the equation (B5) follows the identical equation to (B7). This finally gives,

$$\begin{aligned} & \frac{d}{dt_3}W_{1,2,3}\{O_{1S}(t_1)I[O_{2S}(t_2), A_{\mu S}(t_3)]\} \\ &= \sum_{\lambda} M_{\mu\lambda}W_{1,2,3}\{O_{1S}(t_1)I[O_{2S}(t_2), A_{\lambda S}(t_3)]\} \end{aligned} \quad (\text{B8})$$

3. Irreducible part for four-point function

Set, $O_4 = A_{\mu}$, $t_i < t_4$ with $i = 1, 2, 3$ in the equation (A21) and if we take derivative of the equation (A21) with respect to t_4 then the 1st term of the right hand side will simply give (using equation (8)),

$$\begin{aligned} & \frac{d}{dt_4}(O_{1S}(t_1)O_{2S}(t_2)O_{3S}(t_3)A_{\mu S}(t_4)) \\ &= \sum_{\lambda} M_{\mu\lambda}(O_{1S}(t_1)O_{2S}(t_2)O_{3S}(t_3)A_{\lambda S}(t_4)) \end{aligned} \quad (\text{B9})$$

We can very straight forwardly conclude that the 2nd term will obey the following equation (using equation

The first term of the right hand side of the equation (B5) in the markovian limit is given by (using equation (A26)),

(8)),

$$\begin{aligned} & \frac{d}{dt_4}W_{1,2,3,4}\{I[O_{1S}(t_1), O_{2S}(t_2)]O_{3S}(t_3)A_{\mu S}(t_4)\} \\ &= \sum_{\lambda} M_{\mu\lambda}W_{1,2,3,4}\{I[O_{1S}(t_1), O_{2S}(t_2)]O_{3S}(t_3)A_{\lambda S}(t_4)\} \end{aligned} \quad (\text{B10})$$

By giving the exactly similar argument we can show that the 3rd and 4th term of the right hand side of the equation (A21) will follow the identical to the above equation. Now using equation (B1) we can show that the 5th term of the right hand side of the same equation will give,

$$\begin{aligned} & \frac{d}{dt_4}W_{1,2,3,4}\{I[A_{\mu S}(t_4), O_{3S}(t_3)]O_{2S}(t_2)O_{1S}(t_1)\} \\ &= \sum_{\lambda} M_{\mu\lambda}W_{1,2,3,4}\{I[A_{\lambda S}(t_4), O_{3S}(t_3)]O_{2S}(t_2)O_{1S}(t_1)\} \end{aligned} \quad (\text{B11})$$

Since, in the expression of $I[A_{\mu S}(t_4), O_{3S}(t_3)]$, the t_4 dependency comes from $A_{\mu S}(t_4)$ only and by giving the similar argument we can show that the last two term of the equation (A21) will also follow the identical equation. Then finally we will arrive at the following equation (using equation (A21)),

$$\begin{aligned} & \frac{d}{dt_4}(O_1(t_1)O_2(t_2)O_3(t_3)A_{\mu}(t_4))_S \\ &= \sum_{\lambda} M_{\mu\lambda}(O_1(t_1)O_2(t_2)O_3(t_3)A_{\lambda}(t_4))_S \end{aligned} \quad (\text{B12})$$

Appendix C: Out-of-time-ordered correlators (OTOCs)

In this section we calculate out-of-time-order correlator (OTOC). More specifically we want to derive regression

$$\begin{aligned}
(O_1(t_1)A_\mu(t_2)O_3(t_1)A_\nu(t_2))_S &= O_{1S}(t_1)A_{\mu S}(t_2)O_{3S}(t_1)A_{\nu S}(t_2) + W_{1,2,3,4} \left\{ I[O_{1S}(t_1), A_{\mu S}(t_2)]O_{3S}(t_1)A_{\nu S}(t_2) \right\} \\
&+ W_{1,2,3,4} \left\{ O_{1S}(t_1)I[A_{\mu S}(t_2), O_{3S}(t_1)]A_{\nu S}(t_2) \right\} + W_{1,2,3,4} \left\{ I[O_{1S}(t_1), O_{3S}(t_1)]A_{\mu S}(t_2)A_{\nu S}(t_2) \right\} \\
&+ W_{1,2,3,4} \left\{ I[A_{\nu S}(t_2), O_{3S}(t_1)]A_{\mu S}(t_2)O_{1S}(t_1) \right\} W_{1,2,3,4} \left\{ I[O_{1S}(t_1), A_{\nu S}(t_2)]O_{3S}(t_1)A_{\mu S}(t_2) \right\} \\
&+ W_{1,2,3,4} \left\{ O_{1S}(t_1)I[A_{\mu S}(t_2), A_{\nu S}(t_2)]O_{3S}(t_1) \right\}
\end{aligned} \tag{C1}$$

Differentiate the above equation by t_2 , then the first term of the right hand side gives

$$\begin{aligned}
&\frac{d}{dt_2}(O_{1S}(t_1)A_{\mu S}(t_2)O_{3S}(t_1)A_{\nu S}(t_2)) \\
&= \sum_{\lambda} M_{\mu\lambda}(O_{1S}(t_1)A_{\lambda S}(t_2)O_{3S}(t_1)A_{\nu S}(t_2)) \\
&+ \sum_{\lambda'} M_{\nu\lambda'}(O_{1S}(t_1)A_{\mu S}(t_2)O_{3S}(t_1)A_{\lambda' S}(t_2))
\end{aligned}$$

Similarly, the 2nd term of the right hand side gives,

$$\begin{aligned}
&\frac{d}{dt_2}W_{1,2,3,4} \left\{ I[O_{1S}(t_1), A_{\mu S}(t_2)]O_{3S}(t_1)A_{\nu S}(t_2) \right\} \\
&= \sum_{\lambda} M_{\mu\lambda}W_{1,2,3,4} \left\{ I[O_{1S}(t_1), A_{\lambda S}(t_2)]O_{3S}(t_1)A_{\nu S}(t_2) \right\} \\
&+ \sum_{\lambda'} M_{\nu\lambda'}W_{1,2,3,4} \left\{ I[O_{1S}(t_1), A_{\mu S}(t_2)]O_{3S}(t_1)A_{\lambda' S}(t_2) \right\}
\end{aligned} \tag{C2}$$

All the other terms in (C1) also follow the identical equation except the last term. The last term gives,

$$\begin{aligned}
&\frac{d}{dt_2}W_{1,2,3,4} \left\{ O_{1S}(t_1)I[A_{\mu S}(t_2), A_{\nu S}(t_2)]O_{3S}(t_1) \right\} \\
&= W_{1,2,3,4} \left\{ O_{1S}(t_1)\frac{d}{dt_2}(I[A_{\mu S}(t_2), A_{\nu S}(t_2)])O_{3S}(t_1) \right\}
\end{aligned} \tag{C3}$$

Using expression of I

$$\begin{aligned}
\frac{d}{dt_2}I[A_{\mu S}(t_2), A_{\nu S}(t_2)] &= \sum_{\lambda} M_{\mu\lambda}I[A_{\lambda S}(t_2), A_{\nu S}(t_2)] \\
&+ \sum_{\lambda'} M_{\nu\lambda'}I[A_{\mu S}(t_2), A_{\lambda' S}(t_2)] \\
&+ F[A_{\mu S}(t_2), A_{\nu S}(t_2)]
\end{aligned} \tag{C4}$$

type theorem for OTOC [21]. Now set, $t_1 = t_3, t_2 = t_4, t_2 > t_1$ and $O_2(t_2) = A_{\mu}(t_2)$, $O_4(t_2) = A_{\nu}(t_2)$ in the equation (A21) then we get,

where

$$\begin{aligned}
&F[A_{\mu S}(t_2), A_{\nu S}(t_2)] \\
&= -\lambda^2 \sum_{\omega, \omega'} \sum_{i, j} S_{\omega}^{i\dagger} A_{\mu S}(t_2) S_{\omega'}^j A_{\nu S}(t_2) e^{i(\omega+\omega')t_2} \beta_4^{ij}(\omega') \\
&+ \lambda^2 \sum_{\omega, \omega'} \sum_{i, j} S_{\omega}^{i\dagger} A_{\mu S}(t_2) A_{\nu S}(t_2) S_{\omega'}^j e^{i(\omega-\omega')t_2} \beta_2^{ij}(\omega') \\
&+ \lambda^2 \sum_{\omega, \omega'} \sum_{i, j} A_{\mu S}(t_2) S_{\omega}^i S_{\omega'}^{j\dagger} A_{\nu S}(t_2) e^{-i(\omega-\omega')t_2} \beta_3^{ij}(\omega') \\
&- \lambda^2 \sum_{\omega, \omega'} \sum_{i, j} A_{\mu S}(t_2) S_{\omega}^i A_{\nu S}(t_2) S_{\omega'}^j e^{-i(\omega+\omega')t_2} \beta_1^{ij}(\omega')
\end{aligned} \tag{C5}$$

(C3) becomes

$$\begin{aligned}
&\frac{d}{dt_2}W_{1,2,3,4} \left\{ O_{1S}(t_1)I[A_{\mu S}(t_2), A_{\nu S}(t_2)]O_{3S}(t_1) \right\} \\
&= \sum_{\lambda} M_{\mu\lambda}W_{1,2,3,4} \left\{ O_{1S}(t_1)I[A_{\lambda S}(t_2), A_{\nu S}(t_2)]O_{3S}(t_1) \right\} \\
&+ \sum_{\lambda'} M_{\nu\lambda'}W_{1,2,3,4} \left\{ O_{1S}(t_1)I[A_{\mu S}(t_2), A_{\lambda' S}(t_2)]O_{3S}(t_1) \right\} \\
&+ W_{1,2,3,4} \left\{ O_{1S}(t_1)F[A_{\mu S}(t_2), A_{\nu S}(t_2)]O_{3S}(t_1) \right\}
\end{aligned} \tag{C6}$$

Finally adding all the different contributions we receive,

$$\begin{aligned}
&\frac{d}{dt_2}(O_1(t_1)A_{\mu}(t_2)O_3(t_1)A_{\nu}(t_2))_S \\
&= \sum_{\lambda} M_{\mu\lambda}(O_1(t_1)A_{\lambda}(t_2)O_3(t_1)A_{\nu}(t_2))_S \\
&+ \sum_{\lambda'} M_{\nu\lambda'}(O_1(t_1)A_{\mu}(t_2)O_3(t_1)A_{\lambda'}(t_2))_S \\
&+ W_{1,2,3,4} \left\{ O_{1S}(t_1)F[A_{\mu S}(t_2), A_{\nu S}(t_2)]O_{3S}(t_1) \right\}
\end{aligned} \tag{C7}$$

We observe that regression theorem for OTOC takes a

different form than discussed in previous section.

by,

$$I[\sigma_{xS}(t_1), \sigma_{xS}(t_2)] = \gamma t_1 e^{i\omega'_0(t_2-t_1)} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (\text{D1})$$

Appendix D: Example: Dissipative spin half system

In this appendix we provide the details of the calculation for the dissipative spin 1/2 system as discussed in the main text.

1. Correlation function in Heisenberg picture

We give out expressions for two, three and four-point reduced operators using the Heisenberg picture.

a. Two point function

Here we are going to calculate the irreducible part of the equation (A14) to get the following two-point functions $\langle \sigma_x(t)\sigma_x(t+\tau) \rangle$ and $\langle \sigma_x(t)\sigma_y(t+\tau) \rangle$. Using equation (A16), (A25)-(A27) and by imposing secular approximation we can calculate the irreducible part I and it is given

b. Three and four-point functions

To get three and four point reduced operators we have to calculate all the W terms of the equation (A19) and equation (A21). Here we have written down the expression of all the W terms defined in equation (A19),

$$W_{1,2,3}\{I[\sigma_{xS}(t_1), \sigma_{xS}(t_2)]\sigma_{xS}(t_3)\} = \gamma t_1 \begin{pmatrix} 0 & e^{i\omega'_0(-t_1+t_2+t_3)} \\ e^{i\omega'_0(-t_1+t_2-t_3)} & 0 \end{pmatrix} \quad (\text{D5})$$

$$W_{1,2,3}\{I[\sigma_{xS}(t_1), \sigma_{xS}(t_3)]\sigma_{xS}(t_2)\} = -\gamma t_1 \begin{pmatrix} 0 & e^{i\omega'_0(-t_1+t_2+t_3)} \\ e^{i\omega'_0(-t_1-t_2+t_3)} & 0 \end{pmatrix} \quad (\text{D6})$$

$$W_{1,2,3}\{\sigma_{xS}(t_1)I[\sigma_{xS}(t_2), \sigma_{xS}(t_3)]\} = \gamma t_2 \begin{pmatrix} 0 & e^{i\omega'_0(t_1-t_2+t_3)} \\ e^{i\omega'_0(-t_1-t_2+t_3)} & 0 \end{pmatrix} \quad (\text{D7})$$

and

$$\sigma_{xS}(t_1)\sigma_{xS}(t_2)\sigma_{xS}(t_3) = \left(1 - \frac{\gamma}{2}(t_1 + t_2 + t_3)\right) \begin{pmatrix} 0 & e^{i\omega'_0(t_1-t_2+t_3)} \\ e^{i\omega'_0(-t_1+t_2-t_3)} & 0 \end{pmatrix} \quad (\text{D8})$$

Here we have shown only one W term which is needed to calculate four-point reduced operator and as defined in equation (A21)

$$W_{1,2,3,4}\{I[\sigma_{xS}(t_1), \sigma_{xS}(t_2)]\sigma_{xS}(t_3)\sigma_{xS}(t_4)\} = \gamma t_1 \begin{pmatrix} e^{i\omega'_0(-t_1+t_2+t_3-t_4)} & 0 \\ 0 & e^{i\omega'_0(-t_1+t_2-t_3+t_4)} \end{pmatrix} \quad (\text{D9})$$

Appendix E: Extension of regression theorem for two-point reduced operator

The two-point reduced operator can be written as,

$$(O_1(t_1)O_2(t_2))_S = O_{1S}(t_1)O_{2S}(t_2) + I[O_{1S}(t_1), O_{2S}(t_2)] \quad (\text{E1})$$

Where, $I = I_1 + I_2 + I_3 + I_4$ and we can show that (here we assumed $t_1 > t_2$), (correct upto order λ^2)

Now by differentiating the equation (E2) with respect to t_1 we receive,

$$\begin{aligned}
I_1 &= -\lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) O_{1S}(t_1) \tilde{L}^\dagger(-\tau_1) O_{2S}(t_2) \tilde{L}(-\tau_2), \\
I_2 &= \lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) \tilde{L}^\dagger(-\tau_1) O_{1S}(t_1) O_{2S}(t_2) \tilde{L}(-\tau_2), \\
I_3 &= \lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) O_{1S}(t_1) \tilde{L}^\dagger(-\tau_1) \tilde{L}(-\tau_2) O_{2S}(t_2), \\
I_4 &= -\lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) \tilde{L}^\dagger(-\tau_1) O_{1S}(t_1) \tilde{L}(-\tau_2) O_{2S}(t_2).
\end{aligned} \tag{E2}$$

$$\begin{aligned}
\frac{d}{dt_1} I_1 &= -\lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) (i[H_S, O_{1S}(t_1)] \tilde{L}^\dagger(-\tau_1) O_{2S}(t_2) \tilde{L}(-\tau_2)) - \lambda^2 \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) O_{1S}(t_1) \tilde{L}^\dagger(-t_1) O_{2S}(t_2) \tilde{L}(-\tau_2) \\
\frac{d}{dt_1} I_2 &= \lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) (\tilde{L}^\dagger(-\tau_1) [i[H_S, O_{1S}(t_1)] O_{2S}(t_2) \tilde{L}(-\tau_2)]) + \lambda^2 \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) \tilde{L}^\dagger(-t_1) O_{1S}(t_1) O_{2S}(t_2) \tilde{L}(-\tau_2) \\
\frac{d}{dt_1} I_3 &= \lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) (i[H_S, O_{1S}(t_1)] \tilde{L}^\dagger(-\tau_1) \tilde{L}(-\tau_2) O_{2S}(t_2)) + \lambda^2 \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) O_{1S}(t_1) \tilde{L}^\dagger(-t_1) \tilde{L}(-\tau_2) O_{2S}(t_2) \\
\frac{d}{dt_1} I_4 &= -\lambda^2 \int_0^{t_1} \int_0^{t_2} d\tau_1 d\tau_2 \alpha(\tau_2 - \tau_1) (\tilde{L}^\dagger(-\tau_1) [i[H_S, O_{1S}(t_1)] \tilde{L}(-\tau_2) O_{2S}(t_2)]) - \lambda^2 \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) \tilde{L}^\dagger(-t_1) O_{1S}(t_1) \tilde{L}(-\tau_2) O_{2S}(t_2)
\end{aligned} \tag{E3}$$

and,

$$\frac{d}{dt_1} (O_{1S}(t_1) O_{2S}(t_2)) = \dot{O}_{1S}(t_1) O_{2S}(t_2) \tag{E4}$$

In the Markovian limit, we can show that the 2nd term of the equations (E3) vanishes.

Now assume there exists a set of system operators such that,

$$\frac{d}{dt} A_{\mu S}(t) = \sum_{\lambda} M_{\mu\lambda}(t) A_{\lambda S}(t) \tag{E5}$$

Now set, $O_2 = O$ and $O_1 = A_{\mu}$ then by using equation (E3) and (E5) we get,

$$\begin{aligned}
\frac{d}{dt_1} I_1[A_{\mu S}(t_1), O_S(t_2)] &= \sum_{\lambda} M_{\mu\lambda}(t_1) I_4[A_{\lambda S}(t_1), O_S(t_2)] \\
&- \lambda^2 \int_0^{t_2} d\tau_2 \alpha(\tau_2 - t_1) [A_{\mu S}(t_1) \tilde{L}^\dagger(-t_1) O_{2S}(t_2) \tilde{L}(-\tau_2)]
\end{aligned} \tag{E6}$$

We also get the identical equation for I_2, I_3 and I_4 and then adding all of them we will finally get the equation (51).