

# SPECTRUM OF THE LAMÉ OPERATOR ALONG $\operatorname{Re}\tau = 1/2$ : THE GENUS 3 CASE

ERJUAN FU

ABSTRACT. In this paper, we study the spectrum  $\sigma(L)$  of the Lamé operator

$$L = \frac{d^2}{dx^2} - 12\wp(x + z_0; \tau) \quad \text{in } L^2(\mathbb{R}, \mathbb{C}),$$

where  $\wp(z; \tau)$  is the Weierstrass elliptic function with periods 1 and  $\tau$ , and  $z_0 \in \mathbb{C}$  is chosen such that  $L$  has no singularities on  $\mathbb{R}$ . We prove that a point  $\lambda \in \sigma(L)$  is an intersection point of different spectral arcs but not a zero of the spectral polynomial if and only if  $\lambda$  is a zero of the following cubic polynomial:

$$\frac{4}{15}\lambda^3 + \frac{8}{5}\eta_1\lambda^2 - 3g_2\lambda + 9g_3 - 6\eta_1g_2 = 0.$$

We also study the deformation of the spectrum as  $\tau = \frac{1}{2} + ib$  with  $b > 0$  varying. We discover 7 different types of graphs for the spectrum as  $b$  varies around the double zeros of the spectral polynomial.

## CONTENTS

1. Introduction	1
2. Preliminaries	6
3. Inner intersection points	11
4. Zeros of the spectral polynomial	14
5. Proof of the main theorem	20
References	25

## 1. INTRODUCTION

In this paper, we study the spectrum  $\sigma(L_\tau^g)$  of the classical Lamé operator [15]

$$L_\tau^g = \frac{d^2}{dx^2} - g(g+1)\wp(x + z_0; \tau), \quad x \in \mathbb{R}$$

in  $L^2(\mathbb{R}, \mathbb{C})$ , where  $g \in \mathbb{N}$ ,  $\wp(z; \tau)$  is the Weierstrass elliptic function with basic periods  $\omega_1 = 1$ ,  $\omega_2 = \tau$  and  $z_0 \in \mathbb{C}$  is chosen such that  $\wp(x + z_0; \tau)$  has no singularities on  $\mathbb{R}$ . Remark that  $\sigma(L_\tau^g)$  does not depend on the choice of  $z_0$  due to the fact that the Lamé potential  $-g(g+1)\wp(z; \tau)$  is a Picard

potential in the sense of Gesztesy and Weikard [10] (i.e. all solutions of the Lamé equation

$$\mathcal{L}_{\tau,\lambda}^g : y''(z) = (g(g+1)\wp(z;\tau) + \lambda)y(z), \quad z \in \mathbb{C}$$

are meromorphic in  $\mathbb{C}$ ).

The spectral theory of the Schrödinger operator  $L$  with complex periodic smooth potentials has attracted significant attention and has been studied widely in the literature; see e.g. [1, 2, 10, 12, 13, 19] and references therein. In this theory, it is known [19] that


$$\sigma(L) = \Delta^{-1}([-2, 2]) = \{\lambda \in \mathbb{C} \mid -2 \leq \Delta(\lambda) \leq 2\},$$

where  $\Delta(\lambda)$  denotes the Hill's discriminant which is the trace of the monodromy matrix of  $Ly = \lambda y$  with respect to  $x \rightarrow x + 1$ . Furthermore, it was proved in [10] that  $\sigma(L)$  consists of finitely many analytic arcs if the potential of  $L$  is a Picard potential. In this paper, as in [4, 10] we call the arcs of  $\sigma(L)$  as *spectral arcs*.


For the simplest case  $\operatorname{Re}\tau = 0$ , since the Lamé potential  $-g(g+1)\wp(x + \frac{\omega_2}{2}; \tau)$  is real-valued and smooth on  $\mathbb{R}$ , Ince [14] proved that there are  $2g + 1$  distinct real numbers  $\lambda_0 > \lambda_1 > \dots > \lambda_{2g}$  such that the spectrum

$$(1.1) \quad \sigma(L_\tau^g) = (-\infty, \lambda_{2g}] \cup [\lambda_{2g-1}, \lambda_{2g-2}] \cup \dots \cup [\lambda_1, \lambda_0] \subseteq \mathbb{R}.$$

However, the spectrum  $\sigma(L_\tau^g)$  is no longer of the form (1.1) for general  $\tau$ 's and becomes very complicated; see e.g. [1, 5, 6, 11, 12, 13] and references therein. Gesztesy-Weikard [12] and Haese-Hill et al. [13] concentrated on the  $g = 1$  case, for which the spectrum  $\sigma(L_\tau^1)$  consists of two regular analytic arcs and so there are totally three different types of graphs for different  $\tau$ 's as shown in the following figure (see also [1, 11]).



In particular, different spectral arcs might intersect as shown in Figure (b)-(c). It was pointed out in [13, Section 5] that the rigorous analysis of  $g \geq 2$  cases seems to be difficult since the related explicit formulae quickly become quite complicated as  $g$  grows. Recently, [4, Lemma 4.1] inferred that the spectrum  $\sigma(L_\tau^g)$  is symmetric with respect to the real line  $\mathbb{R}$  if  $\tau = \frac{1}{2} + bi$  with  $b > 0$ . Furthermore, it was proved in [4] that the spectrum  $\sigma(L_{\frac{1}{2}+bi}^2)$  has exactly 9 different types of graphs for different  $b$ 's, and the continuous deformation of the spectrum as  $b$  decreases is as follows.



In this paper, we will focus on the genus  $g = 3$  case. To the best of our knowledge, there seems no more explicit description of  $\sigma(L_\tau^3)$  in the literature. It is well known (see [17, p.569]) that the associated spectral polynomial  $Q_3(\lambda; \tau)$  is given by

$$(1.2) \quad Q_3(\lambda; \tau) = \lambda \prod_{k=1}^3 (\lambda^2 - 6e_k\lambda + 15(3e_k^2 - g_2)),$$

where  $e_k := e_k(\tau) = \wp(\omega_k/2; \tau)$  for  $k = 1, 2, 3$  and  $\omega_3 = 1 + \tau$ , so the associated hyperelliptic curve  $\{(\lambda, C) \mid C^2 = Q_3(\lambda; \tau)\}$  is of genus 3.

By applying [10, Theorem 4.1], we see that the spectrum  $\sigma(L_\tau^3)$  consists of  $\tilde{g} \leq 3$  bounded simple analytic arcs  $\sigma_k$  and one semi-infinite simple analytic arc  $\sigma_\infty$  which tends to  $-\infty + \langle q \rangle$ , with  $\langle q \rangle = \int_{x_0}^{x_0+1} q(x)dx$ , i.e.

$$\sigma(L) = \sigma_\infty \cup \bigcup_{k=1}^{\tilde{g}} \sigma_k, \quad \tilde{g} \leq 3,$$

where the finite endpoints of such arcs must be zeros of the spectral polynomial  $Q_3(\lambda; \tau)$  with odd order.

As shown in the above figures, there are two kinds of intersection points of spectral arcs: One is that this intersection point is also an endpoint of some arc (see Figures (b) and (8)), so it is met by  $2k + 1$  semi-arcs for some  $k \geq 1$ ; the other is that this intersection point is not an endpoint (see Figures (c) and (1), (4)-(7)), so it is met by  $2k$  semi-arcs for some  $k \geq 2$ , i.e. it is an inner point of  $k$  arcs; and we call it an *inner intersection point*.

First, we have to determine completely the inner intersection points of different arcs in order to study the geometry of  $\sigma(L_\tau^3)$ . Our first result is as follows.

**Theorem 1.1.** *Let  $\tau \in \mathbb{H}$  and  $\lambda_0 \in \sigma(L_\tau^3)$  with  $Q_3(\lambda_0; \tau) \neq 0$ . Then  $\lambda_0$  is an inner intersection point if and only if  $\lambda_0$  satisfies the following cubic equation*

$$f(\lambda) := \frac{4}{15}\lambda^3 + \frac{8}{5}\eta_1\lambda^2 - 3g_2\lambda + 9g_3 - 6\eta_1g_2 = 0,$$

where  $g_2, g_3$  denote the elliptic invariants of the Weierstrass elliptic function  $\wp$ .

Now we study the deformation of the spectrum  $\sigma(L_\tau^3)$  as  $\tau = \frac{1}{2} + bi$  deforms. This problem is challenging and we can only obtain some partial results. For this purpose, we need to analysis when some endpoint  $\lambda_0$  (i.e.  $\lambda_0$  is a zero of  $Q_3(\lambda; \tau)$  with odd order) could be an intersection point.

**Theorem 1.2.** *Let  $\tau = \frac{1}{2} + bi$  with  $b > 0$  and  $d(\lambda)$  denote the number of semi-arcs met at  $\lambda$ . Then the zeros of the spectral polynomial  $Q_3(\lambda; \tau)$  are listed as follows*

$$0, \mu, \bar{\mu}, \nu, \bar{\nu}, \vartheta_+, \vartheta_-,$$

where  $\mu \neq \nu$  and  $\mu, \nu \notin \mathbb{R}$ . Furthermore, there is some  $\beta \approx 1.0979 \in \left(\frac{\sqrt{3}}{2}, +\infty\right)$  such that

$$(z1) \quad \vartheta_- = \vartheta_+ = 3e_1 \text{ if and only if } b \in \{\beta, \hat{\beta}\}, \text{ where } \hat{\beta} = \frac{1}{4\beta} \approx 0.2277.$$

$$(z2) \quad \vartheta_-, \vartheta_+ \in \mathbb{R} \text{ and } \vartheta_- < 3e_1 < \vartheta_+ < 0 \text{ if and only if } b \in (0, \hat{\beta}).$$

$$(z3) \quad \vartheta_-, \vartheta_+ \in \mathbb{R} \text{ and } 0 < \vartheta_- < 3e_1 < \vartheta_+ \text{ if and only if } b \in (\beta, +\infty).$$

$$(z4) \quad \vartheta_- = \overline{\vartheta_+} \notin \mathbb{R} \text{ if and only if } b \in (\hat{\beta}, \beta).$$

Moreover,

$$(d1) \quad d(0) = 1 + 2d \text{ for some } d \geq 1 \text{ (i.e. the endpoint } 0 \text{ is an intersection point) if and only if } b = b_g \approx 0.47; \text{ otherwise, } d(0) = 1.$$

$$(d2) \quad d(\mu) = d(\nu) = d(\bar{\mu}) = d(\bar{\nu}) = 1, \text{ i.e. the endpoints } \mu, \nu, \bar{\mu}, \bar{\nu} \text{ can not be intersection points.}$$


$$(d3)$$

$$d(\vartheta_-) = d(\vartheta_+) = \begin{cases} 2 & \text{if } b \in \{\hat{\beta}, \beta\} \\ 1 & \text{if } b \in (\hat{\beta}, \beta). \end{cases}$$

Note that  $Q_3(\lambda; \frac{1}{2} + bi)$  has double zeros at  $b \in \{\beta, \hat{\beta}\}$ . Here we mainly study the deformation of the spectrum  $\sigma(L_\tau^3)$  along  $\tau = \frac{1}{2} + bi$  with  $b$  varying around  $\hat{\beta}$  and  $\beta$ , respectively. We discover 7 different patterns for the spectrum stated in the following theorem. See also the following figure for these 7 rough graphs.

**Theorem 1.3.** *Let  $\tau = \frac{1}{2} + ib$  with  $b > 0$  and denote  $L_b^3 := L_\tau^3$ . Then the spectrum  $\sigma(L_b^3)$  is symmetric with respect to the real line  $\mathbb{R}$ . Furthermore, there exist two sufficient small positive real numbers  $\varepsilon, \delta$  and  $\alpha \approx 0.23217 \in (\hat{\beta}, \frac{1}{2})$  such that the rough graphs of the spectra  $\sigma(L_b^3)$  for  $b \in (\hat{\beta} - \delta, \alpha + \varepsilon) \cup [\beta, +\infty)$  are described as follows.*

Here, the notations  $\sigma_i$  with  $i = 1, 2, 3$  denote simple arcs symmetric with respect to  $\mathbb{R}$  and they are disjoint with each other. The notation  $\sigma_4$  denotes a simple arc in  $\{z \in \mathbb{C} \mid \text{Im } z \geq 0\}$  and  $\bar{\sigma}_4$  denotes the conjugate of  $\sigma_4$ .



- (1) If  $\widehat{\beta} - \delta < b < \widehat{\beta}$ , then  $\vartheta_- < 3e_1 < \vartheta_+ < 0$  and

$$\sigma(L_b^3) = (-\infty, \vartheta_-] \cup [\vartheta_+, 0] \cup \sigma_1 \cup \sigma_2$$

with  $\sigma_1 \cap (0, +\infty) = \{\text{one point}\}$  and  $\sigma_2 \cap (-\infty, \vartheta_-) = \{\lambda_-\}$ , where  $f(\lambda_-) = 0$ .

- (2) If  $b = \widehat{\beta}$ , then  $\vartheta_- = \vartheta_+ = 3e_1 < 0$  and

$$\sigma(L_{\widehat{\beta}}^3) = (-\infty, 0] \cup \sigma_1 \cup \sigma_2$$

with  $\sigma_1 \cap (0, +\infty) = \{\text{one point}\}$  and  $\sigma_2 \cap (-\infty, 0) = \{\lambda_-\}$ , where  $f(\lambda_-) = 0$ .

- (3) If  $\widehat{\beta} < b < \alpha$ , then  $f(\lambda)$  has three real roots  $\lambda_- < \lambda'_- < 0 < \lambda_+$  and

$$\sigma(L_b^3) = (-\infty, 0] \cup \sigma_1 \cup \sigma_2 \cup \sigma_3$$

with  $\sigma_1 \cap (0, +\infty) = \{\text{one point}\}$ ,  $\sigma_2 \cap (-\infty, 0) = \{\lambda_-\}$  and  $\sigma_3 \cap (-\infty, 0) = \{\lambda'_-\}$ , where  $f(\lambda_-) = f(\lambda'_-) = 0$ .

- (4) If  $b = \alpha$ , then

$$\sigma(L_{\alpha}^3) = (-\infty, 0] \cup \sigma_1 \cup \sigma_4 \cup \overline{\sigma_4}$$

with  $\sigma_1 \cap (0, +\infty) = \{\text{one point}\}$  and  $\sigma_4 \cap (-\infty, 0) = \overline{\sigma_4} \cap (-\infty, 0) = \{\lambda_-\}$ , where  $f(\lambda_-) = 0$ .

- (5) If  $\alpha < b < \alpha + \varepsilon$ , then

$$\sigma(L_b^3) = (-\infty, 0] \cup \sigma_1 \cup \sigma_4 \cup \overline{\sigma_4}$$

with  $\sigma_1 \cap (0, +\infty) = \{\text{one point}\}$  and  $\sigma_4 \cap \mathbb{R} = \emptyset$ .

- (6) If  $b = \beta$ , then  $\vartheta_- = \vartheta_+ = 3e_1 > 0$  and

$$\sigma(L_{\beta}^3) = (-\infty, 0] \cup \sigma_1 \cup \sigma_2$$

with  $\sigma_1 \cap \mathbb{R} = \{3e_1\}$  and  $\sigma_2 \cap (-\infty, 0) = \{\lambda_-\}$ , where  $f(\lambda_-) = 0$ .

(7) If  $b > \beta$ , then  $0 < \vartheta_- < 3e_1 < \vartheta_+$  and

$$\sigma(L_b^3) = (-\infty, 0] \cup [\vartheta_-, \vartheta_+] \cup \sigma_1 \cup \sigma_2$$

with  $\sigma_1 \cap (\vartheta_-, \vartheta_+) = \{\lambda_+\}$  and  $\sigma_2 \cap (-\infty, 0) = \{\lambda_-\}$ , where  $f(\lambda_-) = f(\lambda_+) = 0$ .

**Remark 1.4.** The other cases  $b \in (0, \widehat{\beta} - \delta] \cup [\alpha + \varepsilon, \beta)$  seem difficult and remain open.

The rest of this paper is organized as follows. In section 2, we first review the spectral theory of Hill equation from [10] and then review the hyperelliptic curve associated with the Lamé equation, finally, we collect some conclusions about the spectrum of the Lamé operator for later usage. We prove Theorem 1.1 in section 3 and Theorem 1.2 in section 4. In the last section, we give the proof of Theorem 1.3.

## 2. PRELIMINARIES

**2.1. Spectral theory of Hill equation with complex-valued potentials.** Let  $q(x)$  is a complex-valued continuous nonconstant periodic function of period  $\omega$  on  $\mathbb{R}$ . Consider the following Hill equation

$$(2.1) \quad y''(x) + q(x)y(x) = \lambda y(x), \quad x \in \mathbb{R}.$$

This equation has received an enormous amount of consideration due to its ubiquity in applications as well as its structural richness; see e.g. [10, 12] and references therein for historical reviews.

Let  $y_1(x)$  and  $y_2(x)$  be any two linearly independent solutions of (2.1). Then so do  $y_1(x + \omega)$  and  $y_2(x + \omega)$  and hence there is a monodromy matrix  $M(\lambda) \in SL(2, \mathbb{C})$  such that

$$(y_1(x + \omega), y_2(x + \omega)) = (y_1(x), y_2(x))M(\lambda).$$

Define the *Hill's discriminant*  $\Delta(\lambda)$  by

$$(2.2) \quad \Delta(\lambda) := \text{tr}M(\lambda),$$

which is clearly an invariant of (2.1), i.e. does not depend on the choice of linearly independent solutions. This entire function  $\Delta(\lambda)$  encodes all information of the spectrum  $\sigma(L)$  of the operator  $L = \frac{d^2}{dx^2} + q(x)$ ; see e.g. [12] and references therein. Indeed, Rofo-Beketov [19] proved that the spectrum  $\sigma(L)$  of  $L$  can be described as:

$$\sigma(L) = \Delta^{-1}([-2, 2]) = \{\lambda \in \mathbb{C} \mid -2 \leq \Delta(\lambda) \leq 2\}.$$

This important fact will play a key role in this paper.

Clearly  $\lambda$  is a (anti)periodic eigenvalue if and only if  $\Delta(\lambda) = \pm 2$ . Define

$$d(\lambda) := \text{ord}_\lambda(\Delta(\cdot)^2 - 4).$$

Then it is well known (cf. [18, Section 2.3]) that  $d(\lambda)$  equals to *the algebraic multiplicity of (anti)periodic eigenvalues*. Let  $c(\lambda, x, x_0)$  and  $s(\lambda, x, x_0)$  be the special fundamental system of solutions of (2.1) satisfying the initial values

$$c(\lambda, x_0, x_0) = s'(E, x_0, x_0) = 1, \quad c'(\lambda, x_0, x_0) = s(\lambda, x_0, x_0) = 0.$$

Then we have

$$\Delta(\lambda) = c(\lambda, x_0 + \omega, x_0) + s'(\lambda, x_0 + \omega, x_0).$$

Define

$$p(\lambda, x_0) := \text{ord}_\lambda s(\cdot, x_0 + \omega, x_0),$$

$$p_i(\lambda) := \min\{p(\lambda, x_0) : x_0 \in \mathbb{R}\}.$$

It is known that  $p(\lambda, x_0)$  is the algebraic multiplicity of a Dirichlet eigenvalue on the interval  $[x_0, x_0 + \omega]$ , and  $p_i(\lambda)$  denotes the immovable part of  $p(\lambda, x_0)$  (cf. [10]). It was proved in [10, Theorem 3.2] that  $d(\lambda) - 2p_i(\lambda) \geq 0$ .

**2.2. The Weierstrass functions.** Let  $T_\tau := \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$  be a flat torus with  $\tau \in \mathbb{H} = \{\tau \in \mathbb{C} \mid \text{Im } \tau > 0\}$ . Recall that

$$\wp(z; \tau) := \frac{1}{z^2} + \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \left( \frac{1}{(z - m - n\tau)^2} - \frac{1}{(m + n\tau)^2} \right)$$

is the Weierstrass elliptic function with basic periods  $\omega_1 = 1$  and  $\omega_2 = \tau$ . Denote by  $\omega_3 = 1 + \tau$ . It is well known that

$$\wp'(z; \tau)^2 = 4 \prod_{k=1}^3 (\wp(z; \tau) - e_k(\tau)) = 4\wp(z; \tau)^3 - g_2(\tau)\wp(z; \tau) - g_3(\tau),$$

where  $e_k(\tau) = \wp(\frac{\omega_k}{2}; \tau)$ ,  $k = 1, 2, 3$ . The second equality gives us

$$(2.3) \quad e_1 + e_2 + e_3 = 0,$$

$$(2.4) \quad g_2 = 2(e_1^2 + e_2^2 + e_3^2),$$

$$(2.5) \quad g_3 = 4e_1e_2e_3.$$

The Weierstrass zeta function defined by  $\zeta(z) = \zeta(z; \tau) := -\int^z \wp(\xi; \tau) d\xi$  with two quasi-periods  $\eta_j = \eta_j(\tau)$ ,  $j = 1, 2$ :

$$\eta_j(\tau) = 2\zeta(\frac{\omega_j}{2}; \tau) = \zeta(z + \omega_j; \tau) - \zeta(z; \tau), \quad j = 1, 2.$$

The Weierstrass sigma function defined by  $\sigma(z) = \sigma(z; \tau) := \exp \int^z \zeta(\xi) d\xi$ . Notice that  $\zeta(z)$  is an odd meromorphic function with simple poles at  $\Lambda_\tau$  and  $\sigma(z)$  is an odd entire function with simple zeros at  $\Lambda_\tau$ . Moreover,

$$\zeta'(z) = -\wp(z) \quad \text{and} \quad \zeta(z) = \frac{\sigma'(z)}{\sigma(z)}.$$

**2.3. The hyperelliptic curve.** Let  $T_\tau^* := T_\tau \setminus \{0\}$  and denote

$$D_\tau^g := \{ \{z_1, \dots, z_g\} \in \text{Sym}^g T_\tau^* \mid z_i = z_j \text{ for some } i \neq j \}.$$

Define

$$Y_\tau^g := \left\{ \mathbf{a} := \{a_1, \dots, a_g\} \left| \begin{array}{l} \mathbf{a} \in (\text{Sym}^g T_\tau^*) \setminus D_\tau^g, \\ \sum_{j \neq i} (\zeta(a_i - a_j) + \zeta(a_j) - \zeta(a_i)) = 0, \forall i \end{array} \right. \right\}.$$

Clearly  $-\mathbf{a} := \{-a_1, \dots, -a_g\} \in Y_\tau^g$  if  $\mathbf{a} \in Y_\tau^g$ , and  $\mathbf{a} \in Y_\tau^g$  is called a *branch point* of  $Y_\tau^g$  if  $\mathbf{a} = -\mathbf{a}$  in  $T_\tau$ .

**Remark 2.1.** Let  $\mathbf{a} \in Y_\tau^g$  and NOT a branch point. It was proved in [3, Proposition 5.8.3] that

$$(2.6) \quad \sum_{j \neq i} (\zeta(a_i - a_j) + \zeta(a_j) - \zeta(a_i)) = 0, \forall 1 \leq i \leq g,$$

is equivalent to

$$(2.7) \quad \sum_{j=1}^g \wp'(a_j) \wp(a_j)^l = 0, \quad \forall 0 \leq l \leq g-2.$$

For  $\mathbf{a} \in Y_\tau^g$ , define

$$g_{\mathbf{a}}(z) := \sum_{i=1}^g \frac{\wp'(a_i)}{\wp(z) - \wp(a_i)} = \frac{C_{\mathbf{a}}}{\prod_{i=1}^g (\wp(z) - \wp(a_i))}$$

for a constant

$$C_{\mathbf{a}} = \wp'(a_i) \prod_{j \neq i} (\wp(a_i) - \wp(a_j)), \quad \text{which is independent of } i.$$

Remark that  $C_{\mathbf{a}} = 0$  if and only if  $\mathbf{a}$  is a branch point. Here we recall the following important theorem:

**Theorem 2.2.** [3] *There is a so-called spectral polynomial  $Q_g(\lambda; \tau) \in \mathbb{Q}[g_2(\tau), g_3(\tau)][\lambda]$  of degree  $2g+1$  such that if  $\mathbf{a} \in Y_\tau^g$ , then  $C_{\mathbf{a}}^2 = Q_g(\lambda_{\mathbf{a}}; \tau)$ .*

This spectral polynomial  $Q_g(\lambda; \tau)$  is called the *Lamé polynomial* in the literature. Consider the map (See [3, Theorem 7.4])

$$\lambda : Y_\tau^g \rightarrow \mathbb{C}$$

$$(2.8) \quad \mathbf{a} := \{a_1, \dots, a_g\} \mapsto \lambda_{\mathbf{a}} = (2g-1) \sum_{i=1}^g \wp(a_i)$$

which is a ramified covering of degree 2, and Theorem 2.2 implies

$$(2.9) \quad Y_\tau^g \cong \{(\lambda, C) \mid C^2 = Q_g(\lambda; \tau)\}$$

Therefore,  $Y_\tau^g$  is a hyperelliptic curve, known as the *Lamé curve*. And  $\mathbf{a} = -\mathbf{a}$  is a branch point of  $Y_\tau^g$  if and only if  $Q(\lambda_{\mathbf{a}}; \tau) = 0$ .



Next, we will see  $Y_\tau^\mathfrak{g}$  could parametrize the solutions of the Lamé equations. For  $\mathbf{a} = \{a_1, \dots, a_g\} \in (\text{Sym}^\mathfrak{g} T_\tau^*) \setminus D_\tau^\mathfrak{g}$ , we have the *Hermite-Halphen ansatz* (cf. [3, 20]):

$$y_{\mathbf{a}}(z) := e^{z \sum_{i=1}^{\mathfrak{g}} \zeta(a_i)} \frac{\prod_{i=1}^{\mathfrak{g}} \sigma(z - a_i)}{\sigma(z)^\mathfrak{g}}.$$

The following theorem tells us the relation between the Lamé equation  $\mathcal{L}_{\tau, \lambda}^\mathfrak{g}$  and the hyperelliptic curve  $Y_\tau^\mathfrak{g}$ .

**Theorem 2.3.** [3, 20] *Suppose  $\mathbf{a} = \{a_1, \dots, a_g\} \in (\text{Sym}^\mathfrak{g} T_\tau^*) \setminus D_\tau^\mathfrak{g}$ . Then  $y_{\mathbf{a}}(z)$  is a solution of  $\mathcal{L}_{\tau, \lambda}^\mathfrak{g}$  for some  $\lambda$  if and only if  $\mathbf{a} \in Y_\tau^\mathfrak{g}$  and  $\lambda = \lambda_{\mathbf{a}}$ .*

**Corollary 2.4.** [7, p464] *Either  $\mathbf{a} = -\mathbf{a}$  or  $\mathbf{a} \cap -\mathbf{a} = \emptyset$ . Moreover,  $y_{\mathbf{a}}(z)$  and  $y_{-\mathbf{a}}(z)$  are linearly independent if and only if  $\mathbf{a} \cap -\mathbf{a} = \emptyset$ , equivalently, if and only if  $Q_{\mathfrak{g}}(\lambda_{\mathbf{a}}; \tau) \neq 0$ .*

Let  $\mathbf{a} \in Y_\tau^\mathfrak{g}$  in the following discussion. The Legendre relation  $\tau\eta_1 - \eta_2 = 2\pi i$  implies that there is a unique  $(r, s) \in \mathbb{C}^2$  satisfying

$$r + s\tau = a_1 + \dots + a_g \quad \text{and} \quad r\eta_1 + s\eta_2 = \zeta(a_1) + \dots + \zeta(a_g),$$

which is equivalent to

$$\begin{aligned} \zeta(a_1) + \dots + \zeta(a_g) - \eta_1(a_1 + \dots + a_g) &= -2\pi i s, \\ \tau(\zeta(a_1) + \dots + \zeta(a_g)) - \eta_2(a_1 + \dots + a_g) &= 2\pi i r. \end{aligned}$$

Furthermore, the transformation law  $\sigma(z + \omega_j) = -e^{(z + \frac{\omega_j}{2})\eta_j} \sigma(z)$  with  $j = 1, 2$  implies

$$\begin{aligned} y_{\pm \mathbf{a}}(z + 1) &= e^{\pm \sum_{j=1}^{\mathfrak{g}} (\zeta(a_j) - \eta_1 a_j)} y_{\pm \mathbf{a}}(z) = e^{\mp 2\pi i s} y_{\pm \mathbf{a}}(z), \\ y_{\pm \mathbf{a}}(z + \tau) &= e^{\pm \tau \sum_{j=1}^{\mathfrak{g}} (\zeta(a_j) - \eta_2 a_j)} y_{\pm \mathbf{a}}(z) = e^{\pm 2\pi i r} y_{\pm \mathbf{a}}(z), \end{aligned}$$

namely  $y_{\pm \mathbf{a}}(z)$  are elliptic of the second kind. From Theorem 2.3 and  $\lambda_{\mathbf{a}} = \lambda_{-\mathbf{a}}$ , we have  $y_{\pm \mathbf{a}}(z)$  are solutions of  $\mathcal{L}_{\tau, \lambda_{\mathbf{a}}}^\mathfrak{g}$ , thus  $y_{\pm \mathbf{a}}(x + z_0)$  are solutions of  $L_\tau^\mathfrak{g} y = \lambda_{\mathbf{a}} y$ .

**Case 1.** If  $\mathbf{a}$  is not a branch point, i.e.,  $\mathbf{a} \cap -\mathbf{a} = \emptyset$ , then  $y_{\mathbf{a}}(x + z_0)$  and  $y_{-\mathbf{a}}(x + z_0)$  are linearly independent solutions of  $L_\tau^\mathfrak{g} y = \lambda_{\mathbf{a}} y$  and satisfy

$$y_{\pm \mathbf{a}}(x + z_0 + 1) = e^{\mp 2\pi i s} y_{\pm \mathbf{a}}(x + z_0),$$

Thus (2.2) gives that

$$(2.10) \quad \Delta(\lambda) = e^{-2\pi i s} + e^{2\pi i s} = e^{\sum_{j=1}^{\mathfrak{g}} (\zeta(a_j) - \eta_1 a_j)} + e^{-\sum_{j=1}^{\mathfrak{g}} (\zeta(a_j) - \eta_1 a_j)}.$$

**Case 2.** If  $\mathbf{a}$  is a branch point, i.e.,  $\mathbf{a} = -\mathbf{a}$ , then  $y_{\mathbf{a}} = y_{-\mathbf{a}}$ . It was proved in [6, Theorem 2.6] that there is a solution  $y_2(z)$  linearly independent with  $y_{\mathbf{a}}(z)$  such that

$$y_{\mathbf{a}}(z + 1) = e^{2\pi i s} y_{\mathbf{a}}(z), \quad y_2(z + 1) = e^{2\pi i s} y_2(z) + e^{2\pi i s} \chi_{\mathbf{a}} y_{\mathbf{a}}(z),$$

where

$$\begin{aligned}
\chi_{\mathbf{a}} &= -\sum_{i=1}^n c_{a_i} (\wp(a_i) + \eta_1), \\
(2.11) \quad c_{a_i} &= 2\wp''(a_i)^{-1} \prod_{j \neq i} (\wp(a_i) - \wp(a_j))^{-1}, \quad \text{if } a_i = -a_i, \\
c_{a_i} = c_{-a_i} &= \wp'(a_i)^{-2} \prod_{j \neq i, i^*} (\wp(a_i) - \wp(a_j))^{-1}, \quad \text{if } a_i \neq -a_i.
\end{aligned}$$

**2.4. Spectrum of the Lamé operator.** Now we can discuss the spectral of  $L_{\tau}^{\mathfrak{g}}$ . Applying the general result [10, Theorem 4.1] to  $L_{\tau}^{\mathfrak{g}}$ , we obtain

**Theorem 2.5.** [10, Theorem 4.1] *The spectrum  $\sigma(L_{\tau}^{\mathfrak{g}})$  consists of finitely many bounded simple analytic arcs  $\sigma_k$ ,  $1 \leq k \leq \mathfrak{g}$  for some  $\mathfrak{g} \leq g$  and one semi-infinite simple analytic arc  $\sigma_{\infty}$  which tends to  $-\infty + \langle q \rangle$ , with  $\langle q \rangle = \int_{x_0}^{x_0+1} q(x)dx$ , i.e.*

$$\sigma(L_{\tau}^{\mathfrak{g}}) = \sigma_{\infty} \cup \bigcup_{k=1}^{\mathfrak{g}} \sigma_k.$$

Furthermore,

- (1) *The finite end points of such arcs are exactly zeros of  $Q_g(\cdot; \tau)$  with odd order;*
- (2) *There are exactly  $d(\lambda)$ 's semi-arcs of  $\sigma(L_{\tau}^{\mathfrak{g}})$  meeting at each zero  $\lambda$  of  $Q_g(\cdot; \tau)$  and*

$$(2.12) \quad d(\lambda) = \text{ord}_{\lambda} Q_g(\cdot; \tau) + 2p_i(\lambda).$$

Recall another two results about the spectral of  $L_{\tau}^{\mathfrak{g}}$ :

**Theorem 2.6.** [7, Theorem 1.3] *Fix any  $\tau \in \mathbb{H}$  such that the multiplicity of any zero of  $Q_g(\lambda, \tau)$  is at most 2, then the following hold.*

- (1) *For any finite endpoint  $\lambda_{\mathbf{a}}$  of  $\sigma(L_{\tau}^{\mathfrak{g}})$ ,*

$$d(\lambda_{\mathbf{a}}) \geq 3 \quad \text{if and only if } \chi_{\mathbf{a}} = 0,$$

where  $\chi_{\mathbf{a}}$  is defined in (2.11).

- (2)  *$\sigma(L_{\tau}^{\mathfrak{g}})$  has at most one endpoint with  $d(\lambda) \geq 3$ .*

**Theorem 2.7.** *Let  $\tau = \frac{1}{2} + bi$  with  $b > 0$ .*

- (1) [4, Lemma 4.1] *The spectrum  $\sigma(L_{\tau}^{\mathfrak{g}})$  is symmetric with respect to the real line  $\mathbb{R}$ . In particular,  $\sigma_{\infty} \subseteq \mathbb{R}$ .*
- (2) [12, Theorem 2.2] *The complement  $\mathbb{C} \setminus \sigma(L_{\tau}^{\mathfrak{g}})$  is path-connected.*

By Theorem 2.6 and Theorem 2.7, we obtain the following corollary directly.

**Corollary 2.8.** *Fix any  $\tau = \frac{1}{2} + bi$  with  $b > 0$  such that the multiplicity of any zero of  $Q_g(\lambda; \tau)$  is at most 2, any non-real zero  $\lambda$  of  $Q_g(\lambda; \tau)$  with multiplicity 1 has  $d(\lambda) = 1$ .*

## 3. INNER INTERSECTION POINTS

In this and the following sections, we study the spectrum of the  $g = 3$  Lamé operator:

$$L_\tau^3 = \frac{d^2}{dx^2} - 12\wp(x + z_0; \tau), \quad x \in \mathbb{R}.$$

We will prove Theorem 1.1 recalled here.

**Theorem 3.1** (=Theorem 1.1). *Let  $\tau \in \mathbb{H}$ . If  $\lambda \in \sigma(L_\tau^3)$  and  $Q_3(\lambda; \tau) \neq 0$ , then  $\lambda$  is an inner intersection point if and only if  $\lambda$  satisfies*

$$f(\lambda) := \frac{4}{15}\lambda^3 + \frac{8}{5}\eta_1\lambda^2 - 3g_2\lambda + 9g_3 - 6\eta_1g_2 = 0.$$

*Proof.* Let  $\lambda_0 \in \sigma(L_\tau^3)$  with  $Q_3(\lambda_0; \tau) \neq 0$ . By (2.9) there is a small neighborhood  $U \subset \mathbb{C}$  of  $\lambda_0$  such that  $Q_3(\lambda; \tau) \neq 0$  for  $\lambda \in U$  and  $\lambda \in U$  can be a local coordinate for the hyperelliptic curve  $Y_\tau^3$ , namely  $a_1 = a_1(\lambda)$ ,  $a_2 = a_2(\lambda)$  and  $a_3 = a_3(\lambda)$  are holomorphic for  $\lambda \in U$ . For all  $\lambda \in U$ , (2.8) tells us that

$$\lambda = \lambda_{\mathbf{a}} = 5(\wp(a_1) + \wp(a_2) + \wp(a_3))$$

and then

$$\wp'(a_1)a_1'(\lambda) + \wp'(a_2)a_2'(\lambda) + \wp'(a_3)a_3'(\lambda) = \frac{1}{5} \quad \text{for } \lambda \in U$$

and so

$$(3.1) \quad (a_1'(\lambda_0), a_2'(\lambda_0), a_3'(\lambda_0)) \neq (0, 0, 0).$$

Next, note that  $Q_3(\lambda_0; \tau) \neq 0$  implies

$$\{a_1(\lambda), a_2(\lambda), a_3(\lambda)\} \cap \{-a_1(\lambda), -a_2(\lambda), -a_3(\lambda)\} = \emptyset \quad \text{for } \lambda \in U,$$

i.e.,  $\mathbf{a}(\lambda) = \{a_1(\lambda), a_2(\lambda), a_3(\lambda)\}$  is not a branch point of  $Y_\tau^3$  for all  $\lambda \in U$ . Hence

$$(3.2) \quad \wp(a_i(\lambda)) \neq \wp(a_j(\lambda)) \quad \text{for all } \lambda \in U, \quad 1 \leq i < j \leq 3,$$

and (2.7) holds for  $\lambda \in U$ , i.e.,

$$(3.3) \quad \begin{aligned} \wp'(a_1) + \wp'(a_2) + \wp'(a_3) &= 0, \\ \wp'(a_1)\wp(a_1) + \wp'(a_2)\wp(a_2) + \wp'(a_3)\wp(a_3) &= 0. \end{aligned}$$

By taking derivative with respect to  $\lambda$  in (3.3), we obtain from  $(\wp')^2 = 4\wp^3 - g_2\wp - g_3$  and  $\wp'' = 6\wp^2 - \frac{g_2}{2}$  that

$$(3.4) \quad \sum_{i=1}^3 (6\wp_i^2 - \frac{g_2}{2})a_i'(\lambda_0) = 0,$$

$$(3.5) \quad \sum_{i=1}^3 (10\wp_i^3 - \frac{3}{2}g_2\wp_i - g_3)a_i'(\lambda_0) = 0.$$

where  $\wp_i := \wp(a_i(\lambda_0))$  for  $i = 1, 2, 3$ . On the other hand, (3.3) gives us

$$(3.6) \quad \frac{4\wp_1^3 - g_2\wp_1 - g_3}{(\wp_2 - \wp_3)^2} = \frac{4\wp_2^3 - g_2\wp_2 - g_3}{(\wp_1 - \wp_3)^2} = \frac{4\wp_3^3 - g_2\wp_3 - g_3}{(\wp_1 - \wp_2)^2} =: \mathcal{U},$$

which is equivalent to

$$(3.7) \quad \begin{cases} 4\wp_1^3 - g_2\wp_1 - g_3 = \mathcal{U}(\wp_2 - \wp_3)^2, \\ 4\wp_2^3 - g_2\wp_2 - g_3 = \mathcal{U}(\wp_1 - \wp_3)^2, \\ 4\wp_3^3 - g_2\wp_3 - g_3 = \mathcal{U}(\wp_1 - \wp_2)^2. \end{cases}$$

Denote by

$$\begin{cases} s_1 = \wp_1 + \wp_2 + \wp_3, \\ s_2 = \wp_1\wp_2 + \wp_1\wp_3 + \wp_2\wp_3, \\ s_3 = \wp_1\wp_2\wp_3, \end{cases}$$

we have

$$(3.8) \quad (x - \wp_1)(x - \wp_2)(x - \wp_3) = x^3 - s_1x^2 + s_2x - s_3,$$

then (3.7) is equivalent to

$$(3.9) \quad \begin{cases} 4s_1\wp_1^2 - (4s_2 + g_2)\wp_1 + 4s_3 - g_3 = \mathcal{U}(\wp_2 - \wp_3)^2, \\ 4s_1\wp_2^2 - (4s_2 + g_2)\wp_2 + 4s_3 - g_3 = \mathcal{U}(\wp_1 - \wp_3)^2, \\ 4s_1\wp_3^2 - (4s_2 + g_2)\wp_3 + 4s_3 - g_3 = \mathcal{U}(\wp_1 - \wp_2)^2. \end{cases}$$

First, (3.9)+(3.10)+(3.11) gives us

$$(3.12) \quad \mathcal{U} = \frac{4(s_1^3 - 3s_1s_2 + 3s_3) - g_2s_1 - 3g_3}{2s_1^2 - 6s_2}.$$

Note that  $\wp_i \neq \wp_j$  for  $i \neq j$ , then (3.9)-(3.10), (3.9)-(3.11) and (3.10)-(3.11) yield

$$(3.13) \quad \begin{cases} 4s_2 + g_2 - 4s_1(\wp_1 + \wp_2) = \mathcal{U}(\wp_1 + \wp_2 - 2\wp_3), \\ 4s_2 + g_2 - 4s_1(\wp_1 + \wp_3) = \mathcal{U}(\wp_1 + \wp_3 - 2\wp_2), \\ 4s_2 + g_2 - 4s_1(\wp_2 + \wp_3) = \mathcal{U}(\wp_2 + \wp_3 - 2\wp_1). \end{cases}$$

$$(3.14)$$

$$(3.15)$$

Next, (3.13)+(3.14)+(3.15) gives us

$$(3.16) \quad s_2 = \frac{2}{3}s_1^2 - \frac{1}{4}g_2,$$

and (3.13)-(3.14) gives us

$$(3.17) \quad \mathcal{U} = -\frac{4}{3}s_1.$$

Combine (3.12), (3.16) and (3.17), we obtain that

$$(3.18) \quad s_3 = \frac{5}{9}s_1^3 - \frac{1}{3}g_2s_1 + \frac{1}{4}g_3.$$

On the other hand, for any  $\lambda \in U$ , denote by  $A(\lambda) := \sum_{j=1}^3 (\zeta(a_j) - \eta_1 a_j)$ . Since  $\mathbf{a}(\lambda) \cap -\mathbf{a}(\lambda) = \emptyset$ , by (2.10), we have, for  $\lambda \in U$ , that

$$(3.19) \quad \begin{aligned} \Delta(\lambda) &= e^A + e^{-A}, \\ \Delta'(\lambda) &= (e^A - e^{-A})A', \\ \Delta''(\lambda) &= (e^A - e^{-A})A'' + \Delta(A')^2, \\ \Delta'''(\lambda) &= (e^A - e^{-A})(A''' + (A')^3) + 3\Delta A' A''. \end{aligned}$$

**Sufficiency.** Let  $\lambda_0 \in \sigma(L_\tau^3)$  with  $Q_3(\lambda_0; \tau) \neq 0$  be an inner intersection point, then  $\lambda_0$  is met by  $2k \geq 4$  ( $k \in \mathbb{Z}$ ) semi-arcs of the spectrum.

Consider the local behavior of the spectrum at  $\lambda_0 \in \sigma(L_\tau^3)$ :

$$(3.20) \quad \Delta(\lambda) - \Delta(\lambda_0) = c(\lambda - \lambda_0)^k (1 + O(|\lambda - \lambda_0|)), \quad k \geq 1, \quad c \neq 0.$$

If  $\Delta(\lambda_0) \in (-2, 2)$ , it follows from (3.20) and  $\sigma(L_\tau^3) = \{\lambda \mid -2 \leq \Delta(\lambda) \leq 2\}$  that there are precisely  $2k$  semi-arcs of  $\sigma(L_\tau^3)$  meeting at  $\lambda_0$ . If  $\Delta(\lambda_0) = \pm 2$ , then there are precisely  $k$  semi-arcs of  $\sigma(L_\tau^3)$  meeting at  $\lambda_0$ .

If  $\Delta(\lambda_0) = \pm 2$ , then our assumption implies  $k \geq 4$ , i.e.  $\Delta'(\lambda_0) = \Delta''(\lambda_0) = \Delta'''(\lambda_0) = 0$ . Since  $\Delta(\lambda_0) = \pm 2$  implies  $e^A = \pm 1$  at  $\lambda_0$ , we obtain  $A'(\lambda_0) = 0$ .

If  $\Delta(\lambda_0) \in (-2, 2)$ , then our assumption implies  $2k \geq 4$ , i.e.  $k \geq 2$  and so  $\Delta'(\lambda_0) = 0$ . Since  $\Delta(\lambda_0) \neq \pm 2$  implies  $e^A \neq \pm 1$  at  $\lambda_0$ , again we obtain  $A'(\lambda_0) = 0$ .

Therefore, we always have  $A'(\lambda_0) = 0$ , i.e.,

$$(3.21) \quad (\wp_1 + \eta_1) a'_1(\lambda_0) + (\wp_2 + \eta_1) a'_2(\lambda_0) + (\wp_3 + \eta_1) a'_3(\lambda_0) = 0.$$

Noting from (3.1), We conclude from (3.4, 3.5, 3.21) that the determinant of the matrix

$$\Omega := \begin{pmatrix} \wp_1 + \eta_1 & \wp_2 + \eta_1 & \wp_3 + \eta_1 \\ 6\wp_1^2 - \frac{g_2}{2} & 6\wp_2^2 - \frac{g_2}{2} & 6\wp_3^2 - \frac{g_2}{2} \\ 10\wp_1^3 - \frac{3}{2}g_2\wp_1 - g_3 & 10\wp_2^3 - \frac{3}{2}g_2\wp_2 - g_3 & 10\wp_3^3 - \frac{3}{2}g_2\wp_3 - g_3 \end{pmatrix}$$

vanishes, i.e.,

$$(\wp_2 - \wp_1)(\wp_3 - \wp_1)(\wp_3 - \wp_2)(60s_3 + 60\eta_1s_2 + 5g_2s_1 - 6g_3 - 9g_2\eta_1) = 0.$$

By (3.2), we obtain that

$$(3.22) \quad 60s_3 + 60\eta_1s_2 + 5g_2s_1 - 6g_3 - 9g_2\eta_1 = 0.$$

Plug (3.16), (3.18) and

$$(3.23) \quad \lambda_0 = 5(\wp_1 + \wp_2 + \wp_3) = 5s_1$$

into (3.22), we finally obtain that

$$\frac{4}{15}\lambda_0^3 + \frac{8}{5}\eta_1\lambda_0^2 - 3g_2\lambda_0 + 9g_3 - 6\eta_1g_2 = 0.$$

**Necessity.** Suppose  $\lambda_0 \in \sigma(L_\tau^3)$  satisfies  $Q_3(\lambda_0; \tau) \neq 0$  and

$$\frac{4}{15}\lambda_0^3 + \frac{8}{5}\eta_1\lambda_0^2 - 3g_2\lambda_0 + 9g_3 - 6\eta_1g_2 = 0.$$

This, together with (3.16), (3.18) and (3.23) implies  $\det \Omega = 0$ . Since  $\wp_i \neq \wp_j$  for  $i \neq j$ , the second row of  $\Omega$  is nonzero. Suppose that the last two rows of  $\Omega$  are linearly dependent, there is  $c \in \mathbb{C}$  such that

$$10\wp_i^3 - \frac{3}{2}g_2\wp_i - g_3 = c(6\wp_i^2 - \frac{g_2}{2}), \quad i = 1, 2, 3,$$

then

$$(3.24) \quad (x - \wp_1)(x - \wp_2)(x - \wp_3) = x^3 - \frac{3}{5}cx^2 - \frac{3}{20}g_2x + \frac{g_2}{20}c - \frac{1}{10}g_3.$$

Compare (3.8) with (3.24), we have

$$(3.25) \quad \begin{aligned} s_1 &= \frac{3}{5}c, \\ s_2 &= -\frac{3}{20}g_2, \\ s_3 &= \frac{1}{10}g_3 - \frac{c}{20}g_2. \end{aligned}$$

Combine with (3.16) and (3.18), we obtain that

$$\lambda_0 = 5s_1 = \frac{9g_3}{2g_2} \quad \text{and} \quad 27g_3^2 = 5g_2^3.$$

Therefore, if  $\tau \in \mathbb{H} \setminus \{\tau \in \mathbb{H} \mid 27g_3^2(\tau) = 5g_2^3(\tau)\}$ , the last two rows of  $\Omega$  are linearly independent, thus the first row can be linearly spanned by the last two rows. Hence (3.4) and (3.5) yields (3.21). Finally, by the continuity of the left hand side in (3.21), we obtain (3.21) holds for all  $\tau \in \mathbb{H}$ .

If  $\Delta(\lambda_0) \in (-2, 2)$ , then we see from (3.21) and (3.19) that  $\Delta'(\lambda_0) = 0$ , i.e.  $k \geq 2$  in (3.20) and so there are  $2k \geq 4$  semi-arcs of  $\sigma(L_\tau^3)$  meeting at this  $\lambda_0$ . If  $\Delta(\lambda_0) = \pm 2$ , then  $e^A = \pm 1$  at  $\lambda_0$ . From here and (3.21, 3.19), we see that  $\Delta'(\lambda_0) = \Delta''(\lambda_0) = \Delta'''(\lambda_0) = 0$ . This means  $k \geq 4$  in (3.20) and so there are  $k \geq 4$  semi-arcs of  $\sigma(L_\tau^3)$  meeting at this  $\lambda_0$ . Therefore,  $\lambda_0$  is an inner intersection point.  $\square$

#### 4. ZEROS OF THE SPECTRAL POLYNOMIAL

In this and the following sections, we always assume  $\tau = \frac{1}{2} + bi$  with  $b > 0$ . In order to emphasize  $\tau = \frac{1}{2} + bi$ , we will use  $b$  instead of  $\tau$  in notations. Sometimes, we omit the notation  $\tau$  freely. We will first recall some basic properties for the quantities  $e_1, e_2, e_3, g_2, g_3$  and  $\eta_1$  associated with the Weierstrass elliptic function  $\wp(z; \tau)$ .

Note that  $e_1, g_2, \eta_1 \in \mathbb{R}$ ,  $\bar{e}_2 = e_3 \notin \mathbb{R}$  and  $e_1 + e_2 + e_3 = 0$ , in what follows, we set

$$(4.1) \quad e_1 = 2x, \quad e_2 = -x + iy, \quad e_3 = -x - iy \quad \text{with } x, y \in \mathbb{R} \text{ and } y \neq 0.$$

Since  $g_2 = 2(e_1^2 + e_2^2 + e_3^2)$  and  $g_3 = 4e_1e_2e_3$ , we have

$$(4.2) \quad \begin{aligned} g_2 &= 4(3x^2 - y^2), \\ g_3 &= 8x(x^2 + y^2) = 4e_1^3 - e_1g_2. \end{aligned}$$

Since  $e_1 \neq e_2 \neq e_3 \neq e_1$ , it is easy to see,

$$(4.3) \quad g_2 - 3e_k^2 = (e_i - e_j)^2 \neq 0, \quad \text{for } \{i, j, k\} = \{1, 2, 3\}.$$

In particular,

$$(4.4) \quad g_2 - 3e_1^2 = (e_2 - e_3)^2 = -4y^2 < 0, \quad \text{i.e., } g_2 < 3e_1^2.$$

When  $b$  goes to positive infinity, we have

$$(4.5) \quad e_1 \rightarrow \frac{2}{3}\pi^2, \quad \eta_1 \rightarrow \frac{1}{3}\pi^2, \quad g_2 \rightarrow \frac{4}{3}\pi^4, \quad g_3 \rightarrow \frac{8}{27}\pi^6.$$

And the derivatives of  $e_1, g_2$  and  $\eta_1$  with respect to  $b$  are as follows:

$$(4.6) \quad \begin{aligned} e_1'(b) &= \frac{1}{\pi} (e_1^2 - \eta_1 e_1 - \frac{1}{6}g_2), \quad (\text{see [6, (2.15)]}), \\ g_2'(b) &= \frac{1}{\pi} (3g_3 - 2\eta_1 g_2) = \frac{1}{\pi} (12e_1^3 - 3e_1 g_2 - 2\eta_1 g_2), \quad (\text{see [9]}), \\ \eta_1'(b) &= \frac{1}{24\pi} (g_2 - 12\eta_1^2), \quad (\text{see [7, (1.5)]}). \end{aligned}$$

By (4.2), we have

$$(4.7) \quad g_3'(b) = \frac{1}{\pi} (\frac{1}{6}g_2^2 - 3\eta_1 g_3) = \frac{1}{\pi} (\frac{1}{6}g_2^2 + 3e_1 \eta_1 g_2 - 12e_1^3).$$

Moreover, we have the following conclusions about the derivatives.

**Proposition 4.1.** [16, Theorem 1.7] *We have  $e_1(\frac{1}{2}) = 0$  and*

$$e_1'(b) > 0 \quad \text{for all } b > 0.$$

**Proposition 4.2.** [9, Corollary 4.4] *There exists  $b_g \approx 0.47 \in (\frac{1}{2\sqrt{3}}, \frac{1}{2})$  such that*

$$g_2'(b) \begin{cases} < 0 & \text{for } b \in (0, b_g), \\ = 0 & \text{for } b = b_g, \\ > 0 & \text{for } b \in (b_g, \infty). \end{cases}$$

*And  $g_2(b) = 0$  if and only if  $b \in \{\frac{1}{2\sqrt{3}}, \frac{\sqrt{3}}{2}\}$ .*

**Proposition 4.3.** [16, Theorem 1.7] *There exists  $b_\eta \approx 0.24108 < \frac{1}{2\sqrt{3}}$  such that*

$$\eta_1'(b) \begin{cases} > 0 & \text{for } b \in (0, b_\eta), \\ = 0 & \text{for } b = b_\eta, \\ > 0 & \text{for } b \in (b_\eta, +\infty). \end{cases}$$

**Remark 4.4.** All numerical computations use  $q = e^{2\pi i\tau} = -e^{-2\pi b}$  expansions of  $e_1, g_2, \eta_1$  which are recalled here for readers' convenience.

$$\begin{aligned} e_1(b) &= 16\pi^2 \left( \frac{1}{24} + \sum_{k=1}^{\infty} (-1)^k \sigma_0^k e^{-2k\pi b} \right), \quad \text{where } \sigma_0^k = \sum_{d|k, d \text{ is odd}} d, \\ g_2(b) &= 320\pi^4 \left( \frac{1}{240} + \sum_{k=1}^{\infty} (-1)^k \sigma_3^k e^{-2k\pi b} \right), \quad \text{where } \sigma_3^k = \sum_{1 \leq d|k} d^3, \\ \eta_1(b) &= 8\pi^2 \left( \frac{1}{24} - \sum_{k=1}^{\infty} (-1)^k \sigma_1^k e^{-2k\pi b} \right), \quad \text{where } \sigma_1^k = \sum_{1 \leq d|k} d. \end{aligned}$$

Now we will figure out all zeros of the spectral polynomial  $Q_3(\lambda; b)$  and then prove Theorem 1.2.

*Proof.* Recall that [cf. (1.2)]

$$Q_3(\lambda) := Q_3(\lambda; b) = \lambda \prod_{k=1}^3 (\lambda^2 - 6e_k \lambda + 15(3e_k^2 - g_2)).$$

First of all, by (4.3), 0 is not a zero of

$$R_k(\lambda) := \lambda^2 - 6e_k \lambda + 15(3e_k^2 - g_2), \quad k = 1, 2, 3.$$

So 0 is a simple zero of  $Q_3(\lambda)$ .

Next, if  $\lambda \in \mathbb{C}$  satisfies  $Q_3(\lambda) = 0$ , by (2.8) and (2.9), there exists *unique* (cf. [3, Proposition 6.4])  $\mathbf{a} = \{a_1, a_2, a_3\} \in Y_b^3$  with  $\mathbf{a} = -\mathbf{a}$  such that

$$(4.8) \quad \lambda = \lambda_{\mathbf{a}} = 5(\wp(a_1) + \wp(a_2) + \wp(a_3)).$$

Note that  $\mathbf{a} = -\mathbf{a}$  with  $\mathbf{a} \in Y_b^3$  yields that

$$(4.9) \quad \begin{aligned} \mathbf{a} &= \left\{ \frac{\omega_1}{2}, \frac{\omega_2}{2}, \frac{\omega_3}{2} \right\} \quad \text{or} \\ \mathbf{a} &= \left\{ \frac{\omega_k}{2}, a, -a \right\} \quad \text{for some } a \in T_{\tau}^* \setminus \left\{ \frac{\omega_1}{2}, \frac{\omega_2}{2}, \frac{\omega_3}{2} \right\}, \quad k = 1, 2, 3. \end{aligned}$$

First, if  $\mathbf{a} = \left\{ \frac{\omega_1}{2}, \frac{\omega_2}{2}, \frac{\omega_3}{2} \right\}$ , then

$$\lambda_{\left\{ \frac{\omega_1}{2}, \frac{\omega_2}{2}, \frac{\omega_3}{2} \right\}} = 5(e_1 + e_2 + e_3) = 0.$$

By [7, Example 3.4],

$$\chi_{\left\{ \frac{\omega_1}{2}, \frac{\omega_2}{2}, \frac{\omega_3}{2} \right\}} = \frac{12(3g_3 - 2\eta_1 g_2)}{g_2^3 - 27g_3^2} = \frac{12\pi g_2'}{g_2^3 - 27g_3^2},$$

where  $g_2^3 - 27g_3^2 = 16(e_1 - e_2)^2(e_1 - e_3)^2(e_2 - e_3)^2 \neq 0$ . Proposition 4.2 tells us that  $\chi_{\left\{ \frac{\omega_1}{2}, \frac{\omega_2}{2}, \frac{\omega_3}{2} \right\}} = 0$  if and only if  $b = b_g$ . By Theorem 2.6 and the multiplicity of 0 is 1, we obtain that  $d(0) \geq 3$  if and only if  $b = b_g$ , otherwise,  $d(0) = 1$ .

Next, if  $\mathbf{a} = \left\{ \frac{\omega_k}{2}, a, -a \right\}$ , it is well known [17, Theorem 5.3] that  $\lambda_{\mathbf{a}}$  is a zero of  $R_k(\lambda)$  for  $k = 1, 2, 3$  respectively. By the uniqueness of  $\mathbf{a}$ ,  $R_k(\lambda)$  and  $R_{k'}(\lambda)$  cannot have a common zero for any pair  $k \neq k' \in \{1, 2, 3\}$ . Since



$\overline{R_2(\lambda)} = R_3(\lambda)$ , we obtain that the two zeros  $\mu, \nu$  of  $R_2(\lambda)$  are not real, i.e.,  $\mu, \nu \in \mathbb{C} \setminus \mathbb{R}$ , and then  $\overline{\mu}, \overline{\nu} \in \mathbb{C} \setminus \mathbb{R}$  are the zeros of  $R_3(\lambda)$ .

We claim that  $\mu \neq \nu$ , otherwise, the discriminant of  $R_2(\lambda)$  vanishes, i.e.,

$$(6e_2)^2 - 60(3e_2^2 - g_2) = 0,$$

which gives us

$$5g_2 = 12e_2^2 \in \mathbb{R},$$

then (4.1) yields  $x = 0$  and thus  $e_1 = 0, e_3 = -e_2$ . Combine with (2.4), we have

$$10(e_2^2 + e_2^2) = 12e_2^2,$$

which is a contradiction because  $e_2 \neq 0$ !

Therefore, the multiplicity of any zero of  $Q_3(\lambda; b)$  is at most 2, and a zero of  $Q_3(\lambda; b)$  is of multiplicity 2 if and only if it is a zero of  $R_1(\lambda)$ . Denote by  $\vartheta_{\pm} := 3e_1 \pm \frac{1}{2}\sqrt{\Delta_{R_1}}$  the zeros of

$$R_1(\lambda) = \lambda^2 - 6e_1\lambda + 15(3e_1^2 - g_2),$$

where  $\Delta_{R_1}$  denotes the discriminant of  $R_1(\lambda)$ , i.e.,

$$\Delta_{R_1} := 12(5g_2 - 12e_1^2) = 48(3x^2 - 5y^2).$$

Note that  $g_2 < 3e_1^2$ , then  $\vartheta_+$  and  $\vartheta_-$  have the same sign when  $\Delta_{R_1} > 0$ . By Corollary 2.8, we have  $d(\mu) = d(\nu) = d(\overline{\mu}) = d(\overline{\nu}) = 1$  and  $d(\vartheta_{\pm}) = 1$  if  $\Delta_{R_1} < 0$ .

Clearly,  $\Delta_{R_1}$  is holomorphic with respect to  $b$ . By Proposition 4.1 and Proposition 4.2,  $\Delta_{R_1}$  is strictly increasing for  $b \in [b_g, \frac{1}{2}]$ , i.e.,

$$\frac{d\Delta_{R_1}}{db} > 0, \quad \text{for } b \in \left[ b_g, \frac{1}{2} \right].$$

By (4.6), we have

$$\begin{aligned} \frac{d\Delta_{R_1}}{db} &= \frac{12}{\pi} (15g_3 - 10\eta_1g_2 - 24e_1^3 + 24\eta_1e_1^2 + 4e_1g_2) \\ &= \frac{12}{\pi} (36e_1^3 - 11e_1g_2 - 10\eta_1g_2 + 24e_1^2\eta_1). \end{aligned}$$

For  $b \in \left( \frac{1}{2}, \frac{\sqrt{3}}{2} \right]$ , since  $e_1 > 0$  and  $g_2 \leq 0$ , we have

$$\frac{d\Delta_{R_1}}{db} > 0, \quad \text{for } b \in \left( \frac{1}{2}, \frac{\sqrt{3}}{2} \right].$$

For  $b \in \left( \frac{\sqrt{3}}{2}, +\infty \right)$ , note that

$$\begin{aligned} \frac{d\Delta_{R_1}}{db} &= \frac{12}{\pi} (36e_1^3 - 11e_1g_2 - 10\eta_1g_2 + 24e_1^2\eta_1), \\ &= \frac{96}{\pi} (3x^3 + 11xy^2 + (5y^2 - 3x^2)\eta_1), \end{aligned}$$

and  $g_2(b), g_2'(b) > 0$  gives us

$$\eta_1 < \frac{3g_3}{2g_2} = \frac{3x(x^2 + y^2)}{3x^2 - y^2}.$$

If  $5y^2 - 3x^2 \geq 0$ , it is clearly that  $\Delta'_{R_1}(b) > 0$ . If  $5y^2 - 3x^2 < 0$ , we have

$$\begin{aligned} \frac{d\Delta_{R_1}}{db} &= \frac{96}{\pi} (3x^3 + 11xy^2 + (5y^2 - 3x^2) \eta_1), \\ &> \frac{96}{\pi} \left( 3x^3 + 11xy^2 + (5y^2 - 3x^2) \frac{3x(x^2 + y^2)}{3x^2 - y^2} \right) \\ &= \frac{384(9x^2y^2 + y^4)}{\pi(3x^2 - y^2)} > 0. \end{aligned}$$

Hence

$$\frac{d\Delta_{R_1}}{db} > 0, \quad \text{for } b \in \left( \frac{\sqrt{3}}{2}, +\infty \right).$$

Note that

$$\lim_{b \rightarrow +\infty} \Delta_{R_1}(b) = 12 \left( \frac{20\pi^4}{3} - \frac{48\pi^2}{9} \right) = 16\pi^4 > 0,$$

and

$$\Delta_{R_1}(b) = 12(5g_2 - 12e_1^2) < 0 \quad \text{for } b \in \left[ \frac{1}{2\sqrt{3}}, \frac{\sqrt{3}}{2} \right],$$

then there exists unique  $\beta \in \left[ \frac{1}{2\sqrt{3}}, +\infty \right)$  such that

$$\Delta_{R_1}(b) \begin{cases} < 0, & \text{for } b \in \left( \frac{1}{2\sqrt{3}}, \beta \right), \\ = 0, & \text{for } b = \beta, \\ > 0, & \text{for } b \in (\beta, +\infty). \end{cases}$$

In fact,  $\beta \approx 1.0979 > \frac{\sqrt{3}}{2}$  by numerical computation.

In addition, since  $\tau = \frac{1}{2} + ib$ , we have

$$\frac{\tau - 1}{2\tau - 1} = \frac{1}{2} + i\frac{1}{4b} \quad \text{for } \tau = \frac{1}{2} + ib.$$

It is well know that [8, p. 346]

$$(4.10) \quad \begin{aligned} e_1\left(\frac{1}{4b}\right) &= -4b^2 e_1(b), \\ g_2\left(\frac{1}{4b}\right) &= 16b^4 g_2(b), \end{aligned}$$

then

$$\Delta_{R_1}(b) = \frac{1}{16b^4} \Delta_{R_1}\left(\frac{1}{4b}\right),$$

so there exists unique  $\widehat{\beta} = \frac{1}{4\beta} \approx 0.2277 \in \left(0, \frac{1}{2\sqrt{3}}\right)$  such that

$$\Delta_{R_1}(b) \begin{cases} < 0, & \text{for } b \in \left(\widehat{\beta}, \frac{1}{2\sqrt{3}}\right), \\ = 0, & \text{for } b = \widehat{\beta}, \\ > 0, & \text{for } b \in (0, \widehat{\beta}). \end{cases}$$

Moreover,

$$\Delta'_{R_1}(b) < 0 \quad \text{for } b \in \left(0, \frac{1}{2\sqrt{3}}\right).$$

In a word,

$$\Delta_{R_1}(b) \begin{cases} > 0, & \text{for } b \in (0, \widehat{\beta}) \cup (\beta, +\infty), \\ = 0, & \text{for } b \in \{\beta, \widehat{\beta}\}, \\ < 0, & \text{for } b \in (\widehat{\beta}, \beta). \end{cases}$$

For  $b \in (\widehat{\beta}, \beta)$ ,  $\vartheta_- = \overline{\vartheta_+} \notin \mathbb{R}$ . By Corollary 2.8, we have

$$d(\vartheta_+) = d(\vartheta_-) = 1.$$

For  $b \in \{\widehat{\beta}, \beta\}$ , note that  $3e_1 = \vartheta_+ = \vartheta_-$  is of multiplicity 2 and

$$d(3e_1) = 2 + p_i(3e_1).$$

From (4.8) and (4.9), there exists  $\theta \in T_\tau^* \setminus \left\{\frac{\omega_1}{2}, \frac{\omega_2}{2}, \frac{\omega_3}{2}\right\}$  such that

$$3e_1 = \lambda_{\left\{\frac{\omega_1}{2}, \theta, -\theta\right\}} = 5(e_1 + 2\wp(\theta)),$$

thus

$$(4.11) \quad \wp(\theta) = -\frac{1}{5}e_1.$$

At the same time, the discriminant  $\Delta_{R_1} = 0$  gives us

$$(4.12) \quad g_2 = \frac{12}{5}e_1^2,$$

and then

$$(4.13) \quad g_3 = 4e_1^3 - e_1g_2 = \frac{8}{5}e_1^3.$$

Note that  $\wp'^2 = 4\wp^3 - g_2\wp - g_3$  and  $\wp'' = 6\wp^2 - g_2/2$ , by (2.11), we have

$$-\frac{1}{2}(e_1 - \wp(\theta))\chi_{\left\{\frac{\omega_1}{2}, \theta, -\theta\right\}} = \frac{e_1 + \eta_1}{(6e_1^2 - g_2/2)(e_1 - \wp(\theta))} - \frac{\wp(\theta) + \eta_1}{4\wp(\theta)^3 - g_2\wp(\theta) - g_3}.$$

Combine with (4.11, 4.12, 4.13), we obtain that

$$-\frac{3}{5}e_1\chi_{\left\{\frac{\omega_1}{2}, \theta, -\theta\right\}} = \frac{25}{24}\eta_1 > 0,$$

so  $\chi_{\left\{\frac{\omega_1}{2}, \theta, -\theta\right\}} \neq 0$  which means  $d(3e_1) \leq 2$  by Theorem 2.6. Apply (2.12), we obtain that

$$d(3e_1) = 2 \quad \text{for } b \in \{\widehat{\beta}, \beta\}.$$

Consequently, we summary the above discussion into Theorem 1.2.  $\square$

Finally, from Theorem 1.2, Theorem 2.5 and Theorem 2.7, we have the following proposition.

**Proposition 4.5.** For  $b \in (\frac{1}{4\beta}, +\infty)$ , we have  $(-\infty, 0] \subseteq \sigma(L_b^3)$ .

## 5. PROOF OF THE MAIN THEOREM

Recall that

$$f(\lambda) = \frac{4}{15}\lambda^3 + \frac{8}{5}\eta_1\lambda^2 - 3g_2\lambda + 9g_3 - 6\eta_1g_2.$$

By Proposition 4.2, we have

$$f(0) = 3\pi g_2'(b) \begin{cases} < 0 & \text{for } b \in (0, b_g), \\ = 0 & \text{for } b = b_g, \\ > 0 & \text{for } b \in (b_g, +\infty). \end{cases}$$

From here and Proposition 4.5, we obtain the following lemma.

**Lemma 5.1.** If  $b \in (b_g, +\infty)$ , there exists  $\lambda_- < 0$  such that  $\lambda_-$  is an inner intersection point of  $\sigma(L_b^3)$ .


Now, let us prove the main Theorem 1.3.

*Proof. Step1: The spectrum at  $b = \beta$ .* By Theorem 1.2, the zeros of the spectral polynomial  $Q_3(\lambda; b)$  are


$$0, \mu, \bar{\mu}, \nu, \bar{\nu}, \vartheta_+ = \vartheta_- = 3e_1,$$

and  $d(0) = d(\mu) = d(\bar{\mu}) = d(\nu) = d(\bar{\nu}) = 1, d(3e_1) = 2$ .

From Theorem 2.5, Theorem 2.7, Proposition 4.5 and Proposition 5.1, we obtain the rough graph of  $\sigma(L_\beta^3)$  is as follows:



**Step2: The spectrum for  $b > \beta$ .** From Theorem 2.5, Theorem 2.7, Proposition 4.5 and Proposition 5.1, we obtain the graph of  $\sigma(L_b^3)$  consisting of the following two parts.



Next, we need the following lemma.

**Lemma 5.2.** *There exists  $\lambda_+ \in (\vartheta_-, \vartheta_+)$  such that  $f(\lambda_+) = 0$ .*

*Proof.* Recall that

$$\begin{cases} R_1(\lambda) = \lambda^2 - 6e_1\lambda + 15(3e_1^2 - g_2) = (\lambda - \vartheta_-)(\lambda - \vartheta_+), \\ f(\lambda) = \frac{4}{15}\lambda^3 + \frac{8}{5}\eta_1\lambda^2 - 3g_2\lambda + 9g_3 - 6\eta_1g_2, \end{cases}$$

and  $g_3 = 4e_1^3 - e_1g_2$ , we have

$$(5.1) \quad \begin{cases} \vartheta_- + \vartheta_+ = 6e_1, \\ \vartheta_- \vartheta_+ = 15(3e_1^2 - g_2), \end{cases}$$

and

$$\begin{aligned} f(\vartheta_\pm) &= \left( \frac{4}{15}\vartheta_\pm + \frac{8}{5}\eta_1 \right) (6e_1\vartheta_\pm - 15(3e_1^2 - g_2)) - 3g_2\vartheta_\pm + 9g_3 - 6\eta_1g_2 \\ &= \frac{8}{5}e_1(6e_1\vartheta_\pm - 15(3e_1^2 - g_2)) + \left( g_2 + \frac{48}{5}\eta_1e_1 - 12e_1^2 \right) \vartheta_\pm + 9g_3 + 18\eta_1g_2 - 72e_1^2\eta_1 \\ &= \left( g_2 + \frac{48}{5}\eta_1e_1 - \frac{12}{5}e_1^2 \right) \vartheta_\pm + 3(6g_2\eta_1 + 5g_2e_1 - 72\eta_1e_1^2 - 36e_1^3) \\ &:= B(\vartheta_\pm - D), \end{aligned}$$

where

$$(5.2) \quad \begin{aligned} D &= -3 \frac{6g_2\eta_1 + 5g_2e_1 - 72\eta_1e_1^2 - 36e_1^3}{g_2 + \frac{48}{5}\eta_1e_1 - \frac{12}{5}e_1^2} \\ &= -15e_1 + \frac{15}{2} \frac{\eta_1(g_2 - 12e_1^2)}{e_1^2 - 4e_1\eta_1 - \frac{5}{12}g_2}. \end{aligned}$$

Combine (5.1) and (5.2), we have


$$\begin{aligned} &f(\vartheta_-)f(\vartheta_+) \\ &= B^2(\vartheta_- - D)(\vartheta_+ - D) \\ &= B^2(D^2 - 6e_1D + 15(3e_1^2 - g_2)) \\ &= B^2((D - 3e_1)^2 - 3(5g_2 - 12e_1^2)) \\ &= B^2 \left( \left( -18e_1 + \frac{15}{2} \frac{\eta_1(g_2 - 12e_1^2)}{e_1^2 - 4e_1\eta_1 - \frac{5}{12}g_2} \right)^2 - 3(5g_2 - 12e_1^2) \right) \\ &= \frac{3B^2(5g_2 - 12e_1^2)}{4(e_1^2 - 4e_1\eta_1 - \frac{5}{12}g_2)^2} \left( 3(5g_2 - 12e_1^2)(e_1 + \eta_1)^2 - 4(e_1^2 - 4e_1\eta_1 - \frac{5}{12}g_2)^2 \right) \\ &= - \frac{\pi B^2(5g_2 - 12e_1^2)}{24(e_1^2 - 4e_1\eta_1 - \frac{5}{12}g_2)^2} (12(e_1 + \eta_1)g_2' - 5(12e_1^2 - g_2)\eta_1') < 0 \end{aligned}$$

for all  $b > \beta$ , because  $e_1 > 0, \eta_1 > 0, g_2' > 0, \eta_1' < 0$  and  $g_2 < 3e_1^2$  for  $b > \beta$ . Therefore, there exists  $\lambda_+ \in (\vartheta_-, \vartheta_+)$  such that  $f(\lambda_+) = 0$ .  $\square$

From Theorem 1.1 and this lemma,  $\lambda_+$  is an inner intersection point and then

$$\lambda_+ = \ell \cap \mathbb{R} \in (\vartheta_-, \vartheta_+),$$

so the rough graph of spectrum  $\sigma(L_b^3)$  for  $b > \beta$  is as follows.



**Step3: The spectrum for  $b$  around  $\hat{\beta}$ .** Consider

$$\begin{cases} f(\lambda) = \frac{4}{15}\lambda^3 + \frac{8}{5}\eta_1\lambda^2 - 3g_2\lambda - 6\eta_1g_2 + 9g_3 \\ f'(\lambda) = \frac{4}{5}\lambda^2 + \frac{16}{5}\eta_1\lambda - 3g_2 := \frac{4}{5}(\lambda - \lambda_L)(\lambda - \lambda_R), \end{cases}$$

we have

$$\begin{cases} \lambda_L + \lambda_R = -4\eta_1, \\ \lambda_L\lambda_R = -\frac{15}{4}g_2, \end{cases}$$

and

$$\begin{aligned} f(\lambda) &= \frac{\lambda}{3} \left( 3g_2 - \frac{16}{5}\eta_1\lambda \right) + \frac{8}{5}\eta_1\lambda^2 - 3g_2\lambda - 6\eta_1g_2 + 9g_3 \\ &= \frac{8}{15}\eta_1\lambda^2 - 2g_2\lambda - 6\eta_1g_2 + 9g_3 \\ &= \frac{2}{3}\eta_1 \left( 3g_2 - \frac{16}{5}\eta_1\lambda \right) - 2g_2\lambda - 6\eta_1g_2 + 9g_3 \\ &= -2 \left( \frac{16}{15}\eta_1^2 + g_2 \right) \lambda - 4\eta_1g_2 + 9g_3 \\ &:= -\frac{5}{24}\Delta_{f'}(\lambda - H), \end{aligned}$$

where

$$H = \frac{24(9g_3 - 4\eta_1g_2)}{5\Delta_{f'}}.$$

Then

$$\begin{aligned}
f(\lambda_L)f(\lambda_R) &= \frac{25}{24^2} \Delta_{f'}^2 (H - \lambda_L) (H - \lambda_R) \\
&= \frac{25}{24^2} \Delta_{f'}^2 \left( H^2 + 4\eta_1 H - \frac{15}{4} g_2 \right) \\
&= \frac{25}{24^2} \Delta_{f'}^2 \left( (H + 2\eta_1)^2 - \frac{25}{64} \Delta_{f'} \right) \\
&= \frac{25}{24^2} \Delta_{f'}^2 \left( \frac{\left( \frac{64}{15} \eta_1^3 + 9g_3 \right)^2}{\frac{25}{24^2} \Delta_{f'}^2} - \frac{25}{64} \Delta_{f'} \right) \\
&= \left( \frac{64}{15} \eta_1^3 + 9g_3 \right)^2 - \frac{25^2}{24^2 * 64} \Delta_{f'}^3 \\
&= \frac{1}{225} \left( (64\eta_1^3 + 135g_3)^2 - (16\eta_1^2 + 15g_2)^3 \right).
\end{aligned}$$

Clearly, all zeros of  $f$  are real if and only if  $f(\lambda_L)f(\lambda_R) \leq 0$ .


First of all, if  $b = \widehat{\beta}$ , then  $\vartheta_- = \vartheta_+ = 3e_1 < 0$  and  $\Delta_{R_1}(\widehat{\beta}) = 0$ , i.e.,  $5g_2 = 12e_1^2$ , thus  $5g_3 = 8e_1^3$ . By directly computation, we have  $f(3e_1) = 0$  and  $f'(3e_1) = 4e_1\eta_1/5 < 0$  because  $e_1(\widehat{\beta}) < 0$  and  $\eta_1(\widehat{\beta}) > 0$ . Note that  $f(0) < 0$ , then  $f$  has two negative real zeros  $\lambda_-^1 < \lambda_-^2 = 3e_1$  and one positive real zero  $\lambda_+$ , thus  $f(\lambda_L)f(\lambda_R)|_{b=\widehat{\beta}} < 0$ . Note that  $f(\lambda_L)f(\lambda_R)|_{b=b_\eta} > 0$  by numerical computation, we can define

$$\alpha := \sup \left\{ \widetilde{b} > \widehat{\beta} \mid f(\lambda_L)f(\lambda_R)|_{b=b'} < 0 \text{ for all } b' \in [\widehat{\beta}, \widetilde{b}) \right\}.$$


Clearly,  $\widehat{\beta} < \alpha < b_\eta$  and  $f(\lambda_L)f(\lambda_R)|_{b=\alpha} = 0$ . In fact,  $\alpha \approx 0.23217$  by numerical computation and  $f(\lambda_L)f(\lambda_R)|_{\alpha < b < \alpha + \varepsilon} > 0$  with  $\varepsilon > 0$  sufficient small.

For  $b \in [\widehat{\beta}, \alpha)$ , all zeros of  $f$  are real because  $f(\lambda_L)f(\lambda_R)|_{[\widehat{\beta}, \alpha)} < 0$ . Since  $f(0) = 3\pi g_2'(b) < 0$  for  $b \in (0, \gamma_1) \subseteq (0, b_g)$  and  $f$  has two negative real zeros at  $b = \widehat{\beta}$ , by continuity,  $f$  has two negative real zeros  $\lambda_-^1, \lambda_-^2$  for  $b \in [\widehat{\beta}, \alpha)$  and  $\lambda_-^1 = \lambda_-^2$  is a negative real zero of  $f$  with multiplicity 2 at  $b = \alpha$ .


For  $b \in (\widehat{\beta}, \alpha)$ , we have  $(-\infty, 0] \subseteq \sigma(L_b^3)$  and  $\lambda_-^1 < \lambda_-^2 < 0$ , thus  $\lambda_-^1$  and  $\lambda_-^2$  are two inner intersection points. Consequently, the spectrum  $\sigma(L_b^3)$  is one of the following graph.




When  $b$  goes to  $\alpha_1$  from the left hand side, we have (Fa) will goes to (Ga), (Fb) and (Fc) will goes to either (Gb) or (Gc) because all zeros of  $f$  are real.




Note that  $\lambda_-$  can not disappear in (Gb) and (Gc) when  $b$  moves to the right hand side of  $\alpha_1$ , which contradicts with  $f(\lambda_L)f(\lambda_R) > 0$  and  $f(0) < 0$ , i.e.,  $f$  has no negative real zero for  $b \in (\alpha, \alpha + \varepsilon)$ . Therefore, the rough graph of  $\sigma(L_b^3)$  at  $b = \alpha$  is (Ga), and then the rough graph of  $\sigma(L_b^3)$  for  $b \in (\hat{\beta}, \alpha)$  is (Fa). Consequently, the rough graph of  $\sigma(L_b^3)$  for  $b \in (\alpha, \alpha + \varepsilon)$  with  $\varepsilon > 0$  sufficient small is the following:



Now let us go back to  $b = \hat{\beta}$ , we have  $d(3e_1) = 2$  and  $\lambda_-^2 = 3e_1$ . From (Fa), we obtain the rough graph of  $\sigma(L_b^3)$  at  $b = \hat{\beta}$  as follows:



Furthermore, if  $b \in (\hat{\beta} - \delta)$  with  $\delta > 0$  sufficient small, then  $\vartheta_- < 3e_1 < \vartheta_+ < 0$  and  $\lambda_-^1$  still lies on  $\sigma(L_b^3)$ . Hence, the rough graph of  $\sigma(L_b^3)$  is the following.



□



## REFERENCES

- [1] V. Batchenko and F. Gesztesy; *On the spectrum of Schrödinger operators with quasi-periodic algebro-geometric KdV potentials*. J. Anal. Math. **95** (2005), 333-387.
- [2] B. Birnir; *Complex Hill's equation and the complex periodic Korteweg-de Vries equations*. Comm. Pure Appl. Math. **39** (1986), 1-49.
- [3] C.L. Chai, C.S. Lin and C.L. Wang; *Mean field equations, Hyperelliptic curves, and Modular forms: I*. Camb. J. Math. **3** (2015), 127-274.
- [4] Z. Chen, E. Fu and C.S. Lin; *Spectrum of the Lamé operator and application, I: Deformation along  $Re\tau=1/2$* . Adv. Math. **383** (2021), Paper No. 107699, 36 pp.
- [5] Z. Chen, E. Fu and C.S. Lin; *A necessary and sufficient condition for the Darboux-Treibich-Verdier potential with its spectrum contained in  $\mathbb{R}$* . arXiv:2001.11244, to appear in Amer. J. Math.
- [6] Z. Chen and C.S. Lin; *On algebraic multiplicity of (anti)periodic eigenvalues of Hill's equations*. Proc. Am. Math. Soc. **146** (2018), 3039-3047.
- [7] Z. Chen and C.S. Lin; *Critical points of the classical Eisenstein series of weight two*. J. Differ. Geom. **113** (2019), 189-226.
- [8] Z. Chen and C.S. Lin; *Spectrum of the Lamé operator and application, II: When an endpoint is a cusp*. Comm. Math. Phys. **378** (2020), 335-368.
- [9] Z. Chen and C.S. Lin; *Distribution of critical points of the classical Eisenstein series  $E_4$* . preprint.
- [10] F. Gesztesy and R. Weikard; *Picard potentials and Hill's equation on a torus*. Acta Math. **176** (1996), 73-107.
- [11] F. Gesztesy and R. Weikard; *Lamé potentials and the stationary (m)KdV hierarchy*. Math. Nachr. **176** (1995), 73-91.
- [12] F. Gesztesy and R. Weikard; *Floquet theory revisited*. Differential equations and mathematical physics, 67-84, Int. Press, Boston, MA, 1995.
- [13] W. Haese-Hill, M. Hallnäs and A. Veselov; *On the spectra of real and complex Lamé operators*. Symm. Integ. Geom. Meth. Appl. SIGMA. **13** (2017), 049, 23 pages.
- [14] E.L. Ince; *Further investigations into the periodic Lamé equations*. Proc. Roy. Soc. Edinb. **60** (1940), 83-99.
- [15] G. Lamé; *Sur les surfaces isothermes dans les corps homogènes en équilibre de température*. J. Math. Pures Appl. **2** (1837), 147-188.
- [16] C.S. Lin and C.L. Wang; *Elliptic functions, Green functions and the mean field equations on tori*. Ann. Math. **172** (2010), 911-954.
- [17] C.S. Lin and C.L. Wang; *Mean field equations, Hyperelliptic curves, and Modular forms: II*. J. Éc. polytech. Math. **4** (2017), 557-593.
- [18] M.A. Naimark; *Linear differential operators* (Frederick Ungar, New York, 1967).
- [19] F. Rofe-Beketov; *The spectrum of non-selfadjoint differential operators with periodic coefficients*. Soviet Math. Dokl. **4** (1963) 1563-1566.
- [20] E.T. Whittaker and G.N. Watson; *A course of modern analysis, 4th edition*. Cambridge University Press, 1927.

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, 100084, CHINA

Email address: ejfu@tsinghua.edu.cn; 473051732@qq.com