Triple-charm molecular states composed of D^*D^*D and $D^*D^*D^*$

Si-Qiang Luo,^{1,2} Tian-Wei Wu,³ Ming-Zhu Liu,^{4,5} Li-Sheng Geng,^{5,6,7,8,*} and Xiang Liu^{1,8,9,10,†}

¹School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China

²School of mathematics and statistics, Lanzhou University, Lanzhou 730000, China

³School of Fundamental Physics and Mathematical Sciences,

Hangzhou Institute for Advanced Study, UCAS, Hangzhou, 310024, China

⁴School of Space and Environment, Beihang University, Beijing 102206, China

⁵School of Physics, Beihang University, Beijing 102206, China

⁶Beijing Key Laboratory of Advanced Nuclear Materials and Physics, Beihang University, Beijing, 102206, China

School of Physics and Microelectronics, Zhengzhou University, Zhengzhou, Henan 450001, China

⁸Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, China

⁹Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China

¹⁰Key Laboratory of Theoretical Physics of Gansu Province,

and Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, China

Inspired by the newly observed T_{cc}^+ state, we systematically investigate the *S*-wave triple-charm molecular states composed of D^*D^*D and $D^*D^*D^*$. We employ the one-boson-exchange model to derive the interactions between $D(D^*)$ and D^* and solve the three body Schrödinger equations with the Gaussian expansion method. Our results show that the *S*-wave $I(J^P) = \frac{1}{2}(0^-, 1^-, 2^-) D^*D^*D$ and $I(J^P) = \frac{1}{2}(0^-, 1^-, 2^-) D^*D^*D^*$ systems could form bound states, which can be viewed as three-body hadronic molecules. We present not only the binding energies of the three-body bound states, but also the root-mean-square radii of D- D^* and D^* - D^* . These predictions could be tested in the future at LHC or HL-LHC.

I. INTRODUCTION

As important members of the hadron family, exotic states have always interested both theoreticians and experimentalists. By definition, exotic states contain more complex quark and gluon contents than the conventional $q\bar{q}$ mesons and qqqbaryons. Given their peculiar nature, studies of exotic states have been a hot topic in hadron physics.

Among the various exotic states, hadronic molecules are quite distinct. They are loosely bound states composed of two or several conventional hadrons and provide a good laboratory to study hadron structure and non-perturbative strong interactions at hadronic level. In 2003, the Babar Collaboration observed a charmed-strange state $D_{s0}^*(2317)$ in the $D_s\pi^0$ channel [1]. Soon after, the CLEO Collaboration not only confirmed the existence of the $D_{s0}^{*}(2317)$, but also found a new charmed-strange state $D'_{s1}(2460)$ in the $D^*_s\pi^0$ mass spectrum [2]. In the same year, the Belle Collaboration reported a hidden-charm state X(3872) in the $J/\psi \pi^+\pi^-$ channel [3]. The $D_{c0}^{*}(2317)$, $D_{c1}^{\prime}(2460)$, and X(3872) states have two significant features. The first is that their masses are about 100 MeV below the potential model predictions, which implies that it is difficult to categorize them as conventional mesons. The second feature is that $D_{s0}^{*}(2317)$, $D_{s1}^{\prime}(2460)$, and X(3872) are close to and lower than the DK, D^*K , and $D\bar{D}^*$ thresholds, which strongly hints at their molecular nature. Although there are still many controversies, hadronic molecules are one of the most popular interpretations of these exotic hadrons [4-13]. The observations of $D_{s0}^*(2317)$, $D_{s1}'(2460)$, and X(3872)opened a new era in searches for exotic sates. In the following years, a plethora of hidden-charm XYZ and P_c states were observed in experiments (for reviews, see Refs. [14–20]).

Very recently, the LHCb Collaboration observed a T_{cc}^+ state in the $D^0 D^0 \pi^+$ channel [21, 22], whose mass and width obtained from a Breit-Wigner fit are

$$m_{\rm BW} = (m_{D^{*+}} + m_{D^0}) - 273 \pm 61 \pm 5^{+11}_{-14} \text{ keV},$$

$$\Gamma_{\rm BW} = 410 \pm 165 \pm 43^{+18}_{-38} \text{ keV}.$$
(1)

On the other hand, the pole position is given as [22]

$$m_{\text{pole}} = (m_{D^{*+}} + m_{D^0}) - 360 \pm 40^{+4}_{-0} \text{ keV},$$

$$\Gamma_{\text{pole}} = 48 \pm 2^{+0}_{-14} \text{ keV}.$$
(2)



FIG. 1: Various types of hadronic molecular candidates. Here, we choose $D_{s0}^*(2317)$, X(3872), and T_{cc} as examples.

From its decay products, one can infer its minimum quark component to be $cc\bar{u}d$. Since the mass of the T_{cc}^+ is very close to the D^*D threshold, it could well be interpreted as a D^*D molecular state [23–25] as predicted in many previous works [26–29]. It should be noted that the T_{cc}^+ is the first observed double-charm exotic state. It is interesting to note that the decay width obtained from the Breit-Wigner fit and that derived from the pole position are quite different. The latter strongly supports its nature being a hadronic molecule of DD^* , as stressed, e.g., in Ref. [30].

^{*}Electronic address: lisheng.geng@buaa.edu.cn

[†]Electronic address: xiangliu@lzu.edu.cn

The single-charm, hidden-charm, and double-charm molecular candidates have been established in experiments. In Fig. 1, we choose the $D_{s0}^*(2317)$, X(3872), and T_{cc}^+ states as examples and present the corresponding possible substructures. However, until now, there are no signal of triple-charm molecular states. In the future, experimental searches for triple-charm molecular states will be an interesting research issue in exploring exotic hadrons.

In this work, we investigate the likely existences of triplecharm molecular states composed of D^*D^*D and $D^*D^*D^*$. There are three reasons for studying such systems. First, we notice that single and double-charm molecular candidates in Fig. 1 contain one and two charmed mesons, respectively. Thus it is natural to ask whether there exist hadronic molecular states composed of three charmed mesons. Second, the observation of the T_{cc}^+ state provides a way to fix the interaction between two charmed mesons. In Ref. [29], we successfully reproduced the binding energy of the T_{cc}^+ state, with the DD^* interaction provided by the one-boson-exchange (OBE) model. This makes the numerical results more reliable when dealing with systems containing more charmed mesons in the following study. Third, in Ref. [29], we have studied the DDD* system and found that it has a $I(J^P) = \frac{1}{2}(1^{-})$ bound state solution. Compared with the DDD* system, the D^*D^*D and $D^*D^*D^*$ systems can have more spin configurations. Therefore, it is likely that there exist more hadronic molecular states in the D^*D^*D and $D^*D^*D^*$ systems.

In the past several years, the LHCb Collaboration has achieved great success in discovering exotic states, including several P_c states [31, 32], P_{cs} [33], $X_{0,1}(2900)$ [34, 35], and X(6900) [36]. These observations demonstrated the capacity of the LHCb detector in searching for exotic states. With the upgrade of the LHCb detector [37], one can expect that more exotic states be observed in the future. The predictions of molecular states with the configurations of D^*D^*D and $D^*D^*D^*$ may inspire more experimental works along this line.

This paper is organized as follows. In Sec. II, we introduce the interactions between D^* and $D^{(*)}$ and present the details of the Gussian expansion method (GEM). Next in Sec. III, we present the binding energies and root-mean-square radii of the D^*D^*D and $D^*D^*D^*$ systems. Finally, this paper is ended with a short summary in Sec. IV.

II. FORMALISM

In order to study the D^*D^*D and $D^*D^*D^*$ systems, we should first derive the effective potentials of the D^*-D^* and D^-D^* pairs. For this purpose, we adopt the one-boson-exchange (OBE) model of Ref. [27]. In the OBE model, the $D^*-D^{(*)}$ interactions occur by exchanging π , σ , ρ , and ω mesons. To take into the finite sizes of the exchanged mesons, a monopole type form factor is introduced

$$\mathcal{F}(q^2, m_E^2) = \frac{\Lambda^2 - m_E^2}{\Lambda^2 - q^2},\tag{3}$$



FIG. 2: Jacobi coordinates of the $D^*D^*D^{(*)}$ systems.

where q and m_E are the mass and momentum of the exchanged meson, respectively.

To solve the three-body Schrödinger equation, we employ the Gaussian expansion method (GEM) [38, 39], which was widely used in studies of baryon systems [40–42], multi-quark states [43–47], and multi-body hadronic molecular states [29, 48–51] (for reviews on this latter topic, see, e.g., Refs. [52, 53]). The three-body Schrödinger equation reads

$$\left[\hat{T} + V(r_1) + V(r_2) + V(r_3)\right] \Psi_{JM} = E \Psi_{JM}, \tag{4}$$

where \hat{T} is the kinetic energy operator, $V(r_1)$, $V(r_2)$, and $V(r_3)$ are pairwise potentials. Ψ_{JM} is the total wave function, which can be written as

$$\Psi_{JM} = \sum_{c,\alpha} C_{c,\alpha} \Psi_{JM}^{(c,\alpha)},\tag{5}$$

where

$$\Psi_{JM}^{(c,\alpha)} = H_{t,T}^c \left[\chi_{s,S} \left[\phi_{nl}(\mathbf{r}_c) \phi_{NL}(\mathbf{R}_c) \right]_{\lambda} \right]_{JM}$$
(6)

is the basis, $C_{c,\alpha}$ is the coefficient of the corresponding basis, which can be obtained by the Rayleigh-Ritz variational method. The *c* (*c* = 1, 2, 3) represents the three channels in Fig. 2 and $\alpha = \{tT, sS, nN, lL\lambda\}$ is the quantum number of the basis. $H_{t,T}^c$ is the flavor wave function, where *t* is isospin in the \mathbf{r}_c -degree-of-freedom and *T* is the total isospin. χ_{s,S,M_s} is the spin wave function, where the *s*, *S* are spin in the \mathbf{r}_c degree-of-freedom and total spin, respectively. $\phi_{nlm_l}(\mathbf{r}_c)$ and $\phi_{NLM_l}(\mathbf{R}_c)$ are spatial wave functions, which read

$$\phi_{nlm_l}(\mathbf{r}_c) = N_{nl} r_c^l e^{-\nu_n r_c^2} Y_{lm}(\hat{\mathbf{r}}_c),$$

$$\phi_{NLM_L}(\mathbf{R}_c) = N_{NL} R_c^L e^{-\lambda_N R_c^2} Y_{LM}(\hat{\mathbf{R}}_c),$$
(7)

where N_{nl} and N_{NL} are normalization constants. In Eq. (7), The \mathbf{r}_c and \mathbf{R}_c are Jacobi coordinates, and ν_n and λ_N are Gaussian ranges.

For convenience, in the calculation of the Hamiltonian matrix elements $\langle \Psi_{JM}^{(a,\alpha)} | V(r_c) | \Psi_{JM}^{(b,\alpha')} \rangle$ and $\langle \Psi_{JM}^{(a,\alpha)} | \hat{T} | \Psi_{JM}^{(b,\alpha')} \rangle$, we rotate channels *a* and *b* into channel *c*:

$$|H^{a}_{t,T}\rangle = \sum_{t''} \langle H^{c}_{t'',T} | H^{a}_{t,T} \rangle | H^{c}_{t'',T} \rangle,$$

$$|H^{b}_{t',T'}\rangle = \sum_{t''} \langle H^{c}_{t'',T'} | H^{b}_{t',T'} \rangle | H^{c}_{t'',T'} \rangle,$$

$$|\chi^{a}_{s,S}\rangle = \sum_{s''} \langle \chi^{c}_{s'',S} | \chi^{a}_{s,S} \rangle | \chi^{c}_{s'',S} \rangle,$$

$$|\chi^{b}_{s',S'}\rangle = \sum_{s''} \langle \chi^{c}_{s'',S'} | \chi^{b}_{s',S'} \rangle | \chi^{c}_{s'',S'} \rangle,$$
(9)

$$\mathbf{r}_{a} = \alpha_{ac}\mathbf{r}_{c} + \beta_{ac}\mathbf{R}_{c}, \quad \mathbf{R}_{a} = \gamma_{ac}\mathbf{r}_{c} + \delta_{ac}\mathbf{R}_{c},$$

$$\mathbf{r}_{b} = \alpha_{bc}\mathbf{r}_{c} + \beta_{ac}\mathbf{R}_{c}, \quad \mathbf{R}_{b} = \gamma_{ac}\mathbf{r}_{c} + \delta_{ac}\mathbf{R}_{c}.$$
 (10)

In Eq. (9), $\langle H^c_{t',T} | H^a_{t,T} \rangle$ and $\langle H^c_{t',T'} | H^b_{t',T'} \rangle$ are rotational coefficients of the flavor wave functions. Using the 6 - j symbol, $\langle H^c_{t',T^{(i)}} | H^{a(b)}_{t^{(i)},T^{(i)}} \rangle$ could be expressed as

$$\langle H_{t,T}^{1} | H_{t'',T}^{3} \rangle = \langle H_{t'',T}^{3} | H_{t,T}^{1} \rangle$$

$$= (-1)^{t''} \sqrt{(2t+1)(2t''+1)} \begin{cases} 1/2 & 1/2 & t \\ 1/2 & T & t'' \end{cases} .$$

$$(12)$$

For the D^*D^*D system, the spin wave function rotational coefficients $\langle \chi^c_{s'',S^{(\prime)}} | \chi^{a(b)}_{s^{(\prime)},S^{(\prime)}} \rangle$ in Eq. (9) are

$$\langle \chi_{s,S}^{1} | \chi_{s'',S}^{2} \rangle = \langle \chi_{s'',S}^{2} | \chi_{s,S}^{1} \rangle$$

= $(-1)^{s} \sqrt{(2s+1)(2s''+1)} \left\{ \begin{array}{cc} 1 & 1 & s \\ 0 & S & s'' \end{array} \right\},$ (13)

$$\langle \chi^{1}_{s,s} | \chi^{3}_{s'',S} \rangle = \langle \chi^{3}_{s'',S} | \chi^{1}_{s,S} \rangle$$

= $-(-1)^{s''} \sqrt{(2s+1)(2s''+1)} \left\{ \begin{array}{cc} 1 & 1 & s \\ 0 & S & s'' \end{array} \right\},$ (14)

$$\langle \chi^2_{s,S} | \chi^3_{s'',S} \rangle = \langle \chi^3_{s'',S} | \chi^2_{s,S} \rangle$$

= $-(-1)^s \sqrt{(2s+1)(2s''+1)} \left\{ \begin{array}{cc} 1 & 0 & s \\ 1 & S & s'' \end{array} \right\}.$ (15)

For the $D^*D^*D^*$ system, we have

$$\langle \chi_{s,S}^{1} | \chi_{s'',S}^{2} \rangle = \langle \chi_{s'',S}^{2} | \chi_{s,S}^{1} \rangle = \langle \chi_{s,S}^{2} | \chi_{s'',S}^{3} \rangle = \langle \chi_{s'',S}^{3} | \chi_{s,S}^{2} \rangle = (-1)^{s} \sqrt{(2s+1)(2s''+1)} \begin{cases} 1 & 1 & s \\ 1 & S & s'' \end{cases} ,$$
 (16)

$$\langle \chi^{1}_{s,S} | \chi^{3}_{s'',S} \rangle = \langle \chi^{3}_{s'',S} | \chi^{1}_{s,S} \rangle$$

$$= (-1)^{s''} \sqrt{(2s+1)(2s''+1)} \left\{ \begin{array}{cc} 1 & 1 & s \\ 1 & S & s'' \end{array} \right\}.$$

$$(17)$$

In Eq. (10), α , β , γ , δ are rotational matrices of the Jacobi coordinates, which read

$$\alpha = \begin{pmatrix} 1 & -\frac{m_3}{m_1 + m_3} & \frac{-m_3}{m_2 + m_3} \\ -\frac{m_2}{m_1 + m_2} & 1 & -\frac{m_2}{m_2 + m_3} \\ -\frac{m_1}{m_1 + m_2} & -\frac{m_1}{m_1 + m_3} & 1 \end{pmatrix},$$
 (18)

$$\beta = \begin{pmatrix} 0 & -1 & 1\\ 1 & 0 & -1\\ -1 & 1 & 0 \end{pmatrix}, \tag{19}$$



$$\delta = \begin{pmatrix} 1 & -\frac{m_2}{m_1 + m_2} & -\frac{m_1}{m_1 + m_2} \\ -\frac{m_3}{m_1 + m_3} & 1 & -\frac{m_1}{m_1 + m_3} \\ -\frac{m_3}{m_2 + m_3} & -\frac{m_2}{m_2 + m_3} & 1 \end{pmatrix},$$
 (21)

where m_1 , m_2 , and m_3 are masses of the particles in the c = 1 channel. With Eqs. (11-17), all flavor and spin wave functions could be written by unified expressions in channel c. In this way, the two-body flavor- and spin-dependent factors deduced in Refs. [26, 27] could be directly used in this work.

For an *S*-wave state, i.e., l = L = 0, we obtain a simple expression for the interaction matrix element [38]

$$\langle \phi_{n00}(\mathbf{r}_{a})\phi_{N00}(\mathbf{R}_{a})|V(r_{c})|\phi_{n'00}(\mathbf{r}_{b})\phi_{N'00}(\mathbf{R}_{b})\rangle$$

$$= 32 \frac{(v_{n}\lambda_{N}v_{n'}\lambda_{N'})^{3/4}}{\sqrt{\pi}B^{3/2}} \int_{0}^{\infty} e^{-(A-\frac{C^{2}}{4B})r_{c}^{2}}r_{c}^{2}\mathrm{d}r_{c},$$
(22)

where

$$A = \alpha_{ac}^{2} \nu_{n} + \gamma_{ac}^{2} \lambda_{N} + \alpha_{bc}^{2} \nu_{n'} + \gamma_{bc}^{2} \lambda_{N'},$$

$$B = \beta_{ac}^{2} \nu_{n} + \delta_{ac}^{2} \lambda_{N} + \beta_{bc}^{2} \nu_{n'} + \delta_{bc}^{2} \lambda_{N'},$$

$$C = 2(\alpha_{ac} \beta_{ac} \nu_{n} + \gamma_{ac} \delta_{ac} \lambda_{N} + \alpha_{bc} \beta_{bc} \nu_{n'} + \gamma_{bc} \delta_{bc} \lambda_{N'}).$$

(23)

In addition, the matrix element of the kinetic term (see Ref. [54] for details) could also be calculated analytically

$$\langle \phi_{n00}(\mathbf{r}_{a})\phi_{N00}(\mathbf{R}_{a})|\hat{T}|\phi_{n'00}(\mathbf{r}_{b})\phi_{N'00}(\mathbf{R}_{b})\rangle$$

= $-48 \frac{(\nu_{n}\lambda_{N}\nu_{n'}\lambda_{N'})^{3/4}}{\left[\det(A^{a}+A^{b})\right]^{3/2}} \operatorname{tr}\left[(A^{a}+A^{b})^{-1}A^{a}\hat{T}_{c}A^{b}\right],$ (24)

where

$$A^{a} = \begin{pmatrix} \alpha_{ac}^{2} \nu_{n} + \gamma_{ac}^{2} \lambda_{N} & \alpha_{ac} \beta_{ac} \nu_{n} + \gamma_{ac} \delta_{ac} \lambda_{N} \\ \alpha_{ac} \beta_{ac} \nu_{n} + \gamma_{ac} \delta_{ac} \lambda_{N} & \beta_{ac}^{2} \nu_{n} + \delta_{ac}^{2} \lambda_{N} \end{pmatrix}, \quad (25)$$

$$A^{b} = \begin{pmatrix} \alpha_{bc}^{2} \nu_{n'} + \gamma_{bc}^{2} \lambda_{N'} & \alpha_{bc} \beta_{bc} \nu_{n'} + \gamma_{bc} \delta_{bc} \lambda_{N'} \\ \alpha_{bc} \beta_{bc} \nu_{n'} + \gamma_{bc} \delta_{bc} \lambda_{N'} & \beta_{bc}^{2} \nu_{n'} + \delta_{bc}^{2} \lambda_{N'} \end{pmatrix}, \quad (26)$$

$$\hat{T}_c = \begin{pmatrix} -\frac{1}{2\mu_c^r} & 0\\ 0 & -\frac{1}{2\mu_c^R} \end{pmatrix} \quad \text{with} \quad \begin{cases} \mu_c^r = \frac{m_i m_j}{m_i + m_j},\\ \mu_c^R = \frac{(m_i + m_j) m_k}{m_i + m_j + m_k}. \end{cases}$$
(27)

III. NUMERICAL RESULTS

With the effective potentials of Ref. [27], we could solve the three-body Schrödinger equation with the Gaussian expansion method. We not only calculate the binding energy, but also the root-mean-square radii of D^*-D^* and $D-D^*$. In general, orbitally excited hadronic molecular states are more difficult to form because of the repulsive centrifugal potential. It is more likely to find bound state solutions in the *S*-wave (l = L = 0) configurations. Although *S*-*D* mixing may affect the binding energies, its contribution is usually small and therefore it is unlikely to affect the results in a qualitative way. Therefore in this work, we only consider *S*-wave two-body interactions.

In our approach, the only undetermined parameter is the cutoff Λ , which is crucial in investigating the existence of bound state solutions. With the measured binding energy of the T_{cc}^+ state, we obtained $\Lambda = 0.976$, 0.998, and 1.013 GeV in Ref. [29], which are close to the suggested values in previous works [26, 55–59]. The D^*D^*D and $D^*D^*D^*$ systems can be related to the *DDD*^{*} system via heavy quark spin symmetry. As a result, in this work we will take the same strategy as that of Ref. [29], i.e., we scan the range of Λ from 0.95 to 3.0 GeV to search for bound states of D^*D^*D and $D^*D^*D^*$. If a system has a bound state solution with $\Lambda \approx 1$ GeV, we view this state as a good molecular candidate.

In the study of hadronic molecular candidates composed of two charmed mesons [26], it was found that systems with lower isospins are more likely to bind. In this work, we find that this is also true for systems composed of three charmed mesons.

A. *S*-wave D^*D^*D system

I	J^P	с	t	Т	s	S	l	L	λ
		1	1	$\frac{1}{2}$	0	0	0	0	0
	0-	2, 3	0	$\frac{1}{2}$	1	0	0	0	0
		2, 3	1	$\frac{1}{2}$	1	0	0	0	0
		1	0	$\frac{1}{2}$	1	1	0	0	0
$\frac{1}{2}$	1-	2, 3	0	$\frac{1}{2}$	1	1	0	0	0
		2, 3	1	$\frac{1}{2}$	1	1	0	0	0
		1	1	$\frac{1}{2}$	2	2	0	0	0
	2-	2, 3	0	$\frac{1}{2}$	1	2	0	0	0
		2, 3	1	$\frac{1}{2}$	1	2	0	0	0
	0-	1	1	$\frac{3}{2}$	0	0	0	0	0
$\frac{3}{2}$	0	2, 3	1	$\frac{3}{2}$	1	0	0	0	0
	1-	2, 3	1	$\frac{3}{2}$	1	1	0	0	0
	2-	1	1	$\frac{3}{2}$	2	2	0	0	0
	2	2, 3	1	$\frac{3}{2}$	1	2	0	0	0

TABLE I: Bases of the D^*D^*D system.

For the *S*-wave D^*D^*D system, allowed spin-parities are 0^- , 1^- , and 2^- with $I = \frac{1}{2}$ and $\frac{3}{2}$. The corresponding quantum numbers $\{t, T, s, S, l, L, \lambda\}$ of the Jacobi channels are tabulated in Table I. For the $D^* - D^*$ pair (i.e., channel 1), the sum of t + s + l should be odd. With the Gaussian expansion method, we calculate the binding energies and root-mean-square radii of the *S*-wave D^*D^*D with $I(J^P) = \frac{1}{2}(0^-, 1^-, 2^-)$ and $I(J^P) = \frac{3}{2}(0^-, 1^-, 2^-)$. The numerical results are shown in Table II.

TABLE II:	Binding	energies	and	root-mean-sq	uare r	adii o	D	$^{*}D^{*}$	D
system.									

Ι	J^P	Λ (GeV)	B (MeV)	$r_{D^*D^*}$ (fm)	r_{D^*D} (fm)
		0.976	0.361	8.009	6.362
	0-	0.998	1.324	5.891	4.565
		1.013	2.289	4.769	3.686
		0.976	0.631	4.026	4.338
$\frac{1}{2}$	1-	0.998	2.512	2.289	2.618
		1.013	4.332	1.846	2.147
		0.976	1.336	3.628	3.059
	2^{-}	0.998	3.481	2.501	2.128
		1.013	5.431	2.114	1.806
		1.767	0.347	5.466	4.624
<u>3</u> 2	0-	1.792	0.965	4.111	3.427
		1.817	1.822	3.340	2.758
		1.857	0.375	11.799	8.741
	1-	1.882	1.106	11.485	8.363
		1.907	2.063	11.255	8.117
		1.486	0.351	2.820	2.857
	2-	1.511	1.902	1.796	1.823
		1.536	4.127	1.412	1.435

For the *S*-wave D^*D^*D system with $I(J^P) = \frac{1}{2}(0^-)$, we can obtain bound state solutions when the cutoff Λ reaches about 0.976 GeV. The binding energy is on the order of MeV and the root-mean-square radii are several fm. If we increase the cutoff Λ , the binding energy increases while the root-mean-square radii decrease. Since the binding energy and root-mean-square radii are reasonable from the perspective of hadronic molecules, this state could be viewed as a good hadronic molecular candidate.

For the *S*-wave D^*D^*D system with $I(J^P) = \frac{1}{2}(1^-)$, we can also obtain bound state solutions with the cutoff $\Lambda = 0.976$, 0.998, and 1.013 GeV. According to the calculated binding energy and root-mean-square radii, this state could also be treated as an ideal hadronic molecular candidate.

For the *S*-wave D^*D^*D system with $I(J^P) = \frac{1}{2}(2^-)$, one can also find weakly bound states for the same cutoff Λ as that of $I(J^P) = \frac{1}{2}(0^-, 1^-)$. Similar to the case of $P_c(4440)$ and $P_c(4457)$, for the same cutoff, the $0^-, 1^-, 2^-$ states have different binding energies at the order of sevel hundred keV. With increased experimental precision, it is likely that these states can be distinguished from each other in future experiments.

We also study the D^*D^*D system with $I(J^P) = \frac{3}{2}(0^-, 1^-, 2^-)$. There are no bound state solutions for a cutoff Λ below 1.013 GeV. In order to obtain bound state solutions with $J^P = 0^-$, 1^- , and 2^- , we increase the cutoff Λ to $1.767 \sim 1.817$ GeV, $1.857 \sim 1.907$ GeV, and $1.486 \sim 1.536$ GeV, respectively. Here, we increase the cutoff in steps of 0.025 GeV when scanning the cutoff Λ . Considering that the used cutoff Λ are out of the range of $0.976 \sim 1.013$ GeV (the range determined in Ref. [29]), we are a bit reluctant to view the $I(J^P) = \frac{3}{2}(0^-, 1^-, 2^-) D^*D^*D$ bound states as good hadronic molecular candidates.

How to search for D^*D^*D molecular candidates is also an interesting question. One possible decay mode is that the triple-charm molecules decay into a double-charm molecular state and a charmed meson. The other possible mode is that they directly decay into multi-body finial states escaping intermediate sates. Here, we summarize these channels as follows.

- If the binding energies are extremely small, they could first decay into T⁺_{cc}D^{*}, and then T⁺_{cc} can decay into DDπ, and D^{*} could be seen in the Dπ and Dγ channels. In this case, the molecular candidates may be observed in the DDDππ and DDDπγ final states.
- If the masses of the molecular candidates are below the $T_{cc}^+D^*$ threshold, the kinematically allowed channel is T_{cc}^+D . The D^*D^*D molecular candidates could be studied in the $DDD\pi$ channel.
- If the D*D* hadronic molecular state exists, the D*D*D molecular candidates may decay into a D*D* molecular ular sate and D. The D*D* molecular state could be observed in the D*D, DDπ, and DDγ final states.
- In the above three scenarios, the D^*D^*D molecular candidates ultimately decay into three charmed mesons together with π and γ . We should also emphasize that these final states can not only originate from the intermediate double-charm molecular states with $D^{(*)}$, but also from non-resonant processes.
- In addition, the *D***D***D* molecular candidates can also decay into three charmed mesons via fall apart or quark rearrangement mechanisms. The typical channels are *D***DD* and *DDD*.

According to the discussions above, the D^*D^*D molecular candidates could be studied with the three-, four-, and five-body finial states in future experiments.

B. *S*-wave $D^*D^*D^*$ system

Since the $D^*D^*D^*$ system contains three identical mesons, the c = 1, 2, 3 channels share the same bases. In addition, for all channels, $(-1)^{t+s+l+1} = 1$, which restricts allowed combinations of t, s, and l. In Table III, we present the bases of the S-wave $D^*D^*D^*$ system. For the S-wave $D^*D^*D^*$ system with $I = \frac{1}{2}$, the allowed spin-parities are 0^- , 1^- , 2^- , and 3^- . For the $I = \frac{3}{2}$, the allowed spin-parities are 1^- , 2^- , and 3^- . The binding energies and root-mean-square radii are presented in Table IV.

For the *S*-wave $D^*D^*D^*$ system with $I(J^P) = \frac{1}{2}(0^-)$, we find a bound state solution for a cutoff Λ larger than 1.013 GeV. The root-mean-square radii decrease slowly with the increase of the cutoff Λ . The radius $r_{D^*D^*}$ is estimated to be about 9 fm with $\Lambda \sim 1$ GeV, which is a bit larger than that of T_{cc}^+ but similar to those of the *DDD*^{*} states. Judging from the

TABLE III: Bases of the $D^*D^*D^*$ system.

Ι	J^P	с	t	Т	S	S	l	L	λ
	0-	1, 2, 3	0	$\frac{1}{2}$	1	0	0	0	0
	1-	1, 2, 3	1	$\frac{1}{2}$	0	1	0	0	0
		1, 2, 3	0	$\frac{1}{2}$	1	1	0	0	0
$\frac{1}{2}$		1, 2, 3	1	$\frac{1}{2}$	2	1	0	0	0
	2-	1, 2, 3	0	$\frac{1}{2}$	1	2	0	0	0
		1, 2, 3	1	$\frac{1}{2}$	2	2	0	0	0
	3-	1, 2, 3	1	$\frac{1}{2}$	2	3	0	0	0
$\frac{3}{2}$	0-					-			
	1-	1, 2, 3	1	$\frac{3}{2}$	0	1	0	0	0
		1, 2, 3	1	$\frac{3}{2}$	2	1	0	0	0
	2-	1, 2, 3	1	$\frac{3}{2}$	2	2	0	0	0
	3-	1, 2, 3	1	$\frac{3}{2}$	2	3	0	0	0

TABLE IV: Binding energies and root-mean-square radii of $D^*D^*D^*$ system.

I	J^P	Λ (GeV)	B (MeV)	$r_{D^*D^*}$ (fm)
		1.013	0.329	9.958
	0^{-}	1.038	1.769	9.397
		1.063	3.800	9.119
		0.992	0.317	6.538
	1-	1.017	1.738	3.922
1		1.042	4.080	2.730
2		0.976	0.166	5.558
	2^{-}	0.998	1.782	2.774
		1.013	3.485	2.170
		1.834	0.349	9.841
	3-	1.859	1.109	9.472
		1.884	2.111	9.227
		1.809	0.357	7.470
	1-	1.834	0.920	6.591
		1.859	1.714	5.871
		1.834	0.346	9.852
$\frac{3}{2}$	2^{-}	1.859	1.106	9.486
		1.884	2.108	9.244
		1.482	0.336	2.757
	3-	1.507	2.021	1.729
		1.532	4.434	1.361

binding energy and root-mean-square radius, this state could be viewed as a good hadronic molecular candidate.

For the *S*-wave $D^*D^*D^*$ system with $I(J^P) = \frac{1}{2}(1^-)$, we obtain a binding energy in the range of 0.341 ~ 4.141 MeV for a cutoff Λ between 0.992 and 1.042 GeV. While the root-mean-square radius decreases from 6.459 to 2.716 fm with the increase of the cutoff Λ .

For the *S*-wave $D^*D^*D^*$ system with $I(J^P) = \frac{1}{2}(2^-)$, the system becomes bound when the cutoff Λ reaches about 0.976

GeV. Since the obtained binding energy is on the order of MeV and the root-mean-square radius is several fm, this state is also an ideal hadronic molecular candidate.

For the *S*-wave $I(J^P) = \frac{1}{2}(3^-)$ and $I(J^P) = \frac{3}{2}(1^-, 2^-, 3^-)$ $D^*D^*D^*$ systems, a larger cutoff is needed for them to bind. More specifically, we only find bound state solutions when the cutoff Λ reaches around 1.8 GeV. We also obtain shallow bound states in the *S*-wave $I(J^P) = \frac{3}{2}(3^-) D^*D^*D^*$ system if the cutoff Λ is close to 1.5 GeV. Since the needed cutoff Λ is much larger than our expectation, we prefer not to view these states as good hadronic molecular candidates.

Since the $D^*D^*D^*$ molecular candidates have larger masses than that of the D^*D^*D system, much more complex decay modes can be anticipated. Here, the decay channels are summarized as the following:

- In principle, all decay channels of the *D***D***D* molecular candidates are also kinematically allowed for the *D***D***D***D** system.
- There are also some channels specific for the $D^*D^*D^*$ system. For example, the channel of a D^*D^* molecular candidate with a D^* meson is only kinematically allowed for the $D^*D^*D^*$ system.

However, although there more decay channels for the $D^*D^*D^*$ system, the decay patterns are similar for the DDD^* [29], D^*D^*D , and $D^*D^*D^*$ molecular states. In the future, these states could be searched for by measuring three charmed mesons together with pions and photons in the finial states.

IV. SUMMARY

In recent years, the LHCb Collaboration achieved remarkable success in discovering new hadronic states, including many of exotic ones, which cannot fit into the conventional quark model. These observations enriched the members of the exotic hadronic family and improved our understandings of non-perturbative strong interactions. Very recently, the LHCb Collaboration observed a new state T_{cc}^+ in the $D^0 D^0 \pi^+$ channel [22]. The T_{cc}^+ state could well be interpreted as a DD^* molecular state and it is the first double-charm exotic state ever observed.

The observation of the T_{cc}^+ state enabled us to derive the interaction between two charmed mesons. In Ref. [29], by

- [1] B. Aubert *et al.* [BaBar], Observation of a narrow meson decaying to $D_s^+ \pi^0$ at a mass of 2.32 GeV/ c^2 , Phys. Rev. Lett. **90**, 242001 (2003).
- [2] D. Besson *et al.* [CLEO], Observation of a narrow resonance of mass 2.46 GeV/ c^2 decaying to $D_s^{*+}\pi^0$ and confirmation of the $D_{s,j}^*(2317)$ state, Phys. Rev. D **68**, 032002 (2003), [erratum: Phys. Rev. D **75**, 119908 (2007)].
- [3] S. K. Choi *et al.* [Belle], Observation of a narrow charmoniumlike state in exclusive $B^{\pm} \rightarrow K^{\pm}\pi^{+}\pi^{-}J/\psi$ decays, Phys. Rev. Lett. **91**, 262001 (2003).

reproducing the binding energy of T_{cc}^+ , we determined the only parameter in the OBE model, the cutoff Λ , which then fixes the DD^* potential. This allows us to study hadronic molecular states composed of several charmed mesons. In this work, we studied the existence of triple-charm molecular states composed of D^*D^*D and $D^*D^*D^*$. Using the cutoff Λ fixed by the binding energy of the T_{cc}^+ , we find that the $I(J^P) = \frac{1}{2}(0^-, 1^-, 2^-) D^*D^*D$ and $I(J^P) = \frac{1}{2}(0^-, 1^-, 2^-) D^*D^*D$ and $I(J^P) = \frac{1}{2}(0^-, 1^-, 2^-) D^*D^*D$ have loosely bound state solutions, which could be viewed as good hadronic molecular candidates. We suggest to search for the D^*D^*D and D^*D^*D molecular states in the following decay modes:

- a. a double-charm molecular state and a charmed meson,
- b. three charmed mesons,
- c. three charmed mesons together with a number of pions and photons.

On the other hand, we find that the $I(J^P) = \frac{3}{2}(0^-, 1^-, 2^-)$ D^*D^*D , $I(J^P) = \frac{1}{2}(3^-)$ and $I(J^P) = \frac{1}{2}(1^-, 2^-, 3^-)$ $D^*D^*D^*$ systems are more difficult to form bound states.

It is no doubt that the LHCb Collaboration has played an important role in searches for exotic states. The observation of the T_{cc}^+ once again shows the capability of the LHCb detector in this area. With anticipated data accumulation [37], more exotic states can be expected in the future.

Acknowledgement

This work is partly supported by the National Natural Science Foundation of China under Grants No.11735003, No.11975041, No.11961141004, and the fundamental Research Funds for the Central Universities. XL is supported by the China National Funds for Distinguished Young Scientists under Grant No. 11825503, National Key Research and Development Program of China under Contract No. 2020YFA0406400, the 111 Project under Grant No. B20063, the National Natural Science Foundation of China under Grant No. 12047501, and the Fundamental Research Funds for the Central Universities.

- [4] T. Barnes, F. E. Close and H. J. Lipkin, Implications of a DK molecule at 2.32 GeV, Phys. Rev. D 68, 054006 (2003).
- [5] Z. X. Xie, G. Q. Feng and X. H. Guo, Analyzing $D_{s0}^*(2317)^+$ in the *DK* molecule picture in the Beth-Salpeter approach, Phys. Rev. D **81**, 036014 (2010).
- [6] A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, Strong and radiative decays of the D^{*}_{s0}(2317) meson in the DKmolecule picture, Phys. Rev. D 76, 014005 (2007).
- [7] G. Q. Feng, X. H. Guo and Z. H. Zhang, Studying the D^*K molecular structure of $D^+_{s1}(2460)$ in the Bethe-Salpeter ap-

proach, Eur. Phys. J. C 72, 2033 (2012).

- [8] A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, D^{*}K molecular structure of the D_{s1}(2460) meson, Phys. Rev. D 76, 114008 (2007).
- [9] Y. J. Zhang, H. C. Chiang, P. N. Shen and B. S. Zou, Possible S-wave bound-states of two pseudoscalar mesons, Phys. Rev. D 74, 014013 (2006).
- [10] E. S. Swanson, Short range structure in the *X*(3872), Phys. Lett. B 588, 189-195 (2004).
- [11] C. Y. Wong, Molecular states of heavy quark mesons, Phys. Rev. C 69, 055202 (2004).
- [12] X. Liu, Z. G. Luo, Y. R. Liu and S. L. Zhu, *X*(3872) and Other Possible Heavy Molecular States, Eur. Phys. J. C **61**, 411-428 (2009).
- [13] I. W. Lee, A. Faessler, T. Gutsche and V. E. Lyubovitskij, X(3872) as a molecular DD* state in a potential model, Phys. Rev. D 80, 094005 (2009).
- [14] H. X. Chen, W. Chen, X. Liu and S. L. Zhu, The hiddencharm pentaquark and tetraquark states, Phys. Rept. 639, 1-121 (2016).
- [15] X. Liu, An overview of XYZ new particles, Chin. Sci. Bull. 59, 3815-3830 (2014).
- [16] C. Z. Yuan, The XYZ states revisited, Int. J. Mod. Phys. A 33, no.21, 1830018 (2018).
- [17] S. L. Olsen, T. Skwarnicki and D. Zieminska, Nonstandard heavy mesons and baryons: Experimental evidence, Rev. Mod. Phys. 90, no.1, 015003 (2018).
- [18] F. K. Guo, C. Hanhart, U. G. Meißner, Q. Wang, Q. Zhao and B. S. Zou, Hadronic molecules," Rev. Mod. Phys. 90, no.1, 015004 (2018).
- [19] A. Hosaka, T. Iijima, K. Miyabayashi, Y. Sakai and S. Yasui, Exotic hadrons with heavy flavors: X, Y, Z, and related states, PTEP 2016, no.6, 062C01 (2016).
- [20] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C. P. Shen, C. E. Thomas, A. Vairo and C. Z. Yuan, The *XYZ* states: experimental and theoretical status and perspectives, Phys. Rept. 873, 1-154 (2020).
- [21] R. Aaij *et al.* [LHCb], Observation of an exotic narrow doubly charmed tetraquark, [arXiv:2109.01038 [hep-ex]].
- [22] R. Aaij *et al.* [LHCb], Study of the doubly charmed tetraquark T_{cc}^+ , [arXiv:2109.01056 [hep-ex]].
- [23] N. Li, Z. F. Sun, X. Liu and S. L. Zhu, Perfect DD^* molecular prediction matching the T_{cc} observation at LHCb, Chin. Phys. Lett. **38**, 092001 (2021).
- [24] H. Ren, F. Wu and R. Zhu, Hadronic molecule interpretation of T_{cc}^+ and its beauty-partners, [arXiv:2109.02531 [hep-ph]].
- [25] R. Chen, Q. Huang, X. Liu and S. L. Zhu, Another doubly charmed molecular resonance T^{'+}_{cc} (3876), [arXiv:2108.01911 [hep-ph]].
- [26] N. Li, Z. F. Sun, X. Liu and S. L. Zhu, Coupled-channel analysis of the possible $D^{(*)}\overline{B}^{(*)}\overline{B}^{(*)}$ and $D^{(*)}\overline{B}^{(*)}$ molecular states, Phys. Rev. D **88**, no.11, 114008 (2013).
- [27] M. Z. Liu, T. W. Wu, M. Pavon Valderrama, J. J. Xie and L. S. Geng, Heavy-quark spin and flavor symmetry partners of the *X*(3872) revisited: What can we learn from the one boson exchange model? Phys. Rev. D **99**, no.9, 094018 (2019).
- [28] H. Xu, B. Wang, Z. W. Liu and X. Liu, *DD** potentials in chiral perturbation theory and possible molecular states, Phys. Rev. D 99, no.1, 014027 (2019).
- [29] T. W. Wu, Y. W. Pan, M. Z. Liu, S. Q. Luo, X. Liu and L. S. Geng, Discovery of the doubly charmed T⁺_{cc} state implies a triply charmed H_{ccc} hexaquark state, [arXiv:2108.00923 [hepph]].

- [30] X. Z. Ling, M. Z. Liu, L. S. Geng, E. Wang and J. J. Xie, Can we understand the decay width of the T⁺_{cc} state? [arXiv:2108.00947 [hep-ph]].
- [31] R. Aaij *et al.* [LHCb], Observation of $J/\psi p$ Resonances Consistent with Pentaquark States in $\Lambda_b^0 \rightarrow J/\psi K^- p$ Decays, Phys. Rev. Lett. **115**, 072001 (2015).
- [32] R. Aaij *et al.* [LHCb], Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$, Phys. Rev. Lett. **122**, no.22, 222001 (2019).
- [33] R. Aaij *et al.* [LHCb], Evidence of a $J/\psi\Lambda$ structure and observation of excited Ξ^- states in the $\Xi_b^- \rightarrow J/\psi\Lambda K^-$ decay, Sci. Bull. **66**, 1278-1287 (2021).
- [34] R. Aaij *et al.* [LHCb], A model-independent study of resonant structure in $B^+ \rightarrow D^+D^-K^+$ decays, Phys. Rev. Lett. **125**, 242001 (2020).
- [35] R. Aaij *et al.* [LHCb], Amplitude analysis of the $B^+ \rightarrow D^+D^-K^+$ decay, Phys. Rev. D **102**, 112003 (2020).
- [36] R. Aaij *et al.* [LHCb], Observation of structure in the J/ψ -pair mass spectrum, Sci. Bull. **65**, no.23, 1983-1993 (2020).
- [37] R. Aaij *et al.* [LHCb], Physics case for an LHCb Upgrade II -Opportunities in flavour physics, and beyond, in the HL-LHC era, [arXiv:1808.08865 [hep-ex]].
- [38] E. Hiyama, Y. Kino and M. Kamimura, Gaussian expansion method for few-body systems, Prog. Part. Nucl. Phys. 51, 223-307 (2003).
- [39] E. Hiyama, Gaussian expansion method for few-body systems and its applications to atomic and nuclear physics, PTEP **2012**, 01A204 (2012).
- [40] T. Yoshida, E. Hiyama, A. Hosaka, M. Oka and K. Sadato, Spectrum of heavy baryons in the quark model, Phys. Rev. D 92, no.11, 114029 (2015).
- [41] G. Yang, J. Ping, P. G. Ortega and J. Segovia, Triply heavy baryons in the constituent quark model, Chin. Phys. C 44, no.2, 023102 (2020).
- [42] G. Yang, J. Ping and J. Segovia, The S- and P-Wave Low-Lying Baryons in the Chiral Quark Model, Few Body Syst. 59, no.6, 113 (2018).
- [43] G. Yang, J. Ping and J. Segovia, $sQ\bar{q}\bar{q}$ (q = u, d; Q = c, b) tetraquarks in the chiral quark model, Phys. Rev. D **103**, no.7, 074011 (2021).
- [44] G. Yang, J. Ping and J. Segovia, QQss tetraquarks in the chiral quark model, Phys. Rev. D 102, no.5, 054023 (2020).
- [45] G. J. Wang, L. Meng and S. L. Zhu, Spectrum of the fully-heavy tetraquark state $QQ\bar{Q}'\bar{Q}'$, Phys. Rev. D **100**, no.9, 096013 (2019).
- [46] Q. F. Lü, D. Y. Chen, Y. B. Dong and E. Santopinto, Triplyheavy tetraquarks in an extended relativized quark model, Phys. Rev. D 104, no.5, 054026 (2021).
- [47] Q. F. Lü, D. Y. Chen and Y. B. Dong, Masses of fully heavy tetraquarks QQQQ in an extended relativized quark model, Eur. Phys. J. C 80, no.9, 871 (2020).
- [48] T. W. Wu, M. Z. Liu, L. S. Geng, E. Hiyama and M. P. Valderrama, *DK*, *DDK*, and *DDDK* molecules–understanding the nature of the D^{*}_{\$9}(2317), Phys. Rev. D 100, no.3, 034029 (2019).
- [49] T. W. Wu, M. Z. Liu, L. S. Geng, E. Hiyama, M. P. Valderrama and W. L. Wang, Quadruply charmed dibaryons as heavy quark symmetry partners of the *DDK* bound state, Eur. Phys. J. C 80, no.9, 901 (2020).
- [50] T. W. Wu, M. Z. Liu and L. S. Geng, Excited K meson, $K_c(4180)$, with hidden charm as a $D\bar{D}K$ bound state, Phys. Rev. D **103**, no.3, L031501 (2021).
- [51] T. W. Wu, Y. W. Pan, M. Z. Liu, J. X. Lu, L. S. Geng and X. H. Liu, Hidden charm hadronic molecule with strangeness $P_{cs}^*(4739)$ as a $\Sigma_c \bar{D}\bar{K}$ bound state, Phys. Rev. D 104, no.9,

094032 (2021).

- [52] T. W. Wu and L. S. Geng, Study on Triple-Hadron Bound States with Gaussian Expansion Method, Few Body Syst. 62, no.4, 89 (2021).
- [53] T. W. Wu, M. Z. Liu and L. S. Geng, One Way to Verify the Molecular Picture of Exotic Hadrons: From from DK to $DDK/D\bar{D}^{(*)}K$, Few Body Syst. **62**, no.3, 38 (2021).
- [54] D. M. Brink and F. Stancu, Tetraquarks with heavy flavors, Phys. Rev. D 57, 6778-6787 (1998).
- [55] R. Chen, X. Liu, X. Q. Li and S. L. Zhu, Identifying exotic hidden-charm pentaquarks, Phys. Rev. Lett. 115, no.13, 132002 (2015).
- [56] R. Chen, Z. F. Sun, X. Liu and S. L. Zhu, Strong LHCb evidence supporting the existence of the hidden-charm molecular pentaquarks, Phys. Rev. D 100, no.1, 011502 (2019).
- [57] R. Chen, F. L. Wang, A. Hosaka and X. Liu, Exotic triple-charm deuteronlike hexaquarks, Phys. Rev. D 97, no.11, 114011 (2018).
- [58] R. Chen, A. Hosaka and X. Liu, Prediction of triple-charm molecular pentaquarks, Phys. Rev. D 96, no.11, 114030 (2017).
- [59] S. Yasui and K. Sudoh, Exotic nuclei with open heavy flavor mesons, Phys. Rev. D 80, 034008 (2009).