

# Quasi-toposes as elementary quotient completions

Maria Emilia Maietti\*    Fabio Pasquali†    Giuseppe Rosolini‡

## Abstract

The elementary quotient completion of an elementary doctrine in the sense of Lawvere was introduced in previous work by the first and third authors. It generalises the exact completion of a category with finite products and weak equalisers. In this paper we characterise when an elementary quotient completion is a quasi-topos. We obtain as a corollary a complete characterisation of when an elementary quotient completions is an elementary topos. As a byproduct we determine also when the elementary quotient completion of a tripos is equivalent to the doctrine obtained via the tripos-to-topos construction.

Our results are reminiscent of other works regarding exact completions and put those under a common scheme: in particular, Carboni and Vitale’s characterisation of exact completions in terms of their projective objects, Carboni and Rosolini’s characterisation of locally cartesian closed exact completions, also in the revision by Emmenegger, and Menni’s characterisation of the exact completions which are elementary toposes.

The paper contains results presented by the authors at several international meetings in the past years, in particular at Logic Colloquium 2016 and Category Theory 2017, and during the Trimester devoted to Types, Homotopy Type Theory and Verification at the Hausdorff research Institute for Mathematics in 2018. We would like to thank the organisers of these events who gave us the opportunity to present our results.

## 1 Introduction

The study of constructions for completing a category with quotients is a central topic not only in mathematics but also in computer science. In category theory a well-known related notion is that of exact completion of a category with finite limits, and that of a regular category, see [CC82, CV98], which has been widely studied and applied.

---

\*Dipartimento di Matematica “Tullio Levi Civita”, Università di Padova, via Trieste 63, 35121 Padova, Italy, email: [maietti@math.unipd.it](mailto:maietti@math.unipd.it)

†Dipartimento di Matematica “Tullio Levi Civita”, Università di Padova, via Trieste 63, 35121 Padova, Italy, email: [pasquali@dima.unige.it](mailto:pasquali@dima.unige.it)

‡DIMA, Università di Genova, via Dodecaneso 35, 16146 Genova, Italy, email: [rosolini@unige.it](mailto:rosolini@unige.it)

In [MR13b] the first and third authors generalized the notion of exact completion on a category with weak finite limits to that of an *elementary quotient completion* of a Lawvere’s elementary doctrine [Law69, Law70] as an universal construction to close such a doctrine with respect to a suitable notion of quotient.

The exact completion of a category  $\mathcal{C}$  with finite products and weak limits is an instance of such a construction in the sense that its subobject doctrine is the elementary quotient completion doctrine of the doctrine of variations of  $\mathcal{C}$  [Gra00].

In the paper we study elementary quotient completions performed on the special class of Lawvere’s elementary doctrines called *triposes*, introduced in [HJP80], to build elementary toposes by means of what is now known as the tripos-to-topos construction, see [Fre15]. We then characterize those triposes whose elementary quotient completion is an *arithmetic quasi-topos*—*i.e.* a quasi-topos equipped with a natural number object—as base category.

To obtain the characterization, we extend some known results about exact completions such as Carboni and Vitale’s characterization of exact completions in terms of its projective objects in [CV98], Menni’s characterization of the exact completions which are toposes in [Men03] and Carboni and Rosolini’s characterization of the locally cartesian closed exact completions [CR00]. In particular, we show that

- an elementary doctrine  $P: \mathbb{C}^{op} \rightarrow \mathbf{InfSL}$  closed under effective quotients is the elementary quotient completion of the doctrine determined by the restriction of  $P$  to the full subcategory of  $\mathbb{C}$  on its projective objects;
- the base category of the elementary quotient completion of  $P$  turns weak universal properties of  $\mathbb{C}$  into (strong) universal properties of the base of the elementary quotient completion. Those include binary coproducts, a natural number object, a parametrized list object, a subobject classifier, a cartesian closed structure, a locally cartesian structure.
- by using results in [MR16] we characterize when an elementary quotient completion is an elementary topos;
- by using results in [MPR17] we characterize when an elementary quotient completion is a tripos-to-topos construction.

We conclude by pointing out some relevant examples of arithmetic quasi-toposes arising as non-exact elementary quotient completions. Most notably they include the category of equilogical spaces of [Sco76, Sco96, BBS04], that of assemblies over a partial combinatory algebra (see [Hyl82, vO08]), and the category of total setoids, in the style of E. Bishop, over Coquand and Paulin’s Calculus of Inductive Constructions which is the theory at the base of the proof-assistant Coq.

## 2 Preliminary definitions

This section collects the necessary definitions to introduce the *elementary quotient completion* of an elementary doctrine and related properties of a doctrine. Recall from [MR13a], see also [EPR20], that a **primary doctrine** is an indexed inf-semilattice  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  where  $\mathcal{C}$  is a category with a terminal object  $T$  and with binary products

$$C_1 \xleftarrow{\text{pr}_1} C_1 \times C_2 \xrightarrow{\text{pr}_2} C_2$$

The category  $\mathcal{C}$  is often called **base** of the doctrine. We say that  $\alpha$  is over  $A$  if  $\alpha$  is an element of  $P(A)$ . The top element over an object  $A$  of  $\mathcal{C}$  is denoted by  $\top_A$ . Given  $\alpha$  and  $\beta$  over  $A$ , their meet is  $\alpha \wedge_A \beta$  (we may drop subscripts when these are clear from the context).

**2.1 Examples.** (a) An example of primary doctrine comes directly from first order logic is the Lindenbaum-Tarski algebras of well-formed formulas of a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$ . The base category is the category  $\mathcal{V}$  of lists of distinct variables and term substitutions, and the primary doctrine  $LT: \mathcal{V}^{\text{op}} \rightarrow \mathbf{ISL}$  on  $\mathcal{V}$  is given on a list of variables  $\bar{x}$  by taking  $LT(\bar{x})$  as the Lindenbaum-Tarski algebra of well-formed formulas with free variables in  $\bar{x}$ . Meets in  $LT(\bar{x})$  are given by conjunctions while the top element by any true formula. See [MR13a] for more details.

(b) Let  $\mathcal{H}$  be an inf-semilattice. The functor  $\mathcal{P}_{\mathcal{H}}: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{ISL}$  sending a set  $A$  to  $\mathcal{H}^A$  and a function  $f: A \rightarrow B$  to  $\mathcal{P}_{\mathcal{H}}(f) = - \circ f$  is a primary doctrine

(c) If  $\mathcal{C}$  has finite limits the functor  $\text{Sub}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is a primary doctrine.

(d) Another categorical example is given by a category  $\mathcal{C}$  with binary products and weak pullbacks, by defining the doctrine functor of *variations*  $\Psi_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  which evaluates as the poset reflection of each comma category  $\mathcal{C}/A$  at each object  $A$  of  $\mathcal{C}$ , introduced in [Gra00].

Primary doctrines are the objects of the 2-category **PD** where

**the 1-morphisms** in **PD** are pairs  $(F, b)$  where  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a functor and  $b: P \rightarrow R \circ F^{\text{op}}$  is a natural transformation as in the diagram

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} & \xrightarrow{P} & \mathbf{ISL} \\ F^{\text{op}} \downarrow & \begin{array}{c} \searrow b \\ \downarrow \cdot \\ \searrow R \end{array} & \\ \mathcal{D}^{\text{op}} & \xrightarrow{R} & \mathbf{ISL} \end{array}$$

where the functor  $F$  preserves products

**the 2-morphisms** are natural transformations  $\theta: F \rightarrow G$  such that

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} & \xrightarrow{P} & \mathbf{ISL} \\ \left( \begin{array}{c} \downarrow F^{\text{op}} \\ \leftarrow \cdot \\ \downarrow \theta^{\text{op}} \end{array} \right) G^{\text{op}} \cdot \left( \begin{array}{c} \cdot \\ \leq \\ \cdot \\ \downarrow c \end{array} \right) & \xrightarrow{R} & \mathbf{ISL} \\ \mathcal{D}^{\text{op}} & \xrightarrow{R} & \mathbf{ISL} \end{array}$$

so that, for every  $A$  in  $\mathcal{C}$  and every  $\alpha$  in  $P(A)$ , one has  $b_A(\alpha) \leq_{F(A)} R_{\theta_A}(c_A(\alpha))$ .

**2.2 Example.** A set-theoretic model for a first order theory  $\mathcal{T}$  an 1-arrow from  $LT: \mathcal{V}^{\text{op}} \rightarrow \mathbf{ISL}$  to  $\text{Sub}_{\text{Set}}: \text{Set}^{\text{op}} \rightarrow \mathbf{ISL}$  in **ED**. And a homomorphism between two set-theoretic models of  $\mathcal{T}$  determines a 2-arrow.

Given a doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$ , a category  $\mathcal{D}$  with finite products and a functor  $F: \mathcal{D} \rightarrow \mathcal{C}$  that preserves products, the composition of  $P$  with  $F^{\text{op}}: \mathcal{D}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$  gives a doctrine  $PF^{\text{op}}: \mathcal{D}^{\text{op}} \rightarrow \mathbf{ISL}$  called *change of base of  $P$  along  $F$* .

An *elementary doctrine on  $\mathcal{C}$*  is a primary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  such that for every object  $A$  in  $\mathcal{C}$ , there is an object  $\delta_A$  over  $A \times A$  such that for every arrow  $e$  of the form  $\langle \text{pr}_1, \text{pr}_2, \text{pr}_2 \rangle: X \times A \rightarrow X \times A \times A$  in  $\mathcal{C}$ , the assignment

$$\mathcal{E}_e(\alpha) := P_{\langle \text{pr}_1, \text{pr}_2 \rangle}(\alpha) \wedge_{X \times A \times A} P_{\langle \text{pr}_2, \text{pr}_3 \rangle}(\delta_A)$$

for  $\alpha$  in  $P(X \times A)$  determines a left adjoint to

$$P_e: P(X \times A \times A) \rightarrow P(X \times A).$$

It follows that the assignment

$$\mathcal{E}_{\langle \text{id}_A, \text{id}_A \rangle}(\alpha) := P_{\text{pr}_1}(\alpha) \wedge_{A \times A} \delta_A$$

for  $\alpha$  in  $P(A)$  determines a left adjoint to

$$P_{\langle \text{id}_A, \text{id}_A \rangle}: P(A \times A) \rightarrow P(A)$$

This means that  $\delta_A$  is determined uniquely for each object  $A$  in  $\mathcal{C}$  and hence we will refer to  $\delta_A$  as the *fibered equality on  $A$* .

An elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is said to have *comprehensive diagonals* if for every pair of parallel arrows  $f, g: X \rightarrow A$  in  $\mathcal{C}$  it holds that  $f = g$  if and only if  $\top_X = P_{\langle f, g \rangle}(\delta_A)$ . The change of base of an elementary doctrine is again elementary, but the new elementary doctrine might fail to have comprehensive diagonals.

**2.3 Examples.** (a) Consider a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$  and the associated primary doctrine  $LT: \mathcal{V}^{\text{op}} \rightarrow \mathbf{ISL}$  as in 2.1-(a). The doctrine  $LT$  is elementary if and only the equality is definable in  $\mathcal{T}$ .

(b) Let  $\mathcal{H}$  be an inf-semilattice. The primary doctrine  $\mathcal{P}_{\mathcal{H}}: \text{Set}^{\text{op}} \rightarrow \mathbf{ISL}$  as in 2.1-(b) is elementary if and only if  $\mathcal{H}$  has a bottom element (see [EPR20]). In this case  $\delta_A$  is the function that maps  $(a, a')$  to  $\top$  if  $a = a'$  and to  $\perp$  otherwise. In this case  $\mathcal{P}_{\mathcal{H}}$  has comprehensive diagonals.

(c) Suppose  $\mathcal{C}$  has finite limits. The doctrine  $\text{Sub}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  as in 2.1-(c) is elementary with comprehensive diagonals where  $\delta_A$  is represented by the diagonal on  $A$ . (d) Suppose  $\mathcal{C}$  has finite limits and weak pullbacks. The doctrine  $\Psi_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  as in 2.1-(d) is elementary with comprehensive diagonals where  $\delta_A$  is represented by the diagonal on  $A$ .

Elementary doctrines are the object of **ED**, the 2-full subcategory **PD** whose 1-morphisms are those 1-morphism  $(F, b): P \rightarrow R$  of **PD** such that for every object  $A$  in  $\mathcal{C}$ , the functor  $b_A: P(A) \rightarrow R(F(A))$  commutes with the left adjoints

$$\begin{array}{ccccc} P(X \times A) & \xrightarrow{b_{X \times A}} & R(F(X \times A)) & \xrightarrow[\sim]{R\langle F\text{pr}1, F\text{pr}2 \rangle} & R(FX \times FA) \\ \downarrow \mathcal{A}_e & & & & \downarrow \mathcal{A}_{e'} \\ P(X \times A \times A) & \xrightarrow{b_{X \times A \times A}} & R(F(X \times A \times A)) & \xrightarrow[\sim]{R\langle F\text{pr}1, F\text{pr}2, F\text{pr}3 \rangle} & R(FX \times FA \times FA) \end{array}$$

where  $e$  is  $\langle \text{pr}_1, \text{pr}_2, \text{pr}_2 \rangle: X \times A \rightarrow X \times A \times A$  and  $e'$  is  $\langle \text{pr}_1, \text{pr}_2, \text{pr}_2 \rangle: FX \times FA \rightarrow FX \times FA \times FA$ .

**2.4 Remark.** We recall from [MR13a] that it is possible to add comprehensive diagonals to an elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  as follows. Consider the category  $\mathcal{X}_P$ , the “*extensional collapse*” of  $P$  whose objects are the objects of  $\mathcal{C}$  and where an arrow  $[f]: A \rightarrow B$  is an equivalence class of morphisms  $f: A \rightarrow B$  in  $\mathcal{C}$  such that  $\delta_A \leq_{A \times A} P_{f \times f}(\delta_B)$  in  $P(A \times A)$  with respect to the equivalence which relates  $f$  and  $f'$  when  $\delta_A \leq_{A \times A} P_{f \times f'}(\delta_B)$ . Composition is given by that of  $\mathcal{C}$  on representatives, and identities are represented by identities of  $\mathcal{C}$ . We then define the doctrine  $P_x: \mathcal{X}_P^{\text{op}} \rightarrow \mathbf{ISL}$  on  $\mathcal{X}_P$  essentially as  $P$  itself.

Let **ExD** denote the full subcategory of **ED** on elementary doctrines with comprehensive diagonals.

**2.5 Proposition.** *There is left biadjoint to the inclusion of **ExD** into **ED** which on objects associates the doctrine  $P_x: \mathcal{X}_P^{\text{op}} \rightarrow \mathbf{ISL}$  to an elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$ .*

An elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{InfsL}$  is *existential* when, for  $A_1$  and  $A_2$  in  $\mathcal{C}$ , for a(ny) projection  $\text{pr}_i: A_1 \times A_2 \rightarrow A_i$ ,  $i = 1, 2$ , the functor  $P_{\text{pr}_i}: P(A_i) \rightarrow P(A_1 \times A_2)$  has a left adjoint  $\mathcal{A}_{\text{pr}_i}$ —we shall call such a left adjoint *existential*—and those left adjoints satisfy the

**Beck-Chevalley Condition:** for any pullback diagram

$$\begin{array}{ccc} X' & \xrightarrow{\text{pr}'} & A' \\ f' \downarrow & & \downarrow f \\ X & \xrightarrow{\text{pr}} & A \end{array}$$

with  $\text{pr}$  a projection (hence also  $\text{pr}'$  a projection), for any  $\beta$  in  $P(X)$ , the natural inequality  $\mathcal{A}_{\text{pr}'} P_{f'}(\beta) \leq P_f \mathcal{A}_{\text{pr}}(\beta)$  in  $P(A')$  is an identity;

**Frobenius Reciprocity:** for  $\text{pr}: X \rightarrow A$  a projection,  $\alpha$  in  $P(A)$ ,  $\beta$  in  $P(X)$ , the natural inequality  $\mathcal{A}_{\text{pr}}(P_{\text{pr}}(\alpha) \wedge \beta) \leq \alpha \wedge \mathcal{A}_{\text{pr}}(\beta)$  in  $P(A)$  is an identity.

**2.6 Examples.** (a) Consider a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$  and the associated primary doctrine  $LT: \mathcal{V}^{\text{op}} \longrightarrow \mathbf{ISL}$  as in 2.1-(a). The doctrine  $LT$  is existential where left adjoints along projections are given by the existential quantification.

(b) Let  $\mathcal{H}$  be an inf-semilattice. The primary doctrine  $\mathcal{P}_{\mathcal{H}}: \text{Set}^{\text{op}} \longrightarrow \mathbf{ISL}$  is existential if and only if  $\mathcal{H}$  is a frame (see [EPR20]).

(c) Suppose  $\mathcal{C}$  has finite limits. The doctrine  $\text{Sub}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  as in 2.1-(c) is existential if and only if  $\mathcal{C}$  is regular (see [Jac99]).

(d) Suppose  $\mathcal{C}$  has finite limits and weak pullbacks. The doctrine  $\Psi_{\mathcal{C}}: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  as in 2.1-(d) is existential where left adjoint are given by composition.

**2.7 Remark.** As shown by Lawvere, in an elementary existential doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  every map of the form  $P_f$  has a left adjoint  $\mathcal{A}_f$ . For  $f: A \rightarrow B$  the maps  $\mathcal{A}_f: P(A) \rightarrow P(B)$  is

$$\mathcal{A}_f(\alpha) := \mathcal{A}_{\text{pr}_2}[P_{f \times \text{id}_B}(\delta_B) \wedge P_{\text{pr}_1}(\alpha)]$$

see also [Pit02]. Moreover these left adjoint satisfy the Frobenius Reciprocity in the sense that for every  $\beta$  in  $P(B)$  it holds  $\mathcal{A}_f(\alpha) \wedge \beta = \mathcal{A}_f(\alpha \wedge P_f(\beta))$ .

This allows to give a quick description of the 2-category **EED**, which is the 2-full subcategory of **ED** on those elementary doctrines that are existential with comprehensive diagonals and whose 1-morphisms are those  $(F, b): P \rightarrow R$  such that for every  $f: X \rightarrow A$  in  $\mathcal{C}$  the following commute

$$\begin{array}{ccc} P(X) & \xrightarrow{b_X} & R(FX) \\ \mathcal{A}_f \downarrow & & \downarrow \mathcal{A}_{Ff} \\ P(A) & \xrightarrow{b_A} & R(FA) \end{array}$$

The full subcategory of **EED** on those existential elementary doctrines with comprehensive diagonals is **EExD**.

A primary doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{InfSL}$  is *implicational* if, for every object  $A$  in  $\mathcal{C}$ , every  $\alpha$  in  $P(A)$ , the functor  $\alpha \wedge -: P(A) \rightarrow P(A)$  has a right adjoint  $\alpha \Rightarrow -: P(A) \rightarrow P(A)$ . A primary doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{InfSL}$  is *disjunctive* if every  $P(A)$  has finite distributive joins and every  $P_f$  preserves them. A primary doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{InfSL}$  is *universal* if, for  $A_1$  and  $A_2$  in  $\mathcal{C}$ , for a (ny) projection  $\text{pr}_i: A_1 \times A_2 \rightarrow A_i$ ,  $i = 1, 2$ , the functor  $P_{\text{pr}_i}: P(A_i) \rightarrow P(A_1 \times A_2)$  has a right adjoint  $V_{\text{pr}_i}$ , and these satisfy the Beck-Chevalley condition: for any pullback diagram

$$\begin{array}{ccc} X' & \xrightarrow{\text{pr}'} & A' \\ f' \downarrow & & \downarrow f \\ X & \xrightarrow{\text{pr}} & A \end{array}$$

with  $\text{pr}$  a projection (hence also  $\text{pr}'$  a projection), for any  $\beta$  in  $P(X)$ , the canonical arrow  $P_f V_{\text{pr}}(\beta) \leq V_{\text{pr}'} P_{f'}(\beta)$  in  $P(A')$  is iso.

**2.8 Definition.** A *first order doctrine* (f.o.d.) on  $\mathcal{C}$  is an existential elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  which is also implicational, disjunctive and universal.

**2.9 Remark.** As for left adjoint in 2.7 Lawvere proved that in a first order doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  every map of the form  $P_f$  has a right adjoint  $V_f$ . For  $f: A \rightarrow B$  the maps  $V_f: P(A) \rightarrow P(B)$  is  $V_f(\alpha) := V_{\text{pr}_2}[P_f \times_{\text{id}_B}(\delta_B) \Rightarrow P_{\text{pr}_1}(\alpha)]$  see also [Pit02].

**2.10 Examples.** (a) Consider a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$  and the associated primary doctrine  $LT: \mathcal{V}^{\text{op}} \rightarrow \mathbf{ISL}$  as in 2.1-(a). If  $\mathcal{L}$  then the doctrine  $LT$  is first order. In each fibre joins are given by disjunctions while the cartesian closure is provided by the implication. Right adjoints are given by universal quantification.

(b) Let  $\mathcal{H}$  be a frame. Then it is a complete Heyting algebra. Hence the doctrine  $\mathcal{P}_{\mathcal{H}}: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{ISL}$  is first order where in each fibre the Heyting algebra operations are computed point-wise and right adjoint are give by arbitrary infima (see [Pit02] for details).

(c) Suppose  $\mathcal{C}$  has finite limits. The doctrine  $\text{Sub}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is first order if and only if  $\mathcal{C}$  is an Heyting category (or a logos) in the sense of [FS90].

(d) Suppose  $\mathcal{C}$  has finite limits. If  $\mathcal{C}$  has finite coproducts and is weakly locally cartesian closed, then  $\Psi_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  as in 2.1-(d) is first order. We shall see later how to generalise the description to the case in which  $\mathcal{C}$  is assumed to have weak pullbacks.

We will often deal with different doctrines on the same base and in several situations one of these is  $\Psi_{\mathcal{C}}$ , thus we find it convenient to adopt a specific notation to distinguish operations between doctrines and in particular operations in  $\Psi_{\mathcal{C}}$ . We will use the following.

**2.11 Notation.** We write  $\Psi_{\mathcal{C}}(f)$  as  $f^*$ . The left adjoint to  $f^*$  will be denoted by  $\Sigma_f$ . If  $f^*$  has a right adjoint this will be denoted by  $\Pi_f$ . The equality predicated over  $A$  will be denoted by  $[\text{id}_A, \text{id}_A]$ . Binary meets in  $\Psi_{\mathcal{C}}(A)$  will be denoted by  $[f] \times_A [g]$  while the top element over  $A$  will be denoted by  $[\text{id}_A]$ . Joins will be denoted by  $[f] +_A [g]$ . The bottom element will be  $0_A$ . We will freely confuse a class with any of its representatives.

In a primary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  an object  $\Omega$  in  $\mathcal{C}$  is a *weak predicate classifier* if there is an element  $\epsilon$  over  $\Omega$  such that for every  $\phi$  in  $P(A)$  there is a (not necessarily unique) morphism  $\chi_{\phi}: A \rightarrow \Omega$  satisfying  $P_{\chi_{\phi}}(\epsilon) = \phi$ .  $P$  has a *strong predicate classifier* if it has a weak predicate classifier and arrows of the form  $\chi_{\phi}$  are unique.  $P$  has *weak power objects* if for every  $\mathcal{C}$ -object  $A$  there exists an object  $\mathbb{P}A$  in  $\mathcal{C}$  and an object  $\epsilon_A$  in  $P(A \times \mathbb{P}A)$  such that for every  $Y$  and relation  $\phi$  in  $P(A \times Y)$  there is a (not necessarily unique) morphism  $\chi_{\phi}: Y \rightarrow \mathbb{P}A$  satisfying  $P_{\text{id}_A \times \chi_{\phi}}(\epsilon_A) = \phi$ .  $P$  has a *strong power object* if it has a weak power objects and arrows of the form  $\chi_{\phi}$  are unique.

If  $P$  as (weak) power objects, then it has also a (weak) predicate classifier (it suffices to choose  $\mathbb{P}1$  as  $\Omega$ ). If the base  $\mathcal{C}$  is weakly cartesian closed, then

$P$  has a (weak) predicate classifier if and only if it has (weak) power objects, where  $\mathbb{P}A$  can be chosen as the weak exponential of  $\Omega$  to the power of  $A$ .

**2.12 Examples.** (a) The doctrine  $LT: \mathcal{V}^{\text{op}} \longrightarrow \mathbf{ISL}$  built out of a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$  has no predicate classifiers.

(b) Let  $\mathcal{H}$  be an inf-semilattice. The underling set of  $\mathcal{H}$  is  $|\mathcal{H}|$  and is a strong predicate classifier of  $\mathcal{P}_{\mathcal{H}}$  where  $\in$  is the identity  $\text{id}_{|\mathcal{H}|}$ .

(c) Suppose  $\mathcal{C}$  has finite limits. The doctrine  $\text{Sub}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  as a predicate classifier if and only if it has a subobject classifier.

(d) Suppose  $\mathcal{C}$  has finite products and weak pullbacks. If  $\mathcal{C}$  has a weak predicate classifier if and only if it has a weak proof classifier in the sense of [Men03].

**2.13 Definition.** A primary doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  is a tripos if  $P$  is a first order doctrine with weak power objects.  $P$  is a **strong tripos** if it is a tripos with strong power objects.

**2.14 Examples.** (a) The doctrine  $LT: \mathcal{V}^{\text{op}} \longrightarrow \mathbf{ISL}$  built out of a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$  is not a tripos as it has no predicate classifiers.

(b) Let  $\mathcal{H}$  be an inf-semilattice. The doctrine  $\mathcal{P}_{\mathcal{H}}$  as in 2.1 is a tripos if and only if  $\mathcal{H}$  is a frame.

(c) Suppose  $\mathcal{C}$  has finite limits. The doctrine  $\text{Sub}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  is a strong tripos if and only if  $\mathcal{C}$  is an elementary topos.

(d) Suppose  $\mathcal{C}$  has finite products and pullbacks. If  $\mathcal{C}$  has a weak predicate classifier if and only if it has a weak proof classifier in the sense of [Men03].

A proof of the following proposition is in [Pas16].

**2.15 Proposition.** *If the doctrine  $P$  is a tripos then  $P_x$  is a tripos, too.*

Elementary doctrines are the cloven Eq-fibrations of [Jac99] and, as explained in *loc.cit.*, there is a deductive calculus associated to those which is that of the  $\wedge$ -fragment over type theory with just a unit type and a binary product type constructor. To fix notation, we will use the following.

**2.16 Notation.** Let  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  be elementary. Write

$$a_1: A_1, \dots, a_k: A_k \mid \phi_1, \dots, \phi_n \vdash \psi$$

in place of

$$\phi_1 \wedge \dots \wedge \phi_n \leq \psi \quad \text{in } P(A_1 \times \dots \times A_k)$$

and call such an expression sequent. Note that, in line with *loc.cit.*,  $\delta_A$  in  $P(A \times A)$  will be written as  $a: A, a': A \mid a =_A a'$ . Also write  $a: A \mid \alpha \dashv\vdash \beta$  to abbreviate  $a: A \mid \alpha \vdash \beta$  and  $a: A \mid \beta \vdash \alpha$ . Say that  $\alpha$  in  $P(A)$  is true over  $A$  if  $\top_A \leq \alpha$ . An element of  $P(1)$  will be called sentence. An arrow  $r: 1 \rightarrow A$  will be called constant (of type  $A$ ) and for  $\alpha$  in  $P(A)$  we write  $\alpha(r)$  in place of  $P_r(\alpha)$ . If  $P$  is existential and  $a: A, x: X \mid \phi$ , *i.e.*  $\phi$  is in  $P(A \times X)$ , write  $a: A \mid \exists_{x: X} \phi$  in place of  $\exists_{\text{pr}_1} \phi$  in  $P(A)$ . Similarly when  $P$  is first order, for  $\alpha, \beta$  in  $P(A)$  and  $\phi$  in  $P(A \times X)$  write  $a: A \mid \alpha \vee \beta$  and  $a: A \mid \alpha \Rightarrow \beta$  and  $a: A \mid \forall_{x: X} \phi$  in place of  $\alpha \vee \beta$  and  $\alpha \rightarrow \beta$  and  $\forall_{\text{pr}_1} \phi$  in  $P(A)$ . If  $P$  is a tripos write  $a: A, U: \mathbb{P}A \mid a \in_A U$  in place of  $\in_A$  in  $P(A \times \mathbb{P}A)$ .

From now on we feel free to employ this logical language in our proofs or definitions whenever we feel that readability is improved.

### 3 Elementary quotient completions

We now recall the definition of *P-equivalence relation* for any elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  and the related notion of *quotients* from [MR13a]:

**3.1 Definition.** A *P-equivalence relation*  $\rho$  over an object  $A$  of  $\mathcal{C}$  is an element in  $P(A \times A)$  such that

- (i)  $\delta_A \leq \rho$
- (ii)  $\rho = P_{\langle \text{pr}_2, \text{pr}_1 \rangle}(\rho)$
- (iii)  $P_{\langle \text{pr}_1, \text{pr}_2 \rangle}(\rho) \wedge P_{\langle \text{pr}_2, \text{pr}_3 \rangle}(\rho) \leq P_{\langle \text{pr}_1, \text{pr}_3 \rangle}(\rho)$

Where in (ii)  $\text{pr}_1, \text{pr}_2: A \times A \rightarrow A$  are the first and the second projection, while in (iii)  $\text{pr}_i$ , with  $i = 1, 2, 3$ , are projections from  $A \times A \times A$  to each of the factors.

When no confusion arises, we shall refer at *P-equivalence relations* simply as *equivalence relations*, without specifying the doctrine  $P$ . Note that in every elementary doctrine, fibered equalities are equivalence relations.

**3.2 Examples.** (a) Recall from 2.1-(a) the syntactic doctrine  $LT: \mathcal{V}^{\text{op}} \rightarrow \mathbf{ISL}$  built out of a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$ . A *LT-equivalence relation* over  $x$  is a formula  $\phi$  of  $\mathcal{L}$  such that  $\mathcal{T} \vdash \forall x \phi(x, x)$  and  $\mathcal{T} \vdash \forall xy \phi(x, y) \rightarrow \phi(y, x)$  and  $\mathcal{T} \vdash \forall xyz (\phi(x, y) \& \phi(y, z)) \rightarrow \phi(x, z)$ .

(b) When  $\mathcal{H}$  is an inf-semilattice, a  $\mathcal{P}_{\mathcal{H}}$ -equivalence relation over a set  $A$  is an  $\mathcal{H}$ -valued ultra-pseudodistance on  $A$ , *i.e.* a function  $\rho: A \times A \rightarrow \mathcal{H}$  such that for all  $a, a', a''$  in  $A$  it is  $\rho(a, a) = \top$  and  $\rho(a, a') = \rho(a', a'')$  and  $\rho(a, a') \wedge \rho(a', a'') \leq \rho(a, a'')$ .

(c) Suppose  $\mathcal{C}$  has finite limits, then  $\rho$  is a  $\text{Sub}_{\mathcal{C}}$ -equivalence relation over  $A$  if and only if  $\rho$  is an equivalence relation of  $\mathcal{C}$  in the usual categorical sense (see for example [MM92]).

(d) Suppose  $\mathcal{C}$  has finite products and weak pullbacks, then  $\rho$  is a  $\Psi_{\mathcal{C}}$ -equivalence relation over  $A$  if and only if it is a pseudo-equivalence relation of  $\mathcal{C}$  in the sense of [Car95].

An elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is said to have *quotients* if for every  $A$  in  $\mathcal{C}$  and every equivalence relation  $\rho$  over  $A$  there exists a morphism  $q: A \rightarrow A/\rho$  such that  $\rho \leq P_{q \times q}(\delta_{A/\rho})$  and for every morphism  $f: A \rightarrow Y$  such that  $\rho \leq P_{f \times f}(\delta_Y)$  there exists a unique  $h: A/\rho \rightarrow Y$  with  $hq = f$ . Maps of the form  $q: A \rightarrow A/\rho$  will be called *quotient arrow* of  $\rho$ . The quotient  $q: A \rightarrow A/\rho$  is *effective* if  $\rho = P_{q \times q}(\delta_{A/\rho})$ .

An elementary doctrine  $P$  is said to have *stable quotients* if for every pullback  $qp = fh$ , if  $q$  is a quotient arrow of  $\rho$ , then  $h$  is a quotient arrow of  $P_{p \times p}(\rho)$ .

Given an elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{InfSL}$  and a  $P$ -equivalence relation  $\rho$  over  $A$  in  $\mathcal{C}$ , the inf-semilattice of **descent data**  $\mathcal{Des}_\rho$  is the sub-inf-semilattice of  $P(A)$  on those  $\alpha$  such that

$$P_{\text{pr}_1}(\alpha) \wedge \rho \leq P_{\text{pr}_2}(\alpha),$$

where  $\text{pr}_1, \text{pr}_2: A \times A \rightarrow A$  are the projections.

For  $f: A \rightarrow B$  in  $\mathcal{C}$  the map  $P_f: P(B) \rightarrow P(A)$  takes values in  $\mathcal{Des}_{P_f \times_f(\delta_B)}$ . We shall say that  $f$  is **of effective descent** if  $P_f: P(B) \rightarrow \mathcal{Des}_{P_f \times_f(\delta_B)}$  is an isomorphism.

In particular this means that the functor  $P_f$  is of *effective descent type* as defined in [BW84].

**3.3 Definition.** An elementary doctrine  $P$  is said to have **descent effective quotients** if  $P$  has stable effective quotients and the quotient arrows are of effective descent.

**3.4 Examples.** (a) In general doctrines of the form  $LT: \mathcal{V}^{\text{op}} \rightarrow \mathbf{ISL}$  do not have quotients.

(b) For  $\mathcal{H}$  an inf-semilattice, the doctrine  $\mathcal{P}_{\mathcal{H}}$  has quotient. Let  $\rho: A \times A \rightarrow \mathcal{H}$  be a  $\mathcal{H}$ -valued ultra-pseudodistance on  $A$  as in 3.2-(b) and define on  $A$  the equivalence relation  $a \sim a'$  generated by  $\rho(a, a') = \top$ . Then canonical surjection  $q: A \rightarrow A/\sim$  is a quotient arrow. These quotients are not effective and hence not of effective descent.

(c) If  $\mathcal{C}$  has finite limits, then  $\text{Sub}_{\mathcal{C}}$  has effective quotient if and only if  $\mathcal{C}$  is exact. In this case quotients are of effective descent (this follows also from the more general situation of 3.6).

(d) In general doctrines of the form  $\Psi_{\mathcal{C}}$  do not have quotients.

**3.5 Lemma.** *If  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is elementary and existential, an effective quotient arrow is of effective descent if and only if it is  $P$ -surjective.*

*Proof.* Let  $q: A \rightarrow A/\rho$  be an effective quotient arrow (so  $\rho = P_{q \times q}(\delta_{A/\rho})$ ). If  $q$  is of effective descent, then  $P_q: P(A/\rho) \rightarrow \mathcal{Des}_\rho$  is an isomorphism. Its inverse is the restriction of  $\mathcal{E}_q$  to  $\mathcal{Des}_\rho$ , whence  $\mathcal{E}_q \top_A = \top_{A/\rho}$  since for all descent data  $\phi$  we have that  $\phi = P_q \mathcal{E}_q \phi$  by Beck-Chevalley conditions applied to  $\mathcal{E}_q$  described as in remark 2.13 p.381 in [MR13b]. Conversely for every  $\psi$  in  $P(B)$  it holds  $\mathcal{E}_q P_q \psi = \psi \wedge \mathcal{E}_q \top_B = \psi$  and hence  $\psi$  is a descent datum.  $\square$

**3.6 Proposition.** *The quotient arrows of an existential  $m$ -variational doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  with stable effective quotients are of effective descent.*

*Proof.* By 4.21 each quotient arrow  $q: A \rightarrow A/\rho$  can be factored as  $q = \{\{\mathcal{E}_q(\top_A)\}\}g$  where  $g: A \rightarrow C$  is  $P$ -surjective. Since  $\{\{\mathcal{E}_q(\top_A)\}\}$  is monic, it is also  $P$ -injective by 4.8, whence  $\rho = P_{q \times q}(\delta_B) = P_{g \times g} P_{\{\{\mathcal{E}_q(\top_A)\}\} \times \{\{\mathcal{E}_q(\top_A)\}\}}(\delta_B) = P_{g \times g}(\delta_C)$ . The universal property of quotients implies the existence of an arrow  $s: A/\rho \rightarrow C$  with  $g = sq$  and hence  $q = \{\{\mathcal{E}_q(\top_A)\}\}sq$ , so  $\{\{\mathcal{E}_q(\top_A)\}\}s = \text{id}_B$  as  $q$  is epic. Since  $\{\{\mathcal{E}_q(\top_A)\}\}$  is a monomorphism with a section it is an isomorphism. Therefore  $q$  is isomorphic to  $g$  and hence  $P$ -surjective. An application of Lemma 3.5 concludes the proof.  $\square$

We recall from [MR13b] the construction called *elementary quotient completion* that freely adds descent effective quotients to any elementary doctrine.

Given an elementary doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  we call  $\mathcal{Q}_P$  the category whose objects are pairs  $(A, \rho)$  in which  $A$  is in  $\mathcal{C}$  and  $\rho$  is an equivalence relation over  $A$ . An arrow  $[f]: (A, \rho) \rightarrow (B, \sigma)$  is an equivalence class of arrows  $f: A \rightarrow B$  in  $\mathcal{C}$  such that  $\rho \leq P_{f \times f}(\sigma)$ , with respect to the equivalence  $f \sim g$  if and only if

$$\top_A = P_{\langle f, g \rangle}(\sigma)$$

The category  $\mathcal{Q}_P$  has finite products: if  $(A, \rho)$  and  $(B, \sigma)$  are objects of  $\mathcal{Q}_P$ , their product is

$$(A, \rho) \xleftarrow{[\text{pr}_1]} (A \times B, \rho \boxtimes \sigma) \xrightarrow{[\text{pr}_2]} (B, \sigma)$$

where  $\rho \boxtimes \sigma = P_{\langle \text{pr}_1, \text{pr}_2 \rangle}(\rho) \wedge P_{\langle \text{pr}_3, \text{pr}_4 \rangle}(\sigma)$ . The elementary quotient completion of  $P$  is the doctrine  $\widehat{P}: \mathcal{Q}_P^{\text{op}} \longrightarrow \mathbf{ISL}$  where

$$\widehat{P}(A, \rho) = \mathcal{D}es_\rho \quad \widehat{P}_{[f]} = P_f$$

It is proved in [MR13b] that the assignment on arrows does not depend on the choice of representatives. The doctrine  $\widehat{P}$  is elementary with  $\delta_{(A, \rho)} = \rho$ . It is immediate to see that  $\widehat{P}$  as stable descent effective quotient: if  $\sigma$  is an equivalence relation over  $(A, \rho)$ , then its quotient arrow  $[\text{id}_A]: (A, \rho) \rightarrow (A, \sigma)$ . Moreover these quotients are stable.

**3.7 Examples.** (a) The elementary quotient completion of doctrines of the form  $LT: \mathcal{V}^{\text{op}} \longrightarrow \mathbf{ISL}$  is connected to elimination of imaginaries as analysed in [EPR20].

(b) Assuming choice, in the sense that epimorphism in *Set* split, the base of  $\widehat{\mathcal{P}}_{\mathcal{H}}: \widehat{\mathcal{S}et}_{\mathcal{P}_{\mathcal{H}}}^{\text{op}} \longrightarrow \mathbf{ISL}$  is equivalent to  $\mathbf{UM}_{\mathcal{H}}$ , the category of  $\mathcal{H}$ -valued ultrametric spaces. Indeed the functor that maps  $f: (A, \rho) \rightarrow (B, \sigma)$  in  $\mathbf{UM}_{\mathcal{H}}$  to  $[f]: (A, \rho) \rightarrow (B, \sigma)$  in  $\widehat{\mathcal{S}et}_{\mathcal{P}_{\mathcal{H}}}$  is full and faithful as the identity of indiscernibles holds. For essential surjectivity, take  $(A, \rho)$  in  $\widehat{\mathcal{S}et}_{\mathcal{P}_{\mathcal{H}}}$  and consider the quotient  $(A, \rho) \rightarrow (A/\sim, \delta_{A/\sim})$  where  $\sim$  is the equivalence relation described in 3.4-(b). Any of its section represent an inverse in  $\widehat{\mathcal{S}et}_{\mathcal{P}_{\mathcal{H}}}$ .

(d) One the motivating examples for the study of the elementary quotient completion is given by doctrines of the form  $\Psi_{\mathcal{C}}$  where  $\mathcal{C}$  has finite products and weak pullbacks. As proved in [MR13b] the doctrine  $\widehat{\Psi}_{\mathcal{C}}$  is  $\text{Sub}_{\mathcal{C}_{\text{ex/lex}}}$ .

It is quite apparent that the elementary structure plays no role in the definitions of  $\widehat{P}$  (see [Pas15, EPR20]) but it is crucial to embed  $\mathcal{C}$  into  $\mathcal{Q}_P$ . The embedding is given by the functor  $\nabla_P: \mathcal{C} \rightarrow \mathcal{Q}_P$  that assigns to each  $f: X \rightarrow Y$  the arrow  $[f]: (X, \delta_X) \rightarrow (Y, \delta_Y)$ . This functor is full and preserves binary products and when  $P$  has comprehensive diagonals it is also faithful. In this case it is immediate to check that  $P$  is the change of base of  $\widehat{P}$  along  $\nabla_P$ .

Denote by **QEExD** the subcategory of **EExD** on those doctrines with effective descent quotients and on those arrows that preserves quotients, *i.e.* on those  $(F, b)$  from  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  to  $R: \mathcal{D}^{\text{op}} \rightarrow \mathbf{ISL}$  such that the action of  $F$  on  $q: A \rightarrow A/\rho$  is  $Fq: FA \rightarrow FA/(b_{A \times A} \rho)$ .

The following theorem is proved in [MR13b] [MR13a].

**3.8 Proposition.** *There is a left biadjoint to the inclusion of **QEExD** into **EExD** which associates the doctrine  $\hat{P}: \mathcal{Q}_P^{\text{op}} \rightarrow \mathbf{ISL}$  to an elementary doctrine  $P$ .*

Let  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  be in **EED**. Using the language in 2.16, a  $F$  in  $P(A \times B)$

**total (or entire)** if  $\exists_{b:B} F(a, b)$  is true over  $A$

**single-valued (or functional)** if  $a: A, b: B, b': B \mid F(a, b) \wedge F(a, b') \vdash b =_B b'$

We say that  $P$  satisfies the **rule of unique choice** (RUC) if for every total and single-valued  $F$  in  $P(A \times B)$  there is  $f: A \rightarrow B$  such that  $a: A, b: B \mid F(a, b) \dashv\vdash f(a) =_B b$ . We say that  $P$  satisfies the **rule of choice** (RC) if for every total  $F$  in  $P(A \times B)$  there is  $f: A \rightarrow B$  such that  $a: A \mid \exists_{b:B} F(a, b) \dashv\vdash F(a, f(a))$ .

An elementary and existential doctrine  $P$  satisfies (RC) if and only if its completion  $\hat{P}$  satisfies (RC) (see [MPR17]).

Let **EExD**<sup>RUC</sup> and **EExD**<sup>RC</sup> be the full subcategories of **EExD** on doctrines satisfying (RUC) and (RC) respectively and similarly **QEExD**<sup>RUC</sup> and **QEExD**<sup>RC</sup> are the full subcategories of **QEExD** on doctrines satisfying (RUC) and (RC) respectively. These categories fit in the following diagram of inclusions.

$$\begin{array}{ccc}
 \mathbf{EExD}^{RC} & \hookrightarrow & \mathbf{EExD} \\
 \uparrow \dashv\vdash & & \uparrow \dashv\vdash \\
 \mathbf{QEExD}^{RUC} & \hookrightarrow & \mathbf{QEExD}
 \end{array} \tag{1}$$

**3.9 Remark.** The vertical left adjoint in diagram (1) are given by the elementary quotient completion. The biadjoint to the inclusion of **QEExD**<sup>RUC</sup> into **QEExD** is the functor that maps  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  to  $P_F: \mathcal{EF}_P^{\text{op}} \rightarrow \mathbf{ISL}$  where  $\mathcal{C}_F$  has the objects of  $\mathcal{C}$  and an arrow  $F: A \rightarrow B$  is a total and single-valued relation in  $P(A \times B)$ . The functor  $P_F$  maps  $A$  to  $P_F(A) = P(A)$  and given  $\beta$  in  $P(B)$  it is  $P_F(\beta)$  is the formula  $a: A \mid \exists_{b:B} [F(a, b) \wedge \beta(b)]$  over  $A$  (see [Pas16]).

## 4 Comprehension and strong monomorphisms

**4.1 Definition.** A primary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is said to have **weak comprehension** if for every  $A$  in  $\mathcal{C}$  and every  $\alpha$  in  $P(A)$  there is an arrow  $\{\alpha\}: X \rightarrow A$  with  $\top_X = P_{\{\alpha\}}(\alpha)$  such that for every arrow  $f: Y \rightarrow A$  with  $\top_Y \leq P_f(\alpha)$  there is a (not necessarily unique) arrow  $k: Y \rightarrow X$  with  $\{\alpha\}k = f$ . We shall say that comprehension is **strong** if the mediating arrow  $k$  is the unique

such arrow. We shall say that comprehension is **full** if for every  $A$  and every  $\alpha, \beta$  over  $A$  it holds that  $\alpha \leq \beta$  if and only if  $\top_X = P_{\{\alpha\}}(\beta)$ .

It is an easy check that a doctrine  $P$  with weak comprehensions has strong comprehension if and only if arrows of the form are monic.

**4.2 Examples.** (a) The doctrine  $LT: \mathcal{V}^{\text{op}} \longrightarrow \mathbf{ISL}$  built out of a theory  $\mathcal{T}$  over a first order language  $\mathcal{L}$  has no comprehension.

(b) When  $\mathcal{H}$  is an inf-semilattice, the doctrine  $\mathcal{P}_{\mathcal{H}}$  has strong comprehension. Given  $\alpha$  in  $\mathcal{P}_{\mathcal{H}}(A)$  the arrow  $\{\alpha\}$  is the inclusion of  $\{a \in A \mid \alpha(a) = \top\}$  into  $A$ . This comprehension is not full as one can see taking  $\alpha': A \rightarrow \mathcal{H}$  that agrees with  $\alpha$  only on those elements  $a$  of  $A$  such that  $\alpha(a) = \top$ . Then  $\alpha$  and  $\alpha'$  have the same comprehension arrow, even if in general they may be different.

(c) Suppose  $\mathcal{C}$  has finite limits. The doctrine  $\text{Sub}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  as full strong comprehension. The comprehension  $\alpha$  in  $\text{Sub}_{\mathcal{C}}(A)$  is any representative of  $\alpha$ .

(d) Suppose  $\mathcal{C}$  has finite products and weak pullbacks. The doctrine  $\Psi_{\mathcal{C}}: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  as full weak comprehension. The comprehension  $\alpha$  in  $\text{Sub}_{\mathcal{C}}(A)$  is any representative of  $\alpha$  (that need not be monic).

**4.3 Remark.** An arrow of the form  $\{\alpha\}$  will be often called comprehension arrow of  $\alpha$ . Observe that an elementary doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  has **comprehensive diagonals** if and only if diagonals in  $\mathcal{C}$  are the strong comprehension arrows of the corresponding fibered equalities, which justifies the terminology for this property introduced in [MR13b]. Indeed take  $f, g: X \rightarrow A$  with  $\top_X = P_{\langle f, g \rangle}(\delta_A)$ . It is an easy check that strong comprehensions are stable under pullback, so the pullback of  $\{\delta_A\}$  along  $\langle f, g \rangle$  is  $\{\top_X\}$ , which is necessarily isomorphic to  $\text{id}_X$ . If  $\{\delta_A\}$  is  $\langle \text{id}_A, \text{id}_A \rangle$  then the pullback of  $\langle \text{id}_A, \text{id}_A \rangle$  along  $\langle f, g \rangle$  is  $\text{id}_X$  so  $f = g$ . The converse is immediate.

**4.4 Proposition.** *If  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  is elementary with comprehensive diagonals and strong comprehensions, then  $\mathcal{C}$  has finite limits.*

*Proof.* The equalizer of  $f, g: X \rightarrow A$  is  $\{P_{\langle f, g \rangle}(\delta_A)\}$ . □

**4.5 Proposition.** *Suppose  $P$  is elementary existential with weak comprehension. Weak comprehension is full if and only if for every  $\{\alpha\}: X \rightarrow A$  it holds  $\mathcal{A}_{\{\alpha\}}(\top_X) = \alpha$ .*

*Proof.* Let  $\mathcal{A}_{\{\alpha\}} \dashv P_{\{\alpha\}}$ , from  $\top_X = P_{\{\alpha\}}(\alpha)$  it follows  $\mathcal{A}_{\{\alpha\}}(\top_X) \leq \alpha$ . Instead  $\alpha \leq \mathcal{A}_{\{\alpha\}}(\top_X)$  follows by fullness from the adjunction unit  $\top_X \leq P_{\{\alpha\}}(\mathcal{A}_{\{\alpha\}}(\top_X))$ .

Conversely, if  $\mathcal{A}_{\{\alpha\}}(\top_X) = \alpha$  from  $\top_X = P_{\{\alpha\}}(\beta)$  it follows  $\alpha = \mathcal{A}_{\{\alpha\}}(\top_X) = \mathcal{A}_{\{\alpha\}}P_{\{\alpha\}}(\beta) \leq \beta$  by the adjunction counit of  $\mathcal{A}_{\{\alpha\}} \dashv P_{\{\alpha\}}$ . □

**4.6 Proposition.** *If  $P$  has full weak comprehension and comprehensive diagonals, then for every weak pullback  $fg = hk$  one has  $P_f \mathcal{A}_h = \mathcal{A}_g P_k$ .*

*Proof.* This is theorem 2.19 in [MPR17]. □

Suppose that  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  has weak comprehensions. For every  $f: Y \rightarrow B$  in  $\mathcal{C}$  and every  $\alpha$  in  $P(B)$  there is a commutative square

$$\begin{array}{ccc} X & \xrightarrow{m} & A \\ \{\!\{P_f(\alpha)\}\!\} \downarrow & & \downarrow \{\!\{\alpha\}\!\} \\ Y & \xrightarrow{f} & B \end{array}$$

where  $m$  is the mediating arrow coming from the universal property of  $\{\!\{\alpha\}\!\}$  as  $P_{f\{\!\{P_f(\alpha)\}\!\}}(\alpha) = P_{\{\!\{P_f(\alpha)\}\!\}}(P_f(\alpha)) = \top_X$ .

**4.7 Proposition.** *If  $k: Z \rightarrow A$  and  $h: Z \rightarrow Y$  are such that  $\{\!\{\alpha\}\!\}k = fh$ , then  $h$  factors through  $\{\!\{P_f(\alpha)\}\!\}$ . If  $\{\!\{\alpha\}\!\}$  is a strong comprehension, the square above is a weak pullback. If both  $\{\!\{\alpha\}\!\}$  and  $\{\!\{P_f(\alpha)\}\!\}$  are strong comprehensions the square is a pullback.*

*Proof.* Consider  $k$  and  $h$  such that  $\{\!\{\alpha\}\!\}k = fh$ . Then  $P_h(P_f(\alpha)) = P_k(P_{\{\!\{\alpha\}\!\}}(\alpha)) = \top_Z$ . Weak universality of  $\{\!\{P_f(\alpha)\}\!\}$  guarantees the existence of  $u: Z \rightarrow X$  with  $\{\!\{P_f(\alpha)\}\!\}u = h$ . If  $\{\!\{\alpha\}\!\}$  is strong, then it is also monic. Hence  $mu = k$  if and only if  $\{\!\{\alpha\}\!\}mu = \{\!\{\alpha\}\!\}k$ , which is true as  $\{\!\{\alpha\}\!\}mu = f\{\!\{P_f(\alpha)\}\!\}u = fh = \{\!\{\alpha\}\!\}k$ . Finally if  $\{\!\{P_f(\alpha)\}\!\}$  is monic the mediating arrow  $u$  is necessarily unique.  $\square$

If  $P$  is elementary, we say that an arrow  $f: X \rightarrow A$  of  $\mathcal{C}$  is  *$P$ -injective* if  $P_{f \times f}(\delta_A) = \delta_X$ . While if  $P$  is also existential we say that  $f$  is  *$P$ -surjective* if  $\mathcal{A}_f(\top_X) = \top_A$ .

It is immediate to show that if  $P$  has comprehensive diagonals, if an arrow is  $P$ -injective, then it is also monic, and if an arrow is  $P$ -surjective, then it is also epic.

**4.8 Proposition.** *Suppose  $P$  is a variational doctrine on  $\mathcal{C}$ . An arrow  $m: X \rightarrow A$  is monic if and only if it is  $P$ -injective.*

*Proof.* One direction is trivial. If  $m$  is monic the square  $\langle \text{id}_A, \text{id}_A \rangle m = (m \times m) \langle \text{id}_X, \text{id}_X \rangle$  is a pullback. Hence 4.6 applies and

$$\delta_X = \mathcal{A}_{\langle \text{id}_X, \text{id}_X \rangle}(\top_X) = \mathcal{A}_{\langle \text{id}_X, \text{id}_X \rangle}(P_m(\top_A)) = P_{m \times m} \mathcal{A}_{\langle \text{id}_A, \text{id}_A \rangle}(\top_A) = P_{m \times m}(\delta_A)$$

$\square$

**4.9 Proposition.** *Suppose  $P$  is an existential  $m$ -variational doctrine on  $\mathcal{C}$ .  $P$  satisfies (RUC) if and only if every arrow which is both  $P$ -injective and  $P$ -surjective is an isomorphism.*

*Proof.* We use the notation in 2.16. Suppose (RUC) holds and take a  $P$ -injective and  $P$ -surjective arrow  $f: A \rightarrow B$ . The formula  $b: B, a: A \mid b =_B f(a)$  is entire and single-valued. By (RUC) there is  $k: B \rightarrow A$  which is the inverse of  $f$ . Conversely take  $F$  in  $P(A \times B)$  and consider  $\{\!\{F\}\!\}: X \rightarrow A \times B$ . If  $F$  is total and single-valued then  $\text{pr}_1 \{\!\{F\}\!\}: X \rightarrow A$  is  $P$ -injective and  $P$ -surjective so it has an inverse  $h$ . The arrow  $\text{pr}_1 h: A \rightarrow B$  is the desired arrow.  $\square$

Primary doctrine with full strong comprehension form the category **CED**, the 2-full sub category of **ED** whose arrows are those arrow  $(F, b): P \rightarrow R$  in **ED** that preserve comprehension, *i.e.* for every  $A$  and  $\alpha$  in  $P(A)$  the arrow  $F\{\alpha\}$  is isomorphic to  $\{b_A\alpha\}$ .

**4.10 Remark.** We recall from [MR13b] [MR13a] that there is a left biadjoint to the forgetful 2-functor from **ED** to **CED** which associates to an elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  the elementary doctrine with full strong comprehensions the doctrine  $P_c: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  whose description is as follows. Objects of  $\mathcal{C}_c$  are pairs  $(A, \alpha)$  where  $\alpha$  is in  $P(A)$  and an arrow  $f: (A, \alpha) \rightarrow (B, \beta)$  is an arrow  $f: A \rightarrow B$  in  $\mathcal{C}$  such that  $\alpha \leq P_f(\beta)$ .  $P_c$  maps each  $(A, \alpha)$  to  $P_c(A, \alpha) = \{\phi \in P(A) \mid \phi \leq \alpha\}$  and each  $f: (A, \alpha) \rightarrow (B, \beta)$  to the function  $P_c(f): P_c(B, \beta) \rightarrow P_c(A, \alpha)$  determined by the assignment  $\psi \mapsto P_f(\psi) \wedge \alpha$ . For  $\phi$  in  $P_c(A, \alpha)$  it is  $\{\phi\} = \text{id}_A: (A, \phi) \rightarrow (A, \alpha)$ .

Doctrines with comprehensive diagonals and full weak comprehension form the class of doctrine which we will mainly concerned with throughout this paper. In [MPR17], taking the terminology from [Gra00], we called them *variational*. While we called *m-variational doctrines* those variational doctrines in which comprehension is strong. We aim at giving a characterisation of variational doctrines in 4.14. To do this we first need some instrumental definitions and propositions.

Existential variational doctrines will be the main mathematical tool that we will employ throughout the whole paper. They form the category **EV** which is the full subcategory of **EExD** on those doctrines that are also existential. We denote by **EMV** the subcategory of **EV** on m-variational doctrines and on those morphisms of doctrine that preserves strong comprehension. The adjoint situation described in 4.10 and in 2.4 compose to give the following

$$\mathbf{EED} \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{\quad} \end{array} \mathbf{EMV} \quad (2)$$

Consider the following two diagrams in **PD**.

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} & \xrightarrow{P} & \mathbf{ISL} \\ \text{Id}_{\mathcal{C}} \downarrow & j \downarrow \cdot & \downarrow \\ \mathcal{C}^{\text{op}} & \xrightarrow{P} & \mathbf{ISL} \end{array} \quad \begin{array}{ccc} \mathcal{C}^{\text{op}} & \xrightarrow{P} & \mathbf{ISL} \\ \text{id}_{\mathcal{C}}^{\text{op}} \downarrow & l \cdot \left( \dashv \right) \cdot r & \downarrow \\ \mathcal{C}^{\text{op}} & \xrightarrow{R} & \mathbf{ISL} \end{array}$$

where the adjointness symbols in the diagram on the right means that for every  $A$  in  $\mathcal{C}$  and every  $\alpha$  in  $P(A)$  and every  $\beta$  in  $R(A)$  the inequalities  $lr\alpha \leq \alpha$  and  $\beta \leq rl\beta$  both hold. We say that that  $P$  and  $R$  form an *adjoint-retraction pair* and we write  $P \triangleleft R$  if they fit in a diagram as the one on the right and moreover  $lr = \text{id}_P$ . An endomorphism of doctrine of the form  $(\text{id}_{\mathcal{C}}, j): P \rightarrow P$  as the one in the diagram on the left is called *topology on  $P$*  if  $j$  is extensive and idempotent, *i.e.* for every  $A$  in  $\mathcal{C}$  and every  $\alpha$  in  $P(A)$  it holds  $\alpha \leq j_A\alpha$  and  $j_A j_A\alpha = j_A\alpha$ . An element  $\alpha$  in  $P(A)$  is  *$j$ -closed* if  $\alpha = j_A(\alpha)$ . Every

topology  $j$  on  $P$  determines a doctrine  $P_j: \mathcal{C}^{op} \longrightarrow \mathbf{ISL}$  where  $P_j(A)$  is the subinfsemilattice of  $P(A)$  on those  $\alpha$  such that  $j_A \alpha = \alpha$  and where  $p_j(f) = P(f)$ . The doctrine  $P_j$  is called **doctrine of  $j$ -closed element of  $P$** .

The following is proved in [Pas16], but see also [MPR19].

**4.11 Proposition.** *If  $P \triangleleft R$  then there is  $L: \mathcal{Q}_R \rightarrow \mathcal{Q}_P$  and  $R: \mathcal{Q}_P \rightarrow \mathcal{Q}_R$  with  $L$  left adjoint to  $R$  and  $R$  is full and faithful.*

*Proof.*  $L$  maps  $[f]: (A, \rho) \rightarrow (B, \sigma)$  to  $[f]: (A, l_{A \times A}(\rho)) \rightarrow (B, l_{B \times B}(\sigma))$ . The functor  $R$  is built analogously.  $\square$

**4.12 Proposition.** *Suppose  $P: \mathcal{C}^{op} \longrightarrow \mathbf{ISL}$  and  $R: \mathcal{C}^{op} \longrightarrow \mathbf{ISL}$  are functors based on  $\mathcal{C}$ . Then  $P \triangleleft R$  if and only if  $P$  is isomorphic to  $R_j$  for some topology  $j$  on  $R$ .*

*Proof.* Suppose  $P \triangleleft R$  as in the diagram above. The composition  $rl$  is clearly extensive and also idempotent as  $lr = \text{id}_P$ . The morphism  $(\text{id}_C, r): P \rightarrow R_{rl}$  is then the inverse of  $(\text{id}_C, l): R_{rl} \rightarrow P$ , whence  $j = rl$  is the desired topology on  $R$ . Conversely let  $j$  be a topology on  $R$ . There is an morphism  $(\text{id}_C, \iota): R_j \rightarrow R$  where  $\iota$  is a family of inclusions and a morphism  $(\text{id}_C, j): R \rightarrow R_j$ . It is immediate to see that  $R_j \triangleleft R$ .  $\square$

**4.13 Proposition.** *Suppose  $P: \mathcal{C}^{op} \longrightarrow \mathbf{ISL}$  is a primary doctrine and  $j$  is a topology on  $P$ . We have the following:*

1. *If  $P$  is elementary, then so is  $P_j$ .*
2. *If  $P$  is existential, then so is  $P_j$ .*
3. *If  $P$  is disjunctive, then so is  $P_j$ .*
4. *If  $P$  is implicative, then so is  $P_j$ .*
5. *If  $P$  is universal, then so is  $P_j$ .*
6. *If  $P$  has a weak predicate classifier, then so does  $P_j$ .*

*Proof.* Standard argument: for  $f: A \rightarrow B$  and  $\alpha, \beta$  in  $P_j(A)$  note that  $\forall_f(\alpha)$  and  $\alpha \rightarrow \beta$  determines the universal and the implicative structure in  $P_j$  (if they already exists in  $P$ ) while for the left adjoints and the disjunctive structure one takes the closure of the corresponding ones of  $P$  so  $j_B(\exists_f(\alpha))$  and  $j_A(\alpha) \vee j_A(\beta)$ . The weak power object of  $A$  is  $\mathbb{P}A$  with membership predicate  $j_{A \times \mathbb{P}A}(\in_A)$ .  $\square$

**4.14 Proposition.** *Suppose  $P: \mathcal{C}^{op} \longrightarrow \mathbf{ISL}$  is elementary existential and  $\mathcal{C}$  has weak pullbacks. The following are equivalent*

1.  *$P$  is isomorphic to  $\Psi_{C_j}$  for some topology  $j$  over  $\Psi_C$*
2.  *$P \triangleleft \Psi_C$*
3.  *$P$  has full weak comprehensions.*

*Proof.* The equivalence between items 1 and 2 is a special case of 4.12, while  $3 \Rightarrow 2$  is proposition 2.3 of [MPR19] It remains  $2 \Rightarrow 3$ . Take  $\alpha$  in  $P(A)$  and let  $\{\alpha\}: X \rightarrow A$  be any representative of  $\mathbf{r}_A(\alpha)$ . We first need prove that  $\top_X \leq P_{\{\alpha\}}(\alpha)$ . Observe that

$$\alpha = \mathbf{r}_A(\alpha) = \mathbf{l}(\Sigma_{\mathbf{r}_A(\alpha)} \text{id}_X) = \exists_{\mathbf{r}_A(\alpha)} \mathbf{l}(\text{id}_X) = \exists_{\mathbf{r}_A(\alpha)} \mathbf{l}(\mathbf{r}(\top_X)) = \exists_{\mathbf{r}(\alpha)} \top_X$$

Hence  $\top_X \leq P_{\mathbf{r}_A(\alpha)}(\exists_{\mathbf{r}_A(\alpha)} \top_X) = P_{\mathbf{r}_A(\alpha)}(\alpha) = P_{\{\alpha\}}(\alpha)$ . Take  $f: Y \rightarrow A$  with  $\top_Y \leq P_f(\alpha)$ . Then  $\text{id}_Y = \mathbf{r}_A(\top_Y) \leq \mathbf{r}_A P_f(\alpha) = f^* \mathbf{r}_A(\alpha) = f^* \{\alpha\}$  from which we conclude that  $\{\alpha\}$  is a full weak comprehension of  $\alpha$ .  $\square$

Since  $\text{id}_X$  factors through the weak pullback of  $\{\alpha\}$  along itself, it is

$$\mathbf{r}_X(\top_X) = [\text{id}_X] \leq \{\alpha\}^* \{\alpha\} = \{\alpha\}^* \mathbf{r}_A(\alpha) = \mathbf{r}_X P_{\{\alpha\}}(\alpha)$$

The equality  $\mathbf{r} \mathbf{l} = \text{id}_P$  implies that components of  $\mathbf{r}$  reflect the order, so  $\top_X \leq P_{\{\alpha\}}(\alpha)$ . Then  $[\text{id}_Y] = \mathbf{l}_Y(\top_Y) \leq \mathbf{l}_Y P_f(\alpha) = f^* \{\alpha\}$ . This means that  $\text{id}_Y$  factors through the weak pullback of  $\{\alpha\}$  along  $f$ , so  $f$  factors through  $\{\alpha\}$ . To show that  $P$  has comprehensive diagonals take  $h, k: X \rightarrow A$  and suppose  $\top_X \leq P_{\langle h, k \rangle}(\delta_A)$ . Using again  $\mathbf{r}$  (and the fact that it is a morphism of elementary doctrines) it holds

$$[\text{id}_X] = \mathbf{r}_X(\top_X) \leq \mathbf{r}_X P_{\langle h, k \rangle}(\delta_A) = \langle h, k \rangle^* \mathbf{r}_A(\delta_A) = \langle h, k \rangle^* [\langle \text{id}_A, \text{id}_A \rangle]$$

so  $h = k$ . The second part of the theorem is analogous once one observes that a comprehension arrow is strong if and only if it is monic.

The following proposition is stated without a proof as its proof is perfectly analogous to the one of 4.14.

**4.15 Proposition.** *Suppose  $P: C^{op} \rightarrow \mathbf{ISL}$  is elementary existential and  $C$  has pullbacks. Then  $P$  is isomorphic to  $\text{Sub}_{C_j}$  for some topology  $j$  over  $\text{Sub}_C$  if and only if  $P \triangleleft \text{Sub}_C$  if and only if  $P$  has full strong comprehension.*

Recall from [BW84] that a topology on a category  $C$  with finite limits is a topology on  $\text{Sub}_C$ . Given such a topology an object is  $j$ -separated if  $j_{A \times A} \delta_A = \delta_A$ .

**4.16 Proposition.** *Suppose  $P: C^{op} \rightarrow \mathbf{ISL}$  is elementary existential and  $C$  has weak pullbacks. The following are equivalent*

1.  $P$  is isomorphic to  $\Psi_{C_j}$  for some topology  $j$  over  $\Psi_C$  whose objects are all  $j$ -separated.
2.  $P \triangleleft \Psi_C$  and the right adjoint is an elementary morphism of doctrines.
3.  $P$  has full weak comprehensions and comprehensive diagonals.

The elementary quotient completion presented in section 5 very well behaves with respect to comprehension, in particular we have the following.

**4.17 Proposition.**  *$P$  is variational if and only if  $\widehat{P}$  is  $m$ -variational.*

*Proof.* The necessary condition is immediate. Suppose  $P$  has weak comprehensions. For an element  $\alpha$  in  $\widehat{P}(A, \rho) \subseteq P(A)$  its comprehension is

$$[\{\alpha\}]: (X, P_{\{\alpha\} \times \{\alpha\}}(\rho)) \rightarrow (A, \rho)$$

where  $\{\alpha\}: X \rightarrow A$  is the weak comprehension of  $\alpha$  in  $P$ . Since  $\widehat{P}$  has comprehensive diagonals, the arrow  $[\{\alpha\}]$  is monic (4.8).  $\square$

**4.18 Proposition.** *If  $P$  is variational, then  $Q_P$  has finite limits.*

*Proof.* By 4.17  $\widehat{P}$  is m-variational, then apply 4.4.  $\square$

Every topology  $j$  on a primary doctrine  $P$  determines a topology  $\widehat{j}$  on the elementary quotient completion  $\widehat{P}$ . The topology  $\widehat{j}$  is simply the restriction of  $j$  to the poset of descent data, *i.e.* for  $\rho$  a  $P$ -equivalence relation over  $A$  and  $\alpha$  in  $\mathcal{D}es_\rho$  it is  $\widehat{j}_{(A, \rho)}(\alpha) = j_A(\alpha)$  (see also [Men01]). Indeed (using notation as in 2.16)

$$a: A, a': A \mid j\alpha(a) \wedge \rho(a, a') \vdash j\alpha(a) \wedge j\rho(a, a') \vdash j(\alpha(a) \wedge \rho(a, a')) \vdash j\alpha(a')$$

After proposition 4.14 we know that every existential variational doctrine  $P$  on a category  $\mathcal{C}$  with weak pullbacks generates a topology on  $\Psi_{\mathcal{C}}$ . We denote this topology by  $j_P$  and call it **canonical topology** on  $P$ .

**4.19 Proposition.** *Suppose  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is an existential variational doctrine. Then  $Q_P$  is the category of  $\widehat{j}_P$ -separated objects for the topology  $\widehat{j}_P$  induced over  $\mathcal{C}_{ex/lex}$ .*

*Proof.* By 4.11 there is a full and faithful  $R: Q_P \rightarrow Q_{\Psi_{\mathcal{C}}}$  which is right adjoint to  $L: Q_{\Psi_{\mathcal{C}}} \rightarrow Q_P$ . Take a  $\Psi_{\mathcal{C}}$ -equivalence relation  $[\rho: X \rightarrow A \times A]$  over  $A$ . The object  $(A, [\rho])$  in  $Q_{\Psi_{\mathcal{C}}}$  is in  $Q_P$  if and only if  $(A, [\rho]) \simeq RL(A, [\rho])$ . From the construction of  $L$  and  $R$  as in 4.11 this happens if and only if

$$(A, \rho) \simeq (A, j_{P_{A \times A}}[\rho]) = (A, \widehat{j}_{P_{(A, \rho) \times (A, \rho)}}(\delta_{(A, \rho)}))$$

Note that  $\widehat{j}_P$  is a topology on  $\widehat{\Psi}_{\mathcal{C}}$  which is  $\text{Sub}_{\mathcal{C}_{ex/lex}}$ , whence the claim.  $\square$

A first order doctrine  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is **boolean** if for every  $A$  and  $\alpha$  in  $P(A)$  it holds  $\top_A = \alpha \vee \neg\alpha$  where  $\neg\alpha$  is short for  $\alpha \rightarrow \perp$ .

**4.20 Proposition.** *A first order variational doctrine  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  on a base  $\mathcal{C}$  with weak pullbacks and an initial object  $0$  such that  $\{\perp\} = un$  where  $un: 0 \rightarrow 1$  in  $\mathcal{C}$ , is boolean if and only if  $P$  is isomorphic to  $\Psi_{\mathcal{C}^{\neg\neg}}$ , the doctrine of  $\neg\neg$ -closed elements of  $\Psi_{\mathcal{C}}$ . A first order m-variational doctrine  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  on a base  $\mathcal{C}$  is boolean if and only if  $P$  is isomorphic to  $\text{Sub}_{\mathcal{C}^{\neg\neg}}$ , the doctrine of  $\neg\neg$ -closed elements of  $\text{Sub}_{\mathcal{C}}$ .*

*Proof.* We show the non-trivial direction. Suppose  $P$  is boolean. By 4.14 there is a topology  $j$  on  $\Psi_C$  such that  $P$  is isomorphic to  $\Psi_{Cj}$  and for  $[f: X \rightarrow A]$  in  $\Psi_C(A)$  it is  $j_A[f] = [\{\mathcal{A}_f \top_X\}]$ . It is

$$[\{\mathcal{A}_f \top_X\}] = [\{\neg V_f \neg \top_X\}] = [\neg \Pi_f \neg \{\top_X\}] = \neg \neg [\Sigma_f \{\top_X\}] = \neg \neg [\Sigma_f(\text{id}_X)]$$

and hence the claim as  $[\Sigma_f(\text{id}_X)] = [f]$ .  $\square$

Suppose  $C$  is a category with weak pullbacks. A **right weak factorization system** is a pair  $(\mathcal{E}, \mathcal{R})$  of classes of arrows of  $C$  such that

- (i) for every  $f$  in  $C$  there is  $e$  in  $\mathcal{E}$  and  $r$  in  $\mathcal{R}$  with  $f = re$
- (ii) for every commutative square  $rf = ge$  with  $e$  in  $\mathcal{E}$  and  $r$  in  $\mathcal{R}$  there is  $k$  with  $rk = g$ .

A right weak factorization system is stable if any weak pullback of an arrow of  $\mathcal{E}$  is in  $\mathcal{E}$ . A proper actorisation system  $(\mathcal{E}, \mathcal{R})$  in the sense of [FK72] is a right weak factorization system such that every arrow in  $\mathcal{R}$  is monic and every arrow in  $\mathcal{E}$  is epic. Every proper stable factorisation system  $(\mathcal{E}, \mathcal{R})$  on a category  $C$  with pullbacks gives rise to an existential m-variational doctrine  $\mathcal{R}_C: C^{\text{op}} \rightarrow \mathbf{ISL}$  where  $\mathcal{R}_C$  is the sub-infsemilattice of  $\Psi_C(A)$  on those arrows represented by arrows in  $\mathcal{R}$ , which are monic and hence also in  $\text{Sub}_C(A)$  [HJ03]. It is proved in *loc.cit.* that the category of existential elementary doctrines with full strong comprehension is equivalent to the category of proper stable factorisation systems. More precisely,

**4.21 Proposition.** *In every m-variational doctrine comprehension arrows and P-surjective arrows form a factorization system. Moreover, the category of existential m-variational doctrines is equivalent to the category of proper stable factorisation systems where diagonals are in the right class.*

Analogously the category **EV** of existential variational doctrines is equivalent to the category **RWFS** of those stable right weak factorization systems where diagonals are in  $\mathcal{R}$ .

**4.22 Proposition.** *Let  $C$  be a category finite products and a stable right weak factorization system  $(\mathcal{E}, \mathcal{R})$ . Suppose that diagonals are in  $\mathcal{R}$ . Then  $\mathcal{R}_C$  is  $\Psi_C$  if and only if  $\mathcal{R}_C$  satisfies (RC). If moreover  $C$  has finite limits and  $(\mathcal{E}, \mathcal{R})$  is proper, then  $\mathcal{R}_C$  is  $\text{Sub}_C$  if and only if  $\mathcal{R}_C$  satisfies (RUC).*

*Proof.* Immediate after characterisation 4.12.  $\square$

Proposition 4.22 shows that  $\mathbf{EV}^{RC}$ , the full subcategory of **EV** on those doctrines satisfying (RC), is formed only by doctrines of the form  $\Psi_C$ . These doctrines are completely determined by the base so  $\mathbf{EV}^{RC}$  is equivalent to the category  $\mathbf{Car}_{\text{wpb}}$  of categories with finite products and weak pullbacks.

Finally we call a right weak factorization system **exact** if  $\mathcal{R}_C$  has descent effective quotients. These form the category **QRWFS**. Note that a category

with finite limits  $\mathcal{C}$  is exact if and only if  $\text{Sub}_{\mathcal{C}}$  is an existential m-variational doctrine with descent effective quotient. Exact categories form the category **Xct**. So diagram 1 can be refined to the following.

$$\begin{array}{ccccc}
 & & \mathbf{EExD}^{RC} & \xrightarrow{\quad} & \mathbf{EExD} \\
 & \nearrow & \uparrow & & \uparrow \\
 \mathbf{Car}_{\text{wpb}} & \xrightarrow{\quad} & \mathbf{RWFS} & & \mathbf{EExD} \\
 \uparrow & & \uparrow & & \uparrow \\
 & & \mathbf{QExD}^{RUC} & \xrightarrow{\quad} & \mathbf{QExD} \\
 & \nearrow & \uparrow & & \uparrow \\
 \mathbf{Xct} & \xrightarrow{\quad} & \mathbf{QRWFS} & & \mathbf{QExD}
 \end{array} \tag{3}$$

where, as for diagram (1) the left adjoints are provided by the elementary quotient completion. In particular, the elementary quotient completion on  $\mathbf{Car}_{\text{wpb}}$  is the elementary quotient completion of the doctrine of weak subobjects, hence the ex/wlex completion.

It is worth noting that if we start with an elementary doctrine  $P$  without comprehensive diagonals and just with weak full comprehensions we can get anyway a m-variational doctrine closed under effective quotients. Moreover, adding comprehensive diagonals to  $P$  before completing it with quotients does produce the same doctrine as that obtained by completing  $P$  itself.

**4.23 Theorem.** *Let  $P$  an elementary doctrine with weak full comprehensions. Then  $\widehat{P}$  is a m-variational doctrine with stable effective quotients. Moreover, the doctrine  $\widehat{P}_x$  is equivalent to  $\widehat{P}$ .*

## 5 Decomposition of elementary quotient completions

In this section we recall from [MR13a] that the construction of the elementary quotient completion of a elementary introduced in [MR13b] is obtained by applying to  $P$  two other free constructions: first we apply the *intensional quotient completion* of an elementary doctrine producing just effective descent quotients, which we recall below, and then the *extensional collapse construction* of an elementary doctrine which freely add **comprehensive diagonals** applied to it defined in the previous sections.

**5.1 Definition** (intensional quotient completion). Given a elementary  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  we call  $Q_{P_i}$  the category whose objects are pairs  $(A, \rho)$  in which  $A$  is in  $\mathcal{C}$  and  $\rho$  is an equivalence relation over  $A$ . An arrow  $f: (A, \rho) \rightarrow (B, \sigma)$  is an arrow  $f: A \rightarrow B$  in  $\mathcal{C}$  such that  $\rho \leq P_{f \times f}(\sigma)$ .

Then we define an elementary doctrine called *intensional quotient completion* of  $P$

$$P_{iq}(A, \rho) = \{\phi \in P(A) \mid P_{\text{pr}_1}(\phi) \wedge \rho \leq P_{\text{pr}_2}(\phi)\}$$

$$(P_{i_q})_{[f]} = P_f$$

where  $\text{pr}_1$  and  $\text{pr}_2$  are the first and the second projection form  $A \times A$ . It is proved in [MR13a] that the assignment on arrows does not depend on the choice of representatives.

**5.2 Theorem.** *There is a left biadjoint to the forgetful 2-functor from the full 2-category of elementary doctrines with stable effective descent quotients into the 2-category **ED** of elementary doctrines which associates the doctrine  $P_{i_q}: Q_{P_i}^{\text{op}} \longrightarrow \mathbf{ISL}$  to an elementary doctrine  $P: C^{\text{op}} \longrightarrow \mathbf{ISL}$ .*

As shown in [MR13a] the intensional quotient completion of a doctrine has stable effective descent quotients: if  $\sigma$  is an equivalence relation over  $(A, \rho)$ , then its quotient is given by

$$[\text{id}_A]: (A, \rho) \longrightarrow (A, \sigma)$$

Observe that in the doctrine  $Q_{P_i}$ , the equality predicate over  $(A, \rho)$  is  $\rho$  itself [MR13a], *i.e.*

$$\delta_{(A, \rho)} = \rho$$

Note that for any elementary doctrine  $Q: C^{\text{op}} \longrightarrow \mathbf{ISL}$  with effective quotients, the doctrine  $Q_x$  has only a weak form of quotients. But when  $Q = Q_{P_i}$  of an elementary doctrine  $P$ , in [MR13a] it is shown that:

**5.3 Proposition.** *Let  $P: C^{\text{op}} \longrightarrow \mathbf{ISL}$  be an elementary doctrine. The morphism  $(K, k): Q_{P_i} \longrightarrow Q_{P_{i_x}}$  preserves quotients and therefore  $Q_{P_{i_x}}$  has effective descent quotients of  $Q_{P_{i_x}}$ -equivalence relations and coincides with the elementary quotient completion of  $P$ .*

## 6 A characterisation of elementary quotient completions

In this section we give a characterisation of those elementary doctrines with effective descent quotients that arise as elementary quotient completions.

To this purpose recall that the ex/lex completion of a category  $\mathcal{C}$  with finite products and weak pullbacks is the base of the elementary quotient completion of the doctrine of variations of  $\mathcal{C}$ .

Our characterisation arises as a generalisation to the framework of doctrines of the well known fact that an exact category with enough regular projectives is equivalent to the ex/lex completion of its full subcategory on projective objects.

Suppose  $\mathcal{E}$  is a class of morphisms of a category  $\mathcal{C}$ . An object  $X$  of  $\mathcal{C}$  is said  **$\mathcal{E}$ -projective** if for every diagram of the form

$$\begin{array}{ccc} & & Y \\ & & \downarrow s \\ X & \xrightarrow{f} & A \end{array}$$

where  $s$  is in  $\mathcal{E}$ , there is an arrow  $k: X \longrightarrow Y$  with  $sk = f$ .

$\mathcal{C}$  has *enough  $\mathcal{E}$ -projectives* if for every  $A$  in  $\mathcal{C}$  there is a  $\mathcal{E}$ -projective object  $X$  and an arrow of  $\mathcal{E}$  from  $X$  to  $A$ .

**6.1 Lemma.** *Denote by  $\mathcal{P}$  the full subcategory of  $\mathcal{C}$  on  $\mathcal{E}$ -projective objects. If  $\mathcal{P}'$  is a full subcategory of  $\mathcal{P}$  closed under binary products and such that every object of  $\mathcal{P}$  is covered by one in  $\mathcal{P}'$ , then  $\mathcal{P}$  is closed under binary products.*

*Proof.* Suppose  $A$  and  $B$  are in  $\mathcal{P}$  and consider the diagram

$$\begin{array}{ccc} & & X \\ & & \downarrow e \\ A \times B & \xrightarrow{f} & Y \end{array}$$

where  $e$  is in  $\mathcal{E}$ . Let  $q_A: A' \rightarrow A$  and  $q_B: B' \rightarrow B$  be arrow of  $\mathcal{E}$  that cover  $A$  and  $B$ , *i.e.*  $A'$  and  $B'$  are in  $\mathcal{P}'$ . Since both  $A$  and  $B$  are in  $\mathcal{P}$  each cover has a section  $s_A$  and  $s_B$ . Since  $A' \times B'$  is in  $\mathcal{P}'$  it is in particular  $\mathcal{E}$ -projective, then there is  $\bar{f}: A' \times B' \rightarrow X$  with  $e\bar{f} = f(q_A \times q_B)$ . Hence  $\bar{f}(s_A \times s_B): A \times B \rightarrow X$  is such that  $e\bar{f}(s_A \times s_B) = f(q_A \times q_B)(s_A \times s_B) = f$  proving that  $A \times B$  is  $\mathcal{E}$ -projective.  $\square$

Suppose  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is an elementary doctrine Denote by  $\mathbf{q}_P$  the class of quotient arrows of  $P$ .

Note that, if  $q: X \rightarrow X/\rho$  and  $q': X' \rightarrow X'/\rho'$  are quotient arrows, and if  $X$  is  $\mathbf{q}_P$ -projective, then for every arrow  $f: X/\rho \rightarrow X'/\rho'$  there is an arrow  $g: X \rightarrow X'$  with  $\top_X = P_{\langle f, q, q' \rangle}(\delta_{X'/\rho'})$ . Obviously if the doctrine has comprehensive diagonals it is  $f q = q' g$ .

**6.2 Proposition.** *Suppose  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is an elementary doctrine with comprehensive diagonals and effective descent quotients. The following are equivalent*

- i)  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is of the form  $\widehat{P}_0: Q_{P_0}^{op} \rightarrow \mathbf{InfSL}$  for some elementary doctrine  $P_0: \mathcal{C}_0^{op} \rightarrow \mathbf{ISL}$  with comprehensive diagonals
- ii)  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  has enough  $\mathbf{q}_P$ -projectives and these are closed under binary products.

*Proof.*  $i \Rightarrow ii$ ) All quotient arrows in  $Q_{P_0}$  are of the form  $[\text{id}_A]: (A, \rho) \rightarrow (A, \sigma)$ , thus objects of the form  $(A, \delta_A)$  are  $\mathbf{q}_P$ -projective and they determine a full subcategory of  $Q_{P_0}$  which is closed under products. Hence  $Q_{P_0}$  has enough  $\mathbf{q}_P$ -projective and these are closed under binary products by 6.1.

$ii \Rightarrow i$ ) Denote by  $\mathcal{C}_0$  the full subcategory of  $\mathcal{C}$  on its  $\mathbf{q}_P$ -projectives and by  $P_0$  the restriction of  $P$  to  $\mathcal{C}_0$  (*i.e.* the change of base of  $P$  along the inclusion of  $\mathcal{C}_0$  into  $\mathcal{C}$ ). Since  $\mathcal{C}_0$  is closed under products  $P_0$  is an elementary doctrine with comprehensive diagonals. We need prove that  $Q_{P_0}$  is equivalent to  $\mathcal{C}$ .

Consider  $[f]: (A, \rho) \rightarrow (B, \sigma)$  in  $\mathcal{Q}_{P_0}$ . Every representative of  $[f]$  determines a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{q} & A/\rho \\ f \downarrow & & \downarrow \bar{f} \\ B & \xrightarrow{e} & B/\sigma \end{array}$$

where  $\bar{f}$  is the map determined by the universal property of quotients. The diagram above extends a functor from  $\mathcal{Q}_{P_0} \rightarrow \mathcal{C}$ . Which is faithful by effectiveness of quotients. Since  $A$  and  $B$  are projective every arrow  $A/\rho \rightarrow B/\sigma$  determines an arrow  $A \rightarrow B$ , then the functor is also full. Essential surjectivity is a straightforward consequence of the hypothesis that  $\mathcal{C}$  has enough  $\mathbf{q}_P$ -projectives. Since quotients are of effective descent, for every  $(A, \rho)$  in  $\mathcal{Q}_{P_0}$  the poset  $\widehat{P_0}(A, \rho)$  is isomorphic to  $P(A/\rho)$ : this completes the proof.  $\square$

In a similar way we obtain a characterization of *intensional quotient completions*:

**6.3 Theorem.** *Suppose  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is an elementary doctrine with effective descent quotients. The following are equivalent*

- i)  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is the intensional elementary quotient of some elementary doctrine  $P_0: \mathcal{C}_0^{op} \rightarrow \mathbf{ISL}$ .
- ii)  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  has enough  $P$ -projectives and these are closed under binary products.

It is worth noting that theorem 6.2 is a generalization of Carboni-Vitale's characterization of exact completions after recalling that the exact completion of a category with binary products and weak pullbacks is an instance of the elementary quotient completion (see 3.7-(d)).

To prove this we need some lemmas:

**6.4 Lemma.** *In a category  $\mathcal{C}$  with finite products and weak pullbacks, an object is projective with respect to the doctrine of variations  $\Psi_{\mathcal{C}}$  of  $\mathcal{C}$  if and only if it is a regular projective.*

*Proof.* The notion of  $\Psi_{\mathcal{C}}$ -effective quotient coincide with that of categorical effective quotient.  $\square$

**6.5 Corollary.** *Let  $\mathcal{C}$  be an exact category. The following are equivalent:*

- i)  $\mathcal{C}$  is an ex/lex completion
- ii)  $\mathcal{C}$  has enough projectives closed under finite limits.

*When one of the conditions holds, then  $\mathcal{C}$  is the ex/lex completion of its full subcategory of projectives.*

*Proof.* Let  $\mathcal{C}_0$  the full subcategory of  $\mathcal{C}$  of regular projectives and let  $P \equiv \text{Sub}_{\mathcal{C}}$  be the subobject doctrine of  $\mathcal{C}$  which by lemma 3.6 is effective descent and it is also elementary since projectives are closed under equalizers. Therefore by ??  $\mathcal{C}$  is equivalent to the base of the elementary quotient completion of  $P$  restricted to  $\mathcal{C}_0$  since the subobject doctrine is m-variational because projectives are closed under finite limits. But  $P \equiv \text{Sub}_{\mathcal{C}}$  restricted to  $\mathcal{C}_0$  is equivalent to the doctrines of variations  $\Psi_{\mathcal{C}_0}: \mathcal{C}_0^{\text{op}} \longrightarrow \mathbf{ISL}$  because the doctrine of variations can be fully and faithfully embedded in  $P$  via the image factorization of any map  $f: A \longrightarrow B$  in  $\mathcal{C}_0$  with a regular epimorphism and the projectivity of their domain  $A$ . Conversely any subobject  $i: C \longrightarrow B$  in  $\mathcal{C}$  over a projective  $B$ , through a projective cover  $p: A' \longrightarrow C$  of  $C$  gives rise to map  $ip: A' \longrightarrow B$  which is in  $\mathcal{C}_0$ . Therefore  $\mathcal{C}$  is equivalent to the base of the elementary quotient completion of the doctrines of variations  $\Psi_{\mathcal{C}_0}: \mathcal{C}_0^{\text{op}} \longrightarrow \mathbf{ISL}$  which coincides with the exact completion of  $\mathcal{C}_0$ .  $\square$

## 7 Structural properties inherited and reflected by the elementary quotient completion

In this section and in the next ones we generalize to elementary quotient completions other well known facts concerning the ex/lex completion.

It is well known that the ex/lex completion brings weak structures of  $\mathcal{C}$  to strong structures in  $\mathcal{C}_{\text{ex/lex}}$  and this holds for elementary quotient completions in an analogous way. In this section we shall focus on local cartesian closure, disjoint stable coproducts and predicate classifier from a given variational elementary doctrine  $P$ . First note the following.

**7.1 Proposition.**  *$P$  is first order doctrine if and only if  $\widehat{P}$  is a first order doctrine.*

*Proof.* Immediate, see also [Pas16].  $\square$

A  $J$ -diagrams in  $\mathcal{C}$  is a functor of the form  $J \longrightarrow \mathcal{C}$ . We say that  $\mathcal{C}$  has  $J$ -indexed limits if every  $J$ -diagram has a limits and that  $\mathcal{C}$  has  $J$ -weak indexed limits if every  $J$ -diagram has the existence property of a limit but not the uniqueness condition. Accordingly we say that  $\mathcal{C}$  has  $J$ -indexed colimits if every  $J$ -diagram has a colimits.

Recall that for a variational doctrine  $P$  on  $\mathcal{C}$  the  $\nabla_P: \mathcal{C} \longrightarrow \mathcal{Q}_P$  ( *i.e.* the functor that maps  $f: A \longrightarrow b$  to  $[f]: (A, \delta_A) \longrightarrow (B, \delta_B)$ ) is full and faithful.

**7.2 Proposition.** *If  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  is a variational doctrine, then for every  $J \longrightarrow \mathcal{C}$  it holds*

1. if  $\mathcal{Q}_P$  has  $J$ -indexed limits, then  $\mathcal{C}$  has  $J$ -indexed weak limits;
2. if  $\mathcal{Q}_P$  has  $J$ -indexed colimits of the form  $(W, \delta_W)$ , then  $\mathcal{C}$  has  $J$ -indexed colimits.

*Proof.* 1. Let  $F$  be a  $J$ -diagram in  $\mathcal{C}$ . If  $(W, \omega)$  is the limit of  $\nabla_P F$  in  $\mathcal{Q}_P$ , then  $W$  is a weak limit of  $F$  in  $\mathcal{C}$ . 2. Consider a diagram  $F: J \rightarrow \mathcal{C}$  and suppose  $(W, \delta_W)$  is the colimit for  $\nabla_P F$ .  $W$  is easily seen to be a weak colimit for  $F$  in  $\mathcal{C}$ . Suppose  $X$  is a cocone and let arrows  $q, p: W \rightarrow X$  be such that they make commute all the relevant triangles. So do  $[q], [p]: (W, \delta_W) \rightarrow (X, \delta_X)$  in  $\mathcal{Q}_P$ . Universality of  $(W, \delta_W)$  ensures that  $[q] = [p]$ , *i.e.*  $\top_W \leq P_{\langle q, p \rangle}(\delta_X)$ , whence  $q = p$  in  $\mathcal{C}$  as diagonals are comprehensive.  $\square$

**7.3 Proposition.** *Suppose  $P$  is a variational doctrine and  $f: X \rightarrow A, g: Y \rightarrow A$  are arrows of  $\mathcal{C}$ . Suppose also that  $\rho$  and  $\sigma$  are  $P$ -equivalence relations over  $X$  and  $Y$  respectively such that  $\rho \leq P_{f \times f}(\delta_A)$  and  $\sigma \leq P_{g \times g}(\delta_A)$ . Consider two arrows  $k: P \rightarrow X$  and  $h: P \rightarrow Y$  and the following two squares*

$$\begin{array}{ccc} S & \xrightarrow{k} & X \\ h \downarrow & & \downarrow f \\ Y & \xrightarrow{g} & A \end{array} \quad \begin{array}{ccc} (S, P_{h \times h}(\sigma) \wedge P_{k \times k}(\rho)) & \xrightarrow{[k]} & (X, \rho) \\ [h] \downarrow & & \downarrow [f] \\ (Y, \sigma) & \xrightarrow{[g]} & (A, \delta_A) \end{array}$$

*If the left square is a weak pullback in  $\mathcal{C}$ , the right square is a pullback in  $\mathcal{Q}_P$ .*

*Proof.* Take any two arrows  $[a]: (C, \gamma) \rightarrow (X, \rho)$  and  $[b]: (C, \gamma) \rightarrow (Y, \sigma)$  with  $[f][a] = [g][b]$ . That is to say  $\top_C = P_{f a \times g b}(\delta_A)$ . Since diagonals are comprehensive we have  $f a = g b$ . Hence there is  $u: C \rightarrow S$  with  $ku = a$  and  $hu = b$ . Since  $\gamma \leq P_{a \times a}(\rho) = P_{u \times u} P_{k \times k}(\rho)$  and  $\gamma \leq P_{b \times b}(\sigma) = P_{u \times u} P_{h \times h}(\sigma)$  one has  $\gamma \leq P_{u \times u}(P_{h \times h}(\sigma) \wedge P_{k \times k}(\rho))$ , showing that  $[u]$  is an arrow in  $\mathcal{Q}_P$ , thus also  $[k][u] = [a]$  and  $[h][u] = [b]$ . If  $[u']$  is such that  $[k][u'] = [a]$  and  $[h][u'] = [b]$ , then also  $[k][u'] = [k][u]$  and  $[h][u'] = [h][u]$ , *i.e.*

$$\top_C \leq P_{u \times u'} P_{k \times k}(\rho) \quad \top_C \leq P_{u \times u'} P_{h \times h}(\sigma)$$

whence  $\top_C \leq P_{u \times u'}(P_{h \times h}(\sigma) \wedge P_{k \times k}(\rho))$  showing that  $[u] = [u']$ .  $\square$

**7.4 Proposition.** *Suppose  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is a variational doctrine and consider two squares of the form*

$$\begin{array}{ccc} S & \xrightarrow{k} & X \\ h \downarrow & & \downarrow f \\ Y & \xrightarrow{g} & A \end{array} \quad \begin{array}{ccc} (S, \theta) & \xrightarrow{[k]} & (X, \delta_X) \\ [h] \downarrow & & \downarrow [f] \\ (Y, \delta_Y) & \xrightarrow{[g]} & (A, \delta_A) \end{array}$$

*If the right square is a pullback in  $\mathcal{Q}_P$ , the left square is a weak pullback in  $\mathcal{C}$*

*Proof.* Suppose  $a: C \rightarrow X$  and  $b: C \rightarrow Y$  are such that  $f a = g b$ . Then  $[a]: (C, \delta_C) \rightarrow (X, \delta_X)$  and  $[b]: (C, \delta_C) \rightarrow (Y, \delta_Y)$  are such that  $[f][a] = [g][b]$ . So there is  $[u]$  with  $[k][u] = [a]$  and  $[h][u] = [b]$ , *i.e.*

$$\top_C = P_{\langle ku, a \rangle}(\delta_X) \quad \top_C = P_{\langle hu, b \rangle}(\delta_Y)$$

Since diagonals are comprehensive,  $u$  is such that  $ku = a$  and  $hu = b$  in  $\mathcal{C}$ .  $\square$

For the rest of the section, for every doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$ , we denote by  $\mathcal{C}_0$  the full subcategory of  $\mathcal{C}$  on  $Q$ -projective objects and by  $P_0$  the change of base of  $P$  along the inclusion of  $\mathcal{C}_0$  into  $\mathcal{C}$ .

### 7.1 Local cartesian closure

Recall that a category  $\mathcal{C}$  with finite products is weakly cartesian closed if for every  $A$  and  $B$  in  $\mathcal{C}$  there is an arrow  $e: A \times W \rightarrow B$  such that for every arrow  $f: A \times C \rightarrow B$  there is  $\bar{f}: C \rightarrow W$  with  $e(\text{id}_A \times \bar{f}) = f$ . This definition has a direct generalization in the context of categories with weak products, provided that one reads  $A \times W$  and  $A \times C$  as weak products and  $\text{id}_A \times \bar{f}$  as the map that derives from the weak universal property of  $A \times W$ .

Here we say that  $\mathcal{C}$  is *weakly cartesian weakly closed* if for every  $X$  and  $Y$  there is  $W$  and an arrow  $e: S \rightarrow Y$ , where  $S$  is a weak product of  $X$  along  $W$ , such that for every  $C$  and every  $k: T \rightarrow Y$ , where  $T$  is a weak product of  $X$  and  $C$ , there is  $\bar{k}: C \rightarrow W$  such that  $e(\text{id}_X \times \bar{k}) = k$ , where  $\text{id}_X \times \bar{k}$  is any of the obvious arrows coming from the weak universal property of  $W$ .

**7.5 Proposition.** *Suppose  $\mathcal{C}$  is a category with finite weak products and weak pullbacks. If  $\mathcal{C}$  is weakly closed the for every  $A$  in  $\mathcal{C}$  and every weak terminal object  $1$  the map  $\Psi_{\mathcal{C}}(!_A): \Psi_{\mathcal{C}}(1) \rightarrow \Psi_{\mathcal{C}}(A)$  as a right adjoint.*

*Proof.* For  $f: X \rightarrow A$  define  $\Pi_{!_A}[f] = [!_{A^X}]$  where  $A^X$  is a weak exponential.  $\square$

If a category  $\mathcal{C}$  has weak pullbacks the definition above can be phrased in each slice. We say that a category  $\mathcal{C}$  with weak pullbacks is *slice-wise weakly cartesian closed*, if for every  $A$  in  $\mathcal{C}$  the slice is weakly cartesian weakly closed.

**7.6 Proposition.** *Suppose  $\mathcal{C}$  is a category with finite products and weak pullbacks. If  $\mathcal{C}$  is slice-wise weakly cartesian closed, then  $\Psi_{\mathcal{C}}$  is universal and implicational.*

*Proof.* For every  $A$  and every projection  $\pi_A: A \times B \rightarrow A$ , the known equivalence of categories between  $\mathcal{C}/(A \times B)$  and  $(\mathcal{C}/A)/\pi_A$  leads to the following commutative diagram

$$\begin{array}{ccc} \Psi_{\mathcal{C}/A}([\text{id}_A]) & \xrightarrow{\Psi_{\mathcal{C}/A}(!_{\pi_A})} & \Psi_{\mathcal{C}/A}(\pi_A) \\ \parallel & & \parallel \\ \Psi_{\mathcal{C}}(A) & \xrightarrow{\Psi_{\mathcal{C}}(\pi_A)} & \Psi_{\mathcal{C}}(A \times B) \end{array}$$


Since  $\mathcal{C}/A$  is weakly cartesian closed, by 7.5 the map  $\Psi_{\mathcal{C}/A}(!_{\pi_A})$  has a right adjoint, so also  $\Psi_{\mathcal{C}}(\pi_A)$  has a right adjoint. The doctrine  $\Psi_{\mathcal{C}}$  is such that for every  $f$  the map  $\Psi_{\mathcal{C}}(f)$  has a left adjoint  $\Sigma_f$  satisfying the Frobenius Reciprocity and the Beck-Chevalley condition on all pullbacks. So for every  $h: Z \rightarrow Y$  and

every  $\alpha$  in  $\Psi_C(X \times Y)$  it is  $\Sigma_{\text{id}_X \times f} \Psi_C(\pi_Z) \Pi_{\pi_Z} \Psi_C(\text{id}_X \times f)(\alpha) \leq \alpha$ . By the Beck-Chevalley on left adjoints this implies  $\Psi_C(\pi_Y) \Sigma_f \Pi_{\pi_Z} \Psi_C(\text{id}_X \times f)(\alpha) \leq \alpha$ , whence  $\Pi_{\pi_Z} \Psi_C(\text{id}_X \times f)(\alpha) \leq \Psi_C(f) \Pi_{\pi_Y}(\alpha)$ , showing that that  $\Psi_C$  is universal. The weak cartesian closure of  $\mathcal{C}/A$  implies that the inf-semilattice  $\Psi_C(A)$  has the Heyting implication. Thus  $\Psi_C$  it is also implicational, indeed it is easy to see that  $\Psi_C(f)(\alpha \Rightarrow \beta) \leq \Psi_C(f)\alpha \Rightarrow \Psi_C(f)\beta$ . Therefore from  $\Psi_C(f)\alpha \wedge (\Psi_C(f)\alpha \Rightarrow \Psi_C(f)\beta) \leq \Psi_C(f)\beta$  it follows  $\alpha \wedge \Sigma_f(\Psi_C(f)\alpha \Rightarrow \Psi_C(f)\beta) \leq \beta$  whence the claim.  $\square$

**7.7 Lemma.** *Suppose  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is a variational doctrine. If  $Q_P$  is locally cartesian closed, then  $\mathcal{C}$  is slice-wise weakly cartesian closed.*

*Proof.* Immediate after 7.4.  $\square$

We now aim at proving the converse of A.1. Let  $\mathcal{C}$  be slice-wise weakly cartesian closed. Let  $P$  be a universal implicational doctrine on  $\mathcal{C}$  and consider  $[f]: (B, \sigma) \rightarrow (A, \delta_A)$  and  $[g]: (B', \sigma') \rightarrow (A, \delta_A)$ . The fact that  $\mathcal{C}$  is slice-wise weakly cartesian closed ensures that the left diagram below is commutative



where the square (A) is a weak pullback and  $t': W' \rightarrow A$  is a weak exponential of  $g$  to the power of  $f$  in  $\mathcal{C}/A$  with weak evaluation  $e'$ . Now consider the element  $\xi$  of  $P(W')$  defined as

$$\bigvee_{\text{pr}_3} P_{\langle p_{W'}, \text{pr}_1, \text{pr}_3 \rangle}(\delta_{W'}) \wedge P_{\langle p_{W'}, \text{pr}_2, \text{pr}_3 \rangle}(\delta_{W'}) \wedge P_{\langle p_B \text{pr}_1, p_B \text{pr}_2 \rangle}(\sigma) \rightarrow P_{\langle e' \text{pr}_1, e' \text{pr}_2 \rangle}(\delta_{B'})$$

where  $\text{pr}_1$ ,  $\text{pr}_2$  and  $\text{pr}_3$  are projections from  $P \times P \times W'$ . The composition  $t'$  with  $\{\xi\}: W \rightarrow W'$  determines  $t = t' \{\xi\}: W \rightarrow A$  in the right diagram above. The square (B) is the weak pullback of  $t$  along  $f$ . Since  $f \pi_B = t \pi_W = t' \{\xi\} \pi_W$ , there is an arrow  $\text{id}_f \times_A \{\xi\}: Z \rightarrow S$  and we denoted by  $e$  the composition  $e'(\text{id}_f \times_A \{\xi\}): Z \rightarrow B'$ . This shows that also the right diagram above is commutative.

Define the  $P$ -equivalence relation  $\theta$  on  $W$  as  $P_{t \times t}(\delta_A) \wedge \omega$  where  $\omega$  is

$$\bigvee_{\langle \text{pr}_2, \text{pr}_3 \rangle} P_{\langle \pi_W \text{pr}_1, \text{pr}_2 \rangle}(\delta_W) \wedge P_{\langle \pi_W \text{pr}_3, \text{pr}_4 \rangle}(\delta_W) \wedge P_{\langle \pi_B \text{pr}_1, \pi_B \text{pr}_3 \rangle}(\sigma) \rightarrow P_{\langle e \text{pr}_2, e \text{pr}_4 \rangle}(\delta_{B'})$$

where projections are from  $Z \times W \times Z \times W$ . We leave as an exercise to verify that

$$P_{\pi_B \times \pi_B}(\sigma) \wedge P_{\pi_W \times \pi_W}(\theta) \leq P_{e \times e}(\sigma')$$

establishing  $e$  as a legitimate representative of the following arrow in  $Q_P$

$$[e]: (Z, P_{\pi_B \times \pi_B}(\sigma) \wedge P_{\pi_W \times \pi_W}(\theta)) \rightarrow (B', \sigma')$$

**7.8 Lemma.** *Suppose  $P$  on  $\mathcal{C}$  is a variational f.o.d. If  $\mathcal{C}$  is slice-wise weakly cartesian closed, then for every  $A$  the slice  $\mathbf{Q}_P/(A, \delta_A)$  is cartesian closed.*

*Proof.* For  $[f]: (B, \sigma) \rightarrow (A, \delta_A)$  and  $[g]: (B', \sigma') \rightarrow (A, \delta_A)$  consider the diagram

$$\begin{array}{ccccc}
 & & (Z, P_{\pi_B \times \pi_B}(\sigma) \wedge P_{\pi_W \times \pi_W}(\theta)) & & \\
 & \swarrow & & \searrow & \\
 (B, \sigma) & & (C) & & (W, \theta) \\
 & \searrow & & \swarrow & \\
 & & (A, \delta_A) & & (B', \sigma')
 \end{array}$$

$[f]$                        $[g]$                        $[e]$                        $[t]$

where  $(C)$  is a pullback by 7.3. Take any  $[h]: (C, \gamma) \rightarrow (A, \delta_A)$  and  $[m]: [f] \times_{(A, \delta_A)} [h] \rightarrow [g]$ . Thanks to 7.3 we know that  $[m]: [f \times_A h] \rightarrow [g]$ . Thus there is  $m': C \rightarrow W'$  with  $e'(\text{id}_f \times_A m') = m$  which is easily seen to be such that  $m' = \{\xi\} \overline{m}$  for some  $\overline{m}: C \rightarrow W$ . Moreover

$$m = e'(\text{id}_f \times_A m') = e'(\text{id}_f \times \{\xi\})(\text{id}_f \times_A \overline{m}) = e(\text{id}_f \times_A \overline{m})$$

showing that  $[e]$  is a weak evaluation in  $\mathbf{Q}_P/(A, \delta_A)$ . It is now an exercise to show that if  $[l]: (C, \gamma) \rightarrow (W, \theta)$  is such that  $[e]([\text{id}_f] \times_{(A, \rho)} [l]) = [m]$  then  $\top_C \leq P_{(l, \overline{m})}(\theta)$  proving uniqueness of exponential transposes.  $\square$

After 7.8 it remains to prove that every slice of  $\mathbf{Q}_P$  is cartesian closed. Proposition 7.8 says that for every  $(A, \delta_A)$  and every  $[f]$  in  $\mathbf{Q}_P/(A, \delta_A)$  there is a right adjoint to the functor  $- \times_{(A, \delta_A)} [f]: \mathbf{Q}_P/(A, \delta_A) \rightarrow \mathbf{Q}_P/(A, \delta_A)$ . We aim at proving that these right adjoints exist for all  $(A, \rho)$ .

To this purpose we follow the line of the proof in [Emm20] by relying on the existence of right adjoints as a consequence of one of Barr's tripability theorems in [BW84].

We first need some instrumental propositions.

**7.9 Proposition.** *In a  $m$ -variational doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  with stable effective quotients each quotient arrow is the coequalizer of its kernel pairs and hence it is a regular epimorphism. Moreover  $\mathcal{C}$  is regular.*

*Proof.* In [MR13b] it is proved that the kernel pairs of each map in  $\mathcal{C}$  has a coequalizer which is an effective quotient arrow and hence  $\mathcal{C}$  is regular.  $\square$

**7.10 Corollary.** *In a  $m$ -variational doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  with stable effective quotients each coequalizer is a quotient arrow and it is stable under pullback.*

*Proof.* Every coequalizer  $e: A \rightarrow B$  is isomorphic to  $q: A \rightarrow A/P_{e \times e}(\delta_B)$ .  $\square$

**7.11 Proposition.** *Let  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  be an existential variational doctrine and  $\widehat{P}: \mathbf{Q}_P^{\text{op}} \rightarrow \mathbf{ISL}$  be its elementary quotient completion. Then, for every quotient arrow  $q: (A, \delta_A) \rightarrow (A, \rho)$  in  $\mathbf{Q}_P$  the functor  $q^*: \mathbf{Q}_P/(A/\rho) \rightarrow \mathbf{Q}_P/(A, \delta_A)$  is monadic.*

*Proof.* The proof is the same as that for exact categories given in [JM95]. Alternatively, just observe that  $\mathcal{Q}_{\mathcal{P}}$  is regular and that any quotient map is a regular epimorphism by lemma 7.9. Therefore it is well known that  $q^*$  is of descent type.

Moreover, any map  $f: (B, \tau) \rightarrow (A, \rho)$  is isomorphic over  $(A, \rho)$  to  $\text{pr}_1 \cdot \llbracket P_{\langle id, f \rangle}(\rho) \rrbracket: (X, P_{\langle \text{pr}_1, \text{pr}_3 \rangle}(\rho) \wedge P_{\langle \text{pr}_1, \text{pr}_3 \rangle}(\tau)) \rightarrow (A, \rho)$  which comes from a descent datum represented by  $\text{pr}_1 \cdot \llbracket P_{\langle id, f \rangle}(\rho) \rrbracket: (X, \delta_X) \rightarrow (A, \delta_A)$  with the obvious action.  $\square$

Theorem 7.11 actually holds even in the more general situation of a generic existential variational doctrine with effective stable quotients with the same proof.

We recall from theorem 3.7.2 in [BW84].

**7.12 Proposition.** *In a situation like the following*

$$\begin{array}{ccc} \mathcal{B} & \xrightarrow{W} & \mathcal{B}' \\ \swarrow U & & \searrow U' \\ & \mathcal{C} & \\ \nwarrow F & & \nearrow F' \end{array}$$

where  $F \dashv U$  and  $F' \dashv U'$ , if furthermore

1.  $WF$  is naturally isomorphic to  $F'$ ;
2.  $U$  is monadic;
3.  $W$  preserves co-equalizers of  $U$ -contractible pairs;

then  $W$  has a right adjoint.

We apply this theorem to our context as follows:

**7.13 Proposition.** *If an  $m$ -variational elementary  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  has stable effective quotients, then for every quotient arrow  $q: A \rightarrow A/\rho$ , if the slice  $\mathcal{C}/A$  is cartesian closed also the slice  $\mathcal{C}/(A/\rho)$  is cartesian closed.*

*Proof.* The cartesian closure of  $\mathcal{C}/A$  ensures that for every  $\alpha$  in  $\mathcal{C}/(A/\rho)$  the functor  $- \times_A q^* \alpha: \mathcal{C}/A \rightarrow \mathcal{C}/A$  has a right adjoint  $R_\alpha$ . Consider the diagram

$$\begin{array}{ccc} \mathcal{C}/(A/\rho) & \xrightarrow{- \times_{(A/\rho)} \alpha} & \mathcal{C}/(A/\rho) \\ \swarrow q^* & & \searrow R_\alpha q^* \\ & \mathcal{C}/A & \\ \nwarrow \Sigma_q & & \nearrow \Sigma_q(- \times_A q^* \alpha) \end{array}$$

Since  $\Sigma_q \dashv q^*$  and  $\Sigma_q(- \times_A q^* \alpha) \dashv R_\alpha q^*$  (by composition of adjoints), the claim is proved if we show that this situation meets the three points of the list above.

1. Take any  $\beta$  in  $\mathcal{C}/A$ . Frobenius reciprocity precisely says that  $\Sigma_q(\beta) \times_{(A/\rho)} \alpha$

is isomorphic to  $\Sigma_q(\beta \times_A q^* \alpha)$ .

2. Every quotient arrow  $q: A \rightarrow A/\rho$  induces a monadic functor  $q^*$  by 7.11.
3. In  $\mathcal{C}/(A/\rho)$  coequalizers of  $q^*$ -contractible pairs are preserved by  $- \times_A q^* \alpha$  thanks to stability of coequalizers in corollary 7.10.  $\square$

We are at last in the position to formulate the main theorem of the section.

**7.14 Theorem.** *Suppose  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is a variational implicational and universal doctrine. Then  $\mathcal{C}$  is slice-wise weakly cartesian closed if and only if  $\mathcal{Q}_P$  is locally cartesian closed.*

*Proof.* The necessary condition is A.1. For the converse note that every slice  $\mathcal{Q}_P/(A, \delta_A)$  is cartesian closed (7.8). For every  $(A, \rho)$  the arrow  $[\text{id}_A]: (A, \delta_A) \rightarrow (A, \rho)$  is a quotient arrow. The claim follows by 7.13.  $\square$

Recall that the exact completion of a category with binary products and weak pullbacks is an instance of the elementary quotient completion. More specifically  $\widehat{\Psi}_{\mathcal{C}} \simeq \text{Sub}_{\mathcal{C}_{\text{ex/lex}}}$ , which has as corollary  $\mathcal{Q}_{\Psi_{\mathcal{C}}} \equiv \mathcal{C}_{\text{ex/lex}}$ .

Therefore Proposition A.5 shows that  $\mathcal{C}$  is slice-wise weakly cartesian closed if and only if  $\mathcal{C}_{\text{ex/lex}}$  is locally cartesian. Hence we recover the characterization in [CR00] and [Emm20].

**7.15 Corollary.** *Suppose  $\mathcal{C}$  a category with finite products and weak pullbacks. Then  $\mathcal{C}$  is slice-wise weakly cartesian closed if and only if  $\mathcal{C}_{\text{ex/lex}}$  is locally cartesian closed.*

*Proof.* It follows from A.5 when applied to the doctrine functor of variations  $\Psi_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$ . Just note that the doctrine of variations is implicational with universal quantifications by 7.6.  $\square$

**7.16 Remark.** The proof of theorem A.5 can be considered a generalization of Carboni-Rosolini characterization of locally cartesian closed exact completions  $\mathcal{C}_{\text{ex/lex}}$  in [CR00] *only* in the case the category has finite products. Related proofs in the special cases of elementary quotient completion of syntactic category out of specific type theories are in [Pal19] and [Mai09].

**7.17 Remark.** We know from lemma 4.19 that  $Q_P$  is the category of  $\widehat{j}_P$ -separated objects of  $C_{\text{ex/lex}}$ . But this relation does not lift to slice categories. Indeed, for an object  $A$  of  $Q_P$  the category  $Q_P/A$  is not necessarily the category of separated objects of  $C_{\text{ex/lex}}/A$ . To show this, first observe that our definition of separated object does coincide with definition point a) of def. 4.3.4 page 189 in [Joh02a], which we call "presheaf separated object". Then recall that the full subcategory of presheaf separated objects over a separated object  $A$  form an exponential ideal of  $C_{\text{ex/lex}}/A$  as shown in ii) of lemma 4.4.3 in [Joh02a]. But our category  $Q_P/A$  is not an exponential ideal of  $C_{\text{ex/lex}}/A$  because the reflector does not preserve equalizers. Therefore we conclude that the category  $Q_P/A$  does not coincide with all the  $\widehat{j}_P$ -separated objects of  $C_{\text{ex/lex}}$ .

## 7.2 Coproducts

In this section we establish the necessary and sufficient conditions under which an elementary quotient completion has stable finite coproducts.

Suppose  $\mathcal{C}$  has finite coproducts. Coproducts are said distributive if the canonical arrow  $(A \times B) + (A \times C) \rightarrow A \times (B + C)$  is an isomorphism.

We are interested in studying those elementary doctrines with comprehensive diagonals whose elementary quotient completion has coproducts. After 7.2 we know that if  $Q_P$  has coproducts then  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  must have coproducts.

For the rest of the section, unless specified otherwise,  $P$  is a variational f.o.d. on a category  $\mathcal{C}$  with binary distributive coproducts.

**7.18 Proposition.** *Canonical injections are  $P$ -injective, i.e.  $P_{i_A \times i_A}(\delta_{A+B}) = \delta_A$  and  $P_{i_B \times i_B}(\delta_{A+B}) = \delta_B$ .*

*Proof.* The idea of the proof is simple whilst there are various calculations involved and this is the following one in the internal language of doctrines. Given an element  $a: A$  then we can define a projection  $\overline{p}: A + B \rightarrow A$  as follows

$$\overline{p}_a(z) \equiv \begin{cases} w & \text{if } z = i_A(w) \\ a & \text{if } z = i_B(w') \end{cases}$$

Hence if  $i_A(x) =_{A+B} i_A(y)$  then  $x =_A \overline{p}_x(i_A(x)) =_A \overline{p}_x(i_A(y)) =_A y$ .

Suppose  $i_A: A \rightarrow A+B$  is a canonical injection and consider the commutative diagram

$$\begin{array}{ccccc} & & A & & \\ & \text{pr}_2 \curvearrowright & \uparrow & \text{pr}_1 \curvearrowleft & \\ & & [pr_2, pr_1] & & \\ A \times A & \xrightarrow{\quad} & (A \times A) + (A \times B) & \xleftarrow{\quad} & A \times B \\ & \text{id}_A \times i_A \searrow & \uparrow j & \swarrow \text{id}_A \times i_B & \\ & & A \times (A + B) & & \end{array}$$

where  $j$  is the isomorphism that comes from distributivity. Denote by  $\text{pr}_i$  ( $i = 1, 2, 3, 4$ ), the projections from  $Z = A \times (A+B) \times A \times (A+B)$ . By Beck-Chevalley

conditions on equality (see [MR13b]). one has both

$$\begin{aligned}\delta_Z &= P_{\langle \text{pr}_1, \text{pr}_3 \rangle}(\delta_A) \wedge P_{\langle \text{pr}_2, \text{pr}_4 \rangle}(\delta_{A+B}) \\ \delta_Z &\leq P_j \times_j P_{[\text{pr}_2, \text{pr}_1] \times [\text{pr}_2, \text{pr}_1]}(\delta_A)\end{aligned}$$

Evaluating both sides on  $P_e$  for

$$e = \langle \text{pr}_1, i_A \text{pr}_1, \text{pr}_1, i_A \text{pr}_2 \rangle: A \times A \rightarrow Z$$

where  $\text{pr}_1, \text{pr}_2: A \times A \rightarrow A$  are projections, we obtain

$$\begin{aligned}P_e(\delta_Z) &= P_{\langle \text{pr}_1, \text{pr}_1 \rangle}(\delta_A) \wedge P_{i_A \times i_A}(\delta_{A+B}) = P_{i_A \times i_A}(\delta_{A+B}) \\ P_e(\delta_Z) &\leq P_e P_j \times_j P_{[\text{pr}_2, \text{pr}_1] \times [\text{pr}_2, \text{pr}_1]}(\delta_A)\end{aligned}$$

Note that  $e$  can be obtained by the following composition

$$\begin{array}{ccc} A \times (A \times A) & \xrightarrow{\langle \text{id}_A, i_A \rangle \times (\text{id}_A \times i_A)} & (A \times (A + B)) \times (A \times (A + B)) \\ \parallel & & \uparrow e \\ (A \times A) \times A & \xleftarrow{\Delta_A \times \text{id}_A} & A \times A \end{array}$$

Therefore  $([\text{pr}_2, \text{pr}_1] \times [\text{pr}_2, \text{pr}_1])(j \times j)e$  is equal to

$$([\text{pr}_2, \text{pr}_1]j \langle \text{id}_A, i_A \rangle \times [\text{pr}_2, \text{pr}_1]j \langle \text{id}_A \times i_A \rangle)(\Delta_A \times \text{id}_A)$$

since  $[\text{pr}_2, \text{pr}_1]j \langle \text{id}_A \times i_A \rangle = \text{pr}_2: A \times A \rightarrow A$  one has

$$([\text{pr}_2, \text{pr}_1]j \langle \text{id}_A, i_A \rangle \times [\text{pr}_2, \text{pr}_1]j \langle \text{id}_A \times i_A \rangle)(\Delta_A \times \text{id}_A) = \text{id}_{A \times A}$$

whence  $P_{i_A \times i_A}(\delta_{A+B}) = P_e(\delta_Z) \leq \delta_A$ .  $\square$

We now aim at proving that the equality on a coproduct is given by the formula

$$\delta_{A+B} = \mathcal{E}_{i_A \times i_A}(\delta_A) \vee \mathcal{E}_{i_B \times i_B}(\delta_B)$$

To this purpose we say that two arrows  $h: A \rightarrow Y$  and  $k: X \rightarrow Y$  are **jointly  $P$ -surjective** if

$$\top_Y = \mathcal{E}_h(\top_A) \vee \mathcal{E}_k(\top_X)$$

**7.19 Proposition.** *Canonical injections are jointly  $P$ -surjective.*

*Proof.* Suppose  $A$  and  $B$  are objects of  $\mathcal{C}$  and consider their coproduct with canonical injections  $i_A: A \rightarrow A + B$  and  $i_B: B \rightarrow A + B$ . Abbreviate by  $k: X \rightarrow A + B$  the weak comprehension

$$\{\!\{ \mathcal{E}_{i_A}(\top_A) \vee \mathcal{E}_{i_B}(\top_B) \}\!\}: X \rightarrow A + B$$

Clearly

$$\top_A \leq P_{i_A}(\mathfrak{A}_{i_A}(\top_A)) \leq P_{i_A}(\mathfrak{A}_{i_A}(\top_A) \vee \mathfrak{A}_{i_B}(\top_B))$$

and analogously for  $i_B$ . By the universal property of comprehensions there are arrows  $t_A: A \rightarrow X$  and  $t_B: B \rightarrow X$  with  $kt_A = i_A$  and  $kt_B = i_B$ . These induce an arrow  $[t_A, t_B]: A + B \rightarrow X$  which is a section of  $k$ . Thus  $\llbracket \top_{A+B} \rrbracket = \text{id}_{A+B}$  factors through  $\llbracket \mathfrak{A}_{i_A}(\top_A) \vee \mathfrak{A}_{i_B}(\top_B) \rrbracket$ . Fullness of comprehensions completes the proof.  $\square$

**7.20 Proposition.** *For every pair of  $P$ -injective arrows  $h: A \rightarrow Y$  and  $k: X \rightarrow Y$ , for every reflexive relation  $\rho$  over  $A$  and every reflexive relation  $\theta$  over  $X$ , the following relation over  $Y$*

$$\mathfrak{A}_{h \times h}(\rho) \vee \mathfrak{A}_{k \times k}(\theta)$$

*is reflexive if and only if  $h$  and  $k$  are jointly surjective.*

*Proof.* Suppose  $h$  and  $k$  are jointly surjective and consider the relation  $\mathfrak{A}_{h \times h}(\rho)$ . By Beck-Chevalley conditions it is

$$P_{\Delta_Y}(\mathfrak{A}_{h \times h}(\rho)) = \mathfrak{A}_h(P_{\Delta_A}(\rho)) = \mathfrak{A}_h(\top_A)$$

Analogously  $P_{\Delta_Y}(\mathfrak{A}_{k \times k}(\theta)) = \mathfrak{A}_k(\top_X)$  whence

$$P_{\Delta_Y}(\mathfrak{A}_{h \times h}(\rho) \vee \mathfrak{A}_{k \times k}(\theta)) = \mathfrak{A}_h(\top_A) \vee \mathfrak{A}_k(\top_X)$$

whence the claim, i.e.

$$\top_Y = \mathfrak{A}_h(\top_A) \vee \mathfrak{A}_k(\top_X)$$

iff

$$\top_Y = \mathfrak{A}_h(\top_A) \vee \mathfrak{A}_k(\top_X)$$

$\square$

**7.21 Proposition.** *For every  $A$  and  $B$  in  $\mathcal{C}$  it is*

$$\delta_{A+B} = \mathfrak{A}_{i_A \times i_A} \delta_A \vee \mathfrak{A}_{i_B \times i_B} \delta_B$$

*where  $i_A: A \rightarrow A + B$  and  $i_B: B \rightarrow A + B$  are canonical injections.*

*Proof.* From  $\delta_A \leq P_{i_A \times i_A}(\delta_{A+B})$  and  $\delta_B \leq P_{i_B \times i_B}(\delta_{A+B})$  one obtains a canonical inequality

$$\mathfrak{A}_{i_A \times i_A} \delta_A \vee \mathfrak{A}_{i_B \times i_B} \delta_B \leq \delta_{A+B}$$

This is actually an equality, since injections are  $P$ -injective (7.18) and jointly surjective (7.19) hence  $\mathfrak{A}_{i_A \times i_A} \delta_A \vee \mathfrak{A}_{i_B \times i_B} \delta_B$  is reflexive by 7.20.  $\square$

We leave to the readers the proof of the following instrumental lemma before the main statement of the section.

**7.22 Proposition.** *If  $\rho$  in  $P(A \times A)$  is transitive and if  $h: A \rightarrow B$  is  $P$ -injective, then  $\mathfrak{A}_{h \times h}(\rho)$  is transitive.*

*Proof.* Immediate.  $\square$

We can finally focus on the elementary quotient completion of a doctrine whose base has distributive binary coproducts.

**7.23 Theorem.** *Suppose  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is a variational first order doctrine.  $\mathcal{C}$  has distributive binary coproducts if and only if  $\mathcal{Q}_P$  has distributive binary coproducts.*

*Proof.* Suppose  $\mathcal{C}$  has binary distributive coproducts. Consider  $(A, \rho)$  and  $(B, \sigma)$  in  $\mathcal{Q}_P$  and the coproduct  $A + B$  with canonical injections  $i_A$  and  $i_B$ . Define

$$\rho \boxplus \sigma = \mathcal{E}_{i_A \times i_A} \rho \vee \mathcal{E}_{i_B \times i_B} \sigma$$

by lemmas 7.20 and 7.19 the relation  $\rho \boxplus \sigma$  is reflexive. It is trivially symmetric. Transitivity follows from 7.22 and 7.18. Thus

$$(A, \rho) \xrightarrow{[i_A]} (A + B, \rho \boxplus \sigma) \xleftarrow{[i_B]} (B, \sigma)$$

is a diagram in  $\mathcal{Q}_P$ . We claim that it is a coproduct diagram of  $(A, \rho)$  and  $(B, \sigma)$ . It is immediate to see that it is a weak coproduct. We now prove that it is a strong coproduct. Suppose that  $f: A \rightarrow T$  and  $g: B \rightarrow T$  represent two arrows  $[f]: (A, \rho) \rightarrow (T, \theta)$  and  $[g]: (B, \sigma) \rightarrow (T, \theta)$ . If  $k$  and  $l$  represents two arrows  $[k], [l]: (A + B, \rho \boxplus \sigma) \rightarrow (T, \theta)$  with

$$\begin{aligned} [k][i_A] &= [f] & \text{and} & & [k][i_B] &= [g] \\ [l][i_A] &= [f] & \text{and} & & [l][i_B] &= [g] \end{aligned}$$

*i.e.*  $k$  and  $l$  are such that

$$\begin{aligned} \top_A &\leq P_{\langle ki_A, f \rangle}(\theta) & \text{and} & & \top_B &\leq P_{\langle ki_B, g \rangle}(\theta) \\ \top_A &\leq P_{\langle li_A, f \rangle}(\theta) & \text{and} & & \top_B &\leq P_{\langle li_B, g \rangle}(\theta) \end{aligned}$$

then we have also  $\rho \leq P_{ki_A \times f}(\theta) \wedge P_{li_A \times f}(\theta) \leq P_{i_A \times i_A} P_{k \times l}(\theta)$  and similarly  $\sigma \leq P_{i_B \times i_B} P_{k \times l}(\theta)$ . Therefore

$$\mathcal{E}_{i_A \times i_A} \rho \leq P_{k \times l}(\theta) \quad \text{and} \quad \mathcal{E}_{i_B \times i_B} \sigma \leq P_{k \times l}(\theta)$$

from which  $\rho \boxplus \sigma \leq P_{k \times l}(\theta)$ , *i.e.*  $[k] = [l]$ .

For the converse, consider  $A$  and  $B$  in  $\mathcal{C}$ . The coproduct in  $\mathcal{Q}_P$  of  $(A, \delta_A)$  and  $(B, \delta_B)$  is  $(A + B, \delta_A \boxplus \delta_B)$ . By 7.23 this is  $(A + B, \delta_{A+B})$ . The claim follows by 7.2-2.  $\square$

We say  $A + B$  is  $P$ -disjoint if

$$P_{i_A \times i_B}(\delta_{A+B}) = \perp_{A \times B}$$

**7.24 Proposition.**  *$\mathcal{C}$  has  $P$ -disjoint distributive binary coproducts if and only if  $\mathcal{Q}_P$  has  $\widehat{P}$ -disjoint distributive binary coproducts.*

*Proof.* After 7.23 we only have to prove that coproducts in  $\mathcal{C}$  are  $P$ -disjoint if and only if coproduct in  $\mathcal{Q}_P$  are  $\hat{P}$ -disjoint. The necessary condition is immediate after 7.21. Consider the coproduct diagram in  $\mathcal{Q}_P$

$$(A, \rho) \xrightarrow{[i_A]} (A + B, \rho \boxplus \sigma) \xleftarrow{[i_B]} (B, \sigma)$$

it holds

$$P_{i_A \times i_B}(\rho \boxplus \sigma) = P_{i_A \times i_B}(\mathcal{A}_{i_A \times i_A} \rho \vee \mathcal{A}_{i_B \times i_B} \sigma)$$

but, denoting by  $\text{pr}_i$  the projections from  $A \times B \times (A+B) \times (A+B)$ ,  $P_{i_A \times i_B}(\mathcal{A}_{i_A \times i_A} \rho)$  is equal to

$$\mathcal{A}_{\langle \text{pr}_1, \text{pr}_2 \rangle} [P_{\langle \text{pr}_1, \text{pr}_3 \rangle}(P_{i_A \times i_A}(\delta_{A+B})) \wedge P_{\langle \text{pr}_2, \text{pr}_4 \rangle}(P_{i_A \times i_B}(\delta_{A+B})) \wedge P_{\langle \text{pr}_3, \text{pr}_4 \rangle}(\rho)]$$

which is equal to  $\perp_{A \times B}$  under the assumption that  $P_{i_A \times i_B}(\delta_{A+B})$  is so. Analogously  $P_{i_A \times i_B}(\mathcal{A}_{i_B \times i_B} \sigma) = \perp_{A \times B}$  whence the claim.  $\square$

**7.25 Corollary.** *Suppose  $\mathcal{C}$  is such that for every  $A$  the domain of  $\{\perp_A\}$  is an initial object. Then  $\mathcal{C}$  has disjoint distributive binary coproducts if and only if  $\mathcal{Q}_P$  has disjoint distributive binary coproducts.*

### 7.3 Classifiers

In this section we show how the elementary quotient completion inherits a predicate classifier as defined in 2.13 for strong monomorphisms.

We first need to establish the following characterisation of epimorphisms as  $P$ -surjective arrows.

**7.26 Lemma.** *Suppose  $P$  is an existential variational doctrine with a strong predicate classifier. An arrow  $e: X \rightarrow A$  is epic if and only if it is  $P$ -surjective.*

*Proof.* We have already observed that  $P$ -surjective arrows are epimorphism. So let  $e: X \rightarrow A$  be epic. Consider the arrows  $\chi_{\mathcal{A}_e(\top_X)}, \chi_{\top_A}: A \rightarrow \Omega$ . Then

$$P_e(P_{\chi_{\mathcal{A}_e(\top_X)}}(\epsilon)) = P_e(\mathcal{A}_e(\top_X)) = \top_X = P_e(\top_A) = P_e(P_{\chi_{\top_A}}(\epsilon))$$

Since the classifier is strong  $\chi_{\top_A} e = \chi_{\mathcal{A}_e(\top_X)} e$  and then  $\chi_{\top_A} = \chi_{\mathcal{A}_e(\top_X)}$  as  $e$  is epic, whence  $\top_A = P_{\chi_{\top_A}}(\epsilon) = P_{\chi_{\mathcal{A}_e(\top_X)}}(\epsilon) = \mathcal{A}_e(\top_X)$ .  $\square$

Recall that given two factorization system  $(E, M)$  and  $(E', M')$  on the same category  $\mathcal{C}$ , we have that  $E = E'$  if and only if  $M = M'$ . Thus the following is an immediate corollary of 7.26 and 4.21.

**7.27 Proposition.** *In every  $m$ -variational existential doctrine with a strong predicate classifier comprehension arrows are the strong monomorphisms of the base.*

*Proof.* By 4.21 comprehension arrows are the class of arrows which are right orthogonal to  $P$ -surjective arrows, but these coincide with epimorphisms by 7.26, whence the claim.  $\square$

**7.28 Proposition.** *In every  $m$ -variational existential doctrine with a strong predicate classifier the domain  $T$  of  $\{\!\|\in\!\|\}: T \rightarrow \Omega$  is a terminal object.*

*Proof.* Let  $1$  be terminal. The arrow  $\chi_{\top_1}: 1 \rightarrow \Omega$  is such that  $P_{\chi_{\top_1}}(\in) = \top_1$ . The universal property of  $\{\!\|\in\!\|\}$  produces an arrow  $k: 1 \rightarrow T$  with  $\{\!\|\in\!\|\}k = \chi_{\top_1}$ . The universal property of  $1$  ensures that  $!_T k = \text{id}_1$ . Moreover from

$$\{\!\|\in\!\|\}(\in) = \top_T = P_{!_T}(\top_1) = P_{!_T}P_{\chi_{\top_1}}(\in)$$

we have  $\chi_{\top_1}!_T = \{\!\|\in\!\|\}$ , so  $\{\!\|\in\!\|\}k!_T = \{\!\|\in\!\|\}$ , whence  $k!_T = \text{id}_T$  as  $\{\!\|\in\!\|\}$  is monic.  $\square$

**7.29 Corollary.** *If  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is a  $m$ -variational existential doctrine, then  $P$  has a strong predicate classifier if and only if  $\mathcal{C}$  has a classifier of strong monomorphism.*

*Proof.* Suppose  $\Omega$  is a strong predicate classifier in  $P$ . After 7.27 it suffices to show that  $\Omega$  is a classifier for the class of comprehension arrows. Note that every  $\alpha$  in  $P(A)$  the following is a pullback

$$\begin{array}{ccc} X & \xrightarrow{!} & T \\ \{\!\|\alpha\!\|\} \downarrow & & \downarrow \{\!\|\in\!\|\} \\ A & \xrightarrow{\chi_\alpha} & \Omega \end{array}$$

where  $!: X \rightarrow T$  is the arrow produced by the universal property of  $\{\!\|\in\!\|\}$  as  $\top_X = P_{\{\!\|\alpha\!\|\}}(\alpha) = P_{\{\!\|\alpha\!\|\}}P_{\chi_\alpha}(\in) = P_{\chi_\alpha\{\!\|\alpha\!\|\}}(\in)$ . The object  $T$  is a terminal (7.28). If  $f: A \rightarrow \Omega$  makes the square  $\{\!\|\in\!\|\}! = f\{\!\|\alpha\!\|\}$  a pullback, then  $\{\!\|\alpha\!\|\}$  is isomorphic to  $\{\!\|P_f(\in)\!\|\}$  (4.7). By fullness of comprehension  $P_f(\in) = \alpha = P_{\chi_\alpha}(\in)$ , hence  $f = \chi_\alpha$ .

Conversely suppose  $t: 1 \rightarrow \Omega$  is a strong monomorphism classifier. Define  $\in = \mathcal{A}_t(\top_1)$ . For  $\alpha$  in  $P(A)$  the arrow  $\{\!\|\alpha\!\|\}: X \rightarrow A$  is a strong monic by 7.27. Thus there is a unique  $\chi_\alpha: A \rightarrow \Omega$  that makes the following

$$\begin{array}{ccc} X & \longrightarrow & 1 \\ \{\!\|\alpha\!\|\} \downarrow & & \downarrow t \\ A & \xrightarrow{\chi_\alpha} & \Omega \end{array}$$

a pullback. By 4.6 and 4.5 it is  $P_{\chi_\alpha}(\in) = P_{\chi_\alpha}\mathcal{A}_t(\top_1) = \mathcal{A}_{\{\!\|\alpha\!\|\}}P_{!_X}(\top_1) = \mathcal{A}_{\{\!\|\alpha\!\|\}}(\top_X) = \alpha$ . Suppose  $f: A \rightarrow \Omega$  is such that  $P_f(\in) = \alpha$ . Since  $t: 1 \rightarrow \Omega$  is a strong monomorphism classifier its pullback along  $f$  classifies a strong monomorphism. By 7.27 this is of the form  $\{\!\|\beta\!\|\}: X' \rightarrow A$ . Moreover, by 4.6  $\alpha = P_f(\in) \equiv P_f(\mathcal{A}_t(\top_1)) = \mathcal{A}_{\{\!\|\beta\!\|\}}(\top_X) = \beta$ . So  $\{\!\|\alpha\!\|\} = \{\!\|\beta\!\|\}$  and hence  $\chi_\alpha$  and  $f$  classifies the same strong monomorphisms showing that  $f = \chi_\alpha$ .  $\square$

**7.30 Proposition.** *Let  $P$  be a variational first order doctrine on  $\mathcal{C}$ . The following are equivalent*

1.  $P$  has a weak predicate classifier
2.  $\widehat{P}$  has a strong predicate classifier
3.  $Q_P$  has a classifier for strong monomorphisms.

*Proof.* 1  $\Leftrightarrow$  2. The necessary condition is immediate, while if  $\Omega$  is a predicate weak classifier in  $\mathcal{C}$ , then  $(\Omega, \lambda)$  is a predicate classifier in  $Q_P$  where

$$\lambda = P_{\text{pr}_1}(\epsilon) \leftrightarrow P_{\text{pr}_2}(\epsilon)$$

2  $\Leftrightarrow$  3. By 4.17 the doctrine  $\widehat{P}$  is m-variational. Then apply 7.29. □

## 8 The quasi-topos completion

In this section we are going to show how the elementary quotient completion of a suitable tripos gives rise to a quasi-topos completion.

We start by recalling the definition of quasi-topos which is a generalisation of that of topos:

**8.1 Definition.** A *quasi-topos* is a category  $\mathcal{C}$

- (i) with finite limits
- (ii) with finite co-limits
- (iii) locally cartesian closed
- (iv) there is a classifier for strong monomorphisms

Then recall from [Wyl91] that Every quasi-topos has effective quotients of strong equivalence relations, namely a  $\text{Sub}_{\mathcal{C}}$ -equivalence relation represented by a strong monomorphism. Therefore it makes sense to try to characterise those quasi-toposes which arise as elementary quotient completions.

In the following when we refer to an equivalence relation in a category  $\mathcal{C}$  we mean a  $\text{Sub}_{\mathcal{C}}$ -equivalence relation, while for a strong equivalence relation we mean a  $\text{Sub}_{\mathcal{C}}$ -equivalence relation represented by a strong monomorphism.

If  $\mathcal{C}$  is a quasi-topos, then both  $\Psi_{\mathcal{C}}$  and  $\text{Sub}_{\mathcal{C}}$  are first order. There is another doctrine based on a quasi-topos  $\mathcal{C}$  the doctrine of strong subobjects of  $\mathcal{C}$ . We denote it by  $\text{Stg}_{\mathcal{C}}^{\text{op}} \longrightarrow \mathbf{ISL}$ .

**8.2 Definition.** A *intensional hyper tripos* is an elementary existential doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  with with a weak predicate classifier and full weak comprehension such that  $\mathcal{C}$  has weak pullbacks, is slice-wise weakly cartesian closed and has finite distributive coproducts. A *hyper tripos* is a intensional hyper tripos with comprehensive diagonals.

**8.3 Proposition.** A *intensional hyper tripos* is a tripos.

*Proof.* Suppose  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  is a intensional hyper tripos. Then it has full weak comprehension. By 4.14 there is a topology  $j$  on  $\Psi_{\mathcal{C}}$  such that  $P$  is  $\Psi_{\mathcal{C}j}$ . Since  $\mathcal{C}$  has weak pullbacks, is slice-wise weakly cartesian closed and has finite distributive coproducts, the doctrine  $\Psi_{\mathcal{C}}$  is first order, so is  $P$  by 4.13.  $\mathcal{C}$  is weakly cartesian closed and  $P$  has a weak classifier, so  $P$  has weak power objects.  $\square$

**8.4 Proposition.** *A hyper tripos has  $P$ -disjoint coproducts.*

*Proof.* take two objects  $A, B$  and consider  $\chi_{\top_A}: A \rightarrow \Omega$  and  $\chi_{\perp_B}: B \rightarrow \Omega$ . Abbreviate with  $d: A + B \rightarrow \Omega$  the arrow  $[\chi_{\top_A}, \chi_{\perp_B}]$ . From  $\delta_{A+B} \leq P_{d \times d}(\delta_{\Omega})$  we get  $P_{i_A \times i_B}(\delta_{A+B}) \leq P_{di_A \times di_B}(\delta_{\Omega}) = \chi_{\top_A} \times \chi_{\perp_B}(\delta_{\Omega})$ . Finally observe that  $\delta_{\Omega} \leq P_{p_1}(\epsilon) \leftrightarrow P_{p_2}(\epsilon)$  from which we deduce  $\chi_{\top_A} \times \chi_{\perp_B}(\delta_{\Omega}) \leq P_{\pi_1}(P_{\chi_{\top_A}}(\epsilon)) \leftrightarrow P_{\pi_2}(P_{\chi_{\top_A}}(\epsilon)) = P_{p_1}(\top_A) \leftrightarrow P_{p_2}(\perp_B) = \top_{A \times B} \leftrightarrow \perp_{A \times B} = \perp_{A \times B}$ . One concludes that  $P_{i_A \times i_B}(\delta_{A+B}) \leq \perp_{A \times B}$ .  $\square$

**8.5 Proposition.** *Let  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  be a tripos. For every  $A$  and every  $\rho$  in  $P(A \times A)$  there is a  $P$ -equivalence relation  $\bar{\rho}$  over  $A$  such that  $\rho \leq \bar{\rho}$  and for every  $P$ -equivalence relation  $\sigma$  over  $A$ , if  $\rho \leq \sigma$  then  $\sigma \leq \bar{\rho}$ .*

*Proof.* We shall employ the language introduced in 2.16. Take  $A$  in  $\mathcal{C}$  and define the formulas in  $U: \mathbb{P}A \times A$

$$\mathbf{r}(U) := \forall_{a:A} (a, a) \in_A U$$

$$\mathbf{s}(U) := \forall_{a:A} \forall_{a':A} [(a, a') \in_A U \Rightarrow (a', a) \in_A U]$$

$$\mathbf{t}(U) := \forall_{a:A} \forall_{a':A} \forall_{a'':A} [(a, a') \in_A U \wedge (a', a'') \in_A U \Rightarrow (a, a'') \in_A U]$$

$$\mathbf{eq}(U) := \mathbf{r}(U) \wedge \mathbf{s}(U) \wedge \mathbf{t}(U)$$

For every formula  $\rho$  over  $A \times A$  define  $\bar{\rho}$  to be the following

$$a: A, a': A \mid \forall_{U: \mathbb{P}(A \times A)} [\mathbf{eq}(U) \wedge \forall_{x:A} \forall_{x':A} (\rho(x, x') \Rightarrow (x, x') \in_A U) \Rightarrow (a, a') \in_A U]$$

It is an easy exercise in first order logic to check that  $\bar{\rho}$  is a  $P$ -equivalence relation over  $A \times A$ . Take any  $\sigma$  in  $P(A \times A)$  and consider  $\chi_{\sigma}: 1 \longrightarrow \mathbb{P}(A \times A)$ . Recall that  $\chi_{\sigma}$  has the property that  $(a, a') \in_A \chi_{\sigma} \dashv\vdash \sigma(a, a')$ , so in  $a: A, a': A$  it holds

$$\bar{\rho}(a, a') \vdash \mathbf{eq}(\chi_{\sigma}) \wedge \forall_{x:A} \forall_{x':A} (\rho(x, x') \Rightarrow \sigma(x, x')) \Rightarrow \sigma(a, a')$$

If  $\sigma$  is a  $P$ -equivalence relation over  $A$  then  $\mathbf{eq}(\chi_{\sigma})$  is a true sentence. If moreover  $\rho \leq \sigma$  also  $\forall_{x:A} \forall_{x':A} (\rho(x, x') \Rightarrow \sigma(x, x'))$  is a true sentence. Whence the sequent above reduces to  $\bar{\rho}(a, a') \vdash \sigma(a, a')$ , which proves the claim.  $\square$

**8.6 Lemma.** *The base of a tripos with effective quotients and comprehensive diagonals has coequalizers.*

*Proof.* Let  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  be a tripos with quotients. We shall employ the language introduced in 2.16. Take two arrows  $f, g: Y \rightarrow A$  in  $\mathcal{C}$  and define  $\rho$  in  $P(A \times A)$  to be

$$a: A, a': A \mid \exists_{y:Y} (f(y) =_A a \wedge g(y) =_A a')$$

By 8.5 there is the smallest  $P$ -equivalence relation  $\bar{\rho}$  over  $A$  that contains  $\rho$ . Consider the quotient  $q: A \rightarrow A/\bar{\rho}$ . It is clear that

$$y': Y \mid \exists_{y:Y} f(y) =_A f(y') \wedge g(y) =_A g(y')$$

is a true formula, that is to say  $\top_Y \leq P_{\langle f, g \rangle}(\rho)$ . So also  $\top_Y \leq P_{\langle f, g \rangle}(\bar{\rho})$  and by effectiveness of quotient  $\top_Y \leq P_{\langle qf, qg \rangle}(\delta_{A/\bar{\rho}})$ . So  $qf = qg$  as  $P$  has comprehensive diagonals.

Suppose now  $k: A \rightarrow Z$  is such that  $kf = kg$ , then in  $y: Y, a: A, a': A$  it holds

$$f(y) =_A a \wedge g(y) =_A a' \vdash kf(y) =_A k(a) \wedge kg(y) =_A k(a') \vdash k(a) =_A k(a')$$

Therefore  $\exists_{y:Y} (f(y) =_A a \wedge g(y) =_A a' \vdash k(a) =_A k(a'))$ . That is to say that  $\rho \leq P_{k \times k}(\delta_A)$ . By 8.5 also  $\bar{\rho} \leq P_{k \times k}(\delta_A)$ . By the universal property of quotients there is  $h: A/\bar{\rho} \rightarrow Z$  with  $hq = k$ .  $\square$

Every quasi-topos  $\mathcal{C}$  is the base of a strong tripos  $Stg_{\mathcal{C}}: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  where  $Stg_{\mathcal{C}}$  is the subposet of  $\text{Sub}_{\mathcal{C}}(A)$  on those subobjects represented by a strong monomorphism. These are called strong subobjects.

**8.7 Proposition.** *A quasi-topos  $\mathcal{C}$  is a topos if and only if  $Stg_{\mathcal{C}}$  satisfies (RUC).*

*Proof.* A quasi-topos  $\mathcal{C}$  is a topos if and only if it is balanced. By 4.8 a monic arrow in  $\mathcal{C}$  is  $Stg_{\mathcal{C}}$ -injective and by 7.26 epimorphisms are  $Stg_{\mathcal{C}}$ -surjective. The claim follows by 4.9.  $\square$

**8.8 Theorem.** *A doctrine  $P: \mathcal{C}^{\text{op}} \longrightarrow \mathbf{ISL}$  is a hyper tripos if and only if  $Q_P$  is a quasi-topos and  $\hat{P}$  is the doctrine of strong subobjects.*

*Proof.* Points (i) and (iii) of the definition of quasi-topos comes from 4.18 and A.5, while point (iv) comes from 7.30. Finite coproducts come directly from 7.24 and the existence of co-equalizers from 8.6. In particular the initial object in  $Q_P$  is the initial object of  $\mathcal{C}$  with the total relation. By 4.17 the doctrine  $\hat{P}: Q_P^{\text{op}} \longrightarrow \mathbf{ISL}$  is m-variational. Hence  $\hat{P}$  is the doctrine of strong monomorphisms of  $Q_P$  (7.27).

For the converse note that  $Stg_{\mathcal{C}}$  is a tripos, so is  $P$  first order (7.1) on a slice-wise weakly cartesian closed category (A.5) with a weak predicated classifier (7.30). Whence  $P$  is a tripos. Coproducts follows from 7.24.  $\square$

**8.9 Corollary.** *If  $P$  is a intensional hyper tripos then  $Q_P$  is a quasi-topos.*

*Proof.* The same as in 8.8 as comprehensive diagonals in  $P$  play no role in the proof.  $\square$

Every quasi-topos  $\mathcal{C}$  contains a full subcategory  $\mathbf{Crs}_{\mathcal{C}}$  which is a topos. Its objects are called **coarse** and are precisely those orthogonal to the class of arrows that are monic and epic. So an object  $A$  in a quasi-topos  $\mathcal{C}$  is coarse if for every morphism  $f: C \rightarrow B$  which is both monic and epic and every  $g: C \rightarrow A$  there is a unique  $t: B \rightarrow A$  such that  $g = tf$ .

The topos  $\mathbf{Crs}_{\mathcal{C}}$  is reflective and the reflector is build as follows. For any object  $A$  note that the diagonal  $\delta_A$  in  $\mathit{Stg}_{\mathcal{C}}(A \times A)$  is represented by the diagonal (being the diagonal a strong monomorphism). Its classifying arrow  $\chi_{\delta_A}$  factors as a strong monic followed by an epimorphism as

$$\begin{array}{ccc} A & \xrightarrow{\chi_{\delta_A}} & \mathbb{P}A \\ & \searrow \eta_A & \nearrow \{\!\!\{ \mathcal{A}_{\chi_{\delta_A}} \top_A \}\!\!\} \\ & & \mathcal{S}_A \end{array}$$

The object  $\mathcal{S}_A$  will be called *object of  $A$ -singletons* and we call  $\eta_A: A \rightarrow \mathcal{S}_A$  the *singleton arrow* of  $A$ . Note that  $\chi_{\delta_A}$  is monic, whence the singleton arrow of  $A$  is both epic and monic. Strong monic are strong comprehension arrows in  $\mathit{Stg}_{\mathcal{C}}$  so it is easy to see that every arrow  $f: A \rightarrow B$  determines a unique arrow  $\mathcal{S}_f: \mathcal{S}_A \rightarrow \mathcal{S}_B$  with  $\mathcal{S}_f i_A = i_B f$ . This determines a functor  $\mathcal{S}: \mathbf{Crs}_{\mathcal{C}} \rightarrow \mathcal{C}$  where  $\mathbf{Crs}_{\mathcal{C}}$  is the full subcategory of  $\mathcal{C}$  on coarse objects.

As shown in [Joh02b, prop 2.6.12] the functor  $\mathcal{S}$  is left adjoint to the inclusion of  $\mathbf{Crs}_{\mathcal{C}}$  into  $\mathcal{C}$  with singleton arrows as unite. So in particular an object  $A$  is coarse if and only if it is isomorphic to its own singletons, *i.e.* if  $A \simeq \mathcal{S}_A$ .

Recall from 3.9 the construction that maps an existential elementary doctrine  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  to the existential elementary doctrine  $P_F: \mathcal{E}\mathcal{F}_P^{\text{op}} \rightarrow \mathbf{ISL}$  satisfying (RUC).

**8.10 Proposition.** *Let  $\mathcal{C}$  be a quasi-topos. The change of base of  $\mathit{Stg}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  along the inclusion of  $\mathbf{Crs}_{\mathcal{C}}$  is equivalent to  $\text{Sub}_{\mathcal{C}}: \mathcal{E}\mathcal{F}_{\mathit{Stg}_{\mathcal{C}}}^{\text{op}} \rightarrow \mathbf{ISL}$ .*

*Proof.* Any arrow  $m$  in  $\mathcal{C}$  can be written as the composite  $m = se$  where  $s$  is strong monic and  $e$  is epic. If  $m$  is monic then  $e$  is monic too. So if  $m$  is monic in  $\mathbf{Crs}_{\mathcal{C}}$  then  $e$  is an isomorphism. So every monic in  $\mathbf{Crs}_{\mathcal{C}}$  so the change of base of  $\mathit{Stg}_{\mathcal{C}}$  along the inclusion of  $\mathbf{Crs}_{\mathcal{C}}$  into  $\mathcal{C}$  is  $\text{Sub}_{\mathbf{Crs}_{\mathcal{C}}}: \mathbf{Crs}_{\mathcal{C}}^{\text{op}} \rightarrow \mathbf{ISL}$ . So it suffices to show that  $\mathbf{Crs}_{\mathcal{C}}$  is equivalent to  $\mathcal{E}\mathcal{F}_{\mathit{Stg}_{\mathcal{C}}}$ .


We want to find a functor  $S': \mathcal{E}\mathcal{F}_{\mathit{Stg}_{\mathcal{Q}}} \rightarrow \mathbf{Crs}_{\mathcal{Q}}$  naturally inverse to the composition

$$\mathbf{Crs}_{\mathcal{C}} \hookrightarrow \mathcal{C} \xrightarrow{\Gamma_{\mathit{Stg}_{\mathcal{C}}}} \mathcal{E}\mathcal{F}_{\mathit{Stg}_{\mathcal{C}}}$$

where  $\Gamma_{\mathit{Stg}_{\mathcal{C}}}$  acts as the identity on objects and maps  $f: A \rightarrow B$  to the formula  $a: A, b: B \mid f(a) =_B b$  in  $\mathit{Stg}_{\mathcal{C}}(A \times B)$ . On the object we define  $S'$  as the action of the reflector  $S: \mathcal{Q} \rightarrow \mathbf{Crs}_{\mathcal{Q}}$  in lemma 8.10 *i.e.* for any object  $A$  we put  $S'A = \mathcal{S}_A$ . To define the inverse on morphisms take a total and single-valued relation  $F \in \mathit{Stg}_{\mathcal{Q}}(A \times B)$ , this is a strong monic  $F = \langle F_1, F_2 \rangle: X \rightarrow A \times B$  in  $\mathcal{Q}$  where  $F_1: X \rightarrow A$  is monic and epic. Thus  $SF_1: SX \rightarrow \mathcal{S}_A$  is an isomorphism. Finally define  $S'F = SF_2(SF_1)^{-1}$ .  $\square$

**8.11 Remark.** Note that by 8.7 and 8.10 a quasi topos  $\mathcal{C}$  is a topos if and only if  $Stg_{\mathcal{C}}$  is  $Sub_{\mathcal{C}}$ . So 8.8 establishes that a category of the form  $\mathcal{C}_{ex/lex}$  is a topos if and only if  $\Psi_{\mathcal{C}}$  is a hyper tripos. But  $\Psi_{\mathcal{C}}$  is a hyper tripos precisely when  $\mathcal{C}$  is slice-wise cartesian closed and has a weak proof classifier. So Menni's theorem in [Men03] is an instance of 8.8 for  $P = \Psi_{\mathcal{C}}$ . A category with finite products, weak pullbacks, which are slice-wise cartesian closed and have a weak proof classifier form the category  $\mathbf{M}$ . It is easily seen that  $\mathbf{M}$  is equivalent to the full subcategory of  $\mathbf{HTr}$  of those doctrines satisfying (RC).

Theorem 8.10 explains the known adjunction between categories of quasi-topos and categories of toposes in terms of universal constructions of doctrines. In other words we have the following refinement of diagram (3).



where the inclusion of  $\mathbf{QT}$  into  $\mathbf{QRWFS}$  is obtained sending a quasitopos  $\mathcal{C}$  to the factorization system whose right class is that of strong subobjects.

Denote by  $\mathcal{T}_P$  the topos that comes from the tripos  $P$  under the tripos to topos construction. If  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  is a hyper tripos (intensional hyper tripos), then  $\mathcal{T}_P \equiv \mathcal{EF}_{\hat{P}}$  (this is theorem 3.5 in [MPR17]). Thus for every hyper tripos  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  the topos  $\mathcal{T}_P$  is the topos of coarse objects of  $Q_P$ .

Note that from the proof of theorem 8.8 that comprehensive diagonals are not necessary to get a quasi-topos out of the elementary quotient completion.

Recall from [MPR19] that the base of  $P$  has a natural number objects if and only if  $Q_P$  has a natural number object. We call *arithmetic* those quasi-toposes that have a natural number object.

**8.12 Proposition.** *If  $P$  is a intensional hyper tripos (with a natural numbers object) then  $Q_P$  is a (arithmetic) quasi-topos.*

**8.13 Proposition.** *If  $P$  is a intensional hyper tripos then  $P_x$  is a hyper tripos.*

*Proof.* From theorem 8.12 we know that  $Q_P$  is a quasi-topos and it is the base category of  $\widehat{P}$ . From theorem 4.23 we know that  $\widehat{P}$  is equivalent to  $\widehat{P}_x$  and in particular its base category and hence from theorem 8.8 we conclude that  $P_x$  is hyper tripos.  $\square$

## 9 Applications

Suppose  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  is a intensional hyper tripos. The category  $Q_P$  is a quasi-topos by 8.9. Its reflective subcategory on coarse objects is the topos obtained from  $P$  via the tripos-to-topos construction. By 4.12 and 4.11 the category  $Q_P$  is also a full reflective subcategory of  $Q_{\Psi_C}$ . We have the picture.

$$Q_{\Psi_C} \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} Q_P \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} \mathcal{T}_P \quad (5)$$

We already know that  $Q_{\Psi_C}$  is equivalent to  $\mathcal{C}_{\text{ex/lex}}$  and that  $\mathcal{T}_P$  is  $\text{Crs}_{Q_P}$ . By 4.12 the intensional hyper tripos  $P$  generates a topology  $j_P$  on  $\Psi_C$  whose extension  $\widehat{j_P}$  is a topology on  $\text{Sub}_{\mathcal{C}_{\text{ex/lex}}}$ . An we have that  $Q$  is equivalent to  $\mathbf{Sep}(\widehat{j_P})$ . So the previous picture can be equivalently described as follows.

$$\begin{array}{ccc} Q_{\Psi_C} & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} & Q_P & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} & \mathcal{T}_P \\ \parallel & & \parallel & & \parallel \\ \mathcal{C}_{\text{ex/lex}} & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} & \mathbf{Sep}(\widehat{j_P}) & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} & \text{Crs}_{Q_P} \end{array}$$

We now instantiate this picture on different choices of  $P$ .

Recall triposes of the form  $\mathcal{P}_{\mathcal{H}}$  as 2.14-(b). As observed in 4.12 these tripos have comprehension but this is not full. We can then consider the completion  $\mathcal{P}_{\mathcal{H}_c}: \text{Set}_c^{\text{op}} \rightarrow \mathbf{ISL}$  described in 4.12. Note that  $\text{Set}_c$  is Goguen's category  $\mathbf{Fuz}(\mathcal{H})$  of  $\mathcal{H}$ -valued fuzzy sets [Gog74]. Whence a quasitopos and therefore slice-wise cartesian closed with finite coproducts. It can be equivalently described as the category  $\mathcal{H}_+$  obtained by freely adding coproducts to  $\mathcal{H}$  [Men03]. The ex/lex completion of  $\mathcal{H}_+$  is the topos  $\mathbf{PreShv}(\mathcal{H})$  of presheaves of  $\mathcal{H}$ , while the tripos to tops construction applied to  $P_c$  gives the topos  $\mathbf{Shv}(\mathcal{H})$  of sheaves over  $\mathcal{H}$ . Thus when  $P$  is  $\mathcal{P}_{\mathcal{H}}$ , picture becomes

$$\mathbf{PreShv}(\mathcal{H}) \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} \mathbf{Sep}(\widehat{j_{\mathcal{P}_{\mathcal{H}_c}}}) \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \perp \rightarrow \\ \xrightarrow{\quad} \end{array} \mathbf{Shv}(\mathcal{H})$$

The change of base of triposes of the form  $\mathcal{P}_{\mathcal{H}}$  along the forgetful functor  $\mathbf{Top} \rightarrow \text{Set}$  is again a tripos as it suffices to endow  $\mathcal{H}^A$  with the indiscrete topology to have the power objects (see [Pas18]). In the special case of  $\mathcal{H} = \{0, 1\}$ , the tripos  $\mathcal{P}_{\mathcal{H}}$  reduces to the contravariant powerset functor  $\mathcal{P}: \text{Set}^{\text{op}} \rightarrow \mathbf{ISL}$  and its change of base along  $\mathbf{Top} \rightarrow \text{Set}$  is a tripos that we call  $\mathcal{T}$ . The tripos  $\mathcal{T}$  has full strong comprehension given by subspace topologies. Top has coproducts and is slice-wise weakly cartesian closed [CR00]. So  $\mathcal{T}: \mathbf{Top}^{\text{op}} \rightarrow \mathbf{ISL}$  is a hyper tripos. The category of generalised equilogical spaces  $\mathbf{Gequ}$  is equivalent to the base  $Q_{\mathcal{P}}$  of the elementary quotient completion of  $\mathcal{P}$ . Since  $\mathcal{T}$

is boolean the topology  $j_T$  is the double negation, whence also  $\widehat{j_T}$ . So picture 5 becomes.

$$\mathbf{Top}_{\text{ex/lex}} \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{\quad} \end{array} \mathbf{Gequ} \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{\quad} \end{array} \mathbf{Set}$$

As a byproduct we have that  $\mathbf{Gequ}$  is the category of  $\neg\neg$ -separated objects of  $\mathbf{Top}_{\text{ex/lex}}$  as shown in [Ros00].

The category  $\mathbf{Asm}$  of assemblies has as objects are pair  $(A, \alpha)$  where  $A$  is a set  $\alpha: A \rightarrow \mathbb{P}\mathbb{N}$  is a function from  $A$  to non-empty subsets of natural numbers. An arrow  $f: (A, \alpha) \rightarrow (B, \beta)$  is a function  $f: A \rightarrow B$  such that there is  $n \in \mathbb{N}$  such that for all  $a$  in  $A$  and all  $p$  in  $\alpha(a)$  the  $r.p$  is defined and it belongs to  $\beta(f(a))$ . The category  $\mathbf{Pasm}$  of partitioned assemblies is the full subcategory of  $\mathbf{Asm}$  on those  $(A, \alpha)$  such that each  $\alpha(a)$  is a singleton, *i.e.*  $\alpha$  can be seen as a function from  $A$  to  $\mathbb{N}$ . The change of base of  $\mathcal{P}$  along the forgetful functor  $\mathbf{Pasm} \rightarrow \mathbf{Set}$  is a tripos as it suffice to chose has weak power object of  $(A, \alpha)$  the partitioned assembly  $(\mathcal{P}A, k_0)$  where  $k_0$  is the constant map to 0. We call such a tripos  $\mathcal{R}: \mathbf{Pasm}^{\text{op}} \rightarrow \mathbf{ISL}$ . It is easy to see that  $\mathcal{R}$  has full strong comprehension where for a partitioned assembly  $(A, \alpha)$  and a subset  $X \subseteq A$  the inclusion of  $X$  into  $A$  determines a morphism of partitioned assembly  $\{\!\{X}\!\}: (X, \alpha|_X) \rightarrow (A, \alpha)$  which is the desired comprehension arrow. Since  $\mathbf{Pasm}$  is slice-wise cartesian closed, the tripos  $\mathcal{R}$  is an hyper tripos. Whence  $Q_{\mathcal{R}}$  is a quasi-topos by 8.8. Recall that  $\mathbf{Pasm}_{\text{ex/lex}}$  is  $\mathcal{E}ff$ . And the quasitopos of  $\neg\neg$ -separated objects of  $\mathcal{E}ff$  is  $\mathbf{Asm}$ . So picture 5 becomes

$$\mathcal{E}ff \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{\quad} \end{array} \mathbf{Asm} \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{\quad} \end{array} \mathbf{Set}$$

As byproduct we have that  $Q_{\mathcal{R}}$  is equivalent to  $\mathbf{Asm}$ . A different proof of this is given in [MPR19].

Another remarkable example is in type theory with the construction of the so called setoid models over Coquand-Huet's Calculus of Inductive Constructions CoC [Coq90]. The setoid model of functional relations over CoC is the tripos-to-topos construction  $\mathcal{T}_{F^{\text{CoC}}}$  with  $F^{\text{CoC}}$  the doctrine of propositions over CoC mentioned in [MR13b]. The topos.  $\mathcal{T}_{F^{\text{CoC}}}$  coincides with the topos of coarse objects within the quasi-topos  $Q_{F^{\text{CoC}}}$  which was one of the inspiring examples to the introduction of the elementary quotient completion in [MR13b].

**9.1 Theorem** (Topos elementary quotient completion). *Let  $P: \mathcal{C}^{\text{op}} \rightarrow \mathbf{ISL}$  be a hyper tripos. Then the following are equivalent:*

1.  $P$  satisfies (RC).
2.  $Q_P$  is a topos and coincides with the base of the regular completion of  $P$  and with the exact completion of the base of  $P$ .

*Proof.* It follows from 8.7 and 8.9. □

Examples of toposes arising in this way are those obtained as exact completions of a category  $\mathcal{C}$  by taking  $\Psi_{\mathcal{C}}$  for  $P$ .

**9.2 Theorem** (tripos-to-topos elementary quotient completion). *Let  $P: \mathcal{C}^{op} \rightarrow \mathbf{ISL}$  be a tripos. Then the following are equivalent:*

1.  $P_c$  satisfies (RC).
2.  $Q_{P_c}$  is a topos and coincides with the tripos-to-topos construction  $\tau_P$  of the tripos  $P$ .

*Proof.* It follows from theorem 5.5 in [MPR17] that  $P_c$  satisfies the rule of choice iff  $P_{c_x}$  satisfies the rule of choice. We also know that  $P_{c_x}$  satisfies the rule of choice iff  $Q_{P_{c_x}}$  is a topos and coincides with the exact completion of the base of  $P_{c_x}$  by theorem 9.1. Moreover  $Q_{P_{c_x}}$  is the exact completion of the base of  $P_{c_x}$  iff  $Q_{P_{c_x}}$  is equivalent to  $\tau_P$  by corollary 6.3 in [MPR17]. From [MR13a] we know that  $Q_{P_{c_x}}$  is equivalent to  $Q_{P_c}$  and this concludes the proof.  $\square$

## 10 Conclusions

We have introduced the notion of *quasi-topos completion* of an *hypertripos* by employing the machinery of the elementary quotient completion introduced in [MR13b], [MR13a].

In doing so we have generalized three theorems regarding exact completions by adopting the approach of elementary quotient completions: Carboni-Vitali's characterization of exact completions, Carboni-Rosolini's characterization of locally cartesian closed exact completions of a category with finite products and weak pullbacks, and Menni's characterization of topoi as exact completions.

In the future we aim to both generalize our quasi-topos completion further to include model of predicative logical theories including the extensional level of the Minimalist Foundation in [Mai09].

## References

- [BBS04] A. Bauer, L. Birkedal, and D.S. Scott. Equilogical spaces. *Theoret. Comput. Sci.*, 315(1):35–59, 2004.
- [BW84] M. Barr and C. Wells. *Toposes Triples and Theories*. Springer-Verlag, 1984.
- [Car95] A. Carboni. Some free constructions in realizability and proof theory. *J. Pure Appl. Algebra*, 103:117–148, 1995.
- [CC82] A. Carboni and R. Celia Magno. The free exact category on a left exact one. *J. Aust. Math. Soc.*, 33(A):295–301, 1982.
- [Coq90] T. Coquand. Metamathematical investigation of a calculus of constructions. In P. Odifreddi, editor, *Logic in Computer Science*, pages 91–122. Academic Press, 1990.
- [CR00] A. Carboni and G. Rosolini. Locally Cartesian closed exact completions. *J. Pure Appl. Algebra*, 154(1-3):103–116, 2000.

- [CV98] A. Carboni and E.M. Vitale. Regular and exact completions. *J. Pure Appl. Algebra*, 125:79–117, 1998.
- [Emm20] J. Emmenegger. On the local cartesian closure of exact completions. *J. Pure Appl. Algebra*, 224(11):106414, 25, 2020.
- [EPR20] J. Emmenegger, F. Pasquali, and G. Rosolini. Elementary doctrines as coalgebras. *J. Pure Appl. Algebra*, 224(12):106445, 16, 2020.
- [FK72] P.J. Freyd and G.M. Kelly. Categories of continuous functors I. *J. Pure Appl. Algebra*, 2:169–191, 1972.
- [Fre15] J. Frey. Triposes, q-toposes and toposes. *Ann. Pure Appl. Logic*, 166(2):232 – 259, 2015.
- [FS90] P.J. Freyd and A. Scedrov. *Categories, Allegories*. ISSN. Elsevier Science, 1990.
- [Gog74] J. A. Goguen. Concept representation in natural and artificial languages: Axioms, extensions and applications for fuzzy sets. *International Journal of Man-Machine Studies*, 6(5):513 – 561, 1974.
- [Gra00] M. Grandis. Weak subobjects and the epi-monic completion of a category. *J. Pure Appl. Algebra*, 154(1-3):193–212, 2000.
- [HJ03] J. Hughes and B. Jacobs. Factorization systems and fibrations: Toward a fibred Birkhoff variety theorem. *Electron. Notes Theor. Comput. Sci.*, 69:156–182, 2003.
- [HJP80] J.M.E. Hyland, P.T. Johnstone, and A.M. Pitts. Triples Theory. *Math. Proc. Camb. Phil. Soc.*, 88:205–232, 1980.
- [Hyl82] J.M.E. Hyland. The effective topos. In A.S. Troelstra and D. van Dalen, editors, *The L.E.J. Brouwer Centenary Symposium*, pages 165–216. North Holland, 1982.
- [Jac99] B. Jacobs. *Categorical Logic and Type Theory*, volume 141 of *Studies in Logic and the foundations of mathematics*. North Holland, 1999.
- [JM95] A. Joyal and I. Moerdijk. *Algebraic Set Theory*, volume 220 of *London Math. Soc. Lecture Note Ser.* Cambridge University Press, 1995.
- [Joh02a] P.T. Johnstone. *Sketches of an elephant: a topos theory compendium. Vol. 1*, volume 43 of *Oxford Logic Guides*. The Clarendon Press Oxford University Press, 2002.
- [Joh02b] P.T. Johnstone. *Sketches of an elephant: a topos theory compendium. Vol. 2*, volume 44 of *Oxford Logic Guides*. The Clarendon Press Oxford University Press, 2002.

- [Law69] F. W. Lawvere. Adjointness in foundations. *Dialectica*, 23:281–296, 1969.
- [Law70] F.W. Lawvere. Equality in hyperdoctrines and comprehension schema as an adjoint functor. In A. Heller, editor, *Proc. New York Symposium on Application of Categorical Algebra*, pages 1–14. Amer.Math.Soc., 1970.
- [Mai09] M.E. Maietti. A minimalist two-level foundation for constructive mathematics. *Ann. Pure Appl. Logic*, 160(3):319–354, 2009.
- [Men01] M. Menni. Closure operators in exact completions. *Theory Appl. Categ.*, 8:522–540, 2001.
- [Men03] M. Menni. A characterization of the left exact categories whose exact completions are toposes. *J. Pure Appl. Algebra*, 177(3):287–301, 2003.
- [MM92] S. Mac Lane and I. Moerdijk. *Sheaves in Geometry and Logic*. Springer-Verlag, 1992.
- [MPR17] M.E. Maietti, F. Pasquali, and G. Rosolini. Tripases, exact completions, and Hilbert’s  $\epsilon$ -operator. *Tbilisi Math. J.*, 10(3):141–166, December 2017.
- [MPR19] M. E. Maietti, F. Pasquali, and G. Rosolini. Elementary Quotient Completions, Church’s Thesis, and Partitioned Assemblies. *Logical Methods in Computer Science*, Volume 15, Issue 2, June 2019.
- [MR13a] M.E. Maietti and G. Rosolini. Elementary quotient completion. *Theory Appl. Categ.*, 27:445–463, 2013.
- [MR13b] M.E. Maietti and G. Rosolini. Quotient completion for the foundation of constructive mathematics. *Log. Univers.*, 7(3):371–402, 2013.
- [MR16] M.E. Maietti and G. Rosolini. Relating quotient completions via categorical logic. In Dieter Probst and Peter Schuster (eds.), editors, *Concepts of Proof in Mathematics, Philosophy, and Computer Science*, pages 229–250. De Gruyter, 2016.
- [Pal19] E. Palmgren. From type theory to setoids and back. <https://arxiv.org/abs/1804.08585>, 2019.
- [Pas15] F. Pasquali. A co-free construction for elementary doctrines. *Appl. Categ. Structures*, 23(1):29–41, Feb 2015.
- [Pas16] F. Pasquali. Remarks on the tripos to topos construction: Comprehension, extensionality, quotients and functional-completeness. *Appl. Categ. Structures*, 24(2):105–119, Apr 2016.
- [Pas18] F. Pasquali. On a generalization of equilogical spaces. *Logica Universalis*, 12(1):129–140, May 2018.

- [Pit02] A.M. Pitts. Tripos theory in retrospect. *Math. Structures Comput. Sci.*, 12(3):265–279, 2002.
- [Ros00] G. Rosolini. Equilogical spaces and filter spaces. *Rend. Circ. Mat. Palermo (2) Suppl.*, 64:157–175, 2000. Categorical studies in Italy (Perugia, 1997).
- [Sco76] D.S. Scott. Data types as lattices. *SIAM J. Comput.*, 5(3):522–587, 1976.
- [Sco96] D.S. Scott. A new category? Domains, spaces and equivalence relations. Available at <http://www.cs.cmu.edu/Groups/LTC/>, 1996.
- [vO08] J. van Oosten. *Realizability: An Introduction to its Categorical Side*, volume 152 of *Studies in Logic and the Foundations of Mathematics*. North Holland, 2008.
- [Wyl91] O. Wyler. *Lecture notes on Topoi and Quasitopoi*. World Scientific, 1991.

## A Another proof of the local cartesian closure of $Q_P$

In this appendix we give a different proof of the characterisation in A.5. The necessary condition is given by Lemma A.1, that we recall below for readability sake.

**A.1 Lemma.** *Suppose  $P: C^{op} \rightarrow \mathbf{ISL}$  is a variational doctrine. If  $Q_P$  is locally cartesian closed, then  $C$  is slice-wise weakly cartesian closed.*

We now aim at proving the converse of A.1. To this purpose the next two instrumental lemmas play a key role.

Suppose  $P$  is an elementary doctrine on  $C$  with full weak comprehension. Let  $f: B \rightarrow A$  represent an arrow  $[f]: (B, \sigma) \rightarrow (A, \rho)$  in  $Q_P$ . Let  $f_\rho: X \rightarrow A$  and  $f^\sigma: X \rightarrow B$  be the following two compositions

$$\begin{array}{ccc}
 X & \xrightarrow{\llbracket P_{f \times \text{id}_A}(\rho) \rrbracket} & B \times A \\
 & \searrow f_\rho & \downarrow \text{pr}_2 \\
 & & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 X & \xrightarrow{\llbracket P_{f \times \text{id}_A}(\rho) \rrbracket} & B \times A \\
 & \searrow f^\sigma & \downarrow \text{pr}_1 \\
 & & B
 \end{array}$$

Note that  $\rho \otimes_f \sigma = P_{f_\rho \times f^\sigma}(\rho) \wedge P_{f^\sigma \times f_\rho}(\sigma)$  is a  $P$ -equivalence on  $X$ . Denote by  $[f_\rho]$  the following arrow of  $Q_P$

$$[f_\rho]: (X, \rho \otimes_f \sigma) \rightarrow (A, \rho)$$

**A.2 Lemma.** *For every  $[f]: (B, \sigma) \rightarrow (A, \rho)$  in  $Q_P$  and every representative  $f: B \rightarrow A$ , the following hold:*

1.  $f$  factors through  $f_\rho$  in  $C$

$$2. \top_X = P_{\langle f f^\sigma, f_\rho \rangle}(\rho)$$

*Proof.* 1. Consider  $\langle \text{id}_B, f \rangle: B \rightarrow B \times A$ . It is  $P_{\langle \text{id}_B, f \rangle} P_{f \times \text{id}_A}(\rho) = P_{\langle f, f \rangle}(\rho) = \top_B$ . Weak universality of comprehension ensures the existence of  $k: B \rightarrow X$  with  $\llbracket P_{f \times \text{id}_A}(\rho) \rrbracket k = \langle \text{id}_B, f \rangle$  hence  $f = \text{pr}_2 \langle \text{id}_B, f \rangle = \text{pr}_2 \llbracket P_{f \times \text{id}_A}(\rho) \rrbracket k = f_\rho k$ . 2. Consider  $f^\sigma: X \rightarrow B$ . It is

$$\begin{aligned} P_{\langle f f^\sigma, f_\rho \rangle}(\rho) &= P_{\langle f \text{pr}_1 \llbracket P_{f \times \text{id}_A}(\rho) \rrbracket, \text{pr}_2 \llbracket P_{f \times \text{id}_A}(\rho) \rrbracket \rangle}(\rho) \\ &= P_{\llbracket P_{f \times \text{id}_A}(\rho) \rrbracket} P_{f \times \text{id}_A}(\rho) \\ &= \top_X \end{aligned}$$

where the last equality holds by definition of comprehension arrow.  $\square$

Note that in the proof of A.2 we obtain  $k$  such that  $f_\rho k = f$  and  $f^\sigma k = \text{id}_B$ . This allows to prove the following.

**A.3 Corollary.**  $[f]$  is isomorphic to  $[f_\rho]$  in  $\mathbf{Q}_P/(A, \rho)$ .

*Proof.* Consider  $k$  obtained in the proof of A.2. One immediately has  $[f^\sigma][k] = [\text{id}_B]$ . For the other composition it is (recall that  $f_\rho k = f$ )

$$P_{\langle k f^\sigma, \text{id}_X \rangle}(P_{f_\rho \times f_\rho}(\rho)) = P_{\langle f_\rho k f^\sigma, f_\rho \rangle}(\rho) = P_{\langle f f^\sigma, f_\rho \rangle}(\rho) = \top_X$$

where the last equality holds by A.2. Moreover

$$P_{\langle k f^\sigma, \text{id}_X \rangle}(P_{f^\sigma \times f^\sigma}(\sigma)) = P_{\langle f^\sigma k f^\sigma, f^\sigma \rangle}(\sigma) = P_{\langle f^\sigma, f^\sigma \rangle}(\sigma) = \top_X$$

whence  $\top_X = P_{\langle k f^\sigma, \text{id}_X \rangle}(\rho \otimes_f \sigma)$ , i.e.  $[k][f^\sigma] = [\text{id}_X]$ .  $\square$

Note that, although  $[f]$  and  $[f_\rho]$  are isomorphic in  $\mathbf{Q}_P/(A, \rho)$ , they need not factor through each other in  $\mathcal{C}$ . Lemma A.2 shows that  $f$  factors through  $f_\rho$  in  $\mathcal{C}$ , but the other factorization need not occur.

Suppose  $P$  is elementary existential doctrine on  $\mathcal{C}$  with comprehensive diagonals and full weak comprehension. Then  $\mathbf{Q}_P$  has pullbacks. Let  $f: B \rightarrow A$  in  $\mathcal{C}$  represents  $[f]: (B, \sigma) \rightarrow (A, \rho)$  in  $\mathbf{Q}_P$ . Suppose  $q: W \rightarrow A$  in  $\mathcal{C}$  and let  $\theta$  be a  $P$ -equivalence relation over  $W$  such that  $\theta \leq P_{q \times q}(\rho)$ . Consider the following three diagrams

$$\begin{array}{ccccc} Z & \xrightarrow{t_W} & W & & (Z, P_{t_X \times t_X}(\rho \otimes_f \sigma) \wedge P_{t_W \times t_W}(\theta)) & \xrightarrow{[t_W]} & (W, \theta) \\ t_X \downarrow & & \downarrow q & & \downarrow [t_X] & & \downarrow [q] \\ X & \xrightarrow{f_\rho} & A & & (X, \rho \otimes_f \sigma) & \xrightarrow{[f_\rho]} & (A, \rho) \end{array}$$

$$\begin{array}{ccc} (T, \eta) & \xrightarrow{[e]} & (X \times W, P_{f_\rho \times f_\rho}(\rho) \boxtimes \theta) & \xrightarrow{[\pi_W]} & (W, \theta) \\ & \searrow [ \pi_X ] [e] & (C) & & \downarrow [q] \\ & & (X, \rho \otimes_f \sigma) & \xrightarrow{[f_\rho]} & (A, \rho) \end{array}$$

where  $e = \llbracket P_{f_\rho \times q}(\rho) \rrbracket: T \rightarrow X \times W$  and  $\eta = P_{e \times e}(P_{f_\rho \times f_\rho}(\rho) \boxtimes \theta)$  and where we denote by  $\pi_X$  and  $\pi_W$  the projections from  $X \times W$  to distinguish them from  $\text{pr}_1$  and  $\text{pr}_2$  which are projections from  $B \times A$ . The diagram (C) is the pullback in  $\mathcal{Q}_P$  of  $[q]$  along  $[f_\rho]$ .

If the square (A) commutes, so does (B). Since (C) is a pullback, there is

$$[j]: (Z, P_{t_X \times t_X} P_{f_\rho \times f_\rho}(\rho) \boxtimes P_{t_W \times t_W}(\theta)) \rightarrow (T, \eta) \quad (6)$$

with  $[t_W] = [\pi_W][e][j]$  and  $[t_X] = [\pi_X][e][j]$ .

Consider  $(f^\sigma \times q)e: T \rightarrow B \times A$  and note that

$$\begin{aligned} P_e P_{f^\sigma \times q}(P_{f \times \text{id}_A}(\rho)) &= P_e P_{f f^\sigma \times q}(\rho) \\ &= P_e P_{f_\rho \times q}(\rho) \quad (\text{by A.2-2}) \\ &= \top_T \end{aligned}$$

where the last equality comes from the definition of  $e = \llbracket P_{f_\rho \times q}(\rho) \rrbracket$ . Weak universality of  $\llbracket P_{f \times \text{id}_A}(\rho) \rrbracket: X \rightarrow B \times A$  produces an arrow

$$k: T \rightarrow X \quad (7)$$

with  $\llbracket P_{f \times \text{id}_A}(\rho) \rrbracket k = (f^\sigma \times q)e$ . Now observe that

$$q\pi_W e = \text{pr}_2(f^\sigma \times q)e = \text{pr}_2 \llbracket P_{f \times \text{id}_A}(\rho) \rrbracket k = f_\rho k$$

Therefore if the diagram (A) is a pullback, there is an arrow  $j': T \rightarrow Z$  with  $t_W j' = \pi_W e$  and  $t_X j' = k$ . Note that from  $q\pi_W e = f_\rho k$  and  $[f_\rho][\pi_X][e] = [q][\pi_W][e]$ , i.e.  $\top_T = P_{\langle f_\rho \pi_X e, q\pi_W e \rangle}(\rho)$  one has

$$\top_T = P_{\langle q\pi_W e, f_\rho k \rangle}(\rho) = P_{\langle f_\rho \pi_X e, f_\rho k \rangle}(\rho) = P_{\langle \pi_X e, k \rangle}(P_{f_\rho \times f_\rho}(\rho)) \quad (8)$$

and also

$$[f_\rho][\pi_X][e] = [q][\pi_W][e] = [f_\rho][k] = [f_\rho][t_X][j'] \quad (9)$$

From (9) and  $[t_W][j'] = [\pi_W][e]$  we get

$$\eta \leq P_{j \times j'}[P_{t_X \times t_X} P_{f_\rho \times f_\rho}(\rho) \boxtimes P_{t_W \times t_W}(\theta)]$$

which makes

$$[j']: (T, \eta) \rightarrow (Z, P_{t_X \times t_X} P_{f_\rho \times f_\rho}(\rho) \boxtimes P_{t_W \times t_W}(\theta)) \quad (10)$$

a legitimate arrow of  $\mathcal{Q}_P$ . From 8 one also has  $[k] = [\pi_X][e]$  which allows to prove

$$[\pi_X][e][j][j'] = [t_X][j'] = [k] = [\pi_X][e] \quad (11)$$

$$[\pi_W][e][j][j'] = [t_W][j'] = [\pi_W][e] \quad (12)$$

The following lemma will play a crucial role in proving A.5.

**A.4 Lemma.** *If the square (A) is a pullback, also the square (B) is a pullback.*

*Proof.* It suffices to prove that  $[j]$  as in (6) is the inverse of  $[j']$  as in (10). The universal property of pullbacks together with equations (11) and (12) establishes  $[j][j'] = \text{id}_{(T,\eta)}$ . To prove that also  $[j'][j]$  is the identity is to prove that

$$\begin{aligned}\top_Z &= P_{\langle j', \text{id}_Z \rangle} (P_{t_X \times t_X} (\rho \otimes_f \sigma) \wedge P_{t_W \times t_W} (\theta)) \\ &= P_{\langle j', \text{id}_Z \rangle} (P_{t_X \times t_X} (P_{f_\rho \times f_\rho} (\rho) \wedge P_{f^\sigma \wedge f^\sigma} (\sigma)) \wedge P_{t_W \times t_W} (\theta)) \\ &= P_{\langle f_\rho t_X j', f_\rho t_X \rangle} (\rho) \wedge P_{\langle f^\sigma t_X j', f^\sigma t_X \rangle} (\sigma) \wedge P_{\langle t_W j', t_W \rangle} (\theta)\end{aligned}$$

From the construction of the arrow (6) we know that  $[t_X] = [\pi_X][e][j]$  and  $[t_W] = [\pi_W][e][j]$  and hence

$$\begin{aligned}[f_\rho][t_X][j'][j] &= [f_\rho][\pi_X][e][j][j'][j] = [f_\rho][\pi_X][e][j] = [f_\rho][t_X] \\ [f^\sigma][t_X][j'][j] &= [f^\sigma][\pi_X][e][j][j'][j] = [f^\sigma][\pi_X][e][j] = [f^\sigma][t_X] \\ [t_W][j'][j] &= [\pi_W][e][j][j'][j] = [\pi_W][e][j] = [t_W]\end{aligned}$$

which means that  $\top_Z = P_{\langle f_\rho t_X j', f_\rho t_X \rangle} (\rho)$ ,  $\top_Z = P_{\langle f^\sigma t_X j', f^\sigma t_X \rangle} (\sigma)$  and  $\top_Z = P_{\langle t_W j', t_W \rangle} (\theta)$ , whence the claim.  $\square$

We are now ready to prove the main theorem of the section. To do this we first consider a universal implicational doctrine  $P$  with full weak comprehension. Suppose also the base  $\mathcal{C}$  to be weakly cartesian closed. Consider  $[f]: (B, \sigma) \rightarrow (A, \rho)$  and  $[g]: (B', \sigma') \rightarrow (A, \rho)$  in the slice  $\mathcal{Q}_P/(A, \rho)$ . Denote by  $q': W' \rightarrow A$  the weak exponential in  $\mathcal{C}$  of  $g_\rho: X' \rightarrow A$  to the power of  $f_\rho: X \rightarrow A$ . Let arrows  $t'_X: Z' \rightarrow X$  and  $t_{W'}: Z' \rightarrow W'$  be such that  $q' t_{W'} = f_\rho t'_X$  is a pullback and let  $e': Z' \rightarrow X'$  be a weak evaluation. Define  $\xi = \mathbb{V}_{\text{pr}_3}(\xi_1 \rightarrow \xi_2)$  in  $P(W')$  as

$$\begin{aligned}\xi_1 &= P_{\langle t_{W'}, \text{pr}_1, \text{pr}_3 \rangle} (\delta_{W'}) \wedge P_{\langle t_{W'}, \text{pr}_2, \text{pr}_3 \rangle} (\delta_{W'}) \\ \xi_2 &= P_{\langle t'_X, \text{pr}_1, t'_X, \text{pr}_2 \rangle} (\rho \otimes_f \sigma) \rightarrow P_{\langle e', \text{pr}_1, e', \text{pr}_2 \rangle} (\rho \otimes_g \sigma')\end{aligned}$$

where  $\text{pr}_1, \text{pr}_2, \text{pr}_3$  are projections from  $Z' \times Z' \times W'$ . Consider  $\{\xi\} = W \rightarrow W'$  and let

$$\begin{array}{ccc} Z & \xrightarrow{t_W} & W \\ t_X \downarrow & & \downarrow q \\ X & \xrightarrow{f_\rho} & A \end{array}$$

be a pullback where  $q = q'\{\xi\}$ . From  $f_\rho t_X = q\{\xi\} = q'\{\xi\}t_W$  there is an arrow  $w: Z \rightarrow Z'$  with  $t_{W'} w = \{\xi\}t_W$  and  $t'_X w = t_X$ . Define

$$e = e'w: Z \rightarrow X' \tag{13}$$

Finally consider the  $\theta$  in  $P(W \times W)$  defined as follows

$$\theta = P_{qt_W \times qt_W} (\rho) \wedge \mathbb{V}_{\langle \pi_2, \pi_4 \rangle} (\theta_1 \rightarrow \theta_2) \tag{14}$$

where

$$\theta_1 = P_{\langle t_W \pi_1, \pi_3 \rangle} (\delta_W) \wedge P_{\langle t_W \pi_2, \pi_4 \rangle} (\delta_W)$$

$$\theta_2 = P_{\langle t_X \pi_1, t_X \pi_2 \rangle}(\rho \otimes_f \sigma) \rightarrow P_{\langle e \pi_1, e \pi_2 \rangle}(\rho \otimes_g \sigma')$$

and where  $\pi_i$  with  $i = 1, 2, 3, 4$  are projections from  $Z \times Z \times W \times W$ . We leave as an exercise to show that  $\theta$  defines a  $P$ -equivalence relation over  $W$  such that

$$\theta \leq P_{e \times e}(\rho \otimes_g \sigma') \quad (15)$$

**A.5 Proposition.** *Suppose  $P$  is an elementary existential doctrine with comprehensive diagonals and full weak comprehension on a category  $\mathcal{C}$  with pullbacks.  $\mathcal{C}$  is weakly locally cartesian closed if and only if  $\mathcal{Q}_P$  is locally cartesian closed.*

*Proof.* The necessary direction is A.1. Conversely note that if  $\mathcal{C}$  is weakly locally cartesian closed,  $\Psi_{\mathcal{C}}$  is universal and implicational, hence also  $P$  is such by 4.13, as  $P$  is the doctrine of  $j$ -closed elements of  $\Psi_{\mathcal{C}}$  with respect to the canonical topology. For  $[f]:(B, \sigma) \rightarrow (A, \rho)$  and  $[g]:(B', \sigma') \rightarrow (A, \rho)$  in  $\mathcal{Q}_P$  consider  $e: Z \rightarrow X'$  defined as in (13) and  $\theta$  as in (14). The condition in (15) guarantees that  $e$  lifts to an arrow

$$[e]:(Z, \theta) \rightarrow (X', \rho \otimes_g \sigma')$$

in  $\mathcal{Q}_P$  which is also an arrow in  $\mathcal{Q}_P/(A, \rho)$  from  $[f_\rho]$  to  $[g_\rho]$ . Moreover A.4 ensures that  $[e]$  goes to the pullback of  $[g]$  along  $[f_\rho]$  to  $[g_\rho]$ . Hence, after A.3, to prove the claim it suffices to prove that  $[q]$  is the exponential in  $\mathcal{Q}_P/(A, \rho)$  of  $[g_\rho]$  and  $[f_\rho]$  with evaluation  $[e]$ .

Consider  $[h]:(C, \gamma) \rightarrow (A, \rho)$ . Suppose  $[m]$  is an arrow from the pullback of  $[h]$  along  $[f_\rho]$  to  $[g_\rho]$ . By A.4 we think of  $[m]$  as an arrow of the form

$$[m]:(Q, P_{l_X \times l_X} P_{f_\rho \times f_\rho}(\rho) \boxtimes P_{l_C \times l_C}(\gamma)) \rightarrow (X', \rho \otimes_g \sigma')$$

where

$$\begin{array}{ccc} Q & \xrightarrow{l_C} & C \\ l_X \downarrow & & \downarrow h \\ X & \xrightarrow{f_\rho} & A \end{array}$$

is the pullback in  $\mathcal{C}$  of  $h$  along  $f_\rho$ . Thus  $m: Q \rightarrow X'$  determines  $\bar{m}: C \rightarrow W'$  with  $e'(\text{id}_X \times_A \bar{m}) = m$  and  $q' \bar{m} = h$ . Using the universal property of comprehension is again an exercise to show that  $\bar{m} = \mu\{\xi\}$  for some  $\mu: X \rightarrow W$ . This  $\mu$  represents an arrow  $[\mu]:(C, \gamma) \rightarrow (W, \theta)$  which is the existential transpose of  $[m]$  in  $\mathcal{Q}_P/(A, \rho)$ . Uniqueness is a direct consequence of the definition of  $\theta$ .  $\square$