

Dark Energy from Quantum Uncertainty with P-symmetry

Long Huang,^{a,d} Xiaofeng Yang,^{a,b,c} Xiang Liu^{a,b,c}

^a*Xinjiang Astronomical Observatory, Chinese Academy of Sciences, Urumqi 830011, China*

^b*Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, Nanjing 210008, China*

^c*Key Laboratory of Radio Astrophysics in Xinjiang Province, Urumqi 830011, China*

^d*University of Chinese Academy of Sciences, Beijing, 10039, China*

ABSTRACT: We explore the theoretical possibility that dark energy density is derived from massless scalar bosons in vacuum and present a physical model for dark energy. By assuming massless scalar bosons fall into the horizon boundary of the cosmos with the expansion of the universe, we can deduce the uncertainty in the relative position of scalar bosons based on the quantum fluctuation of space-time and the assumption that scalar bosons satisfy P-symmetry under the parity transformation $P\varphi(r) = -\varphi(r)$, which can be used to estimate scalar bosons and dark energy density. Furthermore, we attempt to explain the origin of negative pressure from the increasing entropy density of the Boltzmann system and derive the equation for the state parameter, which is consistent with the specific equations of state for dark energy. Finally, we employ the SNIa Pantheon sample and Planck 2018 CMB angular power spectra to constrain the models and provide statistical results for the cosmology parameters.

Contents

1	Introduction	1
2	The quantum fluctuation of space-time and P-symmetry	2
3	The density and negative pressure for Boltzmann system	4
4	Dark energy density and pressure	5
4.1	The dark energy density and the evolution of ground state energy	5
4.2	The invariable ground state energy for the scalar bosons and energy density	7
5	The Used observation data	7
5.1	SNIa Pantheon sample and the Plank 2018 CMB angular power spectra	7
5.2	SALT2 calibration for Pantheon sample	7
6	Using the Pantheon sample to fit the models parameters	8
6.1	The fitting of model I parameters	8
6.2	The fitting of model II parameters	10
7	Using CMB angular power spectra to constrain the models parameters	11
8	CONCLUSIONS	11
9	Acknowledgments	13

1 Introduction

Since 1998, when Riess et al. used SNIa data to statistically indicate that the expansion of universe is accelerating [1], physicists have been providing various theories to explain this acceleration, including $f(R)$ theory [2], Brans-Dicke theory [3], and dark energy theory [4]. At present, the dark energy theory can be used to effectively explain the cosmic microwave background (CMB) anisotropies [5]. However, this study mainly focuses on the physical nature of dark energy. Dark energy can be studied using two main approaches. The first is to focus on the properties of dark energy, investigating whether or not its density evolves with time; this can be verified by reconstructing the equation of state $w(z)$ for dark energy, which is independent of physical models. The reconstruction of the equation of state involves parametric and non-parametric methods [6], the latter including the Principal Component Analysis [7, 8], Gaussian Processes [9, 10, 11], PCA with the smoothness prior method [12, 13, 14], and PCA based on the Ridge Regression Approach [15]. The second involves dark energy physical models that are presented from the physical origin of its density and

pressure, including scalar field models [16], pseudo-Nambu-Goldstone bosons for cosmology [17], holographic dark energy [18], age dark energy [19] and so on.

Currently, it may be difficult to judge which model, method or result is more persuasive, however, a model of dark energy that concerns its physical nature is essential. From the point of view of the models, Maziashvili presented a method that uses the Krolyhazy relation and time-energy uncertainty relation to estimate the density of dark energy [19, 20], and the result is consistent with astronomical data if the unique numerical parameter in the dark energy model is taken to be on the order of one.

Based on this, to further explore the origin of dark energy density and pressure, we present the possibility that dark energy density is derived from massless scalar bosons in a vacuum. If the scalar boson field is the radiation field that satisfies the Bose-Einstein distribution, positive pressure would be generated, we first exclude this possibility based on the negative pressure of dark energy. Therefore, we can deduce the uncertainty in the position of scalar bosons based on the quantum fluctuation of space-time and the assumption that scalar bosons satisfy P-symmetry under a parity transformation, which can be used to estimate the scalar bosons and dark energy density.

2 The quantum fluctuation of space-time and P-symmetry

The quantum fluctuation of space-time relates to the quantum properties of objects. Using the Heisenberg position-momentum uncertainty relations, Wigner derived a quantum limit on the measurability of a certain length [21]. If t is the time required by the measurement procedure, the uncertainty in the length measurement is

$$\delta t \sim \sqrt{\frac{t}{M_c}} \quad (2.1)$$

where M_c is the mass of the clock, we take the units $\hbar = c = 1$ throughout this study.

As a result of the above relation, when $M_c \rightarrow \infty$, $\delta t \rightarrow 0$. To solve this situation, the quantum fluctuations of space-time itself is presented [22, 23, 24, 25]. It also results in uncertainty in distance measurements. Thus, at very short distance scales, space-time is foamy, and the limitation of space-time distance measurements can be given by

$$\delta t \sim \begin{cases} (t_p t)^{\frac{1}{2}} & r > t_p \\ t_p & r \leq t_p \end{cases} \quad (2.2)$$

where t_p is the Planck time and $t_p = \sqrt{\hbar G}$. This limitation of space-time measurements can be interpreted as the result of quantum fluctuations of space-time. Meanwhile, Eq. (2.2) can be derived for massless particles in the framework of κ -deformed Poincare symmetries [26, 27].

Krolyhazy derived another method for describing the quantum fluctuations of space-time, know as the Krolyhazy relation

$$\delta t \sim t_p^{2/3} t^{1/3}. \quad (2.3)$$

We consider that this quantum fluctuation effect is also applicable to the expanding universe. If the scalar boson field is the radiation field that obeys the Bose-Einstein distribution, positive pressure will be generated, we first exclude this possibility based on the negative pressure of dark energy. By assuming massless scalar bosons fall into the horizon boundary of the cosmos, with the expansion of the universe, scalar bosons satisfy P-symmetry under parity transformation, which can be expressed as

$$P\varphi(r) = -\varphi(r), \quad (2.4)$$

where Ψ_0 is the wave function of the scalar bosons. If r, t are the horizon size and age of the universe, from P-symmetry and quantum fluctuation equation (2.2) or (2.3) of space-time, the former can correlate the uncertainty in the relative position of two scalar bosons with the horizon size or age of the universe; hence, the uncertainty in the position of scalar boson is

$$\delta t_{scalar} \sim \delta t. \quad (2.5)$$

We hypothesise that the mean distance between scalar bosons is no less than the uncertainty relation δt_{scalar} , as a result, we obtain the number density of massless scalar bosons

$$N = \delta t^{-3}. \quad (2.6)$$

If the quantum fluctuation of space-time provides scalar bosons with nonzero energy, we can consider the wave function Ψ_0 for the massless scalar bosons is non-stationary state. Its wave length can be determined by the uncertainty relation δt ; hence, Ψ_0 can be written as the superposition of at least n stationary states $\varphi_1, \varphi_2, \dots, \varphi_n$

$$\Psi_0 = \sum_k^n \alpha_k \varphi_k e^{-i\omega_k t'}, \quad (2.7)$$

where $\omega_k = 2\pi k/\delta t$. It is easy to see that the state components of $\varphi, e^{-i\omega t'}$ are orthogonal in space and time, respectively.

Alternatively, the wave length of scalar bosons can be determined by the age of universe t ; hence, the wave function of scalar bosons Ψ_0 can be written as

$$\Psi_0 = \sum_k^n \bar{\alpha}_k \bar{\varphi}_k e^{-i\bar{\omega}_k t'}, \quad (2.8)$$

where $\bar{\omega}_k = 2\pi k/t$. The wave function Ψ_0 will be further discussed using Higgs potential or Yukawa interaction.

By assuming scalar bosons or Goldstone bosons exist in a vacuum, Higgs proposed the existence of the Higgs scalar boson field which can be described by the field equation [31]

$$\nabla_\mu \nabla^\mu \Phi + \bar{V}'(\Phi\Phi^*)\Phi = 0, \quad (2.9)$$

and When

$$\bar{V}'(\Phi_0\Phi_0^*) = 0, \quad (2.10)$$

the massless-zero spin scalar boson field equation becomes

$$\square\Phi_0 = 0. \quad (2.11)$$

We consider this massless scalar field to contain zero-spin and zero-charge bosons, and the scalar boson couples or it undergoes Yukawa interaction with itself, which can be quantified in the simple potential energy form $-f^2\Phi_0\Phi_0^*$, resulting in their massless state. If the Higgs singlet potential Φ_0 is substituted with the Yukawa potential U , the massless scalar bosons can interact with each other through propagator bosons provided they have weak isospins, which can be described by the Yukawa potential. Using Eq. (2.7) or (2.8), we incorporate a separation of the variable U

$$U = U'(r)e^{-i(w_k-w_l)t}, \quad (2.12)$$

and the Yukawa potential $U'(r)$ satisfies

$$\{\Delta - (m_0^2 - w_{kl}^2)\} U' = -4\pi g\varphi_k\varphi_l, \quad (2.13)$$

where $w_{kl} = w_k - w_l$, m_0 is the mass of the Higgs or scalar boson, and g is the coupling constant. Solving the above equation yields

$$U'(r) = g \int \frac{e^{-\mu|r-r'|}}{|r-r'|} \varphi_k(r')\varphi_l(r') dr', \quad (2.14)$$

where $\mu = \sqrt{m_0^2 - w_{kl}^2}$.

3 The density and negative pressure for Boltzmann system

When all scalar bosons are in the ground state ε_0 , that is

$$\varepsilon_0 = \delta t^{-1} \text{ or } t^{-1}, \quad (3.1)$$

we can use Eq. (2.2) or (2.3), (2.6) and (3.1) to obtain the internal energy per unit volume for massless scalar bosons

$$u = \varepsilon_0 N \sim (t_p t)^{-2}. \quad (3.2)$$

Next we use the increase in entropy to derive the existence of negative pressure.

We consider the Boltzmann system to be composed of massless scalar bosons in vacuum; the bosons have a quantum state number W_p for an energy level at the ground state ε_0 , where W_p is considered the momentum degree of freedom. The momentum is $p_0 = \sqrt{\sum_{i=1}^{W_p} p_i^2}$. When $W_p = 3$, $p_0 = \sqrt{p_x^2 + p_y^2 + p_z^2}$, and it is clear that the microstate numbers of the Boltzmann system are 3^N ; hence, the system has statistical entropy per unit volume

$$s = k_B \ln 3^N, \quad (3.3)$$

where k_B is the Boltzmann constant.

Using the thermodynamic entropy definition $TdS = dU + pdV$, where T is temperature, and the equations $dS = sdV$, $k; and dU = udV$; with Eq. (2.2), (2.6), (3.2) and (3.3), we get

$$T = (\ln T_c)^{-1} k_B^{-1} (t_p t)^{-1/2}. \quad (3.4)$$

Because $u(t_p t)^2$ is constant, T_c is also constant.

We assume the degeneracy of energy W_p is increasing gradually with the expanding universe, which can be expressed as

$$W_p : 1 \rightarrow 2 \rightarrow 3, \quad (3.5)$$

Hence, with the rise in W_p , the entropy density will increase, which can generate the negative pressure. From Eq. (3.2) and (3.3) (3.5), it is easy to find that the negative pressure $p_u = \omega_u u$ satisfies

$$T \ln W_p^N = (1 + \omega_u)u, \quad (3.6)$$

solving above equation yields

$$\omega_u = -1 + \frac{\ln W_p}{\ln T_c}. \quad (3.7)$$

This is the equation of state for scalar field in the vacuum. It has an interesting feature. We set $T_c = 3$, when $W_p \rightarrow 1$, one has $\omega_u \rightarrow -1$, if $W_p \rightarrow 2$, $\omega_u \rightarrow -1/3$. As a result of above relation, it implies that the vacuum energy is from ordered to disordered state along with time, and it can't violate the principle of entropy increase. Namely, the negative pressure can be derived from the increasing entropy, which avoids the universe keeping expansion in the future.

4 Dark energy density and pressure

4.1 The dark energy density and the evolution of ground state energy

We assume that dark energy is composed of massless scalar boson in vacuum, using Eq. (3.2), and obtain the dark energy density $\rho_{de} = (t_p t)^{-2}$. We introduce a numerical constant to ensure the equals sign satisfies Eq. (2.2) or (2.3), and choose the conformal time η as the time scale t [28, 29]. Hence, the energy density can be written as

$$\rho_{de} = \frac{3n^2 M_p^2}{\eta^2}, \quad (4.1)$$

where $M_p^{-2} = 8\pi G$, n^2 is an introduced numerical constant, η is the conformal time, which is given by

$$\eta = \int_0^t \frac{dt}{a}. \quad (4.2)$$

where a is the scale factor of our universe. We take the present scale factor $a_0 = 1$, t is the age of universe.

Suppose the universe be spatially flat, define the fraction energy density of dark matter as $\Omega_m = \rho_m/3M_p^2 H^2$, and $\Omega_{de} = \rho_{de}/3M_p^2 H^2$, one has $\Omega_m = 1 - \Omega_{de}$ from the Friedmann equation. With the Eq. (4.1), Ω_{de} can be written

$$\Omega_{de} = \frac{n^2}{H^2 \eta^2}, \quad (4.3)$$

where $H = \dot{a}/a$ is the Hubble parameter. Using the energy conservation equation $\dot{\rho}_{de} + 3H(\rho_{de} + p_{de}) = 0$, with Eq. (4.1) (4.3), one can get the equation of state of energy density for $\omega_{de} = p_{de}/\rho_{de}$ as

$$\omega_{de} = -1 + \frac{2}{3n} \frac{\sqrt{\Omega_{de}}}{a}. \quad (4.4)$$

Meanwhile using the Friedmann equations with equation Eq. (4.4), as well as $\rho_m \propto a^{-3}$, one can get Ω_{de} that satisfies [29]

$$\Omega'_{de} = \frac{\Omega_{de}}{a} (1 - \Omega_{de}) \left(3 - \frac{2}{n} \frac{\sqrt{\Omega_{de}}}{a} \right), \quad (4.5)$$

where the prime represents the derivative with respect to $\ln a$.

It has some interesting features for the equation of state of the dark energy. In the dark energy dominated phase, the energy density can drive the universe to accelerated expansion if $\omega_{de} < -1/3$. From Eq. (4.4), it is easy to see that when $a \rightarrow \infty$, $\Omega_{de} \rightarrow 1$, thus $\omega_{de} \rightarrow -1$ at later time. As well, in the matter dominated epoch, it has $a \propto t^{2/3}$, from Eq. (4.1)(4.2), one has $\rho_{de} \propto a^{-1}$. Then from the dark energy density conservation equation, one can obtain $\omega_{de} = -2/3$.

For constant ω_{de} , the deceleration factor q_0 is given by $q_0 = 0.5 + 1.5(1 - \Omega_m)\omega_{de}$, we fix $\Omega_m = 0.3$ of the current universe which is from Λ CDM cosmology with SNIa Pantheon samples [34]. It is easy to see that when $\omega_{de} \lesssim -1/2$, $q_0 < 0$, which implies that the energy density can drive the universe to accelerated expansion if $\omega_{de} \lesssim -1/2$ for the current universe.

Meanwhile we also consider the case that the time scale t is replaced by the future event horizon R_h , which was proposed by Li [18], the future event horizon is

$$R_h = a \int_t^\infty \frac{dt}{a}, \quad (4.6)$$

so the dark energy density is

$$\rho_{de} = \frac{3n^2 M_p^2}{R_h^2}. \quad (4.7)$$

Then combining with the dark energy density conservation equation, the equation of state of energy density can be given by [18]

$$\omega_{de} = -\frac{1}{3} - \frac{2}{3n} \sqrt{\Omega_{de}}. \quad (4.8)$$

In the same way, at earlier time where $\Omega_{de} \rightarrow 0$, it has $\omega_{de} \rightarrow -1/3$, and at later time where $\Omega_{de} \rightarrow 1$, $\omega_{de} \rightarrow -1$ for $n = 1$.

4.2 The invariable ground state energy for the scalar bosons and energy density

If we also consider other case that the expected value of energy for scalar boson ε_c does not change with time, ε_c is constant, so Eq. (3.2) becomes

$$u = \varepsilon_c \delta t^{-3}. \quad (4.9)$$

When consider the dark energy density is constrained by the expected value of the Higgs potential, the Eq. (4.9) seems more persuasive.

From Eq. (2.3) and Eq. (4.9), and introduce numerical constant n^2 , we have

$$\Omega_{de} = \frac{n^2}{t H^2}. \quad (4.10)$$

Then we can use SNIa Pantheon sample and the Planck 2018 CMB angular power spectra to constrain the parameters of dark energy models based on Eq. (4.3) and (4.10).

5 The Used observation data

5.1 SNIa Pantheon sample and the Plank 2018 CMB angular power spectra

For SNIa data, the Pantheon sample is the combination of SNe Ia from the Pan-STARRS1 (PS1), the Sloan Digital Sky Survey (SDSS), SNLS, and various low- z and Hubble Space Telescope samples. The Panoramic Survey Telescope & Rapid Response System (Pan-STARRS or PS1) is a wide-field imaging facility built by the University of Hawaii's Institute for Astronomy, which is used for a variety of scientific studies from the nearby to the very distant Universe, and it has provided 279 SNe Ia for Pantheon sample [35]. The Supernova Legacy Survey Program detected approximately 2000 high-redshift Supernovae between 2003 and 2008, and Pantheon sample contains about 236 SNe Ia based on the first three years of data, which can be used to investigate the expansion history of the universe, and improve the constraint of cosmological parameters, as well as dark energy study [32].

In 2014 SDSS Survey released a large catalogue which contains light curves, spectra, classifications, and ancillary data of 10,258 variable and transient sources [39, 33, 38, 40, 34, 41]. The release generated the largest sample of supernova candidates, and 500 of sample have been confirmed as SNe Ia by the spectroscopic follow-up. 335 SNe Ia in Pantheon sample are taken from this spectroscopic sample. The rest of Pantheon sample are from the CfA1 – 4, CSP, and Hubble Space Telescope (HST) SN surveys [32]. This extended sample of 1048 SNe Ia is called the Pantheon sample.

The Planck 2018 CMB angular power spectra data is based on observations obtained with Planck (<http://www.esa.int/Planck>), an ESA science mission with instruments and contributions directly funded by ESA Member States, NASA, and Canada.

5.2 SALT2 calibration for Pantheon sample

When correcting the apparent magnitude of Pantheon sample, considering that the prior dark energy equation of state is unknown, so we use SALT2 and Taylor expansion of $d_H - z$ relation to directly calibrate the distance modulus, which can simplify the problem.

The Taylor expansion of $d_H - z$ relation can be given by

$$d_{H,th} = \frac{1}{1-y} \left\{ y - \frac{q_0 - 1}{2} y^2 + \left[\frac{3q_0 - 2q_0 - j_0}{6} + \frac{-\Omega_{k_0} + 2}{6} \right] y^3 \right\}, \quad (5.1)$$

where $y = z/(1+z)$. In order to reduce calculation error for high redshift data, we take this variable substitution. q_0 is the deceleration parameter, j_0 is the jerk parameters, and Ω_{k_0} is the curvature term.

Meanwhile, the relation of distance modulus μ and luminosity distance d_H can be written as

$$\mu = 5\log_{10}d_H + 25 - 5\log_{10}H_0. \quad (5.2)$$

We use SALT2 and Taylor expansion of $d_H - z$ relation to directly calibrate Pantheon sample. The distance modulus μ_{ob} correction formula is given by SALT2 model [36, 37]

$$\mu_{B,ob} = m_B - M_B + \alpha \times x_1 + \beta \times c + \Delta B, \quad (5.3)$$

where m_B corresponds to the observed peak magnitude in rest frame B band, x_1 describes the time stretching of the light curve, c describes the SN colour at maximum brightness, ΔB is a bias correction based on previous simulations, and α, β are nuisance parameters in the distance estimate. M_B is the absolute B-band magnitude, which depends on the host galaxy properties [34]. Notice that M_B is related to the host stellar mass ($M_{stellar}$) by a simple step function

$$M_B = \begin{cases} M_B^1 & \text{if } M_{stellar} < 10^{10} M_\odot \\ M_B^1 + \Delta_M & \text{otherwise} \end{cases} \quad (5.4)$$

Here M_\odot is the mass of the Sun.

From Eq. (5.2) and (5.3), the χ^2 of Pantheon data can be calculated as

$$\chi^2 = \Delta\mu^T C_{\mu_{ob}}^{-1} \Delta\mu, \quad (5.5)$$

where $\Delta\mu = \mu - \mu_{th}$. C_μ is the covariance matrix of the distance modulus μ , we only consider statistical error, and

$$C_{\mu,stat} = V_{m_B} + \alpha^2 V_{x_1} + \beta^2 V_c + 2\alpha V_{m_B, x_1} - 2\beta V_{m_B, c} - 2\alpha\beta V_{x_1, c} \quad (5.6)$$

From Eq. (5.5), and in combination with Pantheon sample, we obtain the statistical average and error for distance modulus μ and the parameters q_0, j_0 . Then we use the calibrated Pantheon sample to constrain the dark energy models parameters.

6 Using the Pantheon sample to fit the models parameters

6.1 The fitting of model I parameters

For model I $\Omega_{de} = n^2 H^{-2}/t$, we firstly choose the universe age T as the time scale t , and consider Taylor expansion of $T - z$ relation in nearly flat space, Which is

Table 1: Statistical mean values of the cosmological parameters from SN Ia Pantheon sample observation data combined with Model I.

$\Omega_{de} = n^2 H_0^{-2}/t$	Ω_m	$n^2 H_0^{-1}$	q_0	j_0	$t_0 H_0 (t_0 = \eta_{T_0}, T_0)$	$\chi_{\min}^2/d.o.f.$
$t = \eta_T$	0.23 ± 0.013	3.3 ± 0.5	-0.57 ± 0.28	-0.2 ± 2.7	4.3 ± 0.7	1040/1050
$t = T$	0.236 ± 0.012	2.84 ± 0.56	-0.44 ± 0.31	-0.95 ± 2.8	3.7 ± 0.74	1044/1050

$$T - T_0 = -\frac{1}{H_0} \left(y - \frac{q_0}{2} y^2 + \frac{2q_0^2 - j_0}{6} y^3 \right), \quad (6.1)$$

where $y = z/(1+z)$. In order to reduce calculation error for high redshift data, we take this variable substitution. T_0 is the present age of the universe.

Then the Hubble parameter in nearly flat space can be written as

$$E(z) = \sqrt{\Omega_m(1+z)^3 + \Omega_R(1+z)^4 + \frac{n^2}{T H_0^2}}. \quad (6.2)$$

where $E(z) = H(z)/H_0$. From Eq. (6.1) and (6.2), one has $T_0 = n^2 H_0^{-2}/(1 - \Omega_m)$ when $\Omega_R \ll \Omega_m$.

We use χ^2 statistic fitting method to constrain the parameters

$$\chi^2_{Pantheon} = \Delta\mu^T C_\mu^{-1} \Delta\mu + \Delta q_0^2 \sigma_{q_0}^{-2} + \Delta j_0^2 \sigma_{j_0}^{-2}, \quad (6.3)$$

where $\Delta\mu = \mu - \mu_{th}$, $\Delta q_0 = q_0 - q_{0,prior}$, $\Delta j_0 = j_0 - j_{0,prior}$, $q_{0,prior}$, $j_{0,prior}$ are given by SALT2 calibration method in Section 5.2.

Following this, we combine Pantheon data with Eq. (6.2) to constrain the parameters, and then use MCMC technology and χ^2 statistic fitting method to obtain the statistical mean values and the minimum chi-square without systematic errors of the parameters $\Omega_m, n^2 H_0^{-1}, q_0, j_0, t_0 H_0$, which are shown in Table (1), The confidence region of $(\Omega_m, n^2 H_0^{-1})$ plane are 68.3%, 95.4% and 99.7%. (see Figs. (1))

Moreover, we can also choose the conformal age η_T as the time scale t ; and consider Taylor expansion of $\eta_T - z$ relation in near flat space, Which is

$$\eta_T - \eta_{T_0} = \frac{1}{H_0} \left\{ y - \frac{q_0 - 1}{2} y^2 + \left[\frac{3q_0 - 2q_0 - j_0}{6} \right] y^3 \right\}, \quad (6.4)$$

Where η_{T_0} is the currently conformal age of the universe.

The Hubble parameter in near flat space can be expressed as

$$E(z) = \sqrt{\Omega_m(1+z)^3 + \Omega_R(1+z)^4 + \frac{n^2}{\eta_T H_0^2}}. \quad (6.5)$$

Then from Eq. (6.4) and (6.5), η_{T_0} satisfies $\eta_{T_0} = n^2 H_0^{-2}/(1 - \Omega_m)$.

In the same way, we use Eq. (6.5) with Pantheon sample to fit the model parameters, and the statistical results are shown in Table (1), and Figs. (1) shows the confidence region of $(\Omega_m, n^2 H_0^{-1})$ plane are 68.3%, 95.4% and 99.7%.




Figure 1: 68.3%, 95.4%, and 99.7% confidence region of the $(\Omega_m, n^2 H_0^{-1})$ plane from Pantheon observation data combined with Model I, the + dots in responding color represent the best fitting values for $\Omega_m, n^2 H_0^{-1}$.

Table 2: Statistical mean values of the cosmological parameters from SN Ia Pantheon sample observation data combined with Model II.

$\Omega_{de} = n^2 H^{-2} / t^2$	Ω_m	n	q_0	j_0	$t_0 H_0 (t_0 = \eta_{T_0}, T_0)$	$\chi^2_{\min} / d.o.f.$
$t = \eta_T$	0.19 ± 0.016	4.4 ± 0.44	-0.64 ± 0.27	-1.1 ± 2.6	5.5 ± 0.62	1042/1050
$t = T$	0.22 ± 0.01	4.5 ± 0.4	-0.48 ± 0.3	-0.7 ± 2.8	5.7 ± 0.7	1044/1050

6.2 The fitting of model II parameters

When considering model II, in which $\Omega_{de} = n^2 H^{-2} / t^{-2}$, we can also select the universe age T as the time scale t , and the Hubble parameter in near flat space can be written as

$$E(z) = \sqrt{\Omega_m(1+z)^3 + \Omega_R(1+z)^4 + \frac{n^2}{T^2 H_0^2}}. \quad (6.6)$$

Alternatively, if we select the conformal age η_T as the time scale t , the Hubble parameter in near flat space satisfies

$$E(z) = \sqrt{\Omega_m(1+z)^3 + \Omega_R(1+z)^4 + \frac{n^2}{\eta_T^2 H_0^2}}. \quad (6.7)$$

Using the pointing models Eq. (6.6) and (6.7), we obtain the statistical results for the parameters which are shown in Table (2), and Figs. (2) shows the confidence region of (Ω_m, n) plane are 68.3%, 95.4% and 99.7%.

If we consider the current age of the universe to be $T_0 = 13.5 \pm 0.5 Gyr$ which is inferred from globular clusters [42], and Hubble constant is $H_0 = 73.5 \pm 1.4 km s^{-1} Mpc^{-1}$ using the NGC 4258 distance measurement [43]. By combining with the fitting results obtained from Pantheon sample in Table (1) and (2), we discover the choice of η_T as time scale may be more persuasive. Furthermore, for the mean value of Ω_m , Conley et al. provided a the




Figure 2: 68.3%, 95.4%, and 99.7% confidence region of the (Ω_m, n) plane from Pantheon observation data combined with Model II. The + dots in responding color represent the best fitting values for Ω_m, n .

statistical result of matter density for constant w CDM model $\Omega_m = 0.19^{+0.08}_{-0.10}$ from the combination of SNLS, HST, and low- z , SDSS data [32], our result is in agreement with it.

If only Pantheon data are used to investigate dark energy, the statistical results indicate the age dark energy models including $\Omega_{de} = n^2 H^{-2} / \eta_T$ and $\Omega_{de} = n^2 H^{-2} / \eta_T^2$ have no evident superiority compared to Λ CDM using minimum chi-square.

7 Using CMB angular power spectra to constrain the models parameters

In addition, we can use Planck 2018 CMB angular power spectra data to constrain the models parameters. When $z \geq 2.5$, $\Omega_m(1+z)^3 / \Omega_{de}(z) \gg 1$, so we think the Taylor expansion for $\eta_T - z$ relation is also applicable to calculate CMB angular power spectra. A calculation of $C_{TT,l}^s$ that ignores the Sunyaev-Zeldovich (SZ) effect can refer to Weinberg 2008 [44], and SZ-effect can refer to Bond et al. 2005 [45]. We use Planck 2018 data together with Model I and II to constrain the cosmology parameters, and the statistical mean values of the parameters are shown in Table (3), and the Figs. (3) shows the CMB theoretical TT power spectra by the best fitting values from the Planck 2018.

In Table (3), we provide the statistical mean values of Hubble constant $H_0 = 73.2 \pm 1.3 \text{ km s}^{-1} \text{ Mpc}^{-1}$, which is consistent with the result obtained using NGC 4258 distance measurement [43].

For the dark energy density, when using Planck 2018 CMB data, the statistical results indicate the age dark energy models including $\Omega_{de} = n^2 H^{-2} / \eta_T$ and $\Omega_{de} = n^2 H^{-2} / \eta_T^2$ have evident superiority, compared to Λ CDM using a minimum chi-square of $l(l+1)C_{TT,l}^s$ ($l = 30 \sim 1500$).

8 CONCLUSIONS

Understanding the physical nature of dark energy is important for our universe. In addition to the study of particle physics, dark energy may also enable us to further explore the

Table 3: The statistical mean values of the cosmological parameters from The Planck 2018 TT power spectra data combined with the Model I and II.

	$\Omega_{de} = n^2 H^{-2} / \eta_T$	$\Omega_{de} = n^2 H^{-2} / \eta_T^2$
$\Omega_b h^2$	0.0214 ± 0.00012	0.023 ± 0.0002
$\Omega_c h^2$	0.11 ± 0.0014	0.107 ± 0.0014
$10^{10} A_s e^{-2\tau}$	1.29 ± 0.016	1.316 ± 0.027
n_s	0.93 ± 0.004	0.97 ± 0.0025
N_{eff}	3.45 ± 0.13	3.5 ± 0.35
h_0	0.742 ± 0.012	0.732 ± 0.013
Ω_m	0.237 ± 0.018	0.243 ± 0.022
$n^2 H_0^{-1}, n$	5.5 ± 0.33	4.85 ± 0.54
q_0	-0.38 ± 0.24	-0.7 ± 0.59
j_0	1.3 ± 0.22	4.7 ± 1.2
σ_8^{sz}	1.1 ± 0.011	1.06 ± 0.012




Figure 3: Blue dots in responding color represent Planck 2018 CMB TT angular power spectra. The red, yellow and cyan lines are the CMB theoretical values of angular power spectra from Λ CDM, Model I and II, respectively, using the best fitting values, which are constrained by Planck 2018 data.

nature of vacuum. Whether or not the dark energy is derived from scalar bosons, other particle, or neither, still needs to be further verified.

We explore a theoretical possibility that dark energy density is derived from massless scalar bosons in vacuum. Assuming massless scalar bosons fall into the horizon boundary with the expansion of the universe, scalar bosons satisfy P-symmetry under the parity transformation. P-symmetry with the quantum fluctuation of space-time enable us to

estimate dark energy density. Meanwhile, to explain the physical nature of negative pressure, this is deduced from the increasing entropy density with the rise in degeneracy in the Boltzmann system.

Next, we used the SNIa Pantheon sample and Planck 2018 CMB angular power spectra to constrain the specified models. The statistical results indicate that age dark energy models have evident superiority, when only using CMB data compared to Λ CDM using minimum chi-square. Furthermore, we obtain the statistical mean value of the Hubble constant $H_0=73.2 \pm 1.3 \text{ km s}^{-1} \text{ Mpc}^{-1}$, which is consistent with the result obtained using NGC 4258 distance measurement.

Finally, we extend our discussion to the future of universe. From Eq. (3.7), if W_p satisfies $W_p : 3 \rightarrow 2 \rightarrow 1$, the universe may continue expanding in the future; however, if $W_p : 1 \rightarrow 2 \rightarrow 3$, this will change from expansion to contraction. Thus, the property of dark energy will dominate the future of universe, and it will similarly determine the future of humanity.

9 Acknowledgments

This work was supported by Xiaofeng Yang's Xinjiang Tianchi Bairen project and CAS Pioneer Hundred Talents Program. This work was also partly supported by the National Key R&D Program of China under grant No.2018YFA0404602.

References

- [1] A. Riess et al., *Astron. J.* **116**, 1009 (1998)
- [2] S. Capozziello and M. Francaviglia, *General Relativity and Gravitation*, **40**, 357 (2008).
- [3] C. Brans and R. H. Dicke, *Phys. Rev.*, **124**, 925 (1961).
- [4] P. J. E. Peebles and B. Ratra, *Rev. Mod. Phys.*, **75**, 559 (2003).
- [5] W. Hu, S. Dodelson, *Annu. Rev. Astron. Astr.*, **40**: 171 (2002).
- [6] E. V. Linder, *Phys. Rev. Lett.*, **90**, 091301 (2003)
- [7] D. Huterer and G. Starkman, *Phys. Rev. Lett.*, **90**, 031301 (2003)
- [8] C. Clarkson and C. Zunckel, *Phys. Rev. Lett.*, **104**, 211301 (2010)
- [9] T. Holsclaw and U. Alam, B. Sanso, H. Lee, K. Heitmann, S. Habib, D. Higdon, *Phys. Rev. Lett.*, **105**, 241302 (2010)
- [10] A. Shafieloo, A. G. Kim, E. V. Linder, *Phys. Rev. D*, **85**, 123530 (2012)
- [11] M. Seikel, C. Clarkson and M. Smitha, *JCAP*, **06**, 036 (2012)
- [12] R. G. Crittenden and L. Pogosian et al., *JCAP*, **0912**, 025 (2009)
- [13] R. G. Crittenden and G. B. Zhao et al., *JCAP*, **1202**, 048 (2012)

- [14] G. B. Zhao and R. G. Crittenden, L. Pogosian, X. Zhang, *Phys. Rev. Lett.*, **109**, 171301 (2012)
- [15] L. Huang, X. F. Yang, X. Liu, *Astrophys. J.*, **913**: 24 (2021).
- [16] B. Ratra and P. J. E. Peebles, *Phys. Rev. D.*, **37**, 3406 (1988).
- [17] J. A. Frieman, C. T. Hill, A. Stebbins et al, *Phys. Rev. Lett.*, **75**, 2077 (1995).
- [18] M. Li, *Phys. Lett. B*, **603**, 1 (2004).
- [19] M. Maziashvili, *Int. J. Mod. Phys. D*, **16**, 1531 (2007).
- [20] M. Maziashvili, *Phys. Lett. B*, **652**, 165 (2007).
- [21] E. P. Wigner, *Rev. Mod. Phys*, **29**, 255 (1957).
- [22] F. Karolyhazy, *Il Nuovo Cimento A*, **42**, 390 (1966).
- [23] N. Y. Jack and D. H. van, *Mod. Phys. Lett. A*, **9**, 335 (1994).
- [24] C. G. Amelino, *Mod. Phys. Lett. A*, **9**, 3415 (1994).
- [25] C. G. Amelino, *Nature*, **398**, 216 (1999).
- [26] C. G. Amelino and J. Lukierski, arXiv preprint hep-th/9706031, (1997).
- [27] C. G. Amelino, *Mod. Phys. Lett. A*, **13**, 1319 (1998).
- [28] R. G. Cai, *Phys. Lett. B*, **657**, 228 (2007).
- [29] H. Wei and R. G. Cai, *Phys. Lett. B*, **660**, 113 (2008).
- [30] M. Betoule and R. Kessler et al., *Astron. Astrophys.*, **568**, A22 (2014)
- [31] P. W. Higgs, *Phys. Rev. Lett.*, **13**, 508 (1964).
- [32] A. Conley et al., *Astrophys. J. Suppl. S.*, **192**, 1 (2010).
- [33] D. G. York et al., *Astron. J.*, **120**, 1579 (2000).
- [34] M. Betoule et al., *Astron. Astrophys.*, **568**, A22 (2014).
- [35] D. M. Scolnic, D. O. Jones, A. Rest et al., *Astrophys. J.*, **859**, 101 (2018).
- [36] J. Guy et al., *Astron. Astrophys.*, **443**, 781 (2005).
- [37] J. Guy et al., *Astron. Astrophys.*, **466**, 11 (2007).
- [38] J. E. Gunn et al., *Astron. J.*, **131**, 2332 (2006).
- [39] J. E. Gunn et al., *Astron. J.*, **116**, 3040 (1998).
- [40] M. Sako et al., *Astron. J.*, **135**, 348 (2007).
- [41] M. Sako et al. , *Publ. Astron. Soc. Pac.*, **130**, 064002 (2014).
- [42] D. Valcin et al., *JCAP*, **12**, 002 (2020).

- [43] M. J. Reid, D. W. Pesce, A. G. Riess, *Astron. J. Lett.*, **886**, L27 (2019).
- [44] S. Weinberg, Oxford University Press, (2008).
- [45] J. R. Bond, C. R. Contaldi, U. L. Pen et al., *Astron. J.*, **626**, 12 (2005).