An inequality for the compositions of convex functions with convolutions and an alternative proof of the Brunn–Minkowski–Kemperman inequality *

Takashi Satomi

February 21, 2023

Abstract

Let m(G) be the infimum of the volumes of all open subgroups of a unimodular locally compact group G. Suppose integrable functions $\phi_1, \phi_2 \colon G \to [0, 1]$ satisfy $\|\phi_1\| \leq \|\phi_2\|$ and $\|\phi_1\| + \|\phi_2\| \leq m(G)$, where $\|\cdot\|$ denotes the L^1 -norm with respect to a Haar measure dg on G. We have the following inequality for any convex function $f \colon [0, \|\phi_1\|] \to \mathbb{R}$ with f(0) = 0:

$$\int_{G} f \circ (\phi_{1} * \phi_{2})(g) dg \leq 2 \int_{0}^{\|\phi_{1}\|} f(y) dy + (\|\phi_{2}\| - \|\phi_{1}\|) f(\|\phi_{1}\|)$$

As a corollary, we have a slightly stronger version of Brunn–Minkowski–Kemperman inequality. That is, we have

$$\operatorname{vol}_*(B_1B_2) \ge \operatorname{vol}(\{g \in G \mid 1_{B_1} * 1_{B_2}(g) > 0\}) \ge \operatorname{vol}(B_1) + \operatorname{vol}(B_2)$$

for any non-null measurable sets $B_1, B_2 \subset G$ with $\operatorname{vol}(B_1) + \operatorname{vol}(B_2) \leq m(G)$, where vol_* denotes the inner measure and 1_B the characteristic function of B.

Keywords: convolution, convexity, locally compact group, combinatorial inequality, geometric measure theory.

MSC2020: Primary 43A05; Secondary 05A20, 22D05, 28C10, 39B62.

1 Introduction

Let \mathcal{A} be the set of all open subgroups of a unimodular locally compact group G, and we set

$$m(G) := \inf_{G' \in \mathcal{A}} \operatorname{vol}(G')$$

 $(m(G) \text{ may be } \infty)$. Let $\|\cdot\|$ be the L^1 -norm with respect to a Haar measure dg on G. For two measurable functions $\phi_1, \phi_2 \colon G \to \mathbb{R}_{\geq 0}$, we define the convolution $\phi_1 * \phi_2$ as

$$\phi_1 * \phi_2(g) := \int_G \phi_1(g') \phi_2(g'^{-1}g) dg'.$$

The main result of this paper is the following theorem.

Theorem 1.1. Suppose integrable functions $\phi_1, \phi_2 \colon G \to [0, 1]$ on a unimodular locally compact group G satisfy $\|\phi_1\| \leq \|\phi_2\|$ and

$$\|\phi_1\| + \|\phi_2\| \le m(G). \tag{1.1}$$

^{*}The final version of the paper is published in Proc. Steklov Inst. Math. 319 (2022): http://dx.doi.org/10.1134/S0081543822050182

Then, for any convex function $f: [0, ||\phi_1||] \to \mathbb{R}$ with f(0) = 0, one has

$$\int_{G} f \circ (\phi_1 * \phi_2)(g) dg \le 2 \int_{0}^{\|\phi_1\|} f(y) dy + (\|\phi_2\| - \|\phi_1\|) f(\|\phi_1\|).$$
(1.2)

For example, by using the L^1 -norm (and L^{∞} -norm) of ϕ_1 and ϕ_2 , Theorem 1.1 bounds the L^p -norm of the convolution $\phi_1 * \phi_2$ when $f(y) = y^p$ holds (Example 2.1 (2) and (3)), and the entropy of $\phi_1 * \phi_2$ from below when $f(y) = y \log y$ (Example 2.1 (4)). Actually, Theorem 1.1 is best possible for $G = \mathbb{R}$ (Example 2.1 (1)).

Here are some other examples of Theorem 1.1. When $t \ge 0$ is fixed,

$$f_t(y) := \begin{cases} y - t & \text{if } y \ge t \\ 0 & \text{if } y \le t \end{cases}$$
(1.3)

is convex. Since

$$2\int_0^{\|\phi_1\|} f_t(y)dy + (\|\phi_2\| - \|\phi_1\|)f_t(\|\phi_1\|) = (\|\phi_1\| - t)(\|\phi_2\| - t)$$

holds for any $0 \le t \le ||\phi_1||$, we have

$$||f_t \circ (\phi_1 * \phi_2)|| \le (||\phi_1|| - t)(||\phi_2|| - t)$$
(1.4)

by applying $f = f_t$ to Theorem 1.1. Actually, (1.4) holds even when the assumption (1.1) is weakened by

$$\|\phi_1\| + \|\phi_2\| \le m(G) + t. \tag{1.5}$$

That is, the following lemma holds.

Lemma 1.2. One has (1.4) for any integrable functions $\phi_1, \phi_2 \colon G \to [0, 1]$ on a unimodular locally compact group G and for any $0 \leq t \leq \min\{\|\phi_1\|, \|\phi_2\|\}$ with (1.5).

For any characteristic functions ϕ_1 and ϕ_2 on a compact connected group G, Lemma 1.2 was essentially proved by Tao [Tao12, Theorem 3.2.3] and explicitly given by Christ [Chr19, Section 6]. Christ also proved Lemma 1.2 for any characteristic functions ϕ_1 and ϕ_2 on $G = \mathbb{R}^n$, and discussed a relation between Lemma 1.2 and the Riesz–Sobolev inequality [Rie30] [Sob38].

A claim similar to Lemma 1.2 for any cyclic group of prime order was proved by Pollard [Pol74], and for any finite abelian group by Green and Ruzsa [GR05, Proposition 6.1]. By using this claim, Green and Ruzsa explicitly computed the maximal cardinality of all sum-free subsets for any finite abelian group [GR05, Theorem 1.5].

In addition, when ϕ_1 and ϕ_2 are characteristic functions, we obtain the following corollary by applying Theorem 1.1 to

$$f(y) := \begin{cases} 0 & \text{if } y = 0 \\ -1 & \text{if } y > 0 \end{cases}$$
(1.6)

Corollary 1.3. Let $B_1, B_2 \subset G$ be measurable sets on a unimodular locally compact group G with $vol(B_1), vol(B_2) > 0$. If $vol(B_1) + vol(B_2) \leq m(G)$ holds, then one has

$$\operatorname{vol}(\{g \in G \mid 1_{B_1} * 1_{B_2}(g) > 0\}) \ge \operatorname{vol}(B_1) + \operatorname{vol}(B_2), \tag{1.7}$$

where 1_B denotes the characteristic function of B. In particular,

$$\operatorname{vol}_{*}(B_{1}B_{2}) \ge \operatorname{vol}(B_{1}) + \operatorname{vol}(B_{2})$$
 (1.8)

holds, where vol_* denotes the inner measure.

Unimodular locally compact group G	Authors and references	
\mathbb{R}^n (B_1 and B_2 are convex)	Brunn [Bru87], Minkowski [Min96]	
\mathbb{R}^n	Lusternik [Lus35],	
IK.	Henstock–Macbeath [HM53, Theorem 1]	
S^1	Raikov [Rai39]	
$(S^1)^n$	Macbeath [Mac53, Theorem 1]	
Compact connected second countable abelian	Shields [Shi55]	
Abelian	Kneser [Kne56]	
General	Kemperman [Kem64, Theorem 1.2]	
Connected or with $m(G) = \infty$	Ruzsa [Ruz92, Theorem 2]	

Table 1: The authors who proved (1.8) for various cases

The Brunn–Minkowski–Kemperman inequality (1.8) follows from (1.7) and the fact that $\{g \in G \mid 1_{B_1} * 1_{B_2}(g) > 0\} \subset B_1B_2$. Corollary 1.3 was proved by Brascamp and Lieb for $G = \mathbb{R}^n$ [BL76, Theorem A.1], and by Tao for any compact connected group G by using Lemma 1.2 [Tao12, Section 3.2]. This argument is essentially equivalent to applying Theorem 1.1 to (1.6). The inequality (1.8) was proved by Kemperman for any general unimodular locally compact group G, and a number of people for particular cases (see Table 1).

For $G = \mathbb{R}^n$, (1.8) can be improved to

$$\operatorname{vol}_*(B_1B_2)^{1/n} \ge \operatorname{vol}(B_1)^{1/n} + \operatorname{vol}(B_2)^{1/n}.$$
 (1.9)

For any convex sets B_1 and B_2 on $G = \mathbb{R}^n$, Brunn proved (1.9), and Minkowski proved that the equality is attained in (1.9) if and only if B_1 and B_2 are equal up to dilatation. For arbitrary measurable sets B_1 and B_2 on $G = \mathbb{R}^n$, the inequality (1.9) was stated by Lusternik. However, Lusternik's proof is defective, and it was corrected by Henstock and Macbeath. A generalization of (1.9) for any locally compact group G was studied by McCrudden [McC69, Theorem 1.2] and Jing, Tran, and Zhang [JTZ21, Theorem 1.1].

When G is connected or $m(G) = \infty$ holds, Ruzsa gave an alternative proof of (1.8). Kemperman's idea is to study the sets B_1 and B_2 such that the maximum of $\operatorname{vol}(B_1) + \operatorname{vol}(B_2) - \operatorname{vol}_*(B_1B_2)$ is attained. Ruzsa's idea is to show that $T(I) := \inf_{\operatorname{vol}(B_1)=I} \operatorname{vol}_*(B_1B_2)$ is a concave function for any fixed B_2 .

For any compact connected abelian group G, Tao studied a property of measurable sets for which the equality of Corollary 1.3 is almost attained [Tao18, Theorem 1.3]. This result is generalized for disconnected groups by Griesmer [Gri19, Theorem 1.1], and for non-abelian groups by Jing and Tran [JT20, Theorem 1.2]. A sharper result for any compact abelian group is given by Christ and Iliopoulou [CI22, Theorem 1.4].

Here is the strategy of proof of Theorem 1.1. As mentioned above, Lemma 1.2 under the assumption (1.1) is an example of Theorem 1.1. However, we will show Lemma 1.2 without using Theorem 1.1 at first, and obtain Theorem 1.1 by using Lemma 1.2. We will show Lemma 1.2 by generalizing the argument of Tao [Tao12, Theorem 3.2.3], and Theorem 1.1 by approximating f by a linear combination of f_t .

The organization of this paper is as follows. In Section 2, we will give some examples and remarks concerning Theorem 1.1. Sections 3-6 are devoted to the proof of Lemma 1.2. In Section 3, we will discuss the necessity of (1.1). In Section 4, we will give a key lemma (Lemma 4.1) to prove Lemma 1.2. By using the arguments in Sections 3 and 4, we will show Lemma 1.2 in the case of $m(G) = \infty$ in Section 5, and in the case of $m(G) < \infty$ in Section 6. In Section 7, we will complete the proof of Theorem 1.1 by using Lemma 1.2.

2 Some examples and remarks related to Theorem 1.1

In this section, we give some examples and remarks concerning Theorem 1.1 (and related results). At first, we give some examples of application of Theorem 1.1.

Example 2.1. (1) One has $m(G) = \infty$ for $G = \mathbb{R}$. The function $\phi_{(I)} : \mathbb{R} \to \{0, 1\}$ denotes the characteristic function $\phi_{(I)} := \mathbb{1}_{(-I/2, I/2)}$ of the open interval (-I/2, I/2) for I > 0. Let $\psi_{I_1, I_2} := \phi_{(I_1)} * \phi_{(I_2)}$ for $I_1, I_2 > 0$. If $I_1 \leq I_2$, then one has

$$\psi_{I_1,I_2}(x) = \begin{cases} I_1 & \text{if } |x| \le c \\ c' - |x| & \text{if } c \le |x| \le c' , \quad c := \frac{I_2 - I_1}{2}, \quad c' := \frac{I_1 + I_2}{2} \\ 0 & \text{if } c' \le |x| \end{cases}$$

Thus, if $\phi_1 = \phi_{(I_1)}$ and $\phi_2 = \phi_{(I_2)}$, then one has

$$\int_{\mathbb{R}} f \circ \psi_{I_1, I_2}(x) dx = \int_{-c'}^{-c} f\left(x + c'\right) dx + \int_{-c}^{c} f(I_1) dx + \int_{c}^{c'} f\left(c' - x\right) dx$$
$$= 2 \int_{0}^{I_1} f(y) dy + (I_2 - I_1) f(I_1).$$
(2.1)

Since $I_1 = ||\phi_1||$ and $I_2 = ||\phi_2||$, the equality is attained in (1.2). That is, (1.2) can be expressed as

$$\int_G f \circ (\phi_1 * \phi_2)(g) dg \leq \int_{\mathbb{R}} f \circ \psi_{\|\phi_1\|, \|\phi_2\|}(x) dx.$$

(We note that ϕ_1 and ϕ_2 are not the same as $\phi_1 = \phi_{(I_1)}$ and $\phi_2 = \phi_{(I_2)}$ stated before and are already general functions on a general unimodular locally compact group G.) In particular, (1.4) can be expressed as

$$||f_t \circ (\phi_1 * \phi_2)|| \le ||f_t \circ \psi_{||\phi_1||, ||\phi_2||}||.$$
(2.2)

(2) The function $f(y) := y^p$ is convex for $p \ge 1$. Thus, one has

$$\int_{G} (\phi_1 * \phi_2(g))^p dg \le 2 \int_0^{\|\phi_1\|} y^p dy + (\|\phi_2\| - \|\phi_1\|) \|\phi_1\|^p = \|\phi_1\|^p \left(\|\phi_2\| - \frac{(p-1)\|\phi_1\|}{p+1} \right)$$

by applying Theorem 1.1 in this case.

(3) The function $f(y) := -y^p$ is convex for 0 . Thus, one has

$$\int_{G} (\phi_1 * \phi_2(g))^p dg \ge 2 \int_{0}^{\|\phi_1\|} y^p dy + (\|\phi_2\| - \|\phi_1\|) \|\phi_1\|^p = \|\phi_1\|^p \left(\|\phi_2\| + \frac{(1-p)\|\phi_1\|}{p+1} \right)$$

by applying Theorem 1.1 in this case.

(4) The function $f(y) := y \log y$ is convex. Thus, one has

$$\begin{split} \int_{G} \phi_{1} * \phi_{2}(g) \log((\phi_{1} * \phi_{2})(g)) dg &\leq 2 \int_{0}^{\|\phi_{1}\|} y \log y dy + (\|\phi_{2}\| - \|\phi_{1}\|) \|\phi_{1}\| \log \|\phi_{1}\| \\ &= \|\phi_{1}\| \left(\|\phi_{2}\| \log \|\phi_{1}\| - \frac{\|\phi_{1}\|}{2} \right) \end{split}$$

by applying Theorem 1.1 in this case.

(5) Since $f_t(y) = y$ holds for t = 0, one has $\|\phi_1 * \phi_2\| \le \|\phi_1\| \cdot \|\phi_2\|$ by Lemma 1.2. In addition, $-\|\phi_1 * \phi_2\| \le -\|\phi_1\| \|\phi_2\|$ holds by applying Theorem 1.1 to the convex function f(y) := -y. Thus, one has

$$\|\phi_1 * \phi_2\| = \|\phi_1\| \cdot \|\phi_2\|.$$
(2.3)

By Fubini's theorem, (2.3) holds for any $\phi_1, \phi_2 \colon G \to \mathbb{R}_{\geq 0}$ even when the assumption (1.1) is not satisfied. That is, the following lemma holds.

Lemma 2.2. One has (2.3) for any integrable functions $\phi_1, \phi_2 \colon G \to \mathbb{R}_{\geq 0}$ on a unimodular locally compact group G.

The equalities are attained in Lemma 1.2 and Example 2.1 (2), (3) and (4) for $G = \mathbb{R}$, $\phi_1 = \phi_{(||\phi_1||)}$ and $\phi_2 = \phi_{(||\phi_2||)}$ by Example 2.1 (1).

Next, we define the notion of convex function used in Theorem 1.1.

Definition 2.3. For a function $f: A \to \mathbb{R}$ on an interval A and $y_1, y_2 \in A$ with $y_1 < y_2$, the functions $\tilde{f}_{y_1}^{y_2}, \tilde{f}_{y_1}^{y_2}: A \to \mathbb{R}$ are defined as

$$\tilde{f}_{y_1}^{y_2}(y) := \frac{(y_2 - y)f(y_1) + (y - y_1)f(y_2)}{y_2 - y_1} \quad \text{and} \quad \hat{f}_{y_1}^{y_2}(y) := \tilde{f}_{y_1}^{y_2}(y) - f(y).$$

A function $f: A \to \mathbb{R}$ is called convex if $\hat{f}_{y_1}^{y_2}(y) \ge 0$ (that is, $f(y) \le \tilde{f}_{y_1}^{y_2}(y)$) for any $y_1, y_2, y \in A$ with $y_1 < y_2$ and $y_1 \le y \le y_2$.

Here are some remarks on Theorem 1.1 and Corollary 1.3.

Remark 2.4. (1) The composition $f \circ (\phi_1 * \phi_2)$ in Theorem 1.1 is well-defined because

$$\phi_1 * \phi_2(g) = \int_G \phi_1(g')\phi_2(g'^{-1}g)dg' \le \int_G \phi_1(g')dg' = \|\phi_1\|$$
(2.4)

holds for any $g \in G$.

- (2) When f is of C²-class, f is convex if and only if $f'' \ge 0$ holds. However, Theorem 1.1 does not require f to be differentiable or even continuous (at 0 and $||\phi_1||$).
- (3) The following three conditions are equivalent for any locally compact group G:
 - (i) One has $m(G) < \infty$;
 - (ii) The locally compact group G has an open compact subgroup;
 - (iii) The identity component G_0 of G is compact.

The equivalence of (i) and (ii) is obtained by the fact that $\operatorname{vol}(G') < \infty$ holds if and only if G' is compact [Wei40]. We show that (ii) implies (iii). There is an open compact subgroup $G' \subset G$ by (ii). Since $G_0 \subset G'$ holds by the connectedness of G_0 , one has (iii) by the compactness of G'. We show that (iii) implies (ii). Since G/G_0 is a totally disconnected locally compact group [HR79, Theorem 7.3], there is an open compact subgroup $H \subset G/G_0$ [VD31] [HR79, Theorem 7.7]. Then $\pi^{-1}(H) \subset G$ is an open subgroup, where $\pi: G \to G/G_0$ is the projection. Since $\pi^{-1}(H)$ is compact by (iii) [HR79, Note 5.24 (a)], we obtain (ii).

In addition,

$$G_0 = \bigcap_{G' \in \mathcal{A}} G'$$

holds for any G [HR79, Theorem 7.8]. Thus, if the equivalent conditions (i), (ii) and (iii) are satisfied, then we have $m(G) = vol(G_0)$. Thus, (1.1) holds if and only if at least one of the following two conditions is satisfied:

- (a) One has $\|\phi_1\| + \|\phi_2\| \le \operatorname{vol}(G_0);$
- (b) The identity component G_0 is not compact.

In particular, if G_0 is open (e.g., G is a Lie group), then (1.1) and (a) are equivalent.

(4) A result similar to Theorem 1.1 holds in the case of $\|\phi_2\| \le \|\phi_1\|$ as follows. For integrable functions $\phi_1, \phi_2 \colon G \to [0, 1]$ with $\|\phi_2\| \le \|\phi_1\|$ and (1.1),

$$\phi_1 * \phi_2(g) = \int_G \phi_1(g')\phi_2(g'^{-1}g)dg' \le \int_G \phi_2(g'^{-1}g)dg' = \|\phi_2\|$$
(2.5)

holds and hence $f \circ (\phi_1 * \phi_2)$ is well-defined for any function $f : [0, \|\phi_2\|] \to \mathbb{R}$. Then, one has

$$\int_{G} f \circ (\phi_1 * \phi_2)(g) dg \le 2 \int_{0}^{\|\phi_2\|} f(y) dy + (\|\phi_1\| - \|\phi_2\|) f(\|\phi_2\|)$$

for any convex function $f: [0, \|\phi_2\|] \to \mathbb{R}$ with f(0) = 0 by the argument similar to Theorem 1.1.

(5) In Corollary 1.3, we set $U := \{g \in G \mid 1_{B_1} * 1_{B_2}(g) > 0\}$. In fact, there is a case where $\operatorname{vol}_*(B_1B_2) > \operatorname{vol}(U)$. For example, when $G = \mathbb{R}$, $B_1 = (0,1) \cup \{2\}$ and $B_2 = (0,1)$ hold, we have $B_1B_2 = (0,3) \setminus \{2\}$ and U = (0,2). Since $\operatorname{vol}(B_1) = \operatorname{vol}(B_2) = 1$, $\operatorname{vol}_*(B_1B_2) = 3$ and $\operatorname{vol}(U) = 2$ hold, we have

$$\operatorname{vol}_*(B_1B_2) > \operatorname{vol}(U) = \operatorname{vol}(B_1) + \operatorname{vol}(B_2).$$

(6) If $\|\phi_1\| = 0$, then we have $\phi_1 * \phi_2 = 0$ and hence Theorem 1.1 and Lemma 1.2 hold. Thus, it suffices to show Theorem 1.1 and Lemma 1.2 for $\|\phi_1\| > 0$. In this case, one can reduce the proof to the case of $\|\phi_1\| = 1$ by replacing dg and f(y) with $dg/\|\phi_1\|$ and $f(y/\|\phi_1\|)$, respectively.

3 Necessity of the assumption (1.1)

In this section, we discuss the necessity of the assumption (1.1) when G is connected. That is, if (1.1) does not hold, then (1.4) does not hold for any $0 < t < \|\phi_1\| + \|\phi_2\| - m(G)$ by the following lemma.

Lemma 3.1. Let $\phi_1, \phi_2 \colon G \to [0,1]$ be integrable functions on a unimodular locally compact group G with $G = G_0$ (that is, G is connected).

- (1) One has $\phi_1 * \phi_2 \ge \|\phi_1\| + \|\phi_2\| m(G)$.
- (2) If $\|\phi_1\| + \|\phi_2\| m(G) \ge 0$ holds, then one has $\|f_t \circ (\phi_1 * \phi_2)\| = \|\phi_1\| \|\phi_2\| tm(G)$ for any $0 \le t \le \|\phi_1\| + \|\phi_2\| m(G)$.

Proof. (1) For any $g \in G$,

$$\phi_1 * \phi_2(g) = \int_G \phi_1(g')\phi_2(g'^{-1}g)dg'.$$

One has

$$\phi_1(g')\phi_2(g'^{-1}g) = \phi_1(g') + \phi_2(g'^{-1}g) - 1 + (1 - \phi_1(g'))(1 - \phi_2(g'^{-1}g))$$

$$\geq \phi_1(g') + \phi_2(g'^{-1}g) - 1$$

by $\phi_1(g'), \phi_2(g'^{-1}g) \leq 1$, and hence

$$\int_{G} \phi_1(g')\phi_2(g'^{-1}g)dg' \ge \int_{G} (\phi_1(g') + \phi_2(g'^{-1}g) - 1)dg' = \|\phi_1\| + \|\phi_2\| - \operatorname{vol}(G).$$

Since m(G) = vol(G) holds by $G = G_0$ and Remark 2.4 (3), we obtain

$$\phi_1 * \phi_2(g) = \int_G \phi_1(g')\phi_2(g'^{-1}g)dg' \ge \|\phi_1\| + \|\phi_2\| - \operatorname{vol}(G) = \|\phi_1\| + \|\phi_2\| - m(G).$$

(2) Since $f_t \circ (\phi_1 * \phi_2) = \phi_1 * \phi_2 - t$ holds by (1), we obtain

$$||f_t \circ (\phi_1 * \phi_2)|| = ||\phi_1 * \phi_2 - t|| = ||\phi_1|| ||\phi_2|| - tm(G)$$

by Lemma 2.2.

By Lemma 3.1, the inequality (1.4) does not hold for any $0 < t < \|\phi_1\| + \|\phi_2\| - m(G)$ when G is connected and (1.1) does not hold. In fact,

$$||f_t \circ (\phi_1 * \phi_2)|| = ||\phi_1|| \cdot ||\phi_2|| - tm(G) = (||\phi_1|| - t)(||\phi_2|| - t) + t(||\phi_1|| + ||\phi_2|| - m(G) - t)$$

holds by Lemma 3.1 (2) and hence we have $(\|\phi_1\| - t)(\|\phi_2\| - t) < \|f_t \circ (\phi_1 * \phi_2)\|$.

4 First step of the proof of Lemma 1.2

In this section, we give the key lemma (Lemma 4.1) to prove Lemma 1.2. This idea is based on the submodularity inequality by Pollard [Pol74], Ruzsa [Ruz92, Lemma 2.2], Green and Ruzsa [GR05, Proposition 6.1] and Tao [Tao12, Theorem 3.2.3] [Tao13, Lemma 2.1].

It suffices to show Lemma 1.2 for $\|\phi_1\| = 1$ by Remark 2.4 (6). We fix ϕ_1 with $\|\phi_1\| = 1$ and simply write f for f_t in (1.3). The function $S: [0, \operatorname{vol}(G)] \to \mathbb{R}_{\geq 0}$ is defined as

$$S(I) := \sup_{\phi \in \mathcal{B}(I)} \|\xi_{\phi}\|,$$

where $\mathcal{B}(I) := \{\phi \colon G \to [0,1] \mid \|\phi\| = I\}$ and $\xi_{\phi} := f \circ (\phi_1 * \phi)$. Lemma 1.2 can be expressed as

$$S(I) \le (1-t)(I-t)$$
 (4.1)

for any $t \leq I \leq m(G) + t - 1$. We show the following lemma to prove Lemma 1.2.

- **Lemma 4.1.** (1) One has $0 \le S(I') S(I) \le I' I$ for any $0 \le I \le I' \le \text{vol}(G)$. In particular, S is Lipschitz continuous.
 - (2) The function S is convex on [0, m(G)] (or $[0, \infty)$ in the case of $m(G) = \infty$).
 - (3) One has S(t) = 0.
 - (4) One has $S(I) \leq (1-t)I$ for any $0 \leq I \leq \operatorname{vol}(G)$.

Lemma 4.1(2) is deduced from the following lemma.

Lemma 4.2. Let I > 0 and $\phi \in \mathcal{B}(I)$. We define $R_g\phi: G \to [0,1]$ as $R_g\phi(g') := \phi(g'g)$ for $g \in G$. Let $I^2/m(G) < I'_1 \leq I$ and $I'_2 := 2I - I'_1$.

- (1) We define a continuous function $h: G \to \mathbb{R}_{\geq 0}$ as $h(g) := \|\phi R_g \phi\|$. If $I'_1 \leq h(e)$ holds, then there is $g \in G_0$ with $h(g) = I'_1$, where $e \in G$ is the identity.
- (2) There are $g \in G_0$ and $\nu_1 \in \mathcal{B}(I'_1)$ such that $\phi R_g \phi \leq \nu_1 \leq \min\{\phi, R_g \phi\}$ holds.

- (3) Let $g \in G_0$ and $\nu_1 \in \mathcal{B}(I'_1)$ be as in (2). One has $\max\{\phi, R_g\phi\} \le \nu_2 := \phi + R_g\phi \nu_1 \le 1$ and $\nu_2 \in \mathcal{B}(I'_2)$.
- (4) Let $g \in G_0$ and $\nu_1 \in \mathcal{B}(I'_1)$ be as in (2), and $\nu_2 \in \mathcal{B}(I'_2)$ be as in (3). One has $\xi_{\phi} + \xi_{R_g\phi} \leq \xi_{\nu_1} + \xi_{\nu_2}$.
- (5) Let $\nu_1 \in \mathcal{B}(I'_1)$ be as in (2), and $\nu_2 \in \mathcal{B}(I'_2)$ be as in (3). One has $2\|\xi_{\phi}\| \le \|\xi_{\nu_1}\| + \|\xi_{\nu_2}\|$. In particular, $2S(I) \le S(I'_1) + S(I'_2)$ holds.
- *Proof.* (1) Since h is continuous and G_0 is connected, it suffices to show that there is $g' \in G_0$ with $h(g') \leq I'_1$. We divide the proof into the following two cases.
 - (i) The case of $m(G) < \infty$. Then

$$||h||_{G_0} = \int_{G_0} \int_G \phi(g'')\phi(g''g')dg''dg' \le I^2$$

holds, where $||h||_{G_0}$ denotes the L^1 -norm of h on G_0 . Since $vol(G_0) = m(G)$ by Remark 2.4 (3), there is $g' \in G_0$ with

$$h(g') \le \frac{I^2}{m(G)} \le I_1'.$$

(ii) The case of $m(G) = \infty$. There is a compact set $K \subset G$ with $\|\phi\|_K \ge I - I'_1/2$. Since G_0 is not compact by Remark 2.4 (3), there is $g' \in G_0 \setminus K^{-1}K$. Then

 $h(g') = \|\phi R_{g'}\phi\|_{K} + \|\phi R_{g'}\phi\|_{G\setminus K} \le \|R_{g'}\phi\|_{K} + \|\phi\|_{G\setminus K}$

holds by $\phi, R_{g'}\phi \leq 1$. Since $Kg' \subset G \setminus K$ by $g' \in G_0 \setminus K^{-1}K$, one has

$$||R_{g'}\phi||_K = ||\phi||_{Kg'} \le ||\phi||_{G\setminus K}.$$

Thus, we obtain

$$h(g') \le \|R_{g'}\phi\|_{K} + \|\phi\|_{G\setminus K} \le 2\|\phi\|_{G\setminus K} \le I'_{1}.$$

(2) Let $e \in G$ be the identity. When $I'_1 \leq h(e)$ holds, we obtain

$$\nu_1 := \phi R_g \phi \le \min\{\phi, R_g \phi\}$$

by taking $g \in G_0$ in (1). When $I'_1 \ge h(e)$ holds, there is $\nu_1 \in \mathcal{B}(I'_1)$ with $\phi^2 \le \nu_1 \le \phi$. Thus,

$$\phi R_g \phi = \phi^2 \le \nu_1 \le \min\{\phi, R_g \phi\} = \phi$$

holds by taking g = e.

(3) Since $\nu_1 \leq \min\{\phi, R_g\phi\}$, one has

$$\nu_2 \ge \phi + R_g \phi - \min\{\phi, R_g \phi\} = \max\{\phi, R_g \phi\}.$$

As

$$\nu_2 \le \phi + R_g \phi - \phi R_g \phi = 1 - (1 - \phi)(1 - R_g \phi)$$

holds by $\nu_1 \ge \phi R_g \phi$, one has $\nu_2 \le 1$ by $\phi, R_g \phi \le 1$. Thus, we obtain $\nu_2 \in \mathcal{B}(I'_2)$ by

$$\|\nu_2\| = \|\phi\| + \|R_g\phi\| - \|\nu_1\| = I'_2.$$

(4) We fix $g' \in G$ and define $u(\nu) := \phi_1 * \nu(g')$ for an integrable function $\nu : G \to [0, 1]$. Since $\nu_1 \le \phi \le \nu_2$ and $\nu_1 \le R_g \phi \le \nu_2$ by (2) and (3), one has

$$u(\nu_1) \le u(\phi) \le u(\nu_2), \quad u(\nu_1) \le u(R_g \phi) \le u(\nu_2).$$
 (4.2)

If $u(\nu_1) = u(\nu_2)$, then the equalities are attained in (4.2) and hence one has $\xi_{\phi}(g') + \xi_{R_g\phi}(g') = \xi_{\nu_1}(g') + \xi_{\nu_2}(g')$. Thus, it suffices to show the result when $u(\nu_1) < u(\nu_2)$ holds. One has

$$\xi_{\phi}(g') \leq \tilde{f}_{u(\nu_{1})}^{u(\nu_{2})} \circ u(\phi) = \frac{u(\nu_{2} - \phi)\xi_{\nu_{1}}(g') + u(\phi - \nu_{1})\xi_{\nu_{2}}(g')}{u(\nu_{2} - \nu_{1})},$$

$$\xi_{R_{g}\phi}(g') \leq \tilde{f}_{u(\nu_{1})}^{u(\nu_{2})} \circ u(R_{g}\phi) = \frac{u(\nu_{2} - R_{g}\phi)\xi_{\nu_{1}}(g') + u(R_{g}\phi - \nu_{1})\xi_{\nu_{2}}(g')}{u(\nu_{2} - \nu_{1})}$$

by (4.2) and the convexity of f. Since $\nu_2 = \phi + R_g \phi - \nu_1$, one has

$$\xi_{\phi}(g') + \xi_{R_g\phi}(g') \le \frac{u(2\nu_2 - \phi - R_g\phi)\xi_{\nu_1}(g') + u(\phi + R_g\phi - 2\nu_1)(g')\xi_{\nu_2}(g')}{u(\nu_2 - \nu_1)}$$
$$= \xi_{\nu_1}(g') + \xi_{\nu_2}(g').$$

- (5) Since $\phi_1 * R_g \phi(g') = \phi_1 * \phi(g'g)$ holds for any $g' \in G$, one has $\xi_{R_g \phi}(g') = \xi_{\phi}(g'g)$. Thus, $\|\xi_{R_g \phi}\| = \|\xi_{\phi}\|$ holds and hence we obtain $2\|\xi_{\phi}\| \leq \|\xi_{\nu_1}\| + \|\xi_{\nu_2}\|$ by (4). In particular, $2S(I) \leq S(I'_1) + S(I'_2)$ holds by taking the upper bound on $\phi \in \mathcal{B}(I)$.
- Proof of Lemma 4.1. (1) For any $\phi \in \mathcal{B}(I)$, there is $\bar{\phi} \in \mathcal{B}(I')$ with $\phi \leq \bar{\phi}$. Thus, one has $S(I) \leq S(I')$ by $\xi_{\phi} \leq \xi_{\bar{\phi}}$. On the other hand, for any $\bar{\phi} \in \mathcal{B}(I')$, there is $\phi \in \mathcal{B}(I)$ with $\phi \leq \bar{\phi}$. Since $\xi_{\bar{\phi}} \xi_{\phi} \leq \phi_1 * (\bar{\phi} \phi)$ holds, one has

$$\|\xi_{\bar{\phi}}\| - \|\xi_{\phi}\| \le \|\phi_1 * (\bar{\phi} - \phi)\| = I' - I$$

by Lemma 2.2. Thus, $S(I') - S(I) \leq I' - I$ holds. In particular, S is Lipschitz continuous.

(2) Let $0 \leq J_1 < J_2 \leq m(G)$. Since $[J_1, J_2]$ is compact, there is

$$M := \min_{J_1 \le I \le J_2} \hat{S}_{J_1}^{J_2}(I)$$

by (1). It suffices to show that M = 0. The set

$$\mathcal{C} := \left\{ J_1 \le I \le J_2 \; \middle| \; \hat{S}_{J_1}^{J_2}(I) = M \right\}$$

is closed by (1).

We will show that $\mathcal{C} \cap (J_1, J_2)$ is open. Suppose $I \in \mathcal{C}$,

$$\max\{J_1, I^2/m(G), 2I - J_2\} < I'_1 \le I$$
(4.3)

and $I'_{2} := 2I - I'_{1}$. It suffices to show that $\hat{S}^{J_{2}}_{J_{1}}(I'_{1}) = \hat{S}^{J_{2}}_{J_{1}}(I'_{2}) = M$. Since $J_{1} < I'_{1} \le I'_{2} < J_{2}$ holds by (4.3), one has

$$2M \leq \hat{S}_{J_1}^{J_2}(I_1') + \hat{S}_{J_1}^{J_2}(I_2') = \tilde{S}_{J_1}^{J_2}(I_1') - S(I_1') + \tilde{S}_{J_1}^{J_2}(I_2') - S(I_2').$$
(4.4)

As

$$\tilde{S}_{J_1}^{J_2}(I_1') + \tilde{S}_{J_1}^{J_2}(I_2') = \frac{(2J_2 - I_1' - I_2')S(J_1) + (I_1' + I_2' - 2J_1)S(J_2)}{J_2 - J_1} = 2\tilde{S}_{J_1}^{J_2}(I)$$

holds by $I'_1 + I'_2 = 2I$, one has

$$\tilde{S}_{J_1}^{J_2}(I_1') - S(I_1') + \tilde{S}_{J_1}^{J_2}(I_2') - S(I_2') \le 2(\tilde{S}_{J_1}^{J_2}(I) - S(I)) = 2\hat{S}_{J_1}^{J_2}(I)$$
(4.5)

by Lemma 4.2 (5). Since $\hat{S}_{J_1}^{J_2}(I) = M$ holds by $I \in \mathcal{C}$, the equalities are attained in (4.4) and (4.5). Thus, one has $\hat{S}_{J_1}^{J_2}(I_1') = \hat{S}_{J_1}^{J_2}(I_2') = M$ and hence $\mathcal{C} \cap (J_1, J_2)$ is open.

Since $\mathcal{C} \cap (J_1, J_2)$ is clopen on the connected set (J_1, J_2) , one has either $\mathcal{C} \cap (J_1, J_2) = \emptyset$ or $(J_1, J_2) \subset \mathcal{C}$. In the case of $\mathcal{C} \cap (J_1, J_2) = \emptyset$, one has either $M = \hat{S}_{J_1}^{J_2}(J_1)$ or $M = \hat{S}_{J_1}^{J_2}(J_2)$, and hence we obtain M = 0 by

$$\hat{S}_{J_1}^{J_2}(J_1) = \hat{S}_{J_1}^{J_2}(J_2) = 0.$$
(4.6)

In the case of $(J_1, J_2) \subset C$, one has $C = [J_1, J_2]$ because C is closed. Thus, we obtain M = 0 by (4.6).

- (3) Since $\xi_{\phi} = 0$ holds for any $\phi \in \mathcal{B}(t)$ by (2.5), one has S(t) = 0.
- (4) As $\phi_1 * \phi \leq 1$ holds by (2.4), one has $\xi_{\phi} \leq \tilde{f}_0^1 \circ (\phi_1 * \phi) = (1-t)(\phi_1 * \phi)$ by the convexity of f. Since $\|\xi_{\phi}\| \leq (1-t)\|\phi_1 * \phi\| = (1-t)I$ holds by Lemma 2.2, we obtain $S(I) \leq (1-t)I$. \Box

5 Proof of Lemma 1.2 when $m(G) = \infty$

In this section, we show (4.1) when $m(G) = \infty$ holds. It suffices to show

$$S(I_2) \le (1-t)(I_2 - t) \tag{5.1}$$

for any $I_2 \ge t$ by (4.1). If $I \ge I_2$ holds, then

$$S(I_2) \le \tilde{S}_t^I(I_2) = \frac{(I - I_2)S(t) + (I_2 - t)S(I)}{I - t}$$

holds by Lemma 4.1 (2). In addition, one has

$$\frac{(I-I_2)S(t) + (I_2 - t)S(I)}{I - t} \le \frac{(1-t)(I_2 - t)I}{I - t}$$

by Lemma 4.1 (3) and (4). Thus,

$$S(I_2) \le \lim_{I \to \infty} \frac{(1-t)(I_2-t)I}{I-t} = (1-t)(I_2-t)$$

holds because $I_2 \leq I < \infty (= m(G))$ is arbitrary. Therefore, when $m(G) = \infty$ holds, we obtain (5.1) and hence Lemma 1.2.

6 Proof of Lemma 1.2 when $m(G) < \infty$

In this section, we show (4.1) when $m(G) < \infty$ holds. We simply write f for f_t in (1.3). It suffices to show (4.1) when $\|\phi_1\| = 1$ and $\|\phi_2\| > 0$ hold by Remark 2.4 (6). In this case, $G_0 \subset G$ is an open subgroup because

$$0 \le 1 - t \le m(G) - \|\phi_2\| < m(G) = \operatorname{vol}(G_0)$$

holds by Remark 2.4 (3) and (1.5) (we note that G' is open if and only if vol(G') > 0, for any closed subgroup $G' \subset G$). Thus, we can take the Haar measure of G_0 to be the same as of G. That is,

$$\|\phi\| = \sum_{gG_0 \in G/G_0} \|\phi\|_{gG_0}$$
(6.1)

holds for any integrable function $\phi \colon G \to \mathbb{R}_{>0}$.

We give a lemma (Lemma 6.2) to prove Lemma 1.2. For a measurable function $\phi \colon G \to \mathbb{R}_{\geq 0}$, we define a function $\phi^* \colon \mathbb{R} \times G/G_0 \to \{0,1\}$.

Definition 6.1. For a measurable function $\phi \colon G \to \mathbb{R}_{\geq 0}$, the function $\phi^* \colon \mathbb{R} \times G/G_0 \to \{0, 1\}$ is defined as

$$\phi^*(x, gG_0) := \phi_{(P(g))}(x), \quad P(g) := \|\phi\|_{gG_0} = \int_{gG_0} \phi(g') dg'$$

where $\phi_{(I)}$ is the same as in Example 2.1 (1).

Lemma 1.2 is deduced by the following lemma.

Lemma 6.2. If integrable functions $\phi_1, \phi_2: G \to [0,1]$ satisfy (1.5), then

$$\|f \circ (\phi_1 * \phi_2)\|_{gG_0} \le \|f \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{gG_0\}}$$

holds for any $g \in G$. In particular, one has $||f \circ (\phi_1 * \phi_2)|| \le ||f \circ (\phi_1^* * \phi_2^*)||$.

The estimate $||f \circ (\phi_1 * \phi_2)|| \le ||f \circ (\phi_1^* * \phi_2^*)||$ follows from

$$\|f \circ (\phi_1 * \phi_2)\| = \sum_{gG_0 \in G/G_0} \|f \circ (\phi_1 * \phi_2)\|_{gG_0}$$
$$\leq \sum_{gG_0 \in G/G_0} \|f \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{gG_0\}}$$
$$= \|f \circ (\phi_1^* * \phi_2^*)\|$$

by (6.1).

Example 6.3. If $G = G_0$, then one has $\phi_1^* = \phi_{(1)}$ and $\phi_2^* = \phi_{(\|\phi_2\|)}$ by identifying $\mathbb{R} \times G/G_0$ with \mathbb{R} . Thus, $\|f \circ (\phi_1^* * \phi_2^*)\| = \|f \circ \psi_{1,\|\phi_2\|}\|$ holds and hence Lemma 6.2 is equivalent to Lemma 1.2 and (4.1) by (2.2).

Since
$$m(\mathbb{R} \times G/G_0) = \infty$$
, $\|\phi_1\| = \|\phi_1^*\| = 1$ and $\|\phi_2\| = \|\phi_2^*\|$, one has
 $\|f \circ (\phi_1 * \phi_2)\| \le \|f \circ (\phi_1^* * \phi_2^*)\| \le (1-t)(\|\phi_2\| - t)$

by the argument in Section 5. Thus, we obtain Lemma 1.2 from Lemma 6.2.

Lemma 6.2 is shown in two steps. We will show Lemma 6.2 in the case of $G = G_0$ (that is, G is connected) in Subsection 6.1 at first, and for arbitrary G in Subsection 6.2 by using the case of $G = G_0$.

6.1 Proof of Lemma 6.2 when $G = G_0$

In this subsection, we show Lemma 6.2 when $G = G_0$ holds. It suffices to show (4.1) by Example 6.3. We have

$$S(I) \le \tilde{S}_t^{I'}(I) = \frac{(I' - I)S(t) + (I - t)S(I')}{I' - t}$$

by Lemma 4.1 (2) for I' := m(G) + t - 1. Since $G = G_0$ and t = 1 + I' - m(G), it follows that

$$S(I') = I' - tm(G) = (1 - t)(I' - t)$$

by Lemma 3.1 (2). Thus,

$$S(I) \le \frac{(I'-I)S(t) + (I-t)S(I')}{I'-t} = (1-t)(I-t)$$

holds by Lemma 4.1 (3). Therefore, we obtain (4.1) and Lemma 6.2 when $G = G_0$ holds.

6.2 Proof of Lemma 6.2 for any G

In this subsection, we show Lemma 6.2 for any G. It suffices to show it when g = e by the following lemma and remark.

Lemma 6.4. One has $(R_g\phi)^* = R_{(0,gG_0)}(\phi^*)$ for any integrable function $\phi: G \to \mathbb{R}_{\geq 0}$ and $g \in G$, where $R_g\phi$ is the same as in Lemma 4.2.

Proof. Let P(g) be the same as in Definition 6.1. One has

$$(R_g\phi)^*(x,g'G_0) = \phi_{(\|R_g\phi\|_{g'G_0})}(x) = \phi_{(P(g'g))}(x) = \phi^*(x,g'gG_0) = R_{(0,gG_0)}(\phi^*)(x,g'G_0)$$

for any $x \in \mathbb{R}$ and $g' \in G$.

Remark 6.5. Under the hypothesis of Lemma 6.2,

$$||f \circ (\phi_1 * \phi_2)||_{gG_0} = ||f \circ R_g(\phi_1 * \phi_2)||_{G_0} = ||f \circ (\phi_1 * R_g \phi_2)||_{G_0}$$

and similarly

$$\|f \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{gG_0\}} = \|f \circ (\phi_1^* * R_{(0,gG_0)}(\phi_2^*))\|_{\mathbb{R} \times \{G_0\}}$$

hold. Thus, one has

$$\|f \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{gG_0\}} = \|f \circ (\phi_1^* * (R_g \phi_2)^*)\|_{\mathbb{R} \times \{G_0\}}$$

by Lemma 6.4, and hence it suffices to show Lemma 6.2 in the case of g = e by replacing ϕ_2 with $R_g \phi_2$.

We show Lemma 6.2 when g = e holds. We define

$$\phi_1 *_{G_0} \phi_2(g') := \int_{G_0} \phi_1(g'') \phi_2(g''^{-1}g') dg''$$

for $g' \in G_0$. We also define integrable functions $L_{g''}\phi, R_{g''}\phi: G \to \mathbb{R}_{\geq 0}$ as

$$L_{g''}\phi(g') := \phi(g''^{-1}g'), \quad R_{g''}\phi(g') := \phi(g'g'')$$

for $g'' \in G$ and an integrable function $\phi \colon G \to \mathbb{R}_{\geq 0}$. We give the following lemma to prove Lemma 6.2.

Lemma 6.6. For any $g' \in G_0$, one has

$$\phi_1 * \phi_2(g') = \sum_{g''G_0 \in G/G_0} (L_{g''}\phi_1) *_{G_0} (R_{g''}\phi_2)(g''g'g''^{-1}).$$

Proof. Since $g''G_0g''^{-1} = G_0$ holds for any $g'' \in G$,

$$\begin{split} \phi_1 * \phi_2(g') &= \int_G \phi_1(g''') \phi_2(g'''^{-1}g') dg''' \\ &= \sum_{g''G_0 \in G/G_0} \int_{G_0} \phi_1(g''^{-1}g'''') \phi_2(g''''^{-1}g''g') dg'''' \\ &= \sum_{g''G_0 \in G/G_0} \int_{G_0} L_{g''} \phi_1(g''') R_{g''} \phi_2(g''''^{-1}g''g'g''^{-1}) dg'''' \\ &= \sum_{g''G_0 \in G/G_0} (L_{g''}\phi_1) *_{G_0} (R_{g''}\phi_2)(g''g'g''^{-1}) \end{split}$$

is obtained by (6.1).

Example 6.7. Let $P_1(g'') := \|\phi_1\|_{g''G_0}$ and $P_2(g'') := \|\phi_2\|_{g''G_0}$ for $g'' \in G$. The identity component of $\mathbb{R} \times G/G_0$ is $\mathbb{R} \times \{G_0\}$, and hence

$$\phi_1^* * \phi_2^*(x, G_0) = \sum_{g''G_0 \in G/G_0} \phi_{(P_1(g''^{-1}))} * \phi_{(P_2(g''))}(x) = \sum_{g''G_0 \in G/G_0} \psi_{P_1(g''^{-1}), P_2(g'')}(x)$$

for any $x \in \mathbb{R}$ by Lemma 6.6, where ψ_{I_1,I_2} is the same as in Example 2.1 (1). Since $\psi_{P_1(g''^{-1}),P_2(g'')}(x)$ is an even and monotonically decreasing function on $x \ge 0$, the convolution $\phi_1^* * \phi_2^*(x,G_0)$ is also an even function with respect to $x \in \mathbb{R}$ that is monotonically decreasing on $x \ge 0$.

We show Lemma 6.2 by Example 6.3, Remark 6.5, Lemma 6.6 and Example 6.7. It suffices to show it when g = e holds by Remark 6.5. We set $t_0 := \phi_1^* * \phi_2^*(0, G_0)$ and divide the proof into the following two cases.

(1) The case of $t \leq t_0$. Since $\phi_1^* * \phi_2^*(1 + \|\phi_2\|, G_0) = 0 \leq t \leq t_0$ holds and $\phi_1^* * \phi_2^*$ is continuous, there is $x_0 \geq 0$ with $\phi_1^* * \phi_2^*(x_0, G_0) = t$ by the intermediate value theorem. We set $t_{g''} := \psi_{P_1(g''^{-1}), P_2(g'')}(x_0)$ for $g'' \in G$. Then

$$t = \sum_{g''G_0 \in G/G_0} t_{g'}$$

holds by Example 6.7, and hence one has

$$\phi_1 * \phi_2(g') - t = \sum_{g''G_0 \in G/G_0} (\omega_{g''}(g''g'g''^{-1}) - t_{g''})$$

for any $g' \in G_0$ by Lemma 6.6, where we write $\omega_{q''} := (L_{q''}\phi_1) *_{G_0} (R_{q''}\phi_2)$. Thus, one has

$$\|f \circ (\phi_1 * \phi_2)\|_{G_0} = \sum_{g'' G_0 \in G/G_0} \int_{\{g' \in G_0 | \phi_1 * \phi_2(g') \ge t\}} (\omega_{g''}(g''g'g''^{-1}) - t_{g''})dg'.$$
(6.2)

Lemma 6.8. For any $g'' \in G$, one has

$$\int_{\{g'\in G_0|\phi_1*\phi_2(g')\geq t\}} (\omega_{g''}(g''g'g''^{-1}) - t_{g''})dg' \leq \|\psi_{P_1(g''^{-1}),P_2(g'')} - t_{g''}\|_{(-x_0,x_0)}$$

Proof. Since $\omega_{g''} - t_{g''} \leq f_{t_{g''}} \circ \omega_{g''}$ and $g'' G_0 g''^{-1} = G_0$ hold, we have

$$\int_{\{g'\in G_0|\phi_1*\phi_2(g')\geq t\}} (\omega_{g''}(g''g'g''^{-1}) - t_{g''})dg' \leq \|f_{t_{g''}} \circ \omega_{g''}\|_{G_0}.$$
(6.3)

Then $||L_{g''}\phi_1||_{G_0} \le ||\phi_1|| = 1$ and $||R_{g''}\phi_2||_{G_0} \le ||\phi_2||$ hold, and hence one has

$$||L_{g''}\phi_1||_{G_0} + ||R_{g''}\phi_2||_{G_0} \le 1 + ||\phi_2|| \le m(G) + t$$

by (1.5). By Example 6.3 and Subsection 6.1, we have

$$\|f_{t_{g''}} \circ \omega_{g''}\|_{G_0} \le \|f_{t_{g''}} \circ \psi_{P_1(g''^{-1}), P_2(g'')}\|_{\mathbb{R}} = \|\psi_{P_1(g''^{-1}), P_2(g'')} - t_{g''}\|_{(-x_0, x_0)},$$

where the last equality is due to the monotonicity of $\psi_{P_1(g''^{-1}),P_2(g'')}$ and the definition of $t_{g''}$. Thus, we obtain

$$\int_{\{g'\in G_0|\phi_1*\phi_2(g')\geq t\}} (\omega_{g''}(g''g'g''^{-1}) - t_{g''})dg' \leq \|f_{t_{g''}} \circ \omega_{g''}\|_{G_0}$$
$$\leq \|\psi_{P_1(g''^{-1}),P_2(g'')} - t_{g''}\|_{(-x_0,x_0)}$$

by (6.3).

Since

$$\sum_{g''G_0 \in G/G_0} \|\psi_{P_1(g''^{-1}), P_2(g'')} - t_{g''}\|_{(-x_0, x_0)} = \|f \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{G_0\}}$$

holds by Example 6.7, we obtain

$$\|f \circ (\phi_1 * \phi_2)\|_{G_0} \le \|f \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{G_0\}}$$

by (6.2) and Lemma 6.8.

(2) The case of $t \ge t_0$. One has $||f \circ (\phi_1 * \phi_2)||_{G_0} \le ||f_{t_0} \circ (\phi_1 * \phi_2)||_{G_0}$. In addition,

$$\|f_{t_0} \circ (\phi_1 * \phi_2)\|_{G_0} \le \|f_{t_0} \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{G_0\}}$$

holds according to (1). One has $\phi_1^* * \phi_2^*(x, G_0) \leq t_0$ for any $x \in \mathbb{R}$ by Example 6.7 and hence

$$\|f \circ (\phi_1 * \phi_2)\|_{G_0} = \|f \circ (\phi_1^* * \phi_2^*)\|_{\mathbb{R} \times \{G_0\}} = 0.$$

Thus, we obtain $||f \circ (\phi_1 * \phi_2)|| \le ||f \circ (\phi_1^* * \phi_2^*)||$.

7 Proof of Theorem 1.1 for any convex function f

In this section, we show Theorem 1.1 by Lemma 1.2. It suffices to show it when $\|\phi_1\| = 1$ holds by Remark 2.4 (6). The function $f_{(n)}: [0,1] \to \mathbb{R}$ is defined as

$$f_{(n)}(y) := n\left(f\left(\frac{1}{n}\right)y + 2\sum_{k=1}^{n-1}\hat{f}_{(k-1)/n}^{(k+1)/n}\left(\frac{k}{n}\right)f_{k/n}(y)\right)$$

for $n \in \mathbb{Z}_{\geq 1}$ and function $f: [0,1] \to \mathbb{R}$, where $\hat{f}_{y_1}^{y_2}$ is the same as in Definition 2.3 and f_t is as defined in (1.3). We show the following lemma to prove Theorem 1.1.


Lemma 7.1. Let $f: [0,1] \to \mathbb{R}$ be a convex function with f(0) = 0.


- (1) One has $f_{(n)}(y) = \tilde{f}_{(j-1)/n}^{j/n}(y)$ for any $j = 1, \dots, n$ and $(j-1)/n \le y \le j/n$, where $\tilde{f}_{y_1}^{y_2}$ is the same as in Definition 2.3. In particular, $f(y) \le f_{(n)}(y)$ holds for any $0 \le y \le 1$, and the equality is attained at $y = 0, 1/n, 2/n, \dots, 1 1/n, 1$ (Figure 1).
- (2) For any 0 < y < 1, one has

$$f_{(n)}(y) \le f(y) + \frac{\hat{f}_0^1(y)}{4ny(1-y)}.$$

- (3) One has $\lim_{n\to\infty} f_{(n)}(y) = f(y)$ for any $0 \le y \le 1$. That is, $f_{(n)}$ converges pointwise to f.
- (4) One has $f_{(n)}(y) \leq f(1)y$ for any $n \in \mathbb{Z}_{\geq 1}$ and $0 \leq y \leq 1$.
- (5) For any integrable function $\psi \colon G \to [0,1]$, one has

$$\lim_{n\to\infty}\int_G f_{(n)}\circ\psi(g)dg=\int_G f\circ\psi(g)dg.$$





Proof. (1) Since $(j-1)/n \le y \le j/n$ holds, one has

$$f_{k/n}(y) = \begin{cases} y - \frac{k}{n} & \text{if } k \le j - 1, \\ 0 & \text{if } k \ge j \end{cases}$$

for any $k = 1, \dots, n$. Then

$$\hat{f}_{(k-1)/n}^{(k+1)/n}\left(\frac{k}{n}\right) = \frac{1}{2}\left(f\left(\frac{k-1}{n}\right) + f\left(\frac{k+1}{n}\right)\right) - f\left(\frac{k}{n}\right)$$

holds, and hence one has

$$f_{(n)}(y) = nf\left(\frac{1}{n}\right)y + \sum_{k=1}^{j-1} \left(f\left(\frac{k-1}{n}\right) + f\left(\frac{k+1}{n}\right) - 2f\left(\frac{k}{n}\right)\right)(ny-k)$$

Since

$$\sum_{k=1}^{j-1} f\left(\frac{k-1}{n}\right) (ny-k) = f\left(\frac{j-1}{n}\right) (j-ny) + \sum_{k=1}^{j-1} f\left(\frac{k}{n}\right) (ny-k-1)$$

holds by f(0) = 0 and

$$nf\left(\frac{1}{n}\right)y + \sum_{k=1}^{j-1} f\left(\frac{k+1}{n}\right)(ny-k) = f\left(\frac{j}{n}\right)(ny-j+1) + \sum_{k=1}^{j-1} f\left(\frac{k}{n}\right)(ny-k+1)$$

holds, one has

$$f_{(n)}(y) = \tilde{f}_{(j-1)/n}^{j/n}(y) + \sum_{k=1}^{j-1} f\left(\frac{k}{n}\right) (ny - k - 1 + ny - k + 1 - 2(ny - k)) = \tilde{f}_{(j-1)/n}^{j/n}(y).$$

In particular, $f(y) \leq f_{(n)}(y)$ holds for any $0 \leq y \leq 1$ by the convexity of f, and the equality is attained at $y = 0, 1/n, 2/n, \dots, 1 - 1/n, 1$.

(2) It suffices to show the result when $j = 1, \dots, n$ and $(j - 1)/n \le y \le j/n$ hold. One has

$$f_{(n)}(y) \le \tilde{f}_{(j-1)/n}^{j/n}(y) = (j-ny)f\left(\frac{j-1}{n}\right) + (ny-j+1)f\left(\frac{j}{n}\right)$$
(7.1)

by (1). Then

$$f\left(\frac{j-1}{n}\right) \le \tilde{f}_0^y\left(\frac{j-1}{n}\right) = \frac{(j-1)f(y)}{ny}$$

holds by $0 \le (j-1)/n \le y$, f(0) = 0 and the convexity of f, and similarly

$$f\left(\frac{j}{n}\right) \le \tilde{f}_y^1\left(\frac{j}{n}\right) = \frac{(n-j)f(y) + (j-ny)f(1)}{n(1-y)}$$

holds by $y \leq j/n \leq 1$. Thus, one has

$$\begin{split} &f_{(n)}(y) \\ &\leq \frac{(j-ny)(j-1)f(y)}{ny} + \frac{(ny-j+1)((n-j)f(y)+(j-ny)f(1))}{n(1-y)} \\ &= \frac{f(y)-f(1)y}{ny(1-y)} \cdot j^2 + \frac{(2ny+1)(f(1)y-f(y))}{ny(1-y)} \cdot j + f(y) + \frac{(ny+1)(f(y)-yf(1))}{1-y} \\ &= f(y) + \frac{f(y)-f(1)y}{ny(1-y)} \left(\left(j-ny-\frac{1}{2} \right)^2 - \frac{1}{4} \right) \\ &= f(y) + \frac{\hat{f}_0^1(y)(1-(2ny+1-2j)^2)}{4ny(1-y)}. \end{split}$$

Since $\hat{f}_0^1(y) \ge 0$ holds by the convexity of f, we obtain

$$f_{(n)}(y) \le f(y) + \frac{\hat{f}_0^1(y)(1 - (2ny + 1 - 2j)^2)}{4ny(1 - y)} \le f(y) + \frac{\hat{f}_0^1(y)}{4ny(1 - y)}$$

(3) If y = 0, 1 holds, then one has $f_{(n)}(y) = f(y)$ by (1). Thus, it suffices to show the result when 0 < y < 1. One has

$$f(y) \le f_{(n)}(y) \le f(y) + \frac{a}{n}, \quad a := \frac{\hat{f}_0^1(y)}{4y(1-y)}$$

by (1) and (2). Since

$$\lim_{n \to \infty} \left(f(y) + \frac{a}{n} \right) = f(y),$$

we obtain

$$\lim_{n \to \infty} f_{(n)}(y) = f(y).$$

(4) It suffices to show the result when $j = 1, \dots, n$ and $(j-1)/n \le y \le j/n$ hold. One has

$$f_{(n)}(y) \le (j - ny)f\left(\frac{j - 1}{n}\right) + (ny - j + 1)f\left(\frac{j}{n}\right)$$

by (7.1). In addition,

$$f\left(\frac{j-1}{n}\right) \le \tilde{f}_0^1\left(\frac{j-1}{n}\right) = \frac{(j-1)f(1)}{n}, \quad \text{and} \quad f\left(\frac{j}{n}\right) \le \tilde{f}_0^1\left(\frac{j}{n}\right) = \frac{jf(1)}{n}$$

hold by f(0) = 0 and the convexity of f. Thus, one has

$$f_{(n)}(y) \le \frac{(j-ny)(j-1)f(1)}{n} + \frac{(ny-j+1)jf(1)}{n} = f(1)y.$$

(5) Since $f_{(n)} \ge f$ by (1), one has

$$\liminf_{n \to \infty} \int_G f_{(n)} \circ \psi(g) dg \ge \int_G f \circ \psi(g) dg.$$
(7.2)

One has $f_{(n)} \circ \psi(g) \leq f(1)\psi(g)$ for any $n \in \mathbb{Z}_{\geq 1}$ by (4). Since ψ is integrable,

$$\limsup_{n \to \infty} \int_G f_{(n)} \circ \psi(g) dg \le \int_G f \circ \psi(g) dg$$
(7.3)

holds by applying Fatou's lemma to $f(1)\psi - f_{(n)} \circ \psi$. Thus, we obtain

$$\lim_{n \to \infty} \int_G f_{(n)} \circ \psi(g) dg = \int_G f \circ \psi(g) dg$$

by (7.2) and (7.3).

Proof of Theorem 1.1. It suffices to show the result when $\|\phi_1\| = 1$ holds by Remark 2.4 (6). Since

$$\|\phi_1 * \phi_2\| = \|\psi_{1,\|\phi_2\|}\| = \|\phi_2\| < \infty$$
(7.4)

holds by Lemma 2.2, one has

$$\int_{G} f \circ (\phi_1 * \phi_2)(g) dg = \lim_{n \to \infty} \int_{G} f_{(n)} \circ (\phi_1 * \phi_2)(g) dg$$
(7.5)

by Lemma 7.1 (5). By the definition of $f_{(n)}$,

$$\int_{G} f_{(n)} \circ (\phi_1 * \phi_2)(g) dg = n \left(f\left(\frac{1}{n}\right) \|\phi_1 * \phi_2\| + 2 \sum_{k=1}^{n-1} \hat{f}_{(k-1)/n}^{(k+1)/n}\left(\frac{k}{n}\right) \|f_{k/n} \circ (\phi_1 * \phi_2)\| \right)$$
(7.6)

holds. Since (2.2) holds by Lemma 1.2 and $\hat{f}_{(k-1)/n}^{(k+1)/n}(k/n) \ge 0$ holds by the convexity of f, one has

$$\hat{f}_{(k-1)/n}^{(k+1)/n}\left(\frac{k}{n}\right)\|f_{k/n}\circ(\phi_1*\phi_2)\| \le \hat{f}_{(k-1)/n}^{(k+1)/n}\left(\frac{k}{n}\right)\|f_{k/n}\circ\psi_{1,\|\phi_2\|}\|$$

for any $k = 1, \dots, n-1$. Thus,

$$\int_{G} f_{(n)} \circ (\phi_{1} * \phi_{2})(g) dg \leq n \left(f\left(\frac{1}{n}\right) \|\psi_{1,\|\phi_{2}\|}\| + 2 \sum_{k=1}^{n-1} \hat{f}_{(k-1)/n}^{(k+1)/n} \left(\frac{k}{n}\right) \|f_{k/n} \circ \psi_{1,\|\phi_{2}\|}\| \right) \\
= \int_{\mathbb{R}} f_{(n)} \circ \psi_{1,\|\phi_{2}\|}(x) dx$$
(7.7)

holds by (7.4) and (7.6). Since

$$\lim_{n \to \infty} \int_{\mathbb{R}} f_{(n)} \circ \psi_{1, \|\phi_2\|}(x) dx = \int_{\mathbb{R}} f \circ \psi_{1, \|\phi_2\|}(x) dx \tag{7.8}$$

holds by Lemma 7.1 (5), we obtain

$$\int_{G} f \circ (\phi_{1} * \phi_{2})(g) dg \leq \int_{\mathbb{R}} f \circ \psi_{1, \|\phi_{2}\|}(x) dx = 2 \int_{0}^{1} f(y) dy + (\|\phi_{2}\| - 1) f(1)$$
(7.5) (7.7) and (7.8)

by (2.1), (7.5), (7.7), and (7.8).

Acknowledgement

The author would like to thank his advisor Toshiyuki Kobayashi for his support. The author is also grateful to Yuichiro Tanaka and Toshihisa Kubo for helpful comments. The author also would like to thank the reviewers for the careful comments.

Funding

This work was supported by JSPS KAKENHI Grant Number JP19J22628 and Leading Graduate Course for Frontiers of Mathematical Sciences and Physics (FMSP).

References

- [BL76] H. J. Brascamp and E. H. Lieb. "On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation". J. Functional Analysis 22.4 (1976), pp. 366– 389.
- [Bru87] H. Brunn. Über Ovale und Eiflächen. Akademische Buchdruckerei von R. Straub, 1887.
- [Chr19] M. Christ. "Near equality in the Riesz-Sobolev inequality". Acta Math. Sin. (Engl. Ser.) 35.6 (2019), pp. 783–814.
- [CI22] M. Christ and M. Iliopoulou. "Inequalities of Riesz-Sobolev type for compact connected Abelian groups". Amer. J. Math. 144.5 (2022), pp. 1367–1435.
- [GR05] B. Green and I. Z. Ruzsa. "Sum-free sets in abelian groups". Israel J. Math. 147 (2005), pp. 157–188.
- [Gri19] J. T. Griesmer. "Semicontinuity of structure for small sumsets in compact abelian groups". *Discrete Anal.* (2019), Paper No. 18, 46.
- [HM53] R. Henstock and A. M. Macbeath. "On the measure of sum-sets. I. The theorems of Brunn, Minkowski, and Lusternik". Proc. London Math. Soc. (3) 3 (1953), pp. 182– 194.
- [HR79] E. Hewitt and K. A. Ross. Abstract harmonic analysis. Vol. I. Second. Vol. 115. Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin-New York, 1979.
- [JT20] Y. Jing and C.-M. Tran. "Minimal and nearly minimal measure expansions in connected unimodular groups". *arXiv preprint arXiv:2006.01824* (2020).
- [JTZ21] Y. Jing, C.-M. Tran, and R. Zhang. "A nonabelian Brunn-Minkowski inequality". arXiv preprint arXiv:2101.07782 (2021).
- [Kem64] J. H. B. Kemperman. "On products of sets in a locally compact group". Fund. Math. 56 (1964), pp. 51–68.
- [Kne56] M. Kneser. "Summenmengen in lokalkompakten abelschen Gruppen". Math. Z. 66 (1956), pp. 88–110.
- [Lus35] L. A. Lusternik. "Die brunn-minkowskische ungleichung für beliebige messbare mengen". *CR Acad. Sci. URSS* 8 (1935), pp. 55–58.
- [Mac53] A. M. Macbeath. "On measure of sum sets. II. The sum-theorem for the torus". *Proc. Cambridge Philos. Soc.* 49 (1953), pp. 40–43.
- [McC69] M. McCrudden. "On the Brunn-Minkowski coefficient of a locally compact unimodular group". *Proc. Cambridge Philos. Soc.* 65 (1969), pp. 33–45.

- [Min96] H. Minkowski. *Geometrie der Zahlen*. Bibliotheca Mathematica Teubneriana, Band 40. Johnson Reprint Corp., New York-London, 1896, pp. vii+256.
- [Pol74] J. M. Pollard. "A generalisation of the theorem of Cauchy and Davenport". J. London Math. Soc. (2) 8 (1974), pp. 460–462.
- [Rai39] D. Raikov. "On the addition of point-sets in the sense of Schnirelmann". Rec. Math. [Mat. Sbornik] N.S. 5(47) (1939), pp. 425–440.
- [Rie30] F. Riesz. "Sur Une Inegalite Integarale". J. London Math. Soc. 5.3 (1930), pp. 162– 168.
- [Ruz92] I. Z. Ruzsa. "A concavity property for the measure of product sets in groups". Fund. Math. 140.3 (1992), pp. 247–254.
- [Shi55] A. Shields. "Sur la mesure d'une somme vectorielle". Fund. Math. 42 (1955), pp. 57–60.
- [Sob38] S. L. Sobolev. "On a theorem of functional analysis". *Mat. Sbornik* 4 (1938), pp. 471–497.
- [Tao12] T. Tao. Spending symmetry. 2012.
- [Tao13] T. Tao. "Noncommutative sets of small doubling". European J. Combin. 34.8 (2013), pp. 1459–1465.
- [Tao18] T. Tao. "An inverse theorem for an inequality of Kneser". *Proc. Steklov Inst. Math.* 303.1 (2018), pp. 193–219.
- [VD31] D. Van Dantzig. "Studien over topologische Algebra". PhD thesis. HJ Paris Amsterdam, 1931.
- [Wei40] A. Weil. L'intégration dans les groupes topologiques et ses applications. Actualités Scientifiques et Industrielles, No. 869. Hermann et Cie., Paris, 1940.

Takashi Satomi: Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba Meguro-ku Tokyo 153-8914, Japan.

E-mail: tsatomi@ms.u-tokyo.ac.jp