Analytic resurgence in the O(4) model

Zoltán Bajnok,¹ János Balog² and István Vona³

Holographic QFT Group, Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics H-1525 Budapest 114, P.O.B. 49, Hungary

ABSTRACT: We study the perturbative expansion of the ground state energy in the presence of an external field coupled to a conserved charge in the integrable two-dimensional O(4) nonlinear sigma model. By solving Volin's algebraic equations for the perturbative coefficients we study the large order asymptotic behaviour of the perturbative series analytically. We confirm the previously numerically found leading behaviour and study the nearest singularities of the Borel transformed series and the associated alien derivatives. We find a "resurgence" behaviour: the leading alien derivatives can be expressed in terms of the original perturbative series. A simplified 'toy' model is also considered: here the perturbative series can be found in a closed form and the resurgence properties are very similar to that found in the real problem.

¹bajnok.zoltan@wigner.hu

²balog.janos@wigner.hu

³vona.istvan@wigner.hu

Contents

1	Introduction	2
	1.1 TBA calculation of the ground-state energy density	3
	1.2 Perturbative expansion of the TBA equation, Volin's method [1, 2]	4
	1.3 $O(4)$ Volin equations	5
2	Summary	7
	2.1 Properties of the $\varepsilon_p(z)$ series	7
	2.2 "Derivation" of the naive ansatz	10
	2.3 The correct ansatz	11
	2.4 Asymptotic expansion of ρ	13
3	Conclusion	13
A	Simplifying Volin's equations	15
	A.1 Properties of the a_n coefficients	15
	A.2 $c_{n,m}$ equations	15
в	Analytic proof of the asymptotic expansion (2.6)	16
	B.1 Some definitions, notations, relations	16
	B.2 $E_{a,N}$ resurgence	17
	B.3 Proof of the statements	18
С	Consistency of the conjectured asymptotics	21
D	Resurgence	24
\mathbf{E}	Asymptotic expansion of ρ	26
\mathbf{F}	Alien derivatives	28
G	The $\eta_k = 1$ model	29
	G.1 <i>a</i> -dependence	31
	G.2 The alien derivatives of ϵ_p for the $\eta_k = 1$ model	32

1 Introduction

The two dimensional $O(N) \sigma$ models provide an ideal playground for theoretical particle and statistical physics, where non-perturbative phenomena can be analyzed exactly without relying on approximate methods. These models share many features with QCD including asymptotic freedom in perturbation theory and the dynamical generation of a mass scale, which breaks the classical scale invariance in the quantum theory. These models were the first, where the exact S-matrix was determined [3]. By analysing the behaviour of the groundstate energy density in a magnetic field [4] the perturbative scale was the first time exactly related to the dynamically generated mass [5, 6]. This energy density then can be expanded in perturbation theory and its asymptotic series revealed non-perturbative sectors [1, 7–9] related to renormalons, which exist also in QCD [10]. The theory which describes how the large order asymptotics coefficients "resurge" themselves into the non-perturbative contributions is called resurgence¹, which aims at constructing the full non-perturbative ambiguity free trans-series for the physical observables. The exact solvability of the $O(N) \sigma$ models gives a hope to establish the complete resurgence structure exactly and to provide the sought trans-series, which could indicate similar behaviours for analogous realistic theories, such as for QCD.

The resurgence program has been analyzed in many two dimensional and related quantum field theories including also supersymmetric ones [14–21] and some of them focuses on the semiclassical domain of O(N) and principal chiral models [22–25]. Recently there have been active research and significant progress on relativistic integrable quantum field theories [7, 26, 27] and their non-relativistic statistical physical counterparts [28–30]. Resurgence proved to be a useful tool also in non-integrable quantum field theories, such as in the ϕ^4 theory [31, 32] and also in analysing the hydrodynamics of the Yang Mills plasma [33, 34]. There is no asymptotically free QFT, however, where the full resurgence structure was derived. In this paper we would like to make the first step into this direction by analysing the O(4) model, which is the simplest amongst the O(N) theories.

In the integrable $O(N) \sigma$ models we can couple one of the O(N) conserved charges to a magnetic field and analyse the groundstate energy density in two different ways [5, 6]. On the one hand it can be calculated perturbatively in terms of $\Lambda_{\overline{MS}}$ for large magnetic fields. On the other hand, integrability enables to write an exact integral equation for the energy density, which can be expanded using Wiener-Hopf technics. The comparison of the first order terms provided the celebrated mass/coupling relation. Further perturbative terms turned to be decisive in the AdS/CFT correspondence, where the O(6) model plays a crucial role [35]. The calculation of higher order terms are cumbersome and the breakthrough came from the integrable side, where Volin could manage to translate the integral equation into a system of algebraic ones [1, 2]. These recursive equations could be solved for the first 20-50 terms analytically, which allowed to observe a factorial growth [1, 2, 7]. It also helped to identify the closest singularities on the Borel plane, which corresponds typically to UV and IR renormalons, diagrams in perturbation theory, where the asymptotic growths arise from

¹See [11–13] for introductions and references.

integrating over UV or IR domains in specific renormalon diagrams [36]. In the large N limit the leading of such diagrams were identified and their contributions were calculated exactly, which agreed with the solution of the algebraic equations [26].

In our recent works [8, 9], we focused on the O(4) model, where the algebraic equations simplified drastically. We solved these equations numerically with very high precision, from which we conjectured analytical expressions for their leading and subleading factorial behaviours. We numerically identified logarithmic cuts on the Borel plane. We investigated the functions multiplying these cuts (alien derivatives) perturbatively from the asymptotics of the original perturbative series. By using high precision numerics we could identify and characterise further cuts on the real line and with their help we could manage to write the first few terms of an ambiguity free non-perturbative trans-series, which we confronted with the numerical solution of the original integral equation and we found complete agreement. The full power of the resurgence theory can be used, however, when we can bridge the various non-perturbative expansions to each other, as then we would have relations between low order perturbative coefficients. Based on our numerics we conjectured such relations for the nearest cuts on the Borel plane, i.e. for the leading non-perturbative corrections. The aim of the present paper is to elevate these numerical observations to an analytically proved level and to prepare the ground for systematic analytical studies, which could reveal the full resurgence structure.

The paper is organized as follows: in the rest of the introduction we collect the needed formulae from the literature. In particular, we introduce the exact linear integral equation for the ground-state energy density and Volin's approach for the corresponding resolvent, which turns the integral equations into algebraic ones for the systematic perturbative expansion. We simplify these recursive equations in the O(4) case and emphasize that the only input comes from the leading behaviour of the resolvent's Laplace transformed, which can be determined from the Wiener-Hopf solution. In the next section we summarize the main results of this paper. Technically, we manage to formulate recursive equations directly for the parametrizations of the Laplace transform of the resolvent without relying on its coordinate space form. We then calculate the asymptotic form of the perturbative expansions, which provides the analytical solution for the resolvent's Laplace transform around the origin. We then learn how to extend the asymptotic form selfconsistently, which is valid on the unit circle and gives the complete analytical information on leading singularities of the Borel plane, i.e. the leading alien derivates. We summarize the main findings in the conclusion where we also present an outlook for further research. All the technical details are relegated to appendices.

1.1 TBA calculation of the ground-state energy density

The central object of study in this paper is the ground-state energy in a magnetic field h [4–6], when the Hamiltonian is modified as $H(h) = H(0) - hQ_{12}$, where H(0) is the original sigma model Hamiltonian and Q_{12} is the conserved charge associated to the (internal) rotation symmetry in the 12 plane. Particles with largest Q_{12} charge condense into the vacuum if h is larger than m, the mass of the particles. Due to the integrability of the model this condensate

can be described by the Thermodynamic Bethe Ansatz (TBA) equation [4–6]

$$\chi(\theta) - \int_{-B}^{B} \frac{d\theta'}{2\pi} K(\theta - \theta') \chi(\theta') = \cosh\theta$$
(1.1)

where the kernel is determined by the scattering matrix $S(\theta)$ for the condensed particles:

$$2\pi K(\theta) = -2\pi i \partial_{\theta} \log S(\theta) = \sum_{k=\{\frac{1}{2},\Delta\}} \{\Psi(k+\frac{1}{2}-\frac{i\theta}{2\pi}) - \Psi(k+\frac{i\theta}{2\pi})\} + \text{cc.}$$
(1.2)

Here $\Delta = \frac{1}{N-2}$ and Ψ is the digamma function: $\Psi(\theta) = \partial_{\theta} \log \Gamma(\theta)$. The particle density mU and the energy density m^2W are obtained simply from

$$U = \int_{-B}^{B} \frac{d\theta}{2\pi} \chi(\theta) \quad ; \qquad W = \int_{-B}^{B} \frac{d\theta}{2\pi} \cosh\theta \,\chi(\theta). \tag{1.3}$$

These equations depend on B, which can be related to the magnetic field h. Large magnetic fields correspond to large densities and large B, and in the following we will study the 1/B expansion of the solution and the densities.

1.2 Perturbative expansion of the TBA equation, Volin's method [1, 2]

The calculations can be simplified by introducing the resolvent

$$R(\theta) = \int_{-B}^{B} d\theta' \, \frac{\chi(\theta')}{\theta - \theta'} \tag{1.4}$$

and its Laplace transform

$$\hat{R}(s) = \int_{-i\infty+0}^{i\infty+0} \frac{dz}{2\pi i} e^{sz} R(B+z/2).$$
(1.5)

The latter is related to the Fourier transform of $\chi(\theta)$:

$$\hat{R}(s) = 2e^{-2sB}\tilde{\chi}(2is) \quad ; \qquad \tilde{\chi}(\omega) = \int_{-B}^{B} e^{-i\omega\theta}\chi(\theta)d\theta.$$
(1.6)

Then, using the $\chi(\theta) = \chi(-\theta)$ symmetry

$$W = \int_{-B}^{B} \cosh\theta\chi(\theta)\frac{d\theta}{2\pi} = \int_{-B}^{B} e^{\theta}\chi(\theta)\frac{d\theta}{2\pi} = \frac{1}{2\pi}\tilde{\chi}(i) = \frac{e^{B}}{4\pi}\hat{R}(1/2).$$
(1.7)

Although it is possible to proceed for generic O(N) models, in the following we restrict our attention to the O(4) model where $\Delta = 1/2$ since formulas are much simpler there.

Volin made the following two seminal observations. First, both $R(\theta)$ and $\hat{R}(s)$ can be expanded for large B and the form of the two asymptotic expansions are determined by the TBA equation and the analytic properties of its solution. In rapidity space

$$R(\theta) = 2\tilde{A}\sqrt{B} \sum_{n,m=0}^{\infty} \frac{c_{n,m}}{B^{m-n}(\theta^2 - B^2)^{n+1/2}},$$
(1.8)

where the $c_{n,m}$ coefficients are numerical constants, but the overall constant \tilde{A} may depend on B. The density is obtained from the residue of the resolvent at infinity

$$U = \tilde{A} \frac{\sqrt{B}}{\pi} \sum_{m=0}^{\infty} c_{0,m} B^{-m}.$$
 (1.9)

For the Laplace transform study of its analytic properties leads to the following ansatz

$$\hat{R}(s) = \frac{A}{\sqrt{s}} \frac{\Gamma(1+s)}{\Gamma(\frac{1}{2}+s)} \left(\frac{1}{s+\frac{1}{2}} + \frac{1}{Bs} \sum_{n,m=0}^{\infty} \frac{Q_{n,m}}{B^{n+m}s^n} \right) \quad ; \qquad A = \frac{e^B \sqrt{\pi}}{2\sqrt{2}} \tag{1.10}$$

with constant coefficients $Q_{n,m}$.

The second seminal observation was that by re-expanding $R(\theta)$ and performing the Laplace transform, the result can be compared to $\hat{R}(s)$. This gives $\tilde{A} = A$, and by matching the two parametrizations, all the unknown numerical coefficients $Q_{n,m}$ and $c_{n,m}$ can be determined completely algebraically in a recursive manner. This method was used by Volin to calculate the perturbative coefficients for generic O(N) models. The calculations in the particular case of the O(4) model take a simpler form.

1.3 O(4) Volin equations

To describe the algebraic recursion equations for the O(4) model we have to introduce a few definitions.

The input parameters of the recursion are the coefficients a_n of the expansion

$$\sum_{n=0}^{\infty} a_n x^n = g(x) = \frac{e^{ax}}{1+x} \frac{\Gamma^2(1+x/2)}{\Gamma(1+x)}, \qquad a = \ln 2, \qquad a_{-1} = 0, \tag{1.11}$$

and the rational constants

$$E_{n,k} = (-1)^k \frac{1}{2^k k!} \prod_{\ell=1}^k \left[n^2 - \left(\frac{2\ell - 1}{2}\right)^2 \right], \qquad E_{n,0} = 1.$$
(1.12)

Using the Gamma function representation

$$\Gamma(1+z) = \exp\left\{-\gamma z + \sum_{k=2}^{\infty} \frac{\zeta_k}{k} (-z)^k\right\}$$
(1.13)

we can re-express the basic input function (1.11) in terms of Dirichlet's eta coefficients defined by

$$\eta_k = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n^k}.$$
(1.14)

They are simply related to the zeta coefficients by

$$\eta_k = (1 - 2^{1-k})\zeta_k, \quad k \ge 2, \qquad \eta_1 = \ln 2.$$
 (1.15)

Also using the expansion

$$\ln(1+x) = \sum_{k=1}^{\infty} (-1)^{k-1} \frac{x^k}{k}$$
(1.16)

g(x) can be recast in the form

$$\sum_{n=0}^{\infty} a_n x^n = g(x) = \exp\left\{\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} (\eta_k - 1) x^k\right\}.$$
(1.17)

For later convenience we introduce the new two-index variable $p_{n,m}$ by

$$Q_{k,m} = \frac{p_{m+k,k}}{2^{m+2k+1}}.$$
(1.18)

We now (recursively in m) find the unknowns $c_{n,m}$ $(m = 0, 1, \ldots, n = 0, 1, \ldots)$ and $p_{a,b}$ $(a = 0, 1, \ldots, b = 0, 1, \ldots, a)$, starting from

$$c_{n,0} = a_n.$$
 (1.19)

After having completed the recursion up to step m-1, we first have to solve the set of equations $r = 1, \ldots, m$

$$\sum_{k=r}^{m} E_{k-r,k} c_{k-r,m-k} = \frac{1}{2^m} \sum_{k=r}^{m} (a_{k-r} + a_{k-r-1}) p_{m-1,k-1}$$
(1.20)

for $p_{m-1,k-1}$, k = 1, ..., m and then use this solution to express

$$c_{n,m} = -\sum_{k=1}^{m} E_{n+k,k} c_{n+k,m-k} + \frac{1}{2^m} \sum_{k=1}^{m} (a_{n+k} + a_{n+k-1}) p_{m-1,k-1}$$
(1.21)

 $n = 0, 1, \ldots$ To calculate the energy density W(B) and the particle density U(B) we only need $c_{0,m}$ and

$$W_n = \sum_{k=0}^{n} p_{n,k}.$$
 (1.22)

Explicitly,

$$W(B) = \frac{1}{2\pi} \int_{-B}^{B} \mathrm{d}\theta \,\chi(\theta) \cosh\theta = \frac{\mathrm{e}^{2B}}{16} \epsilon(B), \qquad \epsilon(B) = 1 + 2\sum_{n=1}^{\infty} \frac{W_{n-1}}{(2B)^n}, \tag{1.23}$$

$$U(B) = \frac{1}{2\pi} \int_{-B}^{B} \mathrm{d}\theta \,\chi(\theta) = \mathrm{e}^{B} \sqrt{\frac{B}{8\pi}} \rho(B), \qquad \rho(B) = 1 + \sum_{n=1}^{\infty} \frac{c_{0,n}}{B^{n}}.$$
 (1.24)

Later we will study the expansion of the generalized charges

$$W_{p}(B) = \frac{1}{2\pi} \int_{-B}^{B} d\theta \,\chi(\theta) \cosh p\theta = \frac{e^{pB}}{4\pi} \hat{R}(p/2) = \frac{e^{(p+1)B}}{4\pi\sqrt{p}} \,2^{p} \,\frac{\Gamma^{2}(1+p/2)}{\Gamma(2+p)} \,\epsilon_{p},$$
(1.25)

where

$$\epsilon_p(z) = 1 + \frac{p+1}{p} \sum_{n=1}^{\infty} \frac{H_n(p)}{(pz)^n}, \qquad z = 2B,$$
(1.26)

and

$$H_n(p) = \sum_{k=1}^n p_{n-1,n-k} p^k.$$
 (1.27)

For the original energy density

$$W(B) = W_1(B),$$
 $\epsilon(B) = \epsilon_1(B),$ $H_n(1) = W_{n-1}.$ (1.28)

An alternative form of $\epsilon_p(z)$, in terms of the variable q = 1/p, is

$$\epsilon_p(z) = 1 + (1+q) \sum_{n=1}^{\infty} \frac{E_n(q)}{z^n}, \qquad E_n(q) = \sum_{k=0}^{n-1} p_{n-1,k} q^k.$$
 (1.29)

The input parameters a_n are polynomial expressions (with rational coefficients) in ζ_2 , ζ_3, \ldots and $a = \ln 2$ of degree n ($a = \ln 2$ counts as degree 1). Previously we observed that W_n and $c_{0,m}$ do not depend on the even zeta functions ζ_2, ζ_4, \ldots Since they are proportional to powers of π , we call this property π -independence. We now observed that not only the sum W_n but all $p_{a,b}$ coefficients individually are π -independent. However, the even zeta functions do not completely disappear: the variables $c_{n,m}, n \geq 1$ do depend on them.

In [8, 9] we studied the solution of the recursive equations (1.20) and (1.21) with high precision numerics, up to 2000th order, studied the resurgence properties of the asymptotic expansions, and compared the Borel resummation of the asymptotic series for both U(B)and W(B) to the results obtained from the numerical solution of the original TBA equation. In this paper we reproduce many of those previous results analytically. We hope that our methods may be used later in related similar models too.

2 Summary

In this section we summarize the analytic solution of the recursive equations and study the resurgence properties of the asymptotic expansions analytically. We not only confirm our previous findings but also discover some new properties of the asymptotic series.

2.1 Properties of the $\varepsilon_p(z)$ series

The second set of Volin's equations, (1.21), can be solved for $c_{n,m}$ in closed form and using this solution in (1.20) we can obtain closed equations for the $p_{a,b}$ type variables. The details of the derivation are given in appendix A. The simplified equations are of the form

$$\mathcal{L}_{r}^{(M)} = \mathcal{R}_{r}^{(M)}, \qquad M = 1, 2, \dots, \qquad r = 1, \dots, M,$$
(2.1)

where

$$\mathcal{R}_{r}^{(M)} = \sum_{k=0}^{M-r} (a_{k} + a_{k-1}) p_{M-1,k+r-1}$$

$$= a_{M-r} p_{M-1,M-1} + \sum_{k=0}^{M-r-1} a_{k} [p_{M-1,k+r-1} + p_{M-1,k+r}]$$
(2.2)

and

$$\mathcal{L}_{r}^{(M)} = \sum_{n=0}^{M-r} X_{r,n} Y_{n}^{(M-r)}.$$
(2.3)

Here $X_{r,n}$ are *M*-independent rational coefficients:

$$X_{r,n} = 2^r (-2)^n \sum_{p=0}^n (-1)^p E_{p,p+r} E_{p+1,n-p}$$
(2.4)

and the symbols $Y_n^{(M)}$ are

$$Y_n^{(M)} = \sum_{k=1}^{M-n} (a_{n+k} + a_{n+k-1}) p_{M-n-1,k-1}, \quad n = 0, \dots, M-1, \qquad Y_M^{(M)} = a_M.$$
(2.5)

The equations (2.1) can be solved recursively in M.

In appendix B, using this simplified form of Volin's equations we prove the existence of the asymptotic (ASY) expansions

$$\frac{p_{N,N-j}}{\Gamma(N+1)} = \frac{1}{\pi} \left\{ \beta^{(j)} + \frac{\alpha_0^{(j)}}{N} + \frac{\alpha_1^{(j)}}{N(N-1)} + \frac{\alpha_2^{(j)}}{N(N-1)(N-2)} + \dots \right\} \qquad j = 0, 1, \dots$$
(2.6)

Here the coefficients are given by generating functions:

$$\sum_{j=0}^{\infty} \beta^{(j)} p^{j+1} = u(p) \qquad \qquad \sum_{j=0}^{\infty} \alpha_m^{(j)} p^{j+1} = u(p) A_m(p) \quad m = 0, 1, \dots$$
(2.7)

where

$$u(p) = \frac{p}{1+p} \frac{g(-p)}{g(p)} \qquad A_m(p) = (p-1) \sum_{k=0}^m p_{m,k} (-1)^k p^{m-k}$$
(2.8)

These results can be used to study the $\epsilon_p(z)$ asymptotic series and related alien derivatives².

First we note that

$$u(x) = \frac{x}{(1-x)2^{2x}} \frac{\Gamma(1+x)}{\Gamma(1-x)} \frac{\Gamma^2(1-x/2)}{\Gamma^2(1+x/2)}$$
(2.9)

²The definitions related to alien derivatives are summarized in appendix \mathbf{F} .

and using the representation of the Gamma function given by (1.13) we see that we have π -independence for the leading terms $\beta^{(j)}$ and also for $\alpha_m^{(j)}$ (at least for the first few *m* values, for which the π -indepence of $p_{m,k}$ is known by explicit construction). We also note that u(x) can alternatively be written as

$$u(x) = \frac{x}{2^{2x}} \frac{\Gamma(1+x)}{\Gamma(2-x)} \frac{\Gamma^2(1-x/2)}{\Gamma^2(1+x/2)},$$
(2.10)

hence for |x| < 2 the only singular point of u(x) is x = -1:

$$(x \approx -1):$$
 $u(x) \approx -\frac{1}{2}\frac{1}{1+x} = -\frac{1}{2}\sum_{j=0}^{\infty}(-x)^j.$ (2.11)

Therefore for large j

$$\beta^{(j)} \approx \frac{1}{2} (-1)^j \qquad \qquad \alpha_m^{(j)} \approx \frac{1}{2} A_m (-1) (-1)^j \qquad \qquad A_m (-1) = 2 (-1)^{m+1} W_m.$$
 (2.12)

The central object of study is

$$\epsilon_p(z) = 1 + \frac{p+1}{p} \sum_{m=0}^{\infty} \frac{H_{m+1}(p)}{(pz)^{m+1}}.$$
 (2.13)

For -p we have

$$\epsilon_{-p}(z) = 1 + \sum_{m=0}^{\infty} \frac{A_m(p)}{(pz)^{m+1}}.$$
(2.14)

We can write alternatively

$$\epsilon_p(z) = 1 + \frac{p+1}{p} \sum_{m=0}^{\infty} \frac{\Gamma(m+1)c_m^{(\epsilon_p)}}{z^{m+1}p^{m+1}}, \qquad c_m^{(\epsilon_p)} = \frac{H_{m+1}(p)}{\Gamma(m+1)}.$$
 (2.15)

These expansion coefficients define the Borel space function

$$B(x) = \sum_{m=0}^{\infty} c_m^{(\epsilon_p)} x^m \,.$$
(2.16)

The Borel transform of ϵ_p is given by

$$\hat{\epsilon}_p(t) = \frac{p+1}{p^2} B\left(\frac{t}{p}\right).$$
(2.17)

Now it is easy to establish the ASY expansion of the Borel transform coefficients (for |p| < 1).

$$c_{M}^{(\epsilon_{p})} = \frac{H_{M+1}(p)}{\Gamma(M+1)} = \sum_{j=0}^{M} \frac{p_{M,M-j}}{\Gamma(M+1)} p^{j+1} \approx \sum_{j=0}^{\infty} \frac{p^{j+1}}{\pi} \Big\{ \beta^{(j)} + \sum_{m=0}^{\infty} \frac{\alpha_{m}^{(j)}}{M_{[m+1]}} \Big\}$$

$$= \frac{u(p)}{\pi} \Big\{ 1 + \sum_{m=0}^{\infty} \frac{A_{m}(p)}{M_{[m+1]}} \Big\}.$$
(2.18)

We have extended the upper limit of the summation from M to ∞ . This is possible since the error we make by this extension is exponentially small for |p| < 1. Above we used the notation

$$M_{[r]} = M(M-1)(\dots)(M-r+1),$$
 $M_{[1]} = M,$ $M_{[0]} = 1.$ (2.19)

From here we can read off the singular part of the Borel space function B(x):

$$B^{\rm sing}(x) = \frac{u(p)}{\pi} \left\{ \frac{1}{1-x} - \ln(1-x) \sum_{m=0}^{\infty} A_m(p) \frac{(x-1)^m}{m!} \right\}.$$
 (2.20)

The singular part of $\hat{\epsilon}_p(t)$ is

$$\hat{\epsilon}_p^{\text{sing}}(t) = \frac{p+1}{p} \frac{u(p)}{\pi} \left\{ \frac{1}{p-t} - \ln(p-t) \sum_{m=0}^{\infty} \frac{A_m(p)}{p^{m+1}} \frac{(t-p)^m}{m!} \right\}.$$
(2.21)

Expressed in terms of alien derivatives we have

$$(\Delta_p \epsilon_p)(z) = -2i \frac{p+1}{p} u(p) \left\{ 1 + \sum_{m=0}^{\infty} \frac{A_m(p)}{(pz)^{m+1}} \right\} = -2i \frac{p+1}{p} u(p) \epsilon_{-p}(z).$$
(2.22)

We can make it look simpler by introducing the rescaled function³

$$\tilde{\epsilon}_p(z) = g(p)\epsilon_p(z). \tag{2.23}$$

In terms of the rescaled functions the alien derivative is simply

$$\Delta_p \tilde{\epsilon}_p = -2i\tilde{\epsilon}_{-p}.\tag{2.24}$$

We also note that since

$$\epsilon_{-1}(z) \equiv 1, \qquad u(1) = 1$$
 (2.25)

in the $p \to 1$ limit we have

$$\Delta_1 \epsilon = -4i, \tag{2.26}$$

which is what we found earlier numerically in [8, 9]. It means that the only singularity of $\hat{\epsilon}(t)$ around t = 1 is a simple pole with residue $-2/\pi$.

2.2 "Derivation" of the naive ansatz

Using (2.12) we see that the asymptotics if j is also large simplifies to

$$\frac{p_{N,N-j}}{\Gamma(N+1)} = \frac{(-1)^j}{2\pi} \left\{ 1 + \frac{A_0(-1)}{N} + \frac{A_1(-1)}{N(N-1)} + \frac{A_2(-1)}{N(N-1)(N-2)} + \dots \right\}$$
(2.27)

³Note that $W_p = e^{(p+1)B}/(4\pi\sqrt{p})\tilde{\epsilon_p}$. See (1.25).

The precise meaning of the above formula is that first we consider the large N asymptotics of $p_{N,N-j}$ for fixed j and next we let the fixed j to be large. If both N and j are very large but finite, the requirement is that

$$N \gg j \gg 1. \tag{2.28}$$

We will now make the following bold assumption: (2.27) still holds if we take j = N - k with fixed k. N and j are both large in this case but the above requirement is not satisfied. Making this (illegal) substitution we obtain

$$\frac{p_{N,k}}{\Gamma(N+1)} = \frac{(-1)^N}{2\pi} (-1)^k \left\{ 1 + \frac{\tilde{R}_0}{N} + \frac{\tilde{R}_1}{N(N-1)} + \frac{2\tilde{R}_2}{N(N-1)(N-2)} + \dots \right\} \qquad k = 0, 1, \dots$$
(2.29)

with

$$\tilde{R}_m = \frac{2}{m!} (-1)^{m+1} W_m \tag{2.30}$$

We tried to check numerically if our bold assumption is correct. We started from a somewhat more general ansatz:

$$\frac{p_{N,k}}{\Gamma(N+1)} = \frac{(-1)^N}{2\pi} (-1)^k \left\{ b^{(k)} + \frac{\tilde{R}_0^{(k)}}{N} + \frac{\tilde{R}_1^{(k)}}{N(N-1)} + \frac{2\tilde{R}_2^{(k)}}{N(N-1)(N-2)} + \dots \right\} \qquad k = 0, 1, \dots$$
(2.31)

and found the first few coefficients numerically:

$$b^{(k)} = 1 \qquad k = 0, 1, \dots$$

$$\tilde{R}_{0}^{(0)} = -\frac{3}{4} \qquad \tilde{R}_{0}^{(k)} = \tilde{R}_{0} = -\frac{1}{2} = -2W_{0} \qquad k = 1, 2, \dots$$

$$\tilde{R}_{1}^{(0)} = \frac{49}{32} - \frac{3a}{2} \qquad \tilde{R}_{1}^{(1)} = \frac{27}{32} - a \qquad \tilde{R}_{1}^{(k)} = \tilde{R}_{1} = \frac{9}{8} - a = 2W_{1} \qquad k = 2, 3, \dots$$

$$\tilde{R}_{2}^{(0)} = -\frac{585}{256} + \frac{49a}{16} - \frac{3a^{2}}{2} \qquad \tilde{R}_{2}^{(1)} = -\frac{45}{32} + \frac{27a}{16} - a^{2} \qquad \tilde{R}_{2}^{(2)} = -\frac{531}{256} + \frac{9a}{4} - a^{2}$$

$$\tilde{R}_{2}^{(k)} = \tilde{R}_{2} = -\frac{57}{32} + \frac{9a}{4} - a^{2} = -W_{2} \qquad k = 3, 4, \dots$$

$$(2.32)$$

We see that the leading term is correctly given by our bold assumption. (2.29) is not completely correct but we found the following interesting structure. There are irregular and regular coefficients:

irregular:
$$\tilde{R}_m^{(k)}$$
 $k = 0, ..., m$ regular: $\tilde{R}_m^{(k)} = \tilde{R}_m = \frac{2}{m!} (-1)^{m+1} W_m$ $k \ge m+1.$
(2.33)

We see that with the exception of a few irregular coefficients the generic coefficients are correctly given by (2.29).

2.3 The correct ansatz

Motivated by the naive "derivation" and based on our numerical studies we now start from the ansatz (2.31). Unlike the case for our earlier ansatz (2.6) we cannot analytically prove the

existence of the asymptotic expansion (2.31). However, taking this form (with undetermined expansion coefficients $b^{(k)}$, $\tilde{R}^{(k)}$) for granted, we can analytically prove the special structure we found numerically.

The starting point is (2.1) and using (2.31) and comparing the asymptotic expansions of the two sides of the equations first of all we can establish the generic structure

$$\begin{aligned} b^{(k)} &= b \\ \tilde{R}_m^{(k)} &= \tilde{R}_m \qquad k \ge m+1 \end{aligned} \} \text{ with some generic } b, \tilde{R}_m. \end{aligned} (2.34)$$

Moreover, all irregular $\tilde{R}_m^{(k)}$ coefficients (k = 0, ..., m) can be expressed linearly in terms of the generic parameters b, \tilde{R}_m . In particular, up to NNNLO we find the result

$$\tilde{R}_{0}^{(0)} = \tilde{R}_{0} - p_{0,0}b,
\tilde{R}_{1}^{(0)} = \tilde{R}_{1} - p_{0,0}\tilde{R}_{0} + p_{1,0}b,
\tilde{R}_{1}^{(1)} = \tilde{R}_{1} - p_{1,1}b,
\tilde{R}_{2}^{(0)} = \tilde{R}_{2} - \frac{1}{2}p_{0,0}\tilde{R}_{1} + \frac{1}{2}p_{1,0}\tilde{R}_{0} - \frac{1}{2}p_{2,0}b,
\tilde{R}_{2}^{(1)} = \tilde{R}_{2} - \frac{1}{2}p_{1,1}\tilde{R}_{0} + \frac{1}{2}p_{2,1}b,
\tilde{R}_{2}^{(2)} = \tilde{R}_{2} - \frac{1}{2}p_{2,2}b.$$
(2.35)

The proof of the generic structure (2.34) and the technical details of the calculation leading to (2.35) can be found in appendix C.

The above findings suggest a product form of $\Delta_{-1}\epsilon_p$:

$$\Delta_{-1}\epsilon_p = iX\epsilon_p \qquad \qquad X = b + \sum_{m=0}^{\infty} \frac{X_m}{z^{m+1}}.$$
(2.36)

The consequences of this relation are discussed in appendix D. Writing out (2.36) explicitly we obtain

$$\tilde{R}_{m}^{(k)} = \tilde{R}_{m} = \frac{(-1)^{m+1}}{m!} X_{m}, \quad k \ge m+1,$$
(2.37)

$$\tilde{R}_{m}^{(m)} = \tilde{R}_{m} - \frac{p_{m,m}}{m!}b,$$
(2.38)

$$\tilde{R}_{m}^{(k)} = \tilde{R}_{m} + \frac{(-1)^{m+k+1}}{m!} \left\{ bp_{m,k} + \sum_{r=0}^{m-1-k} X_{r} p_{m-r-1,k} \right\}, \quad k \le m-1.$$
(2.39)

This gives the same structure as above and up to NNNLO explicitly reproduces (2.35) taking into account the identification (2.37). The "naive" identification (2.30) gives b = 1 and $X_m = 2W_m$, i.e. $X = \epsilon$. Finally we get

$$\Delta_{-1}\epsilon_p = i\epsilon\epsilon_p. \tag{2.40}$$

This is the generalization of the relation

$$\Delta_{-1}\epsilon = i\epsilon^2 \tag{2.41}$$

found numerically in [8, 9].

2.4 Asymptotic expansion of ρ

From the inversion formula we obtain the coefficients of the Borel transform of ρ :

$$c_N^{(\rho)} = \sum_{n=0}^{N+1} \omega_n \frac{Y_n^{(N+1)}}{\Gamma(N+1)}, \qquad \omega_n = (-2)^n E_{1,n}.$$
(2.42)

For the asymptotic expansion of $c_N^{(\rho)}$ up to NNNLO we need the asymptotics of

$$\frac{Y_n^{(N+1)}}{\Gamma(N+1)} \qquad \text{for} \qquad n = 0, 1, 2, 3$$

To calculate the above expansions the parameters

$$b^{(k)}, \ \tilde{R}_m^{(k)} \qquad m = 0, 1, 2 \qquad \qquad k = 0, 1, 2, 3$$

are needed. We already expressed all these in terms of the generic coefficients. The alien derivative of ρ to this order is calculated in appendix E. The result is

$$\Delta_{-1}\rho = i\left\{b + \frac{X_0 + \tilde{u}_1 b}{z} + \frac{X_1 + \tilde{u}_1 X_0 + \tilde{u}_2 b}{z^2} + \frac{X_2 + \tilde{u}_1 X_1 + \tilde{u}_2 X_0 + \tilde{u}_3 b}{z^3} + \dots\right\}.$$
 (2.43)

Here we use the variables

$$\tilde{u}_M = 2^M c_{0,M},$$
(2.44)

which appear in the ρ asymptotic series:

$$\rho = 1 + \sum_{M=1}^{\infty} \frac{c_{0,M}}{B^M} = \sum_{M=0}^{\infty} \frac{\tilde{u}_M}{z^M}.$$
(2.45)

The meaning of (2.43) is that (to this order)

$$\Delta_{-1}\rho = iX\rho = i\epsilon\rho. \tag{2.46}$$

3 Conclusion

In this paper we investigated the perturbative expansion of the ground-state energy ϵ and the generalized charges ϵ_p of the O(4) model in a magnetic field. We derived a closed system of equations for these perturbative coefficients, which we solved asymptotically by calculating the leading factorial growth together with all the subsequent subfactorial behaviour. From this analytical solution we could show that the asymptotic coefficients contain only odd zeta

numbers, a result similar to what was observed before for the original perturbative coefficients by explicit calculation [1, 2]. Our results also show that the leading singularity of the Borel transform of ϵ_p is a logarithmic cut and the corresponding alien derivative resurges to the same set of charges as

$$\Delta_p \epsilon_p = -2i \frac{p+1}{p} u(p) \epsilon_{-p} \tag{3.1}$$

with u(p) an explicitly known function (2.9). This, in particular, implies our previous finding $\Delta_1 \epsilon = -4i$ for the energy density [8, 9]. We could also selfconsistently determine the behaviour around the next singularity on the Borel plane, which manifested itself as a logarithmic cut starting at -1. The corresponding alien derivative again showed resurgence in the form

$$\Delta_{-1}\epsilon_p = i\epsilon\epsilon_p,\tag{3.2}$$

which is the generalization of what we found earlier numerically: $\Delta_{-1}\epsilon = i\epsilon^2$. We extended this analysis also for the density and showed that $\Delta_{-1}\rho = i\epsilon\rho$, again conforming previous numerical observation. By this we completed the investigations of the leading singularities. Our approach, however, is quite general and is capable of investigating subsequent, exponentially suppressed asymptotics, i.e. further cuts on the Borel plane. We are planning to advance into this direction later.

The recursive equations have their inputs from the asymptotic form of the Laplace transform of the resolvent, g(x), and the functions $E_{n,m}$ playing a crucial role in moving between a function and its Laplace transform, and which itself shows a kind of resurgence property (B.10). As the model-specific input comes only from g(x) we analyzed a toy model, where we replaced all η_n (see (1.15)) with 1. This was also motivated by the fact that numerically they are not far from each other. We could solve this model explicitly, see appendix G, and observed a similar resurgence pattern to what we found for the real problem. This sheds some light on their origins. This toy model then can be systematically dressed up, by including corrections in $\eta_n - 1$. In particular, the exact dependence on η_1 is found in appendix G.

In the present paper we analysed the density and the generalized charges as a function of the TBA parameter B. The change for the running coupling is well understood and the composition formula for alien derivatives [37] allows a direct translation [8, 9].

So far we analyzed only the O(4) model, but the technics we developed here can also be extended for other O(N) models and their supersymmetric generalizations as well as for the principal chiral and Gross-Neveu models, all of which were analysed recently in [38] by directly investigating the integral equation. They may be useful also for non-relativistic two dimensional integrable theories relevant for statistical physics such as [28–30].

Acknowledgments

We thank Gergő Nemes for useful discussions, the Isaac Newton Institute programme 'Applicable resurgent asymptotics: towards a universal theory', and the NKFIH Grant 134946 for support.

A Simplifying Volin's equations

A.1 Properties of the a_n coefficients

The input a_n parameters are obtained from the Taylor expansion of (1.11). The first few coefficients are

$$a_0 = 1,$$
 $a_1 = a - 1,$ $a_2 = 1 - a + \frac{a^2}{2} - \frac{\pi^2}{24}.$ (A.1)

g(x) can also be represented as

$$g(x) = 2^{x+1} \frac{\Gamma^2(2+x/2)}{\Gamma(3+x)} \frac{1}{1+x/2} = \frac{1}{2+x} + \operatorname{conv}_4(x), \tag{A.2}$$

where the second term in the last formula is some function having a power series expansion convergent for |x| < 4. This form implies that for large n

$$a_n \sim \frac{(-1)^n}{2^{n+1}}.$$
 (A.3)

A.2 $c_{n,m}$ equations

For later purposes we note that

$$E_{1,k} = \frac{(-1)^k}{2^k} \omega_k, \tag{A.4}$$

where

$$\omega_k = \frac{1}{k!} \prod_{\ell=1-k}^k \left(\ell + \frac{1}{2}\right), \quad k \ge 1, \qquad \omega_0 = 1.$$
 (A.5)

We introduce the notation

$$c_{n,m} = w_{n,n+m}.\tag{A.6}$$

With this notation the $c_{n,m}$ equations for fixed $M \ge 1$ and $n = 0, 1, \ldots, M - 1$ read

$$\sum_{k=n}^{M} E_{k,k-n} w_{k,M} = 2^{n-M} Y_n^{(M)}, \qquad (A.7)$$

where

$$Y_n^{(M)} = \sum_{k=1}^{M-n} (a_{n+k} + a_{n+k-1}) p_{M-n-1,k-1}, \qquad n = 0, \dots, M-1.$$
(A.8)

The unknowns in these equations are

$$w_{j,M}, \qquad j = 0, \dots, M - 1$$
 (A.9)

since for the last component

$$w_{M,M} = c_{M,0} = a_M, (A.10)$$

which is an input. However, we can formally extend (A.7) for n = M as well, if we define

$$Y_M^{(M)} = a_M. (A.11)$$

This way we have M + 1 equations for M + 1 unknowns and the equations can be written in matrix form:

$$\sum_{k=0}^{M} \mathcal{M}_{n,k} w_{k,M} = 2^{n-M} Y_n^{(M)}, \qquad (A.12)$$

where

$$\mathcal{M}_{n,k} = \begin{cases} E_{k,k-n} & k \ge n \\ 0 & k < n \end{cases}$$
(A.13)

The solution is given by the inverse matrix:

$$w_{a,M} = \sum_{n=0}^{M} \mathcal{M}_{a,n}^{-1} \, 2^{n-M} \, Y_n^{(M)}. \tag{A.14}$$

We are mostly interested in

$$c_{0,M} = w_{0,M} = \sum_{n=0}^{M} \mathcal{M}_{0,n}^{-1} 2^{n-M} Y_n^{(M)}, \qquad (A.15)$$

because these are the ones appearing in the *B*-expansion of $\rho(B)$.

The inverse matrix is also triangular and we found the simple result

$$\mathcal{M}_{a,n}^{-1} = (-1)^{n-a} E_{a+1,n-a}, \qquad (n \ge a).$$
 (A.16)

In particular,

$$\mathcal{M}_{0,n}^{-1} = (-1)^n E_{1,n} = \frac{\omega_n}{2^n}.$$
 (A.17)

For some purposes it is useful to express the variables $w_{a,m}$ from the second set of Volin equations and if we use

$$w_{a,M} = \sum_{n=a}^{M} (-1)^{n+a} E_{a+1,n-a} 2^{n-M} Y_n^{(M)}$$
(A.18)

we can obtain closed equations for the $p_{a,b}$ type variables, which can be solved recursively in M. These equations are given by (2.1)-(2.4).

B Analytic proof of the asymptotic expansion (2.6)

B.1 Some definitions, notations, relations

The definition of the expansion coefficients of the basic input function g(x) is extended to negative indices as

$$a_{-n} = 0, \qquad n = 1, 2, \dots$$
 (B.1)

If a function $\mathcal{F}(M)$ has asymptotic series (ASY) for large M, the coefficients of the individual terms in this expansion will be denoted by $\mathcal{F}^{[r]}$:

$$\mathcal{F}(M) = \sum_{r=0}^{\infty} \frac{\mathcal{F}^{[r]}}{M_{[r]}} = \mathcal{F}^{[0]} + \frac{\mathcal{F}^{[1]}}{M} + \frac{\mathcal{F}^{[2]}}{M(M-1)} + \frac{\mathcal{F}^{[3]}}{M(M-1)(M-2)} + \dots$$
(B.2)

In this appendix M is always assumed to be asymptotically large. For simplicity, the M dependence will not always be explicit. For example, we will use the notations

$$\Omega^{(k)} = \frac{p_{M,M-k}}{\Gamma(M+1)}, \qquad k = 0, 1, \dots$$
(B.3)

and

$$\bar{\mathcal{R}}_j = \frac{\mathcal{R}_{M+1-j}^{(M+1)}}{\Gamma(M+1)}, \qquad \bar{\mathcal{L}}_j = \frac{\mathcal{L}_{M+1-j}^{(M+1)}}{\Gamma(M+1)}, \qquad j = 0, 1, \dots$$
(B.4)

This appendix is about the analytic proof of the following two statements. Statement $\underline{1}$

 $\Omega^{(j)}~(j=0,1,\dots)$ has ASY expansion

$$\frac{p_{M,M-j}}{\Gamma(M+1)} = \frac{1}{\pi} \Big\{ \beta^{(j)} + \frac{\alpha_0^{(j)}}{M} + \frac{\alpha_1^{(j)}}{M(M-1)} + \frac{\alpha_2^{(j)}}{M(M-1)(M-2)} + \dots \Big\}$$

$$= \frac{1}{\pi} \Big\{ \beta^{(j)} + \sum_{m=0}^{\infty} \frac{\alpha_m^{(j)}}{M_{[m+1]}} \Big\}.$$
(B.5)

Here we state the existence of the ASY expansion and define the expansion coefficients.

Statement 2

The expansion coefficients are given with the help of generating functions as follows.

$$\sum_{j=0}^{\infty} \beta^{(j)} p^{j+1} = u(p), \qquad \sum_{j=0}^{\infty} \alpha_m^{(j)} p^{j+1} = u(p) A_m(p), \qquad m = 0, 1, \dots$$
(B.6)

Here

$$u(p) = \frac{p}{p+1} \frac{g(-p)}{g(p)}.$$
 (B.7)

B.2 $E_{a,N}$ resurgence

For N > a an alternative representation of $E_{a,N}$ is

$$E_{a,N} = (-1)^a \frac{\Gamma(2N - 2a + 1)\Gamma(2N + 2a + 1)}{2^{5N}\Gamma(N+1)\Gamma(N-a+1)\Gamma(N+a+1)}.$$
 (B.8)

The following combination has ASY expansion for large N, which can be calculated using the Stirling series.

$$\frac{\pi}{\Gamma(N)}(-1)^{a}2^{N}E_{a,N} = \pi \frac{\Gamma(2N-2a+1)\Gamma(2N+2a+1)}{2^{4N}\Gamma(N)\Gamma(N+1)\Gamma(N-a+1)\Gamma(N+a+1)}$$
$$= 1 + \sum_{m=1}^{\infty} \frac{1}{m!} \prod_{\ell=1}^{m} \frac{a^{2} - (\ell-1/2)^{2}}{N-\ell} = 1 + \sum_{m=1}^{\infty} \frac{(-2)^{m}E_{a,m}}{(N-1)(\dots)(N-m)}.$$
(B.9)

For later purpose we write the above expansion as

$$2^{N} E_{a,N} = \frac{(-1)^{a} \Gamma(N)}{\pi} \left\{ 1 + \sum_{m=1}^{\infty} \frac{(-2)^{m} E_{a,m}}{(N-1)(\dots)(N-m)} \right\},$$
(B.10)

which can be interpreted as a kind of "resurgence" of the function $E_{a,N}$.

B.3 Proof of the statements

We start from (2.1) in the form

$$\bar{\mathcal{L}}_j = \bar{\mathcal{R}}_j, \qquad j = 0, 1, \dots \tag{B.11}$$

The two sides of the equations are

$$\bar{\mathcal{R}}_j = \sum_{k=0}^j (a_k + a_{k-1}) \Omega^{(j-k)}$$
(B.12)

and

$$\bar{\mathcal{L}}_j = \sum_{n=0}^j \frac{X_{M+1-j,n}}{\Gamma(M+1)} Y_n^{(j)}.$$
(B.13)

First we consider the large M asymptotics of the left hand side.

$$\frac{X_{M+1-j,n}}{\Gamma(M+1)} = 2^{M+1-j} (-2)^n \sum_{p=0}^n (-1)^p \frac{E_{p,p+M+1-j}}{\Gamma(M+1)} E_{p+1,n-p}
= (-2)^n \sum_{p=0}^n 2^{-p} E_{p+1,n-p} \frac{1}{\pi} \frac{1}{M_{[j-p]}} \left\{ 1 + \sum_{m=1}^\infty \frac{(-2)^m E_{p,m}}{(p+M-j)(\dots)(p+M-j+1-m)} \right\}$$
(B.14)

$$= \frac{(-2)^n}{\pi} \sum_{p=0}^n 2^{-p} E_{p+1,n-p} \sum_{m=0}^\infty \frac{(-2)^m E_{p,m}}{M_{[j-p+m]}}.$$

The r^{th} component of the ASY expansion is given by the m = r + p - j term (which is nonzero if $n \ge p \ge j - r$):

$$\left(\frac{X_{M+1-j,n}}{\Gamma(M+1)}\right)^{[r]} = \frac{(-2)^n}{\pi} \sum_{p=\mathrm{Max}(0,j-r)}^n 2^{-p} E_{p+1,n-p}(-2)^{r+p-j} E_{p,r+p-j}.$$
(B.15)

If $j \ge r$ we can recognize here the triangular matrix (A.13) and its inverse (A.16) and the above can be written as

$$\frac{(-1)^{r-j}}{\pi} 2^{r-j+n} \sum_{p=j-r}^{n} \mathcal{M}_{p,n}^{-1} \mathcal{M}_{j-r,p}.$$
(B.16)

Since

$$\mathcal{M}_{p,n}^{-1} = 0$$
 if $p > n$ and $\mathcal{M}_{j-r,p} = 0$ if $p < j - r$, (B.17)

we can extend the summation limits to $\sum_{p=0}^{M}$ and the summation becomes the matrix product of the matrix and its inverse and the result is simply the Kronecker delta

$$\delta_{n,j-r}.\tag{B.18}$$

Using this in (B.13) we get

$$\bar{\mathcal{L}}_{j}^{[r]} = \frac{(-1)^{j-r}}{\pi} Y_{j-r}^{(j)}.$$
(B.19)

Explicitly,

$$\bar{\mathcal{L}}_{j}^{[0]} = \frac{(-1)^{j}}{\pi} a_{j},$$

$$\bar{\mathcal{L}}_{j}^{[r]} = \frac{(-1)^{j-r}}{\pi} \sum_{k=0}^{r-1} (a_{j-r+k+1} + a_{j-r+k}) p_{r-1,k}, \qquad j \ge r \ge 1.$$
(B.20)

For j < r (B.15) gives

$$\left(\frac{X_{M+1-j,n}}{\Gamma(M+1)}\right)^{[r]} = \frac{(-2)^{n+r-j}}{\pi} \sum_{p=0}^{n} (-1)^p E_{p+1,n-p} E_{p,r+p-j} = \frac{(-1)^{r-j}}{\pi} X_{r-j,n}.$$
 (B.21)

Using this in (B.13) we get

$$\bar{\mathcal{L}}_{j}^{[r]} = \frac{(-1)^{r-j}}{\pi} \sum_{n=0}^{j} X_{r-j,n} Y_{n}^{(j)} = \frac{(-1)^{r-j}}{\pi} \mathcal{L}_{r-j}^{(r)}.$$
(B.22)

Using (2.1) once more in this case we have

$$\bar{\mathcal{L}}_{j}^{[r]} = \frac{(-1)^{j-r}}{\pi} \mathcal{R}_{r-j}^{(r)} = \frac{(-1)^{j-r}}{\pi} \sum_{k=0}^{j} (a_{k} + a_{k-1}) p_{r-1,k+r-j-1}$$

$$= \frac{(-1)^{j-r}}{\pi} \sum_{k=r-j-1}^{r-1} (a_{k+j-r+1} + a_{k+j-r}) p_{r-1,k}.$$
(B.23)

Finally, using the convention (B.1) the summation limits can be extended:

$$\bar{\mathcal{L}}_{j}^{[r]} = \frac{(-1)^{j-r}}{\pi} \sum_{k=0}^{r-1} (a_{j-r+k+1} + a_{j-r+k}) p_{r-1,k}, \qquad (B.24)$$

which is miraculously the same as the formula for $j \ge r!$

To summarize, the final result of the calculation is (for any j = 0, 1, ...)

$$\bar{\mathcal{L}}_{j}^{[0]} = \frac{(-1)^{j}}{\pi} a_{j},$$

$$\bar{\mathcal{L}}_{j}^{[r]} = \frac{(-1)^{j-r}}{\pi} \sum_{k=0}^{r-1} (a_{j-r+k+1} + a_{j-r+k}) p_{r-1,k}, \qquad r \ge 1.$$
(B.25)

This way we have proven by explicit calculation that the LHS of (B.11) has ASY expansion. But then also the RHS has ASY expansion. In particular

$$\Omega^{(0)} = \bar{\mathcal{R}}_0 = \bar{\mathcal{L}}_0 \tag{B.26}$$

has ASY expansion, then

$$\Omega^{(1)} = -(1+a_1)\Omega^{(0)} + \bar{\mathcal{L}}_1 \tag{B.27}$$

has ASY expansion, and so on: we can prove recursively that all

$$\Omega^{(j)} = -\sum_{k=1}^{j} (a_k + a_{k-1})\Omega^{(j-k)} + \bar{\mathcal{L}}_j$$
(B.28)

have ASY expansion. Thus we have proven Statement 1.

The proof of Statement 2 is now easy. We calculate the infinite sums

$$\sum_{j=0}^{\infty} p^j \bar{\mathcal{L}}_j^{[0]} = \frac{1}{\pi} \sum_{j=0}^{\infty} (-p)^j a_j = \frac{g(-p)}{\pi}$$
(B.29)

and (for $r \ge 1$)

$$\sum_{j=0}^{\infty} p^{j} \bar{\mathcal{L}}_{j}^{[r]} = \frac{(-1)^{r}}{\pi} \sum_{j=0}^{\infty} (-p)^{j} \sum_{k=0}^{r-1} (a_{k+j-r+1} + a_{k+j-r}) p_{r-1,k}$$
$$= \frac{(-1)^{r}}{\pi} \sum_{k=0}^{r-1} p_{r-1,k} \sum_{i=0}^{\infty} a_{i} [(-p)^{i+r-k-1} + (-p)^{i+r-k}]$$
$$= \frac{(p-1)g(-p)}{\pi} \sum_{k=0}^{r-1} p_{r-1,k} (-1)^{k} p^{r-1-k} = \frac{g(-p)}{\pi} A_{r-1}(p).$$
(B.30)

For the RHS

$$\Omega^{(j)[r]} = \frac{1}{\pi} \alpha_{r-1}^{(j)}, \tag{B.31}$$

where we have temporarily introduced the short hand notation

$$\alpha_{-1}^{(j)} = \beta^{(j)}. \tag{B.32}$$

The infinite summation for the RHS gives

$$\sum_{j=0}^{\infty} p^j \bar{\mathcal{R}}_j^{[r]} = \sum_{j=0}^{\infty} p^j \sum_{k=0}^{j} (a_k + a_{k-1}) \frac{1}{\pi} \alpha_{r-1}^{(j-k)} = \frac{1+p}{p} \frac{g(p)}{\pi} \sum_{s=0}^{\infty} p^{s+1} \alpha_{r-1}^{(s)}.$$
 (B.33)

Comparing the infinite sums at the two sides we obtain

$$\sum_{s=0}^{\infty} p^{s+1} \beta^{(s)} = \frac{p}{1+p} \frac{g(-p)}{g(p)} = u(p),$$

$$\sum_{s=0}^{\infty} p^{s+1} \alpha_{r-1}^{(s)} = \frac{p}{1+p} \frac{g(-p)}{g(p)} A_{r-1}(p) = u(p) A_{r-1}(p), \quad r \ge 1.$$
(B.34)

This is Statement 2.

C Consistency of the conjectured asymptotics

In this appendix our starting point is the ansatz (2.31). However, for technical reasons we will use its (equivalent) index-shifted version:

$$\frac{p_{M-1,k}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} (-1)^{k+1} \left\{ b^{(k)} + \frac{R_0^{(k)}}{M} + \frac{R_1^{(k)}}{M(M-1)} + \frac{2R_2^{(k)}}{M(M-1)(M-2)} + \dots \right\} \qquad k = 0, 1, \dots$$
(C.1)

This asymptotic expansion is assumed valid for fixed k = 0, 1, ... in the large M limit. Using this ansatz we can calculate the asymptotic expansion of both sides of the Volin equations (2.1).

The large M asymptotics of $Y_n^{(M)}$ can be written as

$$\frac{Y_n^{(M)}}{\Gamma(M)} \approx \frac{(-1)^{M-1}(-1)^n}{2\pi(M-1)(\cdots)(M-n)} \sum_{k=0}^{M-n-1} (a_{n+k+1} + a_{n+k})(-1)^k \left\{ b^{(k)} + \frac{R_0^{(k)}}{M-n} + \frac{R_1^{(k)}}{(M-n)(M-n-1)} + \dots \right\}.$$
(C.2)

(C.1) is only valid for $k \ll M$, but for large k the coefficients a_{n+k} are exponentially small and the asymptotic expansion in powers of 1/M is not affected by this condition. Using the same reasoning, we can extend the k-summation to infinity:

$$\frac{Y_n^{(M)}}{\Gamma(M)} \approx \frac{(-1)^{M-1}(-1)^n}{2\pi(M-1)(\cdots)(M-n)} \sum_{k=0}^{\infty} (a_{n+k+1} + a_{n+k})(-1)^k \\ \left\{ b^{(k)} + \frac{R_0^{(k)}}{M-n} + \frac{R_1^{(k)}}{(M-n)(M-n-1)} + \dots \right\}.$$
(C.3)

We work up to NNNLO and rearrange the summation as follows.

$$\frac{Y_n^{(M)}}{\Gamma(M)} = \frac{(-1)^{M-n}}{2\pi(M-1)(\dots)(M-n)} \left\{ -a_n \left[b^{(0)} + \frac{R_0^{(0)}}{M-n} + \frac{R_1^{(0)}}{(M-n)(M-n-1)} + \frac{2R_2^{(0)}}{(M-n)(M-n-1)(M-n-2)} \right] + \sum_{k=1}^{\infty} a_{n+k} (-1)^k \left[b^{(k-1)} - b^{(k)} + \frac{R_0^{(k-1)} - R_0^{(k)}}{M-n} + \frac{R_1^{(k-1)} - R_1^{(k)}}{(M-n)(M-n-1)} + \frac{2[R_2^{(k-1)} - R_2^{(k)}]}{(M-n)(M-n-1)(M-n-2)} \right] + \dots \right\}.$$
(C.4)

and similarly

$$\frac{Y_n^{(M-r)}}{\Gamma(M)} = \frac{(-1)^{M-n-r}}{2\pi(M-1)(\dots)(M-n-r)} \left\{ -a_n \left[b^{(0)} + \frac{R_0^{(0)}}{M-n-r} + \frac{R_1^{(0)}}{(M-n-r)(M-n-r-1)} + \frac{2R_2^{(0)}}{(M-n-r)(M-n-r-1)(M-n-r-2)} \right] + \frac{\sum_{k=1}^{\infty} a_{n+k}(-1)^k \left[b^{(k-1)} - b^{(k)} + \frac{R_0^{(k-1)} - R_0^{(k)}}{M-n-r} + \frac{R_1^{(k-1)} - R_1^{(k)}}{(M-n-r)(M-n-r-1)} + \frac{2[R_2^{(k-1)} - R_2^{(k)}]}{(M-n-r)(M-n-r-1)(M-n-r-2)} \right] + \dots \right\}.$$
(C.5)

Proceeding analogously for the RHS, up to NNNLO we find

$$\frac{\mathcal{R}_{r}^{(M)}}{\Gamma(M)} = \frac{(-1)^{r}(-1)^{M}}{2\pi} \sum_{k=0}^{\infty} a_{k}(-1)^{k} \left\{ b^{(k+r-1)} - b^{(k+r)} + \frac{R_{0}^{(k+r-1)} - R_{0}^{(k+r)}}{M} + \frac{R_{1}^{(k+r-1)} - R_{1}^{(k+r)}}{M(M-1)} + \frac{2[R_{2}^{(k+r-1)} - R_{2}^{(k+r)}]}{M(M-1)(M-2)} + \dots \right\}.$$
(C.6)

The crucial observation is that

$$\frac{Y_n^{(M-r)}}{\Gamma(M)} = \mathcal{O}(M^{-n-r}) \tag{C.7}$$

and so

$$\frac{\mathcal{L}_r^{(M)}}{\Gamma(M)} = \mathcal{O}(M^{-r}), \tag{C.8}$$

and consequently we must also have

$$\frac{\mathcal{R}_r^{(M)}}{\Gamma(M)} = \mathcal{O}(M^{-r}). \tag{C.9}$$

But this is possible only if

$$b^{(k)} = b, \qquad k = 0, 1, \dots$$
 (C.10)

and

$$R_m^{(k)} = R_m, \qquad k \ge m+1.$$
 (C.11)

This is precisely the structure we observed numerically.

Using the above simplifications, to a given order in the asymptotic expansion, both sides of the equations contain a finite number of nonzero summands only. We work at NNNLO and we need the following expressions.

$$\frac{\mathcal{R}_{1}^{(M)}}{\Gamma(M)} = \frac{(-1)^{M-1}}{2\pi} \left\{ \frac{R_{0}^{(0)} - R_{0}}{M} + \frac{R_{1}^{(0)} - R_{1}^{(1)} + a_{1}[R_{1} - R_{1}^{(1)}]}{M(M-1)} + \frac{2[R_{2}^{(0)} - R_{2}^{(1)}] + 2a_{1}[R_{2}^{(2)} - R_{2}^{(1)}] + 2a_{2}[R_{2}^{(2)} - R_{2}]}{M(M-1)(M-2)} + \dots \right\},$$
(C.12)

$$\frac{\mathcal{R}_2^{(M)}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} \Big\{ \frac{R_1^{(1)} - R_1}{M(M-1)} + \frac{2[R_2^{(1)} - R_2^{(2)}] + 2a_1[R_2 - R_2^{(2)}]}{M(M-1)(M-2)} + \dots \Big\},$$
(C.13)

$$\frac{\mathcal{R}_3^{(M)}}{\Gamma(M)} = \frac{(-1)^{M-1}}{2\pi} \Big\{ \frac{2[R_2^{(2)} - R_2]}{M(M-1)(M-2)} + \dots \Big\},$$
(C.14)

$$\frac{Y_0^{(M-1)}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} \left\{ \frac{b}{M} + \frac{b + R_0^{(0)} + a_1 [R_0^{(0)} - R_0]}{M(M-1)} + \frac{R_0^{(0)} + R_1^{(0)} + a_1 [R_0^{(0)} - R_0] + a_1 [R_1^{(0)} - R_1^{(1)}] + a_2 [R_1 - R_1^{(1)}]}{M(M-1)(M-2)} + \dots \right\},$$
(C.15)

$$\frac{Y_1^{(M-1)}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} \Big\{ \frac{-ba_1}{M(M-1)} - \frac{2ba_1 + a_1 R_0^{(0)} + a_2 [R_0^{(0)} - R_0]}{M(M-1)(M-2)} + \dots \Big\},$$
(C.16)

$$\frac{Y_2^{(M-1)}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} \Big\{ \frac{ba_2}{M(M-1)(M-2)} + \dots \Big\},\tag{C.17}$$

$$\frac{Y_0^{(M-2)}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} \Big\{ \frac{-b}{M(M-1)} - \frac{2b + R_0^{(0)} + a_1[R_0^{(0)} - R_0]}{M(M-1)(M-2)} + \dots \Big\},$$
(C.18)

$$\frac{Y_1^{(M-2)}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} \Big\{ \frac{ba_1}{M(M-1)(M-2)} + \dots \Big\},\tag{C.19}$$

$$\frac{Y_0^{(M-3)}}{\Gamma(M)} = \frac{(-1)^M}{2\pi} \Big\{ \frac{b}{M(M-1)(M-2)} + \dots \Big\}.$$
 (C.20)

Working up to NNNLO order, we need the asymptotic expansion of the combinations

$$\mathcal{L}_{1}^{(M)} = X_{1,0}Y_{0}^{(M-1)} + X_{1,1}Y_{1}^{(M-1)} + X_{1,2}Y_{2}^{(M-1)} + \dots,$$

$$\mathcal{L}_{2}^{(M)} = X_{2,0}Y_{0}^{(M-2)} + X_{2,1}Y_{1}^{(M-2)} + \dots,$$

$$\mathcal{L}_{3}^{(M)} = X_{3,0}Y_{0}^{(M-3)} + \dots$$
(C.21)

Comparing these expansions to $\mathcal{R}_1^{(M)}$, $\mathcal{R}_2^{(M)}$, and $\mathcal{R}_3^{(M)}$ and matching 3, 2, and 1 asymptotic expansion coefficients respectively, we can write down 6 equations. These can be used to express 6 leading coefficients, $\mathcal{R}_0^{(0)}$, $\mathcal{R}_1^{(0)}$, $\mathcal{R}_1^{(1)}$, $\mathcal{R}_2^{(0)}$, $\mathcal{R}_2^{(1)}$, and $\mathcal{R}_2^{(2)}$, in terms of the "generic" ones, b, \mathcal{R}_0 , \mathcal{R}_1 , and \mathcal{R}_2 . The result is

$$\begin{aligned} R_0^{(0)} &= R_0 - \frac{b}{4}, \\ R_1^{(0)} &= R_1 - \frac{R_0}{4} + \left(\frac{1}{32} - \frac{a}{2}\right)b, \\ R_1^{(1)} &= R_1 - \frac{9b}{32}, \\ R_2^{(0)} &= R_2 - \frac{R_1}{8} + \left(\frac{1}{64} - \frac{a}{4}\right)R_0 + \left(-\frac{3}{256} + \frac{a}{16} - \frac{a^2}{2}\right)b, \\ R_2^{(1)} &= R_2 - \frac{9R_0}{64} + \left(\frac{3}{128} - \frac{9a}{16}\right)b, \\ R_2^{(2)} &= R_2 - \frac{75}{256}b. \end{aligned}$$
(C.22)

Rewriting the above result in terms of the original "tilde" variables appearing in (2.31) we obtain (2.35).

D Resurgence

The results (2.35) show that the first few small-index $p_{a,b}$ quantities appear in the expression of the first few asymptotic coefficients and this structure strongly suggests a product form for the Δ_{-1} alien derivative.

In this appendix we will use the alternative representation of $\epsilon_p(z)$ given by (1.29) and write the Taylor expansion coefficients of its Borel transform as

$$c_n^{(\epsilon_p)} = (1+q)\frac{E_{n+1}(q)}{\Gamma(n+1)}, \qquad n = 0, 1, \dots$$
 (D.1)

We will assume that |q| < 1. Using the ansatz (2.31) we can write down the asymptotic expansion of the Borel transform coefficients:

$$c_n^{(\epsilon_p)} = (1+q)\frac{(-1)^n}{2\pi} \sum_{k=0}^{\infty} (-q)^k \left\{ b^{(k)} + \frac{\tilde{R}_0^{(k)}}{n} + \frac{\tilde{R}_1^{(k)}}{n(n-1)} + \frac{2\tilde{R}_2^{(k)}}{n(n-1)(n-2)} + \dots \right\}.$$
 (D.2)

Here, again, we extended the upper limit of the summation to infinity. Comparing this expansion with the generic forms (F.3) and (F.5) we get

$$\pi \tilde{B}_o = (1+q) \sum_{k=0}^{\infty} (-q)^k b^{(k)}, \tag{D.3}$$

$$\pi \tilde{B}_o \tilde{q}_m = (1+q) \sum_{k=0}^{\infty} (-q)^k \tilde{R}_m^{(k)}, \qquad m = 0, 1, \dots,$$
 (D.4)

$$\Delta_{-1}\epsilon_p = i(1+q)\sum_{k=0}^{\infty} (-q)^k b^{(k)} + i(1+q)\sum_{m=0}^{\infty} \frac{m!(-1)^{m+1}}{z^{m+1}}\sum_{k=0}^{\infty} (-q)^k \tilde{R}_m^{(k)}.$$
 (D.5)

We can compare this expression to the assumed product form

$$\Delta_{-1}\epsilon_p = iX\epsilon_p,\tag{D.6}$$

where

$$X = b + \sum_{m=0}^{\infty} \frac{X_m}{z^{m+1}}.$$
 (D.7)

Before making the comparison it is useful to recast (D.6) in the form

$$X\epsilon_p = X + (1+q)\sum_{n=0}^{\infty} \frac{\phi_{n+1}(q)}{z^{n+1}},$$
 (D.8)

where

$$X\sum_{n=0}^{\infty} \frac{E_{n+1}(q)}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{\phi_{n+1}(q)}{z^{n+1}}.$$
 (D.9)

The comparison of (D.5) with (D.6) gives

$$\sum_{k=0}^{\infty} (-q)^k b^{(k)} = \frac{b}{1+q} = b \sum_{k=0}^{\infty} (-q)^k,$$
(D.10)

$$\sum_{k=0}^{\infty} (-q)^k \tilde{R}_m^{(k)} = \frac{(-1)^{m+1}}{m!} \left\{ \frac{X_m}{1+q} + \phi_{m+1}(q) \right\}.$$
 (D.11)

The first of these gives

$$b^{(k)} = b, \qquad k = 0, 1, \dots$$
 (D.12)

This is what we obtained already from the consistency of asymptotic expansions. Before using the second comparison (D.11) it is useful to write out $\phi_{m+1}(q)$ explicitly:

$$\phi_{m+1}(q) = bE_{m+1}(q) + \sum_{r=0}^{m-1} X_r E_{m-r}(q) = b\sum_{k=0}^m p_{m,k}q^k + \sum_{k=0}^{m-1} \sum_{r=0}^{m-1-k} X_r p_{m-r-1,k}q^k.$$
 (D.13)

We can now read off from (D.11) the relations (2.37)-(2.39). The structure (2.37) is precisely the same as we already found. Writing out (2.38) and (2.39) explicitly for m = 0, 1, 2 we get

$$\begin{split} \tilde{R}_{0}^{(0)} &= \tilde{R}_{0} - p_{0,0}b, \\ \tilde{R}_{1}^{(0)} &= \tilde{R}_{1} + bp_{1,0} + X_{0}p_{0,0} = \tilde{R}_{1} - p_{0,0}\tilde{R}_{0} + p_{1,0}b, \\ \tilde{R}_{1}^{(1)} &= \tilde{R}_{1} - p_{1,1}b, \\ \tilde{R}_{2}^{(0)} &= \tilde{R}_{2} - \frac{1}{2} \left\{ bp_{2,0} + X_{0}p_{1,0} + X_{1}p_{0,0} \right\} \\ &= \tilde{R}_{2} - \frac{1}{2}p_{0,0}\tilde{R}_{1} + \frac{1}{2}p_{1,0}\tilde{R}_{0} - \frac{1}{2}p_{2,0}b, \\ \tilde{R}_{2}^{(1)} &= \tilde{R}_{2} + \frac{1}{2} \left\{ bp_{2,1} + X_{0}p_{1,1} \right\} = \tilde{R}_{2} - \frac{1}{2}p_{1,1}\tilde{R}_{0} + \frac{1}{2}p_{2,1}b, \\ \tilde{R}_{2}^{(2)} &= \tilde{R}_{2} - \frac{1}{2}p_{2,2}b. \end{split}$$
(D.14)

Here we used the identifications (2.37). We see that (D.14) exactly reproduces the earlier result (2.35).

The actual values of the generic coefficients are given by (2.33):

$$X_m = 2W_m, \qquad m = 0, 1, \dots$$
 (D.15)

Thus the coefficient function is

$$X = 1 + \sum_{m=0}^{\infty} \frac{2W_m}{z^{m+1}} = \epsilon$$
 (D.16)

and

$$\Delta_{-1}\epsilon_p = i\epsilon\epsilon_p. \tag{D.17}$$

E Asymptotic expansion of ρ

The asymptotic expansion (C.4) can also be used to obtain the asymptotic series for ρ . Using the inversion formula

$$w_{0,M} = \sum_{n=0}^{M} (-1)^n E_{1,n} 2^{n-M} Y_n^{(M)}$$
(E.1)

we have

$$\tilde{u}_M = 2^M c_{0,M} = 2^M w_{0,M} = \sum_{n=0}^M \omega_n Y_n^{(M)}, \qquad \omega_n = (-2)^n E_{1,n}.$$
(E.2)

Here are the first few ω_n coefficients:

$$\omega_0 = 1, \qquad \omega_1 = \frac{3}{4}, \qquad \omega_2 = -\frac{15}{32}, \qquad \omega_3 = \frac{105}{128}$$
 (E.3)

and the first few \tilde{u}_n values:

$$\tilde{u}_0 = 1, \qquad \tilde{u}_1 = a - \frac{3}{4}, \qquad \tilde{u}_2 = -\frac{15}{32} + \frac{3a}{4} - \frac{a^2}{2},$$
(E.4)

$$\tilde{u}_3 = -\frac{105}{128} + \frac{45a}{32} - \frac{9a^2}{8} + \frac{a^3}{2} + \frac{3}{8}\zeta_3.$$
(E.5)

We also recall the first few input parameters:

$$a_0 = 1,$$
 $a_1 = a - 1,$ $a_2 = 1 - a + \frac{a^2}{2} - \frac{1}{4}\zeta_2,$ (E.6)

$$a_3 = a - 1 - \frac{a^2}{2} + \frac{a^3}{6} + \frac{1 - a}{4}\zeta_2 + \frac{1}{4}\zeta_3.$$
 (E.7)

The Taylor coefficients of the Borel transform of ρ are

$$c_{M-1}^{(\rho)} = \frac{\tilde{u}_M}{\Gamma(M)} = \sum_{n=0}^M \omega_n \frac{Y_n^{(M)}}{\Gamma(M)}.$$
 (E.8)

To a given order of the asymptotic expansion (here we again work up to NNNLO) only a finite number of terms contribute to (C.4):

$$\frac{Y_n^{(M)}}{\Gamma(M)} = \frac{(-1)^{M-n}}{2\pi(M-1)(\dots)(M-n)} \left\{ -a_n \left[b + \frac{R_0^{(0)}}{M-n} + \frac{R_1^{(0)}}{(M-n)(M-n-1)} + \frac{2R_2^{(0)}}{(M-n)(M-n-1)(M-n-2)} \right] - a_{n+1} \left[\frac{R_0^{(0)} - R_0}{M-n} + \frac{R_1^{(0)} - R_1^{(1)}}{(M-n)(M-n-1)} + \frac{2[R_2^{(0)} - R_2^{(1)}]}{(M-n)(M-n-1)} \right] + a_{n+2} \left[\frac{R_1^{(1)} - R_1}{(M-n)(M-n-1)} + \frac{2[R_2^{(1)} - R_2^{(2)}]}{(M-n)(M-n-1)(M-n-2)} \right] - a_{n+3} \left[\frac{2[R_2^{(2)} - R_2]}{(M-n)(M-n-1)(M-n-2)} \right] + \dots \right\}.$$
(E.9)

To NNNLO we will need the n=0,1,2,3 cases and taking into account the shift of the index M by 1 unit we have

$$\frac{Y_0^{(M+1)}}{\Gamma(M+1)} = \frac{(-1)^M}{2\pi} \left\{ b + \frac{(1+a_1)R_0^{(0)} - a_1R_0}{M} + \frac{1}{M(M-1)} \left[(1+a_1)(R_1^{(0)} - R_0^{(0)}) - (a_1+a_2)R_1^{(1)} + a_1R_0 + a_2R_1 \right] + \frac{2}{M(M-1)(M-2)} \left[(1+a_1)(R_2^{(0)} - R_1^{(0)} + R_0^{(0)}) - (a_1+a_2)(R_2^{(1)} - R_1^{(1)}) + (a_2+a_3)R_2^{(2)} - a_1R_0 - a_2R_1 - a_3R_2 \right] + \dots \right\},$$
(E.10)

$$\frac{Y_1^{(M+1)}}{\Gamma(M+1)} = \frac{(-1)^M}{2\pi} \Big\{ -\frac{ba_1}{M} + \frac{a_2 R_0 - (a_1 + a_2) R_0^{(0)}}{M(M-1)} \\
+ \frac{(a_1 + a_2) (R_0^{(0)} - R_1^{(0)}) + (a_2 + a_3) R_1^{(1)} - a_2 R_0 - a_3 R_1}{M(M-1)(M-2)} + \dots \Big\},$$
(E.11)

$$\frac{Y_2^{(M+1)}}{\Gamma(M+1)} = \frac{(-1)^M}{2\pi} \Big\{ \frac{ba_2}{M(M-1)} + \frac{(a_2+a_3)R_0^{(0)} - a_3R_0}{M(M-1)(M-2)} + \dots \Big\},$$
 (E.12)

$$\frac{Y_3^{(M+1)}}{\Gamma(M+1)} = \frac{(-1)^M}{2\pi} \bigg\{ -\frac{ba_3}{M(M-1)(M-2)} + \dots \bigg\}.$$
 (E.13)

Comparing the expansion of $c_n^{(\rho)}$ to the generic case (F.3) we find for the constants determining the alien derivative of ρ :

$$\begin{aligned} \pi \tilde{B}_{o} &= b, \\ \pi \tilde{B}_{o} \tilde{q}_{0} &= (1+a_{1})R_{0}^{(0)} - a_{1}R_{0} - ba_{1}\omega_{1}, \\ \pi \tilde{B}_{o} \tilde{q}_{1} &= (1+a_{1})(R_{1}^{(0)} - R_{0}^{(0)}) - (a_{1}+a_{2})R_{1}^{(1)} + a_{1}R_{0} + a_{2}R_{1} \\ &+ \omega_{1}[a_{2}R_{0} - (a_{1}+a_{2})R_{0}^{(0)}] + ba_{2}\omega_{2}, \\ 2\pi \tilde{B}_{o} \tilde{q}_{2} &= 2[(1+a_{1})(R_{2}^{(0)} - R_{1}^{(0)} + R_{0}^{(0)}) - (a_{1}+a_{2})(R_{2}^{(1)} - R_{1}^{(1)}) \\ &+ (a_{2}+a_{3})R_{2}^{(2)} - a_{1}R_{0} - a_{2}R_{1} - a_{3}R_{2}] \\ &+ \omega_{1}[(a_{1}+a_{2})(R_{0}^{(0)} - R_{1}^{(0)}) + (a_{2}+a_{3})R_{1}^{(1)} - a_{2}R_{0} - a_{3}R_{1}] \\ &+ \omega_{2}[(a_{2}+a_{3})R_{0}^{(0)} - a_{3}R_{0}] - ba_{3}\omega_{3}. \end{aligned}$$

$$(E.14)$$

Using the solution (C.22) this simplifies to⁴

$$\pi \tilde{B}_{o} \tilde{q}_{0} = \tilde{R}_{0} + \left(\frac{3}{4} - a\right) b = -X_{0} - \tilde{u}_{1} b,$$

$$\pi \tilde{B}_{o} \tilde{q}_{1} = \tilde{R}_{1} + \left(\frac{3}{4} - a\right) \tilde{R}_{0} + \left(-\frac{15}{32} + \frac{3a}{4} - \frac{a^{2}}{2}\right) b = X_{1} + \tilde{u}_{1} X_{0} + \tilde{u}_{2} b,$$

$$2\pi \tilde{B}_{o} \tilde{q}_{2} = 2\tilde{R}_{2} + \left(\frac{3}{4} - a\right) \tilde{R}_{1} + \left(-\frac{15}{32} + \frac{3a}{4} - \frac{a^{2}}{2}\right) \tilde{R}_{0}$$

$$+ \left(\frac{105}{128} - \frac{45a}{32} + \frac{9a^{2}}{8} - \frac{a^{3}}{2} - \frac{3}{8}\zeta_{3}\right) b = -X_{2} - \tilde{u}_{1} X_{1} - \tilde{u}_{2} X_{0} - \tilde{u}_{3} b.$$
(E.15)

From here we can read off the alien derivative of ρ , given by (2.43).

F Alien derivatives

In this appendix we summarize our conventions for alien derivatives. For details, see [11-13].

Let us recall that we start from a formal asymptotic series

$$\Psi(z) = 1 + \sum_{n=1}^{\infty} \frac{\Gamma(n)c_{n-1}}{z^n}$$
(F.1)

and define its Borel transform as

$$\hat{\Psi}(t) = \sum_{n=0}^{\infty} c_n t^n.$$
(F.2)

Here we assume that the coefficients have the asymptotic expansion

$$c_{n} = \frac{\tilde{A}_{o}}{2} \left\{ 1 + \frac{\tilde{p}_{0}}{n} + \frac{\tilde{p}_{1}}{n(n-1)} + \frac{2\tilde{p}_{2}}{n(n-1)(n-2)} + \dots \right\}$$

$$+ (-1)^{n} \frac{\tilde{B}_{o}}{2} \left\{ 1 + \frac{\tilde{q}_{0}}{n} + \frac{\tilde{q}_{1}}{n(n-1)} + \frac{2\tilde{q}_{2}}{n(n-1)(n-2)} + \dots \right\}.$$
(F.3)

In this case the singular part of the Borel transform is

$$\hat{\Psi}^{\text{sing}}(t) = \frac{\tilde{A}_o}{2} \left\{ \frac{1}{1-t} - \ln(1-t) \sum_{m=0}^{\infty} \tilde{p}_m (t-1)^m \right\} + \frac{\tilde{B}_o}{2} \left\{ \frac{1}{1+t} - \ln(1+t) \sum_{m=0}^{\infty} (-1)^m \tilde{q}_m (t+1)^m \right\}$$
(F.4)

and the alien derivatives are given by

$$\Delta_{1}\Psi(z) = -i\pi\tilde{A}_{o}\left\{1 + \sum_{m=0}^{\infty} \frac{m!\tilde{p}_{m}}{z^{m+1}}\right\},$$

$$\Delta_{-1}\Psi(z) = i\pi\tilde{B}_{o}\left\{1 - \sum_{m=0}^{\infty} \frac{m!(-1)^{m}\tilde{q}_{m}}{z^{m+1}}\right\}.$$
(F.5)
$$\overline{R_{v} - R_{0}}, \tilde{R}_{0} = R_{0} - R_{1} + R_{0}$$

⁴Note that $\tilde{R}_0 = R_0$, $\tilde{R}_1 = R_1 - R_0$, $\tilde{R}_2 = R_2 - R_1 + R_0$.

G The $\eta_k = 1$ model

Since all η_k are algebraically independent, it makes sense to study the simplest model

$$\eta_k = 1, \qquad k = 1, 2, \dots$$
 (G.1)

For this model

$$g(x) = 1, \qquad a_n = \delta_{n,0}.$$
 (G.2)

Let us make the following definitions.

$$K_n = p_{n-1,0}, \quad n = 1, 2, \dots, \qquad K_0 = 1$$
 (G.3)

and

$$S_n = 2^n E_{0,n} = \frac{1}{n!} \prod_{\ell=1}^n \left(\ell - \frac{1}{2}\right)^2, \quad n = 1, 2..., \qquad S_0 = 1.$$
 (G.4)

In this simple model the building blocks simplify enormously:

$$Y_n^{(M)} = \delta_{n,0} K_M, \quad n = 0, \dots, M,$$
 (G.5)

$$w_{a,M} = \delta_{a,0} 2^{-M} K_M, \quad M = 0, 1, \dots, \quad a = 0, \dots, M.$$
 (G.6)

The left and right sides of the Volin equations simplify accordingly to

$$\mathcal{R}_1^{(1)} = p_{0,0} \tag{G.7}$$

and

$$M \ge 2 \quad \begin{cases} \mathcal{R}_r^{(M)} = p_{M-1,r-1} + p_{M-1,r}, & r = 1, \dots, M-1, \\ \mathcal{R}_M^{(M)} = p_{M-1,M-1}, \end{cases}$$
(G.8)

$$\mathcal{L}_r^{(M)} = S_r K_{M-r}.$$
 (G.9)

Equating the two sides we obtain

$$M \ge 1 \qquad p_{M-1,M-1} = S_M,$$
 (G.10)

$$M \ge 2$$
 $p_{M-1,r-1} + p_{M-1,r} = S_r K_{M-r}, \quad r = 1, \dots, M-1.$ (G.11)

It is easy to find the solution of this recursion:

$$p_{M-1,M-k} = \sum_{p=0}^{k-1} (-1)^{k-p+1} K_p S_{M-p}, \qquad k = 1, \dots, M.$$
 (G.12)

In particular, the most interesting unknowns K_n are determined recursively as

$$K_{M+1} = \sum_{p=0}^{M} (-1)^{M-p} K_p S_{M+1-p}, \qquad M = 0, 1, \dots$$
(G.13)

Summing up (G.11) from r = 1 to r = M - 1 we obtain

$$2W_{M-1} = \sum_{r=0}^{M} S_r K_{M-r}, \qquad (G.14)$$

which also holds for M = 1. If we define

$$\xi_M = \sum_{r=0}^M S_r K_{M-r}, \tag{G.15}$$

we can write

$$\epsilon(x) = \sum_{n=0}^{\infty} \xi_n x^n. \tag{G.16}$$

Here x = 1/z. Similarly

$$\rho(x) = \sum_{n=0}^{\infty} K_n x^n \tag{G.17}$$

and we also introduce

$$B(x) = \sum_{n=0}^{\infty} S_n x^n.$$
(G.18)

The solution of the recursion (G.13) is summarized by

$$\rho(x)B(-x) = 1 \tag{G.19}$$

and (G.15) is equivalent to

$$\epsilon(x) = B(x)\rho(x). \tag{G.20}$$

Let us introduce

$$B(x) = 1 + b(x),$$
 $b(x) = \sum_{n=1}^{\infty} S_n x^n.$ (G.21)

The Borel transform of b(x) is

$$\hat{b}(t) = \sum_{n=0}^{\infty} \beta_n t^n, \qquad (G.22)$$

where

$$\beta_n = \frac{S_{n+1}}{\Gamma(n+1)}, \quad n = 0, 1, \dots$$
 (G.23)

We also define

$$D(x) = B(-x) = 1 + d(x), \qquad d(x) = b(-x)$$
 (G.24)

and see that

$$\hat{d}(t) = -\hat{b}(-t).$$
 (G.25)

By inspecting the Taylor expansion coefficients (G.23) we recognize that

$$\hat{b}(t) = \frac{1}{4^2} F_1\left(\frac{3}{2}, \frac{3}{2}; 2; t\right).$$
(G.26)

Using Mathematica we can verify that in the vicinity of t = 1

$$\hat{b}(t) = -\frac{1}{\pi(t-1)} - \frac{\ln(1-t)}{\pi}\hat{d}(t-1) + \Phi(t-1),$$
(G.27)

where Φ is regular around the origin. This can be written in the language of alien derivatives as

$$\Delta_1 B = -2i - 2id = -2iD. \tag{G.28}$$

Similarly we have

$$\Delta_{-1}D = -2iB. \tag{G.29}$$

From

$$\rho = \frac{1}{D}, \qquad \epsilon = \frac{B}{D}$$
(G.30)

we calculate

$$\Delta_1 \rho = 0, \qquad \Delta_1 \epsilon = \frac{-2iD}{D} = -2i, \qquad (G.31)$$

$$\Delta_{-1}\rho = -\frac{1}{D^2}\Delta_{-1}D = \frac{2iB}{D^2} = 2i\epsilon\rho,$$
(G.32)

$$\Delta_{-1}\epsilon = -\frac{B}{D^2}\Delta_{-1}D = \frac{2iB^2}{D^2} = 2i\epsilon^2.$$
 (G.33)

This structure of alien derivatives is very similar to what was found in the main part of the paper for the full model, it is only that Δ_1 is by a factor 2 too small and Δ_{-1} is by a factor 2 too large. However, the overall factors can be changed easily, as we will see in the next subsection.

G.1 *a*-dependence

The alien derivatives transform under a change of variables. If the original expansion parameter z is changed to x, where

$$z = z(x), \tag{G.34}$$

then the asymptotic expansions $\gamma(z)$ are changed to C(x) by the simple substitution rule

$$C(x) = \gamma(z(x)). \tag{G.35}$$

Let us denote the original alien derivative (obtained from the Borel transform of $\gamma(z)$) at ω by $\Delta_{\omega}\gamma$ and the alien derivative in the new variable (obtained from the Borel transform of C(x)) by $D_{\omega}C$. The transformation rule $\Delta_{\omega} \longrightarrow D_{\omega}$ is given by [37]

$$D_{\omega}\gamma(z(x)) = e^{-\omega(z(x)-x)}(\Delta_{\omega}\gamma)(z(x)) + \gamma'(z(x))(D_{\omega}z)(x).$$
(G.36)

In the case of a simple shift by a constant m,

$$z(x) = x + m, \tag{G.37}$$

(G.36) reduces to a constant rescaling:

$$D_{\omega}C = \mathrm{e}^{-m\omega}\Delta_{\omega}\gamma. \tag{G.38}$$

If we treat the symbol a in (1.11) as a generic parameter (rather then a shorthand for its true numerical value $a = \ln 2$) then

$$\epsilon(a, z)$$
 and $s(a, z) = z\rho^2(a, z)$ (G.39)

have the following remarkable property⁵:

$$s(a, z - 2a) = s_o(z), \qquad \epsilon(a, z - 2a) = \epsilon_o(z), \qquad (G.40)$$

where $s_o(z)$ and $\epsilon_o(z)$ are independent of the parameter a.

In the simplified model we calculated $\epsilon(1, z)$ and $\rho(1, z)$. For example for ϵ we then have

$$\epsilon(a, x) = \epsilon(1, x + 2a - 2) \tag{G.41}$$

and

$$D_{\omega}\epsilon = e^{(2-2a)\omega}\Delta_{\omega}\epsilon = \left(\frac{e}{2}\right)^{2\omega}\Delta_{\omega}\epsilon.$$
 (G.42)

The transformation for the alien derivatives of ρ is the same.

We see that after restoring $a = \ln 2$ the alien derivatives of ϵ change to

$$D_1 \epsilon = -i(e^2/2), \qquad D_{-1} \epsilon = i(8/e^2)\epsilon^2.$$
 (G.43)

The changes for the alien derivatives of ρ are the same. It is remarkable that these coefficients are numerically quite close to the "true" values (4 and 1):

$$D_1 \epsilon = -3.69i,$$
 $D_{-1} \epsilon = 1.08i\epsilon^2.$ (G.44)

G.2 The alien derivatives of ϵ_p for the $\eta_k = 1$ model

We calculate ϵ_p for $\eta_k = 1$ as follows. Let us multiply (G.11) by q^r and sum over r from 1 to M - 1. In this way we obtain the relation (for $M \ge 2$)

$$\sum_{r=0}^{M} S_r K_{M-r} q^r = (1+q) E_M(q).$$
 (G.45)

The above relation also holds for M = 1 and we can write

$$\epsilon_p = 1 + (1+q) \sum_{M=1}^{\infty} \frac{E_M(q)}{z^M} = \sum_{M=0}^{\infty} \frac{1}{z^M} \sum_{r=0}^M S_r K_{M-r} q^r$$

$$= \sum_{r=0}^{\infty} \sum_{k=0}^{\infty} x^{r+k} S_r K_k q^r = \rho(x) B(qx).$$
(G.46)

⁵These hold for arbitrary η_k , not just for the simplified model $\eta_k = 1$. We have verified this property perturbatively up to 16th order using Mathematica.

For clarity let us define

$$P(x) = B(qx) = 1 + p(x) = 1 + \sum_{n=1}^{\infty} S_n q^n x^n.$$
 (G.47)

The Borel transform of p(x) is

$$\hat{p}(t) = q\hat{b}(qt) \tag{G.48}$$

and therefore its singular part, using (G.27), is

$$\hat{p}(t) \approx \frac{-q}{\pi(qt-1)} - \frac{q\ln(1-qt)}{\pi} \hat{d}(qt-1) + \dots$$

$$\approx \frac{-1}{\pi(t-p)} - \frac{\ln(p-t)}{\pi} q \hat{d}(q(t-p)) + \dots$$
(G.49)

From here we can read off the alien derivative

$$\Delta_p P = -2i - 2id(qx) = -2iD(qx). \tag{G.50}$$

Now we can calculate

$$\Delta_p \epsilon_p = \rho(x) \Delta_p P(x) = -2i\rho(x)D(qx) = -2i\epsilon_{-p} \tag{G.51}$$

and

$$\Delta_{-1}\epsilon_p = B(qx)\Delta_{-1}\rho = 2i\epsilon\rho B(qx) = 2i\epsilon\epsilon_p. \tag{G.52}$$

References

- D. Volin, From the mass gap in O(N) to the non-Borel-summability in O(3) and O(4) sigma-models, Phys. Rev. D 81 (2010) 105008, [arXiv:0904.2744].
- [2] D. Volin, Quantum integrability and functional equations: Applications to the spectral problem of AdS/CFT and two-dimensional sigma models. PhD thesis, 2009. arXiv:1003.4725.
- [3] A. B. Zamolodchikov and A. B. Zamolodchikov, Relativistic Factorized S Matrix in Two-Dimensions Having O(N) Isotopic Symmetry, JETP Lett. 26 (1977) 457.
- [4] A. M. Polyakov and P. B. Wiegmann, Theory of Nonabelian Goldstone Bosons, Phys. Lett. B 131 (1983) 121–126.
- [5] P. Hasenfratz, M. Maggiore, and F. Niedermayer, The Exact mass gap of the O(3) and O(4) nonlinear sigma models in d = 2, Phys. Lett. B 245 (1990) 522–528.
- [6] P. Hasenfratz and F. Niedermayer, The Exact mass gap of the O(N) sigma model for arbitrary $N \ge 3$ in d = 2, Phys. Lett. B **245** (1990) 529–532.
- [7] M. Mariño and T. Reis, Renormalons in integrable field theories, JHEP 04 (2020) 160, [arXiv:1909.12134].
- [8] M. C. Abbott, Z. Bajnok, J. Balog, and A. Hegedűs, From perturbative to non-perturbative in the O (4) sigma model, Phys. Lett. B 818 (2021) 136369, [arXiv:2011.09897].

- M. C. Abbott, Z. Bajnok, J. Balog, A. Hegedűs, and S. Sadeghian, Resurgence in the O(4) sigma model, JHEP 05 (2021) 253, [arXiv:2011.12254].
- [10] C. Bauer, G. S. Bali, and A. Pineda, Compelling Evidence of Renormalons in QCD from High Order Perturbative Expansions, Phys. Rev. Lett. 108 (2012) 242002, [arXiv:1111.3946].
- [11] M. Mariño, Lectures on non-perturbative effects in large N gauge theories, matrix models and strings, Fortsch. Phys. 62 (2014) 455–540, [arXiv:1206.6272].
- [12] D. Dorigoni, An Introduction to Resurgence, Trans-Series and Alien Calculus, Annals Phys. 409 (2019) 167914, [arXiv:1411.3585].
- [13] I. Aniceto, G. Basar, and R. Schiappa, A Primer on Resurgent Transseries and Their Asymptotics, Phys. Rept. 809 (2019) 1–135, [arXiv:1802.10441].
- [14] I. Aniceto, R. Schiappa, and M. Vonk, The Resurgence of Instantons in String Theory, Commun. Num. Theor. Phys. 6 (2012) 339–496, [arXiv:1106.5922].
- [15] I. Aniceto, J. G. Russo, and R. Schiappa, Resurgent Analysis of Localizable Observables in Supersymmetric Gauge Theories, JHEP 03 (2015) 172, [arXiv:1410.5834].
- [16] I. Aniceto and R. Schiappa, Nonperturbative Ambiguities and the Reality of Resurgent Transseries, Commun. Math. Phys. 335 (2015), no. 1 183–245, [arXiv:1308.1115].
- [17] I. Aniceto, The Resurgence of the Cusp Anomalous Dimension, J. Phys. A 49 (2016) 065403, [arXiv:1506.03388].
- [18] G. Arutyunov, D. Dorigoni, and S. Savin, Resurgence of the dressing phase for AdS5xS5, JHEP 01 (2017) 055, [arXiv:1608.03797].
- [19] S. Demulder, D. Dorigoni, and D. C. Thompson, Resurgence in η-deformed Principal Chiral Models, JHEP 07 (2016) 088, [arXiv:1604.07851].
- [20] D. Dorigoni and Y. Hatsuda, Resurgence of the Cusp Anomalous Dimension, JHEP 09 (2015) 138, [arXiv:1506.03763].
- [21] D. Dorigoni and P. Glass, The grin of Cheshire cat resurgence from supersymmetric localization, SciPost Phys. 4 (2018), no. 2 012, [arXiv:1711.04802].
- [22] G. V. Dunne and M. Ünsal, What is QFT? Resurgent trans-series, Lefschetz thimbles, and new exact saddles, PoS LATTICE2015 (2016) 010, [arXiv:1511.05977].
- [23] G. V. Dunne and M. Unsal, Resurgence and Dynamics of O(N) and Grassmannian Sigma Models, JHEP 09 (2015) 199, [arXiv:1505.07803].
- [24] G. V. Dunne and M. Ünsal, New Nonperturbative Methods in Quantum Field Theory: From Large-N Orbifold Equivalence to Bions and Resurgence, Ann. Rev. Nucl. Part. Sci. 66 (2016) 245-272, [arXiv:1601.03414].
- [25] T. Fujimori, S. Kamata, T. Misumi, M. Nitta, and N. Sakai, Bion non-perturbative contributions versus infrared renormalons in two-dimensional CP^{N-1} models, JHEP 02 (2019) 190, [arXiv:1810.03768].
- [26] M. Marino, R. Miravitllas, and T. Reis, *Testing the Bethe ansatz with large N renormalons*, arXiv:2102.03078.

- [27] L. Di Pietro, M. Mariño, G. Sberveglieri, and M. Serone, Resurgence and 1/N Expansion in Integrable Field Theories, JHEP 10 (2021) 166, [arXiv:2108.02647].
- [28] M. Marino and T. Reis, Exact perturbative results for the Lieb-Liniger and Gaudin-Yang models, arXiv:1905.09575.
- [29] M. Marino and T. Reis, *Three roads to the energy gap*, arXiv:2010.16174.
- [30] M. Marino and T. Reis, *Resurgence and renormalons in the one-dimensional Hubbard model*, arXiv:2006.05131.
- [31] M. Serone, G. Spada, and G. Villadoro, $\lambda \phi^4$ Theory I: The Symmetric Phase Beyond NNNNNNNLO, JHEP 08 (2018) 148, [arXiv:1805.05882].
- [32] M. Serone, G. Spada, and G. Villadoro, $\lambda \phi_2^4$ theory Part II. the broken phase beyond NNNN(NNN)LO, JHEP **05** (2019) 047, [arXiv:1901.05023].
- [33] I. Aniceto and M. Spaliński, Resurgence in Extended Hydrodynamics, Phys. Rev. D 93 (2016), no. 8 085008, [arXiv:1511.06358].
- [34] I. Aniceto, B. Meiring, J. Jankowski, and M. Spaliński, The large proper-time expansion of Yang-Mills plasma as a resurgent transseries, JHEP 02 (2019) 073, [arXiv:1810.07130].
- [35] Z. Bajnok, J. Balog, B. Basso, G. Korchemsky, and L. Palla, Scaling function in AdS/CFT from the O(6) sigma model, Nucl. Phys. B 811 (2009) 438-462, [arXiv:0809.4952].
- [36] M. Beneke, *Renormalons*, *Phys. Rept.* **317** (1999) 1–142, [hep-ph/9807443].
- [37] E. Delabaere and F. Pham, Resurgent methods in semi-classical asymptotics, Annales de l'I.H.P. Physique théorique 71 (1999), no. 1 1–94.
- [38] M. Marino, R. Miravitllas, and T. Reis, *New renormalons from analytic trans-series*, arXiv:2111.11951.