

The Scalar Potential of Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model

M. C. Rodriguez

Grupo de Física Teórica e Matemática Física

Departamento de Física

Universidade Federal Rural do Rio de Janeiro (UFRRJ)

BR 465 Km 7, 23890-000 Seropédica - RJ

Brasil

Abstract

We will study in details the scalar potential of the Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model where we introduce the anti-sextet, and without the anti-sextet, in our analyses of all masses of the usual scalar sector.

PACS number(s): 12.60. Jv

Keywords: Supersymmetric models

1 Introduction

Models with the following gauge symmetry

$$SU(3)_C \otimes SU(3)_L \otimes U(1)_N \quad (1)$$

are known as $3 - 3 - 1$ for short. They are interesting possibilities for the physics at the TeV scale [1, 2, 3, 4].

The charge operator in the $SU(3)_L$ generators,

$$\frac{Q}{e} = \frac{1}{2}(\lambda_3 - \vartheta\lambda_8) + N I_{3 \times 3}, \quad (2)$$

where the ϑ and N are parameters defining different representation contents and λ_3, λ_8 are the diagonal generators of $SU(3)$ given by

$$\lambda_3 = \begin{pmatrix} +1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} +1 & 0 & 0 \\ 0 & +1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \quad (3)$$

The model presented in Ref. [2], with $\vartheta = \sqrt{3}$, the Eq.(2) become

$$\frac{Q}{e} = \begin{pmatrix} N & 0 & 0 \\ 0 & N-1 & 0 \\ 0 & 0 & N+1 \end{pmatrix}, \quad (4)$$

and its supersymmetric version has already been considered in Refs. [5, 6, 7, 8]. While the 3-3-1 model of Refs. [3] which corresponds the the case when $\vartheta = (1/\sqrt{3})$, the Eq.(2) become

$$\frac{Q}{e} = \begin{pmatrix} N + \frac{1}{3} & 0 & 0 \\ 0 & N - \frac{2}{3} & 0 \\ 0 & 0 & N + \frac{1}{3} \end{pmatrix}, \quad (5)$$

his supersymmetric version was built in [9, 10].

In fact, this may be the last symmetry involving the lightest elementary particles: leptons. The lepton sector is exactly the same as in the Standard Model (SM) [11] but now there is a symmetry, at large energies among, say e^- , ν_e and e^+ [2] or between e^- , ν_e and ν_e^c [3], therefore there are many model we can construct [4]. Once this symmetry is imposed on the lightest generation and extended to the other leptonic generations it follows that the quark sector must be enlarged by considering heavy exotic charged quarks. It means that some gauge bosons carry lepton and baryon quantum number.

There are two main versions of the 3-3-1 models depending on the embedding of the charge operator in the $SU(3)_L$ generators is defined at Eq.(2). In the minimal version, with $\vartheta = \sqrt{3}$, the charge conjugation of the right-handed charged lepton for each generation is combined with the usual $SU(2)_L$ doublet left-handed leptons components to form an $SU(3)$ triplet $(\nu_i, l_i, l_i^c)_L$. No extra leptons are needed in the mentioned model, and we shall call such model as minimal 3-3-1 model. There are also another possibility where the triplet $(\nu_i, l_i, L_i^c)_L$ where L is an extra charged leptons which do not mix with the known leptons [12]. We want to remind that there is no right-handed (RH) neutrino in both model. There exists another interesting possibility, where ($\vartheta = 1/\sqrt{3}$) a left-handed anti-neutrino to each usual $SU(2)_L$ doublet is added to form an $SU(3)$ triplet $(\nu_i, l_i, \nu_i^c)_L$, and this model is called the 3-3-1 model with RH neutrinos. The 3-3-1 models have been studied extensively over the last decade.

Although this model coincides at low energies with the SM it explains some fundamental questions that are accommodated, but not explained, in the SM. These questions are:

1. The family number must be a multiple of three in order to cancel anomalies [2, 3];
2. Why $\sin^2 \theta_W < \frac{1}{4}$ is observed;
3. The models have a scalar sector similar to the two Higgs doublets Model, we will review it at App.(A);
4. It is the simplest model that includes bileptons of both types: scalars and vectors ones;
5. It solves the strong CP -problem;
6. The model has several sources of CP violation [13, 14].

The supersymmetric version of the 3-3-1 minimal model [2] was done Refs. [5, 6, 7, 8, 15] (MSUSY331). The scalar sector of the minimal 331 model [2] was studied on [16, 17]. When we have presented the MSUSY we have shown the lightest higgs has mass around 124.5 GeV [7].

The scalar sector of 331 model with heavy leptons [12] was studied on [13, 17] and we have studied the Higgs potential with only triplets in supersymmetric case in [15]. The scalar potential in several 331 models was considered in [18, 19, 20, 21].

Recently it was reported a light Higgs boson at the LHC and its value is given by [22]

$$M_H = 125.09 \pm 0.21 \text{ (stat.)} \pm 0.11 \text{ (syst.) GeV.} \quad (6)$$

Our goal is to show that our results presented at [7, 15] is in agreement with the actual experimental data on scalar sector.

This paper is organized as follows. In Sec. 2 we review the minimal Supersymmetric 3-3-1 Model with R -Parity Conservation. Our results for the scalar potential with the anti-Sextet is presented in Sec. 3. We done similar analyses with only triplets scalars in Sec. 4. Finally, the last section is devoted to our conclusions. In App.(A) we review the mass spectrum of 3-3-1 Model without Supersymmetry, while in App.(B) we show how to get an analytical expression to our scalar potential in order to compare its expression with the ones used at 3-3-1 Model.

2 Minimal Supersymmetric 3-3-1 model (MSUSY331).

This supersymmetric version of the 3-3-1 model given at [2] was presented at [5, 6, 7, 8]. In the minimal 331 model, denoted as m331, the charged leptons gain mass from a triplet and a anti-sextet [2].

In the m331 the scalar sector is composed of three scalar triplets:

$$\begin{aligned} \eta &= \begin{pmatrix} \eta^0 \\ \eta_1^- \\ \eta_2^+ \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, 0), \quad \rho = \begin{pmatrix} \rho^+ \\ \rho^0 \\ \rho^{++} \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, +1), \\ \chi &= \begin{pmatrix} \chi^- \\ \chi^{--} \\ \chi^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, -1). \end{aligned} \quad (7)$$

In parenthesis it appears the transformations properties under the respective factors ($SU(3)_C, SU(3)_L, U(1)_N$). Those scalars, defined at Eq.(7), give masses for all the quarks and the charged leptons are massless. In order to give masses for the charged leptons, at tree level, we have to introduce the following anti-sextet

$$S = \begin{pmatrix} \sigma_1^0 & \frac{h_2^+}{\sqrt{2}} & \frac{h_1^-}{\sqrt{2}} \\ \frac{h_2^+}{\sqrt{2}} & H_1^{++} & \frac{\sigma_2^0}{\sqrt{2}} \\ \frac{h_1^-}{\sqrt{2}} & \frac{\sigma_2^0}{\sqrt{2}} & H_2^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{6}^*, 0). \quad (8)$$

The scalars get the following vacuum expectation values (VEVs) [7, 8]: When we break the 331 symmetry to the $SU(3)_C \otimes U(1)_{EM}$, the scalars get the following vacuum expectation values (VEVs):

$$\langle \eta \rangle = \begin{pmatrix} v \\ 0 \\ 0 \end{pmatrix}, \quad \langle \rho \rangle = \begin{pmatrix} 0 \\ u \\ 0 \end{pmatrix}, \quad \langle \chi \rangle = \begin{pmatrix} 0 \\ 0 \\ w \end{pmatrix}, \quad (9)$$

and

$$\langle S \rangle = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \frac{z}{\sqrt{2}} \\ 0 & \frac{z}{\sqrt{2}} & 0 \end{pmatrix}, \quad (10)$$

where $v = v_\eta/\sqrt{2}$, $u = v_\rho/\sqrt{2}$, $w = v_\chi/\sqrt{2}$ and $z = v_{\sigma_2}/\sqrt{2}$.

$$\begin{aligned}
v &= \frac{v_\eta}{\sqrt{2}} \left(1 + \frac{H_\eta + \imath F_\eta}{|v_\eta|} \right), & u &= \frac{v_\rho}{\sqrt{2}} \left(1 + \frac{H_\rho + \imath F_\rho}{|v_\rho|} \right), \\
w &= \frac{v_\chi}{\sqrt{2}} \left(1 + \frac{H_\chi + \imath F_\chi}{|v_\chi|} \right), & z &= \frac{v_{\sigma_2^0}}{\sqrt{2}} \left(1 + \frac{H_{\sigma_2^0} + \imath F_{\sigma_2^0}}{|v_{\sigma_2^0}|} \right).
\end{aligned} \tag{11}$$

Besides, in order to to cancel chiral anomalies generated by the superpartners of the scalars, we have to add the following higgsinos in the respective anti-multiplets,

$$\begin{aligned}
\tilde{\eta}' &= \begin{pmatrix} \tilde{\eta}'^0 \\ \tilde{\eta}'^+ \\ \tilde{\eta}'^- \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}^*, 0), & \tilde{\rho}' &= \begin{pmatrix} \tilde{\rho}'^- \\ \tilde{\rho}'^0 \\ \tilde{\rho}'^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}^*, -1), \\
\tilde{\chi}' &= \begin{pmatrix} \tilde{\chi}'^+ \\ \tilde{\chi}'^{++} \\ \tilde{\chi}'^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}^*, +1),
\end{aligned} \tag{12}$$

$$\tilde{S}' = \begin{pmatrix} \frac{\tilde{\sigma}_1'^0}{\sqrt{2}} & \frac{\tilde{h}_2'^-}{\sqrt{2}} & \frac{\tilde{h}_1'^+}{\sqrt{2}} \\ \frac{\tilde{h}_2'^-}{\sqrt{2}} & \tilde{H}_1'^{-} & \frac{\tilde{\sigma}_2'^0}{\sqrt{2}} \\ \frac{\tilde{h}_1'^+}{\sqrt{2}} & \frac{\tilde{\sigma}_2'^0}{\sqrt{2}} & \tilde{H}_2'^{++} \end{pmatrix}_L \sim (\mathbf{1}, \mathbf{6}, 0). \tag{13}$$

There are also the scalar partners of the Higgsinos defined in Eqs. (12,13) and we will denote them η', ρ', χ', S' [7, 8]. This model was presented at [5, 6, 7, 8].

The usual scalars get the vacuum expectation values (VEVs) defined at Eqs.(10,14). In similar way for the new scalars we can write [7, 8]:

$$\begin{aligned}
v' &= \frac{v_{\eta'}}{\sqrt{2}} \left(1 + \frac{H_{\eta'} + \imath F_{\eta'^0}}{|v_{\eta'}|} \right), & u' &= \frac{v_{\rho'}}{\sqrt{2}} \left(1 + \frac{H_{\rho'} + \imath F_{\rho'}}{|v_{\rho'}|} \right), \\
w' &= \frac{v_{\chi'}}{\sqrt{2}} \left(1 + \frac{H_{\chi'} + \imath F_{\chi'}}{|v_{\chi'}|} \right), & z' &= \frac{v_{\sigma_2'^0}}{\sqrt{2}} \left(1 + \frac{H_{\sigma_2'^0} + \imath F_{\sigma_2'^0}}{|v_{\sigma_2'^0}|} \right).
\end{aligned} \tag{14}$$

It is the complete set of fields in the minimal supersymmetric 3-3-1 model (MSUSY331).

2.1 Genral Superpotential at MSUSY331

The superpotential of our model is given by

$$W = \frac{W_2 + \bar{W}_2}{2} + \frac{W_3 + \bar{W}_3}{3}, \quad (15)$$

with W_2 having only two chiral superfields while W_3 has three chiral superfields. The terms allowed by our symmetry are

$$W_2 = \sum_{i=1}^3 \mu_{0i}(\hat{L}_i \hat{\eta}') + \mu_\eta(\hat{\eta} \hat{\eta}') + \mu_\rho(\hat{\rho} \hat{\rho}') + \mu_\chi(\hat{\chi} \hat{\chi}') + \mu_S Tr[(\hat{S} \hat{S}'), \quad (16)$$

all the coefficients in the equation above have mass dimension [23] and we also have

$$\begin{aligned} W_3 = & \sum_{i=1}^3 \left[\sum_{j=1}^3 \left(\sum_{k=1}^3 \lambda_{1ijk} (\epsilon \hat{L}_i \hat{L}_j \hat{L}_k) + \lambda_{2ij} (\epsilon \hat{L}_i \hat{L}_j \hat{\eta}) + \lambda_{3ij} (\hat{L}_i \hat{S} \hat{L}_j) \right) + \lambda_{4i} (\epsilon \hat{L}_i \hat{\chi} \hat{\rho}) \right] \\ & + f_1(\epsilon \hat{\rho} \hat{\chi} \hat{\eta}) + f_2(\hat{\chi} \hat{S} \hat{\rho}) + f_3(\hat{\eta} \hat{S} \hat{\eta}) + f_4 \epsilon_{ijk} \epsilon_{lmn} \hat{S}_{il} \hat{S}_{jm} \hat{S}_{kn} + f'_1(\epsilon \hat{\rho}' \hat{\chi}' \hat{\eta}') + f'_2(\hat{\chi}' \hat{S}' \hat{\rho}') \\ & + f'_3(\hat{\eta}' \hat{S}' \hat{\eta}') + f'_4 \epsilon_{ijk} \epsilon_{lmn} \hat{S}'_{il} \hat{S}'_{jm} \hat{S}'_{kn} + \sum_{i=1}^3 \left[\kappa_{1i} (\hat{Q}_3 \hat{\eta}') \hat{u}_i^c + \kappa_{2i} (\hat{Q}_3 \hat{\rho}') \hat{d}_i^c \right] + \kappa_3 (\hat{Q}_3 \hat{\chi}') \hat{J}^c \\ & + \sum_{\alpha=1}^2 \left[\sum_{i=1}^3 \left(\kappa_{4\alpha i} (\hat{Q}_\alpha \hat{\eta}) \hat{d}_i^c + \kappa_{5\alpha i} (\hat{Q}_\alpha \hat{\rho}) \hat{u}_i^c \right) + \sum_{\beta=1}^2 \kappa_{6\alpha\beta} (\hat{Q}_\alpha \hat{\chi}) \hat{J}_\beta^c + \sum_{i=1}^3 \sum_{j=1}^3 \kappa_{7\alpha ij} (\hat{Q}_\alpha \hat{L}_i) \hat{d}_j^c \right] \\ & + \sum_{i=1}^3 \left\{ \sum_{j=1}^3 \left[\sum_{k=1}^3 \xi_{1ijk} \hat{d}_i^c \hat{d}_j^c \hat{u}_k^c + \sum_{\beta=1}^2 \left(\xi_{2ij\beta} \hat{u}_i^c \hat{u}_j^c \hat{J}_\beta^c \right) + \xi_{3i\beta} \hat{d}_i^c \hat{J}_\beta^c \right] \right\}. \quad (17) \end{aligned}$$

all the coefficients in W_3 are dimensionless [23]. The terms proportional for f_4 and f'_4 were not introduced in our previous works done at [7, 8].

2.2 R-Parity charges at MSUSY331

Some terms in the superpotential W given in Eq. (15) violate the conservation of the new quantum number defined as

$$\mathcal{F} = B + L \quad (18)$$

quantum number, where B is the baryon number while L is the lepton number [12]. For instance, if we allow the ξ_1 term it implies in proton decay [6].

However, if we assume the global $U(1)_{\mathcal{F}}$ symmetry, it allows us to introduce the R -conserving symmetry, defined as

$$R = (-1)^{3\mathcal{F}+2S}. \quad (19)$$

The \mathcal{F} number attribution is

$$\begin{aligned} \mathcal{F}(U^{--}) &= \mathcal{F}(V^-) = -\mathcal{F}(J) = \mathcal{F}(j_{1,2}) = \mathcal{F}(\rho^{--}) \\ &= \mathcal{F}(\chi^{--}) = \mathcal{F}(\chi^-) = \mathcal{F}(\eta_2^-) = \mathcal{F}(\sigma_1^0) = 2, \end{aligned} \quad (20)$$

As in the Minimal Supersymmetric Standard Model this definition implies that all known standard model's particles have even R -parity while their supersymmetric partners have odd R -parity. We can choose the following R -charges Choosing the following R -charges [23, 28]

$$\begin{aligned} n_\eta &= n_{\rho'} = n_S = -1, \quad n_\rho = n_{\eta'} = n_{S'} = 1, \quad n_\chi = n_{\chi'} = 0, \\ n_L &= n_{Q_i} = n_{d_i} = 1/2, \quad n_{J_i} = -1/2, \quad n_u = -3/2, \end{aligned} \quad (21)$$

The terms which are proportional to the following constants: $\mu_0; \lambda_1, \lambda_4, f_{3,4}, f'_{3,4}, \kappa_7, \xi_{1,2,3}$ violate the R -parity defined at Eq.(21).

The terms which satisfy the defined above symmetry (21) the term allowed by this R -charges in our superpotential, given at Eq.(17), are given by

$$W = W_{2RC} + W_{3RC}, \quad (22)$$

where

$$\begin{aligned} W_{2RC} &= \mu_\eta(\hat{\eta}\hat{\eta}') + \mu_\rho(\hat{\rho}\hat{\rho}') + \mu_\chi(\hat{\chi}\hat{\chi}') + \mu_S Tr[(\hat{S}\hat{S}'), \\ W_{3RC} &= \lambda_{2ij}(\epsilon\hat{L}_i\hat{L}_j\hat{\eta}) + \lambda_{3ij}(\hat{L}_i\hat{S}\hat{L}_j) + \lambda_{4i}(\epsilon\hat{L}_i\hat{\chi}\hat{\rho}) + f_1(\epsilon\hat{\rho}\hat{\chi}\hat{\eta}) + f_2(\hat{\chi}\hat{S}\hat{\rho}) \\ &+ f'_1(\epsilon\hat{\rho}'\hat{\chi}'\hat{\eta}') + f'_2(\hat{\chi}'\hat{S}'\hat{\rho}') + \kappa_{1i}(\hat{Q}_3\hat{\eta}')\hat{u}_i^c + \kappa_{2i}(\hat{Q}_3\hat{\rho}')\hat{d}_i^c + \kappa_3(\hat{Q}_3\hat{\chi}')\hat{J}^c \\ &+ \kappa_{4\alpha i}(\hat{Q}_\alpha\hat{\eta})\hat{d}_i^c + \kappa_{5\alpha i}(\hat{Q}_\alpha\hat{\rho})\hat{u}_i^c + \kappa_{6\alpha\beta}(\hat{Q}_\alpha\hat{\chi})\hat{J}_\beta^c. \end{aligned} \quad (23)$$

The soft terms to break SUSY¹ and conserve our R -Parity is written as

$$\begin{aligned} \mathcal{L}_{scalar}^{soft} &= -m_\eta^2(\eta^\dagger\eta) - m_\rho^2(\rho^\dagger\rho) - m_\chi^2(\chi^\dagger\chi) - m_S^2 Tr[(S^\dagger S)] - m_{\eta'}^2(\eta'^\dagger\eta') - m_{\rho'}^2(\rho'^\dagger\rho') \\ &- m_{\chi'}^2(\chi'^\dagger\chi') - m_{S'}^2 Tr[(S'^\dagger S')] + [k_1(\epsilon\rho\chi\eta) + k_2(\chi S\rho) + k'_1(\epsilon\rho'\chi'\eta')] \\ &+ k'_2(\chi'S'\rho') + hc], \end{aligned} \quad (24)$$

¹We are not considering the Gaugino Mass terms neither sleptons and squarks, about the soft terms you can see [23].

$k_{1,2}, k'_1$ and k'_2 has mass dimension.

The pattern of the symmetry breaking of the model is given by the following scheme

$$\begin{aligned} \text{MSUSY331} & \xrightarrow{\mathcal{L}_{soft}} \text{SU}(3)_C \otimes \text{SU}(3)_L \otimes \text{U}(1)_N \xrightarrow{\langle \chi \rangle \langle \chi' \rangle} \text{SU}(3)_C \otimes \text{SU}(2)_L \otimes \text{U}(1)_Y \\ & \xrightarrow{\langle \rho, \eta, S \rho', \eta', S' \rangle} \text{SU}(3)_C \otimes \text{U}(1)_Q. \end{aligned} \quad (25)$$

3 The scalar potential at MSUSY331

The scalar potential is written as

$$V_{331} = V_D + V_F + V_{soft} \quad (26)$$

where

$$\begin{aligned} V_D &= -\mathcal{L}_D, \\ V_F &= -\mathcal{L}_F = \sum_m F_m^\dagger F_m, \\ V_{soft} &= -\mathcal{L}_{soft}^{\text{scalar}}. \end{aligned} \quad (27)$$

How to get an expression to compare our scalar potential with the scalar potential in m331 model see App.(B).

3.1 Numerical Analyses at MSUSY331

We will use below the following set of parameters in the scalar potential:

$$f_1 = 1, \quad f_3 = 1.07, \quad f'_1 = f'_3 = 10^{-6}, \quad (\text{dimensionless}) \quad (28)$$

and

$$-k_1 = k'_1 = 10, \quad k_3 = k'_3 = -100, \quad -\mu_\eta = \mu_\rho = -\mu_s = \mu_\chi = 1000, \quad (\text{in GeV}), (29)$$

we also use the constraint

$$V_\eta^2 + V_\rho^2 + 2V_2^2 = (246 \text{ GeV})^2 \quad (30)$$

coming from M_W , where, we have defined $V_\eta^2 = v_\eta^2 + v_\eta'^2$ and $V_\rho^2 = v_\rho^2 + v_\rho'^2$ and $V_2^2 = v_{\sigma_2}^2 + v_{\sigma_2'}^2$. Assuming that $v_\eta = 20$, $v_\chi = 1500$, $v_{\sigma_2} = 10$, $v'_\eta = v'_\rho = v'_{\sigma_2} = v'_\chi = 1$ in GeV, the value of v_ρ is fixed by the constraint above.

With this set of values for the parameters the real mass eigenstates H_i are (in GeV) $M_{H_i^0} = 125.5, 358.3, 506.3, 1207.4, 1496.5, 1818.1, 4811.5, 5531.4$, $i = 1, \dots, 8$ and $M_{H_j} > M_{H_i}$ with $j > i$.

In the pseudoscalar sector we have verified analytically that the mass matrix, has two Goldstone bosons as it should. The other six physical pseudoscalars have the following masses, with the same parameters as before, in GeV, $M_{A_i} = 277.7, 505.5, 964.6, 1221.1, 3928.5, 4516.43$.

In the Single charged Scalars we have two Goldstone bosons and nine massive states, in GeV. We get two unrelated bases. The first base is defined as

$$\left(\eta_1^-, \rho^-, \eta_1'^-, \rho'^-, h_1^-, h_1'^-\right)^T, \quad (31)$$

we get the following masses $M_{H_i^\pm} = 284.6, 963.1, 1222.3, 3928.7, 4516.1, 4516.1$. The second base is defined as

$$\left(\eta_2^-, \chi^-, \eta_2'^-, \chi'^-, h_2^-, h_2'^-\right)^T, \quad (32)$$

we get the following masses $M_{H_i^\pm} = 372.9, 601.4, 925.7, 3935.3, 4504.5$

In the Double charged Scalars we have one Goldstone bosons and seven massive states, in GeV, $M_{H_i^{\pm\pm}} = 226.5, 596.8, 927.4, 993.5, 1205.1, 184.9, 3922.0, 3935.1$.

We present next some plots to light scalar, first we fix some values for $v_{\chi'}$ and present its mass in term of v_χ , see Fig.(1) and the same for pseudo-scalar see Fig.(3), single charged higgses at first base, see Eq.(31), Fig.(5), second base, see Eq.(32), at Fig.(7) and double charged Higgs Fig.(9) Next we fixed some values for v_χ and it is shown at Fig.(2) and for pseudo-scalar at Fig.(4), single charged higgses at first base, see Eq.(31), Fig.(6), second base, see Eq.(32), at Fig.(8) and double charged Higgs Fig.(10).

4 R -Parity charges at SUSY331RV

In the 3-3-1 model with heavy charged lepton [12] only the triplets are needed for generate masses for all the charged fermions.

However, if we want to allow neutrinos to get their masses and at the same time avoid the fast nucleon decay we can choose the following R -charges

$$\begin{aligned} n_L &= n_\eta = n_{\eta'} = n_\rho = n_{\rho'} = -n_\chi = n_{\chi'} = 0, \\ n_{Q_i} &= 1, \quad n_u = n_{d_i} = n_{J_i} = -1. \end{aligned} \quad (33)$$




Figure 1: The masses of light scalars in terms of v_χ using the parameters defined at Eqs.(28,29).

In this case, the terms allow in our superpotential are

$$\begin{aligned}
W = & \mu_{0i}(\hat{L}_{\alpha L}\hat{\eta}') + \mu_\eta(\hat{\eta}\hat{\eta}') + \mu_\rho(\hat{\rho}\hat{\rho}') + \mu_\chi(\hat{\chi}\hat{\chi}') + \lambda_{1abc}(\epsilon\hat{L}_i\hat{L}_j\hat{L}_k) + \lambda_{2ij}(\epsilon\hat{L}_i\hat{L}_j\hat{\eta}) \\
& + \lambda_{3ij}(\hat{L}_i\hat{S}\hat{L}_j) + \lambda_{4i}(\epsilon\hat{L}_i\hat{\chi}\hat{\rho}) + f_1(\epsilon\hat{\rho}\hat{\chi}\hat{\eta}) + f'_1(\epsilon\hat{\rho}'\hat{\chi}'\hat{\eta}') + \kappa_{1i}(\hat{Q}_3\hat{\eta}')\hat{u}_i^c + \kappa_{2i}(\hat{Q}_3\hat{\rho}')\hat{d}_i^c \\
& + \kappa_3(\hat{Q}_3\hat{\chi}')\hat{J}^c + \kappa_{4\alpha i}(\hat{Q}_\alpha\hat{\eta})\hat{d}_i^c + \kappa_{5\alpha i}(\hat{Q}_\alpha\hat{\rho})\hat{u}_i^c + \kappa_{6\alpha\beta}(\hat{Q}_\alpha\hat{\chi})\hat{j}_\beta^c + \kappa_{7\alpha ij}(\hat{Q}_\alpha\hat{L}_i)\hat{d}_j^c. \quad (34)
\end{aligned}$$

In our superpotential, we can generate mass to neutrinos without introduce the anti-sextet [24].

In the case non supersymmetric we have model with three triplets only if we introduce heavy charged lepton as done at [12].

4.1 The scalar potential at SUSY331RV

In the SUSY331RV there are R -violating interactions give the correct to neutrinos [24].

4.2 Numerical Analyses at SUSY331RV

We will use below the following set of parameters in the scalar potential:

$$f_1 = 1.43, \quad f'_1 = 10^{-6}, \quad (\text{dimensionless}) \quad (35)$$

and

$$-k_1 = k'_1 = 10, \quad -\mu_\eta = \mu_\rho = \mu_\chi = 1000, \quad (\text{in GeV}), \quad (36)$$




Figure 2: The masses of light scalars in terms of $v_{\chi'}$ using the parameters defined at Eqs.(28,29).

we also use the constraint

$$V_{\eta}^2 + V_{\rho}^2 + 2V_2^2 = (246 \text{ GeV})^2 \quad (37)$$

coming from M_W , where, we have defined $V_{\eta}^2 = v_{\eta}^2 + v_{\eta}'^2$ and $V_{\rho}^2 = v_{\rho}^2 + v_{\rho}'^2$. Assuming that $v_{\eta} = 20$, $v_{\chi} = 1500$, $v_{\eta}' = v_{\rho}' = v_{\chi}' = 1$ in GeV, the value of v_{ρ} is fixed by the constraint above.

With this set of values for the parameters the real mass eigenstates H_i are (in GeV) $M_{H_i^0} = 125.5, 304.7, 765.6, 1818.2, 2251.4, 6620.3$, $i = 1, \dots, 6$ and $M_{H_j} > M_{H_i}$ with $j > i$.

In the pseudoscalar sector we have verified analytically that the mass matrix, has two Goldstone bosons as it should. The other six physical pseudoscalars have the following masses, with the same parameters as before, in GeV, $M_{A_i} = 304.4, 765.6, 2251.4, 6620.3$.

In the Single charged Scalars we have two Goldstone bosons and nine massive states, in GeV. We get two unrelated bases. The first base is defined as $\eta_1^-, \rho^-, \eta_1'^-, \rho'^-$ and we get the following masses $M_{H_i^{\pm}} = 306.9, 2252.1, 6620.0$. The second base is defined as $\eta_2^-, \chi^-, \eta_2'^-, \chi'^-$ and we get the following masses $M_{H_i^{\pm}} = 298.8, 908.5, 6602.0$.

In the Double charged Scalars we have one Goldstone bosons and seven massive states, in GeV, $M_{H_i^{\pm\pm}} = 114.6, 906.2, 2195.2$.

We present next some plots to light scalar, first we fix some values for $v_{\chi'}$ and present its mass in term of v_{χ} , see Fig.(11). Next we fixed some values for v_{χ} and it is shown at Fig.(12).




Figure 3: The masses of light pseudo-scalars in terms of v_χ using the parameters defined at Eqs.(28,29).

5 Conclusions

We have studied the scalar potential of supersymmetric 3-3-1 model with the anti-sextet and without it and we show that our results are in agreement with the experimental constraints.

Acknowledgments

We would like to thank V. Pleitez, J. C. Montero and B. L. Schánchez-Vega for useful discussions about 3-3-1 models. We also thank IFT for the nice hospitality during my several visits to perform my studies about the several 3-3-1 Models and also for doing this article.

A Review Scalar Potential at m331

In the m331 the scalar sector is composed for the scalars defined at Eqs.(7,8). We can write the $SU(3)$ multiplets above in terms of the $SU(2)_L$ in the following way

$$\frac{Q^{SM}}{e} = \frac{1}{2}(\sigma_3 + Y I_{2 \times 2}) = \frac{1}{2} \begin{pmatrix} 1+Y & 0 \\ 0 & Y-1 \end{pmatrix}, \quad (38)$$




Figure 4: The masses of light pseudo-scalars in terms of $v_{\chi l}$ using the parameters defined at Eqs.(28,29).

in the Standard Model the Higgs field has hypercharge ($Y=+1$), together with Eq.(38), it is given by

$$\Phi = \begin{pmatrix} h^+ \\ h^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{2}, +1). \quad (39)$$

Now we can decompose our Scalars fields in $SU(2) \otimes U(1)$ representations and the from group theory we can write [25]:

- $\mathbf{3}_0 \rightarrow \mathbf{2}_{-1} \oplus \mathbf{1}_{+2}$

$$\begin{aligned} \eta &= \begin{pmatrix} \Phi_\eta \\ \eta_2^+ \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, 0), \\ \Phi_\eta &= \begin{pmatrix} \eta^0 \\ \eta_1^- \end{pmatrix} \sim (\mathbf{1}, \mathbf{2}, -1), \quad \eta_2^+ \sim (\mathbf{1}, \mathbf{1}, +2), \end{aligned} \quad (40)$$

where the singlet η_2^+ was introduced some years ago by Zee at [26]

- $\mathbf{3}_1 \rightarrow \mathbf{2}_1 \oplus \mathbf{1}_4$

$$\begin{aligned} \rho &= \begin{pmatrix} \Phi_\rho \\ \rho^{++} \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, +1), \\ \Phi_\rho &= \begin{pmatrix} \rho^+ \\ \rho^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{2}, +1), \quad \rho^{++} \sim (\mathbf{1}, \mathbf{1}, +4) \end{aligned} \quad (41)$$




Figure 5: The masses of single scalars at first base, see Eq.(31), in terms of v_χ using the parameters defined at Eqs.(28,29).

- $\mathbf{3}_{-1} \rightarrow \mathbf{2}_{-3} \oplus \mathbf{1}_0$

$$\begin{aligned}\chi &= \begin{pmatrix} \Phi_\chi \\ \chi^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, -1), \\ \Phi_\chi &= \begin{pmatrix} \chi^- \\ \chi^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{2}, -3), \quad \chi^0 \sim (\mathbf{1}, \mathbf{1}, 0)\end{aligned}\quad (42)$$

- $\mathbf{6}_0 \rightarrow \mathbf{3}_2 \oplus \mathbf{2}_{-1} \oplus \mathbf{1}_{-4}$

$$\begin{aligned}S &= \begin{pmatrix} T & \frac{\Phi_S}{\sqrt{2}} \\ \frac{\Phi_S^T}{\sqrt{2}} & H_2^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{6}^*, 0), \\ \Phi_S &= \begin{pmatrix} h_2^+ \\ \sigma_2^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{2}, -1), \quad H_2^{--} \sim (\mathbf{1}, \mathbf{1}, -4),\end{aligned}\quad (43)$$

Φ_S is the Higgs doublet of the Standard Model, see Eq.(39), and the triplet

$$T = \begin{pmatrix} \sigma_1^0 & \frac{h_1^+}{\sqrt{2}} \\ \frac{h_1^+}{\sqrt{2}} & H_1^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, +2),\quad (44)$$

was considered at [27].

Because of the anti-sextet, the scalar potential in the m331 become more complicated. The most general scalar potential involving triplets and the




Figure 6: The masses of single scalars at second base, see Eq.(32), in terms of $v_{\chi'}$ using the parameters defined at Eqs.(28,29).

sextet is [25]

$$V(\eta, \rho, \chi, S) = V^{(2)} + V^{(3)} + V^{(4a)} + V^{(4b)} + V^{(4c)} + V^{(4d)} + V^{(4e)}, \quad (45)$$

where

$$\begin{aligned}
V^{(2)} &= \mu_1^2(\eta^\dagger\eta) + \mu_2^2(\rho^\dagger\rho) + \mu_3^2(\chi^\dagger\chi) + \mu_4^2\text{Tr}[(S^\dagger S)], \\
V^{(3)} &= \frac{f_1}{3!}\epsilon_{ijk}\eta_i\rho_j\chi_k + f_2\rho_i\chi_j S^{\dagger ij} + f_3\eta_i\eta_j S^{\dagger ij} + \frac{f_4}{3!}\epsilon_{ijk}\epsilon_{lmn}S_{il}S_{jm}S_{kn} + H.c., \\
V^{(4a)} &= a_1(\eta^\dagger\eta)^2 + a_2(\rho^\dagger\rho)^2 + a_3(\chi^\dagger\chi)^2 + (\chi^\dagger\chi) [a_4(\eta^\dagger\eta) + a_5(\rho^\dagger\rho)] + a_6(\eta^\dagger\eta)(\rho^\dagger\rho) \\
&\quad + a_7(\chi^\dagger\eta)(\eta^\dagger\chi) + a_8(\chi^\dagger\rho)(\rho^\dagger\chi) + a_9(\eta^\dagger\rho)(\rho^\dagger\eta) + [a_{10}(\chi^\dagger\eta)(\rho^\dagger\eta) + H.c.], \\
V^{(4b)} &= b_1(\chi^\dagger S)(\hat{\chi}\eta) + b_2(\rho^\dagger S)(\hat{\rho}\eta) + b_3(\eta^\dagger S) [(\hat{\chi}\rho) - (\hat{\rho}\chi)] + H.c., \\
V^{(4c)} &= c_1\text{Tr}[(\hat{\eta}S)(\hat{\eta}S)] + c_2\text{Tr}[(\hat{\rho}S)(\hat{\rho}S)] + H.c., \\
V^{(4d)} &= d_1(\chi^\dagger\chi)\text{Tr}[(S^\dagger S)] + d_2\text{Tr}[(\chi^\dagger S)(S^\dagger\chi)] + d_3(\eta^\dagger\eta)\text{Tr}[(S^\dagger S)] + d_4\text{Tr}[(\eta^\dagger S)(S^\dagger\eta)] \\
&\quad + d_5(\rho^\dagger\rho)\text{Tr}[(S^\dagger S)] + d_6\text{Tr}[(\rho^\dagger S)(S^\dagger\rho)], \\
V^{(4e)} &= e_1(\text{Tr}[(S^\dagger S)])^2 + e_2\text{Tr}[(S^\dagger S)(S^\dagger S)], \quad (46)
\end{aligned}$$

and we have defined the $V^{(4b)}$ and $V^{(4c)}$ terms

$$\hat{x}_{ij} \equiv \epsilon_{ijk}x_k, \quad (47)$$

with $x = \eta, \rho, \chi$. The constants f_i , $i = 1, 2, 3$ and 4 have dimension of mass.




Figure 7: The masses of single scalars at second base, see Eq.(32), in terms of v_χ using the parameters defined at Eqs.(28,29).

	η	χ	ρ	S
\mathcal{Z}_7	w^3	w^6	w^5	w^4
\mathcal{Z}_3	w	1	w	w

Table 1: Transformation properties of the scalar fields under $\mathcal{Z}_7 \otimes \mathcal{Z}_3$, taken from [25].

In order to simplify the scalar potential, we impose the following discrete symmetry $\mathcal{Z}_7 \otimes \mathcal{Z}_3$, defined at Tab.(1), which forbids the following coupling $a_{10}, f_3, f_4, b_1, b_2, b_3, c_1$ and c_2 in the scalar potential given at Eq.(46) as presented at [25].

The vevs of our scalars are given at Eqs.(10). We will present in the following sub-sections the main results presented at [17].

We denote the $SU(3)_L$ gauge bosons by V_m^a ($a = 1, 2, \dots, 8$) and since

$$\mathbf{8}_0 \rightarrow \mathbf{3}_0 \oplus \mathbf{2}_3 \oplus \mathbf{2}_{-3} \oplus \mathbf{1}_0, \quad (48)$$

we get a triplet, V_m^1, V_m^2, V_m^3 , has hypercharge 0, plus the doublet of bileptons, (U^{++}, V^+) , its hypercharge is $Y = 3$, and a singlet V_m^8 with $Y = 0$.




Figure 8: he masses of single scalars at base 2 in terms of $v_{\chi'}$ using the parameters defined at Eqs.(28,29).

A.1 CP-Even at MSUSY331

The square mass matrix coming from the base

$$(H_{\eta}, H_{\rho}, H_{\chi}, H_{\sigma_2})^T. \quad (49)$$

The mass spectrum and the eigenstates are get, see our Eq.(75), when we suppose

$$|f_1| \sim w, \quad |f_2| \sim w, \quad (50)$$

we obtain the following squares masases

$$\begin{aligned} m_{H_1^0}^2 &\approx -4\lambda(v^2 + u^2 + z^2), \\ m_{H_2^0, H_3^0}^2 &\approx \frac{w^2}{4vuz} \left\{ (su^2 + rvz)w \pm \left[(su^2 - rvz)^2 w^2 - \frac{4vu^2z}{w^2} [f_1 f_2 u^2 - (f_1^2 + f_2^2)uz] \right]^{(1/2)} \right\}, \\ m_{H_4^0}^2 &\approx -4\lambda_3 w^2, \end{aligned} \quad (51)$$

where we are defining

$$\begin{aligned} r &\equiv \frac{1}{w^2}(f_1 v - wz) \approx -\frac{2}{uw}(\lambda_5 v^2 + \lambda_6 u^2 + 2\lambda_{14} z^2), \\ s &\equiv \frac{1}{w^2}(f_1 z - vw). \end{aligned} \quad (52)$$

This approximation implies

$$\lambda \equiv 4(2\lambda_{10} + \lambda_{11}) \approx \lambda_1 \approx \lambda_2 \approx \lambda_{12} \approx \lambda_{13} \approx \frac{\lambda_4}{2}. \quad (53)$$




Figure 9: The masses of double scalars in terms of v_χ using the parameters defined at Eqs.(28,29).

In this case the Standard Model Higgs Boson correspond to H_1^0 , in the sense that its mass has no dependence in the w parameter.

A.2 CP-Odd at MSUSY331

The square mass matrix coming from the base

$$(F_\eta, F_\rho, F_\chi, F_{\sigma_2})^T, \quad (54)$$

the mass spectrum and the eigenstates are get, see our Eq.(75), to impose

$$f_2 > 0, \quad (55)$$

then the mass is given by:

$$m_{A_1^0, A_2^0}^2 \approx \frac{w^2}{4vuz} \left\{ (su^2 + rvz)w \pm \left[(su^2 - rvz)^2 w^2 - \frac{4vu^2 z}{w^2} [f_1 f_2 u^2 - (f_1^2 + -f_2^2)uz] \right]^{(1/2)} \right\} \quad (56)$$

A.3 Singly Charged at MSUSY331

The square mass matrix coming from the base

$$(\eta_1^+, \rho^+, \eta_2^+, \chi^+, h_1^+, h_2^+)^T. \quad (57)$$




Figure 10: he masses of double scalars in terms of $v_{\chi'}$ using the parameters defined at Eqs.(28,29).

In this approximation two of the masses of the singly charged scalar are given

$$\begin{aligned} m_{H_1^+, H_2^+}^2 &= m_{A_1^0, A_2^0}^2, \\ m_{H_3^+, H_4^+}^2 &\approx f_2 \frac{uw}{2z}. \end{aligned} \quad (58)$$

A.4 Double Charged at MSUSY331

The square mass matrix coming from the base

$$\left(\rho^{++}, \chi^{++}, H_1^+, H_2^{++} \right)^T. \quad (59)$$

The doubly charged scalar masses are

$$\begin{aligned} m_{H_1^{++}, H_2^{++}}^2 &\approx \frac{w^2}{4z} \left\{ u - 2\lambda_9 z \pm \left[(2\lambda_9 z + u)^2 + 4z^2 \right]^{(1/2)} \right\}, \\ m_{H_3^{++}, H_4^{++}}^2 &\approx f_2 \frac{uw}{2z}. \end{aligned} \quad (60)$$

B Construction Scalar Potential

To get the scalar potential of our model we have to eliminate the auxiliary fields F and D that appear in our model. We are going to pick up the F and




Figure 11: The masses of light scalars in terms of v_χ using the parameters defined at Eqs.(35,36).

D - terms we get

$$\begin{aligned}
\mathcal{L}_D^{Gauge} &= \frac{1}{2}D^a D^a + \frac{1}{2}DD \ , \\
\mathcal{L}_F^{Scalar} &= |F_\eta|^2 + |F_\rho|^2 + |F_\chi|^2 + |F_S|^2 + |F_{\eta'}|^2 + |F_{\rho'}|^2 + |F_{\chi'}|^2 + |F_{S'}|^2, \\
\mathcal{L}_D^{Scalar} &= \frac{g}{2} \left[\bar{\eta}\lambda^a\eta + \bar{\rho}\lambda^a\rho + \bar{\chi}\lambda^a\chi + \bar{S}\lambda^a S - \bar{\eta}'\lambda^{*a}\eta' - \bar{\rho}'\lambda^{*a}\rho' - \bar{\chi}'\lambda^{*a}\chi' - \bar{S}'\lambda^{*a}S' \right] D^a \\
&\quad + \frac{g'}{2} [\bar{\rho}\rho - \bar{\chi}\chi - \bar{\rho}'\rho' + \bar{\chi}'\chi'] D, \\
\mathcal{L}_F^{W2} &= \frac{1}{2} \left[\mu_\eta(\eta F_{\eta'} + \eta' F_\eta + \bar{\eta}\bar{F}_{\eta'} + \bar{\eta}'\bar{F}_\eta) + \mu_\rho(\rho F_{\rho'} + \rho' F_\rho + \bar{\rho}\bar{F}_{\rho'} + \bar{\rho}'\bar{F}_\rho) \right. \\
&\quad \left. + \mu_\chi(\chi F_{\chi'} + \chi' F_\chi + \bar{\chi}\bar{F}_{\chi'} + \bar{\chi}'\bar{F}_\chi) \right], \\
\mathcal{L}_F^{W3} &= |F_\eta|^2 + |F_\rho|^2 + |F_\chi|^2 + |F_S|^2 + |F_{\eta'}|^2 + |F_{\rho'}|^2 + |F_{\chi'}|^2 + |F_{S'}|^2.
\end{aligned} \tag{61}$$

From the equation described above we can construct

$$\begin{aligned}
\mathcal{L}_F &= \mathcal{L}_F^{Scalar} + \mathcal{L}_F^{W2} + \mathcal{L}_F^{W3} \\
&= |F_\eta|^2 + |F_\rho|^2 + |F_\chi|^2 + |F_{\eta'}|^2 + |F_{\rho'}|^2 + |F_{\chi'}|^2 \\
&\quad + \frac{1}{2} \left[\mu_\eta(\eta F_{\eta'} + \eta' F_\eta + \bar{\eta}\bar{F}_{\eta'} + \bar{\eta}'\bar{F}_\eta) + \mu_\rho(\rho F_{\rho'} + \rho' F_\rho + \bar{\rho}\bar{F}_{\rho'} + \bar{\rho}'\bar{F}_\rho) \right. \\
&\quad \left. + \mu_\chi(\chi F_{\chi'} + \chi' F_\chi + \bar{\chi}\bar{F}_{\chi'} + \bar{\chi}'\bar{F}_\chi) \right] + \frac{1}{3} [f_1\epsilon(F_\rho\chi\eta + \rho F_\chi\eta + \rho\chi F_\eta
\end{aligned}$$




Figure 12: The masses of light scalars in terms of $v_{\chi'}$ using the parameters defined at Eqs.(35,36).

$$\begin{aligned}
& + \bar{F}_\rho \bar{\chi} \bar{\eta} + \bar{\rho} \bar{F}_\chi \bar{\eta} + \bar{\rho} \bar{\chi} \bar{F}_\eta + f_1 \epsilon (F_{\rho'} \chi' \eta' + \rho' F_{\chi'} \eta' + \rho' \chi' F_{\eta'} \\
& + \bar{F}_{\rho'} \bar{\chi}' \bar{\eta}' + \bar{\rho}' \bar{F}_{\chi'} \bar{\eta}' + \bar{\rho}' \bar{\chi}' \bar{F}_{\eta'}) \quad , \\
\mathcal{L}_D & = \mathcal{L}_D^{Gauge} + \mathcal{L}_D^{Scalar} \\
& = \frac{1}{2} D^a D^a + \frac{1}{2} D D + \frac{g}{2} [\bar{\eta} \lambda^a \eta + \bar{\rho} \lambda^a \rho + \bar{\chi} \lambda^a \chi - \bar{\eta}' \lambda^{*a} \eta' - \bar{\rho}' \lambda^{*a} \rho' \\
& - \bar{\chi}' \lambda^{*a} \chi'] D + \frac{g'}{2} [\bar{\rho} \rho - \bar{\chi} \chi - \bar{\rho}' \rho' + \bar{\chi}' \chi'] D. \tag{62}
\end{aligned}$$

We will now show that these fields can be eliminated through the Euler-Lagrange equations

$$\frac{\partial \mathcal{L}}{\partial \phi} - \partial_m \frac{\partial \mathcal{L}}{\partial (\partial_m \phi)} = 0 \quad , \tag{63}$$

where $\phi = \eta, \rho, \chi, \eta', \rho', \chi'$. Formally auxiliary fields are defined as fields having no kinetic terms. Thus, this definition immediately yields that the Euler-Lagrange equations for auxiliary fields simplify to $\frac{\partial \mathcal{L}}{\partial \phi} = 0$.

$$\begin{aligned}
V_D & = -\mathcal{L}_D = \frac{1}{2} (D^a D^a + D D) \\
& = \frac{g^2}{2} (\rho^\dagger \rho - \rho'^\dagger \rho' - \chi^\dagger \chi + \chi'^\dagger \chi')^2 + \frac{g^2}{8} \sum_{i,j} \left(\eta_i^\dagger \lambda_{ij}^a \eta_j + \rho_i^\dagger \lambda_{ij}^a \rho_j + \chi_i^\dagger \lambda_{ij}^a \chi_j + S_{ij}^\dagger \lambda_{jk}^a S_{kl} \right. \\
& \left. - \eta_i^\dagger \lambda_{ij}^{*a} \eta'_j - \rho_i^\dagger \lambda_{ij}^{*a} \rho'_j - \chi_i^\dagger \lambda_{ij}^{*a} \chi'_j - S_{ij}^\dagger \lambda_{jk}^{*a} S'_{kl} \right)^2, \tag{64}
\end{aligned}$$

We can use the following relations

$$\lambda_{ij}^a \lambda_{kl}^a = \frac{(-2)}{3} \delta_{ij} \delta_{kl} + 2\delta_{il} \delta_{jk}. \quad (65)$$

Then

$$\eta_i^\dagger \eta_j \eta_k^\dagger \eta_l \lambda_{ij}^a \lambda_{kl}^a = -\frac{2}{3} (\eta^\dagger \eta) (\eta^\dagger \eta) + 2 (\eta^\dagger \eta) (\eta^\dagger \eta) = \frac{4}{3} (\eta^\dagger \eta)^2, \quad (66)$$

in similar way

$$\begin{aligned} \rho_i^\dagger \rho_j \rho_k^\dagger \rho_l \lambda_{ij}^a \lambda_{kl}^a &= \frac{4}{3} (\rho^\dagger \rho)^2, \\ \chi_i^\dagger \chi_j \chi_k^\dagger \chi_l \lambda_{ij}^a \lambda_{kl}^a &= \frac{4}{3} (\chi^\dagger \chi)^2, \\ S_{il}^\dagger S_{mj} S_{kn}^\dagger S_{op} \lambda_{lm}^a \lambda_{no}^a &= -\frac{2}{3} \text{Tr} [(S^\dagger S) (S^\dagger S)] + 2 \text{Tr} [(S^\dagger S)(S^\dagger S)]. \end{aligned} \quad (67)$$

By another hand

$$\begin{aligned} \eta_i^\dagger \eta_j \rho_k^\dagger \rho_l \lambda_{ij}^a \lambda_{kl}^a &= -\frac{2}{3} (\eta^\dagger \eta) (\rho^\dagger \rho) + 2 (\rho^\dagger \eta) (\eta^\dagger \rho), \\ \eta_i^\dagger \eta_j \chi_k^\dagger \chi_l \lambda_{ij}^a \lambda_{kl}^a &= -\frac{2}{3} (\eta^\dagger \eta) (\chi^\dagger \chi) + 2 (\chi^\dagger \eta) (\eta^\dagger \chi), \\ \rho_i^\dagger \rho_j \chi_k^\dagger \chi_l \lambda_{ij}^a \lambda_{kl}^a &= -\frac{2}{3} (\rho^\dagger \rho) (\chi^\dagger \chi) + 2 (\chi^\dagger \rho) (\rho^\dagger \chi), \end{aligned} \quad (68)$$

We can rewrite V_D in the following way

$$\begin{aligned} V_D &= \frac{g'^2}{2} [(\rho^\dagger \rho)^2 + (\chi^\dagger \chi)^2 - 2(\rho^\dagger \rho)(\chi^\dagger \chi) + \dots] \\ &+ \frac{g^2}{8} \left\{ \frac{(-4)}{3} [(\eta^\dagger \eta)^2 + (\rho^\dagger \rho)^2 + (\chi^\dagger \chi)^2 + (\eta^\dagger \eta)(\rho^\dagger \rho) + (\eta^\dagger \eta)(\chi^\dagger \chi) \right. \\ &+ (\eta^\dagger \eta) \text{Tr}[(S^\dagger S)] + (\rho^\dagger \rho)(\chi^\dagger \chi) + (\rho^\dagger \rho) \text{Tr}[(S^\dagger S)] + (\chi^\dagger \chi) \text{Tr}[(S^\dagger S)] \\ &+ 4 [(\eta^\dagger \rho)(\rho^\dagger \eta) + (\eta^\dagger \chi)(\chi^\dagger \eta) + \text{Tr}[(S^\dagger \eta)(\eta^\dagger S)] + (\rho^\dagger \chi)(\chi^\dagger \rho) \\ &+ \text{Tr}[(S^\dagger \rho)(\rho^\dagger S)] + \text{Tr}[(S^\dagger \chi)(\chi^\dagger S)]] - \frac{2}{3} (\text{Tr}[(S^\dagger S)])^2 \\ &\left. + 2 \text{Tr}[(S^\dagger S)(S^\dagger S)] + \dots \right\}. \end{aligned} \quad (69)$$

where ... include the new fields due the SUSY algebra. As we want to compare our potential with the m331 we omitt those terms.

$$\begin{aligned}
V_F &= -\mathcal{L}_F = \sum_m F_m^* F_m \\
&= \sum_{i,j,k} \left[\left| \frac{\mu_\eta}{2} \eta'_i + \frac{f_1}{3} \epsilon_{ijk} \rho_j \chi_k + \frac{2f_2}{3} \eta_i S_{ij} \right|^2 + \left| \frac{\mu_\rho}{2} \rho'_i + \frac{f_1}{3} \epsilon_{ijk} \chi_j \eta_k + \frac{f_3}{3} \chi_i S_{ij} \right|^2 \right. \\
&+ \left| \frac{\mu_\chi}{2} \chi'_i + \frac{f_1}{3} \epsilon_{ijk} \rho_j \eta_k + \frac{f_3}{3} \rho_i S_{ij} \right|^2 + \left| \frac{\mu_s}{2} S'_{ij} + \frac{f_2}{3} \eta_i \eta_j + \frac{f_3}{3} \chi_i \rho_j \right|^2 \\
&+ \left| \frac{\mu_\eta}{2} \eta_i + \frac{f'_1}{3} \epsilon_{ijk} \rho'_j \chi'_k + \frac{2f'_2}{3} \eta'_i S'_{ij} \right|^2 + \left| \frac{\mu_\rho}{2} \rho_i + \frac{f'_1}{3} \epsilon_{ijk} \chi'_j \eta'_k + \frac{f'_3}{3} \chi'_i S'_{ij} \right|^2 \\
&+ \left. \left| \frac{\mu_\chi}{2} \chi_i + \frac{f'_1}{3} \epsilon_{ijk} \rho'_j \eta'_k + \frac{f'_3}{3} \rho'_i S'_{ij} \right|^2 + \left| \frac{\mu_s}{2} S_{ij} + \frac{f'_2}{3} \eta'_i \eta'_j + \frac{f'_3}{3} \chi'_i \rho'_j \right|^2 \right]
\end{aligned} \tag{70}$$

We can use the following relation

$$\epsilon_{ijk} \epsilon_{kmn} = \delta_{im} \delta_{jn} - \delta_{in} \delta_{jm}. \tag{71}$$

In both case $f_3 = 0 = f'_3$, then we get

$$\begin{aligned}
\left| \frac{\mu_\eta}{2} \eta_i + \dots \right|^2 &= \frac{\mu_\eta}{4} (\eta^\dagger \eta), \quad \left| \frac{f_1}{3} \epsilon_{ijk} \rho_j \chi_k + \dots \right|^2 = \frac{f_1^2}{9} [(\rho^\dagger \rho)(\chi^\dagger \chi) - (\rho^\dagger \chi)(\chi^\dagger \rho)], \\
\left| \frac{\mu_\rho}{2} \rho_i + \dots \right|^2 &= \frac{\mu_\rho^2}{4} (\rho^\dagger \rho), \quad \left| \frac{\mu_\chi}{2} \chi_i + \dots \right|^2 = \frac{\mu_\chi^2}{4} (\chi^\dagger \chi), \\
\left| \frac{f_1}{3} \epsilon_{ijk} \chi_j \eta_k + \frac{f_3}{3} \chi_i S_{ij} + \dots \right|^2 &= \frac{f_1^2}{9} [(\chi^\dagger \chi)(\eta^\dagger \eta) - (\chi^\dagger \eta)(\eta^\dagger \chi)] + \frac{f_1 f_2}{9} [\epsilon(\chi^\dagger \eta^\dagger)(\chi S) + H.c.] \\
&+ \frac{f_2^2}{9} \text{Tr} [(S^\dagger \chi)(\chi^\dagger S)], \\
\left| \frac{f_1}{3} \epsilon_{ijk} \rho_j \eta_k + \frac{f_3}{3} \rho_i S_{ij} + \dots \right|^2 &= \frac{f_1^2}{9} [(\rho^\dagger \rho)(\eta^\dagger \eta) - (\rho^\dagger \eta)(\eta^\dagger \rho)] + \frac{f_1 f_2}{9} [\epsilon(\rho^\dagger \eta^\dagger)(\rho S) + H.c.] \\
&+ \frac{f_2^2}{9} \text{Tr} [(S^\dagger \rho)(\rho^\dagger S)], \\
\left| \frac{\mu_s}{2} S_{ij} + \dots \right|^2 &= \frac{\mu_s^2}{4} (\text{Tr}[(S^\dagger S)])^2, \\
\left| \frac{f_1}{3} \chi_i \rho_j + \dots \right|^2 &= \frac{f_1^2}{9} [(\rho^\dagger \rho)(\chi^\dagger \chi) - (\rho^\dagger \chi)(\chi^\dagger \rho)].
\end{aligned}$$

Therefore

$$\begin{aligned}
V_F &= \frac{\mu_\eta^2}{4}(\eta^\dagger\eta)^2 + \frac{\mu_\rho^2}{4}(\rho^\dagger\rho)^2 + \frac{\mu_\chi^2}{4}(\chi^\dagger\chi)^2 + \frac{\mu_s^2}{4}(Tr[(S^\dagger S)])^2 \\
&+ \frac{f_1^2}{9} [(\eta^\dagger\eta)(\rho^\dagger\rho) - (\rho^\dagger\eta)(\eta^\dagger\rho) + (\eta^\dagger\eta)(\chi^\dagger\chi) \\
&- (\chi^\dagger\eta)(\eta^\dagger\chi) + (\rho^\dagger\rho)(\chi^\dagger\chi) - (\rho^\dagger\chi)(\chi^\dagger\rho)] + \frac{f_2^2}{9} \{Tr[(S^\dagger\rho)(\rho^\dagger S)] + Tr[(S^\dagger\chi)(\chi^\dagger S)]\} \\
&+ \left\{ \frac{f_1 f_2}{9} [\epsilon(\rho^\dagger\eta^\dagger)(\rho S) + \epsilon(\chi^\dagger\eta^\dagger)(\chi S)] + H.c. \right\} + \dots, \tag{73}
\end{aligned}$$

The scalar potential in this model is

$$\begin{aligned}
V_{scalar} &= m_\eta^2(\eta^\dagger\eta) + m_\rho^2(\rho^\dagger\rho) + m_\chi^2(\chi^\dagger\chi) + m_S^2 Tr[(S^\dagger S)] - [k_1\epsilon\rho\chi\eta + k_2\chi\rho S^\dagger + H.c.] \\
&+ \left(\frac{\mu_\eta^2}{4} - \frac{g^2}{6}\right)(\eta^\dagger\eta)^2 + \left(\frac{\mu_\rho^2}{4} + \frac{g^2}{2} - \frac{g^2}{6}\right)(\rho^\dagger\rho)^2 + \left(\frac{\mu_\chi^2}{4} + \frac{g^2}{2} - \frac{g^2}{6}\right)(\chi^\dagger\chi)^2 \\
&+ \left(\frac{f_1^2}{9} - \frac{g^2}{6}\right)(\eta^\dagger\eta)(\rho^\dagger\rho) + \left(\frac{f_1^2}{9} - \frac{g^2}{6}\right)(\eta^\dagger\eta)(\chi^\dagger\chi) + \left(\frac{f_1^2}{9} - g^2 - \frac{g^2}{6}\right)(\rho^\dagger\rho)(\chi^\dagger\chi) \\
&+ \left(\frac{g^2}{2} - \frac{f_1^2}{9}\right)(\eta^\dagger\rho)(\rho^\dagger\eta) + \left(\frac{g^2}{2} - \frac{f_1^2}{9}\right)(\eta^\dagger\chi)(\chi^\dagger\eta) + \left(\frac{g^2}{2} - \frac{f_1^2}{9}\right)(\rho^\dagger\chi)(\chi^\dagger\rho) \\
&- \frac{g^2}{6} [(\eta^\dagger\eta)Tr[(S^\dagger S)] + (\rho^\dagger\rho)Tr[(S^\dagger S)] + (\chi^\dagger\chi)Tr[(S^\dagger S)]] \\
&+ \frac{g^2}{2} Tr[(S^\dagger\eta)(\eta^\dagger S)] + \left(\frac{g^2}{2} + \frac{f_2^2}{9}\right) \{Tr[(S^\dagger\rho)(\rho^\dagger S)] + Tr[(S^\dagger\chi)(\chi^\dagger S)]\} \\
&+ \left(\frac{\mu_s^2}{4} - \frac{g^2}{2}\right) (Tr[(S^\dagger S)])^2 + g^2 Tr[(S^\dagger S)(S^\dagger S)] \\
&+ \left\{ \frac{f_1 f_2}{9} [\epsilon(\rho^\dagger\eta^\dagger)(\rho S) + \epsilon(\chi^\dagger\eta^\dagger)(\chi S)] + H.c. \right\} + \dots \tag{74}
\end{aligned}$$

and comparing it with Eq.(46) we can see

$$\begin{aligned}
\mu_1^2 &= m_\eta^2, & \mu_2^2 &= m_\rho^2, & \mu_3^2 &= m_\chi^2, & \mu_4^2 &= m_S^2, \\
f_1 &= -3!k_1, & f_2 &= -3!k_2, \\
a_1 &= \frac{\mu_\eta^2}{4} - \frac{g^2}{6}, & e_1 &= \frac{\mu_s^2}{4} - \frac{g^2}{2}, & e_2 &= g^2,
\end{aligned}$$

$$\begin{aligned}
a_2 &= \frac{\mu_\rho^2}{4} + \frac{g'^2}{2} - \frac{g^2}{6}, a_3 = \frac{\mu_\chi^2}{4} + \frac{g'^2}{2} - \frac{g^2}{6}, \\
a_4 &= -\frac{g^2}{6} + \frac{f_1^2}{9} = a_6 = a_7 = a_8 = a_9, \\
a_5 &= -g'^2 - \frac{g^2}{6} + \frac{f_1^2}{9}, \\
d_1 &= -\frac{g^2}{6} = d_3 = d_4 = d_5, \\
d_2 &= d_4 = d_6 = \frac{g^2}{2}, \\
a_{10} &= 0.
\end{aligned} \tag{75}$$

We get analytical two Goldstone bosons at CP-odd sector, two Goldstone bosons at Singly charged sector and one Goldstone boson at Doubly charged sector. We will present our numerical results in Sec.(3.1).

References

- [1] M. Singer, J. W. F. Valle, and J. Schechter, *Canonical neutral-current predictions from the weak-electromagnetic gauge group $SU(3) \otimes U(1)$* , *Phys. Rev. D* **22**, 738, (1980).
- [2] F. Pisano and V. Pleitez, *An $SU(3) \otimes U(1)$ model for electroweak interactions*, *Phys. Rev. D* **46**, 410, (1992); R. Foot, O. F. Hernandez, F. Pisano and V. Pleitez, *Lepton masses in an $SU(3)_L \otimes U(1)_N$ gauge model*, *Phys. Rev. D* **47**, 4158, (1993).
- [3] R. Foot, H.N. Long and Tuan A. Tran, *$SU(3)_L \otimes U(1)_N$ and $SU(4)_L \otimes U(1)_N$ gauge models with right-handed neutrinos*, *Phys. Rev. D* **50**, 34, (1994); J. C. Montero, F. Pisano and V. Pleitez, *Phys. Rev. D* **47**, 2918, (1993); H.N. Long, *$SU(3)_L \otimes U(1)_N$ model for right-handed neutrino neutral currents*, *Phys. Rev. D* **54**, 4691, (1996); H.N. Long, *$SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ model with right-handed neutrinos*, *Phys. Rev. D* **53**, 437, (1996).
- [4] V. Pleitez, *New fermions and a vector - like third generation in $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ models*, *Phys. Rev. D* **53**, 514, (1996).

- [5] T. V. Duong and E. Ma, *Supersymmetric $SU(3) \otimes U(1)$ Gauge Model: Higgs Structure at the Electroweak Energy Scale*, *Phys. Lett.***B316**, 307 (1993).
- [6] H. N. Long and P. B. Pal, *Nucleon instability in a supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)$ model*, *Mod. Phys. Lett.***A13**, 2355, (1998).
- [7] J. C. Montero, V. Pleitez and M. C. Rodriguez, *A Supersymmetric 3-3-1 model*, *Phys. Rev.* **D65**, 035006, (2002).
- [8] M. Capdequi-Peyranère and M.C. Rodriguez, *Charginos and neutralinos production at 3-3-1 supersymmetric model in e^-e^- scattering*, *Phys. Rev.* **D 65**, 035001 (2002); M. C. Rodriguez, *Double Chargino Production in e^-e^- scattering*, *Int. J. Mod. Phys.* **A 22**, 6080, (2008).
- [9] J. C. Montero, V. Pleitez and M. C. Rodriguez, *Supersymmetric 3-3-1 model with right-handed neutrinos*, *Phys. Rev.* **D 70**, 075004, (2004).
- [10] D. T. Huong, M. C. Rodriguez and H. N. Long, *Scalar sector of supersymmetric $SU(3)_C \otimes U(1)_N$ model with right-handed neutrinos*, arXiv:hep-ph/0508045.
- [11] S. J. L. Rosner, *Resource letter: The Standard model and beyond*, *Am. J. Phys.* **71**, 302, (2003); A. S. Kronfeld and C. Quigg, *Resource Letter: Quantum Chromodynamics*, *Am. J. Phys.***78**, 1081, (2010).
- [12] V. Pleitez and M. D. Tonasse, *Heavy charged leptons in an $SU(3)_L \otimes U(1)_N$ model*, *Phys. Rev.* **D 48**, 2353, (1993).
- [13] J. C. Montero, V. Pleitez and O. Ravinez, *Soft superweak CP violation in a 331 model*, *Phys. Rev.***D 60**, 076003, (1999).
- [14] J. C. Montero, C. C. Nishi, V. Pleitez, O. Ravinez and M. C. Rodriguez, *Soft CP violation in K meson systems*, *Phys. Rev.***D73**, 016003, (2006).
- [15] M. C. Rodriguez, *Scalar sector in the minimal supersymmetric 3-3-1 model*, *Int. J. Mod. Phys.***A21**, 4303, (2006).
- [16] V. Pleitez and M. D. Tonasse, *Neutrinoless double beta decay in an $SU(3)_L \times U(1)_N$ model*, *Phys. Rev.***D48**, 5274, (1993), [arXiv:hep-ph/9302201 [hep-ph]].

- [17] M. D. Tonasse, *The scalar sector of $3 - 3 - 1$ models*, *Phys. Lett.***B381**, 191, (1996).
- [18] R. A. Diaz, R. Martinez and F. Ochoa, *The Scalar sector of the $SU(3)_L \otimes U(1)_X$ model*, *Phys. Rev.***D69**, 095009, (2004), [arXiv:hep-ph/0309280 [hep-ph]].
- [19] T. A. Nguyen, N. A. Ky and H. N. Long, *The Higgs sector of the minimal $3 - 3 - 1$ model revisited*, *Int. J. Mod. Phys.***A15**, 283, (2000), [arXiv:hep-ph/9810273 [hep-ph]].
- [20] Y. Giraldo and W. A. Ponce, *Scalar Potential Without Cubic Term in $3-3-1$ Models Without Exotic Electric Charges*, *Eur. Phys. J.***C71**, 1693 (2011), [arXiv:1107.3260 [hep-ph]].
- [21] A. E. Cárcamo Hernández, R. Martinez and J. Nisperuza, *S_3 discrete group as a source of the quark mass and mixing pattern in 331 models*, *Eur. Phys. J.***C75**, no.2, 72, (2015), [arXiv:1401.0937 [hep-ph]].
- [22] G. Aad *et al.* [ATLAS and CMS Collaborations], *Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments*, *Phys. Rev. Lett.***114**, 191803, (2015), [arXiv:1503.07589 [hep-ex]].
- [23] M. C. Rodriguez, *The Minimal Supersymmetric Standard Model (MSSM) and General Singlet Extensions of the MSSM (GSEMSSM), a short review*, arXiv:1911.13043 [hep-ph].
- [24] J. C. Montero, V. Pleitez and M. C. Rodriguez, *Lepton masses in a supersymmetric $3-3-1$ model*, *Phys. Rev.***D 65**, 095008, (2002).
- [25] G. De Conto, A. C. B. Machado and V. Pleitez, *Minimal $3-3-1$ model with a spectator sextet*, *Phys. Rev.* **D92**, 075031, (2015), [arXiv:1505.01343 [hep-ph]].
- [26] A. Zee, *A Theory of Lepton Number Violation, Neutrino Majorana Mass, and Oscillation*, *Phys. Lett.***B93**, 389, (1980); [erratum: *Phys. Lett.***B95**, 461, (1980)].
- [27] G. B. Gelmini and M. Roncadelli, *Left-Handed Neutrino Mass Scale and Spontaneously Broken Lepton Number*, *Phys. Lett.***B99**, 411 (1981).

- [28] M. C. Rodriguez, *Mass Spectrum in the Minimal Supersymmetric 3-3-1 model*, *J. Mod. Phys.***2**, 1193, (2011).