

# Cyclic inequalities involving Cater cyclic function

JiaJin Wen

*College of Mathematics and Computer Science, Chengdu University, Chengdu, Sichuan, 610106, P. R. China.*

*wenjiajin623@163.com*

TianYong Han

*College of Mathematics and Computer Science, Chengdu University, Chengdu, Sichuan, 610106, P. R. China.*

*hantian123\_123@163.com*

Jun Yuan

*(Corresponding author)*

*School of Information Engineering, Nanjing Xiaozhuang University, Nanjing, Jiangsu, 211171, P. R. China.*

*yuanjun@njxzc.edu.cn*

---

## Abstract

By means of the mathematical analysis theory, inequality theory, mathematical induction and the dimension reduction method, under the proper hypotheses, we establish the following cyclic inequalities:

$$\sum_{i=1}^n a_i^{a_{n+1-i}} \leq \sum_{\text{cyc: } n}^{1 \leq i \leq n} a_i^{a_{i+1}} \leq \sum_{i=1}^n a_i^{a_i}, \quad \forall n \geq 2.$$

*Keywords:* cyclic function, Cater cyclic function, Cater inequality, mean.

*2008 MSC:* 26D15, 26E60

---

## 1. Introduction

We will use the following hypotheses and notations throughout the paper:

$$\mathbb{N} \triangleq \{1, 2, \dots, m, \dots\}, \quad \mathbb{Z} \triangleq \{0, \pm 1, \dots, \pm m, \dots\},$$

$$\mathbb{R} \triangleq (-\infty, \infty), \quad (a_1, a_2, \dots, a_n) \in \mathbb{R}^n, \quad E_n \subseteq \mathbb{R}^n, \quad n \in \mathbb{N}, \quad n \geq 2.$$

If the function  $f : E_n \rightarrow \mathbb{R}$  satisfy the conditions:

$$f(a_{k+1}, a_{k+2}, \dots, a_{k+n}) = f(a_1, a_2, \dots, a_n), \quad \forall k \in \mathbb{Z},$$

where

$$a_k = a_i \Leftrightarrow k \equiv i \pmod{n}, \quad \forall k \in \mathbb{Z}, \quad i = 1, 2, \dots, n, \quad (1.1)$$

then we say that  $f$  is a **cyclic function**.

According to the above definition, we know that, under the hypotheses in (1.1), for any function  $\chi : E_m \rightarrow \mathbb{R}$ ,  $2 \leq m \leq n$ , the function

$$f : E_n \rightarrow \mathbb{R}, f(a_1, a_2, \dots, a_n) \triangleq \sum_{i=1}^n \chi(a_i, a_{i+1}, \dots, a_{i+m-1}) = \sum_{\text{cyc: } n}^{1 \leq i \leq n} \chi(a_i, a_{i+1}, \dots, a_{i+m-1})$$

is a cyclic function, where and in the future,  $\sum_{\text{cyc: } n}^{1 \leq i \leq n}$  represent the **cyclic summation**. For example,

$$\sum_{\text{cyc: } n}^{1 \leq i \leq n} a_i^{a_{i+1}} = \sum_{i=1}^{n-1} a_i^{a_{i+1}} + a_n^{a_1} = \sum_{i=1}^{n-1} a_i^{a_{i+1}} + a_n^{a_1}, \quad \forall n \geq 2.$$

In particular, the function  $f(a_1, a_2, \dots, a_n) \triangleq \sum_{i=1}^n a_i$  is a cyclic function. In general, if  $f : E_n \rightarrow \mathbb{R}$  is a symmetric function, then  $f$  is a cyclic function.

Let  $(a_1, a_2, \dots, a_n) \in (0, \infty)^n$ ,  $n \geq 2$ . Then we say that the function

$$C : (0, \infty)^n \rightarrow \mathbb{R}, C(a_1, a_2, \dots, a_n) \triangleq \sum_{\text{cyc: } n}^{1 \leq i \leq n} a_i^{a_{i+1}}$$

is a **Cater cyclic function** which is a cyclic function, and

$$C_* : (0, \infty)^n \rightarrow \mathbb{R}, C_*(a_1, a_2, \dots, a_n) \triangleq \sum_{i=1}^n a_i^{a_i}$$

with

$$C^* : (0, \infty)^n \rightarrow \mathbb{R}, C^*(a_1, a_2, \dots, a_n) \triangleq \sum_{i=1}^n a_i^{a_{n+1-i}}$$

are **Cater-type cyclic functions** which are also the cyclic functions.

Assume that  $f : E_n \rightarrow \mathbb{R}$  and  $g : E_n \rightarrow \mathbb{R}$  are two cyclic functions. Then we say that the inequalities

$$f(a_1, a_2, \dots, a_n) \geq g(a_1, a_2, \dots, a_n) \quad \text{and} \quad f(a_1, a_2, \dots, a_n) \leq g(a_1, a_2, \dots, a_n)$$

are **cyclic inequalities**.

In 1980, F. S. Cater established an interesting cyclic inequality involving power exponents as follows [1]:

$$C(a_1, a_2, \dots, a_n) > 1 + (n-2) \min \{a_1^{a_2}, a_2^{a_3}, \dots, a_{n-1}^{a_n}, a_n^{a_1}\}, \quad \forall n \geq 2, \quad (1.2)$$

which is called as **Cater inequality**.

Like the periodic function, the cyclic function is also a class of important functions. Since the analytic expression of a cyclic function is very special and complex, so we need to estimate the bounds of the cyclic function or establish a cyclic inequality.

Cater inequality is a cyclic inequality, which is a typical representative of the cyclic inequalities. Since the analytic expression of the Cater cyclic function is of extreme particularity and complexity, it is an interesting topic that to establish a new Cater-type cyclic inequality.

In this paper, we will study the lower with upper bounds of the Cater cyclic function and establish two Cater-type cyclic inequalities, and display the applications of the dimension reduction method [2, 3, 5, 7] in the inequality theory.

The research methods of this paper are based on the mathematical induction and the dimension reduction method. The research tools of this paper include mathematical analysis theory, inequalities theory and the mean theory [2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18].

Our main result are the following Theorems 1.1 and 1.2.

**Theorem 1.1.** *Let  $(a_1, a_2, \dots, a_n) \in (0, \infty)^n$ , where  $n \geq 2$ . If*

$$0 < a_1 \leq a_2 \leq \dots \leq a_n \text{ with } a_1^{a_n} \geq e^{-1}, \quad (1.3)$$

*then we have the following **Cater-type cyclic inequality**:*

$$C(a_1, a_2, \dots, a_n) \geq C^*(a_1, a_2, \dots, a_n). \quad (1.4)$$

*Equality in (1.4) holds if  $n = 2$  or  $a_1 = a_2 = \dots = a_n$ .*

**Theorem 1.2.** *Let  $(a_1, a_2, \dots, a_n) \in (0, \infty)^n$ , where  $n \geq 2$ . If*

$$0 < a_1 \leq a_2 \leq \dots \leq a_n, \quad (1.5)$$

*then we have the following **Cater-type cyclic inequality**:*

$$C(a_1, a_2, \dots, a_n) \leq C_*(a_1, a_2, \dots, a_n). \quad (1.6)$$

*Equality in (1.6) holds if and only if  $a_1 = a_2 = \dots = a_n$ .*

## 2. Proof of Theorem 1.1

*Proof.* Let  $j_1 j_2 \dots j_n$  be a permutation [19] of  $1, 2, \dots, n$ , and let

$$1 \leq i < k \leq n \text{ and } 1 \leq j_i < j_k \leq n. \quad (2.1)$$

By (1.3) and (2.1), we have

$$0 < a_1 \leq a_i \leq a_k \leq a_n \text{ and } 0 < a_1 \leq a_{j_i} \leq a_{j_k} \leq a_n. \quad (2.2)$$

We first prove that

$$a_i^{a_{j_i}} + a_k^{a_{j_k}} \geq a_i^{a_{j_k}} + a_k^{a_{j_i}}. \quad (2.3)$$

Indeed, if  $a_i = a_k$  or  $a_{j_i} = a_{j_k}$ , then (2.3) is an equation. Now we assume that

$$0 < a_1 \leq a_i < a_k \leq a_n \text{ and } 0 < a_1 \leq a_{j_i} < a_{j_k} \leq a_n. \quad (2.4)$$

We define a auxiliary function as follows:

$$f(t) \triangleq t^{a_{j_i}} - t^{a_{j_k}}, \quad 0 < a_i \leq t \leq a_k.$$

Then inequality (2.3) can be rewritten as

$$f(a_i) \geq f(a_k). \quad (2.5)$$

Since

$$\frac{df(t)}{dt} < 0, \quad \forall t \in [a_i, a_k] \Rightarrow f(a_i) \geq f(a_k),$$

so we just need to prove that

$$\frac{df(t)}{dt} < 0, \quad \forall t \in [a_i, a_k]. \quad (2.6)$$

Since  $t \geq a_i \geq a_1 > 0$ ,

$$\begin{aligned} \frac{df(t)}{dt} &= a_{j_i} t^{a_{j_i}-1} - a_{j_k} t^{a_{j_k}-1} \\ &= a_{j_k} t^{a_{j_i}-1} \left( \frac{a_{j_i}}{a_{j_k}} - t^{a_{j_k}-a_{j_i}} \right) \\ &\leq a_{j_k} t^{a_{j_i}-1} \left( \frac{a_{j_i}}{a_{j_k}} - a_i^{a_{j_k}-a_{j_i}} \right) \\ &\leq a_{j_k} t^{a_{j_i}-1} \left( \frac{a_{j_i}}{a_{j_k}} - a_1^{a_{j_k}-a_{j_i}} \right), \end{aligned}$$

$$\frac{a_{j_i}}{a_{j_k}} - a_1^{a_{j_k}-a_{j_i}} < 0 \Rightarrow (2.6),$$

and

$$\frac{a_{j_i}}{a_{j_k}} - a_1^{a_{j_k}-a_{j_i}} < 0 \Leftrightarrow \log a_1 > -\frac{\log a_{j_k} - \log a_{j_i}}{a_{j_k} - a_{j_i}},$$

so we just need to prove that

$$\log a_1 > -\frac{\log a_{j_k} - \log a_{j_i}}{a_{j_k} - a_{j_i}}. \quad (2.7)$$

By (1.3), we have

$$a_1^{a_n} \geq e^{-1} \Leftrightarrow \log a_1 \geq -\frac{1}{a_n}. \quad (2.8)$$

According to (2.4), (2.8) and the Lagrange mean value theorem, there exist  $\zeta \in (a_{j_i}, a_{j_k})$  such that

$$-\frac{\log a_{j_k} - \log a_{j_i}}{a_{j_k} - a_{j_i}} = -\frac{d \log \zeta}{d\zeta} = -\frac{1}{\zeta} < -\frac{1}{a_{j_k}} \leq -\frac{1}{a_n} \leq \log a_1 \Rightarrow (2.7) \Rightarrow (2.6).$$

Hence inequality (2.3) is proved.

Based on the above proof, we know that equality in (2.3) holds if and only if  $a_i = a_k$  or  $a_{j_i} = a_{j_k}$ .

Next, we prove inequality (1.4) as follows.

We define a auxiliary function as follows:

$$F(j_1 j_2 \cdots j_n) \triangleq \sum_{i=1}^n a_i^{a_{j_i}}, \quad (2.9)$$

where  $j_1 j_2 \cdots j_n$  is a permutation of  $1, 2, \dots, n$ . Now we prove that

$$F(j_1 j_2 \cdots j_n) \geq F(n(n-1) \cdots (n-m) j_{m+2}^* j_{m+3}^* \cdots j_n^*), \quad \forall m : 0 \leq m \leq n-1, \quad (2.10)$$

where  $j_{m+2}^* j_{m+3}^* \cdots j_n^*$  is a permutation of  $1, 2, \dots, n-m-1$ .

We use the mathematical induction for  $m$ .

(I) Let  $m = 0$ . If  $j_1 = n$ , then (2.10) is an equality. Assume that  $j_1 < n$ . Then exists  $k : 2 \leq k \leq n$ , such that  $j_k = n$ . Since

$$F(j_1 j_2 \cdots j_k \cdots j_n) = a_1^{a_{j_1}} + a_k^{a_{j_k}} + \sum_{1 \leq i \leq n, i \neq j_1, j_k} a_i^{a_{j_i}},$$

by inequality (2.3), we have

$$a_1^{a_{j_1}} + a_k^{a_{j_k}} \geq a_1^{a_{j_k}} + a_k^{a_{j_1}} = a_1^{a_n} + a_k^{a_{j_1}},$$

that is,

$$F(j_1 j_2 \cdots j_n) = F(j_1 j_2 \cdots j_k \cdots j_n) \geq F(j_k j_2 \cdots j_1 \cdots j_n) = F(n j_2^* j_3^* \cdots j_n^*),$$

where  $j_2^* j_3^* \cdots j_n^* = j_2 \cdots j_1 \cdots j_n$  is a permutation of  $1, 2, \dots, n-1$ . Hence inequality (2.10) holds when  $m = 0$ .

(II) Suppose that inequality (2.10) holds, where  $0 \leq m \leq n-2$ . Now we prove that

$$F(j_1 j_2 \cdots j_n) \geq F(n(n-1) \cdots (n-m-1) j_{m+3}^{**} j_{m+4}^{**} \cdots j_n^{**}), \quad (2.11)$$

where  $j_{m+3}^{**} j_{m+4}^{**} \cdots j_n^{**}$  is a permutation of  $1, 2, \dots, n-m-2$ .

We first prove that

$$F(n(n-1) \cdots (n-m) j_{m+2}^* j_{m+3}^* \cdots j_n^*) \geq F(n \cdots (n-m-1) j_{m+3}^{**} j_{m+4}^{**} \cdots j_n^{**}), \quad (2.12)$$

where  $j_{m+3}^{**} j_{m+4}^{**} \cdots j_n^{**}$  is a permutation of  $1, 2, \dots, n-m-2$ .

Indeed, if  $j_{m+2}^* = n-m-1$ , then (2.12) is an equality. Assume that  $j_{m+2}^* < n-m-1$ . Then there exists  $k : m+3 \leq k \leq n$ , such that  $j_k^* = n-m-1$ . By inequality (2.3) and the proof of (I), we have

$$\begin{aligned} & F(n(n-1) \cdots (n-m) j_{m+2}^* j_{m+3}^* \cdots j_n^*) \\ &= F(n(n-1) \cdots (n-m) j_{m+2}^* j_{m+3}^* \cdots j_k^* \cdots j_n^*) \\ &\geq F(n(n-1) \cdots (n-m) j_k^* j_{m+3}^* \cdots j_{m+2}^* \cdots j_n^*) \\ &= F(n(n-1) \cdots (n-m) (n-m-1) j_{m+3}^* \cdots j_{m+2}^* \cdots j_n^*) \end{aligned}$$

$$= F(n(n-1)\cdots(n-m)(n-m-1)j_{m+3}^{**}j_{m+4}^{**}\cdots j_n^{**}),$$

where  $j_{m+3}^{**}j_{m+4}^{**}\cdots j_n^{**} = j_{m+3}^* \cdots j_{m+2}^* \cdots j_n^*$  is a permutation of  $1, 2, \dots, n-m-2$ . Thus, inequality (2.12) is proved.

By inequalities (2.10) and (2.12), we get the inequality (2.11). This ends the proof.

In inequality (2.10), set  $m = n-1$ , we get

$$F(j_1 j_2 \cdots j_n) \geq F(n(n-1)\cdots 321),$$

that is,

$$\sum_{i=1}^n a_i^{a_{j_i}} \geq \sum_{i=1}^n a_i^{a_{n+1-i}} = C^*(a_1, a_2, \dots, a_n). \quad (2.13)$$

Equality in (2.13) holds if  $j_1 j_2 \cdots j_n = n(n-1)\cdots 321$  or  $a_1 = a_2 = \cdots = a_n$ .

In inequality (2.13), set

$$j_1 j_2 \cdots j_n = 23 \cdots (n-1)n1.$$

Then inequality (2.13) can be rewritten as inequality (1.4). Hence inequality (1.4) is proved.

Based on the above proof, we know that equality in (1.4) holds if  $n = 2$  or  $a_1 = a_2 = \cdots = a_n$ .

The proof of Theorem 1.1 is completed.  $\square$

**Remark 2.1.** Let  $n \geq 2$ ,  $1 \leq a_1 \leq a_2 \leq \cdots \leq a_n$  or  $e^{-1} \leq a_1 \leq a_2 \leq \cdots \leq a_n \leq 1$ . Then (1.3) holds. According to Theorem 1.1, inequality (1.4) holds.

**Remark 2.2.** Let  $a_i = \varepsilon + (i-1)/n$ ,  $1 \leq i \leq n$ ,  $n \geq 2$ , where  $\varepsilon = 0.5173446105249118 \cdots$  is the root of the equation  $x^{x+1} = e^{-1}$ , where  $x \in (0, 1)$ . Then  $0 < a_1 \leq a_2 \leq \cdots \leq a_n$  and

$$a_1^{a_n} = \varepsilon^{\varepsilon+1-n^{-1}} > \varepsilon^{\varepsilon+1} = e^{-1}.$$

According to Theorem 1.1, inequality (1.4) holds. The calculation of  $\varepsilon$  is based on the **Mathematica** software. The relevant literatures on proving inequalities by means of the mathematical software can be see [3, 6, 10, 12].

**Remark 2.3.** Based on the proof of Theorem 1.1 we know that: Under the hypotheses of Theorem 1.1, we have

$$C^*(a_1, a_2, \dots, a_n) = \sum_{i=1}^n a_i^{a_{n+1-i}} \leq \sum_{i=1}^n a_i^{a_{j_i}} \leq \sum_{i=1}^n a_i^{a_i} = C_*(a_1, a_2, \dots, a_n), \quad (2.14)$$

where  $j_1 j_2 \cdots j_n$  is a permutation of  $1, 2, \dots, n$ .

**Remark 2.4.** For the Cater-type cyclic function  $C^*(a_1, a_2, \dots, a_n)$ , we have

$$C^*(a_1, a_2, \dots, a_n) > \frac{n}{2}, \quad \forall n \geq 2, \quad (2.15)$$

$$\inf \{C^*(a_1, a_2, \dots, a_{2m})\} = C^*(0I_m, I_m) = m \quad (2.16)$$

and

$$\inf \{C^*(a_1, a_2, \dots, a_{2m+1})\} = C^*(0I_m, e^{-1}, I_m) = m + e^{-e^{-1}}, \quad (2.17)$$

where  $m \in \mathbb{N}$ ,  $I_m = (1, 1, \dots, 1) \in \mathbb{R}^m$ , and  $e^{-e^{-1}} = 0.6922006275553464 \dots > 0.5$ .

Indeed, in inequality (1.2), set  $n = 2$ , we get

$$C(a_1, a_2) = a_1^{a_2} + a_2^{a_1} > 1, \quad \forall a_1 > 0, \quad \forall a_2 > 0.$$

Hence

$$C^*(a_1, a_2, \dots, a_n) = \frac{1}{2} \sum_{i=1}^n C(a_i, a_{n+1-i}) > \frac{1}{2} \sum_{i=1}^n 1 = \frac{n}{2}. \quad (2.18)$$

Since

$$C(0, 1) = C(1, 0) = 1, \quad \inf_{t>0} \{C(t, t)\} = 2 \inf_{t>0} \{t^t\} = C(e^{-1}, e^{-1}) = 2e^{-e^{-1}},$$

by (2.18), we get (2.16) and (2.17).

### 3. Proof of Theorem 1.2

In order to prove Theorem 1.2, we need to establish several lemmas as follows.

**Lemma 3.1.** *Let  $(a_1, \dots, a_2) \in (0, \infty)^n$  and  $n = 2$ . Then inequality (1.6) holds. Equality in (1.6) holds if and only if  $a_1 = a_2$ .*

*Proof.* If  $a_1 = a_2$ , then  $C(a_1, a_2) = C_*(a_1, a_2)$ . Assume that  $a_1 \neq a_2$ . Without losing of generality, we may assume that  $a = a_1 > a_2 = b > 0$ .

If  $a \geq 1$ , then

$$\begin{aligned} a^{a-b} > b^{a-b} &\Rightarrow b^b > \frac{a^b b^a}{a^a} \\ &\Rightarrow C_*(a_1, a_2) - C(a_1, a_2) = a^a + b^b - (a^b + b^a) \\ &> a^a + \frac{a^b b^a}{a^a} - (a^b + b^a) = \frac{(a^a - a^b)(a^a - b^a)}{a^a} \geq 0 \\ &\Rightarrow (1.6). \end{aligned}$$

Now we assume that  $1 > a = a_1 > a_2 = b > 0$ . By the A-G inequality [14]

$$\sum_{i=1}^n p_i x_i \geq \prod_{i=1}^n x_i^{p_i},$$

where  $p_i > 0$ ,  $x_i > 0$ ,  $i = 1, \dots, n$ ,  $n \geq 2$ ,  $\sum_{i=1}^n p_i = 1$ , and the logarithm inequalities

$$\frac{x}{1+x} < \log(1+x) < x, \quad \forall x : -1 < x \neq 0,$$

we have

$$\frac{d}{dt} \left( \frac{\log t}{1-t} \right) = (1-t)^2 \left[ \log(1+t-1) - \frac{t-1}{1+t-1} \right] > 0, \quad \forall t : 0 < t < 1$$

$$\begin{aligned}
&\Rightarrow \frac{\log a}{1-a} > \frac{\log b}{1-b} \Leftrightarrow a^{1-b} - b^{1-a} > 0 \\
&\Rightarrow b^{a-1} - a^{b-1} > 0 \\
&\Rightarrow \frac{ab(b^{a-1} - a^{b-1})}{a^b + b^a} = a \times \frac{b^a}{a^b + b^a} - b \times \frac{a^b}{a^b + b^a} > 0 \\
&\Rightarrow \frac{C_*(a_1, a_2)}{C(a_1, a_2)} = \frac{b^a}{a^b + b^a} \left(\frac{a}{b}\right)^a + \frac{a^b}{a^b + b^a} \left(\frac{b}{a}\right)^b \\
&\geq \left(\frac{a}{b}\right)^{a \times \frac{b^a}{a^b + b^a}} \left(\frac{b}{a}\right)^{b \times \frac{a^b}{a^b + b^a}} = \left(\frac{a}{b}\right)^{a \times \frac{b^a}{a^b + b^a} - b \times \frac{a^b}{a^b + b^a}} > 1 \\
&\Rightarrow (1.6).
\end{aligned}$$

Hence inequality (1.6) holds when  $n = 2$ , and equality in (1.6) holds if and only if  $a_1 = a_2$ . This ends the proof of Lemma 3.1.  $\square$

According to the theory of mathematical analysis, we have the following Lemma 3.2.

**Lemma 3.2.** *Let the function  $f : (\alpha, \beta) \rightarrow \mathbb{R}$  be continuous, and let*

$$f(\alpha) \triangleq f(\alpha + 0), \quad f(\beta) \triangleq f(\beta - 0).$$

*If  $f$  has no any minimum points, then we have*

$$\inf_{t \in (\alpha, \beta)} \{f(t)\} = \min \{f(\alpha), f(\beta)\}; \quad (3.1)$$

*If  $t_1, \dots, t_k$ ,  $k \geq 1$ , are all the minimum points of the function  $f$ , then we have*

$$\inf_{t \in (\alpha, \beta)} \{f(t)\} = \min \{f(\alpha), f(\beta), f(t_1), \dots, f(t_k)\}. \quad (3.2)$$

In the following discussion, we define an auxiliary function as follows:

$$F : (0, \infty)^2 \rightarrow \mathbb{R}, \quad F(x, y) \triangleq (y - x) \log x + \log y - \log x. \quad (3.3)$$

**Lemma 3.3.** *Let  $F$  be defined by (3.3). Then we have*

$$F(x, y) > 0, \quad \forall x, y : 0 < x < y < 1. \quad (3.4)$$

*Proof.* We arbitrarily fix  $x \in (0, 1)$ , that is,  $x \in (0, 1)$  is a constant. Since  $y \in (x, 1)$ , and

$$\frac{\partial F(x, y)}{\partial y} = \log x + \frac{1}{y}, \quad \frac{\partial^2 F(x, y)}{\partial y^2} = -\frac{1}{y^2} < 0,$$

the function  $F : (x, 1) \rightarrow \mathbb{R}$  is a concave function for the variable  $y$ , which has no any minimum points. So, by Lemma 3.2, we have

$$F(x, y) > \inf_{x < y < 1} \{F(x, y)\} = \min \{F(x, x), F(x, 1)\} = \min \{0, -x \log x\} = 0. \quad (3.5)$$

This ends the proof of Lemma 3.3.  $\square$



In the following discussion, we define an auxiliary function as follows:

$$\phi : \Omega \rightarrow \mathbb{R}, \phi(x, y, z) \triangleq y^y + z^x - (z^y + y^x), \quad (3.6)$$

where

$$\Omega \triangleq \{(x, y, z) \in (0, \infty)^3 : \max\{x, z\} \leq y\}. \quad (3.7)$$

**Lemma 3.4.** *Let the function  $\phi : \Omega \rightarrow \mathbb{R}$  be define by (3.6), and let  $(x, y, z) \in \Omega$ . If  $y \geq 1$ , then we have the inequality*

$$\phi(x, y, z) \geq 0. \quad (3.8)$$

*Equality in (3.8) holds if and only if  $y = z$  or  $y = x$ .*

*Proof.* The inequality (3.8) can be rewritten as

$$y^x (y^{y-x} - 1) \geq z^x (z^{y-x} - 1). \quad (3.9)$$

If  $z \leq 1 \leq y$ , by  $0 < \max\{x, z\} \leq y$ , we have

$$y^x (y^{y-x} - 1) \geq 0 \geq z^x (z^{y-x} - 1) \Rightarrow (3.9) \Rightarrow (3.8);$$

If  $1 \leq z \leq y$ , from  $0 < \max\{x, z\} \leq y$ , we have

$$y^x \geq z^x > 0, \quad y^{y-x} - 1 \geq z^{y-x} - 1 \geq 0 \Rightarrow y^x (y^{y-x} - 1) \geq z^x (z^{y-x} - 1) \Rightarrow (3.9) \Rightarrow (3.8).$$

Hence (3.8) is proved.

Based on the above proof, we see that equality in (3.8) holds if and only if  $y = z$  or  $y = x$ . Lemma 3.4 is proved.  $\square$

**Lemma 3.5.** *Let the function  $\phi : \Omega \rightarrow \mathbb{R}$  be define by (3.6), and let  $(x, y, z) \in \Omega$ . If  $0 < x \leq z \leq y < 1$ , then inequality (3.8) also holds. Equality in (3.8) holds if and only if  $y = z$ .*

*Proof.* By the proof of Lemma 3.4, we just need to prove (3.9).

Indeed, if  $y = z$ , then (3.9) is an equation. Let's assume that  $0 < x \leq z < y < 1$ . Then  $0 < x < y < 1$ .

Now we proved that

$$y^x (y^{y-x} - 1) > z^x (z^{y-x} - 1). \quad (3.10)$$

Since

$$\begin{aligned} (3.10) &\Leftrightarrow y^x (1 - y^{y-x}) < z^x (1 - z^{y-x}) \\ &\Leftrightarrow x \log y + \log(1 - y^{y-x}) < x \log z + \log(1 - z^{y-x}) \\ &\Leftrightarrow x(\log y - \log z) < -\log(1 - y^{y-x}) + \log(1 - z^{y-x}) \\ &\Leftrightarrow x < \frac{-\log(1 - y^{y-x}) + \log(1 - z^{y-x})}{\log y - \log z}, \end{aligned}$$

we see that inequality (3.10) can be rewritten as

$$x < \frac{f(y) - f(z)}{g(y) - g(z)}, \quad (3.11)$$

where

$$f(t) \triangleq -\log(1 - t^{y-x}), \quad g(t) \triangleq \log t, \quad 0 < x \leq z \leq t \leq y < 1, \quad (3.12)$$

and  $t$  is independent of  $x, y$  and  $z$ . By (3.12), we have

$$\frac{f'(t)}{g'(t)} = \frac{(y-x)t^{y-x}}{1-t^{y-x}} = (y-x) \left( \frac{1}{1-t^{y-x}} - 1 \right), \quad 0 < x \leq z \leq t \leq y < 1. \quad (3.13)$$

By (3.13), the function  $\omega : (0, 1) \rightarrow \mathbb{R}$ ,  $\omega(t) \triangleq f'(t)/g'(t)$ , is strictly incremental. So we have

$$\frac{f'(t)}{g'(t)} > \frac{f'(z)}{g'(z)} \geq \frac{f'(x)}{g'(x)}, \quad \forall t : 0 < x \leq z < t < y < 1. \quad (3.14)$$

According to the Cauchy mean value theorem, (3.12), (3.13) and (3.14), there exists a  $\zeta \in (z, y) \subset (0, 1)$  such that

$$\frac{f(y) - f(z)}{g(y) - g(z)} = \frac{f'(\zeta)}{g'(\zeta)} > \frac{f'(x)}{g'(x)} = \frac{(y-x)x^{y-x}}{1-x^{y-x}}. \quad (3.15)$$

Noting that

$$\begin{aligned} \frac{(y-x)x^{y-x}}{1-x^{y-x}} > x &\Leftrightarrow yx^{y-x} - x^{y-x+1} > x - x^{y-x+1} \\ &\Leftrightarrow yx^{y-x} > x \\ &\Leftrightarrow (y-x)\log x + \log y - \log x > 0, \end{aligned}$$

that is,

$$\frac{(y-x)x^{y-x}}{1-x^{y-x}} > x \Leftrightarrow F(x, y) > 0. \quad (3.16)$$

By  $0 < x < y < 1$ , (3.15), (3.16) and Lemma 3.3, we have

$$\begin{aligned} &F(x, y) > 0, \quad \forall x, y : 0 < x < y < 1 \\ \Rightarrow &\frac{(y-x)x^{y-x}}{1-x^{y-x}} > x, \\ \Rightarrow &\frac{f(y) - f(z)}{g(y) - g(z)} > \frac{(y-x)x^{y-x}}{1-x^{y-x}} > x, \\ \Rightarrow &\frac{f(y) - f(z)}{g(y) - g(z)} > x \\ \Rightarrow &(3.11) \Rightarrow (3.10). \end{aligned}$$

Hence inequality (3.10) is proved.

Since (3.10)  $\Rightarrow$  (3.9)  $\Rightarrow$  (3.8), so the inequality (3.8) is also proved.

Based on the above proof, we see that equality in (3.8) holds if and only if  $y = z$ . Lemma 3.5 is proved.  $\square$

Now we turn to the proof of Theorem 1.2.

*Proof.* We use the mathematical induction for  $n$ .

When  $n = 2$ , by Lemma 3.1, Theorem 1.2 is true.

Suppose that Theorem 1.2 is true for  $n$ ,  $n \geq 2$ . Then

$$C(a_1, a_2, \dots, a_n) - C_*(a_1, a_2, \dots, a_n) \leq 0, \quad (3.17)$$

and equality in (3.17) holds if and only if  $a_1 = a_2 = \dots = a_n$ , where

$$0 < a_1 \leq a_2 \leq \dots \leq a_n.$$

Now we prove that

$$C(a_1, a_2, \dots, a_{n+1}) \leq C_*(a_1, a_2, \dots, a_{n+1}), \quad (3.18)$$

and equality in (3.18) holds if and only if  $a_1 = a_2 = \dots = a_{n+1}$ , where

$$0 < a_1 \leq a_2 \leq \dots \leq a_{n+1}.$$

Since  $0 < a_1 \leq a_2 \leq \dots \leq a_{n+1}$ , we have

$$\max\{a_1, a_2, \dots, a_{n+1}\} = a_{n+1}. \quad (3.19)$$

For ease of expression, by (3.19), we may assume that

$$(x, y, z) \triangleq (a_1, a_{n+1}, a_n) \in \Omega, \quad (3.20)$$

where  $\Omega$  is defined by (3.7). Since

$$\begin{aligned} & C(a_1, a_2, \dots, a_{n+1}) - C_*(a_1, a_2, \dots, a_{n+1}) \\ &= \sum_{i=1}^n a_i^{a_i+1} + a_{n+1}^{a_1} - \left( \sum_{i=1}^n a_i^{a_i} + a_{n+1}^{a_{n+1}} \right) \\ &= \sum_{i=1}^{n-1} a_i^{a_i+1} + a_n^{a_n+1} + a_{n+1}^{a_1} - \left( \sum_{i=1}^n a_i^{a_i} + a_{n+1}^{a_{n+1}} \right) \\ &= \sum_{\text{cyc: } n}^{1 \leq i \leq n} a_i^{a_i+1} - a_n^{a_1} + a_n^{a_{n+1}} + a_{n+1}^{a_1} - \left( \sum_{i=1}^n a_i^{a_i} + a_{n+1}^{a_{n+1}} \right) \\ &= \left( \sum_{\text{cyc: } n}^{1 \leq i \leq n} a_i^{a_i+1} - \sum_{i=1}^n a_i^{a_i} \right) - a_n^{a_1} + a_n^{a_{n+1}} + a_{n+1}^{a_1} - a_{n+1}^{a_{n+1}} \\ &= \left( \sum_{\text{cyc: } n}^{1 \leq i \leq n} a_i^{a_i+1} - \sum_{i=1}^n a_i^{a_i} \right) - [y^y + z^x - (z^y + y^x)] \\ &= C(a_1, a_2, \dots, a_n) - C_*(a_1, a_2, \dots, a_n) - \phi(x, y, z), \end{aligned}$$

we have,

$$C(a_1, \dots, a_{n+1}) - C_*(a_1, \dots, a_{n+1}) = C(a_1, \dots, a_n) - C_*(a_1, \dots, a_n) - \phi(x, y, z), \quad (3.21)$$

where  $\phi(x, y, z)$  is defined by (3.6).

Since  $0 < a_1 \leq a_2 \leq \dots \leq a_{n+1}$ , we see that (3.19) with (3.20) hold and  $x \leq z \leq y$ . According to the inductive hypothesis, we see that (3.17) holds. By Lemmas 3.4 and 3.5, we know that (3.8) holds. According to the (3.8), (3.17) and (3.21), we have

$$C(a_1, a_2, \dots, a_{n+1}) - C_*(a_1, a_2, \dots, a_{n+1})$$

$$\begin{aligned}
&= C(a_1, a_2, \dots, a_n) - C_*(a_1, a_2, \dots, a_n) - \phi(x, y, z) \\
&\leq -\phi(x, y, z) \leq 0 \Rightarrow (3.18).
\end{aligned}$$

Hence inequality (3.18) is proved.

According to the inductive hypothesis and Lemmas 3.4 with 3.5, we see that equality in (3.18) holds if and only if  $a_1 = a_2 = \dots = a_n$  and  $a_{n+1} = y = z = a_n$ , i.e.  $a_1 = a_2 = \dots = a_{n+1}$ .

This completes the proof of Theorem 1.2.  $\square$

**Remark 3.1.** *The proof method of Theorem 1.2 is called the dimensionality reduction method. The relevant literatures on proving inequalities by means of the dimensionality reduction method can be see [2, 3, 5, 7]. The dimension reduction process of the proof is as follows.*

**(A)** *Inequality (3.18) contain  $n + 1$  variables and inequality (1.6) contain  $n$  variables. We transform inequality (3.18) into inequality (1.6) by mean of the mathematical induction. This transformation process is based on the inequality (3.8).*

**(B)** *Inequality (3.8) contain three variables  $x, y, z$ . By mean of Lemmas 3.2-3.5, we transform inequality (3.8) into inequality (3.4), which contain only two variables.*

**(C)** *Lemmas 3.2 and 3.3 transform inequality (3.4) into inequality (3.5). There is only one variable at the right end of the inequality (3.5).*

**(D)** *For an inequality with only one variable, we use mathematical analysis theory to deal with it.*

**Remark 3.2.** *Let the function  $f : [0, 1] \rightarrow (0, \infty)$  be continuous and incremental, and let  $\lim_{n \rightarrow \infty} n^{-1} C(a_1, a_2, \dots, a_n)$  exists, where  $a_i \triangleq f(i/n)$ ,  $i = 1, 2, \dots, n$ ,  $n \geq 2$ , and  $n^{-1} C(a_1, a_2, \dots, a_n)$  is the mean of the positive real numbers  $a_1^{a_2}, a_2^{a_3}, \dots, a_{n-1}^{a_n}, a_n^{a_1}$ . Then, by Theorem 1.2 and the mathematical analysis theory, we have the following inequality:*

$$\lim_{n \rightarrow \infty} n^{-1} C(a_1, a_2, \dots, a_n) \leq \int_0^1 [f(t)]^{f(t)} dt, \quad (3.22)$$

where  $\int_0^1 [f(t)]^{f(t)} dt$  is the mean of the function  $f^f$ . The relevant literatures on mean theory can be see [2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18].

**Competing interests.** The authors declare that they have no conflicts of interest in this joint work.

**Authors contributions.** All authors contributed equally and significantly in this paper. All authors read and approved the final manuscript.

**Acknowledgements.** The authors would like to acknowledge the support from the National Natural Science Foundation of China (No. 11161024).

## References

## References

- [1] F.S. Cater. Problems, Solutions, Amer. Math. Monthly, **87**(4)(1980), 302–303.

- [2] J. J. Wen, J. E. Pečarić and Y. H. Tian. Weak monotonicity and Chebyshev type inequality, *Math. Inequal. Appl.*, **18**(1)(2015), 217–231.
- [3] J. J. Wen and W. L. Wang. The optimization for the inequalities of power means, *J. Inequal. Appl.*, **2006**, Article ID 46782, 25 pages.
- [4] W. L. Wang. Approaches to prove inequalities, Harbin institute of technology, 2011. (In Chinese)
- [5] C. B. Gao and J. J. Wen. A dimensionality reduction principle on the optimization of function, *J. Math. Inequal.*, **7**(3)(2013), 357–375.
- [6] C. B. Gao and J. J. Wen. Theory of surround system and associated inequalities, *Comput. Math. Appl.*, **63**(2012), 1621-1640.
- [7] J. J. Wen, T. Y. Han and S. S. Cheng. Inequalities involving Dresher variance mean, *J. Inequal. Appl.*, **2013**: 366, 29 pages.
- [8] J. J. Wen, C. B. Gao and W. L. Wang. Inequalities of J-P-S-F type, *J. Math. Inequal.*, **7**(2)(2013), 213–225.
- [9] J. J. Wen and Z. H. Zhang. Jensen type inequalities involving homogeneous polynomials, *J. Inequal. Appl.*, **2010**, Article ID 850215, 21 pages.
- [10] J. J. Wen, J. Yuan and S. H. Wu. Isoperimetric inequalities in surround system and space science, *J. Inequal. Appl.*, **2016**: 74, 28 pages.
- [11] J. J. Wen, J. Yuan, S. H. Wu and T. Y. Han. Gravity inequalities and the mean temperature on a planet, *J. Inequal. Appl.*, **2016**: 264, 18 pages.
- [12] J. J. Wen, S. H. Wu, J. Yuan and T. Y. Han. Mean central distance–central distance inequalities, *J. Math. Inequal.*, **11**(4)(2017), 1131–1149.
- [13] T. X. Wang and J. Q. Wei. Mean-variance portfolio selection under a non-Markovian regime-switching model, *J. Comput. Appl. Math.*, **350** (2019), 442–455.
- [14] P. S. Bullen, D. S. Mitrinović and P. M. Vasić. Means and Their Inequalities, Reidel, Dordrecht /Boston /Lancaster /Tokyo, 1988.
- [15] Y. L. Chang, C. H. Huang, J. S. Chen and C. C. Hu. Some inequalities for means defined on the Lorentz cone, *Math. Inequal. Appl.*, **21**(4)(2018), 1015–1028.
- [16] J. E. Pečarić, J. J. Wen, W. L. Wang and T. Lu. A generalization of Maclaurin’s inequalities and its applications, *Math. Inequal. Appl.*, **8**(4)(2005), 583–598.
- [17] J. J. Wen, T. Y. Han and J. Yuan. Stability inequalities involving gravity norm and temperature, *J. Math. Inequal.*, **14**(4)(2020), 1007-1037.
- [18] X. X. You, M. Adil Khan and H. R. Moradi. Bounds for the Jensen gap in terms of power means with applications, *J. Funct. Spaces*, **2021**, Art. ID 1388843, 11 pages.

- [19] J. J. Wen and W. L. Wang. Chebyshev type inequalities involving permanents and their applications, *Linear Algebra Appl.*, **422**(1)(2007), 295-303.