

Article

Relativistic Rational Extended Thermodynamics of Polyatomic Gases with a New Hierarchy of Moments

Takashi Arima¹, Maria Cristina Carrisi², Sebastiano Pennisi², Tommaso Ruggeri³

¹ Department of Engineering for Innovation, National Institute of Technology, Tomakomai College, Tomakomai, Japan; arima@tomakomai-ct.ac.jp

² Department of di Mathematics and Informatics , University of Cagliari, Cagliari, Italy; mariacri.carrisi@unica.it, spennisi@unica.it

³ Department of Mathematics and Alma Mater Research Center on Applied Mathematics AM², University of Bologna, Bologna, Italy; tommaso.ruggeri@unibo.it

Version December 1, 2021 submitted to Entropy

Abstract: A relativistic version of the rational extended thermodynamics of polyatomic gases based on a new hierarchy of moments that takes into account the total energy composed by the rest energy and the energy of the molecular internal mode is proposed. The moment equations associated with the Boltzmann-Chernikov equation are derived, and the system for the first 15 equations is closed by the procedure of the maximum entropy principle and by using an appropriate BGK model for the collisional term. The entropy principle with a convex entropy density is proved in a neighborhood of equilibrium state, and, as a consequence, the system is symmetric hyperbolic and the Cauchy problem is well-posed. The ultra-relativistic and classical limits are also studied. The theories with 14 and 6 moments are deduced as principal subsystems. Particularly interesting is the subsystem with 6 fields in which the dissipation is only due to the dynamical pressure. This simplified model can be very useful when bulk viscosity is dominant and might be important in cosmological problems. Using the Maxwellian iteration, we obtain the parabolic limit, and the heat conductivity, shear viscosity, and bulk viscosity are deduced and plotted.

Keywords: Relativistic extended thermodynamics; Rarefied polyatomic gas; Causal theory of relativistic fluids

1. Introduction

Rational extended thermodynamics (RET) is a theory applicable to nonequilibrium phenomena out of local equilibrium. It is expressed by an hyperbolic system of field equations with local constitutive equations and is strictly related to the kinetic theory with the closure method of the hierarchies of moment equations in both classical and relativistic framework [1,2].

The first relativistic version of the modern RET was given by Liu, Müller and Ruggeri (LMR) [3] considering the Boltzmann-Chernikov relativistic equation [4–6]:

$$p^\alpha \partial_\alpha f = Q, \quad (1)$$

in which the distribution function f depends on (x^α, p^β) , where x^α are the space-time coordinates, p^α is the four-momentum, $\partial_\alpha = \partial/\partial x^\alpha$, c denotes the light velocity, m is the particle mass in the rest frame, Q is the

collisional term and $\alpha, \beta = 0, 1, 2, 3$. For monatomic gases, the relativistic moment equations associated with (1), truncated at tensorial index $N + 1$, are ¹:

$$\partial_\alpha A^{\alpha\alpha_1\cdots\alpha_n} = I^{\alpha_1\cdots\alpha_n} \quad \text{with } n = 0, \dots, N \quad (2)$$

with

$$A^{\alpha\alpha_1\cdots\alpha_n} = \frac{c}{m^{n-1}} \int_{\mathbb{R}^3} f p^\alpha p^{\alpha_1} \cdots p^{\alpha_n} d\mathbf{P}, \quad I^{\alpha_1\cdots\alpha_n} = \frac{c}{m^{n-1}} \int_{\mathbb{R}^3} Q p^{\alpha_1} \cdots p^{\alpha_n} d\mathbf{P}, \quad (3)$$

and

$$d\mathbf{P} = \frac{dp^1 dp^2 dp^3}{p^0}.$$

When $N = 1$, we have the relativistic Euler system

$$\partial_\alpha A^\alpha = 0, \quad \partial_\alpha A^{\alpha\beta} = 0, \quad (4)$$

where, also in the following, $A^\alpha \equiv V^\alpha$ and $A^{\alpha\beta} \equiv T^{\alpha\beta}$ have the physical meaning, respectively, of the particle number vector and the energy-momentum tensor. Instead, when $N = 2$, we have the LMR theory of a relativistic gas with 14 fields:

$$\partial_\alpha A^\alpha = 0, \quad \partial_\alpha A^{\alpha\beta} = 0, \quad \partial_\alpha A^{\alpha\beta\gamma} = I^{\beta\gamma}, \quad (\gamma = 0, 1, 2, 3; I^\alpha_\alpha = 0). \quad (5)$$

Recently, Pennisi and Ruggeri first constructed a relativistic ET theory for polyatomic gases with (2) in the case of $N = 2$ [7] (see also [8,9]) whose moments are given by

$$\begin{aligned} A^\alpha &= mc \int_{\mathbb{R}^3} \int_0^\infty f p^\alpha \phi(I) dI d\mathbf{P}, \\ A^{\alpha\beta} &= \frac{1}{mc} \int_{\mathbb{R}^3} \int_0^\infty f p^\alpha p^\beta (mc^2 + I) \phi(I) dI d\mathbf{P}, \\ A^{\alpha\beta\gamma} &= \frac{1}{m^2 c} \int_{\mathbb{R}^3} \int_0^{+\infty} f p^\alpha p^\beta p^\gamma (mc^2 + 2I) \phi(I) dI d\mathbf{P}, \end{aligned} \quad (6)$$

19 where the distribution function $f(x^\alpha, p^\beta, I)$ depends on the extra energy variable I , similar to the classical one
20 (see [2] and references therein), and $\phi(I)$ is the state density of the internal mode.

In [7], by taking the traceless part of the third order tensor, i.e., $A^{\alpha(\beta\gamma)}$, as a field instead of $A^{\alpha\beta\gamma}$ in (5)₃, the relativistic theory with 14 fields (ET₁₄) was proposed. It was also shown that its classical limit coincides with the classical ET₁₄ based on the binary hierarchy [2,10,11]. The beauty of the relativistic counterpart is that there exists a single hierarchy of moments, but, as was noticed by the authors, to obtain the classical theory of ET₁₄, it was necessary to put the factor 2 in front of I in the last equation of (6)! This was also more evident in the theory with any number of moments where Pennisi and Ruggeri generalized (6) considering the following moments [12]:

$$\begin{aligned} A^{\alpha\alpha_1\cdots\alpha_n} &= \frac{1}{m^n c} \int_{\mathbb{R}^3} \int_0^{+\infty} f p^\alpha p^{\alpha_1} \cdots p^{\alpha_n} (mc^2 + nI) \phi(I) dI d\mathbf{P}, \\ I^{\alpha_1\cdots\alpha_n} &= \frac{1}{m^n c} \int_{\mathbb{R}^3} \int_0^{+\infty} Q p^{\alpha_1} \cdots p^{\alpha_n} (mc^2 + nI) \phi(I) dI d\mathbf{P}. \end{aligned} \quad (7)$$

21 In this case, we need a factor nI in (7) to obtain, in the classical limit, the binary hierarchy.

¹ When $n = 0$, the tensor reduces to A^α . Moreover, the production tensor in the right-side of (3) is zero for $n = 0, 1$, because the first 5 equations represent the conservation laws of the particle number and of the energy-momentum, respectively.

To avoid this unphysical situation, Pennisi first noticed that $(mc^2 + n\mathcal{I})$ appearing in (7) are the first two terms of the Newton binomial formula for $(mc^2 + \mathcal{I})^n / (mc^2)^{n-1}$. Therefore he proposed in [13] to modify, in the relativistic case, the definition of the moments by using the substitution:

$$(mc^2)^{n-1}(mc^2 + n\mathcal{I}) \quad \text{with} \quad (mc^2 + \mathcal{I})^n,$$

i.e., instead of (7), the following moments are proposed:

$$\begin{aligned} A^{\alpha\alpha_1 \dots \alpha_n} &= \left(\frac{1}{mc}\right)^{2n-1} \int_{\mathbb{R}^3} \int_0^{+\infty} f p^\alpha p^{\alpha_1} \dots p^{\alpha_n} (mc^2 + \mathcal{I})^n \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P}, \\ I^{\alpha_1 \dots \alpha_n} &= \left(\frac{1}{mc}\right)^{2n-1} \int_{\mathbb{R}^3} \int_0^{+\infty} Q p^{\alpha_1} \dots p^{\alpha_n} (mc^2 + \mathcal{I})^n \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P}. \end{aligned} \quad (8)$$

Such definitions are more physical because now the full energy (the sum of the rest frame energy and the energy of internal modes) $mc^2 + \mathcal{I}$ appears in the moments.

The aim of this paper is to consider the system (5) with moments given by (8). In this way, for the case with $N = 2$ also by taking the trace part of $A^{\alpha\beta\gamma}$ as a field, we have 15 field equations and to close the system we adopt the molecular procedure of RET based on the Maximum Entropy Principle.

The paper is organized as follows. In Section 2, the values of generic moments in an equilibrium state are estimated in the general case. In Section 3, the ET theory for 15 fields (ET₁₅) is proposed, and the constitutive quantities are closed near the equilibrium state. By adopting a variant of BGK model appropriate for polyatomic gases proposed by Pennisi and Ruggeri [8], the production tensor is derived. In Section 4 the four-dimensional entropy flux and the entropy production are deduced within the second order with respect to the nonequilibrium variables. Then, we show the condition of convexity of the entropy density and the positivity of the entropy production which ensure the well-posedness of the Cauchy problem and the entropy principle as a result. We also discuss in Section 5 the case of the diatomic gases for which all coefficients are expressed in closed form in terms of the ratio of two Bessel functions similar to the case of monatomic gases. In Section 6, we study the ultra-relativistic limit. In Section 7, the principal subsystems of ET₁₅ are studied. First, we obtain ET₁₄ in which all field variables have physical meaning. Then, at the same level as ET₁₄ in the sense of the principal subsystem, there also exists the subsystem with 6 fields in which the dissipation is only due to the dynamical pressure. This system is important in the case that the bulk viscosity is dominant compared to the shear viscosity and heat conductivity and must be particularly interesting in cosmological problems. The simplest subsystem is the Euler non dissipative case with 5 fields. In Section 8, we use the Maxwellian iteration and, as a result, the phenomenological coefficients of the Eckart theory, that is, the heat conductivity, shear viscosity and bulk viscosity are determined with the present model. Finally, in Section 9, we show that the classic limit of the present model coincides with the classical ET₁₅ studied in [14].

2. Distribution function and moments at equilibrium

The equilibrium distribution function f_E of polyatomic gas that generalizes the Jüttner one of monatomic gas was evaluated in [7] with the variational procedure of Maximum Entropy Principle (MEP) [1,15,16]. Considering the first 5 balance equations of (5) in equilibrium state:

$$A_E^\alpha \equiv V_E^\alpha = m n U^\alpha, \quad A_E^{\alpha\beta} \equiv T_E^{\alpha\beta} = p h^{\alpha\beta} + \frac{e}{c^2} U^\alpha U^\beta.$$

MEP requires that the appropriate distribution function $f \equiv f(x^\alpha, p^\alpha, \mathcal{I})$ is the one which maximizes the entropy density

$$\rho S = h_E = h_E^\alpha U_\alpha = -k_B c U_\alpha \int_{\mathbb{R}^3} \int_0^{+\infty} f \ln f p^\alpha \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P},$$

under the constraints that the temporal part $V^\alpha U_\alpha$ and $T^{\alpha\beta} U_\beta$ are prescribed. Here, $k_B, n, \rho (= nm), U^\alpha, h^{\alpha\beta}, p, e, S$ are respectively the Boltzmann constant, the particle number, the mass density, the

48 four-velocity ($U^\alpha U_\alpha = c^2$), the projector tensor ($h^{\alpha\beta} = U^\alpha U^\beta / c^2 - g^{\alpha\beta}$), the pressure, the energy and
49 the entropy density and $g^{\alpha\beta} = \text{diag}(1, -1, -1, -1)$ is the metric tensor.

The equilibrium distribution function for a rarefied polyatomic gas that maximizes the entropy has the following expression [7]:

$$f_E = \frac{n}{\bar{A}(\gamma)} \frac{1}{4\pi m^3 c^3} e^{-\frac{1}{k_B T} \left[\left(1 + \frac{\mathcal{I}}{mc^2}\right) U_\beta p^\beta \right]}, \quad \bar{A}(\gamma) = \int_0^{+\infty} J_{2,1}^* \phi(\mathcal{I}) d\mathcal{I} \quad (9)$$

with T being the absolute temperature,

$$J_{m,n}^* = J_{m,n}(\gamma^*), \quad \gamma^* = \gamma \left(1 + \frac{\mathcal{I}}{mc^2}\right), \quad \gamma = \frac{mc^2}{k_B T},$$

and

$$J_{m,n}(\gamma) = \int_0^{+\infty} e^{-\gamma \cosh s} \sinh^m s \cosh^n s ds,$$

subjected to the following recurrence relations [3,7]:

$$J_{m+2,n}(\gamma) = J_{m,n+2}(\gamma) - J_{m,n}(\gamma), \quad (10)$$

$$-\gamma J_{m+2,n}(\gamma) = n J_{m,n-1}(\gamma) - (n+m+1) J_{m,n+1}(\gamma). \quad (11)$$

The pressure and the energy compatible with the equilibrium distribution function (9) are [7]:

$$p = \frac{k_B}{m} \rho T, \quad e = \rho c^2 \omega(\gamma), \quad (12)$$

$$\text{with } \omega(\gamma) = \frac{\int_0^{+\infty} J_{2,2}^* \left(1 + \frac{\mathcal{I}}{mc^2}\right) \phi(\mathcal{I}) d\mathcal{I}}{\int_0^{+\infty} J_{2,1}^* \phi(\mathcal{I}) d\mathcal{I}}.$$

Taking into account that $e = \rho c^2 + \rho \varepsilon$, where ε is the internal energy, we deduce from (12):

$$\varepsilon = c^2(\omega - 1). \quad (13)$$

50 Therefore the internal energy is a function only of γ or, it is the same, of T as in the classical case for rarefied
51 gases.

The moments in equilibrium state $A_E^{\alpha_1 \dots \alpha_j}$ for $j \geq 2$ were deduced in [13]:

$$A_E^{\alpha_1 \dots \alpha_{j+1}} = \sum_{k=0}^{\lfloor \frac{j+1}{2} \rfloor} \rho c^{2k} \theta_{k,j} h^{(\alpha_1 \alpha_2 \dots \alpha_{2k-1} \alpha_{2k} U^{\alpha_{2k+1}} \dots U^{\alpha_{j+1}})}, \quad (14)$$

where

$$\theta_{k,j} = \frac{1}{2k+1} \binom{j+1}{2k} \frac{\int_0^{+\infty} J_{2k+2, j+1-2k}^* \left(1 + \frac{\mathcal{I}}{mc^2}\right)^j \phi(\mathcal{I}) d\mathcal{I}}{\int_0^{+\infty} J_{2,1}^* \phi(\mathcal{I}) d\mathcal{I}} \quad (15)$$

are dimensionless functions depending only on γ . Taking into account (12) and (15), we obtain $\theta_{0,0} = 1$, $\theta_{0,1} = \omega(\gamma)$ and using the recurrence formula (10) and (11), in [13], the following recurrence relations hold:

$$\begin{aligned} \theta_{0,0} &= 1, \\ \theta_{0,j+1} &= \omega(\gamma) \theta_{0,j} - \theta'_{0,j} && \text{with } ' = \frac{d}{d\gamma}, \\ \theta_{h,j+1} &= \frac{j+2}{\gamma} \left(\theta_{h,j} + \frac{j+3-2h}{2h} \theta_{h-1,j} \right) && \text{for } h = 1, \dots, \left\lfloor \frac{j+1}{2} \right\rfloor, \\ \theta_{\frac{j+2}{2},j+1} &= \frac{1}{\gamma} \theta_{\frac{j}{2},j} && \text{for } j \text{ even.} \end{aligned} \quad (16)$$

It is interesting to see that all the scalar coefficients can be expressed in terms of the function $\omega(\gamma)$ and of its derivatives with respect to γ (or with respect to the temperature T) and ω is strictly related to the internal energy ε by (13). A similar situation is studied in the article [14] for the non relativistic case.

The values of $\theta_{h,j}$ can be determined, by using the recurrence formula (16), according to the following diagram:

$$\begin{array}{ccccccc} \theta_{0,0} & \Rightarrow & \theta_{0,1} & \Rightarrow & \theta_{0,2} & \Rightarrow & \theta_{0,3} \dots \\ & & \searrow & & & & \\ & & & \theta_{1,1} & \rightarrow & \theta_{1,2} & \rightarrow & \theta_{1,3} \dots \\ & & & & & \searrow & & \\ & & & & & & \theta_{2,3} & \rightarrow & \theta_{2,4} \dots \end{array}$$

We see that all the $\theta_{0,j}$ can be obtained from $\theta_{0,0}$ by using eq. (16)₂ and the other $\theta_{h,j}$ with $j \geq h$ can be obtained from eqs. (16)_{3,4}. In particular, we can evaluate the following ones that are needed to know for the model with 15 fields in the subsequent sections:

$$\begin{aligned} \theta_{0,0} &= 1, & \theta_{0,1} &= \omega, & \theta_{0,2} &= \omega^2 - \omega', \\ \theta_{0,3} &= \omega^3 + \omega'' - 3\omega\omega', & \theta_{0,4} &= \omega^4 - \omega''' + 4\omega\omega'' + 3\omega'^2 - 6\omega^2\omega', \\ \theta_{1,1} &= \frac{1}{\gamma}, & \theta_{1,2} &= \frac{3}{\gamma^2}(\gamma\omega + 1), & \theta_{1,3} &= \frac{6}{\gamma^3} \left[\gamma^2(\omega^2 - \omega') + 2(\gamma\omega + 1) \right], \\ \theta_{1,4} &= \frac{10}{\gamma^4} \left\{ 3\gamma[\omega(\gamma\omega + 2) - \gamma\omega'] + 6 + \gamma^3(\omega^3 + \omega'' - 3\omega\omega') \right\}, \\ \theta_{2,3} &= \frac{3}{\gamma^3}(\gamma\omega + 1), & \theta_{2,4} &= \frac{15}{\gamma^4} \left[\gamma^2(\omega^2 - \omega') + 3(\gamma\omega + 1) \right]. \end{aligned} \quad (17)$$

52 3. The closure for the 15 moments model

In this section, we consider the simplest and physical case, that is, the system (2) for $n = 0, 1, 2$ with the moments given by (8):

$$\partial_\alpha V^\alpha = 0, \quad \partial_\alpha T^{\alpha\beta} = 0, \quad \partial_\alpha A^{\alpha\beta\gamma} = I^{\beta\gamma}, \quad (\beta, \gamma = 0, 1, 2, 3). \quad (18)$$

with

$$\begin{aligned} V^\alpha &= mc \int_{\mathbb{R}^3} \int_0^{+\infty} f p^\alpha \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P}, & T^{\alpha\beta} &= c \int_{\mathbb{R}^3} \int_0^{+\infty} f p^\alpha p^\beta \left(1 + \frac{\mathcal{I}}{mc^2}\right) \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P}, \\ A^{\alpha\beta\gamma} &= \frac{c}{m} \int_{\mathbb{R}^3} \int_0^{+\infty} f p^\alpha p^\beta p^\gamma \left(1 + \frac{\mathcal{I}}{mc^2}\right)^2 \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P}, \\ I^{\beta\gamma} &= \frac{c}{m} \int_{\mathbb{R}^3} \int_0^{+\infty} Q p^\beta p^\gamma \left(1 + \frac{\mathcal{I}}{mc^2}\right)^2 \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P}. \end{aligned} \quad (19)$$

To close the system (19), we adopt the MEP which requires to find the distribution function that maximizes the non-equilibrium entropy density:

$$h = h^\alpha U_\alpha = -k_B c U_\alpha \int_{\mathbb{R}^3} \int_0^{+\infty} f \ln f p^\alpha \phi(\mathcal{I}) d\mathcal{I} d\mathbf{P}, \quad \rightarrow \quad \max \quad (20)$$

under the constraints that the temporal part $V^\alpha U_\alpha$, $T^{\alpha\beta} U_\alpha$ and $A^{\alpha\beta\gamma} U_\alpha$ are prescribed. Proceeding in the usual way as indicates in previous papers of RET (see [2,7]), we obtain:

$$f_{15} = e^{-1 - \frac{\chi}{k_B}}, \quad \text{with} \quad \chi = m \lambda + \lambda_\mu p^\mu \left(1 + \frac{\mathcal{I}}{m c^2}\right) + \frac{1}{m} \lambda_{\mu\nu} p^\mu p^\nu \left(1 + \frac{\mathcal{I}}{m c^2}\right)^2, \quad (21)$$

53 where $\lambda, \lambda_\mu, \lambda_{\mu\nu}$ are the Lagrange multipliers.

Hereafter, recalling the following decomposition of the particle number vector and the energy-momentum tensor

$$V^\alpha = \rho U^\alpha, \quad T^{\alpha\beta} = \frac{e}{c^2} U^\alpha U^\beta + (p + \Pi) h^{\alpha\beta} + \frac{1}{c^2} (U^\alpha q^\beta + U^\beta q^\alpha) + t^{\langle\alpha\beta\rangle_3}, \quad (22)$$

we can choose as fields, as usual, 14 physical variables; $\rho, T, U^\alpha, \Pi, q^\alpha, t^{\langle\alpha\beta\rangle_3}$, where Π is the dynamic pressure, $q^\alpha = -h_\mu^\alpha U_\nu T^{\mu\nu}$ is the heat flux and $t^{\langle\alpha\beta\rangle_3} = T^{\mu\nu} \left(h_\mu^\alpha h_\nu^\beta - \frac{1}{3} h^{\alpha\beta} h_{\mu\nu}\right)$ is the deviatoric shear viscous stress tensor. We also recall the constraints:

$$U^\alpha U_\alpha = c^2, \quad q^\alpha U_\alpha = 0, \quad t^{\langle\alpha\beta\rangle_3} U_\alpha = 0, \quad t^{\langle\alpha}_{\alpha>_3} = 0,$$

and we choose as the 15th variable:

$$\Delta = \frac{4}{c^2} U_\alpha U_\beta U_\gamma (A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}). \quad (23)$$

54 The pressure p and the energy e as function of (ρ, T) are given in (12).

Remark 1. : For any symmetric tensor $M^{\alpha\beta}$, we can define its traceless part $M^{\langle\alpha\beta\rangle}$ and its 3-dimensional traceless part $M^{\langle\alpha\beta\rangle_3}$ that is the traceless part of its projection in the 3-dimensional space orthogonal to U^α as follows

$$\begin{aligned} M^{\langle\alpha\beta\rangle} &= \left(g_\mu^\alpha g_\nu^\beta - \frac{1}{4} g^{\alpha\beta} g_{\mu\nu}\right) M^{\mu\nu} = M^{\alpha\beta} - \frac{1}{4} g_{\mu\nu} M^{\mu\nu} g^{\alpha\beta}, \\ M^{\langle\alpha\beta\rangle_3} &= \left(h_\mu^\alpha h_\nu^\beta - \frac{1}{3} h^{\alpha\beta} h_{\mu\nu}\right) M^{\mu\nu}, \end{aligned}$$

which are different except for the case in which $M^{\mu\nu} U_\mu = 0$, and $M^{\mu\nu} g_{\mu\nu} = 0$. In fact, these conditions indicate that

$$M^{\langle\alpha\beta\rangle} = M^{\langle\alpha\beta\rangle_3}.$$

55 Moreover in the following a parenthesis between two indexes indicate the symmetric part.

56 *3.1. The linear deviation from equilibrium*

The thermodynamical definition of the equilibrium according with Müller and Ruggeri [1] is the state in which the entropy production vanishes and hence attains its minimum value. Using this definition, the theorem was proved [17,18] that the components of the Lagrange multipliers of the balance laws of nonequilibrium variables vanish, and only the five Lagrange multipliers corresponding to the equilibrium conservation laws (Euler System) remain. In the present case, we have:

$$\lambda_E = -\frac{1}{T}(g + c^2), \quad \lambda_{\mu E} = \frac{U_\mu}{T}, \quad \lambda_{\mu\nu E} = 0, \quad (24)$$

57 where $g = \varepsilon + p/\rho - TS$ is the equilibrium chemical potential. We remark that $\lambda_E, \lambda_{\mu E}$ are the components of
58 the *main field* that symmetrize the relativistic Euler system as was first proved by Ruggeri and Strumia (see [19]).

In the molecular ET approach, we consider, as usual, the processes near equilibrium. For this reason, we expand (21) around an equilibrium state as follows:

$$\begin{aligned} f_{15} &\simeq f_E \left(1 - \frac{1}{k_B} \tilde{\chi}\right), \\ \tilde{\chi} &= m(\lambda - \lambda_E) + (\lambda_\mu - \lambda_{\mu E}) p^\mu \left(1 + \frac{\mathcal{I}}{m c^2}\right) + \frac{1}{m} \lambda_{\mu\nu} p^\mu p^\nu \left(1 + \frac{\mathcal{I}}{m c^2}\right)^2. \end{aligned} \quad (25)$$

Inserting the distribution function (25) into the moments (19), we obtain the following system:

$$\begin{aligned} 0 &= V^\alpha - V_E^\alpha = -\frac{m}{k_B} \left[V_E^\alpha (\lambda - \lambda_E) + T_E^{\alpha\mu} (\lambda_\mu - \lambda_{\mu E}) + A_E^{\alpha\mu\nu} \lambda_{\mu\nu} \right], \\ t^{<\alpha\beta>_3} + \Pi h^{\alpha\beta} + \frac{2}{c^2} U^{(\alpha} q^{\beta)} &= -\frac{m}{k_B} \left[T_E^{\alpha\beta} (\lambda - \lambda_E) + A_E^{\alpha\beta\mu} (\lambda_\mu - \lambda_{\mu E}) + A_E^{\alpha\beta\mu\nu} \lambda_{\mu\nu} \right], \\ A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma} &= -\frac{m}{k_B} \left[A_E^{\alpha\beta\gamma} (\lambda - \lambda_E) + A_E^{\alpha\beta\gamma\mu} (\lambda_\mu - \lambda_{\mu E}) + A_E^{\alpha\beta\gamma\mu\nu} \lambda_{\mu\nu} \right], \end{aligned} \quad (26)$$

where the equilibrium values of the tensors $A_E^{\alpha\beta\mu}, A_E^{\alpha\beta\mu\nu}$ and $A_E^{\alpha\beta\mu\nu\gamma}$ can be obtained by (14) taking $j = 2, 3, 4$:

$$\begin{aligned} A_E^{\alpha\beta\gamma} &= \rho \theta_{0,2} U^\alpha U^\beta U^\gamma + \rho c^2 \theta_{1,2} h^{(\alpha\beta} U^{\gamma)}, \\ A_E^{\alpha\beta\mu\nu} &= \rho \theta_{0,3} U^\alpha U^\beta U^\mu U^\nu + \rho c^2 \theta_{1,3} h^{(\alpha\beta} U^\mu U^{\nu)} + \rho c^4 \theta_{2,3} h^{(\alpha\beta} h^{\mu\nu)}, \\ A_E^{\alpha\beta\gamma\mu\nu} &= \rho \theta_{0,4} U^\alpha U^\beta U^\gamma U^\mu U^\nu + \rho c^2 \theta_{1,4} h^{(\alpha\beta} U^\gamma U^\mu U^{\nu)} + \rho c^4 \theta_{2,4} h^{(\alpha\beta} h^{\gamma\mu} U^{\nu)}, \end{aligned} \quad (27)$$

59 with the θ 's given in (17).

60 The system (26) permits to deduce the 15 Lagrange multipliers in terms of the 15 field variables including
61 Δ given in (23), and then we can obtain the remaining part of the tensor $A^{\alpha\beta\gamma}$.

To solve this system, we consider first eq. (26)₁ contracted with U_α , eq. (26)₂ contracted with $U_\alpha U_\beta$, eq. (26)₃ contracted with $U_\alpha U_\beta U_\gamma / c^3$, eq. (26)₂ contracted with $h_{\alpha\beta} / 3$ and (26)₃ contracted with $U_\alpha h_{\beta\gamma} / (3 c^2)$, obtaining the system

$$\begin{aligned}
\theta_{0,0} (\lambda - \lambda_E) + \theta_{0,1} U^\mu \left(\lambda_\mu - \frac{U_\mu}{T} \right) + \theta_{0,2} U^\mu U^\nu \lambda_{\mu\nu} + \frac{c^2}{3} \theta_{1,2} h^{\mu\nu} \lambda_{\mu\nu} &= 0, \\
\theta_{0,1} (\lambda - \lambda_E) + \theta_{0,2} U^\mu \left(\lambda_\mu - \frac{U_\mu}{T} \right) + \theta_{0,3} U^\mu U^\nu \lambda_{\mu\nu} + \frac{c^2}{6} \theta_{1,3} h^{\mu\nu} \lambda_{\mu\nu} &= 0, \\
\theta_{0,2} (\lambda - \lambda_E) + \theta_{0,3} U^\mu \left(\lambda_\mu - \frac{U_\mu}{T} \right) + \theta_{0,4} U^\mu U^\nu \lambda_{\mu\nu} + \frac{c^2}{10} \theta_{1,4} h^{\mu\nu} \lambda_{\mu\nu} &= -\frac{k_B}{4 m^2 n c^4} \Delta, \\
\theta_{1,1} (\lambda - \lambda_E) + \frac{1}{3} \theta_{1,2} U^\mu \left(\lambda_\mu - \frac{U_\mu}{T} \right) + \frac{1}{6} \theta_{1,3} U^\mu U^\nu \lambda_{\mu\nu} + \frac{5}{9} c^2 \theta_{2,3} h^{\mu\nu} \lambda_{\mu\nu} &= -\frac{k_B}{m^2 n c^2} \Pi, \\
\frac{1}{3} \theta_{1,2} (\lambda - \lambda_E) + \frac{1}{6} \theta_{1,3} U^\mu \left(\lambda_\mu - \frac{U_\mu}{T} \right) + \frac{1}{10} \theta_{1,4} U^\mu U^\nu \lambda_{\mu\nu} + \frac{c^2}{9} \theta_{2,4} h^{\mu\nu} \lambda_{\mu\nu} &= \\
&= -\frac{k_B}{3 m^2 c^4 n} \left(A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma} \right) U_\alpha h_{\beta\gamma}.
\end{aligned} \tag{28}$$

This is a system of 5 equations in the 4 unknowns $\lambda - \lambda_E$, $U^\mu \left(\lambda_\mu - \frac{U_\mu}{T} \right)$, $U^\mu U^\nu \lambda_{\mu\nu}$, $h^{\mu\nu} \lambda_{\mu\nu}$; in order to have solutions, the determinant of the complete matrix must be zero, i.e.,

$$0 = \begin{vmatrix} \theta_{0,0} & \theta_{0,1} & \theta_{0,2} & \frac{1}{3} \theta_{1,2} & 0 \\ \theta_{0,1} & \theta_{0,2} & \theta_{0,3} & \frac{1}{6} \theta_{1,3} & 0 \\ \theta_{0,2} & \theta_{0,3} & \theta_{0,4} & \frac{1}{10} \theta_{1,4} & -\frac{k_B}{4 m c^4} \Delta \\ \theta_{1,1} & \frac{1}{3} \theta_{1,2} & \frac{1}{6} \theta_{1,3} & \frac{5}{9} \theta_{2,3} & -\frac{k_B}{m c^2} \Pi \\ \frac{1}{3} \theta_{1,2} & \frac{1}{6} \theta_{1,3} & \frac{1}{10} \theta_{1,4} & \frac{1}{9} \theta_{2,4} & -\frac{k_B}{3 m c^4} \left(A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma} \right) U_\alpha h_{\beta\gamma} \end{vmatrix}. \tag{29}$$

By defining

$$D_4 = \begin{vmatrix} \theta_{0,0} & \theta_{0,1} & \theta_{0,2} & \frac{1}{3} \theta_{1,2} \\ \theta_{0,1} & \theta_{0,2} & \theta_{0,3} & \frac{1}{6} \theta_{1,3} \\ \theta_{0,2} & \theta_{0,3} & \theta_{0,4} & \frac{1}{10} \theta_{1,4} \\ \theta_{1,1} & \frac{1}{3} \theta_{1,2} & \frac{1}{6} \theta_{1,3} & \frac{5}{9} \theta_{2,3} \end{vmatrix},$$

$$N^\Pi = - \begin{vmatrix} \theta_{0,0} & \theta_{0,1} & \theta_{0,2} & \frac{1}{3} \theta_{1,2} \\ \theta_{0,1} & \theta_{0,2} & \theta_{0,3} & \frac{1}{6} \theta_{1,3} \\ \theta_{0,2} & \theta_{0,3} & \theta_{0,4} & \frac{1}{10} \theta_{1,4} \\ \frac{1}{3} \theta_{1,2} & \frac{1}{6} \theta_{1,3} & \frac{1}{10} \theta_{1,4} & \frac{1}{9} \theta_{2,4} \end{vmatrix}, \quad N^\Delta = \begin{vmatrix} \theta_{0,0} & \theta_{0,1} & \theta_{0,2} & \frac{1}{3} \theta_{1,2} \\ \theta_{0,1} & \theta_{0,2} & \theta_{0,3} & \frac{1}{6} \theta_{1,3} \\ \theta_{1,1} & \frac{1}{3} \theta_{1,2} & \frac{1}{6} \theta_{1,3} & \frac{5}{9} \theta_{2,3} \\ \frac{1}{3} \theta_{1,2} & \frac{1}{6} \theta_{1,3} & \frac{1}{10} \theta_{1,4} & \frac{1}{9} \theta_{2,4} \end{vmatrix},$$

the (29) gives:

$$\frac{1}{3c^2} (A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) U_\alpha h_{\beta\gamma} = -\frac{N^\Pi}{D_4} \Pi - \frac{N^\Delta}{D_4} \frac{1}{4c^2} \Delta. \quad (30)$$

We contract now eq. (26)₁ with h_α^δ , eq. (26)₂ with $U_\alpha h_\beta^\delta$, eq. (26)₃ with $U_\alpha U_\beta h_\gamma^\delta / c^3$ and (26)₃ with $h_\alpha^\delta h_{\beta\gamma} / (3c^2)$, obtaining the system

$$\begin{aligned} c^2 \theta_{1,1} h^{\delta\mu} (\lambda_\mu - \lambda_{\mu E}) + \frac{2}{3} c^2 \theta_{1,2} U^\mu h^{\delta\nu} \lambda_{\mu\nu} &= 0, \\ c^2 \theta_{1,2} h^{\delta\mu} (\lambda_\mu - \lambda_{\mu E}) + c^2 \theta_{1,3} U^\mu h^{\delta\nu} \lambda_{\mu\nu} &= -\frac{3k_B}{m^2 c^2 n} q^\delta, \\ c^2 \theta_{1,3} h^{\delta\mu} (\lambda_\mu - \lambda_{\mu E}) + \frac{18}{15} c^2 \theta_{1,4} U^\mu h^{\delta\nu} \lambda_{\mu\nu} &= \frac{6k_B}{m^2 c^4 n} (A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) U_\alpha U_\beta h_\gamma^\delta, \\ \frac{5}{3} c^4 \theta_{2,3} h^{\delta\mu} (\lambda_\mu - \lambda_{\mu E}) + \frac{2}{3} c^4 \theta_{2,4} U^\mu h^{\delta\nu} \lambda_{\mu\nu} &= \frac{k_B}{m^2 n} (A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) h_{\alpha\beta} h_\gamma^\delta. \end{aligned} \quad (31)$$

By eliminating the parameters $h^{\delta\mu} (\lambda_\mu - \lambda_{\mu E})$ and $U^\mu h^{\delta\nu} \lambda_{\mu\nu}$ from these equations, we obtain

$$\begin{aligned} (A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) U_\alpha U_\beta h_\gamma^\delta &= -c^2 \frac{N_3}{D_3} q^\delta, \\ (A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) h_{\alpha\beta} h_\gamma^\delta &= -\frac{N_{31}}{D_3} q^\delta, \end{aligned} \quad (32)$$

with

$$D_3 = \begin{vmatrix} \theta_{1,1} & \theta_{1,2} \\ \theta_{1,2} & \frac{3}{2} \theta_{1,3} \end{vmatrix}, \quad N_3 = \frac{1}{2} \begin{vmatrix} \theta_{1,1} & \theta_{1,2} \\ \theta_{1,3} & \frac{9}{5} \theta_{1,4} \end{vmatrix}, \quad N_{31} = \begin{vmatrix} \theta_{1,1} & \theta_{1,2} \\ 5\theta_{2,3} & 3\theta_{2,4} \end{vmatrix}.$$

We contract now eq. (26)₂ with $h_\alpha^{<\delta} h_\beta^{>3}$ and (26)₃ with $h_\alpha^{<\delta} h_\beta^{>3} U_\gamma$, obtaining

$$\begin{aligned} -\frac{k_B}{m} t^{<\delta\theta>3} &= \frac{2}{3} m n c^4 \theta_{2,3} h^{\mu<\delta} h^{\theta>3\nu} \lambda_{\mu\nu}, \\ (A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) h_\alpha^{<\delta} h_\beta^{>3} U_\gamma &= -\frac{2}{15} \frac{m}{k_B} m n c^6 \theta_{2,4} h^{\mu<\delta} h^{\theta>3\nu} \lambda_{\mu\nu}, \end{aligned} \quad (33)$$

from which it follows

$$(A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) h_\alpha^{<\delta} h_\beta^{>3} U_\gamma = C_5 c^2 t^{<\delta\theta>3} \quad \text{with} \quad C_5 = \frac{1}{5} \frac{\theta_{2,4}}{\theta_{2,3}}. \quad (34)$$

Finally, (26)₃ contracted with $h_\alpha^{<\delta} h_\beta^\theta h_\gamma^{\psi>3}$ gives

$$(A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma}) h_\alpha^{<\delta} h_\beta^\theta h_\gamma^{\psi>3} = 0.$$

This result, jointly with (30), (32), (34), gives the decomposition of the triple tensor $A^{\alpha\beta\gamma}$:

$$\begin{aligned} A^{\alpha\beta\gamma} - A_E^{\alpha\beta\gamma} &= \frac{1}{4c^4} \Delta U^\alpha U^\beta U^\gamma - \frac{3}{4c^2} \frac{N^\Delta}{D_4} \Delta h^{(\alpha\beta} U^{\gamma)} - 3 \frac{N^\Pi}{D_4} \Pi h^{(\alpha\beta} U^{\gamma)} \\ &+ \frac{3}{c^2} \frac{N_3}{D_3} q^{(\alpha} U^\beta U^{\gamma)} + \frac{3}{5} \frac{N_{31}}{D_3} h^{(\alpha\beta} q^{\gamma)} + 3C_5 t^{<\alpha\beta>3} U^{\gamma)}. \end{aligned}$$

Thanks to eq. (27)₁, we have the closure of the triple tensor in terms of the physical variables:

$$A^{\alpha\beta\gamma} = \left(\rho \theta_{0,2} + \frac{1}{4c^4} \Delta \right) U^\alpha U^\beta U^\gamma + \left(\rho c^2 \theta_{1,2} - \frac{3}{4c^2} \frac{N^\Delta}{D_4} \Delta - 3 \frac{N^\Pi}{D_4} \Pi \right) h^{(\alpha\beta} U^{\gamma)} \\ + \frac{3}{c^2} \frac{N_3}{D_3} q^{(\alpha} U^\beta U^{\gamma)} + \frac{3}{5} \frac{N_{31}}{D_3} h^{(\alpha\beta} q^{\gamma)} + 3C_5 t^{(\alpha\beta\gamma>3} U^{\gamma)}. \quad (35)$$

62 3.2. Inversion of the Lagrange Multipliers

In this section, we present the explicit expression of the Lagrange Multipliers in terms of the 15 physical independent variables. From the representation theorems, they are expressed as follows:

$$\lambda - \lambda_E = a_1 \Pi + a_2 \Delta, \\ \lambda_\mu - \lambda_{\mu E} = (b_1 \Pi + b_2 \Delta) U_\mu + b_3 q_\mu, \\ \lambda_{\mu\nu} = (\alpha_1 \Pi + \beta_1 \Delta) U_\mu U_\nu + (\alpha_2 \Pi + \beta_2 \Delta) h_{\mu\nu} + \alpha_3 (q_\mu U_\nu + q_\nu U_\mu) + \alpha_4 t_{\langle\mu\nu\rangle>3}, \quad (36)$$

63 where λ_E and $\lambda_{\mu E}$ can be found in eq. (24), and the coefficients $a_{1,2}, b_{1,2,3}, \alpha_{1,2,3,4}$ and $\beta_{1,2}$ are functions of ρ and
64 γ . By using eqs. (28), (31) and (33), it is possible to obtain the explicit expressions of these coefficients.

For convenience, let us denote by D_4^{ij} the minor determinant obtained from D_4 by deleting its i th row and j th column. From system (28), we obtain

$$\lambda - \lambda_E = -\frac{k_B}{mc^4 \rho D_4} \left(-\Pi c^2 D_4^{41} + \frac{\Delta}{4} D_4^{31} \right), \\ U^\mu (\lambda_\mu - \lambda_{\mu E}) = -\frac{k_B}{mc^4 \rho D_4} \left(\Pi c^2 D_4^{42} - \frac{\Delta}{4} D_4^{32} \right), \\ U^\beta U^\gamma \lambda_{\beta\gamma} = -\frac{k_B}{mc^4 \rho D_4} \left(-\Pi c^2 D_4^{43} + \frac{\Delta}{4} D_4^{33} \right), \\ h^{\beta\gamma} \lambda_{\beta\gamma} = -\frac{k_B}{mc^4 \rho D_4} \left(\Pi D_4^{44} - \frac{\Delta}{4c^2} D_4^{34} \right). \quad (37)$$

From system (31) we obtain

$$h^{\delta\mu} (\lambda_\mu - \lambda_{\mu E}) = \frac{3k_B \theta_{1,2}}{mc^4 \rho D_3} q^\delta \quad \text{and} \quad U^\beta h^{\gamma\delta} \lambda_{\beta\gamma} = -\frac{9k_B \theta_{1,1}}{2mc^4 \rho D_3} q^\delta. \quad (38)$$

Finally, from eq. (33) we have

$$h^{\beta<\delta} h^{\theta>3\gamma} \lambda_{\beta\gamma} = -\frac{3k_B}{2mc^4 \rho \theta_{2,3}} t^{\langle\delta\theta\rangle>3},$$

that, multiplied by $t_{\langle\delta\theta\rangle>3}$, gives

$$t^{\langle\beta\gamma\rangle>3} \lambda_{\beta\gamma} = -\frac{3k_B}{2mc^4 \rho \theta_{2,3}} t^{\langle\beta\gamma\rangle>3} t_{\langle\beta\gamma\rangle>3}. \quad (39)$$

By comparing eqs. (36)₁ with (37)₁ we have

$$a_1 = \frac{k_B}{mc^2 \rho D_4} D_4^{41}, \quad a_2 = -\frac{k_B}{4mc^4 \rho D_4} D_4^{31}. \quad (40)$$

By multiplying eq. (36)₂ respectively times U^μ and $h^{\mu\delta}$, and using eqs. (37)₂ and (38)₁ we have

$$b_1 = -\frac{k_B}{mc^4\rho D_4}D_4^{42}, \quad b_2 = \frac{k_B}{4mc^6\rho D_4}D_4^{32}, \quad b_3 = \frac{3k_B\theta_{1,2}}{mc^4\rho D_3}. \quad (41)$$

Finally, by multiplying equation (36)₃ respectively times $U^\mu U^\nu$, $h^{\mu\nu}$, $U^\nu h^{\mu\delta}$, $h^{\mu<\delta}h^{\theta>\nu}$ and using eqs. (37)-(39) we obtain that

$$\begin{aligned} \alpha_1 &= \frac{k_B}{mc^6\rho D_4}D_4^{43}, & \alpha_2 &= -\frac{k_B}{3mc^4\rho D_4}D_4^{44}, \\ \alpha_3 &= \frac{9k_B\theta_{1,1}}{2mc^6\rho D_3}, & \alpha_4 &= -\frac{3k_B}{2mc^4\rho\theta_{2,3}}, \\ \beta_1 &= -\frac{k_B}{4mc^8\rho D_4}D_4^{33}, & \beta_2 &= \frac{k_B}{12mc^6\rho D_4}D_4^{34}. \end{aligned} \quad (42)$$

3.3. Production term with a variant BGK model

To complete the closure of the system (18), we need to have the expression of the production tensor $I^{\beta\gamma}$. It depends on the collisional term Q (see (19)₂), and obtaining the expression of Q is an hard task in relativity. Usually, for monatomic gas it is adopted the relativistic generalization of the BGK approximation first made by Marle [20,21] and successively by Anderson and Witting [22]. The Marle model is an extension of the classical BGK model in the Eckart frame [6,23], and the Anderson-Witting model obtains such extension using the Landau-Lifshitz frame [6,24]. There are some weak points for Marle model, and the Anderson-Witting model uses the Landau-Lifshitz four velocity. Starting from these considerations, Pennisi and Ruggeri proposed a variant of Anderson-Witting model in the Eckart frame both for monatomic and polyatomic gases, and proved that the conservation laws of particle number and energy-momentum are satisfied and the H-theorem holds [8] (see also [2]). In the polyatomic case, the following collision term has been proposed:

$$Q = \frac{U^\alpha p_\alpha}{c^2\tau} \left(f_E - f - f_E p^\mu q_\mu \frac{1 + \frac{I}{mc^2}}{bmc^2} \right), \quad (43)$$

where $3b$ is the coefficient of $h^{(\alpha\beta}U^{\gamma)}$ in eq. (27)₁, i.e $3b = \rho c^2\theta_{1,2}$.

The most general expression of a nonequilibrium double tensor as a linear function of Δ , Π , $t^{<\mu\nu>3}$ and q^μ is the following:

$$I^{\beta\gamma} = (B_1^\Delta \Delta + B_1^\Pi \Pi) U^\beta U^\gamma + (B_2^\Delta \Delta + B_2^\Pi \Pi) h^{\beta\gamma} + B^q U^{(\beta} q^{\gamma)} + B^t t^{<\beta\gamma>3}.$$

In order to determine the coefficients in $I^{\alpha\beta}$, we have to substitute eq. (43) into eq. (19)₄, obtaining

$$\begin{aligned} I^{\beta\gamma} &= \frac{c}{m} \int_{\mathbb{R}^3} \int_0^{+\infty} \frac{U^\alpha p_\alpha}{c^2\tau} \left(f_E - f - f_E p^\mu q_\mu \frac{1 + \frac{I}{mc^2}}{bmc^2} \right) p^\beta p^\gamma \left(1 + \frac{I}{mc^2} \right)^2 \phi(I) dI d\mathbf{P} = \\ &= \frac{U_\alpha}{c^2\tau} (A_E^{\alpha\beta\gamma} - A^{\alpha\beta\gamma}) - 3 \frac{U_\alpha q_\mu}{\theta_{1,2} m^2 n c^6 \tau} A_E^{\alpha\beta\gamma\mu}, \end{aligned}$$

then we have

$$\begin{aligned} B_1^\Delta &= -\frac{1}{4c^4\tau}, & B_1^\Pi &= 0, & B_2^\Delta &= \frac{1}{4c^2\tau} \frac{N^\Delta}{D_4}, & B_2^\Pi &= \frac{1}{\tau} \frac{N^\Pi}{D_4} \\ B^q &= \frac{1}{c^2\tau} \left(\frac{\theta_{1,3}}{\theta_{1,2}} - 2 \frac{N_3}{D_3} \right), & B^t &= -\frac{1}{\tau} C_5. \end{aligned} \quad (44)$$

Therefore the final expression of the production term $I^{\beta\gamma}$ is

$$I^{\beta\gamma} = \frac{1}{\tau} \left\{ -\frac{1}{4c^4} \Delta U^\beta U^\gamma + \left(\frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta + \frac{N^\Pi}{D_4} \Pi \right) h^{\beta\gamma} + \left(-\frac{2}{c^2} \frac{N_3}{D_3} + \frac{\theta_{1,3}}{\theta_{1,2}} \frac{1}{c^2} \right) q^{(\beta} U^{\gamma)} - C_5 t^{<\beta\gamma>_3} \right\} \quad (45)$$

66 We summarize the results of this section as:

67 **Statement 1.** *The closed system (18) obtained via MEP is the one for which $V^\alpha, T^{\alpha\beta}, A^{\alpha\beta\gamma}, I^{\beta\gamma}$ are given*
 68 *explicitly in terms of the 15 fields $(\rho, \gamma, \Pi, \Delta, U^\alpha, q^\alpha, t^{<\alpha\beta>_3})$ using the expressions (22), (35) and (45). All*
 69 *coefficients are completely determined in terms of a single function $\omega(\gamma)$ given by eq. (12)₃ and its derivatives*
 70 *up to the order 3. Observe, by taking into account (13), that the coefficients θ 's given in (17) can be formally*
 71 *written in terms of the internal energy ε and its derivatives.*

72 3.4. Closed system of the field equations and material derivative

It is possible to write now explicitly the differential system for the field variables using the material derivative. The relativistic material derivative of a function f is defined as the derivative with respect to the proper time $\bar{\tau}$ along the path of the particle:

$$\dot{f} = \frac{df}{d\bar{\tau}} = \frac{df}{dt} \frac{dt}{d\bar{\tau}} = \Gamma(\partial_t f + v^j \partial_j f) = U^\alpha \partial_\alpha f, \quad (46)$$

where Γ is the Lorentz factor and we take into account that

$$U^\alpha = \frac{dx^\alpha}{d\bar{\tau}} \equiv (\Gamma c, \Gamma v^j),$$

where v^j is the velocity. Now we observe that for any balance law we can have the following identity:

$$\begin{aligned} I^{\alpha_1 \dots \alpha_n} &= \partial_\alpha A^{\alpha \alpha_1 \dots \alpha_n} = g_\alpha^\beta \partial_\beta A^{\alpha \alpha_1 \dots \alpha_n} = \left(-h_\alpha^\beta + \frac{U^\beta U_\alpha}{c^2} \right) \partial_\beta A^{\alpha \alpha_1 \dots \alpha_n} = \\ &= \frac{U_\alpha}{c^2} \dot{A}^{\alpha \alpha_1 \dots \alpha_n} - h_\alpha^\beta \partial_\beta A^{\alpha \alpha_1 \dots \alpha_n}. \end{aligned}$$

In our case with $n = 0, 1, 2$, these equations are written as follows:

$$\begin{aligned} \partial_\alpha (\rho U^\alpha) &= 0, \quad h_{\delta\beta} \left(\frac{U_\alpha}{c^2} \dot{T}^{\alpha\beta} - h_\alpha^\mu \partial_\mu T^{\alpha\beta} \right) = 0, \quad U_\beta \left(\frac{U_\alpha}{c^2} \dot{T}^{\alpha\beta} - h_\alpha^\mu \partial_\mu T^{\alpha\beta} \right) = 0, \\ h_{\delta\beta} h_{\theta\gamma} \left(\frac{U_\alpha}{c^2} \dot{A}^{\alpha\beta\gamma} - h_\alpha^\mu \partial_\mu A^{\alpha\beta\gamma} - I^{\beta\gamma} \right) &= 0, \\ h_{\delta\beta} U_\gamma \left(\frac{U_\alpha}{c^2} \dot{A}^{\alpha\beta\gamma} - h_\alpha^\mu \partial_\mu A^{\alpha\beta\gamma} - I^{\beta\gamma} \right) &= 0, \quad U_\beta U_\gamma \left(\frac{U_\alpha}{c^2} \dot{A}^{\alpha\beta\gamma} - h_\alpha^\mu \partial_\mu A^{\alpha\beta\gamma} - I^{\beta\gamma} \right) = 0. \end{aligned}$$

By using the expressions (22), (35) and (45), respectively for V^α , $T^{\alpha\beta}$, $A^{\alpha\beta\gamma}$ and $I^{\beta\gamma}$, we see that these become

$$\begin{aligned}
& \dot{\rho} + \rho \partial_\alpha U^\alpha = 0, \\
& -\frac{e+p+\Pi}{c^2} \dot{U}^\delta + \frac{1}{c^2} h_\beta^\delta \dot{q}^\beta + \frac{1}{c^2} t^{<\alpha\delta>_3} \dot{U}_\alpha - h^{\delta\mu} \partial_\mu (p+\Pi) - \frac{1}{c^2} q^\mu \partial_\mu U^\delta - \frac{1}{c^2} q^\delta \partial_\alpha U^\alpha - h_\beta^\delta h_\alpha^\mu \partial_\mu t^{<\alpha\beta>_3} = 0, \\
& \dot{e} + 2 \frac{U_\alpha}{c^2} \dot{q}^\alpha + (e+p+\Pi) \partial_\alpha U^\alpha - h_\alpha^\mu \partial_\mu q^\alpha - t^{<\alpha\beta>_3} \partial_\alpha U_\beta = 0, \\
& h_{\delta\beta} \left(\frac{1}{3} \rho c^2 \theta_{1,2} - \frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta - \frac{N^\Pi}{D_4} \Pi \right)^\bullet + C_5 h_{\delta\gamma} h_{\theta\beta} i^{<\theta\gamma>_3} + t_{<\delta\beta>_3} \dot{C}_5 - \frac{2}{c^2} \left(\frac{N_3}{D_3} + \frac{1}{5} \frac{N_{31}}{D_3} \right) q_{(\delta} h_{\beta)\gamma} \dot{U}^\gamma - \\
& \quad \frac{1}{5c^2} \frac{N_{31}}{D_3} h_{\beta\delta} q^\alpha \dot{U}_\alpha + \left(-\frac{1}{3} \rho c^2 \theta_{1,2} + \frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta + \frac{N^\Pi}{D_4} \Pi \right) \left[-h_{\delta\beta} \partial_\alpha U^\alpha + 2 h_{\theta(\delta} h_{\beta)}^\mu \partial_\mu U^\theta \right] + \\
& \quad \frac{1}{5} \left(q^\mu h_{\delta\beta} + 2 q_{(\delta} h_{\beta)}^\mu \right) \partial_\mu \left(\frac{N_{31}}{D_3} \right) - \frac{1}{5} \frac{N_{31}}{D_3} \left[h_{\delta\beta} h_\alpha^\mu \partial_\mu q^\alpha + 2 h_{\theta(\delta} h_{\beta)}^\mu \partial_\mu q^\theta \right] + \\
& \quad C_5 \left[t_{<\delta\beta>_3} \partial_\alpha U^\alpha + 2 t^{<\mu\gamma>_3} h_{\gamma(\beta} h_{\delta)\theta} \partial_\mu U^\theta \right] = \\
& \quad \frac{1}{\tau} \left(\frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta + \frac{N^\Pi}{D_4} \Pi \right) h_{\delta\beta} - \frac{1}{\tau} C_5 t_{<\delta\beta>_3}, \tag{47} \\
& h_{\beta\delta} \dot{U}^\beta \left(\rho \theta_{0,2} c^2 + \frac{2}{3} \rho c^2 \theta_{1,2} + \frac{1}{4c^2} \Delta - \frac{1}{2c^2} \frac{N^\Delta}{D_4} \Delta - 2 \frac{N^\Pi}{D_4} \Pi \right) + h_{\beta\delta} \frac{N_3}{D_3} \dot{q}^\beta - q_\delta \left(\frac{N_3}{D_3} \right)^\bullet + (2C_5 - 1) t^{<\delta\gamma>_3} \dot{U}_\gamma - \\
& \quad h_\delta^\mu \partial_\mu \left(\frac{1}{3} \rho c^4 \theta_{1,2} - \frac{1}{4} \frac{N^\Delta}{D_4} \Delta - \frac{N^\Pi}{D_4} c^2 \Pi \right) - \left(\frac{N_3}{D_3} + \frac{1}{5} \frac{N_{31}}{D_3} \right) \left(q^\mu \partial_\mu U_\delta + q_\delta \partial_\alpha U^\alpha \right) + \\
& \quad \frac{1}{5} \frac{N_{31}}{D_3} h_\delta^\mu q^\gamma \partial_\mu U_\gamma + h_\alpha^\mu \partial_\mu \left(C_5 c^2 t^{<\alpha\delta>_3} \right) = \frac{1}{\tau} \left(\frac{N_3}{D_3} - \frac{\theta_{1,3}}{2\theta_{1,2}} \right) q_\delta, \\
& \left(\rho \theta_{0,2} c^4 + \frac{1}{4} \Delta \right)^\bullet - 3 \frac{N_3}{D_3} q^\alpha \dot{U}_\alpha + \partial_\alpha U^\alpha \cdot \left(\rho \theta_{0,2} c^4 + \frac{2}{3} \rho c^4 \theta_{1,2} + \frac{1}{4} \Delta - \frac{1}{2} \frac{N^\Delta}{D_4} \Delta - 2 \frac{N^\Pi}{D_4} \Pi c^2 \right) - \\
& \quad h_\alpha^\mu \partial_\mu \left(\frac{N_3}{D_3} c^2 q^\alpha \right) - 2 C_5 c^2 t^{<\mu\gamma>_3} \partial_\mu U_\gamma = -\frac{1}{4\tau} \Delta.
\end{aligned}$$

It may be useful to decompose (47)₄ into the trace and spatial traceless parts. The trace part is given by

$$\begin{aligned}
& \left(\rho c^2 \theta_{1,2} - \frac{3}{4c^2} \frac{N^\Delta}{D_4} \Delta - 3 \frac{N^\Pi}{D_4} \Pi \right)^\bullet + C_5 h_{\theta\gamma} i^{<\theta\gamma>_3} + \frac{1}{c^2} \left(2 \frac{N_3}{D_3} - \frac{1}{5} \frac{N_{31}}{D_3} \right) q_\gamma \dot{U}^\gamma - \\
& \quad \left(-\frac{1}{3} \rho c^2 \theta_{1,2} + \frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta + \frac{N^\Pi}{D_4} \Pi \right) \partial_\alpha U^\alpha + q^\mu \partial_\mu \left(\frac{N_{31}}{D_3} \right) - \\
& \quad \frac{N_{31}}{D_3} h_\alpha^\mu \partial_\mu q^\alpha - 2 C_5 t^{<\mu\gamma>_3} \partial_\mu U_\gamma = \frac{3}{\tau} \left(\frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta + \frac{N^\Pi}{D_4} \Pi \right), \tag{48}
\end{aligned}$$

and the spatial traceless part is:

$$\begin{aligned}
& C_5 h_{\gamma<\delta} h_{\beta>_3} \theta i^{<\gamma\theta>_3} + t_{<\delta\beta>_3} \dot{C}_5 + \frac{2}{c^2} \left(\frac{N_3}{D_3} + \frac{1}{5} \frac{N_{31}}{D_3} \right) q_{<\delta} \dot{U}_{\beta>_3} + \\
& \quad + 2 \left(-\frac{1}{3} \rho c^2 \theta_{1,2} + \frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta + \frac{N^\Pi}{D_4} \Pi \right) h_{\gamma<\delta} h_{\beta>_3}^\mu \partial_\mu U^\gamma + \\
& \quad + \frac{2}{5} \left(q_{<\delta} h_{\beta>_3}^\mu \right) \partial_\mu \left(\frac{N_{31}}{D_3} \right) - \frac{2}{5} \frac{N_{31}}{D_3} \left(h_{\gamma<\delta} h_{\beta>_3}^\mu \partial_\mu q^\gamma \right) + \\
& \quad + C_5 \left[t_{<\delta\beta>_3} \partial_\alpha U^\alpha + 2 t^{<\mu\gamma>_3} h_{\gamma<\beta} h_{\delta>_3\gamma} \partial_\mu U^\gamma \right] = -\frac{1}{\tau} C_5 t_{<\delta\beta>_3}. \tag{49}
\end{aligned}$$

73 The system formed by the 15 equations (47)_{1,2,3}, (48), (49) and (47)_{5,6} is a closed systems for the 15 unknown
74 $(\rho, U_\delta, T, \Pi, t_{<\alpha\beta>_3}, q_\delta, \Delta)$.

75 4. Entropy density, Convexity, Entropy Principle and well-posedness of Cauchy problem

76 In this section we evaluate the entropy law and we want to prove that every solutions are entropic with an
77 entropy density that is a convex function.

78 *4.1. Entropy density*

By substituting the distribution function (25) with (36) into (20), we can evaluate the four-dimensional entropy flux. In this procedure, it is needed to be careful concerning the order of the nonequilibrium variables. The present linear constitutive equations is related to the entropy with the second order of the nonequilibrium variables. By taking into account up to the second order in the expansion of the distribution function and of the constitutive equations, we may evaluate as follows

$$h^\alpha = h_E^\alpha + h_{(1)}^\alpha + h_{(2)}^\alpha, \quad (50)$$

where $h_{(1)}^\alpha$ and $h_{(2)}^\alpha$ are, respectively, the contribution of the first and second order terms of the nonequilibrium variables, which can be derived as follows (see Appendix A for details):

$$\begin{aligned} h_{(1)}^\alpha &= -\frac{c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \chi_E \tilde{\chi}_{(1)} \varphi(I) dI d\mathbf{P}, \\ h_{(2)}^\alpha &= -\frac{c}{2k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \tilde{\chi}_{(1)}^2 \varphi(I) dI d\mathbf{P}, \end{aligned} \quad (51)$$

where $\tilde{\chi}_{(1)}$ is $\tilde{\chi}$ defined in (25) with the linear constitutive equations studied in the previous. After cumbersome calculations, we obtain explicit expression of them as follows:

$$\begin{aligned} h_{(1)}^\alpha &= \lambda_E (V^\alpha - V_E^\alpha) + \frac{U_\mu}{T} (T^{\alpha\mu} - T_E^{\alpha\mu}) = \frac{q^\alpha}{T}, \\ h_{(2)}^\alpha &= -\frac{m}{2k_B} \left\{ [(\lambda - \lambda^E)]^2 V_E^\alpha + (\lambda_\mu - \lambda_\mu^E)(\lambda_\nu - \lambda_\nu^E) A_E^{\alpha\mu\nu} + (\lambda_{\mu\nu}) (\lambda_{\psi\theta}) A_E^{\alpha\mu\nu\psi\theta} + \right. \\ &\quad \left. + 2(\lambda - \lambda^E)(\lambda_\mu - \lambda_\mu^E) T_E^{\alpha\mu} + 2(\lambda - \lambda^E)(\lambda_{\mu\nu}) A_E^{\alpha\mu\nu} + 2(\lambda_\theta - \lambda_\theta^E)(\lambda_{\mu\nu}) A_E^{\alpha\theta\mu\nu} \right\} \\ &= -\frac{1}{c^2} U^\alpha \left\{ -\frac{c^2 \alpha_4 C_5}{2} t^{<\mu\nu>_3} t^{<\mu\nu>_3} - \left(c^2 \alpha_3 \frac{N_3}{D_3} + \frac{b_3}{2} \right) q_\mu q^\mu + L_1 \Pi^2 + L_2 \Delta^2 + 2L_3 \Pi \Delta \right\} \\ &\quad + \frac{1}{2} \left(b_1 - b_3 + c^2 \frac{N_3}{D_3} \alpha_1 + \frac{N_{31}}{D_3} \alpha_2 + 2\alpha_3 c^2 \frac{N^\Pi}{D_4} \right) \Pi q^\alpha + \frac{1}{2} \left(b_2 + c^2 \frac{N_3}{D_3} \beta_1 + \frac{N_{31}}{D_3} \beta_2 + \frac{1}{2} \alpha_3 \frac{N^\Delta}{D_4} \right) \Delta q^\alpha \\ &\quad + \frac{1}{2} \left(b_3 + 2c^2 \alpha_3 C_5 - \frac{2}{5} \alpha_4 \frac{N_{31}}{D_3} \right) t^{<\alpha\mu>_3} q_\mu, \end{aligned} \quad (52)$$

where

$$L_1 = \frac{3c^2}{2} \alpha_2 \frac{N^\Pi}{D_4}, \quad L_2 = \frac{1}{8} \left(3\beta_2 \frac{N^\Delta}{D_4} - c^2 \beta_1 \right), \quad L_3 = \frac{1}{4} \left(\frac{3\alpha_2}{4} \frac{N^\Delta}{D_4} + 3c^2 \beta_2 \frac{N^\Pi}{D_4} - \frac{c^2 \alpha_1}{4} \right).$$

In particular, for the entropy density $h = h^\alpha U_\alpha$, we have

$$h = h_E + \frac{c^2 \alpha_4 C_5}{2} t^{<\mu\nu>_3} t^{<\mu\nu>_3} + \left(c^2 \alpha_3 \frac{N_3}{D_3} + \frac{b_3}{2} \right) q_\mu q^\mu - (\Pi \quad \Delta) \begin{pmatrix} L_1 & L_3 \\ L_3 & L_2 \end{pmatrix} \begin{pmatrix} \Pi \\ \Delta \end{pmatrix}. \quad (53)$$

79 We emphasize that the convexity of the entropy density is satisfied because from (52)₁ we have $h_{(1)}^\alpha U_\alpha = 0$ and
80 from (51) we have $h_{(2)}^\alpha U_\alpha < 0$ everywhere and zero only at equilibrium. Therefore the following inequalities are
81 automatically satisfied:

- 82 • $\alpha_4 C_5 < 0$,
83 • $2c^2 \alpha_3 \frac{N_3}{D_3} + b_3 > 0$ because $q_\alpha q^\alpha < 0$,
84 • $L_1 > 0$,
85 • $L_1 L_2 - (L_3)^2 > 0$.

86 **4.2. Entropy production**

According with the theorem proved by Boillat and Ruggeri [17] (see also [1,2]), the procedure of MEP at molecular level it is equivalent to the closure using the entropy principle and the Lagrange multipliers coincide with the *main field* for which the original system become symmetric hyperbolic [2]. Therefore the closed system satisfy the entropy balance law

$$\partial_\alpha h^\alpha = \Sigma, \quad (54)$$

where the entropy four-vector is given by (50), (52). For what concerns the entropy production Σ according with the result of Ruggeri and Strumia [2] this is given by the scalar product between the main field components and the production terms [19]. In the present case, we have

$$\Sigma = I^{\beta\gamma} \lambda_{\beta\gamma}. \quad (55)$$

By using eq. (45) we have

$$\Sigma = \frac{1}{\tau} \left\{ -\frac{1}{4c^4} \Delta U^\beta U^\gamma \lambda_{\beta\gamma} + \left(\frac{1}{4c^2} \frac{N^\Delta}{D_4} \Delta + \frac{N^\Pi}{D_4} \Pi \right) I^{\beta\gamma} \lambda_{\beta\gamma} + \left(-\frac{2}{c^2} \frac{N_3}{D_3} + \frac{\theta_{1,3}}{\theta_{1,2}} \frac{1}{c^2} \right) q^{(\beta} U^{\gamma)} \lambda_{\beta\gamma} - C_5 t^{<\beta\gamma>3} \lambda_{\beta\gamma} \right\}. \quad (56)$$

By substituting eqs. (37)-(39) into eq. (56) and reminding that $q^\beta U^\gamma \lambda_{\beta\gamma} = -q_\alpha h^{\alpha\beta} U^\gamma \lambda_{\beta\gamma}$, we obtain Σ in a quadratic form as follows:

$$\Sigma = \frac{3k_B C_5}{2\tau m c^4 \rho \theta_{2,3}} t^{<\beta\gamma>3} t^{<\beta\gamma>3} + \frac{9k_B \theta_{1,1}}{2\tau m^2 n c^6 D_3} \left(-2 \frac{N_3}{D_3} + \frac{\theta_{1,3}}{\theta_{1,2}} \right) q^\alpha q_\alpha + (\Delta \quad \Pi) \begin{pmatrix} M_1 & M_2 \\ M_2 & M_3 \end{pmatrix} \begin{pmatrix} \Delta \\ \Pi \end{pmatrix}, \quad (57)$$

where

$$M_1 = \frac{k_B}{16 c^8 \tau m^2 n D_4} \left(D_4^{33} + \frac{N^\Delta}{D_4} D_4^{34} \right), \quad M_2 = -\frac{k_B}{4 c^6 \tau m^2 n D_4} \left(D_4^{43} + \frac{N^\Delta}{D_4} D_4^{44} - \frac{N^\Pi}{D_4} D_4^{34} \right),$$

$$M_3 = -\frac{k_B}{c^4 \tau m^2 n D_4} \frac{N^\Pi}{D_4} D_4^{44}.$$

87 The Sylvester criteria allows us to state that the quadratic form is positive definite iff all the following conditions
88 hold:

- 89 1. $\frac{3k_B C_5}{2\tau m c^4 \rho \theta_{2,3}} > 0$.
- 90 2. $\left(-2 \frac{N_3}{D_3} + \frac{\theta_{1,3}}{\theta_{1,2}} \right) \frac{9k_B \theta_{1,1}}{2\tau m^2 n c^6 D_3} < 0$, because $q^\alpha q_\alpha < 0$.
- 91 3. $M_1 > 0$.
- 92 4. $M_1 M_3 - (M_2)^2 > 0$.

The first condition is automatically satisfied because of the definition of the functions involved.

In order to prove the second condition, we can consider a space like vector X^β and the following function that is defined to be positive for each value of X^β .

$$g(X^\beta) = \frac{U_\alpha}{c \tau k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} f_{EP}^\alpha \left[X_\beta p^\beta \left(\frac{\theta_{1,3}}{\theta_{1,2}} \left(1 + \frac{I}{mc^2} \right) - \frac{2}{mc^2} \left(1 + \frac{I}{mc^2} \right)^2 U_\nu p^\nu \right) \right]^2 \phi(I) dI d\mathbf{P}.$$

By exploiting the calculation in the above integral and by using eqs. (27), we have

$$g(X^\beta) = \frac{m^2 n c^2}{\tau k_B} \left[\frac{1}{3} \frac{(\theta_{1,3})^2}{\theta_{1,2}} - \frac{2}{5} \theta_{1,4} \right] X^\beta X_\beta.$$

If we choose, as a particular value,

$$X^\beta = -\frac{1}{D_3} \frac{9k_B}{2m^2 n c^4} \theta_{1,1} q^\beta,$$

we obtain

$$g(X^\beta) = \frac{9k_B \theta_{1,1}}{2\tau m^2 n c^6 D_3} \left(-2 \frac{N_3}{D_3} + \frac{\theta_{1,3}}{\theta_{1,2}} \right) q^\alpha q_\alpha > 0.$$

It proves that also the second condition is satisfied.

Conditions 3 and 4 can be proved showing that they are coefficients of a quadratic form that is definite positive. In order to obtain the entropy production up to the second order, we have to substitute eq. (19)₄ into (55) and take the collisional term (43) up to the first order. Then,

$$\Sigma^{(2)} = \frac{c}{m} \int_{\mathbb{R}^3} \int_0^{+\infty} Q^{(1)} p^\beta p^\gamma \lambda_{\beta\gamma} \left(1 + \frac{I}{mc^2} \right)^2 \phi(I) dI d\mathbf{P},$$

with

$$Q^{(1)} = \frac{f_E}{c^2 \tau k_B} U_\alpha p^\alpha \left[\tilde{\chi} - \frac{k_B}{bmc^2} p^\mu q_\mu \left(1 + \frac{I}{mc^2} \right) \right].$$

If we substitute to $\lambda_{\beta\gamma}$ its expression obtained from eq. (25)₂, we obtain

$$\Sigma^{(2)} = c \int_{\mathbb{R}^3} \int_0^{+\infty} Q^{(1)} \tilde{\chi} \phi(I) dI d\mathbf{P}.$$

In the state where $q^\beta = 0$ and $t^{<\alpha\beta>_3} = 0$ the Lagrange multipliers and the Entropy production assume particular values that we denote with a *, in particular

$$\Sigma^{(2*)} = \frac{c}{m} \int_{\mathbb{R}^3} \int_0^{+\infty} Q^{(1*)} \tilde{\chi}^* \phi(I) dI d\mathbf{P} = \frac{c}{m} \int_{\mathbb{R}^3} \int_0^{+\infty} \frac{f_E}{c^2 \tau k_B} U_\alpha p^\alpha [\tilde{\chi}^*]^2 \phi(I) dI d\mathbf{P},$$

that is clearly a positive quantity. Moreover, we have

$$\Sigma^{(2*)} = I^{\beta\gamma*} \lambda_{\beta\gamma*}$$

that corresponds to the quadratic form

$$\begin{pmatrix} \Delta & \Pi \end{pmatrix} \begin{pmatrix} M_1 & M_2 \\ M_2 & M_3 \end{pmatrix} \begin{pmatrix} \Delta \\ \Pi \end{pmatrix},$$

93 which, therefore, turns to be definite positive. Therefore proved the

94 **Statement 2.** *The entropy density (53) is a convex function and is maximum at equilibrium. The solutions*
 95 *satisfies the entropy principle (54) with an entropy production (57) that is always non negative. According with*
 96 *the general theory of symmetrization given first in covariant formulation in [19], and the equivalence between*
 97 *Lagrange multipliers and main field [17], the closed system is symmetric hyperbolic in the neighborhood of*
 98 *equilibrium if we chose as variables the main field variables (36), with coefficients given in (40), (41), (42) and*
 99 *the Cauchy problem is well posed locally in time.*

100 5. Diatomic gases

The system (47) is very complex, in particular, because it is not simple to evaluate the function $\omega(\gamma)$ that involves two integrals (12)₃ that cannot have analytical expression for a generic polyatomic gas. Taking into account the relations [7]

$$J_{2,1}(\gamma) = \frac{1}{\gamma} K_2(\gamma), \quad J_{2,2}(\gamma) = \frac{1}{\gamma} \left(K_3(\gamma) - \frac{1}{\gamma} K_2(\gamma) \right),$$

where K_n denotes the modified Bessel function, we can rewrite ω given in (12)₃ in terms of the modified Bessel functions [7]:

$$\omega(\gamma) = \frac{1}{\gamma} \left(\frac{\int_0^{+\infty} K_3(\gamma^*) \phi(\mathcal{I}) d\mathcal{I}}{\int_0^{+\infty} \frac{K_2(\gamma^*)}{\gamma^*} \phi(\mathcal{I}) d\mathcal{I}} - 1 \right).$$

Moreover, to calculate the integrals, we need to prescribe the measure $\phi(\mathcal{I})$. In [7], the measure $\phi(\mathcal{I})$ was assumed as

$$\phi(\mathcal{I}) = \mathcal{I}^a, \quad a = \frac{D-5}{2},$$

because it is the one for which the macroscopic internal energy in the classical limit, when $\gamma \rightarrow \infty$, converges to the one of a classical polyatomic gas, where D indicates the degree of freedom of molecule. As was observed by Ruggeri, Xiao and Zhao [25] in the case of $a = 0$, i.e., $D = 5$ corresponding to diatomic gas, the energy e have an explicit expression similar to the monatomic gas:

$$e = p \left(\frac{\gamma K_0(\gamma)}{K_1(\gamma)} + 3 \right).$$

Therefore, from (12), we have

$$\omega_{\text{diat}}(\gamma) = \frac{K_0(\gamma)}{K_1(\gamma)} + \frac{3}{\gamma}.$$

Using the following recurrence formulas of the Bessel functions

$$K_n(\gamma) = \frac{\gamma}{2n} \left(K_{n+1}(\gamma) - K_{n-1}(\gamma) \right), \quad (58)$$

we can express ω in terms of

$$G(\gamma) = \frac{K_3(\gamma)}{K_2(\gamma)}.$$

In fact we can obtain immediately the following expression:

$$\omega_{\text{diat}}(\gamma) = \frac{1}{\gamma} + \frac{\gamma}{\gamma G - 4}, \quad (59)$$

which is a simple function similar to the one of monatomic gas for which we have [3]:

$$\omega_{\text{mono}}(\gamma) = -1 + \gamma G.$$

101 Taking into account that the derivative of Bessel function are known, all coefficients appearing in the differential
102 system (47) can be written explicitly in terms of $G(\gamma)$, by using (59) and the recurrence formula (58). This is
103 simple by using a symbolic calculus like Mathematica®.

104 6. Ultra-relativistic limit

In the ultra-relativistic limit where $\gamma \rightarrow 0$, it was proved in [26] and [27] that the energy converges to

$$e = (\alpha + 1) \frac{nm c^2}{\gamma}, \quad \text{with} \quad \alpha = \begin{cases} 2 & \text{if } a \leq 2 \\ a & \text{if } a \geq 2. \end{cases} \quad (60)$$

This implies

$$\omega_{\text{ultra}} = \frac{(\alpha + 1)}{\gamma}, \quad \text{with} \quad \alpha = \begin{cases} 2 & \text{if } a \leq 2 \\ a & \text{if } a \geq 2. \end{cases} \quad (61)$$

By means of this expression, we can evaluate the coefficients $\theta_{h,j}$ in (17) that become:

$$\begin{aligned} \{\theta_{0,0}, \theta_{0,1}, \theta_{0,2}, \theta_{0,3}, \theta_{0,4}\} &= \\ &\left\{ 1, \frac{\alpha + 1}{\gamma}, \frac{(\alpha + 1)(\alpha + 2)}{\gamma^2}, \frac{(\alpha + 1)(\alpha + 2)(\alpha + 3)}{\gamma^3}, \frac{(\alpha + 1)(\alpha + 2)(\alpha + 3)(\alpha + 4)}{\gamma^4} \right\}, \\ \{\theta_{1,1}, \theta_{1,2}, \theta_{1,3}, \theta_{1,4}\} &= \\ &\left\{ \frac{1}{\gamma}, \frac{3(\alpha + 2)}{\gamma^2}, \frac{6(\alpha + 2)(\alpha + 3)}{\gamma^3}, \frac{10(\alpha + 2)(\alpha + 3)(\alpha + 4)}{\gamma^4} \right\}, \\ \{\theta_{2,3}, \theta_{2,4}\} &= \\ &\left\{ \frac{3(\alpha + 2)}{\gamma^3}, \frac{15(\alpha + 2)(\alpha + 4)}{\gamma^4} \right\}. \end{aligned}$$

It follows that, in the ultra-relativistic limit, we have

$$\frac{N_3}{D_3} = \frac{2(\alpha + 3)}{\gamma}, \quad \frac{N_{31}}{D_3} = \frac{10}{\gamma}, \quad C_5 = \frac{\alpha + 4}{\gamma},$$

and

$$\frac{N^\Pi}{D_4} = -\frac{\alpha + 4}{\gamma}, \quad \frac{N^\Delta}{D_4} = -\frac{1}{\alpha + 1}, \quad (62)$$

105 where the last two equations holds for $\alpha \neq 2$ (i.e., $a \neq 2$). For $a = 2$, the ultra-relativistic limit of $\frac{N^\Pi}{D_4}$ and of $\frac{N^\Delta}{D_4}$
 106 gives the indeterminate form $\left[\frac{0}{0}\right]$. We show (see Appendix B for details) that it can be solved by considering
 107 higher order terms for the energy e , allowing to prove that eqs. (62) are valid also with $a = 2$ and, hence, that the
 108 closure of the present model is continuous with respect to the parameter α , at the ultra-relativistic limit.

109 7. Principal subsystems of ET_{15}

110 For a general system of the hyperbolic system of balance laws, the system with smaller set of the field
 111 equations can be deduced retaining the property that the convexity of the entropy and the positivity of the entropy
 112 production is preserved as a principal subsystem [18]. The principal subsystem is obtained by putting some
 113 components of the main field as a constant, and the corresponding balance laws are deleted.

114 Let us recall the system (18). The balance law of $A^{\alpha\beta\gamma}$ is divided into the trace part $A_\beta^{\alpha\beta}$ and the traceless
 115 part $A^{\alpha<\beta\gamma>}$. As we study below, by deleting the trace part and putting the corresponding component of main
 116 field as zero, we obtain the theory with 14 fields (ET_{14}). On the other hand, by conducting the same procedure
 117 on the traceless part, we obtain the theory with 6 fields (ET_6). It is remarkable that ET_{14} and ET_6 is the same
 118 order in the sense of the principle subsystem differently from the classical case in which the classical ET_6 is a
 119 principal subsystem of classical ET_{14} . Moreover, the relativistic Euler theory is deduced as a principal subsystem
 120 by deleting the balance laws of $A^{\alpha\beta\gamma}$ and putting the corresponding component of main field as zero

121 7.1. ET_{14} : 14 fields theory

The ET_{14} is obtained as a principal subsystem of ET_{15} under the condition $\lambda_\alpha^\alpha = 0$. From (36)₃, this
 condition provides Δ expressed by Π as follows:

$$\Delta^{(14)} = -\frac{c^2\alpha_1 - 3\alpha_2}{c^2\beta_1 - 3\beta_2}\Pi = 4\frac{N_a}{D_a}c^2\Pi, \quad (63)$$

where $N_a = D_4^{44} + D_4^{43}$ and $D_a = D_4^{34} + D_4^{33}$. Then, the independent fields are the following 14 fields; $(\rho, \gamma, \Pi, U^\alpha, q^\alpha, t^{<\alpha\beta>3})$. By deleting the balance equation corresponding to λ_a^α , that is, the one of $A_\beta^{\alpha\beta}$, the present system of the balance equations are the following:

$$\partial_\alpha V^\alpha = 0, \quad \partial_\alpha T^{\alpha\beta} = 0, \quad \partial_\alpha A^{\alpha<\beta\gamma>} = I^{<\beta\gamma>}. \quad (64)$$

With (63), the constitutive equation is modified in this subsystem. For the comparison with the ET₁₄ theory studied in [7], let us denote

$$\frac{N_1^\pi}{D_1^\pi} = -\frac{1}{3} \frac{N_a}{D_a}, \quad \frac{N_{11}^\pi}{D_1^\pi} = \frac{1}{D_4} \left(\frac{N_a}{D_a} N^\Delta + N^\Pi \right).$$

We can prove the following identity:

$$\frac{N_b}{D_a} = -\frac{1}{D_4} \left(\frac{N_a}{D_a} N^\Delta + N^\Pi \right), \quad \text{with } N_b = N^{\Delta 34} + N^{\Delta 33},$$

where $N^{\Delta 33}$ and $N^{\Delta 34}$ are the minor determinants of N^Δ which deletes the third row and third column, and the third row and fourth column, respectively. Then, as a result, instead of (35), the closure for $A^{\alpha\beta\gamma}$ in the present principal subsystem is given by

$$\begin{aligned} A^{\alpha\beta\gamma} = & \left(\rho \theta_{0,2} - \frac{3}{c^2} \frac{N_1^\pi}{D_1^\pi} \Pi \right) U^\alpha U^\beta U^\gamma + \left(\rho c^2 \theta_{1,2} - 3 \frac{N_{11}^\pi}{D_1^\pi} \Pi \right) U^{(\alpha} t^{\beta\gamma)} + \\ & + \frac{3}{c^2} \frac{N_3}{D_3} q^{(\alpha} U^\beta U^\gamma) + \frac{3}{5} \frac{N_{31}}{D_3} q^{(\alpha} t^{\beta\gamma)} + 3 C_5 t^{<\alpha\beta>3} U^\gamma). \end{aligned} \quad (65)$$

122 This result is formally same as the result of [7] (eq. (56) of the paper). However, there are the difference in the
123 coefficients due to the presence of $(mc^2 + \mathcal{I})^n$ instead of $mc^2 + n\mathcal{I}$ in the integrals.

Similarly, we obtain the production term in this principal subsystem as follows:

$$I^{<\beta\gamma>} = -\frac{1}{c^2 \tau} \frac{3N_1^\pi + N_{11}^\pi}{D_1^\pi} \Pi U^{<\beta} U^{\gamma>} + \frac{1}{c^2 \tau} \left(\frac{\theta_{1,3}}{\theta_{1,2}} - 2 \frac{N_3}{D_3} \right) q^{(\beta} U^{\gamma)} - \frac{1}{\tau} C_5 t^{<\beta\gamma>3}. \quad (66)$$

124 This expression (66) is formally the same with the result of [8] (eq (16) of the paper) in [8], except that now we
125 have $\frac{\theta_{1,3}}{\theta_{1,2}}$ instead of $\frac{B_2}{B_4}$ defined in [8] and the difference of the integral in the coefficients is similar with the case
126 for $A^{\alpha\beta\gamma}$.

127 The system (64) is symmetric hyperbolic in the main field $(\lambda, \lambda_\alpha, \lambda_{<\mu\nu>})$ given respectively by (36) with
128 $\Delta = \Delta^{(14)}$ given by (63).

129 7.2. ET₆: 6 fields theory

We consider the principal subsystem with $\lambda_{<\mu\nu>} = \lambda_{\mu\nu} - \frac{1}{4} \lambda_\alpha^\alpha g_{\mu\nu} = 0$, then we have

$$\lambda_{\mu\nu} = \frac{1}{4} \lambda_\alpha^\alpha g_{\mu\nu}. \quad (67)$$

By comparing it with (36), we have

$$\left(\alpha_1 + \frac{\alpha_2}{c^2} \right) \Pi + \left(\beta_1 + \frac{\beta_2}{c^2} \right) \Delta = 0, \quad q_\mu = 0, \quad t_{<\mu\nu>3} = 0.$$

The first equation indicates that, in this principal subsystem, Δ is expressed with Π as follows:

$$\Delta^{(6)} = -\frac{c^2 \alpha_1 + \alpha_2}{c^2 \beta_1 + \beta_2} \Pi = w \Pi \quad (68)$$

where

$$w = 4c^2 \frac{D_4^{44} - 3D_4^{43}}{D_4^{34} - 3D_4^{33}}.$$

130 It should be mentioned that the relation between Δ and Π is different from the case of ET₁₄.

The independent fields are now the 6 fields; $(n, \gamma, U^\alpha, \Pi)$, and the balance equations are the following:

$$\partial_\alpha V^\alpha = 0, \quad \partial_\alpha T^{\alpha\beta} = 0, \quad \partial_\alpha A^{\alpha\beta} = I^\beta_\beta. \quad (69)$$

where the energy-momentum tensor is now given, instead of (22), by

$$T^{\alpha\beta} = \frac{e}{c^2} U^\alpha U^\beta + (p + \Pi) h^{\alpha\beta}. \quad (70)$$

and, from (35),

$$A^{\alpha\beta} = \{\rho c^2 (\theta_{0,2} - \theta_{1,2}) + A_1\} \Pi U^\alpha, \quad (71)$$

where

$$A_1 = -\frac{1}{4c^2} \left\{ \left(1 + 3 \frac{N^\Delta}{D_4} \right) \frac{c^2 \alpha_1 + \alpha_2}{c^2 \beta_1 + \beta_2} - 12c^2 \frac{N^\Pi}{D_4} \right\} = \frac{D_4^{44} - 3D_4^{43} + 3N^{\Delta 34} - 9N^{\Delta 33}}{D_4^{34} - 3D_4^{33}}.$$

Similarly, from (45), we obtain

$$I^\beta_\beta = -\frac{A_1}{\tau} \Pi. \quad (72)$$

The corresponding Lagrange multiplier to $A^{\alpha\beta}$ is $\mu = \frac{1}{4} \lambda_\alpha^\alpha$ which is obtained from (67) as follows:

$$\mu = c^2 \frac{\alpha_1 \beta_2 - \alpha_2 \beta_1}{c^2 \beta_1 + \beta_2} \Pi. \quad (73)$$

The system (69) with (70) and (71) is symmetric hyperbolic in the main field $(\lambda, \lambda_\alpha, \mu)$ given respectively by (see (36)_{1,2}):

$$\lambda = -\frac{g + c^2}{T} + (a_1 + a_2 w) \Pi, \quad \lambda_\alpha = \frac{1}{T} (1 + (b_1 + b_2 w) \Pi) U_\alpha, \quad (74)$$

131 and μ given by (73).

The closed field equations with the material derivative are obtained as follows:

$$\begin{aligned} \dot{\rho} + \rho \partial_\alpha U^\alpha &= 0, \\ \frac{e + p + \Pi}{c^2} \dot{U}_\delta + h_\delta^\mu \partial_\mu (p + \Pi) &= 0, \\ \dot{e} + (e + p + \Pi) \partial_\alpha U^\alpha &= 0, \\ \dot{\Pi} + \frac{\rho c^2 (\theta'_{0,2} - \theta'_{1,2})}{A_1} \dot{\gamma} + \frac{\dot{A}_1}{A_1} \Pi + \Pi \partial_\alpha U^\alpha &= -\frac{\Pi}{\tau}. \end{aligned} \quad (75)$$

Taking into account

$$h_\delta^\mu \partial_\mu (p + \Pi) = U_\delta \frac{\dot{p} + \dot{\Pi}}{c^2} - \partial_\delta (p + \Pi), \quad (76)$$

and from (12):

$$\dot{e} = c^2 (\dot{\rho} \omega + \rho \omega' \dot{\gamma}), \quad \dot{p} = \frac{c^2}{\gamma^2} (\gamma \dot{\rho} - \rho \dot{\gamma}), \quad (77)$$

the system (75) can be put in the normal form:

$$\begin{aligned} \dot{\rho} + \rho \partial_\alpha U^\alpha &= 0, \\ \left(\rho + \frac{\rho \mathcal{E} + p + \Pi}{c^2} \right) \dot{U}_\delta - \partial_\delta(p + \Pi) - \frac{(p + \Pi)}{c^2} \left[1 - \frac{1}{A_1 \omega'} \left(A_1' \frac{\Pi}{\rho c^2} + \frac{A_1}{\gamma^2} + \theta'_{0,2} - \theta'_{1,2} \right) \right] U_\delta \partial_\alpha U^\alpha &= \frac{\Pi}{\tau c^2} U_\delta, \\ \rho c^2 \omega' \dot{\gamma} + (p + \Pi) \partial_\alpha U^\alpha &= 0, \\ \dot{\Pi} + \left\{ \Pi - \frac{p + \Pi}{\rho c^2 A_1 \omega'} \left[A_1' \Pi + \rho c^2 (\theta'_{0,2} - \theta'_{1,2}) \right] \right\} \partial_\alpha U^\alpha &= -\frac{\Pi}{\tau}. \end{aligned} \quad (78)$$

132 It is extremely interesting that in the relativistic theory the acceleration is influenced by the relaxation time trough
133 the right hand side of (78)₂ and this may be important for the application to the problems of cosmology.

134 7.3. *ET*₅: Euler 5 fields theory

Let us consider the principal subsystem with $\lambda_{\mu\nu} = 0$. This indicates that any nonequilibrium variables are set to be zero, i.e.,

$$\Pi = \Delta = 0, \quad t_{\langle \mu\nu \rangle_3} = 0, \quad q_\alpha = 0. \quad (79)$$

The independent fields are the 5 fields (n, U^α, γ) , and the balance equations are

$$\partial_\alpha V^\alpha = 0, \quad \partial_\alpha T^{\alpha\beta} = 0, \quad (80)$$

with

$$T^{\alpha\beta} = \frac{e}{c^2} U^\alpha U^\beta + p h^{\alpha\beta}. \quad (81)$$

135 The deduced system is the one of the relativistic Euler theory and the system (80) become symmetric in the main
136 field ($\lambda = -(g + c^2)/T$, $\lambda_\alpha = U_\alpha/T$) obtained first by Ruggeri and Strumia in [19].

137 8. Maxwellian Iteration and phenomenological coefficients

138 In order to find the parabolic limit of system (47) and to obtain the corresponding Eckart equations, we
139 adopt the Maxwellian iteration [28] on (47) in which only the first order terms with respect to the relaxation time
140 are retained. The phenomenological coefficients, that is, the heat conductivity χ , the shear viscosity μ and the
141 bulk viscosity ν are identified with the relaxation time.

The method of the Maxwellian iteration is based on putting to zero the nonequilibrium variables on the left side of equations (47):

$$\begin{aligned} \dot{\rho} - \rho h^{\beta\alpha} \partial_\beta U_\alpha &= 0, \\ \frac{e + p}{c^2} h_{\delta\beta} \dot{U}^\beta - h_\delta^\mu \partial_\mu p &= 0, \\ \dot{e} - (e + p) h_\alpha^\mu \partial_\mu U^\alpha &= 0, \\ \frac{c^2}{3} h_{\delta\beta} \left(\dot{\rho} \theta_{1,2} + \rho \theta'_{1,2} \dot{\gamma} \right) - \frac{1}{3} \rho c^2 \theta_{1,2} \left[h_{\delta\beta} h_\alpha^\mu \partial_\mu U^\alpha + 2 h_{\theta(\delta} h_{\beta)}^\mu \partial_\mu U^\theta \right] &= \\ &= \frac{1}{\tau} \left(\frac{1}{4} \frac{N^\Delta}{D_4} \frac{\Delta}{c^2} + \frac{N^\Pi}{D_4} \Pi \right) h_{\delta\beta} - \frac{1}{\tau} C_5 t_{\langle \delta\beta \rangle_3}, \\ h_{\delta\beta} \dot{U}^\beta \left(\rho \theta_{0,2} c^2 + 2 \frac{1}{3} \rho c^2 \theta_{1,2} \right) - h_\delta^\mu c^2 \partial_\mu \frac{1}{3} \rho c^2 \theta_{1,2} &= \frac{1}{\tau} \left(\frac{N_3}{D_3} - \frac{\theta_{1,3}}{2 \theta_{1,2}} \right) q_\delta, \\ c^4 \left(\dot{\rho} \theta_{0,2} + \rho \theta'_{0,2} \dot{\gamma} \right) - \rho c^4 \left(\theta_{0,2} + \frac{2}{3} \theta_{1,2} \right) h_\alpha^\mu \partial_\mu U^\alpha &= -\frac{1}{4\tau} \Delta. \end{aligned} \quad (82)$$

From the first three equations of (82) and taking into account $p = \rho c^2 / \gamma$, $e = \rho c^2 \omega(\gamma)$ (see (12)), we can deduce

$$\dot{\rho} = \rho h^{\mu\alpha} \partial_\mu U_\alpha, \quad h_\delta^\mu \partial_\mu \rho = \rho \frac{\omega\gamma + 1}{c^2} h_{\delta\beta} U^\mu \partial_\mu U^\beta + \frac{\rho}{\gamma} h_\delta^\mu \partial_\mu \gamma, \quad \dot{\gamma} = \frac{1}{\gamma\omega'} h^{\mu\alpha} \partial_\mu U_\alpha. \quad (83)$$

Putting (83) in the remaining equations (82)_{4,5,6}, we obtain the solution

$$\begin{aligned} q_\beta &= -\chi h_\beta^\alpha \left[\partial_\alpha T - \frac{T}{c^2} U^\mu \partial_\mu U^\alpha \right], \\ \Pi &= -\nu \partial_\alpha U^\alpha, \\ t_{\langle\beta\delta\rangle_3} &= 2\mu h_\beta^\alpha h_\delta^\mu \partial_{\langle\alpha} U_{\mu\rangle}, \\ \Delta &= \sigma \partial_\alpha U^\alpha, \end{aligned} \quad (84)$$

with

$$\begin{aligned} \chi &= -\frac{2\rho c^2}{3B^q T} [3\theta_{0,2} + \theta_{1,2}(1 - \omega\gamma)], \\ \nu &= -\frac{\rho c^2}{3B_2^\Pi} \left\{ \frac{2}{3}\theta_{1,2} - \frac{\theta'_{1,2}}{\gamma\omega'} + 3\frac{N^\Delta}{D_4} \left(\frac{2}{3}\theta_{1,2} - \frac{\theta'_{0,2}}{\gamma\omega'} \right) \right\}, \\ \mu &= -\frac{\rho c^2}{3B^t} \theta_{1,2}, \end{aligned} \quad (85)$$

and

$$\sigma = \frac{\rho}{B_1^\Delta} \left(\frac{2}{3}\theta_{1,2} - \frac{\theta'_{0,2}}{\gamma\omega'} \right),$$

142 where B_2^Π , B^q , B^t are explicitly given by (44) with the relaxation time τ .

143 As the first three equations in (84) are the Eckart equations, we deduce that χ , ν , μ are respectively the heat
144 conductivity, the bulk and shear viscosities. In addition, we have a new phenomenological coefficient σ but as Δ
145 doesn't appear neither in V^α and $T^{\alpha\beta}$ (see the equation (22) or the first three equations in (47)), we arrive to the
146 conclusion that the present theory converges to the Eckart one formed in the first three block equations of (47)
147 with constitutive equations (84) in which the heat conductivity, bulk viscosity and shear viscosity are explicitly
148 given by (85)_{1,2,3}.

We introduce as in [9] the dimensionless variables as follows:

$$\begin{aligned} \bar{\chi} &= \frac{\rho T \chi}{p^2 \tau} = -\frac{2}{3} \gamma^2 \frac{3\theta_{0,2} + \theta_{1,2}(1 - \omega\gamma)}{\frac{\theta_{1,3}}{\theta_{1,2}} - 2\frac{N_3}{D_3}}, \\ \bar{\nu} &= \frac{\nu}{p\tau} = -\frac{1}{3} \frac{\gamma}{\frac{N^\Pi}{D_4}} \left\{ \frac{2}{3}\theta_{1,2} - \frac{\theta'_{1,2}}{\gamma\omega'} + 3\frac{N^\Delta}{D_4} \left(\frac{2}{3}\theta_{1,2} - \frac{\theta'_{0,2}}{\gamma\omega'} \right) \right\}, \\ \bar{\mu} &= \frac{\mu}{p\tau} = \frac{\gamma}{3C_5} \theta_{1,2}, \end{aligned} \quad (86)$$

149 that are functions only of γ .

150 8.1. Ultra-relativistic and classical limit of the phenomenological coefficients

Taking into account eqs. (61) - (62), it is simple to obtain the limit of (86) when $\gamma \rightarrow 0$:

$$\bar{\chi}_{\text{ultra}} = 0, \quad \bar{\nu}_{\text{ultra}} = \frac{2}{3} \frac{\alpha^2 - 4}{(1 + \alpha)(4 + \alpha)}, \quad \bar{\mu}_{\text{ultra}} = \frac{2 + \alpha}{4 + \alpha}.$$

In particular in the most significant case in which $a \leq 2$ for which $\alpha = 2$ we have

$$\bar{\chi}_{\text{ultra}} = 0, \quad \bar{v}_{\text{ultra}} = 0, \quad \bar{\mu}_{\text{ultra}} = \frac{2}{3}. \quad (87)$$

Instead, in the classical limit for which $\gamma \rightarrow \infty$, it was proved in [7] that the internal energy ε converges to the classical internal energy of polytropic gas: $\varepsilon = (D/2)(k_B/m)T$. Therefore, from (13), ω converges to

$$\omega_{\text{class}} = 1 + \frac{D}{2\gamma}. \quad (88)$$

In the present case, using (88), it is not difficult to find $\theta_{h,j}$ deduced in (17) in the limit $\gamma \rightarrow \infty$ as follows:

$$\begin{aligned} \{\theta_{0,0}, \theta_{0,1}, \theta_{0,2}, \theta_{0,3}, \theta_{0,4}\} &= \left\{1, 1 + \frac{D}{2\gamma}, 1 + \frac{D}{\gamma}, 1 + \frac{3D}{2\gamma}, 1 + \frac{2D}{\gamma}\right\}, \\ \{\theta_{1,1}, \theta_{1,2}, \theta_{1,3}, \theta_{1,4}\} &= \left\{\frac{1}{\gamma}, \frac{3}{\gamma}, \frac{6}{\gamma}, \frac{10}{\gamma}\right\}, \\ \{\theta_{2,3}, \theta_{2,4}\} &= \left\{\frac{3}{\gamma^2}, \frac{15}{\gamma^2}\right\}. \end{aligned} \quad (89)$$

Therefore, in the classical limit, we have

$$\frac{N_3}{D_3} = 2, \quad \frac{N_{31}}{D_3} = \frac{10}{2+D}, \quad C_5 = 1, \quad \frac{N^\Pi}{D_4} = -1, \quad \frac{N^\Delta}{D_4} = -\frac{2}{D}, \quad (90)$$

and we find from (86)

$$\bar{\chi}_{\text{class}} = \frac{D+2}{2}, \quad \bar{v}_{\text{class}} = \frac{2(D-3)}{3D}, \quad \bar{\mu}_{\text{class}} = 1, \quad (91)$$

151 which are in perfect agreement with the phenomenological coefficients of the classical RET theory [2].

152 8.2. Phenomenological coefficients in ET_{14} and ET_6

By conducting the Maxwellian iteration to ET_{14} as a principal subsystems of ET_{15} , we may expect that the different bulk viscosity appears. This is because Δ is related to Π by (63) and it affects the balance laws corresponding to Π in ET_{14} . In fact, from (65) and (66), we can obtain the closed field equations for Π , and then, through the Maxwellian iteration as it has been done in [9], we obtain the bulk viscosity for ET_{14} as follows:

$$\bar{v}_{14} = \frac{\frac{1}{\omega'}(\theta'_{0,2} + \frac{1}{3}\theta'_{1,2}) - \frac{8}{9}\gamma\theta_{1,2}}{\left(-1 + \frac{N^\Delta}{D_4}\right)\frac{N_a}{D_a} + \frac{N^\Pi}{D_4}}. \quad (92)$$

153 We remark that the heat conductivity and the shear viscosity is the same between ET_{15} and ET_{14} .

Similarly, from (78)₄, we obtain the bulk viscosity estimated by ET_6 as follows:

$$\bar{v}_6 = -\frac{\theta'_{0,2} - \theta'_{1,2}}{\omega'A_1}. \quad (93)$$

154 It should be noted that, in the classical case studied in [14], the bulk viscosities of ET_{15} , ET_{14} and ET_6
155 are the same. In fact, in the classical limit, \bar{v}_{14} and \bar{v}_6 coincides with \bar{v}_{class} . However, due to the mathematical
156 structure of the relativity, i.e., the scalar fields Π and Δ appear together in the triple tensor, the method of the
157 principal subsystem dictates the difference of the subsystems.

158 8.3. Heat conductivity, Bulk viscosity and Shear viscosity in diatomic gases

159 Inserting (59), after cumbersome calculations (easy with Mathematica), we can obtain the phenomenological
160 coefficients in the diatomic case:

$$\bar{\kappa} = -\frac{\gamma(\gamma^2 + 2\gamma G - 8) \left\{ \gamma^4(G^2 - 1) + 2\gamma^2(G^2 + 2) - 5\gamma^3 G - 16\gamma G + 32 \right\}^2}{(\gamma G - 4)^3 \left\{ \gamma \left[-\gamma^5 + 5\gamma^3 + 48\gamma + (\gamma^4 - 6\gamma^2 - 12)\gamma G^2 + (-5\gamma^4 + 12\gamma^2 + 96)G \right] - 192 \right\}},$$

$$\bar{\mu} = \frac{(\gamma^2 + 2\gamma G - 8)^2}{(\gamma G - 4) \left\{ 4(\gamma^2 - 8) + \gamma(\gamma^2 + 8)G \right\}},$$

$$\bar{\nu} = \frac{g_1}{3(\gamma G - 4)g_2},$$

with

$$g_1 = 4\gamma^{15}G(G^2 - 1)^2 + 81920\gamma^3G(7G^2 + 20) - 196608\gamma^2(7G^2 + 4) + 1024\gamma^5G(21G^4 + 660G^2 - 392) -$$

$$4096\gamma^4(35G^4 + 348G^2 - 56) + 4\gamma^{14}(G^6 - 17G^4 + 21G^2 - 5) + \gamma^{13}G(7G^6 - 86G^4 + 435G^2 - 256) +$$

$$4\gamma^{12}(-40G^6 + 193G^4 - 331G^2 + 48) + 4\gamma^{11}G(-14G^6 + 422G^4 - 943G^2 + 500) + 16\gamma^{10}(77G^6 - 660G^4 + 677G^2 - 84) +$$

$$16\gamma^9G(7G^6 - 714G^4 + 2560G^2 - 1108) - 64\gamma^8(45G^6 - 910G^4 + 1472G^2 - 204) +$$

$$64\gamma^7G(G^6 + 492G^4 - 2800G^2 + 1760) - 256\gamma^6(7G^6 + 740G^4 - 1344G^2 + 192) + 1835008\gamma G - 1048576,$$

$$g_2 = \gamma^4(G^2 - 1) + \gamma^2(G^2 + 4) - 5\gamma^3G - 8\gamma G + 16 \left(\gamma(2\gamma^9G^2(G^2 - 1) + 5\gamma^8G(1 - 3G^2)) + \right.$$

$$40\gamma^6G(6 - 5G^2) + 64\gamma^4G(11G^2 - 25) + 512\gamma^2G(G^2 + 14) - 1024\gamma(3G^2 + 5) +$$

$$\left. \gamma^7(19G^4 - 17G^2 + 28) - 4\gamma^5(13G^4 - 198G^2 + 60) - 32\gamma^3(G^4 + 108G^2 - 52) + 8192G \right) - 8192.$$

161

162

163

164

165

166

167

168

169

170

Let us compare the phenomenological coefficients with the ones for the monatomic case obtained in [9]. In Fig. 1, we plot the dependence of the dimensionless heat conductivity and shear viscosity on γ for both diatomic and monatomic cases. Concerning ν , we also plot the dimensionless bulk viscosity of ET₁₄ derived in (92) in the Fig. 2. We observe that in the ultra-relativistic limit and in classical limit, the figures are in perfect agreement with the limits (87) and (91) (for $D = 3, 5$). We remark, as it is evidently shown in Fig. 2, how small is the bulk viscosity in monatomic gas with respect to the one of diatomic case.

It is also remarkable that the value of the bulk viscosity of ET₆ given by (93) is quite near to the one of ET₁₅. For this reason, we omit the plot of $\bar{\nu}^{(6)}$ in the figure. This indicates that ET₆ captures the effect of the dynamic pressure in consistent with ET₁₅.

171

9. Classic limit of the relativistic theory

We want to perform now the classical limit $\gamma \rightarrow \infty$ of the closed relativistic system (47). For this purpose we recall the limits of the coefficients given in (89) and (90). Moreover, taking into account the decomposition $U^\alpha \equiv (\Gamma c, v^i)$, where Γ is the Lorentz factor, we have $\partial_\alpha U^\alpha = \frac{1}{c} \partial_t(\Gamma c) + \partial_k(\Gamma v^k)$ whose limit is $\partial_t v^i$ because $\partial_t \Gamma = -\Gamma^3 \frac{v_i}{c^2} \partial_t v^i$ has zero limit, and the similar evaluation applies to $\partial_k \Gamma$. Then,

$$\frac{1}{c^2} U^\mu \partial_\mu U^0 = \frac{1}{c^2} \Gamma c \frac{1}{c} \partial_t(\Gamma c) + \frac{1}{c^2} \Gamma v^k \partial_k(\Gamma c) \quad \text{has 0 limit,}$$

$$\frac{1}{c^2} U^\mu \partial_\mu U^i = \frac{1}{c^2} \Gamma c \frac{1}{c} \partial_t(\Gamma v^i) + \frac{1}{c^2} \Gamma v^k \partial_k(\Gamma v^i) \quad \text{has 0 limit.}$$




Figure 1. Dependence of $\bar{\chi}$ (left) and $\bar{\mu}$ (right) for diatomic (red solid line) and monatomic (black dashed line) gases on γ . The dotted line indicates the corresponding value in the classical limit. In the ultra-relativistic limit ($\gamma \rightarrow 0$), $\bar{\chi}_{\text{ultra}} = 0, \bar{\mu}_{\text{ultra}} = 2/3$ both for monatomic and diatomic gases. In the classical limit ($\gamma \rightarrow \infty$), $\bar{\chi}_{\text{class}} = 2.5, \bar{\mu}_{\text{class}} = 1$ for monatomic gas, and $\bar{\chi}_{\text{class}} = 3.5, \bar{\mu}_{\text{class}} = 1$ for diatomic gas.




Figure 2. Dependence of \bar{v} for diatomic (red solid line) monatomic (black dashed line) gases on γ . The prediction by ET_{14} as a principal subsystem of ET_{15} is also shown with the dotted line. In the ultra-relativistic limit ($\gamma \rightarrow 0$), $\bar{v}_{\text{ultra}} = 0$ both for monatomic and diatomic gases. In the classical limit ($\gamma \rightarrow \infty$), $\bar{v}_{\text{class}} = 0$ for monatomic gas, and $\bar{v}_{\text{class}} = 4/15$ for diatomic gas.

Concerning the projection operator in the limit, it is necessary to remind that, with our choice of the metric, $v_j = -v^j$, then

$$h^{\beta\alpha} = -g^{\beta\alpha} + \frac{U^\beta U^\alpha}{c^2} \quad \rightarrow \quad h^{ij} = -g^{ij} + \Gamma^2 \frac{v^i v^j}{c^2} \quad \rightarrow \quad \lim_{c \rightarrow +\infty} h^{ij} = -g^{ij} = \delta^{ij},$$

$$\lim_{c \rightarrow +\infty} h_j^i = -g_j^i = -\text{diag}(1, 1, 1).$$

While from

$$0 = U_\alpha h^{i\alpha} = \Gamma c h^{i0} + \Gamma v_k h^{ik} \quad \rightarrow \quad h^{i0} = -\frac{v_k}{c} h^{ik},$$

$$0 = U_\alpha h^{0\alpha} = \Gamma c h^{00} + \Gamma v_k h^{0k} \quad \rightarrow \quad h^{00} = -\frac{v_k}{c} h^{0k} = -\frac{v_a v_b}{c^2} h^{ab}.$$

¹⁷² The last two relations hold also without taking the non relativistic limit. As a consequence, we have that
¹⁷³ $\lim_{c \rightarrow +\infty} h^{i0} = 0$ and $\lim_{c \rightarrow +\infty} h^{00} = 0$.

The relativistic material derivative (46) of a function f converges to the classical material derivative where we continue to indicate it with a dot. Then, the system (47) becomes in the classical limit:

$$\begin{aligned}
\dot{\rho} + \rho \frac{\partial v_l}{\partial x_l} &= 0, \\
\rho \dot{v}_i + \frac{\partial p}{\partial x_i} + \frac{\partial \Pi}{\partial x_i} - \frac{\partial \sigma_{\langle ik \rangle}}{\partial x_k} &= 0, \\
\dot{T} + \frac{2T}{Dp} \left\{ (p + \Pi) \frac{\partial v_l}{\partial x_l} - \sigma_{\langle ik \rangle} \frac{\partial v_k}{\partial x_i} + \frac{\partial q_l}{\partial x_l} \right\} &= 0, \\
\dot{\Pi} + \frac{2}{3} \frac{D-3}{D} p \frac{\partial v_l}{\partial x_l} + \frac{5D-6}{3D} \Pi \frac{\partial v_l}{\partial x_l} - \frac{2}{3} \frac{D-3}{D} \sigma_{\langle ik \rangle} \frac{\partial v_l}{\partial x_k} + \frac{4(D-3)}{3D(D+2)} \frac{\partial q_l}{\partial x_l} &= -\frac{1}{\tau} \Pi, \\
\dot{\sigma}_{\langle ij \rangle} + \sigma_{\langle ij \rangle} \frac{\partial v_l}{\partial x_l} + 2\sigma_{\langle l(ij)} \frac{\partial v_j \rangle}{\partial x_l} - 2(p + \Pi) \frac{\partial v_{\langle j}}{\partial x_i \rangle} - \frac{4}{D+2} \frac{\partial q_{\langle i}}{\partial x_j \rangle} &= -\frac{1}{\tau} \sigma_{\langle ij \rangle}, \\
\dot{q}_i + \frac{D+4}{D+2} q_i \frac{\partial v_l}{\partial x_l} + \frac{D+4}{D+2} q_l \frac{\partial v_i}{\partial x_l} + \frac{2}{D+2} q_l \frac{\partial v_l}{\partial x_i} \\
+ \frac{D+2}{2} \frac{p}{\rho T} \left\{ (p + \Pi) \delta_{il} - \sigma_{\langle il \rangle} \right\} \frac{\partial T}{\partial x_l} - \frac{p}{\rho^2} (\Pi \delta_{il} - \sigma_{\langle il \rangle}) \frac{\partial \rho}{\partial x_l} \\
+ \frac{1}{\rho} \left\{ (p - \Pi) \delta_{il} + \sigma_{\langle il \rangle} \right\} \left(\frac{\partial \Pi}{\partial x_l} - \frac{\partial \sigma_{\langle rl \rangle}}{\partial x_r} \right) + \frac{1}{2D} \frac{\partial \Delta}{\partial x_i} &= -\frac{1}{\tau} q_i, \\
\dot{\Delta} + \left(\frac{D+4}{D} \Delta + 8 \frac{p}{\rho} \Pi \right) \frac{\partial v_l}{\partial x_l} - 8 \frac{p}{\rho} \sigma_{\langle ik \rangle} \frac{\partial v_i}{\partial x_k} - \frac{8}{\rho} q_i \frac{\partial p}{\partial x_i} \\
+ 4(D+4) \frac{p}{\rho T} q_l \frac{\partial T}{\partial x_l} + \frac{8p}{\rho} \frac{\partial q_l}{\partial x_l} - \frac{8}{\rho} q_i \frac{\partial \Pi}{\partial x_i} + \frac{8}{\rho} q_i \frac{\partial \sigma_{\langle il \rangle}}{\partial x_l} &= -\frac{1}{\tau} \Delta,
\end{aligned} \tag{94}$$

174 where $\sigma_{\langle ij \rangle} = -t_{\langle ij \rangle}$. The system (94) coincides perfectly with the one obtained recently in [14].

175 We remark that, as it has been studied in [14], for classical polytropic gases, ET₁₄ is derived as a principal
176 subsystem of ET₁₅ by setting $\Delta = 0$. Moreover, ET₆ is derived from ET₁₄ as a principal subsystem of ET₁₄ by
177 setting $\sigma_{\langle ij \rangle} = 0$ and $q_i = 0$. This fact corresponds that, in the classical limit, both of $\Delta^{(14)}$ defined in (63) and
178 $\Delta^{(6)}$ defined in (68) become zero.

179 **Acknowledgments:** The work has been partially supported by JSPS KAKENHI Grant Numbers JP18K13471 (TA), by the
180 Italian MIUR through the PRIN2017 project "Multiscale phenomena in Continuum Mechanics: singular limits, off-equilibrium
181 and transitions" Project Number: 2017YBKNCE (SP) and GNFM/INdAM (MCC, SP and TR).

182 **Conflicts of Interest:** The authors declare no conflict of interest.

183 Appendix A. Entropy-entropy flux density

In order to evaluate the entropy density from eq. (20), we need the expression of $f \ln f$ up to the second order with respect to the nonequilibrium variables.

The expansion of the distribution function around an equilibrium state is

$$\begin{aligned}
f &= f_E e^{\frac{-1}{k_B} \tilde{\chi}} = f_E \left[1 - \frac{1}{k_B} \tilde{\chi} + \frac{1}{2k_B^2} (\tilde{\chi})^2 + (\tilde{\chi})^3 (\dots) \right], \\
\text{with } \tilde{\chi} &= (\tilde{\chi})^{(1)} + (\tilde{\chi})^{(2)} + (\tilde{\chi})^{(3)} (\dots),
\end{aligned}$$

defined in (25)₂ and the notation $\eta^{(i)}$ represents the homogeneous part of the generic quantity η at the order i with respect to the nonequilibrium variables. With this notation, the quantities $(\lambda - \lambda^E)^{(1)}$, $(\lambda_\beta - \lambda_\beta^E)^{(1)}$, $(\lambda_{\beta\gamma})^{(1)}$ are those of eq. (36).

By composing the above expressions, we see that the distribution function up to the second order is

$$f = f_E \left\{ 1 - \frac{1}{k_B} \left[(\tilde{\chi})^{(1)} + (\tilde{\chi})^{(2)} \right] + \frac{1}{2k_B^2} \left[(\tilde{\chi})^{(1)} \right]^2 \right\}.$$

and

$$\begin{aligned} f \ln f &= f \left(-1 - \frac{1}{k_B} \tilde{\chi} \right) = f_E \left\{ 1 - \frac{1}{k_B} \left[(\tilde{\chi})^{(1)} + (\tilde{\chi})^{(2)} \right] + \frac{1}{2k_B^2} \left[(\tilde{\chi})^{(1)} \right]^2 + \dots \right\} \\ &\quad \left\{ -1 - \frac{1}{k_B} \chi_E - \frac{1}{k_B} \left[(\tilde{\chi})^{(1)} + (\tilde{\chi})^{(2)} + \dots \right] \right\} \\ &= f_E \ln f_E + \frac{1}{k_B^2} f_E \chi_E (\tilde{\chi})^{(1)} + \frac{1}{k_B^2} f_E \chi_E \left\{ (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} + \\ &\quad + \frac{1}{2k_B^2} f_E \left[(\tilde{\chi})^{(1)} \right]^2. \end{aligned}$$

It follows that

$$h^\alpha = -k_B c \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f \ln f \varphi(I) dI d\mathbf{P} = h_E^\alpha + h_{(1)}^\alpha + h_{(2)}^\alpha,$$

where

$$\begin{aligned} h_{(1)}^\alpha &= -\frac{c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \chi_E (\tilde{\chi})^{(1)} \varphi(I) dI d\mathbf{P} = \\ &= -\frac{c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \left[m \lambda_E + \left(1 + \frac{I}{m c^2} \right) \frac{U_\mu}{T} p^\mu \right] (\tilde{\chi})^{(1)} \varphi(I) dI d\mathbf{P}, \\ h_{(2)}^\alpha &= -\frac{c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \chi_E \left\{ (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} \varphi(I) dI d\mathbf{P} - \\ &\quad \frac{c}{2k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \left[(\tilde{\chi})^{(1)} \right]^2 \varphi(I) dI d\mathbf{P}. \end{aligned}$$

Moreover, we have that the moments appearing in system (18) up to the second order as follows:

$$\begin{aligned} \underline{V^\alpha} &= \underline{V_E^\alpha} - \frac{m c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \left\{ (\tilde{\chi})^{(1)} + (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} \varphi(I) dI d\mathbf{P}, \\ \underline{T^{\alpha\beta}} &= \underline{T_E^{\alpha\beta}} - \frac{c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha p^\beta \left(1 + \frac{I}{m c^2} \right) f_E \left\{ (\tilde{\chi})^{(1)} + (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} \varphi(I) dI d\mathbf{P}, \\ \underline{\frac{U_\alpha U_\beta U_\gamma}{c^4} A^{\alpha\beta\gamma}} &= \underline{\frac{U_\alpha U_\beta U_\gamma}{c^4} A_E^{\alpha\beta\gamma}} - \\ &\quad \underline{\frac{U_\alpha U_\beta U_\gamma}{c^4} \frac{c}{m k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha p^\beta p^\gamma \left(1 + \frac{I}{m c^2} \right)^2 f_E \left\{ (\tilde{\chi})^{(1)} + (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} \varphi(I) dI d\mathbf{P}}. \end{aligned}$$

The underlined terms gives 0 for eqs. (36) and there remain

$$\begin{aligned} & - \frac{m c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \left\{ (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} \varphi(I) dI d\mathbf{P} = 0, \\ & - \frac{c}{k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha p^\beta \left(1 + \frac{I}{m c^2} \right) f_E \left\{ (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} \varphi(I) dI d\mathbf{P} = 0, \\ & \underline{\frac{U_\alpha U_\beta U_\gamma}{c^4} \frac{c}{m k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha p^\beta p^\gamma \left(1 + \frac{I}{m c^2} \right)^2 f_E \left\{ (\tilde{\chi})^{(2)} - \frac{1}{2k_B} \left[(\tilde{\chi})^{(1)} \right]^2 \right\} \varphi(I) dI d\mathbf{P} = 0}. \end{aligned}$$

The first two of them allows prove eq. (52)₁ and to write

$$h_{(2)}^\alpha = -\frac{c}{2k_B} \int_{\mathbb{R}^3} \int_0^{+\infty} p^\alpha f_E \left[(\tilde{\chi})^{(1)} \right]^2 \varphi(I) dI d\mathbf{P}.$$

184 It is sufficient to substitute the expression of $\tilde{\chi}$ to obtain eq. (52)₂

185 **Appendix B. Continuity of the ultra relativistic limit for $a = 2$**

From (12)₂, and by using the recurrence relations (11) and (10), we have

$$\frac{\gamma e}{n m c^2} = \frac{\gamma \int_0^{+\infty} J_{2,2}^* \left(1 + \frac{I}{m c^2}\right) \phi(I) dI}{\int_0^{+\infty} J_{2,1}^* \phi(I) dI} = 3 + \frac{\int_0^{+\infty} J_{0,1}^* \phi(I) dI}{\int_0^{+\infty} J_{2,1}^* \phi(I) dI}.$$

By introducing the Ruggeri's numbers R_k and using eq. (32)₁ of [27] we have

$$\frac{\gamma e}{n m c^2} = 3 + \frac{3}{-\ln \gamma} R_{-4} = 3 - \frac{1}{\ln \gamma}$$

or

$$e = \frac{n m c^2}{\gamma} \left(3 - \frac{1}{\ln \gamma}\right). \quad (\text{A1})$$

186 So we have to calculate D_4 , N^{II} and N^Δ with (A1) instead of (60).

In particular, for D_4 we can add to its fourth line the second one pre-multiplied by $-\frac{1}{3}$ so that it becomes

$$\frac{1}{\gamma \ln \gamma} \left(\frac{1}{3}, \frac{4}{3\gamma}, \frac{20}{3\gamma^2}, \frac{4}{3} \frac{c^2}{\gamma^2}\right).$$

It follows that, after cumbersome calculations that we do not report here for brevity, we have

$$\lim_{\gamma \rightarrow 0} \gamma^9 \ln \gamma D_4 = \begin{vmatrix} 1 & 3 & 12 & 4 \\ 3 & 12 & 60 & 20 \\ 12 & 60 & 360 & 120 \\ \frac{1}{3} & \frac{4}{3} & \frac{20}{3} & \frac{4}{3} \end{vmatrix} = -64.$$

Similarly, for N^{II} we can add to its fourth line the third one multiplied by $-\frac{1}{3}$ so that it becomes

$$\frac{1}{\gamma^2 \ln \gamma} \left(\frac{4}{3}, \frac{20}{3\gamma}, \frac{40}{\gamma^2}, \frac{8}{\gamma^2} c^2\right).$$

It follows that

$$\lim_{\gamma \rightarrow 0} \gamma^{10} \ln \gamma N^{\text{II}} = - \begin{vmatrix} 1 & 3 & 12 & 4 \\ 3 & 12 & 60 & 20 \\ 12 & 60 & 360 & 120 \\ \frac{4}{3} & \frac{20}{3} & 40 & 8 \end{vmatrix} = 384.$$

Finally, for N^Δ we can add to its third line the second one multiplied by $-\frac{1}{3}$ so that its third line becomes

$$\frac{1}{\gamma \ln \gamma} \left(\frac{1}{3}, \frac{4}{3\gamma}, \frac{20}{3\gamma^2}, \frac{4}{3} \frac{c^2}{\gamma^2}\right).$$

It follows that

$$\lim_{\gamma \rightarrow 0} \gamma^9 \ln \gamma N^\Delta = \begin{vmatrix} 1 & 3 & 12 & 4 \\ 3 & 12 & 60 & 20 \\ \frac{1}{3} & \frac{4}{3} & \frac{20}{3} & \frac{4}{3} \\ 4 & 20 & 120 & 40 \end{vmatrix} = \frac{64}{3}.$$

By joining all these results we obtain

$$\lim_{\gamma \rightarrow 0} \frac{\gamma N^\Pi}{D_4} = -6, \quad \lim_{\gamma \rightarrow 0} \frac{N^\Delta}{D_4} = -\frac{1}{3},$$

187 which confirms (62) also for $a = 2$.

188

- 189 1. Müller, I.; Ruggeri, T. *Rational Extended Thermodynamics*, 2nd ed.; Springer: New York, NY, USA, 1998.
- 190 2. Ruggeri, T.; Sugiyama, M. *Classical and Relativistic Rational Extended Thermodynamics of Gases*, Springer Heidelberg
191 New York Dordrecht London, ISBN:978-3-030-59143-4, 2021.
- 192 3. Liu, I-S.; Müller, I.; Ruggeri, T. Relativistic Thermodynamics of Gases, *Ann. Phys.* **1986**, *169*, 191-219.
- 193 4. Chernikov, N. A. Microscopic foundation of relativistic hydrodynamics, *Acta Phys. Polonica.* **1964**, *27*, 465–489.
- 194 5. Synge, J.L. *The Relativistic Gas*. Amsterdam North Holland, 1957.
- 195 6. Cercignani, C.; Kremer, G. M. *The Relativistic Boltzmann Equation: Theory and Applications*. Basel-Boston: Birkhäuser
196 Verlag, 2002.
- 197 7. Pennisi, S.; Ruggeri, T. Relativistic extended thermodynamics of rarefied polyatomic gas. *Ann. Phys.* **2017**, *377*,
198 415–445.
- 199 8. Carrisi, M.C.; Pennisi, S.; Ruggeri, T. Production terms in relativistic extended thermodynamics of gas with internal
200 structure via a new BGK model, *Ann. Phys.* **2019**, *405*, 298–307.
- 201 9. Carrisi, M.C.; Pennisi, S.; Ruggeri, T. Maxwellian iteration of a causal relativistic model of polyatomic gases and
202 evaluation of bulk, shear viscosity and heat conductivity, *Ann. Phys.* **2021**, *428*, 168447.
- 203 10. Arima, T., Taniguchi, S., Ruggeri, T.; Sugiyama, M. Extended thermodynamics of dense gases. *Continuum Mech.*
204 *Thermodyn.* **2011**, *24*, 271–292.
- 205 11. Pavić, M.; Ruggeri, T.; Simić, S. Maximum entropy principle for rarefied polyatomic gases. *Phys. A* **2013**, *392*,
206 1302–1317.
- 207 12. Pennisi, S.; Ruggeri, T. Classical Limit of Relativistic Moments Associated with Boltzmann-Chernikov Equation:
208 Optimal Choice of Moments in Classical Theory. *J. Stat. Phys.* **2020**, *179*, 231–246.
- 209 13. Pennisi, S. Consistent Order Approximations in Extended Thermodynamics of Polyatomic Gases, *J. Nat. Sci. Tech.*
210 **2021**, *2*, 12-21.
- 211 14. Arima T.; Carrisi M.C.; Pennisi S.; Ruggeri T., Which moments are appropriate to describe gases with internal structure
212 in Rational Extended Thermodynamics?, *Int. J. Non-Lin. Mech.* **2021**, *137*, (in press)
- 213 15. Jaynes, E. T.: Information Theory and Statistical Mechanics. *Phys. Rev.* **106**, 620-630 (1957). Jaynes, E. T.: Information
214 Theory and Statistical Mechanics II, *Phys. Rev.*, **1957**, *108*, 171–190 .
- 215 16. Müller, I., Ruggeri, T.: *Extended Thermodynamics*. Springer, New York, (1993).
- 216 17. Boillat, G.; Ruggeri, T. On the Shock Structure Problem for Hyperbolic System of Balance Laws and Convex Entropy.
217 *Continuum Mech. Thermodyn.*, **1998**, *10* 285–292.
- 218 18. Boillat, G.; Ruggeri, T. Hyperbolic principal subsystems: entropy convexity and subcharacteristic conditions, *Arch. Rat.*
219 *Mech. Anal.* **1997**, *137*, 305–320.
- 220 19. Ruggeri, T.; Strumia, A. Main field and convex covariant density for quasi-linear hyperbolic systems: Relativistic fluid
221 dynamics. *Ann. l'IHP Sec. A* **1981**, *34*, 65–84.
- 222 20. Marle, C., Sur l'établissement des equations de l'hydrodynamique des fluides relativistes dissipatifs. II. Méthodes de
223 résolution approchée de l'équation de Boltzmann relativiste. *Ann. Inst. Henri Poincaré* **1969**, *10*, 127–194.

- 224 21. Marle, C.; Modele cinétique pour l'établissement des lois de la conduction de la chaleur et de la viscosité en théorie de
225 la relativité. *C. R. Acad. Sci. Paris.* **1965**, *260*, 6539–6541.
- 226 22. Anderson, J. L.; Witting, H. R. A relativistic relaxation-time model for the Boltzmann equation. *Physica.* **1974**, *74*,
227 466–488.
- 228 23. Eckart, C.; The Thermodynamics of Irreversible Processes. III. Relativistic Theory of the Simple Fluid. *Phys. Rev.*,
229 **1940**, *58*, 919–924.
- 230 24. Landau, L. D.; Lifshitz, E. M. *Fluid Mechanics*. Pergamon Press, 1959.
- 231 25. Ruggeri, T.; Xiao, Q.; Zhao, H. Nonlinear Hyperbolic Waves in Relativistic Gases of Massive Particles with Sygne
232 Energy, *Arch. Rational Mech. Anal.* **2021**, *239*, 1061–1109.
- 233 26. Pennisi, S.; Ruggeri, T. Relativistic Eulerian rarefied gas with internal structure, *J. Math. Phys.* **2018** *59*, 043102-1 -
234 043102-12.
- 235 27. Pennisi, S.; Ruggeri, T. Ultra-relativistic Limit of Extended Thermodynamics of Rarefied Polyatomic Gas, *Le*
236 *Matematiche.* **2019**, *LXXIV*, 369–395.
- 237 28. Ikenberry, E.; Truesdell, C. On the pressure and the flux of energy in a gas according to Maxwell's kinetic theory.
238 *J. Ration. Mech. Anal.* **1956**, *5*, 1–54.

239 © 2021 by the authors. Submitted to *Entropy* for possible open access publication under the terms and conditions of the
240 Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).