

Efficiency and Robustness of Rosenbaum's Regression (Un)-Adjusted Rank-based Estimator in Randomized Experiments

Aditya Ghosh¹, Nabarun Deb², Bikram Karmakar^{*3}, and Bodhisattva Sen^{†4}

¹Department of Statistics, Stanford University

²Department of Mathematics, University of British Columbia

³Department of Statistics, University of Florida

⁴Department of Statistics, Columbia University

June 16, 2023

Abstract

Mean-based estimators of the causal effect in a completely randomized experiment (e.g., the difference-in-means estimator) may behave poorly if the potential outcomes have a heavy-tail, or contain outliers. We study an alternative estimator by Rosenbaum that estimates the constant additive treatment effect by inverting a randomization test using ranks. By investigating the breakdown point and asymptotic relative efficiency of this rank-based estimator, we show that it is provably robust against heavy-tailed potential outcomes, and has variance that is asymptotically, in the worst case, at most about 1.16 times that of the difference-in-means estimator; and its variance can be much smaller when the potential outcomes are not light-tailed. We further derive a consistent estimator of the asymptotic standard error for Rosenbaum's estimator which yields a readily computable confidence interval for the treatment effect. Further, we study a regression adjusted version of Rosenbaum's estimator to incorporate additional covariate information in randomization inference. We prove gain in efficiency by this regression adjustment method under a linear regression model. We illustrate through synthetic and real data that, unlike the mean-based estimators, these rank-based estimators (both unadjusted or regression adjusted) are efficient and robust against heavy-tailed distributions, contamination, and model misspecification.

Keywords: Breakdown point, causal inference, covariate adjustment, Hodges-Lehmann 0.864 lower bound, local asymptotic normality, randomization inference, Wilcoxon rank-sum statistic.

^{*}Supported by NSF grant DMS-2015250

[†]Supported by NSF grant DMS-2015376

1 Introduction

When a treatment is randomly assigned to the units in a study, [Fisher \(1935\)](#) showed that this randomization can be used to provide valid statistical inference about the treatment effect without making strong assumptions regarding the outcome-generating model. In particular, we can calculate confidence intervals for the treatment effect based on an estimator using its randomization distribution. A standard choice for the estimator is the difference-in-means estimator, which takes the difference of the averages of outcomes for the treated and control units. Conveniently for the practitioner, the usual large sample confidence interval based on two independent samples from infinite populations for the difference in two population means with unequal variances gives an asymptotically valid, conservative confidence interval under the randomization distribution ([Splawa-Neyman, 1990](#); [Freedman et al., 2007](#)). [Freedman \(2008a\)](#) establishes its validity under a finite limiting fourth moment assumption on the sequence of potential outcomes, and [Li and Ding \(2017\)](#) relax the assumption to any finite limiting moment of more than two; also see [Lin \(2013\)](#) and [Aronow et al. \(2014\)](#) for further developments.

Despite these advances, this mean-based confidence interval can be very wide, with volatile standard errors, when the potential outcomes are heavy-tailed, have a few extreme values, or contaminated by outliers. [Athey et al. \(2021\)](#) note that in many modern randomized experiments, including in experiments in the digital space, heavy-tailed or extreme outcomes are often likely. An important class of estimators used in practice in such situations use the *ranks* of the potential outcomes as opposed to their exact values (see e.g., [Imbens and Rubin \(2015, Section 5.5.4\)](#)). In particular, [Rosenbaum \(1993\)](#) proposed to use a Hodges-Lehmann type point estimate ([Hodges and Lehmann, 1963](#)) based on the Wilcoxon rank-sum (abbreviated as WRS) test statistic ([Wilcoxon, 1945](#)) for estimating a constant additive treatment effect under randomization inference. The constant treatment effect assumption is often a convenient starting point in answering a causal question ([Rosenbaum, 2002a](#); [Ho and Imai, 2006](#); [Athey et al., 2021](#)). Further, in some situations, the identification of a constant treatment effect has immediate practical use (see [Rosenbaum, 2002b](#), Section 2.4.5).

[Rosenbaum \(2002b](#), see Sections 4.6.7, 4.6.9) provided a numerical method to construct a confidence interval based on the above estimator, as follows. Under randomization, the upper and lower limits of the $100(1-\alpha)\%$ two-sided confidence interval are calculated by solving for the hypothesized treatment effects that equate the corresponding standardized WRS test statistic value to the $\alpha/2$ and $(1-\alpha/2)$ quantiles of the standard normal distribution. Unfortunately, this numerical method does not shed light on the efficiency gain/loss of this rank-based approach compared to the usual differences-in-means estimator, thereby precluding the practitioner from making an informed decision as to when such a rank-based method should be preferred.

In this paper, we undertake a systematic and rigorous study of the robustness and efficiency properties of Rosenbaum’s rank-based estimator based on the WRS test statistic, under randomization and the constant treatment effect assumption. Our asymptotic results have important implications. In particular, we show that the asymptotic variance of Rosenbaum’s estimator is always at most $125/108$ (≈ 1.16) times that of the difference-in-means estimator, and is often much smaller (see [Section 2.4](#) for a detailed discussion). We also provide an ana-

lytically computable, asymptotically valid confidence interval for the treatment effect based on this rank-based estimator, which is more robust to outliers or heavy-tailed potential outcomes than the mean-based confidence intervals. This is also evident in our simulation studies where the rank-based method gives much shorter confidence intervals than those obtained from the difference-in-means estimator, when the distribution of the potential outcomes has thicker tails.

Next we consider randomization inference for the treatment effect with regression adjustment. In modern randomized experiments it is increasingly common to collect additional data on covariates. This covariate information is typically used with an ANCOVA model, aiming to improve the unadjusted difference-in-means type estimator (Fisher, 1935; Cox and Reid, 2000; Freedman, 2008a,b). Lin (2013) showed that including the interaction of the treatment indicator and covariates in the model results in a regression adjusted estimator which is asymptotically at least as efficient as the unadjusted difference-in-means estimator. Recently, regression adjusted or model assisted estimators have been devised for various study designs (Fogarty, 2018; Li and Ding, 2020; Liu and Yang, 2020; Su and Ding, 2021; Zhao and Ding, 2022). But, these methods based on linear models which typically use robust standard errors (Huber, 1967; White, 1980), are still sensitive to heavy-tailed distributions or extreme values of the potential outcomes (MacKinnon and White, 1985; Young, 2018).

We study the asymptotic properties of a regression adjusted version of Rosenbaum’s rank-based estimator, originally suggested by Rosenbaum (2002a) for observational studies. To calculate this estimator, we first regress the control potential outcomes under the hypothesis of a specified treatment effect on the covariates using least squares. Then, we calculate the WRS statistic using the residuals of this regression. Finally, we invert this test statistic to compute the estimator. Unlike some of the regression adjustment methods cited above, we do not use the treatment indicator in the regression model; an approach advocated by Tukey (1993) (also see Gail et al. (1988)). Similar to the unadjusted case, our result on the asymptotic distribution of this estimator does not require any moment condition on the sequence of potential outcomes, and thus also holds for many heavy-tailed distributions of the potential outcomes. Our simulation studies illustrate notable gains in efficiency for this regression adjustment method over its unadjusted counterpart when the potential outcomes are not light-tailed.

1.1 Summary of our contributions

All the theoretical results in this paper are derived under the finite population asymptotics framework where the only source of randomization is the completely randomized assignment of the treatment; the potential outcomes are considered fixed. We formally define Rosenbaum’s rank-based estimator without regression adjustment in (2.11), and study its asymptotic properties in Section 2, under the constant additive treatment effect model (see Rosenbaum (2002b)). In particular, we illustrate the robustness of Rosenbaum’s rank-based estimator against extreme potential outcomes or contamination using the concept of breakdown point (see Proposition 2.2). Next, Theorem 2.2 gives the asymptotic distribution of this estimator under a completely randomized design as the sample size grows to infinity, under a mild regularity condition, namely Assumption 1 (which holds for many heavy-tailed distributions used in practice; see Remark 2.3 for a detailed discussion). Further, Corollary 2.1 gives a readily computable,

asymptotically valid confidence interval for the treatment effect.

Our proof of [Theorem 2.2](#) is novel and does not follow from existing results in the literature. As Rosenbaum’s rank-based estimator is computed by (essentially) equating the WRS statistic for the adjusted responses to its expected null value under randomization (see [\(2.9\)](#)), following [Hodges and Lehmann \(1963\)](#) we require the asymptotic distribution of this statistic under a sequence of *local alternatives* to study the asymptotic properties of this solution. The analysis is complicated by a number of reasons. First, under a completely randomized assignment, the observed outcomes of different units are neither independent nor identically distributed. Second, to study the behavior of the WRS statistic under local alternatives under randomization inference, the classical results from Le Cam’s local asymptotic normality theory (see [van der Vaart \(1998, Chapter 7\)](#) for details) do not apply. Recently, [Li and Ding \(2017\)](#) have established a few useful asymptotic finite population central limit theorems, including the null distribution of the Wilcoxon statistic; see [Proposition 2.1](#) and the accompanying discussion. Some of the results in [Li and Ding \(2017\)](#) have been generalized and used by [Li et al. \(2018\)](#), [Lei and Ding \(2021\)](#) and [Wu and Ding \(2021\)](#). However, these papers only consider statistics that are linear in the observations, and their results do not apply to rank-based methods (which are highly non-linear). Thus, our first technical innovation in [Section 2](#) is deriving the local asymptotic normality of the Wilcoxon statistic; see [Theorem 2.1](#).

Next, we establish in [Theorem 2.4](#) the asymptotic efficiency of Rosenbaum’s estimator relative to the difference-in-means estimator (note that both of these estimators target the same estimand under the constant additive treatment effect model). We note, in particular, that when the potential outcomes are realizations from a normal distribution, Rosenbaum’s estimator has about 95% asymptotic relative efficiency compared to the difference-in-means estimator (which is most efficient in this case). For potential outcome distributions that are not light-tailed, the efficiency gain is often substantial, see [Table 2.1](#). In fact, this result is parallel to the lower bound due to [Hodges and Lehmann \(1956\)](#) on the asymptotic efficiency of the WRS test relative to the Student’s t -test under an infinite population model.

In [Section 3](#), we define our regression adjusted rank-based estimator in [\(3.4\)](#) and study its asymptotic properties. In particular, [Theorem 3.3](#) gives the asymptotic distribution of our regression adjusted rank-based estimator. Previously, the asymptotic distributions of regression adjusted difference-in-means type estimators were derived under different randomization strategies ([Fogarty, 2018](#); [Li and Ding, 2020](#); [Liu and Yang, 2020](#); [Su and Ding, 2021](#); [Zhao and Ding, 2022](#)). These results build on the results of [Li and Ding \(2017\)](#) and do not apply to rank-based estimators, as already discussed above. We prove [Theorem 3.3](#) by deriving the local asymptotic behavior of the regression adjusted rank statistic, which is provided in [Theorem 3.2](#). The proof of this theorem is more involved than the proof of the analogous result in the unadjusted case (i.e., [Theorem 2.2](#)); we refer the reader to [Appendix B](#) for a summary of our proof techniques.

In [Appendix A](#), we propose consistent estimators of the standard error of this regression adjusted estimator which yields analytically computable confidence intervals based on it. The performance of these confidence intervals are included in our simulation studies in [Section 4](#) along with two applications of the methods on real data — one on the electoral impact of *Pro-*

gesa, Mexico’s conditional cash transfer program (De La O, 2013; Imai, 2018), and another on housing prices (Linden and Rockoff, 2008). These numerical examples illustrate that the performance of the rank-based estimators is never substantially worse than the ANOVA-based estimators, and is often considerably better when the potential outcomes are not light-tailed. Moreover, in most cases the regression adjusted rank-based estimator provides shorter confidence intervals than the unadjusted estimator. The proofs of our main results, additional technical results and discussions are relegated to Appendices C–E. Implementation of our proposed methods in **R** are available from <https://github.com/ghoshadi/RARE2021>.

2 Inference without regression adjustment

We assume throughout that the treatment group is formed by choosing $m = m(N)$ out of the N subjects in a randomized trial by *simple random sampling without replacement* (SRSWOR). We work in the Neyman-Rubin potential outcomes framework (Splawa-Neyman, 1990; Rubin, 1974, 1977; Imbens and Rubin, 2015). For the i -th subject, let a_i and b_i denote the potential outcomes under the treatment and control respectively; the observed response Y_i is given by

$$Y_i = Z_i a_i + (1 - Z_i) b_i, \quad (2.1)$$

where Z_i equals 1 if the i -th subject is assigned to the treatment group, and 0 otherwise. In our model Z_i is the only source of randomness in Y_i ; we will assume throughout this paper that $\{a_i\}_{i=1}^N$ and $\{b_i\}_{i=1}^N$ are fixed constants. We also make the assumption of a *constant additive treatment effect* (Rosenbaum, 1993, 2002a; Athey et al., 2021), i.e., $a_i - b_i = \tau$, for all $1 \leq i \leq N$, for some unknown real number τ . We are interested in point estimation and confidence interval for the constant treatment effect τ . In the sequel we will denote vectors by boldface letters, e.g., $\mathbf{Z} := (Z_1, \dots, Z_N)^\top$, $\mathbf{b} := (b_1, \dots, b_N)^\top$ and so forth. To formally state our asymptotic results as the sample size N grows to infinity, we will use an additional index N in our notation for the later sections. For example, the potential outcomes will be denoted by $\{(a_{N,j}, b_{N,j})\}_{1 \leq j \leq N}$, the vector of treatment indicators by $\mathbf{Z}_N = (Z_{N,1}, \dots, Z_{N,N})^\top$, and so on.

We estimate τ by inverting a testing problem. Towards that, consider the problem of testing

$$H_0 : \tau = \tau_0 \quad \text{versus} \quad H_1 : \tau \neq \tau_0. \quad (2.2)$$

When H_0 is true, the vector $\mathbf{Y} - \tau_0 \mathbf{Z}$, called the vector of *adjusted responses*, equals \mathbf{b} , which is non-random. Hence any statistic $t(\mathbf{Z}, \mathbf{Y} - \tau_0 \mathbf{Z})$, which is a function of the treatment indicators and the adjusted responses, can be used to make randomization inference about H_0 , since the null distribution of $t(\mathbf{Z}, \mathbf{Y} - \tau_0 \mathbf{Z}) = t(\mathbf{Z}, \mathbf{b})$ is completely specified by the randomization distribution of \mathbf{Z} . For example, one can consider the difference-in-means test statistic:

$$t_D(\mathbf{Z}, \mathbf{v}) := \frac{1}{m} \sum_{i: Z_i=1} v_i - \frac{1}{N-m} \sum_{i: Z_i=0} v_i, \quad (2.3)$$

where \mathbf{v} is a vector of observations of length N , and $m = \sum_{i=1}^N Z_i$.

Next, consider the problem of constructing valid confidence sets for τ based on the test statistic $t(\mathbf{Z}, \mathbf{Y} - \tau_0 \mathbf{Z})$. Using randomization, one can construct an exact, finite sample confidence interval for the treatment effect by appealing to the duality of testing and confidence intervals (Lehmann and Romano, 2005, Section 3.5). Note that we can simulate the randomization distribution of the test statistic $t(\mathbf{Z}', \mathbf{Y} - \tau_0 \mathbf{Z}')$ under $H_0 : \tau = \tau_0$, by evaluating the test statistic at each $\mathbf{Z}' = (Z'_1, \dots, Z'_N) \in \{0, 1\}^N$ such that $\sum_{i=1}^N Z'_i = m$. Then, the $100(1 - \alpha)\%$ confidence interval for τ is the set of hypothesized values of the parameter not rejected by the level α test using this null distribution. This method is slightly impractical even for moderately large N and m as it requires a lot of computation. In practice, one of the following methods are used to approximate the null distribution. One method evaluates the test statistic at a randomly selected set of \mathbf{Z}' 's. Another method uses the asymptotic null distribution of the test statistic from the infinite population literature (Conover, 1999; Lehmann, 1975). Although some authors have advocated for the first method (Bind and Rubin, 2020; Luo et al., 2021), the second method is easier to implement and thus more popular. But, to ensure that this method is approximately valid, it is important to understand if and when the asymptotic null distribution of the test statistic derived under the infinite population assumption also holds under randomization inference. This indeed holds for the difference-in-means statistic defined in (2.3), see Freedman (2008a) for details. However, the confidence intervals based on the difference-in-means statistic have the disadvantage that they can be very wide with volatile standard errors when the potential outcomes are heavy-tailed, or contaminated by outliers.

On the other hand, test statistics that use the *ranks* of the potential outcomes, as opposed to their exact values, are in general less sensitive to thick-tailed or skewed distributions and hence can lead to more powerful tests (see, e.g., Imbens and Rubin (2015, Section 5.5.4)). There are various popular choices for the rank-based statistic $t(\cdot, \cdot)$; Rosenbaum (1993) recommended using the WRS statistic (Wilcoxon, 1945), defined as

$$t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) := \mathbf{Z}^\top \mathbf{q}^{(\tau)} = \sum_{j: Z_j=1} q_j^{(\tau)}, \quad (2.4)$$

where $q_j^{(\tau)}$ is the rank of $Y_j - \tau Z_j$ among $\{Y_i - \tau Z_i : 1 \leq i \leq N\}$. Considering the possibility of ties in the data, we take the following definition for the ranks, known as *up-ranks*:

$$q_j^{(\tau)} := \sum_{i=1}^N \mathbf{1}\{Y_i - \tau Z_i \leq Y_j - \tau Z_j\}, \quad \text{for } 1 \leq j \leq N. \quad (2.5)$$

Note, under $H_0 : \tau = \tau_0$, the ranks $\{q_j^{(\tau_0)} : 1 \leq j \leq N\}$ can equivalently be written as

$$q_j^{(\tau_0)} = \sum_{i=1}^N \mathbf{1}\{b_i \leq b_j\}, \quad 1 \leq j \leq N. \quad (2.6)$$

We present the asymptotic null distribution of the WRS statistic in the following result (see Appendix C.1 for a proof).

Proposition 2.1 (Asymptotic null distribution of t_N). *Let $t_N := t(\mathbf{Z}_N, \mathbf{Y}_N - \tau_0 \mathbf{Z}_N)$ be the WRS statistic for a sample of size N , with $t(\cdot, \cdot)$ as defined in (2.4). Assume that $m/N \rightarrow \lambda \in (0, 1)$ as $N \rightarrow \infty$, and that the ranks $\{q_j^{(\tau_0)}\}_{1 \leq j \leq N}$ in (2.6) satisfy*

$$\frac{1}{N} \sum_{j=1}^N \left(q_j^{(\tau_0)} - \bar{q}_N^{(\tau_0)} \right)^2 = \frac{N^2 - 1}{12} + o(N^2), \quad (2.7)$$

as $N \rightarrow \infty$, where $\bar{q}_N^{(\tau_0)} := N^{-1} \sum_{j=1}^N q_j^{(\tau_0)}$. Then, under $\tau = \tau_0$,

$$N^{-3/2} \left(t_N - m \bar{q}_N^{(\tau_0)} \right) \xrightarrow{d} \mathcal{N} \left(0, \frac{\lambda(1-\lambda)}{12} \right). \quad (2.8)$$

Remark 2.1 (Null distribution in presence of ties). *The asymptotic null distribution in (2.8) also holds when there are ties, provided (2.7) holds. If instead of using up-ranks we break ties by comparing indices or by randomization, then the ranks being a permutation of $\{1, 2, \dots, N\}$ satisfy (2.7), and consequently (2.8) holds without any assumption. Moreover, if ties are broken using average ranks, then also (2.8) holds provided that the blocks of the ties are not large; see Lemma F.1 in Appendix F for a precise result in this direction.*

Remark 2.2. *Li and Ding (2017) give a finite population CLT for linear rank statistics, under the null hypothesis of no treatment effect, and assuming that there are no ties (see also Lehmann (1975, Appendix 4)). Moreover, they show that the conditions they need in the finite population framework are satisfied almost surely when the responses are realizations from a distribution having finite moments of any order more than two. In contrast, we derive the null distribution of t_N in this setting of constant treatment effect under the milder condition (2.7) that also allows ties and does not require finiteness of moments of any order.*

The limiting distribution in (2.8) is indeed the asymptotic null distribution of the WRS statistic under the infinite population framework (Lehmann, 1975, Section 1.3). Proposition 2.1 thus justifies Rosenbaum’s proposal of deriving a confidence interval for τ by numerically solving for the hypothesized treatment effects that equate the standardized WRS statistic to the $\alpha/2$ and $(1-\alpha/2)$ quantiles of the standard normal distribution (Rosenbaum, 2002b, Section 4.6). Unfortunately, this method of constructing confidence sets does not provide much insight about the efficiency of this rank-based approach over the difference-in-means estimator. In the subsequent sections, we develop the theory for a point estimator $\hat{\tau}^R$ of τ based on the WRS statistic, and establish the robustness and efficiency properties of this estimator.

2.1 Rosenbaum’s estimator based on the WRS statistic

To derive a point estimator for τ , Rosenbaum (2002a) suggested inverting the testing problem (2.2) by equating $t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z})$ to its expectation under the randomization distribution and solve for τ . Thus, the estimator for τ advocated by Rosenbaum (2002a) is a solution of the equation

$$t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) = \mathbb{E}_\tau t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}). \quad (2.9)$$

Note that $\mathbb{E}_\tau t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z})$ is same as $\mathbb{E} t(\mathbf{Z}, \mathbf{b})$, which does not depend on τ . We provide below a concrete example to illustrate this estimation strategy.

Example 1. Consider the difference-in-means test statistic defined in (2.3). Since the random treatment assignment is based on a SRSWOR sample of size m from the N subjects, we have $\mathbb{E} Z_i = m/N$ for each $1 \leq i \leq N$. Observing that $Z_i Y_i = Z_i a_i$ and $(1 - Z_i) Y_i = (1 - Z_i) b_i$, we deduce that $\mathbb{E}_\tau t_D(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) = 0$, and consequently, the solution to (2.9) for τ is given by

$$\hat{\tau}_{\text{dm}} := \frac{1}{m} \sum_{i=1}^N Z_i Y_i - \frac{1}{N - m} \sum_{i=1}^N (1 - Z_i) Y_i. \quad (2.10)$$

Thus, Rosenbaum's suggestion for $\hat{\tau}$ coincides with the difference-in-means estimator, for the above choice of the test statistic $t_D(\cdot, \cdot)$.

Next we implement Rosenbaum's suggestion in (2.9) with the WRS statistic $t(\cdot, \cdot)$ defined in (2.4). Assume for the moment that ties are not present in the ranks, so that

$$\mathbb{E}_\tau t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) = \mathbb{E}_\tau \mathbf{Z}^\top \mathbf{q}^{(\tau)} = \frac{m}{N} \sum_{j=1}^N q_j^{(\tau)} = \frac{m}{N} \cdot \frac{N(N+1)}{2} = \frac{m(N+1)}{2}.$$

Notice that the equation $t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) = m(N+1)/2$ might not always admit a solution for τ , for instance, when $m(N+1)$ is odd. To circumvent this technical issue, we slightly modify the definition of this estimator, following the footsteps of Hodges and Lehmann (1963). We set

$$\hat{\tau}^* := \sup \{ \tau : t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) > \mu \}, \quad \text{and} \quad \hat{\tau}^{**} := \inf \{ \tau : t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) < \mu \},$$

where $\mu := \mathbb{E}_{\tau_0} t(\mathbf{Z}, \mathbf{Y} - \tau_0 \mathbf{Z})$, and define

$$\hat{\tau}^{\text{R}} := \frac{\hat{\tau}^* + \hat{\tau}^{**}}{2}. \quad (2.11)$$

We show in the following section that $\hat{\tau}^{\text{R}}$ is robust to outliers (unlike the difference-in-means estimator $\hat{\tau}_{\text{dm}}$ defined in (2.3)), and study its asymptotic efficiency relative to $\hat{\tau}_{\text{dm}}$ in Section 2.4.

2.2 Robustness of Rosenbaum's estimator

In classical nonparametrics, a natural way to quantify the robustness of an estimator is via its (asymptotic) breakdown point (see, e.g., Hettmansperger and McKean (2011)). As in our setting the indices corresponding to the treatment and control groups are not deterministic but are rather chosen by randomization, we define the (asymptotic) breakdown point of an estimator of the constant treatment effect τ in the following fashion.

Definition 1 (Breakdown point). Consider any estimator $\hat{\tau} = \hat{\tau}(Y_1, \dots, Y_N; Z_1, \dots, Z_N)$ of the constant treatment effect τ . The finite sample breakdown point of $\hat{\tau}$ is defined as:

$$\text{BP}_N(\hat{\tau}) := \frac{1}{N} \min \left\{ 1 \leq k \leq N : \forall z_1, \dots, z_N \in \{0, 1\}, \sum_{i=1}^N z_i = m, \right. \\ \left. \min_{1 \leq i_1 < i_2 < \dots < i_k \leq N} \sup_{y_{i_1}, \dots, y_{i_k} \in \mathbb{R}} |\hat{\tau}(y_1, \dots, y_N; z_1, \dots, z_N)| = \infty \right\}. \quad (2.12)$$

Further, the asymptotic breakdown point of $\hat{\tau}$ is defined as $\text{ABP}(\hat{\tau}) := \lim_{N \rightarrow \infty} \text{BP}_N(\hat{\tau})$.

Intuitively, the above notion of breakdown point formalizes the following question: “What is the minimum proportion of responses that, if replaced with arbitrarily extreme values, will cause the estimator to be arbitrarily large (in absolute value), for all configurations of treatment assignments?” Note the emphasis on the phrase “for all configurations of treatment assignments”, which is reflected in the minimum taken over i_1, i_2, \dots, i_k in (2.12). Another alternative would have been to replace the minimum by a maximum in (2.12), which leads to a different and more adversarial notion of breakdown point. Also note that when we are taking the supremum in (2.12) over the responses, we are changing both the potential outcomes, keeping the assumption of constant additive treatment effect intact.

It is easy to see that changing a single response arbitrarily can lead to $\hat{\tau}_{\text{dm}} = \pm\infty$ irrespective of the treatment assignment, implying that $\text{BP}_N(\hat{\tau}_{\text{dm}}) = 1/N$ and consequently $\text{ABP}(\hat{\tau}_{\text{dm}}) = 0$. In contrast, the following result (see [Appendix C.2](#) for a proof) illustrates that the asymptotic breakdown point of $\hat{\tau}^{\text{R}}$ is strictly positive for all values of the proportion $\lambda \approx m/N$.

Proposition 2.2 (Asymptotic breakdown point of $\hat{\tau}^{\text{R}}$). *Let $\hat{\tau}^{\text{R}}$ be the Rosenbaum’s estimator based on the WRS statistic, as defined in (2.11). Assume that $m/N \rightarrow \lambda \in [0, 1]$, where m is the size of the treatment group. Then the asymptotic breakdown point of $\hat{\tau}^{\text{R}}$ is given by*

$$\text{ABP}(\hat{\tau}^{\text{R}}) = \begin{cases} (1 - \lambda)/2 & \text{if } \lambda < 1/3 \\ 1 - \sqrt{2\lambda(1 - \lambda)} & \text{if } 1/3 \leq \lambda \leq 2/3 \\ \lambda/2 & \text{if } \lambda > 2/3. \end{cases}$$

It is well-known (see, for instance, [Hettmansperger and McKean \(2011, Section 6.5.3\)](#)) that the asymptotic breakdown point of the one sample Hodges-Lehmann location estimator ([Hodges and Lehmann \(1963\)](#)) is $1 - 2^{-1/2} \approx 0.29$. [Proposition 2.2](#) illustrates that the asymptotic breakdown point of $\hat{\tau}^{\text{R}}$ is never less than $1 - 2^{-1/2}$. In words, irrespective of the relative size of the treatment group, more than 29% of the responses need to be altered to force $\hat{\tau}^{\text{R}}$ to be arbitrarily large for any treatment assignment.

2.3 Asymptotic distribution of Rosenbaum’s estimator

In this section, we establish the asymptotic distribution of Rosenbaum’s estimator $\hat{\tau}^{\text{R}}$ defined in (2.11) based on the WRS statistic introduced in (2.4). For notational clarity, we index vectors and matrices by subscript N (the total sample size). Although $\hat{\tau}^{\text{R}}$ can be expressed as the median of the pairwise differences between the responses from the treatment and the control pool, we follow the strategy of [Hodges and Lehmann \(1963\)](#) that works in more generality (in

particular, for the regression adjustment case that we study later). Before presenting our main results, we briefly discuss this method of obtaining the asymptotic distribution of $\hat{\tau}^R$ from that of the test statistic $t_N := t(\mathbf{Z}_N, \mathbf{Y}_N - \tau_0 \mathbf{Z}_N)$. [Lemma D.1](#) in [Appendix D](#) connects the distribution of $\hat{\tau}^R$ to the distribution of t_N under a sequence of local alternatives $\tau_N := \tau_0 - hN^{-1/2}$ for some fixed $h \in \mathbb{R}$. In particular, we show that if

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\tau_N} (N^{-3/2}(t_N - \mathbb{E}_{\tau_0} t_N) \leq x) = \Phi \left(\frac{x + hB}{A} \right), \quad \text{for every } x \in \mathbb{R}, \quad (2.13)$$

where $\Phi(\cdot)$ is the standard normal distribution function, and A, B are positive constants, then

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\tau_0} \left(\sqrt{N}(\hat{\tau}^R - \tau_0) \leq x \right) = \Phi \left(\frac{xB}{A} \right), \quad \text{for every } x \in \mathbb{R}. \quad (2.14)$$

In view of the above, it suffices to establish (2.13) for the WRS statistic $t_N := t(\mathbf{Z}_N, \mathbf{Y}_N - \tau_0 \mathbf{Z}_N)$, with $t(\cdot, \cdot)$ as defined in (2.4). Although [Li and Ding \(2017, Corollary 1\)](#) give the asymptotic null distribution of t_N , their proof technique does not work for finding the asymptotic distribution of t_N under local alternatives. The classical results from Le Cam's local asymptotic normality theory (cf. [van der Vaart \(1998, Chapter 7\)](#)) also do not apply in our fixed design setting.

We present the asymptotic distribution of t_N under a sequence of local alternatives in [Theorem 2.1](#) below, and this is indeed one of our major contributions. To concisely state an assumption we need here, we introduce the following notation. Define, for any $h, x \in \mathbb{R}$,

$$I_{h,N}(x) := \begin{cases} \mathbf{1}\{0 \leq x < hN^{-1/2}\} & \text{if } h \geq 0, \\ -\mathbf{1}\{hN^{-1/2} \leq x < 0\} & \text{if } h < 0, \end{cases} \quad (2.15)$$

where $\mathbf{1}\{\cdot\}$ denotes the indicator of a set.

Assumption 1. *Let $\{b_{N,j} : 1 \leq j \leq N\}$ be the potential control outcomes. We assume that, for $I_{h,N}$ as in (2.15), the following holds:*

$$\lim_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N I_{h,N}(b_{N,j} - b_{N,i}) = h\mathcal{I}_b, \quad (2.16)$$

for some fixed constant $\mathcal{I}_b \in (0, \infty)$ and for every $h \in \mathbb{R}$.

Remark 2.3 (On [Assumption 1](#)). *Intuitively, [Assumption 1](#) roughly says that the proportion of the pairwise differences $(b_{N,j} - b_{N,i})$ falling into small intervals (shrinking at the rate of $N^{-1/2}$) scales with the Lebesgue measure of those intervals. Indeed, if $b_{N,j}$'s are realizations from a distribution with density $f_b(\cdot)$ that satisfies $\int_{\mathbb{R}} f_b^2(x) dx < \infty$, then [Assumption 1](#) holds almost surely with $\mathcal{I}_b = \int_{\mathbb{R}} f_b^2(x) dx$ (see [Lemma D.4](#) in [Appendix D](#) for a formal statement). Not requiring finiteness of moments, this condition holds for many heavy-tailed distributions (e.g., Pareto). Moreover, [Lemma D.5](#) in [Appendix D](#) shows that [Assumption 1](#) implies (2.7), and consequently the asymptotic null distribution of t_N in (2.8) also holds.*

The following theorem is pivotal; it provides the local asymptotic normality for t_N (see [Appendix C.3](#) for a proof), which will aid us to study the limiting behavior of $\hat{\tau}^R$.

Theorem 2.1 (Local asymptotic normality of t_N). *Let $t_N := t(\mathbf{Z}_N, \mathbf{Y}_N - \tau_0 \mathbf{Z}_N)$ be the WRS test statistic, with $t(\cdot, \cdot)$ as in (2.4). Assume that $m/N \rightarrow \lambda \in (0, 1)$, and suppose that [Assumption 1](#) holds. Fix $h \in \mathbb{R}$ and let $\tau_N = \tau_0 - hN^{-1/2}$. Then it follows that under $\tau = \tau_N$,*

$$N^{-3/2} \left(t_N - \frac{m(N+1)}{2} \right) \xrightarrow{d} \mathcal{N} \left(-h\lambda(1-\lambda)\mathcal{I}_b, \frac{\lambda(1-\lambda)}{12} \right),$$

where \mathcal{I}_b is defined in [Assumption 1](#).

With the above result, we are now prepared to present the asymptotic distribution of the estimator $\hat{\tau}^R$. This is the content of our next theorem (see [Appendix C.4](#) for a proof).

Theorem 2.2 (CLT for the estimator $\hat{\tau}^R$). *Let $\hat{\tau}^R$ be as in (2.11), with $t(\cdot, \cdot)$ as in (2.4). Suppose that [Assumption 1](#) holds, and that $m/N \rightarrow \lambda \in (0, 1)$. If τ_0 is the true value of τ , then*

$$\sqrt{N} (\hat{\tau}^R - \tau_0) \xrightarrow{d} \mathcal{N} (0, (12\lambda(1-\lambda)\mathcal{I}_b^2)^{-1}),$$

where \mathcal{I}_b is defined in [Assumption 1](#).

[Theorem 2.2](#) tells us that the problem of estimating the asymptotic variance of $\hat{\tau}^R$ reduces to the problem of estimating the limiting quantity \mathcal{I}_b , which is unknown in general. Naturally, our next target is to estimate \mathcal{I}_b based on the observed responses $\{Y_{N,j}\}_{1 \leq j \leq N}$. We propose an estimator of \mathcal{I}_b and prove its consistency under the same set of assumptions as in [Theorem 2.2](#). This is stated in the following theorem (see [Appendix C.5](#) for a proof).

Theorem 2.3 (Consistent estimation of \mathcal{I}_b). *Suppose that [Assumption 1](#) holds and let $m/N \rightarrow \lambda \in (0, 1)$ as $N \rightarrow \infty$. Define*

$$\hat{\mathcal{I}}_N := \left(1 - \frac{m}{N}\right)^{-2} N^{-3/2} \sum_{j=1}^N \sum_{i=1, i \neq j}^N (1 - Z_{N,i})(1 - Z_{N,j}) \mathbf{1}\{0 \leq Y_{N,j} - Y_{N,i} < N^{-1/2}\}. \quad (2.17)$$

Then $\hat{\mathcal{I}}_N \xrightarrow{P} \mathcal{I}_b$ as $N \rightarrow \infty$.

The above theorem in conjunction with [Theorem 2.2](#) readily yields an asymptotically valid confidence interval for τ_0 (the true value of τ). This is formally stated below.

Corollary 2.1 (Confidence interval for τ_0). *Let $\hat{\mathcal{I}}_N$ be as defined in (2.17). Under the assumptions of [Theorem 2.2](#), an approximate $100(1 - \alpha)\%$ confidence interval for τ_0 is given by*

$$\hat{\tau}^R \pm \frac{z_{\alpha/2}}{\sqrt{N}} \left(12 \frac{m}{N} \left(1 - \frac{m}{N} \right) \hat{\mathcal{I}}_N^2 \right)^{-1/2},$$

where z_α denotes the upper α -th quantile of the standard normal distribution.

2.4 Efficiency of Rosenbaum's estimator

In this section we aim to assess the asymptotic relative efficiency of Rosenbaum's estimator $\hat{\tau}^R$ with respect to the difference-in-means estimator $\hat{\tau}_{\text{dm}}$. In classical statistics, a framework for comparing two consistent, asymptotically normal estimators is through the ratio of their asymptotic variances. A precise definition is given below.

Definition 2 (Asymptotic relative efficiency). *Let $\hat{\tau}_{N,1}$ and $\hat{\tau}_{N,2}$ be two asymptotically normal estimators of τ , in the sense that there exist positive sequences $\sigma_{N,1}^2$ and $\sigma_{N,2}^2$ such that, $\frac{\hat{\tau}_{N,1} - \tau}{\sigma_{N,1}} \xrightarrow{d} \mathcal{N}(0, 1)$ and $\frac{\hat{\tau}_{N,2} - \tau}{\sigma_{N,2}} \xrightarrow{d} \mathcal{N}(0, 1)$. Then the asymptotic relative efficiency of $\hat{\tau}_{N,1}$ with respect to $\hat{\tau}_{N,2}$ is defined as*

$$\text{eff}(\hat{\tau}_{N,1}, \hat{\tau}_{N,2}) := \lim_{N \rightarrow \infty} \frac{\sigma_{N,2}^2}{\sigma_{N,1}^2}.$$

Our next theorem provides a lower bound on the asymptotic relative efficiency of $\hat{\tau}^R$ with respect to $\hat{\tau}_{\text{dm}}$ (see [Appendix C.6](#) for a proof).

Theorem 2.4 (Efficiency lower bound). *Assume that the distribution F_N putting equal mass on the potential control outcomes $\{b_{N,j}\}_{1 \leq j \leq N}$ converges weakly to a distribution F with density f (w.r.t. the Lebesgue measure on \mathbb{R}) such that (a) $N^{-1} \sum_{j=1}^N f(b_{N,j}) \rightarrow \int_{\mathbb{R}} f^2(x) dx < \infty$, and (b) for every fixed h , $\sup_{x \in \mathbb{R}} \left| \sqrt{N} \left(F_N(x + \frac{h}{\sqrt{N}}) - F_N(x) \right) - hf(x) \right| \rightarrow 0$ as $N \rightarrow \infty$. Then,*

$$\sqrt{N} (\hat{\tau}^R - \tau_0) \xrightarrow{d} \mathcal{N} \left(0, (12\lambda(1-\lambda))^{-1} \left(\int_{\mathbb{R}} f^2(x) dx \right)^{-2} \right).$$

Moreover, if (c) $N^{-1} \sum_{i=1}^N (b_{N,i} - \bar{b}_N)^2 \rightarrow \sigma^2$, and (d) $N^{-1} \max_{1 \leq i \leq N} (b_{N,i} - \bar{b}_N)^2 \rightarrow 0$, the asymptotic efficiency of $\hat{\tau}^R$ relative to $\hat{\tau}_{\text{dm}}$ is given by

$$\text{eff}(\hat{\tau}^R, \hat{\tau}_{\text{dm}}) = 12\sigma^2 \left(\int_{\mathbb{R}} f^2(x) dx \right)^2. \quad (2.18)$$

Further, when σ^2 is the variance of the limiting distribution F , it holds that $\text{eff}(\hat{\tau}^R, \hat{\tau}_{\text{dm}}) \geq 0.864$.

Remark 2.4 (On the assumptions of [Theorem 2.4](#)). When $F_N \xrightarrow{d} F$, the assumption (a) holds under mild conditions on f , e.g., if $\{f(B_N)\}_{N \geq 1}$ is uniformly integrable where $B_N \sim F_N$ (this holds, in particular, when f is bounded). Next, it follows from [Chernozhukov et al. \(2014, Corollary 5.1\)](#) that the assumption (b) holds when $\{b_{N,j}\}_{1 \leq j \leq N}$ are realizations from a distribution with density f . Lastly, the conditions (c) and (d) are identical to the assumptions in [Li and Ding \(2017, Theorem 5\)](#), and provide the necessary asymptotic distribution of $\hat{\tau}_{\text{dm}}$.

[Theorem 2.4](#) shows that, the limiting distribution of the rank-based estimator $\hat{\tau}^R$ has a variance that is never more than 1.16 times the same for the difference-in-means estimator. [Table 2.1](#) records the values of the relative efficiency $\text{eff}(\hat{\tau}^R, \hat{\tau}_{\text{dm}})$ for some standard probability

distributions. As can be seen from the table, even when the $\{b_{N,j}\}$ are realizations from a normal distribution, the rank-based estimator only suffers a 5% loss of efficiency. Whereas, for the exponential distribution, the rank-based estimator gives a 2-fold gain in efficiency. It is easy to construct examples where the rank-based estimator will exhibit infinite gains over the difference-in-means estimator, e.g., in the Pareto(α) example in Table 2.1, the efficiency approaches ∞ as $\alpha \rightarrow 2$. We also refer the reader to our simulation results in Section 4 which highlights how these efficiencies manifest themselves in moderate sample sizes.

Table 2.1: Asymptotic efficiency of $\hat{\tau}^R$ relative to $\hat{\tau}_{\text{dm}}$ for different choices of F in Theorem 2.4.

Distribution (F)	Density (f)	$\text{eff}(\hat{\tau}^R, \hat{\tau}_{\text{dm}})$
Normal	$(2\pi)^{-1/2} \exp(-x^2/2)$	$3/\pi \approx 0.955$
Uniform	$\mathbf{1}\{0 \leq x \leq 1\}$	1
Laplace	$2^{-1} \exp(- x)$	$3/2$
t_3	$c(x^2/3 + 1)^{-2}$	$75/(4\pi^2) \approx 1.9$
Exponential	$\exp(-x)\mathbf{1}\{x \geq 0\}$	3
Pareto(α)	$\alpha x^{-(\alpha+1)}\mathbf{1}\{x \geq 1\}$	$\begin{cases} \frac{\alpha^5}{(\alpha-1)^2(2\alpha+1)^2(\alpha-2)} & \text{if } \alpha > 2 \\ +\infty & \text{if } \alpha \in (0, 2] \end{cases}$

Remark 2.5 (The 0.864 lower bound). *The efficiency lower bound given in Theorem 2.4 coincides with a celebrated efficiency lower bound due to Hodges and Lehmann (1956) in the context of two sample testing under location shift alternatives. They showed that the Pitman efficiency (see, e.g., van der Vaart (1998, Section 14.3)) of the WRS test relative to Student's t -test never falls below 0.864. Here, $\hat{\tau}^R$ is the estimator that inverts the WRS test, and thus, in a way, our Theorem 2.4 mimics the efficiency result of Hodges and Lehmann (1956).*

Definition 2 above provides a principled framework for comparing estimators under a randomization inference. In practice, this framework allows us to use a pilot sample to choose an estimator that is expected to be more efficient in the final study. Pilot sampling is a frequently used tool to plan an empirical study (Wittes and Brittain, 1990). In this context, we can estimate f using a pilot sample, which may then be used with (2.18) to choose between the difference-in-means and Rosenbaum's estimator by estimating their relative efficiency.

3 Inference with regression adjustment

We now study a method advocated by Rosenbaum (2002a) for drawing randomization inference on the treatment effect with regression adjustment for pre-treatment covariates, in the setup of completely randomized experiments as in Section 2. Assume that along with the responses $\{Y_i\}_{1 \leq i \leq N}$ (as in (2.1)) we also collect data on p covariates, denoted by \mathbf{x}_i for the i -th subject. Also denote by \mathbf{X} the $N \times p$ matrix of covariates. To avoid notational clutter, we suppress the use of the index N in the notation until Section 3.1.

To motivate our estimation strategy, consider again the testing problem (2.2). The method suggested by Rosenbaum is to apply a randomization test on the residuals, as follows. First we calculate the adjusted responses $\mathbf{Y} - \tau_0 \mathbf{Z}$ and regress it on the covariates \mathbf{X} using the least squares criterion. Define the residuals obtained from this fit as:

$$\mathbf{e}_0 := \widehat{\varepsilon}(\mathbf{Y} - \tau_0 \mathbf{Z}, \mathbf{X}) = (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{Y} - \tau_0 \mathbf{Z}), \quad (3.1)$$

where \mathbf{I} is the identity matrix of order $N \times N$, and $\mathbf{P}_{\mathbf{X}}$ is the projection matrix onto the column space of \mathbf{X} . Now we let the residuals \mathbf{e}_0 play the role of the adjusted responses $\mathbf{Y} - \tau_0 \mathbf{Z}$ while performing the randomization test. Define the WRS test statistic based on the residuals \mathbf{e}_0 as $t_{\text{adj}} := t(\mathbf{Z}, \mathbf{e}_0) = \mathbf{q}^\top \mathbf{Z}$, with $t(\cdot, \cdot)$ as in (2.4) and the up-ranks \mathbf{q} in (2.5) being calculated on the residuals instead of the adjusted responses. That is,

$$t_{\text{adj}} := t(\mathbf{Z}, \mathbf{e}_0) = \sum_{i=1}^N Z_i \sum_{j=1}^N \mathbf{1}\{e_{0,j} \leq e_{0,i}\}. \quad (3.2)$$

To find a point estimator for τ based on the above test procedure, Rosenbaum (2002a) suggested calculating residuals \mathbf{e}_τ as

$$\mathbf{e}_\tau := \widehat{\varepsilon}(\mathbf{Y} - \tau \mathbf{Z}, \mathbf{X}) = (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{Y} - \tau \mathbf{Z}).$$

We then equate $t(\mathbf{Z}, \mathbf{e}_\tau)$ with its expectation under the randomization distribution of \mathbf{Z} and solve for τ . Thus, the regression adjusted estimator advocated by Rosenbaum (2002a) is a solution to the equation

$$t(\mathbf{Z}, \mathbf{e}_\tau) = \mathbb{E}_\tau t(\mathbf{Z}, \mathbf{e}_\tau). \quad (3.3)$$

Note that the right hand side of (3.3) is free of τ , because $\mathbb{E}_\tau t(\mathbf{Z}, \mathbf{e}_\tau) = \mathbb{E}_{\tau_0} t(\mathbf{Z}, \mathbf{e}_0)$ and under $\tau = \tau_0$, $\mathbf{e}_0 = (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{Y} - \tau_0 \mathbf{Z}) = (\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{b}$, whose distribution does not depend on τ .

Again, due to the discreteness of the ranks, (3.3) might not always have a solution. So we modify the estimator using the idea of Hodges and Lehmann (1963), as we did in the previous section for the unadjusted estimator $\widehat{\tau}^R$. We denote the modified Rosenbaum's estimator as $\widehat{\tau}_{\text{adj}}^R$, which is defined using (2.11) in the same manner as $\widehat{\tau}^R$, only with the difference that here $t(\mathbf{Z}, \mathbf{e}_\tau)$ plays the role of $t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z})$. We set

$$\widehat{\tau}_{\text{adj}}^* := \sup\{\tau : t(\mathbf{Z}, \mathbf{e}_\tau) > \mathbb{E}_\tau t(\mathbf{Z}, \mathbf{e}_\tau)\}, \quad \text{and} \quad \widehat{\tau}_{\text{adj}}^{**} := \inf\{\tau : t(\mathbf{Z}, \mathbf{e}_\tau) < \mathbb{E}_\tau t(\mathbf{Z}, \mathbf{e}_\tau)\},$$

and define

$$\widehat{\tau}_{\text{adj}}^R := \frac{\widehat{\tau}_{\text{adj}}^* + \widehat{\tau}_{\text{adj}}^{**}}{2}. \quad (3.4)$$

In the unadjusted case, the WRS statistic $t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z})$, with $t(\cdot, \cdot)$ as defined in (2.4), can easily be seen to be a monotonic function of τ . This is not immediate in the regression adjustment case. However, we will assume throughout that $t(\mathbf{Z}, \mathbf{e}_\tau)$ is a monotonic function of τ (cf. Hodges and Lehmann (1963, p. 599)), so that the above definition of $\widehat{\tau}_{\text{adj}}^R$ makes sense. In our simulation studies in Section 4 and applications to real data in Section 5, we found that this monotonicity

assumption is always satisfied.

Analyzing $\hat{\tau}_{\text{adj}}^{\text{R}}$ presents significant new challenges compared to $\hat{\tau}^{\text{R}}$. The chief reason being that $t(\mathbf{Z}, \mathbf{e}_\tau)$, which determines $\hat{\tau}_{\text{adj}}^{\text{R}}$, involves indicators of the form $\mathbf{1}(e_{\tau,j} \leq e_{\tau,i})$, and each of these indicators depends on the entire random treatment assignment vector \mathbf{Z} . This is not the situation in the unadjusted case, where the corresponding indicator only depends on Z_i and Z_j . We relegate the discussion on how we were able to resolve this technical challenge to [Appendix B](#) so as not to impede the flow of the paper.

3.1 Asymptotic distribution of $\hat{\tau}_{\text{adj}}^{\text{R}}$

Consider now a sequence of completely randomized experiments as in [Section 2.3](#). Denote the residuals obtained from the least squares fitting of $\mathbf{Y}_N - \tau_0 \mathbf{Z}_N$ on the covariates \mathbf{X}_N by $\mathbf{e}_N := \hat{\varepsilon}(\mathbf{Y}_N - \tau_0 \mathbf{Z}_N, \mathbf{X}_N)$, with $\hat{\varepsilon}(\cdot, \cdot)$ as in (3.1), and define the regression adjusted WRS statistic as

$$t_{N,\text{adj}} := t(\mathbf{Z}_N, \mathbf{e}_N) \quad (3.5)$$

with $t(\cdot, \cdot)$ as in (3.2). Finally, define $\hat{\tau}_{\text{adj}}^{\text{R}}$ as in (3.4) using the statistic $t_{N,\text{adj}}$.

As in the without regression adjustment case (see (2.13) and (2.14)), we invoke [Lemma D.1](#) to reduce the problem of deriving the asymptotic distribution of the estimator $\hat{\tau}_{\text{adj}}^{\text{R}}$ to the problem of finding the asymptotic distribution of the test statistic $t_{N,\text{adj}}$ under the sequence of local alternatives $\tau_N = \tau_0 - hN^{-1/2}$ for a fixed $h \in \mathbb{R}$. As a precursor to our result on the local asymptotic normality of $t_{N,\text{adj}}$, we first present its limiting null distribution. Recall the notation $\{b_{N,j}\}$ for the potential control outcomes. Define the regression adjusted potential control outcomes as

$$\tilde{b}_{N,j} := b_{N,j} - \mathbf{p}_{N,j}^\top \mathbf{b}_N, \quad 1 \leq j \leq N, \quad (3.6)$$

where $\mathbf{p}_{N,j}$ is the j -th column of the projection matrix $\mathbf{P}_{\mathbf{X}_N}$ that projects onto the column space of \mathbf{X}_N . Note that under $\tau = \tau_0$, these $\tilde{b}_{N,j}$'s are identical to the residuals obtained by regressing $\mathbf{Y}_N - \tau_0 \mathbf{Z}_N$ on \mathbf{X}_N . The following assumption mimics [Assumption 1](#) of [Section 2.3](#).

Assumption 2. *Let $\{\tilde{b}_{N,j} : 1 \leq j \leq N\}$ be the regression adjusted potential control outcomes as defined in (3.6). We assume that, for $I_{h,N}$ as in (2.15), the following holds:*

$$\lim_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N I_{h,N}(\tilde{b}_{N,j} - \tilde{b}_{N,i}) = h\mathcal{J}_b,$$

for some fixed constant $\mathcal{J}_b \in (0, \infty)$ and for every $h \in \mathbb{R}$.

We show in [Lemma D.8](#) in [Appendix D](#) that [Assumption 2](#) holds (in probability) when the sequence of potential control outcomes are sampled according to the regression model $\mathbf{b}_N = \mathbf{X}_N \boldsymbol{\beta}_N + \boldsymbol{\varepsilon}_N$, where $\varepsilon_{N,1}, \dots, \varepsilon_{N,N}$ are i.i.d. from $\mathcal{N}(0, \sigma^2)$ for some $\sigma^2 > 0$. In fact, the limiting quantity \mathcal{J}_b has a neat formula in this special case, namely, $\mathcal{J}_b = (2\sqrt{\pi}\sigma)^{-1}$.

Our next theorem provides the limiting null distribution of the test statistic $t_{N,\text{adj}}$ under [Assumption 2](#) (see [Appendix C.7](#) for a proof).

Theorem 3.1 (Asymptotic null distribution of $t_{N,\text{adj}}$). *Let $t_{N,\text{adj}}$ be the WRS statistic based on the residuals obtained from the least squares fitting of $\mathbf{Y}_N - \tau_0 \mathbf{Z}_N$ on \mathbf{X}_N , as defined in (3.5). Assume that $m/N \rightarrow \lambda \in (0, 1)$, and suppose that [Assumption 2](#) holds. Then, under $\tau = \tau_0$,*

$$N^{-3/2} \left(t_{N,\text{adj}} - \frac{m(N+1)}{2} \right) \xrightarrow{d} \mathcal{N} \left(0, \frac{\lambda(1-\lambda)}{12} \right). \quad (3.7)$$

It is noteworthy that the limiting null distribution of the regression adjusted statistic $t_{N,\text{adj}}$ given in (3.7) and that of the unadjusted statistic t_N given in (2.8) are identical. However, this is not unexpected, since both \mathbf{b}_N and \mathbf{e}_N are deterministic vectors under the null, the ranks always take values in $\{1, 2, \dots, N\}$, and \mathbf{Z}_N is the only source of randomness.

We next focus on deriving the asymptotic normality of the statistic $t_{N,\text{adj}}$ under the sequence of local alternatives $\tau_N = \tau_0 - hN^{-1/2}$ as $N \rightarrow \infty$. This is addressed in [Theorem 3.2](#) below (see [Appendix C.8](#) for a proof), and it immediately yields a CLT for $\hat{\tau}_{\text{adj}}^R$, which is presented in [Theorem 3.3](#). Although these results are parallel to their unadjusted counterparts ([Theorems 2.1](#) and [2.2](#), respectively), their proofs are substantially different, and more involved. A more detailed description of the technical challenges is provided in [Appendix B](#).

Theorem 3.2 (Local asymptotic normality of $t_{N,\text{adj}}$). *Assume the setting of [Theorem 3.1](#). Fix $h \in \mathbb{R}$ and let $\tau_N = \tau_0 - hN^{-1/2}$. Then, under $\tau = \tau_N$,*

$$N^{-3/2} \left(t_{N,\text{adj}} - \frac{m(N+1)}{2} \right) \xrightarrow{d} \mathcal{N} \left(-h\lambda(1-\lambda)\mathcal{J}_b, \frac{\lambda(1-\lambda)}{12} \right),$$

where \mathcal{J}_b is defined in [Assumption 2](#).

We are now ready to see a central limit theorem for the estimator $\hat{\tau}_{\text{adj}}^R$, which is the content of the following theorem (see [Appendix C.9](#) for a proof).

Theorem 3.3 (CLT for the estimator $\hat{\tau}_{\text{adj}}^R$). *Assume that $m/N \rightarrow \lambda \in (0, 1)$, and suppose that [Assumption 2](#) holds. If τ_0 is the true value of τ , then*

$$\sqrt{N} (\hat{\tau}_{\text{adj}}^R - \tau_0) \xrightarrow{d} \mathcal{N} \left(0, (12\lambda(1-\lambda)\mathcal{J}_b^2)^{-1} \right),$$

where \mathcal{J}_b is defined in [Assumption 2](#).

The problem of estimating the asymptotic variance of $\hat{\tau}_{\text{adj}}^R$ is more challenging than in the unadjusted case, and is deferred until [Appendix A](#) (see [Corollary A.2](#) for details, and [Sections 4](#) and [5](#) for its empirical performance).

3.2 Efficiency gain by regression adjustment

As discussed in [Section 2](#), appropriate randomization inference can be drawn ignoring the information available on the covariates. However, it is a popular belief ([Rosenbaum, 2002a](#)) that regression adjustment may increase the efficiency of the inference based on the rank-based

statistic, although any formal result supporting this belief was missing. We provide one result in this direction in [Theorem 3.4](#) below.

In view of [Theorems 2.2](#) and [3.3](#), a comparison between the asymptotic variances of the adjusted estimator $\hat{\tau}_{\text{adj}}^{\text{R}}$ and the unadjusted estimator $\hat{\tau}^{\text{R}}$ is essentially a comparison of the quantities \mathcal{J}_b and \mathcal{I}_b . In the following theorem (see [Appendix C.10](#) for a proof) we show that $\text{eff}(\hat{\tau}_{\text{adj}}^{\text{R}}, \hat{\tau}^{\text{R}}) \geq 1$ under the assumption that the regression model is correctly specified (recall the notion of asymptotic relative efficiency from [Section 2.4](#)). Thus, in this setting, the regression adjustment provides more efficient estimates than in the unadjusted case.

Theorem 3.4 (Efficiency gain by regression adjustment). *Assume that the potential control outcome sequence \mathbf{b}_N are realizations from the regression model $\mathbf{b}_N = \mathbf{X}_N \boldsymbol{\beta}_N + \boldsymbol{\varepsilon}_N$, where $\varepsilon_{N,i}$'s are i.i.d. from $\mathcal{N}(0, \sigma^2)$. Then [Assumption 2](#) holds, with $\mathcal{J}_b = (2\sqrt{\pi}\sigma)^{-1}$. Further, denote $\mathbf{v}_N := \mathbf{X}_N \boldsymbol{\beta}_N$ and assume that $\lim_{N \rightarrow \infty} N^{-2} \sum_{j=1}^N \sum_{i=1}^N \exp(-(v_{N,j} - v_{N,i})^2 / 4\sigma^2) = \ell$. Then, [Assumption 1](#) holds with $\mathcal{I}_b = \ell \mathcal{J}_b$, and consequently,*

$$\mathcal{J}_b \geq \mathcal{I}_b, \quad \text{i.e.,} \quad \text{eff}(\hat{\tau}_{\text{adj}}^{\text{R}}, \hat{\tau}^{\text{R}}) \geq 1.$$

Moreover, if it holds that $\liminf_{N \rightarrow \infty} N^{-1} \sum_{j=1}^N (v_{N,j} - \bar{v}_N)^2 > 0$, where $\bar{v}_N := N^{-1} \sum_{i=1}^N v_{N,i}$, then $\mathcal{J}_b > \mathcal{I}_b$, and consequently $\text{eff}(\hat{\tau}_{\text{adj}}^{\text{R}}, \hat{\tau}^{\text{R}})$ is strictly greater than 1.

In [Sections 4](#) and [5](#), we illustrate using both simulated and real datasets that the regression adjusted estimators lead to shorter confidence intervals than the unadjusted estimators, while not compromising on the desired significance level. We also observe in our simulations that, even when the regression model is not correctly specified, the regression adjusted estimator $\hat{\tau}_{\text{adj}}^{\text{R}}$ is at least as efficient as the unadjusted estimator $\hat{\tau}^{\text{R}}$, suggesting that [Theorem 3.4](#) might hold in a wider generality, akin to a result by [Lin \(2013, Corollary 1.1\)](#) on the efficiency gain by regression adjustment for ANOVA-based estimators.

4 Simulations

In this section we illustrate, through numerical experiments, the performance of $\hat{\tau}^{\text{R}}$ and $\hat{\tau}_{\text{adj}}^{\text{R}}$. We also compare their performance with those of some conventional estimators, namely, the difference-in-means estimator $\hat{\tau}_{\text{dm}}$ (see [\(2.10\)](#)), the simple regression adjusted estimator $\hat{\tau}_{\text{adj}}$ ([Freedman, 2008a,b](#)), and the estimator $\hat{\tau}_{\text{interact}}$ proposed by [Lin \(2013\)](#).

We take the total sample size as $N = 1000$, choose m subjects by SRSWOR and assign them to the treatment group. We experimented with various values of the proportion m/N ; here we only report the results for $m/N = 0.75, 0.50$, and 0.25 . Consider the settings listed in [Table 4.1](#). Setting 1 is an example where the potential outcomes are independent of the covariates. Setting 2 is an example where the regression model is correctly specified. Finally, Setting 3 is a slight modification of an example of [Lin \(2013, Setting 4.2.3\)](#), and helps us illustrate the performance of the estimators under misspecification of the regression model. In each of these settings, the i.i.d. errors ε_i 's are drawn from: (a) the standard normal distribution, (b) the t_1 distribution,

or (c) the t_3 distribution. Thus, we have 9 simulation settings in total, and we call the sub-settings as Setting 1b, 2a, etc. The potential control outcomes are generated as $b_i = a_i - \tau_0$, where $\tau_0 = 2$ is the true value of the constant treatment effect τ .

Table 4.1: Simulation settings.

Setting 1	$x_i \stackrel{i.i.d.}{\sim} \text{Unif}(-4, 4)$	$a_i = v_i + \varepsilon_i, v_i \stackrel{i.i.d.}{\sim} \text{Exp}(1/10)$
Setting 2	$x_i \stackrel{i.i.d.}{\sim} \text{Unif}(-4, 4)$	$a_i = 3x_i + \varepsilon_i$
Setting 3	$x_i = e^{u_i}, u_i \stackrel{i.i.d.}{\sim} \text{Unif}(-4, 4)$	$a_i = \frac{1}{4}(x_i + \sqrt{x_i}) + \varepsilon_i$

The confidence intervals based on $\hat{\tau}^R$ and $\hat{\tau}_{\text{adj}}^R$ are computed using the plug-in estimators of standard errors that we propose in [Appendix A](#). We also report the results on the oracle confidence intervals obtained by plugging in the true asymptotic variances. The confidence intervals constructed using [Corollary 2.1](#) are not reported here, as they are almost identical, in terms of lengths and coverages, to the confidence intervals constructed using the plug-in estimator of the asymptotic variance of $\hat{\tau}^R$ (see [Corollary A.1](#) in [Appendix A](#)).

We repeat the above experiment 1000 times for each of the sub-settings, and report the coverage and average lengths of the confidence intervals based on the five estimators mentioned above. The results are summarized in [Tables 4.2–4.4](#), and some observations are listed below.

- *Efficiency of $\hat{\tau}^R$ w.r.t. $\hat{\tau}_{\text{dm}}$* : In all the settings except Setting 2c, the confidence intervals constructed using the unadjusted estimator $\hat{\tau}^R$ have shorter lengths than those constructed using $\hat{\tau}_{\text{dm}}$, yet maintaining the desired significance level. Even for Setting 2c the differences in the lengths are within a margin dictated by [Theorem 2.4](#).
- *Regression adjustment improves precision*: For Setting 1, where the responses are drawn independently of the covariates, the confidence intervals based on $\hat{\tau}^R$ and $\hat{\tau}_{\text{adj}}^R$ have similar lengths, as we expect. For Setting 2a, where the model is correctly specified and the errors are Gaussian, the regression adjusted confidence intervals are shorter than the unadjusted ones, validating [Theorem 3.4](#) and reinforcing the popular belief (see, e.g., [Rosenbaum \(2002a\)](#)) that regression adjustment generally improves the precision over the unadjusted case.
- *Comparison of $\hat{\tau}_{\text{adj}}^R$ with $\hat{\tau}_{\text{adj}}$ and $\hat{\tau}_{\text{interact}}$* : In all of the settings, the lengths of our confidence intervals constructed using $\hat{\tau}_{\text{adj}}^R$ are either smaller than or (almost) equal to the lengths of the confidence intervals constructed using $\hat{\tau}_{\text{adj}}$ or $\hat{\tau}_{\text{interact}}$, without compromising on coverage.
- *Robustness against heavy-tails*: In each of the settings, when the errors are drawn from the t_1 distribution, the confidence intervals based on $\hat{\tau}_{\text{dm}}$, $\hat{\tau}_{\text{adj}}$ or $\hat{\tau}_{\text{interact}}$ are too wide. In contrast, the rank-based confidence intervals have reasonably shorter lengths, and provide coverage close to the nominal level.
- *Robustness against model misspecification*: In Setting 3, the regression model is misspecified. Yet, the performance of the rank-based confidence intervals is quite satisfactory, illustrating that the estimators $\hat{\tau}^R$ and $\hat{\tau}_{\text{adj}}^R$ are robust against model misspecification. Note also that in

this setting, despite the regression model being misspecified, the confidence intervals based on $\hat{\tau}_{\text{adj}}^{\text{R}}$ are shorter than those constructed using $\hat{\tau}^{\text{R}}$, suggesting that [Theorem 3.4](#) holds in more generality than what we assumed for our proof of the result.

Table 4.2: Results for simulation Setting 1 (potential outcomes independent of the covariates).

Estimator		(a) Gaussian errors			(b) t_1 errors			(c) t_3 errors		
		0.75	0.50	0.25	0.75	0.50	0.25	0.75	0.50	0.25
$\hat{\tau}^{\text{R}}$	coverage	0.94	0.96	0.95	0.95	0.96	0.96	0.94	0.95	0.95
	length	1.85	1.60	1.85	2.29	1.98	2.29	1.93	1.67	1.93
$\hat{\tau}^{\text{R}}$ (oracle)	coverage	0.94	0.96	0.95	0.95	0.96	0.96	0.94	0.95	0.95
	length	1.85	1.60	1.85	2.29	1.98	2.29	1.93	1.67	1.93
$\hat{\tau}_{\text{dm}}$	coverage	0.95	0.95	0.96	0.97	0.98	0.97	0.95	0.96	0.94
	length	2.88	2.49	2.88	57.62	46.47	43.10	2.91	2.52	2.91
$\hat{\tau}_{\text{adj}}^{\text{R}}$	coverage	0.94	0.96	0.95	0.94	0.96	0.96	0.95	0.95	0.95
	length	1.86	1.61	1.86	3.35	3.03	3.38	1.94	1.68	1.94
$\hat{\tau}_{\text{adj}}^{\text{R}}$ (oracle)	coverage	0.94	0.96	0.95	0.95	0.96	0.96	0.95	0.95	0.95
	length	1.86	1.61	1.86	3.48	3.01	3.48	1.94	1.68	1.94
$\hat{\tau}_{\text{adj}}$	coverage	0.95	0.95	0.96	0.97	0.98	0.97	0.95	0.96	0.94
	length	2.88	2.49	2.87	57.05	45.88	43.04	2.90	2.52	2.91
$\hat{\tau}_{\text{interact}}$	coverage	0.94	0.95	0.96	0.97	0.98	0.97	0.95	0.96	0.94
	length	2.87	2.49	2.87	56.41	45.85	42.90	2.90	2.52	2.90

Table 4.3: Results for simulation Setting 2 (regression model correctly specified).

Estimator		(a) Gaussian errors			(b) t_1 errors			(c) t_3 errors		
		0.75	0.50	0.25	0.75	0.50	0.25	0.75	0.50	0.25
$\hat{\tau}^{\text{R}}$	coverage	0.95	0.95	0.96	0.95	0.96	0.95	0.95	0.94	0.95
	length	2.08	1.81	2.08	2.43	2.11	2.43	2.13	1.85	2.13
$\hat{\tau}^{\text{R}}$ (oracle)	coverage	0.95	0.95	0.96	0.95	0.96	0.95	0.95	0.93	0.95
	length	2.09	1.81	2.09	2.43	2.11	2.43	2.13	1.85	2.13
$\hat{\tau}_{\text{dm}}$	coverage	0.95	0.95	0.96	0.98	0.97	0.98	0.95	0.94	0.96
	length	2.01	1.73	2.00	63.52	70.98	70.32	2.04	1.77	2.04
$\hat{\tau}_{\text{adj}}^{\text{R}}$	coverage	0.95	0.94	0.95	0.94	0.94	0.95	0.96	0.94	0.95
	length	0.29	0.25	0.29	2.30	1.97	2.31	0.36	0.31	0.36
$\hat{\tau}_{\text{adj}}^{\text{R}}$ (oracle)	coverage	0.95	0.94	0.95	0.94	0.94	0.95	0.96	0.95	0.95
	length	0.29	0.25	0.29	2.28	1.98	2.28	0.36	0.31	0.36
$\hat{\tau}_{\text{adj}}$	coverage	0.95	0.95	0.95	0.98	0.98	0.98	0.95	0.93	0.95
	length	0.29	0.25	0.29	63.53	70.51	69.85	0.48	0.42	0.49
$\hat{\tau}_{\text{interact}}$	coverage	0.95	0.95	0.95	0.98	0.98	0.98	0.95	0.93	0.95
	length	0.29	0.25	0.29	63.48	70.50	69.61	0.48	0.42	0.49

Table 4.4: Results for simulation Setting 3 (model misspecification).

Estimator		(a) Gaussian errors			(b) t_1 errors			(c) t_3 errors		
		0.75	0.50	0.25	0.75	0.50	0.25	0.75	0.50	0.25
$\hat{\tau}^R$	coverage	0.94	0.95	0.94	0.95	0.95	0.96	0.96	0.95	0.96
	length	0.56	0.48	0.56	0.89	0.77	0.89	0.65	0.56	0.65
$\hat{\tau}^R$ (oracle)	coverage	0.94	0.95	0.94	0.95	0.95	0.96	0.95	0.95	0.96
	length	0.56	0.48	0.56	0.89	0.77	0.89	0.65	0.56	0.65
$\hat{\tau}_{dm}$	coverage	0.95	0.94	0.94	0.99	0.98	0.97	0.95	0.95	0.95
	length	1.02	0.88	1.02	37.74	37.56	42.99	1.09	0.95	1.09
$\hat{\tau}_{adj}^R$	coverage	0.95	0.96	0.95	0.94	0.95	0.95	0.95	0.95	0.95
	length	0.30	0.26	0.30	0.71	0.61	0.71	0.36	0.31	0.36
$\hat{\tau}_{adj}^R$ (oracle)	coverage	0.96	0.96	0.95	0.94	0.95	0.94	0.95	0.95	0.95
	length	0.30	0.26	0.30	0.71	0.61	0.71	0.36	0.32	0.36
$\hat{\tau}_{adj}$	coverage	0.95	0.95	0.95	0.98	0.98	0.98	0.95	0.95	0.94
	length	0.29	0.25	0.29	37.86	37.66	43.07	0.49	0.43	0.49
$\hat{\tau}_{interact}$	coverage	0.95	0.95	0.96	0.98	0.97	0.98	0.95	0.95	0.95
	length	0.29	0.25	0.29	37.82	37.62	43.03	0.49	0.43	0.49

5 Data analysis

We demonstrate below the performance of the rank-based estimators $\hat{\tau}^R$ and $\hat{\tau}_{adj}^R$, and compare it with that of the regression-based estimators $\hat{\tau}_{dm}$, $\hat{\tau}_{adj}$ and $\hat{\tau}_{interact}$, on two real datasets, namely: (i) Mexico’s conditional cash transfer program (Progresa) data (De La O, 2013; Imai, 2018), and (ii) the house price data from the replication files of Linden and Rockoff (2008).

5.1 Progresa data

We analyze the data from a randomized trial that aims to study the electoral impact of *Progresa*, Mexico’s conditional cash transfer program (CCT program) (De La O, 2013; Imai, 2018). In this experiment, eligible villages were randomly assigned to receive the program either 21 months (early Progresa, “treated”) or 6 months (late Progresa, “control”) before the 2000 Mexican presidential election. The data contains 417 observations, each representing a precinct, and for each precinct we have information about its treatment status, outcomes of interest, socioeconomic indicators, and other precinct characteristics.

Following De La O (2013), we use the support rates for the incumbent party as shares of the eligible voting population in the 2000 election (*pri2000s*) as the outcome, and regress it on the following covariates: the average poverty level in a precinct (*avgpoverty*), the total precinct population in 1994 (*pobtot1994*), the total number of voters who turned out in the previous election (*votos1994*), and the total number of votes cast for each of the three main competing parties in the previous election (*pri1994*, *pan1994*, and *prd1994*), and *villages* (as

factors). [Table 5.1](#) gives the point estimates and approximate 95% confidence intervals for the five methods. The confidence intervals suggest that the CCT program led to a significant positive increase in support for the incumbent party. It is noteworthy that the standard errors of Rosenbaum’s estimators are substantially smaller than that of $\hat{\tau}_{\text{dm}}$, $\hat{\tau}_{\text{adj}}$, or $\hat{\tau}_{\text{interact}}$. Further, the standard error for $\hat{\tau}_{\text{adj}}^{\text{R}}$ is slightly less than that of $\hat{\tau}^{\text{R}}$, as suggested by [Theorem 3.4](#).

Table 5.1: Different estimates of the effect of early Progresa on PRI support rates with the corresponding standard errors, 95% approximate confidence intervals and their lengths.

	estimate	std.error	95% confidence interval	length
$\hat{\tau}^{\text{R}}$	1.834	0.446	[0.960, 2.707]	1.747
$\hat{\tau}_{\text{dm}}$	3.622	1.728	[0.235, 7.010]	6.774
$\hat{\tau}_{\text{adj}}^{\text{R}}$	2.185	0.411	[1.380, 2.989]	1.610
$\hat{\tau}_{\text{adj}}$	3.671	1.510	[0.712, 6.630]	5.917
$\hat{\tau}_{\text{interact}}$	4.214	1.462	[1.348, 7.079]	5.731

5.2 House price data

Our second example is the house price data from the replication files of [Linden and Rockoff \(2008\)](#), which contain data on property sales for Mecklenburg County (North Carolina) between January 1994 and December 2004. Dropping the sales below \$5,000 and above \$1,000,000, the dataset comes with 170,239 observations. Exploratory data analysis shows that even after taking logarithm of house prices, the distribution is heavily skewed on the right side with kurtosis equal to 5.1, so we anticipate that the rank-based estimators will yield shorter confidence intervals than the regression-based estimators.

We create synthetic data sets from this data set to compare the different estimators. Following [Athey et al. \(2021\)](#), we draw subsamples from the dataset and randomly assign exactly half of each sample to the treatment group and the remaining half to the control group. Thus, we know apriori that the treatment effect is zero, and hence can assess the performance of the five estimators in a different way than the previous example. In particular, we repeat this experiment and evaluate the performance of the estimators based on the coverage and average lengths of the confidence intervals.

In our simulations, we draw subsamples of size $n = 1000$ in each replication, take the logarithm of the house prices as the outcome variable, and use several characteristics of the houses (e.g., sales year, age of the house (in years), number of bedrooms) as covariates. The fit of these models are found to be quite satisfactory, with adjusted R^2 values near to 0.7. The estimates, along with their standard errors and approximate 95% confidence intervals obtained from a single simulation are shown in [Table 5.2](#). Repeating this experiment $B = 1000$ times, we report the coverage and average lengths of the confidence intervals in [Table 5.3](#).

Observe that the confidence intervals obtained using $\hat{\tau}_{\text{adj}}^{\text{R}}$ are the shortest among the five, having about half the length than those obtained using $\hat{\tau}_{\text{adj}}$ or $\hat{\tau}_{\text{interact}}$, without any compromise in coverage. Also, the confidence intervals obtained using $\hat{\tau}^{\text{R}}$ are shorter than those obtained

using $\hat{\tau}_{\text{dm}}$, and are wider than those obtained using $\hat{\tau}_{\text{adj}}^{\text{R}}$. These observations are completely aligned with our main theoretical contributions.

Table 5.2: Results from a single simulation from the house price data.

	estimate	std.error	95% confidence interval	length
$\hat{\tau}^{\text{R}}$	-0.04	0.03	[-0.09, 0.02]	0.12
$\hat{\tau}_{\text{dm}}$	-0.05	0.04	[-0.12, 0.03]	0.15
$\hat{\tau}_{\text{adj}}^{\text{R}}$	-0.01	0.01	[-0.03, 0.02]	0.04
$\hat{\tau}_{\text{adj}}$	-0.02	0.02	[-0.06, 0.03]	0.08
$\hat{\tau}_{\text{interact}}$	-0.02	0.02	[-0.06, 0.03]	0.08

Table 5.3: Coverage and average lengths of the approximate 95% confidence intervals obtained from different estimators by repeated simulations from the house price data.

	$\hat{\tau}^{\text{R}}$	$\hat{\tau}_{\text{dm}}$	$\hat{\tau}_{\text{adj}}^{\text{R}}$	$\hat{\tau}_{\text{adj}}$	$\hat{\tau}_{\text{interact}}$
coverage	0.944	0.959	0.954	0.952	0.949
average length	0.121	0.143	0.041	0.073	0.073

6 Discussion

Randomization remains the gold standard method in causal inference for balancing covariates across treatment groups. Large randomization experiments are becoming increasingly popular, see recent discussions in [Banerjee et al. \(2016\)](#) and [Deaton and Cartwright \(2018\)](#). In this paper, we studied Rosenbaum’s unadjusted and regression adjusted rank-based estimators (see [Rosenbaum \(2002a\)](#)) for estimating the constant treatment effect under randomization inference. Our results highlight many benefits of these estimators in terms of their robustness properties and efficiency relative to the standard regression (un)-adjusted mean-based estimators. In particular, we showed that the asymptotic relative efficiency of Rosenbaum’s unadjusted estimator is in the worst case only $\sim 15\%$ lower than the unadjusted difference-in-means estimator, but often the efficiency is much higher. We also showed that regression adjustment gives provable efficiency gain for Rosenbaum’s estimator under a linear outcome model. Additionally, our detailed simulation study illustrates the superior performance of these estimators for normal and heavy-tailed errors, and for linear and non-linear models.

It would be interesting to investigate the properties of the rank-based estimators when the constant treatment effect assumption is violated. There are statistical methods for testing this assumption (see, e.g., [Ding et al. \(2016\)](#)). When there is evidence for heterogeneity of the treatment effect, additional methods and/or studies are needed to understand how and to whom the treatment is more beneficial. We may consider relaxing the constant treatment effect assumption in the following ways. First, it may happen that our data or our existing knowledge indicates that the treatment interacts with a covariate. In that case, we can incorporate this

information in our inference by: (a) changing the design (e.g., using blocked randomization that blocks on the interacting covariate instead of complete randomization as studied in this paper), or (b) changing the model for the adjusted responses (see [Rosenbaum, 2021](#), Section 1.8 for general adjustment). Our results may be extended in these situations to construct confidence intervals for the treatment effect after conditioning on covariates. Second, [Caughey et al. \(2021\)](#) provide methods to construct confidence intervals for quantiles of individual treatment effects by inverting randomization based tests. Following their approach, it will be interesting to generalize our methods to target the quantiles of individual treatment effects.

Finally, the average treatment effect across the units is a popular choice as a causal estimand. But, this estimand may be undesirable, for example, when only a fraction of the units have a large positive treatment effect and others have no effect or negative effect; see, e.g., [Athey et al. \(2021\)](#) and [Rosenbaum \(2021, Section 1.3.4\)](#). In future, we plan to study a more appropriate estimand under such situations and develop rank-based statistical inference procedures targeting this estimand.

Acknowledgement

Bodhisattva Sen would like to thank Michael Sobel for his valuable feedback on an early draft.

References

- Aronow, P. M., D. P. Green, and D. K. K. Lee (2014). Sharp bounds on the variance in randomized experiments. *The Annals of Statistics* 42(3), 850–871.
- Athey, S., P. J. Bickel, A. Chen, G. Imbens, and M. Pollmann (2021). Semiparametric estimation of treatment effects in randomized experiments. Working Paper 29242, National Bureau of Economic Research.
- Banerjee, A. V., E. Duflo, and M. Kremer (2016). The influence of randomized controlled trials on development economics research and on development policy. *The State of Economics, The State of the World*, 482–488. Edited by Kaushik Basu.
- Bind, M.-A. C. and D. B. Rubin (2020). When possible, report a Fisher-exact P value and display its underlying null randomization distribution. *Proceedings of the National Academy of Sciences of the United States of America* 117(32), 19151–19158.
- Boucheron, S., O. Bousquet, G. Lugosi, and P. Massart (2005). Moment inequalities for functions of independent random variables. *The Annals of Probability* 33(2), 514–560.
- Caughey, D., A. Dafoe, X. Li, and L. Miratrix (2021). Randomization Inference beyond the Sharp Null: Bounded Null Hypotheses and Quantiles of Individual Treatment Effects. *arXiv preprint arXiv:2101.09195*.
- Chernozhukov, V., D. Chetverikov, and K. Kato (2014). Gaussian approximation of suprema of empirical processes. *The Annals of Statistics* 42(4), 1564–1597.

- Conover, W. J. (1999). *Practical Nonparametric Statistics* (3rd ed.). John Wiley & Sons.
- Cox, D. R. and N. Reid (2000). *The Theory of The Design of Experiments*. Chapman and Hall/CRC.
- De La O, A. L. (2013). Do conditional cash transfers affect electoral behavior? Evidence from a randomized experiment in Mexico. *American Journal of Political Science* 57(1), 1–14.
- Deaton, A. and N. Cartwright (2018). Understanding and misunderstanding randomized controlled trials. *Social Science & Medicine* 210, 2–21.
- Ding, P., A. Feller, and L. Miratrix (2016). Randomization inference for treatment effect variation. *Journal of the Royal Statistical Society. Series B. Statistical Methodology* 78(3), 655–671.
- Efron, B. and C. Stein (1981). The jackknife estimate of variance. *The Annals of Statistics* 9(3), 586–596.
- Fisher, R. A. (1935). *The Design of Experiments*. Oliver and Boyd, Edinburgh.
- Fogarty, C. B. (2018). Regression-assisted inference for the average treatment effect in paired experiments. *Biometrika* 105(4), 994–1000.
- Freedman, D., R. Pisani, and R. Purves (2007). *Statistics* (4th ed.). W.W. Norton & Company.
- Freedman, D. A. (2008a). On regression adjustments in experiments with several treatments. *The Annals of Applied Statistics* 2(1), 176–196.
- Freedman, D. A. (2008b). On regression adjustments to experimental data. *Advances in Applied Mathematics* 40(2), 180–193.
- Gail, M. H., W. Y. Tan, and S. Piantadosi (1988). Tests for no treatment effect in randomized clinical trials. *Biometrika* 75(1), 57–64.
- Hettmansperger, T. P. and J. W. McKean (2011). *Robust nonparametric statistical methods* (2nd ed.), Volume 119 of *Monographs on Statistics and Applied Probability*. CRC Press, Boca Raton, FL.
- Ho, D. E. and K. Imai (2006). Randomization inference with natural experiments: An analysis of ballot effects in the 2003 California recall election. *Journal of the American Statistical Association* 101(475), 888–900.
- Hodges, Jr., J. L. and E. L. Lehmann (1956). The efficiency of some nonparametric competitors of the t -test. *The Annals of Mathematical Statistics* 27, 324–335.
- Hodges, Jr., J. L. and E. L. Lehmann (1963). Estimates of location based on rank tests. *The Annals of Mathematical Statistics* 34, 598–611.
- Hoeffding, W. (1951). A combinatorial central limit theorem. *The Annals of Mathematical Statistics* 22, 558–566.
- Huber, P. J. (1967). The behavior of maximum likelihood estimates under nonstandard conditions. In *Proc. Fifth Berkeley Sympos. Math. Statist. and Probability, (Berkeley, Calif., 1965/66), Vol. I: Statistics*, pp. 221–233.

- Imai, K. (2018). *Quantitative social science: An introduction*. Princeton University Press.
- Imbens, G. W. and D. B. Rubin (2015). *Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction*. Cambridge University Press, New York.
- Lehmann, E. L. (1975). *Nonparametrics: Statistical methods based on ranks*. Holden-Day Series in Probability and Statistics. Holden-Day, Inc., San Francisco, Calif.; McGraw-Hill International Book Co., New York-Düsseldorf. With the special assistance of H. J. M. d’Abrera.
- Lehmann, E. L. and J. P. Romano (2005). *Testing statistical hypotheses* (3rd ed.). Springer Texts in Statistics. Springer, New York.
- Lei, L. and P. Ding (2021). Regression adjustment in completely randomized experiments with a diverging number of covariates. *Biometrika* 108(4), 815–828.
- Li, X. and P. Ding (2017). General forms of finite population central limit theorems with applications to causal inference. *Journal of the American Statistical Association* 112(520), 1759–1769.
- Li, X. and P. Ding (2020). Rerandomization and regression adjustment. *Journal of the Royal Statistical Society. Series B. Statistical Methodology* 82(1), 241–268.
- Li, X., P. Ding, and D. B. Rubin (2018). Asymptotic theory of rerandomization in treatment-control experiments. *Proceedings of the National Academy of Sciences of the United States of America* 115(37), 9157–9162.
- Lin, W. (2013). Agnostic notes on regression adjustments to experimental data: Reexamining Freedman’s critique. *The Annals of Applied Statistics* 7(1), 295–318.
- Linden, L. and J. E. Rockoff (2008). Estimates of the impact of crime risk on property values from Megan’s laws. *American Economic Review* 98(3), 1103–27.
- Liu, H. and Y. Yang (2020). Regression-adjusted average treatment effect estimates in stratified randomized experiments. *Biometrika* 107(4), 935–948.
- Luo, X., T. Dasgupta, M. Xie, and R. Y. Liu (2021). Leveraging the Fisher randomization test using confidence distributions: Inference, combination and fusion learning. *Journal of the Royal Statistical Society. Series B. Statistical Methodology* 83(4), 777–797.
- MacKinnon, J. G. and H. White (1985). Some heteroskedasticity-consistent covariance matrix estimators with improved finite sample properties. *Journal of Econometrics* 29(3), 305–325.
- Rosenbaum, P. R. (1993). Hodges-Lehmann point estimates of treatment effect in observational studies. *Journal of the American Statistical Association* 88(424), 1250–1253.
- Rosenbaum, P. R. (2002a). Covariance adjustment in randomized experiments and observational studies. *Statistical Science* 17(3), 286–327. With comments and a rejoinder by the author.
- Rosenbaum, P. R. (2002b). *Observational studies* (2nd ed.). Springer Series in Statistics. Springer-Verlag, New York.

- Rosenbaum, P. R. (2021). *Replication and Evidence Factors in Observational Studies*. Chapman and Hall/CRC.
- Rubin, D. B. (1974). Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology* 66(5), 688–701.
- Rubin, D. B. (1977). Assignment to treatment group on the basis of a covariate. *Journal of Educational Statistics* 2(1), 1–26.
- Splawa-Neyman, J. (1990). On the application of probability theory to agricultural experiments. Essay on principles. Section 9. *Statistical Science* 5(4), 465–472. Translated from the Polish and edited by D. M. Dąbrowska and T. P. Speed.
- Su, F. and P. Ding (2021). Model-assisted analyses of cluster-randomized experiments. *Journal of the Royal Statistical Society. Series B. Statistical Methodology* 83(5), 994–1015.
- Tukey, J. W. (1993). Tightening the clinical trial. *Controlled clinical trials* 14(4), 266–285.
- van der Vaart, A. W. (1998). *Asymptotic Statistics*. Cambridge University Press, Cambridge.
- White, H. (1980). A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica* 48(4), 817–838.
- Wilcoxon, F. (1945). Individual comparisons by ranking methods. *Biometrics Bulletin* 1(6), 80–83.
- Wittes, J. and E. Brittain (1990). The role of internal pilot studies in increasing the efficiency of clinical trials. *Statistics in Medicine* 9(1-2), 65–72.
- Wu, J. and P. Ding (2021). Randomization tests for weak null hypotheses in randomized experiments. *Journal of the American Statistical Association* 116(536), 1898–1913.
- Young, A. (2018). Channeling Fisher: Randomization Tests and the Statistical Insignificance of Seemingly Significant Experimental Results. *The Quarterly Journal of Economics* 134(2), 557–598.
- Zhao, A. and P. Ding (2022). Regression-based causal inference with factorial experiments: estimands, model specifications and design-based properties. *Biometrika* 109(3), 799–815.

Appendices [A–F](#)

This supplementary material begins with a plug-in method we propose in [Appendix A](#) to consistently estimate the asymptotic variance of the regression adjusted estimator $\hat{\tau}_{\text{adj}}^{\text{R}}$ studied in our main paper. In [Appendix B](#) we briefly outline the main ideas of the proofs of our major results. The subsequent sections ([Appendices C–E](#)) contain the proofs of all results in the main paper, other auxiliary results (with their proofs) and further discussions. Finally, [Appendix F](#) talks about the analogous results that hold for average ranks instead of the up-ranks we used in the main paper.

If $\{x_n\}_{n \geq 1}$ and $\{y_n\}_{n \geq 1}$ are two sequence of positive real numbers, we write $x_n \sim y_n$ to denote that $\lim_{n \rightarrow \infty} x_n/y_n = 1$, and $x_n \lesssim y_n$ to denote that $x_n \leq C y_n$ holds for all sufficiently large n , for some constant $C > 0$.

A Consistent plug-in estimators of standard errors

In [Theorems 2.2](#) and [3.3](#) of the main paper, we showed that $\sqrt{N}(\hat{\tau}^R - \tau_0)$ and $\sqrt{N}(\hat{\tau}_{\text{adj}}^R - \tau_0)$ both have centered normal limits with appropriate variances. But these limiting variances depend on certain properties of the unobserved (fixed) potential outcomes through the quantities \mathcal{I}_b and \mathcal{J}_b (see [Assumptions 1](#) and [2](#)) and hence, are unknown. The main challenge in obtaining confidence intervals, therefore, would be to estimate these unknown variances consistently. Unfortunately, our proposed method for estimating \mathcal{I}_b (see [Theorem 2.3](#) of the main paper) does not have a natural extension in the regression-adjusted case. In this section, we propose an alternate method that applies to both the unadjusted and the adjusted cases.

A.1 Consistent estimation of the standard error of $\hat{\tau}^R$

In this section, we propose a plug-in estimator of \mathcal{I}_b that uses all the observed data and not just the control outcomes, and can also be generalized to estimate \mathcal{J}_b in the regression adjusted case. Using the relation $b_{N,j} = Y_{N,j} - \tau_0 Z_{N,j}$, we define a plug-in estimator of $b_{N,j}$ as

$$\hat{b}_{N,j} := Y_{N,j} - \hat{\tau}_N Z_{N,j} = b_{N,j} - (\hat{\tau}_N - \tau_0) Z_{N,j}, \quad (\text{A.1})$$

where $\hat{\tau}_N$ is an estimator of τ . At this point, the next intuitive step would be to replace $(b_{N,1}, \dots, b_{N,N})$ with $(\hat{b}_{N,1}, \dots, \hat{b}_{N,N})$ in [Assumption 1](#) of the main paper, i.e., replace the term $I_{h,N}(b_{N,j} - b_{N,i})$ on the left hand side of [\(2.16\)](#), with $I_{h,N}(\hat{b}_{N,j} - \hat{b}_{N,i})$ to construct the estimator. However this leads to some technical problems in proving consistency as $I_{h,N}(\cdot)$ is a discontinuous function which itself changes with N . This requires us to refine our estimation strategy further and impose a slightly stronger condition than that in [Assumption 1](#). Towards this, for $\nu > 0$ and $N \geq 1$, define

$$I_{h,N,\nu}(x) := \begin{cases} \mathbf{1}\{0 \leq x < hN^{-\nu}\} & \text{if } h \geq 0, \\ -\mathbf{1}\{hN^{-\nu} \leq x < 0\} & \text{if } h < 0. \end{cases} \quad (\text{A.2})$$

Assumption 3. For $h \in \mathbb{R}$ let $I_{h,N,\nu}$ be defined in [\(A.2\)](#). We assume that there exists $0 < \nu < 1/2$, the following holds for every $u \in [\nu, 1/2]$ that

$$N^{-(2-u)} \sum_{j=1}^N \sum_{i=1}^N I_{h,N,u}(b_{N,j} - b_{N,i}) \rightarrow h\mathcal{I}_b$$

for some constant $\mathcal{I}_b \in (0, \infty)$, which is the same \mathcal{I}_b as in [Assumption 1](#).

Remark A.1 (On [Assumption 3](#)). Note that if we were to set $\nu = 1/2$ in [Assumption 3](#), then $I_{h,N,\nu}(\cdot) \equiv I_{h,N,1/2}(\cdot)$ is exactly the same function as $I_{h,N}(\cdot)$ from (2.15), and consequently [Assumption 3](#) would then imply [Assumption 1](#). If we define

$$T_{h,N,u} := N^{-(2-u)} \sum_{j=1}^N \sum_{i=1}^N I_{h,N,u}(b_{N,j} - b_{N,i}), \quad (\text{A.3})$$

then [Assumption 1](#) requires $T_{h,N,1/2} \rightarrow h\mathcal{I}_b$ whereas [Assumption 3](#) requires the mildly stronger condition $T_{h,N,u} \rightarrow h\mathcal{I}_b$ for u varying in any non-degenerate interval with right endpoint at $1/2$. We firmly believe that this is a reasonable assumption. For instance, one of the ways we justified [Assumption 1](#) was by showing that it is satisfied when $b_{N,i}$'s are randomly sampled from an absolutely continuous distribution (see [Remark 2.3](#)). The same is also true for [Assumption 3](#), see [Lemma D.16](#) for a formal result.

Finding a consistent estimator of \mathcal{I}_b is now quite intuitive. With $T_{h,N,u}$ defined as in (A.3), [Assumption 3](#) implies that $T_{1,N,\nu}$ converges to \mathcal{I}_b , where ν is as in [Assumption 3](#). Consequently, we construct an estimator \widehat{V}_N from $T_{1,N,\nu}$ using the plug-in principle as described earlier in this section, by replacing $b_{N,i}$'s in (A.3) with $\widehat{b}_{N,i}$'s (see (A.1)). The following result makes it precise.

Theorem A.1 (Consistent estimation of \mathcal{I}_b). Define $\widehat{b}_{N,j}$ as in (A.1), with $\widehat{\tau}_N \equiv \widehat{\tau}^R$. Suppose that [Assumption 3](#) holds for some $\nu \in (0, 1/2)$, and define

$$\widehat{V}_N := N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{0 \leq \widehat{b}_{N,j} - \widehat{b}_{N,i} < N^{-\nu}\right\}. \quad (\text{A.4})$$

Then $\widehat{V}_N \xrightarrow{P} \mathcal{I}_b$ as $N \rightarrow \infty$.

Combining [Theorems 2.2](#) and [A.1](#) we get an asymptotically valid confidence interval for τ_0 . This is formally stated in the following corollary.

Corollary A.1 (Confidence interval for τ_0 based on $\widehat{\tau}^R$). Under [Assumptions 1](#) and [3](#), an approximate $100(1 - \alpha)\%$ confidence interval for τ_0 is given by

$$\widehat{\tau}^R \pm \frac{z_{\alpha/2}}{\sqrt{N}} \left(12 \frac{m}{N} \left(1 - \frac{m}{N}\right) \widehat{V}_N^2\right)^{-1/2}, \quad (\text{A.5})$$

where \widehat{V}_N is defined in (A.4).

We refer the reader to [Remark A.2](#) below for a discussion on the choice of ν in the above result.

A.2 Consistent estimation of the standard error of $\widehat{\tau}_{\text{adj}}^{\text{R}}$

For estimating τ_0 (the true value of τ) under regression adjustment, we follow the same road map as in the unadjusted case in the previous section. Once again, due to technical reasons we require a mildly stronger condition than [Assumption 2](#) of the main paper to come up with a consistent estimator for \mathcal{J}_b . To lay the groundwork, define

$$\widetilde{T}_{h,N,u} := N^{-(2-u)} \sum_{j=1}^N \sum_{i=1}^N I_{h,N,u}(\widetilde{b}_{N,j} - \widetilde{b}_{N,i}), \quad (\text{A.6})$$

where $\widetilde{b}_{N,j}$'s are defined as in (3.6) and $I_{h,N,u}(\cdot)$ is defined as in (A.2). Observe the direct correspondence between $\widetilde{T}_{h,N,u}$ and $T_{h,N,u}$ from (A.3). The only difference is that the potential control outcomes $b_{N,j}$'s are replaced by the regression adjusted potential control outcomes $\widetilde{b}_{N,j}$'s. Note also that [Assumption 2](#) can be restated as $\widetilde{T}_{h,N,1/2} \rightarrow h\mathcal{J}_b$. With this observation, in the same vein as [Assumption 3](#), we now state our mildly stronger condition (compared to [Assumption 2](#)):

Assumption 4. *There exists $0 < \nu < 1/2$, such that for every $u \in [\nu, 1/2]$ and $h \in \mathbb{R}$, $\widetilde{T}_{h,N,u} \rightarrow h\mathcal{J}_b$, for some constant $\mathcal{J}_b \in (0, \infty)$, which is the same \mathcal{J}_b as in [Assumption 2](#).*

One can show that [Assumption 4](#) holds under the same regression model assumption as in [Theorem 3.4](#) (see the arguments used in [Lemma D.8](#)).

We are now in the position to construct our consistent estimator for \mathcal{J}_b . Towards this direction, recall the definition of $\widehat{b}_{N,j}$ from (A.1) and set $\widehat{\mathbf{b}}_N := (\widehat{b}_{N,1}, \dots, \widehat{b}_{N,N})$. To carry out the plug-in approach, we define the empirical analogues of $\widetilde{b}_{N,j}$'s, as follows:

$$\widehat{\widetilde{b}}_{N,j} := \widehat{b}_{N,j} - \mathbf{p}_{N,j}^\top \widehat{\mathbf{b}}_N = Y_{N,j} - \widehat{\tau}_{\text{adj}}^{\text{R}} Z_{N,j} - \mathbf{p}_{N,j}^\top (\mathbf{Y} - \widehat{\tau}_{\text{adj}}^{\text{R}} \mathbf{Z}), \quad j = 1, 2, \dots, N. \quad (\text{A.7})$$

The plug-in (consistent) estimator for \mathcal{J}_b is now constructed in the same manner as we did for \mathcal{I}_b , only with the modification that here $\widehat{\widetilde{b}}_{N,j}$'s will play the role of $\widehat{b}_{N,j}$'s. This is the content of the following theorem.

Theorem A.2 (Consistent estimation of \mathcal{J}_b). *Define $\widehat{\widetilde{b}}_{N,j}$ as in (A.7), and suppose that [Assumption 4](#) holds for some $\nu \in (0, 1/2)$. Define*

$$\widehat{W}_N := N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ 0 \leq \widehat{\widetilde{b}}_{N,j} - \widehat{\widetilde{b}}_{N,i} < N^{-\nu} \right\}. \quad (\text{A.8})$$

Then $\widehat{W}_N \xrightarrow{P} \mathcal{J}_b$ as $N \rightarrow \infty$.

[Theorem A.2](#) in conjunction with [Theorem 3.3](#) readily yields an asymptotically valid confidence interval for τ_0 based on $\widehat{\tau}_{\text{adj}}^{\text{R}}$, which is formally stated below.

Corollary A.2 (Confidence interval for τ_0 based on $\hat{\tau}_{\text{adj}}^{\text{R}}$). Under [Assumptions 2 and 4](#), an approximate $100(1 - \alpha)\%$ confidence interval for τ_0 is given by

$$\hat{\tau}_{\text{adj}}^{\text{R}} \pm \frac{z_{\alpha/2}}{\sqrt{N}} \left(12 \frac{m}{N} \left(1 - \frac{m}{N} \right) \widehat{W}_N^2 \right)^{-1/2}, \quad (\text{A.9})$$

where \widehat{W}_N is defined in [\(A.8\)](#).

The following remark highlights that the choice of the tuning parameter ν is inconsequential for both the regression unadjusted and regression adjusted confidence intervals that we proposed above.

Remark A.2 (Choice of ν). Our plug-in estimators proposed in [\(A.4\)](#) and [\(A.8\)](#) both require a choice of the tuning parameter $\nu \in (0, 1/2)$. We have observed through extensive simulations that the confidence intervals are not very sensitive to the choice of ν . In fact, it seems that the choice $\nu = 1/2$ also works in practice, although our proof of [Theorem A.1](#) precludes that choice.

For an illustration, consider our simulation setting 3a described in [Section 4](#) (see [Table 4.1](#)). We calculate the plug-in estimator of the asymptotic variances of $\hat{\tau}^{\text{R}}$ and $\hat{\tau}_{\text{adj}}^{\text{R}}$ for different values of ν , ranging from $1/4$ to $1/2$. Then fixing one particular value of ν as a reference, say $\nu = 1/3$, we calculate the ratios of these estimates for $\nu = 1/3$ to those for the other values of ν . Repeating this 100 times, we draw the boxplots of the ratios thus obtained, which is provided in [Fig. A.1](#).

The plots suggest that the estimates of the asymptotic variance do not differ much with the choice of ν . We observed similar phenomenon for the other simulation settings as well. Furthermore, in our simulations we also varied ν while calculating the confidence intervals for τ_0 proposed above, and found that the results do not vary considerably with the choice of ν . Thus the choice of ν seems to be unimportant, and in practice one may use any number in $(0, 1/2)$ as the value of ν ; in our simulation studies in [Section 4](#) we use $\nu = 1/3$.




Figure A.1: Boxplots of the ratios of the plug-in estimators of asymptotic variances of $\hat{\tau}^{\text{R}}$ (left panel) and $\hat{\tau}_{\text{adj}}^{\text{R}}$ (right panel) for $\nu = 1/3$ and various other values of ν , in simulation setting 4a (see [Section 4](#) for details on the simulation setup).

Remark A.3 (Scaling the outcome variable). *Since our method involves counts of differences in the outcome variable (or the fitted residuals) in shrinking intervals, we might not get a reasonable estimate if the responses are typically large in magnitude. This is why we suggest to scale (or modify) the outcome variable appropriately while using our variance estimators in practice. For instance, in [Section 5.2](#) of the main paper, where we apply our method on a house price data, we use the logarithm of the house prices as the outcome variable since the house prices are typically large in magnitude.*

B Main technical tools

In this section, we discuss the main tools used in the proofs of our main results in [Sections 2](#) and [3](#) of the main paper and [Appendix A](#). We begin with [Proposition 2.1](#) where $\tau = \tau_0$. In this case, it is easy to check that

$$t_N \equiv t_N^0 = m + \sum_{j=1}^N Z_{N,j} \sum_{i=1, i \neq j}^N \mathbf{1}\{b_{N,i} \leq b_{N,j}\}.$$

Note that t_N^0 is a weighted linear combination of $Z_{N,j}$'s, where the weights are non-random. We use the classical Hoeffding's combinatorial CLT (see [Hoeffding \(1951\)](#); also see [Appendix C.1](#) for more details) to complete the proof of [Proposition 2.1](#). The case of local alternatives, i.e., where $\tau = \tau_N := \tau_0 - h/\sqrt{N}$, is more challenging. This is the subject of [Theorem 2.1](#) of the main paper. Note that under $\tau = \tau_N$, we have:

$$t_N \equiv t_N^h = m + \sum_{j=1}^N Z_{N,j} \sum_{i=1, i \neq j}^N \mathbf{1}\{b_{N,i} - hN^{-1/2}Z_{N,i} \leq b_{N,j} - hN^{-1/2}Z_{N,j}\}.$$

When $h \neq 0$, t_N^h is a weighted linear combination of $Z_{N,j}$'s with random weights depending on the $Z_{N,j}$'s themselves (unlike what we observed for t_N^0). Therefore classical combinatorial CLTs do not apply. Further, as we are in the fixed design setting, the models under $\tau = \tau_0$ and $\tau = \tau_N$ lack the desirable forms for us to use the standard contiguity arguments due to Le Cam for deriving local asymptotic normality (see [van der Vaart \(1998, Chapter 7\)](#) for details). We tackle these challenges with the aid of a special decomposition which is the subject of the following proposition (see [Appendix C.13](#) for a proof).

Proposition B.1. *Let $t_N = t_N(\mathbf{Z}_N, \mathbf{Y}_N - \tau_0 \mathbf{Z}_N)$ be the Wilcoxon rank-sum statistic based on any random treatment assignment, where $t(\cdot, \cdot)$ is as in [\(2.4\)](#) of the main paper. Fix $h \in \mathbb{R}$ and let $\tau_N = \tau_0 - hN^{-1/2}$. It holds under $\tau = \tau_N$ that*

$$t_N \equiv t_N^h \stackrel{d}{=} t_N^0 - \gamma_N^h, \quad \text{where} \quad \gamma_N^h := \sum_{j=1}^N \sum_{i=1, i \neq j}^N (1 - Z_{N,i}) Z_{N,j} I_{h,N}(b_{N,j} - b_{N,i})$$

where $I_{h,N}$ is defined in (2.15) of the main paper.

Based on Proposition B.1, the proof of Theorem 2.1 can be completed via Slutsky's Theorem if we show the following:

$$N^{-3/2} \left(t_N^0 - \frac{m(N+1)}{2} \right) \xrightarrow{d} \mathcal{N} \left(0, \frac{\lambda(1-\lambda)}{12} \right) \quad \text{and} \quad \gamma_N^h \xrightarrow{P} -h\lambda(1-\lambda)\mathcal{I}_b,$$

where \mathcal{I}_b is defined in Assumption 1 of the main paper. The asymptotic fluctuation of t_N^0 , in the above display, follows directly from Proposition 2.1 as already discussed above. To prove the limit of γ_N^h , we use the standard Markov's inequality coupled with Assumption 1 (see Appendix C.3 for more details). Once the asymptotic distribution of $t_N \equiv t_N^h$ is obtained, the same for $\hat{\tau}^R$ as stated in Theorem 2.2, follows as a direct consequence of a classical Hodges and Lehmann argument (see Hodges and Lehmann (1963); also see Theorem 2.1 and Lemma D.1).

The next important question is obtaining asymptotically honest confidence intervals for τ_0 (as presented in Theorem A.1 of the main paper). Based on Theorem 2.2, this reduces to obtaining a consistent estimator for \mathcal{I}_b , which by Assumption 3, is the limit of u_N where

$$u_N := N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}(Y_{N,j} - Y_{N,i} - \tau_0(Z_{N,j} - Z_{N,i}) \geq 0) \right. \\ \left. - \mathbf{1}(Y_{N,j} - Y_{N,i} - \tau_0(Z_{N,j} - Z_{N,i}) \geq N^{-\nu}) \right).$$

The natural plug-in estimator can then be constructed by replacing τ_0 by $\hat{\tau}^R$ above, to get a plug-in analogue of u_N , say \hat{u}_N^R . While \hat{u}_N^R should intuitively be consistent, there are two technical issues: (a) indicators are not continuous functions, and (b) the sets in the indicators used above are not fixed, but effectively shrinking with N (they are of the form $[0, N^{-\nu}]$). These technical issues preclude the possibility of using a continuous mapping type argument to establish consistency. We circumvent this by approximating the indicators in u_N with Gaussian mollifiers with an appropriately chosen variance parameter. In particular, setting $\Phi(\cdot)$ as the standard Gaussian distribution function and $\sigma_{N,\nu} := \log N/N^\nu$, define

$$\hat{u}_N^G := \frac{1}{N^{3/2}} \sum_{i,j=1}^N \left(\Phi(\sigma_{N,\nu}^{-1}(Y_{N,j} - Y_{N,i} - \hat{\tau}^R(Z_{N,j} - Z_{N,i}))) \right. \\ \left. - \Phi(\sigma_{N,\nu}^{-1}(Y_{N,j} - Y_{N,i} - \hat{\tau}^R(Z_{N,j} - Z_{N,i} - N^{-\nu}))) \right).$$

Using some careful truncation arguments, we then show that $\hat{u}_N^G - \hat{u}_N^R \xrightarrow{P} 0$ and $\hat{u}_N^G - u_N \xrightarrow{P} 0$, which when combined, yield $\hat{u}_N^R \xrightarrow{P} \mathcal{I}_b$.

Our next important result in the unadjusted setting is the efficiency lower bound for $\hat{\tau}^R$ when compared with $\hat{\tau}_{\text{dm}}$, as presented in Theorem 2.4 of the main paper. Fortunately, the heavy lifting is done by Theorem 2.2, Li and Ding (2017, Theorem 5), and Hodges and Lehmann (1956, Theorem 1); see Appendix C.6 for details.

We now move on to our results from [Section 3](#) of the main paper. The major technical difference between the adjusted and unadjusted cases is perhaps reflected in the proof of [Theorem 3.2](#), which requires substantially different techniques compared to its unadjusted counterpart in [Theorem 2.1](#). To understand this difference, recall from (3.5) of the main paper that under $\tau = \tau_0 - hN^{-1/2}$, the regression adjusted Wilcoxon rank-sum statistic is given by:

$$t_{N,\text{adj}} \equiv t_{N,\text{adj}}^h = \sum_{i,j=1}^N Z_{N,j} \mathbf{1} \left(hN^{-1/2} (\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} - hN^{-1/2} \right),$$

where $\mathbf{p}_{N,j}$ is the j -th column of the projection matrix $\mathbf{P}_{\mathbf{X}_N}$ that projects onto the column space of \mathbf{X}_N , and $\tilde{b}_{N,i} := Y_{N,i} - \tau_0 Z_{N,i} - \mathbf{p}_{N,i}^\top \mathbf{b}_N$. Therefore, for every pair (i, j) , the indicators in $t_{N,\text{adj}}$ depend on the entire random vector $(Z_{N,1}, \dots, Z_{N,N})$. This is in sharp contrast to t_N^h above, where for every pair (i, j) , the indicators depend only on $(Z_{N,i}, Z_{N,j})$. As a result, even studying the mean and variance of $t_{N,\text{adj}}^h$ raises significantly more technical challenges than those of t_N^h . We circumvent this by leveraging the properties of the projection matrix and showing that the term on the left hand side of the indicators in the above display can be replaced by 0, in a suitable asymptotic sense. To be specific, define

$$\tilde{A}_{N,\text{adj}}^h := \sum_{i,j=1}^N Z_{N,j} \mathbf{1} \left(0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} - hN^{-1/2} \right).$$

Our main contribution lies in showing that

$$N^{-3/2} |t_{N,\text{adj}}^h - \tilde{A}_{N,\text{adj}}^h| = o_p(1).$$

The proof of this step is technical and we defer the reader to [Appendix D](#) for details of these computations. However once the above display is established, it suffices to obtain the asymptotic distribution of $N^{-3/2} \tilde{A}_{N,\text{adj}}^h$. Note that the indicators in $\tilde{A}_{N,\text{adj}}^h$ are deterministic and do not depend on \mathbf{Z}_N . This puts us in a similar situation to the analysis of t_N^0 , which we have already discussed above.

Once [Theorem 3.2](#) is established, the proofs of [Theorem 3.3](#) and [Corollary A.2](#) follow using ideas similar to the proofs of [Theorem 2.2](#) and [Corollary A.1](#) respectively.

C Proofs of the main results

C.1 Proof of [Proposition 2.1](#)

Proof. We begin by recalling that $t_N := \hat{\mathbf{q}}_N^\top \mathbf{Z}_N$, where

$$\hat{q}_{N,j} := \sum_{i=1}^N \mathbf{1} \{ Y_{N,i} - \tau_0 Z_{N,i} \leq Y_{N,j} - \tau_0 Z_{N,j} \}, \quad 1 \leq j \leq N.$$

It follows under $\tau = \tau_0$ that

$$t_N \stackrel{d}{=} \mathbf{q}_N^\top \mathbf{Z}_N, \quad q_{N,j} = \sum_{i=1}^N \mathbf{1}\{b_{N,i} \leq b_{N,j}\}. \quad (\text{C.1})$$

Further, under $\tau = \tau_0$,

$$\mathbf{q}_N^\top \mathbf{Z}_N \stackrel{d}{=} \sum_{i=1}^m q_{N,\Pi_N(i)} = \sum_{i=1}^N c_{N,i} \cdot q_{N,\Pi_N(i)}$$

where $c_{N,i} = \mathbf{1}\{i \leq m\}$ and Π_N is a random permutation of $\{1, 2, \dots, N\}$. Now, $\bar{c}_N := N^{-1} \sum_{i=1}^N c_{N,i} = m/N$, and $\max_{1 \leq i \leq N} (q_{N,i} - \bar{q}_N)^2 = O(N^2)$. Hence

$$\begin{aligned} & \lim_{N \rightarrow \infty} N \cdot \frac{\max_{1 \leq i \leq N} (c_{N,i} - \bar{c}_N)^2}{\sum_{i=1}^N (c_{N,i} - \bar{c}_N)^2} \cdot \frac{\max_{1 \leq i \leq N} (q_{N,i} - \bar{q}_N)^2}{\sum_{i=1}^N (q_{N,i} - \bar{q}_N)^2} \\ &= \lim_{N \rightarrow \infty} \frac{\max\{\bar{c}_N^2, (1 - \bar{c}_N)^2\}}{\frac{m}{N}(1 - \bar{c}_N)^2 + (1 - \frac{m}{N})\bar{c}_N^2} \cdot \frac{O(N^2)}{\frac{N(N^2-1)}{12} + o(N^3)} \\ &= \frac{\max\{\lambda^2, (1 - \lambda)^2\}}{\lambda(1 - \lambda)} \cdot \lim_{N \rightarrow \infty} \frac{O(N^2)}{\frac{N(N^2-1)}{12} + o(N^3)} \\ &= 0. \end{aligned}$$

In view of the above, Hoeffding's combinatorial CLT [Hoeffding \(1951, Theorem 4\)](#) implies that under $\tau = \tau_0$,

$$\frac{t_N - \mathbb{E}(t_N)}{\sqrt{\text{Var}(t_N)}} \xrightarrow{d} \mathcal{N}(0, 1) \text{ as } N \rightarrow \infty.$$

Now invoking [Lemma D.6](#), we have $\text{Var}_{\tau_0}(t_N) \sim \frac{\lambda(1-\lambda)}{12} N^3$, which completes the proof. \square

C.2 Proof of [Proposition 2.2](#)

Proof. It follows from the algebraic manipulations provided in [Hodges and Lehmann \(1963, Section 4\)](#) that our modified estimator $\hat{\tau}^R$ in [\(2.11\)](#), with $t(\cdot, \cdot)$ as in [\(2.4\)](#) of the main paper, is given by

$$\hat{\tau}^R = \text{median}\{Y_i - Y_j : Z_i = 1, Z_j = 0, 1 \leq i, j \leq N\}.$$

Now for any $1 \leq k \leq N$, fix any k indices $1 \leq i_1 < i_2 < \dots < i_k \leq N$, and any treatment assignment $(Z_1, \dots, Z_N) = (z_1, \dots, z_N)$. Set $n_1 := |\{1 \leq j \leq k : z_{i_j} = 1\}|$. Note that changing the responses y_{i_1}, \dots, y_{i_k} arbitrarily can make the pairwise differences $\{y_i - y_j : z_i = 1, z_j = 0\}$ blow up to $+\infty$ except for $(m - n_1)(N - m - (k - n_1))$ many pairs (i, j) . As long as this count is greater than or equal to half of the total number of pairwise differences, i.e., $m(N - m)/2$, the median cannot be made arbitrarily large. Note, this needs to be satisfied for every treatment assignment $z_1, \dots, z_N \in \{0, 1\}$ such that $\sum_{i=1}^N z_i = m$. By symmetry, we can assume without loss of generality that $(i_1, i_2, \dots, i_k) = (1, 2, \dots, k)$. Thus, we can

make $\hat{\tau}^R(y_1, \dots, y_N; z_1, \dots, z_N)$ arbitrarily large by altering k responses only if k satisfies the following:

$$\forall z_1, \dots, z_N \in \{0, 1\}, (m - n_1)(N - m - (k - n_1)) < \frac{m(N - m)}{2}, \text{ where } n_1 = \sum_{j=1}^k z_j. \quad (\text{C.2})$$

Moreover, if k satisfies this condition, then for any $z_1, \dots, z_N \in \{0, 1\}$ such that $\sum_{i=1}^N z_i = m$, and any choice of indices $1 \leq i_1 < i_2 < \dots < i_k \leq N$, we can let $y_{i_j} \rightarrow \infty$ if $z_{i_j} = 1$ and $y_{i_j} \rightarrow -\infty$ if $z_{i_j} = 0$, which results in $\hat{\tau}^R(y_1, \dots, y_N; z_1, \dots, z_N) \rightarrow \infty$. Therefore,

$$\text{BP}_N(\hat{\tau}^R) = \frac{1}{N} \min \{1 \leq k \leq N : (\text{C.2}) \text{ holds}\}.$$

It is elementary to observe that $f(x) = (m - x)(N - m - k + x)$ has a global maximum at $x = m - (N - k)/2$, and the maximum value is $(N - k)^2/4$. However, we want to maximize $f(x)$ over $[0, k]$ only. Consider the following cases:

- **Case 1:** $m > N/3$. Note, $0 \leq m - (N - k)/2 \leq k \iff k \geq 2m - N$. Thus, for $2m - N \leq k \leq N$,

$$(\text{C.2}) \text{ holds} \iff \frac{(N - k)^2}{4} < \frac{m(N - m)}{2} \iff k > N \left(1 - \sqrt{2\frac{m}{N} \left(1 - \frac{m}{N}\right)}\right) \geq 2m - N.$$

In other words, for $2m - N \leq k \leq N$, the condition (C.2) is automatic. On the other hand, for $1 \leq k < 2m - N$, the maximum of the function $f(x) = (m - x)(N - m - k + x)$ is attained at $x = k$. So, for $1 \leq k < 2m - N$,

$$(\text{C.2}) \text{ holds} \iff (m - k)(N - m) < \frac{m(N - m)}{2} \iff k > \frac{m}{2}.$$

Hence we can write

$$\begin{aligned} & \{1 \leq k \leq N : (\text{C.2}) \text{ holds}\} \\ &= \{1 \leq k < 2m - N : (\text{C.2}) \text{ holds}\} \cup \{2m - N \leq k < N : (\text{C.2}) \text{ holds}\} \\ &= \{1 \leq k < 2m - N : k > m/2\} \cup \{2m - N \leq k < N\}. \end{aligned}$$

Since in this case $m/2 < 2m - N$, it follows that for $m > N/2$,

$$\text{BP}_N(\hat{\tau}^R) = \frac{1}{N} \left\lceil \frac{m}{2} \right\rceil.$$

- **Case 2:** $m < N/3$. This case is similar to the previous case. In this case we get

$$\text{BP}_N(\hat{\tau}^R) = \frac{1}{N} \left\lceil \frac{N - m}{2} \right\rceil.$$

- **Case 3:** $N/3 \leq m \leq 2N/3$. First, consider $N/2 \leq m \leq 2N/3$. We note that

$$\begin{aligned} & \{1 \leq k \leq N : (\text{C.2}) \text{ holds}\} \\ &= \{1 \leq k < 2m - N : (\text{C.2}) \text{ holds}\} \cup \{2m - N \leq k < N : (\text{C.2}) \text{ holds}\} \\ &= \left\{1 \leq k < 2m - N : \frac{k}{N} > \frac{m}{2N}\right\} \cup \left\{2m - N \leq k < N : \frac{k}{N} > 1 - \sqrt{2\frac{m}{N}\left(1 - \frac{m}{N}\right)}\right\}. \end{aligned}$$

Now a little algebra shows that $m/2 < N - \sqrt{2m(N-m)}$. Hence in this case we get

$$\text{BP}_N(\hat{\tau}^R) = \frac{1}{N} \left\lfloor N \left(1 - \sqrt{2\frac{m}{N}\left(1 - \frac{m}{N}\right)}\right) \right\rfloor.$$

Similarly one can deal with the other sub-case, namely, $N/3 \leq m \leq N/2$.

Since $m/N \rightarrow \lambda \in [0, 1]$, the desired conclusion follows. \square

C.3 Proof of Theorem 2.1

Proof. We start with the decomposition stated in Proposition B.1. Under $\tau = \tau_N$, it holds that

$$t_N \stackrel{d}{=} \underbrace{m + \sum_{j=1}^N Z_{N,j} \sum_{i=1, i \neq j}^N \mathbf{1}\{b_{N,i} \leq b_{N,j}\}}_{\text{call this } W_N} - \underbrace{\sum_{j=1}^N \sum_{i=1, i \neq j}^N (1 - Z_{N,i}) Z_{N,j} I_{h,N}(b_{N,j} - b_{N,i})}_{\text{call this } S_N},$$

where $I_{h,N}(\cdot)$ is as in (2.15) of the main paper. We now observe that the randomization distribution of W_N is identical to the null distribution of t_N . Thus, invoking Proposition 2.1 and Lemma D.5, we deduce that

$$N^{-3/2} \left(W_N - \frac{m(N+1)}{2} \right) \xrightarrow{d} \mathcal{N} \left(0, \frac{\lambda(1-\lambda)}{12} \right).$$

On the other hand, Lemma D.7 tells us that $N^{-3/2} S_N \xrightarrow{P} h\lambda(1-\lambda)\mathcal{I}_b$. Combining these using Slutsky's theorem we conclude that under $\tau = \tau_N$,

$$\begin{aligned} N^{-3/2} \left(t_N - \frac{m(N+1)}{2} \right) &\stackrel{d}{=} N^{-3/2} \left(W_N - \frac{m(N+1)}{2} \right) - N^{-3/2} S_N \\ &\xrightarrow{d} \mathcal{N} \left(-h\lambda(1-\lambda)\mathcal{I}_b, \frac{\lambda(1-\lambda)}{12} \right), \end{aligned} \tag{C.3}$$

which completes the proof. \square

C.4 Proof of Theorem 2.2

Proof. Recall from Theorem 2.1 that under $\tau = \tau_0 + hN^{-1/2}$ we have

$$N^{-3/2} \left(t_N - \frac{m(N+1)}{2} \right) \xrightarrow{d} \mathcal{N} \left(-h\lambda(1-\lambda)\mathcal{I}_b, \frac{\lambda(1-\lambda)}{12} \right),$$

where \mathcal{I}_b is defined in Assumption 1. Now we can invoke Lemma D.1 to complete the proof. \square

C.5 Proof of Theorem 2.3

Proof. First note that under any value of τ ,

$$B_N \stackrel{d}{=} \sum_{j=1}^N \sum_{i=1, i \neq j}^N (1 - Z_{N,i})(1 - Z_{N,j}) I_{1,N}(b_{N,j} - b_{N,i}).$$

Invoking Assumption 1, we get

$$\mathbb{E} N^{-3/2} B_N = \frac{(N-m)(N-m-1)}{N(N-1)} N^{-3/2} \sum_{j=1}^N \sum_{i=1, i \neq j}^N I_{1,N}(b_{N,j} - b_{N,i}) \rightarrow (1-\lambda)^2 \mathcal{I}_b.$$

Consequently, $\mathbb{E}((1-m/N)^{-2} N^{-3/2} B_N) \xrightarrow{P} \mathcal{I}_b$. Thus it suffices to show that $\text{Var}(B_N) = o(N^3)$. For brevity, we denote by $I_N(i, j)$ the indicator $I_{1,N}(b_{N,j} - b_{N,i})$ in the rest of the proof. Note that

$$\text{Var}(B_N) = \sum_{i=1}^N \sum_{j=1, j \neq i}^N \sum_{k=1}^N \sum_{l=1, l \neq k}^N \text{Cov}((1 - Z_{N,i})(1 - Z_{N,j}), (1 - Z_{N,k})(1 - Z_{N,l})) I_N(i, j) I_N(k, l).$$

Since $2 \leq |\{i, j, k, l\}| \leq 4$, we consider the following cases.

- (a) $|\{i, j, k, l\}| = 2$, i.e., $(i, j) = (k, l)$ or (l, k) . Since $\text{Var}((1 - Z_{N,i})(1 - Z_{N,j})) = p_N(1 - p_N)$ where $p_N = P(Z_{N,i} = 0, Z_{N,j} = 0) = \frac{(N-m)(N-m-1)}{N(N-1)} \sim (1-\lambda)^2$, the contribution of these terms in $\text{Var}(B_N)$ is given by

$$\sum_{i=1}^N \sum_{j=1, j \neq i}^N 2p_N(1 - p_N) I_N^2(i, j) \lesssim 2(1-\lambda)^2 \sum_{i=1}^N \sum_{j=1, j \neq i}^N I_N(i, j) \lesssim N^{3/2}.$$

- (b) $|\{i, j, k, l\}| = 4$, i.e., all 4 indices are distinct. Note that in this case

$$\begin{aligned} & \text{Cov}((1 - Z_{N,i})(1 - Z_{N,j}), (1 - Z_{N,k})(1 - Z_{N,l})) \\ &= P(Z_{N,i} = 0, Z_{N,j} = 0, Z_{N,k} = 0, Z_{N,l} = 0) - p_N^2 \end{aligned}$$

$$\begin{aligned}
&= p_N \left(\frac{(N-m-2)(N-m-3)}{(N-2)(N-3)} - \frac{(N-m)(N-m-1)}{N(N-1)} \right) \\
&= -p_N \frac{2m(2N^2 - 2mN - 6N + 3 + 3m)}{N(N-1)(N-2)(N-3)} \sim -4\lambda(1-\lambda)^3 N^{-1}.
\end{aligned}$$

Hence the contribution u_N of these terms in $\text{Var}(B_N)$ satisfies the following.

$$u_N \lesssim N^{-1} \sum_{i=1}^N \sum_{j=1, j \neq i}^N \sum_{k=1}^N \sum_{l=1, l \neq k}^N I_N(i, j) I_N(k, l) \lesssim N^{-1+3/2+3/2} = N^2.$$

(c) $|\{i, j, k, l\}| = 3$. Here we have 4 sub-cases: $i = k, j \neq l$; $i \neq k, j = l$; $i = l, j \neq k$; $i \neq l, j = k$. In each of the subcases,

$$\begin{aligned}
\text{Cov}((1 - Z_{N,i})(1 - Z_{N,j}), (1 - Z_{N,k})(1 - Z_{N,l})) &= P(Z_{N,1} = Z_{N,2} = Z_{N,3} = 0) - p_N^2 \\
&\sim (1 - \lambda)^3 - (1 - \lambda)^4 = \lambda(1 - \lambda)^3.
\end{aligned}$$

Hence, if v_n be the contribution of these terms in $\text{Var}(B_N)$, then

$$\begin{aligned}
v_N &\lesssim \sum_{i,j,l \text{ distinct}} I_N(i, j) I_N(i, l) + \sum_{i,j,k \text{ distinct}} I_N(i, j) I_N(k, j) \\
&\quad + \sum_{i,j,k \text{ distinct}} I_N(i, j) I_N(k, i) + \sum_{i,j,l \text{ distinct}} I_N(i, j) I_N(j, l) \\
&\leq 4N \sum_{i,j \neq i} I_N(i, j) \lesssim N^{1+3/2} = N^{5/2}.
\end{aligned}$$

Combining the above cases, we conclude that $N^{-3}\text{Var}(B_N) \lesssim N^{-3}(N^{3/2} + N^2 + N^{5/2}) = o(1)$, which completes the proof. \square

C.6 Proof of Theorem 2.4

Proof. Let $B_{N,1}, B_{N,2}$ be two independent samples from F_N . Then the quantity on the LHS of Assumption 1 reads

$$\sqrt{N} \mathbb{P} \left(0 \leq B_{N,2} - B_{N,1} < h/\sqrt{N} \right) = \sqrt{N} \mathbb{E} \left(F_N(B_{N,1} + h/\sqrt{N}) - F_N(B_{N,1}) \right). \quad (\text{C.4})$$

Since

$$\sup_{x \in \mathbb{R}} \left| \sqrt{N} \left(F_N \left(x + \frac{h}{\sqrt{N}} \right) - F_N(x) \right) - hf(x) \right| \rightarrow 0,$$

the quantity in (C.4) is equal to $h\mathbb{E}f(B_{N,1}) + o(1)$ which tends to $h \int_{\mathbb{R}} f^2(x)dx$ as $N \rightarrow \infty$ by assumption. Thus, [Assumption 1](#) holds with $\mathcal{I}_b = \int_{\mathbb{R}} f^2(x)dx$, and hence [Theorem 2.2](#) gives

$$\sqrt{N}(\hat{\tau}^R - \tau_0) \xrightarrow{d} \mathcal{N}\left(0, (12\lambda(1-\lambda))^{-1} \left(\int_{\mathbb{R}} f^2(x)dx\right)^{-2}\right).$$

(Note, since $F_N \xrightarrow{d} F$, the assumption $\mathbb{E}f(B_{N,1}) \rightarrow \int_{\mathbb{R}} f^2(x)dx < \infty$ holds under mild conditions on f ; e.g., when $\{f(B_N)\}_{N \geq 1}$ is uniformly integrable where $B_N \sim \mathbb{P}_N$. For bounded f , this is trivial.)

Next, we consider the difference-in-means estimator $\hat{\tau}_{\text{dm}}$. When $N^{-1} \sum_{j=1}^N (b_{N,j} - \bar{b}_N)^2 \rightarrow \sigma^2$ and $N^{-1} \max_{1 \leq j \leq N} (b_{N,j} - \bar{b}_N)^2 \rightarrow 0$, it follows from [Li and Ding \(2017, Theorem 5\)](#) that

$$\sqrt{N}(\hat{\tau}_{\text{dm}} - \tau) \xrightarrow{d} \mathcal{N}(0, \sigma^2/\lambda(1-\lambda)).$$

Therefore, the asymptotic efficiency of $\hat{\tau}^R$ relative to $\hat{\tau}_{\text{dm}}$ (see [Section 2.4](#) of the main paper for definition) is given by

$$\text{eff}(\hat{\tau}^R, \hat{\tau}_{\text{dm}}) = 12\sigma^2 \left(\int_{\mathbb{R}} f_b^2(x)dx\right)^2.$$

The desired lower bound then follows from [Hodges and Lehmann \(1956, Theorem 1\)](#). (Note that since $F_N \xrightarrow{d} F$, σ^2 is indeed the variance of the distribution with density f under mild regularity conditions; e.g., when $\{B_N^2\}_{N \geq 1}$ is uniformly integrable where $B_N \sim \mathbb{P}_N$.) \square

Remark C.1. Let \mathbb{P}_N and \mathbb{P} respectively denote the probability measures with distribution functions F_N and F . Also, let $\mathbb{P}g := \int g d\mathbb{P}$, and define $\mathbb{P}_N g$ similarly. Note that

$$\sup_{x \in \mathbb{R}} \left| \sqrt{N} \left(F_N \left(x + \frac{h}{\sqrt{N}} \right) - F_N(x) \right) - hf(x) \right| = \sup_{g \in \mathcal{F}_N} \left| \sqrt{N}(\mathbb{P}_N - \mathbb{P})g \right|,$$

where \mathcal{F}_N is the class of indicator functions of intervals of the form $[x, x + h/\sqrt{N})$. It follows from [Chernozhukov et al. \(2014, Corollary 5.1\)](#) that the above supremum goes to 0 as $N \rightarrow \infty$, for any fixed h , when $\{b_{N,j}\}_{1 \leq j \leq N}$ are realizations from a distribution with density f .

C.7 Proof of [Theorem 3.1](#)

Proof. Denote by $\mathbf{p}_{N,i}$ the i -th row of $\mathbf{P}_{\mathbf{X}_N}$. Observe that

$$e_{N,i} \leq e_{N,j} \iff Y_{N,i} - \tau_0 Z_{N,i} - \mathbf{p}_{N,i}^\top (\mathbf{Y}_N - \tau_0 \mathbf{Z}_N) \leq Y_{N,j} - \tau_0 Z_{N,j} - \mathbf{p}_{N,j}^\top (\mathbf{Y}_N - \tau_0 \mathbf{Z}_N).$$

So under $\tau = \tau_0$, we have $t_{N,\text{adj}} \stackrel{d}{=} \sum_{j=1}^N q_{N,j} Z_{N,j}$, where

$$q_{N,j} := \sum_{i=1}^N \mathbf{1}\{b_{N,i} - \mathbf{p}_{N,i}^\top \mathbf{b}_N \leq b_{N,j} - \mathbf{p}_{N,j}^\top \mathbf{b}_N\}, \quad j = 1, 2, \dots, N.$$

Since the ranks $q_{N,j}$'s are deterministic, the asymptotic normality of $\sum_{j=1}^N q_{N,j} Z_{N,j}$ can be derived in the same way as for the without regression adjustment case. A closer look at the proofs of [Lemma D.5](#) and [Proposition 2.1](#) in the without regression adjustment case reveals that the following results hold in this case as well, since [Assumption 2](#) plays the role of [Assumption 1](#).

- (a) As $N \rightarrow \infty$, $\sum_{j=1}^N (q_{N,j} - \bar{q}_N)^2 = \frac{1}{12}N(N^2 - 1) + o(N^3)$.
- (b) $\lim_{N \rightarrow \infty} \max_{1 \leq j \leq N} (q_{N,j} - \bar{q}_N)^2 / \sum_{j=1}^N (q_{N,j} - \bar{q}_N)^2 = 0$.
- (c) Under $\tau = \tau_0$, $\text{Var}(t_{N,\text{adj}}) \sim \frac{1}{12}\lambda(1 - \lambda)N^3$ as $N \rightarrow \infty$.

Equipped with (b) above, we apply Hoeffding's combinatorial CLT ([Hoeffding, 1951](#), Theorem 4) to say that under $\tau = \tau_0$,

$$\frac{t_{N,\text{adj}} - \mathbb{E}(t_{N,\text{adj}})}{\sqrt{\text{Var}(t_{N,\text{adj}})}} \xrightarrow{d} \mathcal{N}(0, 1) \text{ as } N \rightarrow \infty.$$

This, in conjunction with (c) above, completes the proof. □

C.8 Proof of [Theorem 3.2](#)

Proof. Recall the notation $\tilde{b}_{N,i} = b_{N,i} - \mathbf{p}_{N,i}^\top \mathbf{b}_N$ ($1 \leq i \leq N$) from [\(3.6\)](#) of the main paper. Under $\tau = \tau_N$, we have

$$\mathbf{Y}_N - \tau_0 \mathbf{Z}_N \stackrel{d}{=} \mathbf{b}_N + (\tau_N - \tau_0) \mathbf{Z}_N = \mathbf{b}_N - \frac{h}{\sqrt{N}} \mathbf{Z}_N,$$

which implies that

$$\begin{aligned} t_{N,\text{adj}} &\stackrel{d}{=} \sum_{j=1}^N Z_{N,j} \sum_{i=1}^N \mathbf{1}\left\{\tilde{b}_{N,i} - \frac{h}{\sqrt{N}} Z_{N,i} + \frac{h}{\sqrt{N}} \mathbf{p}_{N,i}^\top \mathbf{Z}_N \leq \tilde{b}_{N,j} - \frac{h}{\sqrt{N}} Z_{N,j} + \frac{h}{\sqrt{N}} \mathbf{p}_{N,j}^\top \mathbf{Z}_N\right\} \\ &= \sum_{j=1}^N Z_{N,j} \sum_{i=1}^N (1 - Z_{N,i}) \mathbf{1}\left\{\tilde{b}_{N,i} + \frac{h}{\sqrt{N}} \mathbf{p}_{N,i}^\top \mathbf{Z}_N \leq \tilde{b}_{N,j} + \frac{h}{\sqrt{N}} \mathbf{p}_{N,j}^\top \mathbf{Z}_N - \frac{h}{\sqrt{N}}\right\} + C_N, \end{aligned}$$

where

$$C_N := \sum_{j=1}^N \sum_{i=1}^N Z_{N,j} Z_{N,i} \mathbf{1}\left\{\tilde{b}_{N,i} + \frac{h}{\sqrt{N}} \mathbf{p}_{N,i}^\top \mathbf{Z}_N \leq \tilde{b}_{N,j} + \frac{h}{\sqrt{N}} \mathbf{p}_{N,j}^\top \mathbf{Z}_N\right\}. \quad (\text{C.5})$$

Thus under $\tau = \tau_N$ we have $t_{N,\text{adj}} \stackrel{d}{=} \mathbf{I}_N - \mathbf{II}_N + C_N$, where

$$\mathbf{I}_N := \sum_{j=1}^N \sum_{i=1}^N Z_{N,j} \xi_{N,i,j}, \quad \text{and} \quad \mathbf{II}_N := \sum_{j=1}^N \sum_{i=1}^N Z_{N,j} Z_{N,i} \xi_{N,i,j}, \quad (\text{C.6})$$

and

$$\begin{aligned} \xi_{N,i,j} &:= \mathbf{1}\{b_{N,i} - \mathbf{p}_{N,i}^\top (\mathbf{b}_N - hN^{-1/2} \mathbf{Z}_N) \leq b_{N,j} - \mathbf{p}_{N,j}^\top (\mathbf{b}_N - hN^{-1/2} \mathbf{Z}_N) - hN^{-1/2}\} \\ &= \mathbf{1}\{hN^{-1/2} (\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} - hN^{-1/2}\}. \end{aligned}$$

The indicators $\xi_{N,i,j}$ are quite complicated to handle, since it depends on the entire random vector $(Z_{N,1}, \dots, Z_{N,N})$, for every pair (i, j) . To circumvent this technical hurdle, we replace $\xi_{N,i,j}$ with $\tilde{\xi}_{N,i,j}$, where

$$\tilde{\xi}_{N,i,j} := \mathbf{1}\{0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} - hN^{-1/2}\}.$$

Define

$$\tilde{\mathbf{I}}_N := \sum_{j=1}^N Z_{N,j} \sum_{i=1}^N \tilde{\xi}_{N,i,j}, \quad \text{and} \quad \tilde{\mathbf{II}}_N := \sum_{j=1}^N \sum_{i=1}^N Z_{N,j} Z_{N,i} \tilde{\xi}_{N,i,j}. \quad (\text{C.7})$$

Thus we decompose $t_{N,\text{adj}}$ under $\tau = \tau_N$ as

$$t_{N,\text{adj}} \stackrel{d}{=} \underbrace{(\tilde{\mathbf{I}}_N - \tilde{\mathbf{II}}_N + C_N)}_{\tilde{A}_{N,\text{adj}}^h} + \underbrace{(\mathbf{I}_N - \tilde{\mathbf{I}}_N)}_{D_n} - \underbrace{(\mathbf{II}_N - \tilde{\mathbf{II}}_N)}_{Q_n}. \quad (\text{C.8})$$

We first focus on $\tilde{A}_{N,\text{adj}}^h$. Note that in (C.5) we are summing up the ranks of the m numbers

$$\{\tilde{b}_{N,j} + hN^{-1/2} \mathbf{p}_{N,j}^\top \mathbf{Z}_N : 1 \leq j \leq N, Z_{N,j} = 1\}$$

within this set. Unfortunately, due to possibility of ties, we cannot directly equate it with $m(m+1)/2$. However, invoking [Assumption 2](#) we can show that $N^{-3/2}(C_N - m(m+1)/2) = o_P(1)$ as $N \rightarrow \infty$. Towards that, observe that for any $\delta > 0$,

$$\begin{aligned} \left| C_N - \frac{m(m+1)}{2} \right| &= \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{\tilde{b}_{N,j} - \tilde{b}_{N,i} + hN^{-1/2} (\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N = 0\} \\ &\leq \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} + hN^{-1/2} (\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N < \delta N^{-1/2}\} \\ &\leq \sum_{(i,j): |(\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N| < \delta} \mathbf{1}\{0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} + hN^{-1/2} (\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N < \delta N^{-1/2}\} \\ &\quad + \left| \{(i, j) : |(\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N| \geq \delta\} \right| \end{aligned}$$

$$\begin{aligned}
&\leq \sum_{(i,j): |(\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N| < \delta} \mathbf{1}\left\{-h\delta N^{-1/2} \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < (h+1)\delta N^{-1/2}\right\} \\
&\quad + \left|\left\{(i,j) : |(\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N| \geq \delta\right\}\right| \\
&\leq \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{\left|\tilde{b}_{N,j} - \tilde{b}_{N,i}\right| \leq (h+1)\delta N^{-1/2}\right\} + \delta^{-2} \sum_{j=1}^N \sum_{i=1}^N ((\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N)^2. \tag{C.9}
\end{aligned}$$

Now we invoke [Lemma E.1](#) to say that

$$N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbb{E} \left[((\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N)^2 \right] = O(N^{-1/2}) = o(1),$$

for every fixed $\delta > 0$. This, in conjunction with (C.9) and Markov inequality, tells us that for any fixed $\delta > 0$,

$$N^{-3/2} \left| \left\{ (i,j) : |(\mathbf{p}_{N,j} - \mathbf{p}_{N,i})^\top \mathbf{Z}_N| \geq \delta \right\} \right| = o_P(1).$$

On the other hand, we can use [Lemma D.9](#) to conclude that under [Assumption 2](#),

$$\lim_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{\left|\tilde{b}_{N,j} - \tilde{b}_{N,i}\right| \leq (h+1)\delta N^{-1/2}\right\} = 2(h+1)\delta.$$

Appealing to (C.9) we can now conclude, by letting $N \rightarrow \infty$ first, and then $\delta \rightarrow 0$, that

$$N^{-3/2}(C_N - m(m+1)/2) = o_P(1), \text{ as } N \rightarrow \infty.$$

This allows us to write

$$\tilde{A}_{N,\text{adj}}^h = \sum_{j=1}^N Z_{N,j} \sum_{i=1}^N \mathbf{1}\left\{0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} - hN^{-1/2}\right\} + \frac{m(m+1)}{2} + o_P(1).$$

In view of the above display, the asymptotic normality of $\tilde{A}_{N,\text{adj}}^h$ under $\tau = \tau_N$ can be derived in the same manner as we proved the local asymptotic normality of t_N in the without regression adjustment case. To be precise, it follows by mimicking the proof of [Theorem 2.1](#) (in the same manner as we proved [Theorem 3.1](#) by mimicking the proof of [Proposition 2.1](#)) that under $\tau = \tau_N$,

$$N^{-3/2}(\tilde{A}_{N,\text{adj}}^h - \frac{m(N+1)}{2}) \xrightarrow{d} N\left(-h\lambda(1-\lambda)\mathcal{J}_b, \frac{\lambda(1-\lambda)}{12}\right), \tag{C.10}$$

where \mathcal{J}_b is as defined in [Assumption 2](#). The proof of the fact that D_N and Q_N defined in (C.8) are asymptotically negligible, is split into a couple of lemmas in [Appendix D](#). [Lemmas D.11](#) and [D.12](#) give upper bounds on the second moments of D_N and Q_N , respectively. Then [Lemma D.13](#) shows that under [Assumption 2](#), $N^{-3/2}D_N = o_P(1)$ and $N^{-3/2}Q_N = o_P(1)$ as $N \rightarrow \infty$. This, in conjunction with (C.10) completes the proof of [Theorem 3.2](#). \square

C.9 Proof of Theorem 3.3

Proof. Recall from Theorem 3.2 that under $\tau = \tau_0 + hN^{-1/2}$, we have

$$N^{-3/2} \left(t_{N,\text{adj}} - \frac{m(N+1)}{2} \right) \xrightarrow{d} \mathcal{N} \left(-h\lambda(1-\lambda)\mathcal{J}_b, \frac{\lambda(1-\lambda)}{12} \right),$$

where \mathcal{J}_b is defined in Assumption 2. Now we invoke Lemma D.1 to complete the proof. \square

C.10 Proof of Theorem 3.4

Proof. Under the assumptions of this theorem, Lemma D.8 tells us that Assumption 2 holds in probability, with $\mathcal{J}_b = (2\sqrt{\pi}\sigma)^{-1}$. Next, fix any $h \in \mathbb{R}$ and define

$$\mathcal{I}_{h,N} := N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N I_{h,N}(b_j - b_i),$$

where $I_{h,N}$ is defined in (2.15) of the main paper. Without loss of generality, we take $h > 0$ (the other case will be similar). For simplicity in notation, we shall omit the index N in this proof for $\mathbf{b}_N, \mathbf{X}_N$ etc. From $\mathbf{b} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$ we write $b_j - b_i = \varepsilon_j - \varepsilon_i + v_j - v_i$ where v_i denotes the i -th coordinate of $\mathbf{v} = \mathbf{X}\boldsymbol{\beta}$. It then follows that

$$\begin{aligned} \mathbb{E}\mathcal{I}_{h,N} &= N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N P(0 \leq \varepsilon_j - \varepsilon_i + v_j - v_i < hN^{-1/2}) \\ &= N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left[\Phi \left(\frac{v_i - v_j + hN^{-1/2}}{\sqrt{2}\sigma} \right) - \Phi \left(\frac{v_i - v_j}{\sqrt{2}\sigma} \right) \right] \\ &= N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left[\frac{h}{\sqrt{2}\sigma\sqrt{N}} \phi \left(\frac{v_i - v_j}{\sqrt{2}\sigma} \right) + \frac{h^2}{4N\sigma^2} \phi'(\xi_{i,j}) \right] \\ &= \frac{h}{2\sqrt{\pi}\sigma} N^{-2} \sum_{j=1}^N \sum_{i=1}^N e^{-(v_j - v_i)^2/4\sigma^2} + (4\sigma^2)^{-1} h^2 N^{-5/2} \sum_{j=1}^N \sum_{i=1}^N \phi'(\xi_{i,j}), \end{aligned}$$

where for each (i, j) , $\xi_{i,j}$ is a number between $v_i - v_j$ and $v_i - v_j + N^{-1/2}$. Since ϕ' is bounded, we can show (by proceeding in the same manner as in the proof of Lemma D.8) that the second sum in the above display is asymptotically negligible. Hence

$$\lim_{N \rightarrow \infty} \left(\mathbb{E}\mathcal{I}_{h,N} - \frac{h}{2\sqrt{\pi}\sigma} N^{-2} \sum_{j=1}^N \sum_{i=1}^N e^{-(v_j - v_i)^2/4\sigma^2} \right) = 0. \quad (\text{C.11})$$

Since $0 \leq N^{-2} \sum_{j=1}^N \sum_{i=1}^N e^{-(v_j - v_i)^2 / 4\sigma^2} \leq 1$, it follows that for any $h > 0$,

$$\limsup_{N \rightarrow \infty} \mathbb{E} \mathcal{I}_{h,N} / h \leq (2\sqrt{\pi}\sigma)^{-1} = \mathcal{J}_b.$$

Next we show that $\text{Var}(\mathcal{I}_{h,N}) \rightarrow 0$. Towards that, we first write

$$\mathbb{E} \mathcal{I}_{h,N}^2 = N^{-3} \sum_{i,j,k,l} P(0 \leq b_j - b_i < hN^{-1/2}, 0 \leq b_l - b_k < hN^{-1/2}).$$

Now, as mentioned earlier, [Assumption 2](#) holds in this setting (cf. [Lemma D.8](#)); hence appealing to arguments similar to those given in [Appendix C.5](#), we argue that the contribution from the terms with repeated indices are negligible. So we are left with terms with distinct indices i, j, k, l . For such indices, note that we have $P(0 \leq b_j - b_i < hN^{-1/2}, 0 \leq b_l - b_k < hN^{-1/2}) = P(0 \leq b_j - b_i < hN^{-1/2})P(0 \leq b_l - b_k < hN^{-1/2})$, since the errors ε_i 's are independent and v_i 's non-stochastic. We can thus write

$$\mathbb{E} \mathcal{I}_{h,N}^2 = \left(N^{-3/2} \sum_{i,j \text{ distinct}} P(0 \leq b_j - b_i < hN^{-1/2}) \right)^2 + o(1) = (\mathbb{E} \mathcal{I}_{h,N})^2 + o(1),$$

from which the conclusion follows. Moreover, [\(C.11\)](#) tells us that [Assumption 1](#) holds, with $\lim_{N \rightarrow \infty} \mathbb{E} \mathcal{I}_{h,N} / h = \ell(2\sqrt{\pi}\sigma)^{-1} = \ell \mathcal{J}_b$. Finally, for any $x \in \mathbb{R}$,

$$e^{-x^2} \leq \frac{1}{1+x^2} \leq \max \left\{ 1 - \frac{x^2}{2}, \frac{1}{2} \right\}. \quad (\text{C.12})$$

Hence

$$\begin{aligned} N^{-2} \sum_{j=1}^N \sum_{i=1}^N e^{-(v_j - v_i)^2 / 4\sigma^2} &\leq \max \left\{ 1 - \frac{1}{2N^2} \sum_{j=1}^N \sum_{i=1}^N (v_j - v_i)^2, \frac{1}{2} \right\} \\ &= \max \left\{ 1 - \frac{1}{N} \sum_{j=1}^N (v_j - \bar{v})^2, \frac{1}{2} \right\}. \end{aligned}$$

Therefore, if $\liminf_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N (v_j - \bar{v})^2 > 0$, then

$$\begin{aligned} \limsup_{N \rightarrow \infty} \mathbb{E} \mathcal{I}_{h,N} / h &\leq \frac{1}{2\sqrt{\pi}\sigma} \limsup_{N \rightarrow \infty} \max \left\{ 1 - \frac{1}{N} \sum_{j=1}^N (v_j - \bar{v})^2, \frac{1}{2} \right\} \\ &\leq \frac{1}{2\sqrt{\pi}\sigma} \max \left\{ 1 - \liminf_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N (v_j - \bar{v})^2, \frac{1}{2} \right\} \\ &< \frac{1}{2\sqrt{\pi}\sigma} = \mathcal{J}_b. \end{aligned}$$

Hence the proof is complete. \square

C.11 Proof of Theorem A.1

Proof. Recall the notations from [Appendix A.1](#). In this proof, we replace $\hat{\tau}^R$ by $\hat{\tau}_N$, where $\hat{\tau}_N$ is any \sqrt{N} -consistent estimator of τ_0 . For brevity, we shall skip the index N for $b_{N,i}$'s and $Z_{N,i}$'s in this proof. Define

$$V_N := N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{0 \leq b_j - b_i < N^{-\nu}\},$$

and note that [Assumption 3](#) yields that $V_N \rightarrow \mathcal{I}_b$ as $N \rightarrow \infty$. So, it suffices to show that $\hat{V}_N - V_N \xrightarrow{P} 0$ as $N \rightarrow \infty$. We write

$$\begin{aligned} \hat{V}_N - V_N &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{0 \leq \hat{b}_j - \hat{b}_i\} - \mathbf{1}\{0 \leq b_j - b_i\} \right) \\ &\quad - N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{N^{-\nu} \leq \hat{b}_j - \hat{b}_i\} - \mathbf{1}\{N^{-\nu} \leq b_j - b_i\} \right). \end{aligned} \quad (\text{C.13})$$

Since $\hat{b}_j - \hat{b}_i = b_j - b_i - (\hat{\tau}_N - \tau_0)(Z_j - Z_i)$, we can write

$$\begin{aligned} \Delta_{1,N} &:= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{0 \leq \hat{b}_j - \hat{b}_i\} - \mathbf{1}\{0 \leq b_j - b_i\} \right) \\ &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{(\hat{\tau}_N - \tau_0)(Z_j - Z_i) \leq b_j - b_i\} - \mathbf{1}\{0 \leq b_j - b_i\} \right) \\ &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{(\hat{\tau}_N - \tau_0)(Z_j - Z_i) \leq b_j - b_i < 0\} - \mathbf{1}\{0 \leq b_j - b_i < (\hat{\tau}_N - \tau_0)(Z_j - Z_i)\} \right). \end{aligned}$$

Now pick $\nu < \nu' < 1/2$ and let $E_N = \{|\hat{\tau}_N - \tau_0| > N^{-\nu'}\}$. Since $\hat{\tau}_N - \tau_0 = O_P(N^{-1/2})$ and $\nu' < 1/2$, we get $\hat{\tau}_N - \tau_0 = o_P(N^{-\nu'})$, implying that $P(E_N) \rightarrow 0$ as $N \rightarrow \infty$. Therefore for any $\varepsilon > 0$,

$$P(|\Delta_{1,N} \mathbf{1}_{E_N}| > \varepsilon) \leq P(\mathbf{1}_{E_N} = 1) = P(E_N) \rightarrow 0.$$

On the other hand, we have $|(\hat{\tau}_N - \tau_0)(Z_j - Z_i)| \leq N^{-\nu'}$ on E_N^c , hence

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{(\hat{\tau}_N - \tau_0)(Z_j - Z_i) \leq b_j - b_i < 0\} \mathbf{1}_{E_N^c}$$

$$\begin{aligned}
&\leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{-N^{-\nu'} \leq b_j - b_i < 0\} \mathbf{1}_{E_N^c} \\
&\leq N^{-(\nu'-\nu)} \left(N^{-(2-\nu')} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{-N^{-\nu'} \leq b_j - b_i < 0\} \right).
\end{aligned}$$

[Assumption 3](#) tells us that the term in the above parentheses converges to \mathcal{I}_b . Since $\nu' > \nu$, we conclude that

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{(\widehat{\tau}_N - \tau_0)(Z_j - Z_i) \leq b_j - b_i < 0\} \mathbf{1}_{E_N^c} \xrightarrow{P} 0.$$

Similarly,

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{0 \leq b_j - b_i < (\widehat{\tau}_N - \tau_0)(Z_j - Z_i)\} \mathbf{1}_{E_N^c} \xrightarrow{P} 0.$$

Thus, $\Delta_{1,N} \mathbf{1}_{E_N^c} \xrightarrow{P} 0$ and consequently $\Delta_{1,N} \xrightarrow{P} 0$, as $N \rightarrow \infty$.

We next focus on the second summand in the RHS of (C.13). The key idea to deal with this sum is to replace the indicators with smooth functions, such as a Gaussian CDF. Define $\sigma_N := N^{-\nu}(\log N)^{-1}$, $r_N(i, j) := b_j - b_i - N^{-\nu}$, and $\widehat{r}_N(i, j) := \widehat{b}_j - \widehat{b}_i - N^{-\nu}$. Then

$$\begin{aligned}
\widehat{V}_N - V_N - \Delta_{1,N} &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{N^{-\nu} \leq \widehat{b}_j - \widehat{b}_i\} - \mathbf{1}\{N^{-\nu} \leq b_j - b_i\} \right) \\
&= \Delta_{3,N} - \Delta_{2,N} + \Delta_{4,N} + \Delta_{5,N},
\end{aligned}$$

where

$$\begin{aligned}
\Delta_{2,N} &:= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{r_N(i, j) > 0\} - \Phi\left(\frac{r_N(i, j)}{\sigma_N}\right) \right), \\
\Delta_{3,N} &:= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{\widehat{r}_N(i, j) > 0\} - \Phi\left(\frac{\widehat{r}_N(i, j)}{\sigma_N}\right) \right), \\
\Delta_{4,N} &:= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\Phi\left(\frac{\widehat{r}_N(i, j)}{\sigma_N}\right) - \Phi\left(\frac{r_N(i, j)}{\sigma_N}\right) \right),
\end{aligned}$$

and

$$\Delta_{5,N} := N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N (\mathbf{1}\{\widehat{r}_N(i, j) = 0\} - \mathbf{1}\{r_N(i, j) = 0\}).$$

Showing $\Delta_{2,N} \rightarrow 0$, $\Delta_{k,N} \xrightarrow{P} 0$ for $k = 3, 4, 5$, will complete the proof.

(a) First we deal with $\Delta_{5,N}$. Note that

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{r_N(i, j) = 0\} \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{N^{-\nu} \leq b_j - b_i < (1 + \delta)N^{-\nu}\} \rightarrow \delta \mathcal{I}_b.$$

Now let $\delta \rightarrow 0$ to get

$$\lim_{N \rightarrow \infty} N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{r_N(i, j) = 0\} = 0.$$

For $\widehat{r}_N(i, j)$ we proceed just as in the proof of $\Delta_{1,N} \rightarrow 0$. Define $E_N = \{|\widehat{\tau}_N - \tau_0| > N^{-\nu'}\}$ for some $\nu < \nu' < 1/2$. Then

$$\begin{aligned} & N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{\widehat{r}_N(i, j) = 0\} \mathbf{1}_{E_N^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{N^{-\nu} \leq \widehat{b}_j - \widehat{b}_i < (1 + \delta)N^{-\nu}\} \mathbf{1}_{E_N^c} \\ & = N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{N^{-\nu} \leq b_j - b_i - (\widehat{\tau}_N - \tau_0)(Z_j - Z_i) < (1 + \delta)N^{-\nu}\} \mathbf{1}_{E_N^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{-|\widehat{\tau}_N - \tau_0| + N^{-\nu} \leq b_j - b_i < (1 + \delta)N^{-\nu} + |\widehat{\tau}_N - \tau_0|\} \mathbf{1}_{E_N^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{-N^{-\nu'} + N^{-\nu} \leq b_j - b_i < (1 + \delta)N^{-\nu} + N^{-\nu'}\}. \end{aligned}$$

Since $\nu' > \nu$, it holds for all sufficiently large N that $N^{-\nu'} \leq \delta N^{-\nu}$, and consequently

$$\begin{aligned} & \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \mathbb{E} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{\widehat{r}_N(i, j) = 0\} \mathbf{1}_{E_N^c} \\ & \leq \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{(1 - \delta)N^{-\nu} \leq b_j - b_i < (1 + 2\delta)N^{-\nu}\} = 3\delta \mathcal{I}_b. \end{aligned}$$

Letting $\delta \rightarrow 0$ here, and using $P(E_N) \rightarrow 0$, we conclude that

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{\widehat{r}_N(i, j) = 0\} \xrightarrow{P} 0, \text{ as } N \rightarrow \infty.$$

(b) Next we deal with $\Delta_{2,N}$, for which the bound provided by [Lemma E.2](#) is crucial. Recall that

$$\Delta_{2,N} = N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{r_N(i,j) > 0\} - \Phi\left(\frac{r_N(i,j)}{\sigma_N}\right) \right),$$

where $\sigma_N = N^{-\nu}(\log N)^{-1}$ and $r_N(i,j) = b_j - b_i - N^{-\nu}$. The key idea is to use the bound in [Lemma E.2](#) only for those i, j for which $r_N(i,j)$ is at least as large as $\delta N^{-\nu}$. Towards that, fix $\delta > 0$ and define

$$\mathcal{S}_{N,\delta} = \{(i,j) : |r_N(i,j)| \leq \frac{\delta}{N^\nu}, 1 \leq i, j \leq N\}.$$

Note that

$$\begin{aligned} (i,j) \in \mathcal{S}_{N,\delta} &\implies \delta N^{-\nu} \geq |r_N(i,j)| \geq ||b_j - b_i| - N^{-\nu}| \\ &\implies |b_j - b_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}]. \end{aligned}$$

Therefore [Lemma D.14](#) implies that

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} N^{-(2-\nu)} |\mathcal{S}_{N,\delta}| = 0. \quad (\text{C.14})$$

On the other hand, $(i,j) \notin \mathcal{S}_{N,\delta} \implies |r_N(i,j)| > \delta N^{-\nu}$, and then [Lemma E.2](#) yields that

$$\begin{aligned} \left| \mathbf{1}\{r_N(i,j) > 0\} - \Phi\left(\frac{r_N(i,j)}{\sigma_N}\right) \right| &\leq \sigma_N |r_N(i,j)|^{-1} \exp(-r_N^2(i,j)/2\sigma_N^2) \\ &\leq \sigma_N \delta^{-1} N^\nu \exp(-\delta^2 N^{-2\nu}/2\sigma_N^2) \\ &= (\log N)^{-1} \delta^{-1} \exp(-\delta^2 (\log N)^2/2) \end{aligned}$$

Combining the above bounds, the following chain of inequalities emerges.

$$\begin{aligned} |\Delta_{2,N}| &\leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left| \mathbf{1}\{r_N(i,j) > 0\} - \Phi\left(\frac{r_N(i,j)}{\sigma_N}\right) \right| \\ &\leq N^{-(2-\nu)} \sum_{(i,j) \in \mathcal{S}_{N,\delta}} 2 + N^{-(2-\nu)} \sum_{(i,j) \notin \mathcal{S}_{N,\delta}} \left| \mathbf{1}\{r_N(i,j) > 0\} - \Phi\left(\frac{r_N(i,j)}{\sigma_N}\right) \right| \\ &\leq 2N^{-(2-\nu)} |\mathcal{S}_{N,\delta}| + N^{-(2-\nu)} N^2 (\log N)^{-1} \delta^{-1} \exp(-\delta^2 (\log N)^2/2) \\ &= 2N^{-(2-\nu)} |\mathcal{S}_{N,\delta}| + (\log N)^{-1} \delta^{-1} \exp(\nu \log N - \delta^2 (\log N)^2/2). \end{aligned}$$

Therefore,

$$\limsup_{N \rightarrow \infty} |\Delta_{2,N}| \leq 2 \limsup_{N \rightarrow \infty} N^{-(2-\nu)} |\mathcal{S}_{N,\delta}|.$$

Letting $\delta \downarrow 0$ and invoking [\(C.14\)](#) we get the desired conclusion.

(c) Recall that

$$\Delta_{3,N} = N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\{\widehat{r}_N(i,j) > 0\} - \Phi\left(\frac{\widehat{r}_N(i,j)}{\sigma_N}\right) \right).$$

We can proceed just as in the previous proof. First we argue that for any fixed $\delta > 0$,

$$N^{-(2-\nu)} \sum_{(i,j): |\widehat{r}_N(i,j)| > \delta N^{-\nu}} \left| \mathbf{1}\{\widehat{r}_N(i,j) > 0\} - \Phi\left(\frac{\widehat{r}_N(i,j)}{\sigma_N}\right) \right| \xrightarrow{P} 0, \text{ as } N \rightarrow \infty.$$

Proof of the above part is exactly same as what we did in part (b). For the other part, it suffices to show that for any $\varepsilon > 0$,

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} P \left(N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{ \left| \widehat{b}_j - \widehat{b}_i \right| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} > \varepsilon \right) = 0 \quad (\text{C.15})$$

because the above implies that

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} P \left(N^{-(2-\nu)} \left| \{(i,j) : |\widehat{r}_N(i,j)| \leq \delta N^{-\nu}\} \right| > \varepsilon \right) = 0,$$

which further implies that

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} P(|\Delta_{3,N}| > \varepsilon) = 0.$$

To prove (C.15), we again use the event $\widetilde{E}_{N,K} = \{|\widehat{\tau}_N - \tau_0| > N^{-\nu'}\}$ where $\nu < \nu' < 1/2$.

$$\begin{aligned} & N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{ \left| \widehat{b}_j - \widehat{b}_i \right| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{E_N^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{ (1-\delta)N^{-\nu} \leq \widehat{b}_j - \widehat{b}_i < (1+\delta)N^{-\nu} \right\} \mathbf{1}_{E_N^c} \\ & = N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{ (1-\delta)N^{-\nu} \leq b_j - b_i - (\widehat{\tau}_N - \tau_0)(Z_j - Z_i) < (1+\delta)N^{-\nu} \right\} \mathbf{1}_{E_N^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{ -|\widehat{\tau}_N - \tau_0| + (1-\delta)N^{-\nu} \leq b_j - b_i < (1+\delta)N^{-\nu} + |\widehat{\tau}_N - \tau_0| \right\} \mathbf{1}_{E_N^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\left\{ -N^{-\nu'} + (1-\delta)N^{-\nu} \leq b_j - b_i < (1+\delta)N^{-\nu} + N^{-\nu'} \right\}. \end{aligned}$$

Since $\nu' > \nu$, it holds for all sufficiently large N that $N^{-\nu'} \leq \delta N^{-\nu}$, and consequently

$$\begin{aligned} & \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \mathbb{E} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \left| \widehat{b}_j - \widehat{b}_i \right| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{E_N^c} \\ & \leq \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ (1-2\delta)N^{-\nu} \leq b_j - b_i < (1+2\delta)N^{-\nu} \right\} = 4\delta \mathcal{I}_b. \end{aligned}$$

Letting $\delta \rightarrow 0$ here, we conclude that

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \mathbb{E} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \left| \widehat{b}_j - \widehat{b}_i \right| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{E_N^c} = 0. \quad (\text{C.16})$$

Finally, for any $\varepsilon > 0$,

$$\begin{aligned} & P \left(N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \left| \widehat{b}_j - \widehat{b}_i \right| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} > \varepsilon \right) \\ & \leq P \left(N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \left| \widehat{b}_j - \widehat{b}_i \right| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{E_N^c} > \varepsilon/2 \right) + P(E_N) \\ & \leq (\varepsilon/2)^{-1} N^{-(2-\nu)} \mathbb{E} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \left| \widehat{b}_j - \widehat{b}_i \right| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{E_N^c} + P(E_N). \end{aligned}$$

Invoking (C.16) and $\lim_{N \rightarrow \infty} P(E_N) = 0$, we complete the proof of (C.15).

(d) Next let us focus on $\Delta_{4,N}$. We first write

$$|\Delta_{4,N}| \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left| \Phi \left(\frac{\widehat{r}_N(i,j)}{\sigma_N} \right) - \Phi \left(\frac{r_N(i,j)}{\sigma_N} \right) \right|.$$

Now we break the last summation into three parts: (i) (i,j) such that $|r_N(i,j)| \leq \delta N^{-\nu}$, i.e., $(i,j) \in \mathcal{S}_{N,\delta}$; (ii) (i,j) such that $|\widehat{r}_N(i,j)| \leq \delta N^{-\nu}$; and (iii) remaining (i,j) 's, for which $|r_N(i,j)| \wedge |\widehat{r}_N(i,j)| > \delta N^{-\nu}$. For indices (i,j) in the cases (i) and (ii), we can use the crude bound $|\Phi(x) - \Phi(y)| \leq 2$, since equations (C.14) and (C.15) tell us that these contributions will be asymptotically negligible. For the case (iii), we use the following bound: for any $x, y \in \mathbb{R}$,

$$|\Phi(x) - \Phi(y)| \leq |x - y| \sup_{z \in [x \wedge y, x \vee y]} \phi(z) \leq |x - y| \exp \left(-\frac{x^2 \wedge y^2}{2} \right).$$

Thus, if $\mathcal{T}_{N,\delta}$ denote the (random) set of indices in case (iii), we have

$$\begin{aligned}
& N^{-(2-\nu)} \sum_{(i,j) \in \mathcal{T}_{N,\delta}} \left| \Phi \left(\frac{\widehat{r}_N(i,j)}{\sigma_N} \right) - \Phi \left(\frac{r_N(i,j)}{\sigma_N} \right) \right| \\
& \leq \sigma_N^{-1} N^{-(2-\nu)} \sum_{(i,j) \in \mathcal{T}_{N,\delta}} |\widehat{r}_N(i,j) - r_N(i,j)| \exp \left(-\frac{r_N(i,j)^2 \wedge \widehat{r}_N(i,j)^2}{2\sigma_N^2} \right) \\
& \leq \sigma_N^{-1} N^{-(2-\nu)} \sum_{(i,j) \in \mathcal{T}_{N,\delta}} |\widehat{r}_N(i,j) - r_N(i,j)| \exp \left(-\frac{\delta^2 N^{-2\nu}}{2\sigma_N^2} \right) \\
& \leq \sigma_N^{-1} N^{-(2-\nu)} \exp \left(-\frac{\delta^2 N^{-2\nu}}{2\sigma_N^2} \right) \sum_{j=1}^N \sum_{i=1}^N \left| (\widehat{b}_j - \widehat{b}_i) - (b_j - b_i) \right| \\
& = \sigma_N^{-1} N^{-(2-\nu)} \exp \left(-\frac{\delta^2}{2} (\log N)^2 \right) \sum_{j=1}^N \sum_{i=1}^N |(\widehat{\tau}_N - \tau_0)(Z_j - Z_i)| \\
& \leq \sigma_N^{-1} N^\nu |\widehat{\tau}_N - \tau_0| \exp \left(-\frac{\delta^2}{2} (\log N)^2 \right) \\
& = N^{2\nu-1/2} \log N \cdot \sqrt{N} |\widehat{\tau}_N - \tau_0| \exp \left(-\frac{\delta^2}{2} (\log N)^2 \right).
\end{aligned}$$

Since $\widehat{\tau}_N - \tau_0 = O_P(N^{-1/2})$, we get the desired conclusion that $\Delta_{4,N} \xrightarrow{P} 0$ as $N \rightarrow \infty$.

Combining the above steps, the proof is now complete. \square

C.12 Proof of Theorem A.2

Proof. Recall the notations from [Appendix A.2](#). In this proof, we replace $\widehat{\tau}_{\text{adj}}^R$ by $\widehat{\tau}_N$, where $\widehat{\tau}_N$ is any \sqrt{N} -consistent estimator of τ_0 . For brevity, we shall skip the index N for $b_{N,i}$, $Z_{N,i}$, \mathbf{X}_N , and $\mathbf{p}_{N,i}$ in this proof. To start with, we define

$$W_N := N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ 0 \leq \widetilde{b}_j - \widetilde{b}_i < N^{-\nu} \right\},$$

and note that $W_N \rightarrow \mathcal{J}_b$, by [Assumption 4](#). We are to show that $\widehat{W}_N - W_N \xrightarrow{P} 0$. Let I and J be two random indices chosen with replacement from $\{1, 2, \dots, N\}$. We replace the event $E_N = \{|\widehat{\tau}_N - \tau_0| > N^{-\nu'}\}$ in [Appendix C.11](#) with the following event:

$$\widetilde{E}_{N,K} = \{|\widehat{\tau}_N - \tau_0| > N^{-\nu'}\} \cup \{\sqrt{N} \|\mathbf{p}_J - \mathbf{p}_I\| > K\},$$

where $\nu < \nu' < 1/2$, and $K > 0$ is a fixed positive real number. Observe that

$$\begin{aligned} P(\sqrt{N}\|\mathbf{p}_J - \mathbf{p}_I\| > K) &\leq K^{-2}N \cdot \mathbb{E}_{J,I}\|\mathbf{p}_J - \mathbf{p}_I\|^2 \\ &= K^{-2}N^{-1} \sum_{j=1}^N \sum_{i=1}^N \|\mathbf{p}_j - \mathbf{p}_i\|^2 \\ &= 2(\text{rank}(\mathbf{X}) - 1)K^{-2}, \end{aligned}$$

where the last step follows from the fact that \mathbf{p}_j 's are columns of an idempotent matrix (see (E.4) for a proof). Thus,

$$P(\tilde{E}_{N,K}) \leq P(\sqrt{N}\|\mathbf{p}_J - \mathbf{p}_I\| > K) \lesssim 2(p-1)K^{-2},$$

where p is the number of covariates (i.e., number of columns of \mathbf{X}). Also note that,

$$\widehat{\tilde{b}}_j = Y_j - \widehat{\tau}_N Z_j - \mathbf{p}_j^\top (\mathbf{Y} - \widehat{\tau}_N \mathbf{Z}) = \tilde{b}_j - (\widehat{\tau}_N - \tau_0) (Z_j - \mathbf{p}_j^\top \mathbf{Z}).$$

Thus $\widehat{\tilde{b}}_j - \widehat{\tilde{b}}_i = \tilde{b}_j - \tilde{b}_i - (\widehat{\tau}_N - \tau_0)u_N(i, j)$, where $u_N(i, j) := Z_j - Z_i - (\mathbf{p}_j - \mathbf{p}_i)^\top \mathbf{Z}$. Now write

$$\begin{aligned} \widehat{W}_N - W_N &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\left\{0 \leq \widehat{\tilde{b}}_j - \widehat{\tilde{b}}_i\right\} - \mathbf{1}\left\{0 \leq \tilde{b}_j - \tilde{b}_i\right\} \right) \\ &\quad - N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\left\{N^{-\nu} \leq \widehat{\tilde{b}}_j - \widehat{\tilde{b}}_i\right\} - \mathbf{1}\left\{N^{-\nu} \leq \tilde{b}_j - \tilde{b}_i\right\} \right). \end{aligned}$$

For the first part, we imitate the proof of $\Delta_{1,N} \xrightarrow{P} 0$, as follows.

$$\begin{aligned} \tilde{\Delta}_{1,N} &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\left\{0 \leq \widehat{\tilde{b}}_j - \widehat{\tilde{b}}_i\right\} - \mathbf{1}\left\{0 \leq \tilde{b}_j - \tilde{b}_i\right\} \right) \\ &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\left\{(\widehat{\tau}_N - \tau_0)u_N(i, j) \leq \tilde{b}_j - \tilde{b}_i\right\} - \mathbf{1}\left\{0 \leq \tilde{b}_j - \tilde{b}_i\right\} \right) \\ &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1}\left\{(\widehat{\tau}_N - \tau_0)u_N(i, j) \leq \tilde{b}_j - \tilde{b}_i < 0\right\} - \mathbf{1}\left\{0 \leq \tilde{b}_j - \tilde{b}_i < (\widehat{\tau}_N - \tau_0)u_N(i, j)\right\} \right). \end{aligned}$$

Now fix any $\varepsilon > 0$ and note that on $\tilde{E}_{N,K}^c$ we have $|\widehat{\tau}_N - \tau_0| \leq N^{-\nu'}$, and

$$|u_N(i, j)| \leq |Z_j - Z_i| + |(\mathbf{p}_j - \mathbf{p}_i)^\top \mathbf{Z}| \leq 1 + \|\mathbf{p}_j - \mathbf{p}_i\| \cdot \|\mathbf{Z}\| \leq 1 + \sqrt{N}\|\mathbf{p}_j - \mathbf{p}_i\| \leq 1 + K.$$

Hence

$$\begin{aligned}
& N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ (\widehat{\tau}_N - \tau_0) u_N(i, j) \leq \widetilde{b}_j - \widetilde{b}_i < 0 \right\} \mathbf{1}_{\widetilde{E}_{N,K}^c} \\
& \leq N^\nu \cdot N^{-2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ -N^{-\nu'} (1+K) \leq \widetilde{b}_j - \widetilde{b}_i < 0 \right\} \mathbf{1}_{\widetilde{E}_{N,K}^c} \\
& \leq N^{-(\nu'-\nu)} \left(N^{-(2-\nu')} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ -N^{-\nu'} (1+K) \leq \widetilde{b}_j - \widetilde{b}_i < 0 \right\} \right).
\end{aligned}$$

Now [Assumption 4](#) tells us that the term in the above parentheses converges to $(1+K)\mathcal{J}_b$. Since $\nu' > \nu$, we conclude that

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ (\widehat{\tau}_N - \tau_0) u_N(i, j) \leq \widetilde{b}_j - \widetilde{b}_i < 0 \right\} \mathbf{1}_{\widetilde{E}_{N,K}^c} \xrightarrow{P} 0.$$

Similarly,

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ 0 \leq \widetilde{b}_j - \widetilde{b}_i < (\widehat{\tau}_N - \tau_0) u_N(i, j) \right\} \mathbf{1}_{\widetilde{E}_{N,K}^c} \xrightarrow{P} 0.$$

Thus $\widetilde{\Delta}_{1,N} \mathbf{1}_{\widetilde{E}_{N,K}^c} \xrightarrow{P} 0$. Hence

$$\begin{aligned}
\limsup_{N \rightarrow \infty} P \left(\left| \widetilde{\Delta}_{1,N} \right| > \varepsilon \right) & \leq \limsup_{N \rightarrow \infty} P \left(\left| \widetilde{\Delta}_{1,N} \mathbf{1}_{\widetilde{E}_{N,K}^c} \right| > \varepsilon/2 \right) \\
& \leq \limsup_{N \rightarrow \infty} P \left(\mathbf{1}_{\widetilde{E}_{N,K}^c} = 1 \right) \\
& = \limsup_{N \rightarrow \infty} P(\widetilde{E}_{N,K}) \leq 2(p-1)K^{-2}.
\end{aligned}$$

Letting $K \rightarrow \infty$ here, we conclude that $\widetilde{\Delta}_{1,N} \xrightarrow{P} 0$, as $N \rightarrow \infty$. Let us next focus on the second part of $\widehat{W}_N - W_N$, namely,

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ N^{-\nu} \leq \widehat{\widehat{b}}_j - \widehat{\widehat{b}}_i \right\} - \mathbf{1} \left\{ N^{-\nu} \leq \widetilde{b}_j - \widetilde{b}_i \right\} \right).$$

Once again we approximate the indicators in the above display using the CDF of $\mathcal{N}(0, \sigma_N^2)$ where $\sigma_N := N^{-\nu}(\log N)^{-1}$. Define

$$\widetilde{r}_N(i, j) := \widetilde{b}_j - \widetilde{b}_i - N^{-\nu}, \quad \text{and} \quad \widehat{\widehat{r}}_N(i, j) := \widehat{\widehat{b}}_j - \widehat{\widehat{b}}_i - N^{-\nu}.$$

The rest of the proof is essentially same as the proof of [Theorem A.1](#). We write

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ N^{-\nu} \leq \widehat{b}_j - \widehat{b}_i \right\} - \mathbf{1} \left\{ N^{-\nu} \leq \widetilde{b}_j - \widetilde{b}_i \right\} \right) = \widetilde{\Delta}_{3,N} - \widetilde{\Delta}_{2,N} + \widetilde{\Delta}_{4,N} + \widetilde{\Delta}_{5,N},$$

where

$$\begin{aligned} \widetilde{\Delta}_{2,N} &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ \widetilde{r}_N(i, j) > 0 \right\} - \Phi \left(\frac{\widetilde{r}_N(i, j)}{\sigma_N} \right) \right), \\ \widetilde{\Delta}_{3,N} &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ \widehat{r}_N(i, j) > 0 \right\} - \Phi \left(\frac{\widehat{r}_N(i, j)}{\sigma_N} \right) \right), \\ \widetilde{\Delta}_{4,N} &= N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\Phi \left(\frac{\widehat{r}_N(i, j)}{\sigma_N} \right) - \Phi \left(\frac{\widetilde{r}_N(i, j)}{\sigma_N} \right) \right), \end{aligned}$$

and

$$\widetilde{\Delta}_{5,N} = N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ \widehat{r}_N(i, j) = 0 \right\} - \mathbf{1} \left\{ \widetilde{r}_N(i, j) = 0 \right\} \right).$$

The proof of $\widetilde{\Delta}_{2,N} \rightarrow 0$ is exactly same as the proof of $\Delta_{2,N} \rightarrow 0$ given in [Appendix C.11](#), hence omitted. Next, in order to prove $\widetilde{\Delta}_{3,N} \xrightarrow{P} 0$, we first argue that for any fixed $\delta > 0$,

$$N^{-(2-\nu)} \sum_{(i,j): |\widehat{r}_N(i,j)| > \delta N^{-\nu}} \left| \mathbf{1} \left\{ \widehat{r}_N(i, j) > 0 \right\} - \Phi \left(\frac{\widehat{r}_N(i, j)}{\sigma_N} \right) \right| \xrightarrow{P} 0, \text{ as } N \rightarrow \infty.$$

Proof of the above display is omitted, for being completely analogous to the proof of part (b) in the proof of [Theorem A.1](#). For the other part of $\widetilde{\Delta}_{3,N}$, it suffices to show that for any $\varepsilon > 0$,

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} P \left(N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\widehat{b}_j - \widehat{b}_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} > \varepsilon \right) = 0 \quad (\text{C.17})$$

because the above implies that

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} P \left(N^{-(2-\nu)} \left| \{(i, j) : |\widehat{r}_N(i, j)| \leq \delta N^{-\nu}\} \right| > \varepsilon \right) = 0,$$

which further implies that

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} P \left(\left| \widetilde{\Delta}_{3,N} \right| > \varepsilon \right) = 0.$$

In order to prove (C.17), we once again use the event $\tilde{E}_{N,K}$ and the trick with I and J . For brevity, we use the notation

$$\kappa_{j,i} := (1 + \sqrt{N} \|\mathbf{p}_j - \mathbf{p}_i\|) |\hat{\tau}_N - \tau_0|.$$

Observe that,

$$\begin{aligned} & N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\hat{b}_j - \hat{b}_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ (1-\delta)N^{-\nu} \leq \hat{b}_j - \hat{b}_i < (1+\delta)N^{-\nu} \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \\ & = N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ (1-\delta)N^{-\nu} \leq \tilde{b}_j - \tilde{b}_i - (\hat{\tau}_N - \tau_0)u_N(i,j) < (1+\delta)N^{-\nu} \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ -\kappa_{j,i} + (1-\delta)N^{-\nu} \leq \tilde{b}_j - \tilde{b}_i < (1+\delta)N^{-\nu} + \kappa_{j,i} \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \\ & = N^\nu \cdot \mathbb{E}_{J,I} \left[\mathbf{1} \left\{ -\kappa_{J,I} + (1-\delta)N^{-\nu} \leq \tilde{b}_J - \tilde{b}_I < (1+\delta)N^{-\nu} + \kappa_{J,I} \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \right] \\ & \leq N^\nu \cdot \mathbb{E}_{J,I} \left[\mathbf{1} \left\{ -(1+K)|\hat{\tau}_N - \tau_0| + (1-\delta)N^{-\nu} \leq \tilde{b}_J - \tilde{b}_I \right. \right. \\ & \quad \left. \left. < (1+\delta)N^{-\nu} + (1+K)|\hat{\tau}_N - \tau_0| \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \right] \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ -(1+K)|\hat{\tau}_N - \tau_0| + (1-\delta)N^{-\nu} \leq \tilde{b}_j - \tilde{b}_i \right. \\ & \quad \left. < (1+\delta)N^{-\nu} + (1+K)|\hat{\tau}_N - \tau_0| \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \\ & \leq N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ -(1+K)N^{-\nu'} + (1-\delta)N^{-\nu} \leq \tilde{b}_j - \tilde{b}_i < (1+\delta)N^{-\nu} + (1+K)N^{-\nu'} \right\}. \end{aligned}$$

Since $\nu' > \nu$, it holds for all sufficiently large N that $(1+K)N^{-\nu'} \leq \delta N^{-\nu}$, and consequently

$$\begin{aligned} & \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \mathbb{E} \left(\sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\hat{b}_j - \hat{b}_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{\tilde{E}_{N,K}^c} \right) \\ & \leq \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ (1-2\delta)N^{-\nu} \leq \tilde{b}_j - \tilde{b}_i < (1+2\delta)N^{-\nu} \right\} = 4\delta \mathcal{J}_b. \end{aligned}$$

Letting $\delta \rightarrow 0$ here, we conclude that

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} N^{-(2-\nu)} \mathbb{E} \left(\sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\widehat{b}_j - \widehat{b}_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{\widetilde{E}_{N,K}^c} \right) = 0. \quad (\text{C.18})$$

Finally, for any $\varepsilon > 0$,

$$\begin{aligned} & P \left(N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\widehat{b}_j - \widehat{b}_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} > \varepsilon \right) \\ & \leq P \left(N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\widehat{b}_j - \widehat{b}_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{\widetilde{E}_{N,K}^c} > \varepsilon/2 \right) + P(\widetilde{E}_{N,K}) \\ & \leq (\varepsilon/2)^{-1} N^{-(2-\nu)} \mathbb{E} \left(\sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\widehat{b}_j - \widehat{b}_i| \in [(1-\delta)N^{-\nu}, (1+\delta)N^{-\nu}] \right\} \mathbf{1}_{\widetilde{E}_{N,K}^c} \right) + P(\widetilde{E}_{N,K}). \end{aligned}$$

Invoking (C.18) and using $\lim_{K \rightarrow \infty} \limsup_{N \rightarrow \infty} P(\widetilde{E}_{N,K}) = 0$, we complete the proof of (C.17), which, in turn, proves that $\widetilde{\Delta}_{3,N} \xrightarrow{P} 0$.

For $k = 4$ or 5 , the proof of $\widetilde{\Delta}_{k,N} \xrightarrow{P} 0$ follows by making similar changes in the proof of $\Delta_{k,N} \xrightarrow{P} 0$ in Appendix C.11, just as we did for $k = 1$ or 3 . \square

C.13 Proof of Proposition B.1

Proof. The statistic t_N is defined as

$$t_N = \widehat{\mathbf{q}}_N^\top \mathbf{Z}_N, \quad \widehat{q}_N(j) = \sum_{i=1}^N \mathbf{1} \{Y_{N,i} - \tau_0 Z_{N,i} \leq Y_{N,j} - \tau_0 Z_{N,j}\}, \quad 1 \leq j \leq N.$$

Note, the distribution of t_N under $\tau = \tau_N$ depends on τ_N only through the vector $\widehat{\mathbf{q}}_N$. Under $\tau = \tau_N$ we can write

$$(Y_{N,1} - \tau_0 Z_{N,1}, \dots, Y_{N,N} - \tau_0 Z_{N,N}) \stackrel{d}{=} (b_{N,1} + (\tau_N - \tau_0)Z_{N,1}, \dots, b_{N,N} + (\tau_N - \tau_0)Z_{N,N})$$

and hence the distribution of $\widehat{\mathbf{q}}_N^\top \mathbf{Z}_N$ under $\tau = \tau_N$ is same as the randomization distribution of $\mathbf{q}_N^\top \mathbf{Z}_N$ where

$$q_{N,j} = \sum_{i=1}^N \mathbf{1} \{b_{N,i} + (\tau_N - \tau_0)Z_{N,i} \leq b_{N,j} + (\tau_N - \tau_0)Z_{N,j}\}, \quad 1 \leq j \leq N.$$

In order words, we have $t_N \stackrel{d}{=} t_N^*$ under $\tau = \tau_N$, where

$$\begin{aligned}
t_N^* &= \sum_{j=1}^N Z_{N,j} \sum_{i=1}^N \mathbf{1}\{b_{N,i} + (\tau_N - \tau_0)Z_{N,i} \leq b_{N,j} + (\tau_N - \tau_0)Z_{N,j}\} \\
&= \sum_{j=1}^N Z_{N,j} \sum_{i=1}^N (Z_{N,i} \mathbf{1}\{b_{N,i} \leq b_{N,j}\} + (1 - Z_{N,i}) \mathbf{1}\{hN^{-1/2} \leq b_{N,j} - b_{N,i}\}) \\
&= \sum_{j=1}^N Z_{N,j} + \sum_{j=1}^N \sum_{i=1, i \neq j}^N Z_{N,j} Z_{N,i} \mathbf{1}\{b_{N,i} \leq b_{N,j}\} \\
&\quad + \sum_{j=1}^N \sum_{i=1, i \neq j}^N Z_{N,j} (1 - Z_{N,i}) \mathbf{1}\{hN^{-1/2} \leq b_{N,j} - b_{N,i}\}.
\end{aligned}$$

Now when $h \geq 0$,

$$\mathbf{1}\{hN^{-1/2} \leq b_{N,j} - b_{N,i}\} = \mathbf{1}\{0 \leq b_{N,j} - b_{N,i}\} - \mathbf{1}\{0 \leq b_{N,j} - b_{N,i} < hN^{-1/2}\}$$

and thus

$$t_N^* = m + \sum_{j=1}^N \sum_{i=1, i \neq j}^N Z_{N,j} \mathbf{1}\{b_{N,i} \leq b_{N,j}\} - \sum_{j=1}^N \sum_{i=1, i \neq j}^N (1 - Z_{N,i}) Z_{N,j} \mathbf{1}\{0 \leq b_{N,j} - b_{N,i} < hN^{-1/2}\}.$$

Similarly, for $h < 0$,

$$\mathbf{1}\{hN^{-1/2} \leq b_{N,j} - b_{N,i}\} = \mathbf{1}\{0 \leq b_{N,j} - b_{N,i}\} + \mathbf{1}\{hN^{-1/2} \leq b_{N,j} - b_{N,i} < 0\}$$

which gives us the desired expression. \square

D Some Technical Lemmas

Lemma D.1. *Let $t(\mathbf{Z}, \mathbf{Y} - \tau_0 \mathbf{Z})$ be a test statistic for testing $H_0 : \tau = \tau_0$ and $\hat{\tau}^R$ be the estimator of τ based on $t(\cdot, \cdot)$, as defined in (2.11) of the main paper. Assume that for all values of \mathbf{y} and \mathbf{z} , $t(\mathbf{z}, \mathbf{y} - \tau \mathbf{z})$ is a non-increasing function of τ . Let $\{c_N\}_{N \geq 1}$ and $\{d_N\}_{N \geq 1}$ be sequences of positive real numbers and μ_N be the mean of $t_N = t(\mathbf{Z}_N, \mathbf{Y}_N - \tau_0 \mathbf{Z}_N)$ under $\tau = \tau_0$. Fix $h \in \mathbb{R}$ and define $\tau_N := \tau_0 - h/c_N$. Suppose that*

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\tau_N}(d_N(t_N - \mu_N) \leq x) = G((x + hB)/A) \quad (\text{D.1})$$

holds for every $x \in \mathbb{R}$, where G is a distribution function of a continuous random variable with mean 0 and variance 1, and $A, B > 0$ are constants. Then it holds that

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\tau_0} (c_N(\hat{\tau}^R - \tau_0) \leq h) = G(hB/A). \quad (\text{D.2})$$

Proof. Denote $\tau'_N = \tau_0 + h/c_N$. We use [Lemma D.2](#) to write the following:

$$\begin{aligned} \mathbb{P}_{\tau_0} (t_N(\mathbf{Z}, \mathbf{Y} - \tau'_N \mathbf{Z}) < \mu_N) &\leq \mathbb{P}_{\tau_0} (\hat{\tau}^* \leq \tau'_N) \\ &\leq \mathbb{P}_{\tau_0} (\hat{\tau}^R \leq \tau'_N) \\ &\leq \mathbb{P}_{\tau_0} (\hat{\tau}^{**} \leq \tau'_N) \leq \mathbb{P}_{\tau_0} (t_N(\mathbf{Z}, \mathbf{Y} - \tau'_N \mathbf{Z}) \leq \mu_N). \end{aligned} \quad (\text{D.3})$$

In view of [Lemma D.3](#), the distribution of $t_N(\mathbf{Z}, \mathbf{Y} - \tau'_N \mathbf{Z})$ under $\tau = \tau_0$ is same as the distribution of $t_N = t_N(\mathbf{Z}, \mathbf{Y} - \tau_0 \mathbf{Z})$ under $\tau = \tau_N$, which we denote by G_N . Then equation (D.1) tells us that $G_N(x/d_N + \mu_N) \rightarrow G((x+hB)/A)$ as $n \rightarrow \infty$, for every $x \in \mathbb{R}$. In particular, for $x = 0$ we can say that $G_N(\mu_N) \rightarrow G(hB/A)$ and that $G_N(\mu_N -) \rightarrow G(hB/A -)$ which is also equal to $G(hB/A)$ since G is continuous. Therefore both the extreme sides of (D.3) converge to $G(hB/A)$ and thus by sandwich principle, equation (D.2) holds. \square

Lemma D.2. Let $t(\cdot, \cdot)$ be a test statistic such that for all values of \mathbf{y} and \mathbf{z} , $t(\mathbf{z}, \mathbf{y} - \tau \mathbf{z})$ is a non-increasing function of τ . Then for any $h \in \mathbb{R}$,

$$\begin{aligned} \mathbb{P}_{\tau} (t(\mathbf{Z}, \mathbf{Y} - a \mathbf{Z}) < \mu) &\leq \mathbb{P}_{\tau} (\hat{\tau}^* \leq a) \leq \mathbb{P}_{\tau} (\hat{\tau}^R \leq a) \\ &\leq \mathbb{P}_{\tau} (\hat{\tau}^{**} \leq a) \leq \mathbb{P}_{\tau} (t(\mathbf{Z}, \mathbf{Y} - a \mathbf{Z}) \leq \mu). \end{aligned}$$

Proof. The proof is straightforward from the definitions of $\hat{\tau}^*$ and $\hat{\tau}^{**}$. First observe that $\tau^* \leq \hat{\tau}^R \leq \tau^{**}$ holds almost surely. Now, if a is such that $t(\mathbf{Z}, \mathbf{Y} - a \mathbf{Z}) < \mu$, then

$$h \geq \sup\{\tau : t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) < \mu\} = \hat{\tau}^{**} \implies h \geq \hat{\tau}^*,$$

so the left-most inequality follows. Next,

$$\hat{\tau}^* \leq a \implies \hat{\tau}^R \leq a \implies \hat{\tau}^{**} \leq a.$$

Finally, suppose that a is such that $t(\mathbf{Z}, \mathbf{Y} - a \mathbf{Z}) > \mu$. We split it into two cases:

(a) when $\hat{\tau}^* < \hat{\tau}^{**}$, we have

$$t(\mathbf{Z}, \mathbf{Y} - a \mathbf{Z}) > \mu \implies a \leq \sup\{\tau : t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) > \mu\} = \hat{\tau}^* < \hat{\tau}^{**}.$$

(b) When $\hat{\tau}^* = \hat{\tau}^{**}$, we have

$$t(\mathbf{Z}, \mathbf{Y} - a \mathbf{Z}) > \mu \implies h < \sup\{\tau : t(\mathbf{Z}, \mathbf{Y} - \tau \mathbf{Z}) > \mu\} = \hat{\tau}^* = \hat{\tau}^{**}.$$

This finishes the proof. \square

Lemma D.3. *The distribution of $t(\mathbf{Z}, \mathbf{Y} - (\tau + \delta)\mathbf{Z})$ under $\tau = \tau_0$ is identical to the distribution of $t(\mathbf{Z}, \mathbf{Y} - \tau\mathbf{Z})$ under $\tau = \tau_0 - \delta$.*

Proof. For any $x \in \mathbb{R}$,

$$\begin{aligned}\mathbb{P}_{\tau_0}(t(\mathbf{Z}, \mathbf{Y} - (\tau + \delta)\mathbf{Z}) \leq x) &= \mathbb{P}_{\tau_0}(t(\mathbf{Z}, \tau_0\mathbf{Z} + \mathbf{b} - (\tau + \delta)\mathbf{Z}) \leq x) \\ &= P(t(\mathbf{Z}, (\tau_0 - \delta)\mathbf{Z} + \mathbf{b} - \tau\mathbf{Z}) \leq x) \\ &= \mathbb{P}_{\tau_0 - \delta}(t(\mathbf{Z}, \mathbf{Y} - \tau\mathbf{Z}) \leq x),\end{aligned}$$

where the second step is due to the fact that the randomization distribution of \mathbf{Z} is free of τ . \square

Lemma D.4. *If $\{b_{N,j} : 1 \leq j \leq N\}$ be i.i.d. from a distribution with density $f_b(\cdot)$ satisfying $\int_{\mathbb{R}} f_b^2(x) dx < \infty$, and $I_{h,N}$ be as in (2.15) of the main paper, then*

$$N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N I_{h,N}(b_{N,j} - b_{N,i}) \xrightarrow{a.s.} h \int_{\mathbb{R}} f_b^2(x) dx.$$

Proof. Let us denote $S_N = S_N(b_{N,1}, \dots, b_{N,N}) = \sum_{j=1}^N \sum_{i=1}^N I_{h,N}(b_{N,j} - b_{N,i})$. Observe that for any fixed $i \neq j$, and $h > 0$,

$$\mathbb{E}(I_{h,N}(b_{N,j} - b_{N,i})) = P(0 \leq b_{N,2} - b_{N,1} < hN^{-1/2}) = g(hN^{-1/2}) - g(0)$$

where

$$g(x) = P(b_{N,2} - b_{N,1} \leq x) = \int_{-\infty}^x \int_{\mathbb{R}} f_b(u+t)f_b(u) du dt, \quad x \in \mathbb{R}.$$

Using the DCT for integrals, we argue that $g'(x) = \int_{\mathbb{R}} f_b(u+x)f_b(u)du$. Hence

$$\begin{aligned}\lim_{N \rightarrow \infty} N^{-3/2} \mathbb{E}(S_N) &= \lim_{N \rightarrow \infty} N^{-3/2} N^2 (g(hN^{-1/2}) - g(0)) \\ &= \lim_{N \rightarrow \infty} h \cdot \frac{g(hN^{-1/2}) - g(0)}{hN^{-1/2}} \\ &= hg'(0) \\ &= h \int_{\mathbb{R}} f_b(u)^2 du.\end{aligned} \tag{D.4}$$

Now we bound $\mathbb{E}(S_N - \mathbb{E}S_N)^2$ using the Efron-Stein inequality (Efron and Stein, 1981). For each $1 \leq k \leq N$, let $b'_{N,k}$ be an i.i.d. copy of $b_{N,k}$, independent of everything else, and define

$$S_N^{(k)} = S_N(b_{N,1}, \dots, b_{N,k-1}, b'_{N,k}, b_{N,k+1}, \dots, b_{N,N}), \quad 1 \leq k \leq N.$$

Note that

$$\begin{aligned} S_N - S_N^{(k)} &= \sum_{j=1}^N (I_{h,N}(b_{N,j} - b_{N,k}) - I_{h,N}(b_{N,j} - b'_{N,k})) \\ &\quad + \sum_{j=1}^N (I_{h,N}(b_{N,k} - b_{N,j}) - I_{h,N}(b'_{N,k} - b_{N,j})). \end{aligned}$$

An application of the Cauchy-Schwarz inequality yields the following.

$$\begin{aligned} &\mathbb{E} \left[\left(\sum_{j=1}^N I_{h,N}(b_{N,j} - b_{N,k}) - I_{h,N}(b_{N,j} - b'_{N,k}) \right)^2 \mid b_{N,1}, \dots, b_{N,N} \right] \\ &\leq N \mathbb{E} \left[\sum_{j=1}^N (I_{h,N}(b_{N,j} - b_{N,k}) - I_{h,N}(b_{N,j} - b'_{N,k}))^2 \mid b_{N,1}, \dots, b_{N,N} \right] \\ &\leq 2N \mathbb{E} \left[\sum_{j=1}^N (I_{h,N}(b_{N,j} - b_{N,k}) + I_{h,N}(b_{N,j} - b'_{N,k})) \mid b_{N,1}, \dots, b_{N,N} \right] \\ &= 2N \sum_{j=1}^N (I_{h,N}(b_{N,j} - b_{N,k}) + \mathbb{E}[I_{h,N}(b_{N,j} - b_{N,k}) \mid b_{N,j}]). \end{aligned}$$

Applying the same argument to the second part yields

$$\begin{aligned} &\mathbb{E} \left[\left(\sum_{j=1}^N I_{h,N}(b_{N,k} - b_{N,j}) - I_{h,N}(b'_{N,k} - b_{N,j}) \right)^2 \mid b_{N,1}, \dots, b_{N,N} \right] \\ &\leq 2N \sum_{j=1}^N (I_{h,N}(b_{N,k} - b_{N,j}) + \mathbb{E}[I_{h,N}(b_{N,k} - b_{N,j}) \mid b_{N,j}]). \end{aligned}$$

Using the above bounds and a generalized Efron-Stein inequality (see [Proposition E.1](#)),

$$\begin{aligned} &\mathbb{E}(N^{-3/2}(S_N - \mathbb{E}S_N))^6 \\ &\lesssim N^{-9} \mathbb{E} \left[\sum_{k=1}^N \mathbb{E} \left[(S_N - S_N^{(k)})^2 \mid b_{N,1}, \dots, b_{N,N} \right] \right]^3 \\ &\lesssim N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N (I_{h,N}(b_{N,j} - b_{N,k}) + \mathbb{E}[I_{h,N}(b_{N,j} - b_{N,k}) \mid b_{N,j}]) \right]^3 \\ &\quad + N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N (I_{h,N}(b_{N,k} - b_{N,j}) + \mathbb{E}[I_{h,N}(b_{N,k} - b_{N,j}) \mid b_{N,j}]) \right]^3 \end{aligned}$$

$$\begin{aligned}
&\lesssim N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N I_{h,N}(b_{N,j} - b_{N,k}) \right]^3 + N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N \mathbb{E} [I_{h,N}(b_{N,j} - b_{N,k}) \mid b_{N,j}] \right]^3 \\
&\quad + N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N \mathbb{E} [I_{h,N}(b_{N,k} - b_{N,j}) \mid b_{N,j}] \right]^3.
\end{aligned}$$

We now show that the contribution of each of last three sums is at most of the order of $N^{-3/2}$, which would complete the proof, with the aid of Borel-Cantelli lemma. We start with

$$\begin{aligned}
&N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N I_{h,N}(b_{N,j} - b_{N,k}) \right]^3 \\
&= N^{-6} \sum_{1 \leq i_1, j_1, i_2, j_2, i_3, j_3 \leq N} \mathbb{E} [I_{h,N}(b_{N,j_1} - b_{N,i_1}) I_{h,N}(b_{N,j_2} - b_{N,i_2}) I_{h,N}(b_{N,j_3} - b_{N,i_3})]
\end{aligned}$$

When $|\{i_1, j_1, i_2, j_2, i_3, j_3\}| \leq 4$, the number of ways to choose such a set of indices would be $O(N^4)$, and the summands being bounded above by 1, the contributions from these terms is $O(N^{-2})$. If $|\{i_1, j_1, i_2, j_2, i_3, j_3\}| = 6$, i.e., the indices are all distinct, we use the independence of the $b_{N,i}$'s to split the joint probability as the product of marginal probabilities, and thus the contribution from such terms becomes

$$\begin{aligned}
&N^{-6} \sum_{i_1, j_1, i_2, j_2, i_3, j_3 \text{ distinct}} \mathbb{E} I_{h,N}(b_{N,j_1} - b_{N,i_1}) \mathbb{E} I_{h,N}(b_{N,j_2} - b_{N,i_2}) \mathbb{E} I_{h,N}(b_{N,j_3} - b_{N,i_3}) \\
&\leq N^{-6} \left(\sum_{1 \leq i, j \leq N} \mathbb{E} I_{h,N}(b_{N,j} - b_{N,i}) \right)^3 = N^{-3/2} (\mathbb{E} N^{-3/2} S_N)^3 = O(N^{-3/2}).
\end{aligned}$$

Finally, consider the case where $|\{i_1, j_1, i_2, j_2, i_3, j_3\}| = 5$, i.e., exactly one index is repeated. Then at least two of the sets $\{i_1, j_1\}, \{i_2, j_2\}, \{i_3, j_3\}$ will make four distinct indices altogether. If the set left-out is $\{i_3, j_3\}$, we can just upper bound $I_{h,N}(b_{N,j_3} - b_{N,i_3})$ by 1. Also, the number of ways of choosing the indices $\{i_3, j_3\}$ will be $O(N)$, since one of them is repeated within $\{i_1, j_1, i_2, j_2\}$. Using this idea,

$$\begin{aligned}
&N^{-6} \sum_{|i_1, j_1, i_2, j_2, i_3, j_3|=5} \mathbb{E} [I_{h,N}(b_{N,j_1} - b_{N,i_1}) I_{h,N}(b_{N,j_2} - b_{N,i_2}) I_{h,N}(b_{N,j_3} - b_{N,i_3})] \\
&\lesssim N^{-5} \sum_{i_1, j_1, i_2, j_2 \text{ distinct}} \mathbb{E} I_{h,N}(b_{N,j_1} - b_{N,i_1}) \mathbb{E} I_{h,N}(b_{N,j_2} - b_{N,i_2}) \\
&\leq N^{-5} \left(\sum_{1 \leq i, j \leq N} \mathbb{E} I_{h,N}(b_{N,j} - b_{N,i}) \right)^2 = N^{-2} (\mathbb{E} N^{-3/2} S_N)^2 = O(N^{-2}).
\end{aligned}$$

Combining the above cases we conclude that $N^{-6} \mathbb{E} [\sum_{k=1}^N \sum_{j=1}^N I_{h,N}(b_{N,j} - b_{N,k})]^3 = O(N^{-3/2})$.

We next look at

$$\begin{aligned}
& N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N \mathbb{E} [I_{h,N}(b_{N,j} - b_{N,k}) \mid b_{N,j}] \right]^3 \\
&= N^{-6} \sum_{1 \leq i_1, j_1, i_2, j_2, i_3, j_3 \leq N} \mathbb{E} [I^*(b_{N,j_1}, b_{N,i_1}) I^*(b_{N,j_2}, b_{N,i_2}) I^*(b_{N,j_3}, b_{N,i_3})]
\end{aligned}$$

where $I^*(b_{N,j}, b_{N,i}) := \mathbb{E} [I_{h,N}(b_{N,j} - b_{N,i}) \mid b_{N,j}]$, which is really a function of $b_{N,j}$. Thus, when all the 6 indices are distinct, the joint probability is split into marginals, and once again we can see that the contribution of these terms is $O(N^{-3/2})$. When $|\{i_1, j_1, i_2, j_2, i_3, j_3\}| \leq 4$, the same argument applies as we gave earlier, and tells us that contribution from these terms would be $O(N^{-2})$. Finally, when $|\{i_1, j_1, i_2, j_2, i_3, j_3\}| = 5$ we play the same trick applied in the previous case to conclude that contribution from these terms is

$$\begin{aligned}
& N^{-6} \sum_{|i_1, j_1, i_2, j_2, i_3, j_3|=5} \mathbb{E} [I^*(b_{N,j_1}, b_{N,i_1}) I^*(b_{N,j_2}, b_{N,i_2}) I^*(b_{N,j_3}, b_{N,i_3})] \\
& \lesssim N^{-5} \sum_{i_1, j_1, i_2, j_2 \text{ distinct}} \mathbb{E} [I^*(b_{N,j_1}, b_{N,i_1}) I^*(b_{N,j_2}, b_{N,i_2})] \\
&= N^{-5} \sum_{i_1, j_1, i_2, j_2 \text{ distinct}} \mathbb{E} [I_{h,N}(b_{N,j_1} - b_{N,i_1})] \mathbb{E} [I_{h,N}(b_{N,j_2} - b_{N,i_2})] \\
& \leq N^{-5} \left(\sum_{1 \leq i, j \leq N} \mathbb{E} I_{h,N}(b_{N,j} - b_{N,i}) \right)^2 = N^{-2} (\mathbb{E} N^{-3/2} S_N)^2 = O(N^{-2}).
\end{aligned}$$

Thus, $N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N \mathbb{E} [I_{h,N}(b_{N,j} - b_{N,k}) \mid b_{N,j}] \right]^3 = O(N^{-3/2})$. In a similar manner one can show that $N^{-6} \mathbb{E} \left[\sum_{k=1}^N \sum_{j=1}^N \mathbb{E} [I_{h,N}(b_{N,k} - b_{N,j}) \mid b_{N,j}] \right]^3 = O(N^{-3/2})$. Hence the proof. \square

Lemma D.5. *Suppose that the potential control outcomes $\{b_{N,j}\}_{1 \leq j \leq N}$ satisfy [Assumption 1](#). Define*

$$q_{N,j} = \sum_{i=1}^N \mathbf{1}\{b_{N,i} \leq b_{N,j}\}, \quad 1 \leq j \leq N.$$

Then, as $N \rightarrow \infty$,

$$\bar{q}_N := \frac{1}{N} \sum_{j=1}^N q_{N,j} = \frac{N+1}{2} + o(N^{1/2}), \quad \text{and} \quad \frac{1}{N} \sum_{j=1}^N (q_{N,j} - \bar{q}_N)^2 = \frac{N^2 - 1}{12} + o(N^2).$$

Proof. We first break the ties in an arbitrary manner, and obtain $b_{N(1)} \leq b_{N(2)} \leq \dots \leq b_{N(N)}$.

Define $r_{N,j} = \sum_{i=1}^N \mathbf{1}\{b_{N,i} \leq b_{N(j)}\}$. Note that for each $1 \leq j \leq N$,

$$|r_{N,j} - j| \leq \sum_{i=1}^N \mathbf{1}\{b_{N,i} = b_{N(j)}\}.$$

The last equation in conjunction with [Assumption 1](#) yields the following bounds.

$$\begin{aligned} \sum_{j=1}^N |r_{N,j} - j| &\leq \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{b_{N,i} = b_{N,j}\} \\ &\leq \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{0 \leq b_{N,j} - b_{N,i} < N^{-1/2}\} \lesssim N^{3/2}, \end{aligned}$$

and

$$\begin{aligned} \sum_{j=1}^N (r_{N,j} - j)^2 &\leq \sum_{j=1}^N N \sum_{i=1}^N \mathbf{1}\{b_{N,i} = b_{N,j}\} \\ &\leq N \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{0 \leq b_{N,j} - b_{N,i} < N^{-1/2}\} \lesssim N^{5/2}. \end{aligned}$$

Now,

$$\begin{aligned} \left| \sum_{i=1}^N r_{N,i}^2 - \sum_{i=1}^N i^2 \right| &= \sum_{i=1}^N (r_{N,i} - i)^2 + \left| \sum_{i=1}^N 2i(r_{N,i} - i) \right| \\ &\leq \sum_{i=1}^N (r_{N,i} - i)^2 + 2 \left(\sum_{i=1}^N i^2 \sum_{i=1}^N (r_{N,i} - i)^2 \right)^{1/2} \\ &\lesssim N^{5/2} + N^{11/4} = o(N^3). \end{aligned}$$

On the other hand,

$$\begin{aligned} N \left| \bar{r}_N^2 - ((N+1)/2)^2 \right| &= \frac{1}{N} \left| \sum_{i=1}^N (r_{N,i} - i) \sum_{i=1}^N (r_{N,i} + i) \right| \\ &\leq \frac{1}{N} \sum_{i=1}^N |r_{N,i} - i| \sum_{i=1}^N (|r_{N,i} - i| + 2i) \\ &\lesssim N^{-1} N^{3/2} (N^{3/2} + N^2) = o(N^3). \end{aligned}$$

Combining the above with the fact that $\{r_{N,1}, \dots, r_{N,N}\}$ is same as $\{q_{N,1}, \dots, q_{N,N}\}$,

$$\sum_{i=1}^N (q_{N,i} - \bar{q}_N)^2 = \sum_{i=1}^N r_{N,i}^2 - N\bar{r}_N^2 = \sum_{i=1}^N \left(i - \frac{N+1}{2}\right)^2 + o(N^3) = \frac{N(N^2-1)}{12} + o(N^3).$$

This finishes the proof of the second assertion. To prove the first assertion, we fix any $\delta > 0$ and note that

$$\begin{aligned} \left| \sum_{j=1}^N q_{N,j} - \frac{N(N+1)}{2} \right| &= \left| \sum_{j=1}^N (r_{N,j} - j) \right| \\ &\leq \sum_{j=1}^N |r_{N,j} - j| \\ &\leq \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{b_{N,i} = b_{N,j}\} \\ &\leq \sum_{j=1}^N \sum_{i=1}^N \mathbf{1}\{0 \leq b_{N,j} - b_{N,i} < \delta N^{-1/2}\}. \end{aligned}$$

Invoking [Assumption 1](#) we can say that

$$\limsup_{N \rightarrow \infty} \left| \sum_{j=1}^N q_{N,j} - \frac{N(N+1)}{2} \right| \leq \delta \mathcal{I}_b.$$

Now letting $\delta \rightarrow 0$ completes the argument. \square

Lemma D.6. *Suppose that ranks $\{q_{N,j}\}$ satisfy $\sum_{j=1}^N (q_{N,j} - \bar{q}_N)^2 = \frac{N(N^2-1)}{12} + o(N^3)$ as $N \rightarrow \infty$. Let t_N be the Wilcoxon rank-sum statistic when the treatments are assigned by an m -out-of- N SRSWOR sample where $m/N \rightarrow \lambda \in (0, 1)$ as $N \rightarrow \infty$. Then, under $\tau = \tau_0$,*

$$\text{Var}(t_N) \sim \frac{\lambda(1-\lambda)}{12} N^3 \text{ as } N \rightarrow \infty.$$

Proof. Recall from [\(C.1\)](#) that under $\tau = \tau_0$, we can write $t_N \stackrel{d}{=} \sum_{j=1}^N q_{N,j} Z_{N,j}$, where $q_{N,j} = \sum_{i=1}^N \mathbf{1}\{b_{N,i} \leq b_{N,j}\}$. Note that each $Z_{N,j}$ is a Bernoulli(m/N) random variable, and for $i \neq j$ we have

$$\begin{aligned} \text{Cov}(Z_{N,i}, Z_{N,j}) &= P(Z_{N,i} = 1, Z_{N,j} = 1) - P(Z_{N,i} = 1)P(Z_{N,j} = 1) \\ &= \frac{m(m-1)}{N(N-1)} - \left(\frac{m}{N}\right)^2 = -\frac{1}{N-1} \cdot \frac{m}{N} \left(1 - \frac{m}{N}\right). \end{aligned}$$

Hence we deduce that, under $\tau = \tau_0$,

$$\begin{aligned}\text{Var}(t_N) &= \sum_{j=1}^N \text{Var}(q_{N,j} Z_{N,j}) + 2 \sum_{1 \leq i < j \leq N} \text{Cov}(q_{N,i} Z_{N,i}, q_{N,j} Z_{N,j}) \\ &= \sum_{j=1}^N q_{N,j}^2 \frac{m}{N} \left(1 - \frac{m}{N}\right) - 2 \sum_{1 \leq i < j \leq N} q_{N,i} q_{N,j} \frac{m}{N} \left(1 - \frac{m}{N}\right) \frac{1}{N-1} \\ &= \frac{m}{N} \left(1 - \frac{m}{N}\right) \frac{N}{N-1} \sum_{j=1}^N (q_{N,j} - \bar{q}_N)^2.\end{aligned}$$

In light of the given condition on the ranks, the above implies that

$$\text{Var}(t_N) = \frac{m}{N} \left(1 - \frac{m}{N}\right) \frac{N}{N-1} \left(\frac{N(N^2-1)}{12} + o(N^3) \right) \sim \lambda(1-\lambda) \frac{N^3}{12},$$

which completes the proof. \square

Lemma D.7. *Let \mathbf{Z}_N be the vector of treatment indicators when the treatments are assigned by m -out-of- N SRSWOR where $m(N)/N \rightarrow \lambda \in (0, 1)$ as $N \rightarrow \infty$. Define*

$$S_N = \sum_{j=1}^N \sum_{i=1, i \neq j}^N (1 - Z_{N,i}) Z_{N,j} I_{h,N}(b_{N,j} - b_{N,i})$$

where $I_{h,N}$ is defined in (2.15) of the main paper. Then under [Assumption 1](#) it holds that

$$N^{-3/2} S_N \xrightarrow{P} h\lambda(1-\lambda)\mathcal{I}_b.$$

Proof. We imitate the proof of [Theorem 2.3](#). The first step is to note that

$$N^{-3/2} \mathbb{E} S_N = \frac{m}{N} \left(1 - \frac{m-1}{N-1}\right) N^{-3/2} \sum_{j=1}^N \sum_{i=1, i \neq j}^N I_{h,N}(b_{N,j} - b_{N,i}) \rightarrow h\lambda(1-\lambda)\mathcal{I}_b.$$

It therefore remains to show that $N^{-3} \text{Var}(S_N) \rightarrow 0$. For brevity, let us abuse the notation $I_{h,N}(b_{N,j} - b_{N,i})$ and write $I_{h,N}(i, j)$ instead. We have

$$\text{Var}(S_N) = \sum_{i=1}^N \sum_{j=1, j \neq i}^N \sum_{k=1}^N \sum_{l=1, l \neq k}^N \text{Cov}((1 - Z_{N,i}) Z_{N,j}, (1 - Z_{N,k}) Z_{N,l}) I_{h,N}(i, j) I_{h,N}(k, l).$$

Note, $2 \leq |\{i, j, k, l\}| \leq 4$. Consider the following cases.

- (a) $|\{i, j, k, l\}| = 2$, i.e., $(i, j) = (k, l)$. Note that $\text{Var}((1 - Z_{N,i}) Z_{N,j}) = p_N(1 - p_N)$ where $p_N = P(Z_{N,i} = 0, Z_{N,j} = 1) = \frac{m}{N} \frac{N-m}{N-1} \sim \lambda(1-\lambda)$. Hence the contribution of these terms

in $\text{Var}(S_N)$ is given by

$$\sum_{i=1}^N \sum_{j=1, j \neq i}^N p_N(1-p_N) I_{h,N}^2(i, j) \lesssim \lambda(1-\lambda) \sum_{i=1}^N \sum_{j=1, j \neq i}^N I_{h,N}(i, j) \lesssim N^{3/2}.$$

(b) $|\{i, j, k, l\}| = 4$, i.e., all 4 indices are distinct. Note that in this case

$$\begin{aligned} & \text{Cov}((1 - Z_{N,i})Z_{N,j}, (1 - Z_{N,k})Z_{N,l}) \\ &= P(Z_{N,i} = 0, Z_{N,j} = 1, Z_{N,k} = 0, Z_{N,l} = 1) - p_N^2 \\ &= \frac{\binom{N-4}{m-2}}{\binom{N}{m}} - \left(\frac{m}{N} \frac{N-m}{N-1} \right)^2 \\ &= \frac{m}{N} \frac{N-m}{N-1} \left(\frac{(m-1)(N-m-1)}{(N-2)(N-3)} - \frac{m}{N} \frac{N-m}{N-1} \right) \\ &= \frac{m}{N} \frac{N-m}{N-1} \frac{(4mN^2 - 4m^2N - N^3 + 6m^2 - 6mN + 2N^2 - N)}{N(N-1)(N-2)(N-3)} \\ &\sim -\lambda(1-\lambda)(1-2\lambda)^2 N^{-1}. \end{aligned}$$

Hence the contribution u_N of these terms in $\text{Var}(S_N)$ satisfies the following.

$$u_N \lesssim N^{-1} \sum_{i=1}^N \sum_{j=1, j \neq i}^N \sum_{k=1}^N \sum_{l=1, l \neq k}^N I_{h,N}(i, j) I_{h,N}(k, l) \lesssim N^{-1+3/2+3/2} = N^2.$$

(c) $|\{i, j, k, l\}| = 3$. Here we have 4 sub-cases:

sub-case	$\mathbb{E}(1 - Z_{N,i})Z_{N,j}(1 - Z_{N,k})Z_{N,l}$	$\text{Cov}((1 - Z_{N,i})Z_{N,j}, (1 - Z_{N,k})Z_{N,l})$
$i = k, j \neq l$	$P(Z_{N,i} = 0, Z_{N,j} = 1, Z_{N,l} = 1)$	$\frac{m(m-1)(N-m)}{N(N-1)(N-2)} - p_N^2 \sim \lambda^3(1-\lambda)$
$i \neq k, j = l$	$P(Z_{N,i} = 0, Z_{N,j} = 1, Z_{N,k} = 0)$	$\frac{m(N-m)(N-m-1)}{N(N-1)(N-2)} - p_N^2 \sim \lambda(1-\lambda)^3$
$i = l, j \neq k$	0	$-p_N^2 \sim -\lambda^2(1-\lambda)^2$
$i \neq l, j = k$	0	$-p_N^2 \sim -\lambda^2(1-\lambda)^2$

Hence, if v_N be the contribution these terms in $\text{Var}(S_N)$, then

$$\begin{aligned} v_N &\lesssim \sum_{i,j,l \text{ distinct}} I_{h,N}(i, j) I_{h,N}(i, l) + \sum_{i,j,k \text{ distinct}} I_{h,N}(i, j) I_{h,N}(k, j) \\ &\quad + \sum_{i,j,k \text{ distinct}} I_{h,N}(i, j) I_{h,N}(k, i) + \sum_{i,j,l \text{ distinct}} I_{h,N}(i, j) I_{h,N}(j, l) \\ &\leq 4N \sum_{i,j \neq i} I_{h,N}(i, j) \lesssim N^{1+3/2} = N^{5/2}. \end{aligned}$$

Combining the three cases, we can say that $N^{-3}\text{Var}(S_N) \lesssim N^{-3}(N^{3/2} + N^2 + N^{5/2}) = o(1)$. \square

Lemma D.8. Suppose that $\mathbf{b}_N = \mathbf{X}_N \boldsymbol{\beta}_N + \boldsymbol{\varepsilon}_N$, where $\varepsilon_{N,1}, \dots, \varepsilon_{N,N}$ are i.i.d. from $\mathcal{N}(0, \sigma^2)$. Define, for any fixed h ,

$$J_N := N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N I_{h,N}(\tilde{b}_{N,j} - \tilde{b}_{N,i}), \quad (\text{D.5})$$

where $\tilde{b}_{N,j}$ is defined in (3.6), and $I_{h,N}$ is defined in (2.15) of the main paper. Then $\mathbb{E}J_N \rightarrow h(2\sqrt{\pi}\sigma)^{-1}$, and $\text{Var}(J_N) \rightarrow 0$, implying that [Assumption 2](#) holds in probability, with $\mathcal{J}_b = (2\sqrt{\pi}\sigma)^{-1}$.

Proof. We do the proof for $h > 0$, the other case will be similar. For simplicity in notation, we omit the index N in $\tilde{b}_{N,j}, Z_{N,j}, \mathbf{p}_{N,j}$, etc. throughout this proof. Observe that under the linear model, we have $\tilde{\mathbf{b}} = (\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{b} = (\mathbf{I} - \mathbf{P}_\mathbf{X})\boldsymbol{\varepsilon}$. Hence for any pair (i, j) of distinct indices, $\tilde{b}_j - \tilde{b}_i \sim \mathcal{N}(0, \sigma_{ij}^2)$, where $\sigma_{ij}^2 := \sigma^2(2 - \|\mathbf{p}_j - \mathbf{p}_i\|^2)$ (this follows from the fact that $\mathbf{P}_\mathbf{X}$ is a projection matrix). Thus

$$\mathbb{E}J_N = N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N P\left(0 \leq \tilde{b}_j - \tilde{b}_i < hN^{-1/2}\right) = N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left(\Phi(\sigma_{ij}^{-1}hN^{-1/2}) - \Phi(0)\right).$$

For any $\delta > 0$ and $N \in \mathbb{N}$, define

$$S_{N,\delta} := \{(i, j) : 1 \leq i, j \leq N, \|\mathbf{p}_j - \mathbf{p}_i\| \leq \delta\}. \quad (\text{D.6})$$

Invoking [Lemma E.1](#), we get

$$|S_{N,\delta}^c| = |\{(i, j) : 1 \leq i, j \leq N, \|\mathbf{p}_j - \mathbf{p}_i\| > \delta\}| \leq \delta^{-2} \sum_{j=1}^N \sum_{i=1}^N \|\mathbf{p}_j - \mathbf{p}_i\|^2 = O(N).$$

Consequently,

$$N^{-3/2} \sum_{(i,j) \in S_{N,\delta}^c} \left(\Phi(\sigma_{i,j}^{-1}hN^{-1/2}) - \Phi(0)\right) = O(N^{-1/2}).$$

For $(i, j) \in S_{N,\delta}$ we apply the Taylor theorem to conclude that there exists $\psi_{i,j} \in (0, \sigma_{i,j}^{-1}hN^{-1/2})$ such that

$$\Phi(\sigma_{i,j}^{-1}hN^{-1/2}) - \Phi(0) = \sigma_{i,j}^{-1}hN^{-1/2}\phi(0) + \sigma_{i,j}^{-2}h^2N^{-1}\phi'(\psi_{i,j}).$$

Since ϕ' is bounded on \mathbb{R} , we can say that

$$\Delta_N := \left| \mathbb{E}J_N - N^{-3/2} \sum_{(i,j) \in S_{N,\delta}} \sigma_{i,j}^{-1}hN^{-1/2}\phi(0) \right| \lesssim N^{-1/2} + N^{-3/2} \sum_{(i,j) \in S_{N,\delta}} \sigma_{i,j}^{-2}h^2N^{-1}$$

$$\begin{aligned}
&\lesssim N^{-1/2} + N^{-5/2} \sum_{(i,j) \in S_{N,\delta}} (2 - \|\mathbf{p}_j - \mathbf{p}_i\|^2)^{-1} \\
&\lesssim N^{-1/2} + N^{-5/2} \sum_{(i,j) \in S_{N,\delta}} (2 - \delta^2)^{-1} \lesssim N^{-1/2}.
\end{aligned}$$

Hence $\Delta_N \rightarrow 0$ as $N \rightarrow \infty$, and thus

$$\limsup_{N \rightarrow \infty} \mathbb{E}J_N = \limsup_{N \rightarrow \infty} N^{-3/2} \sum_{(i,j) \in S_{N,\delta}} \sigma_{i,j}^{-1} h N^{-1/2} \phi(0) \leq h \sigma^{-1} \phi(0) (2 - \delta^2)^{-1/2}.$$

Since the above holds for every $\delta > 0$, we can say that $\limsup_{N \rightarrow \infty} \mathbb{E}J_N \leq \frac{h}{2\sqrt{\pi}\sigma}$. On the other hand, the fact that $\Delta_N \rightarrow 0$ also yields the following:

$$\begin{aligned}
\liminf_{N \rightarrow \infty} \mathbb{E}J_N &= \liminf_{N \rightarrow \infty} N^{-3/2} \sum_{(i,j) \in S_{N,\delta}} (2 - \|\mathbf{p}_j - \mathbf{p}_i\|)^{-1/2} h \sigma^{-1} N^{-1/2} \phi(0) \\
&\geq \liminf_{N \rightarrow \infty} N^{-2} |S_{N,\delta}| \cdot h \sigma^{-1} \phi(0) \cdot 2^{-1/2} = h \sigma^{-1} \phi(0) \cdot 2^{-1/2},
\end{aligned}$$

where in the last step we again used the fact that $|S_{N,\delta}^c| = O(N)$. We thus conclude that

$$\lim_{N \rightarrow \infty} \mathbb{E}J_N = \frac{h}{2\sqrt{\pi}\sigma}.$$

Next we show that $\text{Var}(J_N) \rightarrow 0$ as $N \rightarrow \infty$. First,

$$\mathbb{E}J_N^2 = N^{-3} \mathbb{E} \sum_{i,j,k,l} \mathbf{1} \left\{ 0 \leq \tilde{b}_j - \tilde{b}_i < hN^{-1/2}, 0 \leq \tilde{b}_l - \tilde{b}_k < hN^{-1/2} \right\}.$$

Now, as in the proof of [Theorem 2.3](#), observe that the contribution of the terms with repeated indices in the above summation is negligible (since [Assumption 2](#) holds and we already have shown that $\mathbb{E}J_N$ converges). To analyze the terms with distinct indices, note that for any 4 distinct indices i, j, k, l ,

$$\begin{pmatrix} \tilde{b}_j - \tilde{b}_i \\ \tilde{b}_l - \tilde{b}_k \end{pmatrix} \sim \mathcal{N}_2 \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \sigma^2 \begin{pmatrix} 2 - \|\mathbf{p}_j - \mathbf{p}_i\|^2 & -(\mathbf{p}_j - \mathbf{p}_i)^\top (\mathbf{p}_l - \mathbf{p}_k) \\ -(\mathbf{p}_j - \mathbf{p}_i)^\top (\mathbf{p}_l - \mathbf{p}_k) & 2 - \|\mathbf{p}_l - \mathbf{p}_k\|^2 \end{pmatrix} \right).$$

Let $\rho_{i,j,k,l} := \text{corr}(\tilde{b}_j - \tilde{b}_i, \tilde{b}_l - \tilde{b}_k)$. Now we again play the trick of splitting the sum into two groups, such that in one group $|\rho_{i,j,k,l}|$ is small, whereas for the other group the number of summands is small. Fix any $\delta \in (0, 1)$, and consider the set $S_{N,\delta}$ defined in [\(D.6\)](#). If (i, j) or (k, l) does not belong to $S_{N,\delta}$, then that brings down the count for such summands. To be precise,

$$N^{-3} \sum_{(i,j) \in S_{N,\delta}^c \text{ or } (k,l) \in S_{N,\delta}^c} P(0 \leq \tilde{b}_j - \tilde{b}_i \leq hN^{-1/2}, 0 \leq \tilde{b}_l - \tilde{b}_k \leq hN^{-1/2})$$

$$\begin{aligned}
&\leq 2N^{-3} \sum_{1 \leq j, i \leq N} \sum_{(k, l) \in S_{N, \delta}^c} P(0 \leq \tilde{b}_j - \tilde{b}_i \leq hN^{-1/2}) \\
&\lesssim N^{-2} \sum_{1 \leq j, i \leq N} P(0 \leq \tilde{b}_j - \tilde{b}_i \leq hN^{-1/2}) = N^{-1/2} \mathbb{E} J_N.
\end{aligned}$$

Since $\mathbb{E} J_N$ converges, the above shows that the contribution from such indices are also negligible. Finally, for $(i, j), (k, l) \in S_{N, \delta}$, we have

$$|\rho_{i,j,k,l}| = \frac{|(\mathbf{p}_j - \mathbf{p}_i)^\top (\mathbf{p}_l - \mathbf{p}_k)|}{\sqrt{2 - \|\mathbf{p}_j - \mathbf{p}_i\|^2} \sqrt{2 - \|\mathbf{p}_l - \mathbf{p}_k\|^2}} \leq \frac{\delta}{2 - \delta^2} \leq \delta,$$

and hence [Lemma E.3](#) tells us that

$$\begin{aligned}
&|P(0 \leq \tilde{b}_j - \tilde{b}_i \leq hN^{-1/2}, 0 \leq \tilde{b}_l - \tilde{b}_k \leq hN^{-1/2}) \\
&\quad - P(0 \leq \tilde{b}_j - \tilde{b}_i \leq hN^{-1/2}) P(0 \leq \tilde{b}_l - \tilde{b}_k \leq hN^{-1/2})| \\
&\leq |\rho_{i,j,k,l}| ((1 - \rho_{i,j,k,l}^2)^{-1/2} + (1 - |\rho_{i,j,k,l}|)^{-2}) h^2 N^{-2} \sigma_{i,j}^{-1} \sigma_{k,l}^{-1} \\
&\leq h^2 N^{-2} \sigma^{-2} \delta ((1 - \delta^2)^{-1/2} + (1 - \delta)^{-2}).
\end{aligned}$$

Thus we have shown that

$$\begin{aligned}
&\left| \mathbb{E} J_N^2 - \left(N^{-3/2} \sum_{(i,j) \in S_{N,\delta}} P(0 \leq \tilde{b}_j - \tilde{b}_i \leq hN^{-1/2}) \right)^2 \right| \\
&\leq h^2 \sigma^{-2} \delta ((1 - \delta^2)^{-1/2} + (1 - \delta)^{-2}) + O(N^{-1/2}).
\end{aligned}$$

Now for every $\delta \in (0, 1)$, we have $|S_{N,\delta}^c| = O(N)$, so it follows that

$$\limsup_{N \rightarrow \infty} \text{Var}(J_N) \leq h^2 \sigma^{-2} \delta ((1 - \delta^2)^{-1/2} + (1 - \delta)^{-2}).$$

Since $\delta \in (0, 1)$ is arbitrary here, letting $\delta \rightarrow 0$ completes the proof. \square

Lemma D.9. *Suppose that [Assumption 2](#) holds. Then for any $h \geq \delta > 0$,*

$$\lim_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\tilde{b}_{N,j} - \tilde{b}_{N,i}| \in [(h - \delta)N^{-1/2}, (h + \delta)N^{-1/2}] \right\} = 4\delta \mathcal{J}_b$$

where \mathcal{J}_b is defined in [Assumption 2](#) of the main paper.

Proof. First we show that for any $h \in \mathbb{R}$,

$$\lim_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \tilde{b}_{N,j} - \tilde{b}_{N,i} = hN^{-1/2} \right\} = 0. \quad (\text{D.7})$$

To show this, fix $h \geq 0$. Note that for any $\delta > 0$,

$$\begin{aligned} 0 &\leq N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \tilde{b}_{N,j} - \tilde{b}_{N,i} = hN^{-1/2} \right\} \leq N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ hN^{-1/2} \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < \frac{h+\delta}{\sqrt{N}} \right\} \\ &\leq N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ 0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < \frac{h+\delta}{\sqrt{N}} \right\} - N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ 0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < hN^{-1/2} \right\}. \end{aligned}$$

As $N \rightarrow \infty$, the above RHS converges to $\delta \mathcal{J}_b$. Then letting $\delta \rightarrow 0$ we finish the proof of (D.7) for $h \geq 0$. The case $h < 0$ is similar.

Next, fix any $h \geq \delta > 0$. We write

$$\begin{aligned} &N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ \left| \tilde{b}_{N,j} - \tilde{b}_{N,i} \right| \in \left[\frac{h-\delta}{\sqrt{N}}, \frac{h+\delta}{\sqrt{N}} \right] \right\} \\ &= N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ \left| \tilde{b}_{N,j} - \tilde{b}_{N,i} \right| \leq \frac{h+\delta}{\sqrt{N}} \right\} - \mathbf{1} \left\{ \left| \tilde{b}_{N,j} - \tilde{b}_{N,i} \right| \leq \frac{h-\delta}{\sqrt{N}} \right\} \right) \\ &= N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ 0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < \frac{h+\delta}{\sqrt{N}} \right\} + \mathbf{1} \left\{ -\frac{h+\delta}{\sqrt{N}} \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < 0 \right\} \right) \\ &\quad - N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ 0 \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < \frac{h-\delta}{\sqrt{N}} \right\} + \mathbf{1} \left\{ -\frac{h-\delta}{\sqrt{N}} \leq \tilde{b}_{N,j} - \tilde{b}_{N,i} < 0 \right\} \right) \\ &\quad + N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \left(\mathbf{1} \left\{ \tilde{b}_{N,j} - \tilde{b}_{N,i} = \frac{h+\delta}{\sqrt{N}} \right\} - \mathbf{1} \left\{ \tilde{b}_{N,j} - \tilde{b}_{N,i} = \frac{h-\delta}{\sqrt{N}} \right\} \right). \end{aligned}$$

Appealing to [Assumption 2](#) and (D.7), we conclude that as $N \rightarrow \infty$, the above display converges to $2(h+\delta)\mathcal{J}_b - 2(h-\delta)\mathcal{J}_b = 4\delta\mathcal{J}_b$. \square

Lemma D.10. *Suppose that [Assumption 1](#) holds. Then for any $h \geq \delta > 0$,*

$$\lim_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |b_{N,j} - b_{N,i}| \in \left[\frac{h-\delta}{\sqrt{N}}, \frac{h+\delta}{\sqrt{N}} \right] \right\} = 4\delta\mathcal{I}_C$$

where \mathcal{I}_C is defined in [Assumption 1](#) of the main paper.

Proof. The proof is essentially same as the proof of [Lemma D.9](#), hence omitted. \square

Lemma D.11. *Define D_N as in (C.8), and let*

$$\theta_{N,i,j} := \mathbb{E} \left[|\xi_{N,i,j} - \tilde{\xi}_{N,i,j}| \mid Z_{N,j} = 1 \right]. \quad (\text{D.8})$$

Then the following holds:

$$\sqrt{\mathbb{E}(N^{-3/2}D_N^2)} \leq N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \sqrt{\theta_{N,i,j}}.$$

Proof. Recall the notations \mathbf{I}_N and $\tilde{\mathbf{I}}_N$ from (C.6) and (C.7), respectively. We use the simple result $\sqrt{\mathbb{E}(\sum_{j=1}^N V_j)^2} \leq \sum_{j=1}^N \sqrt{\mathbb{E}(V_j^2)}$ to derive the following.

$$\begin{aligned} \sqrt{\mathbb{E}(D_N)^2} &= \sqrt{\mathbb{E}(\mathbf{I}_N - \tilde{\mathbf{I}}_N)^2} \\ &= \left[\mathbb{E} \left(\sum_{j=1}^N Z_{N,j} \sum_{i=1}^N (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \right]^{1/2} \\ &\leq \sum_{j=1}^N \left[\mathbb{E} \left(Z_{N,j} \sum_{i=1}^N (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \right]^{1/2} \\ &= \sum_{j=1}^N \left[\mathbb{E} \left(Z_{N,j} \mathbb{E} \left(\left(\sum_{i=1}^N (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \mid Z_{N,j} \right) \right) \right]^{1/2} \\ &= \sum_{j=1}^N \left[\mathbb{E} \left(Z_{N,j} \mathbb{E} \left(\left(\sum_{i=1}^N (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \mid Z_{N,j} = 1 \right) \right) \right]^{1/2} \\ &= \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \left[\mathbb{E} \left(\left(\sum_{i=1}^N (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &\leq \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \sum_{i=1}^N \left[\mathbb{E} \left((\xi_{N,i,j} - \tilde{\xi}_{N,i,j})^2 \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &= \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \sum_{i=1}^N \left[\mathbb{E} \left(|\xi_{N,i,j} - \tilde{\xi}_{N,i,j}| \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &\leq \sum_{j=1}^N \sum_{i=1}^N \sqrt{\theta_{N,i,j}}. \end{aligned}$$

This completes the proof. □

Lemma D.12. Define Q_N as in (C.8), and let

$$\gamma_{N,i,j} := \mathbb{E} \left[|\xi_{N,i,j} - \tilde{\xi}_{N,i,j}| \mid Z_{N,i} = 1, Z_{N,j} = 1 \right] \quad (\text{D.9})$$

Then it holds that,

$$\sqrt{\mathbb{E}(N^{-3/2}Q_N)^2} \leq N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \sqrt{\gamma_{N,i,j}}.$$

Proof. This proof mimics the proof of [Lemma D.11](#). Recall the notations \mathbf{II}_N and $\tilde{\mathbf{II}}_N$ from [\(C.6\)](#) and [\(C.7\)](#), respectively. Note that,

$$\begin{aligned} \sqrt{\mathbb{E}(Q_N^2)} &= \sqrt{\mathbb{E}(\mathbf{II}_N - \tilde{\mathbf{II}}_N)^2} \\ &= \left[\mathbb{E} \left(\sum_{j=1}^N Z_{N,j} \sum_{i=1}^N Z_{N,i} (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \right]^{1/2} \\ &\leq \sum_{j=1}^N \left[\mathbb{E} \left(Z_{N,j} \sum_{i=1}^N Z_{N,i} (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \right]^{1/2} \\ &= \sum_{j=1}^N \left[\mathbb{E} \left(Z_{N,j} \mathbb{E} \left(\left(\sum_{i=1}^N Z_{N,i} (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \mid Z_{N,j} \right) \right) \right]^{1/2} \\ &= \sum_{j=1}^N \left[\mathbb{E} \left(Z_{N,j} \mathbb{E} \left(\left(\sum_{i=1}^N Z_{N,i} (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \mid Z_{N,j} = 1 \right) \right) \right]^{1/2} \\ &= \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \left[\mathbb{E} \left(\left(\sum_{i=1}^N Z_{N,i} (\xi_{N,i,j} - \tilde{\xi}_{N,i,j}) \right)^2 \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &\leq \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \sum_{i=1}^N \left[\mathbb{E} \left(Z_{N,i} (\xi_{N,i,j} - \tilde{\xi}_{N,i,j})^2 \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &= \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \sum_{i=1}^N \left[\mathbb{E} \left(Z_{N,i} \mathbb{E} \left((\xi_{N,i,j} - \tilde{\xi}_{N,i,j})^2 \mid Z_{N,i}, Z_{N,j} = 1 \right) \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &= \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \sum_{i=1}^N \left[\mathbb{E} \left(Z_{N,i} \mathbb{E} \left((\xi_{N,i,j} - \tilde{\xi}_{N,i,j})^2 \mid Z_{N,i} = 1, Z_{N,j} = 1 \right) \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &= \sum_{j=1}^N \sqrt{\mathbb{E}(Z_{N,j})} \sum_{i=1}^N \sqrt{\mathbb{E}(Z_{N,i} \mid Z_{N,j} = 1)} \left[\mathbb{E} \left(|\xi_{N,i,j} - \tilde{\xi}_{N,i,j}| \mid Z_{N,i} = 1, Z_{N,j} = 1 \right) \right]^{1/2} \\ &\leq \sum_{j=1}^N \sum_{i=1}^N \left[\mathbb{E} \left(|\xi_{N,i,j} - \tilde{\xi}_{N,i,j}| \mid Z_{N,i} = 1, Z_{N,j} = 1 \right) \right]^{1/2} \\ &= \sum_{j=1}^N \sum_{i=1}^N \sqrt{\gamma_{N,i,j}}. \end{aligned}$$

This completes the proof. \square

Lemma D.13. Define D_N and Q_N as in (C.8). It holds under [Assumption 2](#) that $N^{-3/2}D_N = o_P(1)$ and $N^{-3/2}Q_N = o_P(1)$ as $N \rightarrow \infty$.

Proof. We first focus on D_N . In view of [Lemma D.11](#), it suffices to show that

$$\lim_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \sqrt{\theta_{N,i,j}} = 0,$$

where $\theta_{N,i,j}$ is defined in (D.8). Fix h for the moment, and introduce the notation

$$r_N(i, j) := \tilde{b}_{N,j} - \tilde{b}_{N,i} - hN^{-1/2}, \quad 1 \leq i, j \leq N.$$

Note,

$$\begin{aligned} \xi_{N,i,j} - \tilde{\xi}_{N,i,j} &= \mathbf{1}\{hN^{-1/2}(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \leq r_N(i, j)\} - \mathbf{1}\{0 \leq r_N(i, j)\} \\ &= \mathbf{1}\{hN^{-1/2}(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \leq r_N(i, j) < 0\} \\ &\quad - \mathbf{1}\{hN^{-1/2}(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N > r_N(i, j) \geq 0\} \\ &= \mathbf{1}\{hN^{-1/2}|(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N| \geq |r_N(i, j)|\} (\mathbf{1}\{r_N(i, j) < 0\} - \mathbf{1}\{r_N(i, j) \geq 0\}). \end{aligned}$$

Now applying the Markov inequality, we get

$$\begin{aligned} \sqrt{\theta_{N,i,j}} &= \mathbb{E}^{1/2} \left[|\xi_{N,i,j} - \tilde{\xi}_{N,i,j}| \mid Z_{N,j} = 1 \right] \\ &= \left[P \left(hN^{-1/2} |(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N| \geq |r_N(i, j)| \mid Z_{N,j} = 1 \right) \right]^{1/2} \\ &\leq h^2 N^{-1} r_N(i, j)^{-2} \mathbb{E}^{1/2} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^4 \mid Z_{N,j} = 1 \right]. \end{aligned}$$

To upper bound the above RHS, we use [Lemma E.1](#). The key idea is to use this bound only for those i, j for which $r_N(i, j)$ is at least as large as $\delta N^{-1/2}$. Towards that, fix $\delta > 0$ and define

$$\mathcal{S}_{N,\delta} = \{(i, j) : |r_N(i, j)| \leq \delta N^{-1/2}, 1 \leq i, j \leq N\}.$$

Then

$$(i, j) \notin \mathcal{S}_{N,\delta} \implies \sqrt{\theta_{N,i,j}} \leq a^2 \delta^{-2} \mathbb{E}^{1/2} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^4 \mid Z_{N,j} = 1 \right].$$

Therefore

$$N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \sqrt{\theta_{N,i,j}}$$

$$\begin{aligned}
&\leq N^{-3/2} \sum_{(i,j) \in \mathcal{S}_{N,\delta}} \sqrt{\theta_{N,i,j}} + N^{-3/2} \sum_{(i,j) \notin \mathcal{S}_{N,\delta}}^N \sqrt{\theta_{N,i,j}} \\
&\leq N^{-3/2} \sum_{(i,j) \in \mathcal{S}_{N,\delta}} 1 + N^{-3/2} \sum_{(i,j) \notin \mathcal{S}_{N,\delta}}^N a^2 \delta^{-2} \mathbb{E}^{1/2} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^4 \mid Z_{N,j} = 1 \right] \\
&\leq N^{-3/2} |\mathcal{S}_{N,\delta}| + a^2 \delta^{-2} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \mathbb{E}^{1/2} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^4 \mid Z_{N,j} = 1 \right] \\
&= N^{-3/2} |\mathcal{S}_{N,\delta}| + a^2 \delta^{-2} (\text{rank}(\mathbf{X}_N) - 1) \cdot O(N^{-1/2}).
\end{aligned}$$

In the last step we used [Lemma E.1](#). Letting $N \rightarrow \infty$ and then $\delta \downarrow 0$, it follows that

$$\limsup_{N \rightarrow \infty} N^{-3/2} \sum_{j=1}^N \sum_{i=1}^N \sqrt{\theta_{N,i,j}} \leq \lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} N^{-3/2} |\mathcal{S}_{N,\delta}|.$$

Now

$$\begin{aligned}
(i, j) \in \mathcal{S}_{N,\delta} &\implies \frac{\delta}{\sqrt{N}} \geq |r_N(i, j)| \geq \left| (b_{N,j} - \mathbf{p}_{N,j}^\top \mathbf{b}_N) - (b_{N,i} - \mathbf{p}_{N,i}^\top \mathbf{b}_N) \right| - \frac{|a|}{\sqrt{N}} \\
&\implies |(b_{N,j} - \mathbf{p}_{N,j}^\top \mathbf{b}_N) - (b_{N,i} - \mathbf{p}_{N,i}^\top \mathbf{b}_N)| \in \left[\frac{|a| - \delta}{\sqrt{N}}, \frac{|a| + \delta}{\sqrt{N}} \right].
\end{aligned}$$

We now invoke [Lemma D.9](#) to arrive at

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} N^{-3/2} |\mathcal{S}_{N,\delta}| = 0,$$

which finishes the proof for D_N . The proof for Q_N can be done in the same manner, using [Lemma D.12](#), Markov inequality, and [Lemma E.1](#). \square

Lemma D.14. *Suppose that [Assumption 3](#) holds. Then for any $h \geq \delta > 0$,*

$$\lim_{N \rightarrow \infty} N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |b_{N,j} - b_{N,i}| \in \left[\frac{h - \delta}{N^\nu}, \frac{h + \delta}{N^\nu} \right] \right\} = 4\delta \mathcal{I}_b$$

where \mathcal{I}_b is defined in [Assumption 1](#) of the main paper.

Proof. The proof is analogous to those of [Lemmas D.9](#) and [D.10](#), and hence omitted. \square

Lemma D.15. *Suppose that [Assumption 4](#) holds. Then for any $h \geq \delta > 0$,*

$$\lim_{N \rightarrow \infty} N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N \mathbf{1} \left\{ |\tilde{b}_{N,j} - \tilde{b}_{N,i}| \in \left[\frac{h - \delta}{N^\nu}, \frac{h + \delta}{N^\nu} \right] \right\} = 4\delta \mathcal{J}_b$$

where \mathcal{J}_b is defined in [Assumption 2](#) of the main paper.

Proof. The proof is analogous to those of [Lemmas D.9](#) and [D.10](#), and hence omitted. \square

Lemma D.16. *For each $N \geq 1$, let $b_{N,1}, \dots, b_{N,N}$ be i.i.d. from an unknown distribution F_b with density $f_b(\cdot)$, and $I_{h,N,\nu}$ be defined in [\(A.2\)](#) of the main paper, then for any $0 < \nu \leq 1/2$,*

$$N^{-(2-\nu)} \sum_{j=1}^N \sum_{i=1}^N I_{h,N,\nu}(b_{N,j} - b_{N,i}) \xrightarrow{P} h \int_{\mathbb{R}} f_b^2(x) dx.$$

Proof. We proved the above for $\nu = 1/2$ in [Lemma D.4](#), so assume now that $\nu \in (0, 1/2)$. Define

$$S_N := S_N(b_{N,1}, \dots, b_{N,N}) := \sum_{j=1}^N \sum_{i=1}^N I_{h,N,\nu}(b_{N,j} - b_{N,i}).$$

Observe that for any fixed $i \neq j$, and $h > 0$,

$$\mathbb{E}(I_{h,N,\nu}(b_{N,j} - b_{N,i})) = P(0 \leq b_{N,2} - b_{N,1} < hN^{-\nu}) = g(hN^{-\nu}) - g(0)$$

where

$$g(x) = P(b_{N,2} - b_{N,1} \leq x) = \int_{-\infty}^x \int_{\mathbb{R}} f_b(u+t) f_b(u) du dt, \quad x \in \mathbb{R}.$$

Using the DCT for integrals, we argue that

$$g'(x) = \int_{\mathbb{R}} f_b(u+x) f_b(u) du.$$

Hence

$$\begin{aligned} \lim_{N \rightarrow \infty} N^{-(2-\nu)} \mathbb{E}(S_N) &= \lim_{N \rightarrow \infty} N^{-(2-\nu)} N^2 (g(hN^{-\nu}) - g(0)) \\ &= \lim_{N \rightarrow \infty} h \cdot \frac{g(hN^{-\nu}) - g(0)}{hN^{-\nu}} \\ &= hg'(0) \\ &= h \int_{\mathbb{R}} f_b(u)^2 du. \end{aligned} \tag{D.10}$$

The case $h < 0$ can be handled similarly, and the case $h = 0$ is straight-forward. Next we bound $\mathbb{E}(S_N - \mathbb{E}S_N)^2$ using the Efron-Stein inequality ([Efron and Stein, 1981](#)). For each $1 \leq k \leq N$, let $b'_{N,k}$ be an i.i.d. copy of $b_{N,k}$, independent of everything else, and define

$$S_N^{(k)} = S_N(b_{N,1}, \dots, b_{N,k-1}, b'_{N,k}, b_{N,k+1}, \dots, b_{N,N}), \quad 1 \leq k \leq N.$$

Note,

$$\begin{aligned} S_N - S_N^{(k)} &= \sum_{j=1}^N (I_{h,N,\nu}(b_{N,j} - b_{N,k}) - I_{h,N,\nu}(b_{N,j} - b'_{N,k})) \\ &\quad + \sum_{j=1}^N (I_{h,N,\nu}(b_{N,k} - b_{N,j}) - I_{h,N,\nu}(b'_{N,k} - b_{N,j})). \end{aligned}$$

Hence $|S_N - S_N^{(k)}| \leq 4N$ almost surely. The Efron-Stein inequality tells us that

$$\begin{aligned} \mathbb{E}(N^{-(2-\nu)}(S_N - \mathbb{E}S_N))^2 &\leq N^{-(4-2\nu)} \mathbb{E} \left[\sum_{k=1}^N \mathbb{E} \left[(S_N - S_N^{(k)})^2 \mid b_{N,1}, \dots, b_{N,N} \right] \right] \\ &\leq N^{-(4-2\nu)} \cdot 16N^3 = 16N^{-(1-2\nu)}. \end{aligned}$$

Since $\nu < 1/2$, we can now invoke (D.10) to get the desired conclusion. \square

E Auxiliary results

Lemma E.1. *As $N \rightarrow \infty$, the following results hold, for $r = 1, 2$.*

$$\sum_{j=1}^N \sum_{i=1}^N \mathbb{E}^{1/r} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^{2r} \right] = (\text{rank}(\mathbf{X}_N) - 1) \cdot O(N), \quad (\text{E.1})$$

$$\sum_{j=1}^N \sum_{i=1}^N \mathbb{E}^{1/r} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^{2r} \mid Z_{N,j} = 1 \right] = (\text{rank}(\mathbf{X}_N) - 1) \cdot O(N), \quad (\text{E.2})$$

$$\sum_{j=1}^N \sum_{i=1}^N \mathbb{E}^{1/r} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^{2r} \mid Z_{N,i} = 1, Z_{N,j} = 1 \right] = (\text{rank}(\mathbf{X}_N) - 1) \cdot O(N). \quad (\text{E.3})$$

Proof. Using the fact that $\mathbf{P}_{\mathbf{X}_N}$ is idempotent, we derive the following identity.

$$\begin{aligned} \sum_{j=1}^N \sum_{i=1}^N \|\mathbf{p}_{N,i} - \mathbf{p}_{N,j}\|^2 &= \sum_{j=1}^N \sum_{i=1}^N (\mathbf{P}_{\mathbf{X}_N}(i, i) + \mathbf{P}_{\mathbf{X}_N}(j, j) - 2\mathbf{P}_{\mathbf{X}_N}(i, j)) \\ &= 2N \sum_{i=1}^N \mathbf{P}_{\mathbf{X}_N}(i, i) - 2 \sum_{i=1}^N \mathbf{p}_{N,i}^\top \mathbf{1} \\ &= 2N (\text{Tr}(\mathbf{P}_{\mathbf{X}_N}) - 1) = 2N (\text{rank}(\mathbf{X}_N) - 1). \end{aligned} \quad (\text{E.4})$$

Now $\mathbf{1}$ belongs to the column space of \mathbf{X}_N , so $\mathbb{E}[(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N] = 0$ for each i, j . Also

note that $\text{Var}(\mathbf{Z}_N)$ is of the form $\alpha \mathbf{I} + \beta \mathbf{1}\mathbf{1}^\top$, where $\alpha = \frac{m}{N}(1 - \frac{m}{N})$. Consequently,

$$\text{Var}[(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N] = (\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top (\alpha \mathbf{I} + \beta \mathbf{1}\mathbf{1}^\top) (\mathbf{p}_{N,i} - \mathbf{p}_{N,j}) = \frac{m}{N} \left(1 - \frac{m}{N}\right) \|\mathbf{p}_{N,i} - \mathbf{p}_{N,j}\|^2.$$

Hence

$$\sum_{j=1}^N \sum_{i=1}^N \mathbb{E} \left[((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N)^2 \right] \leq \sum_{j=1}^N \sum_{i=1}^N \|\mathbf{p}_{N,i} - \mathbf{p}_{N,j}\|^2.$$

Thus, (E.1) follows for $r = 1$.

To prove it for $r = 2$, fix i and j for the moment and write $(\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N = \sum_{k=1}^N v_k Z_k$. Then

$$\begin{aligned} \mathbb{E} \left(\sum_{k=1}^N v_k Z_k \right)^4 &= \sum_{k_1=1}^N v_{k_1}^4 \mathbb{E} Z_1 + \sum_{k_1, k_2 \text{ distinct}} (3v_{k_1}^2 v_{k_2}^2 + 4v_{k_1}^3 v_{k_2}) \mathbb{E} Z_1 Z_2 \\ &\quad + \sum_{k_1, k_2, k_3 \text{ distinct}} 6v_{k_1}^2 v_{k_2} v_{k_3} \mathbb{E} Z_1 Z_2 Z_3 + \sum_{k_1, k_2, k_3, k_4 \text{ distinct}} v_{k_1} v_{k_2} v_{k_3} v_{k_4} \mathbb{E} Z_1 Z_2 Z_3 Z_4. \end{aligned}$$

Now $\sum_{k=1}^N v_k = (\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{1} = 0$, so we obtain

$$\begin{aligned} \sum_{k_1, k_2, k_3, k_4 \text{ distinct}} v_{k_1} v_{k_2} v_{k_3} v_{k_4} &= \sum_{k_1, k_2, k_3 \text{ distinct}} v_{k_1} v_{k_2} v_{k_3} (-v_{k_1} - v_{k_2} - v_{k_3}) = \sum_{k_1, k_2, k_3 \text{ distinct}} -3v_{k_1}^2 v_{k_2} v_{k_3}, \\ \sum_{k_1, k_2, k_3 \text{ distinct}} v_{k_1}^2 v_{k_2} v_{k_3} &= \sum_{k_1, k_2 \text{ distinct}} v_{k_1}^2 v_{k_2} (-v_{k_1} - v_{k_2}) = \sum_{k_1, k_2 \text{ distinct}} -(v_{k_1}^3 v_{k_2} + v_{k_1}^2 v_{k_2}^2), \\ \sum_{k_1, k_2 \text{ distinct}} v_{k_1}^3 v_{k_2} &= \sum_{k_1=1}^N v_{k_1}^3 (-v_{k_1}) = -\sum_{k_1=1}^N v_{k_1}^4. \end{aligned}$$

Since $\mathbb{E} |Z_1 \cdots Z_v| \leq 1$ for $1 \leq v \leq 4$, it follows from the above identities that

$$\begin{aligned} \mathbb{E}^{1/2} \left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^4 &\leq \sqrt{D'} \left(\sum_{k=1}^N v_k^4 + \sum_{k \neq l} v_k^2 v_l^2 \right)^{1/2} \\ &\leq \sqrt{2D'} \sum_{k=1}^N v_k^2 = \sqrt{2D'} \|\mathbf{p}_{N,i} - \mathbf{p}_{N,j}\|^2 \end{aligned}$$

for some constant $D' > 0$ (which is free of i, j and N). The rest follows again from (E.4).

Next, we deal with the identities where we condition upon $Z_{N,j} = 1$. Observe that

$$\mathbb{E} \left[((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N)^2 \mid Z_{N,j} = 1 \right] = (\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \text{Var}(\mathbf{Z}_N \mid Z_{N,j} = 1) (\mathbf{p}_{N,i} - \mathbf{p}_{N,j}).$$

For any $k \neq l$, $\text{Var}(Z_{N,k} \mid Z_{N,j} = 1) = (\alpha_1 - \alpha_1^2)\mathbf{1}\{k \neq j\}$, and $\text{Cov}(Z_{N,k}, Z_{N,l} \mid Z_{N,j} = 1) = (\alpha_2 - \alpha_1^2)\mathbf{1}\{k \neq j\}$, where $\alpha_1 = \frac{m-1}{N-1}$, $\alpha_2 = \frac{m-1}{N-1} \cdot \frac{m-2}{N-2}$. We thus obtain

$$\mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} = 1 \right] = \left[\alpha_1(1 - \alpha_1) \|\mathbf{p}_{N,i} - \mathbf{p}_{N,j}\|^2 - (\alpha_1 - 2\alpha_2 + \alpha_1^2)(\mathbf{P}_{\mathbf{X}_N}(i, j) - \mathbf{P}_{\mathbf{X}_N}(j, j))^2 \right].$$

But $(\mathbf{P}_{\mathbf{X}_N}(i, j) - \mathbf{P}_{\mathbf{X}_N}(j, j))^2 \leq \|\mathbf{p}_{N,i} - \mathbf{p}_{N,j}\|^2$. So it follows that

$$\mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} = 1 \right] \leq D \|\mathbf{p}_{N,i} - \mathbf{p}_{N,j}\|^2$$

for some $D > 0$ which is free of i, j and N . Invoking (E.4) again, we finish the proof of (E.2) for $r = 1$. To prove (E.2) for $r = 2$, note the following.

$$\begin{aligned} & \sqrt{\mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^4 \mid Z_{N,j} = 1 \right]} \\ & \leq \sqrt{\text{Var} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} = 1 \right]} + \mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} = 1 \right] \\ & \leq \sqrt{\frac{N}{m}} \sqrt{\mathbb{E} \left[\text{Var} \left(\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} \right) \right]} + \mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} = 1 \right] \\ & \leq \sqrt{\frac{N}{m}} \sqrt{\text{Var} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \right]} + \mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} = 1 \right] \\ & \leq \sqrt{\frac{N}{m}} \sqrt{\mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^4 \right]} + \mathbb{E} \left[\left((\mathbf{p}_{N,i} - \mathbf{p}_{N,j})^\top \mathbf{Z}_N \right)^2 \mid Z_{N,j} = 1 \right]. \end{aligned}$$

Since $m/N \rightarrow \lambda \in (0, 1)$, the desired conclusion follows from (E.1) (with $r = 2$) and (E.2) (with $r = 1$). The proof of (E.3) is completely analogous to the proofs of (E.1) and (E.2), hence omitted. \square

Lemma E.2. For $x \neq 0$ it holds that $|\Phi(x/\sigma) - \mathbf{1}\{x \geq 0\}| \leq \sigma |x|^{-1} \exp(-x^2/2\sigma^2)$, where Φ is the standard Normal CDF.

Proof. Let ϕ be the density of standard Normal. For $x > 0$, we have

$$|\Phi(x/\sigma) - \mathbf{1}\{x \geq 0\}| = 1 - \Phi(x/\sigma) \leq \sigma \cdot x^{-1} \phi(x/\sigma),$$

using a standard inequality. For $x < 0$, we can write

$$|\Phi(x/\sigma) - \mathbf{1}\{x \geq 0\}| = \Phi(x/\sigma) = 1 - \Phi(-x/\sigma) \leq \sigma \cdot (-x)^{-1} \phi(-x/\sigma).$$

Hence the result. \square

Lemma E.3. *Suppose that (X, Y) follows the bivariate Normal distribution with $\mathbb{E}X = \mathbb{E}Y = 0$, $\text{Var}(X) = \text{Var}(Y) = 1$, and $\text{corr}(X, Y) = \rho \in (-1, 1)$. Then for any $h_1, h_2 \in (0, 1)$, the following bound holds:*

$$|P(0 \leq X \leq h_1, 0 \leq Y \leq h_2) - P(0 \leq X \leq h_1)P(0 \leq Y \leq h_2)| \leq C_\rho h_1 h_2,$$

where $C_\rho = |\rho| ((1 - \rho^2)^{-1/2} + (1 - |\rho|)^{-2})$.

Proof. Define

$$g_{x,y}(\rho) := \exp \left(-\frac{1}{2(1 - \rho^2)}(x^2 + y^2 - 2\rho xy) \right).$$

Note that for any fixed $x, y \in [0, 1]$,

$$\left| \frac{\partial g_{x,y}(\rho)}{\partial \rho} \right| = g_{x,y}(\rho) \left| \frac{xy}{1 - \rho^2} - \frac{\rho}{1 - \rho^2} \frac{x^2 + y^2 - 2\rho xy}{1 - \rho^2} \right| \leq \frac{1}{(1 - |\rho|)^2}.$$

Hence

$$\begin{aligned} & |P(0 \leq X \leq h_1, 0 \leq Y \leq h_2) - P(0 \leq X \leq h_1)P(0 \leq Y \leq h_2)| \\ &= \left| \int_{[0, h_1] \times [0, h_2]} \left((2\pi\sqrt{1 - \rho^2})^{-1} g_{x,y}(\rho) - (2\pi)^{-1} g_{x,y}(0) \right) dx dy \right| \\ &\leq (2\pi)^{-1} |(1 - \rho^2)^{-1/2} - 1| \int_{[0, h_1] \times [0, h_2]} g_{x,y}(\rho) dx dy \\ &\quad + (2\pi)^{-1} \int_{[0, h_1] \times [0, h_2]} |g_{x,y}(\rho) - g_{x,y}(0)| dx dy \\ &\leq \rho^2 (1 - \rho^2)^{-1/2} h_1 h_2 + \int_{[0, h_1] \times [0, h_2]} |\rho| \int_0^1 \left| \frac{\partial g_{x,y}(t\rho)}{\partial \rho} \right| dt dx dy \\ &\leq |\rho| ((1 - \rho^2)^{-1/2} + (1 - |\rho|)^{-2}) h_1 h_2. \end{aligned}$$

This completes the proof. \square

Proposition E.1 (A generalized Efron-Stein inequality). *Given a sequence of independent real-valued random variables W_1, W_2, \dots, W_n and $F : \mathbb{R}^n \rightarrow \mathbb{R}$ be a measurable function. For each $1 \leq i \leq n$, let W'_i be an independent copy of W_i , independent of the other W_j 's. Define $S := F(W_1, \dots, W_n)$ and $S_i := F(W_1, \dots, W_{i-1}, W'_i, W_{i+1}, \dots, W_n)$. Then for all integers $q \geq 2$, there exists a constant c_q (depending only on q) such that*

$$\mathbb{E} |S - \mathbb{E}S|^q \leq c_q \mathbb{E} \left| \mathbb{E} \left[\sum_{i=1}^n (S - S_i)^2 \mid (W_1, W_2, \dots, W_n) \right] \right|^{q/2}.$$

Proof. See [Boucheron et al. \(2005, Theorem 2\)](#) for a proof. \square

The special case $q = 2$ yields the Efron-Stein inequality (see [Efron and Stein \(1981\)](#)).

F Average ranks

In the main paper, we used *up-ranks* to break ties in the responses (see (2.5) in the main paper for a definition). We discuss in this section the analogous results when ties are broken using average ranks. Suppose that while sorting $\{b_{N,j} : 1 \leq j \leq N\}$ in ascending order, the first t_1 many are equal, then the next t_2 many are equal, and so on. Assume that we get k such blocks of sizes t_1, \dots, t_k , where the t_i 's can equal to 1 as well. Set $s_0 = 0$ and $s_j = \sum_{i=1}^j t_i$. Then for each $1 \leq j \leq k$, define

$$q_N^{\text{avg}}(s_{j-1} + 1) = q_N^{\text{avg}}(s_{j-1} + 2) = \dots = q_N^{\text{avg}}(s_j) := \frac{s_{j-1} + 1 + s_j}{2} = s_{j-1} + \frac{1 + t_j}{2}$$

where the last quantity is nothing but the average of the ranks $\{s_{j-1} + 1, \dots, s_{j-1} + t_j\}$. With the above notion of average ranks, we define the Wilcoxon rank-sum statistic as

$$t_N^{\text{avg}} := \sum_{j=1}^N q_N^{\text{avg}}(j) Z_{N,j}.$$

We make the following assumption on the block-sizes, which essentially tells us that the blocks formed by the ties are not too large.

Assumption 5. *Let t_1, \dots, t_k denote the block sizes, as defined above. Then assume that $\max_{1 \leq i \leq k} t_i = o(N)$, as $N \rightarrow \infty$.*

The following lemma shows that if [Assumption 5](#) holds, then the set of the average ranks do not differ much from $\{1, 2, \dots, N\}$. This implies, in view of [Lemma D.5](#) and [Proposition 2.1](#), that t_N^{avg} has the same asymptotic null distribution as the statistic t_N that we studied in [Section 2](#) of the main paper.

Lemma F.1. *It holds under [Assumption 5](#) that*

$$\sum_{i=1}^N (q_N(i) - \bar{q}_N)^2 = \frac{N(N^2 - 1)}{12} + o(N^3),$$

as $N \rightarrow \infty$, where $\bar{q}_N = N^{-1} \sum_{i=1}^N q_N(i)$.

Proof. Since for each j , $\sum_{i=s_{j-1}+1}^{s_j} (q_N(i) - i) = 0$, and thus $\bar{q}_N = N^{-1} \sum_{i=1}^N i = (N+1)/2$ we

obtain the following.

$$\begin{aligned}
\sum_{i=s_{j-1}+1}^{s_j} ((q_N(i) - \bar{q}_N)^2 - (i - \bar{q}_N)^2) &= \sum_{i=s_{j-1}+1}^{s_j} (q_N(i)^2 - 2\bar{q}_N (q_N(i) - i) - i^2) \\
&= - \sum_{i=s_{j-1}+1}^{s_j} \left(i^2 - \left(\frac{s_{j-1} + 1 + s_j}{2} \right)^2 \right) \\
&= - \sum_{i=s_{j-1}+1}^{s_j} \left(i - \frac{s_{j-1} + 1 + s_j}{2} \right)^2 \\
&= - \frac{t_j (t_j^2 - 1)}{12}.
\end{aligned}$$

Therefore

$$\sum_{i=1}^N (q_N(i) - \bar{q}_N)^2 = \sum_{i=1}^N (i - \bar{q}_N)^2 - \sum_{j=1}^k \frac{t_j (t_j^2 - 1)}{12} = \frac{N(N^2 - 1)}{12} - \sum_{j=1}^k \frac{t_j (t_j^2 - 1)}{12}.$$

Notice that

$$0 \leq \sum_{j=1}^k \frac{t_j (t_j^2 - 1)}{12} \leq \frac{\sum_{i=1}^k t_i}{12} \max_{1 \leq i \leq k} t_i^2 = \frac{N}{12} \max_{1 \leq i \leq k} t_i^2.$$

Hence [Assumption 5](#) completes the proof. \square

Our next lemma extends the last result, by showing that if [Assumption 5](#) is satisfied, then the statistic t_N^{avg} has the same asymptotic distribution as the statistic t_N , for any value of the parameter τ . In particular, this implies that [Theorem 2.1](#) also holds if we replace t_N by t_N^{avg} . Consequently, Rosenbaum's estimator for τ constructed using t_N^{avg} instead of t_N must also satisfy [Theorem 2.2](#).

Lemma F.2. *It holds under [Assumption 5](#) that $N^{-3/2}(t_N^{\text{avg}} - t_N) = o_P(1)$ as $N \rightarrow \infty$. Consequently, t_N^{avg} has the same asymptotic distribution as t_N .*

Proof. Recall that $t_N = \sum_{j=1}^N q_N(j) Z_{N,j}$ where $q_N(j) = \sum_{i=1}^N \mathbf{1}\{b_{N,i} \leq b_{N,j}\}$. Observe that when $s_{j-1} < j \leq s_j$,

$$0 \leq q_N(j) - q_N^{\text{avg}}(j) \leq t_j - \frac{t_j + 1}{2} = \frac{t_j - 1}{2}.$$

Hence $t_N \geq t_N^{\text{avg}}$ a.s., and therefore it suffices to show that the following converges to 0, as $N \rightarrow \infty$.

$$\mathbb{E} N^{-3/2} (t_N - t_N^{\text{avg}}) = \frac{m}{N} N^{-3/2} \sum_{j=1}^N (q_N(j) - q_N^{\text{avg}}(j)) = \frac{m}{N} N^{-3/2} \sum_{i=1}^k \frac{t_i (t_i - 1)}{2}.$$

We split the above sum into two parts, as follows. Fix any $\varepsilon > 0$ and define $\mathcal{S}_\varepsilon = \{1 \leq j \leq k : t_j - 1 \geq \varepsilon N^{1/2}\}$. Then

$$\frac{m}{N} N^{-3/2} \sum_{i \notin \mathcal{S}_\varepsilon} \frac{t_i(t_i - 1)}{2} \leq \varepsilon N^{-1} \sum_{i \notin \mathcal{S}_\varepsilon} t_i \leq \varepsilon.$$

On the other hand, for $i \in \mathcal{S}_\varepsilon$, and $s_{i-1} < j \leq s_i$ we have $q_N(j) - q_N^{\text{avg}}(j) = t_i(t_i - 1)/2 \geq \varepsilon^2 N/2$. Hence, if J be an index chosen uniformly at random from $\{1, 2, \dots, N\}$ (independent of everything else), then

$$\begin{aligned} N^{-3/2} \sum_{i \in \mathcal{S}_\varepsilon} \frac{t_i(t_i - 1)}{2} &= N^{-3/2} \sum_{i \in \mathcal{S}_\varepsilon} \sum_{j=s_{i-1}+1}^{s_i} (q_N(j) - q_N^{\text{avg}}(j)) \\ &= N^{-1/2} P(q_N(J) - q_N^{\text{avg}}(J) \geq \varepsilon^2 N/2) \\ &\leq (\varepsilon^2/2)^{-2} N^{-1/2} N^{-2} \mathbb{E}(q_N(J) - q_N^{\text{avg}}(J))^2 \\ &= (\varepsilon^2/2)^{-2} N^{-7/2} \sum_{j=1}^N (q_N(j) - q_N^{\text{avg}}(j))^2. \end{aligned}$$

Therefore it suffices to show that as $N \rightarrow \infty$,

$$\sum_{j=1}^N (q_N^{\text{avg}}(j) - q_N(j))^2 = O(N^3). \quad (\text{F.1})$$

Note that for each $1 \leq j \leq k$, $\sum_{i=s_{j-1}+1}^{s_j} (q_N^{\text{avg}}(i) - i) = 0$. Thus $\bar{q}_N^{\text{avg}} = N^{-1} \sum_{i=1}^N i = (N+1)/2$, and we obtain the following.

$$\begin{aligned} \sum_{i=s_{j-1}+1}^{s_j} \left((q_N^{\text{avg}}(i) - \bar{q}_N^{\text{avg}})^2 - (i - \bar{q}_N^{\text{avg}})^2 \right) &= \sum_{i=s_{j-1}+1}^{s_j} (q_N^{\text{avg}}(i)^2 - 2\bar{q}_N^{\text{avg}}(q_N^{\text{avg}}(i) - i) - i^2) \\ &= - \sum_{i=s_{j-1}+1}^{s_j} \left(i^2 - \left(\frac{s_{j-1} + 1 + s_j}{2} \right)^2 \right) \\ &= - \sum_{i=s_{j-1}+1}^{s_j} \left(i - \frac{s_{j-1} + 1 + s_j}{2} \right)^2 \\ &= -\frac{1}{12} t_j (t_j^2 - 1). \end{aligned}$$

Therefore

$$\sum_{i=1}^N (q_N^{\text{avg}}(i) - \bar{q}_N^{\text{avg}})^2 = \sum_{i=1}^N (i - \bar{q}_N^{\text{avg}})^2 - \frac{1}{12} \sum_{j=1}^k t_j (t_j^2 - 1)$$

Notice that

$$0 \leq \frac{1}{12} \sum_{j=1}^k t_j (t_j^2 - 1) \leq \frac{1}{12} \left(\sum_{i=1}^k t_i \right) \max_{1 \leq i \leq k} t_i^2 = \frac{N}{12} \max_{1 \leq i \leq k} t_i^2$$

and therefore [Assumption 5](#) tells us that

$$\sum_{i=1}^N (q_N^{\text{avg}}(i) - \bar{q}_N^{\text{avg}})^2 = \frac{N(N^2 - 1)}{12} + o(N^3).$$

On the other hand, it follows from the proof of [Lemma D.5](#) that

$$\sum_{i=1}^N \left(q_N(i) - \frac{N+1}{2} \right)^2 = \frac{N(N^2 - 1)}{12} + o(N^3)$$

Combining these and doing some algebraic manipulations, [\(F.1\)](#) follows. Hence the proof. \square