

Distribution-free tests of multivariate independence based on center-outward quadrant, Spearman, Kendall, and van der Waerden statistics

Hongjian Shi^{*} Mathias Drton[†] Marc Hallin[‡] and Fang Han[§]

Abstract

Due to the lack of a canonical ordering in $\mathbb{R}^d$ for $d > 1$, defining multivariate generalizations of the classical univariate ranks has been a long-standing open problem in statistics. Optimal transport has been shown to offer a solution in which multivariate ranks are obtained by transporting data points to a grid that approximates a uniform reference measure (Chernozhukov et al., 2017; Hallin, 2017; Hallin et al., 2021), thereby inducing ranks, signs, and a data-driven ordering of $\mathbb{R}^d$. We take up this new perspective to define and study multivariate analogues of the sign covariance/quadrant statistic, Spearman’s rho, Kendall’s tau, and van der Waerden covariances. The resulting tests of multivariate independence are fully distribution-free, hence uniformly valid irrespective of the actual (absolutely continuous) distribution of the observations. Our results provide the asymptotic distribution theory for these new test statistics, with asymptotic approximations to critical values to be used for testing independence between random vectors, as well as a power analysis of the resulting tests in an extension of the so-called (bivariate) Konijn model. This power analysis includes a multivariate Chernoff–Savage property guaranteeing that, under elliptical generalized Konijn models, the asymptotic relative efficiency of our van der Waerden tests with respect to Wilks’ classical (pseudo-)Gaussian procedure is strictly larger than or equal to one, where equality is achieved under Gaussian distributions only. We similarly provide a lower bound for the asymptotic relative efficiency of our Spearman procedure with respect to Wilks’ test, thus extending the classical result by Hodges and Lehmann on the asymptotic relative efficiency, in univariate location models, of Wilcoxon tests with respect to the Student ones.

Keywords: Center-outward ranks and signs, elliptical Chernoff–Savage property, multivariate independence test, Pitman asymptotic relative efficiency.

^{*}Department of Mathematics, TUM School of Computation, Information and Technology, Technical University of Munich, 85748 Garching bei München, Germany; e-mail: hongjian.shi@tum.de

[†]Department of Mathematics, TUM School of Computation, Information and Technology, Technical University of Munich, 85748 Garching bei München, Germany; e-mail: mathias.drton@tum.de

[‡]ECARES and Department of Mathematics, Université Libre de Bruxelles, Brussels, Belgium; email: mhallin@ulb.ac.be

[§]Department of Statistics, University of Washington, Seattle, WA 98195, USA; e-mail: fanghan@uw.edu

1 Introduction

Testing independence between two observed random variables X_1 and X_2 is a fundamental problem in statistical inference and has important applications, basically, in all domains. In a bivariate context, when both X_1 and X_2 are univariate Gaussian, the classical tests are based on empirical correlations. These tests remain asymptotically valid under non-Gaussian distributions with finite variances and, therefore, are widely used in practice as pseudo-Gaussian tests. Rank-based alternatives, that do not require any moment assumptions, have been proposed at a very early stage: the Spearman and Kendall correlation coefficients (Spearman, 1904, Kendall, 1938), actually, were among the first applications of rank-based methods in statistical inference, well before Wilcoxon (1945) gave his rank sum and signed rank tests for location. The general opinion, however, was that the (high) price to be paid for the extended validity of these rank tests was their poor performance in terms of power ... until Hodges and Lehmann (1956) and Chernoff and Savage (1958) disproved this fact by showing that, in univariate location models, the asymptotic relative efficiency (ARE) of Wilcoxon tests relative to Student's t ones never falls below 0.864 while the same ARE, for the van der Waerden (normal score) version of rank tests is uniformly not smaller than one. Originally proved in the context of univariate location models, these Hodges–Lehmann and Chernoff–Savage results have been extended to the context of bivariate independence and the so-called Konijn model (Konijn (1956); see also Chapter III.6.1 in Hájek and Šidák (1967)) by Hallin and Paindaveine (2008).

In this paper, we consider the problem of testing *multivariate* independence, that is, independence between random vectors $\mathbf{X}_1$ and $\mathbf{X}_2$ with dimensions d_1 and d_2 such that $\max(d_1, d_2) > 1$. The classical pseudo-Gaussian test for this problem is Wilks' Gaussian likelihood ratio test (Wilks, 1935). Based on the concepts of multivariate *center-outward ranks* and *signs* recently introduced by Hallin et al. (2021), we construct fully distribution-free tests of the quadrant, Spearman, Kendall, and van der Waerden type and assess their asymptotic performance against generalized versions of the Konijn alternatives. We also provide (against elliptical generalized Konijn alternatives) a Hodges–Lehmann lower bound for the ARE of our Spearman tests, and establish a Chernoff–Savage property for our van der Waerden tests, which makes Wilks' pseudo-Gaussian ones Pitman-nonadmissible in this context.

1.1 Testing multivariate independence

The problem of testing independence between two random vectors $\mathbf{X}_1$ and $\mathbf{X}_2$ with dimensions d_1 and d_2 and unspecified densities is significantly harder for $\max(d_1, d_2) > 1$ than for $d_1 = d_2 = 1$ —due, mainly, to the difficulty of defining a multivariate counterpart to univariate ranks. The first attempt to provide a rank-based alternative to the Gaussian likelihood ratio method of Wilks (1935) was developed in Chapter 8 of Puri and Sen (1971) and, for almost thirty years, has remained the only rank-based approach to the problem. The proposed tests, however, are based on componentwise ranks, which are not distribution-free—unless, of course, both vectors have dimension one, in which case we are back to the traditional context of bivariate independence (e.g., Chapter III.6 of Hájek and Šidák, 1967). This issue persists in more recent work, e.g., that of Lin (2017), Weihs, Drton, and Meinshausen (2018), and Moon and Chen (2022). Other types of multivariate independence tests include Pillai (1955), Friedman and Rafsky (1983), Gretton et al.

(2005), Székely, Rizzo, and Bakirov (2007), Heller, Gorfine, and Heller (2012), Heller, Heller, and Gorfine (2013), Azadkia and Chatterjee (2021), and Lin and Han (2023), to name a few; see Shi, Drton, and Han (2022) and Shi et al. (2022) for a more complete review.

Alternatives to the Puri and Sen tests have appeared with the developments of multivariate concepts of signs and ranks. Based on Randles (1989)’s concept of *interdirections*, Gieser (1993) and Gieser and Randles (1997) proposed a sign test extending the univariate quadrant test (Blomqvist, 1950). Taskinen, Oja, and Randles (2005) also proposed, based on the so-called standardized *spatial signs*, a sign test which, under elliptical symmetry assumptions, is asymptotically equivalent to the Gieser and Randles one. *Spatial ranks* are introduced, along with *spatial signs*, in Taskinen, Oja, and Randles (2005), where multivariate analogs of Spearman’s *rho* and Kendall’s *tau* are considered; the Spearman tests (involving Wilcoxon scores) are extended, in Taskinen, Kankainen, and Oja (2004), to the case of arbitrary square-integrable score functions, which includes van der Waerden (normal score) tests. All these multivariate rank-based tests are enjoying, under elliptical symmetry, many of the attractive properties of their traditional univariate counterparts. In particular, under the assumptions of elliptical symmetry, they are asymptotically distribution-free. Local powers are obtained in Taskinen, Kankainen, and Oja (2004) against elliptical extensions of the *Konijn alternatives* (Konijn, 1956). Hodges–Lehmann and Chernoff–Savage results have been established by Hallin and Paindaveine (2008); the latter entail the Pitman-nonadmissibility, in this elliptical context, of Wilks’ classical procedure.

The validity of all these tests, however, is limited to subclasses of distributions—essentially, the family of elliptically symmetric ones—which, from the perspective we are taking here, are overly restrictive; the assumption of elliptical symmetry, in particular, is extremely strong, and unlikely to hold in most applications. Moreover, there is a crucial difference between finite-sample and asymptotic distribution-freeness. Indeed, one should be wary that a sequence of tests $\psi^{(n)}$ with asymptotic size $\lim_{n \rightarrow \infty} \mathbb{E}_P[\psi^{(n)}] = \alpha$ under any element P in a class $\mathcal{P}$ of distributions does not necessarily have asymptotic size α under unspecified $P \in \mathcal{P}$: the convergence of $\mathbb{E}_P[\psi^{(n)}]$ to α , indeed, typically is not uniform over $\mathcal{P}$, so that, in general, $\lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} \mathbb{E}_P[\psi^{(n)}] \neq \alpha$. This is the case for most pseudo-Gaussian procedures, including Wilks’ test. Genuinely distribution-free tests $\psi^{(n)}$, where $\mathbb{E}_P[\psi^{(n)}]$ does not depend on P , do not suffer that problem, and this is why finite-sample distribution-freeness is a fundamental property.

Building on the concept of center-outward ranks and signs recently proposed by Hallin et al. (2021), Shi et al. (2022) are introducing fully distribution-free rank-based versions of a class of *generalized symmetric covariances* which includes, among others, the sophisticated measures of multivariate dependence proposed by Székely, Rizzo, and Bakirov (2007); Székely and Rizzo (2014), Weihs, Drton, and Meinshausen (2018), Zhu et al. (2017), and Kim, Balakrishnan, and Wasserman (2020) (none of which is distribution-free). These rank-based generalized symmetric covariances are cumulating the advantages of distribution-freeness with those of “universal” consistency. However, they also are relatively complex, with tricky non-Gaussian asymptotics involving (Proposition 5.1 of Shi et al. (2022)) the eigenvalues of an integral equation. Practitioners, therefore, may prefer simpler, more familiar and easily interpretable extensions of the classical bivariate quadrant, Spearman, Kendall, and van der Waerden tests. Defining such extensions and studying their performance is the objective of this paper.

1.2 Center-outward signs and ranks

For dimension $d > 1$, the real space $\mathbb{R}^d$ lacks a canonical ordering. As a result, defining, in dimension $d > 1$, concepts of signs and ranks enjoying the properties that make the traditional ranks successful tools in univariate inference has been an open problem for more than half a century. One of the most important properties is exact distribution-freeness (for i.i.d. samples from absolutely continuous distributions). In an important new development involving optimal transport, the concept of center-outward ranks and signs was proposed recently by Chernozhukov et al. (2017), Hallin (2017), and Hallin et al. (2021) and, contrary to earlier concepts such as marginal ranks (Puri and Sen, 1971), spatial ranks (Oja, 2010), depth-based ranks (Liu and Singh, 1993; Zuo and He, 2006), or Mahalanobis ranks and signs (Hallin and Paindaveine, 2002a,b), enjoys a property of *maximal ancillarity*—intuitively, “*maximal distribution-freeness*”—under i.i.d. observations with unspecified absolutely continuous distribution.

Center-outward ranks and signs are based on measure transportation to the unit ball equipped with a spherical uniform distribution. They have been used, quite successfully, in a variety of statistical problems: rank tests and R-estimation for VARMA models (Hallin, La Vecchia, and Liu, 2022, 2023), rank tests for multiple-output regression and MANOVA (Hallin, Hlubinka, and Hudecová, 2023), and nonparametric multiple-output quantile regression (del Barrio, González-Sanz, and Hallin, 2022).

Other authors (Carlier, Chernozhukov, and Galichon, 2016, 2017, Ghosal and Sen, 2022, Deb and Sen, 2023, and Deb, Bhattacharya, and Sen, 2021) are considering measure transportation to the Lebesgue uniform over the unit cube rather than the spherical uniform over the unit ball; the finite-sample impact of that choice has been studied in Hallin and Mordant (2023) who show that it is relatively modest. The inverse of such a transport (a quantile function), however, does not preserve the symmetry features of the underlying distribution (for instance, the quantile function of a spherically symmetric distribution is not spherically symmetric), while the quantile function resulting from a transport to the unit ball does.

1.3 A motivating example

The importance and advantages of center-outward sign- and rank-based tests of independence are illustrated with the following stock market data example. The dataset, collected from Yahoo! Finance (finance.yahoo.com), contains prices for stocks in the Standard & Poor’s 100 (S&P 100) index for the years 2003–2012. We will analyze the daily log returns of adjusted closing prices on the last trading day of the first month of each quarter. The total number of observations, thus, is $n = 40$; the time lag between two consecutive observations is three months, in a financial series where autocorrelations typically are small (see Figure S.1 in the supplement). Treating these observations as independent and identically distributed, thus, is unlikely to affect the validity of the tests.

The S&P 100 stocks are classified into ten sectors by Global Industry Classification Standard (GICS). To illustrate the advantages of the proposed tests, we focus on detecting dependence between two such sectors: (1) Telecommunications, including “AT&T Inc [T]” and “Verizon Communications [VZ]” stocks; and (2) Materials, including “Du Pont (E.I.) [DD]”, “Dow Chemical [DOW]”, “Freeport-McMoran Cp & Gld [FCX]”, and “Monsanto Co. [MON]” stocks.




Figure 1: Univariate histograms (on the diagonal), bivariate scatter plots (below the diagonal), and the lag-zero Pearson correlation coefficients with 99% confidence intervals (above the diagonal) for the forty observations of daily log returns of six stocks: “AT&T Inc [T]”, “Verizon Communications [VZ]”, “Du Pont (E.I.) [DD]”, “Dow Chemical [DOW]”, “Freeport-McMoran Cp & Gld [FCX]”, and “Monsanto Co. [MON].”

Figure 1 presents univariate histograms and bivariate scatterplots for the forty observations (separated by three months) of the daily log returns of the six stocks mentioned above. We notice that all univariate marginals are skewed and/or heavy-tailed. We also notice that the lag-zero Pearson correlations between any stock in $\{T, VZ\}$ and any stock in $\{DD, DOW, FCX, MON\}$ are relatively small, and that most correlations are non-significant at significance level 1%.

We applied the four rank-based tests (quadrant (sign), Spearman, Kendall, van der Waerden) proposed in Section 3.3 to the forty bivariate observations of (T, VZ) and either (DD, DOW) , (DD, FCX) , (DD, MON) , (DOW, FCX) , (DOW, MON) , or (FCX, MON) . These tests are based on the matrices defined in (5)–(8). As a benchmark, we also include Wilks’ Gaussian likelihood ratio test (LRT). The p -values for all these tests are reported in Table 1. One observes that the rank-based tests perform as well as Wilks’ LRT in the first three columns while, in the last three, they yield strong evidence against independence whereas Wilks is unable to reject independence at 5% significance level. The case of Telecommunications and Materials is particularly clear, with p -values of 0.006 for van der Waerden and 0.007 for Spearman in testing independence between (T, VZ) and (DOW, FCX) —to be compared with Wilks’ p -value 0.174.

Table 1: P-values of the quadrant, Spearman, Kendall, van der Waerden, and Wilks tests of the hypotheses of independence between the bivariate vector (T, VZ) and each of the six bivariate vectors (DD, DOW) , (DD, FCX) , (DD, MON) , (DOW, FCX) , (DOW, MON) , or (FCX, MON) .

		(DD, DOW)	(DD, FCX)	(DD, MON)	(DOW, FCX)	(DOW, MON)	(FCX, MON)
quadrant (sign)		0.002	0.050	0.027	0.013	0.008	0.023
Spearman		< 0.001	0.005	0.004	0.007	0.001	0.011
Kendall	(T,VZ)	0.002	0.013	0.016	0.011	0.002	0.036
van der Waerden		< 0.001	0.012	0.011	0.006	< 0.001	0.010
Wilks (LRT)		0.011	0.016	0.009	0.174	0.072	0.074

Rank-based sign, Spearman, Kendall, and van der Waerden tests, thus, are able to detect dependencies that Wilks’ traditional pseudo-Gaussian test cannot.

1.4 Outline of the paper

The paper is organized as follows. Section 2 briefly reviews the notions of center-outward ranks and signs. Section 3 introduces our rank tests of multivariate independence. In Section 4, we establish a Chernoff–Savage property and a Hodges–Lehmann result for our center-outward van der Waerden and Spearman tests, respectively. We briefly conclude in Section 5. Due to the importance, all the proofs are given in Section 6. The auxiliary results and numerical experiments are deferred to the supplement (Shi et al., 2024).

2 Center-outward distribution functions, ranks, and signs

2.1 Definitions

Denoting by $\mathbb{S}_d$ and $\mathcal{S}_{d-1}$, respectively, the open unit ball and the unit hypersphere in $\mathbb{R}^d$, let U_d stand for the spherical¹ uniform distribution over $\mathbb{S}_d$. Let $\mathcal{P}_d$ denote the class of Lebesgue-absolutely continuous distributions over $\mathbb{R}^d$. For any P in $\mathcal{P}_d$, the main result in [McCann \(1995\)](#) then implies the existence of a P -a.s. unique gradient $\nabla\phi$ of a convex (and lower semi-continuous) function $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $\nabla\phi$ pushes P forward to U_d , i.e., $\nabla\phi(\mathbf{Z}) \sim U_d$ under $\mathbf{Z} \sim P$. Call *center-outward distribution function* of P any version $\mathbf{F}_\pm$ of this a.e. unique gradient.

Turning to sample versions, denote by $\mathbf{Z}^{(n)} := (\mathbf{Z}_1^{(n)}, \dots, \mathbf{Z}_n^{(n)})$, $n \in \mathbb{N}$ a triangular array of i.i.d. d -dimensional random vectors with distribution P . The *empirical center-outward distribution function* $\mathbf{F}_\pm^{(n)}$ of $\mathbf{Z}^{(n)}$ maps the n -tuple $\mathbf{Z}_1^{(n)}, \dots, \mathbf{Z}_n^{(n)}$ to a “regular” grid $\mathfrak{G}_n$ of the unit ball $\mathbb{S}_d$.

This regular grid $\mathfrak{G}_n$ is expected to approximate the spherical uniform distribution U_d over the unit ball $\mathbb{S}_d$. The only mathematical requirement needed for the asymptotic results below is the weak convergence, as $n \rightarrow \infty$, of the uniform discrete distribution over $\mathfrak{G}_n$ to the spherical uniform distribution U_d . A spherical uniform i.i.d. sample of n points over $\mathbb{S}_d$ (almost surely) satisfies such a requirement. Since the spherical uniform is highly symmetric, further symmetries can be imposed on $\mathfrak{G}_n$, though, which only can improve the convergence to U_d . In the sequel, we throughout assume that the grids $\mathfrak{G}_n$ are symmetric with respect to the origin, i.e., $\mathbf{u} \in \mathfrak{G}_n$ implies $-\mathbf{u} \in \mathfrak{G}_n$, which considerably simplifies formulas. More symmetry, however, can be assumed: see [Section 2.2](#) below. It should be insisted, however, that imposing such symmetry assumptions does not restrict the generality of the results since the construction of the grid is entirely under control.

The empirical counterpart $\mathbf{F}_\pm^{(n)}$ of $\mathbf{F}_\pm$ is defined as the bijective mapping from $\mathbf{Z}_1^{(n)}, \dots, \mathbf{Z}_n^{(n)}$ to the grid $\mathfrak{G}_n$ that minimizes $\sum_{i=1}^n \|\mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)}) - \mathbf{Z}_i^{(n)}\|^2$, where $\|\cdot\|$ denotes the Euclidean norm. That mapping is unique with probability one; in practice, it is obtained via a simple optimal assignment (pairing) algorithm—a linear program; see [Shi, Drton, and Han \(2022, Section 5.1\)](#) for a review and references therein. Call *rescaled center-outward rank* of $\mathbf{Z}_i^{(n)}$ the modulus

$$\tilde{R}_{i;\pm}^{(n)} := \|\mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)})\|, \quad i = 1, \dots, n \quad (1)$$

and *center-outward sign* of $\mathbf{Z}_i^{(n)}$ the unit vector

$$\mathbf{S}_{i;\pm}^{(n)} := \mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)}) / \|\mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)})\| \quad \text{for } \mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)}) \neq \mathbf{0}; \quad (2)$$

put $\mathbf{S}_{i;\pm}^{(n)} = \mathbf{0}$ for $\mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)}) = \mathbf{0}$. When the grid $\mathfrak{G}_n$ is obtained, as in [Section 2.2](#) below, with a factorization of n into $n_R n_S + n_0$, call $R_{i;\pm}^{(n)} := (n_R + 1) \tilde{R}_{i;\pm}^{(n)}$, which takes values² $(0), 1, \dots, n_R$, the *center-outward rank* of $\mathbf{Z}_i^{(n)}$.

2.2 Grid selection

The way (rescaled) ranks and signs in (1) and (2) are constructed, thus, depends on the way

¹Namely, U_d is the spherical distribution with uniform (over $[0, 1]$) radial density—equivalently, the product of a uniform over the distances to the origin and a uniform over the unit sphere $\mathcal{S}_{d-1}$. For $d = 1$, U_1 coincides with the Lebesgue uniform over $(-1, 1)$.

²The value $R_{i;\pm}^{(n)} = 0$ is attained only if $n_0 \neq 0$.

the grid $\mathfrak{G}_n$ is selected. As already mentioned, the only requirement for asymptotic results (as in Proposition 2.3 below) is the weak convergence to the spherical uniform U_d of the empirical distribution, $U_{\mathfrak{G}}^{(n)}$, say, over $\mathfrak{G}_n$. The closer $U_{\mathfrak{G}}^{(n)}$ to U_d , the better. Imposing on $\mathfrak{G}_n$ some of the many symmetries of U_d only can improve the finite-sample performance of $U_{\mathfrak{G}}^{(n)}$ as an approximation of U_d . Since U_d is the product of a uniform over the distances to the origin and a uniform over the unit sphere $\mathcal{S}_{d-1}$, it is natural to select $\mathfrak{G}_n$ such that $U_{\mathfrak{G}}^{(n)}$ similarly factorizes. This can be obtained as follows:

- (a) first factorize n into $n = n_R n_S + n_0$, with $0 \leq n_0 < \min(n_R, n_S)$ and $\min(n_R, n_S) \rightarrow \infty$ as $n \rightarrow \infty$;³
- (b) next consider a “regular array” $\mathfrak{G}_{n_S} := \{\mathbf{s}_1^{n_S}, \dots, \mathbf{s}_{n_S}^{n_S}\}$ of n_S points on the sphere $\mathcal{S}_{d-1}$ (see Remark 2.1 below);
- (c) construct the grid $\mathfrak{G}_n$ consisting in the collection of the $n_R n_S$ gridpoints of the form

$$(r/(n_R + 1))\mathbf{s}_s^{n_S}, \quad r = 1, \dots, n_R, \quad s = 1, \dots, n_S,$$

along with (n_0 copies of) the origin in case $n_0 \neq 0$: in total $n - (n_0 - 1)$ or n distinct points, thus, according as $n_0 > 0$ or $n_0 = 0$.

Remark 2.1. By “regular” array $\mathfrak{G}_{n_S}$ over $\mathcal{S}_{d-1}$ in (b) above, we mean “as regular as possible” an array $\mathfrak{G}_{n_S}$ —in the sense, for example, of the *low-discrepancy sequences* of the type considered in numerical integration, Monte-Carlo methods, and experimental design.⁴ The asymptotic results in Proposition 2.3 below only require the weak convergence, as $n_S \rightarrow \infty$, of the uniform discrete distribution over $\mathfrak{G}_{n_S}$ to the uniform distribution over $\mathcal{S}_{d-1}$. A uniform i.i.d. sample of points over $\mathcal{S}_{d-1}$ (almost surely) satisfies that requirement. However, one can easily construct arrays that are “more regular” than an i.i.d. one. In particular, in order for the grid $\mathfrak{G}_n$ described in (a)–(c) above to be symmetric with respect to the origin, one could select an even value of n_S and see that $\mathbf{s}_s^{n_S} \in \mathfrak{G}_{n_S}$ implies $-\mathbf{s}_s^{n_S} \in \mathfrak{G}_{n_S}$, so that $\sum_{i=1}^n \mathbf{S}_{i;\pm}^{(n)} = n_R \sum_{s=1}^{n_S} \mathbf{s}_s^{n_S} = \mathbf{0}$.

Remark 2.2. Some desirable finite-sample properties, such as strict independence between the ranks and the signs, hold with grids constructed as in (a)–(c) above provided, however, that $n_0 = 0$ or 1. This is due to the fact that the mapping from the sample to the grid is no longer injective for $n_0 \geq 2$. This fact, which has no asymptotic consequences (since the number n_0 of tied values involved is $o(n)$ as $n \rightarrow \infty$), is easily taken care of by performing the following tie-breaking device in step (c) of the construction of $\mathfrak{G}_n$: (i) randomly select n_0 directions $\mathbf{s}_1^0, \dots, \mathbf{s}_{n_0}^0$ in $\mathfrak{G}_{n_S}$, then (ii) replace the n_0 copies of the origin with the new gridpoints $[1/2(n_R + 1)]\mathbf{s}_1^0, \dots, [1/2(n_R + 1)]\mathbf{s}_{n_0}^0$. This new grid (for simplicity, the same notation $\mathfrak{G}_n$ is used as for the original one) no longer has multiple points: the optimal pairing between the sample and the grid is bijective and the resulting (rescaled) ranks and signs are mutually independent.

2.3 Main properties

This section summarizes some of the main properties of the concepts defined in Section 2.1; further properties and the proofs can be found in Hallin et al. (2021), Hallin, Hlubinka, and Hudcová

³Note that this implies that $n_0/n = o(1)$ as $n \rightarrow \infty$. See Mordant (2021, Section 7.4) for a discussion of the selection of n_R and n_S .

⁴See also Hallin and Mordant (2023) for a spherical version of the so-called *Halton sequences*.

(2023), and Hallin (2022). In the following propositions, the grids $\mathfrak{G}_n$ used in the construction of the empirical center-outward distribution function $\mathbf{F}_\pm^{(n)}$ are only required to satisfy the minimal assumption of an empirical distribution $U_{\mathfrak{G}}^{(n)}$ converging weakly to U_d .

Proposition 2.3. *Let $\mathbf{F}_\pm$ denote the center-outward distribution function of $P \in \mathcal{P}_d$. Then,*

- (i) $\mathbf{F}_\pm$ is a probability integral transformation of $\mathbb{R}^d$, that is, $\mathbf{F}_\pm(\mathbf{Z}) \sim U_d$ if and only if $\mathbf{Z} \sim P$; under $\mathbf{Z} \sim P$, $\|\mathbf{F}_\pm(\mathbf{Z})\|$ is uniform over $[0, 1]$, $\mathbf{F}_\pm(\mathbf{Z})/\|\mathbf{F}_\pm(\mathbf{Z})\|$ is uniform over the sphere $\mathcal{S}_{d-1}$, and they are mutually independent.

Let $\mathbf{Z}_1^{(n)}, \dots, \mathbf{Z}_n^{(n)}$ be i.i.d. with distribution $P \in \mathcal{P}_d$ and empirical center-outward distribution function $\mathbf{F}_\pm^{(n)}$. Then, if there are no ties in the grid $\mathfrak{G}_n$,

- (ii) $(\mathbf{F}_\pm^{(n)}(\mathbf{Z}_1^{(n)}), \dots, \mathbf{F}_\pm^{(n)}(\mathbf{Z}_n^{(n)}))$ is uniformly distributed over the $n!$ permutations of $\mathfrak{G}_n$;
- (iii) the n -tuple $(\mathbf{F}_\pm^{(n)}(\mathbf{Z}_1^{(n)}), \dots, \mathbf{F}_\pm^{(n)}(\mathbf{Z}_n^{(n)}))$ is strongly essentially maximal ancillary;⁵
- (iv) (pointwise convergence) for any $i = 1, \dots, n$,

$$\|\mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)}) - \mathbf{F}_\pm(\mathbf{Z}_i^{(n)})\| \rightarrow 0 \text{ a.s. as } n \rightarrow \infty.$$

Assume, moreover, that P is in the so-called class $\mathcal{P}_d^+ \subset \mathcal{P}_d$ of distributions with nonvanishing densities—namely, the class of distributions with density $f := dP/d\mu_d$ (μ_d the d -dimensional Lebesgue measure) such that, for all $D \in \mathbb{R}^+$, there exist constants $\lambda_{D;P}^-$ and $\lambda_{D;P}^+$ satisfying

$$0 < \lambda_{D;P}^- \leq f(\mathbf{z}) \leq \lambda_{D;P}^+ < \infty \quad (3)$$

for all $\mathbf{z}$ with $\|\mathbf{z}\| \leq D$. Then,

- (v) there exists a version of $\mathbf{F}_\pm$ defining a homeomorphism between the punctured unit ball $\mathbb{S}_d \setminus \{\mathbf{0}\}$ and $\mathbb{R}^d \setminus \mathbf{F}_\pm^{-1}(\{\mathbf{0}\})$; that version has a continuous inverse $\mathbf{Q}_\pm$ (with domain $\mathbb{S}_d \setminus \{\mathbf{0}\}$), which naturally qualifies as P 's center-outward quantile function;
- (vi) (Glivenko–Cantelli)

$$\max_{1 \leq i \leq n} \|\mathbf{F}_\pm^{(n)}(\mathbf{Z}_i^{(n)}) - \mathbf{F}_\pm(\mathbf{Z}_i^{(n)})\| \rightarrow 0 \text{ a.s. as } n \rightarrow \infty.$$

Results (v) and (vi) are due to Figalli (2018), Hallin (2017), del Barrio et al. (2018), Hallin et al. (2021) and can be extended (del Barrio, González-Sanz, and Hallin, 2020) to a more general⁶ class $\mathcal{P}_d^{\text{conv}}$ of absolutely continuous distributions, with density f supported on a open convex set of $\mathbb{R}^d$ but not necessarily the whole space, while the definition of $\mathbf{F}_\pm$ given in Hallin et al. (2021) aims at selecting, for each $P \in \mathcal{P}_d$, a version of $\nabla\phi$ which, whenever $P \in \mathcal{P}_d^{\text{conv}}$, is yielding the homeomorphism mentioned in (v). For the sake of simplicity, since we are not interested in quantiles, we stick here to the P-a.s. unique definition given above for $P \in \mathcal{P}_d$.

Center-outward distribution functions, ranks, and signs also inherit, from the invariance of squared Euclidean distances, elementary but quite remarkable invariance and equivariance properties under shifts, orthogonal transformations, and global rescaling (see Hallin, Hlubinka, and Hudecová (2023)). Denote by $\mathbf{F}_\pm^{\mathbf{Z}}$ the center-outward distribution function of $\mathbf{Z}$ and by $\mathbf{F}_\pm^{\mathbf{Z}; \mathfrak{G}_n}$ the empirical center-outward distribution function of an i.i.d. sample $\mathbf{Z}_1, \dots, \mathbf{Z}_n$ associated with a grid $\mathfrak{G}_n$.

⁵See Section 2.4 and Appendices D.1 and D.2 of Hallin et al. (2021) for a precise definition and a proof of this essential property.

⁶Namely, $\mathcal{P}_d^+ \subsetneq \mathcal{P}_d^{\text{conv}} \subsetneq \mathcal{P}_d$.

Proposition 2.4. Let $\boldsymbol{\mu} \in \mathbb{R}^d$, $k \in \mathbb{R}^+$, and denote by $\mathbf{O}$ a $d \times d$ orthogonal matrix. Then,

- (i) $\mathbf{F}_{\pm}^{\boldsymbol{\mu}+k\mathbf{O}\mathbf{Z}}(\boldsymbol{\mu} + \mathbf{O}\mathbf{z}) = \mathbf{O}\mathbf{F}_{\pm}^{\mathbf{Z}}(\mathbf{z})$, $\mathbf{z} \in \mathbb{R}^d$;
- (ii) denoting by $\mathbf{F}_{\pm}^{\boldsymbol{\mu}+k\mathbf{O}\mathbf{Z};\mathbf{O}\mathfrak{G}_n}$ the empirical center-outward distribution function computed from the sample $\boldsymbol{\mu} + k\mathbf{O}\mathbf{Z}_1, \dots, \boldsymbol{\mu} + k\mathbf{O}\mathbf{Z}_n$ and the grid $\mathbf{O}\mathfrak{G}_n := \{\mathbf{O}\mathbf{u} : \mathbf{u} \in \mathfrak{G}_n\}$,

$$\mathbf{F}_{\pm}^{\boldsymbol{\mu}+k\mathbf{O}\mathbf{Z};\mathbf{O}\mathfrak{G}_n}(\boldsymbol{\mu} + k\mathbf{O}\mathbf{Z}_i) = \mathbf{O}\mathbf{F}_{\pm}^{\mathbf{Z};\mathfrak{G}_n}(\mathbf{Z}_i), \quad i = 1, \dots, n. \quad (4)$$

Note that the orthogonal transformations in Proposition 2.4 include the permutations of $\mathbf{Z}$'s components. Invariance with respect to such permutations is an essential requirement for hypothesis testing in multivariate analysis.

3 Rank-based tests for multivariate independence

3.1 Center-outward test statistics for multivariate independence

In this section, we describe the test statistics we are proposing for testing independence between two random vectors. Consider a sample $(\mathbf{X}'_{11}, \mathbf{X}'_{21})', (\mathbf{X}'_{12}, \mathbf{X}'_{22})', \dots, (\mathbf{X}'_{1n}, \mathbf{X}'_{2n})'$ of n i.i.d. copies of some $(d_1 + d_2) = d$ -dimensional random vector $(\mathbf{X}'_1, \mathbf{X}'_2)'$ with Lebesgue-absolutely continuous joint distribution $P \in \mathcal{P}_d$ and density f . We are interested in the null hypothesis under which $\mathbf{X}_1$ and $\mathbf{X}_2$, with unspecified marginal distributions $P_1 \in \mathcal{P}_{d_1}$ (density f_1) and $P_2 \in \mathcal{P}_{d_2}$ (density f_2), respectively, are mutually independent: f then factorizes into $f = f_1 f_2$.

For $k = 1, 2$ and $i = 1, 2, \dots, n$, denote by $R_{ki;\pm}^{(n)}$, $\tilde{R}_{ki;\pm}^{(n)}$, and $\mathbf{S}_{ki;\pm}^{(n)}$ the center-outward rank, rescaled rank, and sign of $\mathbf{X}_{ki}$ computed from $\mathbf{X}_{k1}, \mathbf{X}_{k2}, \dots, \mathbf{X}_{kn}$ and the grid $\mathfrak{G}_n$. Recall that we throughout assume that if $\mathbf{u} \in \mathfrak{G}_n$ then $-\mathbf{u} \in \mathfrak{G}_n$. This implies that $\sum_{i=1}^n \mathbf{S}_{ki;\pm}^{(n)} = \mathbf{0}$ and $\sum_{i=1}^n J_k(\tilde{R}_{ki;\pm}^{(n)}) \mathbf{S}_{ki;\pm}^{(n)} = \mathbf{0}$ for any score function $J_k : [0, 1) \rightarrow \mathbb{R}$, $k = 1, 2$.

Consider now the $d_1 \times d_2$ matrices

$$\mathbf{W}_{\text{sign}}^{(n)} := \frac{1}{n} \sum_{i=1}^n \mathbf{S}_{1i;\pm}^{(n)} \mathbf{S}_{2i;\pm}^{(n)'} \quad (5)$$

$$\mathbf{W}_{\text{Spearman}}^{(n)} := \frac{1}{n} \sum_{i=1}^n \tilde{R}_{1i;\pm}^{(n)} \tilde{R}_{2i;\pm}^{(n)} \mathbf{S}_{1i;\pm}^{(n)} \mathbf{S}_{2i;\pm}^{(n)'} \quad (6)$$

$$\mathbf{W}_{\text{Kendall}}^{(n)} := \binom{n}{2}^{-1} \sum_{i < i'} \text{sign} \left[\left(\tilde{R}_{1i;\pm}^{(n)} \mathbf{S}_{1i;\pm}^{(n)} - \tilde{R}_{1i';\pm}^{(n)} \mathbf{S}_{1i';\pm}^{(n)} \right) \left(\tilde{R}_{2i;\pm}^{(n)} \mathbf{S}_{2i;\pm}^{(n)} - \tilde{R}_{2i';\pm}^{(n)} \mathbf{S}_{2i';\pm}^{(n)} \right)' \right], \quad (7)$$

where $\text{sign}[\mathbf{M}]$ stands for the matrix collecting the signs of the entries of a real matrix $\mathbf{M}$. More generally, let

$$\mathbf{W}_J^{(n)} := \frac{1}{n} \sum_{i=1}^n J_1(\tilde{R}_{1i;\pm}^{(n)}) J_2(\tilde{R}_{2i;\pm}^{(n)}) \mathbf{S}_{1i;\pm}^{(n)} \mathbf{S}_{2i;\pm}^{(n)'} \quad (8)$$

where the score functions $J_k : [0, 1) \rightarrow \mathbb{R}$, $k = 1, 2$ are continuous and square-integrable, with

$$0 < \lim_{n \rightarrow \infty} n^{-1} \sum_{i=1}^n J_k^2(r_i) = \int_0^1 J_k^2(u) du =: \sigma_{J_k}^2 < \infty \quad (9)$$

for any $\{r_1, \dots, r_n\}$ such that the uniform distribution over which converges weakly to the uniform

distribution over $[0, 1]$. This assumption on score functions will be made throughout the remainder of the paper. The matrices defined in (5)–(8) clearly constitute matrices of cross-covariance measurements based on center-outward ranks and signs (for $\mathbf{W}_{\text{sign}}^{(n)}$, signs only). For $d_1 = 1 = d_2$, it is easily seen that $\mathbf{W}_{\text{sign}}^{(n)}$, $\mathbf{W}_{\text{Spearman}}^{(n)}$, and $\mathbf{W}_{\text{Kendall}}^{(n)}$, up to scaling constants, reduce to the classical quadrant, Spearman, and Kendall test statistics, while $\mathbf{W}_J^{(n)}$ yields a matrix-valued score-based extension of Spearman’s correlation coefficient.

3.2 Asymptotic representation and asymptotic normality

Clearly, neither the ranks nor the signs are mutually independent ($n - 1$ rank and sign pairs indeed determine the last pair): traditional central-limit theorems thus do not apply. However, as we show now, each of the rank-based matrices defined in (5)–(8) admits an asymptotic representation in terms of i.i.d. variables. More precisely, defining $\mathbf{S}_{ki;\pm}$ as $\mathbf{F}_{k;\pm}(\mathbf{X}_{ki}) / \|\mathbf{F}_{k;\pm}(\mathbf{X}_{ki})\|$ if $\mathbf{F}_{k;\pm}(\mathbf{X}_{ki}) \neq \mathbf{0}$ and $\mathbf{0}$ otherwise for $k = 1, 2$, let

$$\mathbf{W}_{\text{sign}}^{(n)} := \frac{1}{n} \sum_{i=1}^n \mathbf{S}_{1i;\pm} \mathbf{S}_{2i;\pm}', \quad (10)$$

$$\mathbf{W}_{\text{Spearman}}^{(n)} := \frac{1}{n} \sum_{i=1}^n \mathbf{F}_{1;\pm}(\mathbf{X}_{1i}) \mathbf{F}_{2;\pm}'(\mathbf{X}_{2i}), \quad (11)$$

$$\mathbf{W}_{\text{Kendall}}^{(n)} := \binom{n}{2}^{-1} \sum_{i < i'} \text{sign} \left[\left(\mathbf{F}_{1;\pm}(\mathbf{X}_{1i}) - \mathbf{F}_{1;\pm}(\mathbf{X}_{1i'}) \right) \left(\mathbf{F}_{2;\pm}(\mathbf{X}_{2i}) - \mathbf{F}_{2;\pm}(\mathbf{X}_{2i'}) \right)' \right], \quad (12)$$

and

$$\mathbf{W}_J^{(n)} := \frac{1}{n} \sum_{i=1}^n J_1 \left(\|\mathbf{F}_{1;\pm}(\mathbf{X}_{1i})\| \right) J_2 \left(\|\mathbf{F}_{2;\pm}(\mathbf{X}_{2i})\| \right) \mathbf{S}_{1i;\pm} \mathbf{S}_{2i;\pm}'. \quad (13)$$

The following asymptotic representation results then hold under the null hypothesis of independence (hence, also under contiguous alternatives). As usual, let $\text{vec}(\mathbf{A}) := (\mathbf{a}'_1, \dots, \mathbf{a}'_m)'$ for an $n \times m$ matrix $\mathbf{A} = (\mathbf{a}_1, \dots, \mathbf{a}_m)$ and write $\mathbf{X}^{(n)} = o_{\text{q.m.}}(c^{(n)})$ when, for some sequence $c^{(n)}$ of positive real numbers, $|X_j^{(n)}| \leq Y^{(n)} c^{(n)}$ for each component $X_j^{(n)}$ of $\mathbf{X}^{(n)}$, with $\lim_{n \rightarrow \infty} E[(Y^{(n)})^2] = 0$.

Lemma 3.1. *Under the null hypothesis of independence, as n tends to infinity, $\text{vec}(\mathbf{W}_{\text{sign}}^{(n)} - \mathbf{W}_{\text{sign}}^{(n)})$, $\text{vec}(\mathbf{W}_{\text{Spearman}}^{(n)} - \mathbf{W}_{\text{Spearman}}^{(n)})$, $\text{vec}(\mathbf{W}_{\text{Kendall}}^{(n)} - \mathbf{W}_{\text{Kendall}}^{(n)})$, and $\text{vec}(\mathbf{W}_J^{(n)} - \mathbf{W}_J^{(n)})$ all are $o_{\text{q.m.}}(n^{-1/2})$.*

The asymptotic normality of $\text{vec}(\mathbf{W}_{\text{sign}}^{(n)})$, $\text{vec}(\mathbf{W}_{\text{Spearman}}^{(n)})$, $\text{vec}(\mathbf{W}_{\text{Kendall}}^{(n)})$, and $\text{vec}(\mathbf{W}_J^{(n)})$ follows immediately from the asymptotic representation results and the standard central-limit behavior of $\text{vec}(\mathbf{W}_{\text{sign}}^{(n)})$, $\text{vec}(\mathbf{W}_{\text{Spearman}}^{(n)})$, $\text{vec}(\mathbf{W}_{\text{Kendall}}^{(n)})$, and $\text{vec}(\mathbf{W}_J^{(n)})$.

Proposition 3.2. *Under the null hypothesis of independence, as n tends to infinity, $n^{1/2} \text{vec}(\mathbf{W}_{\text{sign}}^{(n)})$, $n^{1/2} \text{vec}(\mathbf{W}_{\text{Spearman}}^{(n)})$, $n^{1/2} \text{vec}(\mathbf{W}_{\text{Kendall}}^{(n)})$, and $n^{1/2} \text{vec}(\mathbf{W}_J^{(n)})$ are asymptotically normal with mean vectors $\mathbf{0}_{d_1 d_2}$ and covariance matrices*

$$\frac{1}{d_1 d_2} \mathbf{I}_{d_1 d_2}, \quad \frac{1}{9 d_1 d_2} \mathbf{I}_{d_1 d_2}, \quad \frac{4}{9} \mathbf{I}_{d_1 d_2}, \quad \text{and} \quad \frac{\sigma_{J_1}^2 \sigma_{J_2}^2}{d_1 d_2} \mathbf{I}_{d_1 d_2},$$

respectively.

3.3 Center-outward sign, Spearman, Kendall, and score tests

Associated with $\mathbf{W}_{\text{sign}}^{(n)}$, $\mathbf{W}_{\text{Spearman}}^{(n)}$, $\mathbf{W}_{\text{Kendall}}^{(n)}$, and $\mathbf{W}_J^{(n)}$ are the quadrant or sign, Spearman, Kendall, and score test statistics

$$\begin{aligned} T_{\text{sign}}^{(n)} &:= nd_1 d_2 \|\mathbf{W}_{\text{sign}}^{(n)}\|_F^2, & T_{\text{Spearman}}^{(n)} &:= 9nd_1 d_2 \|\mathbf{W}_{\text{Spearman}}^{(n)}\|_F^2, \\ T_{\text{Kendall}}^{(n)} &:= \frac{9n}{4} \|\mathbf{W}_{\text{Kendall}}^{(n)}\|_F^2, & T_J^{(n)} &:= \frac{nd_1 d_2}{\sigma_{J_1}^2 \sigma_{J_2}^2} \|\mathbf{W}_J^{(n)}\|_F^2, \end{aligned} \quad \text{and}$$

respectively, where $\|\mathbf{M}\|_F$ denotes the Frobenius norm of a matrix $\mathbf{M}$, and $\sigma_{J_1}^2$, $\sigma_{J_2}^2$ are defined as in (9). In particular, considering the van der Waerden score functions $J_k^{\text{vdW}}(u) := (F_{\chi_{d_k}^2}^{-1}(u))^{1/2}$, where $F_{\chi_d^2}$ stands for the χ_d^2 distribution function, the van der Waerden test statistic is $T_{\text{vdW}}^{(n)} = T_{J^{\text{vdW}}}^{(n)} = n \|\mathbf{W}_{J^{\text{vdW}}}^{(n)}\|_F^2$.

In view of the asymptotic normality results in Proposition 3.2, the tests $\psi_{\text{sign}}^{(n)}$, $\psi_{\text{Spearman}}^{(n)}$, $\psi_{\text{Kendall}}^{(n)}$, $\psi_{\text{vdW}}^{(n)}$, and $\psi_J^{(n)}$ rejecting the null hypothesis of independence whenever $T_{\text{sign}}^{(n)}$, $T_{\text{Spearman}}^{(n)}$, $T_{\text{Kendall}}^{(n)}$, $T_{\text{vdW}}^{(n)}$ or $T_J^{(n)}$, respectively, exceed the $(1 - \alpha)$ -quantile $\chi_{d_1 d_2; 1-\alpha}^2$ of a chi-square distribution with $d_1 d_2$ degrees of freedom have asymptotic⁷ level α . These tests are, however, strictly distribution-free, and exact critical values can be computed or simulated as well. The tests based on $T_{\text{sign}}^{(n)}$, $T_{\text{Spearman}}^{(n)}$, $T_{\text{Kendall}}^{(n)}$, and $T_{\text{vdW}}^{(n)}$ are multivariate extensions of the traditional quadrant, Spearman, Kendall and van der Waerden tests, respectively, to which they reduce for $d_1 = 1 = d_2$.

4 Local asymptotic powers

While there is only one way for two random vectors $\mathbf{X}_1$ and $\mathbf{X}_2$ to be independent, their mutual dependence can take many forms. The classical benchmark, in testing for bivariate independence, is a “local” form of an independent component analysis model that goes back to Konijn (1956) and Hájek and Šidák (1967). Multivariate elliptical extensions of such alternatives have been considered also by Gieser and Randles (1997), Taskinen, Kankainen, and Oja (2003), Hallin and Paindaveine (2008), and Oja, Paindaveine, and Taskinen (2016).

4.1 Generalized Konijn families

Let $\mathbf{X}^* = (\mathbf{X}_1^*, \mathbf{X}_2^*)'$, where $\mathbf{X}_1^*$ and $\mathbf{X}_2^*$ are mutually independent random vectors, with absolutely continuous distributions P_1 over $\mathbb{R}^{d_1}$ and P_2 over $\mathbb{R}^{d_2}$ and densities f_1 and f_2 , respectively. Then $\mathbf{X}^*$ has density $f = f_1 f_2$ over $\mathbb{R}^d$. Consider

$$\mathbf{X} = \begin{pmatrix} \mathbf{X}_1 \\ \mathbf{X}_2 \end{pmatrix} := \mathbf{M}_\delta \begin{pmatrix} \mathbf{X}_1^* \\ \mathbf{X}_2^* \end{pmatrix} := \begin{pmatrix} (1 - \delta)\mathbf{I}_{d_1} & \delta\mathbf{M}_1 \\ \delta\mathbf{M}_2 & (1 - \delta)\mathbf{I}_{d_2} \end{pmatrix} \begin{pmatrix} \mathbf{X}_1^* \\ \mathbf{X}_2^* \end{pmatrix} \quad (14)$$

where $\delta \in \mathbb{R}$ and $\mathbf{M}_1 \in \mathbb{R}^{d_1 \times d_2}$, $\mathbf{M}_2 \in \mathbb{R}^{d_2 \times d_1}$ are nonzero matrices. For given P_1 , P_2 , $\mathbf{M}_1$, $\mathbf{M}_2$, and δ , the distribution $P_\delta^{\mathbf{X}}(P_1, P_2; \mathbf{M}_1, \mathbf{M}_2)$ of $\mathbf{X}$ belongs to the one-parameter family

$$\mathcal{P}_{P_1, P_2; \mathbf{M}_1, \mathbf{M}_2}^{\mathbf{X}} := \{P_\delta^{\mathbf{X}}(P_1, P_2; \mathbf{M}_1, \mathbf{M}_2) \mid \delta \in \mathbb{R}\};$$

⁷In view of the distribution-freeness of their test statistics, the convergence to α of the actual size of these tests is uniform over the possible distributions $P_1 \in \mathcal{P}_{d_1}$ and $P_2 \in \mathcal{P}_{d_2}$.

call it a *generalized Konijn family*. Independence between $\mathbf{X}_1$ and $\mathbf{X}_2$, in such families, holds iff $\delta = 0$.

Sequences of the form $P_{n^{-1/2}\tau}^{\mathbf{X}}(P_1, P_2; \mathbf{M}_1, \mathbf{M}_2)$ with $\tau \neq 0$, as we shall see, define contiguous alternatives to the null hypothesis of independence in a sample of size n . More precisely, consider the following regularity assumptions on P_1 and P_2 (on their densities f_1 and f_2).

Assumption 4.1. (K1) *The densities f_1 and f_2 are such that*

$$\int_{\mathbb{R}^{d_k}} \mathbf{x} f_k(\mathbf{x}) d\mathbf{x} = \mathbf{0} \quad \text{and} \quad 0 < \int_{\mathbb{R}^{d_k}} \mathbf{x} \mathbf{x}' f_k(\mathbf{x}) d\mathbf{x} =: \Sigma_k < \infty, \quad k = 1, 2,$$

where $0 < \Sigma_k < \infty$, means the covariance matrix Σ_k is finite and positive definite.

(K2) *The functions $\mathbf{x}_k \mapsto (f_k(\mathbf{x}_k))^{1/2}$, $k = 1, 2$ admit quadratic mean partial derivatives⁸*

$$D_\ell[(f_k)^{1/2}], \quad \ell = 1, \dots, d_k, \quad k = 1, 2.$$

(K3) *The score $\boldsymbol{\varphi} := (\boldsymbol{\varphi}'_1, \boldsymbol{\varphi}'_2)'$ with $\varphi_{k;\ell} := -2D_\ell[(f_k)^{1/2}]/(f_k)^{1/2}$ is such that $0 < \int_{\mathbb{R}^{d_k}} (\varphi_{k;\ell}(\mathbf{x}))^2 f_k(\mathbf{x}) d\mathbf{x} < \infty$, $k = 1, 2$, $\ell = 1, \dots, d_k$,⁹ and*

$$\mathcal{J}_k := \text{Var}(\mathbf{X}_k^* \boldsymbol{\varphi}_k(\mathbf{X}_k^*)) = \int_{\mathbb{R}^{d_k}} (\mathbf{x}' \boldsymbol{\varphi}_k(\mathbf{x}) - d_k)^2 f_k(\mathbf{x}) d\mathbf{x}, \quad k = 1, 2$$

is finite and strictly positive.

It should be stressed that these regularity assumptions are not to be imposed on the observations in order for our rank-based tests to be valid; they only are required for deriving local power and ARE results.

The following local asymptotic normality (LAN) property then holds in the vicinity of $\delta = 0$.

Lemma 4.2. *Denote by $P_\delta^{(n)}$, $n \in \mathbb{N}$, the distribution of the triangular array $\mathbf{X}^{(n)} := (\mathbf{X}_1^{(n)}, \dots, \mathbf{X}_n^{(n)})$ of n independent copies of $\mathbf{X} = (\mathbf{X}'_1, \mathbf{X}'_2)' \sim P_\delta^{\mathbf{X}}(P_1, P_2; \mathbf{M}_1, \mathbf{M}_2) \in \mathcal{P}_{P_1, P_2; \mathbf{M}_1, \mathbf{M}_2}^{\mathbf{X}}$, where P_1 and P_2 satisfy Assumption 4.1. Then, the family $\{P_\delta^{(n)} | \delta \in \mathbb{R}\}$ is Locally Asymptotically Normal (LAN) at $\delta = 0$, with root- n contiguity rate, central sequence*

$$\Delta^{(n)}(\mathbf{X}^{(n)}) := n^{-1/2} \sum_{i=1}^n \left[\mathbf{X}_{1i}^{(n)'} \mathbf{M}'_2 \boldsymbol{\varphi}_2(\mathbf{X}_{2i}^{(n)}) + \mathbf{X}_{2i}^{(n)'} \mathbf{M}'_1 \boldsymbol{\varphi}_1(\mathbf{X}_{1i}^{(n)}) - \left(\mathbf{X}_{1i}^{(n)'} \boldsymbol{\varphi}_1(\mathbf{X}_{1i}^{(n)}) - d_1 \right) - \left(\mathbf{X}_{2i}^{(n)'} \boldsymbol{\varphi}_2(\mathbf{X}_{2i}^{(n)}) - d_2 \right) \right],$$

and Fisher information

$$\gamma^2 := \mathcal{J}_1 + \mathcal{J}_2 + \text{vec}'(\Sigma_1) \text{vec}(\mathbf{M}'_2 \mathcal{I}_2 \mathbf{M}_2) + \text{vec}'(\Sigma_2) \text{vec}(\mathbf{M}'_1 \mathcal{I}_1 \mathbf{M}_1) + \text{tr}(\mathbf{M}_1 \mathbf{M}_2) + \text{tr}(\mathbf{M}_2 \mathbf{M}_1)$$

where $\mathcal{I}_k := \int_{\mathbb{R}^{d_k}} \boldsymbol{\varphi}_k(\mathbf{x}) \boldsymbol{\varphi}_k'(\mathbf{x}) f_k(\mathbf{x}) d\mathbf{x}$, $k = 1, 2$. In other words, under $P_0^{(n)}$, as $n \rightarrow \infty$,

$$\Lambda^{(n)}(\mathbf{X}^{(n)}) := \log \frac{dP_{n^{-1/2}\tau}^{(n)}(\mathbf{X}^{(n)})}{dP_0^{(n)}}(\mathbf{X}^{(n)}) = \tau \Delta^{(n)}(\mathbf{X}^{(n)}) - \frac{1}{2} \tau^2 \gamma^2 + o_P(1) \quad (15)$$

and $\Delta^{(n)}(\mathbf{X}^{(n)})$ is asymptotically normal, with mean zero and variance γ^2 .

⁸Existence of quadratic mean partial derivatives is equivalent to quadratic mean differentiability; this was shown in Lind and Roussas (1972) and independently rediscovered by Garel and Hallin (1995, Lemma 2.1).

⁹Integration by parts yields $\int_{\mathbb{R}^{d_k}} \boldsymbol{\varphi}_k(\mathbf{x}) f_k(\mathbf{x}) d\mathbf{x} = \mathbf{0}$, $\int_{\mathbb{R}^{d_k}} \mathbf{x}' \boldsymbol{\varphi}_k(\mathbf{x}) f_k(\mathbf{x}) d\mathbf{x} = d_k$, and $\int_{\mathbb{R}^{d_k}} \mathbf{x} \boldsymbol{\varphi}_k(\mathbf{x})' f_k(\mathbf{x}) d\mathbf{x} = \mathbf{I}_{d_k}$, for $k = 1, 2$; see also Garel and Hallin (1995, page 555).

If P_1 and P_2 are elliptical with mean $\mathbf{0}$, scatter matrices Σ_k , and radial densities ϕ_k , $k = 1, 2$, viz.

$$f_k(\mathbf{x}_k) \propto (\det(\Sigma_k))^{-1/2} \phi_k\left(\sqrt{\mathbf{x}_k' \Sigma_k^{-1} \mathbf{x}_k}\right), \quad k = 1, 2,$$

write $P_\delta^{\text{ell}}(\phi_1, \phi_2, \Sigma_1, \Sigma_2; \mathbf{M}_1, \mathbf{M}_2)$ and $\mathcal{P}_{\phi_1, \phi_2, \Sigma_1, \Sigma_2; \mathbf{M}_1, \mathbf{M}_2}^{\text{ell}}$, respectively, instead of $P_\delta^{\mathbf{X}}(P_1, P_2; \mathbf{M}_1, \mathbf{M}_2)$ and $\mathcal{P}_{P_1, P_2; \mathbf{M}_1, \mathbf{M}_2}^{\mathbf{X}}$. The Konijn alternatives and Konijn families considered in [Gieser and Randles \(1997\)](#), [Taskinen, Kankainen, and Oja \(2003\)](#), and [Hallin and Paindaveine \(2008\)](#) are particular cases, of the form $P_\delta^{\text{ell}}(\phi_1, \phi_2, \mathbf{I}_{d_1}, \mathbf{I}_{d_2}; \mathbf{M}, \mathbf{M}')$ and $\mathcal{P}_{\phi_1, \phi_2, \mathbf{I}_{d_1}, \mathbf{I}_{d_2}; \mathbf{M}, \mathbf{M}'}^{\text{ell}}$: call them *elliptical* Konijn alternatives and *elliptical* Konijn families, respectively.

4.2 Limiting distributions and Pitman efficiencies

For the univariate two-sample location problem, [Chernoff and Savage \(1958\)](#) and [Hodges and Lehmann \(1956\)](#) established their celebrated results on the asymptotic relative efficiency (ARE) of traditional normal score (van der Waerden) and Wilcoxon rank tests with respect to Student's Gaussian procedure. These results were extended by [Hallin \(1994\)](#) and [Hallin and Tribel \(2000\)](#) to ARMA time-series models, by [Hallin and Paindaveine \(2002b\)](#) to Mahalanobis ranks-and-signs-based location tests under multivariate elliptical distributions, by [Hallin and Paindaveine \(2002a\)](#) to VARMA time-series models with elliptical innovations, by [Paindaveine \(2004\)](#) for the shape parameter of elliptical distributions.

Chernoff–Savage and Hodges–Lehmann bounds for the AREs (with respect to Hotelling) of measure-transportation-based center-outward rank and sign tests were first obtained by [Deb, Bhat-tacharya, and Sen \(2021\)](#) in the context of two-sample location models and for the subclasses of elliptical and independent component distributions. In this section, we similarly establish Chernoff–Savage and Hodges–Lehmann results for the AREs, relative to Wilks' test, of our center-outward van der Waerden and Spearman tests, respectively. The two-sample location model here is replaced with an elliptical Konijn model satisfying (in order for ARE values to make sense) Assumptions 4.1.

To this end, we first derive the limiting distributions of $\mathcal{T}_J^{(n)}$ (of which $\mathcal{T}_{\text{sign}}^{(n)}$, $\mathcal{T}_{\text{Spearman}}^{(n)}$, and $\mathcal{T}_{\text{vdW}}^{(n)}$ are particular cases) and $\mathcal{T}_{\text{Kendall}}^{(n)}$ under local sequences of (not necessarily elliptical) Konijn alternatives of the form $P_{n^{-1/2}\tau}^{\mathbf{X}}(P_1, P_2; \mathbf{M}_1, \mathbf{M}_2)$.

Theorem 4.3. *Let P_1 and P_2 satisfy Assumption 4.1. If the observations are n independent copies of $\mathbf{X} \sim P_{n^{-1/2}\tau}^{\mathbf{X}}(P_1, P_2; \mathbf{M}_1, \mathbf{M}_2)$,*

- (i) *the limiting distribution as $n \rightarrow \infty$ of the test statistic $\mathcal{T}_J^{(n)}$ is noncentral chi-square with $d_1 d_2$ degrees of freedom and noncentrality parameter*

$$C_J(\tau) \left\| \mathbb{E}_{H_0} \left[\mathbf{J}_1(\mathbf{F}_{1;\pm}(\mathbf{X}_1)) \mathbf{R} \mathbf{J}_2(\mathbf{F}_{2;\pm}(\mathbf{X}_2))' \right] \right\|_{\mathbf{F}}^2, \quad (16)$$

where $C_J(\tau) := \frac{\tau^2 d_1 d_2}{\sigma_{J_1}^2 \sigma_{J_2}^2}$, $\mathbf{R} := \mathbf{X}_1' \mathbf{M}_2' \varphi_2(\mathbf{X}_2) + \mathbf{X}_2' \mathbf{M}_1' \varphi_1(\mathbf{X}_1)$, $\mathbf{J}_k(\mathbf{u}) := J_k(\|\mathbf{u}\|) \frac{\mathbf{u}}{\|\mathbf{u}\|} \mathbf{1}_{[\|\mathbf{u}\| \neq 0]}$ for $\mathbf{u} \in \mathbb{S}_d$, $\mathbb{E}_{H_0}$ stands for expectations under the null ($\delta = 0$), and recall that $\|\mathbf{M}\|_{\mathbf{F}}$ denotes the Frobenius norm of $\mathbf{M}$;

- (ii) *the limiting distribution of the Kendall test statistic $\mathcal{T}_{\text{Kendall}}^{(n)}$ is noncentral chi-square with $d_1 d_2$*

degrees of freedom and noncentrality parameter

$$9\tau^2 \left\| \mathbb{E}_{H_0} \left[\mathbf{F}_{1;\pm}^\square(\mathbf{X}_1) \mathbf{R} \mathbf{F}_{2;\pm}^\square(\mathbf{X}_2)' \right] \right\|_{\mathbf{F}}^2,$$

where, denoting by $F_{k\ell}$ the marginal cumulative distribution function of $(\mathbf{F}_{k;\pm}(\mathbf{X}_k))_\ell$,

$$(\mathbf{F}_{k;\pm}^\square(\mathbf{X}_k))_\ell := 2F_{k\ell}((\mathbf{F}_{k;\pm}(\mathbf{X}_k))_\ell) - 1, \quad k = 1, 2, \quad \ell = 1, \dots, d_k.$$

The results for $\tilde{T}_{\text{sign}}^{(n)}$, $\tilde{T}_{\text{Spearman}}^{(n)}$, and $\tilde{T}_{\text{vdW}}^{(n)}$ follow as particular cases of (i) with the constant $C_J(\tau)$ taking values $\tau^2 d_1 d_2$, $9\tau^2 d_1 d_2$, and τ^2 , respectively.

Turning to Wilks' test, the test statistic takes the form

$$T_{\text{Wilks}}^{(n)} := n \log V^{(n)} \quad \text{with} \quad V^{(n)} := \frac{\det(\mathbf{S}_1^{(n)}) \det(\mathbf{S}_2^{(n)})}{\det(\mathbf{S}^{(n)})},$$

where $\det(\mathbf{M})$ denotes the determinant of a square matrix $\mathbf{M}$, $\mathbf{S}_k^{(n)}$ is the $d_k \times d_k$ sample covariance matrix of $\mathbf{X}_{k1}, \dots, \mathbf{X}_{kn}$, $k = 1, 2$, and $\mathbf{S}^{(n)}$ the sample covariance matrix of $(\mathbf{X}'_{11}, \mathbf{X}'_{21})', \dots, (\mathbf{X}'_{1n}, \mathbf{X}'_{2n})'$, which is $(d_1 + d_2) \times (d_1 + d_2)$. Under the null hypothesis of independence, $T_{\text{Wilks}}^{(n)}$ is asymptotically chi-square with $d_1 d_2$ degrees of freedom as soon as $\mathbf{X}_1$ and $\mathbf{X}_2$ have finite variances (Hallin and Paindaveine, 2008, pages 187–188; Oja, Paindaveine, and Taskinen, 2016, Section 5.1). Wilks' test $\psi_{\text{Wilks}}^{(n)}$, thus, rejects the null hypothesis at asymptotic level α whenever $T_{\text{Wilks}}^{(n)}$ exceeds the corresponding chi-square quantile of order $(1 - \alpha)$. Under alternatives of the form $\mathbf{X} \sim \mathcal{P}_{n-1/2\tau}^{\text{ell}}(\phi_1, \phi_2, \Sigma_1, \Sigma_2; \mathbf{M}_1, \mathbf{M}_2)$, with f_1 and f_2 satisfying Assumption 4.1, the limiting distribution of Wilks' statistic is noncentral chi-square, still with $d_1 d_2$ degrees of freedom, with noncentrality parameter

$$\tau^2 \left\| \Sigma_1^{1/2} \mathbf{M}_2' \Sigma_2^{-1/2} + \Sigma_1^{-1/2} \mathbf{M}_1 \Sigma_2^{1/2} \right\|_{\mathbf{F}}^2 \quad (17)$$

reducing, under the elliptical Konijn alternative $\mathcal{P}_{n-1/2\tau}^{\text{ell}}(\phi_1, \phi_2, \mathbf{I}_{d_1}, \mathbf{I}_{d_2}; \mathbf{M}, \mathbf{M}')$, to $4\tau^2 \|\mathbf{M}\|_{\mathbf{F}}^2$ (see, e.g., page 919 of Taskinen, Oja, and Randles (2005)). We are now ready to derive asymptotic relative efficiency results for our center-outward rank tests relative to Wilks' pseudo-Gaussian one.

Proposition 4.4. *Let $\mathcal{P}_1$ and $\mathcal{P}_2$ satisfying Assumption 4.1 be elliptically symmetric with radial densities ϕ_k and covariance matrices Σ_k , $k = 1, 2$. Then, the Pitman asymptotic relative efficiency (ARE) in $\mathcal{P}_{\phi_1, \phi_2, \Sigma_1, \Sigma_2; \mathbf{M}_1, \mathbf{M}_2}^{\text{ell}}$ of the center-outward test $\psi_J^{(n)}$ based on the score functions J_k , $k = 1, 2$ relative to Wilks' test $\psi_{\text{Wilks}}^{(n)}$ is*

$$\text{ARE}(\psi_J^{(n)}, \psi_{\text{Wilks}}^{(n)}) = \frac{\left\| D_1 C_2 \Sigma_1^{1/2} \mathbf{M}_2' \Sigma_2^{-1/2} + D_2 C_1 \Sigma_1^{-1/2} \mathbf{M}_1 \Sigma_2^{1/2} \right\|_{\mathbf{F}}^2}{d_1 d_2 \sigma_{J_1}^2 \sigma_{J_2}^2 \left\| \Sigma_1^{1/2} \mathbf{M}_2' \Sigma_2^{-1/2} + \Sigma_1^{-1/2} \mathbf{M}_1 \Sigma_2^{1/2} \right\|_{\mathbf{F}}^2}, \quad (18)$$

where, for $k = 1, 2$,

$$C_k = C_k(J_k, \phi_k) := \mathbb{E}[J_k(U) \rho_k(\tilde{F}_k^{-1}(U))] \quad \text{and} \quad D_k = D_k(J_k, \phi_k) := \mathbb{E}[J_k(U) \tilde{F}_k^{-1}(U)]$$

with $\rho_k := \frac{-\phi'_k}{\phi_k}$, $\tilde{F}_k$ the cumulative distribution function of $\|\Sigma_k^{-1/2} \mathbf{X}_k\|$, and U a random variable uniformly distributed over $(0, 1)$.

If, moreover, $\Sigma_1 \mathbf{M}_2' = \mathbf{M}_1 \Sigma_2$ —hence, in particular, under the family $\mathcal{P}_{\phi_1, \phi_2, \mathbf{I}_{d_1}, \mathbf{I}_{d_2}; \mathbf{M}, \mathbf{M}'}$, we have

- (i) $\text{ARE}(\psi_{\text{vdW}}^{(n)}, \psi_{\text{Wilks}}^{(n)}) \geq 1$, where $\psi_{\text{vdW}}^{(n)} = \psi_{J_{\text{vdW}}}^{(n)}$ with van der Waerden score functions J_k^{vdW} , $k = 1, 2$ and equality under Gaussian P_1 and P_2 only;
- (ii) $\text{ARE}(\psi_{\text{Spearman}}^{(n)}, \psi_{\text{Wilks}}^{(n)}) \geq \Omega(d_1, d_2) \geq 9/16$, where $\psi_{\text{Spearman}}^{(n)} = \psi_{J_{\text{W}}}^{(n)}$ with the Wilcoxon score functions $J_k^{\text{W}}(u) := u$,
- $$\Omega(d_1, d_2) := 9(2c_{d_1}^2 + d_1 - 1)^2(2c_{d_2}^2 + d_2 - 1)^2 / 1024d_1d_2c_{d_1}^2c_{d_2}^2,$$

$$c_d := \inf \{x > 0 \mid (\sqrt{x}B_{\sqrt{2d-1}/2}(x))' = 0\}, \quad \text{where} \quad B_a(x) := \sum_{m=0}^{\infty} \frac{(-1)^m}{m!\Gamma(m+a+1)} \left(\frac{x}{2}\right)^{2m+a}.$$

Note that the ARE values in (18) depend, via Σ_1 and Σ_2 , on the underlying covariance structure of $P_0^{\text{ell}}(\phi_1, \phi_2, \Sigma_1, \Sigma_2; \mathbf{M}_1, \mathbf{M}_2)$; a similar fact was already pointed out by Gieser (1993). Most authors (e.g. Gieser (1993), Gieser and Randles (1997), Taskinen, Kankainen, and Oja (2003, 2004), Taskinen, Oja, and Randles (2005), Hallin and Paindaveine (2008) and Deb, Bhattacharya, and Sen (2021)), therefore, restrict themselves to the particular case of elliptical Konijn families $\mathcal{P}_{\phi_1, \phi_2, \mathbf{I}_{d_1}, \mathbf{I}_{d_2}; \mathbf{M}_1, \mathbf{M}_2}^{\text{ell}}$. In (18), we more generally provide explicit results for the broader class of $\mathcal{P}_{\phi_1, \phi_2, \Sigma_1, \Sigma_2; \mathbf{M}_1, \mathbf{M}_2}^{\text{ell}}$ families with arbitrary covariances Σ_1, Σ_2 and arbitrary matrices $\mathbf{M}_1$ and $\mathbf{M}_2$.

Claim (i) entails the Pitman non-admissibility, within the class $\bigcup_{\phi_1, \phi_2, \mathbf{I}_{d_1}, \mathbf{I}_{d_2}, \mathbf{M}} \mathcal{P}_{\phi_1, \phi_2, \mathbf{I}_{d_1}, \mathbf{I}_{d_2}; \mathbf{M}, \mathbf{M}'}^{\text{ell}}$ of all elliptical Konijn families satisfying Assumption 4.1, of Wilks' test, which is uniformly dominated by our center-outward van der Waerden test. This is comparable with Theorem 4.1 in Deb, Bhattacharya, and Sen (2021), which deals with two-sample location tests. The proof relies on Proposition 1 in Hallin and Paindaveine (2008) (see also Paindaveine (2004)).

Claim (ii) is a multivariate extension of Hodges and Lehmann (1956)'s univariate result. The infimum $9/16 = 0.5625$ of $\Omega(d_1, d_2)$ is achieved for $d_1, d_2 \rightarrow \infty$. Table S.2 in the supplement gives numerical values of $\Omega(d_1, d_2)$ for $d_1, d_2 \leq 10$.

5 Conclusion

Optimal transport provides a new approach to rank-based statistical inference in dimension $d \geq 2$. The new multivariate ranks retain many of the favorable properties one is used to from the classical univariate ranks. Here, we demonstrate how the new multivariate ranks can be used for a definition of multivariate versions of popular rank-based correlation coefficients such as quadrant (sign) correlation, Kendall's tau, or Spearman's rho. We show how the new multivariate rank correlations yield fully distribution-free, yet powerful and computationally efficient tests of vector independence for which we provide explicit local asymptotic powers. We also show that the van der Waerden (normal score) version of our tests enjoy, relative to Wilks' classical pseudo-Gaussian procedure and, under an elliptical generalization of the traditional Konijn alternatives, the Chernoff–Savage property of their classical bivariate counterparts—which makes Wilks' test non-admissible in the context. These results are not specific to our choice of the uniform over the unit ball as the reference measure in our optimal transport approach; transports to the Lebesgue uniform over the unit cube (as in Chernozhukov et al. (2017)) or “direct” transports to, e.g., the spherical Gaussian (as in Deb, Bhattacharya, and Sen (2021)) would yield the same local asymptotic powers and asymptotic relative efficiencies.

6 Proofs

Proof of Lemma 3.1. We only need to prove the result for $\text{vec}(\mathbf{W}_J^{(n)})$ (since $\text{vec}(\mathbf{W}_{\text{sign}}^{(n)})$, $\text{vec}(\mathbf{W}_{\text{Spearman}}^{(n)})$, and $\text{vec}(\mathbf{W}_{\text{vdW}}^{(n)})$ are particular cases) and $\text{vec}(\mathbf{W}_{\text{Kendall}}^{(n)})$.

(a) Starting with $\text{vec}\mathbf{W}_J^{(n)}$, letting

$$\mathbf{Y}_{ki}^{(n)} := J_k\left(\tilde{R}_{ki;\pm}^{(n)}\right)\mathbf{S}_{ki;\pm}^{(n)} \quad \text{and} \quad \mathbf{Y}_{ki} := J_k\left(\|\mathbf{F}_{k;\pm}(\mathbf{X}_{ki})\|\right)\mathbf{S}_{ki;\pm}, \quad k = 1, 2, \quad i = 1, \dots, n,$$

rewrite $n^{1/2}(\mathbf{W}_J^{(n)} - \mathbf{W}_J^{(n)})_{\ell_1\ell_2}$, $\ell_1 = 1, \dots, d_1$, $\ell_2 = 1, \dots, d_2$ as

$$n^{-1/2} \sum_{i=1}^n \left((\mathbf{Y}_{1i}^{(n)})_{\ell_1} (\mathbf{Y}_{2i}^{(n)})_{\ell_2} - (\mathbf{Y}_{1i})_{\ell_1} (\mathbf{Y}_{2i})_{\ell_2} \right).$$

It suffices to show that

$$\mathbb{E}\left[\left\{n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i})_{\ell_1} (\mathbf{Y}_{2i})_{\ell_2}\right\}^2\right], \quad \mathbb{E}\left[\left\{n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i}^{(n)})_{\ell_1} (\mathbf{Y}_{2i}^{(n)})_{\ell_2}\right\}^2\right], \quad (19)$$

and

$$\mathbb{E}\left[\left\{n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i}^{(n)})_{\ell_1} (\mathbf{Y}_{2i}^{(n)})_{\ell_2}\right\} \left\{n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i})_{\ell_1} (\mathbf{Y}_{2i})_{\ell_2}\right\}\right] \quad (20)$$

tend to the same limit as n tends to infinity.

First, consider the left-hand quantity in (19). Due to the independence between $(\mathbf{Y}_{1i})_{\ell_1}$ and $(\mathbf{Y}_{2i})_{\ell_2}$ and the independence between $(\mathbf{Y}_{1i})_{\ell_1}$ and $(\mathbf{Y}_{2i'})_{\ell_2}$, $i \neq i'$, $\ell_1 = 1, \dots, d_1$, $\ell_2 = 1, \dots, d_2$, we have

$$\mathbb{E}\left[\left\{n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i})_{\ell_1} (\mathbf{Y}_{2i})_{\ell_2}\right\}^2\right] = \mathbb{E}[(\mathbf{Y}_{11})_{\ell_1}^2] \cdot \mathbb{E}[(\mathbf{Y}_{21})_{\ell_2}^2].$$

Turn to the right-hand quantity in (19). Since $((\mathbf{Y}_{11}^{(n)})_{\ell_1}, \dots, (\mathbf{Y}_{1n}^{(n)})_{\ell_1})$ and $((\mathbf{Y}_{21}^{(n)})_{\ell_2}, \dots, (\mathbf{Y}_{2n}^{(n)})_{\ell_2})$, for $\ell_1 = 1, \dots, d_1$, $\ell_2 = 1, \dots, d_2$, are independent, we have, in view of Proposition 2.3(ii) in Section 2.3 and Theorem 2 in Hoeffding (1951),

$$\mathbb{E}\left[\left\{n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i}^{(n)})_{\ell_1} (\mathbf{Y}_{2i}^{(n)})_{\ell_2}\right\}^2\right] = \frac{1}{n(n-1)} \sum_{\mathbf{u}_1 \in \mathfrak{G}_n^{d_1}} \left((\mathbf{J}_1(\mathbf{u}_1))_{\ell_1}\right)^2 \sum_{\mathbf{u}_2 \in \mathfrak{G}_n^{d_2}} \left((\mathbf{J}_2(\mathbf{u}_2))_{\ell_2}\right)^2,$$

where the right-hand side, by the properties of the grids $\mathfrak{G}_n^{d_k}$ and the fact that the score functions J_k , $k = 1, 2$, are continuous, square-integrable, and satisfy (9), tends to

$$\mathbb{E}[(\mathbf{J}_1(\mathbf{V}_1))_{\ell_1}^2] \cdot \mathbb{E}[(\mathbf{J}_2(\mathbf{V}_2))_{\ell_2}^2] = \mathbb{E}[(\mathbf{Y}_{11})_{\ell_1}^2] \cdot \mathbb{E}[(\mathbf{Y}_{21})_{\ell_2}^2]$$

with $\mathbf{V}_k \sim \mathbf{U}_{d_k}$, $k = 1, 2$.

Next, we obtain for (20)

$$\begin{aligned} & \mathbb{E}\left[n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i}^{(n)})_{\ell_1} (\mathbf{Y}_{2i}^{(n)})_{\ell_2} \times n^{-1/2} \sum_{i=1}^n (\mathbf{Y}_{1i})_{\ell_1} (\mathbf{Y}_{2i})_{\ell_2}\right] \\ &= n^{-1} \left\{ \sum_{i=1}^n \left(\mathbb{E}[(\mathbf{Y}_{1i}^{(n)})_{\ell_1} (\mathbf{Y}_{1i})_{\ell_1}] \right) \left(\mathbb{E}[(\mathbf{Y}_{2i}^{(n)})_{\ell_2} (\mathbf{Y}_{2i})_{\ell_2}] \right) \right\} \end{aligned}$$

$$+ \sum_{i \neq i'} \left(E[(\mathbf{Y}_{1i'}^{(n)})_{\ell_1} (\mathbf{Y}_{1i})_{\ell_1}] \right) \left(E[(\mathbf{Y}_{2i'}^{(n)})_{\ell_2} (\mathbf{Y}_{2i})_{\ell_2}] \right) \Big\}. \quad (21)$$

Since, for $k = 1, 2$, $\ell_k = 1, \dots, d_k$,

$$\begin{aligned} E[(\mathbf{Y}_{ki}^{(n)})_{\ell_k} (\mathbf{Y}_{ki})_{\ell_k}] + \sum_{i': i' \neq i} E[(\mathbf{Y}_{ki'}^{(n)})_{\ell_k} (\mathbf{Y}_{ki})_{\ell_k}] &= E[(\mathbf{Y}_{ki})_{\ell_k} \sum_{i'=1}^n (\mathbf{Y}_{ki'}^{(n)})_{\ell_k}] \\ &= E[(\mathbf{Y}_{ki})_{\ell_k} \sum_{\mathbf{u} \in \mathfrak{G}_n^{d_k}} (\mathbf{J}_k(\mathbf{u}))_{\ell_k}] = E[(\mathbf{Y}_{ki})_{\ell_k}] \left(\sum_{\mathbf{u} \in \mathfrak{G}_n^{d_k}} \mathbf{J}_k(\mathbf{u}) \right)_{\ell_k} = 0, \end{aligned}$$

and since, by symmetry, for all $i \neq i'$,

$$E[(\mathbf{Y}_{ki}^{(n)})_{\ell_k} (\mathbf{Y}_{ki})_{\ell_k}] = E[(\mathbf{Y}_{k1}^{(n)})_{\ell_k} (\mathbf{Y}_{k1})_{\ell_k}] \quad \text{and} \quad E[(\mathbf{Y}_{ki'}^{(n)})_{\ell_k} (\mathbf{Y}_{ki})_{\ell_k}] = E[(\mathbf{Y}_{k2}^{(n)})_{\ell_k} (\mathbf{Y}_{k1})_{\ell_k}],$$

we deduce that the right-hand side of (21) equals

$$\begin{aligned} n^{-1} \Big\{ n \left(E[(\mathbf{Y}_{11}^{(n)})_{\ell_1} (\mathbf{Y}_{11})_{\ell_1}] \right) \left(E[(\mathbf{Y}_{21}^{(n)})_{\ell_2} (\mathbf{Y}_{21})_{\ell_2}] \right) \\ + n(n-1) \left(- (n-1)^{-1} E[(\mathbf{Y}_{11}^{(n)})_{\ell_1} (\mathbf{Y}_{11})_{\ell_1}] \right) \left(- (n-1)^{-1} E[(\mathbf{Y}_{21}^{(n)})_{\ell_2} (\mathbf{Y}_{21})_{\ell_2}] \right) \Big\} \\ = \frac{n}{n-1} E[(\mathbf{Y}_{11}^{(n)})_{\ell_1} (\mathbf{Y}_{11})_{\ell_1}] \cdot E[(\mathbf{Y}_{21}^{(n)})_{\ell_2} (\mathbf{Y}_{21})_{\ell_2}]. \quad (22) \end{aligned}$$

The Proof of Theorem 3.1 in [del Barrio et al. \(2018\)](#) and the Proof of Proposition 3.3 in [Hallin et al. \(2021\)](#) entails $\mathbf{Y}_{k1}^{(n)} - \mathbf{Y}_{k1} \rightarrow 0$ a.s., while

$$\lim_{n \rightarrow \infty} E[\|\mathbf{Y}_{k1}^{(n)}\|^2] = E[\|\mathbf{Y}_{k1}\|^2] = \int_0^1 J_k^2(u) du < \infty.$$

It then follows (see, e.g., part (iv) of Theorem 5.7 in [Shorack \(2017, Chap. 3\)](#)) that

$$E[\|\mathbf{Y}_{k1}^{(n)} - \mathbf{Y}_{k1}\|^2] \rightarrow 0, \quad k = 1, 2 \quad \text{as } n \rightarrow \infty.$$

In particular, $E[(\mathbf{Y}_{k1}^{(n)} - \mathbf{Y}_{k1})_{\ell_k}^2] \rightarrow 0$ and thus

$$E[(\mathbf{Y}_{k1}^{(n)})_{\ell_k} (\mathbf{Y}_{k1})_{\ell_k}] = 2^{-1} \left(E[(\mathbf{Y}_{k1}^{(n)})_{\ell_k}^2] + E[(\mathbf{Y}_{k1})_{\ell_k}^2] - E[(\mathbf{Y}_{k1}^{(n)} - \mathbf{Y}_{k1})_{\ell_k}^2] \right) \rightarrow E[(\mathbf{Y}_{k1})_{\ell_k}^2].$$

It follows that the right-hand side of (22) tends to $E[(\mathbf{Y}_{11})_{\ell_1}^2] \cdot E[(\mathbf{Y}_{21})_{\ell_2}^2]$, which concludes the proof of the lemma for $\text{vec}(\mathbf{W}_J^{(n)})$.

(b) The case of the Kendall matrix $\text{vec}(\mathbf{W}_{\text{Kendall}}^{(n)})$ is slightly different, although the arguments in the proof are quite similar. We consider the Hájek projection of U-statistics (see, e.g., Proof of Theorem 7.1 in [Hoeffding \(1948\)](#)) for

$$\left(\mathbf{W}_{\text{Kendall}}^{(n)} \right)_{\ell_1 \ell_2} = \binom{n}{2}^{-1} \sum_{i < i'} \text{sign} \left((\mathbf{F}_{1;\pm}(\mathbf{X}_{1i}) - \mathbf{F}_{1;\pm}(\mathbf{X}_{1i'}))_{\ell_1} \right) \times \text{sign} \left((\mathbf{F}_{2;\pm}(\mathbf{X}_{2i}) - \mathbf{F}_{2;\pm}(\mathbf{X}_{2i'}))_{\ell_2} \right).$$

It follows from Application 9(d) in [Hoeffding \(1948\)](#) that

$$\left(\mathbf{W}_{\text{Kendall}}^{(n)} \right)_{\ell_1 \ell_2} = \frac{2}{n} \sum_{i=1}^n \left\{ 2F_{1\ell_1} \left((\mathbf{F}_{1;\pm}(\mathbf{X}_{1i}))_{\ell_1} \right) - 1 \right\} \left\{ 2F_{2\ell_2} \left((\mathbf{F}_{2;\pm}(\mathbf{X}_{2i}))_{\ell_2} \right) - 1 \right\} + o_{\text{q.m.}}(n^{-1/2}), \quad (23)$$

where $F_{k\ell}$ denotes the cumulative distribution functions of $(\mathbf{F}_{k;\pm}(\mathbf{X}_k))_{\ell}$, $k = 1, 2$, $\ell = 1, \dots, d_k$.

We also have the Hájek projection of combinatorial statistics (see, e.g., page 242 of [Barbour and Eagleson \(1986\)](#) and Chapter II.3.1 of [Hájek and Šidák \(1967\)](#)) for

$$\left(\mathbf{W}_{\text{Kendall}}^{(n)}\right)_{\ell_1 \ell_2} = \binom{n}{2}^{-1} \sum_{i < i'} \text{sign}\left(\left(\mathbf{F}_{1;\pm}^{(n)}(\mathbf{X}_{1i}) - \mathbf{F}_{1;\pm}^{(n)}(\mathbf{X}_{1i'})\right)_{\ell_1}\right) \times \text{sign}\left(\left(\mathbf{F}_{2;\pm}^{(n)}(\mathbf{X}_{2i}) - \mathbf{F}_{2;\pm}^{(n)}(\mathbf{X}_{2i'})\right)_{\ell_2}\right),$$

which implies

$$\left(\mathbf{W}_{\text{Kendall}}^{(n)}\right)_{\ell_1 \ell_2} = \frac{2}{n} \sum_{i=1}^n \left\{ 2F_{1\ell_1;\text{mid}}^{(n)}\left(\left(\mathbf{F}_{1;\pm}^{(n)}(\mathbf{X}_{1i})\right)_{\ell_1}\right) - 1 \right\} \left\{ 2F_{2\ell_2;\text{mid}}^{(n)}\left(\left(\mathbf{F}_{2;\pm}^{(n)}(\mathbf{X}_{2i})\right)_{\ell_2}\right) - 1 \right\} + o_{\text{q.m.}}(n^{-1/2}), \quad (24)$$

where $F_{k\ell;\text{mid}}^{(n)}$ denotes the *mid-cumulative distribution function*¹⁰ ([Parzen, 2004](#)) of $(\mathbf{F}_{k;\pm}^{(n)}(\mathbf{X}_k))_{\ell}$, $k = 1, 2$, $\ell = 1, \dots, d_k$.

Finally, it follows along the same lines as in the proof for part (1) that the difference between the right-hand sides of (24) and (23) is $o_{\text{q.m.}}(n^{-1/2})$ as $n \rightarrow \infty$. This concludes that

$$\left(\mathbf{W}_{\text{Kendall}}^{(n)}\right)_{\ell_1 \ell_2} - \left(\mathbf{W}_{\text{Kendall}}^{(n)}\right)_{\ell_1 \ell_2} = o_{\text{q.m.}}(n^{-1/2}),$$

which completes the proof. $\square$

Proof of Lemma 4.2. Write $\mathbf{P}_{\delta}^{\mathbf{X}}$ for the distribution $\mathbf{P}_{\delta}^{\mathbf{X}}(\mathbf{P}_1, \mathbf{P}_2, \mathbf{M}_1, \mathbf{M}_2)$ of $\mathbf{X}$. It follows from the quadratic mean differentiability of $f_1^{1/2}$ and $f_2^{1/2}$ and the differentiability with respect to δ of $\mathbf{M}_{\delta}$ that, denoting by $[\mathbf{V}]_1$ and $[\mathbf{V}]_2$, respectively, the first d_1 and last d_2 components of a d -dimensional vector $\mathbf{V}$,

$$\delta \mapsto \left(\frac{d\mathbf{P}_{\delta}^{\mathbf{X}}}{d\mu_d}(\mathbf{x})\right)^{1/2} = \left(|\det(\mathbf{M}_{\delta})|^{-1}\right)^{1/2} f_1([\mathbf{M}_{\delta}^{-1}\mathbf{x}]_1) f_2([\mathbf{M}_{\delta}^{-1}\mathbf{x}]_2) \quad \delta \in \mathbb{R}, \mathbf{x} \in \mathbb{R}^d$$

also is differentiable in quadratic mean. The quadratic expansion of the log-likelihood ratio

$$\delta \mapsto \log \frac{d\mathbf{P}_{\delta+n^{-1/2}\tau}^{(n)}}{d\mathbf{P}_{\delta}^{(n)}}(\mathbf{X}^{(n)})$$

follows (see, e.g., Theorem 12.2.3 (i) in [Lehmann and Romano \(2005\)](#)), yielding, at $\delta = 0$, the second-order asymptotic representation (15). The explicit forms of the central sequence $\Delta^{(n)}(\mathbf{X}^{(n)})$ and the Fisher information γ^2 for $\delta = 0$ are obtained via elementary differentiation. The asymptotic normality result for $\Delta^{(n)}(\mathbf{X}^{(n)})$ follows from part (ii) of the same Theorem 12.2.3. $\square$

Proof of Theorem 4.3. We only give the proof for $T_J^{(n)}$; the proof for $T_{\text{Kendall}}^{(n)}$ is similar and hence is omitted. Applying the multivariate central limit theorem ([Bhattacharya and Ranga Rao, 1986](#), Equation (18.24)) to the asymptotic form of $\Lambda^{(n)}(\mathbf{X}^{(n)})$ (see Lemma 3.1), we obtain, under the null hypothesis ($\delta = 0$),

$$\left(n^{1/2} \text{vec}(\mathbf{W}_J^{(n)}), \Lambda^{(n)}(\mathbf{X}^{(n)})\right) \rightsquigarrow N_{d_1 d_2 + 1} \left(\begin{pmatrix} \mathbf{0}_{d_1 d_2} \\ -\frac{1}{2} \tau^2 \gamma^2 \end{pmatrix}, \begin{pmatrix} \sigma_J^2 \mathbf{I}_{d_1 d_2} & \tau \mathbf{v} \\ \tau \mathbf{v}' & \tau^2 \gamma^2 \end{pmatrix} \right) \quad \text{as } n \rightarrow \infty$$

¹⁰The *mid-cumulative distribution function* of a random variable X is defined as $F_{\text{mid}}(x) := [\mathbf{P}(X \leq x) + \mathbf{P}(X < x)]/2$.

where $\sigma_J^2 := \sigma_{J_1}^2 \sigma_{J_2}^2 / (d_1 d_2)$ and

$$\begin{aligned} \mathbf{v} &:= \text{Cov}_{H_0} \left[\text{vec} \left(\mathbf{J}_1(\mathbf{F}_{1;\pm}(\mathbf{X}_1)) \mathbf{J}_2(\mathbf{F}_{2;\pm}(\mathbf{X}_2))' \right), \right. \\ &\quad \left. \mathbf{X}_1' \mathbf{M}_2' \boldsymbol{\varphi}_2(\mathbf{X}_2) + \mathbf{X}_2' \mathbf{M}_1' \boldsymbol{\varphi}_1(\mathbf{X}_1) - \left(\mathbf{X}_1' \boldsymbol{\varphi}_1(\mathbf{X}_1) - d_1 \right) - \left(\mathbf{X}_2' \boldsymbol{\varphi}_2(\mathbf{X}_2) - d_2 \right) \right] \\ &= E_{H_0} \left[\text{vec} \left(\mathbf{J}_1(\mathbf{F}_{1;\pm}(\mathbf{X}_1)) \mathbf{J}_2(\mathbf{F}_{2;\pm}(\mathbf{X}_2))' \right) \left(\mathbf{X}_1' \mathbf{M}_2' \boldsymbol{\varphi}_2(\mathbf{X}_2) + \mathbf{X}_2' \mathbf{M}_1' \boldsymbol{\varphi}_1(\mathbf{X}_1) \right) \right] \\ &= \text{vec} \left(E_{H_0} \left[\mathbf{J}_1(\mathbf{F}_{1;\pm}(\mathbf{X}_1)) \mathbf{R} \mathbf{J}_2(\mathbf{F}_{2;\pm}(\mathbf{X}_2))' \right] \right). \end{aligned}$$

Thus, by Lemma 4.2,

$$\left(n^{1/2} \text{vec}(\mathbf{W}_J^{(n)}), \Lambda^{(n)}(\mathbf{X}^{(n)}) \right) \rightsquigarrow N_{d_1 d_2 + 1} \left(\begin{pmatrix} \mathbf{0}_{d_1 d_2} \\ -\frac{1}{2} \tau^2 \gamma^2 \end{pmatrix}, \begin{pmatrix} \sigma_J^2 \mathbf{I}_{d_1 d_2} & \tau \mathbf{v} \\ \tau \mathbf{v}' & \tau^2 \gamma^2 \end{pmatrix} \right) \quad \text{as } n \rightarrow \infty.$$

Le Cam's third lemma (Hájek and Šidák, 1967, Chapter VI.1.4 then yields, under local alternatives ($\delta = n^{-1/2} \tau$),

$$n^{1/2} \text{vec}(\mathbf{W}_J^{(n)}) \rightsquigarrow N_{d_1 d_2}(\tau \mathbf{v}, \sigma_J^2 \mathbf{I}_{d_1 d_2}) \quad \text{as } n \rightarrow \infty.$$

The result follows. $\square$

Proof of Proposition 4.4. Denoting $\mathbf{Y}_k := \boldsymbol{\Sigma}_k^{-1/2} \mathbf{X}_k$, and recalling $\tilde{F}_k$ as the cumulative distribution function of $\|\mathbf{Y}_k\|$, $k = 1, 2$, direct computation yields

$$\begin{aligned} &E_{H_0} \left[\mathbf{J}_1(\mathbf{F}_{1;\pm}(\mathbf{X}_1)) \left(\mathbf{X}_1' \mathbf{M}_2' \boldsymbol{\varphi}_2(\mathbf{X}_2) \right) \mathbf{J}_2(\mathbf{F}_{2;\pm}(\mathbf{X}_2))' \right] \\ &= E_{H_0} \left[\mathbf{J}_1 \left(\frac{\mathbf{Y}_1}{\|\mathbf{Y}_1\|} \tilde{F}_1(\|\mathbf{Y}_1\|) \right) \left(\mathbf{Y}_1' \boldsymbol{\Sigma}_1^{1/2} \mathbf{M}_2' \boldsymbol{\varphi}_2(\boldsymbol{\Sigma}_2^{1/2} \mathbf{Y}_2) \right) \mathbf{J}_2 \left(\frac{\mathbf{Y}_2}{\|\mathbf{Y}_2\|} \tilde{F}_2(\|\mathbf{Y}_2\|) \right)' \right] \\ &= E_{H_0} \left[\frac{\mathbf{Y}_1}{\|\mathbf{Y}_1\|} J_1 \left(\tilde{F}_1(\|\mathbf{Y}_1\|) \right) \left(\mathbf{Y}_1' \boldsymbol{\Sigma}_1^{1/2} \mathbf{M}_2' \boldsymbol{\Sigma}_2^{-1/2} \frac{\mathbf{Y}_2}{\|\mathbf{Y}_2\|} \rho_2(\|\mathbf{Y}_2\|) \right) J_2 \left(\tilde{F}_2(\|\mathbf{Y}_2\|) \right) \frac{\mathbf{Y}_2'}{\|\mathbf{Y}_2\|} \right] \\ &= E_{H_0} \left[\frac{\mathbf{Y}_1}{\|\mathbf{Y}_1\|} J_1 \left(\tilde{F}_1(\|\mathbf{Y}_1\|) \right) \mathbf{Y}_1' \right] \boldsymbol{\Sigma}_1^{1/2} \mathbf{M}_2' \boldsymbol{\Sigma}_2^{-1/2} E_{H_0} \left[\frac{\mathbf{Y}_2}{\|\mathbf{Y}_2\|} \rho_2(\|\mathbf{Y}_2\|) J_2 \left(\tilde{F}_2(\|\mathbf{Y}_2\|) \right) \frac{\mathbf{Y}_2'}{\|\mathbf{Y}_2\|} \right]. \end{aligned}$$

Write $r_k := \|\mathbf{Y}_k\|$ and $\mathbf{U}_k := \mathbf{Y}_k / \|\mathbf{Y}_k\|$, $k = 1, 2$. We have, by the independence between r_k and $\mathbf{U}_k$ (Fang, Kotz, and Ng, 1990, Theorem 2.3),

$$\begin{aligned} E_{H_0} \left[\frac{\mathbf{Y}_1}{\|\mathbf{Y}_1\|} J_1 \left(\tilde{F}_1(\|\mathbf{Y}_1\|) \right) \mathbf{Y}_1' \right] &= E_{H_0} \left[\mathbf{U}_1 J_1 \left(\tilde{F}_1(r_1) \right) r_1 \mathbf{U}_1' \right] \\ &= E_{H_0} \left[J_1 \left(\tilde{F}_1(r_1) \right) r_1 \right] \cdot E_{H_0} \left[\mathbf{U}_1 \mathbf{U}_1' \right] = D_1 d_1^{-1} \mathbf{I}_{d_1}, \end{aligned}$$

and similarly,

$$E_{H_0} \left[\frac{\mathbf{Y}_2}{\|\mathbf{Y}_2\|} \rho_2(\|\mathbf{Y}_2\|) J_2 \left(\tilde{F}_2(\|\mathbf{Y}_2\|) \right) \frac{\mathbf{Y}_2'}{\|\mathbf{Y}_2\|} \right] = E_{H_0} \left[\rho_2(r_2) J_2 \left(\tilde{F}_2(r_2) \right) \right] \cdot E_{H_0} \left[\mathbf{U}_2 \mathbf{U}_2' \right] = C_2 d_2^{-1} \mathbf{I}_{d_2},$$

where $E_{H_0} \left[\mathbf{U}_k \mathbf{U}_k' \right] = d_k^{-1} \mathbf{I}_{d_k}$, $k = 1, 2$; see Theorem 2.7 in Fang, Kotz, and Ng (1990). Therefore,

$$E_{H_0} \left[\mathbf{J}_1(\mathbf{F}_{1;\pm}(\mathbf{X}_1)) \left(\mathbf{X}_1' \mathbf{M}_2' \boldsymbol{\varphi}_2(\mathbf{X}_2) \right) \mathbf{J}_2(\mathbf{F}_{2;\pm}(\mathbf{X}_2))' \right] = d_1^{-1} d_2^{-1} D_1 C_2 \boldsymbol{\Sigma}_1^{1/2} \mathbf{M}_2' \boldsymbol{\Sigma}_2^{-1/2}, \quad (25)$$

and similarly,

$$E_{H_0} \left[\mathbf{J}_1(\mathbf{F}_{1;\pm}(\mathbf{X}_1)) \left(\mathbf{X}_2' \mathbf{M}_1' \boldsymbol{\varphi}_1(\mathbf{X}_1) \right) \mathbf{J}_2(\mathbf{F}_{2;\pm}(\mathbf{X}_2))' \right] = d_1^{-1} d_2^{-1} D_2 C_1 \boldsymbol{\Sigma}_1^{-1/2} \mathbf{M}_1 \boldsymbol{\Sigma}_2^{1/2}. \quad (26)$$

Plugging (25) and (26) into (16) yields that, under the alternative $\mathbf{X} \sim P_{n^{-1/2}\tau}^{\text{ell}}(\phi_1, \phi_2, \Sigma_1, \Sigma_2; \mathbf{M}_1, \mathbf{M}_2)$, the limiting distribution of the test statistic $\tilde{T}_J^{(n)}$ is noncentral chi-square with $d_1 d_2$ degrees of freedom and noncentrality parameter

$$\frac{\tau^2}{d_1 d_2 \sigma_{J_1}^2 \sigma_{J_2}^2} \left\| D_1 C_2 \Sigma_1^{1/2} \mathbf{M}_2' \Sigma_2^{-1/2} + D_2 C_1 \Sigma_1^{-1/2} \mathbf{M}_1 \Sigma_2^{1/2} \right\|_{\text{F}}^2,$$

and recall that the limiting distribution of Wilks' statistic is noncentral chi-square with $d_1 d_2$ degrees of freedom and noncentrality parameter of (17). The first result (18) then follows from the fact that (see, e.g., Hannan (1956, Equation (5))) when two test statistics are asymptotically noncentral chi-squared distributed under a local sequence of alternatives (here, $P_{n^{-1/2}\tau}^{(n)}$), their Pitman asymptotic relative efficiencies are obtained as the ratios of their noncentrality parameters. Claims (i) and (ii) follow from the proofs of Propositions 1 and 2 in Hallin and Paindaveine (2008); see also Theorem 1 in Paindaveine (2004) and Proposition 7 in Hallin and Paindaveine (2002a). $\square$

Funding

Hongjian Shi and Mathias Drton were supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 883818). Marc Hallin acknowledges the support of the Czech Science Foundation grant GAČR22036365. Fang Han was supported by the United States NSF grants DMS-1712536 and SES-2019363.

Supplement

The supplement consists of auxiliary results and numerical experiments.

References

- Azadkia, M. and Chatterjee, S. (2021). A simple measure of conditional dependence. *Ann. Statist.*, 49(6):3070–3102.
- Barbour, A. D. and Eagleson, G. K. (1986). Random association of symmetric arrays. *Stochastic Anal. Appl.*, 4(3):239–281.
- Bhattacharya, R. N. and Ranga Rao, R. (1986). *Normal Approximation and Asymptotic Expansions* (Rpt. ed.). Robert E. Krieger Publishing Co., Inc., Melbourne, FL.
- Blomqvist, N. (1950). On a measure of dependence between two random variables. *Ann. Math. Statist.*, 21(4):593–600.
- Carlier, G., Chernozhukov, V., and Galichon, A. (2016). Vector quantile regression: an optimal transport approach. *Ann. Statist.*, 44(3):1165–1192.
- Carlier, G., Chernozhukov, V., and Galichon, A. (2017). Vector quantile regression beyond the specified case. *J. Multivariate Anal.*, 161:96–102.

- Chernoff, H. and Savage, I. R. (1958). Asymptotic normality and efficiency of certain nonparametric test statistics. *Ann. Math. Statist.*, 29(4):972–994.
- Chernozhukov, V., Galichon, A., Hallin, M., and Henry, M. (2017). Monge-Kantorovich depth, quantiles, ranks and signs. *Ann. Statist.*, 45(1):223–256.
- Deb, N., Bhattacharya, B. B., and Sen, B. (2021). Efficiency lower bounds for distribution-free Hotelling-type two-sample tests based on optimal transport. Available at [arXiv:2104.01986v2](https://arxiv.org/abs/2104.01986v2).
- Deb, N. and Sen, B. (2023). Multivariate rank-based distribution-free nonparametric testing using measure transportation. *J. Amer. Statist. Assoc.*, 118(541):192–207.
- del Barrio, E., Cuesta-Albertos, J. A., Hallin, M., and Matrán, C. (2018). Smooth cyclically monotone interpolation and empirical center-outward distribution functions. Available at [arXiv:1806.01238v1](https://arxiv.org/abs/1806.01238v1).
- del Barrio, E., González-Sanz, A., and Hallin, M. (2020). A note on the regularity of optimal-transport-based center-outward distribution and quantile functions. *J. Multivariate Anal.*, 180:104671, 13.
- del Barrio, E., González-Sanz, A., and Hallin, M. (2022). Nonparametric multiple-output center-outward quantile regression. Available at [arXiv:2204.11756v2](https://arxiv.org/abs/2204.11756v2).
- Fang, K. T., Kotz, S., and Ng, K. W. (1990). *Symmetric Multivariate and Related Distributions*, volume 36 of *Monographs on Statistics and Applied Probability*. Chapman and Hall, Ltd., London, England.
- Figalli, A. (2018). On the continuity of center-outward distribution and quantile functions. *Non-linear Anal.*, 177(part B):413–421.
- Friedman, J. H. and Rafsky, L. C. (1983). Graph-theoretic measures of multivariate association and prediction. *Ann. Statist.*, 11(2):377–391.
- Garel, B. and Hallin, M. (1995). Local asymptotic normality of multivariate ARMA processes with a linear trend. *Ann. Inst. Statist. Math.*, 47(3):551–579.
- Ghosal, P. and Sen, B. (2022). Multivariate ranks and quantiles using optimal transport: consistency, rates and nonparametric testing. *Ann. Statist.*, 50(2):1012–1037.
- Gieser, P. W. (1993). *A new nonparametric test for independence between two sets of variates*. PhD thesis, University of Florida.
- Gieser, P. W. and Randles, R. H. (1997). A nonparametric test of independence between two vectors. *J. Amer. Statist. Assoc.*, 92(438):561–567.
- Gretton, A., Bousquet, O., Smola, A., and Schölkopf, B. (2005). Measuring statistical dependence with Hilbert-Schmidt norms. In *Algorithmic Learning Theory*, volume 3734 of *Lecture Notes in Comput. Sci.*, pages 63–77. Springer-Verlag Berlin Heidelberg, Berlin, Germany.

- Hájek, J. and Šidák, Z. (1967). *Theory of Rank Tests*. Academic Press, New York-London; Academia Publishing House of the Czechoslovak Academy of Sciences, Prague.
- Hallin, M. (1994). On the Pitman non-admissibility of correlogram-based methods. *J. Time Ser. Anal.*, 15(6):607–611.
- Hallin, M. (2017). On distribution and quantile functions, ranks and signs in $\mathbb{R}^d$: a measure transportation approach. Available at <https://ideas.repec.org/p/eca/wpaper/2013-258262.html>.
- Hallin, M. (2022). Measure transportation and statistical decision theory. *Annu. Rev. Stat. Appl.*, 9:401–424.
- Hallin, M., del Barrio, E., Cuesta-Albertos, J., and Matrán, C. (2021). Distribution and quantile functions, ranks and signs in dimension d : A measure transportation approach. *Ann. Statist.*, 49(2):1139–1165.
- Hallin, M., Hlubinka, D., and Hudecová, Š. (2023). Efficient fully distribution-free center-outward rank tests for multiple-output regression and MANOVA. *J. Amer. Statist. Assoc.*, 118(543):1923–1939.
- Hallin, M., La Vecchia, D., and Liu, H. (2022). Center-outward R-estimation for semiparametric VARMA models. *J. Amer. Statist. Assoc.*, 117(538):925–938.
- Hallin, M., La Vecchia, D., and Liu, H. (2023). Rank-based testing for semiparametric VAR models: a measure transportation approach. *Bernoulli*, 29(1):229–273.
- Hallin, M. and Mordant, G. (2023). On the finite-sample performance of measure-transportation-based multivariate rank tests. In Yi, M. and Nordhausen, K., editors, *Robust and Multivariate Statistical Methods: Festschrift in Honor of David E. Tyler*, pages 87–119. Springer International Publishing, Cham.
- Hallin, M. and Paindaveine, D. (2002a). Optimal procedures based on interdirections and pseudo-Mahalanobis ranks for testing multivariate elliptic white noise against ARMA dependence. *Bernoulli*, 8(6):787–815.
- Hallin, M. and Paindaveine, D. (2002b). Optimal tests for multivariate location based on interdirections and pseudo-Mahalanobis ranks. *Ann. Statist.*, 30(4):1103–1133.
- Hallin, M. and Paindaveine, D. (2008). Chernoff-Savage and Hodges-Lehmann results for Wilks’ test of multivariate independence. In *Beyond Parametrics in Interdisciplinary Research: Festschrift in Honor of Professor Pranab K. Sen*, volume 1 of *Inst. Math. Stat. (IMS) Collect.*, pages 184–196. Inst. Math. Statist., Beachwood, OH.
- Hallin, M. and Tribel, O. (2000). The efficiency of some nonparametric rank-based competitors to correlogram methods. In *Game Theory, Optimal Stopping, Probability and Statistics*, volume 35 of *IMS Lecture Notes Monogr. Ser.*, pages 249–262. Inst. Math. Statist., Beachwood, OH.

- Hannan, E. J. (1956). The asymptotic powers of certain tests based on multiple correlations. *J. Roy. Statist. Soc. Ser. B*, 18(2):227–233.
- Heller, R., Gorfine, M., and Heller, Y. (2012). A class of multivariate distribution-free tests of independence based on graphs. *J. Statist. Plann. Inference*, 142(12):3097–3106.
- Heller, R., Heller, Y., and Gorfine, M. (2013). A consistent multivariate test of association based on ranks of distances. *Biometrika*, 100(2):503–510.
- Hodges, Jr., J. L. and Lehmann, E. L. (1956). The efficiency of some nonparametric competitors of the t -test. *Ann. Math. Statist.*, 27(2):324–335.
- Hoeffding, W. (1948). A class of statistics with asymptotically normal distribution. *Ann. Math. Statist.*, 19(3):293–325.
- Hoeffding, W. (1951). A combinatorial central limit theorem. *Ann. Math. Statist.*, 22(4):558–566.
- Kendall, M. G. (1938). A new measure of rank correlation. *Biometrika*, 30(1/2):81–93.
- Kim, I., Balakrishnan, S., and Wasserman, L. (2020). Robust multivariate nonparametric tests via projection averaging. *Ann. Statist.*, 48(6):3417–3441.
- Konijn, H. S. (1956). On the power of certain tests for independence in bivariate populations. *Ann. Math. Statist.*, 27(2):300–323.
- Lehmann, E. L. and Romano, J. P. (2005). *Testing Statistical Hypotheses* (3rd ed.). Springer Texts in Statistics. Springer, New York.
- Lin, J. (2017). *Copula versions of RKHS-based and distance-based criteria*. PhD thesis, Pennsylvania State University.
- Lin, Z. and Han, F. (2023). On boosting the power of Chatterjee’s rank correlation. *Biometrika*, 110(2):283–299.
- Lind, B. and Roussas, G. (1972). A remark on quadratic mean differentiability. *Ann. Math. Statist.*, 43(3):1030–1034.
- Liu, R. Y. and Singh, K. (1993). A quality index based on data depth and multivariate rank tests. *J. Amer. Statist. Assoc.*, 88(421):252–260.
- McCann, R. J. (1995). Existence and uniqueness of monotone measure-preserving maps. *Duke Math. J.*, 80(2):309–323.
- Moon, H. and Chen, K. (2022). Interpoint-ranking sign covariance for the test of independence. *Biometrika*, 109(1):165–179.
- Mordant, G. (2021). *Transporting probability measures: some contributions to statistical inference*. PhD thesis, UCL-Université Catholique de Louvain.

- Oja, H. (2010). *Multivariate Nonparametric Methods with R: An Approach Based on Spatial Signs and Ranks*, volume 199 of *Lecture Notes in Statistics*. Springer, New York.
- Oja, H., Paindaveine, D., and Taskinen, S. (2016). Affine-invariant rank tests for multivariate independence in independent component models. *Electron. J. Stat.*, 10(2):2372–2419.
- Paindaveine, D. (2004). A unified and elementary proof of serial and nonserial, univariate and multivariate, Chernoff-Savage results. *Stat. Methodol.*, 1(1-2):81–91.
- Parzen, E. (2004). Quantile probability and statistical data modeling. *Statist. Sci.*, 19(4):652–662.
- Pillai, K. C. S. (1955). Some new test criteria in multivariate analysis. *Ann. Math. Statist.*, 26:117–121.
- Puri, M. L. and Sen, P. K. (1971). *Nonparametric Methods in Multivariate Analysis*. John Wiley & Sons, Inc., New York-London-Sydney.
- Randles, R. H. (1989). A distribution-free multivariate sign test based on interdirections. *J. Amer. Statist. Assoc.*, 84(408):1045–1050.
- Shi, H., Drton, M., Hallin, M., and Han, F. (2024). Supplement to “Distribution-free tests of multivariate independence based on center-outward quadrant, Spearman, kendall, and van der Waerden statistics”. *Bernoulli*. To appear.
- Shi, H., Drton, M., and Han, F. (2022). Distribution-free consistent independence tests via center-outward ranks and signs. *J. Amer. Statist. Assoc.*, 117(537):395–410.
- Shi, H., Hallin, M., Drton, M., and Han, F. (2022). On universally consistent and fully distribution-free rank tests of vector independence. *Ann. Statist.*, 50(4):1933–1959.
- Shorack, G. R. (2017). *Probability for Statisticians* (2nd ed.). Springer Texts in Statistics. Springer, Cham, Switzerland.
- Spearman, C. (1904). The proof and measurement of association between two things. *Amer. J. Psychol.*, 15(1):72–101.
- Székely, G. J. and Rizzo, M. L. (2014). Partial distance correlation with methods for dissimilarities. *Ann. Statist.*, 42(6):2382–2412.
- Székely, G. J., Rizzo, M. L., and Bakirov, N. K. (2007). Measuring and testing dependence by correlation of distances. *Ann. Statist.*, 35(6):2769–2794.
- Taskinen, S., Kankainen, A., and Oja, H. (2003). Sign test of independence between two random vectors. *Statist. Probab. Lett.*, 62(1):9–21.
- Taskinen, S., Kankainen, A., and Oja, H. (2004). Rank scores tests of multivariate independence. In *Theory and applications of recent robust methods*, Stat. Ind. Technol., pages 329–341. Birkhäuser, Basel.

- Taskinen, S., Oja, H., and Randles, R. H. (2005). Multivariate nonparametric tests of independence. *J. Amer. Statist. Assoc.*, 100(471):916–925.
- Weihs, L., Drton, M., and Meinshausen, N. (2018). Symmetric rank covariances: a generalized framework for nonparametric measures of dependence. *Biometrika*, 105(3):547–562.
- Wilcoxon, F. (1945). Individual comparisons by ranking methods. *Biometrics Bulletin*, 1(6):80–83.
- Wilks, S. S. (1935). On the independence of k sets of normally distributed statistical variables. *Econometrica*, 3(3):309–326.
- Zhu, L., Xu, K., Li, R., and Zhong, W. (2017). Projection correlation between two random vectors. *Biometrika*, 104(4):829–843.
- Zuo, Y. and He, X. (2006). On the limiting distributions of multivariate depth-based rank sum statistics and related tests. *Ann. Statist.*, 34(6):2879–2896.

Supplement to “Distribution-free tests of multivariate independence based on center-outward quadrant, Spearman, Kendall, and van der Waerden statistics”

Hongjian Shi^{*} Mathias Drton[†] Marc Hallin[‡] and Fang Han[§]

S.1 Auxiliary results

S.1.1 Further analysis of the motivating example

Supporting our claim that forty observations three months apart can be considered as independent, Figure S.1 provides correlogram and cross-correlogram plots for the series in the motivating example of Section 1.3. Observe that, except for lag zero, essentially all auto- and cross-correlations are non-significant at level 5%, and that no clear seasonality pattern emerges. Therefore, treating these observations as independent and identically distributed is unlikely to affect the validity of the tests.

S.1.2 Finite-sample size of Wilks’ likelihood ratio tests

Likelihood ratio tests, even asymptotically valid, often suffer from severe size inflation when the observations are not Gaussian, and Wilks’ test is no exception. While this is hardly surprising, the magnitude of this inflation is typically underestimated; Table S.1 provides a brief but quite alarming simulation-based illustration of the phenomenon for various dimensions d_1 and d_2 . Thanks to distribution-freeness, our rank-based tests are not affected, which constitutes another strong argument in their favor.

The simulated data are samples (size $n = 100$) of independent copies of random vectors of the form $\mathbf{X} = (\mathbf{X}'_1, \mathbf{X}'_2)'$ where $\mathbf{X}_1 = \exp(\mathbf{Z}_1)$ (values in $\mathbb{R}^{d_1}$), $\mathbf{X}_2 = \exp(\mathbf{Z}_2)$ (values in $\mathbb{R}^{d_2}$), with $(\mathbf{Z}_1, \mathbf{Z}_2) \sim N(\mathbf{0}, \sigma^2 \mathbf{I}_{d_1+d_2})$ and $\exp(\mathbf{Z}) := (\exp(Z_1), \dots, \exp(Z_d))'$. The empirical sizes of Wilks’ test with nominal level 0.05, based on 10,000 simulations, are reported in Table S.1 for dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$ and $\sigma \in \{0.5, 1.0, 1.5, 2.0\}$. Note that σ , a standard error for the Gaussian $\mathbf{Z}_1$ and $\mathbf{Z}_2$, is a skewness parameter for the lognormal components of $\mathbf{X}_1$ and $\mathbf{X}_2$. As one might expect, the size spectacularly grows beyond the nominal level 0.05 for increasing skewness σ and/or increasing dimensions d_1 and d_2 , reaching 0.317 for $d_1 = d_2 = 7$ and $\sigma = 2.0$!

^{*}Department of Mathematics, TUM School of Computation, Information and Technology, Technical University of Munich, 85748 Garching bei München, Germany; e-mail: hongjian.shi@tum.de

[†]Department of Mathematics, TUM School of Computation, Information and Technology, Technical University of Munich, 85748 Garching bei München, Germany; e-mail: mathias.drton@tum.de

[‡]ECARES and Department of Mathematics, Université Libre de Bruxelles, Brussels, Belgium; email: mhallin@ulb.ac.be

[§]Department of Statistics, University of Washington, Seattle, WA 98195, USA; e-mail: fanghan@uw.edu




Figure S.1: Autocorrelation (on the diagonal) and cross-correlation plots (below the diagonal), with 95% confidence bands, for the forty observations of daily log returns of the stocks “AT&T Inc [T]”, “Verizon Communications [VZ]”, “Du Pont (E.I.) [DD]”, “Dow Chemical [DOW]”, “Freeport-McMoran Cp & Gld [FCX]”, and “Monsanto Co. [MON].”

Table S.1: Sizes of Wilks' LRT for log-normal distributions at nominal significance level 0.05 for various dimensions d_1 and d_2 and various values of the skewness parameter σ .

	$\sigma = 0.5$	$\sigma = 1.0$	$\sigma = 1.5$	$\sigma = 2.0$
$(d_1, d_2) = (2, 2)$	0.059	0.079	0.101	0.102
$(d_1, d_2) = (3, 3)$	0.067	0.101	0.134	0.155
$(d_1, d_2) = (5, 5)$	0.095	0.154	0.211	0.243
$(d_1, d_2) = (7, 7)$	0.133	0.197	0.272	0.317

Wilks' test, thus, is highly unreliable under non-Gaussian distributions and its conclusions should be interpreted with highest precaution.

S.1.3 The elliptical Hodges–Lehmann lower bound $\Omega(d_1, d_2)$

Table S.2 provides, for dimensions $d_1, d_2 \leq 10$, numerical values of the elliptical Hodges–Lehmann lower bounds $\Omega(d_1, d_2)$ (Proposition 4.4) for the AREs of Spearman center-outward rank tests (based on Wilcoxon scores) with respect to Wilks' test.

Table S.2: Some numerical values of the Hodges–Lehmann lower bound $\Omega(d_1, d_2)$ for the Pitman asymptotic relative efficiency, against elliptical Konijn alternatives, of the center-outward Spearman (Wilcoxon score) test with respect to Wilks', for dimensions $d_1, d_2 \leq 10$.

$d_1 \backslash d_2$	1	2	3	4	5	6	7	8	9	10
1	0.856	0.884	0.867	0.850	0.837	0.826	0.817	0.809	0.803	0.797
2	0.884	0.913	0.895	0.878	0.864	0.853	0.844	0.836	0.829	0.823
3	0.867	0.895	0.878	0.861	0.847	0.836	0.827	0.819	0.813	0.807
4	0.850	0.878	0.861	0.845	0.831	0.820	0.811	0.804	0.797	0.792
5	0.837	0.864	0.847	0.831	0.818	0.807	0.799	0.791	0.785	0.779
6	0.826	0.853	0.836	0.820	0.807	0.797	0.788	0.781	0.775	0.769
7	0.817	0.844	0.827	0.811	0.799	0.788	0.779	0.772	0.766	0.761
8	0.809	0.836	0.819	0.804	0.791	0.781	0.772	0.765	0.759	0.754
9	0.803	0.829	0.813	0.797	0.785	0.775	0.766	0.759	0.753	0.748
10	0.797	0.823	0.807	0.792	0.779	0.769	0.761	0.754	0.748	0.742

The bound, which, with a value of 0.913, is quite high for $d_1 = d_2 = 2$, slowly decreases as the dimensions increase. Recall that an ARE of 0.913 means that, in the worst case, Wilks requires 0.913 n observations to achieve the same performance as Spearman with n observations. The worst efficiency loss, thus, is minimal; on the other hand, the same ARE has no upper bound, and Spearman, potentially, can be arbitrarily more powerful than Wilks.

S.2 Numerical experiments

We now present a numerical analysis of the finite-sample performances of the tests that have been introduced in Section 3.3 and some of their competitors.

Eight tests are considered: (i) the center-outward quadrant or sign test $\psi_{\text{sign}}^{(n)}$, (ii) the center-outward Spearman test $\psi_{\text{Spearman}}^{(n)}$, (iii) the center-outward Kendall test $\psi_{\text{Kendall}}^{(n)}$, (iv) the center-

outward van der Waerden test $\psi_{\text{vdW}}^{(n)} = \psi_{\text{JvdW}}^{(n)}$, (v) the center-outward Wilcoxon score version of the distance covariance test (Shi, Drton, and Han, 2022), (vi) the center-outward van der Waerden score version of the distance covariance test (Shi et al., 2022), (vii) Wilks' pseudo-Gaussian test $\psi_{\text{Wilks}}^{(n)}$, and (viii) Székely and Rizzo's permutation-based distance covariance test $\psi_{\text{SzR}}^{(n)}$ (Székely and Rizzo, 2013, 2014).

The data were generated as a sample of n independent copies of the $(d_1 + d_2)$ -dimensional random vector $\mathbf{X} = (\mathbf{X}'_1, \mathbf{X}'_2)' \sim \mathbf{P}_\delta^{\mathbf{X}}(\mathbf{P}_1, \mathbf{P}_2; \mathbf{M}_1, \mathbf{M}_2)$ under the Konijn model (14) with dimensions $d_1 = d_2 = d_0$ with $d_0 \in \{2, 3, 5, 7\}$, matrices $\mathbf{M}_1 = \mathbf{M}_2 = \mathbf{I}_{d_0}$, $\delta = \delta^{(n)} = n^{-1/2}\tau$. The following distributions $\mathbf{P}_1$ and $\mathbf{P}_2$ were considered with $d_1 = d_2$; note that cases (b), (f), (g), (h), (i), and (j) are not elliptical, while the heavy-tailed (d), (e), (g), and (h) fail to satisfy the conditions for the validity of Wilks and other non-distribution-free tests, and thus the computation of AREs.

Gaussian and nearly Gaussian:

- (a) (standard Gaussian) $\mathbf{X}_k^* \stackrel{\text{i.i.d.}}{\sim} N_{d_0}(\mathbf{0}, \mathbf{I}_{d_0})$, $k = 1, 2$;
- (b) (Gaussian mixture) $\mathbf{X}_k^* \stackrel{\text{i.i.d.}}{\sim} 0.5N_{d_0}(\mathbf{0}, \mathbf{I}_{d_0}) + 0.5N_{d_0}(\mathbf{1}, \mathbf{I}_{d_0})$, $k = 1, 2$.

Heavy-tailed:

- (c) (heavy-tailed elliptical t_3) $\mathbf{X}_k^* \stackrel{\text{i.i.d.}}{\sim} t_{3,d_0}(\mathbf{0}, \mathbf{I}_{d_0})$, $k = 1, 2$, where $t_{\nu,d}(\mathbf{0}, \mathbf{\Sigma})$, with $\mathbf{\Sigma} \in \mathbb{R}^{d \times d}$ symmetric positive definite, has density (elliptical distributions with finite second-order moments)

$$\left[\Gamma[(\nu + d)/2] / \Gamma(\nu/2) \nu^{d/2} \pi^{d/2} |\mathbf{\Sigma}|^{1/2} \right] [1 + \nu^{-1} \mathbf{x}' \mathbf{\Sigma}^{-1} \mathbf{x}]^{-(\nu+d)/2}; \quad (\text{S.1})$$

- (d) (heavy-tailed elliptical t_2) $\mathbf{X}_k^* \stackrel{\text{i.i.d.}}{\sim} t_{2,d_0}(\mathbf{0}, \mathbf{I}_{d_0})$, $k = 1, 2$ (elliptical distributions with infinite second-order moments);
- (e) (heavy-tailed elliptical Cauchy) $\mathbf{X}_k^* \stackrel{\text{i.i.d.}}{\sim} t_{1,d_0}(\mathbf{0}, \mathbf{I}_{d_0})$, $k = 1, 2$ (elliptical distributions with infinite second-order moments);
- (f) (heavy-tailed non-elliptical marginal t_3) $X_{k\ell}^* \stackrel{\text{i.i.d.}}{\sim} t_{3,1}(0, 1)$, $k = 1, 2$, $\ell = 1, \dots, d_0$ (heavy-tailed non-elliptical distributions invariant under coordinate permutations with finite second-order moments);
- (g) (heavy-tailed non-elliptical marginal t_2) $X_{k\ell}^* \stackrel{\text{i.i.d.}}{\sim} t_{2,1}(0, 1)$, $k = 1, 2$, $\ell = 1, \dots, d_0$ (heavy-tailed non-elliptical distributions invariant under coordinate permutations with infinite second-order moments);
- (h) (heavy-tailed non-elliptical marginal Cauchy) $X_{k\ell}^* \stackrel{\text{i.i.d.}}{\sim} t_{1,1}(0, 1)$, $k = 1, 2$, $\ell = 1, \dots, d_0$ (heavy-tailed non-elliptical distributions invariant under coordinate permutations with infinite second-order moments).

Skewed:

- (i) (skewed independent χ_2^2 components) $X_{k\ell}^* \stackrel{\text{i.i.d.}}{\sim} \chi_2^2$, $k = 1, 2$, $\ell = 1, \dots, d_0$ (skewed non-elliptical distributions invariant under coordinate permutations with finite second-order moments);
- (j) (skewed independent χ_1^2 components) $X_{k\ell}^* \stackrel{\text{i.i.d.}}{\sim} \chi_1^2$, $k = 1, 2$, $\ell = 1, \dots, d_0$ (skewed non-elliptical distributions invariant under coordinate permutations with finite second-order moments).

Tables S.3–S.12 report empirical powers (rejection frequencies) of the eight tests (i)–(viii) performed at nominal level $\alpha = 0.05$, based on 1,000 simulations of a sample of size $n \in \{432, 864, 1728\}$ from distributions (a)–(j), in dimensions $d_1 = d_2 = d_0 \in \{2, 3, 5, 7\}$, for parameter values $\delta = \delta^{(n)} = n^{-1/2}\tau$, with $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

Overall, the center-outward van der Waerden score test has the best power except when the underlying distribution is Gaussian (case (a)) or nearly Gaussian (case (b)), where Wilks, unsurprisingly, prevails. The performances of the center-outward Spearman and van der Waerden tests and the two distance covariance tests are comparable, while the center-outward sign and Kendall tests perform poorly in some cases, which is not unexpected since the center-outward ranks and signs are not efficiently used. As a benchmark, the permutation-based distance covariance test performs worse than the center-outward van der Waerden test when the sample size is large enough or when the distributions are skewed (case (j)); it may have some advantage, though, when the dimension is large and the sample size relatively small.

Tables S.3 and S.5 confirm the Chernoff–Savage claim of Proposition 4.4(i) that, under ellipticity, the center-outward van der Waerden test uniformly dominates Wilks’ optimal Gaussian procedure, and the Hodges–Lehmann claim of Proposition 4.4(ii) that the center-outward Spearman test has Pitman asymptotic relative efficiency no smaller than 0.77 with respect to Wilks’ test when $d_1, d_2 \leq 7$ (see Table S.2). In the Gaussian case (a) (see Table S.3), the center-outward van der Waerden test is second best out of all competing tests, and the ratio of its rejection frequencies to those of Wilks’ test approaches one as the sample size increases. In the multivariate t_3 case (c) (see Table S.5), the center-outward van der Waerden test is best when the sample size is sufficiently large.

Table S.4 shows the performance of the seven competing tests under normal mixtures, which are non-elliptical but nearly Gaussian. Wilks’ test has the best finite-sample performance compared to all the center-outward sign- and rank-based tests with, however, the center-outward van der Waerden test achieving similar performance for n large enough.

Although cases (d) and (e) are not covered by Proposition 4.4 due to infinite second-order moments, Tables S.6 and S.7 indicate that the center-outward van der Waerden test remains best while, due to heavy-tailedness, Wilks’s test badly suffers from severe over-rejection.

Table S.8 shows the performance of the eight competing tests when the distributions are non-elliptical and heavy-tailed (case (f)). Inspection of the table reveals that, when the sample size is sufficiently large, the center-outward van der Waerden test outperforms all its competitors, while under t_3 marginals, Wilks’s test, although asymptotically valid, suffers from severe size inflation due to heavy-tailedness; the same comment applies in the multivariate t_3 case (c).

Tables S.9–S.10 deal with non-elliptical heavy-tailed distributions with infinite second-order moments. The center-outward van der Waerden test still performs best while Wilks suffers, as in the multivariate t cases (d) and (e) (Tables S.6 and S.7), from severe size inflation due to heavy-tailedness. The only difference between case (e) (multivariate Cauchy; Table S.7) and case (h) (Cauchy marginals; Table S.10), is that the center-outward Kendall test performs poorly under the latter as the dimension increases.

Table S.11–S.12 shows the performance of the eight tests when the distributions are non-elliptical and skewed (cases (i) and (j)). We notice that when the sample size is large enough, the center-outward Spearman and van der Waerden score tests, and the two distance covariance tests perform better than the center-outward sign and Kendall tests; all center-outward sign- and rank-based

tests are much more powerful than Wilks' test. The only exception occurs when the dimension is relatively large and the sample size is small, which can be attributed to the rate of convergence of empirical center-outward ranks and signs, a rate that fastly decreases as the dimension increases.

To conclude this section, we explore mixture alternatives under which the density of $\mathbf{X} = (\mathbf{X}'_1, \mathbf{X}'_2)'$ is of the form

$$f_\delta := (1 - \delta)f_1f_2 + \delta f_0 \quad \delta \in [0, 1].$$

where f_1 and f_2 are the densities of the null distributions P_1 and P_2 of $\mathbf{X}_1$ and $\mathbf{X}_2$, respectively, and f_0 is the density of a distribution P_0 over $\mathbb{R}^{d_1+d_2}$ which does not factorize into the product of its marginals over $\mathbb{R}^{d_1}$ and $\mathbb{R}^{d_2}$. Clearly, the value $\delta = 0$ yields mutually independent vectors $\mathbf{X}_1$ and $\mathbf{X}_2$. For given P_1 , P_2 , and P_0 , the distribution $P^\mathbf{X}$ of $\mathbf{X}$ belongs to a one-parameter family indexed by $\delta \in [0, 1]$. LAN and ARE results for these families could be obtained along the same lines as in Section 4; details are left to the reader. Rather, we performed simulations with the following distributions P_1 , P_2 , and P_0 and the same values of n , $d_1 = d_2 = d_0$, and $\delta^{(n)} = n^{-1/2}\tau$:

- (k) (standard Gaussian) $P_1 = P_2 = N_{d_0}(\mathbf{0}, \mathbf{I}_{d_0})$, and $P_0 = N_{2d_0}(\mathbf{0}, \mathbf{A})$ where $\mathbf{A}$ is a symmetric $2d_0 \times 2d_0$ matrix with entries

$$A_{j\ell} = A_{\ell j} = \begin{cases} 1 & j = \ell, \\ 0.5 & j = \ell + d_0 \text{ or } \ell = j + d_0, \\ 0 & \text{otherwise;} \end{cases}$$

- (l) (heavy-tailed elliptical) $P_1 = P_2 = t_{3,d_0}(\mathbf{0}, \mathbf{I}_{d_0})$, and $P_0 = t_{3,2d_0}(\mathbf{0}, \mathbf{A})$ (recall $t_{\nu,d}(\mathbf{0}, \mathbf{\Sigma})$ stands for the elliptical distribution with the density of (S.1)).

Tables S.13–S.14 report empirical powers (rejection frequencies) of the same seven tests as above, based on 1,000 simulations with nominal significance level $\alpha = 0.05$, sample size $n \in \{432, 864, 1728\}$, dimensions $d_1 = d_2 = d_0 \in \{2, 3, 5, 7\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 1, 2, 3, 4\}$. When the sample size is sufficiently large, the center-outward van der Waerden test exhibits superior performance compared to, or is closely aligned with, both Wilks' and the distance covariance permutation-based test.

References

- Shi, H., Drton, M., and Han, F. (2022). Distribution-free consistent independence tests via center-outward ranks and signs. *J. Amer. Statist. Assoc.*, 117(537):395–410.
- Shi, H., Hallin, M., Drton, M., and Han, F. (2022). On universally consistent and fully distribution-free rank tests of vector independence. *Ann. Statist.*, 50(4):1933–1959.
- Székel, G. J. and Rizzo, M. L. (2013). The distance correlation t -test of independence in high dimension. *J. Multivariate Anal.*, 117:193–213.
- Székel, G. J. and Rizzo, M. L. (2014). Partial distance correlation with methods for dissimilarities. *Ann. Statist.*, 42(6):2382–2412.

Table S.3: Empirical powers, under the standard Gaussian distributions (case (a)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n		432					864					1728				
τ		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.050	0.055	0.086	0.139	0.263	0.045	0.073	0.088	0.149	0.268	0.049	0.061	0.088	0.162	0.262
	Spearman	0.053	0.067	0.107	0.202	0.393	0.044	0.063	0.119	0.215	0.394	0.044	0.060	0.115	0.225	0.409
	Kendall	0.055	0.072	0.105	0.201	0.372	0.053	0.065	0.106	0.206	0.387	0.049	0.058	0.113	0.212	0.403
	vdW	0.051	0.066	0.110	0.207	0.394	0.044	0.063	0.119	0.217	0.404	0.050	0.059	0.117	0.231	0.422
	dCov	0.053	0.067	0.103	0.194	0.372	0.051	0.063	0.111	0.199	0.359	0.044	0.061	0.113	0.216	0.367
	dCvdW	0.054	0.070	0.103	0.195	0.376	0.048	0.068	0.107	0.200	0.368	0.048	0.059	0.111	0.223	0.381
	Wilks	0.052	0.073	0.129	0.243	0.427	0.047	0.066	0.118	0.221	0.427	0.047	0.057	0.116	0.239	0.427
	SzR	0.048	0.061	0.119	0.225	0.379	0.045	0.075	0.104	0.200	0.385	0.046	0.068	0.110	0.200	0.337
$(d_1, d_2) = (3, 3)$	sign	0.045	0.068	0.111	0.163	0.290	0.054	0.059	0.095	0.158	0.284	0.052	0.060	0.107	0.175	0.289
	Spearman	0.050	0.060	0.122	0.210	0.362	0.045	0.067	0.126	0.208	0.386	0.047	0.069	0.108	0.232	0.395
	Kendall	0.047	0.058	0.119	0.215	0.374	0.051	0.066	0.122	0.217	0.369	0.054	0.056	0.116	0.221	0.374
	vdW	0.053	0.064	0.120	0.221	0.395	0.046	0.067	0.128	0.219	0.401	0.049	0.067	0.125	0.239	0.413
	dCov	0.047	0.063	0.106	0.201	0.353	0.054	0.067	0.117	0.200	0.363	0.050	0.066	0.103	0.221	0.355
	dCvdW	0.045	0.067	0.115	0.207	0.371	0.052	0.063	0.120	0.209	0.378	0.053	0.070	0.109	0.225	0.371
	Wilks	0.043	0.071	0.145	0.266	0.483	0.044	0.071	0.139	0.263	0.456	0.057	0.073	0.123	0.259	0.437
	SzR	0.043	0.070	0.113	0.228	0.430	0.051	0.066	0.101	0.219	0.392	0.046	0.059	0.127	0.222	0.427
$(d_1, d_2) = (5, 5)$	sign	0.050	0.060	0.079	0.140	0.243	0.051	0.058	0.087	0.153	0.278	0.045	0.070	0.085	0.167	0.300
	Spearman	0.055	0.061	0.087	0.143	0.264	0.053	0.058	0.100	0.166	0.291	0.047	0.065	0.095	0.187	0.338
	Kendall	0.052	0.065	0.089	0.142	0.266	0.052	0.061	0.095	0.174	0.310	0.057	0.068	0.089	0.178	0.318
	vdW	0.051	0.057	0.085	0.163	0.290	0.053	0.059	0.101	0.178	0.336	0.054	0.063	0.100	0.198	0.371
	dCov	0.055	0.065	0.085	0.138	0.259	0.051	0.060	0.103	0.159	0.289	0.052	0.063	0.089	0.177	0.314
	dCvdW	0.053	0.065	0.081	0.155	0.284	0.049	0.057	0.099	0.176	0.328	0.050	0.063	0.094	0.196	0.355
	Wilks	0.061	0.079	0.138	0.278	0.527	0.045	0.070	0.127	0.245	0.525	0.059	0.064	0.123	0.269	0.501
	SzR	0.047	0.056	0.122	0.237	0.453	0.054	0.058	0.122	0.237	0.451	0.042	0.059	0.127	0.233	0.424
$(d_1, d_2) = (7, 7)$	sign	0.045	0.049	0.080	0.114	0.181	0.041	0.056	0.085	0.126	0.211	0.043	0.053	0.076	0.154	0.245
	Spearman	0.055	0.065	0.091	0.134	0.197	0.059	0.060	0.090	0.139	0.225	0.045	0.059	0.088	0.139	0.228
	Kendall	0.044	0.061	0.081	0.136	0.180	0.047	0.060	0.087	0.132	0.223	0.054	0.059	0.083	0.131	0.230
	vdW	0.041	0.056	0.090	0.135	0.217	0.048	0.055	0.092	0.156	0.259	0.047	0.051	0.085	0.161	0.283
	dCov	0.033	0.059	0.062	0.103	0.158	0.039	0.051	0.075	0.122	0.196	0.043	0.061	0.074	0.118	0.214
	dCvdW	0.043	0.049	0.083	0.123	0.199	0.041	0.055	0.085	0.143	0.249	0.043	0.057	0.078	0.156	0.269
	Wilks	0.056	0.071	0.145	0.298	0.572	0.055	0.067	0.132	0.285	0.523	0.051	0.063	0.135	0.271	0.521
	SzR	0.051	0.057	0.128	0.228	0.478	0.039	0.075	0.140	0.245	0.495	0.040	0.060	0.108	0.234	0.486

Table S.4: Empirical powers, under the Gaussian mixture distributions (case (b)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n	τ	432					864					1728				
		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.050	0.052	0.093	0.145	0.247	0.049	0.058	0.095	0.161	0.262	0.045	0.063	0.093	0.157	0.255
	Spearman	0.045	0.068	0.109	0.217	0.389	0.055	0.069	0.129	0.230	0.403	0.047	0.061	0.114	0.235	0.386
	Kendall	0.049	0.057	0.111	0.213	0.379	0.051	0.065	0.115	0.219	0.380	0.045	0.063	0.116	0.225	0.355
	vdW	0.043	0.066	0.113	0.222	0.394	0.049	0.067	0.121	0.235	0.418	0.041	0.060	0.117	0.243	0.393
	dCov	0.050	0.059	0.098	0.201	0.367	0.047	0.067	0.120	0.224	0.373	0.049	0.061	0.103	0.218	0.345
	dCvdW	0.050	0.057	0.109	0.203	0.374	0.046	0.064	0.127	0.219	0.376	0.045	0.062	0.107	0.216	0.343
	Wilks	0.047	0.064	0.118	0.248	0.426	0.055	0.065	0.130	0.245	0.430	0.051	0.058	0.116	0.235	0.405
	SzR	0.055	0.065	0.121	0.226	0.356	0.030	0.071	0.111	0.214	0.368	0.051	0.066	0.119	0.214	0.374
$(d_1, d_2) = (3, 3)$	sign	0.043	0.054	0.087	0.183	0.294	0.049	0.056	0.096	0.169	0.291	0.045	0.062	0.093	0.179	0.314
	Spearman	0.042	0.069	0.099	0.217	0.369	0.049	0.075	0.129	0.212	0.382	0.047	0.065	0.125	0.239	0.421
	Kendall	0.049	0.066	0.098	0.220	0.357	0.055	0.071	0.113	0.207	0.356	0.044	0.073	0.110	0.218	0.397
	vdW	0.045	0.066	0.105	0.219	0.373	0.048	0.077	0.133	0.214	0.383	0.039	0.064	0.126	0.239	0.439
	dCov	0.043	0.056	0.097	0.201	0.359	0.053	0.073	0.108	0.199	0.353	0.041	0.062	0.105	0.213	0.380
	dCvdW	0.040	0.057	0.093	0.204	0.363	0.050	0.067	0.111	0.200	0.365	0.039	0.058	0.112	0.219	0.391
	Wilks	0.052	0.064	0.122	0.277	0.462	0.058	0.084	0.145	0.263	0.448	0.045	0.057	0.135	0.259	0.463
	SzR	0.042	0.060	0.119	0.249	0.386	0.045	0.066	0.123	0.199	0.379	0.047	0.070	0.099	0.195	0.375
$(d_1, d_2) = (5, 5)$	sign	0.047	0.060	0.082	0.143	0.244	0.049	0.063	0.081	0.151	0.259	0.047	0.058	0.104	0.150	0.328
	Spearman	0.055	0.067	0.086	0.159	0.283	0.046	0.067	0.093	0.167	0.291	0.050	0.059	0.096	0.166	0.341
	Kendall	0.054	0.069	0.087	0.159	0.279	0.054	0.067	0.094	0.173	0.279	0.047	0.053	0.113	0.172	0.321
	vdW	0.049	0.065	0.091	0.163	0.308	0.051	0.071	0.097	0.185	0.315	0.043	0.061	0.113	0.180	0.377
	dCov	0.053	0.061	0.087	0.153	0.258	0.045	0.061	0.097	0.167	0.281	0.048	0.063	0.099	0.165	0.314
	dCvdW	0.055	0.060	0.095	0.169	0.295	0.049	0.063	0.092	0.177	0.299	0.048	0.059	0.115	0.177	0.365
	Wilks	0.059	0.066	0.133	0.291	0.528	0.056	0.069	0.133	0.272	0.491	0.041	0.061	0.121	0.257	0.502
	SzR	0.047	0.070	0.113	0.253	0.410	0.037	0.050	0.109	0.215	0.406	0.038	0.067	0.126	0.221	0.397
$(d_1, d_2) = (7, 7)$	sign	0.047	0.057	0.085	0.097	0.194	0.051	0.053	0.097	0.127	0.237	0.045	0.066	0.073	0.149	0.257
	Spearman	0.055	0.076	0.091	0.109	0.209	0.057	0.070	0.085	0.129	0.227	0.061	0.062	0.086	0.142	0.267
	Kendall	0.055	0.064	0.089	0.113	0.203	0.053	0.064	0.087	0.131	0.226	0.051	0.059	0.083	0.149	0.260
	vdW	0.051	0.062	0.085	0.113	0.225	0.054	0.057	0.095	0.139	0.265	0.051	0.059	0.078	0.165	0.313
	dCov	0.045	0.065	0.071	0.092	0.173	0.044	0.053	0.079	0.125	0.191	0.043	0.052	0.063	0.121	0.248
	dCvdW	0.050	0.062	0.089	0.108	0.215	0.051	0.058	0.093	0.134	0.263	0.049	0.066	0.077	0.163	0.299
	Wilks	0.065	0.081	0.167	0.299	0.553	0.053	0.068	0.131	0.281	0.559	0.051	0.073	0.130	0.270	0.538
	SzR	0.045	0.072	0.130	0.213	0.448	0.048	0.065	0.115	0.218	0.421	0.048	0.061	0.122	0.219	0.411

Table S.5: Empirical powers, under the heavy-tailed elliptical t_3 distributions (case (c)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n	τ	432					864					1728				
		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.046	0.074	0.110	0.237	0.418	0.044	0.062	0.123	0.227	0.408	0.052	0.060	0.124	0.236	0.401
	Spearman	0.055	0.067	0.138	0.319	0.513	0.057	0.071	0.146	0.299	0.526	0.056	0.065	0.155	0.318	0.521
	Kendall	0.054	0.077	0.143	0.316	0.531	0.043	0.065	0.156	0.309	0.520	0.052	0.074	0.149	0.317	0.529
	vdW	0.056	0.073	0.134	0.314	0.538	0.053	0.071	0.149	0.311	0.538	0.053	0.065	0.162	0.333	0.537
	dCov	0.058	0.071	0.137	0.301	0.505	0.053	0.073	0.151	0.289	0.487	0.050	0.073	0.147	0.295	0.508
	dCvdW	0.055	0.072	0.141	0.307	0.529	0.052	0.069	0.148	0.296	0.503	0.049	0.067	0.156	0.311	0.527
	Wilks	0.057	0.069	0.133	0.252	0.464	0.057	0.067	0.143	0.238	0.415	0.055	0.061	0.119	0.255	0.431
	SzR	0.033	0.073	0.161	0.290	0.533	0.054	0.078	0.130	0.285	0.539	0.044	0.073	0.165	0.284	0.526
$(d_1, d_2) = (3, 3)$	sign	0.056	0.060	0.133	0.260	0.466	0.043	0.063	0.119	0.264	0.479	0.055	0.068	0.123	0.284	0.497
	Spearman	0.049	0.063	0.129	0.276	0.489	0.044	0.063	0.127	0.287	0.509	0.052	0.061	0.122	0.275	0.539
	Kendall	0.056	0.066	0.138	0.281	0.523	0.046	0.059	0.132	0.291	0.515	0.054	0.060	0.119	0.303	0.541
	vdW	0.051	0.061	0.139	0.305	0.537	0.044	0.067	0.141	0.322	0.569	0.053	0.065	0.135	0.320	0.582
	dCov	0.053	0.061	0.121	0.273	0.497	0.045	0.059	0.129	0.274	0.505	0.052	0.059	0.117	0.279	0.519
	dCvdW	0.055	0.059	0.129	0.297	0.543	0.043	0.061	0.135	0.309	0.557	0.051	0.059	0.136	0.315	0.575
	Wilks	0.063	0.076	0.135	0.257	0.487	0.064	0.082	0.133	0.253	0.471	0.059	0.088	0.121	0.241	0.464
	SzR	0.041	0.071	0.124	0.316	0.546	0.045	0.067	0.132	0.330	0.590	0.041	0.049	0.140	0.316	0.552
$(d_1, d_2) = (5, 5)$	sign	0.050	0.061	0.093	0.223	0.431	0.049	0.063	0.107	0.249	0.441	0.056	0.063	0.115	0.255	0.492
	Spearman	0.046	0.067	0.107	0.193	0.363	0.053	0.063	0.119	0.219	0.385	0.057	0.067	0.105	0.203	0.432
	Kendall	0.051	0.063	0.107	0.214	0.407	0.053	0.068	0.137	0.234	0.421	0.057	0.067	0.118	0.227	0.453
	vdW	0.047	0.065	0.104	0.242	0.441	0.053	0.060	0.133	0.274	0.480	0.055	0.072	0.118	0.273	0.543
	dCov	0.056	0.063	0.106	0.206	0.375	0.049	0.060	0.125	0.231	0.390	0.060	0.071	0.111	0.218	0.428
	dCvdW	0.055	0.066	0.106	0.259	0.458	0.055	0.060	0.135	0.289	0.492	0.057	0.074	0.125	0.266	0.547
	Wilks	0.084	0.103	0.147	0.307	0.547	0.071	0.088	0.153	0.283	0.513	0.069	0.093	0.131	0.263	0.487
	SzR	0.042	0.055	0.119	0.295	0.551	0.054	0.063	0.115	0.304	0.578	0.046	0.086	0.142	0.336	0.608
$(d_1, d_2) = (7, 7)$	sign	0.046	0.053	0.075	0.145	0.287	0.041	0.057	0.093	0.184	0.341	0.049	0.065	0.121	0.221	0.411
	Spearman	0.058	0.066	0.091	0.147	0.233	0.055	0.061	0.083	0.161	0.277	0.043	0.065	0.096	0.182	0.313
	Kendall	0.053	0.061	0.089	0.148	0.276	0.047	0.072	0.089	0.165	0.306	0.057	0.063	0.102	0.189	0.319
	vdW	0.051	0.054	0.093	0.165	0.307	0.044	0.065	0.100	0.207	0.365	0.046	0.066	0.111	0.226	0.425
	dCov	0.042	0.053	0.081	0.119	0.204	0.043	0.059	0.071	0.147	0.261	0.042	0.068	0.085	0.157	0.287
	dCvdW	0.045	0.058	0.085	0.165	0.295	0.042	0.066	0.095	0.199	0.361	0.047	0.069	0.117	0.237	0.424
	Wilks	0.098	0.119	0.198	0.323	0.559	0.088	0.121	0.167	0.303	0.521	0.087	0.095	0.153	0.291	0.491
	SzR	0.054	0.062	0.100	0.246	0.535	0.046	0.066	0.119	0.304	0.590	0.037	0.061	0.121	0.266	0.591

Table S.6: Empirical powers, under the heavy-tailed elliptical t_2 distributions (case (d)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n	τ	432					864					1728				
		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.046	0.084	0.147	0.334	0.555	0.047	0.083	0.168	0.347	0.608	0.050	0.067	0.167	0.352	0.586
	Spearman	0.047	0.084	0.205	0.445	0.695	0.047	0.075	0.206	0.427	0.711	0.047	0.077	0.201	0.437	0.716
	Kendall	0.047	0.089	0.205	0.445	0.697	0.047	0.083	0.211	0.433	0.709	0.053	0.083	0.215	0.429	0.725
	vdW	0.048	0.087	0.208	0.469	0.738	0.049	0.081	0.216	0.471	0.740	0.049	0.081	0.221	0.481	0.761
	dCov	0.043	0.083	0.201	0.424	0.676	0.051	0.078	0.187	0.421	0.688	0.045	0.069	0.191	0.413	0.707
	dCvdW	0.047	0.085	0.203	0.438	0.697	0.045	0.085	0.203	0.443	0.709	0.046	0.079	0.206	0.437	0.739
	Wilks	0.070	0.089	0.143	0.278	0.509	0.063	0.083	0.139	0.262	0.482	0.061	0.079	0.126	0.248	0.467
	SzR	0.044	0.062	0.163	0.359	0.667	0.048	0.077	0.165	0.381	0.679	0.049	0.077	0.190	0.394	0.700
$(d_1, d_2) = (3, 3)$	sign	0.041	0.059	0.173	0.335	0.635	0.052	0.080	0.175	0.404	0.667	0.046	0.077	0.185	0.393	0.675
	Spearman	0.052	0.082	0.185	0.387	0.708	0.051	0.073	0.192	0.409	0.673	0.052	0.076	0.200	0.421	0.701
	Kendall	0.059	0.077	0.199	0.401	0.701	0.048	0.073	0.193	0.440	0.719	0.047	0.073	0.197	0.433	0.719
	vdW	0.056	0.075	0.203	0.439	0.747	0.054	0.076	0.215	0.489	0.757	0.044	0.077	0.233	0.490	0.784
	dCov	0.059	0.077	0.175	0.373	0.683	0.047	0.077	0.187	0.407	0.673	0.052	0.073	0.191	0.420	0.683
	dCvdW	0.059	0.069	0.201	0.414	0.723	0.044	0.079	0.214	0.458	0.744	0.051	0.073	0.214	0.463	0.754
	Wilks	0.091	0.117	0.164	0.306	0.533	0.073	0.103	0.152	0.255	0.501	0.070	0.091	0.142	0.263	0.465
	SzR	0.042	0.060	0.132	0.285	0.637	0.046	0.057	0.159	0.352	0.655	0.046	0.067	0.169	0.348	0.726
$(d_1, d_2) = (5, 5)$	sign	0.049	0.075	0.128	0.318	0.577	0.037	0.077	0.160	0.345	0.651	0.053	0.076	0.167	0.402	0.709
	Spearman	0.057	0.084	0.135	0.279	0.548	0.050	0.069	0.152	0.312	0.583	0.046	0.060	0.161	0.336	0.624
	Kendall	0.051	0.079	0.136	0.297	0.570	0.057	0.069	0.162	0.328	0.628	0.055	0.073	0.161	0.363	0.665
	vdW	0.047	0.083	0.146	0.339	0.649	0.046	0.077	0.186	0.400	0.727	0.053	0.076	0.190	0.454	0.779
	dCov	0.060	0.078	0.138	0.274	0.541	0.048	0.070	0.157	0.307	0.583	0.051	0.076	0.154	0.328	0.631
	dCvdW	0.053	0.081	0.154	0.335	0.638	0.045	0.077	0.187	0.375	0.709	0.055	0.071	0.186	0.436	0.757
	Wilks	0.127	0.160	0.223	0.326	0.601	0.127	0.139	0.206	0.313	0.529	0.112	0.125	0.172	0.291	0.479
	SzR	0.051	0.056	0.100	0.234	0.499	0.053	0.057	0.121	0.270	0.543	0.050	0.066	0.121	0.281	0.637
$(d_1, d_2) = (7, 7)$	sign	0.050	0.059	0.105	0.208	0.429	0.049	0.061	0.119	0.261	0.545	0.050	0.079	0.134	0.323	0.616
	Spearman	0.062	0.073	0.105	0.208	0.365	0.050	0.075	0.121	0.242	0.448	0.052	0.072	0.120	0.261	0.503
	Kendall	0.061	0.071	0.106	0.208	0.398	0.054	0.065	0.125	0.261	0.491	0.049	0.076	0.128	0.271	0.516
	vdW	0.058	0.063	0.111	0.238	0.463	0.047	0.073	0.135	0.313	0.592	0.045	0.076	0.163	0.359	0.668
	dCov	0.047	0.054	0.090	0.175	0.323	0.042	0.065	0.102	0.205	0.405	0.049	0.056	0.107	0.227	0.455
	dCvdW	0.051	0.059	0.115	0.229	0.451	0.045	0.064	0.126	0.305	0.572	0.046	0.083	0.155	0.358	0.643
	Wilks	0.142	0.188	0.231	0.381	0.614	0.139	0.158	0.198	0.331	0.538	0.133	0.168	0.203	0.296	0.479
	SzR	0.044	0.056	0.092	0.188	0.373	0.045	0.060	0.119	0.207	0.438	0.046	0.073	0.110	0.252	0.499

Table S.7: Empirical powers, under the heavy-tailed elliptical Cauchy distributions (case (e)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n		432					864					1728				
τ		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.047	0.179	0.517	0.819	0.942	0.042	0.204	0.578	0.864	0.965	0.052	0.218	0.613	0.905	0.967
	Spearman	0.040	0.293	0.766	0.966	0.992	0.049	0.311	0.833	0.977	0.993	0.045	0.358	0.863	0.984	0.986
	Kendall	0.049	0.277	0.740	0.948	0.990	0.051	0.285	0.794	0.969	0.992	0.049	0.331	0.817	0.979	0.984
	vdW	0.041	0.367	0.833	0.981	0.994	0.049	0.397	0.897	0.988	0.994	0.043	0.466	0.930	0.987	0.991
	dCov	0.050	0.242	0.713	0.941	0.988	0.041	0.273	0.783	0.971	0.991	0.049	0.313	0.808	0.977	0.985
	dCvdW	0.049	0.251	0.737	0.953	0.989	0.043	0.299	0.813	0.977	0.991	0.050	0.341	0.839	0.983	0.985
	Wilks	0.053	0.242	0.424	0.594	0.817	0.045	0.215	0.397	0.581	0.805	0.041	0.221	0.397	0.577	0.767
	SzR	0.041	0.360	0.588	0.815	0.973	0.047	0.426	0.683	0.910	0.996	0.055	0.506	0.794	0.981	1.000
$(d_1, d_2) = (3, 3)$	sign	0.051	0.183	0.549	0.875	0.976	0.061	0.217	0.636	0.927	0.981	0.042	0.224	0.683	0.939	0.977
	Spearman	0.053	0.265	0.769	0.973	0.995	0.056	0.321	0.859	0.986	0.992	0.043	0.382	0.902	0.973	0.989
	Kendall	0.050	0.244	0.714	0.967	0.994	0.061	0.277	0.807	0.977	0.990	0.047	0.333	0.848	0.969	0.987
	vdW	0.049	0.304	0.822	0.983	0.995	0.053	0.387	0.897	0.989	0.992	0.041	0.463	0.941	0.975	0.989
	dCov	0.055	0.229	0.709	0.956	0.993	0.055	0.278	0.801	0.977	0.991	0.043	0.321	0.847	0.969	0.986
	dCvdW	0.050	0.241	0.739	0.960	0.992	0.058	0.304	0.809	0.976	0.990	0.044	0.345	0.865	0.970	0.985
	Wilks	0.103	0.283	0.485	0.664	0.868	0.089	0.253	0.473	0.661	0.837	0.061	0.245	0.487	0.643	0.830
	SzR	0.052	0.352	0.562	0.799	0.943	0.046	0.402	0.639	0.865	0.982	0.052	0.488	0.765	0.956	0.998
$(d_1, d_2) = (5, 5)$	sign	0.042	0.131	0.431	0.824	0.963	0.049	0.174	0.554	0.881	0.975	0.042	0.187	0.625	0.922	0.972
	Spearman	0.049	0.197	0.641	0.939	0.993	0.048	0.252	0.775	0.981	0.989	0.045	0.296	0.844	0.974	0.988
	Kendall	0.051	0.177	0.560	0.916	0.985	0.053	0.204	0.695	0.953	0.979	0.051	0.259	0.761	0.963	0.982
	vdW	0.049	0.198	0.661	0.955	0.991	0.047	0.276	0.816	0.983	0.988	0.042	0.343	0.883	0.970	0.986
	dCov	0.053	0.170	0.566	0.916	0.985	0.048	0.202	0.704	0.968	0.984	0.041	0.256	0.772	0.967	0.985
	dCvdW	0.048	0.163	0.575	0.923	0.985	0.050	0.217	0.715	0.962	0.982	0.040	0.269	0.792	0.965	0.983
	Wilks	0.155	0.369	0.577	0.757	0.919	0.133	0.340	0.541	0.736	0.891	0.095	0.330	0.509	0.705	0.873
	SzR	0.054	0.331	0.532	0.754	0.896	0.054	0.385	0.662	0.835	0.951	0.041	0.499	0.740	0.936	0.989
$(d_1, d_2) = (7, 7)$	sign	0.046	0.104	0.311	0.641	0.898	0.044	0.136	0.405	0.797	0.955	0.059	0.165	0.539	0.878	0.971
	Spearman	0.058	0.156	0.489	0.844	0.976	0.055	0.196	0.606	0.935	0.988	0.047	0.230	0.765	0.958	0.983
	Kendall	0.055	0.134	0.411	0.777	0.956	0.053	0.163	0.511	0.904	0.983	0.061	0.191	0.651	0.935	0.978
	vdW	0.047	0.144	0.489	0.859	0.975	0.048	0.195	0.621	0.941	0.987	0.059	0.243	0.778	0.960	0.981
	dCov	0.052	0.111	0.372	0.759	0.961	0.046	0.146	0.515	0.889	0.979	0.053	0.173	0.646	0.939	0.979
	dCvdW	0.050	0.120	0.396	0.777	0.957	0.046	0.165	0.533	0.902	0.978	0.062	0.216	0.677	0.940	0.978
	Wilks	0.218	0.448	0.654	0.818	0.931	0.148	0.401	0.623	0.777	0.914	0.141	0.361	0.553	0.761	0.880
	SzR	0.045	0.329	0.582	0.721	0.855	0.048	0.424	0.668	0.810	0.949	0.052	0.475	0.731	0.896	0.983

Table S.8: Empirical powers, under the heavy-tailed marginal t_3 distributions (case (f)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n	τ	432					864					1728				
		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.045	0.064	0.103	0.208	0.358	0.044	0.069	0.108	0.233	0.395	0.045	0.067	0.130	0.214	0.381
	Spearman	0.045	0.073	0.151	0.277	0.505	0.042	0.069	0.144	0.300	0.521	0.047	0.065	0.148	0.285	0.512
	Kendall	0.047	0.082	0.152	0.298	0.496	0.045	0.070	0.146	0.315	0.527	0.045	0.082	0.155	0.286	0.520
	vdW	0.045	0.073	0.158	0.288	0.523	0.045	0.069	0.145	0.311	0.529	0.049	0.073	0.159	0.299	0.535
	dCov	0.040	0.073	0.143	0.272	0.466	0.045	0.069	0.133	0.284	0.501	0.048	0.071	0.151	0.275	0.487
	dCvdW	0.041	0.075	0.144	0.279	0.485	0.046	0.071	0.137	0.299	0.517	0.051	0.077	0.159	0.290	0.493
	Wilks	0.057	0.075	0.136	0.248	0.438	0.055	0.067	0.129	0.249	0.441	0.045	0.065	0.123	0.235	0.409
	SzR	0.038	0.067	0.146	0.291	0.475	0.054	0.077	0.141	0.297	0.513	0.052	0.068	0.152	0.288	0.495
$(d_1, d_2) = (3, 3)$	sign	0.049	0.056	0.109	0.229	0.416	0.052	0.060	0.122	0.248	0.435	0.040	0.062	0.129	0.214	0.444
	Spearman	0.049	0.071	0.131	0.265	0.461	0.051	0.064	0.131	0.292	0.501	0.046	0.071	0.128	0.260	0.489
	Kendall	0.056	0.069	0.135	0.283	0.491	0.056	0.065	0.140	0.293	0.519	0.039	0.068	0.133	0.265	0.512
	vdW	0.046	0.072	0.137	0.286	0.489	0.049	0.063	0.145	0.319	0.534	0.051	0.070	0.141	0.291	0.541
	dCov	0.048	0.059	0.121	0.255	0.443	0.049	0.065	0.130	0.273	0.489	0.045	0.077	0.134	0.253	0.485
	dCvdW	0.041	0.061	0.129	0.273	0.483	0.055	0.062	0.137	0.300	0.526	0.043	0.069	0.129	0.263	0.519
	Wilks	0.061	0.078	0.135	0.274	0.484	0.065	0.069	0.149	0.261	0.465	0.055	0.073	0.141	0.241	0.443
	SzR	0.048	0.066	0.128	0.310	0.518	0.045	0.064	0.133	0.287	0.534	0.039	0.074	0.155	0.276	0.522
$(d_1, d_2) = (5, 5)$	sign	0.043	0.071	0.101	0.176	0.353	0.046	0.058	0.098	0.211	0.412	0.049	0.070	0.107	0.239	0.415
	Spearman	0.042	0.067	0.099	0.179	0.353	0.047	0.066	0.104	0.217	0.391	0.049	0.071	0.115	0.213	0.411
	Kendall	0.043	0.063	0.097	0.186	0.361	0.045	0.074	0.105	0.212	0.392	0.055	0.065	0.103	0.219	0.427
	vdW	0.040	0.069	0.107	0.211	0.408	0.042	0.069	0.113	0.245	0.465	0.051	0.075	0.117	0.245	0.484
	dCov	0.041	0.067	0.091	0.168	0.359	0.056	0.071	0.102	0.209	0.379	0.055	0.066	0.111	0.216	0.397
	dCvdW	0.041	0.078	0.106	0.201	0.389	0.049	0.071	0.109	0.243	0.455	0.051	0.071	0.115	0.259	0.472
	Wilks	0.051	0.085	0.156	0.297	0.564	0.055	0.085	0.127	0.274	0.501	0.063	0.085	0.128	0.287	0.491
	SzR	0.046	0.050	0.129	0.311	0.554	0.042	0.054	0.122	0.288	0.542	0.041	0.080	0.136	0.299	0.568
$(d_1, d_2) = (7, 7)$	sign	0.049	0.061	0.081	0.138	0.260	0.052	0.070	0.090	0.155	0.313	0.049	0.058	0.095	0.193	0.333
	Spearman	0.047	0.068	0.105	0.143	0.236	0.053	0.066	0.091	0.151	0.263	0.051	0.063	0.089	0.175	0.284
	Kendall	0.049	0.074	0.078	0.136	0.224	0.057	0.070	0.093	0.151	0.260	0.050	0.055	0.087	0.181	0.294
	vdW	0.046	0.067	0.091	0.147	0.281	0.051	0.067	0.102	0.166	0.335	0.052	0.059	0.095	0.215	0.379
	dCov	0.037	0.052	0.077	0.120	0.207	0.047	0.050	0.078	0.126	0.223	0.052	0.055	0.077	0.149	0.260
	dCvdW	0.043	0.067	0.084	0.144	0.278	0.055	0.065	0.106	0.167	0.325	0.056	0.059	0.098	0.207	0.372
	Wilks	0.080	0.094	0.173	0.307	0.596	0.059	0.095	0.136	0.285	0.532	0.065	0.084	0.133	0.262	0.505
	SzR	0.038	0.069	0.120	0.289	0.568	0.053	0.071	0.133	0.270	0.588	0.052	0.063	0.129	0.310	0.582

Table S.9: Empirical powers, under the heavy-tailed marginal t_2 distributions (case (g)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n	τ	432					864					1728				
		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.045	0.081	0.145	0.309	0.515	0.055	0.066	0.151	0.308	0.537	0.055	0.065	0.149	0.332	0.535
	Spearman	0.049	0.082	0.185	0.397	0.679	0.053	0.074	0.205	0.423	0.702	0.040	0.075	0.203	0.432	0.689
	Kendall	0.041	0.082	0.195	0.421	0.676	0.053	0.071	0.202	0.425	0.686	0.041	0.077	0.204	0.428	0.691
	vdW	0.043	0.087	0.199	0.434	0.702	0.054	0.087	0.217	0.452	0.732	0.048	0.083	0.211	0.459	0.739
	dCov	0.048	0.075	0.165	0.385	0.648	0.055	0.080	0.183	0.404	0.661	0.049	0.070	0.205	0.411	0.663
	dCvdW	0.041	0.084	0.175	0.401	0.667	0.054	0.079	0.192	0.427	0.677	0.047	0.072	0.201	0.432	0.682
	Wilks	0.062	0.087	0.140	0.295	0.491	0.057	0.082	0.139	0.267	0.452	0.055	0.067	0.126	0.231	0.431
	SzR	0.048	0.058	0.167	0.379	0.639	0.051	0.076	0.159	0.365	0.645	0.041	0.077	0.175	0.432	0.675
$(d_1, d_2) = (3, 3)$	sign	0.053	0.067	0.151	0.337	0.585	0.055	0.079	0.159	0.335	0.628	0.051	0.079	0.163	0.351	0.634
	Spearman	0.056	0.082	0.183	0.378	0.652	0.048	0.081	0.191	0.390	0.703	0.049	0.079	0.195	0.439	0.724
	Kendall	0.055	0.078	0.180	0.359	0.655	0.051	0.079	0.183	0.403	0.711	0.044	0.079	0.191	0.415	0.719
	vdW	0.055	0.080	0.201	0.406	0.706	0.046	0.076	0.200	0.435	0.766	0.047	0.085	0.214	0.476	0.789
	dCov	0.047	0.078	0.173	0.351	0.625	0.056	0.080	0.177	0.367	0.687	0.041	0.078	0.186	0.396	0.701
	dCvdW	0.051	0.076	0.189	0.380	0.677	0.051	0.082	0.187	0.403	0.741	0.039	0.089	0.194	0.437	0.746
	Wilks	0.089	0.105	0.150	0.296	0.543	0.087	0.093	0.140	0.281	0.512	0.081	0.099	0.139	0.260	0.473
	SzR	0.053	0.061	0.138	0.309	0.631	0.039	0.064	0.170	0.353	0.655	0.046	0.073	0.162	0.360	0.696
$(d_1, d_2) = (5, 5)$	sign	0.043	0.066	0.121	0.277	0.502	0.053	0.071	0.128	0.283	0.569	0.051	0.065	0.159	0.347	0.625
	Spearman	0.048	0.071	0.127	0.277	0.511	0.049	0.070	0.147	0.287	0.561	0.053	0.084	0.166	0.349	0.639
	Kendall	0.052	0.067	0.119	0.266	0.471	0.055	0.071	0.131	0.287	0.533	0.048	0.073	0.154	0.328	0.599
	vdW	0.055	0.070	0.145	0.319	0.594	0.048	0.079	0.158	0.348	0.645	0.055	0.091	0.189	0.412	0.735
	dCov	0.045	0.067	0.115	0.265	0.487	0.055	0.075	0.143	0.276	0.548	0.055	0.087	0.155	0.332	0.609
	dCvdW	0.047	0.070	0.133	0.303	0.573	0.052	0.074	0.148	0.328	0.624	0.049	0.076	0.183	0.397	0.708
	Wilks	0.113	0.127	0.199	0.341	0.580	0.107	0.135	0.175	0.279	0.534	0.107	0.109	0.161	0.281	0.501
	SzR	0.047	0.054	0.116	0.258	0.537	0.048	0.060	0.128	0.283	0.590	0.054	0.073	0.124	0.317	0.664
$(d_1, d_2) = (7, 7)$	sign	0.051	0.061	0.094	0.179	0.369	0.042	0.056	0.115	0.226	0.448	0.049	0.059	0.133	0.257	0.529
	Spearman	0.065	0.073	0.107	0.163	0.332	0.057	0.069	0.105	0.229	0.406	0.057	0.063	0.105	0.244	0.479
	Kendall	0.058	0.066	0.098	0.160	0.303	0.043	0.065	0.097	0.204	0.377	0.055	0.062	0.117	0.216	0.427
	vdW	0.051	0.065	0.107	0.197	0.406	0.039	0.067	0.120	0.262	0.505	0.059	0.065	0.137	0.302	0.593
	dCov	0.046	0.063	0.090	0.141	0.279	0.050	0.065	0.100	0.185	0.372	0.046	0.063	0.095	0.223	0.428
	dCvdW	0.050	0.062	0.098	0.183	0.386	0.037	0.066	0.119	0.250	0.501	0.049	0.065	0.133	0.288	0.572
	Wilks	0.139	0.161	0.226	0.381	0.593	0.125	0.149	0.195	0.327	0.539	0.127	0.131	0.195	0.298	0.529
	SzR	0.038	0.056	0.119	0.210	0.471	0.043	0.060	0.113	0.237	0.549	0.058	0.071	0.123	0.268	0.556

Table S.10: Empirical powers, under the heavy-tailed marginal Cauchy distributions (case (h)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n		432					864					1728				
τ		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.053	0.187	0.548	0.857	0.973	0.043	0.229	0.618	0.881	0.974	0.047	0.241	0.665	0.905	0.968
	Spearman	0.050	0.344	0.828	0.975	0.993	0.042	0.375	0.880	0.983	0.992	0.047	0.426	0.905	0.979	0.985
	Kendall	0.052	0.291	0.755	0.955	0.993	0.045	0.319	0.805	0.973	0.989	0.052	0.350	0.834	0.969	0.980
	vdW	0.053	0.404	0.875	0.988	0.995	0.039	0.456	0.922	0.988	0.993	0.046	0.519	0.945	0.983	0.987
	dCov	0.055	0.295	0.755	0.967	0.992	0.037	0.315	0.825	0.974	0.991	0.051	0.355	0.854	0.973	0.982
	dCvdW	0.051	0.311	0.785	0.972	0.992	0.040	0.338	0.851	0.979	0.991	0.048	0.385	0.877	0.974	0.983
	Wilks	0.062	0.224	0.381	0.592	0.782	0.049	0.220	0.401	0.571	0.775	0.041	0.201	0.390	0.567	0.713
	SzR	0.040	0.334	0.580	0.831	0.971	0.059	0.413	0.715	0.923	0.995	0.046	0.510	0.817	0.975	1.000
$(d_1, d_2) = (3, 3)$	sign	0.040	0.194	0.594	0.909	0.985	0.053	0.237	0.687	0.941	0.981	0.063	0.284	0.752	0.943	0.971
	Spearman	0.051	0.325	0.824	0.984	0.994	0.055	0.396	0.908	0.987	0.989	0.055	0.473	0.937	0.976	0.982
	Kendall	0.047	0.219	0.698	0.963	0.992	0.055	0.274	0.787	0.970	0.985	0.050	0.327	0.833	0.963	0.975
	vdW	0.045	0.352	0.860	0.987	0.994	0.055	0.447	0.926	0.987	0.989	0.057	0.531	0.951	0.978	0.983
	dCov	0.058	0.253	0.777	0.980	0.994	0.051	0.329	0.858	0.983	0.986	0.050	0.405	0.899	0.971	0.979
	dCvdW	0.050	0.266	0.783	0.979	0.994	0.049	0.342	0.865	0.984	0.986	0.061	0.417	0.902	0.969	0.980
	Wilks	0.091	0.277	0.477	0.681	0.826	0.073	0.261	0.445	0.641	0.816	0.064	0.243	0.437	0.611	0.798
	SzR	0.065	0.352	0.576	0.795	0.939	0.061	0.416	0.682	0.903	0.983	0.039	0.477	0.791	0.960	1.000
$(d_1, d_2) = (5, 5)$	sign	0.049	0.164	0.535	0.877	0.973	0.033	0.223	0.667	0.930	0.968	0.045	0.238	0.746	0.937	0.959
	Spearman	0.049	0.247	0.732	0.968	0.989	0.065	0.335	0.869	0.979	0.985	0.048	0.429	0.907	0.969	0.979
	Kendall	0.051	0.151	0.474	0.828	0.969	0.052	0.201	0.590	0.903	0.971	0.048	0.225	0.704	0.939	0.961
	vdW	0.049	0.241	0.762	0.976	0.989	0.050	0.355	0.881	0.978	0.983	0.049	0.445	0.917	0.966	0.977
	dCov	0.050	0.193	0.656	0.945	0.989	0.055	0.272	0.796	0.974	0.978	0.058	0.363	0.875	0.967	0.974
	dCvdW	0.046	0.205	0.673	0.953	0.987	0.045	0.290	0.797	0.969	0.976	0.048	0.354	0.863	0.961	0.969
	Wilks	0.160	0.362	0.575	0.750	0.897	0.134	0.335	0.527	0.722	0.858	0.089	0.311	0.503	0.705	0.847
	SzR	0.039	0.341	0.567	0.739	0.871	0.045	0.389	0.631	0.825	0.958	0.048	0.488	0.745	0.936	0.991
$(d_1, d_2) = (7, 7)$	sign	0.052	0.119	0.381	0.717	0.929	0.044	0.149	0.544	0.858	0.963	0.049	0.197	0.659	0.918	0.951
	Spearman	0.068	0.193	0.561	0.884	0.976	0.056	0.233	0.743	0.965	0.983	0.053	0.317	0.849	0.962	0.967
	Kendall	0.059	0.125	0.289	0.571	0.844	0.057	0.125	0.385	0.734	0.936	0.055	0.143	0.487	0.845	0.933
	vdW	0.052	0.176	0.563	0.882	0.980	0.047	0.231	0.747	0.961	0.980	0.055	0.321	0.847	0.955	0.965
	dCov	0.044	0.129	0.428	0.803	0.971	0.042	0.166	0.635	0.929	0.977	0.057	0.233	0.756	0.953	0.962
	dCvdW	0.047	0.144	0.485	0.833	0.968	0.043	0.189	0.669	0.927	0.973	0.053	0.266	0.780	0.946	0.961
	Wilks	0.201	0.425	0.643	0.829	0.904	0.170	0.391	0.575	0.770	0.899	0.128	0.363	0.566	0.733	0.869
	SzR	0.053	0.356	0.528	0.750	0.884	0.048	0.417	0.649	0.838	0.950	0.043	0.473	0.714	0.915	0.988

Table S.11: Empirical powers, under the skewed independent χ_2^2 components (case (i)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n		432					864					1728				
τ		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.049	0.064	0.132	0.244	0.417	0.051	0.058	0.126	0.240	0.433	0.055	0.070	0.111	0.236	0.449
	Spearman	0.053	0.106	0.317	0.655	0.905	0.046	0.108	0.339	0.713	0.935	0.053	0.109	0.340	0.727	0.954
	Kendall	0.055	0.097	0.218	0.459	0.732	0.043	0.079	0.216	0.485	0.756	0.055	0.077	0.217	0.464	0.763
	vdW	0.055	0.108	0.344	0.685	0.922	0.046	0.112	0.395	0.767	0.955	0.051	0.125	0.411	0.793	0.977
	dCov	0.059	0.097	0.274	0.564	0.857	0.043	0.088	0.278	0.625	0.888	0.051	0.093	0.274	0.624	0.903
	dCvdW	0.059	0.097	0.261	0.561	0.847	0.047	0.090	0.275	0.629	0.884	0.050	0.093	0.287	0.634	0.917
	Wilks	0.059	0.080	0.135	0.240	0.422	0.055	0.068	0.137	0.263	0.444	0.056	0.067	0.118	0.233	0.434
	SzR	0.046	0.058	0.129	0.249	0.445	0.051	0.072	0.123	0.245	0.439	0.037	0.073	0.139	0.236	0.412
$(d_1, d_2) = (3, 3)$	sign	0.048	0.067	0.161	0.300	0.530	0.045	0.073	0.150	0.331	0.588	0.058	0.070	0.158	0.319	0.607
	Spearman	0.045	0.098	0.275	0.613	0.887	0.053	0.114	0.343	0.721	0.945	0.049	0.114	0.384	0.789	0.977
	Kendall	0.044	0.088	0.196	0.408	0.692	0.045	0.085	0.208	0.437	0.764	0.049	0.088	0.223	0.476	0.783
	vdW	0.047	0.091	0.259	0.574	0.870	0.054	0.109	0.317	0.673	0.933	0.049	0.109	0.362	0.743	0.961
	dCov	0.045	0.095	0.241	0.559	0.860	0.047	0.100	0.290	0.633	0.924	0.047	0.105	0.333	0.705	0.948
	dCvdW	0.045	0.089	0.228	0.499	0.820	0.044	0.086	0.268	0.571	0.885	0.053	0.100	0.296	0.630	0.919
	Wilks	0.053	0.072	0.143	0.251	0.447	0.046	0.068	0.138	0.273	0.447	0.053	0.071	0.129	0.233	0.436
	SzR	0.047	0.062	0.150	0.241	0.482	0.043	0.056	0.099	0.247	0.439	0.055	0.071	0.120	0.258	0.446
$(d_1, d_2) = (5, 5)$	sign	0.037	0.067	0.119	0.275	0.499	0.039	0.057	0.141	0.299	0.569	0.045	0.071	0.164	0.361	0.676
	Spearman	0.053	0.071	0.132	0.347	0.609	0.051	0.074	0.159	0.445	0.735	0.052	0.081	0.229	0.531	0.867
	Kendall	0.053	0.065	0.112	0.285	0.495	0.044	0.065	0.145	0.307	0.572	0.049	0.061	0.149	0.362	0.669
	vdW	0.052	0.065	0.143	0.361	0.638	0.041	0.071	0.167	0.432	0.748	0.048	0.079	0.220	0.518	0.871
	dCov	0.054	0.066	0.130	0.341	0.594	0.048	0.061	0.165	0.439	0.713	0.060	0.087	0.209	0.515	0.850
	dCvdW	0.052	0.077	0.143	0.345	0.621	0.043	0.062	0.168	0.403	0.714	0.051	0.079	0.205	0.479	0.828
	Wilks	0.053	0.093	0.139	0.268	0.514	0.051	0.073	0.121	0.293	0.501	0.041	0.065	0.140	0.255	0.497
	SzR	0.044	0.057	0.121	0.274	0.465	0.046	0.062	0.110	0.244	0.452	0.043	0.069	0.137	0.283	0.468
$(d_1, d_2) = (7, 7)$	sign	0.043	0.064	0.095	0.171	0.327	0.045	0.062	0.111	0.215	0.447	0.045	0.059	0.121	0.247	0.501
	Spearman	0.057	0.067	0.098	0.179	0.307	0.051	0.065	0.131	0.225	0.434	0.052	0.067	0.135	0.273	0.545
	Kendall	0.053	0.062	0.085	0.169	0.322	0.051	0.059	0.125	0.195	0.402	0.041	0.063	0.102	0.221	0.459
	vdW	0.051	0.066	0.101	0.197	0.381	0.047	0.065	0.140	0.257	0.513	0.043	0.062	0.130	0.302	0.618
	dCov	0.047	0.055	0.079	0.163	0.307	0.042	0.065	0.106	0.193	0.402	0.047	0.061	0.125	0.254	0.517
	dCvdW	0.044	0.061	0.100	0.193	0.384	0.047	0.064	0.125	0.251	0.501	0.039	0.061	0.132	0.289	0.585
	Wilks	0.059	0.080	0.147	0.307	0.559	0.045	0.081	0.141	0.283	0.539	0.051	0.067	0.131	0.285	0.510
	SzR	0.047	0.062	0.124	0.240	0.494	0.052	0.069	0.141	0.255	0.500	0.056	0.070	0.117	0.230	0.493

Table S.12: Empirical powers, under the skewed independent χ_1^2 components (case (j)), of (i) the center-outward quadrant or sign test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test (“vdW”), (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ pseudo-Gaussian test, and (viii) Székely and Rizzo’s permutation-based distance covariance test (“SzR”). Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 0.2, 0.4, 0.6, 0.8\}$.

n		432					864					1728				
τ		0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8	0	0.2	0.4	0.6	0.8
$(d_1, d_2) = (2, 2)$	sign	0.044	0.087	0.250	0.557	0.826	0.048	0.117	0.265	0.565	0.838	0.051	0.097	0.267	0.597	0.843
	Spearman	0.053	0.448	0.946	0.999	1.000	0.059	0.572	0.989	1.000	1.000	0.055	0.659	0.999	1.000	1.000
	Kendall	0.057	0.205	0.712	0.956	0.999	0.055	0.254	0.765	0.987	1.000	0.056	0.263	0.819	0.990	1.000
	vdW	0.058	0.447	0.943	0.999	1.000	0.058	0.587	0.987	1.000	1.000	0.056	0.707	0.999	1.000	1.000
	dCov	0.058	0.385	0.951	0.999	1.000	0.061	0.533	0.991	1.000	1.000	0.057	0.635	1.000	1.000	1.000
	dCvdW	0.057	0.344	0.919	0.999	1.000	0.057	0.502	0.986	1.000	1.000	0.055	0.626	1.000	1.000	1.000
	Wilks	0.063	0.076	0.131	0.249	0.416	0.057	0.087	0.139	0.249	0.425	0.049	0.073	0.127	0.236	0.421
	SzR	0.046	0.077	0.128	0.309	0.567	0.045	0.079	0.134	0.289	0.528	0.051	0.058	0.168	0.305	0.597
$(d_1, d_2) = (3, 3)$	sign	0.054	0.110	0.401	0.781	0.963	0.051	0.139	0.467	0.850	0.985	0.050	0.141	0.547	0.913	0.995
	Spearman	0.057	0.317	0.875	0.995	1.000	0.047	0.473	0.981	1.000	1.000	0.047	0.661	0.998	1.000	1.000
	Kendall	0.063	0.157	0.586	0.928	0.998	0.049	0.201	0.704	0.978	0.999	0.048	0.245	0.809	0.992	1.000
	vdW	0.059	0.286	0.826	0.992	1.000	0.051	0.395	0.949	1.000	1.000	0.049	0.553	0.992	1.000	1.000
	dCov	0.062	0.301	0.871	0.997	1.000	0.048	0.444	0.980	1.000	1.000	0.035	0.616	0.999	1.000	1.000
	dCvdW	0.061	0.237	0.795	0.992	1.000	0.049	0.331	0.925	0.999	1.000	0.045	0.461	0.989	1.000	1.000
	Wilks	0.066	0.075	0.136	0.280	0.467	0.060	0.074	0.138	0.253	0.447	0.055	0.076	0.130	0.255	0.459
	SzR	0.035	0.071	0.120	0.290	0.551	0.038	0.067	0.125	0.253	0.528	0.046	0.070	0.114	0.251	0.517
$(d_1, d_2) = (5, 5)$	sign	0.052	0.094	0.261	0.607	0.915	0.049	0.089	0.335	0.761	0.979	0.052	0.129	0.450	0.863	0.995
	Spearman	0.051	0.105	0.319	0.669	0.939	0.055	0.119	0.477	0.910	0.997	0.059	0.181	0.704	0.983	1.000
	Kendall	0.051	0.095	0.236	0.559	0.885	0.055	0.081	0.285	0.723	0.965	0.051	0.112	0.421	0.853	0.993
	vdW	0.052	0.110	0.341	0.717	0.970	0.051	0.127	0.471	0.912	0.999	0.051	0.181	0.679	0.977	1.000
	dCov	0.056	0.109	0.331	0.711	0.965	0.057	0.125	0.496	0.923	0.999	0.056	0.186	0.720	0.985	1.000
	dCvdW	0.051	0.105	0.340	0.725	0.973	0.055	0.106	0.459	0.893	0.999	0.051	0.163	0.634	0.971	1.000
	Wilks	0.053	0.082	0.158	0.290	0.533	0.058	0.075	0.136	0.287	0.491	0.055	0.086	0.135	0.251	0.483
	SzR	0.054	0.069	0.108	0.266	0.510	0.051	0.066	0.104	0.235	0.509	0.063	0.070	0.109	0.236	0.473
$(d_1, d_2) = (7, 7)$	sign	0.043	0.066	0.129	0.319	0.600	0.054	0.073	0.187	0.441	0.761	0.043	0.075	0.236	0.581	0.902
	Spearman	0.060	0.072	0.122	0.280	0.518	0.055	0.074	0.161	0.420	0.727	0.049	0.091	0.208	0.601	0.898
	Kendall	0.052	0.068	0.121	0.251	0.507	0.047	0.072	0.150	0.369	0.659	0.051	0.070	0.189	0.462	0.814
	vdW	0.050	0.071	0.140	0.357	0.647	0.049	0.081	0.194	0.522	0.835	0.052	0.083	0.257	0.677	0.952
	dCov	0.052	0.071	0.124	0.277	0.541	0.050	0.080	0.178	0.439	0.762	0.037	0.073	0.230	0.596	0.922
	dCvdW	0.047	0.071	0.141	0.361	0.646	0.047	0.073	0.197	0.528	0.841	0.052	0.080	0.263	0.659	0.942
	Wilks	0.087	0.105	0.163	0.313	0.555	0.061	0.077	0.131	0.287	0.520	0.064	0.072	0.128	0.278	0.518
	SzR	0.050	0.068	0.107	0.266	0.504	0.046	0.070	0.124	0.235	0.476	0.043	0.075	0.101	0.256	0.472

Table S.13: Empirical powers, under the Gaussian mixture alternative (case (k)), of (i) the center-outward sign (quadrant) test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test, (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ likelihood ratio test, and (viii) Székely and Rizzo’s permutation-based distance covariance test. Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 1, 2, 3, 4\}$.

n	τ	432					864					1728				
		0	1	2	3	4	0	1	2	3	4	0	1	2	3	4
$(d_1, d_2) = (2, 2)$	sign	0.062	0.063	0.123	0.245	0.409	0.049	0.065	0.137	0.234	0.424	0.052	0.062	0.118	0.263	0.401
	Spearman	0.042	0.060	0.164	0.347	0.561	0.052	0.079	0.177	0.342	0.593	0.039	0.080	0.170	0.356	0.571
	Kendall	0.041	0.062	0.174	0.342	0.568	0.060	0.073	0.183	0.332	0.567	0.048	0.073	0.156	0.350	0.562
	vdW	0.046	0.063	0.172	0.358	0.564	0.061	0.076	0.172	0.351	0.602	0.038	0.072	0.174	0.363	0.594
	dCov	0.045	0.057	0.159	0.322	0.532	0.051	0.072	0.167	0.323	0.555	0.044	0.068	0.173	0.345	0.547
	dCvdW	0.041	0.055	0.171	0.330	0.529	0.051	0.073	0.166	0.325	0.564	0.042	0.074	0.174	0.345	0.554
	Wilks	0.042	0.069	0.194	0.392	0.626	0.058	0.081	0.192	0.367	0.624	0.036	0.076	0.173	0.373	0.605
	SzR	0.048	0.061	0.174	0.352	0.580	0.042	0.068	0.168	0.330	0.582	0.039	0.072	0.175	0.347	0.567
$(d_1, d_2) = (3, 3)$	sign	0.040	0.063	0.125	0.236	0.445	0.056	0.062	0.120	0.263	0.455	0.058	0.058	0.140	0.278	0.477
	Spearman	0.051	0.063	0.128	0.309	0.549	0.057	0.073	0.154	0.336	0.557	0.051	0.065	0.168	0.357	0.575
	Kendall	0.048	0.071	0.127	0.308	0.536	0.053	0.068	0.157	0.311	0.558	0.054	0.064	0.170	0.354	0.564
	vdW	0.042	0.066	0.136	0.313	0.552	0.056	0.068	0.155	0.334	0.578	0.061	0.064	0.174	0.374	0.606
	dCov	0.036	0.070	0.113	0.292	0.518	0.051	0.058	0.161	0.310	0.536	0.057	0.067	0.143	0.319	0.564
	dCvdW	0.038	0.069	0.137	0.296	0.538	0.059	0.064	0.160	0.324	0.562	0.064	0.067	0.159	0.335	0.571
	Wilks	0.041	0.081	0.174	0.393	0.657	0.060	0.071	0.191	0.382	0.650	0.053	0.066	0.177	0.408	0.648
	SzR	0.040	0.074	0.148	0.367	0.618	0.051	0.062	0.171	0.358	0.604	0.065	0.066	0.187	0.377	0.606
$(d_1, d_2) = (5, 5)$	sign	0.044	0.067	0.100	0.217	0.402	0.044	0.059	0.111	0.244	0.431	0.050	0.059	0.118	0.237	0.503
	Spearman	0.058	0.070	0.114	0.216	0.432	0.041	0.070	0.141	0.259	0.458	0.051	0.066	0.152	0.286	0.503
	Kendall	0.059	0.061	0.108	0.242	0.455	0.043	0.052	0.129	0.247	0.454	0.055	0.063	0.137	0.282	0.513
	vdW	0.042	0.067	0.118	0.244	0.477	0.048	0.061	0.136	0.285	0.499	0.051	0.057	0.135	0.305	0.569
	dCov	0.046	0.071	0.105	0.220	0.408	0.049	0.062	0.127	0.261	0.442	0.051	0.065	0.122	0.278	0.503
	dCvdW	0.039	0.066	0.109	0.243	0.463	0.047	0.067	0.121	0.280	0.480	0.048	0.057	0.125	0.296	0.557
	Wilks	0.056	0.079	0.191	0.425	0.748	0.042	0.076	0.192	0.426	0.721	0.060	0.074	0.186	0.409	0.721
	SzR	0.035	0.065	0.176	0.404	0.708	0.043	0.060	0.170	0.384	0.689	0.055	0.063	0.165	0.381	0.698
$(d_1, d_2) = (7, 7)$	sign	0.038	0.056	0.086	0.169	0.305	0.049	0.057	0.110	0.210	0.355	0.049	0.076	0.113	0.220	0.424
	Spearman	0.047	0.084	0.105	0.172	0.310	0.052	0.061	0.108	0.208	0.379	0.054	0.065	0.112	0.216	0.401
	Kendall	0.050	0.070	0.106	0.180	0.305	0.057	0.064	0.113	0.212	0.372	0.051	0.072	0.115	0.213	0.416
	vdW	0.036	0.060	0.096	0.183	0.349	0.048	0.057	0.121	0.219	0.429	0.049	0.068	0.109	0.247	0.477
	dCov	0.035	0.054	0.079	0.146	0.263	0.047	0.059	0.096	0.177	0.326	0.044	0.047	0.094	0.169	0.354
	dCvdW	0.028	0.060	0.105	0.182	0.340	0.051	0.056	0.116	0.215	0.399	0.054	0.066	0.114	0.219	0.458
	Wilks	0.067	0.088	0.208	0.472	0.764	0.059	0.073	0.199	0.449	0.798	0.050	0.061	0.206	0.423	0.755
	SzR	0.049	0.068	0.172	0.427	0.723	0.057	0.063	0.180	0.414	0.769	0.045	0.064	0.178	0.404	0.744

Table S.14: Empirical powers, under the heavy-tailed elliptical mixture alternative (case (I)), of (i) the center-outward sign (quadrant) test, (ii) the center-outward Spearman test, (iii) the center-outward Kendall test, (iv) the center-outward van der Waerden test, (v) the center-outward Wilcoxon score version of the distance covariance test (“dCov”), (vi) the center-outward van der Waerden score version of the distance covariance test (“dCvdW”), (vii) Wilks’ likelihood ratio test, and (viii) Székely and Rizzo’s permutation-based distance covariance test. Based on 1,000 replications of a sample with dimensions $d_1 = d_2 \in \{2, 3, 5, 7\}$, size $n \in \{432, 864, 1728\}$, and parameter values $\delta^{(n)} = n^{-1/2}\tau$, $\tau \in \{0, 1, 2, 3, 4\}$.

n		432					864					1728				
τ		0	1	2	3	4	0	1	2	3	4	0	1	2	3	4
$(d_1, d_2) = (2, 2)$	sign	0.058	0.070	0.102	0.235	0.391	0.058	0.065	0.133	0.230	0.418	0.052	0.064	0.127	0.244	0.410
	Spearman	0.045	0.088	0.186	0.344	0.543	0.047	0.090	0.192	0.329	0.597	0.062	0.073	0.168	0.331	0.562
	Kendall	0.053	0.088	0.166	0.334	0.523	0.052	0.088	0.178	0.315	0.565	0.067	0.076	0.156	0.316	0.532
	vdW	0.039	0.088	0.184	0.357	0.548	0.050	0.087	0.201	0.335	0.602	0.053	0.072	0.176	0.340	0.584
	dCov	0.047	0.079	0.158	0.331	0.517	0.047	0.091	0.173	0.310	0.572	0.063	0.067	0.165	0.316	0.556
	dCvdW	0.049	0.085	0.157	0.322	0.504	0.050	0.080	0.172	0.308	0.564	0.059	0.070	0.169	0.307	0.548
	Wilks	0.068	0.130	0.266	0.435	0.623	0.064	0.124	0.265	0.423	0.627	0.063	0.112	0.252	0.409	0.598
	SzR	0.045	0.116	0.264	0.491	0.717	0.050	0.098	0.248	0.449	0.713	0.060	0.081	0.210	0.430	0.683
$(d_1, d_2) = (3, 3)$	sign	0.048	0.057	0.131	0.253	0.444	0.060	0.084	0.155	0.256	0.481	0.047	0.066	0.139	0.262	0.498
	Spearman	0.042	0.086	0.166	0.359	0.567	0.057	0.098	0.141	0.341	0.604	0.054	0.082	0.170	0.357	0.615
	Kendall	0.053	0.077	0.149	0.332	0.539	0.049	0.093	0.148	0.319	0.575	0.043	0.073	0.160	0.332	0.568
	vdW	0.041	0.077	0.158	0.352	0.561	0.059	0.089	0.159	0.345	0.600	0.055	0.073	0.155	0.356	0.624
	dCov	0.049	0.065	0.142	0.319	0.554	0.046	0.077	0.135	0.321	0.582	0.057	0.075	0.165	0.356	0.604
	dCvdW	0.048	0.071	0.145	0.322	0.524	0.051	0.079	0.152	0.323	0.570	0.047	0.072	0.155	0.345	0.594
	Wilks	0.061	0.156	0.301	0.511	0.696	0.068	0.179	0.286	0.500	0.686	0.084	0.123	0.265	0.447	0.680
	SzR	0.046	0.120	0.308	0.560	0.811	0.054	0.133	0.270	0.538	0.803	0.050	0.100	0.269	0.521	0.790
$(d_1, d_2) = (5, 5)$	sign	0.043	0.069	0.099	0.199	0.382	0.040	0.062	0.111	0.249	0.413	0.050	0.055	0.109	0.237	0.470
	Spearman	0.059	0.076	0.150	0.262	0.482	0.056	0.080	0.154	0.314	0.512	0.051	0.075	0.133	0.312	0.565
	Kendall	0.055	0.069	0.145	0.240	0.455	0.056	0.068	0.126	0.273	0.448	0.058	0.058	0.120	0.258	0.527
	vdW	0.043	0.073	0.138	0.237	0.469	0.049	0.072	0.129	0.299	0.487	0.052	0.055	0.129	0.268	0.550
	dCov	0.056	0.070	0.145	0.247	0.483	0.052	0.066	0.129	0.299	0.543	0.052	0.064	0.140	0.310	0.614
	dCvdW	0.045	0.071	0.119	0.206	0.420	0.046	0.069	0.108	0.265	0.465	0.056	0.058	0.128	0.254	0.546
	Wilks	0.092	0.207	0.417	0.631	0.815	0.095	0.198	0.364	0.597	0.789	0.072	0.182	0.326	0.539	0.778
	SzR	0.046	0.163	0.385	0.640	0.882	0.052	0.133	0.332	0.635	0.880	0.045	0.127	0.291	0.594	0.888
$(d_1, d_2) = (7, 7)$	sign	0.042	0.059	0.084	0.163	0.299	0.052	0.057	0.100	0.178	0.320	0.059	0.060	0.102	0.208	0.421
	Spearman	0.051	0.108	0.131	0.215	0.422	0.059	0.077	0.138	0.242	0.403	0.054	0.082	0.113	0.264	0.496
	Kendall	0.036	0.081	0.112	0.183	0.389	0.057	0.073	0.115	0.200	0.366	0.060	0.076	0.113	0.217	0.413
	vdW	0.041	0.078	0.112	0.169	0.358	0.066	0.069	0.121	0.212	0.392	0.058	0.069	0.101	0.236	0.472
	dCov	0.034	0.075	0.100	0.179	0.429	0.053	0.057	0.119	0.244	0.458	0.064	0.070	0.124	0.289	0.582
	dCvdW	0.036	0.077	0.094	0.157	0.345	0.047	0.056	0.105	0.196	0.349	0.060	0.066	0.099	0.216	0.444
	Wilks	0.095	0.272	0.458	0.728	0.867	0.100	0.242	0.423	0.660	0.852	0.092	0.222	0.417	0.624	0.835
	SzR	0.045	0.170	0.385	0.699	0.904	0.039	0.149	0.380	0.709	0.914	0.060	0.150	0.364	0.682	0.920