HOMOMORPHIC ENCODERS OF PROFINITE ABELIAN GROUPS II

MARÍA V. FERRER AND SALVADOR HERNÁNDEZ

ABSTRACT. Let $\{G_i: i \in \mathbb{N}\}$ be a family of finite Abelian groups. We say that a subgroup $G \leq \prod_{i \in \mathbb{N}} G_i$ is order controllable if for every $i \in \mathbb{N}$ there is $n_i \in \mathbb{N}$ such that

for each $c \in G$, there exists $c_1 \in G$ satisfying that $c_{1|[1,i]} = c_{|[1,i]}$, $supp(c_1) \subseteq [1, n_i]$, and $order(c_1)$ divides $order(c_{|[1,n_i]})$. In this paper we investigate the structure of order controllable group codes. It is proved that if G is an order controllable, shift invariant, group code over a finite abelian group H, then G possesses a finite canonical generating set. Furthermore, our construction also yields that G is algebraically conjugate to a full group shift.

1. Introduction

In coding theory, a code refers to a set of sequences (the codewords), with good error-correcting properties, used to transmit information over nosy channels. In communication technology most codes are linear (that is, vector spaces on a finite field) and there are two main classes of codes: $block\ codes$, in which the codewords are finite sequences all of the same length, and $convolutional\ codes$, in which the codewords can be infinite sequences. However, some very powerful codes that were first thought to be nonlinear can be described as additive subgroups of A^n , where A is a cyclic abelian group (see [1, 9]). This fact motivated the study of a more general class of codes.

Date: today.

Research Partially supported by the Spanish Ministerio de Economía y Competitividad, grant: MTM/PID2019-106529GB-I00 (AEI/FEDER, EU) and by the Universitat Jaume I, grant UJI-B2019-06.

 $^{2010\} Mathematics\ Subject\ Classification.$ Primary 20K25. Secondary 22C05, 20K45, 54H11, 68P30, 37R10

Key Words and Phrases: profinite abelian group, controllable group, order controllable group, group code, generating set, homomorphic encoder.

According to Forney and Trott [9, 10], a group code G is a subgroup of a product

$$X = \prod_{i \in I} G_i,$$

where each G_i is a group and the composition law is the componentwise group operation. The subgroup

$$G_f := G \cap \bigoplus_{i \in \mathbb{Z}} G_i$$

is called the *finite subcode* of G. It may happen that all elements of G has finite support, which means that G coincides with G_f .

If all code symbols are drawn from a common group H, then $G \leq H^I$ and G will be called a group code over H defined on I.

A key point in the study of group codes is the finding of appropriate encoders

Definition 1.1. Given a group code G, a homomorphic encoder is a continuous homomorphism $\Phi \colon \prod_{i \in I} H_i \to G$ that sends a full direct product of (topological) groups onto G. Of special relevance are the so called noncatastrophic encoders, that is, homomorphic continuous encoders α that are one-to-one and such that $\Phi(\bigoplus_{i \in I} H_i) = G_f$ (see [4, 9, 10] for some references).

From here on, we deal with a group shift (or group code) G over a finite abelian group H. That is, G is a closed, shift-invariant subgroup of the full shift group $X = H^{\mathbb{Z}}$. Therefore, if $\sigma: X \to X$ denotes the backward shift operator

$$\sigma[x](i) := x(i+1), \ \forall x \in X, \ i \in \mathbb{Z},$$

we have that $\sigma(G) = G$. For simplicity's sake, we denote the forward shift operator by ρ , that is $\rho[x](i) := x(i-1)$, $\forall x \in X$, $i \in \mathbb{Z}$. A group shift G over a finite abelian group H is irreducible or transitive if there is $x \in G$ such that the partial forward orbit $\{\sigma^n(x): n \geq n_0\}$ is dense in G for all $n_0 \in \mathbb{Z}$. Given two group codes G and \bar{G} if there is a homeomorphism (resp. topological group isomorphism) $\Phi: G \longrightarrow \bar{G}$ so that $\sigma \circ \Phi = \Phi \circ \sigma$ then we say that G and \bar{G} are topologically conjugate (resp. algebraically and topologically conjugate) (see [11, 12, 16]).

In [8], Forney proved that every (linear) convolutional code is conjugate to a full shift, via a linear conjugacy. Subsequently, it was proved by several authors (see [9, 11, 13, 14]) that every irreducible group shift is conjugate to a full shift. In fact, one might expect that the conjugacy was also a group homomorphism (algebraic conjugacy). But, for group shifts, this turns to be false in general (cf. [11, 13]). In this sense, Fagnani [2] has obtained necessary and sufficient conditions for a group shift to be algebraically conjugate to the full shift over a finite group. His approach is based on Pontryagin duality, which let one reduce the question to its discrete dual group that turns out to be a finitely generated module of Laurent polynomials.

We next collect some definitions and basic facts introduced in [6].

Definition 1.2. Let G be a group shift over a finite abelian group H. We have the following notions:

- (1) G is weakly controllable if $G \cap H^{(\mathbb{Z})}$ is dense in G; here $H^{(\mathbb{Z})}$ denotes the subgroup of $H^{\mathbb{Z}}$ consisting of the elements with finite support.
- (2) G is $controllable^1$ (equivalently irreducible or transitive) if there is a positive integer n_c such that for each $g \in G$, there exists $g_1 \in G$ such that $g_{1|(-\infty,0]} = g_{|(-\infty,0]}$ and $g_{1||n_c,+\infty)} = 0$ (we assume that n_c is the least integer satisfying this property). Remark that this property implies the existence of $g_2 := g g_1 \in G$ such that $g = g_1 + g_2$, $supp(g_1) \subseteq (-\infty, n_c]$ and $supp(g_2) \subseteq [1, +\infty[$.

It is easily verified that every controllable group code G is irreducible (see [11]).

(3) G is order controllable if there is a positive integer n_o such that for each $g \in G$, there exists $g_1 \in G$ such that $g_{1|(-\infty,0]} = g_{|(-\infty,0]}$, $supp(g_1) \subseteq (-\infty,n_o]$, and $order(g_{1|[1,n_o]})$ divides $order(g_{|[1,n_o]})$ (we assume that n_o is the least natural number satisfying this property). Again, this implies the existence of $g_2 \in G$ such that $g = g_1 + g_2$, $supp(g_2) \subseteq [1, +\infty[$, and $order(g_2)$ divides order(g). Here, the order of g is taken in the usual sense, as an element of the group G.

We now state our main result.

Theorem 1.3. Let G be an order controllable group shift over a finite abelian group H. Then there is a noncatastrophic isomorphic encoder for G. As a consequence G is algebraically and topologically conjugate to a full group shift.

2. Group shifts

In this section, we apply the result accomplished in [6, Theorem 3.2] in order to prove that order controllable group shifts over a finite abelian group possess canonical generating sets. Furthermore, our construction also yields that they are algebraically conjugate to a full group shift.

In the sequel $H^{(\mathbb{Z})}$ will denote the subgroup of $H^{\mathbb{Z}}$ consisting of all elements with finite support.

Theorem 2.1. Let G be a weakly controllable, group shift over a finite abelian p-group H. If G[p] is weakly controllable, then there is a finite generating subset $B_0 := \{x_j : 1 \le j \le m\} \subseteq G_{f[0,\infty)}[p]$, where $x_j = p^{h_j}y_j$, $y_j \in G_f$, and each x_j is selected with the maximal possible height h_j in G_f with $h_j \ge h_{j+1}$, $1 \le j < m$, such that the following assertions hold true:

(1) There is a canonically defined σ -invariant, onto, group homomorphism

$$\Phi \colon \left(\prod_{1 \le j \le m} \mathbb{Z}(p^{h_j+1})\right)^{\mathbb{Z}} \to G.$$

(2) ((G is weakly rectangular and)) Φ is a noncatastrophic, isomorphic encoder for G if there is a finite block $[0, N] \subseteq \mathbb{N}$ such that the set

$$\{\sigma^n[x_j]_{|[0,N]} \neq 0 : n \in \mathbb{Z}, 1 \leq j \leq m\}$$

is linearly independent.

Proof. (1) Using that G and G[p] are weakly controllable, we can proceed as in [6, Theorem 3.2] in order to define a subset $B_0 := \{x_1, \ldots, x_m\} \subseteq G_f[p]_{[0,\infty)}$ such that $\pi_{[0]}(B_0)$ forms a basis of $\pi_{[0]}(G_{[0,\infty)}[p])$ and for each $x_j \in B_0$ there is a nonnegative integer h_j and an element $y_j \in G_f$ such that $x_j = p^{h_j}y_j$, where each x_j has the maximal possible height h_j in G_f and $h_1 \geq h_2 \geq \cdots \geq h_m$. Now define

$$\varphi_0 \colon \mathbb{Z}(p)^m \to G[p]$$

by

$$\varphi_0[(\lambda_1,\ldots,\lambda_m)] = \lambda_1 x_1 + \cdots + \lambda_m x_m$$

and, for each $n \in \mathbb{Z}$, n > 0, set $B_n := \rho^n(B_0) \subseteq G_f[p]_{[n,\infty)}$ and define

$$\varphi_n \colon Z(p)^m \to G[p]$$

by

$$\varphi_n[(\lambda_1,\ldots,\lambda_m)] = \lambda_1 \rho^n(x_1) + \cdots + \lambda_m \rho^n(x_m).$$

Now, we can define

$$\bigoplus_n \varphi_n : \bigoplus_{n \geq 0} (\mathbb{Z}(p)^m)_n \to G_f[p]_{[0,\infty)}$$

by

$$\bigoplus_{n} \varphi_n[\sum_{n\geq 0} (\lambda_{1n}, \lambda_{2n}, \dots, \lambda_{mn})] := \sum_{n\geq 0} \varphi_n[(\lambda_{1n}, \lambda_{2n}, \dots, \lambda_{mn})],$$

where $(\mathbb{Z}(p)^m)_n = \mathbb{Z}(p)^m$ for all $n \geq 0$.

Remark that all the maps set above are well defined group homomorphisms since each of these maps involves finite sums in its definition. Furthermore, since the range of φ_n is contained in $G_f[p]_{[n,\infty)}$ for all $n \geq 0$, it follows that the map $\bigoplus_n \varphi_n$ is continuous when its domain (and its range) is equipped with the product topology. Therefore, there is a canonical extension of $\bigoplus_n \varphi_n$ to a continuous group homomorphism

$$\Phi_0: \prod_{n>0} (\mathbb{Z}(p)^m)_n \to G[p]_{[0,\infty)}.$$

Now, repeating the same arguments as in [6, Theorem 3.2], it follows that

$$G_f[p]_{[0,\infty)} \subseteq \Phi_0(\prod_{n>0} (\mathbb{Z}(p)^m)_n),$$

which implies that Φ_0 is a continuous onto group homomorphism because $G_f[p]_{[0,\infty)}$ is dense in $G[p]_{[0,\infty)}$. Furthermore, using the σ -invariance of G, we can extend Φ_0 canonically to continuous onto group homomorphism

$$\Phi_N: \prod_{n\geq -N} (\mathbb{Z}(p)^m)_n \longrightarrow G[p]_{[-N,\infty)}$$

by

$$\Phi_N[\sum_{n>-N}(\lambda_{1n},\lambda_{2n},\ldots,\lambda_{mn})] := \sigma^N[\Phi_0[\rho^N[\sum_{n>-N}(\lambda_{1n},\lambda_{2n},\ldots,\lambda_{mn})]]],$$

for every N > 0. Now, if we identify $\prod_{n \geq -N} (\mathbb{Z}(p)^m)_n$ with the subgroup $(\prod_{n \in \mathbb{Z}} (\mathbb{Z}(p)^m)_n)_{[-N,+\infty)}$, remark that $\Phi_{(N+1)}$ restricted to $\prod_{n \geq -N} (\mathbb{Z}(p)^m)_n$ is equal to Φ_N . Therefore, we have defined a map

$$\Phi_{\infty} : \bigcup_{N>0} \prod_{n>-N} (\mathbb{Z}(p)^m)_n \to G[p].$$

Again, because $\bigcup_{N>0} \prod_{n\geq -N} (\mathbb{Z}(p)^m)_n$ is dense in $(\mathbb{Z}(p)^m)^{\mathbb{Z}}$, it follows that we can extend Φ_{∞} to a continuous onto group homomorphism

$$\Phi: (\mathbb{Z}(p)^m)^{\mathbb{Z}} \longrightarrow G[p].$$

Now, taking into account that $\lim_{n\to\pm\infty}\sigma^n(y_j)=0$ for all $1\leq j\leq m$, we proceed as in [6, Theorem 3.2] in order to lift Φ to a continuous onto group homomorphism

$$\Phi \colon \left(\prod_{1 \le j \le m} \mathbb{Z}(p^{h_j+1})\right)^{\mathbb{Z}} \to G.$$

This completes the proof of (1).

(2) First, we remark that repeating the proof accomplished in [6, Theorem 3.2], it follows that the sets $\{\sigma^n[x_j]: n \in \mathbb{Z}, 1 \leq j \leq m\}$ and $\{\sigma^n[y_j]: n \in \mathbb{Z}, 1 \leq j \leq m\}$ are both (linearly) independent.

Furthermore, since all elements x_j $(1 \le j \le m)$ have finite support, it follows that the set $\{\sigma^n[x_j]_{|[0,N]} \ne 0 : n \in \mathbb{Z}, 1 \le j \le m\}$ is finite. Thus, using the σ -invarince of G, we proceed as in [6, Theorem 3.2] to obtain that Φ is one-to-one.

In order to prove that Φ is noncatastrophic, that is $\Phi[(\prod_{1 \leq j \leq m} \mathbb{Z}(p^{h_j+1})^{(\mathbb{Z})}] \subseteq G_f$, first notice that Φ^{-1} is continuous, being the inverse map a continuous one-to-one group homomorphism. Now, reasoning by contradiction, suppose there is $w \in G_f$ such that $(\vec{\lambda_n}) = \Phi^{-1}(w)$ is an infinite sequence, let us say, without loss of generality, an infinite sequence on the right side. Then we have that the sequence $(\sigma^n(w))_{n>0}$ converges to 0 in G. However, since $(\vec{\lambda_n})$ is infinite on the right side, it follows that the sequence $(\Phi^{-1}(\sigma^n(w)))_{n>0} = (\sigma^n[(\vec{\lambda_n})])_{n>0}$ does not converge to 0 in $(\prod_{1 \leq j \leq m} \mathbb{Z}(p^{(h_j+1)}))^{\mathbb{Z}}$. This contradiction completes the proof.

Definition 2.2. In the sequel, a set $\{y_1, \ldots, y_m\}$ (resp. $\{x_1, \ldots, x_m\}$) that satisfies the properties established in Theorem 2.1 is called *topological generating set* of G (resp. G[p]).

Next, we are going to use the preceding results in order to characterize the existence of noncatastrophic, isomorphic encoders. As a consequence, we also characterize when a group shift is algebraically conjugate to a full group shift. First we need the following notions.

Definition 2.3. A group shift $G \subseteq X = H^{\mathbb{Z}}$ is a shift of *finite type* (equivalently, is an *observable* group code) if it is defined by forbidding the appearance a finite list of (finite) blocks. As a consequence, there is $N \in \mathbb{N}$ such that if x_1, x_2 belong to G and they coincide on an N-block $[k, \ldots, k+N]$, then there is $x \in G$ such that $x_{|(-\infty,k+N]} = x_{1|(-\infty,k+N]}$ and $x_{|[k,\infty)} = x_{2|[k,\infty)}$. It is known that if G is an irreducible group shift over a finite group H, then G is also a group shift of finite type (see [11, Prop. 4]). Moreover, since every order controllable group shift G is irreducible, it follows that order controllable group shifts are of finite type.

Given an element $x \in G_f$ with $\operatorname{supp}(x) = \{i \in \mathbb{Z} : x(i) \neq 0\}$, the first index (resp. last index) $i \in \operatorname{supp}(x)$ is denoted by $i_f(x)$ (resp. $i_l(x)$). The lenth of $\operatorname{supp}(x)$ is defined as $|\operatorname{supp}(x)| := i_l(x) - i_f(x) + 1$.

Proposition 2.4. Let G be a weakly controllable, group shift of finite type over a finite abelian p-group H. If exp(H) = p, then there is a noncatastrophic isomorphic encoder for G. As a consequence G is algebraically and topologically conjugate to a full group shift.

Proof. First, remark that G = G[p] in this case. By Theorem 2.1, there is a topological generating subset $\mathcal{B}_0 := \{x_j : 1 \leq j \leq m\} \subseteq G_{f[0,\infty]}[p] = G_{f[0,\infty]}$ such that $\pi_{[0]}(\mathcal{B}_0)$ forms a basis of $\pi_{[0]}(G_{[0,\infty)})$ and there is a canonically defined σ -invariant, onto, group homomorphism

$$\Phi \colon (\mathbb{Z}(p)^m)^{\mathbb{Z}} \to G.$$

Furthermore, we select each element x_j with minimal support in $G_{f[0,\infty)}$ and such that $|\operatorname{supp}(x_1)| \leq \cdots \leq |\operatorname{supp}(x_m)|$.

By Theorem 2.1 (2), it will suffice to verify that there is a finite block $[0, N] \subseteq \mathbb{N}$ such that the set $\{\sigma^n[x_j]_{|[0,N]} \neq 0 : n \in \mathbb{Z}, 1 \leq j \leq m\}$ is linearly independent. Indeed, let N be a natural number such that $\operatorname{supp}(x_j) \subseteq [0, N]$ for all $1 \leq j \leq m$ and satisfying the condition of being a group shift of finite type for G. That is, if ω_1, ω_2 belong to G and they coincide on any N-block $[k, \ldots, k+N]$, then there is $w \in G$ such that $w_{|(-\infty,k+N]} = w_{1|(-\infty,k+N]}$ and $w_{|[k,\infty)} = w_{2|[k,\infty)}$.

Reasoning by contradiction, let us suppose that there is a linear combination

$$\sum \lambda_{nj} \sigma^n(x_j)_{|[0,N]} = 0.$$

Since the set $\{\sigma^n[x_j] : n \in \mathbb{Z}, 1 \leq j \leq m\}$ is linearly independent, there must be an element $u = \sigma^{n_1}[x_{j_1}]$ (for some n_1 and j_1) such that

$$\operatorname{supp}(u) \cap (-\infty, 0) \neq \emptyset.$$

As a consequence, there exist $\{\alpha_{nj}\}\subseteq \mathbb{Z}(p)$ such that

$$u_{|[0,N]} = \sum_{(n \neq n_1, j \neq j_1)} \alpha_{nj} \sigma^n(x_j)_{|[0,N]}.$$

We select u such that $i_f(u)$ is minimal among the elements satisfying this property. Set

$$v := \sum_{(n \neq n_1, j \neq j_1)} \alpha_{nj} \sigma^n(x_j).$$

We have that

$$(u-v)_{|[0,N]}=0.$$

Since G is of finite type for N-blocks, there exists $w \in G$ such that

$$w_{|(-\infty,N]} = (u-v)_{|(-\infty,N]}$$
 and $w_{|[0,\infty)} = 0$.

We have that $i_f(u) \leq i_f(w)$ and $i_l(w) < i_l(u)$. Therefore, we have found an element $w \in G_f$ with |supp(w)| < |supp(u)|. Therefore, we can replace x_{j_1} by $\widetilde{x}_{j_1} := \sigma^{-n_1}(w)$ and $|\text{supp}(\widetilde{x}_{j_1})| < |\text{supp}(x_{j_1})|$. This is a contradiction with our previous selection of the (ordered) set $\{x_j : 1 \leq j \leq m\}$, which completes the proof.

Lemma 2.5. Let G be an order controllable group shift over a finite abelian p-group H. Then G[p] and p^rG are order controllable group shifts for all r with $p^r < exp(H)$. As a consequence, it holds that $(p^rG)_f = p^rG_f$ for all r with $p^r < exp(H)$.

Proof. It is obvious that G[p] is order controllable. Regarding the group p^rG , take an arbitrary element $x = p^ry \in p^rG$. By the order controllability of G, there is $z \in G$ and $n_0 \in \mathbb{N}$ such that $y_{|(-\infty,0]} = z_{|(-\infty,0]}$, $\operatorname{supp}(z) \subseteq (-\infty,n_0]$ and $\operatorname{order}(z_{|[1,n_0]})$ divides $\operatorname{order}(y_{|[1,n_0]})$. Then $p^rz \in p^rG$, $x_{|(-\infty,0]} = p^rz_{|(-\infty,0]}$, $\operatorname{supp}(p^rz) \subseteq (-\infty,n_0]$ and $\operatorname{order}(p^rz_{|[1,n_0]})$ divides $\operatorname{order}(x_{|[1,n_0]})$.

Finally, it is clear that $p^rG_f \subseteq (p^rG)_f$. Next we check the reverse implication.

Let $y \in G$ such that $x = p^r y \in (p^r G)_f$. Then there are two integers m, M such that $x \in G_{[m,M]}$. Assume that $M \geq 0$ without loss of generality. By order controllability,

there is $z \in G$ such that $\sigma^M(y)_{|(-\infty,0]} = z_{|(-\infty,0]}$, $\operatorname{supp}(z) \subseteq (-\infty,n_0]$ and $\operatorname{order}(z_{|[1,n_0]})$ divides $\operatorname{order}(\sigma^M(y)_{|[1,n_0]})$. Hence, if $v = \sigma^{-M}(z)$, we have $y_{|(-\infty,M]} = v_{|(-\infty,M]}$, $\operatorname{supp}(v) \subseteq (-\infty, M+n_0]$ and $\operatorname{order}(v_{|[M+1,M+n_0]})$ divides $\operatorname{order}(y_{|[M+1,M+n_0]})$. Therefore $x = p^r v$ with $v \in G_{(-\infty,M+n_0]}$.

If $m - n_o > 0$, by order controllability, there is $u \in G$ such that $v_{|(-\infty,0]} = u_{|(-\infty,0]}$, $\sup_{v \in G} u(v) \subseteq (-\infty, n_0] \subseteq (-\infty, m-1]$ and $\operatorname{order}(u_{|[1,n_0]})$ divides $\operatorname{order}(v_{|[1,n_0]})$. Set w = v - u. We have that $w \in G_{[1,M+n_o]}$ and $x = p^r w$, which yields $x \in p^r G_f$.

If $m - n_o \leq 0$, set $N = m - n_o - 1$. By order controllability, there is $u_1 \in G$ such that $\sigma^N(v)_{|(-\infty,0]} = u_{1|(-\infty,0]}$, $\operatorname{supp}(u_1) \subseteq (-\infty,n_0]$ and $\operatorname{order}(u_{1|[1,n_0]})$ divides $\operatorname{order}(\sigma^N(v)_{|[1,n_0]})$. Hence, if $u_2 = \sigma^{-N}(u_1)$, we have $v_{|(-\infty,N]} = u_{2|(-\infty,N]}$, $\operatorname{supp}(u_2) \subseteq (-\infty, N + n_0] \subseteq (-\infty, m - 1]$ and $\operatorname{order}(u_{2|[N+1,N+n_0]})$ divides $\operatorname{order}(v_{|[N+1,N+n_0]})$. Set $w = v - u_2$. We have that $w \in G_{[N+1,M+n_o]}$ and $x = p^r w$, which agin yields $x \in p^r G_f$. This completes the proof.

Let G be a group shift over a finite abelian p-group H and let G/pG denote the quotient group defined by the map $\pi: G \to G/pG$. We define the subgroup

$$(G/pG)_f := \{\pi(u) : u \in G \text{ and } u(n) \in pH \text{ for all but finitely many } n \in \mathbb{Z}.\}$$

Lemma 2.6. Let G be an order controllable group shift over a finite abelian p-group H and let $\{x_1, \ldots, x_m\} \subseteq (pG_f)_{[0,\infty)}$ be a topological generating set of pG, where $x_i = py_i$, $y_i \in G_f$, $1 \le i \le m$. If $u \in G_f$ then there exist $v \in G_f[p]$ and $w \in \langle \{\sigma^n(y_j) : n \in \mathbb{Z}, 1 \le j \le m\} \rangle$ such that u = v + w.

Proof. Since $\{x_1, \ldots, x_m\}$ is a topological generating set of pG, we have

$$pu = \sum_{n \in \mathbb{Z}} \sum_{i=1}^{m} \lambda_{in} \sigma^{n}(x_{i}) = \sum_{n \in \mathbb{Z}} \sum_{i=1}^{m} \lambda_{in} p \sigma^{n}(y_{i}) = p \sum_{n \in \mathbb{Z}} \sum_{i=1}^{m} \lambda_{in} \sigma^{n}(y_{i}).$$

Furthermore, since the group shift pG is of finite type and $(pG)_f = p(G_f)$ by Lemma 2.5, we can apply Proposition 2.4 to the group shift pG, in order to obtain that the sum in the equality above only involves non-null terms for a finite subset of indices $F \subseteq \mathbb{Z}$. Therefore

$$pu = p \sum_{n \in F} \sum_{i=1}^{m} \lambda_{in} \sigma^{n}(y_{i}).$$

Set

$$w := \sum_{n \in F} \sum_{i=1}^{m} \lambda_{in} \sigma^{n}(y_{i}) \in G_{f}.$$

Then

$$u = w + (u - w),$$

where $w \in \langle \{\sigma^n(y_j) : n \in \mathbb{Z}, 1 \leq j \leq m\} \rangle$ and p(u-w) = 0. It now suffices to take v := u - w.

Theorem 2.7. Let G be an order controllable group shift (therefore, of finite type) over a finite abelian p-group H. Then there is a noncatastrophic isomorphic encoder for G. As a consequence G is algebraically and topologically conjugate to a full group shift.

Proof. Using induction on the exponent of G, we will prove that there is topological generating set B_0 of G[p], where $B_0 := \{x_1, \ldots, x_m\} \subseteq (pG_f[p])_{[0,\infty)}$ such that $\pi_{[0]}(B_0)$ forms a basis of $\pi_{[0]}((pG[p])_{[0,\infty)})$ and for each $x_j \in B_0$ there is an element $y_j \in G_f$ such that $x_j = p^{h_j}y_j$. Furthermore G is algebraically conjugate to the full group shift generated by $\mathbb{Z}(p^{h_1}) \times \ldots \mathbb{Z}(p^{h_m})$.

The case exp(G) = p has already be done in Proposition 2.4. Now, suppose that the proof has been accomplished if $exp(G) = p^h$ and let us verify it for $exp(G) = p^{h+1}$. We proceed as follows:

First, take the closed, shift invariant, subgroup pG. We have that $exp(pG) = p^h$ and by the induction hypothesis, there is topological generating set B_0 of pG[p], where $B_0 := \{x_1, \ldots, x_m\} \subseteq (pG_f[p])_{[0,\infty)}$ such that $\pi_{[0]}(B_0)$ forms a basis of $\pi_{[0]}((pG[p])_{[0,\infty)})$ and for each $x_j \in B_0$ there is an element $y_j \in pG_f$ such that $x_j = p^{h_j}y_j$.

Since $y_j \in pG_f$, there is $z_j \in G_f$ such that $y_j = pz_j$, $1 \le j \le m$. Furthermore, we may assume that there is a finite block $[0, N_1] \subseteq \mathbb{N}$ such that the set $\{\sigma^n[y_j]_{|[0,N_1]} \ne 0 : n \in \mathbb{Z}, 1 \le j \le m\}$ is linearly independent. As a consequence, using similar arguments as in [6, Theorem 3.2], it follows that the set $\{\sigma^n[z_j]_{|[0,N_1]} \ne 0 : n \in \mathbb{Z}, 1 \le j \le m\}$ also is linearly independent. Therefore there is a canonically defined σ -invariant, onto, group homomorphism

$$\Phi \colon \left(\prod_{1 \le j \le m} \mathbb{Z}(p^{h_j})\right)^{\mathbb{Z}} \to pG.$$

Now, we complete the set $B_0 := \{x_1, \ldots, x_m\} \subseteq (pG)[p]_{f[0,\infty)}$ with a finite set $B_1 := \{u_1, \ldots, u_k\} \subseteq G[p]_{f[0,\infty)}$ such that $\pi_{[0]}(B_0 \cup B_1)$ is a basis of $\pi_{[0]}(G[p])$. Remark that we must have $h(u_i) = 0$ for all $1 \le i \le k$, since $\pi_{[0]}(B_0)$ forms a basis of $\pi_{[0]}(pG[p])$. Furthermore, arguing as in Proposition 2.4, we may assume that there is a finite block $[0, N_2] \subseteq \mathbb{N}$ such that the set

$$E := \{ \sigma^n[u_i]_{|[0,N_2]} : \sigma^n[u_i]_{|[0,N_2]} \neq 0 : n \in \mathbb{Z}, 1 \le i \le k \}$$

is an independent subset of $G[p]_{|[0,N_2]}$.

Now, consider the quotient group homomorphism

$$q: G \to G/pG$$

and remark that G/pG is a group shift over $(H/pH)^{\mathbb{Z}}$. Making use of this quotient map, we select a basis

$$V_1 := \{v_1, \dots, v_k\} \subseteq G_f[p]_{[0, +\infty)}$$

satisfying the following properties:

- (1) $V_{1|[0,N_2]} \subseteq \langle \{\sigma^n[u_i]_{|[0,N_2]} : \sigma^n[v_i]_{|[0,N_2]} \neq 0 : n \in \mathbb{Z}, 1 \leq i \leq k\} \rangle$.
- (2) $\pi_{[0]}(B_0 \cup V_1)$ is a basis of $\pi_{[0]}(G[p])$.
- (3) The set

$$\{\sigma^n[v_i]_{|[0,N_2]}: \sigma^n[v_i]_{|[0,N_2]} \neq 0: n \in \mathbb{Z}, 1 \leq i \leq k\}$$

is independent.

(4) Each $q(v_i)$ has the minimal possible support in $(G/pG)_f$. That is

$$|\operatorname{supp}(q(v_1))| \le \cdots \le |\operatorname{supp}(q(v_k))|$$

where, if
$$supp(q(v_i)) = \{\ldots, l_1, \ldots, l_{p_i}\}$$
, then $|supp(q(v_i))| := l_{p_i} - l_1 + 1$.

It is straightforward to verify that $q(G_f) \subseteq (G/pG)_f$ and, as a consequence, it follows that the group G/pG is controllable and its controllability index is less than or equal to the controllability index of G. As in Theorem 2.1, the topological generating set $\{v_1, \ldots, v_k\} \cup \{z_1, \ldots, z_m\}$ defines a continuous onto group homomorphism

$$\Phi: \left(\mathbb{Z}(p)^k \times \prod_{1 \le j \le m} (\mathbb{Z}_{p^{h_m+1}})^{\mathbb{Z}} \longrightarrow G\right)$$

By Theorem 2.1, in order to proof that Φ is one-to-one, it will suffice to find some block $[0, N] \in \mathbb{Z}$ such that

$$S := \left(\{ \sigma^s[v_i]_{|[0,N]} \neq 0 : s \in \mathbb{Z}, 1 \le i \le k \} \cup \{ \sigma^n[z_j]_{|[0,N_1]} \neq 0 : n \in \mathbb{Z}, 1 \le j \le m \} \right)_{|[0,N]}$$

forms and independent subset of $G_{|[0,N]}$.

Since that this property holds separately for $\{z_1, \ldots, z_m\}$ on the block $[0, N_1]$ and $\{v_1, \ldots, v_k\}$ on the block $[0, N_2]$, it will suffice to verify that if we denote by Y to the group shift generated by $\{z_1, \ldots, z_m\}$ and by U to the group shift generated by $\{v_1, \ldots, v_k\}$, then there is an block $[0, N] \subseteq \mathbb{Z}$ such that

$$(Y \cap U)_{|[0,N]} = \{0\}.$$

Since this implies that $S_{[[0,N]}$ is an independet subset.

Indeed, take $N \ge \max(2N_1, 2N_2)$. Then, reasoning by contradiction, assume we have a sum

$$(\sum \alpha_{in}\sigma^n(v_i) + \sum \beta_{js}\sigma^s(z_j))_{|[0,N]} = \{0\}.$$

Remark that we may assume that this sum is finite without loss of generality since G is order controllable. Then

$$p(\sum \alpha_{in}\sigma^n(v_i) + \sum \beta_{js}\sigma^s(z_j))_{|[0,N]} = (\sum p\alpha_{in}\sigma^n(v_i) + \sum p\beta_{js}\sigma^s(z_j))_{|[0,N]} = \{0\}$$
this yields

$$\sum p\beta_{js}\sigma^{s}(z_{j})_{|[0,N]} = \sum \beta_{js}\sigma^{s}(y_{j})_{|[0,N]} = \{0\}.$$

Since $N \geq N_1$, this implies that

$$\sum \beta_{js} \sigma^s(y_j) = \{0\}.$$

This mans that $\beta_{js} = p\gamma_{js}$ for every index js. Thus we have

$$(\sum \alpha_{in}\sigma^n(v_i) + \sum p\gamma_{js}\sigma^s(z_j))_{|[0,N]} = \{0\}.$$

Now, we select an element $\sigma^n(v_i)$ such that $i_f(q(\sigma^n(u_i)))$ is minimal among the elements satisfying this property. Suppose without loss of generality that $\sigma^n(v_i) = \sigma^{n_1}v_1$ for simplicity's sake. Solving for $\sigma^{n_1}v_1$ in the equality above, we have

$$\sigma^{n_1} v_{1|[0,N]} = \left(\sum_{n \neq n_1, i \neq 1} \alpha'_{in} \sigma^n(v_i) + \sum_{i} p \gamma'_{js} \sigma^s(z_j) \right)_{|[0,N]} = \{0\}.$$

16

Set

$$w := \sum_{n \neq n_1, i \neq 1} \alpha'_{in} \sigma^n(v_i)$$

and set

$$w_1 := \sigma^{n_1} v_1 - w.$$

Remark that $pw_1 = 0$, that is $w_1 \in G[p]$ and

$$w_{1|[0,N]} = \sum p \gamma'_{js} \sigma^s(y_j)_{|[0,N]} \in pH.$$

Therefore

$$\operatorname{supp}(q(w_1)) \cap [0, N] = \emptyset.$$

Since G is a group shift of finite type, there is $w_2 \in G$ such that

$$w_{2|(-\infty,N]} = w_{1|(-\infty,N]}$$
 and $w_{2|[0,+\infty)} = \sum p \gamma'_{js} \sigma^s(y_j)_{|[0,+\infty)}$.

From the way w_2 has been defined, we have that $\sigma^{-n_1}(w_2) \in G_f[p]_{[0,+\infty)}$ satisfies that

$$\sigma^{-n_1}(w_2)_{|[0,N_2]} \in \langle \{\sigma^n[v_i]_{|[0,N_2]} : \sigma^n[v_i]_{|[0,N_2]} \neq 0 : n \in \mathbb{Z}, 1 \le i \le k \} \rangle$$

and

$$|\operatorname{supp}(q(w_2))| \le |\operatorname{supp}(q(\sigma^{n_1}(v_1)))|.$$

This is a contradiction and completes the proof.

We can now proof Theorem 1.3.

Proof of Theorem 1.3. Since every finite abelian group is the direct sum of all its nontrivial p-subgroups, the proof follows from Theorem 2.7, in like manner as [6, Theorem A] follows from [6, Theorem 3.2].

QUESTION: Under what conditions is it possible to extend Theorem 1.3 to non-abelian groups?

References

- [1] A. R. Calderbank, A. Roger Hammons Jr., P. Vijay Kumar, N. J. A. Sloane, Patrick Solé, A linear construction for certain Kerdock and Preparata codes. Bulletin AMS, 29 (1993), 218-222.
- [2] F. Fagnani, Some results on the classification of expansive automorphisms of compact abelian groups, Ergodic Theory Dynam. Systems, 16, No. 5, (1996), 45–50.
- [3] F. Fagnani, Shifts on compact and discrete Lie groups: Algebraic topological invariants and classification problems, Adv. Math., vol. 127, pp. 283-306, 1997.
- [4] F. Fagnani, S. Zampieri, Dynamical Systems and Convolutional Codes Over Finite Abelian groups, IEEE Trans. Inform. Theory, vol. 42 (6), pp. 1892-1912, Nov. 1996.
- [5] M.V. Ferrer, S. Hernández, Subdirect products of finite abelian groups, In: Ferrando J. (eds) Descriptive Topology and Functional Analysis II. TFA 2018. Springer Proceedings in Mathematics and Statistics, vol 286. Springer, (2019), 89–101.
- [6] M.V. Ferrer, S. Hernández, Homomorphic encoders of profinite abelian groups I, Pending (2021),
- [7] M.V. Ferrer, S. Hernández, D. Shakhmatov, Subgroups of direct products closely approximated by direct sums, Forum Mathematicum, 29, No. 5, (2017), 1125-1144.
- [8] G. D. Forney, Convolutional codes I: Algebraic structure, IEEE Trans. Inform. Theory, vol. 16 (1970), 1491-1513.
- [9] G. D. Forney, Jr. and M. D. Trott, *The dynamics of group codes: State spaces, trellis diagrams and canonical encoders*, IEEE Trans. Inform. Theory, vol. 39 (1993), 1491-1513.
- [10] G. D. Forney, Jr. and M. D. Trott, The Dynamics of Group Codes: Dual Abelian Group Codes and Systems, Proc. IEEE Workshop on Coding, System Theory and Symbolic Dynamics (Mansfield, MA), pp. 35-65 (2004).
- [11] B. Kitchens, Expansive dynamics on zero-dimensional groups, Ergodic Theory and Dynamical Systems, 7, no. 2, (1987), 249–261.
- [12] B. P. Kitchens and K. Schmidt, Automorphisms of compact groups, Ergod. Th. Dynam. Sys. 9, (1989), 691–735.
- [13] H-A. Loeliger, T. Mittelholzer, Convolutional codes over groups, IEEE Trans. Inform. Theory, 42, no. 6, (1996), 1660–1686.
- [14] G. Miles, R. K. Thomas, The breakdown of automorphisms of compact topological groups, Academic Press (Studies in probability and ergodic theory), Adv. in Math. Suppl. Stud. vol 2 (1978), 207–218.
- [15] J. Rosenthal, S.M. Schumacher and E.V. York, On Behaviors and Convolutional Codes, IEEE Trans. on Information Theory, 42 (1996), no. 6, 1881–1891.
- [16] K. Schmidt, Automorphisms of compact abelian groups and affine varieties, Proc. London Math. Soc. **61** (1990), 480–496.

Universitat Jaume I, Instituto de Matemáticas de Castellón, Campus de Riu Sec, 12071 Castellón, Spain.

Email address: mferrer@mat.uji.es

Universitat Jaume I, INIT and Departamento de Matemáticas, Campus de Riu Sec, 12071 Castellón, Spain.

Email address: hernande@mat.uji.es